

## **7. REFERENCES**

Ainsworth M and Teokul W (2000). Breaking the silence: setting realistic priorities for AIDS control in less-developed countries. *Lancet* 356(9223): 55-60.

Alioum A, Dabis F, Dequae-Merchadou L, et al. (2001). Estimating the efficacy of interventions to prevent mother-to-child transmission of HIV in breast-feeding populations: development of a consensus methodology. *Stat Med* 20(23): 3539-56.

Alioum A, Cortina-Borja M, Dabis F, et al. (2003). Estimating the efficacy of interventions to prevent mother-to-child transmission of human immunodeficiency virus in breastfeeding populations: comparing statistical methods. *Am J Epidemiol* 158(6): 596-605.

Allen S, Meinzen-Derr J, Kautzman M, et al. (2003). Sexual behavior of HIV discordant couples after HIV counseling and testing. *AIDS* 17(5): 733-740.

Angelsen NK, Vik T, Jacobsen G and Bakketeig LS (2001). Breast feeding and cognitive development at age 1 and 5 years. *Arch Dis Child* 85(3): 183-8.

Anglaret X, Chene G, Attia A, et al. (1999). Early chemoprophylaxis with trimethoprim-sulphamethoxazole for HIV-1-infected adults in Abidjan, Cote d'Ivoire: a randomised trial. Cotrimo-CI Study Group. *Lancet* 353(9163): 1463-8.

Asamoah-Odei E, Garcia Calleja JM and Boerma JT (2004). HIV prevalence and trends in sub-Saharan Africa: no decline and large subregional differences. *Lancet* 364(9428): 35-40.

Bambra CS (1999). Current status of reproductive behaviour in Africa. *Hum Reprod Update* 5(1): 1-20.

Becquart P, Chomont N, Roques P, et al. (2002). Compartmentalization of HIV-1 between breast milk and blood of HIV-infected mothers. *Virology* 300(1): 109-17.

Becquet R, Bequet L, Ekouevi D, et al. (2003a). Mortality in breastfed and formula-fed children born to HIV-infected women in a PMTCT project in Abidjan (Côte d'Ivoire),

Ditrame Plus ANRS 1202. Oral communication N°63. The 2nd IAS Conference on HIV Pathogenesis and Treatment, Paris, France. *Antiviral Therapy* 8(suppl. 1): S200.

Becquet R, Bequet L, Sakarovitch C, et al. (2003b). Mortality in breastfed and formula-fed children born to HIV-infected women. Oral communication N°598285. The 13<sup>th</sup> International Conference on AIDS & STIs in Africa. Nairobi, Kenya.

Becquet R, Rouet F, Bequet L, et al. (2004). Transmission post-natale du VIH en fonction des modalités d'alimentation infantile dans un projet de prévention de la transmission mère-enfant du VIH à Abidjan, Côte d'Ivoire, 2001-2004. Communication orale N°B3-1. Congrès international de l'Association Des Epidémiologistes de Langue Française, Bordeaux, France. *Rev Epidemiol Sante Publique* 52: 1S23.

Becquet R, Castetbon K, Viho I, et al. (2005a). Infant feeding practices before implementing alternatives to prolonged breastfeeding to reduce HIV transmission through breastmilk in Abidjan, Cote d'Ivoire. *J Trop Pediatr* 51(6): 351-5.

Becquet R, Ekouevi DK, Sakarovitch C, et al. (2005b). Knowledge, attitudes, and beliefs of health care workers regarding alternatives to prolonged breast-feeding (ANRS 1201/1202, Ditrame Plus, Abidjan, Cote d'Ivoire). *J Acquir Immune Defic Syndr* 40(1): 102-105.

Becquet R, Ekouevi DK, Viho I, et al. (2005c). Acceptability of exclusive breastfeeding with early cessation to prevent HIV transmission through breastmilk, ANRS 1201/1202 Ditrame Plus, Abidjan, Côte d'Ivoire. *J Acquir Immune Defic Syndr* 40(5): 600-8.

Becquet R and Leroy V (2005d). HIV and infant feeding: a complex issue in resource-limited settings. *AIDS* 19(15): 1717-8.

Becquet R (in press-a). Pratiques contraceptives dans le post-partum immédiat en relation avec les modalités d'alimentation infantile chez des mères infectées par le VIH. Expérience du projet Ditrame Plus ANRS 1201/1202, Abidjan, Côte d'Ivoire. In: Sexualité et procréation confrontées au Sida dans les pays du Sud. Paris, France, Les collections du CEPED.

Becquet R, Leroy V, Ekouevi DK, et al. (in press-b). Complementary feeding adequacy in relation to nutritional status among early weaned breastfed children born to HIV-infected mothers, ANRS 1201/1202 Ditrane Plus, Abidjan, Côte d'Ivoire. *Pediatrics*.

Bedikou G, Viho I, Tonwe-Gold B, et al. (2005). 6-month immunological response with HAART containing Nevirapine in HIV infected women post exposure to single dose of Nevirapine for PMTCT. The MTCT-Plus Initiative in Abidjan, Côte d'Ivoire (2003-2005). Oral communication N°MoOa0203. *The 3<sup>rd</sup> IAS Conference on HIV pathogenesis and treatment*, Rio de Janeiro, Brazil.

Bland RM, Rollins NC, Coutsooudis A and Coovadia HM (2002). Breastfeeding practices in an area of high HIV prevalence in rural South Africa. *Acta Paediat* 91(6): 704-711.

Bland RM, Rollins NC, Solarsh G, Van den Broeck J and Coovadia HM (2003). Maternal recall of exclusive breast feeding duration. *Arch Dis Child* 88(9): 778-83.

Breastfeeding and HIV International Transmission Study Group (BHITS) (2004). Late Postnatal Transmission of HIV-1 in Breast-Fed Children: An Individual Patient Data Meta-Analysis. *J Infect Dis* 189(12): 2154-2166.

Breastfeeding and HIV International Transmission Study Group (BHITS) (2005). Mortality Among HIV-1-Infected Women According to Children's Feeding Modality: An Individual Patient Data Meta-Analysis. *J Acquir Immune Defic Syndr* 39(4): 430-438.

Bryce J, Boschi-Pinto C, Shibuya K and Black RE (2005). WHO estimates of the causes of death in children. *Lancet* 365(9465): 1147-52.

Burger H, Kovacs A, Weiser B, et al. (1997). Maternal serum vitamin A levels are not associated with mother-to-child transmission of HIV-1 in the United States. *J Acquir Immune Defic Syndr Hum Retrovirol* 14(4): 321-6.

Buve A, Carael M, Hayes RJ, et al. (2001). Multicentre study on factors determining differences in rate of spread of HIV in sub-Saharan Africa: methods and prevalence of HIV infection. *AIDS* 15 Suppl 4: S5-14.

Carael M, Van de Perre PH, Lepage PH, et al. (1988). Human immunodeficiency virus transmission among heterosexual couples in Central Africa. *Aids* 2(3): 201-5.

Chaix ML, Montcho C, Ekouevi DK, et al. (2004). Genotypic resistance analysis in women who received Intrapartum Nevirapine associated to a Short Course of Zidovudine to Prevent Perinatal HIV-1 Transmission: The Ditrane Plus ANRS 1201/02 Study, Abidjan, Côte d'Ivoire. Poster N°657. *The 11th Conference on Retroviruses and Opportunistic Infections*, San Francisco, USA.

Chaix ML, Dabis F, Ekouevi DK, et al. (2005). Three days of ZDV+3TC postpartum after a short antenatal course of ZDV+3TC and single-dose Nevirapine (sdNVP) provides low rate of NVP resistance mutations and high efficacy in preventing peripartum HIV-1 transmission: ANRS Ditrane Plus, Abidjan, Côte d'Ivoire. Oral N°74LB. *The 12th Conference on Retroviruses and Opportunistic Infections*, Boston, USA.

Chantry CJ, Morrison P, Panchula J, et al. (2000). Effects of lipolysis or heat treatment on HIV-1 provirus in breast milk. *J Acquir Immune Defic Syndr* 24(4): 325-9.

Chintu PC, Bhat G, Walker A, et al. (2004). Co-trimoxazole as prophylaxis against opportunistic infections in HIV-infected Zambian children (CHAP): a double-blind randomised placebo-controlled trial. *Lancet* 364(9448): 1865-71.

Cogill B (2003). Anthropometric indicators measurement guide. Washington D.C., USA, Food and Nutrition technical Assistance Project, Academy for Educational Development.

Cohen MS (2004). HIV and sexually transmitted diseases: lethal synergy. *Top HIV Med* 12(4): 104-7.

Coutinho SB, de Lira PI, de Carvalho Lima M and Ashworth A (2005). Comparison of the effect of two systems for the promotion of exclusive breastfeeding. *Lancet* 366(9491): 1094-100.

Coutsoudis A, Pillay K, Spooner E, Kuhn L and Coovadia HM (1999). Influence of infant-feeding patterns on early mother-to-child transmission of HIV-1 in Durban, South Africa: a prospective cohort study. South African Vitamin A Study Group. *Lancet* 354(9177): 471-6.

Coutsoudis A, Pillay K, Kuhn L, et al. (2001). Method of feeding and transmission of HIV-1 from mothers to children by 15 months of age: prospective cohort study from Durban, South Africa. *AIDS* 15(3): 379-87.

Coutsoudis A, Goga A, Rollins N and Coovadia H (2002). Free formula milk for infants of HIV-infected women: blessing or curse? *Health Policy Plan* 17(2): 154-160.

Dabis F, Msellati P, Meda N, et al. (1999). 6-month efficacy, tolerance, and acceptability of a short regimen of oral zidovudine to reduce vertical transmission of HIV in breastfed children in Cote d'Ivoire and Burkina Faso: a double-blind placebo-controlled multicentre trial. DITRAME Study Group. DIminution de la Transmission Mere-Enfant. *Lancet* 353(9155): 786-92.

Dabis F, Elenka N, Méda N, et al. (2001). 18-month mortality and perinatal exposure to zidovudine in West Africa. *AIDS* 15: 771-779.

Dabis F and Ekpini ER (2002). HIV-1/AIDS and maternal and child health in Africa. *Lancet* 359(9323): 2097-104.

Dabis F, Bequet L, Ekouevi DK, et al. (2005). Field efficacy of Zidovudine, Lamivudine and single-dose Nevirapine to prevent peripartum transmission of HIV. The ANRS 1201/1202 Ditrane Plus study, Abidjan, Cote d'Ivoire. *AIDS* 19(3): 309-18.

De Cock KM, Zadi F, Adjorolo G, et al. (1994). Retrospective study of maternal HIV-1 and HIV-2 infections and child survival in Abidjan, Cote d'Ivoire. *BMJ* 308(6926): 441-43.

De Cock KM, Fowler MG, Mercier E, et al. (2000). Prevention of mother-to-child HIV transmission in resource-poor countries: translating research into policy and practice. *Jama* 283(9): 1175-82.

Desgrées du Loû A, Msellati P, Yao A, et al. (1999). Impaired fertility in HIV-1 infected pregnant women: a clinic-based survey in Abidjan, Côte d'Ivoire, 1997. *AIDS* 13(4): 517-521.

Desgrées du Loû A (2004). Sida et fécondité. In: Santé de la reproduction au temps du Sida en Afrique. Paris, Les collections du CEPED: 154.

Dunne EF, Angoran-Benie H, Kamelan-Tano A, et al. (2001). Is drinking water in Abidjan, Cote d'Ivoire, safe for infant formula? *J Acquir Immune Defic Syndr* 28(4): 393-8.

Ekouevi DK, Becquet R, Bequet L, et al. (2004a). Obtaining informed consent in HIV-1 infected pregnant women participating in a PMTCT study in Abidjan, Côte d'Ivoire. *AIDS* 18(10): 1486-8.

Ekouevi DK, Leroy V, Viho A, et al. (2004b). Acceptability and uptake of a package to prevent mother-to-child transmission using rapid HIV testing in Abidjan, Cote d'Ivoire. *AIDS* 18(4): 697-700.

Ekouevi DK, Rouet F, Becquet R, et al. (2004c). Immune status and uptake of antiretroviral interventions to prevent mother-to-child transmission of HIV in Africa. *J Acquir Immune Defic Syndr* 36(2): 755-7.

Ekpini ER, Wiktor SZ, Satten GA, et al. (1997). Late postnatal mother-to-child transmission of HIV-1 in Abidjan, Cote d'Ivoire. *Lancet* 349(9058): 1054-1059.

Eshleman SH, Mracna M, Guay LA, et al. (2001). Selection and fading of resistance mutations in women and infants receiving nevirapine to prevent HIV-1 vertical transmission (HIVNET 012). *AIDS* 15(15): 1951-7.

Eshleman SH, Guay LA, Mwatha A, et al. (2004a). Characterization of nevirapine resistance mutations in women with subtype A vs. D HIV-1 6-8 weeks after single-dose nevirapine (HIVNET 012). *J Acquir Immune Defic Syndr* 35(2): 126-30.

Eshleman SH, Guay LA, Mwatha A, et al. (2004b). Comparison of nevirapine (NVP) resistance in Ugandan women 7 days vs. 6-8 weeks after single-dose nvp prophylaxis: HIVNET 012. *AIDS Res Hum Retroviruses* 20(6): 595-9.

EuroHIV (2005). HIV/AIDS Surveillance in Europe. End-year report 2004 - No. 71. Saint-Maurice, France, Institut national de la Veille Sanitaire.

European collaborative study (1999a). Elective caesarean-section versus vaginal delivery in prevention of vertical HIV-1 transmission: a randomised clinical trial. *Lancet* 353(9158): 1035-1039.

European collaborative study (1999b). Maternal viral load and vertical transmission of HIV-1: an important factor but not the only one. The European Collaborative Study. *AIDS* 13(11): 1377-85.

European collaborative study (2001). HIV-infected pregnant women and vertical transmission in Europe since 1986. European collaborative study. *AIDS* 15(6): 761-70.

European collaborative study (2005a). Mother-to-child transmission of HIV infection in the era of highly active antiretroviral therapy. *Clin Infect Dis* 40(3): 458-65.

European collaborative study (2005b). Does exposure to antiretroviral therapy affect growth in the first 18 months of life in uninfected children born to HIV-infected women? *J Acquir Immune Defic Syndr* 40(3): 364-370.

European HIV in obstetrics group (2004). Higher rates of post-partum complications in HIV-infected than in uninfected women irrespective of mode of delivery. *AIDS* 18(6): 933-938.

European Multicentre Study Group for Cabergoline in Lactation Inhibition (1991). Single dose cabergoline versus bromocriptine in inhibition of puerperal lactation: randomised, double blind, multicentre study. European Multicentre Study Group for Cabergoline in Lactation Inhibition. *Bmj* 302(6789): 1367-71.



Fassinou P, Elenga N, Rouet F, et al. (2004). Highly active antiretroviral therapies among HIV-1-infected children in Abidjan, Cote d'Ivoire. *AIDS* 18(14): 1905-13.

Fawzi W, Msamanga G, Spiegelman D, et al. (1998). Randomised trial of effects of vitamin supplements on pregnancy outcomes and T cell counts in HIV-1-infected women in Tanzania. *Lancet* 351(9114): 1477-1482.

Fawzi W, Msamanga G, Spiegelman D, et al. (2002a). Transmission of HIV-1 through breastfeeding among women in Dar es Salaam, Tanzania. *J Acquir Immune Defic Syndr* 31(3): 331-8.

Fawzi W, Msamanga GI, Hunter D, et al. (2002b). Randomized trial of vitamin supplements in relation to transmission of HIV-1 through breastfeeding and early child mortality. *AIDS* 16(14): 1935-1944.

Gaillard P, Piwoz E and Farley TM (2001). Collection of standardized information on infant feeding in the context of mother-to-child transmission of HIV. *Stat Med* 20(23): 3525-37.

Gaillard P, Fowler M, Dabis F, et al. (2004). Use of antiretroviral drugs to prevent HIV-1 transmission through breast-feeding: from animal studies to randomized clinical trials. *J Acquir Immune Defic Syndr* 35(2): 178-187.

Glynn JR, Carael M, Auvert B, et al. (2001). Why do young women have a much higher prevalence of HIV than young men? A study in Kisumu, Kenya and Ndola, Zambia. *AIDS* 15 Suppl 4: S51-60.

Gray RH, Li X, Kigozi G, et al. (2005). Increased risk of incident HIV during pregnancy in Rakai, Uganda: a prospective study. *The Lancet* 366(9492): 1182-1188.

Greenberg BL, Semba RD, Vink PE, et al. (1997). Vitamin A deficiency and maternal-infant transmissions of HIV in two metropolitan areas in the United States. *AIDS* 11(3): 325-32.

Guay LA, Musoke P, Fleming T, et al. (1999). Intrapartum and neonatal single-dose nevirapine compared with zidovudine for prevention of mother-to-child transmission of HIV-1 in Kampala, Uganda: HIVNET 012 randomised trial. *Lancet* 354(9181): 795-802.

Hamers FF and Downs AM (2004). The changing face of the HIV epidemic in western Europe: what are the implications for public health policies? *Lancet* 364(9428): 83-94.

Harmsen MC, Swart PJ, de Bethune MP, et al. (1995). Antiviral effects of plasma and milk proteins: lactoferrin shows potent activity against both human immunodeficiency virus and human cytomegalovirus replication in vitro. *J Infect Dis* 172(2): 380-8.

Iliff P, Piwoz E, Tavengwa N, et al. (2005). Early exclusive breastfeeding reduces the risk of postnatal HIV-1 transmission and increases HIV-free survival. *AIDS* 19(7): 699-708.

Institut National de la Statistique (Côte d'Ivoire) and ORC Macro (2001). Enquête démographique et de santé, Côte d'Ivoire 1998-1999., Calverton, Maryland USA: INS et ORC Macro: 298.

Israel-Ballard K, Chantry C, Dewey K, et al. (2005). Viral, Nutritional, and Bacterial Safety of Flash-Heated and Pretoria-Pasteurized Breast Milk to Prevent Mother-to-Child Transmission of HIV in Resource-Poor Countries: A Pilot Study. *J Acquir Immune Defic Syndr* 40(2): 175-181.

Jackson JB, Musoke P, Fleming T, et al. (2003). Intrapartum and neonatal single-dose nevirapine compared with zidovudine for prevention of mother-to-child transmission of HIV-1 in Kampala, Uganda: 18-month follow-up of the HIVNET 012 randomised trial. *Lancet* 362(9387): 859-868.

Jeffery BS and Mercer KG (2000). Pretoria pasteurisation: a potential method for the reduction of postnatal mother to child transmission of the human immunodeficiency virus. *J Trop Pediatr* 46(4): 219-23.

Jeffery BS, Webber L, Mokhondo KR and Erasmus D (2001). Determination of the effectiveness of inactivation of human immunodeficiency virus by Pretoria pasteurization. *J Trop Pediatr* 47(6): 345-9.

Jeffery BS, Soma-Pillay P, Makin J and Moolman G (2003). The effect of Pretoria Pasteurization on bacterial contamination of hand-expressed human breastmilk. *J Trop Pediatr* 49(4): 240-4.

John GC, Nduati RW, Mbori-Ngacha DA, et al. (2001). Correlates of mother-to-child human immunodeficiency virus type 1 (HIV-1) transmission: association with maternal plasma HIV-1 RNA load, genital HIV-1 DNA shedding, and breast infections. *J Infect Dis* 183(2): 206-212.

John-Stewart G, Mbori-Ngacha D, Ekpini R, et al. (2004). Breast-feeding and Transmission of HIV-1. *J Acquir Immune Defic Syndr* 35(2): 196-202.

Jourdain G, Ngo-Giang-Huong N, Le Coeur S, et al. (2004). Intrapartum exposure to nevirapine and subsequent maternal responses to nevirapine-based antiretroviral therapy. *N Engl J Med* 351(3): 229-40.

Kalter H, Gray R, Black R and Gultiano S (1990). Validation of postmortem interviews to ascertain selected causes of death in children. *Int. J. Epidemiol.* 19(2): 380-386.

Kramer MS, Guo T, Platt RW, et al. (2003). Infant growth and health outcomes associated with 3 compared with 6 mo of exclusive breastfeeding. *Am J Clin Nutr* 78(2): 291-295.

Kuhn L, Kasonde P, Sinkala M, et al. (2005a). Does severity of HIV disease in HIV-infected mothers affect mortality and morbidity among their uninfected infants? *Clin Infect Dis* 41(11): 1654-61.

Kuhn L, Kasonde P, Sinkala M, et al. (2005b). Prolonged breast-feeding and mortality up to two years post-partum among HIV-positive women in Zambia. *AIDS* 19(15): 1677-1681.

Kuhn L, Trabattoni D, Kankasa C, et al. (2005c). Alpha-Defensins in the Prevention of HIV Transmission Among Breastfed Infants. *J Acquir Immune Defic Syndr* 39(2): 138-142.

Lauer JA, Betran AP, Victora CG, de Onis M and Barros AJ (2004). Breastfeeding patterns and exposure to suboptimal breastfeeding among children in developing countries: review and analysis of nationally representative surveys. *BMC Med* 2: 26.

Lepage P, Msellati P, Hitimana DG, et al. (1996). Growth of human immunodeficiency type 1-infected and uninfected children: a prospective cohort study in Kigali, Rwanda, 1988 to 1993. *Pediatr Infect Dis J* 15(6): 479-85.

Lepage P, Spira R, Kalibala S, et al. (1998). Care of human immunodeficiency virus-infected children in developing countries. International Working Group on Mother-to-Child Transmission of HIV. *Pediatr Infect Dis J* 17(7): 581-6.

Leroy V, Newell ML, Dabis F, et al. (1998). International multicentre pooled analysis of late postnatal mother-to-child transmission of HIV-1 infection. Ghent International Working Group on Mother-to-Child Transmission of HIV. *Lancet* 352(9128): 597-600.

Leroy V, Montcho C, Manigart O, et al. (2001). Maternal plasma viral load, zidovudine and mother-to-child transmission of HIV-1 in Africa: DITRAME ANRS 049a trial. *AIDS* 15(4): 517-22.

Leroy V, Karon JM, Alioum A, et al. (2002). Twenty-four month efficacy of a maternal short-course zidovudine regimen to prevent mother-to-child transmission of HIV-1 in West Africa. *AIDS* 16(4): 631-41.

Leroy V, Karon JM, Alioum A, et al. (2003). Postnatal transmission of HIV-1 after a maternal short-course zidovudine peripartum regimen in West Africa. *AIDS* 17(10): 1493-1501.

Leroy V (2004a). L'inégalité des sexes face à l'infection par le VIH-sida en Afrique : Un cercle vicieux anthropologique, sociologique, épidémiologique et clinique, facteur d'entretien de l'épidémie. *Sciences Sociales et Santé* 22(3): 71-85.

Leroy V, Becquet R, Rouet F, et al. (2004b). Postnatal transmission risk according to feeding modalities in children born to HIV-infected mothers in a PMTCT project in Abidjan, Côte d'Ivoire. Ditrane Plus ANRS 1201/1202. Oral Poster N°MoPpB2007. *The XV International AIDS conference*, Bangkok, Thailand.

Leroy V, Sakarovitch C, Cortina-Borja M, et al. (2005). Is there a difference in the efficacy of peripartum antiretroviral regimens in reducing mother-to-child transmission of HIV in Africa? *AIDS* 19(16): 1865-75.

Leroy V (in press). La contraception face au VIH/SIDA dans les pays du Sud : enjeux médicaux, sociaux et comportementaux. In: Sexualité et procréation confrontées au Sida dans les pays du Sud. Paris, France, Les collections du CEPED.

Lewis JJ, Ronsmans C, Ezeh A and Gregson S (2004). The population impact of HIV on fertility in sub-Saharan Africa. *AIDS* 18 Suppl 2: S35-43.

Linkages project (1999). Recommended feeding and dietary practices to improve infant and maternal nutrition. USAID & Academy for Educational Development.

Linkages project (2002). La méthode de l'allaitement maternel et de l'aménorrhée (MAMA) : une option temporaire de contraception en post-partum pour les femmes qui allaitent. USAID & AED: 94.

Magoni M, Bassani L, Okong P, et al. (2005). Mode of infant feeding and HIV infection in children in a program for prevention of mother-to-child transmission in Uganda. *AIDS* 19(4): 433-7.

Manigart O, Crepin M, Leroy V, et al. (2004). Effect of perinatal zidovudine prophylaxis on the evolution of cell-free HIV-1 RNA in breast milk and on postnatal transmission. *J Infect Dis* 190(8): 1422-8.

Marston M, Zaba B, Salomon JA, Brahmbhatt H and Bagenda D (2005). Estimating the net effect of HIV on child mortality in African populations affected by generalized HIV epidemics. *J Acquir Immune Defic Syndr* 38(2): 219-27.

Martinson N, Morris L, Gray G, et al. (2004). HIV resistance and transmission following single-dose nevirapine in a PMTCT cohort. Oral communication N°38. *The 11th Conference on Retroviruses and Opportunistic Infections*, San Francisco, USA.

Mbori-Ngacha D, Nduati R, John G, et al. (2001). Morbidity and mortality in breastfed and formula-fed infants of HIV-1-infected women: A randomized clinical trial. *Jama* 286(19): 2413-20.

Méda N, Leroy V, Viho I, et al. (2002). Acceptability and field efficacy of a routine utilisation of zidovudine to reduce mother-to-child transmission of HIV-1 in West Africa. *AIDS* 16(17): 2323-2328.

Mitka M (2002). MTCT-Plus program has two goals: end maternal HIV transmission + treat mothers. *Jama* 288(2): 153-4.

Mohlala BK, Tucker TJ, Besser MJ, et al. (2005). Investigation of HIV in amniotic fluid from HIV-Infected pregnant women at full term. *J Infect Dis* 192(3): 488-91.

Mondot-Bernard J (1981). Fertility and breast-feeding in Africa. *Afr Environ*(14-16): 131-50.

Moodley D, Moodley J, Coovadia H, et al. (2003). A Multicenter Randomized Controlled Trial of Nevirapine Versus a Combination of Zidovudine and Lamivudine to Reduce Intrapartum and Early Postpartum Mother-to-Child Transmission of Human Immunodeficiency Virus Type 1. *J Infect Dis* 187(5): 725-35.

Msellati P, Newell ML and Dabis F (1995). Rates of mother-to-child transmission of HIV-1 in Africa, America and Europe: results from 13 perinatal studies. *J Acquir Immune Defic Syndr Hum Retrovirol* 8: 506-510.

Msellati P, Hingst G, Kaba F, et al. (2001). Operational issues in preventing mother-to-child transmission of HIV-1 in Abidjan, Cote d'Ivoire, 1998-99. *Bull World Health Organ* 79(7): 641-7.

MTCT-Plus initiative (2005). <http://www.mtctplus.org/>. Accédé le 5 décembre 2005.

Muro E, Droste JA, Hofstede HT, et al. (2005). Nevirapine plasma concentrations are still detectable after more than 2 weeks in the majority of women receiving single-dose Nevirapine: implications for intervention studies. *J Acquir Immune Defic Syndr* 39(4): 419-421.

Nagelkerke NJ, Moses S, Embree JE, Jenniskens F and Plummer FA (1995). The duration of breastfeeding by HIV-1-infected mothers in developing countries: balancing benefits and risks. *J Acquir Immune Defic Syndr Hum Retrovirol* 8(2): 176-81.

Nduati R, John G, Mbori-Ngacha D, et al. (2000). Effect of breastfeeding and formula feeding on transmission of HIV-1: a randomized clinical trial. *Jama* 283(9): 1167-74.

Nduati R, Richardson BA, John G, et al. (2001). Effect of breastfeeding on mortality among HIV-1 infected women: a randomised trial. *Lancet* 357(9269): 1651-5.

Newell ML, Brahmbhatt H and Ghys PD (2004a). Child mortality and HIV infection in Africa: a review. *AIDS* 18: S27-S34.

Newell ML, Coovadia H, Cortina-Borja M, et al. (2004b). Mortality of infected and uninfected infants born to HIV-infected mothers in Africa: a pooled analysis. *Lancet* 364(9441): 1236-43.

Nicolosi A, Correa Leite ML, Musicco M, et al. (1994). The efficiency of male-to-female and female-to-male sexual transmission of the human immunodeficiency virus: a study of 730 stable couples. Italian Study Group on HIV Heterosexual Transmission. *Epidemiology* 5(6): 570-5.

Obimbo EM, Mbori-Ngacha DA, Ochieng JO, et al. (2004). Predictors of early mortality in a cohort of human immunodeficiency virus type 1-infected african children. *Pediatr Infect Dis J* 23(6): 536-43.

Perez F, Mukotekwa T, Miller A, et al. (2004). Implementing a rural programme of prevention of mother-to-child transmission of HIV in Zimbabwe: first 18 months of experience. *Trop Med Int Health* 9(7): 774-83.

Petra study team (2002). Efficacy of three short-course regimens of zidovudine and lamivudine in preventing early and late transmission of HIV-1 from mother to child in Tanzania, South Africa, and Uganda (Petra study): a randomised, double-blind, placebo-controlled trial. *Lancet* 359(9313): 1178-86.

Piwoz EG, Creed de Kanashiro H, Lopez de Romana G, Black RE and Brown KH (1995). Potential for misclassification of infants' usual feeding practices using 24-hour dietary assessment methods. *J Nutr* 125(1): 57-65.

Preble E and Piwoz E (1998). HIV and infant feeding: a chronology of research and policy advances and their implications for programs. Academy for Educational Development, the LINKAGES & SARA projects. Washington DC: 28.

Rankin WW, Brennan S, Schell E, Laviwa J and Rankin SH (2005). The stigma of being HIV-positive in Africa. *PLoS Medicine* 2(8): e247.

Richardson BA, John-Stewart GC, Hughes JP, et al. (2003). Breast-milk infectivity in human immunodeficiency virus type 1-infected mothers. *J Infect Dis* 187(5): 736-40.

Rollins N, Meda N, Becquet R, et al. (2004). Preventing postnatal transmission of HIV-1 through breast-feeding: modifying infant feeding practices. *J Acquir Immune Defic Syndr* 35(2): 188-195.

Rollins N, Coovadia H, Bland R, et al. (2005). Pregnancy outcomes in HIV infected and uninfected women in rural and urban South Africa. Poster N°TuPp0401. *The 3<sup>rd</sup> IAS Conference on HIV Pathogenesis and Treatment*, Rio de Janeiro, Brazil.

Rouet F, Montcho C, Rouzioux C, et al. (2001). Early diagnosis of paediatric HIV-1 infection among African breast-fed children using a quantitative plasma HIV RNA assay. *AIDS* 15(14): 1849-56.



Rouet F, Ekouevi DK, Inwoley A, et al. (2004). Field evaluation of a rapid human immunodeficiency virus (HIV) serial serologic testing algorithm for diagnosis and differentiation of HIV type 1 (HIV-1), HIV-2, and dual HIV-1-HIV-2 infections in West African pregnant women. *J Clin Microbiol* 42(9): 4147-53.

Rouet F, Ekouevi DK, Chaix ML, et al. (2005). Transfer and Evaluation of an Automated, Low-Cost Real-Time Reverse Transcription-PCR Test for Diagnosis and Monitoring of Human Immunodeficiency Virus Type 1 Infection in a West African Resource-Limited Setting. *J Clin Microbiol* 43(6): 2709-17.

Rousseau CM, Nduati RW, Richardson BA, et al. (2003). Longitudinal analysis of human immunodeficiency virus type 1 RNA in breast milk and of its relationship to infant infection and maternal disease. *J Infect Dis* 187(5): 741-7.

Rousseau CM, Nduati RW, Richardson BA, et al. (2004). Association of levels of HIV-1-infected breast milk cells and risk of mother-to-child transmission. *J Infect Dis* 190(10): 1880-1888.

Rouzioux C, Costagliola D, Burgard M, et al. (1995). Estimated timing of mother-to-child human immunodeficiency virus type 1 (HIV-1) transmission by use of a Markov model. The HIV Infection in Newborns French Collaborative Study Group. *Am J Epidemiol* 142(12): 1330-7.

Royce RA, Sena A, Cates W, Jr. and Cohen MS (1997). Sexual transmission of HIV. *N Engl J Med* 336(15): 1072-8.

Sakarovitch C, Msellati P, Leroy V, et al. (2004). Incidence of HIV infection among women in Abidjan, Côte d'Ivoire : estimation from serial antenatal seroprevalence data. Oral communication N°ThOrC1422. *The XV International AIDS conference*, Bangkok, Thailand.

Schmid GP, Buve A, Mugenyi P, et al. (2004). Transmission of HIV-1 infection in sub-Saharan Africa and effect of elimination of unsafe injections. *Lancet* 363(9407): 482-488.

Semba RD, Miotti PG, Chipangwi JD, et al. (1994). Maternal vitamin A deficiency and mother-to-child transmission of HIV-1. *Lancet* 343(8913): 1593-7.

Semba RD, Kumwenda N, Hoover DR, et al. (1999). Human immunodeficiency virus load in breast milk, mastitis, and mother-to-child transmission of human immunodeficiency virus type 1. *J Infect Dis* 180(1): 93-8.

Semrau K, Kuhn L, Vwalika C, et al. (2005). Women in couples antenatal HIV counseling and testing are not more likely to report adverse social events. *AIDS* 19(6): 603-609.

Shapiro RL, Holland DT, Capparelli E, et al. (2005a). Antiretroviral concentrations in breast-feeding infants of women in Botswana receiving antiretroviral treatment. *J Infect Dis* 192(5): 720-7.

Shapiro RL, Ndung'u T, Lockman S, et al. (2005b). Highly Active Antiretroviral Therapy Started during Pregnancy or Postpartum Suppresses HIV-1 RNA, but Not DNA, in Breast Milk. *J Infect Dis* 192(5): 713-9.

Shapiro RL, Thior I, Lockman S, et al. (2005c). Breastfeeding with 6 Months of infant zidovudine prophylaxis vs. formula feeding for reducing HIV-1 transmission and infant mortality: a randomized trial in Southern Africa. *The 12th Conference on Retroviruses and Opportunistic Infections*, Boston, USA.

Songok EM, Fujiyama Y, Tukei PM, et al. (2003). The use of short-course zidovudine to prevent perinatal transmission of human immunodeficiency virus in rural Kenya. *Am J Trop Med Hyg* 69(1): 8-13.

Spira R, Lepage P, Msellati P, et al. (1999). Natural history of human immunodeficiency virus type 1 infection in children: a five-year prospective study in Rwanda. Mother-to-Child HIV-1 Transmission Study Group. *Pediatrics* 104(5): e56.

Stringer JS, Sinkala M, Maclean CC, et al. (2005). Effectiveness of a city-wide program to prevent mother-to-child HIV transmission in Lusaka, Zambia. *AIDS* 19(12): 1309-15.

Taha TE, Kumwenda NI, Gibbons A, et al. (2003). Short postexposure prophylaxis in newborn babies to reduce mother-to-child transmission of HIV-1: NVAZ randomised clinical trial. *Lancet* 362(9391): 1171-1177.

Temmerman M, Quaghebeur A, Mwanyumba F and Mandaliya K (2003). Mother-to-child HIV transmission in resource poor settings: how to improve coverage? *AIDS* 17(8): 1239-42.

Thea DM, Vwalika C, Kasonde P, et al. (2004). Issues in the design of a clinical trial with a behavioral intervention--the Zambia exclusive breast-feeding study. *Control Clin Trials* 25(4): 353-65.

Tonwe-Gold B, Ekouevi DK, Viho I, et al. (2005). HAART for the prevention of perinatal HIV transmission in West Africa. *Personal communication*.

Toure H, Tonwe-Gold B, Becquet R, et al. (2005). Adverse pregnancy outcomes in HIV-infected women treated with HAART in Abidjan Côte d'Ivoire. Oral communication number TuFo0202. *The 3<sup>rd</sup> IAS Conference on HIV pathogenesis and treatment*, Rio de Janeiro, Brazil.

Udall JN, Colony P, Fritze L, et al. (1981). Development of gastrointestinal mucosal barrier. II. The effect of natural versus artificial feeding on intestinal permeability to macromolecules. *Pediatr Res* 15(3): 245-9.

UNAIDS (2004a). AIDS epidemic update. Geneva, Switzerland, United Nations program on HIV/AIDS.

UNAIDS, WHO and UNICEF (2004b). Epidemiological fact sheets on HIV/AIDS and sexually transmitted infections - 2004 update. Geneva, Switzerland, United Nations program on HIV/AIDS, World Health Organisation & United Nations Children's Fund.

UNICEF (2000). Multiple Indicator Cluster Survey (MICS-2000), Côte d'Ivoire.

UNICEF, WHO and UNAIDS (2004). Les enfants au bord du gouffre en 2004, United Nations program on HIV/AIDS, World Health Organisation & United Nations Children's Fund.

Van de Perre P, Simonon A, Msellati P, et al. (1991). Postnatal transmission of human immunodeficiency virus type 1 from mother to infant. A prospective cohort study in Kigali, Rwanda. *N Engl J Med* 325(9): 593-8.

Van der Strate BW, Beljaars L, Molema G, Harmsen MC and Meijer DK (2001). Antiviral activities of lactoferrin. *Antiviral Res* 52(3): 225-39.

Van der Straten A, King R, Grinstead O, Serufilira A and Allen S (1995). Couple communication, sexual coercion and HIV risk reduction in Kigali, Rwanda. *Aids* 9(8): 935-44.

Verani A and Lusso P (2002). Chemokines as natural HIV antagonists. *Curr Mol Med* 2(8): 691-702.

Victora CG, Smith PG, Vaughan JP, et al. (1987). Evidence for protection by breast-feeding against infant deaths from infectious diseases in Brazil. *Lancet* 2(8554): 319-22.

Vyankandondera J, Luchters S, Hassink E, et al. (2003). Reducing risk of HIV-1 transmission from mother to infant through breastfeeding using antiretroviral prophylaxis in infants (SIMBA-study). Oral communication N°LB7. *The 2<sup>nd</sup> IAS Conference on HIV Pathogenesis and Treatment*, Paris, France.

Wahl SM, McNeely TB, Janoff EN, et al. (1997). Secretory leukocyte protease inhibitor (SLPI) in mucosal fluids inhibits HIV-I. *Oral Dis* 3 Suppl 1: S64-9.

Welffens-Ekra C, Desgrees-du-Lou A, Leroy V and Dabis F (2001). L'inégalité homme-femme face au SIDA. In: *Le SIDA au féminin*. Paris, Doin: 242.

WHO (1991). Indicators for assessing breast-feeding practices : report of an informal meeting. Geneva, Switzerland, World Health Organisation.

WHO and UNICEF (1993). Breastfeeding counselling, a training course. Geneva, Switzerland & New York, USA, world Health Organization & United Nations Children's Fund.

WHO, UNICEF and UNAIDS (1998a). HIV and infant feeding: a review of HIV transmission through breastmilk. Geneva, Switzerland, World Health Organisation, United Nations Children's Fund & United Nations program on HIV/AIDS.

WHO, UNICEF and UNAIDS (1998b). HIV and infant feeding: a guide for health care managers and supervisors. Geneva, Switzerland, World Health Organisation, United Nations Children's Fund & United Nations program on HIV/AIDS.

WHO (1999). Integration of vitamin A supplementation with immunization. *Wkly Epidemiol Rec* 74(1): 1-6.

WHO and UNAIDS (2000). WHO/UNAIDS hail consensus on use of cotrimoxazole for prevention of HIV-related infections in Africa. Geneva, Switzerland, World Health Organisation & United Nations program on HIV/AIDS.

WHO and UNICEF (2003). Global strategy for infant and young children feeding. Geneva, Switzerland, World Health Organisation & United Nations Children's Fund.

WHO (2004). Antiretroviral drugs for treating pregnant women and prevention HIV infection in infants: guidelines on care, treatment and support for women living with HIV/AIDS and their children in resource-constrained settings. WC/503.2. Geneva, World Health Organisation: 50.

WHO collaborative study team on the role of breastfeeding on the prevention of infant mortality (2000). Effect of breastfeeding on infant and child mortality due to infectious diseases in less developed countries: a pooled analysis. *Lancet* 355(9202): 451-455.

WHO Working Group on Infant Growth (1995). An evaluation of infant growth: the use and interpretation of anthropometry in infants. WHO Working Group on Infant Growth. *Bull World Health Organ* 73(2): 165-74.

Wiktor SZ, Ekpini E, Karon JM, et al. (1999). Short-course oral zidovudine for prevention of mother-to-child transmission of HIV-1 in Abidjan, Cote d'Ivoire: a randomised trial. *Lancet* 353(9155): 781-785.

Willumsen JF, Filteau SM, Coutsoadis A, et al. (2003). Breastmilk RNA viral load in HIV-infected South African women: effects of subclinical mastitis and infant feeding. *AIDS* 17(3): 407-14.