

Contribution à la résolution numérique des problèmes de Helmholtz

Magdalena Grigoroscuta-Strugaru

INRIA Magique-3D and LMA, University of Pau

Advisors:

Mohamed Amara (INRIA, LMA)

Henri Calandra (TOTAL)

Rabia Djellouli (CSUN, INRIA)



Outline

Outline

- Motivation and Context

Outline

- Motivation and Context
- Solution methodologies for Helmholtz problems

Outline

- Motivation and Context
- Solution methodologies for Helmholtz problems
 - A **new** DG method

Outline

- Motivation and Context
- Solution methodologies for Helmholtz problems
 - A **new** DG method
 - A **modified** DG method

Outline

- Motivation and Context
- Solution methodologies for Helmholtz problems
 - A **new** DG method
 - A **modified** DG method
 - An **improved** modified DG method

- Motivation and Context
- Solution methodologies for Helmholtz problems
 - A **new** DG method
 - A **modified** DG method
 - An **improved** modified DG method
- Summary and perspectives

Motivation and Context

Applications

Applications

- Geophysical exploration

Applications

- Geophysical exploration
- Radar

Applications

- Geophysical exploration
- Radar
- Sonar

Applications


- Geophysical exploration
- Radar
- Sonar
- Medical imaging

Applications

- Geophysical exploration
- Radar
- Sonar
- Medical imaging
- Nondestructive testing


Numerical Difficulties

Numerical Difficulties



$$\color{red}{k} \color{green}{a} = 1$$


Numerical Difficulties



$$ka = 1, \quad \frac{h}{a} = \frac{1}{10}$$


Motivation and Context

Numerical Difficulties




$$\textcolor{red}{k}a = 1, \quad \frac{h}{a} = \frac{1}{10}$$

Numerical Difficulties




$$ka = 1, \quad \frac{h}{a} = \frac{1}{10}$$

Numerical Difficulties




$$\textcolor{red}{k}a = 3$$

Numerical Difficulties




$$\textcolor{red}{k}a = 3, \quad \frac{h}{a} = \frac{1}{10}$$

Numerical Difficulties




$$ka = 3, \quad \frac{h}{a} = \frac{1}{20}$$

Numerical Difficulties




$$ka = 3, \quad \frac{h}{a} = \frac{1}{30}$$

Numerical Difficulties




$$ka = 3, \quad \frac{h}{a} = \frac{1}{30} \implies kh = \frac{1}{10}$$

Numerical Difficulties



Resolution necessary to achieve 10% on the relative error

Numerical Difficulties



Resolution necessary to achieve 10% on the relative error



$kh \neq \text{constant}$


Numerical Difficulties

$$\frac{|u - u_h|_1}{|u|_1} \leq C_1 kh + C_2 k^3 h^2; \quad kh < 1$$

(Babuška et al (95, 00))

Numerical Difficulties

“Realistic” simulation (Tezaur *et al* (02))



$$\textcolor{red}{k}a = 10$$

System of about 9.6 million complex
unknowns


Motivation and Context

Prominent Plane Waves Based Approaches


- Weak Element Method (**Rose** (75))
- Partition of Unity Method
(**Babuška-Melenk** (97), **Laghrouche-Bettes** (00))
- Ultra-Weak Variational Method
(**Cessenat-Desprī;œs** (98))
- Least Squares Method (**Monk-Wang** (99))
- Discontinuous Galerkin Method: DGM
(**Farhat *et al*** (01, 03, 04, 05))

DGM Formulation (*Farhat et al*)


DGM Formulation (Farhat *et al*) Mathematical model




DGM Formulation (Farhat *et al*) Mathematical model



DGM Formulation (Farhat *et al*) Mathematical model




DGM Formulation (Farhat *et al*) Mathematical model



$$\Delta u + k^2 u = 0 \quad \text{in} \quad \Omega$$

DGM Formulation (Farhat *et al*) Mathematical model




$$\Delta \mathbf{u} + \mathbf{k}^2 \mathbf{u} = 0 \quad \text{in} \quad \Omega$$

$$\partial_n \mathbf{u} = i \mathbf{k} \mathbf{u} + \mathbf{g} \quad \text{on} \quad \Sigma$$


DGM Formulation (*Farhat et al*) Approximation



DGM Formulation (*Farhat et al*) Approximation




DGM Formulation (Farhat *et al*) Approximation



$$\boldsymbol{u} \equiv \sum_{l=1}^4 \boldsymbol{u}_l^K \boldsymbol{\phi}_l^K \text{ in } \mathbf{K}$$
$$\boldsymbol{\phi}_l^K = e^{i\mathbf{k}x \cdot d_l}$$

DGM Formulation (Farhat *et al*) Approximation




$$\boldsymbol{u} \equiv \sum_{l=1}^4 \boldsymbol{u}_l^K \boldsymbol{\phi}_l^K \text{ in } \mathbf{K}$$

$$\boldsymbol{\phi}_l^K = e^{ikx \cdot d_l}$$

$$\lambda = \lambda^K \in \mathbb{C} \text{ on } \partial\mathbf{K} \cap \partial\mathbf{K}'$$

DGM Formulation (Farhat *et al*)


Variational Formulation



$$\begin{cases} a(\textcolor{brown}{u}, v) + \textcolor{magenta}{b}(v, \lambda) = F(v) \\ \textcolor{magenta}{b}(\textcolor{brown}{u}, \mu) = 0 \end{cases}$$

DGM Formulation (Farhat *et al*)

Variational Formulation




$$\begin{cases} \mathbf{a}(\mathbf{u}, \mathbf{v}) + \mathbf{b}(\mathbf{v}, \boldsymbol{\lambda}) = \mathbf{F}(\mathbf{v}) \\ \mathbf{b}(\mathbf{u}, \boldsymbol{\mu}) = \mathbf{0} \end{cases}$$

$$\mathbf{a}(\mathbf{u}, \mathbf{v}) = \sum_{\mathbf{K} \in \tau_h} \left(\int_{\mathbf{K}} (\nabla \mathbf{u} \cdot \nabla \bar{\mathbf{v}} - \mathbf{k}^2 \mathbf{u} \bar{\mathbf{v}}) dx - i \mathbf{k} \int_{\partial \mathbf{K} \cap \Sigma} \mathbf{u} \bar{\mathbf{v}} ds \right)$$

DGM Formulation (Farhat *et al*)

Variational Formulation




$$\begin{cases} a(\mathbf{u}, \mathbf{v}) + \mathbf{b}(\mathbf{v}, \boldsymbol{\lambda}) = \mathbf{F}(\mathbf{v}) \\ \mathbf{b}(\mathbf{u}, \boldsymbol{\mu}) = \mathbf{0} \end{cases}$$

$$\mathbf{b}(\mathbf{v}, \boldsymbol{\mu}) = \sum_{\mathbf{K} \in \tau_h} \int_{\partial\mathbf{K} \cap \partial\mathbf{K}'} \boldsymbol{\mu} \bar{\mathbf{v}} ds$$

DGM Formulation (Farhat *et al*)

Variational Formulation




$$\begin{cases} a(\mathbf{u}, \mathbf{v}) + \mathbf{b}(\mathbf{v}, \boldsymbol{\lambda}) = \mathbf{F}(\mathbf{v}) \\ \mathbf{b}(\mathbf{u}, \boldsymbol{\mu}) = 0 \end{cases}$$

$$\mathbf{F}(\mathbf{v}) = \sum_{\mathbf{K} \in \tau_h} \int_{\partial \mathbf{K} \cap \Sigma} g \bar{v} ds$$

DGM Formulation (Farhat *et al*)


Algebraic Formulation



$$\begin{array}{l|l} \mathbf{A}\mathbf{u} + \mathbf{B}\boldsymbol{\lambda} &= \mathbf{f} \\ \mathbf{B}^T\mathbf{u} &= 0 \end{array}$$

DGM Formulation (Farhat *et al*)

Algebraic Formulation



$$\begin{array}{l} | \quad A\boldsymbol{u} + B\boldsymbol{\lambda} = \boldsymbol{f} \\ \quad \quad B^T \boldsymbol{u} = 0 \end{array}$$

$$\implies B^T A^{-1} B \boldsymbol{\lambda} = B^T A^{-1} \boldsymbol{f}$$

DGM Formulation (*Farhat et al*)

Main Features

DGM Formulation (*Farhat et al*)

Main Features

- **Plane waves** for **local** approximation of the field

DGM Formulation (*Farhat et al*)

Main Features

- **Plane waves** for **local** approximation of the field
- **Lagrange multipliers** to enforce continuity

DGM Formulation (*Farhat et al*)

Main Features

- **Plane waves** for **local** approximation of the field
- **Lagrange multipliers** to enforce continuity
- **Analytical** evaluation of the **elementary** matrices

DGM Formulation (*Farhat et al*)

Main Features

- Plane waves for local approximation of the field
- Lagrange multipliers to enforce continuity
- Analytical evaluation of the elementary matrices
- Global system: symmetric and sparse

DGM Formulation (*Farhat et al*)

Main Features

- Plane waves for local approximation of the field
- Lagrange multipliers to enforce continuity
- Analytical evaluation of the elementary matrices
- Global system: symmetric and sparse
- Size of the global system $\equiv \#$ Lagrange multipliers

DGM Formulation (*Farhat et al*)

Performance

DGM Formulation (Farhat *et al*)

Performance

DGM **outperforms** high order FE methods:

DGM Formulation (Farhat *et al*)

Performance

DGM **outperforms** high order FE methods:

- R-4-1, R-8-2 require 5 to 7 times fewer dof than Q2

DGM Formulation (*Farhat et al*)

Performance

DGM **outperforms** high order FE methods:

- R-4-1, R-8-2 require 5 to 7 times fewer dof than Q2
- Q-16-4 requires 6 times fewer dof than Q4

DGM Formulation (*Farhat et al*)

Performance

DGM **outperforms** high order FE methods:

- R-4-1, R-8-2 require 5 to 7 times fewer dof than Q2
- Q-16-4 requires 6 times fewer dof than Q4
- Q-32-8 requires 25 times fewer dof than Q4

DGM Formulation (*Farhat et al*) Issues


DGM Formulation (*Farhat et al*) Issues

Inf-Sup condition: Discrete spaces compatibility

DGM Formulation (Farhat *et al*)

Issues

Inf-Sup condition: Discrete spaces compatibility





plane waves *vs.* # Lagrange multipliers

?

DGM Formulation (Farhat *et al*) Issues

Inf-Sup condition: Discrete spaces compatibility




R-8-2 element

Relative error, $ka=10$


DGM Formulation (Farhat *et al*)

Issues


Inf-Sup condition: Discrete spaces compatibility



Relative error, $ka=10$



R-8-2 element



R-8-3 element

DGM Formulation (*Farhat et al*)



Issues

Inf-Sup condition: Numerical instabilities

DGM Formulation (Farhat *et al*)

Issues

Inf-Sup condition: Numerical instabilities





R-4-2 element

Total relative error, $\mathbf{ka}=1$

DGM Formulation (Farhat *et al*)

Issues

Inf-Sup condition: Numerical instabilities





R-4-2 element

Total relative error, $\mathbf{ka}=1$

DGM Formulation (Farhat *et al*)

Issues

Inf-Sup condition: Numerical instabilities



R-8-2 element

Total relative error, $ka=1$

Our objective: build on top of DGM

Our objective: build on top of DGM

- Preserve the good features of DGM

Our objective: build on top of DGM

- **Preserve** the good features of DGM
- **Overcome** the numerical instabilities

DGM Formulation: Another point of view

$$\begin{cases} \Delta \mathbf{u} + \mathbf{k}^2 \mathbf{u} = 0 & \text{in } \Omega \\ \partial_n \mathbf{u} = i\mathbf{k}\mathbf{u} + \mathbf{g} & \text{on } \Sigma \end{cases}$$

DGM Formulation: Another point of view

$$\begin{cases} \Delta \mathbf{u} + \mathbf{k}^2 \mathbf{u} = 0 & \text{in } \Omega \\ \partial_n \mathbf{u} = i\mathbf{k}\mathbf{u} + \mathbf{g} & \text{on } \Sigma \end{cases}$$

- Split the solution \mathbf{u} :

$$\mathbf{u} = \boldsymbol{\varphi} + \Phi(\boldsymbol{\lambda})$$

DGM Formulation: Another point of view

$$\left\{ \begin{array}{l} \Delta \varphi^K + k^2 \varphi^K = 0 \quad \text{in } K \\ \end{array} \right.$$

$$\left\{ \begin{array}{l} \Delta \Phi^K(\lambda) + k^2 \Phi^K(\lambda) = 0 \quad \text{in } K \\ \end{array} \right.$$

DGM Formulation: Another point of view

$$\begin{cases} \Delta \varphi^K + k^2 \varphi^K = 0 & \text{in } K \\ \partial_n \varphi^K = ik\varphi^K + g & \text{on } \partial K \cap \Sigma \end{cases}$$

$$\begin{cases} \Delta \Phi^K(\lambda) + k^2 \Phi^K(\lambda) = 0 & \text{in } K \\ \partial_n \Phi^K(\lambda) = ik\Phi^K(\lambda) & \text{on } \partial K \cap \Sigma \end{cases}$$

DGM Formulation: Another point of view

$$\begin{cases} \Delta \varphi^K + k^2 \varphi^K = 0 & \text{in } K \\ \partial_n \varphi^K = ik\varphi^K + g & \text{on } \partial K \cap \Sigma \\ \partial_n \varphi^K = 0 & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

$$\begin{cases} \Delta \Phi^K(\lambda) + k^2 \Phi^K(\lambda) = 0 & \text{in } K \\ \partial_n \Phi^K(\lambda) = ik\Phi^K(\lambda) & \text{on } \partial K \cap \Sigma \\ \partial_n \Phi^K(\lambda) = \lambda & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

DGM Formulation: Another point of view

$$\begin{cases} \Delta \varphi^K + k^2 \varphi^K = 0 & \text{in } K \\ \partial_n \varphi^K = ik\varphi^K + g & \text{on } \partial K \cap \Sigma \\ \partial_n \varphi^K = 0 & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

$$\begin{cases} \Delta \Phi^K(\lambda) + k^2 \Phi^K(\lambda) = 0 & \text{in } K \\ \partial_n \Phi^K(\lambda) = ik\Phi^K(\lambda) & \text{on } \partial K \cap \Sigma \\ \partial_n \Phi^K(\lambda) = \lambda & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

$$\lambda^{K'} = -\lambda^K \text{ on } \partial K \cap \partial K'$$

DGM Formulation: Another point of view

$$\begin{cases} \Delta \varphi^K + k^2 \varphi^K = 0 & \text{in } K \\ \partial_n \varphi^K = ik\varphi^K + g & \text{on } \partial K \cap \Sigma \\ \partial_n \varphi^K = 0 & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

$$\begin{cases} \Delta \Phi^K(\lambda) + k^2 \Phi^K(\lambda) = 0 & \text{in } K \\ \partial_n \Phi^K(\lambda) = ik\Phi^K(\lambda) & \text{on } \partial K \cap \Sigma \\ \partial_n \Phi^K(\lambda) = \lambda & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

$$\lambda^{K'} = -\lambda^K \text{ on } \partial K \cap \partial K'$$

$$[u] = [\varphi + \Phi(\lambda)] = 0 \text{ on each interior edge}$$

DGM Formulation: Another point of view

- Solve **local** variational problems in each \mathbf{K} :

$$\int_{\partial \mathbf{K}} (\partial_n \mathbf{v} - i \mathbf{k} \mathbf{v} \chi_\Sigma) \overline{w} \, ds = L(w)$$

DGM Formulation: Another point of view

- Solve **local** variational problems in each \mathbf{K} :

$$\int_{\partial \mathbf{K}} (\partial_n \mathbf{v} - i \mathbf{k} \mathbf{v} \chi_\Sigma) \overline{w} \, ds = L(w)$$

- Solve one **global** variational problem:

$$\sum_{e \subset \mathring{\Omega}} \frac{1}{|e|} \int_e [\Phi(\lambda)] \overline{\mu} = - \sum_{e \subset \mathring{\Omega}} \frac{1}{|e|} \int_e [\varphi] \overline{\mu}$$

New solution methodologies for Helmholtz problems

A new DGM

A new DGM: first idea



Restore the weak continuity of the field in the least-squares sense:

A new DGM: first idea



Restore the weak continuity of the field in the least-squares sense:

$$\sum_{e \subset \mathring{\Omega}} \frac{1}{|e|} \int_e [\Phi(\lambda)] [\overline{\Phi(\mu)}] = - \sum_{e \subset \mathring{\Omega}} \frac{1}{|e|} \int_e [\varphi] [\overline{\Phi(\mu)}]$$

A new DGM: first idea





Restore the weak continuity of the field in the least-squares sense:

$$\sum_{e \subset \mathring{\Omega}} \frac{1}{|e|} \int_e [\Phi(\lambda)] [\overline{\Phi(\mu)}] = - \sum_{e \subset \mathring{\Omega}} \frac{1}{|e|} \int_e [\varphi] [\overline{\Phi(\mu)}]$$

Gain: Hermitian and **positive semi-definite** global matrix



A new DGM: Illustrative example of the improvement for a fixed resolution



R-8-3 element

Relative error, $ka=10$



A new DGM: Illustrative example of the improvement for a fixed resolution



R-8-3 element

Relative error, $ka=30$

A new DGM: Persistance of the numerical instabilities



R-8-2 element

Total relative error, $ka=1$

A new DGM: second idea



Reformulate the local variational problem:

A new DGM: second idea



Reformulate the local variational problem:

$$\int_{\partial \textcolor{violet}{K}} (\partial_n \textcolor{brown}{v} - i \textcolor{red}{k} \textcolor{brown}{v} \chi_\Sigma) (\partial_n \overline{w} + i \textcolor{red}{k} \overline{w} \chi_\Sigma) = L(w)$$

A new DGM: second idea





Reformulate the local variational problem:

$$\int_{\partial \textcolor{violet}{K}} (\partial_n \textcolor{brown}{v} - i \textcolor{red}{k} \textcolor{brown}{v} \chi_\Sigma) (\partial_n \overline{w} + i \textcolor{red}{k} \overline{w} \chi_\Sigma) = L(w)$$

Gain: Hermitian and **positive definite** local matrix

A new DGM: Illustrative example of the improvement for a fixed frequency



R-8-2 element

Total relative error, $\mathbf{k}\mathbf{a}=1$

A new DGM: Solution strategy

- Solve **local** variational problems in each \mathbf{K} :

$$a_K(v, w) = \int_{\partial K} (\partial_n v - i \mathbf{k} v \chi_\Sigma)(\partial_n \bar{w} + i \mathbf{k} \bar{w} \chi_\Sigma)$$

A new DGM: Solution strategy

- Solve **local** variational problems in each \mathbf{K} :

$$a_K(v, w) = \int_{\partial K} (\partial_n v - i \mathbf{k} v \chi_\Sigma)(\partial_n \bar{w} + i \mathbf{k} \bar{w} \chi_\Sigma)$$

- Solve one **global** variational problem:

$$\begin{aligned} A(\lambda, \mu) = & \sum_{e \subset \mathring{\Omega}} (\beta_e \int_e [\Phi(\lambda)] [\overline{\Phi(\mu)}] \\ & + \gamma_e \int_e [[\partial_n \Phi(\lambda)]] [[\partial_n \overline{\Phi(\mu)}]]) \end{aligned}$$

A new DGM: Accomplishment

- A priori error estimates
- Application in geophysical exploration

A new DGM: A priori error estimates

$$\|\textcolor{brown}{u} - \textcolor{brown}{u}_h\|_{0,\Omega} \leq \hat{C} \inf_{\mu_h, v_h} \left[\sum_{K \in \tau_h} \frac{1}{\textcolor{red}{k}^4 \textcolor{green}{h}_K^3} \left(\|\textcolor{magenta}{\lambda} - \mu_h\|_{0,\partial K \cap \bar{\Omega}}^2 + \|\textcolor{magenta}{\lambda} - \partial_n v_h\|_{0,\partial K \cap \bar{\Omega}}^2 \right) \right]^{1/2}$$

A new DGM: A priori error estimates. Application

\tilde{R} - m - n element ($m \geq 2N + 1$):

$$\|\mathbf{u} - \mathbf{u}_h\|_{0,\Omega} \leq \frac{\hat{C}}{\mathbf{k}^2} \left[\mathbf{h}^n |\Phi(\lambda)|_{n+2,\Omega} \right.$$

$$\left. + \mathbf{h}^{n-1} |\Phi(\lambda)|_{n+1,\Omega} \right]$$

$$+ \mathbf{h}^{N-2} \left(\sum_{l=0}^N \mathbf{k}^{N+1-l} |\Phi(\lambda)|_{l,\Omega} \right.$$

$$\left. + |\Phi(\lambda)|_{N+1,\Omega} + \mathbf{h} |\Phi(\lambda)|_{N+2,\Omega} \right) \Big]$$

A new DGM: Application in geophysical exploration

A new DGM: Application in geophysical exploration

- **Objective:** produce images of the subsurface from tomography measurements



A new DGM: Application in geophysical exploration

- **Objective:** produce images of the subsurface from tomography measurements

A new DGM: Application in geophysical exploration

- **Objective:** produce images of the subsurface from tomography measurements
- Wave propagation in **time** domain:
Discrete Fourier Transform


A new DGM: Application in geophysical exploration

- **Objective:** produce images of the subsurface from tomography measurements
- Wave propagation in **time** domain:
Discrete Fourier Transform
- Solve Helmholtz equation (reduced wave equation)

A new DGM: Application in geophysical exploration


- **Objective:** produce images of the subsurface from tomography measurements
- Wave propagation in **time** domain:
Discrete Fourier Transform
- Solve Helmholtz equation (reduced wave equation)
- Build the solution in **time** domain:
Inverse Discrete Fourier Transform

A new DGM: Application in geophysical exploration




Homogeneous medium

A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{30} \leq kh \leq 2$




Homogeneous medium

A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{30} \leq kh \leq 2$




Homogeneous medium

A new DGM: Application in geophysical exploration





R-4-1 element
 $\frac{1}{30} \leq kh \leq 2$



Homogeneous medium



A new DGM: Application in geophysical exploration



R-4-1 element
 $\frac{1}{30} \leq kh \leq 2$

Homogeneous medium



A new DGM: Application in geophysical exploration



R-4-1 element
 $\frac{1}{30} \leq kh \leq 2$

Homogeneous medium



A new DGM: Application in geophysical exploration



R-4-1 element
 $\frac{1}{30} \leq kh \leq 2$

Homogeneous medium



A new DGM: Application in geophysical exploration



R-4-1 element
 $\frac{1}{30} \leq kh \leq 2$

Homogeneous medium


A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{30} \leq kh \leq 2$

Homogeneous medium

A new DGM: Application in geophysical exploration





R-4-1 element
 $\frac{1}{30} \leq kh \leq 2$



Homogeneous medium



A new DGM: Application in geophysical exploration



R-4-1 element
 $\frac{1}{30} \leq kh \leq 2$

Homogeneous medium


A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{30} \leq kh \leq 2$

Homogeneous medium

A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{30} \leq kh \leq 2$





Homogeneous medium

A new DGM: Application in geophysical exploration



Stratified medium


A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{50} \leq kh \leq 2$

Stratified medium

A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{50} \leq kh \leq 2$




Stratified medium

A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{50} \leq kh \leq 2$




Stratified medium

A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{50} \leq kh \leq 2$




Stratified medium

A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{50} \leq kh \leq 2$




Stratified medium

A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{50} \leq kh \leq 2$




Stratified medium

A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{50} \leq kh \leq 2$




Stratified medium

A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{50} \leq kh \leq 2$




Stratified medium

A new DGM: Application in geophysical exploration




R-4-1 element
 $\frac{1}{50} \leq kh \leq 2$




Stratified medium

A new DGM: Application in geophysical exploration





R-4-1 element
 $\frac{1}{50} \leq kh \leq 2$



Stratified medium

A new DGM: Application in geophysical exploration



R-4-1 element
 $\frac{1}{50} \leq kh \leq 2$

Stratified medium

A new DGM: Application in geophysical exploration

- Multi-frequency solver


A new DGM: Application in geophysical exploration

- Multi-frequency solver
- Resolution: 3 to 300 elements per wavelength


A new DGM: Application in geophysical exploration

- Multi-frequency solver
- Resolution: 3 to 300 elements per wavelength
- R-4-1 element: accurate and stable element

A new DGM: Stability of the R-4-1 element



L^2 error, $ka=10$



R-4-1 element

A new DGM: Summary

DGM:

- Solve **local** variational problems in each $\textcolor{violet}{K}$:

$$a_K(\textcolor{brown}{v}, w) = \int_{\partial \textcolor{violet}{K}} (\partial_n \textcolor{brown}{v} - i \textcolor{red}{k} \textcolor{brown}{v} \chi_\Sigma) \overline{w} \, ds$$

- Solve **global** variational problem:

$$A(\lambda, \mu) = \sum_{e \subset \mathring{\Omega}} \frac{1}{|e|} \int_e [\Phi(\lambda)] \overline{\mu}$$

A new DGM: Summary

A new DGM:

- NEW local variational problems in each $\textcolor{violet}{K}$:

$$a_K(\textcolor{brown}{v}, w) = \int_{\partial \textcolor{violet}{K}} (\partial_n \textcolor{brown}{v} - i \textcolor{red}{k} v \chi_\Sigma) (\partial_n \overline{w} + i \textcolor{red}{k} \overline{w} \chi_\Sigma) ds$$

- Solve global variational problem:

$$A(\lambda, \mu) = \sum_{e \subset \mathring{\Omega}} \frac{1}{|e|} \int_e [\Phi(\lambda)] \overline{\mu}$$

A new DGM: Summary

A new DGM:

- NEW local variational problems in each $\textcolor{violet}{K}$:

$$a_K(\textcolor{brown}{v}, w) = \int_{\partial \textcolor{violet}{K}} (\partial_n \textcolor{brown}{v} - i \textcolor{red}{k} v \chi_\Sigma)(\partial_n \overline{w} + i \textcolor{red}{k} \overline{w} \chi_\Sigma) ds$$



- NEW global variational problem:

$$A(\lambda, \mu) = \sum_{e \subset \mathring{\Omega}} \left(\beta_e \int_e [\Phi(\lambda)] [\overline{\Phi(\mu)}] + \right.$$

$$\left. \gamma_e \int_e [[\partial_n \Phi(\lambda)]] [[\partial_n \overline{\Phi(\mu)}]] \right)$$

A new DGM: Summary

- Numerical instabilities: improvement, but not enough





R-8-2 element

Total relative error, $ka=1$

A new DGM: Summary

- Numerical instabilities: improvement, but not enough



R-8-2 element

Total relative error, $ka=1$

- Source of the instabilities: local problems **nearly singular**

New solution methodologies for Helmholtz problems

A modified DGM (mDGM)

mDGM: main idea



Modify the local problems:

$$\begin{cases} \Delta \varphi^K + k^2 \varphi^K = 0 & \text{in } K \\ \partial_n \varphi^K - ik \varphi^K = g & \text{on } \partial K \cap \Sigma \\ \partial_n \varphi^K = 0 & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

$$\begin{cases} \Delta \Phi^K(\lambda) + k^2 \Phi^K(\lambda) = 0 & \text{in } K \\ \partial_n \Phi^K(\lambda) - ik \Phi^K(\lambda) = 0 & \text{on } \partial K \cap \Sigma \\ \partial_n \Phi^K(\lambda) = \lambda & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

mDGM: main idea



Modify the local problems:

$$\begin{cases} \Delta \varphi^K + k^2 \varphi^K = 0 & \text{in } K \\ \partial_n \varphi^K - ik \varphi^K = g & \text{on } \partial K \cap \Sigma \\ \partial_n \varphi^K - i \underline{\alpha \varphi^K} = 0 & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

$$\begin{cases} \Delta \Phi^K(\lambda) + k^2 \Phi^K(\lambda) = 0 & \text{in } K \\ \partial_n \Phi^K(\lambda) - ik \Phi^K(\lambda) = 0 & \text{on } \partial K \cap \Sigma \\ \partial_n \Phi^K(\lambda) - i \underline{\alpha \Phi^K(\lambda)} = \lambda & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

mDGM: main idea



Modify the local problems:

$$\begin{cases} \Delta \varphi^K + k^2 \varphi^K = 0 & \text{in } K \\ \partial_n \varphi^K - ik \varphi^K = g & \text{on } \partial K \cap \Sigma \\ \partial_n \varphi^K - i \alpha \varphi^K = 0 & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

$$\begin{cases} \Delta \Phi^K(\lambda) + k^2 \Phi^K(\lambda) = 0 & \text{in } K \\ \partial_n \Phi^K(\lambda) - ik \Phi^K(\lambda) = 0 & \text{on } \partial K \cap \Sigma \\ \partial_n \Phi^K(\lambda) - i \alpha \Phi^K(\lambda) = \lambda & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

Gain: Well-posed local problems for $\alpha \in \mathbb{R}_+^*$

mDGM: main idea



Modify the local problems:

$$\begin{cases} \Delta \varphi^K + k^2 \varphi^K = 0 & \text{in } K \\ \partial_n \varphi^K - ik \varphi^K = g & \text{on } \partial K \cap \Sigma \\ \partial_n \varphi^K - i \alpha \varphi^K = 0 & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

$$\begin{cases} \Delta \Phi^K(\lambda) + k^2 \Phi^K(\lambda) = 0 & \text{in } K \\ \partial_n \Phi^K(\lambda) - ik \Phi^K(\lambda) = 0 & \text{on } \partial K \cap \Sigma \\ \partial_n \Phi^K(\lambda) - i \alpha \Phi^K(\lambda) = \lambda & \text{on } \partial K \cap \bar{\Omega} \end{cases}$$

Gain: Well-posed local problems for $\alpha \in \mathbb{R}_+^*$

Price: Both u and $\partial_n u$ are discontinuous

mDGM: Solution strategy

- Solve **local** variational problems in each \mathbf{K} :

$$\int_{\partial \mathbf{K}} (\partial_n \mathbf{v} - i \mathbf{k} \mathbf{v}) \bar{w} \, ds = L(w)$$

mDGM: Solution strategy


- Solve **local** variational problems in each $\textcolor{violet}{K}$:

$$\int_{\partial \textcolor{violet}{K}} (\partial_n \textcolor{brown}{v} - i \textcolor{red}{k} v) \bar{w} \, ds = L(w)$$


- Solve one **global** variational problem:

$$\sum_{e \subset \mathring{\Omega}} \left(\beta_e \int_e [\Phi(\lambda)] [\overline{\Phi(\mu)}] + \gamma_e \int_e [[\partial_n \Phi(\lambda)]] [[\partial_n \overline{\Phi(\mu)}]] \right) = \\ - \sum_{e \subset \mathring{\Omega}} \left(\beta_e \int_e [\varphi] [\overline{\Phi(\mu)}] + \gamma_e \int_e [[\partial_n \varphi]] [[\partial_n \overline{\Phi(\mu)}]] \right)$$

mDGM: Performance assessment




Total relative error, $ka=1$




R-4-2 element

mDGM: Performance assessment




Total relative error, $ka=1$




R-4-2 element

mDGM: Performance assessment




Total relative error, $\kappa a = 1$




R-4-2 element

mDGM: Performance assessment




Total relative error, $ka=1$




R-8-2 element

mDGM: Performance assessment




Total relative error, $ka=1$




R-8-3 element

mDGM: Performance assessment




Total relative error, $ka=1$




R-8-4 element

mDGM: Performance assessment



Total relative error, $\mathbf{k}\mathbf{a}=1$



R-8-5 element

mDGM: Implementation issue

$$F_{jl} = \sum_{e \subset \mathring{\Omega}} \left(\beta_e \int_e [\Phi_h(\mu_l)] [\overline{\Phi_h(\mu_j)}] + \gamma_e \int_e [[\partial_n \Phi_h(\mu_l)]] [[\partial_n \overline{\Phi_h(\mu_j)}]] \right)$$



F is Hermitian

mDGM: Implementation issue

$$F_{jl} = \sum_{e \subset \mathring{\Omega}} \left(\beta_e \int_e [\Phi_h(\mu_l)] [\overline{\Phi_h(\mu_j)}] + \gamma_e \int_e [[\partial_n \Phi_h(\mu_l)]] [[\partial_n \overline{\Phi_h(\mu_j)}]] \right)$$

⚠ F is Hermitian, but NOT in practice

mDGM: Implementation issue




**R-8-2
element**



Total relative error, $ka=1$

mDGM: Implementation issue

Before



After




R-8-3
element



Total relative error, $ka=1$

mDGM: Implementation issue

Before



After




R-8-4
element



Total relative error, $ka=1$

mDGM: Implementation issue

Before





After



R-8-5
element

Total relative error, $ka=1$



mDGM: Numerical issue



R-8-2 element

The smallest eigenvalue, $\mathbf{k}\mathbf{a}=1$

mDGM: Numerical issue





R-8-2 element

The smallest eigenvalue, $\mathbf{k}\mathbf{a}=1$




Loss of the linear independence

mDGM: Numerical issue




R-8-2 element




R-7-2 element

The smallest eigenvalue, $ka=1$


mDGM: Numerical issue



Total relative error, $\kappa a = 1$





R-8-2 element



R-7-2 element

mDGM: Numerical issue



R-7-2 element

Total relative error, $ka=1$

mDGM: Summary

Local problems:

- DGM:

$$a_K(v, w) = \int_{\partial K} (\partial_n v - i \mathbf{k} v \chi_\Sigma) \bar{w} \, ds$$

- mDGM:

$$a_K(v, w) = \int_{\partial K} (\partial_n v - i \mathbf{k} v) \bar{w} \, ds$$

mDGM: Summary

Global problems:

- DGM:

$$A(\lambda, \mu) = \sum_{e \subset \mathring{\Omega}} \frac{1}{|e|} \int_e [\Phi(\lambda)] \bar{\mu}$$



- mDGM:

$$A(\lambda, \mu) = \sum_{e \subset \mathring{\Omega}} \left(\beta_e \int_e [\Phi(\lambda)] [\overline{\Phi(\mu)}] \right.$$

$$\left. + \gamma_e \int_e [[\partial_n \Phi(\lambda)]] [[\partial_n \overline{\Phi(\mu)}]] \right)$$

mDGM: Summary

- Numerical instabilities: improvement, but not enough





R-7-2 element

Total relative error, $ka=1$

mDGM: Summary

- Numerical instabilities: improvement, but not enough



R-7-2 element

Total relative error, $ka=1$

- Source of the instabilities: loss of the linear independence

New solution methodologies for Helmholtz problems

An improved modified DGM (imDGM)

imDGM: main idea



Reformulate the local variational problems:

imDGM: main idea



Reformulate the local variational problems:

$$\int_{\partial K} (\partial_n \mathbf{v} - i \mathbf{k} \mathbf{v}) (\partial_n \overline{\mathbf{w}} + i \mathbf{k} \overline{\mathbf{w}}) \, ds = L(\mathbf{w})$$

imDGM: main idea





Reformulate the local variational problems:

$$\int_{\partial K} (\partial_n \mathbf{v} - i \mathbf{k} \mathbf{v}) (\partial_n \overline{\mathbf{w}} + i \mathbf{k} \overline{\mathbf{w}}) \, ds = L(\mathbf{w})$$

Gain: Hermitian and **positive definite** local matrix



imDGM: Performance assessment



R-7-2 element

The smallest eigenvalue, $ka=1$

imDGM: Performance assessment



R-7-2 element

Total relative error, $\mathbf{k}\mathbf{a}=1$

imDGM: Comparison with DGM and LSM (Monk-Wang, 1999)

imDGM: Comparison with DGM and LSM (Monk-Wang, 1999)

- An improved modified DGM (imDGM)

$$A(\lambda, \mu) = \sum_{e \subset \mathring{\Omega}} \left(\beta_e \int_e [\Phi(\lambda)] [\overline{\Phi(\mu)}] + \gamma_e \int_e [[\partial_n \Phi(\lambda)]] [[\partial_n \overline{\Phi(\mu)}]] \right)$$



- Least-Squares Method (LSM)

$$A(u, v) = \sum_{e \subset \mathring{\Omega}} \left(k^2 \int_e [u] [\bar{v}] + \int_e [[\partial_n u]] [[\partial_n \bar{v}]] \right)$$

- Discontinuous Galerkin Method (DGM)

$$A(\lambda, \mu) = \sum_{e \subset \mathring{\Omega}} \left(\beta_e \int_e [\Phi(\lambda)] \bar{\mu} \right)$$



imDGM: Comparison with DGM and LSM



R-7-2 element

Total relative error, $\mathbf{k}\mathbf{a}=1$

imDGM: Comparison with DGM and LSM




R-7-2 element


Total relative error, $\text{ka}=20$

imDGM: Performance assessment for a fixed resolution: $kh=2$

ka R-7-2 R-11-3




R-7-2 element




R-11-3 element

imDGM: Performance assessment for a fixed resolution: $kh=2$

ka	R-7-2	R-11-3
50	28%	0.05%




R-7-2 element




R-11-3 element

imDGM: Performance assessment for a fixed resolution: $kh=2$

ka	R-7-2	R-11-3
50	28%	0.05%
100	51%	0.07%




R-7-2 element




R-11-3 element

imDGM: Performance assessment for a fixed resolution: $kh=2$

ka	R-7-2	R-11-3
50	28%	0.05%
100	51%	0.07%
200	69%	0.20%




R-7-2 element




R-11-3 element

imDGM: Performance assessment for a fixed resolution: $kh=2$

ka	R-7-2	R-11-3
50	28%	0.05%
100	51%	0.07%
200	69%	0.20%




R-7-2 element




R-11-3 element

imDGM: Performance assessment for a fixed resolution: $kh=2$

ka	R-7-2	R-11-3
50	28%	0.05%
100	51%	0.07%
200	69%	0.20%





R-7-2 element



R-11-3 element

Computational cost increased by **50%**
Gain of more than **two orders** of magnitude



imDGM: Comparison with DGM and LSM for a fixed frequency



R-11-3 element

Total relative error, $ka=200$

imDGM: Comparison with DGM and LSM for a fixed frequency



R-11-3 element

Total relative error, $ka=400$

Summary and Perspectives

Accomplishment

Accomplishment

- Small systems: Hermitian and **positive definite**

Accomplishment

- Small systems: Hermitian and **positive definite**
- Global system: Hermitian, sparse and **positive semi-definite**

Accomplishment

- Small systems: Hermitian and **positive definite**
- Global system: Hermitian, sparse and **positive semi-definite**
- Performance: **outperforms** DGM
(accuracy and stability)

Short-term goals

- Assess the performance of higher-order elements

Short-term goals

- Assess the performance of higher-order elements
- Experiment other shape functions

Short-term goals

- Assess the performance of higher-order elements
- Experiment other shape functions
- Unstructured mesh

Short-term goals

- Assess the performance of higher-order elements
- Experiment other shape functions
- Unstructured mesh
- Apply iterative methods for the global system

Short-term goals

- Assess the performance of higher-order elements
- Experiment other shape functions
- Unstructured mesh
- Apply iterative methods for the global system
- Extend the mathematical analysis to imDGM

Mid-term goals

- Three-dimensional acoustic scattering problems

Mid-term goals

- Three-dimensional acoustic scattering problems
- Elasto-acoustic scattering problems

The End...

Thank you for your attention!