
High-energy gamma-ray emission in compact binaries

Benoît Cerutti

Advisors: Guillaume Dubus & Gilles Henri
Laboratoire d'Astrophysique de Grenoble

PhD defense, June 10th 2010, Grenoble


Outline

- 1. Binary systems in the gamma-ray sky!**
- 2. High-energy processes & anisotropies**
- 3. Gamma-ray emission in gamma-ray binaries**
- 4. Pair cascade emission in gamma-ray binaries**
- 5. Gamma-ray emission from relativistic outflow**
- 6. Conclusions & futures directions**

Binary systems in the gamma-ray sky!

Gamma rays

Electromagnetic spectrum:




- High-energy (HE): $\sim \text{GeV}$ (10^9 eV)
- Very-high energy (VHE): $\sim \text{TeV}$ (10^{12} eV)


Gamma-ray Astronomy

Gamma-ray telescopes

From space: **GeV** domain




Fermi, AGILE



From the ground: **TeV** domain




HESS, MAGIC & VERITAS



- Simultaneous coverage of **HE** and **VHE** domains
- Unprecedented **sensitivity** and **angular resolution**

γ -ray astronomy!


The optical sky



© 2000, Axel Mellinger

The gamma-ray sky

1 year of exposure



Extra-Galactic γ -ray sources:

- Active Galactic Nuclei (AGN)
- Gamma-Ray Bursts (GRB)
- Starbursts Galaxies

Galactic γ -ray sources:

- Pulsars & Pulsar wind nebulae
 - Supernovae remnants
 - Many are unidentified
- ... and 4 binary systems!

Gamma-ray emitting binaries

1 year of exposure

- γ -ray binary
- Microquasar

(Cyg X-1?)
LS I +61°303 Cyg X-3 LS 5039 PSR B1259-63 (HESS J0632+057?)

2 kpc


7 kpc

2.5 kpc

1.5 kpc



Credit: NASA/DOE/Fermi LAT Collaboration



Legend

- Massive star
- Compact object orbit


Two classes of binaries

GAMMA-RAY BINARIES

=

Massive star + young pulsar

[Maraschi & Treves 1980, Dubus 2006b]



Credit: F. Mirabel


ROTATION-POWERED

LS I+61°303, LS 5039 & **PSR B1259-63**
(HESS J0632+057?)

MICROQUASAR

=


Massive star + accreting compact object
(neutron star or black hole)



ACCRETION-POWERED

Cyg X-3
(Cyg X-1?)

Why are these systems so interesting?




Gamma-ray binaries

- **Nearby** systems (kpc) and **well constrained physical** conditions (stellar radiation field, orbit)
- Probe **relativistic winds** at **small scale**: 0.1 AU !
- Exploration of **new class** of Galactic objects (population, evolution)
- Application to **pulsars** and **PWN**


[Credit: F.Seward et al, J.Hester & A.Loll, R.Gehrz]

Microquasars

- **Nearby** systems (kpc) and **well constrained physical** conditions (stellar radiation field, orbit)
- Study the **physics of relativistic jets, accretion-ejection** mechanisms at small scales (AU-pc)
- Application to e.g. **AGN & GRBs**




Gamma-ray beacons: the case of LS 5039




Credit: M. de Naurois

- GeV/TeV stable orbital modulation
- Min/Max at conjunctions
- Geometrical effects !?



Conj. Sup. Conj. Inf.



[Aharonian et al. 2006]

What we want to understand

GeV and/or TeV orbital modulation

GeV

- LS I +61°303 [Fermi LAT, 2009a]
- LS 5039 [Fermi LAT, 2009b]
- Cygnus X-3 [Fermi LAT, 2009c]
- (PSR B1259-63 ?)

TeV

- LS I +61°303 [MAGIC coll., 2008]
- LS 5039 [HESS coll., 2006]
- PSR B1259-63 [HESS coll., 2005]

Key questions

- Where does the gamma-ray modulation come from?
- What are the relevant processes in binaries? Acceleration site?
- What can we learn on pulsar winds and relativistic jets?


Goal thesis → Theoretical modeling of temporal & spectral γ -ray variability

High-energy processes & anisotropies


Relevant high-energy processes

Leptonic

- Inverse Compton scattering




- Synchrotron radiation




- Triplet Pair Production

Sarkis Rastikian work




Hadronic

- Proton-proton collision




- Photomeson production




Absorption

- Pair production



Anisotropies



- High UV stellar photon density, **dominant source**
($n_{ph} \approx 10^{14} \text{ ph/cm}^3$ in **LS 5039!**)
- **Anisotropic** and **well constrained** radiation source
→ **Angular dependence** on:
inverse Compton scattering (emission) & **pair production** (absorption)


Anisotropic inverse Compton scattering

Derivation of **new analytical formulae** for anisotropic inverse Compton scattering

Circular orbit with an inclination $i=90^\circ$


Massive star: **Black-body** spectrum

Relativistic electrons with a -2 power-law



Head-on: $e^\pm \rightarrow \gamma \gamma$ Max

Rear-end: $e^\pm \gamma \rightarrow \gamma \gamma$ Min




Strong angular dependence!

Similar dependence for **pair production**

Gamma-ray emission in γ -ray binaries

The scenario: the big picture!



[Maraschi & Treves 1981; Dubus 2006b]

Observer!


Classical pulsar wind model

[Rees & Gunn 1974, Kennel & Coroniti 1984]

- “Unshocked”: cold plasma of relativistic e^+ / e^- pairs: $v_w \approx c$
- “Shocked”: pairs randomized and reaccelerated: $v_w \approx c/3$
- Model applied to PSR B1259-63
[Tavani et al. 1994, Kirk et al. 1999, Ball & Kirk 2000, Khangulyan et al. 2007]

Emission from the “shocked” wind

[Maraschi & Treves 1981; Dubus 2006b]




Observer!

Assumptions

- Isotropized pairs with power-law
- Anisotropic inverse Compton with stellar photons + Synchrotron + Anisotropic pair production


Application to LS 5039

Spectra




- High inclination favored
- compact object = pulsar
- GeV/TeV anti-correlation predicted! (due to absorption)
[see also Bednarek 2006, Khangulyan et. al 2008]

γ -ray flux: absorption & Compton





HESS data
points



[Dubus, Cerutti & Henri, A&A 2008]

Strength & weakness



Strength


- Simple model
- Explains most of the **GeV** and **TeV** modulation in **LS 5039**.
- Robust constraints on physical parameters (B, p, L_p)

Weakness

- **Fermi** observations **not explained!**
Extra-component required.
- **HESS** flux @ superior conjunction not explained in **LS 5039** → **Pair Cascades?**
- Modulation in **LS I +61°303** and **PSR B1259-63** **not explained**

Emission from the “unshocked” pulsar wind

[Maraschi & Treves 1981; Dubus 2006b]




Observer!

Assumptions

- **Mono-energetic** pulsar wind: Lorenz factor γ_0
- **Inverse Compton** with stellar photons, **no synchrotron**
- **Kinetic energy dominated** wind ($\sigma \ll 1$)


Application to γ -ray binaries



- **Strong** line-like spectra! ➔ **Excluded** by *Fermi* and TeV observations
- **Not mono-energetic**, e.g. power-law? [Sierpowska-Bartosik & Torres 2008]
- Conversion **Electromagnetic** to **kinetic** energy **not completed?** Striped wind?

Pair cascade emission in γ -ray binaries


Motivations



[Dubus, Cerutti & Henri, A&A 2008]


- **Model: Huge absorption** at superior conjunction \rightarrow no TeV flux...
- ...but, **excess** measured by HESS!

Solution: Pair cascade emission!



- Absorbed energy **recycled**
 \rightarrow Emission of new generations of pairs
- **Increase the transparency** of the source at superior conjunction.

The physics of pair cascading




Cascade regimes

10^{-8} G

10^{-2} G


10 G Magnetic field

1D Cascade




- Magnetic deviations **negligible** ($B < 10^{-8}$ G)
- **Pairs and photons boosted** in the direction of primary γ -rays

3D “isotropic” Cascade



- Pairs locally **isotropized** ($R_L < \lambda_{IC}$)
- **Extended & anisotropic** emission

Quenched 3D Cascade





- Dominant **Synchrotron** cooling ($t_{syn} < t_{ic}$)
- X-ray emission, **γ -ray emission quenched**


1D cascade

- **No magnetic field** (or $B < 10^{-8}$ Gauss)
- Cascade develops **along the line of sight** → **High anisotropy**
- **Complete and accurate** semi-analytical calculation possible (all generations)

Ex: LS 5039 at sup. conj.



Application to LS 5039



[Cerutti, Dubus & Henri, A&A 2009]

- Contribution too high from the cascade
→ Cascade 1D excluded!
- Upper limit for the cascade contribution @ superior conjunction
→ Cascade 3D not excluded

Cascade regimes


10^{-8} G

10^{-2} G


10 G

Magnetic field


1D Cascade



3D “isotropic” Cascade



Quenched 3D Cascade



- Magnetic deviations **negligible** ($B < 10^{-8}$ G)
- **Pairs and photons boosted** in the direction of primary γ -rays

- Pairs locally **isotropized** ($R_L < \lambda_{IC}$)
- **Extended & anisotropic** emission

- Dominant **Synchrotron** cooling ($t_{syn} < t_{ic}$)
- X-ray emission, **γ -ray emission quenched**

3D cascades: assumptions

- Pairs **isotropized** at creation
- Compton & synchrotron **radiation** and **cooling**
- **No explicit** full analytical calculation possible


→ Decomposition into **discrete generations** of particles

■ Methods:

1. **Semi-analytical (1st generation)**
2. **Monte Carlo (all generations)**

→ Collaboration with Julien Malzac


Both methods **compatible**
Effect of **extra-generations** important



[Cerutti, Malzac, Dubus, & Henri, A&A 2010,
(accepted)]

Spatial distribution 1st generation γ rays




Application to LS 5039: Semi-analytical results





[Cerutti, Malzac, Dubus, & Henri, A&A 2010, (accepted)]

3D cascade modulation in LS 5039

Fermi




HESS



3D cascade modulation in LS 5039

$i=40^\circ$

Total
3D
Cascade
only
More flux @ sup.
Conjunction!




- Rather low inclination: $i \approx 40^\circ$
- TeV source should lie close to the compact object
- Ambient magnetic field $\mathbf{B} < 10 \text{ G}$ (synchrotron radiation from pairs in the cascade)

γ -ray emission from relativistic outflow

The puzzling X-ray modulation in LS 5039 !

Suzaku observations

X-rays




X-ray lightcurve

- **Orbital** modulation
- Min and Max at **conjunctions**:
 - **Intrinsic** variation **unlikely**
 - **Geometrical** explanation


Solution: Doppler Boost!

Observer !



Boosted emission in LS 5039

Deboosted !



- Shocked pulsar wind \approx radial
(stellar wind speed $>>$ orbital motion pulsar)

- Mildly relativistic wind $\beta=1/3$ ($=v/c$)



- TeV modulation \approx unchanged

- X-ray modulation \approx reproduced

$\beta=0$


Boost

$\beta=1/3$



[Dubus, Cerutti, & Henri 2010b]



Boosted emission in LSI +61°303 (1)



■ **Model:** Pulsar wind **trailing** in the Be Keplerian disk

■ Mildly relativistic wind $\beta=1/3$:


X-ray and **TeV** both **correlated** and **maximum close to apastron**



[Dubus, Cerutti, & Henri 2010b]


Boosted emission in LS I +61°303 (2)

MAGIC lightcurve




[Albert et al. 2009]

X- γ correlation



[Anderhub et al. 2009]

Model for $\beta=1/3$



[Dubus, Cerutti, & Henri 2010b]

- The Boost in LS I +61°303 could explain TeV and X-ray modulations !
- Same model for PSR B1259-63
→ small impact


Two classes of binaries

GAMMA-RAY BINARIES

=

Massive star + young pulsar

[Maraschi & Treves 1980, Dubus 2006b]



Credit: F. Mirabel


ROTATION-POWERED

LS I +61°303, LS 5039 & **PSR B1259-63**
(HESS J0632+057?)

MICROQUASAR

=

Massive star + accreting compact object
(neutron star or black hole)




ACCRETION-POWERED



Cyg X-3
(Cyg X-1?)

Cygnus X-3: The first GeV microquasar!

[*Fermi LAT coll., 2009c*]




GeV light curve



- First firm detection of a microquasar at **GeV** by **AGILE & Fermi** [*Tavani et al. 2009, Fermi LAT coll. 2009c*]
- γ -ray flares correlated with radio flares
→ γ -ray emission related to the **jet**
- γ -ray flux modulated at the **orbital period** (4.8 h)

Relativistic Doppler emission in Cygnus X-3



Assumptions


- **Boosted Inverse Compton** stellar photons – pairs in the jet
- The jet is **inclined & relativistic**
- Energetic electrons **located @ specific locations**

Parameters

- **5 free** parameters
- **Exhaustive exploration** of parameter space → $\min \chi^2$

Best fit solution

Best fit



Constraints

[*Dubus, Cerutti, & Henri, MNRAS 2010a*]

- e^\pm not located within the system
- Jet inclined close to the line of sight (μ -blazar!)
- Jet mildly relativistic ($\beta < 0.9$)
- Black-hole favored
- Precession of the jet: strong impact on the modulation

Conclusions & futures directions

Summary

Pulsar wind emission

- **Shocked: Anisotropic IC + absorption: TeV modulation OK in LS 5039.**
- **Unshocked: Strong emission excluded: wind highly magnetized?**

Publications: [Dubus, Cerutti, & Henri, 2008]; [Cerutti, Dubus, & Henri 2008]

Cascades

- **1D cascade:** provides an **upper limit**, **excluded** in LS 5039
- **3D cascades:** **Viable solution** for **TeV** flux @ sup. conj. in LS 5039

Publications: [Cerutti, Dubus, & Henri, 2009]; [Cerutti, Malzac, Dubus, & Henri, 2010]

Boosted emission

- **γ -ray binaries:** changes **significantly** high-energy modulation (X, γ)
- **Cygnus X-3:** **Boosted** Compton scattering → **GeV** modulation **OK**

Publications: [Dubus, Cerutti, & Henri, 2010a]; [Dubus, Cerutti, & Henri, 2010b]

Open questions and looking forward

Open questions

- What is the origin of the **GeV component?** -> **Striped pulsar wind in binaries?**
- What is the origin of the **γ -ray modulation** in **LS I +61°303** and **PSR B1259-63?**
- Particle acceleration in **microquasar jet?**


Looking forward

- **Present and future observational backdrop favorable** (Fermi, CTA, ...)
→ **Discovery of new systems!**
- Exciting **new modeling challenges** with **GeV** observations!
(pulsar emission, model for microquasars ...)
- **Theoretical expertise in high-energy processes** in astrophysics
→ Application to many other objects e.g.:
 - **AGN and GRBs**
 - **Pulsars, PWN... and magnetars!**

Future directions

Post-doc

- **Research associate** position with Pr. **Dmitri Uzdensky** in Theoretical plasma astrophysics @ University of Colorado, Boulder, USA.
- Magnetic reconnection in **High-energy-density astrophysical environment** (*e.g. magnetars & GRBs*)



University of Colorado

The end
