



**HAL**  
open science

## Role of ASF1 in histone deposition during replication

Vivek Tripathi

► **To cite this version:**

Vivek Tripathi. Role of ASF1 in histone deposition during replication. Biochemistry, Molecular Biology. Université de Strasbourg, 2012. English. NNT : 2012STRAJ036 . tel-01081099

**HAL Id: tel-01081099**

**<https://theses.hal.science/tel-01081099>**

Submitted on 7 Nov 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Role of ASF1 in Histone Deposition During Replication

By

Vivek Tripathi A

thesis submitted to the

University of Strasbourg

For the degree of

Doctor of Philosophy

November 7<sup>th</sup>, 2012

Thesis Director

Dr. Ali Hamiche

Rapporteur

Dr. Stefan Dimitrov

Rapporteur

Dr. Mikhail Grigoriev

Examiner

Dr. Christophe Romier

## Table of contents

Acknowledgements	3
Abstract	4
List of abbreviations	5
List of figures and tables	7
<b>1 Introduction</b>	
<b>1.1 Chromatin</b>	<b>9</b>
1.1.1 Nucleosome	9
1.1.2 Chromatin higher order structures	11
1.1.3 Specialized chromatin structures	13
1.1.3.1 Euchromatin	13
1.1.3.2 Heterochromatin	14
1.1.3.3 Heterochromatin proteins	15
<b>1.2 Chromatin Dynamics</b>	<b>18</b>
1.2.1 Histone post Posttranslational modifications	19
1.2.1.1 Acetylation	20
1.2.1.2 Methylation	21
1.2.1.3 Phosphorylation	23
1.2.1.4 Ubiquitination	23
1.2.1.5 Sumoylation	24
1.2.1.6 ADP-ribosylation	24
1.2.2 Histone variants	25
1.2.2.1 Histone H1 variants	25
1.2.2.2 Histone H2A variants	26
1.2.2.3 Histone H3 variants	29
1.2.2.4 H2B variants	30
1.2.3 ATP-dependent chromatin remodeling enzymes	30

1.2.3.1 SWI/SNF family remodelers	31
1.2.3.2 ISWI family remodelers	31
1.2.3.3 CHD family remodelers	32
1.2.3.4 INO80 family remodelers	32
1.2.4 Histone Chaperones	33
1.2.4.1 Histone chaperone N1/N2	36
1.2.4.2 Histone chaperone NAP1	37
1.2.4.3 Histone chaperone HIRA	39
1.2.4.4 DAXX	40
1.2.4.5 Histone chaperone CAF1	41
1.2.4.6 Histone chaperone Asf1	42
1.3 Histone Deposition	44
1.3.1 Replication independent chromatin assembly	47
1.3.2 Replication dependent Chromatin assembly	49
1.4 Dynamic roles of the histone chaperone Asf1	52
1.4.1 Structure of Asf1	52
1.4.2 Asf1-H3/H4 complex structure	54
1.4.3 Asf1 in transcriptional regulation	55
1.4.4 Asf1 in DNA damage response	56
1.4.5 Asf1 in histone modifications	56
2 . Aim of the study	58
3. Results	59
4. Discussion	95
5. References	101

## Acknowledgments

First and foremost, I would like to express my sincerest gratitude to my thesis supervisor **Dr. Ali Hamiche**, for giving me the opportunity to work on this project and for his continual guidance and advice in every aspect of my research. I was privileged to be his PhD student and prepare my thesis under his supervision.

I am extremely grateful to all of my committee members Dr. Stefan Dimitrov, Dr. Mikhail Grigoriev and Dr. Christophe Romier for their valuable feedbacks in evaluating my research. I would also like to sincerely thank Dr. Christian Bronner for his valuable help during my thesis preparation.

I would like to acknowledge my Collaborators Dr. Khalid Quararhni, Dr. Arnaud Depaux and Dr. Christophe Papin for their excellent experimental contribution and scientific interest.

A big thanks to all the members of our laboratory, Dr. Catherine Romain, Dr Philippe Romain, Isabelle for her technical assistance, and Abdul, Arnaud, Shuaib, Julien and Guillaume who have provided support and humor during my PhD. It was a pleasure to work with such a wonderful group.

Additionally, thanks to Surya, Sailendra, Suruchika, Rafi, Abhay, Jorge, Shakti, Arun, and Manohar.

## Abstract

Chromatin acts as a key element that controls access to the DNA. The assembly and disassembly of chromatin structure are of central importance for cellular functions. The main factors that determine chromatin assembly, reconfiguration and dynamics are histone chaperones and chromatin remodeling factors. Asf1 is a highly conserved histone chaperone that facilitates histone deposition and removal during nucleosome assembly and disassembly. Asf1 cooperates with the chromatin assembly factor 1 (CAF-1) to promote replication-dependent chromatin assembly and with HIRA to promote replication-independent chromatin assembly. Mammalian cells possess two isoforms of Asf1; Asf1a and Asf1b. However to date, whether they have individual functional roles has remained elusive. Here, we purified the epitope tagged Asf1a and Asf1b complexes from HeLa cells soluble nuclear extracts and analyzed their specific interacting partners. Mass spectrometry analysis followed by immunoblotting revealed that Asf1a associates exclusively with HP1 $\gamma$  and Asf1b with HP1 $\alpha$ . The selectivity of this interaction can be recapitulated *in vitro* using highly purified recombinant proteins. Mutational analysis revealed that the two Asf1 isoforms discriminate between HP1 $\gamma$  and HP1 $\alpha$  through their divergent C-terminal tail, which selectively recognizes an extended non-conserved Hinge domain encompassing the first 11 amino-acids of either HP1 $\gamma$  or HP1 $\alpha$  chromoshadow domains. The interaction of Asf1 with the Hinge domain has important functional implications and might serve to counteract the binding of non-coding RNA to HP1. Our data elucidate for the first time the functional importance of Asf1 isoforms and reveal specific interactions with the different HP1 isoforms. Asf1a/b may associate with specific HP1 isoforms to promote nucleosome occupancy at defined chromatin loci.

## Abbreviations

ASF1a: anti-silencing function 1a protein

Asf1b: anti-silencing function 1b protein

ATR: Ataxia telangiectasia and Rad3 related

bp : base pare

BrdU: Bromodeoxyuridine

CAF1: chromatin assembly factor 1

CENP-A: centromere protein A

CDK: Cyclin dependent kinase

DAXX: death-domain associated protein

DNA: deoxyribonucleic acid

dNTP: deoxyribonukleotide

DSB: double strand break

DTT: dithiothreitol

FACS: Fluorescence activated cell sorting

GST: Glutathione-S-transferase

HAT 1: histone acetyltransferase 1

HIRA: HIR histone cell cycle regulation defective homolog A

HJURP: holliday junction recognition protein

HP1: heterochromatin protein 1

HSP90: heat shock protein 90

HU: hydroxyurea

IPTG: Isopropyl- $\beta$ -D-thiogalaktopyranosid

IP : Immunoprecipitation

MCM: minichromosome maintenance

NAP-1: nucleosome assembly protein-1

NCP: nucleosome core particle

Np : nucleoplasmin

PCNA: proliferating cell nuclear antigen

PCR: Polymerase chain reaction

PI3K: Phosphoinositide-3 kinase

Pol: Polymerase

Rad53: Radiation sensitive mutant 53

RC: replication-coupled

RI: replication- Independent

Rtt106: regulator of Ty1 transposition protein 106

sNASP: somatic nuclear autoantigenic sperm protein

S.cerevisiae: Saccharomyces cerevisiae

SDS-PAGE: Sodium dodecyl sulfate polyacrylamide gel electrophoresis

S.pombe: Schizosaccharomyces pombe

ssDNA: single-strand DNA

tNASP: testicular nuclear autoantigenic sperm protein

TLK1: Tousled-like kinase 1

TLK2: Tousled-like kinase 2

UBN1: Ubinuclein 1

WT : Wildtype



## List of figures

Figure 1. Nucleosome core particle	2
Figure 2. Hierarchical folding of chromatin	4
Figure 3. Electron Microscope (EM) of cell	6
Figure 4. HP1 domains of interaction with selected partners that are potentially important for the stability of heterochromatin domains	9
Figure 5. Post-translational modifications of core histones	11
Figure 6. Distribution of histone chaperones at mammalian centromeres	28
Figure 7. A view of H3.3-H4 deposition by HIRA	43
Figure 8. A view of H3.1-H4 deposition by CAF-1	46
Figure 9 Structure of the Asf1 N-terminal domain is well-conserved	49

## List of Table

Table1 - Classification of histone chaperones in multiple organisms	30
---	----

# **INTRODUCTION**


## **1.1 Chromatin**

In eukaryotic cells, the genome is packed into chromatin, a hierarchically organized complex composed of DNA, histones and non-histone proteins. Chromatin can exist under different higher structures, in which DNA is more or less accessible to regulatory proteins. The first observations of eukaryotic interphase nuclei have shown the presence of two forms of chromatin; a condensed form named heterochromatin, and a more relaxed one named euchromatin (Lamond and Earnshaw, 1998). Heterochromatin is generally described as transcriptionally inactive, unlike euchromatin, which is associated with the active part of genome. However, all inactive genes and non-transcribed regions of the genome do not appear as heterochromatin. Furthermore, the genome-wide patterns of alternatively condensed chromatin persist through cell generation. Because of this property, chromatin is believed to be a principal carrier of the epigenetic information, which plays an important role in many physiological processes such as development. Aberrations in the maintenance and regulation of alternative chromatin states are being increasingly recognized as important factors in carcinogenesis and other pathologies.

### **1.1.1 Nucleosome**

The fundamental building block of chromatin is the nucleosome, a histone octamer composed of two copies of each core histone protein, H2A, H2B, H3 and H4 with 147bp DNA wrapped in ~1.7 left handed super helical turns (Figure 1.a) (Kornberg, 1974; Luger et al., 1997).

Each core histone displays a characteristic and highly conserved structure formed by three  $\alpha$ -helices linked by two short loops L1 and L2. This “histone fold” is used to form “handshake” arrangements between histones H2A and H2B, and histones H3 and H4 (Figure 1.b).



**Figure 1:** a. Nucleosome core particle: ribbon traces for the eight core histones and 147bp DNA phosphodiester backbones (blue: H3; green: H4; yellow: H2A; red: H2B; brown and turquoise: DNA  
b. Ribbon traces of the H3-H4 histone-fold pair. The  $\langle 1$ -L1- $\langle 2$ -L2- $\langle 3$  structure is shown. (Adapted from Luger et al., 1997)


The protein octamer is divided into four dimers defined by H2A-H2B and H3-H4 pairs. The two H3-H4 dimers interact through a four  $\alpha$ -helices bundle formed by the two H3 histone folds to define the H3-H4 tetramer. Then, each H2A-H2B dimer interacts with the tetramer through a second four  $\alpha$ -helices bundle between H2B and H4 histone-folds (Arents et al., 1991).

All the core histones have N-terminal basically charged tails. These tails represent up to 20% of the mass of the core histones. They show no secondary structures and extend radially beyond the disc-shaped surface of the nucleosome, in position to associate with linker DNA or adjacent nucleosomes (Arents and

Moudrianakis, 1993; Luger et al., 1997). They also form ideal targets for covalent modifications. Two nucleosome core particles are separated by linker DNA varying in length from 10 to 80 bp. Histone H1 locks the linker DNA at the entry and exit points of the nucleosome (Maier et al., 2008; Ramakrishnan, 1997; Widom, 1998). Nucleosomes are connected with one another to form nucleosomal arrays which further fold into less understood higher order chromatin structures.

### **1.1.2 Chromatin higher order structures**

Condensation has been shown to be an intrinsic property of the nucleosome arrays. The presence of linker histone H1 or its variant H5 helps compaction of the chromatin fiber and stabilizes various folding patterns (Carruthers et al., 1998). H1 winds the nucleosomes together into a chromatin fiber that is about 30 nanometers in diameter. These chromatin fibers are looped into a 300 nanometer coiled chromatin fiber, and these coiled chromatin fibers are



**Figure 2:** Hierarchical folding of chromatin (Lodish et.al., 2000)


then wound into 700 nanometer-condensed chromatin. A chromosome that has been observed during cell division has a diameter of 1400 nanometers (Figure 2) (Hamiche and Shuaib, 2012; Woodcock, 2006; Woodcock and Dimitrov, 2001). The chromatin fiber itself undergoes further levels of packaging resulting in compaction by a factor of

~1000 in interphase euchromatin and ~10,000 in heterochromatin (Khorasanizadeh, 2004).

### **1.1.3 Specialized chromatin structures**

#### **1.1.3.1 Euchromatin**

On the basis of microscopic observations, chromatin is divided into two specialized structures, euchromatin and heterochromatin. Euchromatin replicates relatively early in the cell cycle is decondensed during interphase, and is transcriptionally more active than heterochromatin (Figure 3). In general, decondensed euchromatin is hyperacetylated. The acetylation of histone amino-terminal tails reduces the global charge of the tails, and allows them to interact with adjacent nucleosome, which destabilizes chromatin structure and finally facilitates the recruitment of transcription factors and co-factors (Annunziato et al., 1988; Nightingale et al., 1998). Studies have shown striking link between actively transcribed gene in euchromatin, like HSP70, and heterochromatin protein 1 (HP1) (Piacentini et al., 2003). In addition, three euchromatic genes Pros35 (encoding



**Figure 3.** Electron Microscope (EM) of cell, indicating Euchromatin and Heterochromatin regions (Alberts et al., 1994).

the 35-kd subunit of the proteasome), *cdc2* (encoding a cell cycle dependent protein kinase) and *CG5676* (encoding a predicted protein of uncharacterized function), found to be associated with HP1 (Cryderman et al., 2005). Lately, around one hundred euchromatic genes are shown to be associated with HP1, leading towards a new functional link between euchromatin and heterochromatin (Piacentini and Pimpinelli, 2010).

### **1.1.3.2 Heterochromatin**

Heterochromatin is the nuclear material distinguished by its dense pattern of staining throughout the cell cycle. Heterochromatin is not simply a DNA sequence, but rather a chromatin state. Heterochromatin is highly condensed, inaccessible and less transcribed (Figure 3) (Huisinga et al., 2006). Heterochromatin is the layer of different



kinds of information, including chromosomal location, nuclear localization and density of repetitive DNA elements (Birchler et al., 2000; Weiler and Wakimoto, 1995). The chromosomal regions that contain high density of repetitive DNA elements such as clusters of satellite sequences and transposable elements, which are found at centromeres, telomeres and “knobs” are the main targets of heterochromatin formation (Birchler et al., 2000).

The regions remaining condensed through the entire cell cycle are called constitutive heterochromatin. However, heterochromatin is also found at the developmentally regulated loci, where chromatin state can change in response to cellular signals and gene transcription. These regions are referred as facultative heterochromatin. Molecular characterization has led to the identification of several key constituents of heterochromatin. Extensive methylation of histone 3 at Lysine 9 is a characteristic of heterochromatin regions (Richards and Elgin, 2002). Other differences involve protein components of heterochromatin. Heterochromatin protein 1 (HP-1) is found abundantly in heterochromatin regions of fly and mammalian chromosomes (James and Elgin, 1986; Wreggett et al., 1994), and other proteins, such as SU(VAR)3-7 and SU(VAR)3-9 also contribute to the heterochromatic state (Aagaard et al., 1999; Reuter et al., 1990; Wu et al., 2005). However, it should be noted that some genes, termed heterochromatic genes, require the presence of HP1 for their expression (Lu et al., 2000), but the mechanism involved is not yet known.

### **1.1.3.3 Heterochromatin Protein (HP1 $\sphericalangle$ , HP1 $\text{\textcircled{R}}$ , and HP1 $\text{\textcircled{C}}$ )**

Specific modifications of histones are essential epigenetic markers which are heritable changes in gene expression that do not affect the DNA sequence (Strahl and

Allis, 2000). Methylation of lysine 9 in histone H3 is recognized by HP1, which directs the binding of other proteins to control chromatin structure and gene expression (Lachner et al., 2001; Nakayama et al., 2001). Methylation at lysine 9 in histone by Suv39h1 methyltransferase strongly correlates to repressed chromatin state (Rea et al., 2000). HP1 was initially identified in *D. melanogaster* in a screen for nuclear protein that showed a predominant distribution to the chromocentre on polytene chromosomes (James and Elgin, 1986). This highly conserved protein has homologues in various organisms, ranging from *S. pombe* (Swi6) to mammals. In mammals three HP1 isoforms, HP1 $\alpha$ , HP1 $\beta$  (also known as MOD1 or M31) and HP1 $\gamma$  (also known as MOD2 or M32) have been identified (Singh et al., 1991). HP1 proteins are small (around 25 kDa) and contain a conserved N-terminal chromodomain (chromatin organization modifier), followed by variable hinge region and a conserved carboxy-terminal chromoshadow domain (Figure 4).

In general chromodomains are found in numerous proteins, many of them are known to function in chromatin organization and the regulation of gene expression such as polycomb proteins and Suv39h1. The three dimensional structure of both chromodomain (CD) and chromoshadow (CSD) domain of mouse HP1 $\beta$  shows that each domain forms a hydrophobic pocket and can function as protein-binding motif (Ball et al., 1997; Nielsen et al., 2002). HP1 serves as a bridging protein, connecting histones, through interactions with the CD, non-histone chromosomal proteins, through the interactions with the CSD. In mammals, the association of HP1, the retinoblastoma (Rb) protein and Suv39h1 with the cyclin E promoter correlates with gene silencing (Nielsen et al., 2001). Furthermore, HP1 has been implicated in gene repression mediated by Kruppel-associated box (KRAB) zinc finger proteins.




Figure 4. HP1 domains of interaction with selected partners that are potentially important for the stability of heterochromatin domains (Maison et al., 2004).

Taken together these findings suggest that HP1 is recruited to specific gene by protein-protein interactions, resulting in gene silencing by an unknown mechanism. Regarding transcriptional regulators, HP1 interacts with proteins involved in DNA replication and repair. Chromatin assembly factor 1 (CAF1) that assembles histones H3 and H4 into newly replicated DNA in both euchromatic and heterochromatic regions of the genome. Although the significance of CAF1-HP1 interaction is not clear, the data suggest that CAF1 might stabilize heterochromatin structure during chromosomes decondensation and transcription. In mammals, HP1 phosphorylation status changes through the cell cycle. HP1 $\alpha$  and HP1 $\beta$  exhibit increased level of phosphorylation during mitosis (Minc et al., 1999). Clearly, the role of phosphorylation needs further investigation to understand the biological significance of this dynamic process. Recently a detailed study showed that HP1 has preferred binding affinity for trimethylated K9 of histone H3 (H3K9me3) (Fischle et al., 2003b). The histone methyltransferase, Suv39h1 is responsible for (H3K9me3) modification and is able to bind to HP1 (Aagaard et al., 1999). HP1 proteins can bind to RNA *in vitro*, but this binding does not involve the HP1 chromodomain (Muchardt et al., 2002). In addition, the hinge region of HP1 can

bind DNA without obvious sequence specificity (Meehan et al., 2003; Sugimoto et al., 1996). The chromoshadow domain of HP1 functions as a dimerization module and mediates the interaction with either HP1 or HP1 itself (Le Douarin et al., 1996). HP1 interacting partners include the DNA methyltransferases Dnmt1 and Dnmt3a, which are involved in CpG methylation (Fuks et al., 2003). HP1 chromoshadow domain binds the large subunit p150 of chromatin assembly factor-1 (CAF-1) a factor that is implicated in histone deposition during both DNA replication and repair (Mello and Almouzni, 2001; Murzina et al., 1999). Therefore, the role of HP1 is proposed to be dependent on the presence of various binding partners, and it could possibly exchange the partners and contribute to a structure, which can be highly dynamic.

## **1.2 Chromatin dynamics**

The DNA wrapped in nucleosomes is inaccessible to the proteins that bind to it and regulate transcription, replication, recombination, and repair. Therefore remodelers are needed to provide regulated DNA accessibility in packaged regions. There are chromatin-remodeling ATPase, histone modifying enzymes, linker histones, transcription factors and the nature of the DNA sequence are responsible for making the chromatin more or less accessible. Transcription factors, linker histones, and enzymes that act on chromatin may themselves show dynamic interactions with the chromatin template, thus allowing rapid changes of chromatin states (Karpova et al., 2004; Kassabov et al., 2003).

### 1.2.1 Histone post-translational modifications

One major challenge in chromatin biology is connecting particular histone post-translational modifications (HPTMs) with different biological functions and vice versa. Histone post-translational modifications (HPTMs) play a crucial role in changing the chromatin structure by both directly or indirectly activating chromatin remodeling complexes. HPTMs have been shown to control several developmental processes, and are deregulated in many human diseases, including cancer (Mathews et al., 2009). Histones are subjected to a variety of post-translational modifications, including acetylation (Grunstein, 1997; Sterner and Berger, 2000), methylation (Zhang and Reinberg, 2001), ubiquitination (Davie and Murphy, 1990), phosphorylation (Nowak and Corces, 2004), sumoylation (Nathan et al., 2003), ADP ribosylation (Adamietz and Rudolph, 1984), glycosylation (Liebich et al., 1993), biotinylation (Hymes et al., 1995), and carbonylation (Wondrak et al., 2000).



**Figure 5:** Post-translational modifications of core histones (Bhaumik et al., 2007). (ph-Phosphorylation, me-methylation, ac-acetylation, ub- ubiquitination).

One of the best understood histone modifications, in that aspect, is histone acetylation (Grunstein, 1997; Turner, 2000). Most of the post-translational modifications take place within the tail extensions that protrude from the nucleosome. HPTMs might affect electrostatic interactions between the histone tails and DNA to “loosen” chromatin structure. Later, it was proposed that combinations of these modifications might create binding epitopes to recruit other proteins. Recently, more modifications have been found in the structured regions of histones, and these modifications may affect histone-DNA and histone-histone interactions within the nucleosome core (Hansen et al., 1998; Wolffe and Hayes, 1999). Phosphorylation, particularly histones H1 and H3 has long been implicated in chromosome condensation during mitosis (Bradbury, 1992; Koshland and Strunnikov, 1996).

#### **1.2.1.1 Acetylation**

Histone acetylation and deacetylation of the lysine residues present in the histone tails has long been linked to transcriptional activity (Allfrey et al., 1964). Indeed, histone acetylation is strongly associated with active transcription and histone acetylation sites are required for gene activity (Shahbazian and Grunstein, 2007). There are twelve known modification sites two on histone H2A (K5, K9), two on histone H2B (K12, K15), four on histone H3 (K9, K14, K18, K56) and four on histone H4 (K5, K8, K12, K16) (Leszinski et al., 2012; Ohzeki et al., 2012; Roth et al., 2001). It is believed that acetylation may affect chromatin structure as it neutralizes the basic charge of lysine, which may affect the interaction of DNA with histones and nucleosome-nucleosome interactions (Tse et al., 1998). It has been shown that acetylation of H4K16

has a negative effect on the formation of the 30 nm chromatin fiber (Shogren-Knaak et al., 2006). Thus, histone acetylation results in a ‘loosening’ of chromatin structure to allow greater access to transcription factors. Furthermore, histone H3 K9 acetylation in promoter regions is associated with low nucleosome density in the vicinity of transcription start sites (Nishida et al., 2006). Histone acetylation can also function as recognition sites for factors that promote transcription. For example, the bromodomain of BRG1, which is the catalytic subunit of the SWI/SNF chromatin remodeling complex, binds acetylated H4K8, while acetylation of H3K9 and H3K14 is critical for the recruitment of TFIID (Agalioti et al., 2002). Recent studies, which mapped acetylated histones on a genome-wide level, found that acetylation of most lysines in the histone H3 and H4 tails was observed in the 5’ end of coding regions and correlated with active transcription (Kurdistani et al., 2004; Pokholok et al., 2005; Roh et al., 2004; Roh et al., 2007). Furthermore, many inducible genes are marked by histone acetylation even in the inactive state suggesting that the presence of histone acetylation serves to prime these genes for activation at a later stage (Roh et al., 2004).

#### **1.2.1.2 Methylation**

Histones can be methylated either on lysine (K) or arginine (R) residues (Murray, 1964; Patterson and Davies, 1969). Lysine residues can be mono-, di-, or trimethylated (Lachner et al., 2003), whereas arginine residues can only be mono- or dimethylated but di-methylation can occur in a symmetrical or asymmetric configuration (Yuan et al., 2011; Zhang, 2004). Histone H3 can be methylated on a number of lysine sites, which include K4, K9, K27, K36 and K79. Histone H3 can also be methylated on a number of arginine sites: R2, R8, R17 and R26. On histone H4, the main sites of

methylation are K20 and R3. Methylation at H3K4 was first observed in the trout testes (Honda et al., 1975), and several studies have linked it to active gene expression in numerous eukaryotes (Leszinski et al., 2012; Ng et al., 2003; Santos-Rosa et al., 2002; Strahl et al., 1999; von Holt et al., 1989). Heterochromatin in higher eukaryotes is characterized by histone hypoacetylation and H3K9 methylation (Richards and Elgin, 2002). Heterochromatin protein 1 (HP1) was shown to specifically recognize methylated H3K9 via its chromodomain (Lachner et al., 2001; Nakayama et al., 2001). This recognition of H3K9 by HP1 is required for the formation of heterochromatin. However, H3K9 methylation and HP1 binding were recently detected on active genes (Vakoc et al., 2005; Vakoc et al., 2006), suggesting that H3K9 methylation is not limited to inactive regions of chromatin. Methylation of histone H3K27 exhibits some similarities to K9 methylation. Both lysines are found within an ARKS sequence on histone H3 and K27 methylation and are associated with transcriptional silencing (Ringrose et al., 2004). In particular, methylation of H3K27 is characteristic of the inactive X chromosome in female cells (Mak et al., 2002; Wang et al., 2001). Methylation of H3K79 is unusual because this modification lies at the core of the nucleosome rather than on the tail. Global analysis of H3K79 methylation has shown that this modification primarily associates with the coding region of actively transcribed genes (Miao and Natarajan, 2005). H3K79 methylation has also been implicated in DNA repair considering that the checkpoint protein p53BP1 has been shown to bind methylated H3K79 (Martin and Zhang, 2005). H4K20 methylation is connected to transcriptional repression and DNA repair, although very little is known about how it functions in these processes.



### **1.2.1.3 Phosphorylation**

Histone phosphorylation is also linked to the regulation of chromatin structure. The role of histone phosphorylation had been implicated in different processes such as transcription, DNA repair, apoptosis and chromosome condensation (Cheung et al., 2000). Most studies have focused on the role of H3S10 phosphorylation (Johansen and Johansen, 2006). Phosphorylation of this residue was found to occur in tandem with the activation of immediate early genes such as *c-jun* and *c-fos* (Mahadevan et al., 1991), and at activated heat shock genes. Phosphorylation of H3S10 was also observed during chromosome condensation. H3S10 seems to function by regulating a methylation/phosphorylation switch that inhibits HP1 binding to H3K9me3 (Fischle et al., 2003a), and indeed this was shown to be true as phosphorylation of H3S10 is responsible for HP1 dissociation during mitosis (Fischle et al., 2005). These observations suggest a model for how H3S10 phosphorylation functions in the two opposing processes of gene activation and chromosome condensation. During interphase this modification promotes removal of HP1 from specific regions, allowing gene expression. Removal of the phosphorylation mark would therefore promote heterochromatin formation and promote chromatin condensation.

### **1.2.1.4 Ubiquitination**

In human, histones H2A and H2B have been reported to be ubiquitinated, at K119 in H2A and K120 in H2B. Histone H2A was the first ubiquitinated histone to be identified (Goldknopf et al., 1975), and the majority of this modification is the monoubiquitinated form. H2A ubiquitination has been linked to polycomb silencing and

X-chromosome inactivation (Wang et al., 2004). Ubiquitinated H2A at K119 was found on the inactive X-chromosome in females and is correlated with the recruitment of PcG proteins PRC1-like (PRC1-L). The ubiquitin moiety is approximately half the size of a core histone, so it has been suggested that ubiquitination of a nucleosome would impact chromatin folding, thus affecting transcription (Shilatifard, 2006).

#### **1.2.1.5 Sumoylation**

SUMO is a small ubiquitin-related protein of ~100 amino acids, which is capable of being ligated to its target protein. Protein sumoylation is involved in the regulation of transcription factors and components of the transcriptional and often results in transcriptional repression (Hannich et al., 2005; Manza et al., 2004). Histone H4 sumoylation has been reported in mammalian cells and correlates with transcriptional repressive events such as histone deacetylation and HP1 recruitment (Shiio and Eisenman, 2003).

#### **1.2.1.6 ADP-ribosylation**

Mono-ADP ribosylation of histones is linked to DNA repair and cell proliferation (Hassa et al., 2006). Histones are mono-ADP-ribosylated when exposed to DNA damaging agents. This modification has the potential for 'cross-talking' with other modifications such as mono-ADP-ribosylation on H4 occurs preferentially when H4 is acetylated (Golderer and Grobner, 1991). Poly-ADP-ribosylation has not been confirmed on histones but it may play a role in local chromatin compaction.

### **1.2.2 Histone variants**

Additional complexity has recently been added to chromatin dynamics with the renewed interest in alternative versions of the canonical core histones. These paralogues, now known as histone variants, share the same overall structure with the canonical histones, displaying for instance the histone fold, but differ in terms of primary sequence from their canonical relatives. Variant histones have been found in all eukaryotes and seem to play particular roles in the specific regions of the genome. Furthermore, at least some of the variant histones appear to be incorporated at specific sites, replacing canonical histones. The substitution of a canonical histone by a variant will then create a new nucleoprotein complex, with an alternate structure and stability. Some histone variants are interesting candidate for an epigenetic marking of the genome.

#### **1.2.2.1 Histone H1 variants**

Histone H1 has a role in stabilizing both the nucleosome and chromatin higher-order structure. H1 in humans is a family of closely related, single gene encoded proteins, including seven somatic subtypes (H1.1, H1.2, H1.3, H1.4, H1.5, H1.0, and H1X), three testis-specific variants (H1t, H1T2m and H1LS1), and one restricted to oocytes (H1oo) (Henikoff and Ahmad, 2005; Szenker et al., 2011; Yun et al., 2011). Histone H1 is a group of one of the most well studied histone variants till date. They are about 21 to 29 kDa proteins consisting of 194-225 amino acid residues depending on the variants. Among the somatic histone H1 variants, H1.1 to H1.5 are expressed in a

replication-dependent manner, whereas H1.0 and H1X are replication-independent. The H1.1 to H1.5 encoding genes are clustered in a region of chromosome 6 together with the core histone genes, whereas H1X and H1.0 genes are on chromosome 3 and 22 respectively. Change in the relative amounts of the somatic H1 variants and their potential roles during differentiation have been little investigated. The histone H1 consists of a well-conserved hydrophobic GH1 domain but less conserved N- and C-terminal parts. H1 variants are highly similar between human and mouse with at least 90% sequence identity within the GH1 domain (Globular domain of histone H1). GH1 consists of three helices and is primarily involved in DNA binding (Clark et al., 1993; Ramakrishnan et al., 1993; Varga-Weisz et al., 1994). The terminal parts are thought to be involved in binding of non-histone chromatin proteins regulating transcription (Hansen et al., 2006; Zlatanova et al., 2000). The C-terminal tail of the linker histone is involved in chromatin condensation and H1-dependent activation of the apoptotic nucleases (Widlak et al., 2005). The mode of binding of histone H1 to the linker DNA of nucleosome fiber is unclear. Similarly to the core histones, the linker histones are the components of both condensed and transcriptionally active chromatin. Diverse posttranslational modifications such as acetylation, methylation, and phosphorylation are considered to be regulators of histone H1 function in chromatin condensation and chromatin specific proteins.

#### **1.2.2.2 Histone H2A variants**

H2A variants are particularly divergent in regions that are involved in histone-histone contacts or are located as surface residues. Four histone H2A variants have been reported in mammals (H2A.Z, H2A.X, macroH2A, and H2A.Bbd). H2A.Z is highly

conserved throughout evolution and it has a single evolutionary origin and remains distinct from all other H2A variants (Malik and Henikoff, 2003). H2A.Z has a distinct function from all other H2A variants. H2A.Z differs from canonical H2A and other H2A variants mainly in its “docking” domain in the C-terminus and in the L1 loop where two H2A molecules interact (Arents et al., 1991; Luger et al., 1997; Suto et al., 2000). The SWR1 complex, which is an ATP- dependent chromatin-remodeling complex, was the first identified H2A.Z chaperone that mediates H2A.Z deposition. Nap1 mediates the nuclear import of the cytosolic H2A.Z-H2B, whereas Chz1 presents the H2A.Z-H2B dimers to the SWR1 complex for the final deposition. SWR1 replaces H2A-H2B with H2A.Z-H2B in a stepwise manner, one copy of the H2A.Z-H2B dimer at a time, which leads to the formation of intermediate hybrid nucleosomes that contain both H2A and H2A.Z (Luk et al., 2010; Straube et al., 2010). The role of H2A.Z in heterochromatin formation and maintenance is first enriched at pericentromeric heterochromatin and subsequently enriched at other chromatin regions. H2A.Z directly interacts with the pericentromeric heterochromatin protein INCENP *in vivo* (Rangasamy et al., 2003). H2A.Z depletion caused genomic instability and disruption of HP1 localization at the pericentromeric regions, which suggests that HP1 functions and pericentromeric heterochromatin identity are regulated by H2A.Z during the early embryonic development in mice (Rangasamy et al., 2004).

H2A.X variant is present nearly in all eukaryotes and this histone variant has a histone fold domain that is similar to the canonical histone H2A. H2A.X has a unique C-terminal motif termed SQ (E/D)√ (where √ represents a hydrophobic residue) (Talbert and Henikoff, 2010). This motif can become rapidly phosphorylated by the phosphoinositide 3-kinase-like kinases, ataxia telangiectasia mutated (ATM), ataxia telangiectasia and RAD3-related (ATR) and DNA-dependent protein kinase (DNA-PK).

This phosphorylated form is known as  $\text{H2A.X}$ . Phosphorylation of  $\text{H2A.X}$  occurs in response to double-strand breaks (DSBs), at which  $\text{H2A.X}$  helps to recruit DNA repair proteins and histone modifying enzymes (van Attikum and Gasser, 2009).  $\text{H2A.X}$  has also been implicated to play a role in meiosis, growth, tumor suppression, and immune receptor rearrangements (Fernandez-Capetillo et al., 2004; van der Heijden et al., 2007).

MacroH2A is vertebrate specific H2A variant that contains two distinct domains. N-terminal region of macroH2A is similar to canonical H2A and C-terminal region contains large (200 residue) called “the macro domain”, which does not have sequence similarity with any other histones (Malik and Henikoff, 2003; Pehrson and Fried, 1992). In mammals, macro H2A is enriched on the inactive X chromosomes in females and thought to be an epigenetic marker of X-inactivation (Costanzi and Pehrson, 1998; Mermoud et al., 1999; Rasmussen et al., 2000). Several studies have suggested potential mechanisms where macroH2A represses gene expression. It has been shown that the C-terminal macro domain of macroH2A interferes with the binding of transcription factors, and the N-terminal domain is involved in chromatin remodeling via SWI/SNF (Angelov et al., 2003).

H2A.Bbd is the most recently discovered H2A variant, which has a truncated C-terminal docking domain. It was named H2A.Bbd because of its unique genomic distribution, where it is deficient in inactive X chromosomes (bar body deficient). The H2A.Bbd nucleosome binds only 116 base pairs of DNA and is less stable than the canonical nucleosome (Bao et al., 2004; Chadwick and Willard, 2001; Gautier et al., 2004). H2A.Bbd lacks a small acidic region on the nucleosome surface that is involved in transcriptional repression and also lacks K119, which is often ubiquitinated on canonical H2A at transcriptionally inactive regions. The unique subnuclear localization and key features of H2A.Bbd suggest that it is involved in gene activation.

### 1.2.2.3 Histone H3 variants

To date, eight different histone H3 variants have been found in mammals: H3.1, H3.2, H3t, H3.3, H3.5, H3.X, H3.Y, and CENP-A. H3.3, CENP-A, H3.X and H3.Y are somatic histone variants, while H3t and H3.5 are testis specific variants (Hamiche and Shuaib, 2012; Postberg et al., 2010; Wiedemann et al., 2010). On the basis of their incorporation into chromatin they can be categorized into two categories: (i) canonical, replication dependent H3 histones (H3.1 and H3.2) and (ii) replication independent histone H3 variants (H3t, H3.3, CENP-A, H3.X, H3.Y and H3.5) (Hamiche and Shuaib, 2012). The CENP-A histone H3-like variants are centromere specific histones found in all eukaryotes from budding yeast to human. Genetic studies revealed that mammalian CENP-A and its counterparts in other eukaryotes (generally referred to as CenH3) are absolutely required for assembly of the kinetochore to which the spindle microtubules attach at mitosis and meiosis (Blower and Karpen, 2001; Buchwitz et al., 1999; Howman et al., 2000). Centromere is the chromosomal region that directs kinetochore assembly during mitosis in order to facilitate the faithful segregation of sister chromatids. In most eukaryotes, centromeres exist as a single locus on each chromosome, and a chromosome lacking a centromere will fail to segregate properly. Segregation errors lead to aneuploidy, which in turn causes cellular stress and greater genomic instability. The evolutionary conserved mark of centromere is the presence of unique nucleosome in which canonical histone H3 is replaced by CENP-A (Cse4 in budding yeast, Cnp1 in fission yeast, and CID/CenH3 in fruit flies). The essential constitutive centromere components include CENP-A (Howman et al., 2000), CENP-C (or Mif2 in budding yeast) (Fukagawa et al., 1999; Meluh and Koshland, 1995), CENP-

I (or Mis6 in fission yeast) (Goshima et al., 1999; Nishihashi et al., 2002), CENP-H (Fukagawa et al., 2001) and Mis12, which are required for the recruitment of many constitutive centromeric components. Furthermore, heterogeneity of the N-terminal domains in the CENP-A family could be the result of the differences in the chromosome segregation mechanisms between yeast and human.

#### **1.2.2.4 H2B variants**

Unlike the H3 and H2A histones, which have several variants with different functions, no ubiquitously expressed H4 and H2B variants have been reported. Nevertheless, a few tissue-specific H2B isoforms have been reported, including sperm specific H2B variant (spH2B), testis specific H2B variants, TH2B and H2BFWT (Gineitis et al., 2000).

#### **1.2.3 ATP-dependent chromatin remodeling**

ATP-dependent chromatin remodeling enzymes use energy from ATP hydrolysis to remodel nucleosomes (Narlikar et al., 2002). Remodeling enzymes disrupt histone-DNA interactions and they provide nucleosome “sliding”, allowing transcription factors to access the DNA (Becker, 2002). They can also reposition the DNA so that it is accessible on the surface of the histone octamer (Aoyagi et al., 2002). In addition to these activities, remodeling complexes can transfer a histone octamer from one DNA template to another and can cause changes in super-helicity by twisting the DNA, which disrupts histone-DNA interactions (Gavin et al., 2001; Whitehouse et al., 1999). There are four classes of ATP-dependent remodelers, that can be



distinguished by their unique flanking domains: the SWI/SNF (Switch/Sucrose non-fermentable), the CHD (chromodomain and helicase-like domain), the ISWI (imitation SWI), and the INO80 families. All remodeler catalytic subunits share conserved ATPase domain.

#### **1.2.3.1 SWI/SNF family remodelers**

The SWI/SNF family remodelers were initially purified from *S. cerevisiae* and are composed of 8-14 subunits. Most eukaryotes utilize two related SWI/SNF family remodelers, built around two catalytic subunits. The catalytic ATPase includes an HSA (helicase-SANT), a post-HSA, and a C-terminal bromodomain (Mohrmann and Verrijzer, 2005). Other conserved subunits bear additional conserved domains: for example hBAF155/170 (SANT, SWIRM), and hBAF60 (SwiB), and human poly bromo (multiple bromodomains). This family has many activities, and it slides and ejects nucleosomes at many loci for different physiological processes but lacks role in chromatin assembly.

#### **1.2.3.2 ISWI family remodelers**

The ISWI family remodelers contain 2 to 4 subunits. dNURF, dCHRAC, and dACF complexes were initially purified from *D. melanogaster* and later hWICH or hNoRC.(Corona and Tamkun, 2004). Most eukaryotes built multiple ISWI family complexes using one or two different catalytic subunits, with specialized proteins. A set of domains bind to the C-terminus of ISWI family ATPase: aSANT domain (ySWI3, yADA2, hNCoR, hTFIIIB) adjacent to a SLIDE domain (SANT-like ISWI), which

together form a nucleosome recognition module that binds to an unmodified histone tail and DNA (Boyer et al., 2004). These proteins contain many domains including DNA-binding histone fold motifs (hCHRAC 15-17), plant homeodomain (PHD), bromodomains (hBPTF and hACF1) and additional DNA-binding motifs HMGI (Y) for dNURF301.

### **1.2.3.3 CHD family remodelers**

The CHD family remodelers combine 1 to 10 subunits and were first purified from *Xenopus laevis*. Characteristic features include two tandemly arranged chromodomains on the N-terminus of the catalytic subunits. The catalytic subunit is monomeric in lower eukaryotes but can be in large complexes in vertebrates (Marfella and Imbalzano, 2007). Proteins often bear DNA-binding domains and PHD, BRK, CR1-3, and SANT domains. Some CHD remodelers slide or eject nucleosomes to promote transcription.

### **1.2.3.4 INO80 family remodelers**

The INO80 (inositol requiring 80) family remodelers contain more than 10 subunits, include the SWR1-related complexes, and were initially purified from *S. cerevisiae* (Bao and Shen, 2007). Higher orthologs include hINO80, hSRCAP (SNF2-related CREB-activator protein), and p400, which also contain HAT activity. Both yINO80 and ySWRI complexes also contain actin and Arp4. INO80 has diverse functions, including promoting transcriptional activation and DNA repair. Although highly related to INO80, SWRI is unique in its ability to restructure the nucleosome by

removing canonical H2A-H2B dimers and replacing them with H2A.Z-H2B dimers, thereby inserting the histone H2A variant H2A.Z.

#### **1.2.4 Histone chaperones**

Three decades have passed since the discovery of nucleosomes in 1974 and the first isolation of histone chaperone in 1978. Nucleosomes can be reconstituted from DNA and histones by salt gradient dialysis (Axel et al., 1974; Kornberg and Thomas, 1974; Oudet et al., 1975). However the mixtures of histones and DNA are aggregated and precipitated at physiological ionic strength. Laskey et al constructed a cell-free nucleosome assembly system at physiological ionic strength with *Xenopus egg* extracts and used this system to isolate nucleoplasmin as a nucleosome assembly factor that does not require low-molecular-weight cofactors, such as ATP (Laskey et al., 1978). They termed this factor as “histone chaperone” because it prevents nonspecific interactions between DNA and histones and promotes specific interactions that lead to nucleosome assembly (Laskey and Earnshaw, 1980; Laskey et al., 1978). Later work showed that the slow addition of histone made it possible to assemble nucleosomes without the histone chaperone at near physiological ionic strength. Under these conditions, nucleosome assembly could be stimulated by the addition of other chromatin components, like linker histone H1 and non-histone protein HMG (Bonne-Andrea et al., 1984; Drew, 1993), acidic polymers (Stein et al., 1979), and RNA (Nelson et al., 1981).

These studies resulted in the isolation of other histone chaperones and demonstrated that there are various types of histones chaperones (Table 1) (Loyola and Almouzni, 2004). Biochemical studies have revealed that histone chaperones interact not only with histones but also with numerous other chromatin factors.




Figure 6: Distribution of histone chaperones at mammalian centromeres (Boyarchuk et al., 2001).

Histone chaperones include Nucleosome Assembly Protein 1 (NAP1) (Ishimi et al., 1984; Ito et al., 1996), Chromatin Assembly Factor 1 (CAF1), N1/N2 (Bonner, 1975; Kleinschmidt et al., 1986), Nucleoplasmin (Arnan et al., 2003; Laskey et al., 1978; Prado et al., 2004), HIRA (Ray-Gallet et al., 2002), DAXX (Drané et al., 2010) and ASF1 (Tyler et al., 1999; Tyler et al., 2001). Histone chaperones are localized at different sites of the genome (Figure 6). Interactions between histone chaperones and chromatin factors are important for the biochemical and biological functions of histone

Classification of chaperones		Histone chaperones in species of function identification	Histone selectivity	Conservation	Main functions	
Class I: single chaperone	H3-H4	Asf1 ( <i>D.m.</i> ) <sup>126</sup>	Both H3.1-H4 & H3.3-H4	<i>S.c.</i> , Asf1; <i>S.p.</i> , Cia1 <i>A.t.</i> , Sga1 & Sga2; <i>D.m.</i> , Asf1; <i>X.l.</i> , Asf1; <i>M.m.</i> , Asf1a & Asf1b; <i>H.s.</i> , Asf1a & Asf1b	Histone donor for CAF-1 and HIRA	
		Fkbp39p ( <i>S.p.</i> ) <sup>127</sup>	H3-H4	<i>S.c.</i> , Fpr; <i>S.p.</i> , Fkbp39p	rDNA silencing	
		HIRA ( <i>X.l.</i> ) <sup>11</sup>	H3.3-H4	<i>S.c.</i> , HIR1/HIR2; <i>S.p.</i> , Hip1, Sim2 <i>A.t.</i> , HIRA; <i>D.m.</i> , HIRA; <i>X.l.</i> , HIRA; <i>M.m.</i> , HIRA; <i>H.s.</i> , HIRA	Deposition factor independent of DNA synthesis	
		N1/N2 ( <i>X.l.</i> ) <sup>128</sup>	H3-H4 <sup>b</sup>	<i>X.l.</i> , N1/N2; <i>M.m.</i> , tNasp & sNasp; <i>H.s.</i> , tNasp & sNasp	H3-H4 storage in <i>X. laevis</i> oocytes	
		Spt6 ( <i>S.c.</i> ) <sup>129</sup>	H3-H4	<i>S.c.</i> , Spt6; <i>S.p.</i> , Spt6 <i>D.m.</i> , Spt6; <i>X.l.</i> , Spt6; <i>M.m.</i> , Spt6; <i>H.s.</i> , Spt6	Transcription initiation & elongation	
	Rtt106 ( <i>S.c.</i> ) <sup>130</sup>	H3-H4	<i>S.c.</i> , Rtt106; <i>S.p.</i> , SPAC6G9.03c	Heterochromatic silencing		
	H2A-H2B	Nucleoplasmin ( <i>X.l.</i> ) <sup>131</sup> Nucleophosmin (NPM1) ( <i>H.s.</i> ) <sup>132</sup>	H2A-H2B	<i>D.m.</i> , Nip; <i>X.l.</i> , Nucleoplasmin; <i>M.m.</i> , NMP1, NPM2 & NPM3; <i>H.s.</i> , NMP1; NPM2 & NPM3	Storage in <i>X. laevis</i> oocytes, cytosolic-nuclear transport, replication, transcription	
		Chz1 ( <i>S.c.</i> ) <sup>16</sup>	H2AZ-H2B	<i>S.c.</i> , Chz1; <i>H.s.</i> , HIRIP3?	Chz1: H2AZ incorporation by SWR1	
		Nap1 ( <i>X.l.</i> ) <sup>131</sup> Nap1-related proteins: Nap1L2 ( <i>M.m.</i> ) <sup>133</sup> , SET/TAF1b ( <i>H.s.</i> ) <sup>134</sup> , CINAP ( <i>H.s.</i> , <i>M.m.</i> ) <sup>135</sup> and Vps75 ( <i>S.c.</i> ) <sup>136</sup>	H2A-H2B <sup>b</sup>	<i>S.c.</i> , Nap1; <i>S.p.</i> , Nap1 & Nap1.2 <i>A.t.</i> , NRP1 & NRP2; <i>D.m.</i> , Nap1 <i>X.l.</i> , Nap1; <i>M.m.</i> , Nap1; <i>H.s.</i> , Nap1	Cytosolic-nuclear transport Transcription Replication	
		Nucleolin ( <i>H.s.</i> ) <sup>25</sup>	H2A-H2B <sup>c</sup>	<i>S.c.</i> , NSR1 <i>A.t.</i> , Nucleoli; <i>X.l.</i> , Nucleolin <i>M.m.</i> , Nucleolin; <i>H.s.</i> , Nucleolin	Transcription elongation; assist chromatin remodelling	
CAF-1 complex ( <i>H.s.</i> ) <sup>8</sup>		H3-H4	p150 p60 RbAp48 <sup>a</sup>	H3.1-H4	<i>S.c.</i> , Rll2/Cac1; <i>S.p.</i> , SPBC29A10.03C <i>A.t.</i> , Fas1; <i>D.m.</i> , p180; <i>X.l.</i> , p150; <i>M.m.</i> , p150; <i>H.s.</i> , p150	Deposition factor coupled to DNA synthesis: -Replication -Repair
	H3.1-H4			<i>S.c.</i> , Cac2; <i>S.p.</i> , SPAC26H5.03 <i>A.t.</i> , Fas2; <i>D.m.</i> , p150; <i>X.l.</i> , p60; <i>M.m.</i> , p60; <i>H.s.</i> , p60		
	H3-H4			<i>S.c.</i> , Msi1/Cac3; <i>S.p.</i> , Msi16 <i>A.t.</i> , Msi1; <i>D.m.</i> , p55; <i>X.l.</i> , p48; <i>M.m.</i> , RbAp48; <i>H.s.</i> , RbAp48		
FACT complex ( <i>H.s.</i> ) <sup>24</sup>	H2A-H2B	Spt16 SSRP1	H2A-H2B	<i>S.c.</i> , Spt16; <i>S.p.</i> , Spt16 <i>A.t.</i> , Spt16; <i>D.m.</i> , Spt16; <i>X.l.</i> , Spt16; <i>M.m.</i> , Spt16; <i>H.s.</i> , Spt16	Transcription elongation	
			H3-H4 <sup>d</sup>	<i>S.c.</i> , Pob3; <i>S.p.</i> , Pob3 <i>A.t.</i> , SSRP; <i>D.m.</i> , SSRP1; <i>X.l.</i> , SSRP1; <i>M.m.</i> , SSRP1; <i>H.s.</i> , SSRP1		
Class III: within enzymatic complexes	H3-H4	Chaperone	Enzymatic complex			
		Hif1 ( <i>S.c.</i> ) <sup>137</sup>	Hat1p/Hat2 ( <i>S.c.</i> )	H3-H4	<i>S.c.</i> , Hif1	Assist HAT
		Rsf-1 ( <i>H.s.</i> ) <sup>138</sup>	RSF ( <i>H.s.</i> )	H3-H4	<i>H.s.</i> , Rsf-1; <i>M.m.</i> , Rsf-1	Assist RC
	ND/multiple	Arp4 ( <i>S.c.</i> ) <sup>15</sup>	RC INO80 ( <i>S.c.</i> , <i>D.m.</i> , <i>M.m.</i> , <i>H.s.</i> ) RC SWR1 ( <i>S.c.</i> ) HAT NuA4 ( <i>S.c.</i> ) RC+HAT Tip60 ( <i>D.m.</i> , <i>M.m.</i> , <i>H.s.</i> )	ND	<i>S.c.</i> , Arp4; <i>S.p.</i> , Alp5 <i>A.t.</i> , Arp4; <i>D.m.</i> , BAP55; <i>M.m.</i> , BAF53; <i>H.s.</i> , BAF53	Assist RC
			RC SWI/SNF & RSC ( <i>S.c.</i> ): BAF/PBAF & BAP ( <i>D.m.</i> , <i>M.m.</i> , <i>H.s.</i> )	ND	<i>S.c.</i> , Arp7.9 <i>A.t.</i> , Arp7.9; <i>D.m.</i> , BAP55; <i>M.m.</i> , BAF53; <i>H.s.</i> , BAF53	Assist RC
			RC INO80 ( <i>S.c.</i> , <i>D.m.</i> , <i>M.m.</i> , <i>H.s.</i> )	H3-H4	<i>S.c.</i> , Arp8 <i>D.m.</i> , BAP55; <i>M.m.</i> , BAF53; <i>H.s.</i> , BAF53	Assist RC
RC ACF/CHRAC ( <i>S.c.</i> , <i>D.m.</i> , <i>M.m.</i> )			Both H2A-H2B & H3-H4	<i>S.c.</i> , Itc1 <i>D.m.</i> , Act1; <i>M.m.</i> , Act1; <i>H.s.</i> , Act1	Assist RC	
Multiclass chaperone	H3-H4	RbAp48 <sup>a</sup> ( <i>H.s.</i> ) <sup>8</sup>	1. Single chaperone 2. CAF-1 complex 3. RC, HAT, HDAC, PC	Both H3-H4 & CENPA-H4	<i>S.c.</i> , Msi1/Cac3; <i>S.p.</i> , Msi16 <i>A.t.</i> , Msi1; <i>D.m.</i> , p55; <i>X.l.</i> , p48; <i>M.m.</i> , RbAp48; <i>H.s.</i> , RbAp48	1. Centromeric chromatin maintenance 2. Chromatin assembly coupled to DNA synthesis 3. Assist multiple enzymatic activities

Table1 - Classification of histone chaperones in multiple organisms.

chaperones. Histones H3/H4 exist as a dimer when bound to some histone chaperones (Tagami et al., 2004), but they exist as H3/H4 heterotetramers within the nucleosome. The explanation of this apparent discrepancy was explained by the structure of Asf1

bound to a H3/H4 dimer (English et al., 2006; Natsume et al., 2007), showing that Asf1 physically interacts the H3/H4 tetramerization interphase. The fact that Asf1 binds to newly synthesized dimers of H3/H4 indicates that process of chromatin assembly involves the formation of an H3/H4 heterotetramer from two dimers of H3/H4. This has been still a challenging question to get exact mechanism, whether the H3/H4 heterotetramer is formed on the DNA or prior to being incorporated onto the DNA. Chromatin assembly factor 1 (CAF-1) is a histone chaperone acting in cooperation with Asf1. Recent crystal structures of the p55 subunit of *Drosophila melanogaster* CAF-1 and the related protein human RbAp46 has revealed that they interacted with the alpha helix 1 of histone H4 (Murzina et al., 2008; Song et al., 2008). This interacting region of histone H4 is far from the H3/H4 tetramerization interface, rising the possibility that CAF-1 may be able to bind and deposit tetramers of H3/H4 onto DNA.

#### 1.2.4.1 **Histone chaperone N1/N2**

N1 and N2 was isolated in 1975 as proteins that accumulate in the nucleoplasm when *Xenopus* nuclear extracts were injected into *Xenopus laevis* eggs (Bonner, 1975; De Robertis et al., 1978). Later, it was shown that N1 and N2 form a complex with histones and possess histone chaperone activity (Kleinschmidt and Franke, 1982). The DNA sequence of *N1* and *N2* genes are identical, and differences in biochemical and immunological properties have not yet been explained. Mammalian NASP (nuclear autoantigenic sperm protein) and yeast Hif1 (Hat1 interacting factor 1) have sequence similarities to N1/N2 (Ai and Parthun, 2004; O'Rand et al., 1992; Poveda et al., 2004; Welch and O'Rand, 1990). N1/N2 and NASP possess two acidic histone-binding clusters in the N-terminal region and a nuclear localization signal in

the C-terminal region (Batova and O'Rand, 1996; Richardson et al., 2000). There are two types of NASP, a testis-specific form (tNASP) and a somatic form (sNASP). This indicated that NASP is involved in the formation of chromosomes in both somatic and sperm cells. Human NASP is present in the histone H3.1 and H3.3 complexes in HeLa cells (Drane et al., 2010; Tagami et al., 2004), indicating its role in the nucleosome assembly. NASP can bind the linker histone H1 and competes with DNA for histone H1 (Alekseev et al., 2003). In addition, NASP interacts with the heat-shock protein HSP90 and activates its ATPase activity (Alekseev et al., 2005). The DNA binding factors E2F and Sp1 bind to the NASP promoter suggesting that they regulate the expression of NASP in S phase (Richardson et al., 2006; Richardson et al., 2001). During the cell cycle, NASP shuttles between the nucleus and cytoplasm. These results indicate that NASP may transport linker histone H1 into the nucleus in the presence of HSP90 in the S phase and deposit this histone onto replicated DNA (Alekseev et al., 2003).

#### 1.2.4.2 **Histone chaperone NAP1**

Nucleosome assembly protein 1 (Nap1) was originally identified in HeLa cells and is one of few histone chaperones that is capable of recognizing all four core histone proteins (Ishimi et al., 1987). Nap1 is a member of the Nap family of histone chaperone proteins that includes SET/TAF-I $\beta$ , Vps75, as well as the human Nap1 variants Nap1-like proteins 1-5 (Nap1L1-5) and has homologues in yeast, *Drosophila melanogaster* and *Xenopus laevis*. Members of the Nap family of proteins are highly conserved and have a broad range of functions including

transcription activation, assembly and disassembly of nucleosomes, and interaction with other proteins that carry out cellular processes unrelated to chromatin dynamics and transcription activation (Zlatanova et al., 2007). Specifically, Nap1 interacts with cellular proteins that carry out many diverse functions including cell cycle regulation, mediation of nuclear import, control of RNA synthesis and translation, ATPase activity within the mitochondria, lysine biosynthesis, and histone transport and control. In the cell, under physiological conditions, Nap exists as a dimer and does not exist in monomeric form (McBryant and Peersen, 2004). As a chaperone, Nap1 is capable of interacting with both H2A/H2B dimers and H3/H4 tetramers. Generally the ratio of interaction is 1:1, Nap monomer:histone monomer. Specifically, one Nap dimer interacts with one H2A/H2B dimer or two Nap dimers interact with one H3/H4 tetramer (Das et al., 2010; McBryant et al., 2003).

Nap1, have been implicated in maintaining a balance between assembly and disassembly of nucleosomes (Park and Luger, 2006). Previously, Nap1 was known for H2A/H2B dimer removal and exchange (Levchenko and Jackson, 2004; Park et al., 2005). However, evidence suggests a role for Nap1 in disassembly of all four-core histones (Okuwaki et al., 2005). Of significance, the Nap1-dependent disassembly of nucleosomes from the HTLV-1 promoter was the first example of a histone chaperone functioning to disassemble the complete octamer in an acetylation-dependent manner. This reaction occurs in the absence of ATP-dependent chromatin remodeling proteins, transcription initiation, and elongation. This observation was noteworthy as Nap1-mediated disassembly occurs in the absence of ATP, but requires acetyl coenzyme A (acetyl CoA) and the HAT activity of p300 (Sharma and Nyborg, 2008)



### 1.2.4.3 Histone chaperone HIRA

HIRA was first isolated as a protein encoded by a gene within the DiGeorge syndrome critical region of human chromosome 22q11 in 1993 (Halford et al., 1993). HIRA was originally named TUPLE1 because of sequence similarity to the yeast corepressor Tup1 and *Drosophila* E (sp1). Later it was found that TUPLE1 was more similar to Hir1 and Hir2/Spt1, repressor of histone gene transcription in yeast, and therefore it was renamed the protein HIRA (histone regulatory homolog A) (Lamour et al., 1995). HIRA's primary structure is divided into an N-terminal Hir1-like region and C-terminal Hir2-like region. HIR complex is functionally related to the SWI/SNF complex. The ATP-dependent nucleosome remodeling complex SWI/SNF is required for the expression of the histone genes and is recruited to this locus through its interaction with Hir1 and Hir2 (Dimova et al., 1999). Interactions between HIRA and transcription factors like Pax3 and Pax7 (Magnaghi et al., 1998), functional interactions with transcription elongation factors such as Spt4, 5, 6 and 16 (DeSilva et al., 1998), and Spt phenotype of Hir1 and Hir2 mutants (Sherwood et al., 1993) suggest that HIRA, Hir1 and Hir2 are involved in the regulation of the transcription of several classes of genes. HIRA also interacts with CAF-1 p48 and with HDAC2 through the N-terminal WD40 domain and C-terminal LXXLL motif, respectively (Ahmad et al., 2003). HIRA is also known to be required for the recruitment of HP1 to pericentromeres in human cells (Zhang et al., 2007). Human HIRA, yeast Hir1 and Hir2 interact with human and yeast CIA respectively. HIRA selectively binds to histone H3.3 and is required for the deposition of histone H3.3 during sperm nucleus decondensation in fertilized *Drosophila* eggs and mouse

zygote (Loppin et al., 2005; Munakata et al., 2000). These results all indicate that HIRA and CIA are coordinately required for replication-independent nucleosome assembly *in vivo*. Structural and functional analyses of the interaction between HIRA and CIA have identified their interaction surface, which are distinct from the sites of interaction between CIA and the histone H3-H4 dimer (English et al., 2006; Natsume et al., 2007). Further structural and functional analyses of HIRA with the CIA-H3-H4 complex will clarify the mechanisms of nucleosome assembly and disassembly.

#### **1.2.4.4 Histone chaperone DAXX**

DAXX was initially linked to FAS-mediated apoptosis. DAXX was found to colocalize with both the promyelocytic leukaemia (PML) nuclear body and the alpha-thalassemia/mental retardation X-linked syndrome protein (ATRX), which is highly enriched at pericentric heterochromatin (Lewis et al., 2010). Recently, it is shown that DAXX facilitates the deposition of H3.3 with the help of ATRX (Drane et al., 2010; Santenard et al., 2010). DAXX directly interacts with H3.3 both *in vivo* and *in vitro* and mediates deposition of bacterially purified recombinant H3.3-H4 tetramer and naked DNA *in vitro* (Santenard et al., 2010).

Interestingly, the DAXX-ATR complex deposited H3.3 at regions different from the ones that contained H3.3 deposited by HIRA. Genome-wide enrichment studies show HIRA-independent localization of H3.3 at telomeres and transcription factor binding sites (Drane et al., 2010). Surprisingly, CAF1 was found to be associated with H3.3 predeposition complex in the absence of DAXX, suggesting that cells can use replication dependent assembly pathway to counterbalance the loss

of DAXX (Mosammaparast et al., 2002). It seems that DAXX prevents the interaction of H3.3 with CAF-1 complex in order to promote replication independent chromatin assembly of H3.3.

#### **1.2.4.5 Histone chaperone CAF-1**

In 1986, Stillman and colleagues identified an activity that preferentially assembles nucleosomes onto replicating DNA (Stillman, 1986), and purified a factor responsible for this activity and named it “chromatin assembly factor-1” (CAF-1) (Smith and Stillman, 1989). Subsequently, CAF-1 homologues were identified in both *Drosophila* (Kamakaka et al., 1996; Tyler et al., 1996) and yeast (Enomoto et al., 1997). Human CAF-1 (hCAF-1) is composed of three subunits: p150, p60 and p46/p48.

Human p150, *Drosophila* p180, and yeast Cac1/Rlf2 share primary structure homology. The C-terminal region of p150 is involved in a dimer formation and directly interacts with p60, which is essential for CAF-1 activity. Moreover, the N-terminal region of p150 interacts with the DNA polymerase clamp PCNA, and p150 colocalizes at DNA replication foci with PCNA and p60 in the S phase (Krude, 1995; Shibahara and Stillman, 1999). Human p60, which is encoded in the Down’s syndrome region of human chromosome 21, possesses HIRA like amino acid sequence (Blouin et al., 1996; Kaufman et al., 1998; Kirov et al., 1998).

Human p60, *Drosophila* p105, and yeast homologue Cac2 contain a WD40 domain and binds to CIA/Asf1 directly like HIRA (Krawitz et al., 2002; Tang et al., 2006). P60 contains several consensus target sites for various kinases and is phosphorylated by Cyclin/Cdk *in vitro* (Keller and Krude, 2000; Martini et al., 1998).

Active CAF-1 is present in the nucleus during the G1, S, and G2 phases, whereas inactive CAF-1 is present in both nucleus and cytoplasm during the M phase (Marheineke and Krude, 1998). The p60 subunit of CAF-1 is phosphorylated and associates with chromatin in response to UV irradiation, and CAF-1 promotes nucleosome assembly on newly synthesized DNA after UV irradiation. These observations suggest that p60 is a cell cycle and UV-irradiated –dependent regulated subunit of CAF-1.

Human p48, *Drosophila* p55, and *yeast* Cac3/Msi1 contain a WD40 domain like p60. In addition, p48 is present not only in the CAF-1 complex but also in several chromatin-related complexes including HAT1 complex (Martini et al., 1998), the HDAC1 complex (Tyler et al., 1996), the histone methyltransferase complex ESC-E(Z) (Muller et al., 2002), and the ATP-dependent nucleosome remodeling complex NURF (Martinez-Balbas et al., 1998). The p46/p48 subunit of CAF-1 is associated with histone H4 in the absence of other CAF-1 subunits indicating that it links predeposited histones and various chromatin-related complexes (Verreault et al., 1996).

#### **1.2.4.6 Histone chaperone CIA/Asf1**

Sternglanz and colleagues isolated anti-silencing function 1 (Asf1) in yeast genetic screen as a factor whose overexpression disrupts both telomeric and mating-type silencing (Le et al., 1997). Asf1 is called CIA (CCG1-interacting factor A) in *S. pombe*. CIA is composed of two structurally distinct domains, an evolutionarily conserved N-terminal core domain and a divergent C-terminal domain. The N-terminal core domain possesses an immunoglobulin fold structure and is sufficient

for the histone chaperone activity. The C-terminal species-specific region of the yeast homologue is rich in acidic amino acid residues (Le et al., 1997; Padmanabhan et al., 2005; Umehara et al., 2002). However, in the human protein Asf1 the C-terminal region is enriched in serine and threonine residues. Although there is only one gene in yeast and *Drosophila* but there are two CIA genes in vertebrates, plants, and *C. elegans*. The vertebrate proteins are CIA-I/Asf1a and CIA-II/Asf1b (Sillje and Nigg, 2001; Umehara and Horikoshi, 2003), which functionally differ in many aspects. (i) HIRA preferentially interacts with CIA-I (Tagami et al., 2004; Tang et al., 2006; Zhang et al., 2005), (ii) CIA-I is more strongly phosphorylated than CIA-II in the C-terminal serine/threonine-abundant regions (Sillje and Nigg, 2001), and (iii) CIA-I is ubiquitously expressed, but CIA-II is expressed in a tissue-specific manner (Umehara and Horikoshi, 2003). In DNA replication, CIA associates with the histone H3.1 chaperone CAF-1 through its p60 subunit in an evolutionarily conserved manner. In addition, the ATPase complex RFC (replication factor C), which loads the DNA polymerase clamp factor PCNA, directly interacts with CIA and recruits CIA to the DNA *in vitro* (Schulz and Tyler, 2006). During the DNA repair, CIA interacts physically and functionally with the FHA domain of RAD53, which is a conserved checkpoint protein kinase (Emili et al., 2001; Hu et al., 2001). These results all show that CIA is involved in various DNA specific reactions including transcription, DNA replication, and DNA repair. CIA is regulated in the cell cycle dependent manner and phosphorylated by S phase specific kinase Tlk in human, *Arabidopsis*, and *Drosophila* cells (Carrera et al., 2003; Ehsan et al., 2004; Sillje and Nigg, 2001). Disruption of CIA results in a growth defect in *S. cerevisiae* and lethality in *S. pombe* (Tyler et al., 1999). In human fibroblast cells, the formation of senescence-associated heterochromatin foci (SAHF) is driven by CIA-I and HIRA (Ye et al., 2007). Histone

chaperones form extensive interaction network and conjunction with other factors participate in a variety of chromatin transactions and cellular task. Recent studies have emphasized a critical role for histone H3/H4 chaperone ASF1 in orchestrating distinct chromatin modifying activities. Asf1 brings histones to CAF1 and HIRA, thus allowing replication-coupled and replication independent chromatin assembly, respectively (Green et al., 2005; Groth et al., 2007a; Tyler et al., 1999; Tyler et al., 2001). The human protein complex CAF-1, composed of three subunits p150, p60 and p48, is the best characterized histone chaperone to promote nucleosome assembly specifically onto newly replicated DNA (Kaufman et al., 1995). CAF-1 also mediates nucleosome assembly coupled to nucleotide excision repair (NER) (Gaillard et al., 1996) and the repair of single-strand breaks (Moggs et al., 2000). Replication-coupled chromatin assembly also involves a complex of ASF1 with MCM2-7 DNA helicase, which is required for DNA unwinding (Groth et al., 2007a). In conjunction with Rtt109 and CBP/p300 acetyltransferase, ASF1 stimulates chromatin assembly and incorporation of H3K56Ac marks at DNA repair foci (Chen et al., 2008; Das et al., 2009). In addition to chromatin assembly, ASF1 cooperates with SWI/SNF family of ATP-dependent chromatin remodelers and H2A/H2B chaperone FACT to displace nucleosomes from active promoters (Gkikopoulos et al., 2009; Takahata et al., 2009). These findings suggest that CIA/Asf1 is involved in a variety of cell functions including cell proliferations, death, and senescence.

### **1.3 Histone deposition**

The first in chromatin assembly is the deposition of H3/H4 onto the DNA, followed by deposition of the two H2A/H2B dimers and later incorporation of linker

histones such as H1. Incorporation of newly synthesized H3/H4 after DNA replication is mediated by the CAF-1, based on their posttranslational modification pattern (Verreault et al., 1996). A role for CAF-1 in chromatin assembly during replication *in vivo* is suggested by the fact that it binds to the replication-specific histone H3.1 variant but not to the replication-independent histone H3.3 variant (Tagami et al., 2004). Interestingly, depletion of human CAF-1 also activates the DNA damage checkpoint and stalls DNA replication itself (Nabatiyan et al., 2006). The histone chaperone Asf1 appears to deliver the newly synthesized histones to CAF-1 for deposition onto the newly replicated DNA. The transfer of newly synthesized histones from Asf1 to CAF-1 is likely to be mediated via physical interaction between these two histone chaperones (Tyler et al., 2001). The role of Asf1 at the replication fork is clearly important because depletion of human, chicken, and *Drosophila* Asf1 slows down DNA replication (Groth et al., 2005), and slows down DNA unwinding by the MCM helicase in human cells (Groth et al., 2007a). Nucleosome disassembly includes unwinding of the DNA duplex by the MCM2-7 (minichromosome maintenance) helicase complex, followed by passing of the single-stranded DNA through various polymerases. The process of DNA replication is maintained by PCNA (proliferating cell nuclear antigen), a ring like- trimer that encircles the DNA and tethers DNA polymerases to prevent its dissociation. Conceptually, it is difficult to imagine the physical process of threading the DNA through the MCM helicases, PCNA, and DNA polymerases. As such, DNA must be nucleosome free or “naked” while it is replicated. Nearly 250 bp or more of naked DNA resides behind the replication fork (Sogo et al., 1986), while a stretch of approximately 300 bp naked DNA lies ahead of the replication fork (Gasser et al., 1996). Whether this destabilization of nucleosomes ahead of the replication fork is


due to the passage of the replication machinery itself or is mediated by histone chaperones or ATP-dependent chromatin remodelers is not yet known. The loss of the histone octamer from the parental DNA during DNA replication is accompanied by the dissociation of H3/H4 from H2A/H2B. There are candidate histone chaperones for both H2A/H2B and H3/H4 that may disassemble chromatin at the DNA replication fork. However, chromatin assembly and disassembly are so tightly associated with each other during DNA replication that it is very difficult to distinguish between a role for histone chaperone in chromatin assembly and chromatin disassembly. FACT (facilitates chromatin transcription) is an H2A/H2B chaperone during DNA replication. FACT is two subunit complex containing Spt16 (suppressor of Ty) and Pob3 (polymerase one binding) in yeast and SPT16 and SSRP1 (structural specific recognition protein) in humans. FACT is unusual chaperone, which binds with both H2A/H2B and H3/H4 (Stuwe et al., 2008; VanDemark et al., 2008). Human and mouse FACT localize to replication origins (Hertel et al., 1999; Tan et al., 2006), whereas budding yeast FACT interacts with the single-stranded DNA binding protein replication protein A (RPA) (VanDemark et al., 2006), and human FACT interacts with the MCM complex (Tan et al., 2006) The most convincing evidence for a role of Human FACT in chromatin disassembly during DNA replication comes from its ability to promote replication initiation *in vivo* and to promote DNA unwinding by the MCM helicase on nucleosome templates *in vitro* (Tan et al., 2006). Another histone chaperone that could be involved in removing H2A/H2B from DNA is Nap1. Nap1 interacts with H2A/H2B and can assemble chromatin *in vitro* (Ito et al., 1996). Nap1 can also disassemble histone octamers from chromatin in concert with the ATP-dependent chromatin remodeling factor RSC *in vitro* (Lorch et al., 2006). However, Interpretation of *in vivo* studies



becomes complicated when the histone chaperones are involved in multiple pathways, when are they essential, and when are they functionally redundant. In nucleosome disassembly steps, it is clear that first H2A/H2B are removed from DNA and then more stable H3/H4 can be removed. One histone chaperone that has potential to remove the H3/H4 from replicating chromatin is Asf1. Asf1 localizes to replication foci in *Drosophila* S2 cells, but rapidly dissociates when replication is halted in response to hydroxyurea treatment (Schulz and Tyler, 2006). Human Asf1 has been shown to bind to the MCM helicase complex, through histones H3 and H4 and in the absence of Asf1, DNA unwinding by the MCM helicase is greatly reduced *in vivo* (Groth et al., 2007a). To understand the exact mechanism of chromatin disassembly and how the parental pattern of histone variants and histone post-translational modifications are re-established on the daughter strands to ensure epigenetic inheritance, during DNA replication awaits the development of *in vivo* assays for chromatin disassembly during replication.

### **1.3.1 Replication-independent chromatin assembly**

Outside the S phase, histones can be deposited onto DNA by HIRA (HIR complex in *S. cerevisiae* via a replication-independent pathway (Henikoff and Ahmad, 2005). The interaction between Asf1 and Hir was initially found in budding yeast and Asf1 can copurify with all four subunits of HIR (Hir1, Hir2, Hir3 and Hpc2). Similarly, Asf1 forms complexes with histones H3 and H4 as well as HIRA in humans (Figure 7). HIRA preferentially deposits the histone variant H3.3 in nucleosomes (Loppin et al., 2005; Nakayama et al., 2007; Tagami et al., 2004).



**Figure 7:** A view of H3.3-H4 deposition by HIRA. (a) The structure of Asf1-H3/H4 complex shows two binding sites for human ASFa on the histone dimer. Orange triangle indicates HIRA-binding site. (b) hAsf1a-HIRA Ribbon diagram of a HIRA B domain peptide (green) bound to hAsf1a N-terminal domain (gray) (Tang et al, 2006). (c) Asf1a cooperates with HIRA to deposit H3.3.


Since H3.3 is predominantly incorporated into actively transcribed genes (Schwartz and Ahmad, 2005; Wirbelauer et al., 2005), the HIRA/ASF1a complex is thought to mediate transcription-coupled deposition of histone H3.3 (Henikoff, 2008; Tagami et al., 2004). The overexpression of HIRA can also inhibit histone expression and lead to an S-phase arrest (Nelson et al., 2002). The human Asf1 N-terminal domain has been shown to interact with the B-domain of HIRA (Daganzo et al., 2003), in the form of an antiparallel  $\beta$ -hairpin (Figure 7b). The evolutionarily

conserved B-domain of HIRA (425-472) is located in the central portion of the protein. This surface is located on the opposite side of the H3 binding site of Asf1 (Tang et al., 2006). The ASF1 D37R+E39R double mutant disrupts the ASF1a-HIRA interaction, but does not affect the ASF1a-H3 complex (Mousson et al., 2005; Tang et al., 2006).

### **1.3.2 Replication dependent chromatin assembly**

The association of DNA with the histones in the nucleosome makes it difficult to access the DNA by protein molecules. The nucleosome is disassembled into two H2A-H2B dimers and a (H3-H4)<sub>2</sub> tetramer ahead of the moving fork during DNA replication, transcription and repair. Then, the parental histones are relocated behind the replication fork and the full nucleosome density is completed by the deposition of newly synthesized histones (Falbo and Shen, 2006; Groth et al., 2007a; Tagami et al., 2004). In *S. cerevisiae*, passage through S phase in the absence of core histone synthesis results in a loss of viability that cannot be rescued by re-expression of histones in G2 (Kim et al., 1988). The H2A-H2B chaperone FACT has been shown to be associated with the MCM (Minichromosome maintenance) helicase that unwinds DNA in front of the replication fork (Tan et al., 2006). Asf1 is also associated with the MCM helicase, suggesting that Asf1 plays a role in disrupting parental nucleosomes and potentially transferring them onto the nascent DNA behind the fork (Groth et al., 2007a). Asf1 acts in both chromatin assembly and disassembly (Adkins et al., 2004; Adkins et al., 2007b; Korber et al., 2006). A complex of Asf1 in association with the histones H3 and H4 was purified from *Drosophila melanogaster* embryonic extracts as a factor that could assemble nucleosomes onto replicating

DNA in a synergistic manner with CAF-I (Tyler et al., 1999). CAF-I is itself a H3/H4 histone chaperone composed of three subunits that was unified from HeLa cell nuclear extract as a factor that could assemble nucleosomes onto the SV40 DNA *in vitro* (Smith and Stillman, 1989). These biochemical results suggest that Asf1 and CAF-1 cooperate in nucleosome assembly onto newly replicated DNA during the S-phase and Asf1 binds to p60 subunit of CAF-1 in yeast, *Drosophila*, chicken, and human (Krawitz et al., 2002; Mello et al., 2002; Sanematsu et al., 2006; Tang et al., 2006). The way these two histone chaperones cooperate in the nucleosome assembly coupled to DNA replication is still a matter of debate. CAF-1 and Asf1 co-purified in a soluble predeposition complex together with the replication-specific histone H3.1 in HeLa cells (Tagami et al., 2004). CAF-1 is recruited to the replication fork through interaction with cell proliferating nuclear antigen (PCNA) and PCNA is loaded onto DNA by replication factor C (RFC) (Majka and Burgers, 2004; Shibahara and Stillman, 1999). Thus, Asf1 and CAF-I could both be associated with the progressing replication fork and cooperate to deposit H3/H4 onto DNA. Yeast Asf1 and both human isoforms Asf1a and Asf1b can interact with CAF-1 p60, promoting replication-dependent chromatin assembly synergistically with CAF-1 (Figure 8) (Loyola and Almouzni, 2004; Mello et al., 2002; Sharp et al., 2001; Verreault, 2000).



**Figure 8.** A view of H3.1-H4 deposition by CAF-1. (a) The structure of Asf1-H3/H4 complex shows two binding sites for human ASFa on the histone dimer. Orange triangle indicates CAF-1-binding site. (b) Interaction of yeast Asf1 (SpAsf1) with a peptide from the p60 subunit of CAF-1, spCac2 (Malay et al, 2008). (c) Both Asf1a and Asf1b act as histone donors for CAF-1, promoting H3.1 deposition (De Koning et al., 2007).

This pathway ensures that histones are promptly assembled onto newly replicated DNA to minimize the potential for DNA damage, as well as being important for the inheritance of the epigenetic information during DNA replication and repair (Groth et al., 2007a; Henikoff et al., 2004). CAF-1 associates with the replication forks through an interaction with proliferating cell nuclear antigen (PCNA), a component essential for DNA replication and DNA repair (Shibahara and Stillman, 1999; Zhang et al., 2000). CAF-1 is essential in humans, as depletion of p60CAF-1 triggers apoptosis in proliferating cells (Nabatiyan and Krude, 2004). In


contrast to human cells, Asf1 and CAF-1 are not essential for cell viability in *S.cerevisiae*, probably because of the existence of other chaperones for histone H3-H4, such as Rtt106 (Huang et al., 2005). Besides Asf1, CAF-1 has been shown to mediate histone deposition onto DNA, assisted by Rtt106 that binds to CAF-1 as well (Huang et al., 2005). Although CAF-I is not essential in *S. cerevisiae*, its inactivation results in increased sensitivity to UV radiation and reduced silencing of genes adjacent to telomeric DNA (Kaufman et al., 1997).

## **1.4 Dynamic roles of the histone chaperone Asf1**

### **1.4.1 Structure of Asf1**

The Asf1 gene is found at one or two copies per haploid genome in all examined eukaryotes. Asf1a/CIA-I is the major isoform expressed in mammals and appears to be widely expressed, whereas Asf1b/CIA-II shows restricted tissue specific distribution with high-level expression in the testis and thymus, and small level in the intestine and colon (Umehara and Horikoshi, 2003). Both isoforms are similar in sequence (around 70% identical amino acids). In all species, Asf1 consists of a highly conserved 155-amino acid N-terminal domain and a variable C-terminal tail. In human Asf1a, the N-terminal domain contains 23 acidic residues vs only 9 basic ones. Indeed few acidic positions are strictly conserved and are involved in the binding of H3 and HirA (Daganzo et al., 2003; Mousson et al., 2005). Non-conserved residues and the variable C-terminal tail might be involved in species-specific regulation of binding interactions. It is noteworthy that the C-terminal tail of Asf1a,

and to a lesser extent Asf1b, is very rich in serine and threonines, which are the targets of Tousled-like kinases (Sillje and Nigg, 2001). 3-D structure of the functional N-terminal domain of budding yeast Asf1 and human Asf1a were resolved by X-ray crystallography and nuclear magnetic resonance (NMR), respectively. NMR experiments performed on the intact hAsf1a protein revealed that its C-terminal tail is unstructured and highly flexible (Mousson et al., 2005; Padmanabhan et al., 2005). The N-terminal domain containing 155 residues is the highly-conserved core region of Asf1. In contrast, the C-terminal tail is variable, unstructured and flexible (Daganzo et al, 2003; Mousson et al, 2005). In *S. cerevisiae* and *S. pombe*, the Asf1 C-terminal sequence is extremely rich in asparatates and glutamates and this type of tail is common in histone chaperones. In vertebrates, the Asf1 C-terminal sequences are not as rich in acidic residues, but they are phosphorylated by Tousled-like kinases (Tlks). The major Tlks phosphorylation sites are located in the C-terminal part of *Drosophila melanogaster* and human Asf1 within a (D/E)-N-S-(L/M) consensus motif, and both proteins cooperate in control of chromatin dynamics and cell cycle progression. The loss of Tlk activity or mutation of phosphorylation sites of Asf1 results in degradation of Asf1 by both proteasome-dependent and independent pathways (Pilyugin et al., 2009). The N-terminal domain of Asf1 consists of three helical linkers on top of a compact immunoglobulin-like  $\beta$ -sandwich fold. This domain is sufficient for all currently known functions of the full-length protein. Asf1 has a large electro-negative surface potential surrounding one side, and a highly conserved hydrophobic groove that interacts with histone. The 3D structure of the functional N-terminal domain of budding yeast was determined by X-ray crystallography (Figure 9)(Daganzo et al., 2003).



**Figure 9:** Structure of the Asf1 N-terminal domain is well conserved. The superposition between hAsf1a (1-156) (purple) and *S. cerevisiae* Asf1 (green) (Daganzo et al., 2003; Mousson et al., 2005). (b) Ribbon diagram model of the Asf1p N-terminal domain bound to heterodimer histone H3 (cyan) and H4 (green) (English et al, 2006).

Florence Mousson and Françoise Ochsenbein determined the structure of the human Asf1a N-terminal domain by nuclear magnetic resonance spectroscopy (NMR) (Mousson et al., 2005). The structures of yeast and human Asf1 N-terminal domains are quite similar.

#### 1.4.2 Asf1-H3/H4 complex structure

Structure shows that Asf1 binds to a histone H3/H4 heterodimer. The hydrophobic groove of Asf1 binds the histone H3-H4 heterodimer by enveloping the



C-terminus helix of histone H3, thereby blocking the formation of a (H3-H4)<sub>2</sub> heterotetramer (English et al., 2006). Furthermore, the C-terminus of histone H4, that forms a mini- $\beta$  sheet with histone H2A in the nucleosome, undergoes a major conformational change upon binding to Asf1 and adds a  $\beta$  strand to the Asf1  $\beta$  sheet sandwich. Additionally, the non-conserved acidic C-terminal tail of yeast Asf1 may strengthen the interaction between Asf1 and H3/H4 (English et al., 2006; Natsume et al., 2007).

### **1.4.3 Asf1 in transcriptional regulation**

In yeast, Asf1 facilitates chromatin disassembly at the PHO5 promoter to promote transcriptional activation, suggesting that it acts as a histone acceptor (Adkins and Tyler, 2004). Asf1 travels with the transcription machinery and/or rapidly fills in gaps left in nucleosome arrays following passage of RNA polymerase (Schwabish and Struhl, 2006). The nature of Asf1 as an interactor with the TFIID subunit Bdf1 also suggests its participation in transcription control at various RNA polymerase II-dependent gene loci (Zabaronick and Tyler, 2005). Asf1 is also involved in developmental gene expression control by mediating transcriptional repression of NOTCH target genes in *Drosophila melanogaster* (Goodfellow et al., 2007). In yeast, the loss of Asf1 results in impaired cell proliferation and minor defects of gene silencing at telomere and silent mating loci HMR and HML. These effects are greatly enhanced by inactivation of CAF-1, but not by Hir. Asf1 and Hir participate together in a pathway for telomeric silencing that is independent of a pathway dependent on CAF-I (Daganzo et al., 2003). In addition, Asf1 was found to mediate histone H3 eviction and deposition during transcriptional elongation.

Furthermore, Asf1 has been implicated in transcription dependent, but replication-independent histone H3 exchange at promoters, another process which can deposit K56-acetylated H3.

#### **1.4.4 Asf1 in DNA damage response**

Asf1 has been strongly implicated in the cellular response to DNA damage. In budding yeast, Asf1 is bound to the Rad53 DNA damage checkpoint kinase during normal growth but this association is abolished when yeast cells are subjected to genotoxic stress (Emili et al., 2001). Upon genotoxic stress, RAD53 is activated by its phosphorylation on multiple sites through the action of the upstream Mec1 and Tel1 kinases, as well as Rad53 autophosphorylation (Gilbert et al., 2001; Sanchez et al., 1996). This multiple phosphorylated form of Rad53 is not found to be associated with Asf1. Several models have been proposed to explain how Asf1 and Rad51 function in response to DNA damage. The interaction may serve to recruit the Asf1-Rad53 complex to sites of DNA damage. Phosphorylation and activation of Rad53 at these sites would release Asf1 for binding histone H3/H4 and other potential partners.

#### **1.4.5 Asf1 in histone modifications**

Asf1 can affect the PTM state of histones. Asf1 also contributes to the acetylation of H3K9 and can promote trimethylation of H3K36 by Set2 in yeast

(Adkins et al., 2007a). In human U2OS cells, histones bound to Asf1 showed two typical chromatin marks: H4K16Ac and H3K9Me3, giving rise to the hypothesis that Asf1 handles both new and parental histones during DNA replication. Asf1 and H3K56Ac Acetylation of H3K56 is an abundant modification of all newly synthesized H3 in budding yeast, fission yeasts and in *Tetrahymena thermophiles*, but is much less abundant in mammalian cells (Groth et al., 2007a; Jasencakova et al., 2010). H3K56 is located at the DNA entry/exit point on the nucleosome core. Although H3K56 acetylation does not appear to greatly alter the overall structure of the nucleosome, acetylation at this residue increases the plasticity of nucleosomes, which may facilitate access of necessary protein factors to the DNA. Asf1 physically interacts with Rtt109 and is absolutely required for H3K56 acetylation. H3K56 acetylation is required for S-phase chromatin assembly and was proposed to be a critical signal for turning off the DNA damage checkpoint following DNA but this claim is still controversial (Chen et al., 2008). H3K56 acetylation can increase the binding affinity between H3-H4 with CAF-1 and Rtt106 to promote efficient deposition of H3-H4 onto replicating DNA by these two histone chaperones. Asf1 is thought to maintain the integrity of the replisome through H3K56 acetylation. Asf1 has been shown to directly interact with origins of replications and can also associate with components of the replisome (Groth et al., 2007b). It appears that Asf1 and H3K56 acetylation promote the stability of stalled replications forks, contributing to cellular survival upon replication stress.

## **2 Aim of the study**

The presented study addresses the elementary processes that govern the establishment of chromatin structure and its dynamic properties. Since chromatin acts as a key element that controls access to the DNA, the assembly and altering of its structure are of central importance for cellular function. The main factors that determine chromatin assembly, reconfiguration and dynamics are histone chaperones and chromatin remodeling complexes. Here, two of these factors, the chaperone Asf1a and the Asf1b class-remodeling complex were investigated in HeLa cells. Key questions that were addressed include the molecular mechanisms for the biological activity of both factors and their relation to the assembly and dynamics of chromatin in replication independent (RI) and replication coupled (RC) manner.

### 3. RESULTS

# Manuscript

**An extended Hinge domain of heterochromatin protein 1  
discriminates between Asf1a and Asf1b to constitute specific H3-  
H4 deposition complexes**

Vivek Tripathi, Arnaud Depaux, Isabelle Stoll, Christian Bronner and  
Ali Hamiche\*

**An extended Hinge domain of heterochromatin protein 1 discriminates between Asf1a  
and Asf1b to constitute specific H3-H4 deposition complexes**

Vivek Tripathi, Arnaud Depaux, Isabelle Stoll, Christian Bronner and Ali Hamiche\*

Institut de Génétique et de Biologie Moléculaire et Cellulaire, the Centre National de la  
Recherche Scientifique UMR 7104, the Institut National de la Santé et de la Recherche  
Médicale U964, Université de Strasbourg, Parc d'innovation, 1 rue Laurent Fries, 67404  
Illkirch Cedex, France

\*Corresponding author:

Dr Ali Hamiche

IGBMC

1, rue Laurent Fries

67404 Illkirch cedex

E-mail: hamiche@igbmc.fr

Phone: +33 388 653 250

Abbreviations: Asf1, (anti-silencing function 1); CAF-1, chromatin assembly factor 1;  
HP1, heterochromatin protein 1

Keywords: Asf1, CAF-1, chromo domain (CD), chromoshadow domain (CSD), Hinge  
domain (HD), HP1, histone deposition, nucleosome assembly

## Abstract

Asf1 is a highly conserved histone chaperone that facilitates histone deposition and removal during nucleosome assembly and disassembly. Asf1 cooperates with the chromatin assembly factor 1 (CAF-1) to promote replication-dependent chromatin assembly and with HIRA to promote replication-independent chromatin assembly. Mammalian cells possess two isoforms of Asf1; Asf1a and Asf1b. However to date, whether they have individual functional roles has remained elusive. Here, we purified the epitope tagged Asf1a and Asf1b complexes from HeLa cells soluble nuclear extracts and analyzed their specific interacting partners. Mass spectrometry analysis followed by immunoblotting revealed that Asf1a associates exclusively with HP1 $\langle$  and Asf1b with HP1 $\textcircled{R}$ . The selectivity of this interaction can be recapitulated *in vitro* using highly purified recombinant proteins. Mutational analysis revealed that the two Asf1 isoforms discriminate between HP1 $\langle$  and HP1 $\textcircled{R}$  through their divergent C-terminal tail, which selectively recognizes an extended non-conserved Hinge domain encompassing the first 11 amino-acids of either HP1 $\langle$  or HP1 $\textcircled{R}$  chromoshadow domains. The interaction of Asf1 with the Hinge domain has important functional implications and might serve to counteract the binding of non-coding RNA to HP1. Our data elucidate for the first time the functional importance of Asf1 isoforms and reveal specific interactions with the different HP1 isoforms. Asf1a/b may associate with specific HP1 isoforms to promote nucleosome occupancy at defined chromatin loci.

## Introduction

Histone chaperones play essential roles in chromatin dynamics during the different steps of gene transcription, DNA repair and replication. Histone chaperones are responsible for delivery of histones to the site of chromatin assembly and histone deposition onto DNA, histone exchange and removal (Shuaib et al., 2010). The histone chaperone Asf1 (anti-silencing function 1) is important for S-phase progression (Sanematsu et al., 2006; Tyler et al., 1999) and was first identified by its capacity to derepress silencing when overexpressed in yeast (Le et al., 1997). Structural and biochemical studies show that Asf1 binds only to one H3–H4 dimer (English et al., 2006; English et al., 2005; Natsume et al., 2007) thus preventing the formation of H3–H4 tetramer. The role of Asf1 is to provide H3.1-H4 dimers to CAF-1 for replication-dependent nucleosome assembly (Mello et al., 2002; Tyler et al., 1999) and to HIRA to promote replication-independent chromatin assembly (Galvani et al., 2008; Green et al., 2005; Tagami et al., 2004). Asf1 binds histone H3-H4 heterodimers in the cytoplasm and together are imported to the nucleus with thanks to the association with importin-4 with subsequent transfert to downstream chromatin assembly factors (Alvarez et al., 2011; Campos et al., 2010; De Koning et al., 2007; Jasencakova et al., 2010). Asf1 binds canonical S-phase histones H3.1-H4 as well as replacement histones H3.3-H4, which are transferred to CAF-1 and HIRA, respectively (Green et al., 2005; Tagami et al., 2004). However, the precise mechanism by which histones H3.1-H4 and H3.3-H4 are delivered to chromatin are not yet fully understood. For instance, it is not yet known whether Asf1 needs additional factor to fulfill the deposition reaction. However, it is known that



the CAF-1 via its p150 subunit, requires to interact with heterochromatin protein 1 (HP1) for pericentric heterochromatin replication and S-phase progression (Murzina et al., 1999; Quivy et al., 2008).

In yeast, only one form of Asf1 has been identified, in contrast to mammalian cells, which possess two isoforms of Asf1; Asf1a and Asf1b. Although, both isoforms have redundant functions in S-phase histone dynamics, Asf1b appears to be more specific to proliferating cells (Corpet et al., 2011) suggesting that the two isoforms may exert different intimate functions. Asf1a and Asf1b are highly homologous but only Asf1a has been shown to be part of the HIRA-containing complex (Tagami et al., 2004; Zhang et al., 2005) and directly interacts with a central B-domain of HIRA (Tang et al., 2006). Asf1a belongs to a large multicomponent complex containing Ubinuclein 1 (UBN1) and the Calcineurin-binding protein 1 (CABIN1) that is involved in HIRA-dependent H3.3 deposition (Elsaesser and Allis, 2010; Tagami et al., 2004). Therefore, Asf1a appears to be associated to replication-independent nucleosome assembly and thus underlying a possible role outside DNA replication such as gene silencing of proliferating-promoting genes (Galvani et al., 2008; Gazin et al., 2007).

Asf1b is rather specific for proliferating cells and it has been suggested that it acts as the prominent histone acceptor/donor during DNA replication (Corpet et al., 2011). Depleted Asf1b cells show increased nucleus morphology abnormalities, which have been suggested to result from replication defects during mitosis (Corpet et al., 2011; Galvani et al., 2008). Asf1b is a 202 amino-acid long, *i.e.*, 2 amino-acids shorter than Asf1a, but it differs from the latter by 58 amino-acids, most of them being located in the C-terminal part of the protein, suggesting that if there is a difference in functions between both proteins, it will come from the C-terminal part. In spite of this difference, no difference in the partners of Asf1a and Asf1b could be found, which

were NASP, codanin, importin, Ku70, p60CAF-1, p48/p46CAF-1 and HAT1 (Kang et al., 2011). Exceptions are Ubp11, an ubiquitin protease, and S100A8 (a member of the S100 family), that are found to be associated with Asf1b but not with Asf1a (Kang et al., 2011). Recently, data from the literature show some evidence that Asf1 may associate with HP1 to promote nucleosome occupancy at heterochromatic loci (Yamane et al., 2011).

Mammalian cells have three isoforms of HP1 that fulfill different functions: HP1 $\langle$ , HP1 $\otimes$  and HP1 $\odot$  (Kwon and Workman, 2008). HP1 $\langle$  and HP1 $\otimes$  preferentially localize to heterochromatin whereas HP1 $\odot$  is found in euchromatin and functions in gene-specific silencing (Ayyanathan et al., 2003). HP1 $\otimes$  can also be found in euchromatic regions. Indeed, HP1b (*Drosophila* homologue) plays important roles in transcriptional activation and development. It has been suggested that HP1b may counteract HP1a function both in heterochromatin formation and in the transcriptional regulation of euchromatic genes (Kwon and Workman, 2008). HP1 $\langle$  appears to be as an active factor in early DNA damage signaling and repair pathways (Baldeyron et al., 2011). HP1 $\otimes$  explores chromatin for sites of methyl-mark enrichment where it can bind H3 tails from adjacent nucleosomes (Munari et al., 2012).

In search for further, specific partners for each Asf1 isoform, we purified the epitope tagged-Asf1a and the epitope tagged-Asf1b complexes by using the double-immunoaffinity purification method. Among the Asf1a and Asf1b partners we could confirm the already known proteins that co-purified, NASP, HAT1, importin-4 and p150CAF-1. Surprisingly, but highly interesting and relevant, was the discovery that heterochromatin protein 1 alpha (HP1 $\langle$ ) associates with the Asf1a complex and heterochromatin protein 1 beta (HP1 $\otimes$ ) with the Asf1b complex. Furthermore, we observed that each isoform of Asf1 physically interacts with its respective HP1

isoform. We mapped the domain of HP1 $\alpha$  and Asf1a that are involved in this interaction and called them “Extended Hinge” (EH) and HP1 Interacting Domain (HID), respectively. We propose that the interaction between EH and HID domains governs specific roles for Asf1 and HP1 isoforms in the deposition of histone H3-H4 dimers in a spatio-temporal dependent process.

## Results

### *Isolation of the Asf1a and Asf1b specific nuclear soluble complexes and identification of specific partners*

We used the double-immunoaffinity purification method (Drane et al., 2010; Ouararhni et al., 2006) to isolate the epitope tagged Asf1a and Asf1b specific nuclear soluble complexes (e-Asf1a.com, e-Asf1b.com) from HeLa cells. Epitope tagged Asf1a (e-Asf1a) and Asf1b (e-Asf1b) proteins were stably expressed as fusion protein with N-terminal Flag- and HA-epitope-tags in HeLa cells. The epitope tag HA allowed us first to investigate the nuclear localization of each protein. Asf1a expression pattern shows a specific staining throughout the nucleoplasm, forming numerous dots whereas Asf1b shows a more diffuse staining throughout the nucleus (Fig. 1A). Epitope-tagged Asf1a and Asf1b deposition complexes were then purified by double immunoaffinity from HeLa soluble nuclear extracts by sequential immunoprecipitations with anti-Flag antibody followed by anti-HA antibody (Drane et al., 2010; Ouararhni et al., 2006). Proteins associated with Asf1a and Asf1b were separated by SDS-containing 4%-12% polyacrylamide gradient gels and subsequently silver-stained (Fig.1B). Mass spectrometry and immunoblotting analyses allowed the identification of RIF1, MCM2-7, tNASP, sNASP,

MMS22L, NFKBIL2, HAT1, TLK1/2, Codanin-1, RbAp46 (RBBP7), RbAp48 (RBBP4), histones H4 and H3 as components of the Asf1a and Asf1b complexes (Figs. 1B, 1C & 1D). RIF1, Rap1-interacting factor 1 homolog, a critical determinant of the replication-timing program (Cornacchia et al., 2012; Yamazaki et al., 2012). MCM2-7 is the replicative helicase. tNASP and sNASP are two isoforms of NASP, an H1 histone-binding protein that is cell cycle-regulated. tNASP is found in gametes, embryonic cells, and transformed cells whereas sNASP is found in all rapidly dividing somatic cells (Richardson et al., 2000). MMS22L and NFKBIL2 proteins, which are part of the MMS22L-TONSL complex required to maintain genome integrity during DNA replication by promoting homologous recombination-mediated repair of replication fork-associated double-strand breaks (Duro et al., 2010; O'Connell et al., 2010; O'Donnell et al., 2010). HAT1 is a histone acetyltransferase that has been recently reported to preferentially acetylate histone H4 in H3.1-H4 dimer over H3.3-H4 dimer. TLK1/2, serine/threonine-protein kinase tousel-like 1/2, involved in Asf1 phosphorylation (Sillje and Nigg, 2001). Codanin-1, an Asf1 interacting protein regulating S-phase histone supply (Ask et al., 2012). Other proteins have also been identified with very good scores, but surprisingly were more enriched in the Asf1a complex than in the Asf1b complex, such as the HIRA complex (HIRA, Cabin1, Ubinuclein 1 and 2) and the CAF-1 complex (p60CAF-1 and p150CAF-1) (Fig. 1D). To our surprise, we found HP1< in the e-Asf1a complex and HP1® in the e-Asf1b complex, suggesting a specific role for each couple in the histone H3/H4 deposition probably depending on a spatio-temporal process.

*Asf1a and Asf1b physically interacts with HP1< and HP1®, respectively*

Considering that we found the same proteins in each complex, although some of them in different amounts, we hypothesized that each isoform of Asf1 is able to discriminate between the HP1 isoforms through a direct interaction. To address this question, GST-Asf1a or GST-Asf1b fusions together with His-HP1 $\langle$  or His-HP1 $\otimes$  or His-HP1 $\odot$  were coexpressed in bacterial cells. Then, GST-Asf1a or GST-Asf1b, together with their associated proteins, were purified at high salt concentration (1M NaCl) and run on a SDS gel and the gel was stained with colloidal blue. The data obtained clearly show that HP1 $\langle$  binds stoichiometrically to Asf1a (Fig. 2A) but not to Asf1b (Fig. 2B). Conversely, HP1 $\otimes$  does not bind to Asf1a (Fig. 2C) but binds stoichiometrically to Asf1b (Fig. 2D). Asf1a exhibited weak binding to HP1 $\odot$  (Fig. 2E) in contrast to Asf1b, which showed strong binding to HP1  $\odot$  (Fig. 2F). This result is consistent with the sequence alignment of HP1 isoforms (Fig. S1A), showing that HP1 $\odot$  is more closely related to HP1 $\otimes$  than to HP1 $\langle$ , particularly regarding the Hinge domain.

To map the domain of interaction between Asf1a and HP1  $\langle$ , we generated various deletion mutants of HP1 $\langle$  and Asf1a. HP1 $\langle$  (191 amino-acid long) is constituted of a chromodomain (17-71), a Hinge domain (86-108) and a chromoshadow domain (116-176) (Fig. 3A). We constructed five deletion mutants for HP1 $\langle$ ,  $\otimes$ 1(72-191),  $\otimes$ 2(1-115),  $\otimes$ 3(116-191),  $\otimes$ 4(72-176) and  $\otimes$ 5(86-126) (Fig. 3A). These mutants were co-expressed with Asf1a in bacteria and investigated for their binding to the GST-Asf1a fusion protein by a GST pull-down assay. The HP1 $\langle$  mutant  $\otimes$ 1(72-191) was found to be able to bind to Asf1a (Fig. 3B,  $\otimes$ 1) but neither the  $\otimes$ 2(1-116) (Fig. 3B) nor the  $\otimes$ 3(116-191) (Fig. 3B) could bind. Interestingly, when the Hinge domain was present but supplemented with an extension at least up to amino-acid 126, overlapping the beginning of the chromoshadow of HP1, the mutants  $\otimes$ 4(72-176) and  $\otimes$ 5(86-

126) exhibited binding to Asf1a (Figs. 3B, ⊗4 and 3B, ⊗5). These results show that the Hinge domain of HP1< is not sufficient for binding to Asf1a but requires an additional extension up to amino-acid 126, which we call “Extended Hinge” (EH). This finding is consistent with the sequence alignment of the three isoforms of HP1, which show that major difference in the primary sequences lies in the extended Hinge domain (Fig. S1A). Therefore, we suggest that the Hinge domain HP1< has the capacity to discriminate between the Asf1 isoforms. We propose that such mechanism also occurs for the HP1@/Asf1b.

Next, we wanted to map the Asf1a domain involved in the interaction with HP1<. For this, we assumed that the most likely domain is located in the C-terminal part of Asf1a considering that it holds the major difference in the primary sequences of both isoforms. We constructed two deletion mutants of Asf1a namely ⊗1(1-160) and ⊗2(161-204) (Fig. 3F) fused with GST to perform GST pull-down assays. The mutant ⊗1(72-191) lacking the 44 amino-acids of the C-terminal part of Asf1a was unable to bind to HP1<. In contrast, the construct containing the 44 amino-acid fused with the GST exhibits interaction capacity with HP1< showing that it is the C-terminal part of Asf1 that is involved in the interaction. We named this domain of Asf1a the “HP1 Interaction Domain” or “HID” (Fig. S1B).

#### *Reconstitution of the Asf1a, HP1<, CAF-1 and H3/H4 deposition complex*

We next investigated how Asf1a and HP1< are integrated in the CAF-1 complex and assayed to stepwise reconstitute the Asf1a/HP1</CAF-1/H3-H4 deposition complex. We first investigated the ability of Asf1a to bind to the H3.1/H4 dimer and to the H3.3-H4 dimer *in vitro* using a GST pull-down assay. GST-Asf1a was co-expressed in bacteria with H3.1 and H4 or

H3.3 and H4. Glutathione bead-bound proteins were eluted and fractioned on SDS-PAGE and stained with colloidal blue. Asf1a strongly bound to either H3.1-H4 or H3.3-H4 dimers and this interaction resisted to 1M NaCl (Fig. 4A and 4B). When co-expressing additionally HP1 $\zeta$ , we observed that this latter associates with Asf1a, H4 and either with H3.1 (Fig. 4C) or with H3.3 (Fig. 4D). This result shows that HP1 $\zeta$  is a part of the nucleosome assembly complex and is probably involved in an early stage of this process.

The identification of the CAF-1 subunit that interacts with Asf1a and Asf1b, respectively, was the next step. CAF-1 is composed of three subunits, *i.e.* p150, p60 and p48/46. p150CAF-1 and p60CAF-1 were highly specific to the e-H3.1 complex, whereas p46/48 was a component common to e-H3.1 and e-H3.3 complexes (Drane et al., 2010; Tagami et al., 2004). Asf1 is the histone chaperone that is donating new histones to CAF-1. We have assessed the interaction of the three CAF-1 subunits with each Asf1 isoform by using a GST pull-down assays. We have produced and purified GST-Asf1a and GST-Asf1b fusion proteins from bacteria and incubated them with the three different CAF-1 subunits produced in baculovirus (Figs. 5A and 5B). Inputs of GST-Asf1a/b (lane 2), p150 (lane 3), p60 (lane 5) and p48 (lane 7) are shown (Fig. 5A and 5B). Asf1a or Asf1b did not interact with p150 (lane 4) and p48 (lane 8) but did interact with p60 (lane 6). These results show that Asf1a and Asf1b are both able to interact with CAF-1 by binding to the p60 subunit.

Our further challenge was to reconstitute the Asf1a, HP1 $\zeta$ , CAF-1, H3.1-H4 complex in the presence of HP1 $\zeta$  to check whether this latter is a component of the histone predeposition complex. For this, we incubated the three CAF-1 subunits in the presence of GST-Asf1a, HP1 $\zeta$ , H3.1 and H4 and performed a GST pull-down assay. Glutathione bead-bound proteins were eluted and fractioned on SDS-PAGE and stained with colloidal blue (Fig. 5C). The data obtained

show that HP1< associates with the histone predeposition complex, *i.e.*, with H3.1, H4, CAF-1 and Asf1a. Our results suggest that HP1< is associated with Asf1a before being recruited to chromatin.

## **Discussion**

In search for specific partners for each Asf1 isoform, we purified the epitope tagged Asf1a and Asf1b complexes from HeLa cells soluble nuclear extracts and analyzed their specific interacting partners. Our data revealed an unexpected association of Asf1a and Asf1b with the specific HP1 isoforms. Asf1a associates exclusively with HP1< and Asf1b with HP1®. The selectivity of these interactions have important biological implications and could explain how Asf1a/b throughout their association with specific HP1 isoforms could help establishing specialized chromatin structures.

The genome of many examined eucaryotes, with the exception of yeast and fungi, contains two Asf1-encoding genes (Kawai et al., 2001; Munakata et al., 2000). The Asf1 homologues are highly conserved throughout evolution (80% identity) with similar structure; while the carboxyl-termini are very diverse among species (Daganzo et al., 2003; Mousson et al., 2005). The high similarity between Asf1 isoforms causes difficulties in relating functional diversities to sequence variations. The purification and analysis of the two Asf1 isoforms from human cells allowed us to characterize in details their interacting partners. Among the partners, we find some proteins that have already been described as part of the Asf1 complex, *e.g.*, HIRA, HAT1, NASP, MCM2-7, Codanin-1, importin, and CAF-1. No significant differences in the composition of the two complexes could be observed except for one protein, which was HP1 of



which a different isoform was found in each complex. Consistently with this result, it is known that CAF-1 is a partner of Asf1 and of HP1 (Eitoku et al., 2008) thus showing the relevancy of the observation that Asf1 and HP1 are found in the same complex. Highly interesting and novel, we found that HP1< associates with the Asf1a complex and HP1® associates with the Asf1b complex. Considering, that so far the only differences of the Asf1a complex and Asf1b complex are HP1< and HP1®, we hypothesized that each isoform of Asf1 is able to discriminate between the two HP1 isoforms through a putative direct interaction. That is what was investigated and observed *in vitro* by using GST pull down assays. Indeed, we observed that Asf1a through its non-conserved carboxyl-termini directly binds to HP1< whereas Asf1b binds to HP1®. The selectivity of each binding is highlighted by a complete absence of binding of Asf1a to HP1® and *vice versa*. We next asked whether the third HP1 isoform, HP1©, could discriminate between the two Asf1 isoforms. Interestingly, HP1© was found to be able to bind to both Asf1 isoforms with some preference for Asf1b. This result is not surprising given that HP1© is more closely related to HP1® than to HP1< (Fig. S1A).

All HP1 proteins share the same architecture consisting of two structured and conserved domains, a chromodomain (CD) and a chromoshadow domain (CSD), connected by a less conserved Hinge domain (Jacobs and Khorasanizadeh, 2002; Zeng et al., 2010; Fig. S1A). We next, characterized the interaction of Asf1a with the different HP1< domains and identified the specific Asf1 interaction motif. This domain extends from the beginning of the HP1< Hinge domain up to the first 11 amino-acids of the chromoshadow. As a consequence, we called it “Extended Hinge”. To our knowledge, it is the first time that a protein was reported to interact with the Hinge domain of HP1<. So far only RNA has been reported to directly recognize the

Hinge domain. Indeed, it was found that the Hinge domain of HP1< strongly recognized both forward and reverse centromeric RNA probes, whereas the chromo-hinge and the hinge-chromoshadow domains showed lower binding capacities (Maison et al., 2011). Previously, it was suggested that the RNA binding activity of the Hinge domain would be implicated in the targeting of HP1< to chromatin (Muchardt et al., 2002). In *Xenopus laevis*, it has been reported that it is the Hinge domain that is responsible for targeting HP1 to native chromatin (Meehan et al., 2003). Interestingly, the Hinge domain is the site where HP1< can be sumoylated in order to be targeted to pericentromeric chromatin (Maison et al., 2011). This would suggest that sumoylation might be a potent regulatory process for the interaction between Asf1a and HP1<. In the light of our results, we can speculate that HP1 is recruited to chromatin via an Asf1-CAF-1-PCNA mediated chromatin assembly pathway. Our reconstitution experiments of the recombinant CAF-1 complex support this view and suggest that Asf1 is integrated in the CAF-1 complex through its association with p60CAF-1 subunit (Fig. 5). CAF-1 is recruited to replication foci by PCNA through an interaction with p150CAF-1 subunit (Shibahara and Stillman, 1999). We do not exclude that sumoylation would stabilize the Asf1 and HP1< interaction and thus facilitates the recruitment of HP1< to chromatin via PCNA.

We do not yet know the precise significance of the association of Asf1a with HP1< and Asf1b with HP1®. Consistently, with our present data, showing no mixing of HP1< and HP1® in the soluble complexes of Asf1 isoforms, it has been previously shown that HP1< targeting causes the recruitment of endogenous HP1® to the chromatin region and *vice versa* (Verschure et al., 2005). The isoforms of HP1 are all homodimers thanks to the chromoshadow domain, in which each monomer, the chromodomain is binding the H3K9me2 or H3K9me3. HP1< appears to be

as an active factor in early DNA damage signaling and repair pathways (Baldeyron et al., 2011). Considering, that Asf1a appears to be associated to replication-independent nucleosome assembly (Gazin et al., 2007), it is plausible that the Asf1a complex containing HP1<, would be rather involved in DNA damage repair process. It is known that only HP1® is essential for cell viability (Singh, 2010). Thus, we suggest that the Asf1a complex is dispensable for cell viability but indispensable for chromatin repair processes.

It has long been considered that HP1 is stably bound to H3K9me3 epigenetic mark without dynamic exchange, but recent findings have shown that HP1 is highly mobile and forms a stable compartment that is actively renewed (Ayoub et al., 2008; Cheutin et al., 2003; Festenstein et al., 2003). We propose that the renewal of HP1 is governed by the dynamical behavior of the Asf1 complexes and we anticipate that HP1 sumoylation might play an important role in this dynamic. We also suggest that the highly flexible nature of HP1 in its free and bound forms plays an important role in its fundamental biological activity to connect different nuclear processes, as it enables the two other HP1 binding platforms, *i.e.*, the chromo domain and the chromoshadow domain, to be adapted within the dense chromatin architecture to recruit additional proteins.

Given the direct connection between Asf1a/b and HP1</®, it will be interesting to address the role of Asf1 isoforms in tissue specification at different developmental stages. Mouse Asf1a shows a ubiquitous expression pattern, whereas Asf1b is highly expressed in the testis and thymus, with little expression in the small intestine and colon (Umehara and Horikoshi, 2003). These results support the view that the two Asf1 isoforms have different functions. Our results reporting differential interactions between Asf1a/b and HP1</® provide a conceptual framework for understanding the different functions of Asf1a/b in the context of chromatin.

## **Acknowledgments**

This work was supported by grants from CNRS, INSERM, INCA “INCa\_4496”, INCa\_4454, ANR “VariZome”, ANR “Nucleoplat” NT09\_476241, the Association pour la Recherche sur le Cancer and La Fondation pour la Recherche Medicale. V.T. acknowledges the Fondation pour la Recherche Medicale for Financial support.

## **Materials and Methods**

### **Cell lines**

Human Asf1a and Asf1b proteins fused with N-terminal Flag-, HA- and His-epitope tags (e-Asf1a and e-Asf1b) were stably expressed in HeLa cells by retroviral transduction (Ouararhni et al., 2006).

### **Plasmid construction**

Human Asf1a and Asf1b cDNAs were PCR amplified and cloned into the XhoI-NotI sites of pGEX-5X1 (Amersham) to generate pGEX-5X1-Asf1a (1-204) and pGEX-5X1-Asf1b (1-202) constructs. The deletion mutants pGEX-5X1-Asf1a (1-161) and pGEX-5X1-Asf1a (162-204) were obtained by cloning the PCR products amplified from pGEX-5X1-Asf1a (1-204) into the XhoI-NotI sites of the vector.

Human HP1<, HP1®, HP1© cDNAs were PCR amplified from a human cDNA library and cloned into the XhoI-NotI sites of pET28b generating pET28b-HP1<, pET28b-HP1® and pET28b-HP1© full length constructs. The deletion mutants pET28b-HP1< (1-115), pET28b-HP1< (72-191), pET28b-HP1< (116-191), pET28b-HP1< (72-176), pET28b-HP1< (86-126) and

pET28b-HP1< (86-115) were obtained by cloning the PCR products amplified from pET28b-HP1< into the XhoI-NotI sites of the pET28b vector. The deletion mutant pET28b-HP1® (86-115) was obtained by cloning the PCR product amplified from pET28b-HP1® into the XhoI-NotI sites of the pET28b vector.

### **Double-Immunoaffinity purification**

To identify the proteins interacting with Asf1a and Asf1b, we established stable HeLa cell lines expressing a N-terminal FLAG-HA-His epitope tagged Asf1a and Asf1b (e-Asf1a and e-Asf1b). Extracts were prepared using a modification of the Dignam protocol (Dignam, 1990). Briefly, cells were lysed in hypotonic buffer (10 mM Tris-HCl at pH 7.65, 1.5 mM MgCl<sub>2</sub>, 10 mM KCl) and disrupted by Dounce homogenizer. The cytosolic fraction was separated from the pellet by centrifugation at 4°C. The nuclear-soluble fraction was obtained by incubation of the pellet in high-salt buffer in order to get a final NaCl concentration of 300 mM. Tagged proteins were immunoprecipitated with anti-Flag M2-agarose (Sigma), eluted with Flag peptide (0.5 mg/mL), further affinity-purified with anti-HA antibody-conjugated agarose, and eluted with HA peptide (1 mg/mL). The HA and Flag peptides were first buffered with 50 mM Tris-Cl (pH 8.5), then diluted to 4 mg/mL in TGEN 150 buffer (20 mM Tris at pH 7.65, 150 mM NaCl, 3 mM MgCl<sub>2</sub>, 0.1 mM EDTA, 10% glycerol, 0.01% NP40), and stored at -20°C until use. The proteins associated with e-Asf1a and e-Asf1b were separated on 4-12% gradient PAGE containing SDS and silver stained using the Silver Quest kit (Invitrogen). Identification of proteins was carried out using an Orbitrap mass spectrometer (ThermoFinnigan Orbitrap LTQ-Velos) or by Taplin Biological Mass Spectrometry Facility (Harvard Medical School, Boston, MA).

## **Western blotting and antibodies**

Western blotting was carried out using standard protocol and antibodies employed were as follows: monoclonal antibody anti-Flag M2 (Sigma); pan anti-H3 Ct (05-928, Upstate Biotechnologies); polyclonal anti-NASP (ProteinTech Group); anti-MCM (BD Pharmingen); anti-HAT1 (ProteinTech); anti-HP1< and anti-HP1® was produced by the IGBMC facility.

## **Immunofluorescence**

Immunofluorescence was performed using standard procedures on a Leica DMR microscope (Leica) using a 63 x 1.32 NA oil immersion objective. Rat anti-HA antibody (Roche) was used at 1:400 dilution; the secondary antibody used is a goat anti-rat IgG coupled to Alexa Fluor 488 (Molecular Probes) at 1:400 dilution.

## **Expression and purification of proteins**

Appropriate plasmids were transformed in *Escherichia coli* strain BL21-CodonPlus-RIL-pLysS (Stratagene) and induced with 0.5 mM IPTG at 30°C for 3 h. Cells were harvested by centrifugation and processed for protein purification. GST-fusion Asf1a, Asf1b and its deletion mutants were purified as a GST-tagged fusion protein by glutathione-sepharose affinity chromatography (GE healthcare). HP1<, HP1®, HP1© and their deletion mutants containing His-tag was purified using Ni-NTA agarose resin according to the kit protocol (QIAGEN).

Human His-p48CAF-1, FLAG-p60CAF-1 and HA-p150-CAF-1 were cloned in the pFast-bac1 vector and produced in the baculovirus system according to the manufacturer's recommendation (Invitrogen). Recombinant proteins were purified by the respective affinity

chromatography from the cell lysates using standard protocols in TGN 500 buffer (20 mM Tris at pH 7.65, 500 mM NaCl, 3 mM MgCl<sub>2</sub>, 10% glycerol, 0.01% NP40). Purified proteins were gel quantified using BSA as a standard and stored at -80°C until use.

### **Protein–protein interactions**

GST fusion proteins immobilized on glutathione Sepharose were incubated for 2 h at room temperature with recombinant proteins in TGEN buffer (20 mM Tris at pH 7.65, 3 mM MgCl<sub>2</sub>, 0.1 mM EDTA, 10% glycerol, 0.01% NP40) containing 250 mM NaCl. Beads were then washed extensively in TGEN containing 250 mM, 500 mM, or 1 M NaCl. Bound proteins were eluted in SDS sample buffer and fractionated on SDS-PAGE.

To study the interaction between GST-Asf1a and GST-Asf1b with HP1 isoforms, GST-tag constructs were coexpressed with his-tagged HP1 constructs in *Escherichia coli* strain BL21-CodonPlus-RIL-pLysS (Stratagene), as described in (Shuaib et al., 2010).

### **References**

- Alvarez, F., Munoz, F., Schilcher, P., Imhof, A., Almouzni, G., and Loyola, A. (2011). Sequential establishment of marks on soluble histones H3 and H4. *J Biol Chem* 286, 17714-17721.
- Ask, K., Jasencakova, Z., Menard, P., Feng, Y., Almouzni, G., and Groth, A. (2012). Codanin-1, mutated in the anaemic disease CDAI, regulates Asf1 function in S-phase histone supply. *Embo J* 31, 2013-2023.

Ayoub, N., Jeyasekharan, A.D., Bernal, J.A., and Venkitaraman, A.R. (2008). HP1-beta mobilization promotes chromatin changes that initiate the DNA damage response. *Nature* 453, 682-686.

Ayyanathan, K., Lechner, M.S., Bell, P., Maul, G.G., Schultz, D.C., Yamada, Y., Tanaka, K., Torigoe, K., and Rauscher, F.J., 3rd (2003). Regulated recruitment of HP1 to a euchromatic gene induces mitotically heritable, epigenetic gene silencing: a mammalian cell culture model of gene variegation. *Genes Dev* 17, 1855-1869.

Baldeyron, C., Soria, G., Roche, D., Cook, A.J., and Almouzni, G. (2011). HP1alpha recruitment to DNA damage by p150CAF-1 promotes homologous recombination repair. *J Cell Biol* 193, 81-95.

Campos, E.I., Fillingham, J., Li, G., Zheng, H., Voigt, P., Kuo, W.H., Seepany, H., Gao, Z., Day, L.A., Greenblatt, J.F., *et al.* (2010). The program for processing newly synthesized histones H3.1 and H4. *Nat Struct Mol Biol* 17, 1343-1351.

Cheutin, T., McNairn, A.J., Jenuwein, T., Gilbert, D.M., Singh, P.B., and Misteli, T. (2003). Maintenance of stable heterochromatin domains by dynamic HP1 binding. *Science* 299, 721-725.

Cornacchia, D., Dileep, V., Quivy, J.P., Foti, R., Tili, F., Santarella-Mellwig, R., Antony, C., Almouzni, G., Gilbert, D.M., and Buonomo, S.B. (2012). Mouse Rif1 is a key regulator of the replication-timing programme in mammalian cells. *Embo J* 31, 3678-3690.

Corpet, A., De Koning, L., Toedling, J., Savignoni, A., Berger, F., Lemaitre, C., O'Sullivan, R.J., Karlseder, J., Barillot, E., Asselain, B., *et al.* (2011). Asf1b, the necessary Asf1 isoform for proliferation, is predictive of outcome in breast cancer. *Embo J* 30, 480-493.



Daganzo, S.M., Erzberger, J.P., Lam, W.M., Skordalakes, E., Zhang, R., Franco, A.A., Brill, S.J., Adams, P.D., Berger, J.M., and Kaufman, P.D. (2003). Structure and function of the conserved core of histone deposition protein Asf1. *Curr Biol* *13*, 2148-2158.

De Koning, L., Corpet, A., Haber, J.E., and Almouzni, G. (2007). Histone chaperones: an escort network regulating histone traffic. *Nat Struct Mol Biol* *14*, 997-1007.

Dignam, J.D. (1990). Preparation of extracts from higher eukaryotes. *Methods Enzymol* *182*, 194-203.

Drane, P., Ouararhni, K., Depaux, A., Shuaib, M., and Hamiche, A. (2010). The death-associated protein DAXX is a novel histone chaperone involved in the replication-independent deposition of H3.3. *Genes Dev* *24*, 1253-1265.

Duro, E., Lundin, C., Ask, K., Sanchez-Pulido, L., MacArtney, T.J., Toth, R., Ponting, C.P., Groth, A., Helleday, T., and Rouse, J. (2010). Identification of the MMS22L-TONSL complex that promotes homologous recombination. *Mol Cell* *40*, 632-644.

Eitoku, M., Sato, L., Senda, T., and Horikoshi, M. (2008). Histone chaperones: 30 years from isolation to elucidation of the mechanisms of nucleosome assembly and disassembly. *Cell Mol Life Sci* *65*, 414-444.

Elsaesser, S.J., and Allis, C.D. (2010). HIRA and Daxx constitute two independent histone H3.3-containing predeposition complexes. *Cold Spring Harb Symp Quant Biol* *75*, 27-34.

English, C.M., Adkins, M.W., Carson, J.J., Churchill, M.E., and Tyler, J.K. (2006). Structural basis for the histone chaperone activity of Asf1. *Cell* *127*, 495-508.

English, C.M., Maluf, N.K., Tripet, B., Churchill, M.E., and Tyler, J.K. (2005). ASF1 binds to a heterodimer of histones H3 and H4: a two-step mechanism for the assembly of the H3-H4 heterotetramer on DNA. *Biochemistry* *44*, 13673-13682.

Festenstein, R., Pagakis, S.N., Hiragami, K., Lyon, D., Verreault, A., Sekkali, B., and Kioussis, D. (2003). Modulation of heterochromatin protein 1 dynamics in primary Mammalian cells. *Science* 299, 719-721.

Galvani, A., Courbeyrette, R., Agez, M., Ochsenbein, F., Mann, C., and Thuret, J.Y. (2008). In vivo study of the nucleosome assembly functions of ASF1 histone chaperones in human cells. *Mol Cell Biol* 28, 3672-3685.

Gazin, C., Wajapeyee, N., Gobeil, S., Virbasius, C.M., and Green, M.R. (2007). An elaborate pathway required for Ras-mediated epigenetic silencing. *Nature* 449, 1073-1077.

Green, E.M., Antczak, A.J., Bailey, A.O., Franco, A.A., Wu, K.J., Yates, J.R., 3rd, and Kaufman, P.D. (2005). Replication-independent histone deposition by the HIR complex and Asf1. *Curr Biol* 15, 2044-2049.

Jacobs, S.A., and Khorasanizadeh, S. (2002). Structure of HP1 chromodomain bound to a lysine 9-methylated histone H3 tail. *Science* 295, 2080-2083.

Jasencakova, Z., Scharf, A.N., Ask, K., Corpet, A., Imhof, A., Almouzni, G., and Groth, A. (2010). Replication stress interferes with histone recycling and predeposition marking of new histones. *Mol Cell* 37, 736-743.

Kang, B., Pu, M., Hu, G., Wen, W., Dong, Z., Zhao, K., Stillman, B., and Zhang, Z. (2011). Phosphorylation of H4 Ser 47 promotes HIRA-mediated nucleosome assembly. *Genes Dev* 25, 1359-1364.

Kawai, J., Shinagawa, A., Shibata, K., Yoshino, M., Itoh, M., Ishii, Y., Arakawa, T., Hara, A., Fukunishi, Y., Konno, H., *et al.* (2001). Functional annotation of a full-length mouse cDNA collection. *Nature* 409, 685-690.

Kwon, S.H., and Workman, J.L. (2008). The heterochromatin protein 1 (HP1) family: put away a bias toward HP1. *Mol Cells* 26, 217-227.

Le, S., Davis, C., Konopka, J.B., and Sternglanz, R. (1997). Two new S-phase-specific genes from *Saccharomyces cerevisiae*. *Yeast* 13, 1029-1042.

Maison, C., Bailly, D., Roche, D., Montes de Oca, R., Probst, A.V., Vassias, I., Dingli, F., Lombard, B., Loew, D., Quivy, J.P., *et al.* (2011). SUMOylation promotes de novo targeting of HP1alpha to pericentric heterochromatin. *Nat Genet* 43, 220-227.

Meehan, R.R., Kao, C.F., and Pennings, S. (2003). HP1 binding to native chromatin in vitro is determined by the hinge region and not by the chromodomain. *Embo J* 22, 3164-3174.

Mello, J.A., Sillje, H.H., Roche, D.M., Kirschner, D.B., Nigg, E.A., and Almouzni, G. (2002). Human Asf1 and CAF-1 interact and synergize in a repair-coupled nucleosome assembly pathway. *EMBO Rep* 3, 329-334.

Mousson, F., Lautrette, A., Thuret, J.Y., Agez, M., Courbeyrette, R., Amigues, B., Becker, E., Neumann, J.M., Guerois, R., Mann, C., *et al.* (2005). Structural basis for the interaction of Asf1 with histone H3 and its functional implications. *Proc Natl Acad Sci U S A* 102, 5975-5980.

Muchardt, C., Guilleme, M., Seeler, J.S., Trouche, D., Dejean, A., and Yaniv, M. (2002). Coordinated methyl and RNA binding is required for heterochromatin localization of mammalian HP1alpha. *EMBO Rep* 3, 975-981.

Munakata, T., Adachi, N., Yokoyama, N., Kuzuhara, T., and Horikoshi, M. (2000). A human homologue of yeast anti-silencing factor has histone chaperone activity. *Genes Cells* 5, 221-233.

Munari, F., Soeroes, S., Zenn, H.M., Schomburg, A., Kost, N., Schroder, S., Klingberg, R., Rezaei-Ghaleh, N., Stutzer, A., Gelato, K.A., *et al.* (2012). Methylation of Lysine 9 in Histone

H3 Directs Alternative Modes of Highly Dynamic Interaction of Heterochromatin Protein hHP1beta with the Nucleosome. *J Biol Chem* 287, 33756-33765.

Murzina, N., Verreault, A., Laue, E., and Stillman, B. (1999). Heterochromatin dynamics in mouse cells: interaction between chromatin assembly factor 1 and HP1 proteins. *Mol Cell* 4, 529-540.

Natsume, R., Eitoku, M., Akai, Y., Sano, N., Horikoshi, M., and Senda, T. (2007). Structure and function of the histone chaperone CIA/ASF1 complexed with histones H3 and H4. *Nature* 446, 338-341.

O'Connell, B.C., Adamson, B., Lydeard, J.R., Sowa, M.E., Ciccia, A., Bredemeyer, A.L., Schlabach, M., Gygi, S.P., Elledge, S.J., and Harper, J.W. (2010). A genome-wide camptothecin sensitivity screen identifies a mammalian MMS22L-NFKBIL2 complex required for genomic stability. *Mol Cell* 40, 645-657.

O'Donnell, L., Panier, S., Wildenhain, J., Tkach, J.M., Al-Hakim, A., Landry, M.C., Escribano-Diaz, C., Szilard, R.K., Young, J.T., Munro, M., *et al.* (2010). The MMS22L-TONSL complex mediates recovery from replication stress and homologous recombination. *Mol Cell* 40, 619-631.

Ouararhni, K., Hadj-Slimane, R., Ait-Si-Ali, S., Robin, P., Miettton, F., Harel-Bellan, A., Dimitrov, S., and Hamiche, A. (2006). The histone variant mH2A1.1 interferes with transcription by down-regulating PARP-1 enzymatic activity. *Genes Dev* 20, 3324-3336.

Quivy, J.P., Gerard, A., Cook, A.J., Roche, D., and Almouzni, G. (2008). The HP1-p150/CAF-1 interaction is required for pericentric heterochromatin replication and S-phase progression in mouse cells. *Nat Struct Mol Biol* 15, 972-979.

Richardson, R.T., Batova, I.N., Widgren, E.E., Zheng, L.X., Whitfield, M., Marzluff, W.F., and O'Rand, M.G. (2000). Characterization of the histone H1-binding protein, NASP, as a cell cycle-regulated somatic protein. *J Biol Chem* *275*, 30378-30386.

Sanematsu, F., Takami, Y., Barman, H.K., Fukagawa, T., Ono, T., Shibahara, K., and Nakayama, T. (2006). Asf1 is required for viability and chromatin assembly during DNA replication in vertebrate cells. *J Biol Chem* *281*, 13817-13827.

Shibahara, K., and Stillman, B. (1999). Replication-dependent marking of DNA by PCNA facilitates CAF-1-coupled inheritance of chromatin. *Cell* *96*, 575-585.

Shuaib, M., Ouararhni, K., Dimitrov, S., and Hamiche, A. (2010). HJURP binds CENP-A via a highly conserved N-terminal domain and mediates its deposition at centromeres. *Proc Natl Acad Sci U S A* *107*, 1349-1354.

Sillje, H.H., and Nigg, E.A. (2001). Identification of human Asf1 chromatin assembly factors as substrates of Tousled-like kinases. *Curr Biol* *11*, 1068-1073.

Singh, P.B. (2010). HP1 proteins--what is the essential interaction? *Genetika* *46*, 1424-1429.

Tagami, H., Ray-Gallet, D., Almouzni, G., and Nakatani, Y. (2004). Histone H3.1 and H3.3 complexes mediate nucleosome assembly pathways dependent or independent of DNA synthesis. *Cell* *116*, 51-61.

Tang, Y., Poustovoitov, M.V., Zhao, K., Garfinkel, M., Canutescu, A., Dunbrack, R., Adams, P.D., and Marmorstein, R. (2006). Structure of a human ASF1a-HIRA complex and insights into specificity of histone chaperone complex assembly. *Nat Struct Mol Biol* *13*, 921-929.

Tyler, J.K., Adams, C.R., Chen, S.R., Kobayashi, R., Kamakaka, R.T., and Kadonaga, J.T. (1999). The RCAF complex mediates chromatin assembly during DNA replication and repair. *Nature* *402*, 555-560.

Umehara, T., and Horikoshi, M. (2003). Transcription initiation factor IID-interactive histone chaperone CIA-II implicated in mammalian spermatogenesis. *J Biol Chem* 278, 35660-35667.

Verschure, P.J., van der Kraan, I., de Leeuw, W., van der Vlag, J., Carpenter, A.E., Belmont, A.S., and van Driel, R. (2005). In vivo HP1 targeting causes large-scale chromatin condensation and enhanced histone lysine methylation. *Mol Cell Biol* 25, 4552-4564.

Yamane, K., Mizuguchi, T., Cui, B., Zofall, M., Noma, K., and Grewal, S.I. (2011). Asf1/HIRA facilitate global histone deacetylation and associate with HP1 to promote nucleosome occupancy at heterochromatic loci. *Mol Cell* 41, 56-66.

Yamazaki, S., Ishii, A., Kanoh, Y., Oda, M., Nishito, Y., and Masai, H. (2012). Rif1 regulates the replication timing domains on the human genome. *Embo J* 31, 3667-3677.

Zeng, W., Ball, A.R., Jr., and Yokomori, K. (2010). HP1: heterochromatin binding proteins working the genome. *Epigenetics* 5, 287-292.

Zhang, R., Poustovoitov, M.V., Ye, X., Santos, H.A., Chen, W., Daganzo, S.M., Erzberger, J.P., Serebriiskii, I.G., Canutescu, A.A., Dunbrack, R.L., *et al.* (2005). Formation of MacroH2A-containing senescence-associated heterochromatin foci and senescence driven by ASF1a and HIRA. *Dev Cell* 8, 19-30.

## Figure legends

### Figure 1

**Purification and characterization of e-Asf1a and e-Asf1b soluble nuclear complexes from HeLa cells.** (A) Stable expression of e-Asf1a and e-Asf1b in HeLa cells. Cells expressing e-

Asf1a and e-Asf1b and control cells (CTRL) were stained with anti-HA (*right*) and DAPI (*left*). (B) Silver staining of proteins associated with e-Asf1a (lane 3) and e-Asf1b (lane 4) complexes. The complexes containing e-Asf1a (e-Asf1a.com) and e-Asf1b (e-Asf1b.com) were purified by double immunoaffinity from soluble nuclear extracts. Mock, indicate mock purification from non-tagged HeLa cells (*lane 2*) Polypeptides identified by mass spectrometry analysis and the positions of molecular size markers are indicated. (C) Hp1< and Hp1® proteins are respectively specific to the e-Asf1a and e-Asf1b complexes. The e-Asf1a (lane 3) and e-Asf1b (lane 4) complexes were analyzed by immunoblotting with the indicated antibodies. (D) mass spectrometry table indicating proteins identified in the e-Asf1a and e-Asf1b complexes.

## Figure 2

**Selective interactions of Asf1a and Asf1b with the different Hp1 isoforms.** (A) Asf1a physically associates with Hp1< when co-expressed in bacteria. Bacterial cells were either non-induced (NI) or induced (NI) with 0.5 mM IPTG to co-express GST-Asf1a and His-Hp1< (lanes 2 and 3). The Asf1a/Hp1< complex was purified and washed with 1M NaCl (lane 4). The purified material was separated on SDS-PAGE and stained with Colloidal blue. M, indicates the molecular weight marker. (B) Asf1b does not associate with Hp1< when co-expressed in bacteria. Bacterial cells were either non-induced (NI) or induced (I) with 0.5 mM IPTG to co-express GST-Asf1b and His-Hp1< (lanes 2 and 3). The GST-pulled down material was washed with 1 M NaCl, eluted and separated on SDS-PAGE. Different concentrations of Asf1b were loaded on the gel to ascertain the absence of Hp1< in the pull-down (lanes 4-6). M, indicates the

molecular weight marker. (C) Asf1a does not associate with Hp1<sup>®</sup> when co-expressed in bacteria. Bacterial cells were either non-induced (NI) or induced (I) with 0.5 mM IPTG to co-express GST-Asf1a and His-Hp1<sup>®</sup> (lane 2 and 3). The GST-pulled down material was washed with 1 M NaCl, eluted and separated on SDS-PAGE (lane 4). M, indicates the molecular weight marker. (D) Asf1b physically associates with Hp1<sup>®</sup> when co-expressed in bacteria. Bacterial cells were either non-induced (NI) or induced (NI) with 0.5 mM IPTG to co-express GST-Asf1b and His-Hp1<sup>®</sup> (lanes 2 and 3). The Asf1b/Hp1<sup>®</sup> complex was purified and washed with 1M NaCl (lane 4). The purified material was separated on SDS-PAGE and stained with Coloidal blue. M, indicates the molecular weight marker. (E) Asf1a weakly associates with Hp1<sup>©</sup> when co-expressed in bacteria. Bacterial cells were either non-induced (NI) or induced (NI) with 0.5 mM IPTG to co-express GST-Asf1a and His-Hp1<sup>©</sup> (lanes 2 and 3). The Asf1a/Hp1<sup>©</sup> complex was purified and washed with 1M NaCl (lane 4). The purified material was separated on SDS- PAGE and stained with Coloidal blue. M, indicates the molecular weight marker. (E) Asf1b strongly associates with Hp1<sup>©</sup> when co-expressed in bacteria. Bacterial cells were either non- induced (NI) or induced (NI) with 0.5 mM IPTG to co-express GST-Asf1b and His-Hp1<sup>©</sup> (lane 1 and 2). The Asf1b/Hp1<sup>©</sup> complex was purified and washed with 1M NaCl (lane 3). The purified material was separated on SDS-PAGE and stained with Coloidal blue.

### Figure 3

#### **The distal C-terminal tail of Asf1a interacts with the Extended Hinge domain of Hp1<sup>⟨</sup>.**

(A) Schematics of Hp1<sup>⟨</sup> domain organization and mapping. Hp1<sup>⟨</sup> contains three different domains: the N-terminal Chromo domain (CD), the central Hinge Domain (HD) and the C-terminal chromoshadow domain (CSD). (B) Interaction of Asf1a with Hp1<sup>⟨</sup> deletion mutants.



Bacterial cells were either non-induced (NI) or induced (NI) with 0.5 mM IPTG to co-express full-length GST-Asf1a with His-Hp1< deletion mutants ( $\Delta 1$ – $\Delta 5$ ). The purified bound material (lane B) was washed with 1M NaCl, eluted, separated on a SDS-PAGE and stained with colloidal blue. M, indicates the molecular weight marker. (C) Schematics of Asf1a domain mapping. (D) Hp1< interacts with the distal C-terminal tail of Asf1a. Interaction of Hp1< with Asf1a deletion mutants. Bacterial cells were either non-induced (NI) or induced (NI) with 0.5 mM IPTG to co-express full-length His-Hp1< with GST-Asf1a deletion mutants ( $\Delta 1$  and  $\Delta 2$ ). The purified bound material (lane B) was washed with 1M NaCl, eluted, separated on a SDS- PAGE and stained with colloidal blue. M, indicates the molecular weight marker.

#### **Figure 4**

**Reconstitution of the four Asf1a subunits complex.** (A) Asf1a strongly interacts with H3.1. Full-length GST-Asf1a, together with H3.1-H4 histones, were coexpressed and purified from bacteria. The purified material was washed with different salt concentrations, eluted, separated on a SDS-PAGE and stained with colloidal blue. (B) Asf1a strongly interacts with H3.3. Full-length GST-Asf1a, together with H3.3-H4 histones, were coexpressed and purified from bacteria. The purified material was washed with different salt concentrations, eluted, separated on a SDS-PAGE and stained with colloidal blue. (C) Reconstitution of the Asf1a, HP1< and H3.1-H4 deposition complex. Full-length GST-Asf1a, together with Hp1< and H3.1-H4 histones, were coexpressed and purified from bacteria. The purified bound material (bound) was washed with 1M NaCl, eluted, separated on a SDS-PAGE and stained with colloidal blue. M, indicates the molecular weight marker. (D) Reconstitution of the Asf1a, HP1< and H3.3-H4

deposition complex. Full-length GST-Asf1a, together with Hp1 $\gamma$  and H3.3-H4 histones, were coexpressed and purified from bacteria. The purified bound material (bound) was washed with 1M NaCl, eluted, separated on a SDS-PAGE and stained with colloidal blue.

### **Figure 5**

**Reconstitution of the CAF-1 complex.** (A and B) Identification of the CAF-1 subunit interacting with Asf1a and Asf1b. Recombinant GST-Asf1a or GST-Asf1b were incubated with either one of the CAF-1 subunits (p48, p60 and p150). The bound material was purified by GST-pull down, separated on a SDS-PAGE and stained with colloidal blue. M, indicates the molecular weight marker. (B) Reconstitution of the seven CAF-1 subunits complex. The human CAF-1 subunits (p48, p60 and p150) were reconstituted by overexpression of the three subunits in Sf9 cells, mixed with bacterially purified GST-Asf1a four subunits complex (Asf1a, HP1 $\gamma$  and H3.1-H4) and followed by GST purification to reconstitute the seven CAF-1 subunits complex.

### **Figure S1**

(A) Multiple primary sequence alignments of the human Hp1 isoforms. (B) Multiple primary sequence alignments of the human Asf1a and Asf1b isoforms.

Figure 1




Figure 2






Figure 3


A



B



C



D






Figure 4


A



B



C



D




Figure 5





Fig. S1A





Fig. S1B





#### **4. Discussion**

# DISCUSSION AND FUTURE PERSPETIVES

### ***Role of the Interaction of Asf1 with HP1: where does it occur?***

Asf1 is a highly conserved histone chaperone that facilitates histone deposition and removal during nucleosome assembly and disassembly. Histones can be deposited into chromatin either in a replication-dependent manner or in a replication-independent manner. Interestingly, Asf1 is involved in both ways of histone deposition pathways but not well characterized. Asf1 cooperates with the chromatin assembly factor 1 (CAF-1) to promote replication-dependent chromatin assembly and with HIRA to promote replication-independent chromatin assembly. In the nucleus the newly synthesized H3–H4 dimers appeared to remain initially associated with Asf1. Next Asf1 supplies the newly synthesized H3–H4 dimers to CAF-1. The p150 subunit of CAF-1 is recruited to the site of DNA synthesis through direct interaction with proliferating cell nuclear antigen (PCNA) and colocalizes with the replication foci and p60 during S-phase (Moggs et al., 2000). Importantly, CAF-1 was found associated *in vivo* with the replication dependent H3.1 complex and not with the replication independent H3.3 complex, a key finding further demonstrating the direct implication of CAF-1 in replication coupled deposition (Tagami et al., 2004). Asf1 also supplies newly synthesized H3.3-H4 dimers to HIRA for histone deposition in a replication-independent manner. This deposition occurs at gene bodies (Goldberg et al., 2010).

In our thesis work, we could demonstrate that the histone chaperone Asf1a specifically and directly interacts with HP1a and conversely Asf1b specifically and also directly interacts with HP1b. Such specific interaction has never been reported in the literature. This novel and intriguing interaction push us to go deeper in the investigations by identifying the interacting domains on both proteins. All HP1 isoforms exhibit the same characteristic domain organization, but differ in their functions, expression profiles and chromosomal localization. Noteworthy, we could identify a new functional domain in HP1a, that we called “Extended Hinge” (ED). This domain was found to interact with the C-terminal domain of Asf1, which we called

“Heterochromatin protein 1 Interacting Domain” (HID). To our knowledge, it is the first time that a partner of protein nature has been reported to interact with the Hinge domain of HP1a. So far only RNA has been reported to directly recognize the Hinge domain of HP1. Indeed, it was found that the Hinge domain of HP1a strongly recognized both forward and reverse centromeric RNA probes, whereas the chromo-hinge and the hinge-chromoshadow domains showed lower binding capacities (Maison et al., 2011). Previously, it was suggested that the RNA binding activity of the Hinge domain would be implicated in the targeting of Hp1a to chromatin (Muchardt et al., 2002). In *Xenopus laevis*, it has been reported that it is the Hinge domain that is responsible for targeting Hp1 to native chromatin (Meehan et al., 2003). According to our results, one might imagine that the Hinge domain facilitates HP1a to be targeted to chromatin via an interaction with HID domain of Asf1a and probably Asf1b facilitates HP1b to be recruited by the chromatin. Altogether, these results support a key regulatory role for the Hinge domain of HP1 in the regulation of gene transcription and genome integrity.

Asf1a, but not Asf1b has been shown to be a part of the HIRA-containing complex (Tagami et al., 2004; Zhang et al., 2005). This would mean that Asf1b in association with HP1b is never found in HIRA complexes. However, it is known that HP1b co-localises with HIRA complexes in human cells (Zhang et al., 2007). Our results show that Asf1b is found in HIRA complexes, by demonstrating that HIRA is enriched in the Asf1a complex *versus* the Asf1b complex. HIRA mediates H3.3 deposition at pericentromeric regions to ensure, with the help of HP1b, kinetochore structure integrity (Zhang et al., 2007). We suggest that HP1a can also be involved in the maintenance of kinetochore structure integrity considering that we found HIRA and HP1a in the Asf1a complex. Although, HP1a and HP1b localize together at heterochromatic loci, microscopy analysis showed that both isoforms do not exactly overlap (Dialynas et al., 2007) suggesting that they do not associate and that they may have different functions in the pericentromeric chromatin. We propose that these different functions may be set by

Asf1 isoforms, which discriminates between HP1 isoforms during the constitution of the Asf1 complexes.

Concerning HP1g, we could show that HP1g has a preference for Asf1b. HP1g is found in euchromatin and function in gene-specific silencing (Ayyanathan et al., 2003). Our results suggest that HP1g may be brought to euchromatin by the Asf1b complex. It has been observed that HP1g plays an unexpected role in the activation of gene transcription that is dependent on its association with RNA polymerase II (Vakoc et al., 2005). Therefore, our results support the idea that it is the Asf1b complex that might associate with RNA polymerase II in euchromatin to ensure transcription activation of specific genes. However, this requires further investigations.

### ***Interaction of Asf1 with HP1: when does it occur?***

Unfortunately, so far we have not yet deciphered the role of each isoform specific interaction in the histone deposition and transcription regulation during the cell cycle, which would have helped us to understand specific histone deposition mechanisms. Nevertheless, we will discuss now when the Asf1/HP1 association might occur.

The first step in the deposition of the newly synthesized histones is the transport from the cytoplasm to the nucleus, a process, which is assisted by distinct chaperones. The chaperone Asf1 was the first chaperone identified to play a key role in supplying histones H3–H4 to the downstream chaperones, like CAF-1 and HIRA for nucleosome assembly. Structural and biochemical studies show that Asf1 binds only to one H3–H4 dimer, thus preventing the formation of H3–H4 tetramer. In addition, the cytosolic complex of ectopically expressed epitope-tagged H3.1 (e-H3.1) contains only the tagged H3.1 fusion, but not the endogenous H3. These data evidence that Asf1 is associated with a single H3–H4 dimer both *in vitro* and *in vivo* and that the H3–H4 dimer, and not the (H3–H4)<sub>2</sub> tetramer, is further delivered to the nucleus. In two very recent studies, the

biochemical purification of the cytoplasmic H3 complex has allowed both the identification of distinct H3 chaperones and the suggestion of a comprehensive mechanism(s) for the sequential assembly of H3–H4 dimers. The first study, carried out by the group of D. Reinberg, identified six different H3 chaperones (HSC70, HSP90, tNASP, sNASP, RbAp46, and Asf1b) along with histone H4, importin4 and HAT1. This study has not detected Asf1a as a part of the cytosolic H3 complex. Four distinct cytosolic complexes were found to be sequentially involved in the assembly of H3–H4 dimers (Campos et al., 2010). After synthesis, histones H3 and H4 were sheltered from misfolding and aggregation by interaction with chaperones HSC70 and HSP90, respectively. For transport and deposition onto DNA, histones H3–H4 first assembled to form the dimer, a process facilitated by HSP90 and tNASP. Once the H3–H4 dimer was assembled, RbAp46 associates with the H4 carboxyl domain and helps the recruitment of HAT1, which in turn acetylates H4 on Lys5 and Lys12. Then the acetylated histones are transferred to Asf1b and importin4 for nuclear transport (Campos et al., 2010). The other study, carried out by the group of A. Loyola, reported some different results. First, they described an interesting finding concerning the free soluble H3–H4 dimers. Loyola and colleagues showed that the free soluble endogenous histones H3 and H4 were transiently poly-ADPribosylated and that this mark was removed after dimer formation (Alvarez et al., 2011). Moreover, the two distinct translocation complexes contained H3–H4 dimers with distinct pre-deposition marks. The first translocation complex contained Asf1a, H3K9me1, and H3K14ac, while the second complex contained Asf1b and H3K9me1. This study also claimed that both H3 acetylation and methylation do not affect nuclear translocation, but may be instead linked to the nucleosomal histone H3 PTM patterns (Alvarez et al., 2011). Thus, it is possible that HP1a as well as HP1b may associate with the each isoform of Asf1 before nuclear import.

It has been recently observed that Asf1a accumulated inside the nucleoli of proliferating C7 and C8 cells (Jiang et al., 2011). This accumulation was suggested to be

associated with cell growth and the release of Asf1a from nucleoli with growth arrest. Accordingly, we might propose that Asf1a is playing such a role with the help of HP1a. In contrast, Asf1b had much less nucleolar localization than Asf1a. Asf1a is involved in both DNA replication-dependent and DNA replication-independent nucleosome assembly processes whereas Asf1b is only involved in DNA replication-dependent (Tagami et al., 2004; Tang et al., 2006; Zhang et al., 2005; Mello et al., 2002). As a consequence, considering our results showing that each Asf1 isoform interacts with a different HP1 isoform, we propose that HP1a would be involved in both DNA replication-dependent and DNA replication-independent, while HP1b would only be involved in DNA replication-dependent.

### **Concluding remarks**

It has been recently questioned about the factors that allow each isoform of HP1 to ensure a specific function in spite of the fact that they share similar architecture organization (Kwon and Workman, 2011). We suggest that the specificity lies in the role of the “Extended hinge” domain that is able to discriminate between Asf1a and Asf1b. Therefore, we propose that it is the composition of Asf1a and Asf1b complexes that governs the targeting of each HP1 isoform. Of course this requires further investigations and several studies are needed to address this question.

## 5. References

- Aagaard, L., G. Laible, P. Selenko, M. Schmid, R. Dorn, G. Schotta, S. Kuhfittig, A. Wolf, A. Lebersorger, P.B. Singh, G. Reuter, and T. Jenuwein. 1999. Functional mammalian homologues of the *Drosophila* PEV-modifier Su(var)3-9 encode centromere-associated proteins which complex with the heterochromatin component M31. *EMBO J.* 18:1923-1938.
- Adamietz, P., and A. Rudolph. 1984. ADP-ribosylation of nuclear proteins in vivo. Identification of histone H2B as a major acceptor for mono- and poly(ADP-ribose) in dimethyl sulfate-treated hepatoma AH 7974 cells. *J Biol Chem.* 259:6841-6846.
- Adkins, M.W., J.J. Carson, C.M. English, C.J. Ramey, and J.K. Tyler. 2007a. The histone chaperone anti-silencing function 1 stimulates the acetylation of newly synthesized histone H3 in S-phase. *J Biol Chem.* 282:1334-1340.
- Adkins, M.W., S.R. Howar, and J.K. Tyler. 2004. Chromatin disassembly mediated by the histone chaperone Asf1 is essential for transcriptional activation of the yeast PHO5 and PHO8 genes. *Mol Cell.* 14:657-666.
- Adkins, M.W., and J.K. Tyler. 2004. The histone chaperone Asf1p mediates global chromatin disassembly in vivo. *J Biol Chem.* 279:52069-52074.
- Adkins, M.W., S.K. Williams, J. Linger, and J.K. Tyler. 2007b. Chromatin disassembly from the PHO5 promoter is essential for the recruitment of the general transcription machinery and coactivators. *Mol Cell Biol.* 27:6372-6382.
- Agalioti, T., G. Chen, and D. Thanos. 2002. Deciphering the transcriptional histone acetylation code for a human gene. *Cell.* 111:381-392.

- Ahmad, A., Y. Takami, and T. Nakayama. 2003. WD dipeptide motifs and LXXLL motif of chicken HIRA are necessary for transcription repression and the latter motif is essential for interaction with histone deacetylase-2 in vivo. *Biochem Biophys Res Commun.* 312:1266-1272.
- Ai, X., and M.R. Parthun. 2004. The nuclear Hat1p/Hat2p complex: a molecular link between type B histone acetyltransferases and chromatin assembly. *Mol Cell.* 14:195-205.
- Alekseev, O.M., D.C. Bencic, R.T. Richardson, E.E. Widgren, and M.G. O'Rand. 2003. Overexpression of the Linker histone-binding protein tNASP affects progression through the cell cycle. *J Biol Chem.* 278:8846-8852.
- Alekseev, O.M., E.E. Widgren, R.T. Richardson, and M.G. O'Rand. 2005. Association of NASP with HSP90 in mouse spermatogenic cells: stimulation of ATPase activity and transport of linker histones into nuclei. *J Biol Chem.* 280:2904-2911.
- Allfrey, V.G., R. Faulkner, and A.E. Mirsky. 1964. Acetylation and Methylation of Histones and Their Possible Role in the Regulation of Rna Synthesis. *Proc Natl Acad Sci U S A.* 51:786-794.
- Alvarez, F., Munoz, F., Schilcher, P., Imhof, A., Almouzni, G., and Loyola, A. (2011). Sequential establishment of marks on soluble histones H3 and H4. *J Biol Chem* 286, 17714-17721.
- Angelov, D., A. Molla, P.Y. Perche, F. Hans, J. Cote, S. Khochbin, P. Bouvet, and S. Dimitrov. 2003. The histone variant macroH2A interferes with transcription factor binding and SWI/SNF nucleosome remodeling. *Mol Cell.* 11:1033-1041.



- Annunziato, A.T., L.L. Frado, R.L. Seale, and C.L. Woodcock. 1988. Treatment with sodium butyrate inhibits the complete condensation of interphase chromatin. *Chromosoma*. 96:132-138.
- Aoyagi, S., G. Narlikar, C. Zheng, S. Sif, R.E. Kingston, and J.J. Hayes. 2002. Nucleosome remodeling by the human SWI/SNF complex requires transient global disruption of histone-DNA interactions. *Mol Cell Biol*. 22:3653-3662.
- Arents, G., R.W. Burlingame, B.C. Wang, W.E. Love, and E.N. Moudrianakis. 1991. The nucleosomal core histone octamer at 3.1 Å resolution: a tripartite protein assembly and a left-handed superhelix. *Proc Natl Acad Sci U S A*. 88:10148-10152.
- Arents, G., and E.N. Moudrianakis. 1993. Topography of the histone octamer surface: repeating structural motifs utilized in the docking of nucleosomal DNA. *Proc Natl Acad Sci U S A*. 90:10489-10493.
- Arnan, C., N. Saperas, C. Prieto, M. Chiva, and J. Ausio. 2003. Interaction of nucleoplasmin with core histones. *J Biol Chem*. 278:31319-31324.
- Axel, R., W. Melchior, Jr., B. Sollner-Webb, and G. Felsenfeld. 1974. Specific sites of interaction between histones and DNA in chromatin. *Proc Natl Acad Sci U S A*. 71:4101-4105.
- Ayyanathan, K., Lechner, M.S., Bell, P., Maul, G.G., Schultz, D.C., Yamada, Y., Tanaka, K., Torigoe, K., and Rauscher, F.J., 3rd (2003). Regulated recruitment of HP1 to a euchromatic gene induces mitotically heritable, epigenetic gene silencing: a mammalian cell culture model of gene variegation. *Genes Dev* 17, 1855-1869.

- Ball, L.J., N.V. Murzina, R.W. Broadhurst, A.R. Raine, S.J. Archer, F.J. Stott, A.G. Murzin, P.B. Singh, P.J. Domaille, and E.D. Laue. 1997. Structure of the chromatin binding (chromo) domain from mouse modifier protein 1. *EMBO J.* 16:2473-2481.
- Bao, Y., K. Konesky, Y.J. Park, S. Rosu, P.N. Dyer, D. Rangasamy, D.J. Tremethick, P.J. Laybourn, and K. Luger. 2004. Nucleosomes containing the histone variant H2A.Bbd organize only 118 base pairs of DNA. *EMBO J.* 23:3314-3324.
- Bao, Y., and X. Shen. 2007. INO80 subfamily of chromatin remodeling complexes. *Mutat Res.* 618:18-29.
- Batova, I., and M.G. O'Rand. 1996. Histone-binding domains in a human nuclear autoantigenic sperm protein. *Biol Reprod.* 54:1238-1244.
- Becker, P.B. 2002. Nucleosome sliding: facts and fiction. *EMBO J.* 21:4749-4753.
- Birchler, J.A., M.P. Bhadra, and U. Bhadra. 2000. Making noise about silence: repression of repeated genes in animals. *Curr Opin Genet Dev.* 10:211-216.
- Blouin, J.L., G. Duriaux-Sail, H. Chen, A. Gos, M.A. Morris, C. Rossier, and S.E. Antonarakis. 1996. Mapping of the gene for the p60 subunit of the human chromatin assembly factor (CAF1A) to the Down syndrome region of chromosome 21. *Genomics.* 33:309-312.
- Blower, M.D., and G.H. Karpen. 2001. The role of Drosophila CID in kinetochore formation, cell-cycle progression and heterochromatin interactions. *Nat Cell Biol.* 3:730-739.
- Bonne-Andrea, C., F. Harper, J. Sobczak, and A.M. De Recondo. 1984. Rat liver HMG1: a physiological nucleosome assembly factor. *EMBO J.* 3:1193-1199.
- Bonner, W.M. 1975. Protein migration into nuclei. II. Frog oocyte nuclei accumulate a class of microinjected oocyte nuclear proteins and exclude a class of microinjected oocyte cytoplasmic proteins. *J Cell Biol.* 64:431-437.

- Boyer, L.A., R.R. Latek, and C.L. Peterson. 2004. The SANT domain: a unique histone-tail-binding module? *Nat Rev Mol Cell Biol.* 5:158-163.
- Bradbury, E.M. 1992. Reversible histone modifications and the chromosome cell cycle. *Bioessays.* 14:9-16.
- Buchwitz, B.J., K. Ahmad, L.L. Moore, M.B. Roth, and S. Henikoff. 1999. A histone-H3-like protein in *C. elegans*. *Nature.* 401:547-548.
- Campos, E.I., Fillingham, J., Li, G., Zheng, H., Voigt, P., Kuo, W.H., Seepany, H., Gao, Z., Day, L.A., Greenblatt, J.F., *et al.* (2010). The program for processing newly synthesized histones H3.1 and H4. *Nat Struct Mol Biol* 17, 1343-1351.
- Carrera, P., Y.M. Moshkin, S. Gronke, H.H. Sillje, E.A. Nigg, H. Jackle, and F. Karch. 2003. Tousled-like kinase functions with the chromatin assembly pathway regulating nuclear divisions. *Genes Dev.* 17:2578-2590.
- Carruthers, L.M., J. Bednar, C.L. Woodcock, and J.C. Hansen. 1998. Linker histones stabilize the intrinsic salt-dependent folding of nucleosomal arrays: mechanistic ramifications for higher-order chromatin folding. *Biochemistry.* 37:14776-14787.
- Chadwick, B.P., and H.F. Willard. 2001. A novel chromatin protein, distantly related to histone H2A, is largely excluded from the inactive X chromosome. *J Cell Biol.* 152:375-384.
- Chen, C.C., J.J. Carson, J. Feser, B. Tamburini, S. Zabaronick, J. Linger, and J.K. Tyler. 2008. Acetylated lysine 56 on histone H3 drives chromatin assembly after repair and signals for the completion of repair. *Cell.* 134:231-243.
- Cheung, P., C.D. Allis, and P. Sassone-Corsi. 2000. Signaling to chromatin through histone modifications. *Cell.* 103:263-271.

- Clark, K.L., E.D. Halay, E. Lai, and S.K. Burley. 1993. Co-crystal structure of the HNF-3/fork head DNA-recognition motif resembles histone H5. *Nature*. 364:412-420.
- Corona, D.F., and J.W. Tamkun. 2004. Multiple roles for ISWI in transcription, chromosome organization and DNA replication. *Biochim Biophys Acta*. 1677:113-119.
- Costanzi, C., and J.R. Pehrson. 1998. Histone macroH2A1 is concentrated in the inactive X chromosome of female mammals. *Nature*. 393:599-601.
- Cryderman, D.E., S.K. Grade, Y. Li, L. Fanti, S. Pimpinelli, and L.L. Wallrath. 2005. Role of Drosophila HP1 in euchromatic gene expression. *Dev Dyn*. 232:767-774.
- Daganzo, S.M., J.P. Erzberger, W.M. Lam, E. Skordalakes, R. Zhang, A.A. Franco, S.J. Brill, P.D. Adams, J.M. Berger, and P.D. Kaufman. 2003. Structure and function of the conserved core of histone deposition protein Asf1. *Curr Biol*. 13:2148-2158.
- Das, C., M.S. Lucia, K.C. Hansen, and J.K. Tyler. 2009. CBP/p300-mediated acetylation of histone H3 on lysine 56. *Nature*. 459:113-117.
- Das, C., J.K. Tyler, and M.E. Churchill. 2010. The histone shuffle: histone chaperones in an energetic dance. *Trends Biochem Sci*. 35:476-489.
- Davie, J.R., and L.C. Murphy. 1990. Level of ubiquitinated histone H2B in chromatin is coupled to ongoing transcription. *Biochemistry*. 29:4752-4757.
- De Koning, L., A. Corpet, J.E. Haber, and G. Almouzni. 2007. Histone chaperones: an escort network regulating histone traffic. *Nat Struct Mol Biol*. 14:997-1007.
- De Robertis, E.M., R.F. Longthorne, and J.B. Gurdon. 1978. Intracellular migration of nuclear proteins in *Xenopus* oocytes. *Nature*. 272:254-256.
- DeSilva, H., K. Lee, and M.A. Osley. 1998. Functional dissection of yeast Hir1p, a WD repeat-containing transcriptional corepressor. *Genetics*. 148:657-667.

- Dialynas, G.K., Terjung, S., Brown, J.P., Aucott, R.L., Baron-Luhr, B., Singh, P.B., and Georgatos, S.D. (2007). Plasticity of HP1 proteins in mammalian cells. *J Cell Sci* 120, 3415-3424.
- Dimova, D., Z. Nackerdien, S. Furgeson, S. Eguchi, and M.A. Osley. 1999. A role for transcriptional repressors in targeting the yeast Swi/Snf complex. *Mol Cell*. 4:75-83.
- Drane, P., K. Ouararhni, A. Depaux, M. Shuaib, and A. Hamiche. 2010. The death-associated protein DAXX is a novel histone chaperone involved in the replication-independent deposition of H3.3. *Genes Dev*. 24:1253-1265.
- Drew, H.R. 1993. Reconstitution of short-spaced chromatin from the histone octamer and either HMG-14,17 or histone H1. *J Mol Biol*. 230:824-836.
- Ehsan, H., J.P. Reichheld, T. Durfee, and J.L. Roe. 2004. TOUSLED kinase activity oscillates during the cell cycle and interacts with chromatin regulators. *Plant Physiol*. 134:1488-1499.
- Emili, A., D.M. Schieltz, J.R. Yates, 3rd, and L.H. Hartwell. 2001. Dynamic interaction of DNA damage checkpoint protein Rad53 with chromatin assembly factor Asf1. *Mol Cell*. 7:13-20.
- English, C.M., M.W. Adkins, J.J. Carson, M.E. Churchill, and J.K. Tyler. 2006. Structural basis for the histone chaperone activity of Asf1. *Cell*. 127:495-508.
- Enomoto, S., P.D. McCune-Zierath, M. Gerami-Nejad, M.A. Sanders, and J. Berman. 1997. RLF2, a subunit of yeast chromatin assembly factor-I, is required for telomeric chromatin function in vivo. *Genes Dev*. 11:358-370.
- Falbo, K.B., and X. Shen. 2006. Chromatin remodeling in DNA replication. *J Cell Biochem*. 97:684-689.

- Fernandez-Capetillo, O., A. Lee, M. Nussenzweig, and A. Nussenzweig. 2004. H2AX: the histone guardian of the genome. *DNA Repair (Amst)*. 3:959-967.
- Fischle, W., B.S. Tseng, H.L. Dormann, B.M. Ueberheide, B.A. Garcia, J. Shabanowitz, D.F. Hunt, H. Funabiki, and C.D. Allis. 2005. Regulation of HP1-chromatin binding by histone H3 methylation and phosphorylation. *Nature*. 438:1116-1122.
- Fischle, W., Y. Wang, and C.D. Allis. 2003a. Binary switches and modification cassettes in histone biology and beyond. *Nature*. 425:475-479.
- Fischle, W., Y. Wang, S.A. Jacobs, Y. Kim, C.D. Allis, and S. Khorasanizadeh. 2003b. Molecular basis for the discrimination of repressive methyl-lysine marks in histone H3 by Polycomb and HP1 chromodomains. *Genes Dev*. 17:1870-1881.
- Fukagawa, T., Y. Mikami, A. Nishihashi, V. Regnier, T. Haraguchi, Y. Hiraoka, N. Sugata, K. Todokoro, W. Brown, and T. Ikemura. 2001. CENP-H, a constitutive centromere component, is required for centromere targeting of CENP-C in vertebrate cells. *EMBO J*. 20:4603-4617.
- Fukagawa, T., C. Pendon, J. Morris, and W. Brown. 1999. CENP-C is necessary but not sufficient to induce formation of a functional centromere. *EMBO J*. 18:4196-4209.
- Fuks, F., P.J. Hurd, R. Deplus, and T. Kouzarides. 2003. The DNA methyltransferases associate with HP1 and the SUV39H1 histone methyltransferase. *Nucleic Acids Res*. 31:2305-2312.
- Gaillard, P.H., E.M. Martini, P.D. Kaufman, B. Stillman, E. Moustacchi, and G. Almouzni. 1996. Chromatin assembly coupled to DNA repair: a new role for chromatin assembly factor I. *Cell*. 86:887-896.

- Gasser, R., T. Koller, and J.M. Sogo. 1996. The stability of nucleosomes at the replication fork. *J Mol Biol.* 258:224-239.
- Gautier, T., D.W. Abbott, A. Molla, A. Verdel, J. Ausio, and S. Dimitrov. 2004. Histone variant H2ABbd confers lower stability to the nucleosome. *EMBO Rep.* 5:715-720.
- Gavin, I., P.J. Horn, and C.L. Peterson. 2001. SWI/SNF chromatin remodeling requires changes in DNA topology. *Mol Cell.* 7:97-104.
- Gilbert, C.S., C.M. Green, and N.F. Lowndes. 2001. Budding yeast Rad9 is an ATP-dependent Rad53 activating machine. *Mol Cell.* 8:129-136.
- Gineitis, A.A., I.A. Zalenskaya, P.M. Yau, E.M. Bradbury, and A.O. Zalensky. 2000. Human sperm telomere-binding complex involves histone H2B and secures telomere membrane attachment. *J Cell Biol.* 151:1591-1598.
- Gkikopoulos, T., K.M. Havas, H. Dewar, and T. Owen-Hughes. 2009. SWI/SNF and Asf1p cooperate to displace histones during induction of the *saccharomyces cerevisiae* HO promoter. *Mol Cell Biol.* 29:4057-4066.
- Goldberg, A.D., Banaszynski, L.A., Noh, K.M., Lewis, P.W., Elsaesser, S.J., Stadler, S., Dewell, S., Law, M., Guo, X., Li, X., *et al.* (2010). Distinct factors control histone variant H3.3 localization at specific genomic regions. *Cell* 140, 678-691.
- Golderer, G., and P. Grobner. 1991. ADP-ribosylation of core histones and their acetylated subspecies. *Biochem J.* 277 ( Pt 3):607-610.
- Goldknopf, I.L., C.W. Taylor, R.M. Baum, L.C. Yeoman, M.O. Olson, A.W. Prestayko, and H. Busch. 1975. Isolation and characterization of protein A24, a "histone-like" non-histone chromosomal protein. *J Biol Chem.* 250:7182-7187.

- Goodfellow, H., A. Krejci, Y. Moshkin, C.P. Verrijzer, F. Karch, and S.J. Bray. 2007. Gene-specific targeting of the histone chaperone asf1 to mediate silencing. *Dev Cell*. 13:593-600.
- Goshima, G., S. Saitoh, and M. Yanagida. 1999. Proper metaphase spindle length is determined by centromere proteins Mis12 and Mis6 required for faithful chromosome segregation. *Genes Dev*. 13:1664-1677.
- Green, E.M., A.J. Antczak, A.O. Bailey, A.A. Franco, K.J. Wu, J.R. Yates, 3rd, and P.D. Kaufman. 2005. Replication-independent histone deposition by the HIR complex and Asf1. *Curr Biol*. 15:2044-2049.
- Groth, A., A. Corpet, A.J. Cook, D. Roche, J. Bartek, J. Lukas, and G. Almouzni. 2007a. Regulation of replication fork progression through histone supply and demand. *Science*. 318:1928-1931.
- Groth, A., D. Ray-Gallet, J.P. Quivy, J. Lukas, J. Bartek, and G. Almouzni. 2005. Human Asf1 regulates the flow of S phase histones during replicational stress. *Mol Cell*. 17:301-311.
- Groth, A., W. Rocha, A. Verreault, and G. Almouzni. 2007b. Chromatin challenges during DNA replication and repair. *Cell*. 128:721-733.
- Grunstein, M. 1997. Histone acetylation in chromatin structure and transcription. *Nature*. 389:349-352.
- Halford, S., R. Wadey, C. Roberts, S.C. Daw, J.A. Whiting, H. O'Donnell, I. Dunham, D. Bentley, E. Lindsay, A. Baldini, and et al. 1993. Isolation of a putative transcriptional regulator from the region of 22q11 deleted in DiGeorge syndrome, Shprintzen syndrome and familial congenital heart disease. *Hum Mol Genet*. 2:2099-2107.



- Hamiche, A., and M. Shuaib. 2012. Chaperoning the histone H3 family. *Biochim Biophys Acta*. 1819:230-237.
- Hannich, J.T., A. Lewis, M.B. Kroetz, S.J. Li, H. Heide, A. Emili, and M. Hochstrasser. 2005. Defining the SUMO-modified proteome by multiple approaches in *Saccharomyces cerevisiae*. *J Biol Chem*. 280:4102-4110.
- Hansen, J.C., X. Lu, E.D. Ross, and R.W. Woody. 2006. Intrinsic protein disorder, amino acid composition, and histone terminal domains. *J Biol Chem*. 281:1853-1856.
- Hansen, J.C., C. Tse, and A.P. Wolffe. 1998. Structure and function of the core histone N-termini: more than meets the eye. *Biochemistry*. 37:17637-17641.
- Hassa, P.O., S.S. Haenni, M. Elser, and M.O. Hottiger. 2006. Nuclear ADP-ribosylation reactions in mammalian cells: where are we today and where are we going? *Microbiol Mol Biol Rev*. 70:789-829.
- Henikoff, S. 2008. Nucleosome destabilization in the epigenetic regulation of gene expression. *Nat Rev Genet*. 9:15-26.
- Henikoff, S., and K. Ahmad. 2005. Assembly of variant histones into chromatin. *Annu Rev Cell Dev Biol*. 21:133-153.
- Henikoff, S., T. Furuyama, and K. Ahmad. 2004. Histone variants, nucleosome assembly and epigenetic inheritance. *Trends Genet*. 20:320-326.
- Hertel, L., M. De Andrea, G. Bellomo, P. Santoro, S. Landolfo, and M. Gariglio. 1999. The HMG protein T160 colocalizes with DNA replication foci and is down-regulated during cell differentiation. *Exp Cell Res*. 250:313-328.
- Honda, B.M., G.H. Dixon, and E.P. Candido. 1975. Sites of in vivo histone methylation in developing trout testis. *J Biol Chem*. 250:8681-8685.

- Howman, E.V., K.J. Fowler, A.J. Newson, S. Redward, A.C. MacDonald, P. Kalitsis, and K.H. Choo. 2000. Early disruption of centromeric chromatin organization in centromere protein A (Cenpa) null mice. *Proc Natl Acad Sci U S A.* 97:1148-1153.
- Hu, F., A.A. Alcasabas, and S.J. Elledge. 2001. Asf1 links Rad53 to control of chromatin assembly. *Genes Dev.* 15:1061-1066.
- Huang, S., H. Zhou, D. Katzmann, M. Hochstrasser, E. Atanasova, and Z. Zhang. 2005. Rtt106p is a histone chaperone involved in heterochromatin-mediated silencing. *Proc Natl Acad Sci U S A.* 102:13410-13415.
- Huisinga, K.L., B. Brower-Toland, and S.C. Elgin. 2006. The contradictory definitions of heterochromatin: transcription and silencing. *Chromosoma.* 115:110-122.
- Hymes, J., K. Fleischhauer, and B. Wolf. 1995. Biotinylation of histones by human serum biotinidase: assessment of biotinyl-transferase activity in sera from normal individuals and children with biotinidase deficiency. *Biochem Mol Med.* 56:76-83.
- Ishimi, Y., J. Hirosumi, W. Sato, K. Sugawara, S. Yokota, F. Hanaoka, and M. Yamada. 1984. Purification and initial characterization of a protein which facilitates assembly of nucleosome-like structure from mammalian cells. *Eur J Biochem.* 142:431-439.
- Ishimi, Y., M. Kojima, M. Yamada, and F. Hanaoka. 1987. Binding mode of nucleosome-assembly protein (AP-I) and histones. *Eur J Biochem.* 162:19-24.
- Ito, T., M. Bulger, R. Kobayashi, and J.T. Kadonaga. 1996. Drosophila NAP-1 is a core histone chaperone that functions in ATP-facilitated assembly of regularly spaced nucleosomal arrays. *Mol Cell Biol.* 16:3112-3124.

- James, T.C., and S.C. Elgin. 1986. Identification of a nonhistone chromosomal protein associated with heterochromatin in *Drosophila melanogaster* and its gene. *Mol Cell Biol.* 6:3862-3872.
- Jasencakova, Z., A.N. Scharf, K. Ask, A. Corpet, A. Imhof, G. Almouzni, and A. Groth. 2010. Replication stress interferes with histone recycling and predeposition marking of new histones. *Mol Cell.* 37:736-743.
- Jiang, W.Q., Nguyen, A., Cao, Y., Chang, A.C.M., Redde, R.R. HP1-Mediated Formation of Alternative Lengthening of Telomeres-Associated PML Bodies Requires HIRA but Not ASF1a | PLoS One. 2011; 6(2): e17036.
- Johansen, K.M., and J. Johansen. 2006. Regulation of chromatin structure by histone H3S10 phosphorylation. *Chromosome Res.* 14:393-404.
- Kamakaka, R.T., M. Bulger, P.D. Kaufman, B. Stillman, and J.T. Kadonaga. 1996. Postreplicative chromatin assembly by *Drosophila* and human chromatin assembly factor 1. *Mol Cell Biol.* 16:810-817.
- Karpova, T.S., T.Y. Chen, B.L. Sprague, and J.G. McNally. 2004. Dynamic interactions of a transcription factor with DNA are accelerated by a chromatin remodeller. *EMBO Rep.* 5:1064-1070.
- Kassabov, S.R., B. Zhang, J. Persinger, and B. Bartholomew. 2003. SWI/SNF unwraps, slides, and rewraps the nucleosome. *Mol Cell.* 11:391-403.
- Kaufman, P.D., J.L. Cohen, and M.A. Osley. 1998. Hir proteins are required for position-dependent gene silencing in *Saccharomyces cerevisiae* in the absence of chromatin assembly factor I. *Mol Cell Biol.* 18:4793-4806.

- Kaufman, P.D., R. Kobayashi, N. Kessler, and B. Stillman. 1995. The p150 and p60 subunits of chromatin assembly factor I: a molecular link between newly synthesized histones and DNA replication. *Cell*. 81:1105-1114.
- Kaufman, P.D., R. Kobayashi, and B. Stillman. 1997. Ultraviolet radiation sensitivity and reduction of telomeric silencing in *Saccharomyces cerevisiae* cells lacking chromatin assembly factor-I. *Genes Dev*. 11:345-357.
- Keller, C., and T. Krude. 2000. Requirement of Cyclin/Cdk2 and protein phosphatase 1 activity for chromatin assembly factor 1-dependent chromatin assembly during DNA synthesis. *J Biol Chem*. 275:35512-35521.
- Khorasanizadeh, S. 2004. The nucleosome: from genomic organization to genomic regulation. *Cell*. 116:259-272.
- Kim, U.J., M. Han, P. Kayne, and M. Grunstein. 1988. Effects of histone H4 depletion on the cell cycle and transcription of *Saccharomyces cerevisiae*. *EMBO J*. 7:2211-2219.
- Kirov, N., A. Shtilbans, and C. Rushlow. 1998. Isolation and characterization of a new gene encoding a member of the HIRA family of proteins from *Drosophila melanogaster*. *Gene*. 212:323-332.
- Kleinschmidt, J.A., C. Dingwall, G. Maier, and W.W. Franke. 1986. Molecular characterization of a karyophilic, histone-binding protein: cDNA cloning, amino acid sequence and expression of nuclear protein N1/N2 of *Xenopus laevis*. *EMBO J*. 5:3547-3552.
- Kleinschmidt, J.A., and W.W. Franke. 1982. Soluble acidic complexes containing histones H3 and H4 in nuclei of *Xenopus laevis* oocytes. *Cell*. 29:799-809.

- Korber, P., S. Barbaric, T. Luckenbach, A. Schmid, U.J. Schermer, D. Blaschke, and W. Horz. 2006. The histone chaperone Asf1 increases the rate of histone eviction at the yeast PHO5 and PHO8 promoters. *J Biol Chem.* 281:5539-5545.
- Kornberg, R.D. 1974. Chromatin structure: a repeating unit of histones and DNA. *Science.* 184:868-871.
- Kornberg, R.D., and J.O. Thomas. 1974. Chromatin structure; oligomers of the histones. *Science.* 184:865-868.
- Koshland, D., and A. Strunnikov. 1996. Mitotic chromosome condensation. *Annu Rev Cell Dev Biol.* 12:305-333.
- Krawitz, D.C., T. Kama, and P.D. Kaufman. 2002. Chromatin assembly factor I mutants defective for PCNA binding require Asf1/Hir proteins for silencing. *Mol Cell Biol.* 22:614-625.
- Krude, T. 1995. Chromatin assembly factor 1 (CAF-1) colocalizes with replication foci in HeLa cell nuclei. *Exp Cell Res.* 220:304-311.
- Kurdistani, S.K., S. Tavazoie, and M. Grunstein. 2004. Mapping global histone acetylation patterns to gene expression. *Cell.* 117:721-733.
- Kwon, S.H. and Workman, J.L. The changing faces of HP1 : from heterochromatin formation and gene expression silencing to euchromatic gene expression. 2011, *Bioassays* 33 : 280-289.
- Lachner, M., D. O'Carroll, S. Rea, K. Mechtler, and T. Jenuwein. 2001. Methylation of histone H3 lysine 9 creates a binding site for HP1 proteins. *Nature.* 410:116-120.
- Lachner, M., R.J. O'Sullivan, and T. Jenuwein. 2003. An epigenetic road map for histone lysine methylation. *J Cell Sci.* 116:2117-2124.

- Lamond, A.I., and W.C. Earnshaw. 1998. Structure and function in the nucleus. *Science*. 280:547-553.
- Lamour, V., Y. Lecluse, C. Desmaze, M. Spector, M. Bodescot, A. Aurias, M.A. Osley, and M. Lipinski. 1995. A human homolog of the *S. cerevisiae* HIR1 and HIR2 transcriptional repressors cloned from the DiGeorge syndrome critical region. *Hum Mol Genet*. 4:791-799.
- Laskey, R.A., and W.C. Earnshaw. 1980. Nucleosome assembly. *Nature*. 286:763-767.
- Laskey, R.A., B.M. Honda, A.D. Mills, and J.T. Finch. 1978. Nucleosomes are assembled by an acidic protein which binds histones and transfers them to DNA. *Nature*. 275:416-420.
- Le Douarin, B., A.L. Nielsen, J.M. Garnier, H. Ichinose, F. Jeanmougin, R. Losson, and P. Chambon. 1996. A possible involvement of TIF1 alpha and TIF1 beta in the epigenetic control of transcription by nuclear receptors. *EMBO J*. 15:6701-6715.
- Le, S., C. Davis, J.B. Konopka, and R. Sternglanz. 1997. Two new S-phase-specific genes from *Saccharomyces cerevisiae*. *Yeast*. 13:1029-1042.
- Leszinski, G., U. Gezer, B. Siegele, O. Stoetzer, and S. Holdenrieder. 2012. Relevance of histone marks H3K9me3 and H4K20me3 in cancer. *Anticancer Res*. 32:2199-2205.
- Levchenko, V., and V. Jackson. 2004. Histone release during transcription: NAP1 forms a complex with H2A and H2B and facilitates a topologically dependent release of H3 and H4 from the nucleosome. *Biochemistry*. 43:2359-2372.
- Lewis, P.W., S.J. Elsaesser, K.M. Noh, S.C. Stadler, and C.D. Allis. 2010. Daxx is an H3.3-specific histone chaperone and cooperates with ATRX in replication-independent chromatin assembly at telomeres. *Proc Natl Acad Sci U S A*. 107:14075-14080.

- Liebich, H.M., E. Gesele, C. Wirth, J. Woll, K. Jobst, and A. Lakatos. 1993. Non-enzymatic glycation of histones. *Biol Mass Spectrom.* 22:121-123.
- Loppin, B., E. Bonnefoy, C. Anselme, A. Laurencon, T.L. Karr, and P. Couble. 2005. The histone H3.3 chaperone HIRA is essential for chromatin assembly in the male pronucleus. *Nature.* 437:1386-1390.
- Lorch, Y., B. Maier-Davis, and R.D. Kornberg. 2006. Chromatin remodeling by nucleosome disassembly in vitro. *Proc Natl Acad Sci U S A.* 103:3090-3093.
- Loyola, A., and G. Almouzni. 2004. Histone chaperones, a supporting role in the limelight. *Biochim Biophys Acta.* 1677:3-11.
- Lu, B.Y., P.C. Emtage, B.J. Duyf, A.J. Hilliker, and J.C. Eissenberg. 2000. Heterochromatin protein 1 is required for the normal expression of two heterochromatin genes in *Drosophila*. *Genetics.* 155:699-708.
- Luger, K., A.W. Mader, R.K. Richmond, D.F. Sargent, and T.J. Richmond. 1997. Crystal structure of the nucleosome core particle at 2.8 Å resolution. *Nature.* 389:251-260.
- Luk, E., A. Ranjan, P.C. Fitzgerald, G. Mizuguchi, Y. Huang, D. Wei, and C. Wu. 2010. Stepwise histone replacement by SWR1 requires dual activation with histone H2A.Z and canonical nucleosome. *Cell.* 143:725-736.
- Magnaghi, P., C. Roberts, S. Lorain, M. Lipinski, and P.J. Scambler. 1998. HIRA, a mammalian homologue of *Saccharomyces cerevisiae* transcriptional co-repressors, interacts with Pax3. *Nat Genet.* 20:74-77.
- Mahadevan, L.C., A.C. Willis, and M.J. Barratt. 1991. Rapid histone H3 phosphorylation in response to growth factors, phorbol esters, okadaic acid, and protein synthesis inhibitors. *Cell.* 65:775-783.

- Maier, V.K., M. Chioda, and P.B. Becker. 2008. ATP-dependent chromatosome remodeling. *Biol Chem.* 389:345-352.
- Maison, C., Bailly, D., Roche, D., Montes de Oca, R., Probst, A.V., Vassias, I., Dingli, F., Lombard, B., Loew, D., Quivy, J.P., *et al.* (2011). SUMOylation promotes de novo targeting of HP1alpha to pericentric heterochromatin. *Nat Genet* 43, 220-227.
- Majka, J., and P.M. Burgers. 2004. The PCNA-RFC families of DNA clamps and clamp loaders. *Prog Nucleic Acid Res Mol Biol.* 78:227-260.
- Mak, W., J. Baxter, J. Silva, A.E. Newall, A.P. Otte, and N. Brockdorff. 2002. Mitotically stable association of polycomb group proteins eed and enx1 with the inactive x chromosome in trophoblast stem cells. *Curr Biol.* 12:1016-1020.
- Malik, H.S., and S. Henikoff. 2003. Phylogenomics of the nucleosome. *Nat Struct Biol.* 10:882-891.
- Manza, L.L., S.G. Codreanu, S.L. Stamer, D.L. Smith, K.S. Wells, R.L. Roberts, and D.C. Liebler. 2004. Global shifts in protein sumoylation in response to electrophile and oxidative stress. *Chem Res Toxicol.* 17:1706-1715.
- Marfella, C.G., and A.N. Imbalzano. 2007. The Chd family of chromatin remodelers. *Mutat Res.* 618:30-40.
- Marheineke, K., and T. Krude. 1998. Nucleosome assembly activity and intracellular localization of human CAF-1 changes during the cell division cycle. *J Biol Chem.* 273:15279-15286.
- Martin, C., and Y. Zhang. 2005. The diverse functions of histone lysine methylation. *Nat Rev Mol Cell Biol.* 6:838-849.
- Martinez-Balbas, M.A., T. Tsukiyama, D. Gdula, and C. Wu. 1998. Drosophila NURF-55, a WD repeat protein involved in histone metabolism. *Proc Natl Acad Sci U S A.* 95:132-137.



- Martini, E., D.M. Roche, K. Marheineke, A. Verreault, and G. Almouzni. 1998. Recruitment of phosphorylated chromatin assembly factor 1 to chromatin after UV irradiation of human cells. *J Cell Biol.* 143:563-575.
- Mathews, L.A., F. Crea, and W.L. Farrar. 2009. Epigenetic gene regulation in stem cells and correlation to cancer. *Differentiation.* 78:1-17.
- McBryant, S.J., Y.J. Park, S.M. Abernathy, P.J. Laybourn, J.K. Nyborg, and K. Luger. 2003. Preferential binding of the histone (H3-H4)<sub>2</sub> tetramer by NAP1 is mediated by the amino-terminal histone tails. *J Biol Chem.* 278:44574-44583.
- McBryant, S.J., and O.B. Peersen. 2004. Self-association of the yeast nucleosome assembly protein 1. *Biochemistry.* 43:10592-10599.
- Meehan, R.R., C.F. Kao, and S. Pennings. 2003. HP1 binding to native chromatin in vitro is determined by the hinge region and not by the chromodomain. *EMBO J.* 22:3164-3174.
- Mello, J.A., and G. Almouzni. 2001. The ins and outs of nucleosome assembly. *Curr Opin Genet Dev.* 11:136-141.
- Mello, J.A., H.H. Sillje, D.M. Roche, D.B. Kirschner, E.A. Nigg, and G. Almouzni. 2002. Human Asf1 and CAF-1 interact and synergize in a repair-coupled nucleosome assembly pathway. *EMBO Rep.* 3:329-334.
- Meluh, P.B., and D. Koshland. 1995. Evidence that the MIF2 gene of *Saccharomyces cerevisiae* encodes a centromere protein with homology to the mammalian centromere protein CENP-C. *Mol Biol Cell.* 6:793-807.
- Mermoud, J.E., C. Costanzi, J.R. Pehrson, and N. Brockdorff. 1999. Histone macroH2A1.2 relocates to the inactive X chromosome after initiation and propagation of X-inactivation. *J Cell Biol.* 147:1399-1408.

- Miao, F., and R. Natarajan. 2005. Mapping global histone methylation patterns in the coding regions of human genes. *Mol Cell Biol.* 25:4650-4661.
- Minc, E., Y. Allory, H.J. Worman, J.C. Courvalin, and B. Buendia. 1999. Localization and phosphorylation of HP1 proteins during the cell cycle in mammalian cells. *Chromosoma.* 108:220-234.
- Moggs, J.G., P. Grandi, J.P. Quivy, Z.O. Jonsson, U. Hubscher, P.B. Becker, and G. Almouzni. 2000. A CAF-1-PCNA-mediated chromatin assembly pathway triggered by sensing DNA damage. *Mol Cell Biol.* 20:1206-1218.
- Mohrmann, L., and C.P. Verrijzer. 2005. Composition and functional specificity of SWI2/SNF2 class chromatin remodeling complexes. *Biochim Biophys Acta.* 1681:59-73.
- Mosammamarast, N., C.S. Ewart, and L.F. Pemberton. 2002. A role for nucleosome assembly protein 1 in the nuclear transport of histones H2A and H2B. *EMBO J.* 21:6527-6538.
- Mousson, F., A. Lautrette, J.Y. Thuret, M. Agez, R. Courbeyrette, B. Amigues, E. Becker, J.M. Neumann, R. Guerois, C. Mann, and F. Ochsenbein. 2005. Structural basis for the interaction of Asf1 with histone H3 and its functional implications. *Proc Natl Acad Sci U S A.* 102:5975-5980.
- Muchardt, C., M. Guilleme, J.S. Seeler, D. Trouche, A. Dejean, and M. Yaniv. 2002. Coordinated methyl and RNA binding is required for heterochromatin localization of mammalian HP1alpha. *EMBO Rep.* 3:975-981.
- Muller, J., C.M. Hart, N.J. Francis, M.L. Vargas, A. Sengupta, B. Wild, E.L. Miller, M.B. O'Connor, R.E. Kingston, and J.A. Simon. 2002. Histone methyltransferase activity of a *Drosophila* Polycomb group repressor complex. *Cell.* 111:197-208.

- Munakata, T., N. Adachi, N. Yokoyama, T. Kuzuhara, and M. Horikoshi. 2000. A human homologue of yeast anti-silencing factor has histone chaperone activity. *Genes Cells*. 5:221-233.
- Murray, K. 1964. The Occurrence of Epsilon-N-Methyl Lysine in Histones. *Biochemistry*. 3:10-15.
- Murzina, N., A. Verreault, E. Laue, and B. Stillman. 1999. Heterochromatin dynamics in mouse cells: interaction between chromatin assembly factor 1 and HP1 proteins. *Mol Cell*. 4:529-540.
- Murzina, N.V., X.Y. Pei, W. Zhang, M. Sparkes, J. Vicente-Garcia, J.V. Pratap, S.H. McLaughlin, T.R. Ben-Shahar, A. Verreault, B.F. Luisi, and E.D. Laue. 2008. Structural basis for the recognition of histone H4 by the histone-chaperone RbAp46. *Structure*. 16:1077-1085.
- Nabatiyan, A., and T. Krude. 2004. Silencing of chromatin assembly factor 1 in human cells leads to cell death and loss of chromatin assembly during DNA synthesis. *Mol Cell Biol*. 24:2853-2862.
- Nabatiyan, A., D. Szuts, and T. Krude. 2006. Induction of CAF-1 expression in response to DNA strand breaks in quiescent human cells. *Mol Cell Biol*. 26:1839-1849.
- Nakayama, J., J.C. Rice, B.D. Strahl, C.D. Allis, and S.I. Grewal. 2001. Role of histone H3 lysine 9 methylation in epigenetic control of heterochromatin assembly. *Science*. 292:110-113.
- Nakayama, T., K. Nishioka, Y.X. Dong, T. Shimojima, and S. Hirose. 2007. Drosophila GAGA factor directs histone H3.3 replacement that prevents the heterochromatin spreading. *Genes Dev*. 21:552-561.

- Narlikar, G.J., H.Y. Fan, and R.E. Kingston. 2002. Cooperation between complexes that regulate chromatin structure and transcription. *Cell*. 108:475-487.
- Nathan, D., D.E. Sterner, and S.L. Berger. 2003. Histone modifications: Now summoning sumoylation. *Proc Natl Acad Sci U S A*. 100:13118-13120.
- Natsume, R., M. Eitoku, Y. Akai, N. Sano, M. Horikoshi, and T. Senda. 2007. Structure and function of the histone chaperone CIA/ASF1 complexed with histones H3 and H4. *Nature*. 446:338-341.
- Nelson, D.M., X. Ye, C. Hall, H. Santos, T. Ma, G.D. Kao, T.J. Yen, J.W. Harper, and P.D. Adams. 2002. Coupling of DNA synthesis and histone synthesis in S phase independent of cyclin/cdk2 activity. *Mol Cell Biol*. 22:7459-7472.
- Nelson, T., R. Wiegand, and D. Brutlag. 1981. Ribonucleic acid and other polyanions facilitate chromatin assembly in vitro. *Biochemistry*. 20:2594-2601.
- Ng, H.H., F. Robert, R.A. Young, and K. Struhl. 2003. Targeted recruitment of Set1 histone methylase by elongating Pol II provides a localized mark and memory of recent transcriptional activity. *Mol Cell*. 11:709-719.
- Nielsen, P.R., D. Nietlispach, H.R. Mott, J. Callaghan, A. Bannister, T. Kouzarides, A.G. Murzin, N.V. Murzina, and E.D. Laue. 2002. Structure of the HP1 chromodomain bound to histone H3 methylated at lysine 9. *Nature*. 416:103-107.
- Nielsen, S.J., R. Schneider, U.M. Bauer, A.J. Bannister, A. Morrison, D. O'Carroll, R. Firestein, M. Cleary, T. Jenuwein, R.E. Herrera, and T. Kouzarides. 2001. Rb targets histone H3 methylation and HP1 to promoters. *Nature*. 412:561-565.

- Nightingale, K.P., R.E. Wellinger, J.M. Sogo, and P.B. Becker. 1998. Histone acetylation facilitates RNA polymerase II transcription of the *Drosophila hsp26* gene in chromatin. *EMBO J.* 17:2865-2876.
- Nishida, H., T. Suzuki, S. Kondo, H. Miura, Y. Fujimura, and Y. Hayashizaki. 2006. Histone H3 acetylated at lysine 9 in promoter is associated with low nucleosome density in the vicinity of transcription start site in human cell. *Chromosome Res.* 14:203-211.
- Nishihashi, A., T. Haraguchi, Y. Hiraoka, T. Ikemura, V. Regnier, H. Dodson, W.C. Earnshaw, and T. Fukagawa. 2002. CENP-I is essential for centromere function in vertebrate cells. *Dev Cell.* 2:463-476.
- Nowak, S.J., and V.G. Corces. 2004. Phosphorylation of histone H3: a balancing act between chromosome condensation and transcriptional activation. *Trends Genet.* 20:214-220.
- O'Rand, M.G., R.T. Richardson, L.J. Zimmerman, and E.E. Widgren. 1992. Sequence and localization of human NASP: conservation of a *Xenopus* histone-binding protein. *Dev Biol.* 154:37-44.
- Ohzeki, J., J.H. Bergmann, N. Kouprina, V.N. Noskov, M. Nakano, H. Kimura, W.C. Earnshaw, V. Larionov, and H. Masumoto. 2012. Breaking the HAC Barrier: histone H3K9 acetyl/methyl balance regulates CENP-A assembly. *EMBO J.* 31:2391-2402.
- Okuwaki, M., K. Kato, H. Shimahara, S. Tate, and K. Nagata. 2005. Assembly and disassembly of nucleosome core particles containing histone variants by human nucleosome assembly protein I. *Mol Cell Biol.* 25:10639-10651.
- Oudet, P., M. Gross-Bellard, and P. Chambon. 1975. Electron microscopic and biochemical evidence that chromatin structure is a repeating unit. *Cell.* 4:281-300.

- Padmanabhan, B., K. Kataoka, T. Umehara, N. Adachi, S. Yokoyama, and M. Horikoshi. 2005. Structural similarity between histone chaperone Cia1p/Asf1p and DNA-binding protein NF-kappaB. *J Biochem.* 138:821-829.
- Park, Y.J., J.V. Chodaparambil, Y. Bao, S.J. McBryant, and K. Luger. 2005. Nucleosome assembly protein 1 exchanges histone H2A-H2B dimers and assists nucleosome sliding. *J Biol Chem.* 280:1817-1825.
- Park, Y.J., and K. Luger. 2006. Structure and function of nucleosome assembly proteins. *Biochem Cell Biol.* 84:549-558.
- Patterson, B.D., and D.D. Davies. 1969. Specificity of the enzymatic methylation of pea histone. *Biochem Biophys Res Commun.* 34:791-794.
- Pehrson, J.R., and V.A. Fried. 1992. MacroH2A, a core histone containing a large nonhistone region. *Science.* 257:1398-1400.
- Piacentini, L., L. Fanti, M. Berloco, B. Perrini, and S. Pimpinelli. 2003. Heterochromatin protein 1 (HP1) is associated with induced gene expression in *Drosophila* euchromatin. *J Cell Biol.* 161:707-714.
- Piacentini, L., and S. Pimpinelli. 2010. Positive regulation of euchromatic gene expression by HP1. *Fly (Austin).* 4:299-301.
- Pilyugin, M., J. Demmers, C.P. Verrijzer, F. Karch, and Y.M. Moshkin. 2009. Phosphorylation-mediated control of histone chaperone ASF1 levels by Tousled-like kinases. *PLoS One.* 4:e8328.
- Pokholok, D.K., C.T. Harbison, S. Levine, M. Cole, N.M. Hannett, T.I. Lee, G.W. Bell, K. Walker, P.A. Rolfe, E. Herbolsheimer, J. Zeitlinger, F. Lewitter, D.K. Gifford, and R.A.

- Young. 2005. Genome-wide map of nucleosome acetylation and methylation in yeast. *Cell*. 122:517-527.
- Postberg, J., S. Forcob, W.J. Chang, and H.J. Lipps. 2010. The evolutionary history of histone H3 suggests a deep eukaryotic root of chromatin modifying mechanisms. *BMC Evol Biol*. 10:259.
- Poveda, A., M. Pamblanco, S. Tafrov, V. Tordera, R. Sternglanz, and R. Sendra. 2004. Hif1 is a component of yeast histone acetyltransferase B, a complex mainly localized in the nucleus. *J Biol Chem*. 279:16033-16043.
- Prado, A., I. Ramos, L.J. Frehlick, A. Muga, and J. Ausio. 2004. Nucleoplasmin: a nuclear chaperone. *Biochem Cell Biol*. 82:437-445.
- Ramakrishnan, V. 1997. Histone H1 and chromatin higher-order structure. *Crit Rev Eukaryot Gene Expr*. 7:215-230.
- Ramakrishnan, V., J.T. Finch, V. Graziano, P.L. Lee, and R.M. Sweet. 1993. Crystal structure of globular domain of histone H5 and its implications for nucleosome binding. *Nature*. 362:219-223.
- Rangasamy, D., L. Berven, P. Ridgway, and D.J. Tremethick. 2003. Pericentric heterochromatin becomes enriched with H2A.Z during early mammalian development. *EMBO J*. 22:1599-1607.
- Rangasamy, D., I. Greaves, and D.J. Tremethick. 2004. RNA interference demonstrates a novel role for H2A.Z in chromosome segregation. *Nat Struct Mol Biol*. 11:650-655.
- Rasmussen, T.P., M.A. Mastrangelo, A. Eden, J.R. Pehrson, and R. Jaenisch. 2000. Dynamic relocalization of histone MacroH2A1 from centrosomes to inactive X chromosomes during X inactivation. *J Cell Biol*. 150:1189-1198.

- Ray-Gallet, D., J.P. Quivy, C. Scamps, E.M. Martini, M. Lipinski, and G. Almouzni. 2002. HIRA is critical for a nucleosome assembly pathway independent of DNA synthesis. *Mol Cell*. 9:1091-1100.
- Rea, S., F. Eisenhaber, D. O'Carroll, B.D. Strahl, Z.W. Sun, M. Schmid, S. Opravil, K. Mechtler, C.P. Ponting, C.D. Allis, and T. Jenuwein. 2000. Regulation of chromatin structure by site-specific histone H3 methyltransferases. *Nature*. 406:593-599.
- Reuter, G., M. Giarre, J. Farah, J. Gausz, A. Spierer, and P. Spierer. 1990. Dependence of position-effect variegation in *Drosophila* on dose of a gene encoding an unusual zinc-finger protein. *Nature*. 344:219-223.
- Richards, E.J., and S.C. Elgin. 2002. Epigenetic codes for heterochromatin formation and silencing: rounding up the usual suspects. *Cell*. 108:489-500.
- Richardson, R.T., O. Alekseev, O.M. Alekseev, and M.G. O'Rand. 2006. Characterization of the NASP promoter in 3T3 fibroblasts and mouse spermatogenic cells. *Gene*. 371:52-58.
- Richardson, R.T., I.N. Batova, E.E. Widgren, L.X. Zheng, M. Whitfield, W.F. Marzluff, and M.G. O'Rand. 2000. Characterization of the histone H1-binding protein, NASP, as a cell cycle-regulated somatic protein. *J Biol Chem*. 275:30378-30386.
- Richardson, R.T., D.C. Bencic, and M.G. O'Rand. 2001. Comparison of mouse and human NASP genes and expression in human transformed and tumor cell lines. *Gene*. 274:67-75.
- Ringrose, L., H. Ehret, and R. Paro. 2004. Distinct contributions of histone H3 lysine 9 and 27 methylation to locus-specific stability of polycomb complexes. *Mol Cell*. 16:641-653.
- Roh, T.Y., W.C. Ngau, K. Cui, D. Landsman, and K. Zhao. 2004. High-resolution genome-wide mapping of histone modifications. *Nat Biotechnol*. 22:1013-1016.



- Roh, T.Y., G. Wei, C.M. Farrell, and K. Zhao. 2007. Genome-wide prediction of conserved and nonconserved enhancers by histone acetylation patterns. *Genome Res.* 17:74-81.
- Roth, S.Y., J.M. Denu, and C.D. Allis. 2001. Histone acetyltransferases. *Annu Rev Biochem.* 70:81-120.
- Sanchez, Y., B.A. Desany, W.J. Jones, Q. Liu, B. Wang, and S.J. Elledge. 1996. Regulation of RAD53 by the ATM-like kinases MEC1 and TEL1 in yeast cell cycle checkpoint pathways. *Science.* 271:357-360.
- Sanematsu, F., Y. Takami, H.K. Barman, T. Fukagawa, T. Ono, K. Shibahara, and T. Nakayama. 2006. Asf1 is required for viability and chromatin assembly during DNA replication in vertebrate cells. *J Biol Chem.* 281:13817-13827.
- Santenard, A., C. Ziegler-Birling, M. Koch, L. Tora, A.J. Bannister, and M.E. Torres-Padilla. 2010. Heterochromatin formation in the mouse embryo requires critical residues of the histone variant H3.3. *Nat Cell Biol.* 12:853-862.
- Santos-Rosa, H., R. Schneider, A.J. Bannister, J. Sherriff, B.E. Bernstein, N.C. Emre, S.L. Schreiber, J. Mellor, and T. Kouzarides. 2002. Active genes are tri-methylated at K4 of histone H3. *Nature.* 419:407-411.
- Schulz, L.L., and J.K. Tyler. 2006. The histone chaperone ASF1 localizes to active DNA replication forks to mediate efficient DNA replication. *FASEB J.* 20:488-490.
- Schwabish, M.A., and K. Struhl. 2006. Asf1 mediates histone eviction and deposition during elongation by RNA polymerase II. *Mol Cell.* 22:415-422.
- Schwartz, B.E., and K. Ahmad. 2005. Transcriptional activation triggers deposition and removal of the histone variant H3.3. *Genes Dev.* 19:804-814.

- Shahbazian, M.D., and M. Grunstein. 2007. Functions of site-specific histone acetylation and deacetylation. *Annu Rev Biochem.* 76:75-100.
- Sharma, N., and J.K. Nyborg. 2008. The coactivators CBP/p300 and the histone chaperone NAP1 promote transcription-independent nucleosome eviction at the HTLV-1 promoter. *Proc Natl Acad Sci U S A.* 105:7959-7963.
- Sharp, J.A., E.T. Fouts, D.C. Krawitz, and P.D. Kaufman. 2001. Yeast histone deposition protein Asf1p requires Hir proteins and PCNA for heterochromatic silencing. *Curr Biol.* 11:463-473.
- Sherwood, P.W., S.V. Tsang, and M.A. Osley. 1993. Characterization of HIR1 and HIR2, two genes required for regulation of histone gene transcription in *Saccharomyces cerevisiae*. *Mol Cell Biol.* 13:28-38.
- Shibahara, K., and B. Stillman. 1999. Replication-dependent marking of DNA by PCNA facilitates CAF-1-coupled inheritance of chromatin. *Cell.* 96:575-585.
- Shiio, Y., and R.N. Eisenman. 2003. Histone sumoylation is associated with transcriptional repression. *Proc Natl Acad Sci U S A.* 100:13225-13230.
- Shilatifard, A. 2006. Chromatin modifications by methylation and ubiquitination: implications in the regulation of gene expression. *Annu Rev Biochem.* 75:243-269.
- Shogren-Knaak, M., H. Ishii, J.M. Sun, M.J. Pazin, J.R. Davie, and C.L. Peterson. 2006. Histone H4-K16 acetylation controls chromatin structure and protein interactions. *Science.* 311:844-847.
- Sillje, H.H., and E.A. Nigg. 2001. Identification of human Asf1 chromatin assembly factors as substrates of Tousled-like kinases. *Curr Biol.* 11:1068-1073.

- Singh, P.B., J.R. Miller, J. Pearce, R. Kothary, R.D. Burton, R. Paro, T.C. James, and S.J. Gaunt. 1991. A sequence motif found in a *Drosophila* heterochromatin protein is conserved in animals and plants. *Nucleic Acids Res.* 19:789-794.
- Smith, S., and B. Stillman. 1989. Purification and characterization of CAF-I, a human cell factor required for chromatin assembly during DNA replication in vitro. *Cell.* 58:15-25.
- Sogo, J.M., H. Stahl, T. Koller, and R. Knippers. 1986. Structure of replicating simian virus 40 minichromosomes. The replication fork, core histone segregation and terminal structures. *J Mol Biol.* 189:189-204.
- Song, J.J., J.D. Garlick, and R.E. Kingston. 2008. Structural basis of histone H4 recognition by p55. *Genes Dev.* 22:1313-1318.
- Stein, A., J.P. Whitlock, Jr., and M. Bina. 1979. Acidic polypeptides can assemble both histones and chromatin in vitro at physiological ionic strength. *Proc Natl Acad Sci U S A.* 76:5000-5004.
- Sterner, D.E., and S.L. Berger. 2000. Acetylation of histones and transcription-related factors. *Microbiol Mol Biol Rev.* 64:435-459.
- Stillman, B. 1986. Chromatin assembly during SV40 DNA replication in vitro. *Cell.* 45:555-565.
- Strahl, B.D., and C.D. Allis. 2000. The language of covalent histone modifications. *Nature.* 403:41-45.
- Strahl, B.D., R. Ohba, R.G. Cook, and C.D. Allis. 1999. Methylation of histone H3 at lysine 4 is highly conserved and correlates with transcriptionally active nuclei in *Tetrahymena*. *Proc Natl Acad Sci U S A.* 96:14967-14972.
- Straube, K., J.S. Blackwell, Jr., and L.F. Pemberton. 2010. Nap1 and Chz1 have separate Htz1 nuclear import and assembly functions. *Traffic.* 11:185-197.

- Stuwe, T., M. Hothorn, E. Lejeune, V. Rybin, M. Bortfeld, K. Scheffzek, and A.G. Ladurner. 2008. The FACT Spt16 "peptidase" domain is a histone H3-H4 binding module. *Proc Natl Acad Sci U S A*. 105:8884-8889.
- Sugimoto, K., T. Yamada, Y. Muro, and M. Himeno. 1996. Human homolog of Drosophila heterochromatin-associated protein 1 (HP1) is a DNA-binding protein which possesses a DNA-binding motif with weak similarity to that of human centromere protein C (CENP-C). *J Biochem*. 120:153-159.
- Suto, R.K., M.J. Clarkson, D.J. Tremethick, and K. Luger. 2000. Crystal structure of a nucleosome core particle containing the variant histone H2A.Z. *Nat Struct Biol*. 7:1121-1124.
- Szenker, E., D. Ray-Gallet, and G. Almouzni. 2011. The double face of the histone variant H3.3. *Cell Res*. 21:421-434.
- Tagami, H., D. Ray-Gallet, G. Almouzni, and Y. Nakatani. 2004. Histone H3.1 and H3.3 complexes mediate nucleosome assembly pathways dependent or independent of DNA synthesis. *Cell*. 116:51-61.
- Takahata, S., Y. Yu, and D.J. Stillman. 2009. FACT and Asf1 regulate nucleosome dynamics and coactivator binding at the HO promoter. *Mol Cell*. 34:405-415.
- Talbert, P.B., and S. Henikoff. 2010. Histone variants--ancient wrap artists of the epigenome. *Nat Rev Mol Cell Biol*. 11:264-275.
- Tan, B.C., C.T. Chien, S. Hirose, and S.C. Lee. 2006. Functional cooperation between FACT and MCM helicase facilitates initiation of chromatin DNA replication. *EMBO J*. 25:3975-3985.

- Tang, Y., M.V. Poustovoitov, K. Zhao, M. Garfinkel, A. Canutescu, R. Dunbrack, P.D. Adams, and R. Marmorstein. 2006. Structure of a human ASF1a-HIRA complex and insights into specificity of histone chaperone complex assembly. *Nat Struct Mol Biol.* 13:921-929.
- Tse, C., T. Sera, A.P. Wolffe, and J.C. Hansen. 1998. Disruption of higher-order folding by core histone acetylation dramatically enhances transcription of nucleosomal arrays by RNA polymerase III. *Mol Cell Biol.* 18:4629-4638.
- Turner, B.M. 2000. Histone acetylation and an epigenetic code. *Bioessays.* 22:836-845.
- Tyler, J.K., C.R. Adams, S.R. Chen, R. Kobayashi, R.T. Kamakaka, and J.T. Kadonaga. 1999. The RCAF complex mediates chromatin assembly during DNA replication and repair. *Nature.* 402:555-560.
- Tyler, J.K., M. Bulger, R.T. Kamakaka, R. Kobayashi, and J.T. Kadonaga. 1996. The p55 subunit of *Drosophila* chromatin assembly factor 1 is homologous to a histone deacetylase-associated protein. *Mol Cell Biol.* 16:6149-6159.
- Tyler, J.K., K.A. Collins, J. Prasad-Sinha, E. Amriott, M. Bulger, P.J. Harte, R. Kobayashi, and J.T. Kadonaga. 2001. Interaction between the *Drosophila* CAF-1 and ASF1 chromatin assembly factors. *Mol Cell Biol.* 21:6574-6584.
- Umehara, T., T. Chimura, N. Ichikawa, and M. Horikoshi. 2002. Polyanionic stretch-deleted histone chaperone *cia1/Asf1p* is functional both in vivo and in vitro. *Genes Cells.* 7:59-73.
- Umehara, T., and M. Horikoshi. 2003. Transcription initiation factor IID-interactive histone chaperone CIA-II implicated in mammalian spermatogenesis. *J Biol Chem.* 278:35660-35667.

- Vakoc, C.R., S.A. Mandat, B.A. Olenchok, and G.A. Blobel. 2005. Histone H3 lysine 9 methylation and HP1gamma are associated with transcription elongation through mammalian chromatin. *Mol Cell*. 19:381-391.
- Vakoc, C.R., M.M. Sachdeva, H. Wang, and G.A. Blobel. 2006. Profile of histone lysine methylation across transcribed mammalian chromatin. *Mol Cell Biol*. 26:9185-9195.
- van Attikum, H., and S.M. Gasser. 2009. Crosstalk between histone modifications during the DNA damage response. *Trends Cell Biol*. 19:207-217.
- van der Heijden, G.W., A.A. Derijck, E. Posfai, M. Giele, P. Pelczar, L. Ramos, D.G. Wansink, J. van der Vlag, A.H. Peters, and P. de Boer. 2007. Chromosome-wide nucleosome replacement and H3.3 incorporation during mammalian meiotic sex chromosome inactivation. *Nat Genet*. 39:251-258.
- VanDemark, A.P., M. Blanksma, E. Ferris, A. Heroux, C.P. Hill, and T. Formosa. 2006. The structure of the yFACT Pob3-M domain, its interaction with the DNA replication factor RPA, and a potential role in nucleosome deposition. *Mol Cell*. 22:363-374.
- VanDemark, A.P., H. Xin, L. McCullough, R. Rawlins, S. Bentley, A. Heroux, D.J. Stillman, C.P. Hill, and T. Formosa. 2008. Structural and functional analysis of the Spt16p N-terminal domain reveals overlapping roles of yFACT subunits. *J Biol Chem*. 283:5058-5068.
- Varga-Weisz, P., J. Zlatanova, S.H. Leuba, G.P. Schroth, and K. van Holde. 1994. Binding of histones H1 and H5 and their globular domains to four-way junction DNA. *Proc Natl Acad Sci U S A*. 91:3525-3529.
- Verreault, A. 2000. De novo nucleosome assembly: new pieces in an old puzzle. *Genes Dev*. 14:1430-1438.

- Verreault, A., P.D. Kaufman, R. Kobayashi, and B. Stillman. 1996. Nucleosome assembly by a complex of CAF-1 and acetylated histones H3/H4. *Cell*. 87:95-104.
- von Holt, C., W.F. Brandt, H.J. Greyling, G.G. Lindsey, J.D. Retief, J.D. Rodrigues, S. Schwager, and B.T. Sewell. 1989. Isolation and characterization of histones. *Methods Enzymol*. 170:431-523.
- Wang, H., L. Wang, H. Erdjument-Bromage, M. Vidal, P. Tempst, R.S. Jones, and Y. Zhang. 2004. Role of histone H2A ubiquitination in Polycomb silencing. *Nature*. 431:873-878.
- Wang, J., J. Mager, Y. Chen, E. Schneider, J.C. Cross, A. Nagy, and T. Magnuson. 2001. Imprinted X inactivation maintained by a mouse Polycomb group gene. *Nat Genet*. 28:371-375.
- Weiler, K.S., and B.T. Wakimoto. 1995. Heterochromatin and gene expression in Drosophila. *Annu Rev Genet*. 29:577-605.
- Welch, J.E., and M.G. O'Rand. 1990. Characterization of a sperm-specific nuclear autoantigenic protein. II. Expression and localization in the testis. *Biol Reprod*. 43:569-578.
- Whitehouse, I., A. Flaus, B.R. Cairns, M.F. White, J.L. Workman, and T. Owen-Hughes. 1999. Nucleosome mobilization catalysed by the yeast SWI/SNF complex. *Nature*. 400:784-787.
- Widlak, P., M. Kalinowska, M.H. Parseghian, X. Lu, J.C. Hansen, and W.T. Garrard. 2005. The histone H1 C-terminal domain binds to the apoptotic nuclease, DNA fragmentation factor (DFF40/CAD) and stimulates DNA cleavage. *Biochemistry*. 44:7871-7878.
- Widom, J. 1998. Chromatin structure: linking structure to function with histone H1. *Curr Biol*. 8:R788-791.

- Wiedemann, S.M., S.N. Mildner, C. Bonisch, L. Israel, A. Maiser, S. Matheisl, T. Straub, R. Merkl, H. Leonhardt, E. Kremmer, L. Schermelleh, and S.B. Hake. 2010. Identification and characterization of two novel primate-specific histone H3 variants, H3.X and H3.Y. *J Cell Biol.* 190:777-791.
- Wirbelauer, C., O. Bell, and D. Schubeler. 2005. Variant histone H3.3 is deposited at sites of nucleosomal displacement throughout transcribed genes while active histone modifications show a promoter-proximal bias. *Genes Dev.* 19:1761-1766.
- Wolffe, A.P., and J.J. Hayes. 1999. Chromatin disruption and modification. *Nucleic Acids Res.* 27:711-720.
- Wondrak, G.T., D. Cervantes-Laurean, E.L. Jacobson, and M.K. Jacobson. 2000. Histone carbonylation in vivo and in vitro. *Biochem J.* 351 Pt 3:769-777.
- Woodcock, C.L. 2006. Chromatin architecture. *Curr Opin Struct Biol.* 16:213-220.
- Woodcock, C.L., and S. Dimitrov. 2001. Higher-order structure of chromatin and chromosomes. *Curr Opin Genet Dev.* 11:130-135.
- Wreggett, K.A., F. Hill, P.S. James, A. Hutchings, G.W. Butcher, and P.B. Singh. 1994. A mammalian homologue of *Drosophila* heterochromatin protein 1 (HP1) is a component of constitutive heterochromatin. *Cytogenet Cell Genet.* 66:99-103.
- Wu, R., A.V. Terry, P.B. Singh, and D.M. Gilbert. 2005. Differential subnuclear localization and replication timing of histone H3 lysine 9 methylation states. *Mol Biol Cell.* 16:2872-2881.
- Ye, X., B. Zerlanko, R. Zhang, N. Somaiah, M. Lipinski, P. Salomoni, and P.D. Adams. 2007. Definition of pRB- and p53-dependent and -independent steps in HIRA/ASF1a-mediated formation of senescence-associated heterochromatin foci. *Mol Cell Biol.* 27:2452-2465.



- Yuan, W., M. Xu, C. Huang, N. Liu, S. Chen, and B. Zhu. 2011. H3K36 methylation antagonizes PRC2-mediated H3K27 methylation. *J Biol Chem.* 286:7983-7989.
- Yun, M., J. Wu, J.L. Workman, and B. Li. 2011. Readers of histone modifications. *Cell Res.* 21:564-578.
- Zabaronick, S.R., and J.K. Tyler. 2005. The histone chaperone anti-silencing function 1 is a global regulator of transcription independent of passage through S phase. *Mol Cell Biol.* 25:652-660.
- Zhang, R., S.T. Liu, W. Chen, M. Bonner, J. Pehrson, T.J. Yen, and P.D. Adams. 2007. HP1 proteins are essential for a dynamic nuclear response that rescues the function of perturbed heterochromatin in primary human cells. *Mol Cell Biol.* 27:949-962.
- Zhang, R., M.V. Poustovoitov, X. Ye, H.A. Santos, W. Chen, S.M. Daganzo, J.P. Erzberger, I.G. Serebriiskii, A.A. Canutescu, R.L. Dunbrack, J.R. Pehrson, J.M. Berger, P.D. Kaufman, and P.D. Adams. 2005. Formation of MacroH2A-containing senescence-associated heterochromatin foci and senescence driven by ASF1a and HIRA. *Dev Cell.* 8:19-30.
- Zhang, Y. 2004. Molecular biology: no exception to reversibility. *Nature.* 431:637-639.
- Zhang, Y., and D. Reinberg. 2001. Transcription regulation by histone methylation: interplay between different covalent modifications of the core histone tails. *Genes Dev.* 15:2343-2360.
- Zhang, Z., K. Shibahara, and B. Stillman. 2000. PCNA connects DNA replication to epigenetic inheritance in yeast. *Nature.* 408:221-225.
- Zlatanova, J., P. Caiafa, and K. Van Holde. 2000. Linker histone binding and displacement: versatile mechanism for transcriptional regulation. *FASEB J.* 14:1697-1704.

Zlatanova, J., C. Seebart, and M. Tomschik. 2007. Napl: taking a closer look at a juggler protein of extraordinary skills.