

# Investigations éco-épidémiologiques et génétiques des Lyssavirus et des Paramyxovirus chez les micromammifères du sud-ouest de l'océan Indien

Julien Mélade

#### ▶ To cite this version:

Julien Mélade. Investigations éco-épidémiologiques et génétiques des Lyssavirus et des Paramyxovirus chez les micromammifères du sud-ouest de l'océan Indien. Maladies infectieuses. Université de la Réunion, 2015. Français. NNT: 2015LARE0030. tel-01326944

## HAL Id: tel-01326944 https://theses.hal.science/tel-01326944

Submitted on 11 Jul 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.













# Investigations éco-épidémiologiques et génétiques des Lyssavirus et des Paramyxovirus chez les micromammifères du Sud-Ouest de l'Océan Indien

#### **THÈSE**

Pour l'obtention du grade de

DOCTEUR DE L'UNIVERSITE DE LA REUNION

Spécialité : Microbiologie

École Doctorale des Sciences, Technologies et Santé

Présentée et soutenue publiquement par

#### Julien MÉLADE

Le 08 Décembre 2015, devant le Jury composé de :

Pierre-Emmanuel CECCALDI, Professeur, Université de Paris, Institut Pasteur Rapporteur

Serge MORAND, Directeur de recherche, CNRS, Montpellier Rapporteur

Steven M. GOODMAN, Professeur, Université d'Antananarivo, Examinateur

MacArthur Field Biologist, Field Museum de Chicago

Philippe DESPRÈS, Professeur, Université de La Réunion, CNRS, INSERM, IRD Examinateur

Hervé PASCALIS, Docteur, Université de La Réunion, CNRS, INSERM, IRD Co-encadrant

Koussay M. DELLAGI, Professeur de médecine à l'Université de Tunis, Institut Pasteur Directeur de thèse

Cette thèse a reçu le soutient financier de la Région Réunion et de l'Union Européenne (Fonds Européen de Développement Régional – FEDER) dans le cadre du Programme Opérationnel de Coopération Territoriale.

#### Remerciements

Je tiens en premier lieu à remercier mon directeur de thèse, Koussay Dellagi. Je vous remercie de m'avoir guidé, d'avoir été disponible tout le temps, même après votre départ, et de m'avoir apporté toutes les bases et les connaissances scientifiques dans ce domaine qui m'ont grandement été utiles pour l'aboutissement de mes 3 années de recherche. Merci de m'avoir offert l'opportunité d'intégrer le monde de la recherche.

Un grand merci à Patrick Mavingui, directeur de l'UMR PIMIT, pour m'avoir accompagné surtout vers la fin. Je vous remercie de m'avoir motivé, pour avoir été disponible soir et weekend. Pour vos mails et message pour prendre de nos nouvelles. Pour un doctorant, recevoir des encouragements de la part d'un de ses directeurs c'est quelque chose de vraiment remotivant. Pour tout cela, je vous remercie énormément.

Je remercie également Hervé Pascalis pour son encadrement.

J'adresse toute ma reconnaissance envers mes membres du jury de thèse, Serge Morand et Pierre-Emmanuel Ceccaldi qui ont eu l'amabilité d'être tous deux rapporteurs, ainsi que Philipe Desprès et Steve Goodman en tant qu'examinateurs. Je vous remercie d'avoir accepté d'évaluer ces travaux.

Je tiens à remercier Steve Goodman, Beza Ramasindrazana ainsi que toute l'association Vahatra, pour les collaborations que vous avez entretenues depuis toutes ces années avec le laboratoire CRVOI devenu maintenant PIMIT. Merci pour votre aide et surtout votre efficacité sur le terrain, toutes les missions de collectes, les sacrifices etc... Je vous suis sincèrement reconnaissant car, aujourd'hui, si je peux prétendre au titre de Docteur, c'est en grande partie grâce à votre travail de terrain.

Je remercie Wanda Markotter pour m'avoir accueilli à l'Université de Pretoria en Afrique du Sud. Merci de m'avoir montré toute la beauté de l'Afrique du Sud. Merci à Stewart McCulloch et Terence Scott pour m'avoir fait découvrir Pretoria et de m'avoir initié au « cricket sud-africain ». J'adhère!

Je remercie également Christian Drosten de m'avoir accueilli au Centre Hospitalier Universitaire de Bönn en Allemagne. Je remercie Marcel Müller du laboratoire pour m'avoir

accompagné pendant ce séjour et m'avoir fait découvrir Bönn et tous ses coins chaleureux. Je reviendrais!

Un grand merci à Xavier de Lamballerie de la Fac de Médecine à la Timone, qui m'a initié à la virologie et sans qui je ne serais peut-être pas là aujourd'hui. Merci d'avoir continué cette collaboration pendant ces 3 années de thèse. Merci à toi Gégé d'avoir pu garder ce lien que nous avons tissé en Métropole. Merci de m'avoir aidé pour le séquençage de tous ces virus. Je t'avoue que c'était galère mais nous y sommes parvenus!

Merci à toi Pablo pour m'avoir conseillé, motiver et pour avoir partagé ton expérience personnelle du monde de la recherche mais aussi en dehors.

Merci aux anciens post-docs du laboratoire pour m'avoir conseillé, encourager et même parfois supporter! Mes pensées vont particulièrement envers David, Camille et Muriel. David, merci pour tous tes conseils et ton extrême gentillesse. Tes connaissances m'impressionneront toujours. Merci Camille pour tes encouragements surtout vers la fin, moment très difficile. J'espère que cela suffise pour que tu ne me poses pas de question pour ma soutenance! Muriel, merci pour tes précieux conseils et cette balade périlleuse à Johannesburg. Quelle aventure! On est en vie!

Erwan, merci de ta gentillesse, de ta confiance la bonne humeur et la joie que tu as apportées au cours de ces 3 années. Merci pour tout.

Merci Magali pour tes coups de pouce qui m'ont beaucoup aidés.

Yann, mon compagnon « d'infortune », mon acolyte, et surtout mon ami, pour avoir été à mes côtés durant ces 3 années, je te remercie du plus profond de moi. J'ai appris énormément tant sur le côté scientifique que personnel grâce à toi. Merci pour tous ces moments. Je te souhaite le meilleur pour la suite. Cette petite guerre entre l'équipe virus et l'équipe bactérie va vraiment me manquer et tu vas me manquer. Je t'attends à la fin de cette thèse pour un dernier galopin et un dernier « magazine à  $2 \in$  ».

Claire Tessier, je suis désolé pour la frayeur que je t'ai causée la dernière fois, je suis désolé d'en avoir rigolé tout de suite après, et je suis désolé d'en rigoler encore ... Plus sérieusement, Claire, comment pourrais-je t'oublier? Je te suis vraiment reconnaissant d'avoir été là, de m'avoir encouragé, de m'avoir poussé, merci infiniment pour tous ces moments passés

ensemble. Toi aussi, tu vas énormément me manquer par la suite. Et bien sûr, je ne t'oublierais jamais. C'était vraiment « mignon mignon » tous ces petits moments !

Merci à toi Lisa, d'avoir été là tout le temps. Merci de m'avoir boosté. Merci pour ces moments que ce soit ici, ou en Métropole. Merci pour tout! Ta présence nous manque énormément.

Claire B, Catherine, Camille « fille », merci pour tous les moments passés ensembles, pour tous vos conseils. Claire, merci d'avoir pu m'enseigner les bases des statistiques. Je pense qu'on devrait appeler ces moments « les statistiques pour les nuls ».

Léon, Etienne, Miora, futurs docteurs, et amis, je vous remercie d'avoir été là pendant ces quelques années. Miora, merci d'avoir apporté ta présence féminine quand toutes les autres étaient parties. Merci à vous trois d'avoir apporté de la bonne humeur au bureau surtout quand Yann et moi rédigions ce qui n'était pas forcément un moment facile. Je vous souhaite une bonne fin de thèse, entraidez-vous, surtout vers la fin, et je vous souhaite une très bonne continuation pour l'après.

Un grand merci à toute l'équipe du CYROI, à son directeur Christian Meriau pour m'avoir accueilli sur la plateforme. Un grand merci à Jean Turquet, Mayalen Zubia et Régis Roche avec qui m'ont permet d'effectuer mon 1<sup>er</sup> stage sur la plateforme et qui ont attisé la petite flamme pour la Recherche qui était en moi. Merci à toutes les différentes équipes et à toutes les personnes qui y travaillent au quotidien, pour m'avoir encouragé, motivé, et soutenu. Merci Giovédie pour toute ces pauses cafés ou tu as était ma confidente, et m'avoir aidé à surmonter les difficultés. Merci à Imade, Johan, Florian, les 3 acolytes, Merci à Wildriss, Marjo, Alina, Fanny, Aurélie, Brice, Géraldine et Marie pour votre soutien au quotidien. Merci à tous les stagiaires, particulièrement à Anne claire Dorsemans.

Enfin, j'adresse mes plus sincères remerciement à mes parents, ma maman et mon papa qui m'ont soutenu à fond jusqu'au bout (Maman, pour tes appels téléphonique à 5h57 du matin où j'étais encore au labo, merci). A mes frères qui m'ont soutenu. Ce manuscrit ne serait suffisant pour décrire tout ce que vous avez fait pour moi, jusqu'à maintenant. Pour vos sacrifices et votre amour, je me battrais jusqu'au bout! Mi aime zot toute!

Merci à tous ceux que je n'ai pas cités mais qui ont contribué, de près ou de loin à l'aboutissement de ce travail.

# Table des matières

Remerciements	5
Table des matières	9
Liste des abréviations	11
Liste des tableaux	14
Liste des figures	16
Résumé	18
Abstract	20
PARTIE I. ETAT DE L'ART	22
Chapitre 1. Les maladies infectieuses émergentes	24
1.1. Généralités	24
1.1.1. Exemple d'agent émergent : le virus EBOLA	
1.1.2. Exemple d'agent émergent : le SARS-Coronavirus	
1.1.3. Réservoirs et sauts d'espèces	
1.2. Différentes étapes dans l'émergence et facteurs impliqués	
1.2.1. Facteurs anthropogéniques	
1.2.2. Facteurs écologiques	
1.2.3. Facteurs génétiques	
1.2.3.1. Macro-évolution	
1.2.3.2. Micro-évolution.	
Chapitre 2. Micromammifères sauvages un réservoir d'agents zoonotiques	39
2.1. Généralités	40
2.2. Les chauves-souris	
2.2.1. Caractéristiques	
2.2.2. Réservoirs d'agents pathogènes	
2.3. Les petits mammifères terrestres	
2.3.1. Caractéristiques	48
2.3.2 Réservoirs d'agents pathogènes et perte de la biodiversité	
Chapitre 3. Caractéristiques de la région du Sud-ouest de l'Océan Indien	53
3.1. Généralités	53
3.2. Les chauves-souris	54
3.3. Les petits mammifères terrestres	
PARTIE II. TRAVAUX DE RECHERCHE	61
Présentation du sujet, des objectifs et questions de recherche	62
Chapitre 4. Étude de la dynamique d'infection des paramyxovirus chez les petits mammifères et les chauves-souris de la zone Sud-Ouest Océan Indien	66

4.1. Généralités	67
4.2. Approches expérimentales	
4.3 Etude de la dynamique d'infection au sein des micromammifères de la zone SOOI	
4.4. Conclusion du chapitre 4	
Chapitre 5. Les mécanismes moléculaires et les facteurs environnementaux associés à	à la
diversité des paramyxovirus chez les chauves-souris de Madagascar	
5.1. Introduction	. 103
5.2. Etude éco-épidémiologique d'une communauté de chauves-souris de Madagascar : Analyses des processus macro-évolutifs et facteurs écologique associés à la transmission	
virale	. 105
5.3. Conclusion du chapitre 5	. 132
Chapitre 6. Etude des lyssavirus chez les chauves-souris de la zone Sud-Ouest de	
l'Océan Indien	. 136
6.1. Généralités	. 136
6.2. Approches expérimentales	
6.3. Mise en évidence sérologique d'une circulation importante de lyssavirus chez les	
chauves-souris de la zone du Sud-ouest de l'Océan Indien.	. 140
6.4. Conclusion du chapitre 6	. 175
Discussion générale	. 177
Conclusion et perspectives	. 189
Bibliographie	. 193
Annexes	. 211
Résumé	. 285

#### Liste des abréviations

ADN : Acide désoxyribonucléique AIC : Akaike Information Criterion

AICc: Corrected Akaike Information Criterion

ARN: Acide ribonucléique

BEAST: Bayesian Evolutionary Analysis Sampling Trees

cDNA: Complementary DNA

EID: Emerging Infectious Disease

ESS: Effective sample size values

FAO: Food and Agriculture Organization

GLM: Generalized Linear Model

GLMM: Generalized Linear Mixed Model

GPS: Global positioning system

Kb: Kilobase

MCMC: Markov Chain Monte Carlo

MAT: Mean average temperature

MAR: Mean average rainfall

Mya: Million years ago

OIE : Office international des épizooties, devenu maintenant Organisation mondiale de la

santé animale

OTU: Operational taxonomic units

PAR: Pneumovirus Avulavirus Rubulavirus

PCR: Polymerase chain reaction

PPR: Peste des petits ruminants

PVs: Paramyxoviruses

qRT-PCR: Real time reverse transcriptase polymerase chain reaction

RFFIT: Rapid fluorescent focus inhibition test

RMH: Respirovirus Morbillivirus Hénipavirus

RT-PCR: Reverse transcriptase polymerase chain reaction

SOOI: Sud-ouest Océan Indien

SWIO: South Western Indian Ocean islands

UMRVs: Unclassified Morbilli-Related viruses

UHRV: Unclassified Henipa-Related viruses

# Liste des tableaux

Tableau 1. Exemple de virus isolés de chauves-souris	. 45
Tableau 2. Aperçu des différentes maladies et agents pathogènes retrouvés chez	
les rongeurs.	. 50
Tableau 3. Caractéristiques et particularités des chauves-souris collectées à Madagascar	. 56
Tableau 4. Caractéristiques et particularités des chauves-souris collectées	
dans les autres îles de la zone SOOI.	. 57
Tableau 5. Caractéristiques et particularités des petits mammifères terrestres	
collectées à Madagascar.	. 59

# Liste des figures

Figure 1. Place des maladies infectieuses parmi les causes de mortalité/morbidité	
dans le monde.	25
Figure 2. Exemples mondiaux de maladies émergentes ou ré-émergentes	28
Figure 3. Différentes étapes d'émergences d'agents zoonotiques.	30
Figure 4. Continuum entre populations d'hôtes-agents zoonotiques et facteurs d'én	mergences.
	32
Figure 5. Transmission du virus Nipah chez l'homme par les chiroptères via la fau	ine
domestique.	34
Figure 6. Mécanismes évolutifs hôte-parasites.	36
Figure 7. Colonie de Mormopterus francoismoutoui (insectivores) dans la grotte d	e Trois-
Bassins à La Réunion.	43
Figure 8. Colonie de <i>Pteropus scapularis</i> (frugivores) perchés sur un arbre	46
Figure 9. Les îles du Sud-ouest de l'Océan Indien	54

#### Résumé

La faune sauvage a été depuis longtemps incriminée dans la survenue de zoonoses et joue le rôle de réservoir d'agents pathogènes (virus *Nipah*, *Hendra*, *Ebola*, *Hantaan* etc.) pour l'homme. Les îles tropicales et subtropicales du Sud-Ouest de l'Océan Indien (SOOI) constituent l'une des 34 régions reconnues comme « *hotspot* » de biodiversité au niveau mondial. Elles sont caractérisées par un très fort endémisme de la faune sauvage surtout sur l'Ile de Madagascar. Le caractère multi-insulaire de la région du SOOI, la diversité de ses biotopes et ses disparités biogéographiques et humaines offrent un champ d'investigation unique pour explorer « *in natura* » la dynamique évolutive des agents infectieux et les relations hôtes-virus.

Nos travaux de recherche ont porté sur deux modèles de virus à ARN de polarité négative, les paramyxovirus et les lyssavirus. Le premier modèle viral nous a permis d'aborder les questions relatives à la dynamique de transmission virale au sein de communauté d'hôtes, plus particulièrement, les chauves-souris et les petits mammifères terrestres de Madagascar et d'identifier les facteurs agissant sur cette dynamique de transmission et de diversification virale, en particulier les facteurs bio-écologiques associés à leurs hôtes. Le second modèle viral, les lyssavirus, nous a permis de décrire sur l'ensemble des îles du SOOI échantillonés, la circulation virale dans ce système multi-insulaire diversifié, au sein des chauves-souris dont la plupart des espèces sont endémiques à cette région.

Dans l'ensemble, nos investigations ont permis de mettre en évidence des échanges viraux (« host-switch ») importants entre chauves-souris, petits mammifères terrestres endémiques de Madagascar et les rongeurs introduits, le rôle de ces mammifères en tant que réservoir viral majeur et souligner le rôle disséminateur de Rattus rattus. Par ailleurs, nous avons pu identifier ce phénomène de « host-switch » comme étant le mécanisme macro-évolutif prépondérant et l'importance des facteurs biotiques et abiotiques à l'origine de la dynamique de transmission et de la diversification virale observée chez les paramyxovirus de chauves-souris de Madagascar.

**Mots-clés**: faune sauvage, zoonoses, SOOI, biodiversité, lyssavirus, paramyxovirus, diversité virale, réservoir viral, disséminateur viral, « *host-switch* » facteurs écologique

#### **Abstract**

For several decades, the animal have been incriminated as an important reservoir of many zoonotic pathogens (*Nipah*, *Hendra*, *Ebola*, *Hantaan* viruses etc.) at risk for humans. Tropical and subtropical islands of the South Western Indian Ocean (SWIO) are comprise on of 34 areas of the world recognized as biodiversity "hotspot". They are characterized by high levels of animal endemism, particularly on Madagascar. The multi-island structure of the SWIO region, the diversity of its biotopes, and differences in aspects related to human culture, offer a unique opportunity to investigate "*in natura*" the evolutionary dynamics of infectious agents and associated host-virus relationships.

Our research has focused on two models of negative RNA viruses, paramyxoviruses, and lyssaviruses. The first virus model allowed us to address issues related to the dynamics of viral transmission within a host community, in particular bats and Malagasy small terrestrial mammals, and to identify the driving factors, especially bio-ecological aspects associated with their hosts, affecting the dynamic of transmission and viral diversification. The second model allowed us to describe on the islands, the intense circulation of bat lyssaviruses in this system, including bats endemic to this region.

Overall, our investigations highlighted: (i) intense viral exchanges ("host-switch") between bats, endemic terrestrial small mammals, and introduced rodents on Madagascar, (ii) the role of these mammals as major viral reservoir, and (iii) the key role played by *Rattus rattus* as viral spreader. Furthermore, we identified both the phenomenon of "host-switch" as the major macro-evolutionary mechanism among bat paramyxoviruses from Madagascar and the importance of biotic and abiotic factors in shaping the transmission dynamics and viral diversification.

**Keywords**: wild fauna, zoonoses, SWIO, biodiversity, lyssavirus, paramyxoviruses, viral diversity, reservoir, viral spreader, "*host-switch*" ecological factors

#### PARTIE I. ETAT DE L'ART

L'état de l'art sera présenté en 3 chapitres:

Dans le chapitre I nous aborderons les notions générales sur les maladies infectieuses émergentes, les principales étapes d'émergences et les facteurs d'émergences dans le but de mieux comprendre et appréhender leur survenue.

Dans le chapitre II, nous nous focaliserons sur les agents zoonotiques retrouvés chez les chauves-souris et les petits mammifères ainsi que l'importance de leurs caractéristiques biologiques dans le maintien et la transmission d'agents zoonotiques.

Dans le chapitre III, nous détaillerons les caractéristiques bio-écologiques de la zone Sudouest de l'Océan, notamment la biodiversité résidente en micromammifères, en mettant un accent sur la faune sauvage de Madagascar et particulièrement sur les chauves-souris et les petits mammifères endémiques qui y vivent.

# Chapitre 1. Les maladies infectieuses émergentes


### 1.1. Généralités

L'Organisation Mondiale de la Santé (OMS) définit les maladies infectieuses émergentes comme des maladies survenant pour la première fois dans une population ou qui ont déjà sévi mais dont la prévalence augmente à nouveau à un niveau élevé (ré-émergentes) présentant une grave menace pour les populations exposées. Par le passé, les plus dangereuses d'entre elles ont provoqué chez l'homme des pandémies de choléra, de variole, de rougeole ou encore de grippe causant la mort de millions de personnes dans le monde. Les études récentes ont mis en évidence une augmentation significative du nombre d'émergences infectieuses depuis les années 1940, et des auteurs ont recensé en 2008 plus de 300 maladies infectieuses (Jones et al., 2008). Les maladies qui sont naturellement transmises de l'animal à l'homme et vice versa, appelées zoonoses, constituent plus de la moitié des maladies infectieuses émergentes (Jones et al., 2008). Plus de 75% d'entre elles ont pour origine la faune sauvage (Taylor et al., 2001). Par ailleurs, on sait que plus de 25% de ces maladies seraient dues à des pathogènes viraux (Taylor et al., 2001; Jones et al., 2008). A ce jour, il est estimé que sur les 57 millions de morts par an dans le monde, environ 14,9 millions seraient dus aux maladies infectieuses, soit plus de 25% (Muray & Lopez 1996; Fauci et al., 2001) (Figure 1) avec un impact plus important dans les pays en développement (Guerrant & Blackwood, 1999). On estime à plus de 1400 le nombre d'agents infectieux pathogènes pour l'homme (Taylor et al., 2001). En 2001, des auteurs ont rapporté qu'en moyenne, trois à quatre nouveaux virus pathogènes étaient détectés en population humaine chaque année, et la plupart de ces nouveaux agents provenaient des espèces animales (Woolhouse & Gaunt, 2007).

#### 1.1.1. Exemple d'agent émergent : le virus EBOLA

Le virus *Ebola* de la famille des *Filoviridae*, a récemment ré-émergé entre 2013 et 2015 en Afrique de l'Ouest là où le virus est endémique, provoquant une fièvre hémorragique grave avec une issue fatale fréquente chez les humains, (plus de 11000 décès sur 23000 cas [WHO, 2015]) et plusieurs espèces de primates non humains dont les grands singes.

Les chauves-souris frugivores étaient considérées comme le réservoir naturel du virus Ebola en Afrique (Leroy et al., 2005). Cependant, au cours de l'épidémie survenue en 2015, les chauves-souris insectivores ont été fortement incriminées et leur contact avec l'homme serait à l'origine de la ré-émergence récente du virus (Marí Saéz et al., 2014).



**Figure 1.** Place des maladies infectieuses parmi les causes de mortalité/morbidité dans le monde (Morens et al., 2004).

#### 1.1.2. Exemple d'agent émergent : le SARS-Coronavirus

Un second exemple saisissant d'émergence concerne le Syndrome Respiratoire Aigu Sévère (SRAS) provoqué par un virus de la famille des *Coronaviridae* qui a émergé dans les populations humaines en Chine en 2003 faisant plus de 8000 cas et plus de 700 décès (WHO, 2014). Des analyses sur des échantillons d'animaux sauvages vendus comme viande de brousse sur des marchés de la province de Guangdong (Chine) ont détecté la présence de coronavirus du SRAS chez des civettes (Guan et al., 2003) et des chauves-souris insectivores (Lau et al., 2005). Un scénario d'émergence a été proposé incriminant un contact indirect entre l'homme et la faune sauvage par l'intermédiaire d'animaux domestiqués, les civettes.

#### 1.1.3. Réservoirs et sauts d'espèces

De nombreux autres virus illustrés sur la **Figure 2** sont des exemples bien documentés de zoonoses qui ont eu aussi, ou menacent d'avoir, un impact sur la santé humaine (Morens et al., 2004).

Le point commun à toutes ces émergences virales reste le franchissement de la barrière d'espèce des virus de la faune sauvage ou domestique, considérés comme des réservoirs, vers la population humaine. Nous utilisons le terme réservoir selon la définition de Drexler et al. comme un taxon animal naturellement infecté de façon durable, au niveau populationnel et au-delà des limites géographiques, par une diversité d'agents infectieux, et pouvant ou non exprimer la maladie (Drexler et al., 2012). L'émergence d'un pathogène dans une espèce nouvelle (et pas uniquement dans la population humaine) va impliquer un saut de celui-ci depuis son hôte originel « réservoir » dans lequel il est déjà établi, vers un hôte différent. Ce phénomène est communément appelé « franchissement ou saut de la barrière d'espèces » (« host-switch ») et les mécanismes qui en rendent compte ne sont pas encore très bien compris (Morens et al., 2004 ; Longdon et al., 2011).




Figure 2. Exemples mondiaux de maladies émergentes ou ré-émergentes (Morens et al., 2004).

# 1.2. Différentes étapes dans l'émergence et facteurs impliqués

Morse et al. (2012), en s'appuyant sur le modèle de Daszak et al. (2000), ont décomposé le processus pouvant conduire à une zoonose chez l'homme en 3 étapes (**Figure 3**). La première étape, pré-émergence, correspond aux perturbations écologique, sociale, socio-économique ou encore génétique qui modifient la dynamique de transmission d'un agent infectieux à l'intérieur de son réservoir, le poussant à se multiplier et augmenter sa densité, à diffuser dans de nouvelles régions, et à de nouveaux hôtes non humains, le plus souvent de la faune domestique (épizooties). Il s'agit de l'étape à laquelle nous nous sommes particulièrement intéressés aux cours de ces travaux. La seconde étape correspond au phénomène de saut d'espèce du pathogène depuis les réservoirs de la faune sauvage ou domestique vers les populations humaines, c'est l'étape d'émergence. Enfin, la dernière étape correspond à la transmission du pathogène entre populations humaines, à son expansion au niveau mondial par divers moyen de transport, c'est l'étape de pandémie.





Figure 3. Différentes étapes d'émergences d'agents zoonotiques (Morse et al., 2012).

De nombreuses études ont ainsi décrit des facteurs écologiques et des processus évolutifs qui seraient potentiellement impliqués dans le phénomène de sauts d'espèce. L'un des facteurs capitale serait le changement dans l'environnement ou dans le comportement des hôtes réservoirs qui pourrait permettre que de nouvelles niches écologiques soient exploitées par des agents infectieux hébergés *via* des sauts entre espèces voisines compatibles et/ou permissives. Ces changements provoqueraient un déséquilibre entre un hôte et un agent pathogène évoluant dans un écosystème donné (Schrag & Wiener 1995; Daszak et al., 2000; Morand et al., 2006; Plowright et al., 2015) (**Figure 4**).



**Figure 4.** Continuum entre populations d'hôtes-agents zoonotiques et facteurs d'émergences (Daszak et al., 2000).

#### 1.2.1. Facteurs anthropogéniques

modifications des écosystèmes, Les les comportements humains, les activités socioéconomiques et les bouleversements démographiques (développement économique, déplacements massifs de populations, échanges internationaux, intensification des élevages), les résistances aux antibiotiques ou encore des catastrophes naturelles sont autant de facteurs pouvant conduire à ce déséquilibre (Daszak et al., 2000 ; Woolhouse et al., 2005 ; Food and Agriculture Organisation of the United Nations, 2011). C'est typiquement le cas des évènements de cohabitation générés par la levée du filtre de rencontre entre espèces allopatriques, devenues alors sympatriques au sein des mêmes écosystèmes.

L'épidémie du virus Nipah illustre clairement l'influence des comportements humains et des perturbations environnementales sur l'apparition de maladies émergentes infectieuses. Le virus Nipah (Famille: Paramyxoviridae, Genre: Henipavirus) a été décrit pour la première fois suite à une épidémie majeure chez les porcs et les humains dans la péninsule malaise entre septembre 1998 et avril 1999. Plus d'un million de porcs ont été abattus pour contenir l'épidémie. Au moins 265 cas humains et 105 décès ont été signalés (Chua et al., 2000). Le contact direct avec des porcs infectés a été identifié comme le principal mode de contamination chez l'homme. Les chauves-souris du genre Pteropus ont été reconnues comme hôtes naturels du virus (Johara et al., 2001). On sait que c'est l'entassement des cochons dans des enclos situés à proximité des colonies de chauves-souris frugivores du genre Pteropus porteuses qui en est la cause majeure. Cette proximité entre les deux espèces était elle-même la conséquence de la perte par les chauves-souris de leur habitat naturel et des ressources alimentaires frugales détruits par la déforestation malaise et indonésienne en faveur des programmes de plantation intensive de palmiers à huile. Il en a résulté une migration des chauves-souris vers les enclos de porcs à la recherche de nouvelles sources de nourriture (Figure 5). Il a été démontré que les gouttelettes d'urines émises en vol par ces chauvessouris contiennent des traces du virus qui ont pu constituer des sources de contamination des porcs. Le virus est secondairement transmis par aérosol, directement entre porcs et des porcs infectés vers l'homme (Chua et al., 2000).




Figure 5. Transmission du virus Nipah chez l'homme par les chiroptères via la faune domestique.

#### 1.2.2. Facteurs écologiques

A l'échelle de la faune, la richesse et la diversité d'hôtes et d'agents zoonotiques au sein d'une communauté ont été décrites comme des facteurs importants dans le maintien et la transmission d'agents zoonotiques (Ostfeld, 2000; Woolhouse et al., 2005; Keusch et al., 2009). Keesing et al. (2010) ont par ailleurs souligné que les changements en richesse spécifique associés aux modifications que les facteurs essentiellement anthropogéniques imposent à la biodiversité peuvent impacter les évènements d'émergences infectieuses par un effet de dilution ou un effet d'amplification. L'analyse de tels facteurs est déterminante dans la compréhension de la transmission d'agents zoonotiques. Se concentrer ainsi non pas sur une seule espèce, mais toute une communauté d'hôtes permet de comprendre le rôle de chaque espèce dans la séquence des interactions hôtes-pathogènes (Morand et al., 2014). Par ailleurs, de nombreuses études ont montré l'importance des facteurs biotiques et abiotiques dans la transmission d'agents zoonotiques. O'Shea et al. ont décrit par exemple l'importance de l'âge, du sexe, de la saison et du climat dans la circulation du virus de la rage chez une communauté de chauves-souris au Colorado (O'Shea et al., 2014). Streicker et al. ont rapporté le rôle du climat tropical et subtropical favorisant la transmission inter-espèces du virus de la rage chez les chauves-souris (Streicker et al., 2012). George et al. ont montré que l'état de torpeur pendant les périodes hivernales diminuait l'activité virale chez les chauves-souris (Georges et al., 2011).

# 1.2.3. Facteurs génétiques

Plus loin dans leur étude, Morse et al. (2012) identifient un ensemble de facteurs génétiques qui apparaissent liés au phénomène d'émergence, parmi lesquelles on peut citer : i) la parenté phylogénétique avec l'hôte primaire hébergeant un agent zoonotique qui favoriserait le saut d'espèce d'un virus (Streicker et al., 2010), comme c'est le cas du virus de l'immunodéficience humaine (VIH) des chimpanzés, ii) la parenté phylogénétique entre agents pathogènes de la faune sauvage et ceux infectant les populations humaines (Pulliam, 2008) iii) le large spectre d'hôte ainsi qu'une certaine plasticité génétique virale (mutation, recombinaison, réassortiment) génèrent une diversité populationnelle (Woolhouse et al., 2001 ; Woolhouse et al., 2005 ; Parrish et al., 2008), iv) la présence de récepteurs cellulaires « ubiquitaires » qui favoriserait un large tropisme tissulaire ou un changement d'hôte (Pepin et al., 2010 ; Woolhouse et al., 2012), et enfin, v) les relations co-évolutives entre un hôte et son microbiote (Poulin & Morand, 2004) sont des déterminants importants de l'émergence d'un agent zoonotique (Cui et al., 2007; Olival et al., 2014), un parasite exprimant un fort pattern de co-évolution avec son hôte ne fera que très peu de « host-switch », alors qu'un parasite co-évoluant faiblement avec son hôte sera plus à même de sauter la barrière d'espèce. Ces relations passent essentiellement par deux processus évolutifs à savoir, la macroévolution et la micro-évolution.

#### 1.2.3.1. Macro-évolution

Les processus macro-évolutifs sont des mécanismes se déroulant généralement sur une échelle de temps longue et vont être en partie à l'origine de la diversité génétique d'un parasite. Ils regroupent deux mécanismes principaux: la "co-spéciation" et le saut d'espèce ou "host-switch" (Page, 2003). La co-spéciation correspond à une diversification de l'hôte, entraînant à sa suite une co-divergence du parasite qui lui est fortement adapté (**Figure 6**). A l'inverse, le saut d'un parasite correspond à un passage depuis l'hôte ancestral vers un nouvel hôte, passage suivi d'une spécialisation générant une nouvelle combinaison hôte-microorganisme (Agosta et al., 2010; de Vienne et al., 2013). Certaines études qui plaidaient en faveur d'un rôle prépondérant de la co-spéciation dans les processus macro-évolutifs doivent être interpréter avec précaution du fait d'une probable surestimation due à l'utilisation d'outils inadaptés en phylogénie (de Vienne et al., 2013). C'est le cas du modèle hantavirus

souvent cité comme prototypique de l'association des virus à leurs hôtes respectifs présentant une parfaite congruence entre les arbres phylogénétiques des hôtes et des différents lignages viraux (Hughes & Friedman, 2000; Jackson & Charleston 2004; Ramsden et al., 2009). Cependant, il a par la suite été démontré que le moteur réel de leur évolution dépendait plus des multiples évènements de transmission par "host-switch", que d'un mécanisme de cospéciation des hôtes (Ramsden et al., 2009; Lin et al., 2012; Guo et al., 2013). D'autres événements évolutifs, tel que la duplication, correspondant à la spéciation indépendante du parasite dans son hôte ainsi que le « sorting » correspondant le plus généralement à (i) l'absence de colonisation d'un parasite dans les différentes lignées de l'hôte (« Missing the boat ») ou à (ii) l'extinction de la lignée parasitaire après la co-spéciation (« extinction ») (iii) ou à l'absence de spéciation du parasite lors de la spéciation de l'hôte (« failure to speciate ») (Figure 6) (Page, 2003).


#### 1.2.3.2. Micro-évolution

Les processus micro-évolutifs agissent sur des échelles de temps beaucoup plus courtes, et regroupent les interactions hôte-agent infectieux que l'on décrit généralement comme celles qui permettent, suite à un « host-switch », la meilleure adaptation de l'agent infectieux à son hôte. Il s'agit d'une "coévolution" sensu stricto (de Vienne et al., 2013), c'est-à-dire issue de la pression de sélection réciproque entre l'hôte et l'agent infectieux: échappement d'un agent infectieux, fixation des mutations, recombinaisons et réassortiments, etc. (Domingo & Holland, 1997). Cette forte dynamique évolutive entraîne l'apparition d'une grande diversité de variants génétiquement proches et qui vont constituer une population virale particulière appelée « quasi-espèce » (Domingo et al., 1999).

Les processus qui mettent en jeu une micro-évolution ne sont nullement exclusifs de ceux qui dirigent la macro-évolution, et peuvent être sujets à la coopération. Par exemple, après transmission d'un parasite à un nouvel hôte par *host-switch*, il est nécessaire pour ne pas connaître l'extinction, que le pathogène s'adapte au mieux à son nouvel environnement, son nouvel hôte via ces mécanismes évolutifs.

Il est donc intéressant et important d'investiguer ainsi à la fois les facteurs écologiques et les processus macro/micro évolutifs qui influeraient sur la diversité et la transmission d'agents zoonotiques inter et intra populationnelle. Au cours de nos travaux nous nous sommes

intéressés aux animaux réservoirs dont certains hébergent des agents pathogènes présentant des caractéristiques décrites ci-dessus.



**Figure 6.** Mécanismes évolutifs hôte-parasites (issue de Michel Widmann, ENS Lyon, adapté de Paterson, 1997 et Taraschewski, 2006).

# Chapitre 2. Micromammifères sauvages un réservoir d'agents zoonotiques

# 2.1. Généralités

La faune sauvage joue un rôle clé dans l'émergence des agents pathogènes dans les populations d'animaux domestiques et les populations humaines en servant de «réservoir enzootique" source d'agents infectieux pour la plupart antérieurement inconnus qui peuvent s'adapter, se répliquer et émerger dans de nouveaux hôtes (Morse et al., 1993). A cet égard l'exploration des agents infectieux dans la faune sauvage revêt un grand intérêt afin d'anticiper leur émergence potentielle et le déclenchement des maladies infectieuses nouvelles. Durant mes travaux de thèse, je me suis intéressé à deux réservoirs majeurs d'agents zoonotiques particuliers, les chauves-souris et les petits mammifères terrestres (Woolhouse et al., 2012).

# 2.2. Les chauves-souris

# 2.2.1. Caractéristiques

Les chauves-souris (Order Chiroptera) représentent un groupe très diversifié de mammifères que l'on retrouve pratiquement dans l'ensemble du globe à l'exception des régions polaires et quelques îles océaniques. Le terme *chiroptère* est dérivé du grec *kheir* (ἣ χείρ) et désigne la main et le terme *ptéron* (τὸ πτερόν) désignant l'aile. Elles représentent l'un des plus grands ordres de mammifères au monde avec plus de 1200 espèces (UICN pour Union Internationale pour la Conservation de la Nature, 2010), soit 20% des mammifères terrestres, et occupent le second rang après les rongeurs en termes de nombre de genres et d'espèces (Koopman, 1993). Elles sont classées en deux ordres : les Yangochiroptera (anciennement appelées Microchiroptéres) et Yinpterochiroptera (anciennement appelées Megachiroptéres) (Koopman, 1993). L'ordre des Yangochiroptera est constitué de deux sous-ordres et contient le plus grand nombre d'espèces, près de 1000 (UICN, 2010), et a une plus grande diversité écologique. La plupart des familles de cet ordre sont insectivores, bien que certaines familles aient un comportement polyphage.

L'ordre Yinpterochiroptera comprend plus de 200 espèces (UICN, 2010) dont la taille est en moyenne bien supérieure à celle des espèces de l'ordre Yangochiroptera (Mickleburgh & Hutson, 1992). Elles se nourrissent principalement de fruits et de nectar. Les chauves-souris frugivores sont surtout retrouvées dans les régions sub-tropicales et tropicales de l'Ancien Monde; en Méditerranée orientale et la péninsule arabique, en Afrique et en Asie, en Australie et les îles du Pacifique et de l'Océan Indien (Rainey & Pierson, 1992). Bien que l'analyse des fossiles permettant de retracer l'évolution des chauves-souris reste incomplète (Jones et al., 2002), une datation a permis d'estimer l'origine des chiroptères à la période éocène, il y a environ 52 à 50 millions d'années coïncidant avec l'augmentation significative de la température mondiale (Teeling et al., 2005). Généralement longévives, elles sont dotées de grandes capacités de dispersion sur de grandes distances. Elles forment des colonies densément peuplées (**Figure 7**) et présentent des comportements sociaux qui structurent leurs populations. Elles sont capables de rentrer en hibernation et possèdent un système

immunitaire qui leur confère une tolérance remarquable vis-à-vis d'agents infectieux (Calisher et al., 2006 ; Turmelle & Olival, 2009).

Le rôle des chauves-souris dans la transmission de pathogènes et l'émergence des maladies infectieuses d'origine virale est maintenant bien établi dans de nombreuses régions du monde (Sulkin & Allen, 1974; Wong et al., 2007; Wang, 2009). Des études récentes ont par ailleurs montré que les chauves-souris sont des hôtes depuis des temps très anciens de lyssavirus (Badrane & Tordo, 2001; Streicker et al., 2010; Streicker et al 2012; Banyard et al., 2014) et les henipavirus (Calisher et al., 2006; Drexler et al., 2012), suggérant une longue cospéciation entre ces virus et les chauves-souris.



**Figure 7.** Colonie de *Mormopterus francoismoutoui* (insectivores) dans la grotte de Trois-Bassins à La Réunion (Crédits photo : Erwan Lagadec).


# 2.2.2. Réservoirs d'agents pathogènes

Les chauves-souris sont reconnues comme les réservoirs les plus importants d'agents zoonotiques (Calisher et al., 2006) sans doute les plus anciens (Vidgen et al., 2015). Calisher et al. (2006) ont fait état de plus de 66 virus isolés ou détectés à partir de tissus de 74 espèces de chauves-souris (**Tableau 1**). Certains de ces virus sont connus pour causer des maladies humaines et animales. Les chauves-souris possèdent certaines caractéristiques qui peuvent maximiser leur efficacité en tant que réservoir et pourvoyeur de virus (Calisher et al., 2006). Certaines espèces de Yinpterochiroptera souvent perchées dans les arbres et fortement agrégés, forment des colonies considérables (**Figure 8**). L'occupation de ce genre de site peut être saisonnière (par exemple, pendant les périodes d'hibernation ou de maternité) ou pérenne, tout au long de l'année (Kunz & Pierson, 1994). Selon les espèces, la saison, et l'emplacement du site, la taille des colonies va de quelques spécimens à de milliers d'individus. C'est ainsi que les chauves-souris sont connues pour former les plus grands rassemblements comparés à tous les autres mammifères.

Cette propriété est particulièrement remarquable chez les colonies d'insectivores tels les Molossidae et les Vespertilionidae qui ont les plus grands comportements grégaires (McCracken & Gustin, 1991) (**Figure 7**). Plus d'un tiers des espèces de Yinpterochiroptera forme des colonies (Marshall, 1983; Pierson & Rainey, 1992). Des études ont montré que le contact étroit entre chauves-souris ayant ce mode de vie serait favorable la transmission d'agents infectieux (Luis et al., 2015).

Tableau 1. Exemple de virus isolés de chauves-souris (Calisher et al., 2006).

Virus	Bat species (common name) <sup>a</sup>
Family Rhabdoviridae, genus Lyssavirus	
Rabies virus	Numerous bat species, essentially worldwide
-	Eidolon helvum (African straw-colored fruit bat), Micropteropus pusillus (Peters' lesser epauletted fruit bat), Epomops dobsonii (Dobson's epauletted fruit bat), Nycteris gambiensis (Gambian slit-faced bat), Epomophorus wahlbergi (Wahlberg's epauletted fruit bat)
Duvenhage virus	Miniopterus sp., Nyctalus noctula (noctule), Vespertilio murinus (particolored bat), Nycteris thebaica (Egyptian slit-faced bat)
•	Megachiroptera (multiple <i>Pteropus</i> spp.), Microchiroptera sp. from Australia, <i>Saccolaimus</i> flaviventris (yellow-bellied pouched bat)
European bat lyssavirus 1	Eptesicus serotinus (common serotine), Rousettus aegyptiacus (Egyptian rousette)
. ,	Myotis myotis (mouse-eared myotis), Myotis dasycneme (pond myotis), Myotis nattereri (Nattereri's myotis), Miniopterus schreibersii (Schreibersi' long-fingered bat), Rhinolophus ferrumequinum (greater horseshoe bat), Myotis daubentonii (Daubentoni's myotis)
	Myotis blythii (lesser mouse-eared myotis)
Khujand virus	Myotis mystacinus (whiskered myotis)
	Murina leucogaster (greater tube-nosed bat)
West Caucasian bat virus	Miniopterus schreibersii (Schreibers' long-fingered bat)
Family Rhabdoviridae, genus unassigned	
Gossas virus	
Kern Canyon virus	
	Rhinolophus eloquens (eloquent horseshoe bat)
Oita 296 virus	Rhinolophus cornutus (little Japanese horseshoe bat)
Family <i>Orthomyxoviridae</i> , genus <i>Influenzavirus A</i> , influenza A virus	Nyctalus noctula (noctule)
Family Paramyxoviridae, genus Rubulavirus Mapuera virus	Stumira lilium (yellow epauletted bat)
	Pteropus poliocephalus (gray-headed flying fox) Pteropus hypomelanus (variable flying fox)
Family <i>Paramyxoviridae</i> , genus undetermined, a parainfluenzavirus	Rousettus leschenaultia (Leschenault's rousette)
Family Coronaviridae, SARS coronavirus	Rhinolophus sinicus (Chinese horseshoe bat), Rhinolophus pearsonii (Pearson's horseshoe bat), Rhinolophus macrotis (big-eared horseshoe bat), Rhinolophus ferrumequinum (greater horseshoe bat)



**Figure 8.** Colonie de *Pteropus scapularis* (frugivores) perchés sur un arbre (Crédits photo : Harrison Chris).

La durée de vie de nombreuses espèces de petites chauves-souris de l'ordre des Yangochiroptera, particulièrement celles vivant dans des régions tempérées, dépasse 25 ans, avec la plus grande longévité estimée à 35 ans, pour les frugivores. Cette longévité importante par rapport à leur taille relativement petite, est un trait de vie considéré comme favorisant l'infection et associée à des infections persistantes pourrait expliquer en partie le rôle de transmetteur efficace d'agents infectieux (Calisher et al., 2006).

Le régime alimentaire des chauves-souris est très diversifié allant du régime généraliste à très spécialisé. Les insectivores se nourrissent d'un large éventail d'arthropodes, notamment d'insectes mais aussi de mites, d'araignées et de petits crustacés (Kunz & Pierson, 1994). Certaines espèces sont frugivores se nourrissant de fruits et de nectars (von Helversen, 1993) alors que d'autres sont carnivores. Une dernière catégorie de chauves-souris est sanguivore, se nourrissant principalement de sang de mammifères et d'oiseaux (Kunz & Pierson, 1994). Les chauves-souris sanguivores revêtent une grande importance d'un point de vue santé humaine et vétérinaire dans les pays du Nouveaux Monde où la transmission d'agents zoonotiques se produit lors des morsures, notamment pour le virus de la rage (Constantine, 1988).

Enfin, le comportement migratoire de certaines espèces de chauves-souris a aussi été incriminé dans la transmission d'agents zoonotiques. Certaines espèces de chauves-souris sont capables de voler sur de longues distances particulièrement lors des migrations saisonnières (Griffin, 1970). Les espèces de Yangochiroptera voyagent souvent entre 10 à 15 km mais peuvent parcourir de plus longues distances dans leur quête de nourriture (Kunz & Pierson, 1994). Certaines espèces de Yinpterochiroptera sont également connues pour parcourir des distances au-delà de 80 km pour se nourrir (Epstein et al., 2009). Les Yangochiroptera dans les régions tempérées du nord, migrent en période hivernale vers le sud pour bénéficier d'un climat moins extrême (Strelkov et al., 1969).

# 2.3. Les petits mammifères terrestres

# 2.3.1. Caractéristiques

Les rongeurs (ordre : Rodentia) représentent l'ordre le plus abondant et le plus diversifié des mammifères et comporte environ 43% du nombre total d'espèces de mammifères vivant dans le monde (Huchon et al 2002; Wilson & Reeder 2005). Le genre Rattus (sous famille Murinae, famille des Muridae, super famille Muroidae) est le plus important avec plus de 60 espèces inventoriées (Aplin et al., 2003). Rattus norvegicus (rats bruns) et Rattus rattus (rats noirs) sont parmi les deux espèces les plus répandues et sont communément appelées rongeurs de l'Ancien Monde (Wilson & Reeder, 2005). Rattus norvegicus est connu pour être originaire du nord de la Chine (Nowak & Walker, 1991) alors que R. rattus proviendrait de l'Inde et d'Asie du Sud (Bonnefoy et al., 2008; Nagorsen, 2005). Les premiers membres sont apparus il y a environ trois millions d'années, dans le Pliocène tardif et se sont dispersés rapidement à travers l'Asie du Sud (Aplin et al., 2003). Ils ont depuis réussi à coloniser tous types d'habitats et se retrouvent sur tous les continents sauf en Antarctique (Lund, 1994). Leur expansion mondiale a été favorisée par les moyens de transport en particulier le trafic maritime (Bonnefoy et al., 2008; Nagorsen, 2005). Rattus norvegicus est plus fortement associés aux écosystèmes urbains (Yahner, 2001) tandis que R. rattus sera communément retrouvé dans les zones rurales. Ces deux espèces de rongeurs sont connues pour être des ravageurs de cultures causant des destructions importantes de récoltes, notamment en Asie du Sud et du Sud-est (Aplin et al., 2003).

# 2.3.2 Réservoirs d'agents pathogènes et perte de la biodiversité

Les petits mammifères terrestres, notamment les rats tels que *Rattus norvegicus* et *R rattus* sont eux aussi identifiés comme source d'un certain nombre d'agents pathogènes responsables de morbidité et mortalité importantes dans les zones urbaines de différentes régions du monde. En terme de santé publique, *R rattus* et *R norvegicus* sont les deux rongeurs les plus incriminés dans la dissémination d'agents pathogènes responsables de bactérioses (la leptospirose due à *Leptospira*, la peste causée par *Yersinia* et des Rickettsioses comme le typhus murin dues à *Rickettsia*) et de viroses (syndrome pulmonaire à hantavirus) responsables de morbidité et mortalité humaines importantes à travers le monde (Meerburg et al., 2009).

Meerburg et al. (Meerburg et al., 2009) ont fait état de plusieurs agents pathogènes isolés de rongeurs (**Tableau 2**). Parmi eux, les infections à hantavirus du genre *Hantavirus* (Famille : *Bunyaviridae*) sont les plus connues avec le syndrome pulmonaire à hantavirus. Les hantavirus sont aussi retrouvés chez d'autres petits mammifères terrestres, notamment ceux appartenant à la famille des Cricetidae Murnidae, Soricidae et Talpidae (Guo et al., 2014).

Tableau 2. Aperçu des différentes maladies et agents pathogènes retrouvés chez les rongeurs (Meerburg et al., 2009).

		Carrier/Reservoir	Population at-risk	Chance	Severity	
Disease	Agent				Human Health	Economy
Hantavirus Pulmonary Syndrome	Virus, Bunyaviridae	Carrier	2	1	3	1
Hemorrhagic Fever with renal syndrome (+ other hemorrhagic fevers)	Virus, Bunyaviridae	Carrier	2	2	2	2
Nephropathia epidemica	Virus, Bunyaviridae	Carrier	1	1	1	1
Crimean-Congo hemor- rhagic fever	Virus, Bunyaviridae	Reservoir	1	1	3	1
Borna disease	Virus, Bornaviridae	Reservoir	1	1	1	2
Omsk hemorrhagic fever	Virus, Flaviviridae	Reservoir	1	1	1	1
Kyasanur Forest Disease	Virus, Flaviviridae	Reservoir	1	1	1	1
Apoi Virus Disease	Virus, Flaviviridae	Unknown	Unknown	Unknown	Unknown	Unknown
Tick-borne encephalitis	Virus, Flaviviridae	Reservoir	2	1	3	1
Powassan encephalitis	Virus, Flaviviridae	Reservoir	1	1	1	1
Lymphocytic	Virus, Arenaviridae	Reservoir	1	1	1	1

Reservoir: rodents harbor disease-causing organisms and thus serve as potential sources of disease outbreaks, but always via a vector (tick, sand-fly etc.)

Carrier: rodent that shows no or limited symptoms of a disease but harbors the disease-causing agent and is capable of passing it directly onto humans

Population at-risk: focal = 1, regional = 2, more than 2 continents = 3

Chance: chance of contracting the disease (all pathways, not only via rodents): small chance = 1, moderate chance = 2, high chance = 3 Human health: Mortality without treatment <5%=1, 5 to 10%=2, >10%=3. No mortality = 0.

Economy: losses in terms of morbidity combined with other losses (e.g. in animal productivity): small losses = 1, moderate losses = 2, huge losses = 3.

La caractéristique principale des rongeurs, aggravante en terme de risque infectieux est le contact étroit qu'ils établissent avec les populations humaines. La transmission de pathogènes peut alors se produire soit de façon directe, par morsure la plupart du temps, soit de façon indirecte, par contamination des produits alimentaires et du milieu par les fèces ou les urines, par formation d'aérosols contagieux ou médiée par des vecteurs arthropodes ectoparasites (tiques, acariens, puces) (Meerburg et al., 2009). La grande taille des populations, résultant de leur taux de reproduction extrêmement élevé, augmente le risque de contact et de transmission d'agents pathogènes vers les populations humaines. Ces deux espèces de rongeurs sont connues pour être des ravageurs de cultures causant des destructions importantes de récoltes, notamment en Asie du Sud et du Sud-est (Aplin et al., 2003). Rattus rattus et R. norvegicus sont à l'origine d'invasion biologique pouvant avoir des effets dévastateurs sur les écosystèmes naturels en déplaçant et réduisant les populations indigènes, comme les oiseaux, les reptiles ou encore les végétaux (Nowak & Walker, 1991). Cette invasion par des populations de rongeurs peut entraîner l'extinction d'autres espèces animales en particulier endémiques (Clavero et al., 2005) avec une réduction de la biodiversité surtout en territoire insulaire. L'appauvrissement spécifique peut favoriser la transmission épidémique des agents infectieux quand les espèces tampons viennent à disparaître au profit d'une espèce invasive vectrice (Keesing et al., 2006).

# Chapitre 3. Caractéristiques de la région du Sudouest de l'Océan Indien (SOOI)

# 3.1. Généralités

Les îles tropicales pour lesquelles nous avons axé notre étude sont Madagascar, et les archipels des Comores (Mayotte et Anjouan), des Seychelles (Mahé) et des Mascareignes (La Réunion et Maurice) (**Figure 9**). Cet ensemble insulaire est remarquable par la diversité des îles qui le composent: diversité de leur origine géologique, de leur âge, des origines ethniques des populations humaines qui le peuplent, disparités de leurs développements économiques, et des voies d'échanges internationales et commerciales. Parmi les îles du SOOI, Madagascar a plusieurs caractéristiques écologiques exceptionnelles: Six types de végétations, quatre régions bioclimatiques et une échelle altitudinale allant de 0 à 2875 m (Goodman & Raherilalao, 2013).

Typiquement, la faune vertébrée des îles du SOOI proviendrait de l'Afrique ou encore de l'Australasie (Peake et al., 1971). Ces îles font partie des 34 régions de notre planète considérées par l'UICN comme « hotspots » de la biodiversité (Myers et al., 2000).

Nous avons porté notre intérêt sur les chauves-souris et les petits mammifères terrestres de ces îles. La plus grande richesse et le plus grand taux d'endémisme pour ces deux groupes d'animaux sont observés sur la grande île de Madagascar. On y dénombre aujourd'hui le plus d'espèces de petits mammifères non volants, un total de 45 espèces de chauves-souris (3 frugivores et 42 insectivores), dont 36 sont endémiques à l'île et plus de 64 espèces de petits mammifères terrestres (Goodman, 2011 ; Soarimalala & Goodman, 2013).




Figure 9. Les îles du Sud-ouest de l'Océan Indien (adapté de Tortosa et al., 2012).

# 3.2. Les chauves-souris

Dans un récent atlas, Goodman & Ramasindrazana, (2013) ont décrit les familles et espèces de chauves-souris connues à ce jour pour être retrouvées à Madagascar. Par ailleurs, ils ont inscrit différentes caractéristiques écologiques intéressantes et importantes. La plupart des sites connus pour héberger des chauves-souris sont constitués de communautés d'espèces parfois de plusieurs familles différentes vivant en sympatrie, particulièrement chez les chauves-souris de la famille de Miniopteridae qui, avec les Vespertilionidae, représente les deux plus grandes familles de chauves-souris de Madagascar. Les chauves-souris de ces îles vivent majoritairement dans les grottes (cavernicoles) ou dans les forêts ou feuillage (forestière), mais certaines espèces sont rencontrées proche des habitations humaines (synantropique) notamment chez les chauves-souris de la famille des Molossidae. Plusieurs espèces peuvent être retrouvées sur une étendue géographique importante allant du Sud au Nord de l'île (Goodman, 2011). Ces modes de vie contrastés sont intéressants, d'une part, pour analyser la transmission inter-espèces de pathogènes, notamment viraux, et d'autre part, déterminer le risque de transmission infectieuse des chauves-souris vers les populations humaines. Le tableau 3 décrit les caractéristiques connues des chauves-souris retrouvées à Madagascar qui ont fait l'objet de notre étude (Goodman, 2011). Certaines espèces ont été récemment décrites (Goodman et al., 2012a; Goodman et al., 2012b; Goodman & Ramasindrazana, 2015; Goodman et al. 2015). Par ailleurs, la revue d'O'Brien et al. (2011) décrit les différentes espèces de chauves-souris présentes sur les îles de la zone SOOI. Certaines espèces comme Taphozous mauritianus sont retrouvées en commun sur de nombreuses îles.

Le genre *Mormopterus*, dont les espèces sœurs *Mormopterus jugularis*, *Mormopterus francoismoutoui* et *Mormopterus acetabulosus*, est retrouvé sur Madagascar, La Réunion et Maurice, respectivement (Goodman et al., 2008).

Le **tableau 4** décrit quant à lui les différentes chauves-souris des autres îles de la zone du SOOI étudiées (O'Brien, 2011).

Tableau 3. Caractéristiques et particularités des chauves-souris collectées à Madagascar.

Famille	Espèce	Endémicité	Particularité
Emballonuridae	Coleura kibomalandy	Oui	
	Paremballonura tiavato	Oui	
Hipposideridae	Hipposideros commersoni	Oui	
Miniopteridae	Miniopterus aelleni	Non	Sympatrie
	Miniopterus cf. ambohitrensis	Oui	Sympatrie
	Miniopterus gleni	Oui	Sympatrie
	Miniopterus griffithsi	Oui	Sympatrie
	Miniopterus griveaudi	Non	Sympatrie
	Miniopterus mahafaliensis	Oui	Sympatrie
	Miniopterus majori	Oui	Sympatrie
	Miniopterus sororculus	Oui	Sympatrie
Molossidae	Chaerephon atsinanana	Oui	Sympatrie et Synanthropique
	Chaerephon leucogaster	Non	Sympatrie et Synanthropique
	Mops leucostigma	Non	Sympatrie et Synanthropique
	Mops midas	Non	Sympatrie et Synanthropique
	Mormopterus jugularis	Oui	Sympatrie et Synanthropique
	Otomops madagascariensis	Oui	Sympatrie et Synanthropique
Pteropodidae	Eidolon dupreanum	Oui	
1	Pteropus rufus	Oui	
	Rousettus madagascariensis	Oui	
Rhinonycteridae	Paratriaenops furculus	Oui	
J	Triaenops menamena	Oui	
Vespertilionidae	Hypsugo bemainty	Oui	
F	Myotis goudoti	Oui	Sympatrie
	Neoromicia malagasyensis	Oui	r
	Neoromicia matroka	Oui	
	Neoromicia robertsi	Oui	
	Pipistrellus cf. hesperidus	Non	
	Pipistrellus hesperidus	Non	Sympatrie
	Pipistrellus raceyi	Oui	Sympatrie
	Scotophilus marovaza	Oui	Synanthropique

**Tableau 4.** Caractéristiques et particularités des chauves-souris collectées dans les autres îles de la zone SOOI (adapté d'O'Brien, 2011).

Localité	Famille	Espèce Endémicité		Particularité
La Réunion	Molossidae	Mormopterus francoismoutoui	Oui	Synanthropique
Maurice	Molossidae	Mormopterus acetabulosus	Oui	
	Pteropodidae	Pteropus niger	Oui	Synanthropique
Mahé	Pteropodidae	Pteropus seychellensis	Oui	Synanthropique
Anjouan	Miniopteridae	Miniopterus griveaudi	Non	
	Molossidae	Chaerephon pusillus	Non	Synanthropique
Mayotte	Pteropodidae	Pteropus seychellensis	Oui	Synanthropique

# 3.3. Les petits mammifères terrestres

Chaque île du SOOI est par ailleurs connue pour abriter des rongeurs introduits, notamment, *Rattus rattus* et *Rattus norvegicus*. Cependant, Madagascar est l'une des seules îles du SOOI à héberger de nombreuses espèces endémiques. A ce jour 64 espèces de petits mammifères terrestres y ont été décrites dont 59 endémiques (Soarimalala & Goodman, 2013). La sous famille des Oryzorictinae représente la plus grande sous famille dont le genre *Microgale* est le plus diversifié et contient plus de 23 espèces. La plus grande partie des espèces de cette sousfamille est capable de vivre en sympatrie. Le **Tableau 5** décrit les principaux petits mammifères terrestres connus à Madagascar qui ont fait l'objet de notre étude.

Les petits mammifères terrestres indigènes jouent un rôle très important dans l'écosystème. Ils sont prédateurs d'insectes nuisibles ou ravageurs, disséminateurs de graines, ou servent de proies aux rapaces, carnivores et reptiles. A l'inverse, parmi les espèces introduites, les rats (*Rattus rattus*) sont source de nombreux problèmes. Ils sont responsables de la destruction des cultures agricoles et sont vecteurs de nombreuses maladies. A titre d'exemple *Rattus rattus*, *Mus musculus* et *Rattus norvegicus*, ainsi que 2 autres espèces introduites, *Suncus murinus* et *Suncus etruscus*, sont réputés vecteurs de la peste.

**Tableau 5.** Caractéristiques et particularités des petits mammifères terrestres collectées à Madagascar (adapté de Soarimalala & Goodman, 2013).

Famille	Sous-famille	Espèce	Endémicité	Particularité
Tenrecidae	Tenrecinae	Hemicentetes semispinosus	Oui	Sympatrie
		Setifer setosus	Oui	Sympatrie
		Tenrec ecaudatus	Oui	
	Oryzorictinae	Microgale cowani	Oui	Sympatrie
		Microgale dobsoni	Oui	
		Microgale fotsifotsy	Oui	
		Microgale gymnorhyncha	Oui	Sympatrie
		Microgale jobihely	Oui	
		Microgale longicaudata	Oui	Sympatrie
		Microgale majori	Oui	
		Microgale parvula	Oui	
		Microgale principula	Oui	
		Oryzorictes hova	Oui	
Soricidae	Soricinae	Suncus murinus	Non	Synanthropique
Muridae	Murinae	Rattus norvegicus	Non	Synanthropique
		Rattus rattus	Non	Synanthropique
		Mus musculus	Non	
Nesomyidae	Nesomyinae	Eliurus majori	Oui	
-	-	Eliurus minor	Oui	Sympatrie
		Gymnuromys roberti	Oui	
		Nesomys rufus	Oui	Sympatrie

Malgré la diversité très élevée de chauves-souris et de petits mammifères terrestres dans le SOOI, la faune sauvage de cette région a été très peu explorée jusqu'à peu de temps en terme de réservoirs potentiels d'agents zoonotiques. Dans une étude préliminaire réalisée en 2012 par le CRVOI, sur les chauves-souris de Madagascar, des Comores et de La Réunion, une dizaine de nouveaux paramyxovirus a été détectée chez 5 espèces de chauves-souris insectivores. Ces paramyxovirus formaient un phylogroupe à part entière, proche du genre Morbillivirus sans qu'il soit possible de les rattacher complètement sur le plan phylogénétique (Wilkinson et al., 2012). Les taux d'infection élevés par ces nouveaux paramyxovirus Morbilli-related, leurs diversité et variabilité génétique (15 à 30% de variabilité génétique), ainsi que les spectres d'hôtes qui leur ont été associé marqué par un fort endémisme, en comparaison à d'autres études similaires dans le monde, nous ont alors conduit à émettre l'hypothèse des singularités écologiques régionales, agissant au niveau des échanges viraux entre hôtes. Dans ce contexte, l'objectif principal de mes travaux de thèse a été de déterminer quels sont les déterminants génétiques et écologiques qui peuvent rendre compte de tels taux d'infection par les paramyxovirus, d'une telle variabilité génétique et d'un tel spectre d'hôte dans la région du SOOI.

# PARTIE II. TRAVAUX DE RECHERCHE

Dans cette partie, j'introduirais le contexte et les objectifs des programmes de recherche conduits au cours des 3 années (Janvier 2013-Décembre 2015).

Les résultats seront ensuite présentés en 3 chapitres (**chapitres 4 à 6**). Une partie de ces résultats est déjà publiée sous forme d'un article dans une revue internationale à comité de lecture, l'autre partie est structurée en 3 manuscrits soumises ou en cours de publication.

Une discussion générale des résultats obtenus, de leur signification ainsi que des perspectives envisageables par ces travaux clôturera ce manuscrit.

# Présentation du sujet, des objectifs et questions de recherche

#### Contexte de l'étude

Comme abordé dans le **chapitre 3** de l'état de l'art, la région du SOOI est singulière par l'existence d'une multi-insularité fortement contrastée, à la fois au niveau géographique que géologique et de la faune sauvage qui y vit. Celle dernière est particulière dans la plupart des petites îles par une richesse spécifique relativement limitée par rapport à Madagascar ou à certaines zones continentales mais elle possède un niveau d'endémisme extrêmement marqué pour certains groupes d'animaux, notamment les chauves-souris et les petits mammifères terrestres (Goodman et al 2003 ; Yoder et al., 2005 ; Goodman et al., 2007; Goodman et al., 2008 ; Goodman, 2011). Nous avons fait l'hypothèse que cette biodiversité ne devrait pas être limitée aux seuls règnes animal et végétal; mais que le monde microbien, qui leur est inféodé, devrait l'être également. Ainsi, les spécificités écologiques des îles du SOOI associées à leur isolement géographique font de cette région un remarquable champ d'investigation sur (i) la dynamique évolutive des processus infectieux et des flux de pathogènes qui peuvent survenir au sein des communautés d'espèces, et (ii) des facteurs biotiques et abiotiques qui entrent en jeu dans les interactions hôte-pathogène, et des mécanismes macro-évolutifs mobilisés (Tortosa et al., 2012).

L'avènement des méthodes moléculaires a rendu possible des investigations menées aussi bien au niveau de l'écologie des hôtes que des agent infectieux, pour identifier des animaux réservoirs et inventorier les pathogènes potentiels qui leurs sont associés. Etudier la dynamique insulaire de transmission virale au sein d'une communauté d'hôtes pouvant être assimilées à des réservoirs et comprendre les forces évolutives en action au niveau macroévolutif (Moss et al., 2012; Saluzzo et al., 2014) sont considères comme des étapes indispensables pour contextualiser "in natura" un évènement d'émergence s'il devait se produire un jour chez des populations humaines naïves.

#### Objectifs et questions de recherche

Deux modèles viraux ont été étudiés pour leur aptitude à transgresser les barrières d'espèces, un trait directement lié au risque d'émergence.

Les paramyxovirus (*Paramyxoviridae*) et les lyssavirus (*Rhabdoviridae*). Elles représentent deux familles virales très diversifiées et structurellement très proches (Assenberg et al., 2010; Drexler et al., 2012; Banyard et al., 2014). Plus spécifiquement les questions suivantes ont été posées:

# i). Quelle est la dynamique de transmission virale inter-espèces chez les petits mammifères terrestres et les chauves-souris de la zone du SOOI ?

Cette étude a principalement porté sur les petits mammifères terrestres (ordre: Afrosorocidae) et volants (ordre: Chiroptera) endémiques de Madagascar, puis elle a été étendue aux rongeurs (ordre: Rodentia) de Mayotte, La Réunion et Madagascar. Des analyses phylogénétiques ont permis, d'une part, de mieux établir l'appartenance phylogénétique des paramyxovirus détectés aux *Unclassified Morbilli-Related viruses* (*UMRVs*), et d'autre part, de déterminer la façon dont pouvait opérer les flux de transmission au niveau intra- et inter-ordres, définissant ainsi les contours des différents spectres d'hôtes associés à ces *UMRVs* (**Chapitre 4**).

# ii). Quel est le mécanisme macro-évolutif qui prédomine chez les *UMRVs* de chauvessouris et les facteurs environnementaux associés à la dynamique virale?

L'étude de consortia de chauves-souris réparties sur l'ensemble de l'île de Madagascar, représentant un ensemble de 35 espèces différentes et de leurs *UMRVs* inféodés, a permis de dégager, à l'aide d'analyses phylogénétiques (co-phylogénies) et statistiques (modèles linéaires généralisés), le mécanisme évolutif qui sous-tend la transmission interspécifique au sein des communautés de chauves-souris, et d'identifier certains facteurs biotiques et abiotiques qui influencent la relation entre les hôtes et les pathogènes (**Chapitre 5**).

#### iii). Quelle est la situation épidémiologique des lyssavirus de la région SOOI?

Pour investiguer la présence et caractériser la circulation des lyssavirus des chiroptères du SOOI, des chauves-souris de différentes espèces ont été collectées dans l'archipel des Comores, des Mascareignes, et des Seychelles, ainsi qu'à Madagascar. Les échantillons prélevés ont été analysés par la technique de référence RFFIT, et par biologie moléculaire pour la mise en évidence, respectivement, d'anticorps neutralisants anti-lyssavirus et de génomes viraux (**Chapitre 6**).

Chapitre 4. Étude de la dynamique d'infection des paramyxovirus chez les petits mammifères et les chauves-souris de la zone Sud-Ouest Océan Indien

#### 4.1. Généralités

Les paramyxovirus appartiennent à une vaste famille qui ne cesse de grandir et de se diversifier. La famille *Paramyxoviridae* est divisée en deux sous familles: les *Paramyxovirinae* et les *Pneumovirinae* (**Annexe 1**).

La sous famille des *Paramyxovirinae* comprend 31 espèces classées en sept genres (ICTV, 2014): les *Respirovirus*, les *Rubulavirus*, les *Avulavirus*, les *Morbillivirus*, les *Henipavirus* et ceux nouvellement classés, les *Ferlavirus* et les *Aquaparamyxovirus*. Elle comprend entre autre, les virus de la rougeole (*measles*), des oreillons, (*mumps*) de la maladie de Newcastle (*Newcastle disease virus*), les parainfluenza, les virus *Hendra* et *Nipah*. Certains membres des *Paramyxovirinae* récemment émergents (par exemple *Menangle*, *Tioman*, *Beilong*, et *J-virus*) ne sont pas formellement phylogénétiquement reliés à ces 7 genres principaux et sont en phase de reclassification.

La seconde sous famille *Pneumovirinae* se décompose en deux genres, les *Pneumovirus* et les *Métapneumovirus*, qui comprennent cinq espèces. Elle comprend également des anciens et des nouveaux pathogènes pour l'homme et l'animal, tels que les virus respiratoires syncytiaux humains et bovins, ainsi que les métapneumovirus humain et aviaire.

Les paramyxovirus sont mondialement distribués avec un large spectre d'hôte important (Drexler et al., 2012). Les *Morbillivirus* ont un très large spectre d'hôte et sont retrouvés chez l'homme, les dauphins, les chauves-souris, les rongeurs, les dromadaires et les bovins (Drexler et al., 2012), alors que les genres *Avulavirus*, *Aquaparamyxovirus* et *Ferlavirus* n'ont jamais été retrouvés chez l'homme.

Measles, reste à ce jour le *Morbillivirus* le plus dévastateur dans le monde causant des millions de morts, particulièrement dans les pays en développement (Moss & Griffin, 2012).

Les virus *Hendra* et *Nipah* du genre *Henipavirus* ont causé des épidémies mortelles en Australie et en Malaisie en 1994 et 1999 respectivement, provoquant d'innombrables pertes humaines et animales (Chua et al., 2000 ; Field et al., 2001 ; Mahalingam et al., 2012).

Au niveau structural, les paramyxovirus sont des virus à ARN monocaténaire de polarité négative, dont le génome mesure entre de 15 à 18 kb (**Annexe 2**). La nucléocapside est constituée de l'ARN associé à la nucléoprotéine (NP), la phosphoprotéine polymérase (P), et la protéine (L). La protéine L est l'ARN polymérase, la protéine P facilite la synthèse d'ARN

et la protéine NP contribue à maintenir la structure génomique. La nucléocapside associée avec la protéine de matrice (M) garnissant l'intérieur de l'enveloppe du virion. L'enveloppe contient deux glycoprotéines, une protéine de fusion (F), qui favorise la fusion des membranes virales et de l'hôte cellulaire, et une protéine de fixation virale (hémagglutinine-neuraminidase (HN), l'hémagglutinine (H), ou la protéine (G). La protéine F doit être activée par clivage protéolytique, ce qui génère deux glycopeptides F1 et F2 maintenus ensemble par un pont disulfure, et qui expriment une activité de fusion membranaire (Lamb & Parks, 2007).

#### 4.2. Approches expérimentales

Les résultats issus des investigations du CRVOI antérieures à mes travaux de thèse ont mis en évidence de nouveaux paramyxovirus et l'existence potentielle de mécanismes d'échanges viraux entre des hôtes endémiques particuliers à la région du SOOI (Wilkinson et al., 2012). Ces acquis ont été rendus possibles grâce à de nombreuses campagnes de collectes de micromammifères volants et terrestres vivant en isolement ou en communauté d'espèces. Ce sont au total plus de 140 chauves-souris (8 espèces), 333 petits mammifères terrestres endémiques de Madagascar (20 espèces) et 264 rongeurs introduits (4 espèces) qui ont été capturés entre 2010 et 2013, et criblés pour la mise en évidence de paramyxovirus. La détection de ces derniers a été effectuée par RT-PCR à l'aide d'un système pangénérique (Tong et al., 2008) capable d'amplifier la grande majorité des membres des genres *Respirovirus*, *Morbillivirus* et *Henipavirus*. Les analyses phylogénétiques réalisées (Mr Bayes, Beast, Mesquite) ont permis de retracer l'histoire évolutive de ces nouveaux virus et de quantifier les possibles évènements de saut d'espèces par taxon et par origine géographique.

Dans la première publication ci-après (Wilkinson et al., 2014), nous avons appliqué cette méthode pour caractériser en profondeur la dynamique d'infection et démontrer l'existence d'un flux viral intra-ordre particulièrement chez les Chiroptera, Afrosorocidae de Madagascar et Rodentia. Plus surprenant encore d'un point de vue écologique nous avons mis en évidence que ces flux de virus pouvaient être identifiés au niveau inter-ordre, notamment, entre Chiroptera ou Afrosorocidae et Rodentia collectés à Mayotte, La Réunion et Madagascar.

Annexe 5 : Par ailleurs, une étude phylogénétique et phylogéographique a été conduite afin de déterminer l'histoire évolutive des paramyxovirus des rongeurs de Tunisie. Un total de 784 petits mammifères composé de *Psammomys obesus*, *Meriones shawi* et *Ctenodactylus gundi* indigènes du Nord de l'Afrique et de l'espèce introduite *Rattus rattus* ont pu être analysés et comparés aux résultats issus en partie de mes travaux sur les animaux des régions du SOOI. Les résultats de ces travaux constituent la seconde publication ci-dessous et soumise (Ghawar et al., 2015 en soumission).

4.3 Etude de la dynamique d'infection au sein des micromammifères d	le la
zone SOOI. Résultats présentés dans la publication ci-après	

## Journal of Virology

### Highly Diverse Morbillivirus-Related Paramyxoviruses in Wild Fauna of the Southwestern Indian Ocean Islands: Evidence of Exchange between Introduced and Endemic Small Mammals

David A. Wilkinson, Julien Mélade, Muriel Dietrich, Beza Ramasindrazana, Voahangy Soarimalala, Erwan Lagadec, Gildas le Minter, Pablo Tortosa, Jean-Michel Heraud, Xavier de Lamballerie, Steven M. Goodman, Koussay Dellagi and Herve Pascalis

*J. Virol.* 2014, 88(15):8268. DOI: 10.1128/JVI.01211-14. Published Ahead of Print 14 May 2014.

	Updated information and services can be found at: http://jvi.asm.org/content/88/15/8268
SUPPLEMENTAL MATERIAL	These include: Supplemental material
REFERENCES	This article cites 49 articles, 12 of which can be accessed free at: http://jvi.asm.org/content/88/15/8268#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: http://journals.asm.org/site/misc/reprints.xhtml
To subscribe to to another ASM Journal go to: http://journals.asm.org/site/subscriptions/

Journals.ASM.org



### Highly Diverse Morbillivirus-Related Paramyxoviruses in Wild Fauna of the Southwestern Indian Ocean Islands: Evidence of Exchange between Introduced and Endemic Small Mammals

David A. Wilkinson,<sup>a,b</sup> Julien Mélade,<sup>a,b</sup> Muriel Dietrich,<sup>a,b</sup> Beza Ramasindrazana,<sup>a,d</sup> Voahangy Soarimalala,<sup>d</sup> Erwan Lagadec,<sup>a,c</sup> Gildas le Minter,<sup>a,c</sup> Pablo Tortosa,<sup>a,b</sup> Jean-Michel Heraud,<sup>g</sup> Xavier de Lamballerie,<sup>f</sup> Steven M. Goodman,<sup>d,e</sup> Koussay Dellagi,<sup>a,c</sup> Herve Pascalis<sup>a,c</sup>

Centre de Recherche et de Veille sur les Maladies Émergentes Dans l'Océan Indien (CRVOI), Plateforme de Recherche CYROI, La Réunion, France<sup>a</sup>; Université de La Réunion, La Réunion, France<sup>b</sup>; Institut de Recherche pour le Développement (IRD), IRD–BP 50172, La Réunion, France<sup>c</sup>; Association Vahatra, Antananarivo, Madagascar<sup>d</sup>; Field Museum of Natural History, Chicago, Illinois, USA<sup>c</sup>; UMR\_D 190, Aix Marseille University, IRD French Institute of Research for Development, EHESP French School of Public Health, Marseille, France<sup>f</sup>; Virology Unit, Institut Pasteur de Madagascar, Antananarivo, Madagascar<sup>a</sup>

### ARSTRACT

The *Paramyxoviridae* form an increasingly diverse viral family, infecting a wide variety of different hosts. In recent years, they have been linked to disease emergence in many different animal populations and in humans. Bats and rodents have been identified as major animal populations capable of harboring paramyxoviruses, and host shifting between these animals is likely to be an important driving factor in the underlying evolutionary processes that eventually lead to disease emergence. Here, we have studied paramyxovirus circulation within populations of endemic and introduced wild small mammals of the southwestern Indian Ocean region and belonging to four taxonomic orders: Rodentia, Afrosoricida, Soricomorpha, and Chiroptera. We report elevated infection levels as well as widespread paramyxovirus dispersal and frequent host exchange of a newly emerging genus of the *Paramyxoviridae*, currently referred to as the unclassified morbillivirus-related viruses (UMRVs). In contrast to other genera of the *Paramyxoviridae*, where bats have been shown to be a key host species, we show that rodents (and, in particular, *Rattus rattus*) are significant spreaders of UMRVs. We predict that the ecological particularities of the southwestern Indian Ocean, where small mammal species often live in densely packed, multispecies communities, in combination with the increasing invasion of *R. rattus* and perturbations of endemic animal communities by active anthropological development, will have a major influence on the dynamics of UMRV infection.

### **IMPORTANCE**

Identification of the infectious agents that circulate within wild animal reservoirs is essential for several reasons: (i) infectious disease outbreaks often originate from wild fauna; (ii) anthropological expansion increases the risk of contact between human and animal populations and, as a result, the risk of disease emergence; (iii) evaluation of pathogen reservoirs helps in elaborating preventive measures to limit the risk of disease emergence. Many paramyxoviruses for which bats and rodents serve as major reservoirs have demonstrated their potential to cause disease in humans and animals. In the context of the biodiversity hot spot of southwestern Indian Ocean islands and their rich endemic fauna, we show that highly diverse UMRVs exchange between various endemic animal species, and their dissemination likely is facilitated by the introduced *Rattus rattus*. Hence, many members of the *Paramyxoviridae* appear well adapted for the study of the viral phylodynamics that may be associated with disease emergence.

Ild rodents and bats together comprise more than 60% of all Wild rodents and vars together company when we mammalian species and have been found to carry many zoonotic viruses (1). Recently, these groups of mammals have been shown to host a broad spectrum of novel paramyxoviruses (2-8). These data have broadened our knowledge of the host spectrum of Paramyxoviridae, a diverse viral family which is currently divided into two main subfamilies: the Pneumovirinae and the Paramyxovirinae. Members of the Paramyxoviridae include the causative agents of the human diseases mumps, measles and other respiratory tract infections due to Parainfluenza viruses and metapneumoviruses, as well as a large number of viruses associated with disease in animals, such as Newcastle disease virus, canine distemper virus, rinderpest virus, and others. Additionally, paramyxoviruses have been linked to a number of recent emerging or reemerging disease epidemics with high mortality rates (9). Of these, henipaviruses (Paramyxovirinae) have emerged as human pathogens from fruit bat populations since 1994 in Bangladesh, Malaysia, and Australia (10, 11). Additionally, a novel rubulavirus-related virus, similar to those known to be hosted by fruit bats, has been described with the capacity to cause illness in humans (12), showing how as-yet unknown *Paramyxoviridae* in wild animal reservoirs may pose an important health risk.

Received 29 April 2014 Accepted 5 May 2014

Published ahead of print 14 May 2014

Editor: A. García-Sastre

Address correspondence to David A. Wilkinson, dwilkin799@gmail.com, or Herve Pascalis, herve.pascalis@ird.fr.

Supplemental material for this article may be found at http://dx.doi.org/10.1128 /JVI.01211-14.

Copyright © 2014, American Society for Microbiology. All Rights Reserved. doi:10.1128/JVI.01211-14

8268jvi.asm.orgJournal of Virologyp. 8268–8277August 2014Volume 88Number 15

A virus's reservoir consists of often-complex host communities that participate in its propagation and maintenance (13). Dynamics of individual host populations contribute to viral prevalence (14, 15), circulation, and dispersal (16); however, virus adaptation to its host can limit the potential for transmission between species (17, 18). The dynamics of viruses associated with their phylogenetics have been shown to be driven by a large number of complex and interacting factors (19–21), including the genetic diversity, ecology, and population dynamics of their host communities.

The southwestern Indian Ocean (SWIO) region is recognized as part of a biodiversity hot spot (22). In Madagascar, endemic mammalian taxa (including all native terrestrial and a significant percentage of volant species) have colonized the island independently since its separation from the Gondwana supercontinent more than 150 million years ago (23). In contrast, the smaller and much younger oceanic islands (mainly of volcanic origin) located around Madagascar (Mascarene, Comoros, and Seychelles archipelagos) host a limited number of mammalian species, and all of the terrestrial forms have been introduced (24, 25). The islands of the SWIO have been subjected to rapid anthropological expansion resulting in the dispersal and invasion of nonendemic rats and mice of the family Muridae and shrews of the family Soricidae to these isolated island ecosystems (26, 27). On virtually all of the SWIO islands, these introduced small-mammal populations occur in often dense populations, and on Madagascar, they compete with and displace native animal communities (28), potentially promoting the exchange of introduced infectious agents. These aspects make the SWIO an ideal location for the study of pathogen exchange (29).

In a previous study (3), we reported paramyxovirus infection in SWIO bats. Here, we have expanded our investigation into the diversity of this viral family within bats and other small mammals of Madagascar and other SWIO islands. We find that novel and unclassified morbillivirus-related paramyxoviruses (UMRVs) are commonplace within regional small mammals. Important differences between the morbillivirus-related and henipavirus-related virus models are observed, and detailed phylogenetic analyses suggest a major role for introduced *R. rattus* in diffusing UMRVs between different mammalian orders and different islands. These results highlight that UMRVs are a good model for understanding viral dynamics within wild fauna, especially between rodents and bats, two major reservoirs of zoonotic pathogens.

### **MATERIALS AND METHODS**

Field work and sample collection. A total of 597 small, nonflying mammals were captured on Madagascar at two sites over the course of 3 years (2010 to 2012), once on Mahé (Seychelles) (June 2011) and once on Mayotte (Comoros) (July 2012), and from different sites on La Réunion (September 2012 to March 2013). A total of 140 bats were collected from several locations in Madagascar in 2012. Details on animal capture techniques are provided in the Text S1 in the supplemental material, and those on sampling locations are provided in Table S3.

Ethics statement. All animals were manipulated in accordance with guidelines for the handling of wild mammals (30). This study benefitted from sampling efforts conducted in the context of an ongoing international long-term project to catalogue the terrestrial vertebrate fauna of Madagascar based on voucher specimens (31). All protocols followed the terms of research permits (see Acknowledgments) issued by national authorities: Ministère des Forêts et de l'Environnement, Madagascar National Parks, Département de Biologie Animale (Madagascar); Direction

de l'Environnement, de l'Aménagement et du Logement (France); and the Seychelles Bureau of Standards (Seychelles).

Laboratory work. (i) Nucleic acid preparation. Samples of approximately 1 to 2 mm³ of kidney, spleen, and lung tissues were dissected on sterile ice from each individual animal. These tissue samples were then combined in 750  $\mu$ l of Dulbecco's modified medium (Gibco, USA) containing 2- by 3-mm tungsten beads and homogenized for 2 min at 25 Hz in a TissueLyser (Qiagen). Homogenized organ samples were then clarified by centrifugation at  $10,000 \times g$  for 5 min. Two hundred  $\mu$ l of the clarified medium was biologically inactivated in 200  $\mu$ l of AVL buffer (Qiagen), and total nucleic acids were extracted using the EZ1 virus V2 minikit in an EZ1 biorobot (Qiagen). Total nucleic acids were reverse transcribed in the presence of random hexameric primers using the GoScript reverse transcription kit (Promega) according to the manufacturer's instructions.

(ii) PCR screening. cDNA from each sample was screened using a seminested PCR system that targets a partial sequence (~490 bp) of the L polymerase gene of respiroviruses, morbilliviruses, and henipaviruses (RMH) as described by Tong et al. (32) and previously exploited in similar studies (2–6). Negative controls were routinely incorporated, and PCRs were repeated independently to ensure no cross-contamination when using nested PCR protocols.

(iii) Statistical analyses. Differences in detected prevalence (defined as the proportion of animals tested from which the sequence of the RMH PCR product obtained had significant homology to known *Paramyxovirus* sequences in BLAST) were statistically analyzed using a two-tailed Fisher's exact test. P < 0.05 was considered statistically significant.

(iv) Sequencing. PCR cDNA products of the approximate anticipated size (450 to 500 bp) were purified using the Qiagen PCR purification kit and cloned into the pGEM-T vector system (Promega). Cloned PCR products were Sanger sequenced (Genoscreen) using M13 standard sequencing primers. The sequence quality of individual reads was assessed, and all sequences were processed using the Geneious Pro software package (33). DNA sequences obtained from at least three independent bacterial clones were aligned to correct for the majority of sequencing or PCR-introduced errors. Coinfection was defined when sequences from the same animal showed greater than 5% genetic dissimilarity; in these cases, at least three clones of each different sequence were obtained. Primer sequences were trimmed from the finalized sequences.

(v) OTU-based family-level phylogeny. In order to classify the detected paramyxoviruses, viral family-level phylogenetic analyses were performed. The search parameters "Taxonomic classification; Paramyxoviridae" and "Any Field: polymerase" were used in GenBank (15 December 2013) to generate the final comparative sequence data set. In total, 1,816 sequences were assessed for their compatibility via pairwise alignment against a reference sequence from the Henipavirus genus (JN255806). The Geneious Translational alignment tool, using the default ClustalW cost matrix, was used to align all compatible sequences. Sequences were trimmed to remove any free end gaps or entirely removed from the analysis if the obtained alignment did not provide at least 420 bp of non-gap overlap, leaving 1,192 sequences for phylogenetic analysis (final length after trim, excluding gaps, 438 bp). Internal gaps were permitted. Operational taxonomic units (OTU) were defined using mothur (34) with a genetic distance cutoff of 10%, generating 196 consensus sequences that spanned all known, classified, and unclassified paramyxovirus genera. Representative sequences were selected using mothur for each defined OTU, which were then used in the final phylogeny (see Fig. 3).

(vi) Sequence polymorphism. DNAsp v.5 (available at http://www.ub.edu/dnasp/) was used to calculate the estimates of genetic diversity, Pi and Theta, from subsets of the final nucleotide alignments, separated by host species.

(vii) UMRV (and henipavirus) phylogeny. Individual sequences from OTUs falling within the UMRV clade were extracted from the original sequence alignment and used for further phylogenetic analysis (see

Fig. 4 and 5; also see Fig. S2 in the supplemental material). *Salem virus* (JQ697837) was included as an outgroup reference sequence.

(viii) Phylogenetic analysis. All presented phylogenies were constructed using the same methodologies; from aligned sequence data sets, jModelTest v2.1.2 (35) identified GTR+I+G as the most appropriate substitution model for all phylogenetic analyses in BEAST (36). Parameter estimates for different clock models were assessed in Tracer v1.5.0 (36), and a strict molecular clock was used for all further analyses. An additional three replicates of 100,000,000 iterations then were performed using these optimal parameters. Trees and parameter estimate files for all independent replicates then were combined using LogCombiner (BEAST package), assigning a 10% burn-in, which left only converged parameter estimates from each repeat. The obtained effective sample size values for each parameter were all superior to 200. The final phylogeny was generated using TreeAnnotator, and trees were manipulated using FigTree v1.4.

(ix) Cartographic data. Distributions of hosts were downloaded from the IUCN Global Mammal Assessment (http://www.iucnredlist.org/). Maps were processed in the Quantum GIS 2.0.1 (Dufour) software package.

(x) Ancestral-state reconstruction. Bayesian phylogenies from each of the preceding clade-level analyses were resampled, and the resulting 4,000 trees were imported into Mesquite (version 2.75; http://mesquiteproject.org/mesquite/mesquite.html). Geographical origins and mammalian host categories were attributed to each sequence as discrete state characters. The number of reconstructed trait changes was calculated using the unordered parsimony assumption and averaged for each ordinal category and over all trees, as described in reference 4.

**Nucleotide sequence accession numbers.** All sequences used for the present analyses have been deposited in GenBank under the reference numbers KF245939 to KF246061, KF408256 to KF408261, and KF928225 to KF928265.

### **RESULTS**

Sampling across the SWIO islands yielded a wealth of animal samples. A description of the animal species subject to trapping activities is provided in Text S2 in the supplemental material. A molecular epidemiology survey for paramyxoviruses was carried out on 732 samples from four different animal orders, Chiroptera, Afrosoricida, Rodentia, and Soricomorpha, collected over a period of 3 years (2010 to 2012) in the SWIO region, specifically Madagascar, La Réunion, Mayotte (Comoros), and Mahé (Seychelles). All screening data are summarized in Fig. 1.

Madagascar. A total of 391 small nonflying mammals belonging to three orders (Rodentia, Afrosoricida, and Soricomorpha) and 22 different species were sampled at two sites on Madagascar. These included endemic Nesomyidae rodents (Eliurus, Gymnuromys, and Nesomys) and Tenrecidae tenrecs (Microgale, Oryzorictes, Hemicentetes, Setifer, and Tenrec), as well as introduced Muridae rodents (Rattus) and Soricidae shrews (Suncus). The species captured at each of the two sites varied, although the species diversity was comparable. The numbers of taxa captured at each site showed little temporal variation during the different field visits, suggesting homogenous sampling between years; however, a larger number of R. rattus specimens were collected in Lakato in 2012 than in the previous years.

Of the Malagasy nonflying mammals screened for paramyxoviruses, 25% (98/391) tested positive. Shrew tenrecs (Tenrecidae) belonging to the genus Microgale, for which the majority of sampled animals were represented by  $M.\ cowani\ (n=73)$  and  $M.\ dobsoni\ (n=54)$ , showed the highest rate of paramyxovirus detection (40% positive; 69/172). Of these, four animals were coinfected with at least two different paramyxovirus strains (see Table

S1 in the supplemental material). Paramyxoviruses were also detected in rodents, including *R. rattus* (30% positive; 16/54) and *Eliurus minor* (11% positive; 12/111). Variation in paramyxovirus infection levels was observed over the 3 years of collections on Madagascar, most markedly in populations of *M. cowani*, where differences between consecutive years were strongly significant (P < 0.001 for 2010 to 2011 and P = 0.001 for 2011 to 2012) and paramyxovirus infections peaked in 2011, with 85% of animals testing positive (Fig. 2).

Of the 140 Malagasy bats screened, a total of 41 belonging to eight bat species endemic to SWOI (*Chaerephon leucogaster*, *Miniopterus griveaudi*, *Mops leucostigma*, *Mormopterus jugularis*, *Myotis goudoti*, *Otomops madagascariensis*, *Triaenops menamena*, and *Vespertilionidae* spp.) and from four bat families (Hipposideridae, Miniopteridae, Molossidae, and Vespertilionidae) tested positive for paramyxoviruses.

Animals not endemic to other SWIO islands. Nonendemic small mammals collected outside Madagascar included *R. rattus*, *R. norvegicus*, and *Suncus murinus* on La Réunion and *R. rattus* on Mayotte and Mahé; paramyxoviruses were detected at each of these locations (Fig. 1), with *Rattus* spp. showing the highest proportion of positive animals (22%; 42/187).

Genetic and phylogenetic analyses. Phylogenetic analysis based on OTUs was sufficient to generate a well-supported, family-level phylogeny containing monophyletic groups for all main *Pneumovirinae* and *Paramyxovirinae* genera (Fig. 3). From this phylogeny, 10 clades were defined based on well-supported nodes (posterior probability [PP] > 0.8) that provided definitions at the taxonomic level of genus based on current taxonomic classifications. All of the 177 paramyxovirus sequences from the SWIO were members of a single genus-level group (UMRVs).

Absolute levels of genetic divergence were comparable between henipaviruses, morbilliviruses, and the UMRVs (see Fig. S1 in the supplemental material). Partial L-gene sequence divergence was calculated for those host species groups that were well represented in global sequence data and of interest for this study (see Table S2 in the supplemental material). Interestingly, less genetic divergence was observed between viral sequences originating from host species of different bird and mammalian orders (Aves, Afrosoricida, Rodentia, Soricomorpha, and Scandentia) than between viral sequences originating from within the Chiroptera, suggesting that bats can be infected by a broader range of paramyxoviruses than other hosts (Table 1). However, within the UMRV group, the genetic diversity was comparable between viral sequences from rodents and bats (Table 2).

Further phylogenetic analyses of all sequences that fell within the UMRV OTU-defined subgroups indicate that the detected paramyxoviruses comprise a certain level of host specificity at the level of host order (Fig. 4; also see Fig. S2 in the supplemental material). Two well-supported clades contain the majority of sequences originating from rodents, an additional two clades include most sequences originating from bats, and one final clade contains the greater part of sequences from tenrecs. Geographical clustering within the phylogeny was less apparent apart from viruses found in Afrosoricida, which are endemic to Madagascar (Fig. 5; also see Fig. S2). Cartographic analysis of the distributions of those species identified as hosts of the different UMRV groups (Fig. 4) suggested the global dispersal of many of these viral agents.

Members of the UMRVs included the Jeilam viruses (J, Beilong, and Tailam), viruses of rodent origin isolated in Australia

**8270** jvi.asm.org Journal of Virology




FIG 1 Paramyxovirus detection at different sampling sites in Madagascar and the SWIO. Numbers of positive animals, total numbers of sampled animals, and the year of capture are indicated for each sampling site. Species names preceded by an asterisk represent introduced taxa.

and China. Genetic and virological evidence has previously demonstrated that these viruses belong to the same taxonomic group (37–39). Interestingly, a strongly supported phylogenetic substructure separated Beilong and J viruses (see Fig. S2 in the supplemental material). The well-supported monophyletic group uniting these two viruses (PP = 1.0), designated rodent group 2 in Fig. S2 in the supplemental material, contained a further 71 paramyxovirus sequences, which were obtained from rodents and other small mammals from across the globe. Further well-supported substructures within rodent group 2 demonstrated differences in geographical distribution and host species association; for example, one subgroup contained only sequences obtained from the southern African rodent *Rhabdomys pumilio* (Muridae), whereas Beilong and Tailam viruses branched closely with *Rattus* species-derived sequences from the SWIO.

Ancestral-state reconstructions of UMRVs. Ancestral-state reconstructions were used to estimate the number of viral host switches and geographical exchanges based on the phylogeny pre-

sented in Fig. 2. These data are presented in donor/acceptor probability heatmap form in Fig. 4. Host-switching events were predicted to be most frequent between closely related species and between species belonging to the same order. *R. rattus* was predicted to participate in the largest number of host-switching events, being both the principle donor and principal acceptor species. *R. rattus* was also predicted to be involved in the majority of host exchanges that occurred between distantly related animal orders. Notably, elevated numbers of bidirectional viral exchanges between *R. rattus* and the Tenrecidae (*Microgale* spp.) were predicted to have occurred, and bats were also predicted to have received paramyxoviruses from *R. rattus*.

The geographical flux of paramyxoviruses was predicted to have occurred at least five times more frequently between Africa and the SWIO than any other continental regions. Exchange was also predicted to be frequent between SWIO islands, in particular between Madagascar, La Réunion, and the Comoros. It is important to note that our ancestral-state reconstructions could be in-




FIG 2 Temporal variation of detected paramyxovirus infection levels in animals from Madagascar. The number of paramyxovirus-positive animals is indicated in percentages. The total number of animals tested for each year is indicated above the bars. Colors are used to highlight the year of sampling, as described in the legend. P values displayed above each graph indicate the significance of differences between consecutive years (two-tailed Fisher's exact test; ns, not significant).

fluenced by a possible overrepresentation of paramyxovirus sequences from the SWIO; however, it should be noted that other large-scale screening efforts across the world have also contributed considerable data (2–6), originating from both rodents and bats and distributed across most known paramyxovirus genera (see Table S3 in the supplemental material). The limitations of similar analyses have been discussed in detail elsewhere for paramyxoviruses (4) and lyssaviruses (40).

For comparative purposes, similar analyses were performed on sequences from the henipavirus and unclassified henipavirus-related virus (UHRV) genus of the *Paramyxoviridae* (see Fig. S3 in the supplemental material). Here, host exchanges corresponding to human disease emergence via pigs and horses could be reliably predicted from the generated phylogeny, and geographical exchanges predicted an African origin for the henipaviruses, as

well as large-scale circulation on the African continent, in agreement with other published literature on henipavirus biogeography (4, 8).

### DISCUSSION

We have shown that many lineages of *Morbillivirus*-related paramyxovirus are in active circulation between islands in the SWIO. Bats and terrestrial small mammals play host to a number of highly divergent paramyxovirus strains belonging to this UMRV group.

Large numbers of paramyxovirus-positive animals were detected in this study. Endemic *Microgale* spp. and *Triaenops menamena* were identified as important paramyxovirus hosts, where 40% and 63% of animals, respectively, tested positive. In regional populations of introduced *Rattus* spp., this figure was




FIG 3 Paramyxovirus clades of the SWIO. (A) OTU-based family-level phylogeny of paramyxoviruses. The presented phylogeny is a proportional, ordered representation of data generated in BEAST. Relevant posterior probabilities (>0.5) are shown above each branch for clarity. Sequences are grouped into functional clades based on node support and height (see the text). Each functional group is assigned a letter (A to J). All sequences obtained within this study fell within group J, which is highlighted in black.

**8272** jvi.asm.org Journal of Virology

TABLE 1 Genetic diversity of the Paramyxoviridae by host order

	Viral diversity in host ord	er(s):		
Parameter	Aves	Chiroptera	Rodentia	Scandentia, Soricomorpha, and Afrosoricida
Pi (SD) Theta	0.21123 (0.01385) 0.32854	0.38181 (0.00730) 0.35262	0.25410 (0.00947) 0.26267	0.32354 (0.01149) 0.33052

<sup>&</sup>quot; Sequences from mammalian orders Scandentia and Soricomorpha are grouped with sequences from Afrosoricida due to the small number of viral sequences originating from animals of these orders. Pi is an estimation of nucleotide diversity, and Theta is equivalent to the Watterson estimator of the population mutation rate.

20%. A recent study (4) reported that *R. rattus* in Gabon and Thailand and *R. norvegicus* in Germany had low levels of infection (0/113, 1/49, and 0/131, respectively) when employing similar detection methods. However, the same study reported elevated numbers of paramyxoviruses in southern African *Rhabdomys pumilio* populations (17% of 512 animals tested). Additionally, another study conducted in Zambia has recently reported comparably high paramyxovirus infection levels in rodent (19%) and shrew (40%) populations (7). It is also of note that no henipavirus-related viruses were detected as part of this study despite the existence of serological evidence for the circulation of henipaviruses in Madagascar (41); however, members of this viral genus most often are associated with Pteropodidae fruit bats, which were absent from our samples.

The genetic diversity of partial L-gene sequences studied here is notably high, which is suggestive of the circulation of a large number of previously uncategorized viral strains (4). Additionally, multiple viral lineages could be identified within individual hosts, suggesting a level of compatibility between infecting viral strains. The overall observed genetic diversity of all paramyxoviruses per host was highest within the Chiroptera, suggesting that bats host a broader range of paramyxoviruses than terrestrial small mammals, in agreement with a recent hypothesis that bats are more permissive to viral infection than rodents (1). However, within the UMRV genus the level of viral diversity was similar between rodents and bats, showing that the relative importance of each host species likely is viral genus specific.

Different factors that may drive the observed levels of high genetic diversity of paramyxoviruses include the following: (i) the inherent genetic drift associated with many RNA viruses (42), which alone may be insufficient to explain high genetic variation between closely related viral lineages; (ii) coinfection within individual hosts, which is a "prerequisite for genetic exchange between different pathogen species or strains" (43), was observed in this and other studies of novel *Paramyxovirus* genera (2–4), and the possibility of recombination events between highly similar viruses should not be excluded, although their observation is rare for members of the *Paramyxoviridae*; and (iii) viral adaptation due to exchange within multihost systems (18).

The observed phylogenetic structuring of a divergent viral population of Jeilam-related viruses suggests that these closely related lineages have followed separate ancestral paths, generating a population structure similar to that of previously reported viral metapopulations (44, 45). Furthermore, analysis of the overall sequence diversity within well-defined *Paramyxovirus* genera suggests that the studied small fragment (~450 bp) is prone to genetic diversification compared to other loci of the paramyxovirus genome (46). Thus, in the absence of further virological or genetic data, the taxonomy of these viruses remains imprecise. In contrast, the observed genetic diversity, in association with considerable data originating from sources across the globe, has proven efficient for tracing the evolutionary paths of different viral lineages.

Using previously proposed terminology (13), the abundance of observed paramyxovirus infections suggests that animals of the SWIO islands constitute a putative paramyxovirus maintenance community. We predict, based on the reported virus-host interactions, that interspecies contacts are frequent within the natural ecosystems of Madagascar, in turn promoting what has elsewhere been referred to as a disease hot spot (22). For example, within 125 km of our two study sites, 12 *Microgale* species occur in syntopic and densely packed ecosystems (47). Thus, a considerable number of phylogenetically closely related host species, as well as animals of different mammalian orders, interact within dense and closed biotic systems, and host-specific adaptation appears to impose few barriers to viral host switching, resulting in frequent viral spillover and persistence within a multispecies community.


Viral persistence within a classical model source/sink community can be accompanied by genetic adaptation that establishes efficient virus-host interactions (17, 48); this may eventually create an evolutionary barrier that inhibits further host switching and results in the establishment of a new source population. This evolutionary process becomes complex for multihost ecosystems that exhibit frequent bidirectional host switching, such as that observed on Madagascar. Numerous factors will influence the efficiency of viral spreading within any studied community, including (i) viral and host abundance, (ii) host fitness, (iii) infection cost to the host, and (iv) the frequency and mode of transmission events (48). Infection dynamics within reservoir populations also directly influence viral epidemiology, with peaks of infection resulting in increased viral transmission (16). Here, the proportion of paramyxovirus-positive *Microgale cowani* increased 4-fold be-

TABLE 2 Genetic diversity within the UMRV genus per host group<sup>a</sup>

	Genetic diversity in:			
Parameter	Chiroptera	Rattus spp.	Rodentia	Afrosoricida (Tenrecidae)
Pi (SD)	0.28304 (0.00637)	0.24013 (0.01689)	0.28515 (0.00767)	0.22637 (0.00964)
Theta	0.27860	0.23846	0.25717	0.22740

<sup>&</sup>lt;sup>a</sup> Pi is an estimation of nucleotide diversity, and Theta is equivalent to the Watterson estimator of the population mutation rate.

August 2014 Volume 88 Number 15



**FIG 4** Ancestral-state reconstructions: host switching. The number of estimated host-switching events between species, based on parsimony analyses in Mesquite, are represented in tabular heatmap form. The presented phylogenetic tree is identical to that in Fig. 5 and Fig. S2 in the supplemental material and presents the most likely arrangement of the phylogenies used to estimate host-switching numbers. Dots represent nodes with posterior support greater than 0.9. Colors denote host-species order throughout. Red stars indicate positions of *R. rattus* in the phylogeny and the table.

8274 jvi.asm.org Journal of Virology




FIG 5 Ancestral-state reconstructions: geographical exchange. The number of estimated geographical switching events between sites based on parsimony analyses in Mesquite are represented in tabular form and are heatmap colored based on the estimated frequencies. The presented phylogenetic tree is identical to that in Fig. S2 in the supplemental material and Fig. 4 and presents the most likely arrangement of the phylogenies used to estimate geographical exchange numbers. Dots represent nodes with posterior support greater than 0.9. Colors depict continental origins.

tween 2010 and 2011, coinciding with the detection of closely related viral lineages in other host species. Spillover is a critical process for understanding zoonotic transmission but has rarely been documented in wild animal populations.

In addition to circulation within insular ecosystems, viral reservoirs can be established over large host communities, aided by their widespread geographical dispersal. From our data, the estimated levels of UMRV exchange between SWIO islands were high compared to global transmission levels, implying active host-associated dispersal between the African continent, Madagascar, and other SWIO islands.

Bats are theorized to play crucial roles in the processes of viral evolution that may lead to disease emergence (49) and are important hosts for numerous paramyxoviruses. Unlike henipaviruses, whose geographical distribution can be largely explained by the dispersal of Pteropodidae fruit bats (50), the observed geographical distribution of UMRVs is unlikely to be attributed uniquely to the dispersal of those bat species from which these viruses have been detected and most likely relies on interaction with other animal hosts. The data presented here indicate an important role for R. rattus in the transmission of UMRV paramyxoviruses and highlight these animals as potential intermediates in the dissemination of infectious agents that are endemic or otherwise isolated that facilitate disease emergence. The global dispersal of Rattus and their associated biota have been studied in many different contexts (25, 51) and have particularly important implications within the SWIO (52), where many mammal species are endemic, particularly on Madagascar, and at risk from both macro- and microorganism invasions, and where active commercial trade results in considerable and largely uncontrolled exchange (24, 53).

Within the insular ecosystems of the SWIO, paramyxovirus infection provides an excellent model for the study of both viral population dynamics and biogeography. Geographical exchange and host-switching events are common, likely occurring over

short time scales due to some ecological factors that promote gene flow within an expansive viral maintenance community. In this context, *Rattus* play the dual role of a viral maintenance population and spreaders. Further virological investigation into the evolutionary implications of our observations may go a long way to developing an understanding of viral evolution within complex multihost communities.

### **ACKNOWLEDGMENTS**

This work was supported by funding from CRVOI, ERDF/French Government/Regional Council of La Réunion, FEDER POCT (2009–2013) "Pathogènes associés à la Faune Sauvage Océan Indien" 31189, and the European Regional Development Funds FEDER-POCT; Réunion, ParamyxOI project. D. A. Wilkinson, M. Dietrich, and B. Ramasindrazana's postdoctoral fellowships were funded by "RUN-Emerge: European project funded by European Commission under FP7 program." J. Mélade's doctoral stipend was financed by the Conseil Regionale de La Réunion. B. Ramasindrazana received support from the "Fonds de Coopération régionale, Prefecture de La Réunion" and from the RUN Emerge project.

The following animal capture permits were used: (i) La Réunion, no permits required; (ii) Mahé, Research Approval Letter A0347; (iii) Madagascar, 350/10/MEF/SG/DGF/DCB.SAP/SCB, 032/12/MEF/SG/DGF/DCB.SAP/SCBSE, and 194/12/MEF/SG/DGF/DCB.SAP/SCBSE, and 194/12/MEF/SG/DGF/DCB.SAP/SCBS; and (iv) Mayotte, no permits required.

We thank Gerard Rocamora and the Island Conservation Society for help with captures in Mahé, and we also thank Arivony Nomenjanahary for contributing to the work in Madagascar.

We have no conflicts of interest in relation to the submitted work to declare.

### **REFERENCES**

1. Luis AD, Hayman DT, O'Shea TJ, Cryan PM, Gilbert AT, Pulliam JR, Mills JN, Timonin ME, Willis CK, Cunningham AA, Fooks AR, Rupprecht CE, Wood JL, Webb CT. 2013. A comparison of bats and rodents as reservoirs of zoonotic viruses: are bats special? Proc. Biol. Sci. R. Soc. 280:20122753. http://dx.doi.org/10.1098/rspb.2012.2753.

August 2014 Volume 88 Number 15

- Baker KS, Todd S, Marsh G, Fernandez-Loras A, Suu-Ire R, Wood JL, Wang LF, Murcia PR, Cunningham AA. 2012. Co-circulation of diverse paramyxoviruses in an urban African fruit bat population. J. Gen. Virol. 93:850–856. http://dx.doi.org/10.1099/vir.0.039339-0.
- Wilkinson DA, Temmam S, Lebarbenchon C, Lagadec E, Chotte J, Guillebaud J, Ramasindrazana B, Heraud JM, de Lamballerie X, Goodman SM, Dellagi K, Pascalis H. 2012. Identification of novel paramyxoviruses in insectivorous bats of the southwest Indian Ocean. Virus Res. 170:159–163. http://dx.doi.org/10.1016/j.virusres.2012.08.022.
- 4. Drexler JF, Corman VM, Muller MA, Maganga GD, Vallo P, Binger T, Gloza-Rausch F, Rasche A, Yordanov S, Seebens A, Oppong S, Adu Sarkodie Y, Pongombo C, Lukashev AN, Schmidt-Chanasit J, Stocker A, Carneiro AJ, Erbar S, Maisner A, Fronhoffs F, Buettner R, Kalko EK, Kruppa T, Franke CR, Kallies R, Yandoko ER, Herrler G, Reusken C, Hassanin A, Kruger DH, Matthee S, Ulrich RG, Leroy EM, Drosten C. 2012. Bats host major mammalian paramyxoviruses. Nat. Commun. 3:796. http://dx.doi.org/10.1038/ncomms1796.
- Kurth A, Kohl C, Brinkmann A, Ebinger A, Harper JA, Wang LF, Muhldorfer K, Wibbelt G. 2012. Novel paramyxoviruses in free-ranging European bats. PLoS One 7:e38688. http://dx.doi.org/10.1371/journal .pone.0038688.
- Sasaki M, Setiyono A, Handharyani E, Rahmadani I, Taha S, Adiani S, Subangkit M, Sawa H, Nakamura I, Kimura T. 2012. Molecular detection of a novel paramyxovirus in fruit bats from Indonesia. Virol. J. 9:240. http://dx.doi.org/10.1186/1743-422X-9-240.
- Sasaki M, Muleya W, Ishii A, Orba Y, Hangombe BM, Mweene AS, Moonga L, Thomas Y, Kimura T, Sawa H. 2013. Molecular epidemiology of paramyxoviruses in Zambian wild rodents and shrews. J. Gen. Virol. 95(Part 2):325–330. http://dx.doi.org/10.1099/vir.0.058404-0.
- Peel AJ, Sargan DR, Baker KS, Hayman DT, Barr JA, Crameri G, Suu-Ire R, Broder CC, Lembo T, Wang LF, Fooks AR, Rossiter SJ, Wood JL, Cunningham AA. 2013. Continent-wide panmixia of an African fruit bat facilitates transmission of potentially zoonotic viruses. Nat. Commun. 4:2770. http://dx.doi.org/10.1038/ncomms3770.
- Aguilar HC, Lee B. 2011. Emerging paramyxoviruses: molecular mechanisms and antiviral strategies. Expert Rev. Mol. Med. 13:e6. http://dx.doi.org/10.1017/S1462399410001754.
- Halpin K, Hyatt AD, Plowright RK, Epstein JH, Daszak P, Field HE, Wang L, Daniels PW. 2007. Emerging viruses: coming in on a wrinkled wing and a prayer. Clin. Infect. Dis. 44:711–717. http://dx.doi.org/10.1086/ /511078.
- 11. Luby SP, Hossain MJ, Gurley ES, Ahmed BN, Banu S, Khan SU, Homaira N, Rota PA, Rollin PE, Comer JA, Kenah E, Ksiazek TG, Rahman M. 2009. Recurrent zoonotic transmission of Nipah virus into humans, Bangladesh, 2001-2007. Emerg. Infect. Dis. 15:1229–1235. http://dx.doi.org/10.3201/eid1508.081237.
- Albarino CG, Foltzer M, Towner JS, Rowe LA, Campbell S, Jaramillo CM, Bird BH, Reeder DM, Vodzak ME, Rota P, Metcalfe MG, Spiropoulou CF, Knust B, Vincent JP, Frace MA, Nichol ST, Rollin PE, Stroher U. 2014. Novel paramyxovirus associated with severe acute febrile disease, South Sudan and Uganda, 2012. Emerg. Infect. Dis. 20:211–216. http://dx.doi.org/10.3201/eid2002.131620.
- Haydon DT, Cleaveland S, Taylor LH, Laurenson MK. 2002. Identifying reservoirs of infection: a conceptual and practical challenge. Emerg. Infect. Dis. 8:1468–1473. http://dx.doi.org/10.3201/eid0812.010317.
- Gloza-Rausch F, Ipsen A, Seebens A, Gottsche M, Panning M, Drexler JF, Petersen N, Annan A, Grywna K, Muller M, Pfefferle S, Drosten C. 2008. Detection and prevalence patterns of group 1 coronaviruses in bats, northern Germany. Emerg. Infect. Dis. 14:626–631. http://dx.doi.org/10 3201/eid1404.071439.
- Baker KS, Suu-Ire R, Barr J, Hayman DT, Broder CC, Horton DL, Durrant C, Murcia PR, Cunningham AA, Wood JL. 23 September 2013. Viral antibody dynamics in a chiropteran host. J. Anim. Ecol. http://dx.doi.org/10.1111/1365-2656.12153.
- Kallio ER, Begon M, Henttonen H, Koskela E, Mappes T, Vaheri A, Vapalahti O. 2009. Cyclic hantavirus epidemics in humans–predicted by rodent host dynamics. Epidemics 1:101–107. http://dx.doi.org/10.1016/j .epidem.2009.03.002.
- Dennehy JJ, Friedenberg NA, Holt RD, Turner PE. 2006. Viral ecology and the maintenance of novel host use. Am. Nat. 167:429–439. http://dx .doi.org/10.1086/499381.
- Woolhouse ME, Webster JP, Domingo E, Charlesworth B, Levin BR. 2002. Biological and biomedical implications of the co-evolution of

- pathogens and their hosts. Nat. Genet. 32:569–577. http://dx.doi.org/10.1038/ng1202-569.
- Streicker DG, Lemey P, Velasco-Villa A, Rupprecht CE. 2012. Rates of viral evolution are linked to host geography in bat rabies. PLoS Pathog. 8:e1002720. http://dx.doi.org/10.1371/journal.ppat.1002720.
- Streicker DG, Altizer SM, Velasco-Villa A, Rupprecht CE. 2012. Variable evolutionary routes to host establishment across repeated rabies virus host shifts among bats. Proc. Natl. Acad. Sci. U. S. A. 109:19715–19720. http://dx.doi.org/10.1073/pnas.1203456109.
- Volz EM, Koelle K, Bedford T. 2013. Viral phylodynamics. PLoS Comput. Biol. 9:e1002947. http://dx.doi.org/10.1371/journal.pcbi.1002947.
- Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GA, Kent J. 2000. Biodiversity hotspots for conservation priorities. Nature 403:853– 858. http://dx.doi.org/10.1038/35002501.
- Goodman SM, Benstead JP (ed). 2003. The natural history of Madagascar. University of Chicago Press, Chicago, IL.
- 24. Cheke A. 2010. The timing of arrival of humans and their commensal animals on western Indian Ocean oceanic islands. Phelsuma 18:38–69. http://www.academia.edu/3126693/The\_timing\_of\_arrival\_of\_humans\_ and\_their\_commensal\_animals\_on\_Western\_Indian\_Ocean\_oceanic\_is lands.
- 25. Aplin KP, Suzuki H, Chinen AA, Chesser RT, Ten Have J, Donnellan SC, Austin J, Frost A, Gonzalez JP, Herbreteau V, Catzeflis F, Soubrier J, Fang YP, Robins J, Matisoo-Smith E, Bastos AD, Maryanto I, Sinaga MH, Denys C, Van Den Bussche RA, Conroy C, Rowe K, Cooper A. 2011. Multiple geographic origins of commensalism and complex dispersal history of black rats. PLoS One 6:e26357. http://dx.doi.org/10.1371/journal.pone.0026357.
- 26. Hutterer R, Tranier M. 1990. The immigration of the Asian house shrew (Suncus murinus) into Africa and Madagascar, p 309–319. In Peters G, Hutterer R (ed), Vertebrates in the tropics. Museum Alexander Koenig, Bonn, Germany.
- Duplantier JM, Orth A, Catalan J, Bonhomme F. 2002. Evidence for a mitochondrial lineage originating from the Arabian peninsula in the Madagascar house mouse (Mus musculus). Heredity 89:154–158. http://dx .doi.org/10.1038/sj.hdy.6800122.
- Goodman SM. 1995. The spread of *Rattus* on Madagascar: the dilemma of protecting the endemic rodent fauna. Conservation Biol. 9:450–453. http: //dx.doi.org/10.1046/j.1523-1739.1995.9020450.x.
- Tortosa P, Pascalis H, Guernier V, Cardinale E, Le Corre M, Goodman SM, Dellagi K. 2012. Deciphering arboviral emergence within insular ecosystems. Infect. Genet. Evol. 12:1333–1339. http://dx.doi.org/10.1016/j.meegid.2012.03.024.
- Sikes RS, Gannon WL. 2011. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. J. Mammal. 92: 235–253. http://dx.doi.org/10.1644/10-MAMM-F-355.1.
- Goodman SM, Raherilalao MJ. 2013. Atlas of selected land vertebrates of Madagascar. Association Vahatra, Antananarivo, Madagascar.
- Tong S, Chern SW, Li Y, Pallansch MA, Anderson LJ. 2008. Sensitive and broadly reactive reverse transcription-PCR assays to detect novel paramyxoviruses. J. Clin. Microbiol. 46:2652–2658. http://dx.doi.org/10 .1128/JCM.00192-08.
- 33. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics 28:1647–1649. http://dx.doi.org/10.1093/bioinformatics/bts199.
- 34. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF. 2009. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. Appl. Environ. Microbiol. 75:7537–7541. http://dx.doi.org/10.1128/AEM.01541-09.
- Darriba D, Taboada GI, Doallo R, Posada D. 2012. jModelTest 2: more models, new heuristics and parallel computing. Nat. Methods 9:772. http://dx.doi.org/10.1038/nmeth.2109.
- Drummond AJ, Suchard MA, Xie D, Rambaut A. 2012. Bayesian phylogenetics with BEAUti and the BEAST 1.7. Mol. Biol. Evol. 29:1969–1973. http://dx.doi.org/10.1093/molbev/mss075.
- 37. Li Z, Yu M, Zhang H, Magoffin DE, Jack PJ, Hyatt A, Wang HY, Wang LF. 2006. Beilong virus, a novel paramyxovirus with the largest genome of

8276 jvi.asm.org Journal of Virology

- non-segmented negative-stranded RNA viruses. Virology 346:219–228. http://dx.doi.org/10.1016/j.virol.2005.10.039.
- http://dx.doi.org/10.1016/j.virol.2005.10.039.

  38. Woo PC, Lau SK, Wong BH, Wong AY, Poon RW, Yuen KY. 2011.
  Complete genome sequence of a novel paramyxovirus, Tailam virus, discovered in Sikkim rats. J. Virol. 85:13473–13474. http://dx.doi.org/10.1128/JVI.06356-11.
- Magoffin DE, Mackenzie JS, Wang LF. 2007. Genetic analysis of J-virus and Beilong virus using minireplicons. Virology 364:103

  —111. http://dx .doi.org/10.1016/j.virol.2007.01.045.
- Faria NR, Suchard MA, Rambaut A, Streicker DG, Lemey P. 2013. Simultaneously reconstructing viral cross-species transmission history and identifying the underlying constraints. Philos. Trans. R. Soc. Lond. B Biol. Sci. 368:20120196. http://dx.doi.org/10.1098/rstb.2012.0196.
- Iehle C, Razafitrimo G, Razainirina J, Andriaholinirina N, Goodman SM, Faure C, Georges-Courbot MC, Rousset D, Reynes JM. 2007. Henipavirus and Tioman virus antibodies in pteropodid bats, Madagascar. Emerg. Infect. Dis. 13:159–161. http://dx.doi.org/10.3201/eid1301 060791.
- Holmes EC, Grenfell BT. 2009. Discovering the phylodynamics of RNA viruses. PLoS Comput. Biol. 5:e1000505. http://dx.doi.org/10.1371 /journal.pcbi.1000505.
- Malpica JM, Sacristan S, Fraile A, Garcia-Arenal F. 2006. Association and host selectivity in multi-host pathogens. PLoS One 1:e41. http://dx .doi.org/10.1371/journal.pone.0000041.
- Frost SD, Dumaurier MJ, Wain-Hobson S, Brown AJ. 2001. Genetic drift and within-host metapopulation dynamics of HIV-1 infection. Proc. Natl. Acad. Sci. U. S. A. 98:6975–6980. http://dx.doi.org/10.1073/pnas .131056998.
- Bahl J, Nelson MI, Chan KH, Chen R, Vijaykrishna D, Halpin RA, Stockwell TB, Lin X, Wentworth DE, Ghedin E, Guan Y, Peiris JS, Riley S, Rambaut A, Holmes EC, Smith GJ. 2011. Temporally structured

- metapopulation dynamics and persistence of influenza A H3N2 virus in humans. Proc. Natl. Acad. Sci. U. S. A. 108:19359–19364. http://dx.doi.org/10.1073/pnas.1109314108.
- McCarthy AJ, Goodman SJ. 2010. Reassessing conflicting evolutionary histories of the Paramyxoviridae and the origins of respiroviruses with Bayesian multigene phylogenies. Infect. Genet. Evol. 10:97–107. http://dx .doi.org/10.1016/j.meegid.2009.11.002.
- Dammhahn M, Soarimalala V, Goodman SM. 2013. Trophic niche differentiation and microhabitat utilization in a species-rich montane forest small mammal community of eastern Madagascar. Biotropica 45:111– 118. http://dx.doi.org/10.1111/j.1744-7429.2012.00893.x.
- Morse SS. 1995. Factors in the emergence of infectious diseases. Emerg. Infect. Dis. 1:7–15. http://dx.doi.org/10.3201/eid0101.950102.
- Wang LF, Walker PJ, Poon LL. 2011. Mass extinctions, biodiversity and mitochondrial function: are bats "special" as reservoirs for emerging viruses? Curr. Opin. Virol. 1:649–657. http://dx.doi.org/10.1016/j.coviro .2011.10.013.
- Dikid T, Jain SK, Sharma A, Kumar A, Narain JP. 2013. Emerging & re-emerging infections in India: an overview. Indian J. Med. Res. 138: 19–31.
- 51. Ewer RF. 1971. The biology and behaviour of a free-living population of black rats (*Rattus rattus*). Elsevier, New York, NY.
- 52. Brouat C, Rahelinirina S, Loiseau A, Rahalison L, Rajerison M, Laffly D, Handschumacher P, Duplantier JM. 2013. Plague circulation and population genetics of the reservoir *Rattus rattus*: the influence of topographic relief on the distribution of the disease within the Madagascan focus. PLoS Negl. Trop. Dis. 7:e2266. http://dx.doi.org/10.1371/journal.pntd.0002266.
- Četre-Sossah C, Pedarrieu A, Guis H, Defernez C, Bouloy M, Favre J, Girard S, Cardinale E, Albina E. 2012. Prevalence of Rift Valley fever among ruminants, Mayotte. Emerg. Infect. Dis. 18:972–975.

### 1 Supplementary Material:

- 2 Supplementary Figure and Table Legends:
- 3 Supplementary Table S1: Paramyxovirus co-infections. Highlighted are animals from which
- 4 multiple paramyxovirus L-gene sequences could be obtained, that showed clearly divergent
- 5 sequences. Multiple discrete sequences are suggestive of viral co-infections.
- 6 Supplementary Table S2: Genbank accession numbers are presented for paramyxovirus
- 7 sequences originating from bats, rodents or other small mammal species that were of
- 8 interest to this study. Sequences are grouped into the major clades of the paramyxovirus
- 9 family, based on family level phylogenies presented in main text Figure 2. Host-species
- 10 names and the number of sequences originating from that species are shown for each group.
- 11 Sequences generated as part of this study are underlined.
- 12 **Supplementary Table S3:** Description of capture sites, dates and species.
- 13 **Supplementary Figure S1:** Graphical representation of the genetic similarity (% identity)
- 14 between different functional clade groups of the paramyxoviridae (see Figure 2). Numbers
- 15 below the diagonal are the mean identities between all individual members of the compared
- 16 groups. Numbers above the diagonal are the corresponding standard deviations of sequence
- 17 identities. The indicated color scale of green to red highlights strong to weak sequence
- 18 conservation.
- 19 Supplementary Figure S2 (previous page). Phylogeny of all sequences belonging to the
- 20 UMRV group. Groups are defined by majority host-species. Red stars indicate the positions
- 21 of Beilong (DQ100461) and J-viruses (NC007454, AY900001) in the phylogenetic tree.
- 22 Supplementary Figure S3. Geographical distributions of paramyxovirus-positive
- 23 species. The potential extent of each group presented in Supplementary Figure S2 is

calculated from the geographical distribution of its hosts, according to the IUCN database (http://www.iucnredlist.org/), and is indicated in dark red to the right. Light red is used for locations outside of Madagascar for the "Malagasy Afrotheria Group", all sequences within this phylogenetic group originated from Madagascar including the minority of sequences that were detected in R. rattus. There is no evidence that viruses sharing sequence homology to the Malagasy Afrotheria Group exist outside of Madagascar.

Supplementary Figure S4. Ancestral state reconstructions – *Henipaviruses* and *Unclassified Henipa-related Viruses*. Ancestral host-species (A) and geographical (B) states were reconstructed and used to predict switching events in Mesquite based on 4000 raw trees of the *Henipavirus* phylogeny generated in BEAST. Host switching events that are linked with disease emergences in human populations are highlighted in green.

### Supplementary Text 1: Animal Capture Methodology

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

For terrestrial mammals, two different field techniques were used on Madagascar. In montane humid forests, pitfall traps were used as previously described (1, 2). Briefly, a series of 12 I buckets, each separated by 10 m distance, are buried into the ground and with the rim flush with soil level. Each 100 m line commenced and terminated with a bucket. Bisecting each bucket was a continuous vertical plastic fencing, stapled to stakes and partially buried into the ground, which acted to impede the passage of small mammals and guide them towards the buckets. The second technique consisted of two types of live traps: Sherman (22,5 x 8,6 x 7,4 cm) (H. B. Sherman Traps Inc, Tallahassee, Florida) and National (39,2 x 12,3 x 12,3 cm) (Tomahawk Live Trap, Hazelhurst, Wisconsin), the former has a tendency to capture animals of less than 50 g and the latter animals of larger body size. Briefly, traps were baited with a suitable food source (peanut butter), and placed on the ground or in trees. Due to the diurnal nature of some of the studied species in Madagascar, traps were left in situ for six consecutive nights and visited twice per day, just after dawn and in the late afternoon, when bait was removed and replenished. These types of live traps were used for sampling on La Réunion, Mayotte and Mahé, including forested and more urbanized areas and where targeted species are nocturnal. Hence, traps were installed at dusk and inspected the following morning. Bats were capturedat dusk or at night using mist nets or harp traps placed in proximity to cave entrances, across presumed flight pathways, or near water source where these animals come to drink at dusk. On several occasions, a butterfly hand net was used to collect bats inside their day roosts. Captured animals were identified based on different external and cranio-dental characters. All animals were dispatched using humane techniques identified by professional mammalogist committees (3). Dissections were performed using sterile utensils. All tissue

- 61 samples were immediately conserved in liquid nitrogen, and stored at -80°C after transport to
- the laboratory.

### **Supplementary Text 2: Animal Descriptions**

64

65 The family Tenrecidae is best considered endemic to Madagascar and composed of a 66 radiation belonging to the superorder Afrotheria, of African origin and comprising a diversity 67 of mammals (4). The Tenrecidae colonization of Madagascar and divergence from the 68 balance of Afrotheria is dated to the Middle Oligocene, about 30 Mya (5). These largely 69 animalivorous mammals are either terrestrial or arboreal and vary in body size from about 1 70 kg to a few g (6). 71 The second group is an Afro-Malagasy family of rodents, Nesomyidae of which the 72 Nesomyinae are endemic to Madagascar. The nesomyine radiation is dated to have diverged 73 during the early portion of the Miocene, about 20 Mya, and probably of African origin (5). 74 They are frugivores or granivores, either terrestrial or arboreal and vary in body mass from 1 75 kg down to about 20 g (6). 76 The last two groups involved in this study are both introduced to SWIO islands. The first 77 comprises shrews of the genus Suncus (family Soricidae). These animals can be human 78 commensals, as well as living in natural forest settings. Two species occur in the region and at least one, S. murinus, was probably introduced by Arab traders between the 11th to 14th 79 80 centuries (7, 8). The second group includes rodents (family Muridae) and represented in the 81 region by two genera, Rattus and Mus. The earliest known physical remains indicate that Rattus was first introduced to Madagascar also during the 11th to 14th centuries (9). In the 82 83 case of Mus, Malagasy populations are closely related to those from the Arabian Peninsula 84 and probably brought to the island during the Middle Ages (10). 85 In contrast to the Tenrecidae and Nesomyinae, each of which comprises a single 86 colonization by an ancestor that subsequently underwent an adaptive radiation, the bat fauna 87 of Madagascar and neighboring islands involves a considerable number of different 88 colonization events. For Madagascar alone, at least 24 different colonization events took 89 place, largely from African sources, although a few taxa appear to be Asiatic in origin (11).

- 90 The bat fauna of the island includes largely frugivorous or insectivorous species, Ranging in
- 91 body mass from nearly 400 to 4 g (12).

- 93 1. Raxworthy CJ, Nussbaum RA. 1994. A rainforest survey of amphibians, reptiles and
- 94 small mammals at Montagne d'Ambre, Madagascar. Biological Conservation 69:65 -
- 95 73.
- 96 2. Goodman SM, Ganzhom JU, Olson LE, Pidgeon M, V S. 1997. Annual variation in
- 97 species diversity and relative density of rodents and insectivores in the Parc National
- de la Montagne d'Ambre, Madagascar. Ecotropica **3:**109 118.
- 99 3. Sikes RS, Gannon WL. 2011. Guidelines of the American Society of Mammalogists
- for the use of wild mammals in research. J Mammal **92:**235-253.
- 101 4. Kuntner M, May-Collado LJ, Agnarsson I. 2010. Phylogeny and conservation
- priorities of afrotherian mammals (Afrotheria Mammalia). Zoologica Scripta **40:**1 15.
- 103 5. Poux C, Madsen O, Glos J, de Jong WW, Vences M. 2008. Molecular phylogeny
- and divergence times of Malagasy tenrecs: influence of data partitioning and taxon
- sampling on dating analyses. BMC evolutionary biology **8:**102.
- 106 6. Soarimalala V, Goodman SM. 2011. Les petits mammifères de Madagascar,
- 107 Association Vahatra, Antananarivo.
- 108 7. Hutterer R, Tranier M. 1990. The immigration of the Asian house shrew (Suncus
- murinus) into Africa and Madagascar., p. 309 319. In Peters G, Hutterer R (ed.),
- 110 Vertebrates in the tropics. Museum Alexander Koenig, Bonn.
- 111 8. **Rakotozafy LMA.** 1996. Etude de la constitution du régime alimentaire des habitants
- du site de Mahilaka du XIè au XIVè siècle à partir des produits de fouilles
- 113 archéologiques. . Doctorat de troisième cycle. Université d'Antananarivo,
- 114 Antananarivo.

115 9.	Tollenaere C, Brouat C, Duplantier J-M, Rahalison L, Rahelinirina S, Pascal M
116	Moné H, Mouahid G, Leirs H, Cosson J-F. 2010. Phylogeography of the invasive
117	species Rattus rattus in the western Indian Ocean, with special emphasis on the
118	colonization history of Madagascar. Journal of Biogeography 37:398-410.
119 10.	Duplantier JM, Orth A, Catalan J, Bonhomme F. 2002. Evidence for a
120	mitochondrial lineage originating from the Arabian peninsula in the Madagasca
121	house mouse (Mus musculus). Heredity 89:154-158.
122 11.	Samonds KE, Godfrey LR, Ali JR, Goodman SM, Vences M, Sutherland MR
123	Irwin MT, Krause DW. 2012. Spatial and temporal arrival patterns of Madagascar's
124	vertebrate fauna explained by distance, ocean currents, and ancestor type
125	Proceedings of the National Academy of Sciences of the United States of America
126	<b>109</b> :5352-5357.
127 12.	Samonds KE, Godfrey LR, Ali JR, Goodman SM, Vences M, Sutherland MR
128	Irwin MT, Krause DW. 2013. Imperfect isolation: factors and filters shaping
129	Madagascar's extant vertebrate fauna. PloS one 8:e62086.

Animal species with co-infection (identification code)	Origin	Associated paramyxovirus L-gene Accession numbers
Microgale cowani (VS 1912)	Madagascar	KF245972, KF245973
Microgale cowani (VS 1989)	Madagascar	KF245983, KF245984
Microgale cowani (VS 2211)	Madagascar	KF246048, KF246049
Microgale dobsoni (VS 2215)	Madagascar	KF246051, KF246052

1	32
1	33
1	34
1	35
1	36

. . .

**Supplementary Table S1:** Paramyxovirus co-infections. Highlighted are animals from which multiple paramyxovirus L-gene sequences could be obtained, that showed clearly divergent sequences. Multiple discrete sequences are suggestive of viral co-infections.

### 137 Host associations for the studied Paramyxoviridae

	Chiroptera	Rodentia	Scandentia/ Soricomorpha/ Afrosoricida
A. Pneumovirinae	Eidolon helvum (5): FJ971952-3, FJ971956-7, FJ971960	<b>Mus sp. (4)</b> : AY729016, AY743909-10, NC_006579	-
B. Rubulaviridae	Sturnira lilium (2): EF095490, NC_009489		
C. Unclassified Rubulavirus- Related	Eidolon helvum (6): JX051319-20, HQ660092, FJ971943, FJ809195, HQ660109 Rousettus leschenauttii (3): GU128080-2 Pteropus hypomelanus (2): AF298895, NC004074 Pteropus spp. (3): JX112711, AF326114, NC_007620 Rousettus aegypticus (3): HQ660098, HQ660100, HQ660106		
F. Respiroviridae	-	<b>Mus sp. (10)</b> : AB005795-6, AB065186- 9, AB195947-8, DQ219803, NC 001552	-
G.Henipaviridae and Unclassified Henipavirus- Related	Eidolon helvum (79): FJ609194, FJ971936-40, GQ168929, HE647821-2, HE647827, HE647829-34, HE801035-6, HQ6601027, HQ660131-5, HQ660140, HQ660142, HQ660147, HQ660150, JN862567- 70, JN862576, JN862578, JN862584-6, JN862588, JN862590, FJ809191, HE6478236, HE647828, HE647833, HQ660123-5, HQ660130, HQ660136, HQ660141, HQ660143-4, HQ660146, HQ660148-9, HQ660141, HQ660143-4, HQ660146, HQ660148-9, HQ660151, JN862562-6, JN862589, JN862591-4, FJ609198 Myonycteris torquata (2): HQ660118, HQ660137 Hypsignathus monstrosus (2): HQ660119, HQ660152 Epomorphorous sp. (4): HQ660120-2, HQ660128 Rousettus aegypticus (1): HQ660138 Pteropus vampyrus (1): FN869553 Myonycteris torquata (1): HQ660126, Rousettus aegypticus (2): HQ660119, HQ660145 Pteropus vampyrus (1): FN869553 Myonycteris torquata (1): HQ660126, Rousettus aegypticus (2): HQ660139, HQ660145 Pteropus pampili (1): JF828297	Rattus rattus (1): AB844426	Crocidura spp. (9): AB844341, AB844343-50
I. Morbilliviridae	Desdomonus rotundus (2): HQ660188-9	Rhabdomys pumilio (5): HQ660178- 180, HQ660172-3 Apodemus sp. (2): JF828302-3 Mastomys natalensis (3): AB844407, AB AB844395, AB844382	
J. Unclassified Morbillivirus- Related	Hipposideros spp. (8): HQ660153-4, HQ660156-62 Coleura afra (1): HQ660155 Triaenops menamena (21): F928232, KF928233, KF928234, KF928235, KF928236, KF928237, KF928238, KF928239, KF928240, KF928241, KF928242, KF928243, KF928244, KF928245, KF928246, KF928247, KF928248, KF928261, KF928262, JQ886096, JQ886098 Miniopterus girieaudi (9): JQ886097 Miniopterus griveaudi (9): JQ886099-103, KF928257, KF928258, KF928259, KF928260 Miniopterus sororculus (1): JQ886104 Mormopterus acetabulosus (1): JQ886105 Pipistrellus spp. (1): FJ609192, JN086951, JN086954 Hipposideros cafer (1): HQ660168-71, JN086954 Hipposideros cafer (1): HQ660187 Glossophaga soricina (1): HQ660190 Carollia spp. (4): HQ660191-4 Pteronotus parmellii (2): JF828295-6 Chaerephon leucogaster (1): KF9282256 Mops leucostigma (7): KF9282256 Mops leucostigma (7): KF928225, KF928226 Myotis goudoti (2): KF928227, KF928226 Myotis goudoti (2): KF928227, KF928228, KF928229, KF928230, KF928231, KF928221, KF928229, KF928230, KF928231, KF928255	Microtus arvalis (2): HQ660184-5 Apodemus flavicolis (3): JF828298, JF828308, HQ660183 Myodes glarious (11): HQ660181-2, JF828299-301, JF828304-7, HQ660186 Mus musculus (2): NC_007454, AY990001 Rattus norvegicus (13): DQ100461, KF245939-44, KF408257-60, AY286409, NC_005339 Rattus andamanensis (1): JN689227 Rattus rattus (32): KF245947-8, KF245950, KF245953-8, KF245960, KF246903, KF245957-8, KF245960, KF246003-4, KF246023, KF246990, KF246003-4, KF246023, KF246901, KF245081, KF245959, KF245951, KF245081, KF245959, KF245951, KF246034, KF246002, KF246011, KF245968-9, KF246005, AB844427 Rhabdomys pumilio (4): HQ660174, HQ660175-7 Mus minutoides (2): AB844424-5 Mastomys natalensis (15): AB844369, AB844376, AB844401, AB844404, AB844376, AB844398, AB844378, AB844371, AB844413, AB844396, AB844381, AB844422 Zygodontomys brevicauda (2): NC017937, FJ362497	Suncus murinus (2): KF245946, KF245952 Microgale spp. (73): KF245962, KF245966-7, KF245970-5, KF245977-88, KF245991-7, KF246006-10, KF246015-21, KF246024, KF246030-41, KF246035-40, KF246042-56, KF246058-40, KF246045-5, KF246057, KF246041, KF246060 Eliurus spp. (12): KF246011-2, KF246012, KF245998, KF246032-3, KF246013, KF245999-601, KF245989, KF246018 F246018 F246018 F246018 F245999-601, KF245989, KF246018 F245999-601, KF245989, KF246028 F245999-601, KF245999-601, KF245999-601, KF245999-601, KF245989, KF246028 F245999-601, KF245999-601, KF245999-601, KF245999-601, KF245999-601, KF245999-601, KF245999-601, KF245999-601, KF245999-601, KF245999-601, KF24599-601, KF245999-601, KF24599-601,

**Supplementary Table S2:** Genbank accession numbers are presented for paramyxovirus sequences originating from bats, rodents or other small mammal species that were of interest to this study. Sequences are grouped into the major clades of the paramyxovirus family, based on family level phylogenies presented in main text Figure 2. Host-species names and the number of sequences originating from that species are shown for each group. Sequences generated as part of this study are underlined.

0 1	6"	1		A 10'2 - 1		0.1.0.1	
Country Madagascar	Site Ambahibe	Latitude 12°58.060'S	Longitude 49°07.231′E	Altitude 100 m	Description Dry deciduous forest	Capture Dates 12 Sep. 2013	Species Captured  Miniopterus griveaudi
Madagascar	Andranoilovy	24°03'S	43°45'E	50 m	Spiny bush	21-26 Apr. 2012	Triaenops menamena
Madagascar	Androimpano	24°39.012'S	43°57.797'E	110 m	Spiny bush	23 Apr. 2012	Triaenops menamena
Madagascar	Anjohikinakina	19.0099°S	44.7677°E	130 m	Dry deciduous forest	6 Nov. 2012	Myotis goudoti Otomops madagascariensis
Madagascar	Ankarana National Park	12°55.9'S	49°03.4'E	50 m	Dry deciduous forest	15 Sep. 2012	Triaenops menamena
Madagascar	Ankazomivady	20.464948	47.09444E	1675 m	Montane forest	9-14 Dec. 2010 3-8 Nov 2011 16 Mar3 Apr 2012	Eliurus minor Rattus rattus Microgale cowani Microgale dobsoni Microgale gymnorhyncha Microgale longicaudata Microgale parvula Microgale parvula Microgale pricipula Microgale taiva Hemicenteles nigriceps Setifer setosus Tenrec ecaudatus
Madagascar	Antanandava	19.92793°S	44.60.594°E	40 m	Dry deciduous forest	2 May 2012	Mops leucostigma
Madagascar	Bekoaky	22°46'22.8"S	43°43'21.6"E	80 m	Dry deciduous forest	10 Feb. 2012	Myotis goudoti
Madagascar	Kirindy	20.06332°S	44.59679°E	50 m	Dry deciduous forest	1-2 May 2012	Vespertilionidae sp.
Madagascar	Lakato  Grotte des Makis	19.0238S	48.2055E 43°46'14.5"E	1000 m	Lower Montane Forest	23-28 Oct. 2010 19-25 Mar. 2011 6 -21 Mar. 2012 8 Feb. 2012	Eliurus minor Eliurus tanala Gymnuromys roberti Nesomys rufus Rattus rattus Rattus rattus Rattus norvegicus Microgale dobsoni Microgale jobihely Microgale longicaudata Microgale principula Microgale principula Microgale taiva Oryzorictes hova Suncus murinus Hemicentetes semispinosus Setifier setosus
Madagascar	Grotte des Makis	23 28 19.6 5	43 46 14.5 E		Dry	8 Feb. 2012	Mormopterus jugularis
Madagascar	Marofandilia	20°04.046'S	44° 39.480'E	20 m	deciduous forest	1-2 May 2012	Mops leucostigma
Madagascar	Marofototra	20.30244°S	44.39807°E	20 m	Dry deciduous forest	4 May 2012	Chaerephon leucogaster
Madagascar	Tanambao	23°32.933'S	43°46.044'E	10 m	Spiny bush Dry	10-12 Feb. 2012	Otomops madagascariensis Triaenops menamena
Madagascar	Tsimafana	19.92793°S	44.58432°E	40 m	deciduous forest	2 May 2012 9 Nov. 2012	Mops leucostigma
Madagascar	Vintany	24°42'8.5"S	43°42'08.5'E	25 m	Spiny bush	24 Apr. 2012	Triaenops menamena
Mayotte (France)	Sada	12°52.680'S	45°07.339'E	70 m	Urban	12-18 Jul. 2012	Rattus rattus
Mayotte (France)	Combani	12°47.915'S	45°09.386'E	130 m	Forest	12-18 Jul. 2012	Rattus rattus
Mayotte (France)	Tzoundzou	12°48.349'S	45°12.261'E	20 m	Urban	12-18 Jul. 2012	Rattus rattus
Mayotte (France)	Convalesence	12°46.102'S	45°11.415'E	400 m	Forest	12-18 Jul. 2012	Rattus rattus
Mayotte (France)	Mbouzi	12°48.581'S	45°14.191'E	30 m	Forest	12-18 Jul. 2012	Rattus rattus
Mayotte (France)	Mamoudzou	12°46.778'S	45°13.671'E	10 m	Urban	12-18 Jul. 2012	Rattus rattus
Mahé (Seychelles)	Anse aux Pins	4°42.216'S	55°31.188'E	10 m	Residential	4-15 Jun. 2011	Rattus rattus
Mahé (Seychelles)	Zone 21	4°40.119'S	55°30.355'E	1 m	Residential, Coastal	4-15 Jun. 2011	Rattus rattus
Mahé (Seychelles)	La Misère	4°39.992'S	55°28.225'E	425 m	Dense Residential	4-15 Jun. 2011	Rattus rattus
Mahé (Seychelles)	Station forestière de Grande Anse Helvetia Tea	4°40.588'S	55°27.263'E	30 m	Secondary Forest	4-15 Jun. 2011	Rattus rattus
Mahé (Seychelles)	Helvetia Tea Plantation	4°40.446'S	55°27.214'E	75 m	Forest	4-15 Jun. 2011	Rattus rattus
La Réunion (France)	Diverse	-	-	-	-	Sep. 2012-Mar. 2013	Rattus rattus Rattus norvegicus Suncus murinus Mus musculus


Supplementary Table S3: Description of capture sites, locations, dates and species captured.

Std Mean	A	В	с	D	E	F	G	н	1	J	100,0
Α	14.67 73.05	2.07	2.17	1.83	1.61	1.48	2.09	1.30	2.32	2.14	90,00
В	38.07	17.28 78.53	2.18	19.29	1.42	1.48	1.70	1.32	1.22	1.53	80,00
с	38.69	61.60	8.64 72.25	2.03	1.54	1.42	1.75	1.64	1.94	1.74	70,00
D	37.86	53.92	47.94	14.72 82.16	0.94	1.06	1.83	0.99	1.08	1.38	60,00
E	38,06	46.42	48.18	41.20	0.00	0.83	1.27	0.00	1.13	1.58	50,00
F	39,72	46.75	45.95	42.23	58.73	14.00 80.07	1.83	1.64	1.64	1.89	40,00
G	39.77	45.97	45.92	43.16	59.19	57.69	8.44 66.60	1.44	1.99	2.48	30,00
н	36.79	44.09	44.25	41.85	55.70	55.65	58.61	0.00	1.08	1.82	20,00
ı	39.46	44.81	44.58	42.56	58.15			57.05	14.92 79.62	1.63	10,00
J	39.82	46.15	45.28					59.60	61.66	7,25 69.46	0,00

A - Pneumovirinae, B - Rubulaviridae, C - Unclassified Rubulavirus-Related, D - Avulaviridae, E - Ferlaviridae, F - Respiroviridae G - Henipaviridae and Henipa-Related viruses, H - Salem virus, I - Morbilliviridae, J - Unclassified Morbilli-Related viruses


151

**Supplementary Figure S1:** Graphical representation of the genetic similarity (% identity) between different functional clade groups of the *Paramyxoviridae* (see Figure 2). Numbers below the diagonal are the mean identities between all individual members of the compared groups. Numbers above the diagonal are the corresponding standard deviations of sequence identities. The indicated color scale of green to red highlights strong to weak sequence conservation.




12/14

153 154 155	Supplementary Figure S2 (previous page). Phylogeny of all sequences belonging to the UMRV phylogroup. Groups are defined by majority host-species. Red stars indicate the positions of Beilong (DQ100461) and J-viruses (NC007454, AY900001) in the phylogenetic tree.
156	



158

Supplementary Figure S3. Geographical distributions of paramyxovirus-positive species. The potential extent of each group presented in Supplementary Figure S2 is calculated from the geographical distribution of its hosts, according to the IUCN database (http://www.iucnredlist.org/), and is indicated in dark red to the right. Light red is used for locations outside of Madagascar for the "Malagasy Afrotheria Group", all sequences within this phylogenetic group originated from Madagascar including the minority of sequences that were detected in *R. rattus*. There is no evidence that viruses sharing sequence homology to the Malagasy Afrotheria Group exist outside of Madagascar.



**Supplementary Figure S4.** Ancestral state reconstructions – *Henipaviruses* and *Unclassified Henipa-related Viruses*. Ancestral host-species (A) and geographical (B) states were reconstructed and used to predict switching events in Mesquite based on 4000 raw trees of the *Henipavirus* phylogeny generated in BEAST. Host switching events that are linked with disease emergences in human populations are highlighted in green.

### 4.4. Conclusion du chapitre 4

Les analyses phylogénétiques sous BEAST nous ont permis tout d'abord de classer les paramyxovirus détectés dans un nouveau genre, proches des *Morbillivirus*, mais sans y être attaché. Nous les avons nommé «*Unclassified morbilivirus-related viruses* ou *UMRVs*». Une observation intéressante est le niveau de variabilité génétique très élevé de ces *UMRVs*. Nous avons également observé un niveau très élevé de diversité génétique au sein des *URMVs* des Chiroptera et des Rodentia. Par ailleurs, la diversité génétique estimée au sein des *URMVs* des Chiroptera est beaucoup plus importante en comparaison de celui estimé au sein des *URMVs* des Rodentia et Afrosoricidae démontrant le rôle des chauves-souris en tant que réservoir des paramyxovirus.

Une autre observation originale concerne l'étendue du spectre d'hôtes notamment par la mise en évidence de ces virus chez de nouvelles espèces, incluant les chauves-souris insectivores, tels que *Otomops madagascariensis*, les *Eliurus* (*E. minor*) et les microgales (*M. dobsoni, M. principula, M. majori, M. taiva, M. cowani, M. gymnorhyncha* et *M. johibely*). De plus, une certaine spécificité d'hôte a également été démontrée.

Un phénomène de saut d'espèce par lequel deux espèces d'hôtes phylogénétiquement très éloignées (par exemple appartenant à deux ordres d'animaux différents) hébergent des *UMRVs* phylogénétiquement extrêmement proches a été démontré. Cette caractéristique a été montrée entre les 3 ordres de mammifères étudiés.

De plus, nous avons montré que des paramyxovirus génétiquement similaires sont souvent repartis à travers le monde, suggérant la dissémination de ces virus sur de grandes distances (Aplin et al., 2003 ; Lund, 1994). Le rat, plus précisément *Rattus rattus*, est l'espèce qui serait la plus impliquée dans les processus de transmission de ces nouveaux virus, probablement par l'établissement de "pont épidémiologique" entre espèces animales très différentes.

**Annexe 5**: Nos résultats semblent montrer que les *UMRVs* de Rodentia auraient pour origine l'Afrique et auraient disséminé vers le reste du monde *via Rattus rattus*. Cet article est en cours de soumission (Ghawar et al., soumis).

# Chapitre 5. Les mécanismes moléculaires et les facteurs environnementaux associés à la diversité des Paramyxovirus chez les chauves-souris de Madagascar

### 5.1. Introduction

La compréhension des relations hôtes-pathogènes au niveau évolutif, ainsi que les interactions écologiques qui facilitent le maintien du pathogène dans son biotope naturel sont importants dans la prédiction des maladies émergentes (Morse et al., 2012 ; Childs et al., 2007).

Afin d'étudier les mécanismes macro-évolutifs prédominants chez les paramyxovirus de chauves-souris, nous avons conduit une série de campagnes de collecte au sein de consortia d'habitats naturels et anthropisés à Madagascar. Vingt-deux grottes, hébergeant soit une ou plusieurs colonies différentes de chauves-souris vivant en sympatrie pour certaines d'entre elles (voire même en syntonie), 18 habitats synanthropiques, et 12 sites forestiers, distribués sur toute la grande île, représentant un total de 947 individus qui ont été testés pour la présence ou non de paramyxovirus.

Les prévalences des paramyxovirus au niveau intra- et inter-hôtes, intra- et inter-grottes ont été estimées après criblage des individus par RT-PCR. Des analyses phylogénétiques sous Mr Bayes ont été réalisées pour déterminer l'histoire évolutive de ces virus. Des tests de co-phylogénies ont été effectués sous CoRe-PA pour l'identification du (ou des) mécanisme(s) principal (aux) macro-évolutif(s) chez les paramyxovirus.

Par ailleurs, une analyse statistique sous PaRaFit a été conduite pour mettre en évidence les éventuelles associations significatives entre les hôtes et les paramyxovirus. Enfin, identifier certains facteurs biotique et abiotique pouvant jouer un rôle dans la transmission de ces virus, nous avons réalisé une analyse statistique multivariée en passant par la construction modèle linéaire généralisé binomiale (GLM) sous R.

5.2. Etude éco-épidémiologique d'une communauté de chauves-souris de
Madagascar : Analyses des processus macro-évolutifs et facteurs écologique
associés à la transmission virale.

- 1 An eco-epidemiological study of Morbilli-related paramyxovirus infection in Madagascar bats
- 2 reveals host-switching as the dominant macro-evolutionary mechanism.
- Julien Mélade<sup>1,2,3</sup>, Nicolas Wieseke<sup>4#</sup>, Beza Ramazindrazana<sup>1,2,3,5,6#</sup>, Olivier Flores<sup>7,88</sup>, Erwan
- 4 Lagadec<sup>1,2,3\$</sup>, Yann Gomard<sup>1,2,3\$</sup>, Steven M. Goodman<sup>5,9</sup>, Koussay Dellagi<sup>1,2,3</sup> & Hervé Pascalis<sup>1,2,3\*</sup>
- 5 \*\*, Contributed equally.
- 6 \* Corresponding author: Hervé Pascalis: Université de La Réunion, UMR PIMIT "Processus
- 7 Infectieux en Milieu Insulaire Tropical", INSERM U1187, CNRS 9192, IRD 249, Plateforme de
- 8 Recherche CYROI, Saint Denis, La Réunion, France. E-mail: herve.pascalis@ird.fr. Tel: +262 938835
- 9 <sup>1</sup> Centre de Recherche et de Veille sur les Maladies Emergentes dans l'Océan Indien (CRVOI),
- 10 Plateforme de Recherche CYROI, 2 rue Maxime Rivière, 97490 Sainte Clotilde, La Réunion, France.
- 11 <sup>2</sup>Université de La Réunion, UMR PIMIT "Processus Infectieux en Milieu Insulaire Tropical",
- 12 INSERM U1187, CNRS 9192, IRD 249, Plateforme de Recherche CYROI, Saint Denis, La Réunion,
- 13 France
- <sup>3</sup> Institut de Recherche pour le Développement (IRD), IRD BP 50172, 97492 Sainte-Clotilde, La
- 15 Réunion.
- 4 University of Leipzig, Department of Computer Science, Augustusplatz 10, D-04109 Leipzig,
- 17 Germany.
- 18 Sassociation Vahatra, BP 3972, Antananarivo 101, Madagascar.
- 19 <sup>6</sup> Institut Pasteur de Madagascar, BP 1274 Ambohitrakely, Antananarivo 101, Madagascar.
- <sup>7</sup>UMR C53 CIRAD, Peuplements Végétaux et Bioagresseurs en Milieu Tropical, 7 chemin de l'IRAT,
- 21 97410 St Pierre, France.
- 22 <sup>8</sup> Université de La Réunion, 15 Avenue René Cassin, 97400 Saint-Denis, France.
- 23 Field Museum of Natural History, 1400 S. Lake Shore Dr, Chicago, IL 60605-2496, USA.
- 24 Correspondence and requests for materials should be addressed to H.P. (email: herve.pascalis@ird.fr).
- 26 Abstract

- 27 An eco-epidemiological investigation was carried out on Madagascar bat communities to better
- 28 understand the evolutionary mechanisms and environmental factors that affect virus transmission
- 29 among bat species in closely related members of the genus Morbillivirus, currently referred to as
- 30 Unclassified Morbilli-related paramyxoviruses (UMRVs). A total of 947 bats were investigated
- 31 originating from 52 capture sites (22 caves, 18 buildings, and 12 outdoor sites) distributed over
- 32 different bioclimatic zones of the island. Using RT-PCR targeting the L-polymerase gene of the
- 33 Paramyxoviridae family, we found that 10.5% of sampled bats were infected, representing six out of
- 34 seven families and 15 out of 31 species analyzed. Univariate analysis indicates that both abiotic and

biotic factors may promote viral infection. Using generalized linear modeling of *UMRV* infection overlaid on biotic and abiotic variables, we demonstrate that sympatric occurrence of bats is a major factor for virus transmission. Phylogenetic analyses revealed that all paramyxoviruses infecting Malagasy bats are *UMRVs* and showed little host specificity. Analyses using the maximum parsimony reconciliation tool CoRe-PA, indicate that host-switching, rather than co-speciation, is the dominant macro-evolutionary mechanism of *UMRVs* among Malagasy bats.

### **Introduction**

The transgression of the species barrier by pathogens moving from their natural host reservoir to infect a new host species (also referred to as host-switching, host-jumping or host-shifting), may induce an abortive infection in the few infected individuals of the new host, or trigger a short lived outbreak, or an epidemic<sup>1,2</sup>. Co-speciation and host-switching are the two main evolutionary mechanisms generating genetic diversity among micro-organisms. Both are long-term dynamic processes<sup>3</sup>, in contrast to co-evolution *sensu stricto*, which continuously acts on a short-time scale<sup>4</sup>. Co-speciation, the simultaneous speciation of the host and their parasites<sup>3, 5,7</sup>, was considered for many years as the principal macro-evolutionary process generating viral diversity<sup>8,11</sup>. As convincing examples of co-speciation are rare, this mechanism has probably been overestimated. Host-switching refers to a new host-parasite combination that results from the shift of the parasite to a new host and its subsequent specialization, for example, under environmental selection pressure<sup>12</sup>. Colonization by a parasite of a phylogenetically closely related host species, often of the same genus or family, has proven to be the typical macro-evolutionary mechanism for RNA viruses<sup>13</sup>. An excellent example is the evolutionary history of *Hantaviruses* and *Arenaviruses*<sup>14,15</sup>, mostly shaped by multiple host-jumps, followed by adaptive processes within the new host, as demonstrated in bats and other operative host species

The often gregarious roosting behavior of bats and an assortment of different ecological parameters (e.g., climate, season, and migration) are important factors that can shape viral transmission dynamics, which subsequently act upon evolutionary processes<sup>10,11,13,18</sup>. Deciphering such mechanisms helps to understand how a virus hosted in wild animals can emerge as a pathogen in human populations<sup>19</sup>. For example, host-switching of *Ebola virus*, *SARS Coronavirus* or *Nipah virus* have led to major pandemics or epidemics in humans<sup>2,20,21</sup>.

Paramyxoviridae is a large and diverse viral family (Order: Mononegavirales) composed of single-stranded negative RNA viruses<sup>22</sup>. Newly recognized paramyxoviruses (PVs), named Unclassified Morbilli-Related Viruses (UMRVs), have recently been shown to infect small mammals around the world<sup>23</sup>, such as bats and terrestrial small mammals from the southwestern Indian Ocean (SWIO) islands<sup>17,24</sup>, including the biodiversity hotspot of Madagascar<sup>25</sup>. The island is divided into several unique bioclimatic zones, characterized by different meteorological regimes overlaid on elevation and underlying geology<sup>26</sup>, which in turn give rise to distinct vegetation types and highly endemic biotic communities.

After rodents, bats (order Chiroptera), constitute the most abundant, diversified, and geographically wide spread group of mammals in the world<sup>27</sup>. Genetic and fossil studies have estimated the basal split of placental mammals in the superorder Laurasiatheria from their ancestors at approximately 80-90 million years ago (Mya) and a diversification of bat families at approximately 62 Mya<sup>28</sup>. The Chiroptera of Madagascar are placed in eight different families and currently 45 species recognized, of which 36 species (80%) are endemic<sup>29-31</sup>; it is assumed that most originated from Africa. In certain cases, phylogenetic analyses provide evidence for recent periods of diversification. For example, Malagasy *Miniopterus*, a notably speciose genus, colonized the island from an African source population approximately between 4.5 and 2.5 Mya, followed by a second phase between 2.5 and 1 Mya<sup>32</sup>.

An important characteristic of Malagasy bat communities is that species co-occupy day roost sites in caves, buildings or tree cavities (often in forests) in different species combinations and varying numbers. Furthermore, certain bat species may have indirect contact with other wild, introduced or domestic animals, including synanthropic small mammals, which may imply contamination of shared common water sources or fruits by bat urine/saliva<sup>29</sup>. Considering the notable species diversity and high levels of endemism of Malagasy bats, as well as varying community structure and ecological conditions in which they occur, Madagascar provides an excellent context to study virus transmission in these animals. Herein, we examine the factors involved in interspecific transmission of *UMRVs* and try to unravel the macro-evolutionary mechanisms underlying genetic diversification in these viruses.

#### Results

91

92 In total, 947 bats (867 insectivorous and 80 frugivorous), representing seven different families and 31 species, were collected at 52 sites in all six provinces of Madagascar: Antananarivo (n = 44 bats), 93 Antsiranana (n = 125), Fianarantsoa (n = 178), Mahajanga (n = 207), Toamasina (n = 37), and Toliara 94 (n = 356). The sampling sites included 22 different caves (n = 480 bats), 18 buildings (n = 290), and 95 12 different forested areas (n = 177). Thirty-one sites (n = 664 bats) contained at least two species and 96 97 21 sites (n = 283) were monospecific. The sampling sites were in different elevational zones, ranging from low (0 to 800 m, n = 40 sites), mid (801 to 1000 m, n = 6), and high (over 1000 m, n = 6), with 98 761, 101, and 85 bats collected in each zone, respectively. Seventeen sites were sampled in dry (n = 99 100 384 bats), 22 in sub-arid (n = 382), 11 in sub-humid (n = 144), and two in humid (n = 37) bioclimatic 101 zones. Twenty-two sites (n = 377 bats) were visited during the summer (warm, wet) season and 30 102 sites (n = 570) during the winter (cool, dry) season. Ninety-nine of 947 bats (10.5%) tested positive for PVs by RT-PCR, giving a global 103 infection rate of 11.1% in insectivorous bats and 3.8% in frugivorous bats (df (degrees of freedom) = 104 1; n = 947;  $\chi^2 P = 0.02$ ). The infection rates varied according to province, from 4.5% in Antananarivo 105 106 to 15.2% in Antsiranana (df = 5; n = 947;  $\chi^2$ , P = 0.01). The infection rates of PVs for bats living in caves, buildings, and forests were 12.9%, 7.9%, and 7.9%, respectively (df = 2; n = 947  $\chi^2$ , P = 0.041). 107 The fraction of sites hosting PV positive bats among the 31 multispecies sites and the 21 monospecific 108 sites were 70.9% and 61.9%, respectively (df = 1; n = 947;  $\chi^2$ , P > 0.05). The infection rates for PV 109 were 11.4% in multispecies sites and 8.1% in monospecific sites (df = 1; n = 947;  $\chi^2$ , P > 0.05). 110 111 Infection rates at individual sites varied from 2.0% at ANJHB to 38.1% at VINT with no PV positive 112 bat at 17 sites (n = 121) (see Figure 1 for identification of sites and associated acronyms). At low, middle, and high elevation, the fraction of sites hosting PV positive bats was 67.5%, 83.3%, and 113 50.0% (df = 2; n = 947;  $\chi^2$ , P > 0.05), respectively, and the mean positive rates were 11.4%, 8.9%, and 114 3.5%, respectively (df = 2; n = 947;  $\chi^2$ , P > 0.05). In the humid, sub-humid, sub-arid, and dry 115 116 bioclimatic zones, the percentages of sites hosting PV positive bats were 50.0%, 54.5%, 72.7%, and 70.6%, respectively (df = 2; n = 947;  $\chi^2$ , P > 0.05) and the mean positive rates were 5.4%, 6.3%, 117 12.0%, and 10.9%, respectively (df = 2; n = 947;  $\chi^2$ , P > 0.05). PV positive rates were 7.9% and 12.1% 118

for bats captured during the summer and winter seasons, respectively (df = 1; n = 947; χ², P = 0.038).
Sites with UMRVs detection rates higher than 20.0% are indicated on Figure 1.

Six of seven sampled bat families yielded PV positive individuals, with the exception being Hipposideridae, for which the only Malagasy species is *Hipposideros commersoni* (Table 1). The highest PV detection rate was in the family Rhinonycteridae (39.3%) and the lowest in the family Pteropodidae (3.8%) (df = 6; n = 947;  $\chi^2$ , P < 0.001). Half of the sampled species (16/32) contained PV positive individuals. The highest PV infection rate was in *Triaenops menamena* (n = 21/42; 50.0%) and the lowest in *Miniopterus mahafaliensis* (4/89; 4.5%) (df = 31; n = 947;  $\chi^2$ , P < 0.0001). Insectivorous species had significantly higher detection rates (96/867; 11.1%) than frugivorous species (3/80; 3.8%) (df = 1; n = 947;  $\chi^2$ , P = 0.02). No significant difference was found associated with sex and age classes, regardless of diet, habitat or site ( $\chi^2$ , P > 0.05).

Model construction procedure lead to a binomial Generalized Linear Model (GLM) explaining individual infection based on seven different effects (Table 2). Among abiotic factors, Mean Annual Temperature (MAT) had an overall effect where, Mean Annual Rainfall (MAR) did not show any overall relationship with infection. However, relationships between rainfall and infection appeared different across multi- *versus* single-species sites with a quadratic effect observed for MAR. Habitat type and the multispecies characteristics did not show any significant effects, but showed marginal interaction. The multispecies sites show higher infection rates, compared to monospecific sites, for caves compared to buildings and forest capture sites (Figure 2), reinforcing the important role of multispecies bat environments on PV infection. Diet was also associated with viral infection (Table 2), with higher infection among insectivorous bat species, whereas, age and sex did not show any significant relationships. Generalized Linear Mixed Model (GLMM) with species, locality, and province as random factors were tested separately and did not improve the fit, but models with family, species and locality failed to converge due to numerical issues in model estimation.

We conducted a Bayesian analysis on the PV sequences generated from positive Malagasy bats together with PV GenBank sequences from Madagascar and elsewhere in the world. All new PV sequences presented in this study were identified as *UMRVs*, as they appeared more closely related to Morbilliviruses<sup>17</sup> (Supplementary Figure S1), than to any other genera of the family *Paramyxoviridae* 

family. The UMRVs were characterized by a high level of genetic variability and nucleotide sequences varied from 62.0 to 100% sequence identity. Only two sequence pairs of the 99 that tested positive were identical. Although URMVs showed weak exclusivity to their bat host species, two phylogenetic patterns were identified: (i) closely related UMRV sequences were hosted by bat species and families that are phylogenetically closely related, particularly those occupying day roost sites in the same caves i.e., Miniopterus griveaudi and Myotis goudoti at AMBB; Miniopterus gleni and Miniopterus sororculus at BEK; (highlighted in blue in Figure 3). This feature suggests that host-switching events might be favored by physical proximity between phylogenetically closely related bat taxa. (ii) some degree of host-specificity for URMVs was found, with individuals of one host species having closely related UMRVs, independent of other individuals occurring at the same roost site, (i.e., Triaenops menamena at VINT TSP, and ANDRF2) highlighted in green, or distant sites, (i.e.,

147

148

149

150

151

152

153

154 155

156

157

158

159

160

161

162

164

165 166

167

168

169

170

171

172

173

174

Triaenops menamena at VINT and TSP), highlighted in red on Figure 3. In some cases, a correlation was observed between the distance separating capture sites and the degree of nucleotide sequence similarity of the infecting PVs across sites. More specifically, conspecifics living on distant sites host UMRVs that display lower level of nucleotide similarity than

those infecting bats at sites in closer geographical proximity (i.e., Triaenops menamena at VINT and TANA, Miniopterus griveaudi at AMBB and ANJHK1, and Miniopterus griveaudi at ANDFR and 163

AMBB) highlighted in yellow in Figure 3. This suggests that increasing geographical distance favors

virus genetic differentiation and/or low levels of virus migration between bat roosting sites.

Using CoRe-PA, we performed a consensus phylogram for both viruses and bats, presented along with their tanglegram depicting bat-virus associations (Figure 4A-B). By evaluating 5000 random cost schemes, CoRe-PA computed the most parsimonious reconstruction and predicted the frequencies for co-evolutionary events, including co-speciation, host-switching, duplication, and sorting. For the generated 24 OTUs subset (Table 3a), the best quality value obtained was 0.256 for a solution with eight co-speciation events, 21 duplications, 51 sortings, and 19 host-switches. For the 39 OTUs subset, CoRe-PA produced 57 reconstructions (Table 3b), with a quality value of 0.25 and five co-speciation events, 33 duplications, 57 sorting, and 24 host-switches. Hence, for both sets, no clear evidence of co-speciation between UMRVs and bat species was found. The statistical analysis suggests

fewer co-speciation events in the data set than expected by chance (99.0% of randomized data sets showed more than eight co-speciation events) but more host-switching events than expected (100% of randomized data sets showed less than 19 host-switching events) (Figure 5A-B). Thus, notwithstanding the numerous identified duplication and sorting events, host-switching events appear to be the predominant aspect in the evolutionary history in *UMRVs* identified from Malagasy bats, as compared to co-speciation.

We quantified the degree of congruence between bats and UMRVs topologies, and the potential individual associations for each of the two OTU subsets. The hypothesis associated with independent speciation events could not be rejected by ParaFit (ParaFitGlobal = 38.62571; P = 0.067), for the 24 OTUs subset, whereas a significant overall pattern of association (ParaFitGlobal = 46.158; P = 0.002) was calculated for the 39 OTUs subset. Eight of 50 (16.0%) individual host-virus links were significant, based on a P < 0.05 for the 90.0% threshold, and 19 out of 60 (31.7%) for the 98% threshold. Tables S3a and S3b summarize the different associations of UMRVs with their respective hosts and the corresponding P-values for the two OTU subsets. Among the different bat species, Triaenops menamena was the most coupled species for both OTUs subsets, and Miniopterus mahafaliensis for the 39 OTUs subset. Depending on nucleotide identity, we observed a discrepancy of the global association signal, which is related to specificity increasing genetic variability by increasing the number of clades (i. e., increasing the nucleotide acid identity between sequences).

### Discussion

The overall *UMRV* infection rate in Malagasy bats was 10.5%, we also found that in some cases, that certain bat families or species showed higher PV detection rates:. Four bat species had particularly high *UMRV* infection rates: *Triaenops menamena*, *Mops leucostigma*, *Miniopterus griveaudi*, and *Miniopterus gleni*. Except for the latter taxon, all were living at sites where substantial virus circulation was recorded (Figure 1). Whether these species have higher susceptibility to PV infection cannot be discerned based on current data.

Moreover, statistical modeling demonstrated that environments supporting multiple species are positively associated to viral transmission, with a marginal effect of natural habitats (caves) being

more prone to PV infection, whereas habitat type alone was not a significant predictor of infection. As previously reported, the spread of viruses between bat species is promoted by sympatric conditions, specifically multispecies day roost sites 10. Other studies on bat rabies transmission demonstrated the importance of sympatric occurrence for viral infection 16. The high detection rate in multispecies sites likely results from greater species diversity in caves<sup>29</sup>, inducing a proximity effect between individuals, which has been previously shown to promote virus transmission. Further work correlating rates of infection in caves as a function of bat density would help support this hypothesis; however, because of seasonal variation in bat density associated with population cycling and possible dispersal movements, this aspect will be difficult to document based on field studies. Certain climatic factors also seem to promote viral transmission: the probability of PV infection increased at localities with higher mean annual temperature, which favors infection compared to cooler regions. This result supports the importance of warmer temperature on viral transmission 18. Whereas PV infection showed no overall relationship with rainfall, average rainfall conditions favored PV infection in multispecies sites, compared to drier/wetter conditions and to single-species sites. Further analyses need to be conducted to have a greater understanding of the role of climatic factors on infection. Finally, we also note that circulation of UMRVs seems to be much more active among insectivorous than frugivorous bat species, with only 3.8% of the latter tested positive. These results confirm previous studies conducted on SWIO islands 17,24.

203

204

205

206

207208

209

210

211

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

The bat-associated  $\mathit{UMRV}$  phylogeny underlines several points, particularly among the four taxa with the highest infection rates:

(i) bats collected sympatrically or some cases syntopically in the same day roost sites, for example, Miniopterus griveaudi and Myotis goudoti, belong to the families Miniopteridae and Vespertilionidae, respectively, host closely related viruses, suggesting that host-switching events occurred between these species/families;

(ii) bat/virus co-phylogenies, suggest that co-speciation cannot explain the observed patterns. Host-switching is the predominant macro-evolutionary process. In either case, numerous reciprocal selection pressures that act over the short-term scale, such as the *sensu stricto* co-evolutionary process, also drive host-virus interactions. Indeed, micro-evolutionary aspects, including those implying

selection, drift, and dispersal, result in intraspecific co-divergence of viruses<sup>33-36</sup>. Using CoRe-PA, we 231 232 highlight the lack of congruence between bat and UMRV phylogenies. In previous studies it has been 233 shown that a large number of phylogenies, set at the family level, including the Paramyxoviridae, are driven by this mode of macro-evolution<sup>13</sup>. Moreover, a considerable number of spillover events have 234 been reported between rodents and bats<sup>17,23</sup>. Our phylogenetic analyses show that the same *UMRVs* 235 infect different bat species or families, leading to the observed phylogenetic incongruence. This aspect 236 237 can, at least in part be explained by the extremely rapid evolution of some RNA viruses, which as a consequence of their higher mutation rate<sup>37</sup> generate large quasi-species virus populations, allowing 238 for greater chances after a host-jump to adapt to a new host or, in other words, to promote a better 239 adapted variant that can be sustained in the new host<sup>38</sup>. Examples of such macro-evolutionary 240 241 processes, driven by host-switching, have been reported for Puumala virus and a Hantavirus detected in bats from northern Europe and for which no evidence of co-divergence was observed<sup>39</sup>. This 242 243 scenario has also been cited for other Hantaviruses and is probably a general rule for this viral family 14,40; 244 (iii) a viral allopatric process, in which a virus speciates within a host species living in different 245 geographical areas, and giving rise to independent evolution 12,13. This may have occurred for 246 247 Triaenops menamena, Mops leucostigma, and Miniopterus griveaudi; these three taxa have relatively broad distributions on Madagascar<sup>26</sup>. Interestingly, we could observe 7 major but phylogenetically 248 distant viral clades with 5 or more closely related viruses detected in different bat species or families. 249 250 This observation may suggest the circulation of 7 major UMRVs strain across Madagascar infecting a 251 large host range. The CoRe-PA analysis indicates for these three species, 33 events of duplication. 252 Duplication is a virus speciation event that occurs within the same host. This phenomenon can be the 253 consequence of events that affect only the host, i.e., adaptive co-evolution, such as environmental 254 adaptation. Such duplication events can be, for example, an immune pressure selection or virus specialization related to adaptation to different organs of the host. We also disproportionately found 255 256 numerous duplication events in our analysis (21 duplications for the 24 OTUs and 33 duplications for the 39 OTUs subsets). This was anticipated, as CoRe-PA tends to place too many nodes from the virus 257 258 tree near the root of the host tree. Consequently, whenever two descendant parasites (i.e., parasites that

emerged straight from the same ancestral parasite) share the same host, a duplication event is predicted.. One explanation could be associated with the capability of a virus within the host species to replicate independently. Different species of Miniopterus can be found roosting in strict syntopy and this close physical contact between related bat taxa may facilitate host-switching followed by mutation and duplication within the host. These sorting events might have multiple evolutionary causes and several hypotheses can explain these observations: (a) an ancient co-speciation event between the ancestor of the host and the virus, but the viral descendants subsequently went extinct; (b) an unidirectional and irreversible host-jump of the ancestral virus from one host to another; (c) no hostvirus association ever existed between the virus and the respective sibling host; (iv) as indicated by ParaFit Global test, significant associations were observed with the OTUs subsets (Tables S5a and S5b). We found significant linkage associations between species such as Triagnops menamena and both OTUs subsets. These results matched with our virus phylogeny analysis, which indicate some host-specificity, in particular for Triaenops menamena. We hypothesize that this association occurred after multiple host-switching events at some point in the past (macro-evolution) and continued in the form of a co-evolutionary adaptation (micro-evolution) inside the new host. Such phenomenon has also been reported in different Coronaviruses<sup>41</sup>, for which co-evolution with bats seems to be the predominant evolutionary process. Since phylogenetically distant bat families hosted closely related UMRVs, the genetic distance between different groups of bats does not seem to be a major constraint for host-switching. Such results are important since host and virus traits determine the ability for a virus to infect a new host and host-switching events should a priori occur between phylogenetically closely related bat species on Madagascar. Besides, the occurrence of multiple interspecies transmissions, even to genetically distant host species, could be promoted by the existence of ubiquitous or alternative receptors for the virus 42,43. It has been shown that genetic distances between bat species are a key factor for host-switching events 13,16,44. However, our data also indicate that genetically closely related UMRVs infecting different bat species, sometimes occurring in geographically distant areas, may suggest the intervention of a probable vector, capable to connect these different populations<sup>1</sup>. Furthermore, except for regular bat foraging or dispersal movements, the black rat, Rattus rattus, introduced to Madagascar, has been identified as a significant reservoir of

259

260

261262

263264

265

266

267

268269

270

271

272

273274

275

276

277

278

279

280

281

282

283

284

*UMRVs*<sup>17</sup>. This rodent might be the ideal candidate to play this spreading role and establishing epidemiological bridges between different species.

### Methods

**Fieldwork and sampling.** This study used samples collected in the context of a long-term project to document the land vertebrates of Madagascar based on voucher specimens and for a variety of studies<sup>26</sup>. From February 2012 to March 2013, bats were captured in the six different provinces of Madagascar using harp traps, hand nets, and mist nets. Some *Pteropus* fruit bats were purchased in markets. Individual bats were identified to species using external and cranio-dental characters and comparison to museum specimens. For each animal, different parameters, including age, sex, and reproductive condition<sup>45</sup> were recorded and this information deposited in DRYAD (http://www.datadryad.org/).

Bat tissue samples were collected in the field and immediately stored in liquid nitrogen, then transferred to -80°C storage upon arrival in the laboratory. The geographic ranges of the captured bat species were variable, with some having broad distributions nearly across the complete island and others distinctly more restricted. Several species, especially insectivorous bats, occur sympatrically in the same cave systems and in the same forest blocks, or synanthropically in human-built structures. Information on the species, province and specific collection locality, sampling season, geographic coordinates, elevation, habitat type, and the number of bat species found at each site and the associated species composition are presented in Tables S1 and S2. Mean climatic conditions of the sampling sites were extracted from the WorldClim database (http://www.worldclim.org/). We used the resolution proposed by WorldClim as 30 arc seconds (1km). An open-source GIS software, QGIS<sup>46</sup>, was used to generate the map for visualizing Madagascar bioclimatic regions proposed by Cornet<sup>47</sup>.

Ethics statement. Animals used in this study were manipulated in strict accordance with the guidelines for the handling of wild mammals<sup>48</sup>. All protocols strictly followed the terms of research permits and regulations and were approved by licencing authorities: Direction du Système des Aires Protégées et Direction Générale de l'Environnement et des Forêts and Madagascar National Parks

314	under	different	research	pen	mits	(n°194/1	2/MEF/S	G/DGF/DCB.	SAP/SCB,
315	067/12/MEF/	SG/DGF/DCB.	SAP/SCBSE,	and	032/12/N	IEF/SG/D	GF/DCB.	SAP/SCBSE).	Animals
316	were capture	d, manipulated,	and dispatched	l with	thoracic	compressi	on follow	ing procedure	s accepted
317	by the scienti	fic community:	for the handlin	g of v	vild mamı	mals <sup>48</sup> . <i>Pte</i>	ropus wei	re purchased i	n a market
318	and were not	physically coll	ected by the r	esear	ch team ii	n a natural	setting. I	Euthanasia wa	s used for
319	Pteropus and	l not any other	r bat genera.	All f	ieldwork	conducted	on Mad	agascar was	before the
320	creation and	implementation	of an institution	onal a	and/or lice	ensing con	ımittee on	the island to	issue such
321	clearances. A	CITES permi	t from the Ma	alagas	sy nationa	authorit	y was iss	ued for Ptero	pus tissue
322	export (n°243	3C-EA06/MG12	e) to CRVOI or	ı La F	Réunion.				
323	Statistical p	rocedures. Ex	ploratory anal	yses	were per	rformed u	sing Pear	son chi-squa	re $(\chi^2)$ or
324	Fisher's exac	t tests in R sof	tware <sup>49</sup> (95%	confi	dence into	ervals with	n continui	ty correction)	. With the
325	intent of ider	ntifying variable	es potentially o	orrel	ated with	UMRV in	fection, w	e performed	a binomial
326	GLM analys	is <sup>50</sup> . We first	visually inspe	ected	the relat	ionships 1	etween v	variables (me	an annual
327	temperature [	MAT], mean ar	mual rainfall [	MAR	.], habitat	type), and	"multi" a	binary factor	indicating
328	whether a giv	ven site contain	ed multiple (>	2) 01	r one spec	cies of bat	Graphic	inspection su	ggested an
329	overall effect	of MAT and a	no effect of M	AR.	However,	relationsh	ips sugge	sted a linear	interaction
330	between MA	R and habitat ty	pes, and a poss	sible (	quadratic	effect of r	ninfall wit	hin multi- <i>ver</i>	sus single-
331	species sites.	Main and inter	action effects	were	tested se	parately w	hile acco	unting for the	effects of
332	other variable	es. We retained	the best model	acco	rding to A	kaike Info	rmation (	Criterion (AIC	). We then
333	tested the effe	ects of biotic va	riables (sex, di	et, an	ıd age), or	our best	nodel, to	determine a si	gnificance
334	effect while a	accounting for t	he effects of a	biotic	factors.	GLMM <sup>51</sup>	were cons	tructed with u	ınbalanced
335	variables (i.e.	., province, loca	ilities, and spe	cies -	related to	o non-hom	ogenous	sampling) set	as random
336	factors in or	der to be comp	ared to the b	est G	LM fit. S	Statistical	analyses	were conduct	ed with R
337	software pack	kage <sup>49</sup> .							
338	Sample scree	ening. Approxii	nately 1 mm³	of lun	ıg, kidney	, and sple	en collecte	ed from the sa	me animal
339	were pooled	in DMEM medi	ium and homo	geniz	ed in a Ti	ssueLyser	II (Qiage	n, Hilden, Ger	rmany) for
340	2 min at 25	Hz using 3 mr	n tungsten bea	ads.	Total nucl	leic acids	were extr	acted from th	ne mixture

341	supernatant using the viral mini kit $v2.0$ and an EZ1 BioRobot (Qiagen). cDNA products were
342	generated via reverse transcription (cDNA kit, Promega, Madison, Wisconsin, USA). PVs were
343	detected by a semi-nested PCR targeting part of the L-gene polymerase gene, designed as to detect
344	$\textit{Respirovirus}, \textit{Morbillivirus}, \text{ and } \textit{Henipavirus} \text{ (RMH)}^{52}. \text{ The 400-600 bp PCR amplified cDNAs were}$
345	$purified \ using \ the \ Qiagen \ PCR \ purification \ kit \ and \ cloned \ into \ the \ pGEM-T \ vector \ system \ (Promega).$
346	Cloned PCR products were sequenced by the Sanger method (Big Dye sequencing kit, ABI,
347	Genoscreen, Lille, France) using M13 standard sequencing primers.
348	Bioinformatics analysis. The sequences were first compared to the published sequences from the
349	${\it Paramyxoviridae} \ \ {\it and published} \ \ {\it UMRVs} \ \ {\it in GenBank} \ \ ({\it National Center for Biotechnology Information},$
350	Bethesda, Maryland, USA) online (www.ncbi.nih.gov) using BLASTn and BLASTx. The sequence
351	quality of individual reads was assessed, and all sequences were processed using the Geneious Pro
352	software package $v7.1.8^{53}$ . DNA sequences obtained from at least three independent bacterial clones
353	were aligned to correct for most sequencing or PCR introduced errors. M13 Primer sequences were
354	trimmed from the finalized sequences. The resulting partial sequences (~490 bp) of the L-gene
355	$polymerase\ gene\ were\ then\ aligned\ with\ Translation\ Alignment\ using\ the\ default\ ClustalW\ cost\ matrix$
356	in Geneious Pro software package. PVs sequences from bats reported in a previous study by
357	Wilkinson et al. $^{17}$ were used for phylogenetic analysis (GenBank number to KF928225 to KF928256
358	and to K928261 to KF928265). PVs and bat Cytochrome b (Cyt-b) sequences used for the present
359	analyses were deposited in GenBank under the reference numbers in respectively Table S3 and Table
360	S4. Information concerning amplified PV sequences is given in Table S3. In order to classify the
361	detected new paramyxoviruses, viral family-level phylogenetic analyses were performed. A total of
362	209 partial L-gene paramyxovirus sequences collected in Genbank were used. Sequences were
363	trimmed to remove any free end gaps or were entirely removed if the obtained alignment did not
364	provide at least 462 bp of non-gap overlap. Internal gaps were permitted. The tree was performed in
365	5,000,000 iterations in MrBayes with the GTR + G + I evolutionary model and a $10%$ burn-in rooted
366	with an Aquaparamyxovirus sequence (GenBank number EF646380). Genbank accession numbers
367	used for each virus genera are indicated in Table S6.

A best-fit nucleotide substitution model of the alignment was determined using jModelTest<sup>54</sup> with the Corrected Akaike Information Criterion (AICc)<sup>55</sup>, and the most appropriate one for *URMVs* from Malagasy bats was GTR + G. Phylogenetic trees were constructed using MrBayes v3.2<sup>56</sup> employing a Bayesian Markov Chain Monte Carlo (MCMC) method, rooted with a *Mumps* virus sequence (GenBank number AY309060). A minimum of two independent runs were made, with four chains in each run, for a total of 50 000 000, sampling every 5000 generations. The first 5000 trees burn-in were discarded. The obtained effective sample size values (ESS) for each parameter were all superior to 200. Trees obtained after the convergence point were summarized and visualized by FigTree 1.4.2 (http://treebioedacuk/software/figtree).

Available full-length Cyt-b gene sequences corresponding to each bat species that were positive for *UMRV* infection were downloaded from GenBank. When Cyt-b sequences were not available for a given bat species, PCR using primers targeting the Cyt-b gene were performed<sup>57</sup> to generate ~1140 bp sequences. All bat Cyt-b sequences were aligned, the GTR+I+G model was also the most appropriate substitution model, and the phylogenetic relationship among bat species were analyzed using RAxML 7.2.8 Geneious plugin<sup>53</sup> using 1000 generations. Two subsets of operational taxonomic units (OTUs) were defined using Mothur<sup>58</sup>, and based on two genetic distance cutoffs (90.0% and 98.0%), generated 24 and 39 representative sequences, respectively.

To study the history of co-evolution of *UMRVs* with respect to their associated bat hosts, we performed event-based co-phylogenetic reconciliation, using the tool CoRe-PA<sup>59</sup>. CoRe-PA is an event-based maximum parsimony method, which attempts to construct the most parsimonious co-evolutionary history of hosts and associated parasites. A cost is assigned to each type of co-evolutionary event (co-speciation, host-switching, duplication, and sorting) and then, the parasite phylogeny is mapped onto the host phylogeny, while trying to minimize the total costs of all occurring events. In contrast to many other co-evolutionary software packages, CoRe-PA does not require an *a priori* assignment of a cost scheme. It has been shown that the results of such analyses strongly depend on the designed cost scheme, and choosing a biologically meaningful cost scheme in an *a priori mammer* may be difficult<sup>59</sup>. CoRe-PA can assess several random cost schemes and evaluate these, based on the best fit with respect to the resulting reconstructions. In our study, we performed

396	reconstruction between the phylogenetic trees of $\mathit{UMRVs}$ and bats, using 5000 random cost schemes.
397	Furthermore, to test statistical significance, we computed the reconstructions of 100 random data sets,
398	considering the same phylogenetic trees for bats and UMRVs with different bat and virus associations.
399	In this case, the formulated null hypothesis is that there are more co-speciation and less host-switching
400	events in the data set, than compared to data sets with random host-parasite associations.
401	We quantified the degree of congruence between bat and UMRVs topologies, and the
402	potential individual associations leading to a potential co-phylogenetic structure using a global-fit
403	$method, ParaFit^{60}. \ The \ latter \ program \ tests \ the \ independence \ of \ host \ and \ symbiont \ genetic \ or \ patristic$
404	distances, and specifically herein, tests the hypothesis of evolution independence between bats and

405 *UMRVs*, i.e., one partner randomly evolving with respect to the other. Statistical analyses were done using the R software package<sup>49</sup>.

407 408

### Supplementary information

- 409 Table S1. Description of sampled sites in Madagascar.
- 410 Table S2. Number of species samples in each site.
- 411 Table S3. UMRV sequences data set.
- 412 Table S4. Bats cytochrome b sequences data set
- Table S5. Test of host-parasite co-evolution using global fit methods ParaFit for the (a) 24 OTUs and
- 414 (b) 39 OTUs.
- 415 Table S6. Paramyxovirus sequences data set.
- 416 Supplementary Figure S1. Phylogeny of all sequences belonging to the *UMRV* phylogroup.

417

### 418 References

- 419 1 Woolhouse, M. E., Haydon, D. T. & Antia, R. Emerging pathogens: the epidemiology and 420 evolution of species jumps. *Trends Ecol. Evol.* **20**, 238-244, doi:10.1016/j.tree.2005.02.009 421 (2005).
- 422 2 Parrish, C. R. et al. Cross-species virus transmission and the emergence of new epidemic 423 diseases. *Microbiol. Mol. Biol. Rev.* 72, 457-470, doi:10.1128/MMBR.00004-08 (2008).
- 424 3 Page, R. D. M. Tangled Trees. Phylogeny, cospeciation and coevolution. *Ethol*; **110**, 577-578, doi:10.1111/j.1439-0310.2004.00990.x (2003).

- 426 4 Woolhouse, M. E. Population biology of emerging and re-emerging pathogens. <u>Trends</u>
   427 *Microbiol.* 10, S3-7 (2002).
- 428 5 Burland, T. M. & Wilmer, J. W. Seeing in the dark: molecular approaches to the study of bat 429 populations. *Biol. Rev. Camb. Philos. Soc.* **76**, 389-409 (2001).
- 430 6 Johnson, K. P., Adams, R. J., Page, R. D. & Clayton, D. H. When do parasites fail to speciate in response to host speciation? *Syst. Biol.* **52**, 37-47 (2003).
- 432 7 Charleston, M. A. & Perkins, S. L. Traversing the tangle: algorithms and applications for cophylogenetic studies. *J. Biomed. Inform.* **39**, 62-71, doi:10.1016/j.jbi.2005.08.006 (2006).
- 434 8 Turmelle, A. S. & Olival, K. J. Correlates of viral richness in bats (order Chiroptera). *Ecohealth* 435 **6**, 522-539, doi:10.1007/s10393-009-0263-8 (2009).
- 436 9 Wibbelt, G., Moore, M. S., Schountz, T. & Voigt, C. C. Emerging diseases in Chiroptera: why
   437 bats? *Biol. Lett.* 6, 438-440, doi:10.1098/rsbl.2010.0267 (2010).
- 438 10 Luis, A. D. *et al.* A comparison of bats and rodents as reservoirs of zoonotic viruses: are bats special? *Proc. Biol. Sci.* **280**, 20122753, doi:10.1098/rspb.2012.2753 (2013).
- 440 11 Luis, A. D. *et al.* Network analysis of host-virus communities in bats and rodents reveals determinants of cross-species transmission. *Ecol. Lett.* doi:10.1111/ele.12491 (2015).
- 442 12 de Vienne, D. M. *et al.* Cospeciation vs host-shift speciation: methods for testing, evidence 443 from natural associations and relation to coevolution. *New Phytologist.* **198, 347-385,** (2013).
- 444 13 Kitchen, A., Shackelton, L. A. & Holmes, E. C. Family level phylogenies reveal modes of 445 macroevolution in RNA viruses. *Proc. Natl. Acad. Sci. USA* **108**, 238-243, 446 doi:10.1073/pnas.1011090108 (2011).
- 447 14 Ramsden, C., Holmes, E. C. & Charleston, M. A. Hantavirus evolution in relation to its rodent 448 and insectivore hosts: no evidence for codivergence. *Mol. Biol. Evol.* **26**, 143-153, 449 doi:10.1093/molbev/msn234 (2009).
- Coulibaly-N'Golo, D. et al. Novel arenavirus sequences in Hylomyscus sp. and Mus setulosus
   (Nannomys) from Cote d'Ivoire: implications for evolution of arenaviruses in Africa. PLoS One.
   6, e20893, doi:10.1371/journal.pone.0020893 (2011).
- 453 16 Streicker, D. G. *et al.* Host phylogeny constrains cross-species emergence and establishment 454 of rabies virus in bats. *Science* **329**, 676-679, doi:10.1126/science.1188836 (2010).
- 455 17 Wilkinson, D. A. *et al.* Highly diverse Morbillivirus-related paramyxoviruses in wild fauna of 456 the southwestern indian ocean islands: evidence of exchange between introduced and 457 endemic small mammals. *J. Virol.* **88**, 8268-8277, doi:10.1128/JVI.01211-14 (2014).
- Streicker, D. G., Lemey, P., Velasco-Villa, A. & Rupprecht, C. E. Rates of viral evolution are
   linked to host geography in bat rabies. *PLoS Pathog.* 8, e1002720,
   doi:10.1371/journal.ppat.1002720 (2012).
- 461 19 Gortazar, C. et al. Crossing the interspecies barrier: opening the door to zoonotic pathogens.
   462 PLoS Pathog. 10, e1004129, doi:10.1371/journal.ppat.1004129 (2014).
- Pandey, A. *et al.* Strategies for containing Ebola in West Africa. *Science* **346**, 991-995,
   doi:10.1126/science.1260612 (2014).
- 465 21 Morse, S. S. et al. Prediction and prevention of the next pandemic zoonosis. Lancet 380,
   466 1956-1965, doi:10.1016/S0140-6736(12)61684-5 (2012).
- Pringle, C. R. & Easton, A. J. Monopartite negative strand RNA genomes. <u>Sem. Virol.</u> 8, 49-57 (1997).
- 469 23 Drexler, J. F. *et al.* Bats host major mammalian paramyxoviruses. *Nat. Commun.* **3**, 796, 470 doi:10.1038/ncomms1796 (2012).
- Wilkinson, D. A. *et al.* Identification of novel paramyxoviruses in insectivorous bats of the southwest indian ocean. *Virus Res* **170**, 159-163, doi:10.1016/j.virusres.2012.08.022 (2012).
- 473 25 Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. & Kent, J. Biodiversity 474 hotspots for conservation priorities. *Nature* **403**, 853-858, doi:10.1038/35002501 (2000).
- 475 26 Goodman, S. M. & Raherilalao, M. J. *Atlas of selected land vertebrates of Madagascar*.
  476 Association Vahatra, Antananarivo, Madagascar (2013).

- 477 27 Koopman, K. F. A synopsis of the families of bats, part VII. *Bat Research News* **25**, 25-27 478 (1984).
- 479 28 Teeling, E. C. *et al.* A molecular phylogeny for bats illuminates biogeography and the fossil record. *Science* **307**, 580-584, doi:10.1126/science.1105113 (2005).
- 481 29 Goodman, S. M. *Les chauves-souris de Madagascar*. Association Vahatra, Antananarivo, Madagascar (2011).
- 483 30 Goodman, S. M. & Ramasindrazana, B. Description of a new species of the Miniopterus 484 aelleni group (Chiroptera: Miniopteridae) from upland areas of central and northern 485 Madagascar. *Zootaxa* **3936**, 538-558, doi:10.11646/zootaxa.3936.4.4 (2015).
- Goodman, S. M. *et al.* An integrative approach to characterize Malagasy bats of the subfamily Vespertilioninae Gray, 1821, with the description of a new species of *Hypsugo. Zool. J. Linn.*Soc. 173, 988-1018 (2015).
- 489 32 Christidis, L., Goodman, S. M., Naughton, K. & Appleton, B. Insights into the evolution of a 490 cryptic radiation of bats: dispersal and ecological radiation of Malagasy *Miniopterus* 491 (Chiroptera: Miniopteridae). *PLoS One* **9**, e92440, doi:10.1371/journal.pone.0092440 (2014).
- 492 33 Clayton, D. H., Bush, S. E. & Johnson, K. P. Ecology of congruence: past meets present. *Syst.* 493 *Biol.* 53, 165-173 (2004).
- 494 34 Criscione, C. D., Poulin, R. & Blouin, M. S. Molecular ecology of parasites: elucidating 495 ecological and microevolutionary processes. *Mol. Ecol.* **14**, 2247-2257, doi:10.1111/j.1365-496 294X.2005.02587.x (2005).
- 497 35 Nieberding, C. M. & Morand, S. Comparative phylogeography: The use of parasites for 498 insights into host history (eds S. Morand, B.R. Krasnov and R. Poulin). Chapter 15, 277-293. 499 (Micromammals and macroparasites, 2006).
- Nieberding, C. M., Jousselin, E. & Desdevises. The use of co-phylogeographic patterns to predict the nature of host–parasite interactions, and vice versa (eds Serge Morand & Boris Krasnov). Chapter 5, 59-69 (The Biogeography of Host-Parasite Interactions, 2010).
- 503 37 Duffy, S., Shackelton, L. A. & Holmes, E. C. Rates of evolutionary change in viruses: patterns and determinants. *Nat. Rev. Genet.* **9**, 267-276, doi:10.1038/nrg2323 (2008).
- 505 38 Domingo, E. *et al.* Viruses as quasispecies: biological implications. *Curr. Top. Microbiol.* 506 *Immunol.* **299**, 51-82 (2006).
- Nemirov, K., Leirs, H., Lundkvist, A. & Olsson, G. E. Puumala hantavirus and *Myodes glareolus* in northern Europe: no evidence of co-divergence between genetic lineages of virus and
   host. *J. Gen. Virol.* **91**, 1262-1274, doi:10.1099/vir.0.016618-0 (2010).
- Wang, C. Q. et al. Co-circulation of Hantaan, Kenkeme, and Khabarovsk Hantaviruses in
   Bolshoy Ussuriysky Island, China. Virus. Res. 191, 51-58, doi:10.1016/j.virusres.2014.07.021
   (2014).
- 513 41 Cui, J. et al. Evolutionary relationships between bat coronaviruses and their hosts. Emerg. 514 Infect. Dis. 13, 1526 (2007).
- Weir, D. L., Smith, I. L., Bossart, K. N., Wang, L. F. & Broder, C. C. Host cell tropism mediated
  by Australian bat lyssavirus envelope glycoproteins. *Virology* 444, 21-30,
  doi:10.1016/j.virol.2013.06.016 (2013).
- Lawrence, P. *et al.* Surface glycoproteins of the recently identified African Henipavirus promote viral entry and cell fusion in a range of human, simian and bat cell lines. *Virus. Res.* **181**, 77-80, doi:10.1016/j.virusres.2014.01.003 (2014).
- Faria, N. R., Suchard, M. A., Rambaut, A., Streicker, D. G. & Lemey, P. Simultaneously
   reconstructing viral cross-species transmission history and identifying the underlying
   constraints. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 368, 20120196,
   doi:10.1098/rstb.2012.0196 (2013).
- 525 45 Kunz, T. H. & Parsons, S. *Ecological and Behavioral Methods for the Study of Bats*, 2nd Edition. Austral Ecology, 36: e10–e11. doi: 10.1111/j.1442-9993.2010.02172.x. (1988).
- 527 46 Quantum GIS Development Team. Quantum GIS Geographic Information System. Open 528 Source Geospatial Foundation Project. http://qgis.osgeo.org (2015).

529 47 Cornet, A. Essai de cartographie bioclimatique à Madagascar. Notice explicative, ORSTOM, 530 Paris 55, 1-28 (1974). 531 48 Sikes, R. S. & Gannon, W. L. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. J. Mammal. 92, 235-253 (2011). 532 533 49 R Core Team. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, http://www.R-project.org/. 534 535 (2013).536 McCullagh, P. Generalized linear models. Eur. J. Oper. Res. 16, 285-292 (1984). 50 537 Breslow, N. E. & Clayton, D. G. Approximate inference in generalized linear mixed models. J 538 Am. Stat. Assoc. 88, 9-25 (1993). 539 52 Tong, S., Chern, S. W., Li, Y., Pallansch, M. A. & Anderson, L. J. Sensitive and broadly reactive reverse transcription-PCR assays to detect novel paramyxoviruses. J. Clin. Microbiol. 46, 540 541 2652-2658, doi:10.1128/JCM.00192-08 (2008). Kearse, M. et al. Geneious Basic: an integrated and extendable desktop software platform for 542 53 543 the organization and analysis of sequence data. Bioinformatics 28, 1647-1649, 544 doi:10.1093/bioinformatics/bts199 (2012). 545 Posada, D. & Crandall, K. A. MODELTEST: testing the model of DNA substitution. 54 546 Bioinformatics 14, 817-818 (1998). 547 55 Burnham, K. P. & Anderson, D. R. Multimodel inference understanding AIC and BIC in model 548 selection. Sociol. Methods Res. 33, 261-304 (2004). 56 Huelsenbeck, J. P., Ronquist, F., Nielsen, R. & Bollback, J. P. Bayesian inference of phylogeny 549 550 and its impact on evolutionary biology. Science 294, 2310-2314, 551 doi:10.1126/science.1065889 (2001). 552 57 Irwin, D. M., Kocher, T. D. & Wilson, A. C. Evolution of the cytochrome b gene of mammals. J 553 Mol. Evol. 32, 128-144 (1991). 554 58 Schloss, P. D. et al. Introducing mothur: open-source, platform-independent, communitysupported software for describing and comparing microbial communities. Appl. Environ. 555 556 Microbiol. 175, 7537-7541, doi:10.1128/AEM.01541-09 (2009). 557 59 Merkle, D., Middendorf, M. & Wieseke, N. A parameter-adaptive dynamic programming 558 approach for inferring cophylogenies. BMC Bioinformatics 11 Suppl 1, S60, 559 doi:10.1186/1471-2105-11-S1-S60 (2010). 560 60 Desdevises, Y. Cophylogeny: insights from fish-parasite systems. Parassitologia 49, 125-128 561 (2007).562 563 Acknowledgments For kindly providing research permits for our work on Madagascar, we are grateful to Département de 564 Biologie Animale, Université d'Antananarivo; Direction du Système des Aires Protégées et Direction 565 566 Générale de l'Environnement et des Forêts; and Madagascar National Parks. We are indebted to the 567 Institut Pasteur de Madagascar for their help in sample preservation and exportation and Pr. Xavier N. de Lamballerie at Aix Marseille University, IRD, and Dr. David A. Wilkinson from CRVOI for their 568

18

technical assistance. This work was financially supported by CRVOI and from the European Regional

Development Funds FEDER POCT ParamyxOI project (N° 33857). The postdoctoral fellowships of

B. Ramasindrazana were funded by RUN-Emerge, European Commission FP7 Regpot Capacity

569

570

572	program, the Fonds de Coopération régionale, Prefecture de La Réunion, and The Field Museum of
573	$Natural\ History,\ Chicago,\ through\ the\ Dr.\ Ralph\ and\ Marian\ Falk\ Medical\ Research\ Trust.\ N.\ Wieseke$
574	was funded by the German Research Foundation (DFG) (Proj. No. MI439/14-1). J. Mélade is a PhD
575	candidate at The University of La Réunion. His research interests include zoonotic virus and viral
576	evolution mechanisms.
577	Author contributions
578	KD, SMG, and HP conceived and designed the study. JM performed the experiments. JM, NW, OF,
579	and HP analyzed the data. SMG, BR, YG, and EL conducted the bat field sampling and contributed
580	reagents/materials/analysis tools. JM, KD, SMG, and HP wrote the paper.
581	Additional information
582	Supplementary information accompanies this paper at http://www.nature.com/srep
583	Competing financial interests: The authors declare no competing financial interests.
584	Reprints and permission information is available online at http://npg.nature.com/
585	reprints and permissions/
586	How to cite this article: xxxx
587	License: This work is licensed under a Creative Commons Attribution-NonCommercial-
588	No Derivative Works $3.0$ Unported License. To view a copy of this license, visit http://
589	creativecommons.org/licenses/by-nc-nd/3.0/
590	
591	

Family	Species	Total positive/total tested (%)	Grand total for family
Emballonuridae	Coleura kibomalandy	2/6 (33.3%)	2/9 (22.2)
	Paremballomira tiavato	0/3	()
Hipposideridae	Hipposideros commersoni	0/27	0/27
Miniopteridae	Miniopterus aelleni	0/7	30/289 (10.4)
8	Miniopterus cf. ambohitrensis	1/19 (5.3)	
	Miniopterus gleni	4/22 (18.2)	
	Miniopterus griffithsi	0/7	
	Miniopterus griveaudi	18/116 (15.5)	
	Miniopterus mahafaliensis	4/89 (4.5)	
	Miniopterus majori	0/7	
	Miniopterus sororculus	2/22 (9.1)	
Molossidae	Chaerephon atsinanana	0/34	36/406 (8.9)
	Chaerephon leucogaster	6/94 (6.4)	
	Mops leucostigma	11/68 (16.2)	
	Mops midas	1/19 (5.3)	
	Mormopterus jugularis	12/152 (7.9)	
	Otomops madagascariensis	7/39 (17.9)	
Pteropodidae	Eidolon dupreamım	0/11	3/80 (3.8)
\$2,000 for \$2.5 (2000) • 1000 (1000) (1000) (1000)	Pteropus rufus	3/20 (15.0)	5-0000 600 cm - 19-00 600 600 F (s
	Rousettus madagascariensis	0/49	
Rhinonycteridae	Paratriaenops furculus	1/14 (7.1)	22/56 (39.3)
	Triaenops menamena	21/42 (50.0)	
Vespertilionidae	Hypsugo bemainty	0/2	6/80 (7.5)
•	Myotis goudoti	5/48 (10.4)	
	Neoromicia malagasyensis	0/2	
	Neoromicia matroka	0/4	
	Neoromicia robertsi	0/1	
	Pipistrellus cf. hesperidus	0/8	
	Pipistrellus hesperidus	1/11 (9.1)	
	Pipistrellus raceyi	0/3	
	Scotophilus marovaza	0/1	
Grand total	3	ş.	99/947 (10.5%)

Table 2. Summary of the binomial GLM on individual infection (n=947). The model was selected after inspection of bivariate relationships and interactions. Because of unbalance, type III sums-of-squares were used to test the effects. MAT: Mean Annual Temperature, Habitat: habitat type, Multi: multispecies/monospecific site, MAR: mean annual rainfall, Df: degrees of freedom associated with the effect, Deviance: deviance of the model, F value: value of Fisher statistics for the different effects, Pr (>F): P values associated with the tests. Symbols for P values as follows:  $^< 0.1$ ,  $^* < 0.05$ ,  $^* < 0.01$ ,  $^* < 0.01$ .

Effect	Df	Deviance	F value	Pr (>F)
MAT	1	604.1	9.166	0.002**
Habitat	2	598.9	0.4938	0.61
Multi	1	605.7	11.65	0.0006***
Diet	1	602.6	6.803	0.009**
Habitat:Multi	2	601.3	2.423	0.0891^
Multi:MAR	2	604.6	5.006	0.007**
Multi:MAR <sup>2</sup>	2	602.3	3.142	0.043*
Residuals		598.2		


Table 3. Results for event base co-phylogeny obtained with CoRe-PA and number of the different events for sets of (A) 29 OTUs and (B) 39 OTUs. q indicates the quality values of each reconstructions.


			Frequency of events			Total cost					
OTUs	Reconstruction	(q)	Co- speciation	Sorting	Duplication	Host- switch	Co- speciation	Sorting	Duplication	Host- switch	Total
	1A	0.256	8	51	21	19	0.227	0.064	0.146	0.564	18.91
A	2A	0.277	7	47	21	20	0.241	0.071	0.162	0.526	18.93
	3A	0.286	10	47	18	20	0.171	0.076	0.187	0.566	19.96
			× ·	×	Va	*60	š		X:		fe v
В	1B	0.25	5	57	33	21	0.22	0.07	0.13	0.58	21.52
	2B	0.26	5	51	32	22	0.23	0.08	0.13	0.56	21.84
	3B	0.28	6	63	33	20	0.15	0.05	0.08	0.7	21.1


609


### 610 Figures Legends


- Figure 1. PVs detection rates among the sites sampled on Madagascar. Only sites containing positive bats are represented. Abbreviations refer to the names of sampling sites (e.g. ANDRF for
- 613 "Andrafiabe"). n, numerator = the number of individuals that tested positive for PVs and denominator
- ethe number of individuals tested. Provincial capitals are indicated by black squares. QGIS<sup>46</sup>, an
- open-source GIS software (http://qgis.osgeo.org/en/site/), was used to generate the map for
- visualizing bioclimatic regions of Madagascar proposed by Cornet 47.
- Figure 2. Proportion of infected bats depending on species diversity at each sampling site and
- 618 the context of the where the samples were collected. Individual outlying data points are displayed as
- 619 circles.
- 620 Figure 3. Phylogeny of the UMRVs detected in bats from Madagascar. A global phylogeny of 99
- 621 partial L-gene sequences calculated in 50,000,000 iterations in MrBayes with the GTR + G
- 622 evolutionary model and a 10% burn-in rooted with a Mumps virus sequence (GenBank number
- 623 AY309060). Only Bayesian with posterior probabilities > 0.7 were represented. Host switching events
- 624 were highlighted in blue and host-specificity for bats sharing the same sites in green. Bat species
- occurring at distant sites are highlighted in red. Bats living at distant sites and hosting with low level
- of *UMRVs* nucleotide similarity are highlighted in yellow.
- 627 Figure 4. The first preferred reconstruction with the first best-cost model fit of the co-
- 628 evolutionary history for the set of (A) 24 OTUs and (B) 39 OTUs and associated bat-species
- 629 retrieved from CoRe-PA software. Host tree is represented in black; parasite tree is represented in
- 630 grey.
- 631 Figure 5. Frequency of co-speciation (A) and host-switches (B) events for the set of 24 OTUs
- 632 **obtained by random reconstructions.** The number of co-speciation and host-switches events
- 633 expected in the most parsimonious reconstructions by CoRe-PA, 8 and 15, respectively (framed in red;
- also in Table 3) were compared to the random reconstructions events below. The macro-evolutionary
- 635 events showing lower random reconstruction events than expected (8 or 15) was determined as the
- 636 predominant event.














## 5.3. Conclusion du chapitre 5

Nos résultats ont permis de caractériser de nouveaux Paramyxovirus chez une centaine de chauves-souris, appartenant à plus de 16 espèces différentes, et confirment la grande variabilité génétique de ces virus et l'étendue de leur spectre d'hôtes dans la faune sauvage.

L'analyse phylogénétique Bayesienne nous a conduit à classer les nouveaux membres de ces paramyxovirus dans le phylogroupe des *URMVs* permettant de dégager deux patrons phylogénétiques distincts:

- (i) une spécificité d'hôte entre certaines chauves-souris géo-localisées au même endroit,
- (ii) un saut d'espèces évident non seulement entre chauves-souris phylogénétiquement apparentées et physiquement proches, mais également entre chauves-souris phylogénétiquement et physiquement éloignées. Cette dernière observation suggère l'intervention probable d'une espèce intermédiaire jouant le rôle d'un "pont épidémiologique" entre chauves-souris qui ne sont *a priori* pas en contact.

Nous avons montré que le mécanisme macro-évolutif majeur mis en œuvre chez les *UMRVs* repose essentiellement sur le « *host-switch* », et non pas sur la co-spéciation. Ce mécanisme est sans doute à l'origine de la diversité génétique virale observée et du spectre d'hôte important.

Nous avons également caractérisé des associations statistiquement significatives entre certains hôtes et virus (spécificité d'hôtes) qui seraient le résultat, non pas de mécanismes macro-évolutifs (actifs sur le très long terme), mais probablement de mécanismes micro-évolutifs agissant sur des échelles de temps beaucoup plus courtes et en rapport avec l'adaptation de virus à son nouvel hôte postérieure au saut d'espèce.

L'analyse statistique multivariée a permis d'identifier 4 facteurs environnementaux qui favoriseraient soit une transmission virale, soit l'infection virale elle-même. Ils comprennent la température, le niveau de précipitations, un contexte multi-espèces, ainsi que le régime alimentaire insectivores. Le contact physique (proximité) entre chauves-souris dans les grottes (que ce soit entre espèces ou entre familles apparentées) influence la probabilité d'infection.

Ces travaux ont été valorisés sous forme d'un article scientifique soumis et d'une communication orale présentée au congrès international « bats, small mammals and infectious agents » à l'Université de La Réunion (2013).

# Chapitre 6. Etude des lyssavirus chez les chauvessouris de la zone Sud-Ouest de l'Océan Indien

### 6.1. Généralités

La famille des *Rhabdoviridae* est composée de six genres : *Vesiculovirus*, *Ephemerovirus*, *Novirhabdovirus*, *Cytorhabdovirus*, *Nucleorhabdovirus* et *Lyssavirus*. Ce dernier genre comprend 15 virus (ICTV, 2014). Les 15 virus sont regroupés en 3 phylogroupes en fonction de leur antigénicité croisée et de leurs distances phylogénétiques (Badrane et al., 2001).

Les Lyssavirus sont retrouvés sur tous les continents à l'exception de l'Antarctique. La Figure en **annexe 3** illustre les différents réservoirs et la distribution géographique des différents variants de lyssavirus rapportés à ce jour. Les lyssavirus non rabiques ont été trouvés uniquement dans l'Ancien Monde alors qu'ils n'ont jamais été détectés en Amérique. A l'opposé, le continent américain est la seule partie du monde où il est courant de détecter le virus rabique (agent de la rage) chez les chauves-souris. Les seuls lyssavirus signalés en Afrique sont les virus Duvenhage, le virus de Lagos Bat, le virus Mokola, le virus Shimoni et le virus Ikoma.

Le virus rabique, agent de la rage, appartient au génotype 1 des lyssavirus, est l'espèce type du genre et aussi le plus important en termes de santé publique et vétérinaire. La rage est transmise à l'homme par la salive d'un animal infecté en phase d'excrétion virale. Le virus remonte à partir du point d'inoculation le long des terminaisons nerveuses, vers le système nerveux central. Quand ce dernier est atteint, le tableau clinique réalisé est celui d'une encéphalite aigue virale et à ce stade la rage est constamment mortelle (Tordo et al., 2010). Parmi les 14 autres lyssavirus, seuls EBLV-1 et 2, ABLV, DUVV, IRKV et MOKV ont causé des pathologies neurologiques similaires à la rage (Condori-Condori et al., 2013).

Le génome des lyssavirus est d'environ 12000 bases et code cinq gènes monocistroniques: une nucléoprotéine (N), une phosphoprotéine (P), une protéine de matrice (M), une

glycoprotéine (G) et une ARN polymérase ARN-dépendante (L). Les protéines N, P et L forment ensemble avec l'ARN, la nucléocapside (Lyles & Rupprecht, 2007).

Les protéines G forment des spicules qui se projettent à travers la membrane du virion. La protéine M relit la nucléocapside interne aux protéines G membranaires (Finke & Conzelmann, 2005) (annexe 4).

Le diagnostic de l'infection par les lyssavirus est basé i) soit sur la détection des antigènes viraux ou du génome viral (par RT-PCR ou qRT-PCR (Coertse et al., 2010) ou par isolement à partir de tissus infectés ou la salive ii) soit par la détection des anticorps spécifiques qui indique une exposition antérieure au virus. La technique « rapid fluorescent focus inhibition test » (RFFIT) détecte des anticorps neutralisants le virus rabique et est le test sérologique considéré comme le « gold standard ».

# **6.2.** Approches expérimentales

Avant nos travaux, les investigations sur les lyssavirus de la région SOOI n'ont concerné que Madagascar, et encore, de façon très parcellaire. En effet, une seule étude antérieur décrit la présence d'anticorps neutralisants chez deux espèces de chauves-souris frugivores malgaches, *Rousettus madagascariensis* et *Eidolon helvum* (Reynes et al., 2012). La totalité des chauves-souris insectivores (6 espèces pour un total de 92 individus) analysées s'était révélée être séronégatives. L'étude régionale que nous rapportons dans mon travail de thèse apporte une information plus complète en démontrant la circulation intense des lyssavirus dans toutes les îles de la région.

Dans un premier temps, nous avons recherché la trace sérologique d'une infection passée des chauves-souris par des lyssavirus, en ayant recours à la technique RFFIT. Ce ne sont pas moins de 572 chauves-souris, représentant 22 espèces, qui ont pu être criblées pour la présence d'anticorps neutralisants contre 4 lyssavirus de référence: le virus rabique CVS-11, les lyssavirus DUVV, EBLV-1 et ABLV appartenant au phylogroupe 1 et le LBV appartenant au phylogroupe 2. Dans une seconde étape, le broyat de cerveau de chaque individu, à l'exception des chauves-souris de Mayotte, a été analysé par qRT-PCR pour rechercher la présence d'ARN viral. Cette étude sérologique a permis de démontrer la circulation des lyssavirus dans la faune sauvage de la région SOOI et d'établir le phylogroupe auquel ils appartiendraient.

Ces travaux ont été menés conjointement avec le laboratoire du Pr Wanda Markotter de l'Université de Pretoria en Afrique du Sud où j'ai réalisé un stage doctoral de 5 mois.

Annexe 6: Dans un travail complémentaire, nous avons élargi l'enquête sérologique afin d'identifier les traces du passage de virus ARN simple brin dans la faune sauvage de la région SOOI. En utilisant la technique d'immunofluorescence indirect (ou Indirect Immunofluorescence assay IIFA) sur un panel de 15 virus de référence grâce aux lames de microscopie « Euroimmun Mosaic Chip » nous avons pu dresser le profil de réactivité sérologique à ces virus des populations de chauves-souris et de rongeurs de la région. Un total de 643 chauves-souris (29 espèces), de Madagascar, La Réunion, Maurice, Mahé, et Mayotte; ainsi que 199 rongeurs introduits (3 espèces) de Madagascar, Mahé, Mayotte and La Réunion et 57 petits mammifères endémiques de Madagascar (*Eliurus majori* et *Eliurus minor*) ont été analysés pour la présence d'anticorps dirigés contre en 5 familles virales décrites en annexe.

Cette étude nous a permis d'identifier au niveau de la famille virale et du genre, les agents infectieux viraux putatifs auxquels les animaux de la faune sauvage sont exposés. Bien que l'information ne puisse aller jusqu'à la caractérisation de l'agent viral lui-même, elle apporte une donnée précieuse sur la réactivité antigénique croisée entre les virus tests utilisés et les agents viraux putatifs, donnée utile pour l'identification des genres viraux en circulation, l'identification des principaux réservoirs en jeu, l'orientation des investigations complémentaires et l'évaluation du risque potentiel pour l'homme.

Ce travail a été mené conjointement avec le Laboratoire de Virologie du Pr Christian Drosten du Centre Hospitalier Universitaire de Bonn en Allemagne par la réalisation d'un stage doctoral de 3 semaines que j'ai effectué du 6 Septembre 2014 au 27 Septembre 2014.

6.3. Mise en évidence sérologique d'une circulation importante de lyssavirus chez les chauves-souris de la zone du Sud-ouest de l'Océan Indien.

### **PLOS ONE**

# Serological Evidence of Diverse Circulation of Lyssaviruses among Bats on Southwestern Indian Ocean Islands --Manuscript Draft--

Manuscript Number:						
Article Type:	Research Article					
Full Title:	Serological Evidence of Diverse Circulation of Lyssaviruses among Bats on Southwestern Indian Ocean Islands					
Short Title:	Lyssavirus neutralizing antibody in bats from the Southwestern Indian Ocean Islands					
Corresponding Author:	Julien Mélade Universite de la Reunion Ste Clotilde, RÉUNION					
Keywords:	Lyssavirus, Bats, Rabies, South Western Indian Ocean, RFFIT					
Abstract:	We provide serological evidence of diverse lyssavirus circulation among bats on southwestern Indian Ocean (SWIO) islands. A total of 572 bats, belonging to 22 species, from Anjouan and Mayotte (Comoros), Madagascar, Reunion Island, Mauritius, and Mahé (Seychelles) were screened by the Rapid Fluorescent Focus Inhibition test for the presence of neutralizing antibodies against five lyssaviruses representing both phylogroup I (Rabies virus, Duvenhage virus, European bat lyssavirus I, and Australian bat lyssavirus) and phylogroup II (Lagos bat virus). In total, 186 bats were able to neutralize at least one challenge virus, with large variation between islands, roost sites, and bat species. However, no lyssavirus RNA was detected by PCR of bat brains. These results highlight that lyssaviruses belonging to phylogroups I and II circulate in bat populations from the region, a feature that could be of public health importance, considering that certain bat species have contact with humans.					
Order of Authors:	Julien Mélade					
	Stewart McCulloch					
	Beza Ramasindrazana					
	Erwan Lagadec					
	Magali Turpin					
	Hervé Pascalis					
	Steven Michael Goodman					
	Wanda Markotter					
	Koussay Dellagi					
Opposed Reviewers:						
Additional Information:						
Question	Response					
Financial Disclosure  Please describe all sources of funding that have supported your work. A	This work was supported by funding from CRVOI and from the European Regional Development Funds FEDER POCT "Pathogènes associés à la Faune Sauvage Océan Indien" project (N° 31189), and FEDER-POCT ParamyxOI project (N° 33857), as well as the National Research Foundation, South Africa.  Sampling on Mayotte was supported by ECOSAN/CNRS-INEE "Batman".  The funders had no role in study design, data collection and analysis, decision to					
complete funding statement should do the following:	publish, or preparation of the manuscript					
Include grant numbers and the URLs of any funder's website. Use the full name, not acronyms, of funding institutions, and						

Powered by Editorial Manager® and ProduXion Manager® from Aries Systems Corporation

- 1 Serological Evidence of Diverse Circulation of Lyssaviruses among Bats on
- 2 Southwestern Indian Ocean Islands

3

- 4 Julien Mélade<sup>1,2,3</sup>, Stewart McCulloch<sup>4</sup>, Beza Ramasindrazana<sup>1,2,3,5,6</sup>, Erwan Lagadec<sup>1,2,3</sup>,
- 5 Magali Turpin<sup>1,2,3</sup>, Hervé Pascalis<sup>1,2,3</sup>, Steven M. Goodman<sup>5,7</sup>, Wanda Markotter<sup>4</sup>,
- 6 Koussay Dellagi<sup>1,2,3</sup>

7

- 8 1 Centre de Recherche et de Veille sur les Maladies Emergentes dans l'Océan Indien
- 9 (CRVOI), Plateforme de Recherche CYROI, 2 rue Maxime Rivière, 97490 Sainte Clotilde, La
- 10 Réunion, France, 2 Université de La Réunion, UMR PIMIT «Processus Infectieux en Milieu
- 11 Insulaire Tropical», INSERM U1187, CNRS 9192, IRD 249. Plateforme de Recherche
- 12 CYROI, Saint Denis, La Réunion, France, 3 Institut de Recherche pour le Développement
- 13 (IRD), IRD BP 50172, 97492 Sainte-Clotilde, La Réunion, France, 4 Department of
- 14 Microbiology and Plant Pathology, Faculty of Natural and Agricultural Sciences, University
- of Pretoria, Pretoria 0001, South Africa, 5 Association Vahatra, BP 3972, Antananarivo 101,
- 16 Madagascar, 6 Institut Pasteur de Madagascar, BP 1274, Antananarivo 101, Madagascar,
- 7 Field Museum of Natural History, 1400 S. Lake Shore Dr, Chicago, IL 60605-2496, United
- 18 States of America

### 19 Abstract

- 20 We provide serological evidence of diverse lyssavirus circulation among bats on southwestern
- 21 Indian Ocean (SWIO) islands. A total of 572 bats, belonging to 22 species, from Anjouan and
- 22 Mayotte (Comoros), Madagascar, Reunion Island, Mauritius, and Mahé (Seychelles) were
- 23 screened by the Rapid Fluorescent Focus Inhibition test for the presence of neutralizing

antibodies against five lyssaviruses representing both phylogroup I (*Rabies virus*, *Duvenhage virus*, *European bat lyssavirus I*, and *Australian bat lyssavirus*) and phylogroup II (*Lagos bat virus*). In total, 186 bats were able to neutralize at least one challenge virus, with large variation between islands, roost sites, and bat species. However, no lyssavirus RNA was detected by PCR of bat brains. These results highlight that lyssaviruses belonging to phylogroups I and II circulate in bat populations from the region, a feature that could be of public health importance, considering that certain bat species have contact with humans.

# Introduction

33	Lyssaviruses (order: Mononegavirales, family: Rhabdoviridae) are RNA viruses with
34	single-stranded, negative-sense genomes approximately 12 kb in length [1]. Fifteen distinct
35	viral species, classified into three phylogroups, have been identified to date within the
36	Lyssaviruses genus [2, 3]. These are the prototypic Rabies virus (RABV) and the 14
37	genetically related species referred to as rabies-related viruses that cluster into phylogroups
38	according to their antigenicity reaction and phylogenetic distances [4]: Duvenhage virus
39	(DUVV), European bat Lyssavirus types 1 (EBLV-1) and type 2 (EBLV-2), Australian bat
40	lyssavirus (ABLV), Irkut virus (IRKV), Aravan virus (ARAV), Khujand virus (KHUV), and
41	Bokeloh bat lyssavirus (BBLV) form phylogroup I. Lagos bat virus (LBV), Mokola virus
42	(MOKV), and Shimoni bat virus (SHIV) form the phylogroup II. Ikoma virus (IKOV), West
43	Caucasian bat virus (WCBV), and the newly described Lleida Bat lyssavirus (LLEBV) form
44	the phylogroup III. With the exceptions of MOKV and IKOV, all lyssaviruses have been
45	isolated from bats [5, 6].
46	RABV, the type species of the genus, is the principal causative agent of human rabies
47	a fatal acute encephalitis claiming 59 000 lives annually, primarily affecting Africa and Asia
48	[7]. The natural cyclic maintenance of RABV worldwide is principally achieved by dogs and
49	wild canids [8], while bats are seen as a reservoir only in the New World, where the virus is
50	known to infect insectivorous and hematophagous species [9]. Nowadays, the majority of
51	human exposure to rabies in the Americas is caused by contact with rabid bats [10]. In
52	contrast, among the 14 rabies-related viruses, only EBLV-1 and 2, ABLV, DUVV, IRKV
53	(phylogroup I), and MOKV (phylogroup II) have caused sporadic human deaths [9].

Islands of the southwestern Indian Ocean (SWIO) region, are seen as one of the world's biodiversity hotspots [11], hosting a rich chiropteran fauna. For example, on Madagascar with its 45 recognized bat species, 36 are endemic [12-14]; neighboring islands have notably lower measures of species diversity. Only one study conducted on Madagascar, a rabies endemic country where dogs are both maintenance reservoirs and vectors [15], has recorded antibodies to LBV and EBLV-1 in the frugivorous bat species *Eidolon dupreanum* and *Pteropus rufus*, but not six tested insectivorous bat species (n = 92) [16]. These antibodies are the trace indicators of past exposure in bats to a lyssavirus resulting in seroconversion. The other islands of the SWIO are deemed rabies free with no lyssavirus infection being recorded in non-volant mammals [15], although, the locally occurring bats have yet to be investigated. We report the results of a serosurvey conducted on bats from SWIO islands based on the screening of sera for lyssavirus neutralizing antibodies. This study is the first comprehensive analysis of SWIO region bats for exposure to lyssaviruses and aims to strengthen the available regional knowledge.

#### Methods

## Specimen collection

From March 2010 to March 2015, 653 bats were collected from the islands of Anjouan and Mayotte (Comoros Archipelago), Reunion Island and Mauritius (Mascarene Archipelago), Madagascar, and Mahé (Seychelles Archipelago) (Figure 1), representing 22 species from 6 families. Both insectivorous and frugivorous bats were captured using harp traps, hand nets and mist nets. Some of the larger fruit bats were obtained from hunters.

Individual bats were identified by morphological characters (outlined or cited [13]) and by

comparison to voucher specimens in The Field Museum of Natural History and The Département de Biologie Animale, Université d'Antananarivo.

Animals were manipulated in accordance with the guidelines for the handling of wild mammals [17], and all field protocols were designed to strictly follow the terms of the research permits issued by national authorities in the different countries this study took place (S1 Text). Details on ethical clearance are provided (S2 Text). This study has benefited from the sampling efforts conducted in the context of an ongoing international long-term project to catalog the terrestrial vertebrate fauna of Madagascar based on voucher specimens [18]. Bats trapped on Mayotte were released at the site of capture after sampling. Bat specimens sampled on the other islands where taken as voucher specimens housed in the above named museums. Details concerning the sex, the reproduction status [19], and age (wing bone ossification and dental eruption patterns) [20] for each captured bat were recorded. Names of capture sites were abbreviated (e.g. "Mangajou" abbreviate into MGJ) and reported in Table S.

Blood was collected by veinopuncture of the brachial vein or by cardiac puncture and subsequent exsanguinations (in some cases, needle and syringe were rinsed with heparin or EDTA to avoid blood coagulation during the procedure). Sera were separated from blood by centrifugation at 5000 rpm for 5 minutes. Brain was also collected from each of the vouchered individuals. Tissue samples were immediately stored in liquid nitrogen in the field and then transferred to -80°C freezers upon arrival at the laboratory.

Detection of lyssavirus neutralizing antibodies

Detection and titration of lyssavirus neutralizing antibodies were performed using the miniaturized Rapid Fluorescent Focus Inhibition Test (RFFIT) [21]. The challenge viruses used in the RFFIT were selected to represent two of the three phylogroups of lyssaviruses [3]:

RABV (CVS-11), DUVV (isolate DUVVSA06), EBLV-1 (isolate RV20), ABLV (isolate RV634), as representatives of phylogroup I, while LBV (isolate LBVAFR1999 Africa 1999) represented phylogroup II. Sera were tested at 1:5, 1:25, 1:125, and 1:625 dilutions. Each neutralization assay was performed in triplicate and was read by two different individuals. Serum samples producing neutralization above the threshold value of 1:5 with > 50% reduction of fluorescent foci were considered positive. Of the 653 serum samples, 81 were excluded from analysis either because they displayed a cell-adhesion inhibitory-type cytotoxicity (n = 45) or because the volume of sera was insufficient to do the assay in triplicate (n = 36). Hence, RFFIT was performed on 572 samples: all of which were tested against LBV as the phylogroup II challenge virus; 528 were also tested against CVS-11 and 540 also screened against DUVV as phylogroup I representatives. A subset of 220 samples had enough serum to allow additional testing against EBLV-1, and only 14 samples were screened for the presence of ABLV neutralizing antibodies.

## Lyssaviruses RNA Detection

For 550 bats, total RNA was extracted from brain with Trizol (Invitrogen, Carlsbad, California, USA). Lyssavirus RNA detection was performed using a real time reverse transcription PCR (QuantiTect Probe RT-PCR Kit, QIAGEN, Cat number 204443) with specific primers targeting the conserved region of the nucleoprotein genes of lyssaviruses [22].

### Statistical analysis

Statistical analyses were performed using Pearson chi-square ( $\chi^2$ ) or Fisher's exact tests in R software for calculation procedures (95% confidence intervals with a continuity

correction). Probability values < 0.05 were considered statistically significant. All statistical computations were conducted using R version 3.0.0 (RCoreTeam, 2013).

#### Results

123

143

124 Of the 653 serum samples collected, we were able to screen 572 samples by RFFIT for neutralizing antibodies to bat associated lyssaviruses, namely CVS-11, DUVV, EBLV, and 125 126 ABLV all belonging to phylogroup I and LBV from phylogroup II. These samples were from 127 421 insectivorous and 151 frugivorous bats, belonging to six bat families and 22 species. We 128 detected 186 individuals (32.5%) capable of neutralizing at least one of the challenge lyssaviruses above the 1:5 threshold. Seropositivity rates differed significantly between 129 130 islands: being highest on Reunion Island, Mauritius, and Mahé, with 42.9% (n = 52/121) 131 41.8% (n = 28/67), and 32.5% (n = 13/40,) respectively, while on Madagascar, Mayotte, and Anjouan, the seropositivity rates dropped to 27.9% (n = 84/301), 22.7% (n = 5/22), and 19.0%132 (n = 4/21), respectively (p = 0.01) (Table 1). The overall proportion of seropositive 133 134 individuals between females (n = 125/338; 37.0%) and males (n = 61/234; 26.1%) was 135 different (p = 0.005). The seropositivity rates did not significantly differ between 136 insectivorous bats (n = 134/421; 31.8%) or frugivorous bats (n = 52/151; 34.4%) nor between 137 neonatal (n = 0/2; 0.0%), juvenile (n = 16/42; 38.1%), sub-adult (n = 17/75; 22.7%), and adult 138 specimens (n = 153/453; 33.8%). 139 All six bat families represented in the samples had seropositive individuals to at least 140 one of the challenge viruses. At the taxon-level, 12 out of the 17 insectivorous bat species 141 tested seropositive to at least one of the challenge viruses. A significant difference (p < 0.001) 142 between species was observed with Hipposideros commersoni (Hipposideridae), and

Mormopterus francoismoutoui and M. acetabulosus (Molossidae) displaying the highest

seropositivity rates with 50.0% (n = 3/6), 42.9% (n = 52/121), and 38.7% (n = 12/31), respectively. Among the Molossidae, each of the three sister species of *Mormopterus*, M. acetabulosus, M. francoismoutoui, and M. jugularis [23] endemic to Reunion Island, Mauritius, and Madagascar, respectively, had a high proportion of seropositive individuals --42.9% (n = 52/121), 38.7% (n = 12/31), and 34.5% (n = 20/58), respectively. Two of the three frugivorous bats endemic to Madagascar (Pteropus rufus and Rousettus madagascariensis) tested positive (n = 8/12; 66.7% and n = 12/35; 34.3%, respectively), while the third, Eidolon dupreanum, tested negative (n = 0/9). Frugivorous species found on other regional islands were also positive: P. niger on Mauritius (n = 16/36; 44.4%), P. seychellensis from Mayotte (n = 3/19; 15.8%), and P. seychellensis Mahé (Seychelles) (n = 13/40; 32.5%). Six bat species, all from Madagascar, tested seronegative against all challenge viruses: Chaerephon atsinanana (n = 16), Miniopterus mahafaliensis (n = 11), Eidolon dupreanum (n = 9), Mops midas (n = 7), Miniopterus gleni (n = 4), and M. sororculus (n = 3).

Overall, and for both frugivorous and insectivorous bats, antibodies neutralizing DUVV were shown to be the most common: 97 sera out of 540 (17.9%) neutralized DUVV only and 13 out of 220 (5.9%) neutralized EBLV-1 only. Besides this, 42 samples of 572 (7.3%) neutralized exclusively LBV, indicating past exposure to a phylogroup II virus. Many sera reacted with more than one virus representative of phylogroup I: 32 of 220 (14.5%) simultaneously neutralized DUVV and EBLV-1; one sera of 14 (7.1%) neutralized CVS-11 and ABLV, and 1 sera of 218 (0.5%) cross-neutralized CVS-11 and EBLV-1. No serum could neutralize both challenge viruses of phylogroups I and II. In Table 1 we present the reactivity profile with all challenge viruses according to island and bat family and species sampled.

There was no apparent specificity of bat species to one challenge lyssavirus: all bat species, contained individuals that could neutralize either of the three test viruses, with LBV

- being constantly the least reactive. Four exceptions were observed: *Miniopterus* cf.
   *ambohitrensis* (Miniopteridae) reacted only with DUVV, *Mops leucostigma* (Molossidae),
   *Myotis goudoti* (Vespertilionidae), and *Triaenops menamena* (Rhinonycteridae) reacted only
   with LBV.
- 172 Looking at intra-island variation, several patterns emerge. On Madagascar, significant 173 difference were recorded between four bioclimate zones with the highest rate recorded in the 174 sub-humid region (n = 21/58; 36.2%) (p < 0.001) and among the low, middle, and high 175 elevational zones with 27.9% (n = 64/229), 11.5% (n = 3/26), and 36.9% (n = 17/46), 176 respectively, with the highest rates at high elevation (p = 0.05). Significant difference in 177 seroprevalence rates for positive animals was recorded between species, with the highest rate 178 in Pteropus rufus (n = 8/12; 66.7%) and the lowest in Mops leucostigma (n = 2/20; 10.0%). 179 Significant difference was also observed among localities with higher rate in AMBO (n = 180 8/12; 66.6%), the site the *P. rufus* were obtained and lower in AMBB (n = 1/11; 9.1%). 181 Twelve localities out of 33 were negative to the four challenge viruses (n = 70). On 182 Madagascar, seropositivity rates were statistically not different at sites containing multiple 183 species (n = 51/190; 26.8%) as compared to sites with single species (n = 33/111; 29.7%) or 184 between bat populations collected in caves (n = 46/158; 29.1%), in synanthropic roost sites (n = 46/158; 29.1%), in synanthropic roost sites (n = 46/158; 29.1%), in synanthropic roost sites (n = 46/158). 185 = 29/109; 26.6%), and in forests (n = 9/34; 26.5%). Rates were similar between insectivorous 186 bats (n = 64/245; 26.1%) and frugivorous bats (n = 20/56; 35.7%). No statistically significant 187 difference was recorded in seroprevalence rates in bats belonging to a given species collected 188 at different localities, whether captured at sites with multiple or single species (i.e., 189 Miniopterus griveaudi at sites AMBB, ANDRF, ANJHB, ANJHK1, and ANJKK). Moreover, 190 seroprevalence rates were not significantly different in bats common to some islands, such as 191 P. seychellensis on Mayotte (n = 3/19; 15.8%) and on Mahé (n = 13/40; 32.5%). On Reunion

Island, seropositivity rates were higher in females (n = 46/90; 51.1%) than males (6/31; 19.4%) (p = 0.001) and significantly higher seroprevalence rates was observed in bats captured in a maternity cave (50/100; 50.0%) as compared to those from a synanthropic locality (n = 2/21; 9.5%) (p < 0.001). In contrast, on Mahé, seroprevalence rates in *P. seychellensis* were significantly higher in male bats (n = 10/21; 47.6%) as compared to females (n = 3/19; 15.8%) (p = 0.03). No other significant differences were recorded between islands.

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

Various patterns of lyssavirus antibodies detection were observed on the different regional islands (Figure S1-S6). In most cases, positive sites did not show specific patterns. There were, however, some exceptions. For example, on Mayotte, where four positive sites were detected, bats neutralized DUVV only at two sites (n = 2/15; 13.3%) and LBV only at two other sites (n = 3/7; 42.9%). On Reunion Island, one site contained bats that neutralized only DUVV (n = 2/21; 9.5%) and the other one combined animals that neutralized LBV or cross neutralized DUVV and EBLV-1 (n = 50/100; 50.0%). The same feature was observed on Anjouan, where one locality contained bats that neutralized only DUVV (n = 1/6; 16.7%) and the other one combined animals that neutralized DUVV or LBV (n = 3/13; 23.1%). On Mahé, the two positive localities combined animals that neutralized DUVV or LBV, cross neutralized DUVV and EBLV-1 or cross neutralized CVS-11 and EBLV-1 (n = 13/40; 32.5%). On Mauritius, where seven positive sites where detected, bats neutralized DUVV only (n = 5/14; 35.7%) at three localities and the four other positive localities combined animals that neutralized DUVV, EBLV-1 or LBV or cross neutralized DUVV and EBLV-1 lyssaviruses (n = 23/53; 43.4%). On Madagascar, bats neutralized DUVV only (n = 31/105; 29.5%) or LBV only (n = 5/29; 17.2%) at 11 and three localities, respectively. The seven

other positive localities combined animals that neutralized DUVV, EBLV-1 or LBV, cross neutralized DUVV and EBLV or cross neutralized CVS-11 and ABLV (n = 48/97; 49.5%).

No lyssavirus RNAs were detected by PCR in the nucleic acids extracted from the brains of 550 different bats, whether seronegative or seropositive, with either of the test lyssaviruses.

### Discussion

Although several studies have reported on seroprevalence to lyssaviruses in bats in Africa and Asia [6, 24-30], limited information is available for the SWIO islands, which are located between these two continents. Our study reveals serological evidence of wide spread lyssavirus exposure, often at high prevalence rates, among frugivorous and insectivorous bats, on all investigated SWIO islands. At a regional scale, our study considerably expands the geographic coverage and bat species investigated when compared to the single study previously published on two Malagasy fruit bats, *Eidolon dupreanum* and *Pteropus rufus* [15]. Seroprevalence rates reported herein are similar to those reported in Africa, northern Vietnam, Cambodia, and the Philippines [24, 31-33]. Considering the close genetic relationships between lyssaviruses belonging to the same phylogroup, the dual neutralization occasionally observed between DUVV and EBLV-1 was expected [4, 25]. For those sera that reacted with one single challenge virus of phylogroup 1, one should consider the possibility that cross-neutralizing antibodies may have disappeared at the time of testing, long after infection of the bat by lyssavirus.

Our data highlight that 12 species of insectivorous bats and almost all species of frugivorous bats from the SWIO region, are seropositive towards at least one lyssavirus species. Particularly high rates of prevalence exposure was found in *Hipposideros* 

commersoni (Madagascar), Mormopterus francoismoutoui (Reunion Island), and M. acetabulosus (Mauritius). Each of these three species contained high neutralizing antibodies rates to DUVV or DUVV and EBVL-1. In agreement with earlier reports [32] showing higher rates in females, particularly in nursing colonies [33], we were also able to demonstrate higher seroprevalence rates in a maternity cave on Reunion Island, where 89 out of the 100 captured individuals were females. In a previous study [34], it has been shown that this maternity colony on Reunion Island, which harbors tens of thousands of female bats and their offspring, is a hotspot for pathogen transmission between roosting individuals. Longitudinal studies are warranted to analyze the temporal dynamics of antibody responses to lyssaviruses in bats living in stable or intermittent roosting sites, such as maternity colonies. Environmental conditions have been documented to promote exposure to and circulation of bat lyssaviruses [35]. This was also apparent in our study where higher seroprevalence rates were detected in sub-humid region and at higher elevations. However, we did not observe any significant difference in seroprevalences according to age classes of the sampled bats.

Based on the sero-neutralization profiles, at least one circulating lyssavirus should belong to phylogroup I, since most of positive sera neutralize DUVV or EBLV-1 or cross neutralize CVS-11 and ABLV or EBLV-1. Interestingly, at the regional level, if we exclude data obtained with ABLV (because of the limited number of samples tested with this virus), the highest rates of seroreactivity are related to DUVV (17.9%), followed by a dual reactivity with DUVV and EBLV (14.5%), then only with LBV (7.3%) and this reactivity profile is consistent across each island. In contrast to previous studies on African bats [36, 24], where antibodies to DUVV were mainly detected in insectivorous bats and antibodies to LBV in frugivorous bats, our data reveal that antibodies to DUVV were detected at high rates, in both these two groups of bats on SWIO islands. The very low reactivity of sera towards CVS-11

strengthens the hypothesis of a closer phylogenetic relationship between the actually circulating virus with DUVV or EBLV-1 than with CVS-11.

Our results also support the view that a phylogroup II lyssavirus, closely related to, if not LBV itself, is circulating in the region. This is in agreement with the study of Reynes and colleagues [16], who reported that 24% of sera from Malagasy *Eidolon dupreanum* neutralized only LBV. In our series, the highest rates with LBV across the different bat species and islands were recorded on Mayotte and Mauritius (13.6% and 10.4% of the samples, respectively) and in Malagasy *Myotis goudoti* and *Rousettus madagascariensis* (36.4% and 20.0% of the samples, respectively).

Madagascar and islands making up the Comoros Archipelago are closer to the African continent than the more outlying islands of the Mascarene and Seychelles archipelagos. Among lyssavirus belonging to phylogroups I and II, only DUVV, SHIV, and LBV, have been reported to date in African bats. Hence, one cannot exclude that DUVV and LBV are the putative lyssaviruses actually circulating on SWIO islands. Interestingly, the reported geographic distribution of DUVV, is primarily the northeastern part of South Africa, Zimbabwe, Swaziland, and eastern Africa [24], i.e. in close proximity to the Mozambique Channel and facing Madagascar. LBV has a much wider distribution, being isolated in western, central, eastern, and southern Africa [27], not only from frugivorous bats (Eidolon helvum and Epomophorus gambianus) but also from non-volant mammals, which develop dead end infection. However, with the exception of a few species, the bat fauna of Madagascar is largely endemic and holds a different suite of species than mainland Africa.

Alternatively, the putative bat lyssavirus(es) circulating on SWIO islands might be a yet unidentified species. Over geological time, numerous bat groups colonized Madagascar by

over water dispersal from the African continent. The lack of subsequent genetic exchange gave rise to divergence from these ancestral African lineages and the high levels of endemism of Madagascar's bat fauna. This may have generated long-term co-evolution for a range of hosts and their associated pathogens [37].

Our research shows for the first time, evidence of bat lyssavirus circulation on SWIO islands, all previously deemed rabies free. In our samples, we did not detect by RT-PCR, lyssavirus RNA in the brain of any insectivorous and frugivorous bat, whether the animal was seropositive or seronegative. This result is not unanticipated, as no abnormal morbidity or mortality among bats was observed during our study, and, in turn, the chance to isolate viruses from healthy animals is reported to be very low [38]. The failure to detect any active infection among animals, confirm that the seropositive individuals may have been infected during an earlier stage of their life and able to quickly clear the virus once they developed specific antibodies. The potential risk for human health with regard to contamination by bat lyssaviruses is very low, though it should not be considered as nil. In the Old World, very few cases of human rabies due to rabies-related bat lyssaviruses have been reported: only three cases, all fatal, have been reported for DUVV [39] and not one associated with LBV. However, the real number is unknown due to limited surveillance. Hence, an increased awareness of the risk of rabies transmission through accidental or professional contact with bats is recommended and any injury should be followed by a post exposure prophylaxy [40].

## **Supporting Information**

Figures S1-S6. Lyssaviruses neutralized by sera from bats captured on Anjouan (S1), Mayotte (S2), Reunion Island (S3) Mauritius (S4), Mahé (S5), and Madagascar (S6). (DOC)

- 309 Text S1 Authorizations for sampling of bats on different western Indian Ocean islands
- 310 **(DOC)**
- 311 Text S2. Ethical clearance (DOC)
- 312 Table S1. Sites names and corresponding abbreviations (DOC)
- 313 Acknowledgments
- 314 For kindly providing research and permits, we are grateful on Madagascar to Département de 315 Biologie Animale, Université d'Antananarivo; The Direction du Système des Aires Protégées 316 et Direction Générale de l'Environnement et des Forêts; and Madagascar National Parks; on 317 Reunion Island to Direction de l'Environnement, de l'Aménagement et du Logement de La 318 Réunion; on Mayotte to Direction de l'Environnement, de l'Aménagement et du Logement de 319 Mayotte; on Anjouan to Centre National de Documentation et de Recherche Scientifique 320 (CNDRS) and Université des Comores; and on Mahé to Direction of Wildlife, Trade and 321 Conservation Section and the Ministry of Environment and Energy of Republic of Seychelles. 322 We thank the Institut Pasteur de Madagascar for their help for sample conservation and 323 exportation; and the National Park and Conservation on Mauritius for authorization for 324 sampling. We are grateful to Association Naturalistes Environnement et Patrimoine de 325 Mayotte in helping to locate some bat roost sites on Mayotte, and to COOPADEM, which 326 provided the field group laboratory space. We are grateful to Dr. Rajendraprasad Sookhareea 327 for his technical assistance in sampling bats on Mauritius. We thank the Secretary of State for 328 Environment, Food and Rural Affairs, London through the Animal and Plant Health Agency 329 of New Haw, Addlestone, Surrey, representing the European Virus Archive for the supply of 330 EBLV-1, EBLV-2, and ABLV.

- 331 This work was supported by funding from CRVOI and from the European Regional
- 332 Development Funds FEDER POCT "Pathogènes associés à la Faune Sauvage Océan Indien"
- project (N° 31189), and FEDER-POCT ParamyxOI project (N° 33857), as well as the
- 334 National Research Foundation, South Africa. Sampling on Mayotte was supported by
- 335 ECOSAN/CNRS-INEE "Batman". B. Ramasindrazana's postdoctoral fellowship at CRVOI
- was funded by "RUN-Emerge, European Commission FP7 Regpot Capacity program", the
- 337 "Fonds de Coopération régionale, Prefecture de La Réunion", and The Field Museum of
- 338 Natural History, Chicago, through the Dr. Ralph and Marian Falk Medical Research Trust.
- 339 Contributions
- 340 SMG, WM, and KD, conceived and designed the experiments. JM performed the
- experiments. JM and KD analyzed the data. SMG, WM, SMC, BR, EL, and MT contributed
- reagents/materials/analysis tools. JM, SMG, WM, SMC, HP, and KD and contributed to the
- writing of the manuscript.
- 344 All authors reviewed the manuscript

### 345 Competing interests

346 The authors declare no conflict of interest.

#### 347 References

- 1. Bourhy H, Kissi B, Tordo N. Molecular diversity of the *Lyssavirus* genus. Virology. 1993
- 349 May; 194(1):70-81.
- 350 2. International Committee on Taxonomy of Viruses (ICTV).
- 351 ictvonline.org/virustaxonomy.asp
- 352 3. Banyard AC, Evans JS, Luo TR, Fooks AR. Lyssaviruses and bats: Emergence and
- 353 zoonotic threat. Viruses. 2014 Aug 4; 6(8):2974-90.

- 4. Badrane H, Bahloul C, Perrin P, Tordo N. Evidence of two Lyssavirus phylogroups with
- distinct pathogenicity and immunogenicity. J Virol. 2001 Apr; 75(7):3268-76.
- 5. Kgaladi J, Wright N, Coertse J, Markotter W, Marston D, Fooks AR, et al. Diversity and
- epidemiology of *Mokola virus*. PLoS Negl Trop Dis. 2013 Oct 24; 7(10):e2511.
- 6. Marston DA, Horton DL, Ngeleja C, Hampson K, McElhinney LM, Banyard AC, et al.
- 359 *Ikoma Lyssavirus*, highly divergent novel *Lyssavirus* in African civet. Emerg Infect Dis.
- 360 2012 Apr; 18(4):664-7.
- 7. Hampson K, Coudeville L, Lembo T, Sambo M, Kieffer A, Attlan M, et al. Estimating the
- 362 global burden of endemic canine rabies. Global Alliance for Rabies Control Partners for
- Rabies Prevention. PLoS Negl Trop Dis. 2015 May; 9(5):e0003786.
- 8. Knobel DL, Cleaveland S, Coleman PG, Fèvre EM, Meltzer MI, Miranda ME, et al. Re-
- evaluating the burden of rabies in Africa and Asia. Bull World Health Organ. 2005 May;
- 366 83(5):360-8. Epub 2005 Jun 24.
- 9. Condori-Condori RE, Streicker DG, Cabezas-Sanchez C, Velasco-Villa A. Enzootic and
- 368 epizootic rabies associated with vampire bats, Peru. Emerg Infect Dis. 2013; 19(9).
- 369 10. da Rosa ES, Kotait I, Barbosa TF, Carrieri ML, Brandão PE, Pinheiro AS, et al. Bat-
- transmitted human rabies outbreaks, Brazilian Amazon. Emerg Infect Dis. 2006 Aug;
- 371 12(8):1197-202.
- 372 11. Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GA, Kent J. Biodiversity hotspots
- for conservation priorities. Nature. 2000 Feb 24; 403(6772):853-8.
- 12. Cheke, A. The timing of arrival of humans and their commensal animals on western
- Indian Ocean oceanic islands. Phelsuma. 2010; 18:38-69.
- 376 13. Goodman, S.M. The bats of Madagascar. [Les chauves-souris de Madagascar] Association
- 377 Vahatra, Antananarivo. 2011; p. 129.

- 378 14. Goodman SM, Ramasindrazana B. Description of a new species of the *Miniopterus*
- 379 *aelleni* group (Chiroptera: Miniopteridae) from upland areas of central and northern
- 380 Madagascar. Zootaxa. 2015 Mar 23; 3936(4):538-58.
- 381 15. Andriamandimby SF, Héraud JM, Ramiandrasoa R, Ratsitorahina M, Rasambainarivo JH,
- Dacheux L, et al. Surveillance and control of rabies in La Réunion, Mayotte, and
- 383 Madagascar. Vet Res. 2013 Sep 9; 44:77.
- 384 16. Reynes JM, Andriamandimby SF, Razafitrimo GM, Razainirina J, Jeanmaire EM, Bourhy
- 385 H, et al. Laboratory surveillance of rabies in humans, domestic animals, and bats in
- Madagascar from 2005 to 2010. Adv Prev Med. 2011; 2011:727821. Epub 2011 Aug 21.
- 387 17. Gannon WL, Sikes RS, American Society of Mammalogists. Guidelines of the American
- 388 Society of Mammalogists for the use of wild mammals in research. J. Mammal. 2007
- 389 June; 88(3):809-23.
- 390 18. Goodman S.M. and Raherilalao MJ. Atlas of selected land vertebrates of Madagascar
- 391 [Atlas d'une sélection de vertébrés terrestres de Madagascar] Association Vahatra,
- 392 Antananariyo. 2013; 290 pp.
- 393 19. Racey PA. Reproductive assessment in bats. In: Kunz TH, editor. Ecological and
- behavioral methods for the study of bats. Washington, D. C.: Smithsonian Institution
- 395 Press. 1988; pp. 31-45.
- 396 20. Anthony ELP. Age determination in bats. In: Kunz TH, editor. Ecological and behavioral
- methods for the study of bats. Washington, DC.; Smithsonian Institution Press. 1988; pp.
- 398 47-58.
- 399 21. Kuzmin IV, Niezgoda M, Franka R, Agwanda B, Markotter W, Beagley JC, et al. Lagos
- 400 Bat Virus in Kenya. J Clin Microbiol. 2008 Apr; 46(4):1451-61. Epub 2008 Feb 27.
- 401 22. Coertse J, Markotter W, Nel LH. Demonstration of African lyssavirus RNA with real-time
- 402 polymerase chain reaction. In Rupprecht, C & Nagajaran, T, editors. Current laboratory

- 403 techniques in rabies diagnosis, research, and prevention, Volume 1. Amsterdam:
- 404 Academic Press. 2014; pp. 63-73.
- 405 23. Goodman SM, Jansen van Vuuren B, Ratrimomanarivo F, Probst JM, Bowie RCK.
- 406 Specific status of populations in the Mascarene Islands referred to *Mormopterus*
- 407 *acetabulosus* (Chiroptera: Molossidae), with description of a new species. J. Mammal.
- 408 2008 Oct; 89(5):1316-27.
- 409 24. Markotter W, Monadjem A, Nel LH. Antibodies against Duvenhage Virus in
- insectivorous bats in Swaziland. J Wildl Dis. 2013 Oct; 49(4):1000-3.
- 411 25. Kuzmin IV, Turmelle AS, Agwanda B, Markotter W, Niezgoda M, Breiman RF, et al.
- 412 Commerson's leaf-nosed bat (*Hipposideros commersoni*) is the likely reservoir of Shimoni
- 413 bat virus. Vector Borne Zoonotic Dis. 2011 Nov; 11(11):1465-70. Epub 2011 Aug 25.
- 414 26. Kuzmin IV, Niezgoda M, Franka R, Agwanda B, Markotter W, Beagley JC, et al. Possible
- emergence of West Caucasian bat virus in Africa. Emerg Infect Dis. 2008 Dec;
- 416 14(12):1887-9.
- 417 27. Hayman DT, Fooks AR, Horton D, Suu-Ire R, Breed AC, Cunningham AA, et al.
- 418 Antibodies against *Lagos Bat Virus* in Megachiroptera from West Africa. Emerg Infect
- 419 Dis. 2008 Jun; 14(6):926-8.
- 420 28. Kemp GE, Causey OR, Moore DL, Odelola A, Fabiyi A. Mokola virus. Further studies on
- 421 IbAn 27377, a new rabies-related etiologic agent of zoonosis in Nigeria. Am J Trop Med
- 422 Hyg. 1972 May; 21(3):356-9.
- 423 29. Nguyen AT, Nguyen TT, Noguchi A, Nguyen DV, Ngo GC, Thong VD, et al. Bat
- 424 lyssaviruses, Northern Vietnam. Emerg Infect Dis. 2014 Jan; 20(1):161-3.
- 425 30. Reynes JM, Molia S, Audry L, Hout S, Ngin S, Walston J, et al. Serologic evidence of
- 426 Lyssavirus infection in bats, Cambodia. Emerg Infect Dis. 2004 Dec; 10(12):2231-4.

- 427 31. Arguin PM, Murray-Lillibridge K, Miranda ME, Smith JS, Calaor AB, Rupprecht CE.
- Serologic evidence of *Lyssavirus* infections among bats, the Philippines. Emerg Infect
- 429 Dis. 2002 Mar; 8(3):258-62.
- 430 32. Turmelle AS, Allen LC, Jackson FR, Kunz TH, Rupprecht CE, McCracken GF. Ecology
- of rabies virus exposure in colonies of Brazilian free-tailed bats (*Tadarida brasiliensis*) at
- natural and man-made roosts in Texas. Vector Borne Zoonotic Dis. 2010 Mar; 10(2):165-
- 433 75.
- 434 33. Picard-Meyer E, Dubourg-Savage MJ, Arthur L, Barataud M, Bécu D, Bracco S, et al.
- 435 Active surveillance of bat rabies in France: A 5-year study (2004–2009). Vet Microbiol.
- 436 2011 Aug 5; 151(3-4):390-5. Epub 2011 Apr 12.
- 437 34. Dietrich M, Wilkinson DA, Benlali A, Lagadec E, Ramasindrazana B, Dellagi K, et al.
- 438 Leptospira and paramyxovirus infection dynamics in a bat maternity enlightens pathogen
- maintenance in wildlife. Environ Microbiol. E-pub 2015 Jan 11.
- 35. O'Shea TJ, Bowen RA, Stanley TR, Shankar V, Rupprecht CE. Variability in
- 441 seroprevalence of rabies virus neutralizing antibodies and associated factors in a Colorado
- population of big brown bats (*Eptesicus fuscus*).. PLoS One. 2014 Jan 22; 9(1):e86261.
- 36. Markotter W, Van Eeden C, Kuzmin IV, Rupprecht CE, Paweska JT, Swanepoel R, et al.
- Epidemiology and pathogenicity of African bat lyssaviruses. Dev Biol (Basel). 2008;
- 445 131:317-25.
- 446 37. Dietrich M, Wilkinson DA, Soarimalala V, Goodman SM, Dellagi K, Tortosa P.
- 447 Diversification of an emerging pathogen in a biodiversity hotspot: *Leptospira* in endemic
- small mammals of Madagascar. Mol Ecol. 2014 Jun; 23(11):2783-96. Epub 2014 May 17.
- 449 38. Banyard AC, Johnson N, Voller K, Hicks D, Nunez A, Hartley M, et al. Repeated
- detection of European bat lyssavirus type 2 in dead bats found at a single roost site in the
- 451 UK. Arch Virol. 2009; 154(11):1847-50.

452	39. van Thiel PP, de Bie RM, Effimov F, Tepaske R, Zaaijer HL, van Doornum GJ, et al.
453	Fatal human rabies due to Duvenhage virus from a bat in Kenya: Failure of treatment with
454	coma-induction, ketamine, and antiviral drugs. PLoS Negl Trop Dis. 2009 Jul 28;
455	3(7):e428.
456	40. Van Gucht S, Verlinde R, Colyn J, Vanderpas J, Vanhoof R, Roels S, et al. Favourable
457	outcome in a patient bitten by a rabid bat infected with the European bat lyssavirus-1.
458	Acta Clin Belg. 2013 Jan-Feb; 68(1):54-8.
459	
460	Corresponding authors:
461	Koussay Dellagi: Université de La Réunion, UMR PIMIT «Processus Infectieux en Milieu
462	Insulaire Tropical», INSERM U1187, CNRS 9192, IRD 249.Plateforme de Recherche
463	CYROI, Saint Denis, La Réunion, France. koussay.dellagi@ird.fr; tel: +33 610618184
464	Wanda Markotter: Department of Microbiology and Plant Pathology, Faculty of Natural and
465	Agricultural Sciences, University of Pretoria, Pretoria 0001, South Africa.
466	wanda.markotter@up.ac.za; tel: + 27 828246356; fax: + 27 12420 3266;
467	
468	
469	
470	
471	
472	Figure 1. Geographic distribution of bats collected on the SWIO islands of Anjouan
473	(Union of Comoros), Mayotte (France), Madagascar, Reunion Island (France)
474	Mauritius, and Mahé (Seychelles Archipelago). n: number of bats sampled.
475	

476 Figure S1-S6. Lyssaviruses neutralized by sera from bats captured on Anjouan (S1), Mayotte (S2), Reunion Island (S3) Mauritius (S4), Mahé (S5), and Madagascar (S6). 477 478 Lyssaviruses neutralized by sera from bats captured on Anjouan (S1), Mayotte (S2), Reunion 479 Island (S3) Mauritius (S4), Mahé (S5), and Madagascar (S6). Abbreviations in the pie 480 diagrams correspond to each lyssaviruses (see Methods section). Abbreviations next to 481 illustrated islands indicate names of capture sites (e.g. MGJ for "Mangajou") reported in 482 Table S1A and Table S1B. In the pie diagram, n indicates at the numerator the number of sites 483 with bats neutralizing or cross neutralizing (e.g. DUVV + EBLV-1) lyssaviruses and at the 484 denominator the number of investigated sites; next to illustrated islands, n in the numerator 485 indicates the number of individuals tested positive for lyssaviruses and in the denominator the 486 number of individuals tested. For Madagascar, the six black squares indicate the provincial capitals. 487

**Table 1.** Seroreactivity of bats from southwestern Indian Ocean islands as assessed by RFFIT using a panel of challenge lyssaviruses.

		zation cl	ation challenge viruses			Cross neutralization			
	Positive/teste	d CVS-11	DUVV	EBLV-1	ABLV	LBV	CVS-11+ EBLV-1	CVS-11+	EBLV-1
Anjouan	4/21	0/8	2/8	0/0	0/0	2/21	0/0	0/0	0/0
Molossidae									
Chaerephon pusillus	4/19	0/8	2/7	0/0	0/0	2/19	0/0	0/0	0/0
/liniopteridae									
Miniopterus griveaudi	0/2	0/0	0/1	0/0	0/0	0/2	0/0	0/0	0/0
Mayotte	5/22	0/22	2/22	0/22	0/0	3/22	0/22	0/0	0/22
Molossidae				- 1-					
Chaerephon pusillus	2/3	0/3	1/3	0/3	0/0	1/3	0/3	0/0	0/3
Pteropodidae									
Pteropus seychellensis	3/19	0/19	1/19	0/19	0/0	2/19	0/19	0/0	0/19
Madagascar	84/301	0/277	54/286	1/70	0/7	23/301	0/70	1/7	5/70
lipposideridae	2/6	0/6	1/6	1/4	0/0	0/6	0/4	0/0	1/4
Hipposideros commersoni	3/6	0/6	1/6	1/4	0/0	0/6	0/4	0/0	1/4
Miniopteridae	4147	0/45	A14 F	0/4	0/0	0/47	0/4	0/0	0/4
Miniopterus cf. ambohitrensis		0/15	4/15	0/4	0/0	0/17	0/4	0/0	0/4
Miniopterus gleni	0/4	0/4	0/4	0/0	0/0	0/4	0/0	0/0	0/0
Miniopterus griveaudi	12/33	0/23	10/29	0/6	0/0	2/33	0/6	0/0	0/6
Miniopterus mahafaliensis	0/11	0/9	0/11	0/0	0/0	0/11	0/0	0/0	0/0
Miniopterus sororculus	0/3	0/3	0/2	0/1	0/0	0/3	0/1	0/0	0/1
Molossidae				2/2	- 1-	244	2.12	0.10	0.10
Chaerephon atsinanana	0/16	0/16	0/15	0/0	0/0	0/16	0/0	0/0	0/0
Chaerephon leucogaster	10/28	0/26	9/28	0/6	0/0	0/28	0/6	0/0	1/6
Mops leucostigma	2/20	0/20	0/19	0/4	0/2	2/20	0/4	0/2	0/4
Mops midas	0/7	0/7	0/7	0/0	0/0	0/7	0/0	0/0	0/0
Mormopterus jugularis	20/58	0/58	17/58	0/8	0/0	3/58	0/8	0/0	0/8
Otomops madagascariensis	7/20	0/20	4/18	0/11	0/0	3/20	0/11	0/0	0/11
Pteropodidae	0.10	0.10	0.10	0/4	0.10	0.10	0/4	0.10	0.14
Eidolon dupreanum	0/9	0/9	0/9	0/1	0/0	0/9	0/1	0/0	0/1
Pteropus rufus	8/12	0/10	5/11	0/5	0/0	0/12	0/5	0/0	3/5
Rousettus madagascariensis	12/35	0/33	4/33	0/16	0/5	7/35	0/16	1/5	0/16
Rhinonycteridae		0.77	0111	0.10	0.40	0///	0.40	0.10	0.10
riaenops menamena	2/11	0/7	0/11	0/2	0/0	2/11	0/2	0/0	0/2
/espertilionidae	,,,,	0.11.1	0/10	0.10	0/6	.,,,	0/0	0/0	0/0
Myotis goudoti	4/11	0/11	0/10	0/2	0/0	4/11	0/2	0/0	0/2
Reunion Island	52/121	0/119	14/117	9/82	0/0	3/121	0/80	0/0	26/82
∕lolossidae ∕lormopterus francoismoutoui	i 52/121	0/119	14/117	9/82	0/0	3/121	0/0	0/0	26/82
Mauritius (	28/67	0/62	19/67	2/29	0/5	7/67	0/29	0/5	0/29
Molossidae	_5,0,	J. 02						J	J
formopterus acetabulosus	12/31	0/29	6/31	1/8	0/0	5/31	0/8	0/0	0/8
,									
Pteropodidae Pteropus niger	16/36	0/33	13/36	1/21	0/5	2/36	0/21	0/5	0/21
<b>lahé</b> Iteropodidae	13/40	0/40	6/40	1/17	0/2	4/40	1/17	0/2	1/17
Pteropus seychellensis	13/40	0/40	6/40	1/17	0/2	4/40	1/17	0/2	1/17
Succeed Total	400/570	0/500	07/540	40/000	0/4.4	40/570	4/040	4/4.4	20/200
Grand Total	186/572	0/528	97/540	13/220	0/14	42/572	1/218	1/14	32/220

Fig 1. Geographic distribution of bats collected on the SWIO islands of Anjouan (Union of Comoros), Mayotte (France), Madagascar, Reunion Island (France), Mauritius, and Mahé (Seychelles Archipelago). n: number of bats sampled.




Fig S1.




Fig S2.




Fig S3.




Fig S4.





Fig S5.






Table S1. Sites names and corresponding abbreviations

Abbreviations	Island	Site name
AMB	Madagascar	Grotte d'Ambanilia, 3.7 km SSE Sarodrano
AMBB	Madagascar	Parc National d'Ankarana, Ambahibe Cave, 2 km W Mahamasina
AMBO	Madagascar	Ambovondramanesy village near Berivotra, along RN4
AMBT	Madagascar	Réserve Spéciale d'Ambohitantely, Grotte des Chauves-souris
AMBT2	Madagascar	Réserve Spéciale d'Ambohitantely, début Sentier Touristique
ANDA1	Madagascar	CEG Andasibe
ANDOK	Madagascar	Parc National d'Ankarana, Grotte du troisiéme Canyon, along Andokotokana River
ANDRF	Madagascar	Parc National d'Ankarana, Grotte d'Andrafiabe, 3.3 ESE Andrafiabe
ANJHB	Madagascar	Grotte d'Anjohibe, 3.7 km NE Antanamarina
ANJHK1	Madagascar	Grotte d'Anjohikely (south entrance), 1.5 km NE Antanamarina
ANJKK	Madagascar	Parc National de Bemaraha, Anjohikinakina, 15.5 km N Bekopaka
ANJOZ2	Madagascar	Ambohibeloma, 3.2 km W. Anjozorobe
ANK	Madagascar	Commune rurale d'Ankily, west of Ihosy, off RN7
ANKPK	Madagascar	Ankapoka
ANTAN	Madagascar	Antanandava, Eglise FLM, 5.8 km NE Beroboka-Sud
ANTNM	Madagascar	Cascade d'Antanamarina
BEK	Madagascar	Parc National d'Isalo, Grotte de Bekapity
BELO	Madagascar	Belo Tsiribihina, central hospital
BET	Madagascar	Betioky-Sud, new Lutherian church
BRDPC	Mauritius	Black River District, Palma Cave
BVL	Mahé	Beau Vallon
CCC	Mayotte	Carrefour Chiconi
CGS	Madagascar	Parc National d'Ankarana, Grotte des Chauve-souris, 3 km NW Mahamasina
COA	Comoros	Anjouan, Collège d'Ouani
CROCO	Madagascar	Parc National d'Ankarana, 2.2 km ESE Amboandriky, Grotte d'Ambatoharanana (Crocodile Cave)
CSC	Mauritius	Black River District, Cascavelle

Table S1. Continued

Abbreviation	Island	Site name
СТВ	Mauritius	Rivière du Rempart District, Caverne Trois Bras
CTC	Mauritius	Moka District, Camp Thorel, Camp Thorel Cave
EBP	Reunion Island	Eglise de Bras-Panon
EPC	Comoros	Anjouan, Ecole primaire de Chirove
FRL	Mahé	Fairy Land
FYM	Mauritius	Flacq District, northern slope Fayence Mountain
GBK	Comoros	Anjouan, Grotte de Buemokolo
GTB	Reunion Island	Grotte de Trois-Bassin
IHO	Madagascar	Ihosy, Bureau du chef de la Région,
ISALO1	Madagascar	Parc National d'Isalo, 3.8 km NW de Ranohira, along Namaza River
MAH	Madagascar	Mahabo, EPP de Mahabo
MAHA	Madagascar	North of Mahajanga, Petite Plage
MAKI	Madagascar	Grotte de Makis (Mikea), near Hotel la Mangrove, on Toliara-St Augustin Road
MARFA	Madagascar	Marofandilia, Ecole primaire
MGJ	Mayotte	Mangajou
PCR	Mauritius	Black River District, Petite Case Noyale
REE	Mauritius	Grand Port District, Riche en Eau
SAK1	Madagascar	Sakaraha, Direction des Eaux et forêt, Bureau chef de cantonnement
SBP	Mayotte	Sohoa be plage
TANA	Madagascar	Grotte de Tanambao (Bishiko), 0.75 km E St Augustin
TDZ	Mayotte	Tsoundzou
TSMIF	Madagascar	Tsimafana, CEG de Tsimafana
TSP	Madagascar	Parc National de Tsimanampetsotsa, Grotte d'Andranoilovy
VINT	Madagascar	Grotte de Vintane (Vintany), 4.1 km SE Itampolo
VOHI	Madagascar	Vohipoa, CSB II

Text S1. Authorizations for sampling of bats on different western Indian Ocean islands

- 1. Madagascar: Direction du Système des Aires Protégées and Direction Générale de l'Environnement et des Forêts; Madagascar National Parks: Export permit n°194/12/MEF/SG/DGF/DCB.SAP/SCB, 067/12/MEF/SG/DGF/DCB.SAP/SCBSE, and 032/12/MEF/SG/DGF/DCB.SAP/SCBSE. A CITES permit from the Malagasy national authority was issued for tissue export (permit 243C-EA06/MG12) to CRVOI on Reunion Island.
- 2. Reunion Island: Préfecture de La Réunion: Arrêté préfectoral of 11 Février 2013.
- 3. Mayotte: Préfecture de Mayotte: Arrêté préfectoral n°158/DEAL/SEPR/2014.
- Anjouan, Union of Comoros: Centre National de Documentation et de Recherche Scientifique (CNDRS) of Union des Comores: Export permit CNDRS 021/10.
- 5. Mahé, Seychelles Archipelago: Direction of Wildlife, Trade and Conservation Section and the Ministry of Environment and Energy of Republic of Seychelles: Export permits Agreement of 5 March 2014. CITES permit from the Republic of Seychelles was issued for export (permit N°1772) to CRVOI on Reunion Island.
- 6. Mauritius: National Park and Conservation Service for authorization of Mauritius: Memorandum of agreement for the supply of biological material by Government of Mauritius, signed 17 December 2010 and 09 January 2013. CITES permit from the Mauritian national authority was issued for tissue export (permit MU120933) to CRVOI laboratory on Reunion Island.

## 6.4. Conclusion du chapitre 6

Dans cette étude nous avons mis en évidence une circulation sérologique importante des lyssavirus chez les chauves-souris de toutes les îles de la zone du SOOI. Un très grand nombre d'espèces de chauves-souris (plus de 12 espèces insectivores et 2 espèces de frugivores) a été retrouvé séropositif suggérant un large spectre d'hôte pour ces lyssavirus. Les taux d'infection les plus élevés ont été observés à La Réunion, Maurice et Mahé, et des taux les plus faibles à Madagascar, Mayotte et à Anjouan. Les profils de séroneutralisation sont en faveur de la circulation d'au moins deux lyssavirus, l'un appartenant au phylogroupe I (rendant compte des réactivités croisées DUVV, EBLV et ABLV), et l'autre, au phylogroupe II (rendant compte de la réactivité croisée avec LBV).

Cependant, aucune trace d'ARN n'a été mise en évidence chez les chauves-souris séropositives ou séronégatives comme l'indique les résultats de l'analyse moléculaire qRT-PCR sur les cerveaux d'animaux. Cela suggère qu'une exposition antérieur des chauves-souris vis-à-vis d'un lyssavirus du phylogroupe I et / ou du phylogroupe II a eu lieu mais que celui-ci pourrait avoir été éliminé par le système immunitaire ou serait présent mais à très faible concentration. Des travaux complémentaires sont nécessaires, en particulier sur les chauves-souris malades ou morbides ainsi que sur les glandes salivaires pour pouvoir identifier les lyssavirus qui circulent dans la région.

Ces travaux ont été valorisés sous forme d'un article scientifique soumis et de trois présentations orales au congrès international « African Small Mammal Symposium » (ASMS, 2015), et aux Journées scientifiques de la « Fédération Environnement, Biodiversité » et Santé (FedEBS; 2014, Saint-Denis La Réunion), ainsi qu'au Colloque FEDER POCT Biodiversité (2015, Saint-Denis La Réunion).

Annexe 6: Les résultats obtenus sur la sérologie Panvirale nous ont permis de détecter de fortes séroprévalence sen *Paramyxoviridae*, *Bunyaviridae* et *Flaviviridae* chez les chauves-souris, petits mammifères terrestres endémiques et rongeurs. Les séroprévalences en *Cornoraviridae* et *Togaviridae* étaient les plus faibles. Les individus fortement séropositifs appartiennent à la famille des Pteropodidae (chauves-souris frugivores du genre *Eidolon*, *Pteropus* et *Rousettus*).

Enfin, les analyses montrent une plus forte séroprévalence contre les *Rubulavirus*, notamment *Mumps* et *PIV-2*, au sein des individus collectés en particulier chez les chauves-souris frugivores. Cela suggère une circulation active d'un virus putatif proche de ce genre viral en plus des *UMRVs* au sein des colonies de frugivores.

L'ensemble de ces résultats témoigne d'une forte sensibilité des chauves-souris de la zone du SOOI aux virus à ARN potentiellement zoonotiques, et particulièrement, les virus des familles *Paramyxoviridae*, *Flaviviridae* et *Bunyaviridae*.

Ces travaux seront valorisés sous forme d'un article scientifique qui est en cours de soumission.

## Discussion générale

Les déterminants écologiques et les processus évolutifs qui entrent en jeu dans les phénomènes de "host-switch" des agents infectieux zoonotiques issus de la faune sauvage vers les populations humaines et d'animaux domestiques ou de compagnie sont encore mal connus. Etant donné que 80% des virus émergeants qui concernent l'homme proviennent de la faune sauvage (Taylor et al., 2001), une attention croissante leur sont accordés au niveau mondial par les scientifiques et les autorités sanitaires, en raison des enjeux majeurs socio-économiques et de leurs impacts sur les politiques publiques de santés humaine et vétérinaire. Les graves épidémies toujours en cours provoquées par le virus *Ebola* en Afrique de l'Ouest (CDC, 2015) ou du MERS-CoV au Moyen Orient (CDC, 2014), ainsi que les flambées épidémiques de la PPR (Ishag et al., 2015) qui dévastent les cheptels des petits ruminants d'Asie et d'Afrique attestent bien de l'urgence de la situation (OMS, OIE, FAO). Anticiper la survenue de telles émergences infectieuses et comprendre les tenants et les aboutissants des facteurs qui y concourent passe par l'exploration systématique des réservoirs sauvages impliquant à la fois l'étude de leur l'écologie et des mécanismes évolutifs des agents infectieux qui leurs sont associés (Morse et al., 2012).

Mes travaux de thèse s'inscrivent dans le cadre de l'approche éco-systémique conduite par le CRVOI sur les maladies infectieuses émergentes et font suite aux premières informations obtenues lors de l'exploration de certaines îles du SOOI (Madagascar, Mayotte et La Réunion) sur certains réservoirs animaux potentiels tels que les chauves-souris (Wilkinson et al., 2012). Ces travaux avaient mis en évidence une diversité importante de nouveaux paramyxovirus, avec des prévalences élevées en comparaison à d'autres études (Kurth et al., 2012; Sasaki et al., 2012).

Face à ces observations saisissantes, mes travaux de thèse ont abordé la question de la dynamique de transmission virale et ont tenté de mieux caractériser ces virus potentiellement zoonotiques circulant parmi les micromammifères terrestres et volants de la zone du SOOI. Les informations préliminaires laissaient penser que de multiples échanges viraux inter-îles, surviennent entre des hôtes très divers dans la région du SOOI.

Cette région multi-insulaire est singulière de par sa grande biodiversité et un fort endémisme en lien avec des particularités écologiques (Tortosa et al., 2012).

Nous avons ainsi pu mettre en évidence plusieurs résultats originaux : (i) une circulation importante de nouveaux paramyxovirus (*UMRVs*), appartenant probablement à un nouveau genre viral phylogénétiquement très apparenté aux *Morbillivirus*, ayant une aptitude à infecter un large spectre d'hôtes, appuyant ainsi le rôle de réservoir de ces mammifère, incluant Chiroptera, Rodentia et Afrosoricidae ; (ii) la co-divergence génétique des *UMRVs* ne peut pas être le résultat d'un mécanisme de co-spéciation des hôtes, mais résulte principalement du mécanisme du *host-Switch* en terme macroévolutif chez les chauves-souris ; (iii) des facteurs écologiques jouent un rôle important dans la dynamique de transmission virale ; (iv) enfin de façon plus générale, l'analyse sérologique révèle une intense circulation régionale de lyssavirus de chauves-souris dans un très grand nombre d'espèces, virus qui se rattachent très probablement pour l'un au phylogroupe I et pour l'autre au phylogroupe II. Il n'est d'ailleurs pas impossible, considérant l'isolement relatif des îles du SOOI, que l'un ou l'autre se révèlent *in fine* de nouveaux membres de ce genre viral et qu'ils soient spécifiques des espèces endémiques de chiroptères retrouvées uniquement dans le SOOI.

## Les processus évolutifs mis en jeu

#### Macro-évolution

Une caractéristique remarquable commune aux deux modèles viraux, paramyxovirus et lyssavirus, est le large spectre d'hôtes. En effet, plus de 15 espèces de chauves-souris (6 familles) et au moins 10 espèces de petits mammifères terrestres endémiques (2 familles) sont infectées par les *UMRVs* et 14 espèces de chauves-souris (6 familles) sont concernées par un putatif lyssavirus. Ces résultats confirment le rôle des chauves-souris et petits mammifères terrestres en tant que réservoir viral majeur (Calisher et al., 2006 ; Meerburg et al., 2009 ; Drexler et al., 2012 ; Banyard et al., 2014). A ce stade de nos investigations, certaines questions méritent d'être posées : Quelles sont les raisons qui permettent à des virus aussi génétiquement divers, d'infecter des hôtes si diversifiés ? Existe-t-il à l'origine d'un tel spectre, un mécanisme évolutif global au niveau viral, ou est-ce la conséquence de facteurs écologiques ? Le moteur macro-évolutif (à savoir le mécanisme de *host-switch*) que nous avons démontré chez les paramyxovirus de chauves-souris répond en grande partie à cette question.

Ce même mécanisme de co-divergence virale associée à des évènements de sauts d'espèces a été rapporté chez d'autres agents zoonotiques viraux issus de la faune sauvage (Ramsden et al., 2009; Coulibaly-N'Golo et al., 2011), mais aussi pour les Paramyxovirus infectant l'homme (Kitchen et al., 2011).

Au cours de nos travaux, nous avons tiré profit de la diversité d'espèces de chauves-souris présentes à Madagascar afin de déterminer si la diversité des paramyxovirus était issue de celle des hôtes (spéciation d'hôte entraînant une spéciation du virus) ou serait propre aux paramyxovirus eux-mêmes. Ainsi, en comparant 7 familles et 31 espèces de chauves-souris Malgaches différentes, nous avons montré que les multiples phénomènes de sauts d'espèces prennent le pas sur l'hypothèse d'une macro-évolution directement liée à une co-spéciation de l'hôte. Ce dernier mécanisme a longtemps (et le reste encore pour beaucoup d'auteurs) été avancé comme le principal mécanisme macro-évolutif en œuvre chez de nombreux modèles viraux (de Vienne et al., 2013). Il est important de mentionner que la plupart des algorithmes de co-phylogénie ont été mis au point sur des modèles bactériens et de facto ne sont pas ou mal adaptés, à l'analyse des modalités évolutives chez les virus ARN. Ces logiciels s'appuient sur des algorithmes essentiellement basés sur des processus de co-spéciation (Merkle & Middendorf, 2005), et nous pensons que c'est l'une des raisons pour lesquels ils aboutissent à une surreprésentation de ces derniers: derrière une apparente congruence phylogénétique (phénomènes d'associations hôtes-virus), entre des agents infectieux et leurs hôtes, il ne pouvait s'agir que d'une co-divergence virale liée à la co-spéciation de l'hôte (Merkle et al., 2010). En comparaison, si on se réfère à des modèles bactériens, comme chez les Leptospires, ou encore sur des modèles parasitaires plus complexes, tels que les Hémoparasites (plasmodium et filaire), modèles développés par mes collègues au sein du CRVOI à partir du même échantillonnage d'hôtes (chauves-souris) (respectivement thèse de Yann Gomard et post-doctorat de Beza Ramasindrazana), les très fortes associations entre bactéries/hémoparasites et leurs hôtes respectifs dans leurs cas, révèlent que les relations évolutives hôtes/leptospires/hémoparasites sont dominées par le mécanisme de co-spéciation des hôtes, entrainant à chaque étape de spéciation, la co-divergence des agents infectieux correspondants. Quelque part, une telle discordance entre ces modèles était attendue s'agissant de bactéries et de parasites qui n'élaborent pas les mêmes stratégies du point de vue évolutif. Ces agents infectieux diffèrent en termes de structure, de cycles de vie, de mode de réplication/multiplication, et surtout, au niveau de leur horloge moléculaire respective.

Les bactéries et les parasites ont des taux d'évolution très lents (de l'ordre de  $10^{-9}$  à  $10^{-6}$  substitutions/site/an) comparés à ceux des virus à ARN (de l'ordre de  $10^{-5}$  -  $10^{-3}$  substitutions/site/an) (Holland et al., 1982), Ces derniers, accumulent des mutations de manière beaucoup plus importante en raison de leur taux de réplication élevé et du fait de l'absence de l'activité correctrice 3'-5' de leur ARN polymérase ARN-dépendante (Duffy et al., 2008) produisant ainsi des populations virales hétérogènes complexes (quasi-espèces) (Holmes & Moya, 2002 ; Domingo et al., 2005).

#### Microévolution

D'après certains auteurs, la variabilité génétique qui caractérise les virus à ARN, serait intimement liée aux mécanismes micro-évolutifs, c'est-à-dire aux relations d'adaptations complexes qui s'établissent lors des interactions hôtes-pathogènes au sein de communautés d'espèces, par exemple (Page, 2003; Jackson et al., 2004; Woolhouse et al., 2001). Elle pourrait expliquer la diversité virale observée au sein des UMRVs. Une étude récente a par ailleurs incriminé la plasticité des génomes des virus à ARN comme pouvant être à l'origine du « host-switch » (Kreuder Johnson et al., 2015). Comme décrit dans le chapitre I, après chaque « host-switch » viral, il est nécessaire pour le virus de s'adapter à son nouvel hôte, au risque de connaître l'extinction. Les façons de s'adapter à son nouvel environnement sont diverses tels que celui de varier (variant de mutation) ou encore de procéder à des recombinaisons, permettant ainsi d'obtenir des variants qui seraient plus à même de reconnaitre et de se lier à des récepteurs "alternatifs" permettant des infections de nouveaux types cellulaires, ou encore modifier la pathogénicité (Baranowski et al., 2003). Les dispositifs ainsi mis en jeu peuvent permettre au virus d'échapper ou de contourner les défenses immunitaires de l'hôte nouvellement infecté (échappement viral) (Ciurea et al., 2000). Quelles sont ces mutations et à quelles fréquences apparaissent-elles ? Sont-elles toutes les mêmes pour tous les virus ? Sur quelle partie précise du génome se produisentelles? Comment les paramyxovirus infectent ils et se reproduisent ils dans de multiples hôtes? Ya t'il en définitive un phénomène de persistance virale? En effet n'ayant pu conduire d'étude longitudinale sur une cohorte de chauves-souris ou de rongeurs sauvages, nous ne pouvons formellement répondre à cette question.

Les récepteurs étant des éléments essentiels dans les premières étapes d'infection et, les paramyxovirus infectant un très grand nombre d'espèces différentes, ils devraient reconnaitre un récepteur "ubiquitaire" chez les espèces communes concernées qui reste à déterminer pour ce qui concerne les *UMRVs*. Il convient de signaler que la conservation des récepteurs et voies cellulaires entre chauves-souris et différents mammifères est suspectée comme l'élément moteur de la transgression de la barrière d'espèce facilitant la transmission de virus à d'autres taxons (Calisher et al., 2006).

Comme décrit dans la littérature (Anishchenko et al., 2006; Orton et al., 2013), il serait très intéressant de caractériser ces mutations et ces récepteurs cellulaires ubiquitaires qui vont permettre la persistance et l'adaptation virale. Le séquençage complet des génomes des *UMRVs* qui est en cours d'analyse devrait en partie nous aider à répondre à ces interrogations. Quoiqu'il en soit, de tels mécanismes pourraient probablement expliquer la grande diversité virale des *UMRVs*, laquelle résulterait, d'une part, de leur longue histoire évolutive, jalonnée au niveau macro-évolutif, par de multiples sauts d'espèces, et sans cesse compensée, au niveau micro-évolutif, par des mécanismes d'adaptation au sein des nouveaux hôtes.

On sait que les hôtes phylogénétiquement proches entres eux sont plus à risque de hostswitch, notamment, du fait de la forte conservation entre leurs récepteurs cellulaires de surface (Longdon et al., 2011). C'est le cas pour le virus rabique et pour les hantavirus, où des études montrent que le host-switch viral se produit préférentiellement entre des espèces de chauvessouris et rongeurs phylogénétiquement très proches (Ramsden et al., 2009; Steicker et al., 2010). Nos résultats (Wilkinson et al., 2014) démontrent qu'il est aussi possible que des paramyxovirus phylogénétiquement très proches, peuvent infecter des phylogénétiquement éloignés: dynamique infectieuse et échanges viraux inter-ordres, Chiroptera et Rodentia. Un tel phénomène a déjà était décrit par l'équipe de Guo et al. (2014) qui démontré que le phénomène de « host-switch » s'est produit depuis la chauves-souris, décrit comme le réservoir principal des hantavirus dans son étude, vers les rongeurs.

Le rat noir (*Rattus rattus*) et le rat brun (*Rattus norvegicus*), originaires d'Inde et d'Asie rspectivement, se sont ensuite rapidement dispersés dans le monde (Aplin et al., 2003). Ces espèces sont réservoirs de très nombreux virus et surtout décrits comme disséminateurs de parasites et de virus (Wilkinson et al., 2014; Firth et al., 2014; Morand et al., 2015).

On sait notamment qu'à Madagascar, les rongeurs introduits sont présents à proximité des entrées de grottes de chauves-souris. Il est envisageable qu'une transmission virale chauvessouris / rongeurs ait lieu et que celle-ci passe par l'ingestion d'eau ou d'aliments contaminés par l'urine ou les fèces de chauves-souris par exemple. Nos analyses ont également montré un échange de paramyxovirus entre l'Afrique et le SOOI (Wilkinson et al., 2014). Comme décrit dans le chapitre I, Rattus rattus est connu pour être une espèce invasive largement distribuée à travers le monde (Lund et al., 1994; Aplin et al., 2003). Les transports maritimes ont contribué en grande partie à sa dissémination mondiale, et entre pays très éloignés (Bonnefoy et al., 2008). De fait, les échanges commerciaux des îles du SOOI se produisent avec l'Asie, l'Europe ou l'Afrique. Par ailleurs, l'étude conduite sur les paramyxovirus des rongeurs indigènes à la Tunisie, en Afrique du Nord (Psammomys obesus, Meriones shawi et Ctenodactilus gundii) a montré leur proximité phylogénétique avec des paramyxovirus des Iles du SOOI (Ghawar et al., soumis). De plus nous avons pu retracer l'histoire évolutive de ces paramyxovirus, montrant qu'ils seraient originaires d'Afrique australe, et qu'ils ont été disséminés très vraisemblablement par les rongeurs introduits Rattus. Ainsi R. rattus aurait joué le rôle de pont épidémiologique entre les différents hôtes, mais également entre toutes les régions du globe.

Par ailleurs, en plus des *host-switch* fréquemment observés, nous avons mis en évidence quelques cas d'associations entre hôtes et *UMRVs*. Du fait que les virus évoluent rapidement, le système immunitaire de l'hôte doit lui aussi s'adapter, notamment par l'évolution rapide simultanée des gènes de ses récepteurs à l'antigène lui permettant de moduler continuellement sa réponse face à ses « cibles changeantes ». Ces situations conflictuelles créent une course sans fin aux armements entre l'hôte et le virus (van Valen, 1973). Il s'agit en partie de l'hypothèse évolutive de la "Reine Rouge" dans laquelle l'évolution des gènes de l'immunité doit toujours rester au même niveau que celle des virus. L'alternative adoptée par certains virus, serait que, sous certaines pressions de sélection du système immunitaire de l'hôte, certains virus vont être "séquestrés" au niveau de différents compartiments (niches) à l'intérieur d'un hôte (Kitchen et al., 2011) et s'y spécialiser, générant une association entre l'hôte et le virus extrêmement forte et étroite. Ce type d'adaptation peut conduire le pathogène vers une "hyperspécialisation" ou une sanctuarisation dans un tissu ou un type cellulaire donné. Ce phénomène pourrait représenter une barrière évolutive moins "permissive" que celle qui ferait suite à l'infection par *host-switch* d'un pathogène pour un nouvel hôte.

#### Facteurs éco-épidémiologique

De nombreuses études ont montré la présence de paramyxovirus appartenant au genre *Henipavirus* chez les chauves-souris frugivores, principalement du genre *Pteropus* ou *Eidolon* (Field et al., 2012 ; Drexler et al., 2009 ; Yadav et al., 2015 ; Chua et al., 2000). Nos résultats sont différents de ces études puisque nous décrivons une très grande proportion de chauves-souris insectivores infectées, et une très faible proportion de chauve-souris frugivores infectées, et pour celles qui le sont, l'infection est toujours due par un virus appartenant au genre des *UMRVs*. Cette restriction à ce genre unique doit cependant être nuancée dans la mesure ou les autres genres de paramyxovirus (*Pneumovirus*, *Avulavirus* et *Rubulavirus*) ne sont pas détectables par le système de PCR nichée que nous avons utilisée. En tout état de cause, nous ne pouvons que souligner l'absence de détection de virus Henipa dans la région du SOOI.

L'analyse multivariée a mis en évidence une transmission virale plus active au sein des chauves-souris insectivores dans notre étude, puisque sur 87 chauves-souris frugivores, seulement 3 d'entre elles, du genre *Pteropus*, ont été retrouvées positives, démontrant que les *UMRVs* infectent préférentiellement le groupe des insectivores. Les Yinpterochiroptera (chauves-souris insectivores) représentent la sous famille la plus diversifiée à Madagascar et la plupart des individus phylogénétiquement proches, et plus important encore, la plupart des genres Miniopteridae sont retrouvés en sympatrie sur le même site.

Notre analyse multivariée a en effet montré que le comportement grégaire des chauves-souris, et principalement la sympatrie, sont des facteurs clés dans les prévalences fortes en paramyxovirus chez les chauves-souris. Ce patron a été observé à la fois pour des chauves-souris de la même famille ou de familles différentes. Un tel phénomène a déjà été soulevé par l'équipe de Luis et al. (2014) qui démontre que le contact étroit entre différentes espèces de chauves-souris sur un même site, trait écologique caractéristique des chiroptères, favoriserait la transmission virale entre différentes espèces. Dans une étude ultérieure, ces mêmes auteurs ont montré que le comportement grégaire était lui aussi corrélé à la richesse virale au sein des chauves-souris. Par ces résultats nous démontrons l'importance de la biodiversité, à savoir une richesse d'hôtes au sein d'une communauté de chauves-souris, les Yinpterochiroptera, accompagnés des traits de vie écologiques, la sympatrie, en tant que facteur amplificateur de la transmission virale.

Keesing et al ont décrit l'effet favorable ou défavorable que pouvait avoir la biodiversité sur la transmission d'agents pathogènes (Keesing et al., 2010). Dans notre modèle d'étude, nous montrons que la diversité d'espèces est corrélée à la diversité virale, ce qui d'après Morse et al (Morse et al., 1993) favoriserait la génération de "pools zoonotiques". Un dernier facteur écologique important relevé dans notre étude est l'effet de la température plus élevée dans certaines localités de Madagascar qui apparait comme facteur de risque dans la transmission de paramyxovirus chez les chauves-souris de Madagascar. Des études intéressantes avaient déjà révélé l'importance du climat sur le taux d'évolution du virus de la rage (Streicker et al., 2012). Ainsi le virus rabique retrouvé chez des chauves-souris des régions tropicales et subtropicales aurait un taux d'évolution quatre fois plus important que chez les chauves-souris vivant en zone tempérée. Il est apparu que chez les chauves-souris vivant dans les régions plus chaudes, le métabolisme énergétique serait accéléré, augmentant potentiellement le taux de mutations dû à la production massive de radicaux libres (Rohde et al., 1992; Martin & Palumbi, 1993). A l'inverse, les chauves-souris qui entreraient en hibernation/torpeur dans les régions froides présentent des taux d'infections très bas sans doute dû à une diminution du métabolisme cellulaire (Domingo et al., 1997). Au vu du nombre de « host-switch » observé inter- et intra-ordres de micromammifères dans la zone SOOI, il est fort probable qu'en couplant nos analyses écologiques aux analyses génétiques, nos résultats pourraient indiquer un effet positif de la température élevée sur le taux d'évolution viral expliquant in fine, la quantité importante de *host-switch* observée dans la région SOOI.

Enfin, une étude réalisée par l'équipe du CRVOI en 2014 a décrit la saisonnalité, notamment, des périodes de reproduction des chauves-souris, comme étant un facteur important dans la dynamique de transmission d'agent pathogène à La Réunion (Dietrich et al., 2014) en accord avec d'autres études (O'Shea et al., 2014). L'étude menée à La Réunion sur l'espèce endémique *Mormopterus francoismoutoui* retrouvée dans une grotte dédiée à la maternité à La Réunion a montré une dynamique infectieuse saisonnière où la transmission des paramyxovirus semble intervenir de façon active en deux pics; le premier pic d'infection est observé à la fin de la grossesse et concerne préférentiellement les mères gestantes au moment où elles se concentrent à très forte densité dans la grotte maternité et sont le plus en contact entre elles. Le taux d'infection des chauves-souris peut atteindre 50%.

Un second pic d'infection survient deux mois après l'accouchement, et concerne les juvéniles quelques semaines après leur naissance, vraisemblablement après la perte des anticorps maternels passivement transmis lors de la période d'allaitement (Dietrich et al., 2014). Les séquences génétiques correspondant aux UMRVs excrétés dans les urines ont été trouvées similaires des séquences des UMRVs retrouvés dans le surnageant de broyat de pool d'organes (reins, poumon et rate) (données non publiés). Il est envisageable que l'excrétion des paramyxovirus chez les chauves-souris coïncide avec une infection systémique, i.e. avec une virémie transitoire. Cependant une telle possibilité n'est pas exclusive d'une excrétion virale plus prolongée dans le temps et non corrélée avec la virémie, une fois que l'infection rénale chronique se serait établie chez les animaux qui deviennent de ce fait excréteurs de virus, i.e. un hôte réservoir. Par ailleurs, les résultats de cette étude suggèrent que la dynamique virale pourrait être modulée par la réponse immune développée par l'animal infecté. Ces résultats très intéressants permettent d'orienter les futures investigations par biologie moléculaire sur des prélèvements récoltés de manière non invasive telle que l'urine afin d'identifier la fraction des chauves-souris qui jouerait le rôle d'hôtes de maintien du virus et permettant une excrétion virale sur de longues périodes, du moins d'une saison sur l'autre entre deux épisodes d'accouchement, pour des raisons (immunologiques ?) qui restent encore à clarifier.

### Système immunitaire

Un aspect intéressant encore peu étudié, est le système immunitaire des chauves-souris. Les chauves-souris sont des mammifères très anciens, près de 80 Millions d'années il est possible que leurs réponses immunitaires innées et acquises soit différentes des autres mammifères mieux étudiés (rats, souris). Comme décrit dans le chapitre I, on sait que des virus extrêmement pathogènes comme *Ebola* et *SARS* qui provoquent respectivement fièvres hémorragiques et pneumonies, sont retrouvés chez des chauves-souris saines (Biek et al., 2006; Wang et al., 2006). Les réponses immunitaires de celles-ci sont-elles régulées différemment permettant de contrôler le niveau de réplication du virus, sans l'éliminer totalement? (Calisher et al., 2006). Pourquoi certains virus infectent-ils et persistent-ils chez les chauves-souris alors qu'ils se révèlent hautement pathogènes pour les humains et les autres vertébrés? Pourquoi certaines espèces de chauves-souris infectées expriment elles des symptômes et d'autres pas ?

Quel est le rôle du système immunitaire dans le maintien de l'infection virale? Malheureusement très peu de données sont disponibles sur ces points cruciaux.

L'hypothèse d'un mutualisme a été émise dans lequel le système immunitaire imposerait un équilibre qui maintiendrait l'infection et la transmission d'un virus à un niveau basal favorisant à la fois l'hôte et l'agent pathogène (Calisher et al., 2006). Ce trade off expliquerait aussi l'absence de symptômes chez les individus collectés dans cette étude puisque nous n'avons jamais observé d'individu apparemment malade ni de mortalité anormale sur les sites visités. Les lyssavirus qui sont les virus le plus souvent impliqués dans la transgression de la barrière d'espèces, sont capables d'infecter un spectre d'hôte extrêmement élevé (Banyard et al., 2014). L'absence de détection d'ARN ou d'antigène viraux, malgré la très forte séropositivité est une observation courante rapportée dans la plupart des études portant sur les lyssavirus (Arguin et al., 2002; Reynes et al., 2011). Elle indique que chez les animaux séropositifs, les anticorps neutralisants détectés sont le reflet d'une infection ancienne, et que le système immunitaire aurait réussi à éliminer le virus. De fait, de nombreuses études ont montré que l'ARN lyssavirus n'est détecté que chez des individus symptomatiques, présentant des signes cliniques caractéristiques d'une infection neurotrope (déséquilibre, perte d'orientation, agressivité, etc.) (Markotter et al., 2006 ; Aubert, 1999 ; King & Crick, 1988). Il est possible que pour certaines raisons que l'on ignore (perte de fitness de l'hôte?), chez certains individus, l'équilibre entre le système immunitaire et le niveau de réplication virale serait rompu entraînant la multiplication virale et provoquant l'état morbide caractéristique de la neurovirulence et pouvant même provoquer la mort de l'individu (Kuzmin et al., 2008; Kuzmin et al., 2010). Dans d'autres modèles animaux d'infection à Lyssavirus, comme l'infection expérimentale du chien par le virus rabique, tous les intermédiaires peuvent être observés depuis l'infection symptomatique et rapidement mortelle jusqu'à l'infection très lente et progressive sur plusieurs mois en passant par une infection asymptomatique et apparemment contrôlée chez une minorité d'animaux qui survivent (tout en excrétant de façon discontinue le virus dans la salive) (Fekadu, 1988 ; Baer, 2012). Il serait utile de vérifier si l'absence de localisation cérébrale des lyssavirus chez les chauves-souris telle que prouvée dans notre série, est corrélée ou pas à l'absence d'excrétion virale du même virus dans la salive. La dissociation entre les deux tissus serait un argument en faveur de l'établissement d'une persistance virale chez les chiroptères due à un petit pourcentage d'animaux à la fois réservoirs et transmetteurs.

D'une façon générale, les virus doivent se répliquer et se transmettre au-delà du seuil d'extinction (R<sub>0</sub> > 1) afin de ne pas disparaitre et ce maintien doit être amélioré continuellement dans le temps sans enclencher des réponses immunitaires inhibitrices, ou entrainer le décès rapide de l'hôte ce qui interromprait prématurément la transmission (Real & Biek, 2007). Il est aussi fort à penser que la persistance virale chez les chauves-souris dans la plupart des études, n'est peut-être pas le reflet de la balance immunitaire de l'hôte, mais plutôt la conséquence d'une stratégie réplicative virale contrôlée, très opportune et très ancienne de survie des virus. Il est aussi envisageable qu'une infection virale persistante puisse apporter une protection contre d'autres agents pathogènes par le biais d'une interférence virale avec réponse d'Interférons de type I endogène. Une telle interférence virale protectrice a été démontrée dans l'infection expérimentale de *Rattus* par le bacille *Yersinia pestis* agent de la peste. L'infection bactérienne, toujours mortelle est totalement neutralisée par une infection latente des rats par un virus herpes murin (Barton et al., 2007).

L'enquête séro-épidémiologique conduite sur les chauves-souris et les petits mammifères terrestres de la région du SOOI utilisant les chips composites pour la détection d'anticorps en IIF contre 15 virus à ARN monocaténaires appartenant à 5 familles virales a apporté des informations intéressantes. Cinq virus représentent la famille des *Paramyxoviridae* : le virus de la rougeole (genre Morbillivirus), le Parainfluenza virus 2 et le virus des oreillons (genre Rubulavirus), le RSV (genre Pneumovirus), le virus Parainfluenza virus 1 (genre Respirovirus). Cette étude a permis de révéler que les anticorps anti-Paramyxovirus détectés chez les chauves-souris sont principalement dirigés contre les deux virus appartenant au genre Rubulavirus représentés sur la lame chips : oreillons et PIV2 (et beaucoup plus rarement contre le virus de la rougeole, PIV1 et RSV). Il aurait été très intéressant de pouvoir corréler ce résultat sérologique anti-RubulaV avec l'expression d'ARN du genre Rubulavirus afin d'éclairer la relation entre présence d'anticorps et persistance virale. Malheureusement, notre système de détection RT-PCR employé pour les paramyxovirus, utilise des oligonucléotides du système RMH, connu pour amplifier des régions fortement conservées du génome dans le gène codant la polymérase (L) des paramyxovirus des genres Respirovirus, Morbilivirus et Hénipavirus (Tong et al., 2008). Ce système ne détecte pas le genre Rubulavirus. Un autre système amplifiant les genres, Pneumovirus, Avulavirus et Rubulavirus (PAR) a été testé au sein de notre équipe de recherche mais il s'est avéré moins sensible et génère de faux positifs.

De façon surprenante, la seule corrélation négative que nous avons pu caractériser dans notre étude sérologique relie la présence des anticorps anti-rubulavirus à l'absence de détection d'ARN d'URMVs qui phylogénétiquement sont plutôt classés comme Morbilli-related (UMRVs). L'interrogation que pose cette corrélation négative ne pourra être clarifiée qu'une fois notre étude complétée par la détection spécifique des ARN de Rubulavirus par le système PAR. En attendant, si elle devait correspondre à une réalité biologique, la corrélation négative indiquerait une antigénicité croisée entre Rubulavirus et UMRVs, ou la possibilité d'échanges génétiques (recombinaison) entre ces derniers. La séquence complète de quelques UMRVs actuellement très avancée au CRVOI permettrait de répondre à ces questions.

## **Conclusion et perspectives**

Nos résultats apportent des informations totalement nouvelles, jamais rapportées jusqu'à aujourd'hui dans la zone SOOI. Les investigations menées au cours de ces 3 années de recherche, grâce aux échantillonnages intenses, aux traitements des échantillons et *in fine* à l'analyse via des outils de bioinformatiques ont apporté un éclaircissement sur la caractérisation et la circulation de virus hautement diversifiés au sein d'une communauté de chauves-souris et de petits mammifères terrestres de la zone SOOI.

Ces déterminants écologiques et génétiques mis en évidence, font partie des critères qu'il convient d'investiguer de façon prioritaire afin de mieux appréhender le risque d'émergence, dans le futur, de ces agents potentiellement zoonotiques dans les populations animales ou humaines (Woolhouse et al., 2005 ; Keusch et al., 2009 ; Morse et al., 2012).

Plusieurs perspectives sont envisageables pour la suite.

Bien que la sérologie RFFIT soit spécifique et indique une circulation intense d'au moins deux lyssavirus appartenant aux phylogroupes I et II respectivement, nous n'avons pas pu identifier les espèces virales en cause. Cela est rendu d'autant plus complexe par la présence de nombreuses réactions croisées en lyssavirus au sein du même phylogroupe. Dans l'objectif de caractériser les déterminants évolutifs et écologiques des lyssavirus et de leurs hôtes dans la zone du SOOI ou encore les classer phylogénétiquement, il serait souhaitable de réaliser un échantillonnage d'individus présentant des symptômes caractéristiques de la rage ou des individus morbides.

La possibilité de *host-switch* d'agents zoonotiques de la population sauvage vers les populations domestiques, constitue l'une des premières étapes avant l'émergence vers les populations humaines (Jones et al., 2008 ; Morrens et al., 2012). Ainsi, dans la continuité de notre étude, il serait intéressant d'investiguer les populations d'animaux domestiques et les populations humaines susceptibles d'être en contact avec les micromammifères des îles

étudiées. Des analyses sérologiques permettront d'évaluer le potentiel risque de sauts d'espèces de ces paramyxovirus de chauves-souris vers ces différentes populations.

Celles-ci pourraient se faire après analyse des séquences complètes de paramyxovirus que nous avons obtenus qui permettra non seulement de classer "définitivement" phylogénétiquement ces nouveaux virus mais aussi de réaliser des tests de sérologie spécifiques afin d'apprécier le taux d'exposition virale antérieure chez ces individus. Ceci étant, nous avons pu réaliser le séquençage complet des paramyxovirus qui ont pu être isolés confirmant que les *UMRVs* appartiennent tous à un même lignage structuré, et proches des *Morbillivirus*, mais ne peuvent pas être considérés comme des *Morbillivirus*: leur structure génomique et la taille de leurs génomes diffèrent - Les *UMRVs* font environ 18 kb contre 15 kb pour les *Morbillivirus* (analyses en cours de finalisation).

Outre *Rattus rattus*, existent-t-ils d'autres réservoirs/vecteurs capables de véhiculer des paramyxovirus génétiquement très proches entre des gîtes de chauves-souris géographiquement très éloignés? L'équipe de Ci-Xiu Li et al. (2015) a mis en évidence chez plus de 70 espèces d'arthropodes près de 112 nouveaux virus à ARN. Ces virus étaient notamment très proches des arénavirus, filovirus, hantavirus, influenzavirus, lyssavirus et paramyxovirus. Les arthropodes de la zone du SOOI participeraient-ils à la transmission virale au sein de chauves-souris localisées dans des gîtes très éloignés? Pour répondre à cette question, il serait intéressant de déterminer la richesse virale inféodée aux arthropodes du SOOI.

Il est connu que La Réunion, Maurice et Madagascar hébergent chacune une espèce endémique « sœur » de chiroptère, respectivement, *Mormopterus francoismoutoui* (La Réunion), *Mormopterus acetabulosus* (Maurice) et *Mormopterus jugularis* (Madagascar). Sur la base d'éléments moléculaires et morphologiques, *Mormopterus acetabulosus*, qui était jusqu'à présent considérée comme endémique des îles Mascareignes (Maurice et La Réunion), a été subdivisée très récemment en deux taxons étroitement liés, *Mormopterus acetabulosus* et *Mormopterus francoismoutoui* : les deux groupes étant réciproquement monophylétiques, mais séparés par une moyenne de divergence de l'ordre de 5% (Goodman et al., 2008). Par ailleurs, ces deux espèces sont également proches de *Mormopterus jugularis*. Il serait intéressant de déterminer si les virus au sein de ces chauves-souris vivant sur 3 îles différentes avec des écosystèmes contrastés répondraient aux mêmes déterminants génétique

et écologique, au regard de la récente radiation observée. Serait-il possible d'observer cette fois-ci une co-divergence virale associée à de la co-spéciation? Quels sont les facteurs écologiques dans les 3 îles, favorables à la transmission virale?

Finalement, il serait utile de déterminer si les infections des chauves-souris par des virus, bactéries ou parasites les protégeraient contre l'une ou l'autre de ces infections ou contre d'autres pathogènes dans le cadre d'une pathocénose infectieuse. Pour répondre à cette question, nous analyserons en collaboration avec Yann Gomard et Beza Ramasindrazana, l'ensemble des résultats générés sur le même échantillonnage de chauves-souris dans les 3 modèles biologiques explorés au CRVOI : à savoir le modèle viral des paramyxovirus, le modèle bactérien de la leptospirose, et le modèle parasitaire des hémoparasitoses.

## **Bibliographie**

Agosta, S.J., Janz, N. & Brooks, D.R. (2010). How specialists can be generalists: resolving the "parasite paradox" and implications for emerging infectious disease. Zoologica, 27: 151–162. doi: 10.1590/51984-46702010000200001.

Andriamandimby, S.F., Héraud, J.M., Ramiandrasoa, R., Ratsitorahina, M., Rasambainarivo, J.H., Dacheux, L., Lepelletier, A., Goodman, S.M., Reynes, J.M. & Bourhy, H. (2013). Surveillance and control of rabies in La Reunion, Mayotte, and Madagascar.Vet Res; Sep; 44:77. doi: 10.1186/1297-9716-44-77.

Anishchenko, M., Bowen, R.A., Paessler, S., Austgen, L., Greene, I.P. & Weaver, S.C. (2006). Venezuelan encephalitis emergence mediated by a phylogenetically predicted viral mutation. Proc Natl Acad Sci U S A;103(13):4994-9. Epub 2006 Mar 20.

Aplin, K.P., Chesser, T. & ten Have, J. (2003). Evolutionary biology of the genus *Rattus*: profile of an archetypal rodent pest. In: Singleton GR, Hinds LA, Krebs CJ. & Spratt DM (eds) Rats, mice and people: rodent biology and management. ACIAR, Canberra, pp 487–498.

Arguin, P.M., Murray-Lillibridge, K., Miranda, M.E., Smith, J.S., Calaor, A.B. & Rupprecht, C.E. (2002). Serologic evidence of Lyssavirus infections among bats, the Philippines. Emerg Infect Dis.; 8(3):258-62.

Assenberg, R., Delmas, O., Ren, J., Vidalain, P.O., Verma, A., Larrous, F., Graham, S.C., Tangy, F., Grimes, J.M. & Bourhy, H. (2010). Structure of the nucleoprotein binding domain of Mokola virus phosphoprotein. J Virol; 84(2):1089-96. doi: 10.1128/JVI.01520-09. Epub Nov 11, 2009.

Aubert, M. F. (1999). Rabies in individual countries, France. Rabies Bull. Europe 23:6.

Badrane, H., & Tordo. N. (2001). Host switching in lyssavirus history from the Chiroptera to the Carnivora orders. J. Virol. 75:8096–8104.

Baer, G.M. (2012). The Natural History of rabies. Elsevier, Dec 2- Health & Fitness.

Banyard, A.C., Evans, J.S., Luo, T.R. & Fooks, A.R. (2014). Lyssaviruses and bats: Emergence and zoonotic threat. Viruses 6(8):2974-90.

Baranowski, E., Ruiz-Jarabo, C.M., Pariente, N., Verdaguer, N. & Domingo, E. (2003). Evolution of cell recognition by viruses: A source of biological novelty with medical implications. Adv Vir Res; 62, 19-111.

Barton, E.S., White, D.W., Cathelyn, J.S., Brett-McClellan, K.A., Engle, M., Diamond, M.S., Miller V.L.& Virgin, H.W. (2007). Herpesvirus latency confers symbiotic protection from bacterial infection. Nature 447(7142):326-329.

Biek, R., Walsh, P.D., Leroy, E.M., & Real, L.A. (2006). Recent common ancestry of Ebola Zaire virus found in a bat reservoir. PLoS Pathog; 2, e90.

Bonnefoy, X., Kampen, H. & Sweeney, K. (2008). Public health significance of urban pests. World Health Organization, Copenhagen.

Calisher, C.H., Childs, J.E., Field, H.E., Holmes, K.V. & Schountz, T. (2006). Bats: Important reservoir hosts of emerging viruses. Clin. Microbiol. Rev. 19: 531-545.

CDC. (2015). [Online]. http://www.cdc.gov/vhf/ebola/outbreaks/2014-west-africa.

CDC. (2014). [Online]. <a href="http://www.cdc.gov/coronavirus/mers/about/index.html">http://www.cdc.gov/coronavirus/mers/about/index.html</a>.

Childs, J.E., Richt, J.A. & Mackenzie, J.S. (2007). Introduction: conceptualizing and partitioning the emergence process of zoonotic viruses from wildlife to humans. Curr. Top. Microbiol. Immunol. 315:1-31.

Chua, K.B., Bellini, W.J., Rota, P.A., Harcourt, B.H., Tamin, A., Lam, S.K., Ksiazek, T.G., Rollin, P.E., Zaki, S.R., Shieh, W., Goldsmith, C.S., Gubler, D.J., Roehrig, J.T., Eaton, B., Gould, A.R., Olson, J., Field, H., Daniels, P., Ling, A.E., Peters, C.J., Anderson, L.J. & Mahy, B.W. (2000). Nipah virus: a recently emergent deadly paramyxovirus. Science 288: 1432-1435.

Ciurea, A., Klenerman, P., Hunziker, L., Horvath, E., Senn, B.M., Ochsenbein, A.F., Hengartner, H. & Zinkernagel, R.M. (2000). Viral persistence in vivo through selection of neutralizing antibody-escape variants. Proc Natl Acad Sci USA.;97(6): 2749-54.

Clavero, M. & García-Berthou, E. (2005). Invasive species are a leading cause of animal extinctions. Trends Ecol. Evol. 20: 110.

Coertse, J., Markotter, W. & Nel, L.H. (2014). Demonstration of African lyssavirus RNA with real-time polymerase chain reaction. In Rupprecht, C & Nagajaran, T, editors. Current laboratory techniques in rabies diagnosis, research, and prevention, Volume 1. Amsterdam: Academic Press; pp. 63-73.

Condori-Condori, R.E., Streicker, D.G., Cabezas-Sanchez, C.& Velasco-Villa, A. (2013). Enzootic and epizootic rabies associated with vampire bats, Peru. Emerg Infect Dis; 19(9).

Constantine, D.G. (1988). Transmission of pathogenic organisms by vampire bats. In: Greenhall AM, Schmidt U, editors. Natural history of vampire bats. Boca Raton, FL: CRC Press; p. 167–189.

Coulibaly-N'Golo, D., Allali, B., Kouassi, S.K., Fichet-Calvet, E., Becker-Ziaja, B., Rieger, T., Olschläger, S., Dosso, H., Denys, C., Ter Meulen, J, Akoua-Koffi, C. & Günther, S. (2011). Novel Arenavirus sequences in *Hylomyscus sp.* and *Mus (Nannomys) setulosus* from Côte d'Ivoire: implications for evolution of Arenaviruses in Africa. PLoS ONE; 6(6): e20893.

Cui, J., Han, N., Streicker, D., Li, G., Tang, X., Shi, Z., Hu, Z., Zhao G., Fontanet., A, Guan., Y, Wang., L, Jones., G, Field., H.E, Daszak, P. & Zhang, S. (2007). Evolutionary relationships between bat coronaviruses and their hosts. Emerg Infect Dis; (10):1526-32.

Daszak, P. Cunningham, A. & Hyatt, A. (2000). Emerging infectious diseases of wildlife – threats to biodiversity and human health. Science 287: 443-448.

de Vienne, D.M., Refrégier, G., López-Villavicencio, M., Tellier, A., Hood, M.E. & Giraud, T. (2013). Cospeciation vs host-shift speciation: methods for testing, evidence from natural associations and relation to coevolution. New Phytologist; 198: 347-385.

Dietrich, M., Wilkinson, D.A., Benlali, A., Lagadec, E., Ramasindrazana, B., Dellagi, K. & Tortosa, P. (2015). Leptospira and paramyxovirus infection dynamics in a bat maternity enlightens pathogen maintenance in wildlife. Environ Microbiol. (11):4280-9. doi: 10.1111/1462-2920.12766. Epub 2015 Feb 14.

Domingo, E. & Holland, J.J. (1997). RNA virus mutations and fitness for survival. Annu. Rev. Microbiol. 51:151-78.

Domingo, E. (1999). Quasispecies. In Granoff A, Webster, R.G. (eds) Encyclopedia of virology. Academic Press, London, 1431–1436.

Domingo, E., Escarmís, C., Lázaro, E. & Manrubia, S.C. (2005). Quasispecies dynamics and RNA virus extinction. Virus Res.;107(2):129-39.

Drexler, J.F., Corman, V.M., Müller, M.A., Maganga, G.D., Vallo, P;, Binger, T., Gloza-Rausch, F., Cottontail, V.M., Rasche, A., Yordanov, S., Seebens, A., Knörnschild, M., Oppong, S., Adu Sarkodie, Y., Pongombo, C., Lukashev, A.N., Schmidt-Chanasit, J., Stöcker, A., Carneiro, A.J., Erbar, S., Maisner, A., Fronhoffs, F., Buettner, R., Kalko E.K., Kruppa, T., Franke, C.R., Kallies, R., Yandoko, E.R., Herrler, G., Reusken, C., Hassanin, A., Krüger, D.H., Matthee, S., Ulrich, R.G., Leroy, E.M. & Drosten, C. (2012). Bats host major mammalian paramyxoviruses. Nat Commun. 24;3:796.

Duffy, S., Shackelton, L.A. & Holmes, E.C. (2008). Rates of evolutionary change in viruses: patterns and determinants. Nature Reviews Genetics 9, 267-276 | doi:10.1038/nrg2323

Epstein, J.H., Olival, K.J., Pulliam, J.R.C., Smith, C., Westrum, J., Hughes, T., Dobson, A.P., Akbar, Z., Rahman, S.A., Basir, M.M., Field, H.E. & Daszak, P. (2009). *Pteropus vampyrus*, a hunted migratory species with a multinational home-range and a need for regional management. J. Appl. Ecol. 46: 991-1002.

Fauci, A.S. (2001). Infectious diseases: considerations for the 21st century. Clin. Infect. Dis. 32: 675–685. Epub 2001.

Fekadu, M. (1988). Pathogenesis of rabies virus infection in dogs. Rev. Infect. Dis. 10 Suppl 4:S678-83.

Field, H., Young, P., Yob, J.M., Mills, J., Hall, L. & Mackenzie, J. (2001). The natural history of Hendra and Nipah viruses. Microbes Infect 3: 307-314.10.1016/S1286-4579(01)01384-3

Field, H., Crameri, G., Kung, N.Y., Wang, L.F. (2012). Ecological aspects of hendra virus. Curr. Top. Microbiol. Immunol. 359:11-23. doi: 10.1007/82 2012 214.

Finke, S. & Conzelmann, K.K. (2005). Replication strategies of rabies virus. Virus Res., 111(2): 120-131.

Firth, C., Bhat, M., Firth, M.A., Williams, S.H., Frye, M.J., Simmonds, P., Conte, J.M., Ng, J., Garcia, J., Bhuva, N.P., Lee, B., Che, X., Quan, P.L. & Lipkin W.I. (2014). Detection of zoonotic pathogens and characterization of novel viruses carried by commensal *Rattus norvegicus* in New York City. MBio; 14;5(5):e01933-14. doi: 10.1128/mBio.01933-14.

Food and Agriculture Organisation of the United Nations (FAO). (2011). Investigating the role of bats in emerging zoonoses: balancing ecology, conservation and public health interests. Edited by S.H. Newman, H.E. Field, C.E. de Jong and J.H. Epstein. FAO Animal Production and Health Manual No. 12. Rome.

George, D.B., Webb, C.T., Farnsworth, M.L., O'Shea, T.J., Bowen, R.A., Smith, D.L., Stanley, T.R., Ellison, L.E. & Rupprecht C.E. (2011). Host and viral ecology determine bat rabies seasonality and maintenance. Proc. Natl. Acad. Sci. U S A; 108(25):10208-13.

Goodman, S.M., J.U. Ganzhorn, & D. Rakotondravony. (2003). Introduction to the mammals, in the natural history of Madagascar, S.M. Goodman and J.P. Benstead, Editors. The University of Chicago Press, Chicago.

Goodman SM, Buccas W, Naidoo T, Ratrimomanarivo F, Taylor PJ, Lamb J. (2010). Patterns of morphological and genetic variation in western Indian Ocean members of the *Chaerephon 'pumilus'* complex (Chiroptera: Molossidae), with the description of a new species from Madagascar. Zootaxa, 2551:1-36.

Goodman, S.M., van Vuuren, B.J., Ratrimomanarivo, F., Probs,t J-M. & Bowie, R.C.K (2008). Specific status of populations in the Mascarene Islands referred to *Mormopterus acetabulosus* (Chiroptera: Molossidae), with description of a new species. J. Mammal. 89(5):1316-1327.

Goodman, S.M. (2011). Les chauves-souris de Madagascar. Association Vahatra, Antananarivo.

Goodman, S.M., Puechmaille, S.J., Friedli-Weyeneth, N., Gerlach, J., Ruedi, M., Schoeman, M.C., Stanley, W.T. & Teeling, E.C. (2012a). Phylogeny of the Emballonurini

(Emballonuridae) with descriptions of a new genus and species from Madagascar. J. Mammal. 93: 1440–1455.

Goodman, S.M, Taylor, P.J., Ratrimomanarivo, F. & Hoofer, S. (2012b). The genus *Neoromicia* (Family Vespertilionidae) in Madagascar with the description of a new species. Zootaxa 3250: 1–25.

Goodman, S.M. & Raherilalao, M.J. (2013). (eds.). Atlas d'une sélection de vertébrés terrestre de Madagascar. Association Vahatra, Antananarivo.

Goodman, S.M. &, Ramasindrazana, B. (2015). Description of a new species of the *Miniopterus aelleni* group (Chiroptera: Miniopteridae) from upland areas of central and northern Madagascar. Zootaxa. Mar 23;3936(4):538-58. doi: 10.11646/zootaxa.3936.4.4.

Goodman, S.M., Rakotondramanana, C.F., Ramasindrazana, B., Kearney, T., Monadjem, A., M Schoeman, M.C., Taylor, P.J., Naughton, K. & Appleton, B. (2015). An integrative approach to characterize Malagasy bats of the subfamily Vespertilioninae Gray, 1821, with the description of a new species of *Hypsugo*. Zool. J. Linnean. Soc. 04/2015; 173(4):988-1018. DOI:10.1111/zoj.12223

Griffin, D. R. (1970). Migrations and homing in bats, p. 233–264. In W. A. Wimsatt (ed.), Biology of bats. Academic Press, New York, N.Y.

Guan Y., Zheng, BJ., He, Y.Q., Liu, X.L., Zhuang, Z.X., Cheung, C.L., Luo, S.W., Li, P.H., Zhang L.J., Guan, Y.J., Butt, K.M., Wong, K.L., Chan, K.W., Lim, W., Shortridge, K.F., Yuen, K.Y., Peiris, J.S. & Poon, L.L. (2003). Isolation and characterization of viruses related to the SARS coronavirus from animals in southern China. Science 302:276-8.

Guerrant, R. L. & Blackwood, B. L. (1999). Threats to global health and survival: the growing crises of tropical infectious diseases — an "unfinished" agenda. Clin. Infect. Dis; 28, 966–986.

Guo, W.P., Lin, X.D., Wang, W., Tian, J.H., Cong, M.L., Zhang, H.L, Wang, M.R., Zhou, R.H., Wang, J.B., Li, M.H., Xu J., Holmes, E.C. & Zhang, Y.Z. (2013). Phylogeny and origins of Hantaviruses harbored by bats, insectivores, and rodents. PLoS Pathog 9(2): e1003159. doi:10.1371/journal.ppat.1003159.

Holland, J., Spindler, K., Horodyski, F., Grabau, E., Nichol, S. & VandePol, S. (1982). Rapid evolution of RNA genomes. Science 215, 1577–1585. (doi:10.1126/science.7041255)

Holmes, E.C. & Moya, A. (2002). Is the quasispecies concept relevant to RNA viruses? J. Virol. 76, 460–465. (doi:10.1128/JVI.76.1. 460-462.2002).

Huchon, D., Madsen, O., Sibbald, M.J.J.B., Ament, K., Stanhope, M.J., Catzeflis, F., De Jong, W.W. & Douzery, E.J.P. (2002) Rodent phylogeny and a timescale for the evolution of glires: evidence from an extensive taxon sampling using three nuclear genes. Mol. Biol. Evol. 19, 1053, 2002.

Hughes, A.L. & Friedman, R. (2000). Evolutionary diversification of protein-coding genes of hantaviruses. Mol. Biol. Evol. 17(10):1558-68.

ICTV. (2014). [Online] <a href="https://www.ictvonline.org/virustaxonomy.asp">www.ictvonline.org/virustaxonomy.asp</a>.

Ishag, O.M., Saeed, I.K., Ali & Y.H. (2015). Peste des petits ruminants outbreaks in White Nile State, Sudan. J. Vet. Res. 82(1):E1-4. doi: 10.4102/ojvr.v82i1.897.

Jackson, A.P. & Charleston, M.A. (2004). A cophylogenetic perspective of RNA-virus evolution. Mol. Biol. Evol. 21(1): 45-57.

Johara, M.Y., Field, H., Rashdi, A.M., Morrissy, C., van der Heide, B., Rota, P., bin Adzhar, A., White, J., Daniels, P., Jamaluddin, A. & Ksiazek, T. (2001). Nipah virus infection in bats (order Chiroptera) in peninsular Malaysia. Emerg. Infect. Dis; 7(3): 439-441.

Jones, K. E., Purvis, A., MacLarnon, A., Bininda-Emonds, O. R. & Simmons, N. B. (2002). A phylogenetic supertree of the bats (Mammalia: Chiroptera). Biol. Rev. Camb. Philos. Soc. 77:223–259.

Keesing, F., Holt, R.D. & Ostfeld, R.S. (2006). Effects of species diversity on disease risk. Ecol Lett;9(4):485-98.

Keesing F., Belden, L.K., Daszak, P., Dobson, A., Harvell, C.D., Holt, R.D., Hudson, P., Jolles, A., Jones, K.E., Mitchel, C.E., Myers, S.S., Bogich, T. & Ostfeld, R.S. (2010). Impacts of biodiversity on the emergence and transmission of infectious diseases. Nature 468, 647–652.

Keusch, G.T., Pappaioanou, M., Gonzalez, M.C., Scott, K.A. & Tsai, P. (eds). (2009). Sustaining global surveillance and response to emerging zoonotic diseases. Washington, DC: The National Academies Press.

King, A., & Crick, J. (1988). Rabies-related viruses. In "Rabies", (. Charlton, K.M and Campbell, J. B. eds.) 177–200. Kluwer Academic Publishers, Boston.

Kitchen, A., Shackelton, L.A. & Holmes, E.C. Family level phylogenies reveal modes of macroevolution in RNA viruses. Proc. Natl. Acad. Sci. U S A; 108(1): p. 238-43.

Koopman, K.F. (1993). Chiroptera. In D.E. Wilson and D.M. Reeder, eds. Mammalian species of the world,137-241. Washington, DC, Smithsonian Institution Press.

Kreuder Johnson, C., Hitchens, P.L., Smiley Evans, T., Goldstein, T., Thomas, K., Clements A., Joly, D.O., Wolfe, N.D., Daszak, P., Karesh, W.B. & Mazet J.K. (2015). Spillover and pandemic properties of zoonotic viruses with high host plasticity. Sci Rep; Oct 7;5:14830. doi: 10.1038/srep14830.

Kunz, T.H. & Pierson, E.D. (1994). Bats of the world: an introduction. In R.M. Nowak. Walker's bats of the world, 1-46. Baltimore, Maryland, USA and London, Johns Hopkins University Press.

Kurth, A., Kohl, C., Brinkmann, A., Ebinger, A., Harper, J.A., Wang, L.F., Mühldorfer, K. & Wibbelt, G. (2012). Novel paramyxoviruses in free-ranging European bats. PLoS One; 7(6):e38688. doi: 10.1371/journal.pone.0038688. Epub 2012 Jun 21.

Kuzmin, I. V., Niezgoda, M., Franka, R., Agwanda, B., Markotter, W., Beagley, J. C., Urazova, O. Y., Breiman, R. F., & Rupprecht, C. E. (2008). Lagos bat virus in Kenya. J. Clin. Microbiol. 46:1451–1461

Kuzmin, I. V., Mayer, A. E., Niezgoda, M., Markotter, W., Agwanda, B., Breiman, R. F., & Rupprecht, C. E. (2010). Shimoni bat virus, a new representative of the Lyssavirus genus. Virus Res. 149:197–210.

Lamb, R.A. & Parks, G.D. (2007). Paramyxoviridae: The viruses and their replication. In: Fiels Virology, eds. Knipe, D.M., Howley, P.P. 1450-1474. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins, 2007.

Lau, S.K., Woo, P.C., Li, K.S., Huang, Y., Tsoi, H.W., Wong, B.H., Wong, S.S., Leung, S.Y., Chan, K.H. & Yuen, K.Y. (2005). Severe acute respiratory syndrome coronavirus-like virus in Chinese horseshoe bats. Proc. Natl. Acad. Sci. USA; 102:14040-14045.

Lei, B.R. & Olival, K.J. (2014). Contrasting patterns in mammal-bacteria coevolution: bartonella and leptospira in bats and rodents. PLoS Negl Trop Dis; 20;8(3):e2738.

Leroy, E.M., Kumulungui, B., Pourrut, X., Rouquet, P., Hassanin, A., Yaba, P., Délicat, A., Paweska, JT., Gonzalez, JP. & Swanepoel, R. (2005). Fruit bats as reservoirs of Ebola virus. Nature 438, 575–576.

Li, C.X., Shi, M., Tian, J.H., Lin, X.D., Kang, Y.J., Chen, L.J., Qin, X.C., Xu, J., Holmes, E.C., Zhang, Y.Z. (2015). Unprecedented genomic diversity of RNA viruses in arthropods reveals the ancestry of negative-sense RNA viruses. Elife; 29;4. doi: 10.7554/eLife.05378.

Lin, X.D., Wang, W., Guo, W.P., Zhang, X.H., Xing, J.G., Chen, S.Z., Li, M.H., Chen, Y., Xu, J., Plyusnin, A. & Zhang, Y.Z. (2012). Cross-species transmission in the speciation of the currently known murinae-associated hantaviruses. J Virol; 86(20): 11171-82.

Longdon B., Hadfield, J.D., Webster, C.L., Obbard, D.J.& Jiggins, F.M. (2011). Host phylogeny determines viral persistence and replication in novel hosts. PLoS Pathog;;7(9):e1002260.

Luis, A.D., Hayman, D.T., O'Shea, T.J., Cryan, P.M., Gilbert, A.T., Pulliam, J.R., Mills, J.N., Timonin, ME, Willis, CK, Cunningham, AA, Fooks, AR, Rupprecht, CE, Wood, JL & Webb, C.T. (2014). A comparison of bats and rodents as reservoirs of zoonotic viruses: are bats special? Proc. Biol. Sci. 1;280(1756):20122753. doi: 10.1098/rspb.2012.2753.

Luis, A.D, O'Shea, T.J., Hayman, D.T, Wood, J.L., Cunningham, A.A., Gilbert, A.T., Mills, J.N. & Webb, C.T. (2015). Network analysis of host-virus communities in bats and rodents reveals determinants of cross-species transmission. Ecol Lett 24. doi: 10.1111/ele.12491.

Lund, M. (1994) Commensal rodents. In: Buckle A, Smith R (eds) Rodent pests and their control. CABI Publishing, Wallingford,23–43.

Lyles, D.D. & Rupprecht, C.E. (2007). Rhabdoviridae. In: Fiels Virology, eds. Knipe, D.M., Howley, P.P. pp. 1364-1379. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins, ©2007.

Mahalingam, S., Herrero, L.J, Playford, E.G., Spann, K., Herring, B., Rolph, M.S., Middleton, D., McCall, B., Field, H. & Wang, L.F. (2012). Hendra virus: an emerging paramyxovirus in Australia. Lancet Infect Dis; 12(10):799-807. doi: 10.1016/S1473-3099(12)70158-5. Epub 2012 Aug 24. Review.

Marí Saéz, A., Weiss, S., Nowak, K., Lapeyre, V., Zimmermann, F., Düx, A., Kühl, HS., Kaba, M., Regnau, t S., Merkel, K., Sachse, A., Thiesen, U., Villányi, L., Boesch, C., Dabrowski, PW., Radonić, A., Nitsche, A., Leendertz, SA., Petterson, S., Becker, S., Krähling, V., Couacy-Hymann, E., Akoua-Koffi, C., Weber, N., Schaade, L., Fahr, J., Borchert, M., Gogarten, J.F., Calvignac-Spencer, S.& Leendertz, F.H. (2014). Investigating the zoonotic origin of the West African Ebola epidemic. Mol Med; 30;7(1):17-23.

Markotter, W., Randles, J., Rupprecht, C.E., Sabeta, C.T., Taylor, P.J., Wandeler, A.I. & Nel L.H. (2006). Lagos bat virus, South Africa. Emerg Infect Dis. 12(3):504-6.

Marshall, A.G. (1983). Bats, flowers and fruit: evolutionary relationships in the Old World. Biological Journal of the Linnean Society, 20(1): 115-135.

Martin, A.P. & Palumbi, S.R. (1993). Body size, metabolic rate, generation time, and the molecular clock. Proc. Natl. Acad. Sci. U S A; 1;90(9):4087-91.

McCracken, G.F. & Gustin, M.K. (1991). Nursing behavior in Mexican free-tailed bat maternity colonies. Ethology; 89(4): 305-321.

Meerburg, B.G., Singleton, G.R. & Kijlstra, A. (2009). Rodent-borne diseases and their risks for public health. Crit Rev Microbiol; 35(3):221-70. doi: 10.1080/10408410902989837.

Merkle, D. & Middendorf, M. (2005). Reconstruction of the cophylogenetic history of related phylogenetic trees with divergence timing information. Theory Biosci;123(4):277-99. doi: 10.1016/j.thbio.2005.01.003.

Merkle, D., Middendorf, M. & Wieseke, N. (2010). A parameter-adaptive dynamic programming approach for inferring cophylogenies. BMC Bioinformatics. 18;11 Suppl 1:S60. doi: 10.1186/1471-2105-11-S1-S60.

Mickleburgh, S.P., Hutson, A.M. & Racey, P.A. 1992. Old World fruit bats: an action plan for their conservation. Gland, Switzerland, IUCN.

Mickleburgh, S., Waylen, K. & Racey, P. (2009). Bats as bushmeat: a global review. Oryx, 43(2): 217-234.

Morens, D.M., Folkers, G.K. & Fauci, A.S. (2004). The challenge of emerging and remerging infectious diseases. Nature 430(6996): 242-249.

Morand S., Krasnov B., Poulin R. (2006). Global change, biodiversity and the future of mammals-parasite interactions. In Micro-mammals and macroparasites: from evolutionary ecology to management Edited by: Morand S, Krasnov BR, Poulin R. Tokyo: Springer-Verla.;617-635.

Morand, S., Jittapalapong, S., Suputtamongkol, Y., Abdullah, M. T., & Huan, T. B. (2014). Infectious Diseases and Their Outbreaks in Asia-Pacific: Biodiversity and Its Regulation Loss Matter. PLoS ONE; 9(2), e90032. http://doi.org/10.1371/journal.pone.0090032

Morand, S., Bordes, F., Chen, H.W., Claude, J., Cosson, J.F., Galan, M., Czirják, G.Á., Greenwood A.D., Latinne, A., Michaux, J., & Ribas, A. (2015). Global parasite and Rattus rodent invasions: The consequences for rodent-borne diseases. Integr Zool. 10(5):409-23. doi: 10.1111/1749-4877.12143.

Morse, S.S. (1993). Examining the origins of emerging viruses. In: Morse, S.S. (Ed.), Emerging Viruses. Oxford University Press, New York, 10 – 28

Morse, S.S., Mazet, J.A, Woolhouse, M., Parrish, C.R., Carroll D., Karesh, W.B., Zambrana-Torrelio, C, Lipkin, W.I.,& Daszak, P. (2012). Prediction and prevention of the next pandemic zoonosis. Lancet; 380(9857): p. 1956-65.

Moss, W.J., Griffin, D.E. (2012). Measles. Lancet. Jan 14;379(9811):153-64. doi: 10.1016/S0140-6736(10)62352-5.

Muray, C.J.L. & Lopez, A.D, eds. (1996). The Global Burden of Disease: a comprehensive assessment of mortality and disability from diseases, injuries and risk factors in 1990 and projected to 2020. Cambridge, MA, Harvard School of Public Health on behalf of the World Health Organization and the World Bank, (Global Burden of Disease and Injury Series, Vol. I).

Myers N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A. & Kent J. (2000). Biodiversity hotspots for conservation priorities. Nature 24;403(6772):853-8.

Nagorsen, D.W. (2005) Rodents & lagomorphs of British Columbia. Royal British Columbia Museum, Victoria.

Nichol, S.T., Spiropoulou, C.F., Morzunov, S., Rollin, P.E., Ksiazek, T.G., Feldmann, H., Sanchez, A.J., Childs, J.E., Zaki, S.R., & Peters, C.J. (1993). Genetic identification of a hantavirus associated with an outbreak of acute respiratory illness, Science 262(5135), 914.

Nowak, R.M. & Walker, E.P. (1991) Walker's mammals of the world. Johns Hopkins University Press, Baltimore.

O' Brien, J. (2011). Bats of the Western Indian Ocean Islands. Animals; 1, 259-290; doi:10.3390/ani1030259

O'Shea, T.J., Bowen, R.A., Stanley, T.R., Shankar, V. & Rupprecht, C.E. (2014). Variability in seroprevalence of rabies virus neutralizing antibodies and associated in a colorado population of big brown bats (*Eptesicus fuscus*). PLoS ONE 9(1): e86261.

Orton, R.J, Wright, C.F., Morelli, M.J., Juleff, N., Thébaud, G., Knowles, N.J., Valdazo-González, B., Paton, D.J., King, D.P., & Haydon, D.T. (2013). Observing micro-evolutionary

processes of viral populations at multiple scales. Philos Trans R Soc Lond B Biol Sci.19; 368(1614): 20120203. doi: 10.1098/rstb.2012.0203.

Ostfeld, R. &, Keesing, F. (2000). Biodiversity and disease risk: the case of Lyme disease. Conserv Biol; 14:722-728.

Page, R.D.M. (2003). Tangled Trees. Phylogeny, Cospeciation and Coevolution. Ethology, (110): 577-578.

Parrish, C.R., Holmes, E.C., Morens, D.M., Park, E.C., Burke, D.S., Calisher, C.H., Laughlin, C.A., Saif L.J. & Daszak, P. (2008). Cross-species virus transmission and the emergence of new epidemic diseases. Microbiol. Molec. Biol. Rev. 72:457–470.

Paterson, A. (1997) Host - Parasite Evolution: General Principles & Avian Models. Oxford University Press;

Peake, J.F. (1971). The evolution of terrestrial faunas in the western Indian ocean. Phil. Trans. Roy. Soc. Lond. B Biol. Sci;, 260, 581-610.

Pepin, K.M., Lass, S., Pulliam, J.R.C., Read, A.F. & Lloyd-Smith, J.O. (2010). Identifying genetic markers of adaptation for surveillance of viral host jumps. Nat. Rev. Microbiol. 8:802–813.

Pierson, E.D. & Rainey, W.E. (1992). The biology of flying foxes of the genus Pteropus: A review. pp. 1-17 in: Pacific Island Flying Foxes: Proceedings of an international conservation conference (Wilson D. & Graham, G.L.. eds). Biological Report 90 (23). Washington DC, United States Department of the Interior Fish and Wildlife Service.

Plowright R.K., Eby P., Hudson P.J., Smith I.L., Westcott D., Bryden W.L., Middleton D., Reid P.A., McFarlane R.A., Martin G., Tabor G.M., Skerratt L.F., Anderson D.L., Crameri G., Quammen D., Jordan D., Freeman P., Wang L.F., Epstein J.H., Marsh G.A., Kung N.Y. & McCallum H. (2015). Ecological dynamics of emerging bat virus spillover. Proc. R. Soc. B ;282: 20142124.

Poulin, R. & Morand, S. (2004). Parasite biodiversity. Smithsonian Institution Press, Washington D.C.

Pulliam, J.R.C. (2008). Viral host jumps: moving toward a predictive framework. Ecohealth. ;5:80–91.

Rainey, W.E. & Pierson, E.D. (1992). Distribution of Pacific Island flying foxes. 111-122 in: Pacific Island Flying Foxes: Proceedings of an International Conservation Conference (Wilson D. & Graham, G.L. eds). Biological Report 90 (23). Washington DC, United States Department of the Interior Fish and Wildlife Service.

Ramsden, C., Holmes, E.C. & Charleston, M.A. (2009). Hantavirus evolution in relation to its rodent and insectivore hosts: no evidence for codivergence. Mol. Biol. Evol. 26(1): p. 143-53.

Real, L.A.,& Biek, R. (2007). Infectious disease modeling and the dynamics of transmission. Curr Top Microbiol Immunol; 315:33-49.

Reynes, J.M., Andriamandimby, S.F., Razafitrimo, G.M., Razainirina, J., Jeanmaire, E.M., Bourhy, H. & Heraud, J.M. (2011). Laboratory surveillance of rabies in humans, domestic animals, and bats in Madagascar from 2005 to 2010. Adv Prev Med. 2011; 2011:727821. Epub 2011 Aug 21.

Rohde, K. (1992). Latitudinal gradients in species diversity: The search for the primary cause. Oikos, Vol. 65, No. 3; 514-527

Saluzzo, J.F., P. Vidal, & J.P. Gonzales. (2004) Les virus émergents. IRD Éditions.

Sasaki, M., Setiyono, A., Handharyani, E., Rahmadani, I., Taha, S., Adiani, S., Subangkit, M., Sawa, H., Nakamura, I. & Kimura, T. (2012). Molecular detection of a novel paramyxovirus in fruit bats from Indonesia. Virol J; Oct 19;9:240. doi: 10.1186/1743-422X-9-240.

Schrag, S. J. & Wiener P. (1995). Emerging infectious disease: what are the relative roles of ecology and evolution? Trends Ecol. Evol; Aug;10(8):319-24.

Soarimalala, V. & Goodman, S.M. (2013). Les petits mammifères de Madagascar. Association Vahatra, Antananarivo.

Streicker, D.G., Turmelle, A.S., Vonhof, M.J., Kuzmin, I.V., McCracken, G.F. & Rupprecht, C.E. (2010). Host phylogeny constrains cross-species emergence and establishment of rabies virus in bats. Science 329, 676–679.

Streicker, D.G., Lemey, P., Velasco-Villa, A. & Rupprecht, C.E. (2012). Rates of viral evolution are linked to host geography in bat rabies. PLoS Pathog; 8(5): p. e1002720.

Strelkov, P.P. (1969). Migratory and stationary bats (Chiroptera) of the European part of the Soviet Union. Acta zool. 14: 393-439.

Sulkin, S. & Allen, R. 1974. Virus infections in bats. Monographs in Virology, 8: 1-103.

Taraschewski H. (2006). Hosts and parasites as aliens. J Helminthol; 80(2):99-128.

Taylor, L.H., Latham, S.M. & Woolhouse, M.E. (2001) Risk factors for human disease emergence. Philos. Trans. R. Soc. Lond. B. Biol Sci. 29;356(1411):983-9.

Teeling, E. C., Springer, M. S., Madsen, O., Bates, P., O'Brien, S. J., & Murphy, W. J. (2005). A molecular phylogeny for bats illuminates biogeography and the fossil record. Science 307:580–584.

Tong, S., Chern, S.W., Li, Y., Pallansch, M.A. & Anderson, L.J. (2008). Sensitive and broadly reactive reverse transcription-PCR assays to detect novel paramyxoviruses. J Clin Microbiol. 46(8):2652-8. doi: 10.1128/JCM.00192-08. Epub 2008 Jun 25.

Tordo, N., Ceccaldi, P.-E., Gaudin, Y. and Wandeler, A. I. (2010). Rhabdoviruses: Rabies. Topley and Wilson's Microbiology and Microbial Infections.

Tortosa, P., Pascalis, H., Guernier, V., Cardinale, E., Le Corre, M., Goodman, S.M. & Dellagi, K. (2012). Deciphering arboviral emergence within insular ecosystems. Infect Genet Evol. 12(6):1333-9. doi: 10.1016/j.meegid.2012.03.024. Epub 2012 Apr 4.

Turmelle, A.S. & Olival, K.J. (2009). Correlates of viral richness in bats (order Chiroptera). Ecohealth; 6(4): 522-539.

IUCN. (2010). [Online]. http://www.iucn.org/fr/.

van Valen, L. (1973). A new evolutionary law. Evol Theory1, 1-30.

Viralzone. 2015. [Online]. http://viralzone.expasy.org/all by species/22.html

Vidgen, M.E., de Jong, C., Rose, K., Hall, J., Field, H.E., Smith, C.S. (2015). Novel paramyxoviruses in Australian flying-fox populations support host-virus co-evolution. J Gen Virol; Jul;96(Pt 7):1619-25. doi: 10.1099/vir.0.000099. Epub 2015 Feb 20.

von Helversen, O. (1993). Adaptations of flowers to the pollination by glossophagine bats. In C.M.N.W. Barthlott, K. Schmidt-Loske and K.L. Schuchmann. Plant-animal interactions in tropical environments, pp. 41-59. Bonn, Germany, Museum Alexander Koening.

Wang, L.F. (2009). Bats and viruses: a brief review. Virol Sin; 24:93-99.

Wilkinson, D.A., Temmam, S., Lebarbenchon, C., Lagadec, E., Chotte, J., Guillebaud, J., Ramasindrazana, B., Héraud, J.M., de Lamballerie, X., Goodman, S.M., Dellagi, K. & Pascalis, H. (2012). Identification of novel paramyxoviruses in insectivorous bats of the Southwest Indian Ocean. Virus Res; 170(1-2):159-63. doi: 10.1016/j.virusres.2012.08.022. Epub 2012 Sep 7.

Wilkinson, D.A., Mélade, J., Dietrich, M., Ramasindrazana, B., Soarimalala, V., Lagadec, E., le Minter, G., Tortosa, P., Heraud, J.M., de Lamballerie, X., Goodman, S.M., Dellagi, K., Pascalis, H. (2014). Highly diverse morbillivirus-related paramyxoviruses in wild fauna of the southwestern Indian Ocean Islands: evidence of exchange between introduced and endemic small mammals. J Virol; 88(15):8268-77. doi: 10.1128/JVI.01211-14. Epub 2014 May 14.

Wilson, D.E. & Reeder, D.E.M (editors). (2005). Mammal species of the world. A taxonomic and geographic reference (3rd ed), Johns Hopkins University Press, 2,142 pp. (Available from Johns Hopkins University Press, 1-800-537-5487 or (410) 516-6900 <a href="http://www.press.jhu.edu">http://www.press.jhu.edu</a>).

Wong, S., Lau, S., Woo, P. & Yuen, K.Y. (2007). Bats as a continuing source of emerging infections in humans. Rev Med Virol. 17:67-91.

WHO. 2015. [Online]. <a href="http://apps.who.int/ebola/ebola-situation-reports">http://apps.who.int/ebola/ebola-situation-reports</a>.

WHO. 2014. [Online]. <a href="http://www.who.int/topics/sars/en/">http://www.who.int/topics/sars/en/</a>.

Woolhouse, M.E.J., Taylor, L.H. & Haydon, D.T. (2001). Population biology of multi-host pathogens. Science; 292, 1109–1112.

Woolhouse, M.E.J. & Gowtage-Sequeria, S. (2005). Host range and emerging and reemerging pathogens. Emerg. Infect. Dis; 11(12): 1842-1847.

Woolhouse, M. E. J. & Gaunt, E. (2007). Ecological origins of novel human pathogens. Crit. Rev. Microbiol; 33, 1–12.

Woolhouse, M.E.J, Scott, F., Hudson, Z., Howey, R. & Chase-Topping, M. (2012). Human viruses: discovery and emergence. Philos. Trans. R. Soc. Lond. B. Biol. Sci. 367: 2864–2871.


Yahner, R.H. (2001). Fascinating mammals: conservation and ecology in the mid-eastern states. University of Pittsburgh Press, Pittsburgh

Yadav, P.D., Raut, C.G., Shete, A.M., Mishra, A.C., Towner, J.S., Nichol, S.T., Mourya, D.T. (2012). Detection of Nipah virus RNA in fruit bat (*Pteropus giganteus*) from India. Am J Trop Med Hyg; 87(3):576-8. doi: 10.4269/ajtmh.2012.11-0416. Epub 2012 Jul 16.


Yoder, A.D., Olson, L.E., Hanley, C., Heckman, K.L., Rasoloarison, R., Russell, A.L., Ranivo, J., Soarimalala, V., Karanth, K.P., Raselimanana, A.P. & Goodman, S.M. (2005). A multidimensional approach for detecting species patterns in Malagasy vertebrates. Proc. Natl. Acad. Sci. USA; 102 Suppl 1: p. 6587-94.

## Annexes


**Annexe 1.** Phylogénie de la famille des *Paramyxoviridae* (Drexler et al., 2012).




**Annexe 2.** Structure schématique d'un paramyxovirus (Chang et al., 2012).



**Annexe 3.** Phylogénie et distribution géographique des lyssavirus (Banyard et al., 2014).



# **Annexe 4.** Structure générale (A) d'un virion et (B) d'un génome d'un Rhabdovirus (Source : Viralezone)



**Annexe 5.** Insight into the global evolution of Rodentia associated Morbillirelated Paramyxoviruses reveals the role of *Rattus* as a worldwide spreader (soumis)

Insight Into the Global Evolution of Rodentia Associated Morbillirelated Paramyxoviruses Reveals the Role of *Rattus* As A Worldwide Spreader Wissem Ghawar, a,b,c,d,# Hervé Pascalis, a,e,# Jihéne Bettaieb, b,c,d David Wilkinson, a,e Julien Mélade, a,e Adel ElGharbi, b,c,d Mohamed Ali Snoussi, b,c,d Dhafer Laouini, c,d Afif Ben Salah, b,c,d Koussay Dellagi, a,c,e Centre de Recherche et de Veille sur les maladies émergentes dans l'Océan Indien (CRVOI), Plateforme de Recherche CYROI, Sainte Clotilde, Reunion, France<sup>a</sup>; Laboratory of Medical Epidemiology, Institut Pasteur de Tunis (IPT), Tunis, Tunisia<sup>b</sup>; Laboratory of Transmission, Control and Immunobiology of Infections (LTCII), LR11IPT02, Institut Pasteur de Tunis (IPT), Tunis, Tunisia<sup>c</sup>; Université Tunis El Manar, Tunis, Tunisia<sup>d</sup>; UMR PIMIT-Processus Infectieux en Milieu Insulaire Tropical, INSERM 1187, CNRS 9192, IRD 249, Université de La Réunion, Réunion, France<sup>e</sup>. Running Head: Rodentia Associated Morbillirelated Paramyxoviruses #Address correspondence to <a href="mailto:ghawarwissemmed@yahoo.fr">ghawarwissemmed@yahoo.fr</a> or <a href="mailto:herve.pascalis@ird.fr">herve.pascalis@ird.fr</a>. W.G. and H.P. contributed equally to this work. Words count: abstract (196) and text (6674). 

#### Abstract

Paramyxoviruses (PVs) are a large and diverse family of viruses that affect humans and a wide range of animal species. Among this family, a group of *Unclassified Morbilli-Related Viruses* (*UMRVs*) has been recently recognized in wild small mammals, but its evolutionary history is still poorly understood. In the present study, 160 rodents belonging to four species, indigenous to Northern Africa, namely *Psammomys obesus*, *Ctenodactylus gundi*, and *Meriones shawi*, as well as the cosmopolitan *Rattus* (*R*.) *rattus* were captured in Tunisia and analyzed for PV infection using RT-PCR targeting part of the L-polymerase gene. PVs were detected in 49 out of 160 rodents yielding a global positivity rate of 30.6% and analyses allowed their identification as novel *Rodentia UMRVs*. Phylogenetic and phylogeographic analyses were conducted based on the *URMVs* sequences amplified from positive animals together with *URMVs* sequences from rodents originating in all continents and recovered from public databases. These analyses revealed three major different clusters among *Rodentia*-associated *UMRVs* and suggest that strain diversification occurred during geographic migration and host exchanges with purification selection pressure as the principal evolving force. *Rodentia*-associated *UMRVs* likely originated in Southern Africa and were mainly vectored by *R. rattus* to emerge worldwide.

#### **Importance**

Recently, novel Paramyxoviruses known as *Unclassified Morbilli-Related Viruses* (*UMRVs*, *Paramyxovirinae*, *Paramyxoviridae*) were reported to infect at high infection rates, various wild

small mammal orders. The current study shows those novel *UMRVs* are endemic to four wild

rodent species captured in the Center and the South of Tunisia. The highlighting of the history of

these *Rodentia* associated *UMRVs* was made possible because we compared endemic species to

highly localized species in North Africa. The genetic diversity detected among the *Rodentia* 

48 associated *UMRVs* and their phylogeography contributes to the understanding of their molecular

49 epidemiology and their natural evolutionary history. The study highlights and confirms the role

of *Rattus rattus* in the global dispersal of the *UMRVs*.

Our findings provide important knowledge on the epidemiology of the UMRVs and might help

52 understanding why they still worldwide spreading.

53

54

60

62

63

64

65

66

67

68

69

70

## Introduction

55 Rodentia is the most species rich mammalian order and has been found to carry the highest

number of zoonotic pathogens (1). Hence, eco-epidemiological studies on rodents as reservoirs of

57 pathogens are highly relevant to human health (2, 3) and to the design of control programs of

58 rodent-borne diseases.

59 Rattus (R.) rattus is by far the most important invasive rodent and reservoir host which vectors

dissemination of many pathogens globally (4, 5). This species has an Asian origin which dates

back to over one million years ago (middle Pleistocene age) as assessed by molecular clock

estimates (6) and the detection of fossils in Thailand(7, 8) and Java(9). Then, this species had a

global dispersal bringing significant implications to human health and economy.

Paramyxoviruses (PVs) are a family of negative-sense single-stranded RNA viruses that infect a large range of hosts including mammals, birds, reptiles and fish (10). The group includes established pathogens of humans (e.g. *Measles* and *Mumps* viruses), rodents (e.g. *Sendai* virus), avifauna (e.g. *Newcastle Disease* virus), livestock (e.g. *Rinderpest* virus), canids (e.g. *Canine Distemper* virus) and also new emerging zoonotic viruses of medical importance such as *Hendra* or *Nipah* viruses (10, 11). Some other PVs may have significant socio-economic impact such as the equine disease outbreak caused by *Salem* virus (12). Recently, novel PVs known as

Unclassified Morbilli-Related Viruses (UMRVs, Paramyxovirinae, Paramyxoviridae) were reported to infect at high infection rates, various wild small mammal orders (Rodentia, Afrosoricidae, Chiroptera) worldwide (13, 14). UMRVs are phylogenetically closely-related (14) but their molecular evolutionary history is still poorly understood. Heretofore, few UMRVs have been reported to be pathogenic to the animal host, such as the J and Belinga viruses inducing hemorrhagic lesions in rodents and bats respectively (15, 16), but, for most of them, there is a lack of information regarding their pathogenicity. Virus transmission between animals belonging to the same species is mainly horizontal suggesting that contaminated feces, urine or saliva are potentially contagious to other species including humans, living in contact (17).

In this study, we investigated the phylogeography and the evolutionary history of *UMRVs* among reservoir species belonging to the order *Rodentia* and the role played by the *R. rattus* in the global dissemination of these viruses. Our study was focused on PVs infecting rodents from Tunisia: three wildlife species, namely, *Psammomys* (*P.*) *obesus*, *Meriones* (*M.*) *shawi*, and *Ctenodactylus* (*C.*) *gundi*, all indigenous to the North African countries, and one cosmopolitan species, *R. rattus*. Then, using phylogenetic and phylogeographic tools, we compared sequences derived from these Tunisian rodents to those reported in public databases originating from rodents distributed worldwide, to identify the relationship between *UMRVs*, their geographic origins and their animal host reservoirs and to infer their origin as well as the mechanisms of their global evolution.

## Materials and Methods

# **Ethics Statement**

The research protocol was approved by the IRB at Institut Pasteur de Tunis. All animal experimentations in the field to collect animal samples and individual data on captured animals, were conducted in Tunisia, and were in agreement with the guidelines of International Guiding Principles for Biomedical Research Involving Animals. Study of Infection and analysis were performed at Centre de Recherche et de Veille sur les maladies Emergentes dans l'Océan Indien,

# 98 La Réunion, France.

# Sampling of rodents in Tunisia

Between 2009 and 2013, 784 wild rodents were trapped as previously described (18) in five districts belonging to two governorates of Tunisia: the districts of Mnara, Khbina and Ouled Mhamed were located in the governorate of Sidi Bouzid in the subhumid central Tunisia, and the districts of Germessa and Mdhila were located in the governorate of Tataouine in the arid southern Tunisia (Fig. 1). The sites of capture were geo-referenced by GPS and trapped rodents were transferred alive to the laboratory for physical examination and tissue sampling. Relevant parameters were recorded for each animal: species identification, sex and standard morphometric measurements. After cardiac exsanguination, small tissue fragments were cut from heart, liver, kidney, spleen, lungs and brain organs; snap frozen in liquid nitrogen and stored in deep freezer at -80°C until analysis. Rodent age was estimated by measuring the lenses weight as previously described (18).

Pearson Chi-Square and Fisher exact tests allowed comparison of categorical variables.

STATA software version 11 was used to carry out all statistical analysis.

#### Paramyxoviruses detection by RT-PCR

For each animal, total nucleic acids were extracted from approximately 1 mm<sup>3</sup> of lungs, kidneys and spleen using the viral mini kit v2.0 and an EZ1 BioRobot (QIAGEN). Tissues were homogenized in DMEM medium by TissueLyser (QIAGEN) for 2 min at 25 Hz using 3 mm

tungsten beads. cDNA was generated through reverse transcription (Promega cDNA kit). Seminested PCR targeting a partial sequence of the L-gene polymerase locus (~490 bp) of *Paramyxovirinae* subfamilies was carried out using previously described conditions (14, 19). RT-PCR products were purified using the QIAquick PCR purification kit (QIAGEN), cloned into the pGEMt-Easy vector (Promega), according to the manufacturer's instructions and sequenced using chain-termination (Sanger) sequencing (Big Dye Sequencing kit, ABI) on both strands by a commercial service (Genoscreen, Lille, France). Forty-four viral sequences from rodents caught in Tunisia were generated in the present study and were deposited in Genbank with the reference numbers: KJ408384 – KJ408428.

## Origin of the viruses and data sets construction

Phylogenetic and phylogeographic analyses were conducted using the 42 partial L-gene *UMRVs* sequences from rodents caught in Tunisia into a panel of 145 partial *UMRVs* sequences infecting rodents from over 11 countries, recovered from public databases spanning years 1962 to 2013. Only sequences with well-documented collection date and location origin were considered for analysis. Three data sets were used: the first set (Supplementary Table S1 in Supplementary material) was representative of most of PVs genera and allowed classification of Tunisian sequences among *Paramyxoviridae* family, the second set (Supplementary Table S2 in Supplementary material) was devoted to the phylogenetic and the phylogeographic analysis of the *Rodentia* associated *UMRVs*. The third data set recovered from Genbank, was focused only on *Measles* virus (MeV) (Supplementary Table S3 in Supplementary material) and was used to estimate the evolutionary rate of the L-gene polymerase locus.

# Phylogenetic analysis: maximum-likelihood and Bayesian inference analysis

Partial sequences of the polymerase of PVs infecting rodents from Tunisia were compared to published sequences in GenBank (National Center for Biotechnology Information, Bethesda,

cleaned for primers sequences, edited, assembled and compared using Geneious®Pro 7.1.8 (20). The multiple alignments of partial L-gene sequences were performed with MAAFT 7.0.17 algorithm (21). A GTR nucleotide substitution model with rate variation and a proportion of invariant sites (Gamma + I) was determined to best fit the data using Akaike Information Criterion in MEGA6 (22), and in ModelTest via the integrated Geneious Paup tool (23). The same best-fit substitution model was used for all phylogenetic analyses. Phylogenetic relationships among PVs strains based on partial L-gene sequences were analyzed using both Bayesian inference and maximum likelihood (ML) methodologies. ML trees were estimated, with 1000 rapid bootstraps, using RAXML 7.2.8 Geneious plugin, under the best-scoring ML tree parameter, and using Mega6 (22). Bayesian phylogenetic trees were supported by the topology of the ML trees (data not shown). Bayesian inference analysis were performed with the MrBayes 3.2.2 plugin included in Geneious software (24), with constrained and unconstrained branch lengths, four Metropolis-coupled chains for 10 million Markov Chain Monte Carlo (MCMC) generations, sampling of the Markov chain every 200 generations, discarding the 25% as burn-in. The sequences of bat-rabies virus (JQ595353) and rodents PVs (RodPVs) (HQ660195 and AB844426, 2 Respirovirus sequences) were chosen as outgroup in Fig. 1 and 2, respectively. Trees were visualized by FigTree 1.4.2 (www.tree.bio.ed.ac.uk/software/figtree/). The applicability of a molecular clock to the data set was evaluated with a log-likelihood-ratio test using the ML scores, with and without a molecular clock enforced. The null hypothesis of equal evolutionary rate throughout the trees was not rejected at a 5% significance level when assessed using Mega6 (22). As a consequence, all subsequent phylogenetic analyses were performed using a strict-molecular clock. For the Bayesian tree reconstructions no difference was observed between strict- and relaxed-molecular

USA) online (www.ncbi.nih.gov) using BLASTn and BLASTx. Raw sequences were then

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

clocks. However, for the phylogenetic analysis performed with Beast package 1.8.2 (25), we used a relaxed-clock Bayesian MCMC method, as previous studies have shown that it allowed a better fit to the data than a strict clock (25-27).

### Bayesian evolutionary analyses of spatio-temporal UMRVs dynamics

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

To explore the evolutionary relationships and times to the most recent common ancestors (t<sub>MRCAs</sub>) among UMRV lineages circulating at a global level, we reconstructed phylogenetic history using Bayesian MCMC analysis implemented in the Beast package. UMRVs sequences were annotated to one of 11 locations accordingly to the country of origin (Australia, China, Germany, Madagascar, Mayotte, Reunion, Seychelles, South Africa, Trinidad & Tobago, Tunisia and Zambia), and used as a discrete trait during MCMC analysis. We employed a GTR + G + I model of nucleotide substitution with a normally distributed rate variation among sites  $(3.48 \times 10^{-4}, \text{ sd})$  $1.69 \times 10^{-5}$ , values obtained from the MeV data set analysis (See Supplementary Text 1 and Supplementary Table S3 in Supplementary material)) and a relaxed (uncorrelated log-normal) molecular clock model. First, we specified a Bayesian skyline population coalescent model (10 piece-wise constant groups) (26), then, we investigated, with the same prior distribution, a nonparametric skygrid plot model (28). All chains were run for a good enough length and convergence was assessed using Tracer, with statistical uncertainty reflected in values of the 95% highest posterior density, removing at least 10% of the chain as burn-in. The probable locations of each ancestral node and evolutionary time past were summarized using an annotated Maximum Clade Credibility (MCC) phylogenetic tree. Posterior probability values provide an assessment of the degree of support for each node on the tree.

Lemey et al. (29) have implemented a continuous-time Markov chain (CTMC) over discrete sampling locations in the Beast package, making it possible to model spatial diffusion on

a time-scaled phylogenetic tree (29). To determine the different lineage migration patterns, inferred with a Bayesian stochastic search variable selection scheme (BSSVS), allowing the switch rates in the CTMC to be zero with some prior probability, we used either a CTMC symmetric (reversible: bidirectional between locations) or asymmetric (non-reversible: unidirectional route) substitution models for discrete geographic traits. Coalescent models were compared using a modified Akaike information criterion (AICM) in Tracer (30). The BSSVS procedure computes the most parsimonious possible rate values, by assigning a Bayes factor (BF) significance test between transmission routes, explaining the different forms of diffusion. The result has been calculated and visualized by Spread (31), with a BF cutoff greater than 3 taken as significant support, as previously reported (29, 32).

#### **Selection pressure**

First, we used a series of tests provided by MEGA6 (22). We conducted Tajima's test of neutrality, a statistical test to compare the number of segregating sites per site with the nucleotide diversity. The purpose of the test is to determine between sequences randomly evolving (neutrally) and those whose evolution is driven by a non-stochastic process (directional selection, balancing selection, demographic expansion or contraction, etc.). A Tajima's D = 0, means that the population dynamic evolves without evidence of selection. When D < 0, it means that population size is not at equilibrium, but expanding, due to an over representation of infrequent polymorphisms, typical after a bottleneck or a selective sweep and/or a purifying selection. On the opposite, if D > 0, it represents a decrease in population size and/or balancing selection due to an under representation of both low and high frequency polymorphisms. Based on the comparison of the numbers of Synonymous ( $d_S$ ) and Nonsynonymous ( $d_N$ ) substitutions between sequences, a codon-based Fisher's exact test has been conducted, and also an Estimate Selection for each codon (HyPhy). The latter computes the selection robustness, whether negative or

212 positive, acting on each of the corresponding codon and provides statistical estimates. 213 Considering the rather large size of our dataset, we made also a Z-Test to test the null hypothesis 214 that  $H_0$ :  $d_N = d_S$ . The level of significance at which  $H_0$  is rejected depends on the respective alternative hypotheses (HA): (a), (b), or (c) with (a)  $d_N \neq d_S$  (test of neutrality), (b)  $d_N > d_S$ 215 216 (positive selection), (c)  $d_N < d_S$  (purifying selection). Selection pressures were also measured 217 using SLAC/REL/FEL maximum likelihood methods, MEME, FUBAR, BGM, and DEPS/FADE 218 via the Datamonkey facility (33). The recombination detection program (RDP) implemented in 219 the RDP 4.46 software package (34) was also used.

# 220 Accession numbers generated in the present study

Forty-four viral sequences of *UMRV* from rodents caught in Tunisia were generated in the present study and were deposited in Genbank with the reference numbers: KJ408384–KJ408428.

223

224

225

226

227

228

229

230

231

232

233

234

## Results

Rodents belonging to four species namely P. obesus (Muridae), M. shawi (Muridae), R. rattus (Muridae) and C. gundi (Ctenodactylidae) were randomly selected (40 individuals per species) from a large panel of animals captured in Tunisia and tested by RT-PCR for infection with PVs (Table 1). PVs RNAs were detected in 49 out of 160 rodents yielding a global positivity rate of 30.6% (range 2.5 - 50% according to the rodent-species ( $p < 10^{-3}$ ) (Table 2). There were no significant differences in the infection rates according to sex, morphometric parameters, age and sampling site location.

#### PVs infecting wild rodents from Tunisia are UMRVs

(designated hereafter as RodPVs-Tun). In order to correctly assess their evolutionary position, we

Partial L-gene sequences of the PVs were obtained from 45 PVs positive rodents from Tunisia

performed a Bayesian phylogenetic analysis representing most of PVs genera. As shown in Fig. 1, RodPVs-Tun shape a strongly supported sister clade along with morbilliviruses (BP = 0.98) allowing to identify them as UMRVs.

RodPVs-Tun can be further subdivided into two discrete phylogroups (BP = 0.92). The first phylogroup shapes a clade that gathers all RodPVs-Tun hosted by P. obesus and most C. gundi, as well as the unique sequence from M. shawi. On a basal position to this clade, are the Mossman and Tupaia viruses (35, 36) as well as Salem virus (12). The second phylogroup gathers the rest of the RodPVs-Tun, i.e., all those detected in R. rattus and some PVs sequences from C. gundi, together with Tailam, Beilong and J viruses (15, 37, 38). In contrast to the phylogroup 1, where P. obesus and C. gundi clustered separately into two rather well-supported subclades (BP = 0.67 and 1 respectively), phylogroup 2 mixed together PVs sequences from C. gundi and R. rattus.

Altogether our sampling reveals that at least two main PVs lineages are co-circulating among rodents from Tunisia: *C. gundi* are infected by both lineages whereas *P. obesus*, *M. shawi* and *R. rattus* are apparently affected by only one. Interestingly, the viral lineage which infects *C. gundi* and *R. rattus* was found only in these rodents (despite the presence of some other species captured in the same region) and only from *C. gundi* trapped in the governorate of Sidi Bouzid.

#### Phylogeny of Rodentia associated UMRVs

Using a larger set of data available in GenBank, we then extended the phylogenetic analysis of RodPVs-Tun, to *UMRVs* hosted by rodents originating from various countries, worldwide. Rodents belonged to four families (*Nesomyidae*, *Ctenodactylidae*, *Cricetidae* and *Muridae*) and originated from eleven countries distributed on five continents i.e., Africa (Tunisia, Zambia, South Africa), America (Trinidad & Tobago), Asia (China), Australia, and Europe (Germany), as well as multiple islands from the Southwestern Indian Ocean (SWIO) region i.e., Madagascar,

Mayotte, Reunion, Seychelles. Noteworthy, a bias in sample representativeness, inherent to sequence availability in GenBank, is introduced by the fact that *Muridae* family included more than ten different species. The topology of the tree defines, at the global level, four relatively well-supported major paraphyletic clades (Fig. 2; BP = 0.78, 0.93, 1 and 1, respectively for clades I, II, III and IV). Each clade grouped *UMRVs* that were hosted by rodents belonging to various species and originating from distant countries (Fig. 2, branches of the phylogenetic tree are colored according to the geographic origins of the *UMRVs*).

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

Hence, Clade I includes UMRVs sequences infecting C. gundi and R. rattus collected in Tunisia together with sequences from Eliurus (E.) sp. and Rattus sp. from Madagascar and SWIO Islands, respectively. Similarly, UMRVs sequences from Rhabdomys (R.) pumilio, Mastomys (M.) natalensis, Apodemus (A.) flavicollis, Myodes glareolus and Microtus (M.) sp. from Germany, South Africa and Zambia also clustered in this clade I, together with sequences of Jeilong virus group (Rattus sp. group 2). Tailam and Beilong viruses are in basal position of a group comprising only UMRVs sequences from Rattus sp. from some SWIO Islands, and J virus is inserted between M. natalensis group 1 and sequences from A. flavicollis group. Clade II includes UMRVs sequences from P. obesus from Tunisia shaping a phyletic group associated with Rattus sp. sequences from the SWIO Islands (Rattus sp. group 3), M. natalensis group 2 from Zambia clustered with one R. pumilio sequence from South Africa. Most sequences isolated from C. gundi clustered with the only sequence from M. shawi and with a Malagasy R. rattus sequence (C. gundi group 2). This latter is sister group with another mixing group from Madagascar (E. minor group 2 and Rattus sp. group 3). Besides, clade II holds in a separate cluster two Zambian sequences of Mus (M.) Minutoides and one Rattus sequence from Reunion Island. Clade III brings together *Mossman* and *Nariva* viruses and *M. natalensis* group 3 from Zambia while the South African R. pumilio group 2 corresponds to Clade IV.

Obviously, the evolutionary history of <i>UMRVs</i> cannot be explained only on the basis of a
geographical structuring. For example, UMRVs from Tunisia were genetically strongly related to
UMRVs from SWIO Islands. In both clades I and II, closely related PVs infect C. gundi
indigenous to Northern Africa and Eliurus sp. endemic to Madagascar. The same holds true for
UMRVs circulating in Germany (A. flavicolis) and in Zambia (M. natalensis).
However, the topology of the tree reveals some host virus-species associations; for instance
UMRVs infecting Rattus sp. from the SWIO Islands define 3 groups clustered into two well-
supported lineages; Similarly, UMRVs infecting R. pumilio from South Africa feature 2 groups
clustered into two well-supported lineages; UMRVs infecting M. natalensis from Zambia, 3
groups clustered into three well-supported lineages; A. flavicollis, M. glareolus and Microtus sp.
from Germany (each rodents-species with one group) and P. obesus from Tunisia (one group).
Nonetheless, the evolutionary history of UMRVs infecting Rodentia suggests the occurrence of
different events. That closely structured UMRVs sequences are shared between different rodent
species, geographically close or distant, suggests the occurrence of host-jumps or/and spatial
diffusion. The fact that most of them involve Rattus sp. as a partner (represented by an asterisk in
Fig. 2), pinpoints the possible role of this species as diffusing vector. Thus, $UMRVs$ from $C$ .
gundi from Tunisia illustrate a species jump with R. rattus from Tunisia in clade I and another
jump with M. shawi from Tunisia and R. rattus from Madagascar in clade II. In both cases C.
gundi sequences were closely structured with those harbored by E. minor from Madagascar
which presents a spillover with <i>R. rattus</i> from the same Island.

# Phylogeography analysis of Rodentia associated UMRVs

Fig. 3 shows the generated Maximum Clade Credibility (MCC) tree (the branches of the tree have been colored according to the most probable location of their descendent nodes). Two main clades are segregated: clade A and clade C.

307	Clade C includes five UMRVs sequences harbored by R. pumilio from South Africa whereas
308	clade A is resolved into two major Subclades: subclade A1 groups sequences distributed
309	worldwide and subclade B includes only African sequences significantly segregating into four
310	distinct subgroups: Tunisia (B2); South Africa (B3); Zambia (B4) and some SWIO Islands i.e.,
311	Mayotte, Reunion and Madagascar (B5). Subclade A1 contains two major phylogroups: A2 and
312	A'2. The latter groups two Zambian sequences clustered with Tunisian and Malagasy sequences
313	(A'3) as well as with Nariva and Mossman viruses, originating from Trinidad & Tobago and
314	Australia, respectively. In fact, most UMRVs sequences infecting C. gundi from Tunisia clustered
315	with one sequence isolated among R. rattus from Madagascar (A'5) that significantly segregated
316	from Malagasy sequences from E. minor and R. rattus (A'4).
317	Phylogroup A2 segregated into several subclades (A3-A9): The remaining UMRVs sequences
318	identified in C. gundi from Tunisia were clustered with sequences hosted by E. minor from
319	Madagascar and Rattus sp. from SWIO Islands (A9). The latter significantly segregated from
320	another clade (A7), contains sequences of Rattus sp. from Reunion, Mayotte and Seychelles
321	clustered with Beilong and Tailam viruses from China. These sequences from Tunisia and SWIO
322	Islands were significantly clustered with sequences from South Africa isolated from R. pumilio
323	(A6). J virus significantly clustered with four sequences, two of them were from Germany and
324	the two other from Zambia, isolated respectively from M. musculus, A. flavicollis and M.
325	natalensis were significantly assigned with the clade A6 forming the clade A5. UMRVs
326	sequences from Germany identified in different rodents-species such as M. glareolus (forming a
327	monophyletic specific cluster), M. arvalis, M. agrestis and A. flavicollis were significantly
328	clustered with the clade A5 within the clade A4. The clade A3, consisting in two sequences from
329	Zambia and a sequence from Reunion, was clustered with the clade A4 forming the second major
330	phylogroup A2.

Statistical analysis of the phylogeographic tree showed that it most probably rooted in South Africa, with a posterior probability of 1.0. Zambia was also the most probable location of the two main clades A1 and B, and of the two major phylogroups A2 and A'2. Hence, *Rodentia* associated *UMRVs* sequences most probably originated in Africa and their currently circulating sequences have emerged from a common ancestor more than 4000 years ago. Until 2500 years before present, *Rodentia* associated *UMRVs* could be detected only in two African countries: South Africa and Zambia.

The most recent common ancestor (t<sub>MRCAs</sub>) of all *Rodentia* associated *UMRVs* from Australia and Trinidad & Tobago has most probably emerged 3000 years ago (clade A1; 95% HPD = 2121-3897) whereas European *Rodentia-UMRVs* (represented by Germany) emerged more recently (clade A4; 95% HPD = 1524-2794), and Asian (represented by China) was the last to diverge (clade A7; 95% HPD = 785-1563). However it should be noted that the current circulating *UMRVs* on SWIO Islands and Tunisia have a recent emergence, which does not exceed 200 years ago with the exception of other previous introductions in Reunion Island and Madagascar dating back to around 2100 (clade A3; 95% HPD= 1331-2951) and 1200 (clade A'4; 95% HPD= 779-1628) years ago, respectively.

Bayesian phylogeographic analysis showed 46 well-supported route linkages between countries. We used a Bayes factor (BF) cutoff of 3.0 to determine significance (Supplementary Table S4 in Supplementary material). Most roots involved African countries and SWIO Islands, suggesting that they played an important role in the migration of rodent associated *UMRVs*. The root with the strongest support was found between Zambia and Trinidad & Tobago with a BF of 207. Most roots begun with African Countries (17 with 8 roots begin with South Africa) followed by SWIO Islands (15 roots), Trinidad & Tobago (7 roots), Australia (4 roots) and Germany (3 roots). On the other hand, China was found on 10 roots but only in the end. The map with the

rates of transitions constructed using Spread is shown in Supplementary Fig. S1 (Supplementary material).

Analysis of the skygrid plot (Fig. 4a) showed that the effective number of *Rodentia UMRVs* infections at global level increased rapidly until around 1000 years ago. A further sharp increase in the number of infections occurred between 1000 and 600 years ago when it reached the maximum peak, after which the epidemic growth in diversity stopped and even decreased though it remained at a level higher than it was at the beginning.

The effective number of viral lineages depicted in Fig. 4b shows an increasing diversification of *UMRVs* strands over time through three well-defined successive phases. The first one, set between the remote ancestor and the 11<sup>th</sup> century, and the second one, comprised between the 11<sup>th</sup> century and 1900, are indicative of a linear relationship of diversification process over the time, nonetheless with a faster process regarding the first phase. The third and last phase, starting at 1900 to present, is clearly indicative of a nonlinear process over time, and shows a double increase of the whole diversification process in less than one hundred years.

## **Selection pressures**

The selection pressure operating in the *Rodentia UMRVs* studied here, have been estimated by different statistical methods provided by MEGA6 (data available upon request). Overall, these analyses revealed that purification selection is acting as the principal evolving force, with an abundance of negatively selected sites versus a little support for positive selection. Moreover, we computed different codon-specific substitution likelihood-based methods (Table 3) that accounted for discrete individual codon sites compared to *Peste-des-petits-ruminants* virus (PPRV), *Measles* virus (MeV), Canine distemper virus (CDV), and *Feline morbillivirus* (FMV) for the same L-gene region (33).

Codon-specific-analyses showed that none of the studied viruses showed evidence of positive selection (Table 3), except for UMRVs where only three positively selected sites have been inferred with the Mixed Effects Model Evolution (MEME) at a significance value of p < 0.1, and just one at p < 0.05. The latter is most appropriate to detect episodic diversifying selection affecting discrete codon sites. Interestingly, with DEPS method using amino acid sequences to identify directional evolution towards residues at sites, and valuable for the detection of selective sweeps, there is strong evidence that some sites are evolving under a directional selection in UMRVs, but probably not for the considered morbilliviruses, despite the small number of sequences used for testing. Indeed, no evidence of recombination has been detected among those viruses.

## **Discussion**

The infection rate with PVs in four wild rodent-species (*P. obesus*, *M. shawi*, *C. gundi* and *R. rattus*) captured within their natural biotopes in central and southern Tunisia was estimated at 30.6% with the highest rates found in *P. obesus* (50%) and *C. gundi* (47.5%). These rates are to our knowledge, the highest reported so far in rodents (13, 14, 39) other than prevalence of 68% recently reported in the red and gray squirrels in the United Kingdom (40).

Phylogenetic analysis shows that RodPVs-Tun sequences belong to *UMRVs*, as previously reported in other small mammal species (13, 14, 39) confirming the worldwide distribution of these viruses. Until recently, the *Rodentia* taxa harboring *UMRVs* spanned four families *Muridae*, *Cricetidae*, *Sciuridae* and *Nesomyidae* (13, 14, 39, 40). The present study enlarges this range to rodents of the *Ctenodactylidae* family (*C. gundi*) and adds two species of the *Muridae* family (*P. obesus* and *M. shawi*) as potential reservoir hosts.

Phylogenetically, RodPVs-Tun fell into two major clades. The first clade contains all sequences from *P. obesus*, *M. shawi* and most sequences from *C. gundi*, while the second clade includes sequences from *R. rattus* and some sequences from *C. gundi*. In fact, the two viral lineages affect in different ways the four rodent-species. The sedentary behavior of *P. obesus* which lives in special ecologic niches i.e. salty biotopes (18), may account for the restricted specificity of the viral strain to this species, in contrast to the other rodent species that show evidence of virus sharing. However, this species restriction may also just reflect a sampling bias and a larger investigation of *R. rattus* trapped in vicinity of *P. obesus* should help clarifying this issue. These sequences were closely-related to *Beilong* and *J* viruses. As the *Jeilong* viruses have been shown to have interchangable genome replication machineries (41), the same feature might also hold true for *UMRVs* that belong to the same phylogroup; however full genome sequencing are needed to ascertain this point.

At the global level, *UMRVs* are phylogenetically resolved into three major lineages: one is restricted to viruses from African countries including the SWIO Islands and the two others include *UMRVs* that are distributed worldwide. Clustering of viruses originating from distant locations i.e., Tunisia and Madagascar and the SWIO Islands or Trinidad & Tobago and Australia clearly attests to the absence of a strict geographical structuring among the dataset of sequences. Most interestingly, the fact that the phylogenetic tree retains at the global level the same topology observed with *UMRVs* that infect rodents from Tunisia, supports the highly diffusing character of these viruses and their capacity to infect a wide range of rodents, possibly through *R. rattus*. Viral sequences from *C. gundi*, a species endemic to North Africa, clustered in two separate clades with *UMRVs* sequences from *E. minor*, an endemic species from Madagascar. The fact that the reservoir species harboring genetically related *UMRVs* are scattered in distant countries strongly suggests the involvement of a third animal host which would act as a disseminating vector of

UMRVs. Rattus sp. is the ideal candidate to play this spreading role. Indeed, sequences from C.
gundi and E. minor closely cluster with viral sequences detected in R. rattus from Tunisia and
Madagascar. Several infection spillover events are actually detected that involve R. rattus as a
partner i.e., E. minor/R. rattus, C. gundi/R. rattus, M. natalensis/R. rattus, R. norvegicus/R.
rattus. Similar exchanges of UMRVs between R. rattus and other animal species belonging to
other orders such as Chiroptera and Afrosoricida were previously described (14) which further
stress the central role played by Rattus sp. in the transmission and the worldwide dissemination
of these UMRVs. Our phylogeographic model supports this view. We consider that the UMRVs'
distribution became a global phenomenon mainly as a consequence of the development of global
trade (42) that spread infected rats over the five continents as previously described for other
zoonoses (4, 5, 43). On the basis of this spatio-temporal reconstruction, we estimate that rodent
associated UMRVs currently circulating worldwide, originated some 4000 years ago in Southern
Africa. Some 500 years later, they split into two major viral lineages with a conservation of the
ancestral one: clade B which gives origin to some African and SWIO Islands viruses around 2200
years ago and clade A1 which diverged to worldwide strains at around 3000 years before the
present. $J$ and $Mossman$ viruses that were described in Australia, are closely-related to German
and Zambian viruses, and were introduced between 1400 and 1500 years ago, respectively.
Similarly Nariva virus from Trinidad & Tobago is closely-related to Zambian viruses and was
introduced later. Beilong and Tailam viruses from China were related to SWIO Islands except
Madagascar and were introduced about 1000 years ago. German viruses were introduced at about
2150 years ago. Noteworthy, the current circulation of Rodentia associated UMRVs from Tunisia
is much more recent as it does not exceed 200 years ago.
Previous studies have revealed many migrations events in the Murinea subfamily between
African regions and Asia during the last 12 million years (6, 44). Furthermore, dates of virus

introduction differ between the SWIO islands: the t<sub>MRCAs</sub> indicates dates at around 2100 and 1200 years ago for Reunion Island and Madagascar, respectively; but only 150 years ago for Seychelles and Mayotte. These dates are congruent with the evolutionary history of Rattus introductions into the SWIO Islands (6, 45, 46), except maybe for Reunion Island. It has been considered that Rattus introduction on Reunion began with the arrival of the first waves of colonization on the island at around 1600 AD whereas our estimation would place the introduction of Rattus much earlier. Although the latter estimate is plausible in light of oldest description of the island by some travelers (45), it is more likely that this gap reflects a bias in our data. In fact, this cluster is based solely on three sequences (two from M. musculus, whose migratory history is likely identical to that of *Rattus* (46, 47); and a single sequence from *Rattus* from Reunion Island), and consequently, without excluding a possibly poor estimation for that particular introduction event. For Madagascar, our results are consistent with estimates from the literature that places Rattus introduction at around 3000 before present (6, 46). For South Africa, studies on R. rattus did not determine good estimates for the introduction of this species in this country, but some authors believe that many introductions have occurred in this region, the oldest was directly from India and/or the Middle East possibly with early Arab trading activities (6). Outside we find a circulation of *UMRVs* in *R. pumilio* species which date back to around 250-500 years before present.

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

The possibility of different dates of virus introduction in the same location can be explained by the occurrence of multiple introductions of the black rat into the zone as demonstrated, for example, for *R. rattus* in Madagascar (46, 48). Likewise, the evolutionary history of *R. rattus* described by Aplin et al. (6) suggests that commensalism arose multiple times and in different geographic populations of black rats which is closely-related to the human history as well as to the host-pathogen coevolution in this group of small mammals (42, 43). To

our knowledge this is the first observation ever described, where natural history and evolution dynamics for a zoonotic virus, originating from wild fauna and hosted by wildlife species, could have been impacted by commensalism related to human activities.

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

On the basis of phylogeographically reconstructed flow rates, Rodentia-UMRVs migrated from Africa, worldwide. Interestingly, analysis of the population dynamics revealed that UMRVs genetic diversity increased continuously until around 1000 years ago. Then, an exponential increase in the viral diversity started at around the 10-11th century AD to reach a peak at around the 16<sup>th</sup> century. In terms of genetic diversity, this picture could be explained first, by the initiation and then deeper diversification, leading to a faster increase of the diverse polymorphisms, necessary to create new viral lineages from Africa. The decline in the genetic diversity that began on the 16<sup>th</sup> century and worsened over the last two hundred years until today reflects a dramatic overall reduction of viral diversity. This phenomenon could have resulted from several events: (i) an ancient drastic recombination event occurred in the past evolutionary history; (ii) a founder effect with multiple host-jump transmissions, a geographical large-scale expansion and global population bottleneck which purged preexisting genetic diversity, selectively determined. Regarding the first hypothesis, though we cannot exclude a remote infrequent recombination event (49), it is difficult to detect such a phenomenon, as multiple viral generations have taken place and purification selection can obscure the ancient age of viral lineages (50). Moreover, the presence of some breakpoints could be attributed to heterotachy, an important process in protein evolution (51). However, it is not the most suitable explanation, because if recombination can shuffle mutations and purge deleterious variants, generating genotypes better fitting to the new environment, yet, it still does not explain the persisting trend of the whole genetic diversity contraction over a long period.

Regarding the founder effect hypothesis (small population size) it could be a possible explanation, for both the increase, at different time/location points, of the viral diversification, and the boost of genetic drift with mutations accumulation, leading to a faster loss of diversity (52). Actually, the colonization by a few founders throughout periods of expansion, occurring at multiple occasions, at large geographic scale, can create population bottlenecks and an iterative loss of genetic diversity (53), and is in agreement with our data and the positive D Tajima value estimate because of the shift in genetic composition owing to sampling effects. Hence, UMRVs' global evolution would have been driven by a genetic drift compensated by a major purification selection, which is a mode of selection outlined by its effect on frequency distributions of trait values, and the slightly more exotic regimes of balancing selection (frequency-dependent selection versus convergent selection) as a consequence of multiple constraints combinations. Moreover, the weak evidence that some codons could be evolving under positive selection can indicate adaptive fine-tuning, may be as a result of cross-species transmission (54). Most interestingly the 11<sup>th</sup> century represented a dramatic turning point in human history of Europe and the Mediterranean. This is the beginning of the Crusades and Wars with their large population displacement and mixing, and the expansion on a large scale of the international trade (55). Between the 10<sup>th</sup> and the 15<sup>th</sup> century, political and socioeconomic changes sped up the opening up of regions, countries and continents: wars, large urban developments which were flourishing (56) and changes in land use and cultivation methods towards intensive agriculture (57). The recurrent bouts of bubonic Plague in which Rattus has played the role of the disseminating vector through maritime trade, illustrate the health consequences of such global revolution (43, 58). The history of black rat, a native of the Indian subcontinent, refers to one type of lineage (lineage I of the complex R. rattus) that became one of the most widespread species of mammals in the world (48). Rattus diffused worldwide accompanying human

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

displacements and maritime transport (59), including through the Oman Gulf and the Indian Ocean, two important commercial routes between Asia and the Mediterranean via the Eastern African coast (45, 60). In this context, the evolutionary history of the Rodentia UMRVs seems particularly closely connected to the black rat history. Besides, this could be evidenced by the fact that the number of lineages increased concomitantly with the decline in genetic diversity observed among the PVs populations (Fig. 4b). Strikingly, in all the UMRVs studied here, our estimates of the genetic diversity for the partial L-gene, are between 1000 and 5000 and, hence, considerably higher than those previously recorded for the structural N and H genes in MeV (61) and PPRV (62), reported by the respective authors themselves as being an unexpected very low genetic diversity. However, our results are congruent with other viruses such as HIV (63) and hepatitis C (64). From an epidemiological point of view, the increased number of lineages probably reflects the fact that Rattus established epidemiological bridges between different species. Since the 16<sup>th</sup> century, multiple exploratory expeditions from Europe to the rest of the world (such those of Alfonso de Albuquerque, Vasco de Gama, Ferdinand Magellan, James Cook (65), etc.) increased the dispersion of Rattus. In Madagascar, they coincided with an increasing anthropogenic pressure causing major ecological disturbances, the destruction of primary forests and habitat fragmentation (66) that have facilitated Rattus invasion, and the large expansion of this alien species over endemic species (67). In Europe and Africa, there was an intensification of agriculture and modern farming (68, 69). All these elements support the role of Rattus as a pathogen spreader at the local, regional and global scales. Further studies are needed to evaluate how other reservoirs hosts (particularly, Chiroptera and/or Afrosoricida), might have fueled the UMRVs natural history. In term of migration pattern, they will likely bring little to the story as most of the available sequences for these two orders are from species endemic to Madagascar However, it may change the host origin of the ancestral viral lineage as it is not known which

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

among the *Chiroptera*, *Tenrecidae* or *Rodentia* stands as the ancestral reservoir of *UMRVs* in Africa, from which these zoonotic viruses have started their global diffusion. The specificity of interactions between these viruses, their potential vectors and reservoir hosts, the co-infection with other pathogens and the consequences of infections in humans and animals constitute a research priority to predict future emerging infections with epidemic and pandemic potential.

# **Funding information**

This work was supported by Institut Pasteur de Tunis, Centre de Recherche et de Veille sur les maladies Emergentes dans l'Océan Indien and European Program RUN-Emerge made to Reunion University (FP7 capacities, REGPOT to W.G.). The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

#### References

- Meerburg BG, Singleton GR, Kijlstra A. 2009. Rodent-borne diseases and their risks
   for public health. Critical reviews in microbiology 35:221-270.
- 560 2. **Mills JN, Childs JE.** 1998. Ecologic studies of rodent reservoirs: their relevance for human health. Emerg Infect Dis **4:**529-537.
- Ramsey JM, Gutierrez-Cabrera AE, Salgado-Ramirez L, Peterson AT, SanchezCordero V, Ibarra-Cerdena CN. 2012. Ecological connectivity of Trypanosoma cruzi
  reservoirs and Triatoma pallidipennis hosts in an anthropogenic landscape with endemic
  Chagas disease. PloS one 7:e46013.

- 4. Hayman DT, McDonald KD, Kosoy MY. 2013. Evolutionary history of rat-borne
- Bartonella: the importance of commensal rats in the dissemination of bacterial infections
- 568 globally. Ecol Evol **3:**3195-3203.
- 569 5. Gage KL, Kosoy MY. 2005. Natural history of plague: perspectives from more than a
- 570 century of research. Annu Rev Entomol **50:**505-528.
- 6. Aplin KP, Suzuki H, Chinen AA, Chesser RT, Ten Have J, Donnellan SC, Austin J,
- 572 Frost A, Gonzalez JP, Herbreteau V, Catzeflis F, Soubrier J, Fang YP, Robins J,
- 573 Matisoo-Smith E, Bastos AD, Maryanto I, Sinaga MH, Denys C, Van Den Bussche
- RA, Conroy C, Rowe K, Cooper A. 2011. Multiple geographic origins of commensalism
- and complex dispersal history of Black Rats. PloS one **6:**e26357.
- 576 7. **Grayson DK.** 2001. The archaeological record of human impacts on animal populations.
- Journal of World Prehistory **15:**1-68.
- 578 8. **Bellwood P.** 1987. The prehistory of Island Southeast Asia: a multidisciplinary review of
- recent research. Journal of World Prehistory 1:171-224.
- 580 9. Musser G. 1984. Identities of subfossil rats from caves in southwestern Sulawesi. Mod
- Ouat Res Southeast Asia 8:61-94.
- 582 10. Lamb RA, Griffith DP. 2007. Paramyxoviridae: The Viruses and their replication.
- Fields' Virology 1:1449.
- 584 11. Field HE, Mackenzie JS, Daszak P. 2007. Henipaviruses: emerging paramyxoviruses
- associated with fruit bats. Current topics in microbiology and immunology **315:**133-159.
- 586 12. Renshaw RW, Glaser AL, Van Campen H, Weiland F, Dubovi EJ. 2000.
- 587 Identification and phylogenetic comparison of Salem virus, a novel paramyxovirus of
- 588 horses. Virology **270:**417-429.

- 589 13. Drexler JF, Corman VM, Muller MA, Maganga GD, Vallo P, Binger T, Gloza-
- Rausch F, Rasche A, Yordanov S, Seebens A, Oppong S, Adu Sarkodie Y,
- 591 Pongombo C, Lukashev AN, Schmidt-Chanasit J, Stocker A, Carneiro AJ, Erbar S,
- Maisner A, Fronhoffs F, Buettner R, Kalko EK, Kruppa T, Franke CR, Kallies R,
- Yandoko ER, Herrler G, Reusken C, Hassanin A, Kruger DH, Matthee S, Ulrich
- **RG**, Leroy EM, Drosten C. 2012. Bats host major mammalian paramyxoviruses. Nature
- 595 communications **3:**796.
- 596 14. Wilkinson DA, Melade J, Dietrich M, Ramasindrazana B, Soarimalala V, Lagadec
- E, le Minter G, Tortosa P, Heraud JM, de Lamballerie X, Goodman SM, Dellagi K,
- 598 **Pascalis H.** 2014. Highly Diverse Morbillivirus-Related Paramyxoviruses in Wild Fauna
- 599 of the Southwestern Indian Ocean Islands: Evidence of Exchange between Introduced and
- Endemic Small Mammals. J Virol 88:8268-8277.
- 501 15. **Jun MH, Karabatsos N, Johnson RH.** 1977. A new mouse paramyxovirus (J virus). The
- Australian journal of experimental biology and medical science **55:**645-647.
- 603 16. Maganga GD, Bourgarel M, Obame Nkoghe J, N'Dilimabaka N, Drosten C, Paupy
- 604 C, Morand S, Drexler JF, Leroy EM. 2014. Identification of an unclassified
- paramyxovirus in Coleura afra: a potential case of host specificity. PloS one 9:e115588.
- 606 17. Plowright RK, Field HE, Smith C, Divljan A, Palmer C, Tabor G, Daszak P, Foley
- **JE.** 2008. Reproduction and nutritional stress are risk factors for Hendra virus infection in
- 608 little red flying foxes (Pteropus scapulatus). Proc Biol Sci 275:861-869.
- 609 18. Ghawar W, Toumi A, Snoussi MA, Chlif S, Zaatour A, Boukthir A, Hamida NB,
- 610 Chemkhi J, Diouani MF, Ben-Salah A. 2011. Leishmania major infection among
- 611 Psammomys obesus and Meriones shawi: reservoirs of zoonotic cutaneous leishmaniasis
- 612 in Sidi Bouzid (central Tunisia). Vector Borne Zoonotic Dis 11:1561-1568.

- 19. Tong S, Chern SW, Li Y, Pallansch MA, Anderson LJ. 2008. Sensitive and broadly
- 614 reactive reverse transcription-PCR assays to detect novel paramyxoviruses. Journal of
- 615 clinical microbiology **46:**2652-2658.
- 616 20. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S,
- Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A.
- 618 2012. Geneious Basic: an integrated and extendable desktop software platform for the
- organization and analysis of sequence data. Bioinformatics **28:**1647-1649.
- 620 21. Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid
- multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res 30:3059-
- 622 3066.
- 22. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular
- 624 Evolutionary Genetics Analysis version 6.0. Molecular biology and evolution 30:2725-
- 625 2729.
- 626 23. **Posada D, Crandall KA.** 1998. MODELTEST: testing the model of DNA substitution.
- 627 Bioinformatics **14:**817-818.
- 628 24. Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Hohna S, Larget B,
- 629 Liu L, Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: efficient Bayesian
- phylogenetic inference and model choice across a large model space. Systematic biology
- **631 61:**539-542.
- 632 25. Drummond AJ, Rambaut A. 2007. BEAST: Bayesian evolutionary analysis by
- sampling trees. BMC evolutionary biology 7:214.
- 634 26. Drummond AJ, Ho SY, Phillips MJ, Rambaut A. 2006. Relaxed phylogenetics and
- dating with confidence. PLoS Biol 4:e88.

- 636 27. Ho SY, Duchene S. 2014. Molecular-clock methods for estimating evolutionary rates and
- timescales. Molecular ecology **23:**5947-5965.
- 638 28. Gill MS, Lemey P, Faria NR, Rambaut A, Shapiro B, Suchard MA. 2013. Improving
- Bayesian population dynamics inference: a coalescent-based model for multiple loci.
- Molecular biology and evolution **30:**713-724.
- 641 29. Lemey P, Rambaut A, Drummond AJ, Suchard MA. 2009. Bayesian phylogeography
- finds its roots. PLoS computational biology **5:**e1000520.
- 643 30. Baele G, Li WL, Drummond AJ, Suchard MA, Lemey P. 2013. Accurate model
- selection of relaxed molecular clocks in bayesian phylogenetics. Molecular biology and
- 645 evolution **30:**239-243.
- 646 31. Bielejec F, Rambaut A, Suchard MA, Lemey P. 2011. SPREAD: spatial phylogenetic
- reconstruction of evolutionary dynamics. Bioinformatics **27:**2910-2912.
- 648 32. Li WL, Drummond AJ. 2012. Model averaging and Bayes factor calculation of relaxed
- 649 molecular clocks in Bayesian phylogenetics. Molecular biology and evolution 29:751-
- 650 761.
- 651 33. Kosakovsky Pond SL, Frost SD. 2005. Not so different after all: a comparison of
- 652 methods for detecting amino acid sites under selection. Molecular biology and evolution
- **22:**1208-1222.
- 654 34. Martin DP, Lemey P, Lott M, Moulton V, Posada D, Lefeuvre P. 2010. RDP3: a
- flexible and fast computer program for analyzing recombination. Bioinformatics **26:**2462-
- 656 2463.
- 657 35. Miller PJ, Boyle DB, Eaton BT, Wang LF. 2003. Full-length genome sequence of
- Mossman virus, a novel paramyxovirus isolated from rodents in Australia. Virology
- **317:**330-344.

- 660 36. Tidona CA, Kurz HW, Gelderblom HR, Darai G. 1999. Isolation and molecular
- characterization of a novel cytopathogenic paramyxovirus from tree shrews. Virology
- **258:**425-434.
- 663 37. Woo PC, Lau SK, Wong BH, Wong AY, Poon RW, Yuen KY. 2011. Complete
- genome sequence of a novel paramyxovirus, Tailam virus, discovered in Sikkim rats. J
- 665 Virol **85:**13473-13474.
- 666 38. Li Z, Yu M, Zhang H, Magoffin DE, Jack PJ, Hyatt A, Wang HY, Wang LF. 2006.
- Beilong virus, a novel paramyxovirus with the largest genome of non-segmented
- negative-stranded RNA viruses. Virology **346:**219-228.
- 669 39. Sasaki M, Muleya W, Ishii A, Orba Y, Hang'ombe BM, Mweene AS, Moonga L,
- Thomas Y, Kimura T, Sawa H. 2014. Molecular epidemiology of paramyxoviruses in
- Zambian wild rodents and shrews. J Gen Virol 95:325-330.
- 672 40. Brooks F, Wood AR, Thomson J, Deane D, Everest DJ, McInnes CJ. 2014.
- Preliminary characterisation of Pentlands paramyxovirus-1, -2 and -3, three new
- paramyxoviruses of rodents. Vet Microbiol **170:**391-397.
- 675 41. Magoffin DE, Mackenzie JS, Wang LF. 2007. Genetic analysis of J-virus and Beilong
- virus using minireplicons. Virology **364:**103-111.
- 42. Jones EP, Eager HM, Gabriel SI, Johannesdottir F, Searle JB. 2013. Genetic tracking
- of mice and other bioproxies to infer human history. Trends in genetics: TIG 29:298-308.
- 679 43. Green MH, Symes C, Colet A, i Santiveri M, Xavier J, Ruíz J, Saula O, Subirà de
- 680 Galdàcano ME, Jáuregui C, DeWitte SN. 2014. TMG 1 (2014): Pandemic Disease in
- the Medieval World: Rethinking the Black Death, ed. Monica Green.
- 682 44. Lecompte E, Aplin K, Denys C, Catzeflis F, Chades M, Chevret P. 2008. Phylogeny
- and biogeography of African Murinae based on mitochondrial and nuclear gene

- sequences, with a new tribal classification of the subfamily. BMC evolutionary biology
- 685 **8:**199.
- 686 45. Cheke A. 2010. The timing of arrival of humans and their commensal animals on
- 687 Western Indian Ocean oceanic islands. Phelsuma Nature Protection Trust of Seychelles
- **18:**38-69.
- 689 46. Tollenaere C, Brouat C, Duplantier J-M, Rahalison L, Rahelinirina S, Pascal M,
- Mone H, Mouahid G, Leirs H, Cosson J-F. 2010. Phylogeography of the introduced
- 691 species Rattus rattus in the western Indian Ocean, with special emphasis on the
- 692 colonization history of Madagascar. Journal of Biogeography **37**:398-410.
- 693 47. **Duplantier JM, Orth A, Catalan J, Bonhomme F.** 2002. Evidence for a mitochondrial
- lineage originating from the Arabian peninsula in the Madagascar house mouse (Mus
- 695 musculus). Heredity (Edinb) **89:**154-158.
- 696 48. Brouat C, Tollenaere C, Estoup A, Loiseau A, Sommer S, Soanandrasana R,
- Rahalison L, Rajerison M, Piry S, Goodman SM, Duplantier JM. 2014. Invasion
- genetics of a human commensal rodent: the black rat Rattus rattus in Madagascar.
- 699 Molecular ecology **23:**4153-4167.
- 700 49. Chare ER, Gould EA, Holmes EC. 2003. Phylogenetic analysis reveals a low rate of
- 701 homologous recombination in negative-sense RNA viruses. J Gen Virol **84:**2691-2703.
- 702 50. Wertheim JO, Kosakovsky Pond SL. 2011. Purifying selection can obscure the ancient
- age of viral lineages. Molecular biology and evolution **28:**3355-3365.
- 704 51. Lopez P, Casane D, Philippe H. 2002. Heterotachy, an important process of protein
- evolution. Molecular biology and evolution **19:**1-7.
- 706 52. **Kimura M.** 1991. The neutral theory of molecular evolution: a review of recent evidence.
- 707 Jpn J Genet **66:**367-386.

- 708 53. Nei M, Maruyama T, Chakraborty R. 1975. The bottleneck effect and genetic
- variability in populations. Evolution:1-10.
- 710 54. Pepin KM, Lass S, Pulliam JR, Read AF, Lloyd-Smith JO. 2010. Identifying genetic
- 711 markers of adaptation for surveillance of viral host jumps. Nat Rev Microbiol 8:802-813.
- 712 55. **Powell JM.** 2002. The Crusades from the Perspective of Byzantium and the Muslim
- 713 World. The Catholic Historical Review 88.
- 714 56. Davis K. 1955. The origin and growth of urbanization in the world. American Journal of
- 715 Sociology:429-437.
- 716 57. White Jr L. 1967. The Historical Roots of Our Ecologic Crisis. Science 155:1203-1207.
- 717 58. Kruse H, Kirkemo A-M, Handeland K. 2004. Wildlife as source of zoonotic infections.
- 718 Emerging Infectious Diseases **10:**2067-2072.
- 719 59. Searle JB. 2008. The genetics of mammalian invasions: a review. Wildlife Research
- 720 **35:**185-192.
- 721 60. Verin P, Wright H. 1999. Madagascar and Indonesia: new evidence from archaeology
- 722 and linguistics. Bulletin of the Indo-Pacific Prehistory Association 18:35-42.
- 723 61. **Pomeroy LW, Bjornstad ON, Holmes EC.** 2008. The evolutionary and epidemiological
- dynamics of the paramyxoviridae. Journal of molecular evolution **66:**98-106.
- 725 62. **Padhi A, Ma L.** 2014. Genetic and epidemiological insights into the emergence of peste
- des petits ruminants virus (PPRV) across Asia and Africa. Sci Rep **4:**7040.
- 727 63. Robbins KE, Lemey P, Pybus OG, Jaffe HW, Youngpairoj AS, Brown TM, Salemi
- 728 M, Vandamme AM, Kalish ML. 2003. U.S. Human immunodeficiency virus type 1
- 729 epidemic: date of origin, population history, and characterization of early strains. J Virol
- 730 **77:**6359-6366.

- 731 64. Pybus OG, Charleston MA, Gupta S, Rambaut A, Holmes EC, Harvey PH. 2001.
- The epidemic behavior of the hepatitis C virus. Science **292:**2323-2325.
- 733 65. Cook J, Beaglehole JC. 1968. The voyage of the Endeavour, 1768-1771. Published for
- the Hakluyt Society at the University Press, Cambridge [England.
- 735 66. Harper GJ, Steininger MK, Tucker CJ, Juhn D, Hawkins F. 2007. Fifty years of
- deforestation and forest fragmentation in Madagascar. Environmental Conservation
- 737 **34:**325-333.
- 738 67. Goodman SM. 1995. Rattus on Madagascar and the dilemma of protecting the endemic
- rodent fauna. Conservation Biology:450-453.
- 740 68. Bouma J, Varallyay G, Batjes N. 1998. Principal land use changes anticipated in
- Europe. Agriculture, ecosystems & environment **67:**103-119.
- 742 69. Koning N, Smaling E. 2005. Environmental crisis or 'lie of the land'? The debate on soil
- 743 degradation in Africa. Land use policy **22:**3-11.
- 744
- 745

# Figure legends

746

768

747 Fig. 1. Maximum likelihood phylogeny of the partial L-gene of the *Paramyxoviridae* family 748 including the Tunisian paramyxoviruses. PV sequences are distributed according to the 749 geographic location of sampling sites and are colored according to the host species: Psammomys 750 obesus, Meriones shawi, Rattus rattus and Ctenodactylus gundi colored in red, green, blue and 751 purple; respectively. Values at node points indicate Bayesian posterior probabilities. Virus 752 designations were as follow: Virus name or virus affiliation/typical host/accession 753 number/origin/collection year/host Subfamily/host Family. Abbreviations used are detailed in 754 Supplementary Table S1 in Supplementary material. Rabies virus (JQ595353) was used as an 755 outgroup. 756 Fig. 2. Bayesian phylogeny of the partial L-gene of Rodentia-associated UMRVs infecting 757 Tunisian-rodents and sequences on GenBank. The branches colors are represented according to 758 the geographic origin of the sequences as specified in the figure. Groups are identified according 759 to the host species of the UMRVs. Viruses preceded by an asterisk indicate a spillover event 760 between Rattus sp. and other Rodentia species. Values at node points indicate Bayesian posterior 761 probabilities. Virus designations are as follows: Virus name or virus affiliation/typical 762 host/accession number/origin/collection year/host Subfamily/host Family. Abbreviations used are 763 detailed in Supplementary Table S2 in Supplementary material. Two Respirovirus sequences 764 (HQ660195 and AB844426) were used as an outgroup. 765 Fig. 3. Maximum clade credibility tree of Rodentia associated UMRVs. The phylogenetic 766 relationships and temporal evolutionary history have been estimated by molecular clock analysis. 767 Branch lengths are temporally scaled, and the x-axis shows the time unit (years before present).

The branches of the tree are colored on the basis of the most probable location of their descendent

769	nodes as specified in the figure. Geographical groups are identified according to the origin of the
770	UMRVs. Viruses preceded by small Colored Square indicate a spillover event between Rattus sp.
771	and other Rodentia species with the color indicating the location of this spillover. Values at node
772	points indicate Bayesian posterior probabilities and the 95% HPD. Virus designations are as
773	follows: Virus name or virus affiliation/typical host/accession number/origin/collection year/host
774	Subfamily/host Family. Abbreviations used are detailed in Supplementary Table S2 in
775	Supplementary material.
776	Fig. 4. Bayesian skygrid plots (BSP), inferred from partial L-gene, for Rodentia associated
777	UMRVs. BSP depict viral population dynamics and the changing levels of genetic diversity (a)
778	and the effective number of viral lineages (b) (y axis; log10 scale) over time (x axis; calendar
779	years) for Rodentia associated UMRVs lineages, showing the median estimate (solid line) and
780	credibility interval (blue area). The vertical dotted line represents the upper limit of the root
781	height, with the mean t <sub>MRCAs</sub> at the origin.
782	
783	

Table 1. Morphometric and demographic descriptions of rodents from Tunisia: *Psammomys (P.)* obesus, Meriones (M.) shawi, Rattus (R.) rattus and Ctenodactylus (C.) gundi.

	P. obesus	M. shawi	R. rattus	C. gundi
Governorates (n (%))				
Sidi Bouzid	40 (100)	40 (100)	40 (100)	20 (50)
Tataouine	/	/	/	20 (50)
Sex (n (%))				
Male	23 (57.5)	15 (37.5)	22 (55)	20 (50)
Female	17 (42.5)	25 (62.5)	18 (45)	20 (50)
Standard morphometric parameters (Mean±SD)				
Weight (g)	124±24.9	81.83±26.34	98.13±33.67	224.53±54.85
Ear length (mm)	16.15±1.02	18.06±2.78	21.63±1.69	18.08±1.28
Head and body length (mm)	153.15±9.88	163.33±28.31	163.35±24.7	198.1±16.31
Tail length (mm)	128.28±7.81	119.09±64.37	181.43±60.91	28.75±9.87
Hind Feet length (mm)	36.18±1.33	36.23±4.17	33.08±2.55	41.93±2.26
Eyes lenses weight (Mean±SD)				
ELW (g)	39.21±6.77	66.19±20.73	52.05±18.41	92.50±32.57

Table 2. Detection by RT-PCR of paramyxovirus infection in rodents from Tunisia according to
 the sampling site and to the biometric parameters. *Psammomys* (*P*.) *obesus*, *Meriones* (*M*.) *shawi*,
 *Rattus* (*R*.) *rattus* and *Ctenodactylus* (*C*.) *gundi*.

	P. obesus	M. shawi	R. rattus	C. gundi
	(n = 40)	(n = 40)	(n = 40)	(n = 40)
NV 1.6 (1 (0) P	20	1	9	19
PVs infection (%, P)	$(50, < 10^{-3})$	$(2.5, < 10^{-3})$	$(22.5, <10^{-3})$	$(47.5, <10^{-3})$
According to the capture sites (n (%))				
Sidi Bouzid	20 (100)	1 (100)	9 (100)	11 (57.9)
Tataouine	1	/	1	8 (42.1)
P	/	/	1	NS
According to gender (n (%))				
Male	14 (70)	1 (100)	6 (66.7)	8 (42.1)
Female	6 (30)	/	3 (33.3)	11 (57.9)
P	NS	/	NS	NS
According to the standard morphometric p	parameters (Mean±SD (P))			
Weight (g)	129±70 (NS)	35	96.67±28.28 (NS)	227.95±46.30 (NS)
Ear length (mm)	16.35±1.13 (NS)	14.00	21.67±0.86 (NS)	18.21±1.22 (NS)
Head and body length (mm)	155.30±10.09 (NS)	105.00	164.89±16.72 (NS)	199.11±11.65 (NS)
Tail length (mm)	125.95±7.30 (NS)	152.00	199.11±22.72 (NS)	28.16±9.48 (NS)
Hind Feet length (mm)	36.00±1.37 (NS)	27.00	33.44±1.59 (NS)	42.21±1.58 (NS)
According to the eyes lenses weight (Means	±SD (P))			
ELW (g)	41.21±4.39 (NS)	40	47.70±21.31 (NS)	93.86±27.76 (NS)

793 NS: Not Significant.

**Table 3.** Summary of selection pressures acting in *Rodentia UMRVs* versus Morbilliviruses.

	Codon-bas	sed nucleic a	icid gene	level									Protein g	gene level	
Genus	N° of seq (Partial	D Tajima Value		AC <sup>a</sup> Positive		EL <sup>a</sup> vely sele Neg.	ected sites	EL <sup>a</sup> Neg.	MEME <sup>a</sup> (codon position)	FUBAR <sup>b</sup>	Integrative Selection (at least one	BGM <sup>c</sup> (codon position)	N° of seq (Partial L-gene)	DEPS (N° residues)	FADE
Rodent	L-gene)	+4.2839	None	131	None	138	NA	NA	3 (31, 62,	142	method)	3 (30, 84,	80	62 (19)	72
UMRVs PPRV	15	+0.0925	None	12	None	24	None	71	None	22	71	94) None	6	None	4
MeV	17	-1.0352	None	6	None	11	None	None	None	7	11	None	10	None	4
CDV	11	-0.4934	None	6	None	19	None	None	None	14	20	None	6	None	1
FMV	6	-0.2620	None	1 (109)	None	28	None	None	None	12	28	None	5	None	None

a p < 0.1 (no positively selected sites were detected, except for one site (62) with MEME at p < 0.05). b Posterior prob > 0.90. c Posterior prob > 0.95. seq: sequences.
 For nucleic acid level, we used the HKY85 model, except for Rod-UMRVs (GTR) with input neighbor-joining trees. For protein gene level, we used the JTT model. Directional Evolution Protein Sequences (DEPS) in UMRVs (selective sweeps): 23 residues substitutions evolving under a Frequency-dependent selection, 73 residues substitutions evolving under a Convergent evolution and 18 sites evolving under a Balancing selection (Frequency-dependent selection versus Convergent evolution). Except for the D Tajima statistical test (Mega6), all the analyses have been performed via the Datamonkey facility.





Fig. 1. Maximum likelihood phylogeny of the partial L-gene of the Paramyxoviridae family including the Tunisian paramyxoviruses. PV sequences are distributed according to the geographic location of sampling sites and are colored according to the host species: Psammomys obesus, Meriones shawi, Rattus rattus and Ctenodactylus gundi colored in red, green, blue and purple; respectively. Values at node points indicate Bayesian posterior probabilities. Virus designations were as follow: Virus name or virus affiliation/typical host/accession number/origin/collection year/host Subfamily/host Family. Abbreviations used are detailed in Supplementary Table S1 in Supplementary material. Rabies virus (JQ595353) was used as an outgroup.







Fig. 4. Bayesian skygrid plots (BSP), inferred from partial L-gene, for Rodentia associated UMRVs. BSP depict viral population dynamics and the changing levels of genetic diversity (a) and the effective number of viral lineages (b) (y axis; log10 scale) over time (x axis; calendar years) for Rodentia associated UMRVs lineages, showing the median estimate (solid line) and credibility interval (blue area). The vertical dotted line represents the upper limit of the root height, with the mean tMRCAs at the origin.



### 1 Supplementary Material:

**Supplementary Figure and Table Legends:** 

2

Supplementary Figure S1: The hypothetical worldwide dispersion patterns of Rodentia 3 4 associated UMRV isolates as reconstructed on the basis of significant genetic flow rates and 5 dated phylogeny. Only the rates supported by a Bayes Factor (BF) greater than 3 are shown. 6 The map was reconstructed using SPREAD. 7 Supplementary Table S1: Viruses used for the classification of RodPVs-Tun among 8 Paramyxoviridae family in Figure 1. 9 Supplementary Table S2: Viruses used for the phylogenetic and phylogeographic analysis of 10 Rodentia associated UMRV in Figures 2 and 3. 11 Supplementary Table S3: Viruses used for the estimate of the evolutionary rate change for 12 the partial L-gene among the Measles virus. 13 Supplementary Table S4: Significant diffusion rates between countries at Bayesian 14 phylogeographic analysis. Posterior probability of the rates k (pk) and Bayes factor (BF) of 15 the significant migration rates as revealed by the Skygrid symmetric model. 16 17 18 19 20 21 22 23

### 24 Supplementary Text 1: Estimate of the evolutionary rate change for the partial L-gene

### 25 among the Measles virus.

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

As Rodentia PVs presented in this study are very close Morbilli-related viruses (1-3) and for some authors probably could be considered as sister genera (4) we used the Measles virus (MeV) as a reference for the analysis of evolutionary changes of UMRVs. Out of 189 complete genomes, 24 MeV partial L-gene sequences globally distributed have been selected according to the criterion of the date of collection availability, spanning the years between 1989 and 2014 (Supplementary Table S3). Besides, one might think that only 24 sequences of the MeV may lead to inaccurate estimates; however, to compensate this aspect, the L-gene may provide the most reliable evolutionary signal at the family level. The best-fit substitution model was K2 + G, but we used HKY85 + G the nearest close relative best-fit model available in Beauti (Beast package 1.8.2). We used serial samples of MeV for the partial Lgene to estimate the rate of nucleotide substitution and the time to the most recent common ancestor (t<sub>MRCAs</sub>), using the Bayesian MCMC method with a coalescent skyline plot method, allowing the analysis of the distribution branch length among collected viruses at different times sampled from millions of trees. We performed a lognormal relaxed molecular clock (5) and a random local clock (RLC) (6), either with a constant population size or a skyline demographic model. The RLC model submits and analyses a succession of different local molecular clocks, each potentially occurring on any branch and extending over a contiguous part of the phylogeny. Runs were carried out with chain lengths of 100-200 millions. The output from Beast was analyzed using the program Tracer 1.6 (http://beast.bio.ed.ac.uk/Tracer). Obtained mean rate was used to the subsequent UMRVs phylogeographic analysis. The majority of the set comprising the 145 sequences of UMRVs felt in a too small time interval, introducing too much bias to determine empirically the evolutionary history and

migration patterns of these viruses directly from our data set. This is due to the fact that different field campaigns to collect samples have all been conducted between 2009 and 2013. However, it is reasonable to assume that determining the evolutionary rate change a virusrelated of the same genera or from a virus that proceeds to the same common ancestor could remain a good alternative and a reliable approximation in our case (L-gene). Parameter estimates were consistent among models and similar values were obtained for all the analyses. The evolutionary rate of the current circulating MeV for the L-gene was estimated to be 3.48 x 10<sup>-4</sup> substitutions per site per year (95% HPD 3.94 x 10<sup>-5</sup>, 6.74 x 10<sup>-4</sup>), and coalescent estimates place its recent emergence at around 1900 (~116 years from 2014, 95% HPD 30.46 to 245.53). The mean evolutionary rate and the  $t_{MRCAs}$  derived from the partial L-gene were consistent and quietly similar to those already described for other various genes of the current circulating MeV. However, our values were slightly lower than those obtained for the structural H and N genes (7, 8), but it was not unexpected, being considered that the polymerase gene is the most conserved gene among the Paramyxoviridae to maintain crucial structural domains such as multiple functions related to different replicative mechanisms (4, 9). This molecular clock has been used for the phylogeographic analysis.

65

66

67

68

69

70

71

72

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

1. Drexler JF, Corman VM, Muller MA, Maganga GD, Vallo P, Binger T, Gloza-Rausch F, Rasche A, Yordanov S, Seebens A, Oppong S, Adu Sarkodie Y, Pongombo C, Lukashev AN, Schmidt-Chanasit J, Stocker A, Carneiro AJ, Erbar S, Maisner A, Fronhoffs F, Buettner R, Kalko EK, Kruppa T, Franke CR, Kallies R, Yandoko ER, Herrler G, Reusken C, Hassanin A, Kruger DH, Matthee S, Ulrich RG, Leroy EM, Drosten C. 2012. Bats host major mammalian paramyxoviruses. Nature communications 3:796.

- 73 2. Wilkinson DA, Melade J, Dietrich M, Ramasindrazana B, Soarimalala V,
- Lagadec E, le Minter G, Tortosa P, Heraud JM, de Lamballerie X, Goodman
- 75 SM, Dellagi K, Pascalis H. 2014. Highly Diverse Morbillivirus-Related
- 76 Paramyxoviruses in Wild Fauna of the Southwestern Indian Ocean Islands: Evidence
- 77 of Exchange between Introduced and Endemic Small Mammals. J Virol 88:8268-
- 78 8277.
- 79 3. Sasaki M, Muleya W, Ishii A, Orba Y, Hang'ombe BM, Mweene AS, Moonga L,
- Thomas Y, Kimura T, Sawa H. 2014. Molecular epidemiology of paramyxoviruses
- in Zambian wild rodents and shrews. J Gen Virol **95:**325-330.
- 82 4. McCarthy AJ, Goodman SJ. 2010. Reassessing conflicting evolutionary histories of
- the Paramyxoviridae and the origins of respiroviruses with Bayesian multigene
- phylogenies. Infection, genetics and evolution : journal of molecular epidemiology
- and evolutionary genetics in infectious diseases **10:**97-107.
- 86 5. Drummond AJ, Ho SY, Phillips MJ, Rambaut A. 2006. Relaxed phylogenetics and
- dating with confidence. PLoS Biol **4:**e88.
- 88 6. **Drummond AJ, Suchard MA.** 2010. Bayesian random local clocks, or one rate to
- rule them all. BMC Biol 8:114.
- 90 7. Pomerov LW, Bjornstad ON, Holmes EC. 2008. The evolutionary and
- 91 epidemiological dynamics of the paramyxoviridae. Journal of molecular evolution
- 92 **66:**98-106.
- 93 8. Furuse Y, Suzuki A, Oshitani H. 2010. Origin of measles virus: divergence from
- rinderpest virus between the 11th and 12th centuries. Virol J 7:52.
- 95 9. Cox R, Plemper RK. 2015. The paramyxovirus polymerase complex as a target for
- 96 next-generation anti-paramyxovirus therapeutics. Front Microbiol **6:**459.



**Supplementary Figure S1:** The hypothetical worldwide dispersion patterns of *Rodentia* associated *UMRV* isolates as reconstructed on the basis of significant genetic flow rates and dated phylogeny. Only the rates supported by a Bayes Factor (BF) greater than 3 are shown. The map was reconstructed using SPREAD.

**Supplementary Table S1:** Viruses used for the classification of RodPVs-Tun among
108 *Paramyxoviridae* family in Figure 1.

Virus ID <sup>a,b</sup>	Accession	Country	Collection	Host
VIEWS ID	numbers	Country	year	Host
RodPV/C. gundi/KJ408415/Tunisia/2010/nd/Ctenodactylidae	KJ408415	Tunisia	2010	Ctenodactylus gundi
RodPV/C. gundi/KJ408417/Tunisia/2010/nd/Ctenodactylidae	KJ408417	Tunisia	2010	Ctenodactylus gundi
RodPV/C. gundi/KJ408418/Tunisia/2010/nd/Ctenodactylidae	KJ408418	Tunisia	2010	Ctenodactylus gundi
RodPV/C. gundi/KJ408419/Tunisia/2010/nd/Ctenodactylidae	KJ408419	Tunisia	2010	Ctenodactylus gundi
RodPV/R. rattus/KJ408424/Tunisia/2013/Murinae/Muridae	KJ408424	Tunisia	2013	Rattus rattus
RodPV/R. rattus/KJ408422/Tunisia/2013/Murinae/Muridae	KJ408422	Tunisia	2013	Rattus rattus
RodPV/R. rattus/KJ408423/Tunisia/2013/Murinae/Muridae	KJ408423	Tunisia	2013	Rattus rattus
RodPV/R. rattus/KJ408420/Tunisia/2013/Murinae/Muridae	KJ408420	Tunisia	2013	Rattus rattus
RodPV/R. rattus/KJ408421/Tunisia/2013/Murinae/Muridae	KJ408421	Tunisia	2013	Rattus rattus
RodPV/R. rattus/KJ408425/Tunisia/2013/Murinae/Muridae	KJ408425	Tunisia	2013	Rattus rattus
Tailam virus/R. andamanensis/JN689227/China/2009	JN689227	China	2009	Rattus andamanensis
RodPV/R. rattus/KJ408426/Tunisia/2013/Murinae/Muridae	KJ408426	Tunisia	2013	Rattus rattus
Beilong virus/Rod. sp./DQ100461/China/2005	DQ100461	China	2005	Rodent sp.
RodPV/R. rattus/KJ408427/Tunisia/2013/Murinae/Muridae	KJ408427	Tunisia	2013	Rattus rattus
J virus/M. musculus/AY900001/Australia/1970	AY900001	Australia	1970	Mus musculus
RodPV/C. gundi/KJ408410/Tunisia/2010/nd/Ctenodactylidae	KJ408410	Tunisia	2010	Ctenodactylus gundi
RodPV/C. gundi/KJ408409/Tunisia/2010/nd/Ctenodactylidae	KJ408409	Tunisia	2010	Ctenodactylus gundi
RodPV/C. gundi/KJ408403/Tunisia/2011/nd/Ctenodactylidae	KJ408403	Tunisia	2011	Ctenodactylus gundi
RodPV/C. gundi/KJ408404/Tunisia/2011/nd/Ctenodactylidae	KJ408404	Tunisia	2011	Ctenodactylus gundi
RodPV/C. gundi/KJ408405/Tunisia/2011/nd/Ctenodactylidae	KJ408405	Tunisia	2011	Ctenodactylus gundi
RodPV/C. gundi/KJ408406/Tunisia/2011/nd/Ctenodactylidae	KJ408406	Tunisia	2011	Ctenodactylus gundi
RodPV/C. gundi/KJ408407/Tunisia/2011/nd/Ctenodactylidae	KJ408407	Tunisia	2011	Ctenodactylus gundi
RodPV/C. gundi/KJ408408/Tunisia/2011/nd/Ctenodactylidae	KJ408408	Tunisia	2011	Ctenodactylus gundi
RodPV/C. gundi/KJ408413/Tunisia/2010/nd/Ctenodactylidae	KJ408413	Tunisia	2010	Ctenodactylus gundi
RodPV/C. gundi/KJ408416/Tunisia/2010/nd/Ctenodactylidae	KJ408416	Tunisia	2010	Ctenodactylus gundi
RodPV/C. gundi/KJ408414/Tunisia/2010/nd/Ctenodactylidae	KJ408414	Tunisia	2010	Ctenodactylus gundi
RodPV/C. gundi/KJ408411/Tunisia/2010/nd/Ctenodactylidae	KJ408411	Tunisia	2010	Ctenodactylus gundi
RodPV/M. shawi/KJ408402/Tunisia/2010/Gerbillinae/Muridae	KJ408402	Tunisia	2010	Meriones shawi
RodPV/C. gundi/KJ408412/Tunisia/2010/nd/Ctenodactylidae	KJ408412	Tunisia	2010	Ctenodactylus gundi
Mossman virus/Wild_rat/NC005339/Australia/2003	NC005339	Australia	2003	Wild rat
Tupaia virus/Shrew/AF079780/Thailand/1998	AF079780	Thailand	1998	Shrew

FMV/Cat/JQ411015/HongKong/2009	JQ411015	Hong Kong	2009	Felis catus
FMV/Cat/JQ411015/HongKong/2009	JQ411015	Hong Kong	2009	Felis catus
MePV/Human/AB016162/Japan/1998	AB016162	Japan	1998	Human
PPRV/Goat/X74443/Nigeria/1993	X74443	Nigeria	1993	Goat
CDV/Dog/NC001921South Africa/1987	NC001921	South Africa	1987	Dog
PDV/Seal/Y09630/Ireland/1997	Y09630	Ireland	1997	Seal
RodPV/P. obesus/KJ408384/Tunisia/2009/Gerbillinae/Muridae	KJ408384	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408385/Tunisia/2009/Gerbillinae/Muridae	KJ408385	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408386/Tunisia/2009/Gerbillinae/Muridae	KJ408386	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408387/Tunisia/2009/Gerbillinae/Muridae	KJ408387	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408388/Tunisia/2009/Gerbillinae/Muridae	KJ408388	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408389/Tunisia/2009/Gerbillinae/Muridae	KJ408389	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408390/Tunisia/2009/Gerbillinae/Muridae	KJ408390	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408391/Tunisia/2009/Gerbillinae/Muridae	KJ408391	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408392/Tunisia/2009/Gerbillinae/Muridae	KJ408392	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408393/Tunisia/2009/Gerbillinae/Muridae	KJ408393	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408394/Tunisia/2009/Gerbillinae/Muridae	KJ408394	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408396/Tunisia/2009/Gerbillinae/Muridae	KJ408396	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408395/Tunisia/2009/Gerbillinae/Muridae	KJ408395	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408397/Tunisia/2009/Gerbillinae/Muridae	KJ408397	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408399/Tunisia/2009/Gerbillinae/Muridae	KJ408399	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408401/Tunisia/2009/Gerbillinae/Muridae	KJ408401	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408400/Tunisia/2009/Gerbillinae/Muridae	KJ408400	Tunisia	2009	Psammomys obesus
RodPV/P. obesus/KJ408398/Tunisia/2009/Gerbillinae/Muridae	KJ408398	Tunisia	2009	Psammomys obesus
Nipah virus/Bat/NC002728/Bangladesh/2005	NC002728	Bangladesh	2005	Bat
Hendra virus/Horse/AF017149/Australia/1994	AF017149	Australia	1994	Horse
Cedar virus/Bat/JQ001776/Australia/2009	JQ001776	Australia	2009	Bat
Salem virus/Horse/JQ697837/USA/1992	JQ697837	USA	1992	Horse
HPIV-1/Human/NC003461/USA/1964	NC003461	USA	1964	Human
HPIV-3/Human/FJ455842/China/2003	FJ455842	China	2003	Human
Mapuera virus/Bat/EF095490/Brazil/1979	EF095490	Brazil	1979	Bat
Porcine rubulavirus/Swine/NC009640/Mexico/1980	NC009640	Mexico	1980	Swine
MuPV/Human/AB600843/Mongolia/2009	AB600843	Mongolia	2009	Human
Menangle virus/Swine/AF326114/Australia/1997	AF326114	Australia	1997	Swine
Tioman virus/Bat/AF298895/Malaysia/2000	AF298895	Malaysia	2000	Bat
NDV/Duck/JN653339/China/2007	JN653339	China	2007	Duck
PAPMV/Pigeon/GQ429292/Ireland/1996	GQ429292	Ireland	1996	Pigeon

h-RSV/Human/AF013254/USA/1985	AF013254	USA	1985	Human
b-RSV/Bovine/AF092942/GER/1998	AF092942	Germany	1998	Bovine
AMPV/Turkey/FJ977568/USA/2009	FJ977568	USA	2009	Turkey
HMPV/Human/GQ153651/China/2008	GQ153651	China	2008	Human
MPV/M. musculus/AY743910/USA/1944	AY743910	USA	1944	Mus musculus
Rabies Virus/Bat/JQ595353/USA/2005*	JQ595353	USA	2005	Bat

109 a: Virus name or virus affiliation/typical host/ accession number/origin/collection year/host Subfamily/host Family

b: AMPV = Avian Metapneumovirus; b-RSV = Bovine Respiratory Syncitical Virus; C. gundi = Ctenodactylus gundi; CDV = Canine Distemper Virus; FMV = Feline Morbillivirus; HMPV = Human Metapneumovirus; HPIV = Human Parainfluenza Virus; h-RSV = Human Respiratory Syncitical Virus; M. musculus = Mus musculus; M. shawi = Meriones shawi; MePV = Measles Virus; MPV = Murine Pneumonia Virus; MuPV = Mumps Virus; nd = not defined; NDV = Newcastle Disease Virus; P. obesus = Psammomys obesus; PAPMV = Pigeon Paramyxovirus; PDV= Phocine Distemper Virus; PPRV = Peste des Petits Ruminents Virus; R. rattus= Rattus rattus; Rod. = Rodents; RodPV = Rodents Paramyxovirus; sp. = species.

\*: Sequence of rabies virus used as outgroup.

8 /16

- Supplementary Table S2: Viruses used for the phylogenetic and phylogeographic analysis of
- 120 Rodentia associated UMRV in Figures 2 and 3.

Beilong virus/Rodent_origin/DQ100461/China/1998/nd/Muridae  J virus/Mus_musculus/NC007454/Australia/1972/Murinae/Muridae  Mossman virus/Rattus_leucopus/AY286409/Australia/1970/Murinae/Muridae  Nariva virus/Zygodontomys_brevicauda/FJ362497/TrinidadTobago/1962/ Sigmodontinae/Cricetidae  RodPV/Apodemus_flavicollis/HQ660183/Germany/2010/Murinae/Muridae  RodPV/Apodemus_flavicollis/JF828298/Germany/2010/Murinae/Muridae  RodPV/Apodemus_flavicollis/JF828308/Germany/2010/Murinae/Muridae  RodPV/Ctenodactylus_gundi/KJ408403/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408404/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408405/Tunisia/2011/nd/Ctenodactylidae	numbers DQ100461 NC007454 AY286409 FJ362497 HQ660183 JF828298 JF828308 KJ408403 KJ408404 KJ408405 KJ408406 KJ408407	Country  China  Australia  Australia  Trinidad  Tobago  Germany  Germany  Tunisia  Tunisia  Tunisia  Tunisia  Tunisia  Tunisia	year 1998 1972 1970 1962 2010 2010 2010 2011 2011 2011	Rodent origin  Mus musculus  Rattus_leucopus  Zygodontomys brevicauda  Apodemus_flavicollis  Apodemus_flavicollis  Ctenodactylus gundi  Ctenodactylus gundi  Ctenodactylus gundi  Ctenodactylus gundi
J virus/Mus_musculus/NC007454/Australia/1972/Murinae/Muridae  Mossman virus/Rattus_leucopus/AY286409/Australia/1970/Murinae/Muridae  Nariva virus/Zygodontomys_brevicauda/FJ362497/TrinidadTobago/1962/  Sigmodontinae/Cricetidae  RodPV/Apodemus_flavicollis/HQ660183/Germany/2010/Murinae/Muridae  RodPV/Apodemus_flavicollis/JF828298/Germany/2010/Murinae/Muridae  RodPV/Apodemus_flavicollis/JF828308/Germany/2010/Murinae/Muridae  RodPV/Ctenodactylus_gundi/KJ408403/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408404/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408405/Tunisia/2011/nd/Ctenodactylidae	NC007454 AY286409 FJ362497 HQ660183 JF828298 JF828308 KJ408403 KJ408404 KJ408405 KJ408406	Australia Australia Trinidad Tobago Germany Germany Tunisia Tunisia Tunisia Tunisia	1972 1970 1962 2010 2010 2010 2011 2011 2011 2011	Mus musculus  Rattus_leucopus  Zygodontomys brevicauda  Apodemus_flavicollis  Apodemus_flavicollis  Ctenodactylus gundi  Ctenodactylus gundi  Ctenodactylus gundi
Mossman virus/Rattus_leucopus/AY286409/Australia/1970/Murinae/Muridae Nariva virus/Zygodontomys_brevicauda/FJ362497/TrinidadTobago/1962/ Sigmodontinae/Cricetidae RodPV/Apodemus_flavicollis/HQ660183/Germany/2010/Murinae/Muridae RodPV/Apodemus_flavicollis/JF828298/Germany/2010/Murinae/Muridae RodPV/Apodemus_flavicollis/JF828308/Germany/2010/Murinae/Muridae RodPV/Ctenodactylus_gundi/KJ408403/Tunisia/2011/nd/Ctenodactylidae RodPV/Ctenodactylus_gundi/KJ408404/Tunisia/2011/nd/Ctenodactylidae RodPV/Ctenodactylus_gundi/KJ408405/Tunisia/2011/nd/Ctenodactylidae	AY286409  FJ362497  HQ660183  JF828298  JF828308  KJ408403  KJ408404  KJ408405  KJ408406	Australia Trinidad Tobago Germany Germany Tunisia Tunisia Tunisia Tunisia	1970 1962 2010 2010 2010 2011 2011 2011 2011	Rattus_leucopus  Zygodontomys brevicauda  Apodemus_flavicollis Apodemus_flavicollis Ctenodactylus gundi Ctenodactylus gundi
Nariva virus/Zygodontomys_brevicauda/FJ362497/TrinidadTobago/1962/ Sigmodontinae/Cricetidae  RodPV/Apodemus_flavicollis/HQ660183/Germany/2010/Murinae/Muridae  RodPV/Apodemus_flavicollis/JF828298/Germany/2010/Murinae/Muridae  RodPV/Apodemus_flavicollis/JF828308/Germany/2010/Murinae/Muridae  RodPV/Ctenodactylus_gundi/KJ408403/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408404/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408405/Tunisia/2011/nd/Ctenodactylidae	FJ362497  HQ660183  JF828298  JF828308  KJ408403  KJ408404  KJ408405  KJ408406	Trinidad Tobago Germany Germany Tunisia Tunisia Tunisia Tunisia	2010 2010 2010 2010 2011 2011 2011 2011	Zygodontomys brevicauda Apodemus_flavicollis Apodemus_flavicollis Ctenodactylus gundi Ctenodactylus gundi Ctenodactylus gundi
Sigmodontinae/Cricetidae  RodPV/Apodemus_flavicollis/HQ660183/Germany/2010/Murinae/Muridae  RodPV/Apodemus_flavicollis/JF828298/Germany/2010/Murinae/Muridae  RodPV/Apodemus_flavicollis/JF828308/Germany/2010/Murinae/Muridae  RodPV/Ctenodactylus_gundi/KJ408403/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408404/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408405/Tunisia/2011/nd/Ctenodactylidae	HQ660183  JF828298  JF828308  KJ408403  KJ408404  KJ408405  KJ408406	Tobago Germany Germany Tunisia Tunisia Tunisia Tunisia	2010 2010 2010 2011 2011 2011 2011	brevicauda  Apodemus_flavicollis  Apodemus_flavicollis  Apodemus_flavicollis  Ctenodactylus gundi  Ctenodactylus gundi  Ctenodactylus gundi
Sigmodontinae/Cricetidae  RodPV/Apodemus_flavicollis/HQ660183/Germany/2010/Murinae/Muridae  RodPV/Apodemus_flavicollis/JF828298/Germany/2010/Murinae/Muridae  RodPV/Apodemus_flavicollis/JF828308/Germany/2010/Murinae/Muridae  RodPV/Ctenodactylus_gundi/KJ408403/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408404/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408405/Tunisia/2011/nd/Ctenodactylidae	HQ660183  JF828298  JF828308  KJ408403  KJ408404  KJ408405  KJ408406	Germany Germany Tunisia Tunisia Tunisia Tunisia	2010 2010 2010 2011 2011 2011 2011	Apodemus_flavicollis Apodemus_flavicollis Apodemus_flavicollis Ctenodactylus gundi Ctenodactylus gundi Ctenodactylus gundi
RodPV/Apodemus_flavicollis/JF828298/Germany/2010/Murinae/Muridae  RodPV/Apodemus_flavicollis/JF828308/Germany/2010/Murinae/Muridae  RodPV/Ctenodactylus_gundi/KJ408403/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408404/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408405/Tunisia/2011/nd/Ctenodactylidae	JF828298 JF828308 KJ408403 KJ408404 KJ408405 KJ408406	Germany Germany Tunisia Tunisia Tunisia Tunisia	2010 2010 2011 2011 2011 2011	Apodemus_flavicollis Apodemus_flavicollis Ctenodactylus gundi Ctenodactylus gundi Ctenodactylus gundi
RodPV/Apodemus_flavicollis/JF828308/Germany/2010/Murinae/Muridae  RodPV/Ctenodactylus_gundi/KJ408403/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408404/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408405/Tunisia/2011/nd/Ctenodactylidae	JF828308 KJ408403 KJ408404 KJ408405 KJ408406	Germany Tunisia Tunisia Tunisia Tunisia	2010 2011 2011 2011 2011	Apodemus_flavicollis Ctenodactylus gundi Ctenodactylus gundi Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408403/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408404/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408405/Tunisia/2011/nd/Ctenodactylidae	KJ408403 KJ408404 KJ408405 KJ408406	Tunisia Tunisia Tunisia Tunisia	2011 2011 2011 2011	Ctenodactylus gundi Ctenodactylus gundi Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408404/Tunisia/2011/nd/Ctenodactylidae  RodPV/Ctenodactylus_gundi/KJ408405/Tunisia/2011/nd/Ctenodactylidae	KJ408404 KJ408405 KJ408406	Tunisia Tunisia Tunisia	2011 2011 2011	Ctenodactylus gundi Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408405/Tunisia/2011/nd/Ctenodactylidae	KJ408405 KJ408406	Tunisia Tunisia	2011	Ctenodactylus gundi
,,	KJ408406	Tunisia	2011	
RodPV/Ctenodactylus_gundi/K I408406/Tunisia/2011/nd/Ctenodactylidae				Ctenodactylus gundi
Total 1, Cicliodactylus_guildi153700700/14iiisia/2011/iid/Cicliodactylidac	KJ408407	Tunisia	2011	
RodPV/Ctenodactylus_gundi/KJ408407/Tunisia/2011/nd/Ctenodactylidae			2011	Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408408/Tunisia/2011/nd/Ctenodactylidae	KJ408408	Tunisia	2011	Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408409/Tunisia/2010/nd/Ctenodactylidae	KJ408409	Tunisia	2010	Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408410/Tunisia/2010/nd/Ctenodactylidae	KJ408410	Tunisia	2010	Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408411/Tunisia/2010/nd/Ctenodactylidae	KJ408411	Tunisia	2010	Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408412/Tunisia/2010/nd/Ctenodactylidae	KJ408412	Tunisia	2010	Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408413/Tunisia/2010/nd/Ctenodactylidae	KJ408413	Tunisia	2010	Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408414/Tunisia/2010/nd/Ctenodactylidae	KJ408414	Tunisia	2010	Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408415/Tunisia/2010/nd/Ctenodactylidae	KJ408415	Tunisia	2010	Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408416/Tunisia/2010/nd/Ctenodactylidae	KJ408416	Tunisia	2010	Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408417/Tunisia/2010/nd/Ctenodactylidae	KJ408417	Tunisia	2010	Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408418/Tunisia/2010/nd/Ctenodactylidae	KJ408418	Tunisia	2010	Ctenodactylus gundi
RodPV/Ctenodactylus_gundi/KJ408419/Tunisia/2010/nd/Ctenodactylidae	KJ408419	Tunisia	2010	Ctenodactylus gundi
RodPV/Eliurus_minor/KF245989/Madagascar/2011/Nesomyinae/Nesomyidae	KF245989	Madagascar	2011	Eliurus minor
RodPV/Eliurus_minor/KF245998/Madagascar/2011/Nesomyinae/Nesomyidae	KF245998	Madagascar	2011	Eliurus minor
RodPV/Eliurus_minor/KF245999/Madagascar/2011/Nesomyinae/Nesomyidae	KF245999	Madagascar	2011	Eliurus minor
RodPV/Eliurus_minor/KF246000/Madagascar/2011/Nesomyinae/Nesomyidae	KF246000	Madagascar	2011	Eliurus minor
RodPV/Eliurus_minor/KF246001/Madagascar/2011/Nesomyinae/Nesomyidae	KF246001	Madagascar	2011	Eliurus minor
RodPV/Eliurus_minor/KF246011/Madagascar/2011/Nesomyinae/Nesomyidae	KF246011	Madagascar	2011	Eliurus minor

RodPV/Eliurus_minor/KF246012/Madagascar/2011/Nesomyinae/Nesomyidae	KF246012	Madagascar	2011	Eliurus minor
RodPV/Eliurus_minor/KF246013/Madagascar/2011/Nesomyinae/Nesomyidae	KF246013	Madagascar	2011	Eliurus minor
RodPV/Eliurus_minor/KF246022/Madagascar/2011/Nesomyinae/Nesomyidae	KF246022	Madagascar	2011	Eliurus minor
RodPV/Eliurus_minor/KF246032/Madagascar/2012/Nesomyinae/Nesomyidae	KF246032	Madagascar	2012	Eliurus minor
RodPV/Eliurus_minor/KF246033/Madagascar/2012/Nesomyinae/Nesomyidae	KF246033	Madagascar	2012	Eliurus minor
$RodPV/Eliurus\_tanala/KF246028/Madagascar/2012/Nesomyinae/Nesomyidae$	KF246028	Madagascar	2012	Eliurus tanala
RodPV/Mastomys_natalensis/AB844358/Zambia/2010/Murinae/Muridae	AB844358	Zambia	2010	Mastomys natalensis
$RodPV/Mastomys\_natalensis/AB844367/Zambia/2011/Murinae/Muridae$	AB844367	Zambia	2011	Mastomys natalensis
RodPV/Mastomys_natalensis/AB844369/Zambia/2011/Murinae/Muridae	AB844369	Zambia	2011	Mastomys natalensis
RodPV/Mastomys_natalensis/AB844376/Zambia/2011/Murinae/Muridae	AB844376	Zambia	2011	Mastomys natalensis
RodPV/Mastomys_natalensis/AB844378/Zambia/2011/Murinae/Muridae	AB844378	Zambia	2011	Mastomys natalensis
RodPV/Mastomys_natalensis/AB844384/Zambia/2011/Murinae/Muridae	AB844384	Zambia	2011	Mastomys natalensis
RodPV/Mastomys_natalensis/AB844388/Zambia/2011/Murinae/Muridae	AB844388	Zambia	2011	Mastomys natalensis
RodPV/Mastomys_natalensis/AB844391/Zambia/2011/Murinae/Muridae	AB844391	Zambia	2011	Mastomys natalensis
RodPV/Mastomys_natalensis/AB844396/Zambia/2011/Murinae/Muridae	AB844396	Zambia	2011	Mastomys natalensis
RodPV/Mastomys_natalensis/AB844398/Zambia/2011/Murinae/Muridae	AB844398	Zambia	2011	Mastomys natalensis
RodPV/Mastomys_natalensis/AB844401/Zambia/2011/Murinae/Muridae	AB844401	Zambia	2011	Mastomys natalensis
RodPV/Mastomys_natalensis/AB844404/Zambia/2011/Murinae/Muridae	AB844404	Zambia	2011	Mastomys natalensis
$RodPV/Mastomys\_natalensis/AB844406/Zambia/2011/Murinae/Muridae$	AB844406	Zambia	2011	Mastomys natalensis
RodPV/Mastomys_natalensis/AB844413/Zambia/2011/Murinae/Muridae	AB844413	Zambia	2011	Mastomys natalensis
RodPV/Mastomys_natalensis/AB844422/Zambia/2012/Murinae/Muridae	AB844422	Zambia	2012	Mastomys natalensis
RodPV/Meriones_shawi/KJ408402/Tunisia/2010/Gerbillinae/Muridae	KJ408402	Tunisia	2010	Meriones_shawi
RodPV/Microtus_arvalis/HQ660185/Germany/2010/Arvicolinae/Cricetidae	HQ660185	Germany	2010	Microtus arvalis
RodPV/Mus_minutoides/AB844424/Zambia/2010/Murinae/Muridae	AB844424	Zambia	2010	Mus minutoides
RodPV/Mus_minutoides/AB844425/Zambia/2011/Murinae/Muridae	AB844425	Zambia	2011	Mus minutoides
RodPV/Myodes_glareolus/HQ660181/Germany/2010/Arvicolinae/Cricetidae	HQ660181	Germany	2010	Myodes glareolus
RodPV/Myodes_glareolus/HQ660182/Germany/2010/Arvicolinae/Cricetidae	HQ660182	Germany	2010	Myodes glareolus
RodPV/Myodes_glareolus/HQ660186/Germany/2010/Arvicolinae/Cricetidae	HQ660186	Germany	2010	Myodes glareolus
RodPV/Myodes_glareolus/JF828299/Germany/2010/Arvicolinae/Cricetidae	JF828299	Germany	2010	Myodes glareolus
RodPV/Myodes_glareolus/JF828300/Germany/2010/Arvicolinae/Cricetidae	JF828300	Germany	2010	Myodes glareolus
RodPV/Myodes_glareolus/JF828301/Germany/2010/Arvicolinae/Cricetidae	JF828301	Germany	2010	Myodes glareolus
RodPV/Myodes_glareolus/JF828304/Germany/2010/Arvicolinae/Cricetidae	JF828304	Germany	2010	Myodes glareolus
RodPV/Myodes_glareolus/JF828305/Germany/2010/Arvicolinae/Cricetidae	JF828305	Germany	2010	Myodes glareolus
RodPV/Myodes_glareolus/JF828306/Germany/2010/Arvicolinae/Cricetidae	JF828306	Germany	2010	Myodes glareolus
RodPV/Myodes_glareolus/JF828307/Germany/2010/Arvicolinae/Cricetidae	JF828307	Germany	2010	Myodes glareolus
RodPV/Myodes_glareolus/JF828309/Germany/2010/Arvicolinae/Cricetidae	JF828309	Germany	2010	Myodes glareolus
RodPV/Psammomys_obesus/KJ408384/Tunisia/2009/Gerbillinae/Muridae	KJ408384	Tunisia	2009	Psammomys obesus

RodPV/Psammomys_obesus/KJ408385/Tunisia/2009/Gerbillinae/Muridae	KJ408385	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408386/Tunisia/2009/Gerbillinae/Muridae	KJ408386	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408387/Tunisia/2009/Gerbillinae/Muridae	KJ408387	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408388/Tunisia/2009/Gerbillinae/Muridae	KJ408388	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408389/Tunisia/2009/Gerbillinae/Muridae	KJ408389	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408390/Tunisia/2009/Gerbillinae/Muridae	KJ408390	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408391/Tunisia/2009/Gerbillinae/Muridae	KJ408391	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408392/Tunisia/2009/Gerbillinae/Muridae	KJ408392	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408393/Tunisia/2009/Gerbillinae/Muridae	KJ408393	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408394/Tunisia/2009/Gerbillinae/Muridae	KJ408394	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408395/Tunisia/2009/Gerbillinae/Muridae	KJ408395	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408396/Tunisia/2009/Gerbillinae/Muridae	KJ408396	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408397/Tunisia/2009/Gerbillinae/Muridae	KJ408397	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408398/Tunisia/2009/Gerbillinae/Muridae	KJ408398	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408399/Tunisia/2009/Gerbillinae/Muridae	KJ408399	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408400/Tunisia/2009/Gerbillinae/Muridae	KJ408400	Tunisia	2009	Psammomys obesus
RodPV/Psammomys_obesus/KJ408401/Tunisia/2009/Gerbillinae/Muridae	KJ408401	Tunisia	2009	Psammomys obesus
RodPV/Rattus_norvegicus/KF245939/Reunion/2012/Murinae/Muridae	KF245939	Reunion	2012	Rattus norvegicus
RodPV/Rattus_norvegicus/KF245940/Reunion/2012/Murinae/Muridae	KF245940	Reunion	2012	Rattus norvegicus
RodPV/Rattus_norvegicus/KF245942/Reunion/2012/Murinae/Muridae	KF245942	Reunion	2012	Rattus norvegicus
RodPV/Rattus_norvegicus/KF245948/Reunion/2012/Murinae/Muridae	KF245948	Reunion	2012	Rattus norvegicus
RodPV/Rattus_norvegicus/KF245955/Seychelles/2011/Murinae/Muridae	KF245955	Seychelles	2011	Rattus norvegicus
RodPV/Rattus_norvegicus/KF408257/Reunion/2012/Murinae/Muridae	KF408257	Reunion	2012	Rattus norvegicus
RodPV/Rattus_norvegicus/KF408258/Reunion/2012/Murinae/Muridae	KF408258	Reunion	2012	Rattus norvegicus
RodPV/Rattus_norvegicus/KF408259/Reunion/2012/Murinae/Muridae	KF408259	Reunion	2012	Rattus norvegicus
RodPV/Rattus_rattus/HQ660195/Thailand/2007/Murinae/Muridae*	HQ660195	Thailand	2007	Rattus rattus
RodPV/Rattus_rattus/KF245950/Reunion/2012/Murinae/Muridae	KF245950	Reunion	2012	Rattus rattus
RodPV/Rattus_rattus/KF245951/Reunion/2012/Murinae/Muridae	KF245951	Reunion	2012	Rattus rattus
RodPV/Rattus_rattus/KF245953/Seychelles/2011/Murinae/Muridae	KF245953	Seychelles	2011	Rattus rattus
RodPV/Rattus_rattus/KF245954/Seychelles/2011/Murinae/Muridae	KF245954	Seychelles	2011	Rattus rattus
RodPV/Rattus_rattus/KF245956/Seychelles/2011/Murinae/Muridae	KF245956	Seychelles	2011	Rattus rattus
RodPV/Rattus_rattus/KF245958/Mayotte/2012/Murinae/Muridae	KF245958	Mayotte	2012	Rattus rattus
RodPV/Rattus_rattus/KF245960/Mayotte/2012/Murinae/Muridae	KF245960	Mayotte	2012	Rattus rattus
RodPV/Rattus_rattus/KF245963/Madagascar/2010/Murinae/Muridae	KF245963	Madagascar	2010	Rattus rattus
RodPV/Rattus_rattus/KF245968/Madagascar/2010/Murinae/Muridae	KF245968	Madagascar	2010	Rattus rattus
RodPV/Rattus_rattus/KF245969/Madagascar/2010/Murinae/Muridae	KF245969	Madagascar	2010	Rattus rattus
RodPV/Rattus_rattus/KF245976/Madagascar/2011/Murinae/Muridae	KF245976	Madagascar	2011	Rattus rattus
		<u> </u>		

RodPV/Rattus_rattus/KF245990/Madagascar/2011/Murinae/Muridae	KF245990	Madagascar	2011	Rattus rattus
RodPV/Rattus_rattus/KF246002/Madagascar/2011/Murinae/Muridae	KF246002	Madagascar	2011	Rattus rattus
$RodPV/Rattus\_rattus/KF246003/Madagascar/2011/Murinae/Muridae$	KF246003	Madagascar	2011	Rattus rattus
RodPV/Rattus_rattus/KF246004/Madagascar/2011/Murinae/Muridae	KF246004	Madagascar	2011	Rattus rattus
RodPV/Rattus_rattus/KF246005/Madagascar/2011/Murinae/Muridae	KF246005	Madagascar	2011	Rattus rattus
RodPV/Rattus_rattus/KF246014/Madagascar/2011/Murinae/Muridae	KF246014	Madagascar	2011	Rattus rattus
RodPV/Rattus_rattus/KF246023/Madagascar/2011/Murinae/Muridae	KF246023	Madagascar	2011	Rattus rattus
RodPV/Rattus_rattus/KF246025/Madagascar/2012/Murinae/Muridae	KF246025	Madagascar	2012	Rattus rattus
RodPV/Rattus_rattus/KF246026/Madagascar/2012/Murinae/Muridae	KF246026	Madagascar	2012	Rattus rattus
RodPV/Rattus_rattus/KF246027/Madagascar/2012/Murinae/Muridae	KF246027	Madagascar	2012	Rattus rattus
RodPV/Rattus_rattus/KF246029/Madagascar/2012/Murinae/Muridae	KF246029	Madagascar	2012	Rattus rattus
RodPV/Rattus_rattus/KF246034/Madagascar/2012/Murinae/Muridae	KF246034	Madagascar	2012	Rattus rattus
RodPV/Rattus_rattus/KF408256/Reunion/2012/Murinae/Muridae	KF408256	Reunion	2012	Rattus rattus
RodPV/Rattus_rattus/KF408260/Reunion/2012/Murinae/Muridae	KF408260	Reunion	2012	Rattus rattus
RodPV/Rattus_rattus/KJ408420/Tunisia/2013/Murinae/Muridae	KJ408420	Tunisia	2013	Rattus rattus
RodPV/Rattus_rattus/KJ408421/Tunisia/2013/Murinae/Muridae	KJ408421	Tunisia	2013	Rattus rattus
RodPV/Rattus_rattus/KJ408422/Tunisia/2013/Murinae/Muridae	KJ408422	Tunisia	2013	Rattus rattus
RodPV/Rattus_rattus/KJ408423/Tunisia/2013/Murinae/Muridae	KJ408423	Tunisia	2013	Rattus rattus
RodPV/Rattus_rattus/KJ408424/Tunisia/2013/Murinae/Muridae	KJ408424	Tunisia	2013	Rattus rattus
RodPV/Rattus_rattus/KJ408425/Tunisia/2013/Murinae/Muridae	KJ408425	Tunisia	2013	Rattus rattus
RodPV/Rattus_rattus/AB844426/Zambia/2010/Murinae/Muridae*	AB844426	Zambia	2010	Rattus rattus
RodPV/Rattus_rattus/AB844427/Zambia/2010/Murinae/Muridae	AB844427	Zambia	2010	Rattus rattus
RodPV/Rattus_Sp./KF245941/Reunion/2012/Murinae/Muridae	KF245941	Reunion	2012	Rattus sp.
RodPV/Rattus_Sp./KF245944/Reunion/2012/Murinae/Muridae	KF245944	Reunion	2012	Rattus sp.
RodPV/Rattus_Sp./KF245945/Reunion/2012/Murinae/Muridae	KF245945	Reunion	2012	Rattus sp.
RodPV/Rattus_Sp./KF245946/Reunion/2012/Murinae/Muridae	KF245946	Reunion	2012	Rattus sp.
RodPV/Rattus_Sp./KF245947/Reunion/2012/Murinae/Muridae	KF245947	Reunion	2012	Rattus sp.
RodPV/Rattus_SP./KF245949/Reunion/2012/Murinae/Muridae	KF245949	Reunion	2012	Rattus sp.
RodPV/Rattus_Sp./KF245952/Reunion/2012/Murinae/Muridae	KF245952	Reunion	2012	Rattus sp.
RodPV/Rattus_Sp./KF245957/Mayotte/2012/Murinae/Muridae	KF245957	Mayotte	2012	Rattus sp.
RodPV/Rattus_Sp./KF245959/Mayotte/2012/Murinae/Muridae	KF245959	Mayotte	2012	Rattus sp.
RodPV/Rattus_Sp./KF245959/Mayotte/2012/Murinae/Muridae  RodPV/Rattus_Sp./KF245961/Mayotte/2012/Murinae/Muridae	KF245959 KF245961	Mayotte Mayotte	2012 2012	Rattus sp.  Rattus sp.
= .		3.		
RodPV/Rattus_Sp./KF245961/Mayotte/2012/Murinae/Muridae	KF245961 HQ660172	Mayotte	2012	Rattus sp.
RodPV/Rhabdomys_pumilio/HQ660172/South_Africa/2009/Murinae/Muridae  RodPV/Rhabdomys_pumilio/HQ660173/South_Africa/2009/Murinae/Muridae	KF245961	Mayotte  South Africa  South Africa	2012	Rattus sp.  Rhabdomys pumilio  Rhabdomys pumilio
RodPV/Rattus_Sp./KF245961/Mayotte/2012/Murinae/Muridae  RodPV/Rhabdomys_pumilio/HQ660172/South_Africa/2009/Murinae/Muridae	KF245961 HQ660172 HQ660173	Mayotte South Africa	2012 2009 2009	Rattus sp.  Rhabdomys pumilio

RodPV/Rhabdomys_pumilio/HQ660177/South_Africa/2009/Murinae/Muridae	HQ660177	South Africa	2009	Rhabdomys pumilio
RodPV/Rhabdomys_pumilio/HQ660178/South_Africa/2009/Murinae/Muridae	HQ660178	South Africa	2009	Rhabdomys pumilio
RodPV/Rhabdomys_pumilio/HQ660179/South_Africa/2009/Murinae/Muridae	HQ660179	South Africa	2009	Rhabdomys pumilio
RodPV/Rhabdomys_pumilio/HQ660180/South_Africa/2009/Murinae/Muridae	HQ660180	South Africa	2009	Rhabdomys pumilio
RodPV/Microtus_agrestis/HQ660184/Germany/2010/Arvicolinae/Cricetidae	HQ660184	Germany	2010	Microtus agrestis
Tailam virus/Rattus_andamanensis/JN689227/China/2009/Murinae/Muridae	JN689227	China	2009	Rattus andamanensis

<sup>121</sup> a: Virus name or virus affiliation/typical host/ accession number/origin/collection year/host Subfamily/host Family

123 \*: Two Respirovirus used as an outgroup.

b: RodPV = Rodents Paramyxovirus; sp. = species.

#### Supplementary Table S3: Viruses used for the estimate of the evolutionary rate change for

#### the partial L-gene among the Measles virus.

v. mah	Accession	G .	Collection	www.h
Virus ID <sup>a,b</sup>	numbers	Country	year	Host b
Measles_virus/Human/FJ161211/China/2006	FJ16121	China	2006	Human
Measles_virus/ND/HM439386/Sudan/1997	HM439386	Sudan	1997	ND
Measles_virus/Human/JF727649/Russia/2010	JF727649	Russia	2010	Human
Measles_virus/Human/JF727650/Russia/2010	JF727650	Russia	2010	Human
Measles_virus/Human/GQ376026/Japan/1999	GQ376026	Japan	1999	Human
Measles_virus/Human/GQ376027/Japan/1999	GQ376027	Japan	1999	Human
Measles_virus/ND/AB481088/Japan/1989	AB481088	Japan	1989	ND
Measles_virus/ND/AB481087/Japan/1989	AB481087	Japan	1989	ND
Measles_virus/Human/KC164757/Italy/2010	KC164757	Italy	2010	Human
Measles_virus/ND/KJ018971/Canada/2010	KJ018971	Canada	2010	ND
Measles_virus/Human/KJ410048/Germany/2013	KJ410048	Germany	2013	Human
Measles_virus/Human/KC164758/Italy/2011	KC164758	Italy	2011	Human
Measles_virus/ND/KJ018970/Canada/2010	KJ018970	Canada	2010	ND
Measles_virus/ND/KM054581/France/2014	KM054581	France	2014	ND
Measles_virus/Human/JN635406/USA/2008	JN635406	USA	2008	Human
Measles_virus/Human/JN635410/USA/2003	JN635410	USA	2003	Human
Measles_virus/Human/JN635403/USA/2009	JN635403	USA	2009	Human
Measles_virus/Human/JN635408/USA/2005	JN635408	USA	2005	Human
Measles_virus/Human/JN635402/USA/2009	JN635402	USA	2009	Human
Measles_virus/Human/JN635411/USA/2009	JN635411	USA	2009	Human
Measles_virus/Human/JN635407/USA/2007	JN635407	USA	2007	Human
Measles_virus/Human/JN635404/USA/2009	JN635404	USA	2009	Human
Measles_virus/Human/JN635405/USA/2008	JN635405	USA	2008	Human
Measles_virus/Human/JN635409/USA/2004	JN635409	USA	2004	Human

a: Virus name/ typical host/ accession number/origin/collection year

b: ND = not defined.

**Supplementary Table S4:** Significant diffusion rates between countries at Bayesian phylogeographic analysis. Posterior probability of the rates k (pk) and Bayes factor (BF) of the significant migration rates as revealed by the Skygrid symmetric model.

Rate (from/to)	pk	BF
Zambia / TrinidadTobago	0.98	207.27
Australia / Madagascar	0.97	193.02
Zambia / Madagascar	0.97	191.93
Tunisia / Madagascar	0.97	177.40
South_Africa / TrinidadTobago	0.97	160.21
South_Africa / Madagascar	0.97	146.70
TrinidadTobago / Tunisia	0.97	144.16
TrinidadTobago / Mayotte	0.96	127.91
Seychelles / Madagascar	0.96	123.41
Mayotte / Madagascar	0.966	119.23
Madagascar / China	0.96	117.75
TrinidadTobago / China	0.96	116.92
Germany / Madagascar	0.96	115.56
Reunion / TrinidadTobago	0.96	108.65
TrinidadTobago / Germany	0.96	102.95
Reunion / Madagascar	0.96	100.14
South_Africa / Tunisia	0.95	98.66
Tunisia / Mayotte	0.95	96.04
Seychelles / Mayotte	0.95	91.55
South_Africa / Australia	0.95	89.15
TrinidadTobago / Seychelles	0.94	75.17
South_Africa / Seychelles	0.94	73.71
Australia / Mayotte	0.94	70.02
Australia / China	0.93	60.21
Tunisia / China	0.93	57.20
Reunion / Tunisia	0.93	55.90
Zambia / China	0.92	49.00
Reunion / Australia	0.92	48.90
Seychelles / China	0.91	45.77
Seychelles / Germany	0.91	43.80
Australia / Germany	0.91	42.64

Reunion / Germany	0.90	41.77
Reunion / Seychelles	0.90	41.06
Zambia / Reunion	0.90	37.85
TrinidadTobago / Australia	0.89	37.21
Tunisia / Germany	0.89	34.10
Zambia / Germany	0.88	31.14
Germany / Mayotte	0.88	30.75
South_Africa / Germany	0.87	29.01
Reunion / Mayotte	0.87	28.80
Mayotte / China	0.87	28.31
Germany / China	0.85	25.38
Reunion / China	0.83	21.52
South_Africa / China	0.83	20.66
South_Africa / Reunion	0.80	17.01
TrinidadTobago / Madagascar	0.79	15.81

**Annexe 6.** Serological evidence of widespread circulation of single stranded

RNA viruses in wild fauna from the South Western Indian Ocean islands.

(En cours de soumission)

Séroprévalence obtenue pour 643 chauves-souris et 256 petits mammifères terrestres de la

zone SOOI par la technique d'indirect immunofluorescence assay (IIFA) en utilisant un panel

de virus à ARN simple brin appartenant à 5 familles et 10 genres virales.

Matériels et Méthodes

Collecte des individus

De Mars 2010 à Mars 2015, 643 chauves-souris ont été collectés à Mayotte (archipel des

Comores), La Réunion, Maurice (Mascareignes archipel), Madagascar et Mahé (archipel des

Seychelles). Ces individus se répartissaient en 7 familles, 11 genres et 30 espèces dont 4

espèces frugivores et 26 espèces insectivores. D'autre part, 256 petits mammifères terrestres

ont été collectés à Mayotte (archipel des Comores), La Réunion Maurice (Mascareignes

archipel), Madagascar et Mahé (archipel des Seychelles) et qui se répartissaient en 2 familles,

3 genres et 5 espèces.

Détection d'anticorps

La détection des anticorps a été réalisée en utilisant les lames Euroimmun Mosaïque Chip

(EU38, Lübeck, Allemagne). Les lames présentent sur des champs séparés, des cellules

infectées au niveau de chaque champ par un virus ARN donné (avec en tout 15 virus

analysables) représentant 5 familles virales, et 10 genres :

Genre Coronavirus: SARS coronavirus, (SRAS; UE 14, souche Francfort-1); Human

coronavirus (HCoV-229E; UE 169, souche ATCC (VR-740).

Genre Flavivirus: West Nile virus (VNO; (UE 14, souche Ouganda); Yellow Fever

virus (YFV; UE 14, souche vaccinale RKI-17D); Dengue virus (DENV-2; UE 14, souche TH-

36).

Genre *Hantavirus*: DOBV (DOBVII; UE 14, souche DOB / Slov).

Genre Phlébovirus: RFVF (RFVF; UE 14, souche du Rift 12 / 2/01).

Genre *Nairovirus* : CCHF (CCHFV; UE 90 Zellen, glycoprotéine exprimée Gc (GPC-Protein).

Genre Morbillivirus: Measles (UE 38, souche Edmonston).

Genre Rubulavirus: Mumps (UE 38, souche Jones); PIV-2(PIVII; UE 18/9, souche Greer).

Genre Avulavirus: PIV-1(PIV-I; UE 18/9, souche Sendai CPJ-3)

Genre Pneumovirus: RSV (RSV; UE 38, souche RSV B Wash / 18537 / '62 (CH 18537);

Genre *Alphavirus*: CHIKV (CHIKV; UE 14, souche OPY1, LR2006; La *Réunion 2006*); VEEV (VEEV; UE 14, souche vaccinale).

La première incubation s'est faite par ajout de 70μL de sérum d'un individu dilué au 1/40 sur une puce. L'incubation s'est fait à température ambiante pendant 2 heures. Les lames sont ensuite rincées en Phosphate Buffer Saline (PBS) + 0.1% Tween (PBS-T) pendant 5 min et séchées à l'air libre. La seconde incubation s'est faite par ajout d'immunoglobuline (Ig) de chèvre anti-chauves-souris ou anti-souris (Bethyl Laboratories, Montgomery, TX, USA) dilué au 1/1000 dans du tampon Euroimmun. Les lames sont ensuite rincées en PBS-T pendant 5 min. La troisième incubation s'est faite par ajout d'Ig de singe anti-chèvre conjugué à la cyanine 3 (Dianova, Hamburg, Germany) dilué au 1/2 en glycérol puis au 1/50 dans du PBS et incubés pendant 30 minutes à température ambiante. Les lames sont ensuite rincées en PBS-T pendant 10 minutes et séchées. Les noyaux cellulaires sont colorés au 4',6-diamidino-2-phenylindole (DAPI) et les lames recouvertes d'une lamelle. La lecture s'est faite au microscope à fluorescence (Zeiss).

### Résultats

Réactivités sérologique des chauves-souris et petits mammifères terrestres de la zone SOOI testés par IIFA contre 5 familles dont 10 genres viraux. Résultats exprimés en nombres d'individus positifs sur nombre d'individus total testé. Pourcentage de séropositivité indiqué entre parenthèse.

### A. Coronaviridae

		Coronavirus	
Famille	Espèce	SARS-CoV HCoV-229	
Emballuronidae	Emballonura tiavato	0/1	0/1
Hipposideridae	Hipposideros commersoni	0/14	0/14
Miniopteridae	Miniopterus aelleni	0/5	0/5
	Miniopterus cf. ambohitrensis	0/30	2/30 (6.7)
	Miniopterus giffithsi	1/6 (16.7)	0/6
	Miniopterus gleni	0/13	4/13 (30.8)
	Miniopterus griveaudi	1/38 (2.7)	1/38 (2.6)
	Miniopterus mahafaliensis	0/38	0/38
	Miniopterus sororculus	0/8	0/8
Molossidae	Chaerephon atsinanana	0/7	1/7 (14.3)
	Chaerephon leucogaster	1/43 (2.3)	1/43 (2.3)
	Mops leucostigma	0/32	0/32
	Mops midas	0/16	0/16
	Mormopterus acetabulosus	2/36 (5.6)	0/36
	Mormopterus francoismoutoui	0/46	2/46 (4.3)
	Mormopterus jugularis	3/73 (4.1)	2/73 (2.7)
	Otomops madagascariensis	1/31 (3.2)	0/31
Pteropodidae	Eidolon dupreanum	0/11	0/11
	Pteropus niger	0/24	0/24
	Pteropus rufus	0/17	0/17
	Pteropus seychelliensis	1/39 (2.6)	1/39 (2.6)
	Rousettus madagascariensis	1/42 (2.4)	2/42 (4.8)
Rhinonycteridae	Triaenops furculus	0/6	0/6
	Triaenops menamena	0/32	2/32 (6.25)
Vespertilionidae	Myotis goudoti	2/28 (7.1)	1/28 (3.6)
	Neoromicia	0/2	0/2
	Pipistrellus raceyi	0/1	0/1
	Scotophilus marovaza	0/1	0/1
	vesper	0/3	0/3
Chauves-souris		13/643 (2.0)	19/643 (2.9)
Muridae	Mus musculus	0/3	1/3 (33.3)
	Rattus norvegicus	5/41 (12.2)	1/41 (2.4)
	Rattus rattus	0/155	1/155 (0.6)
Nesomyidae	Eliurus majori	0/4	1/4 (25)
	Eliurus minor	2/53 (3.8%)	0/53 (0%)
Mammifères terrestres		7/256 (2.7)	4/256 (1.6)

# B. Flaviviridae

		Flavivirus		
<b>Famille</b>	Espèce	WNV	YFV	DENV-2
Emballuronidae	Emballonura tiavato	0/1	0/1	0/1
Hipposideridae	Hipposideros commersoni	0/14	0/14	0/14
Miniopteridae	Miniopterus aelleni	1/5 (20)	0/5	1/5 (20.0)
	Miniopterus cf. ambohitrensis	4/30 (13.3)	13/30 (43.3)	10/30 (33.3)
	Miniopterus giffithsi	0/6	1/6 (16.7)	0/6
	Miniopterus gleni	2/13 (15.4)	3/13 (23.1)	3/13 (23.1)
	Miniopterus griveaudi	3/38 (7.9)	2/38 (5.3)	1/38 (2.6)
	Miniopterus mahafaliensis	0/38	4/38 (10.5)	5/38 (13.2)
	Miniopterus sororculus	0/8 ()	3/8 (37.5)	2/8 (25.0)
Molossidae	Chaerephon atsinanana	1/7 (14.3)	1/7 (14.3)	1/7 (14.3)
	Chaerephon leucogaster	7/43 (16.3)	12/43 (27.9)	10/43 (23.3)
	Mops leucostigma	0/32	1/32 (3.1)	1/32 (3.0)
	Mops midas	0/16	1/16 (63)	1/16 (6.3)
	Mormopterus acetabulosus	0/36	4/36 (11.1)	5/36 (13.9)
	Mormopterus francoismoutoui	1/46 (2.2)	9/46 (19.6)	10/46 (21.7)
	Mormopterus jugularis	14/73 (19.2)	45/73 (61.6)	35/73 (47.9)
	Otomops madagascariensis	3/31 (9.7)	8/31 (25.8)	6/31 (19.4)
Pteropodidae	Eidolon dupreanum	9/11 (81.8)	11/11 (100)	10/11 (90.9)
	Pteropus niger	1/24 (4.7)	2/24 (8.3)	3/24 (12.5)
	Pteropus rufus	5/17 (29.4)	5/17 (29.4)	6/17 (35.3)
	Pteropus seychelliensis	3/39 (7.7)	11/39 (28.2)	11/39 (28.2)
	Rousettus madagascariensis	2/42 (4.8)	4/42 (9.5)	5/42 (11.9)
Rhinonycteridae	Triaenops furculus	0/6	1/6 (16.7)	0/6
	Triaenops menamena	1/32 (3.125)	1/32 (3.125)	0/32
Vespertilionidae	Myotis goudoti	1/28 (3.6)	1/28 (3.6)	1/28 (3.6)
	Neoromicia	0/2	0/2	0/2
	Pipistrellus raceyi	0/1	0/1	0/1
	Scotophilus marovaza	0/1	0/1	0/1
	vesper	0/3	0/3	0/3
Chauves-souris		58/643 (9.0)	143/643 (22.2)	127/643 (19.6)
Muridae	Mus musculus	0/3	0/3	0/3
	Rattus norvegicus	0/41	1/41 (2.4)	5/41 (12.2)
	Rattus rattus	5/155 (3.2)	52/155 (33.5)	36/155 (23.2)
Nesomyidae	Eliurus majori	0/4	0/4	0/4
	Eliurus minor	0/53 (0%)	0/53(0%)	1/53 (1.9%)
Mammifères terrestres		5/256 (1.9)	53/256 (20.7)	42/256 (16.4)

# C. Bunyaviridae

		Hantavirus	Phlébovirus	Nairovirus
Famille	Espèce	DOBV	RVFV	CCHFV
Emballuronidae	Emballonura tiavato	0/1	0/1	0/1
Hipposideridae	Hipposideros commersoni	0/14	0/14	1/14 (7.1)
Miniopteridae	Miniopterus aelleni	0/5	0/5	1/5 (20)
	Miniopterus cf. ambohitrensis	0/30	0/30	5/30 (16.7)
	Miniopterus giffithsi	0/6	0/6	2/6 (33.3)
	Miniopterus gleni	0/13	0/13	2/13 (15.4)
	Miniopterus griveaudi	1/38 (2.6)	0/38	7/38 (18.4)
	Miniopterus mahafaliensis	0/38	0/38	4/38 (10.5)
	Miniopterus sororculus	0/8 ()	0/8 ()	1/8 (12.5)
Molossidae	Chaerephon atsinanana	0/7	0/7	0/7
	Chaerephon leucogaster	1/43 (2.3)	1/43 (2.3)	2/43 (4.7)
	Mops leucostigma	0/32	3/32 (9.4)	1/32 (3.1)
	Mops midas	0/16	0/16	0/16
	Mormopterus acetabulosus	0/36	1/36 (2.8)	2/36 (5.6)
	Mormopterus francoismoutoui	0/46	0/46	0/46
	Mormopterus jugularis	0/73	5/73 (6.8)	5/73 (6.8)
	Otomops madagascariensis	0/31	2/31 (6.5)	15/31 (48.4)
Pteropodidae	Eidolon dupreanum	0/11	2/11 (18.2)	2/11 (18.2)
	Pteropus niger	2/24 (8.3)	3/24 (12.5)	0/24
	Pteropus rufus	1/17 (5.9)	1/17 (5.9)	0/17
	Pteropus seychelliensis	12/39 (30.8)	17/39 (43.6)	5/39 (12.8)
	Rousettus madagascariensis	0/42	0/42	1/42 (2.4)
Rhinonycteridae	Triaenops furculus	0/6	0/6	0/6
	Triaenops menamena	0/32	0/32	2/32 (6.25)
Vespertilionidae	Myotis goudoti	0/28	0/28	2/28 (7.1)
	Neoromicia	0/2	0/2	0/2
	Pipistrellus raceyi	0/1	0/1	0/1
	Scotophilus marovaza	0/1	0/1	0/1
	vesper	0/3	0/3	0/3
Chauves-souris		17/643 (2.6)	35/643 (5.4)	60/643 (9.3)
Muridae	Mus musculus	0/3	0/3	1/3 (33.3)
	Rattus norvegicus	3/41 (7.3)	5/41 (12.2)	2/41 (4.9)
	Rattus rattus	33/155 (21.2)	45/155 (29.0)	22/155 (14.2)
Nesomyidae	Eliurus majori	0/4	0/4	0/4
	Eliurus minor	1/53 (1.9%)	3/53 (5.7%)	3/53 (5.7%)
Mammifères terrestres		37/256 (14.5)	53/256 (20.7)	28/256 (10.9)

# D. Paramyxoviridae

		Morbillivirus	Rubulavirus		Pneumovirus	Respirovirus
Famille	Espèce	Measles	Mumps	PIV-2	RSV	PIV-1
Emballuronidae	Emballonura tiavato	0/1	0/1	0/1	0/1	0/1
Hipposideridae	Hipposideros commersoni	3/14 (21.4)	4/14 (28.6)	4/14 (28.6)	1/14 (7.1)	2/14 (14.3)
Miniopteridae	Miniopterus aelleni	1/5 (20.0)	0/5	1/5 (20.0)	1/5 (20.0)	0/5
	Miniopterus cf. ambohitrensis	6/30 (20.0)	12/30 (40.0)	9/30 (30.0)	8/30 (26.7)	1/30 (3.3)
	Miniopterus giffithsi	0/6	1/6 (16.7)	3/6 (50.0)	0/6	0/6
	Miniopterus gleni	1/13 (7.7)	1/13 (7.7)	2/13 (15.4)	0/13	3/13 (23.1)
	Miniopterus griveaudi	8/38 (21.1)	11/38 (28.9)	8/38 (21.1)	4/38 (10.5)	7/38 (18.4)
	Miniopterus mahafaliensis	5/38 (13.2)	10/38 (26.3)	4/38 (10.5)	1/38 (2.6)	3/38 (7.9)
	Miniopterus sororculus	2/8 (25.0)	4/8 (50.0)	4/8 (50.0)	2/8 (25.0)	3/8 (37.5)
Molossidae	Chaerephon atsinanana	0/7	0/7	2/7 (28.6)	0/7	1/7 (14.3)
	Chaerephon leucogaster	11/43 (25.6)	11/43 (25.6)	9/43 (20.9)	4/43 (9.3)	1/43 (2.3)
	Mops leucostigma	1/32 (3.1)	1/32 (3.1)	5/32 (15.6)	0/32	3/32 (9.4)
	Mops midas	5/16 (31.3)	2/16 (12.5)	3/16 (18.75)	1/16 (6.3)	0/16
	Mormopterus acetabulosus	0/36	7/36 (19.4)	2/36 (5.6)	0/36	3/36 (8.3)
	Mormopterus francoismoutoui	0/46	5/46 (10.9)	4/46 (8.7)	1/46 (2.1)	4/46 (8.7)
	Mormopterus jugularis	16/73 (21.9)	38/73 (52.1)	21/73 (28.8)	3/73 (4.1)	2/73 (2.7)
	Otomops madagascariensis	10/31 (32.3)	9/31 (29.0)	5/31 (16.1)	5/31 (16.1)	5/31 (16.1)
Pteropodidae	Eidolon dupreanum	0/11	7/11 (63.6)	8/11 (72.7)	0/11	0/11
	Pteropus niger	1/24 (4.7)	3/24 (12.5)	15/24 (62.5)	0/24	1/24 (4.2)
	Pteropus rufus	0/17	3/17 (17.6)	11/17 (64.7)	0/17	0/17
	Pteropus seychelliensis	2/39 (5.1)	12/39 (30.8)	12/39 (30.8)	2/39 (5.1)	0/39
	Rousettus madagascariensis	3/42 (7.1)	19/42 (45.2)	25/42 (59.5)	2/42 (4.7)	3/42 (7.1)
Rhinonycteridae	Triaenops furculus	1/6 (16.7)	1/6 (16.7)	0/6	0/6	0/6
	Triaenops menamena	4/32 (12.5)	2/32 (6.25)	6/32 (18.8)	3/32 (9.4)	2/32 (6.3)
Vespertilionidae	Myotis goudoti	5/28 (17.9)	6/28 (21.4)	7/28 (25.0)	2/28 (7.1)	3/28 (10.7)
•	Neoromicia	1/2 (50)	1/2 (50)	0/2	0/2	0/2
	Pipistrellus raceyi	0/1	0/1	0/1	0/1	0/1
	Scotophilus marovaza	0/1	0/1	0/1	0/1	0/1
	vesper	0/3	0/3	0/3	0/3	0/3
Chauves-souris	•	86/643 (13.4)	170/643 (26.4)	170/643 (26.4)	40/643 (6.2)	47/643 (7.3)
Muridae	Mus musculus	2/3 (66.7)	2/3 (66.7)	3/3(100)	2/3(66.7)	0/3
	Rattus norvegicus	13/41 (31.7)	16/41 (39.0)	9/41(21.9)	7/41(17.1)	4/41(9.8)
	Rattus rattus	34/155 (21.9)	58/155 (37.4)	46/155(29.7)	26/155(16.8)	47/155(30.3)
Nesomyidae	Eliurus majori	0/4	0/4	0/4	0/4	1/4(25)
	Eliurus minor	16/53 (30.2%)	15/53 (28.3%)	6/53 (11.3%)	6/53 (11.3%)	7/53 (13.2%)
Mammifères terrestres		65/256 (25.4)	91/256 (35.5)	64/256(25.0)	41/256(16.0)	59/256(23.0)
		·	·			

# E. Togaviridae

		Alphavirus	
<b>Famille</b>	Espèce	VEEV	CHIKV
Emballuronidae	Emballonura tiavato	0/1	0/1
Hipposideridae	Hipposideros commersoni	0/14	0/14
Miniopteridae	Miniopterus aelleni	0/5	0/5
	Miniopterus cf. ambohitrensis	0/30	1/30 (3.3)
	Miniopterus giffithsi	0/6	0/6
	Miniopterus gleni	1/13 (7.7)	1/13 (7.7)
	Miniopterus griveaudi	0/38	0/38
	Miniopterus mahafaliensis	0/38	0/38
	Miniopterus sororculus	0/8 ()	1/8 (12.5)
Molossidae	Chaerephon atsinanana	0/7	1/7 (14.3)
	Chaerephon leucogaster	0/43	0/43
	Mops leucostigma	0/32	0/32
	Mops midas	0/16	0/16
	Mormopterus acetabulosus	0/36	0/36
	Mormopterus francoismoutoui	0/46	0/46
	Mormopterus jugularis	0/73	1/73 (1.4)
	Otomops madagascariensis	0/31	5/31 (16.1)
Pteropodidae	Eidolon dupreanum	0/11	0/11
	Pteropus niger	0/24	1/24 (4.7)
	Pteropus rufus	0/17	0/17
	Pteropus seychelliensis	0/39	7/39 (17.9)
	Rousettus madagascariensis	0/42	1/42 (2.4)
Rhinonycteridae	Triaenops furculus	0/6	0/6
	Triaenops menamena	0/32	0/32
Vespertilionidae	Myotis goudoti	1/28 (3.6)	1/28 (3.6)
	Neoromicia	0/2	0/2
	Pipistrellus raceyi	0/1	0/1
	Scotophilus marovaza	0/1	0/1
	vesper	0/3	0/3
Chauves-souris		2/643 (0.3)	20/643 (3.1)
Muridae	Mus musculus	0/3	0/3
	Rattus norvegicus	0/41	3/41 (7.3)
	Rattus rattus	0/155	10/155 (6.5)
Nesomyidae	Eliurus majori	0/4	/4
-	Eliurus minor	0/53 (0%)	3/53 (5.7%)
Mammifères terrestres		0/256	0/256 (6.3)

### Annexe 7. Liste des publications

### Contribution à d'autres travaux: publication acceptée

Salez, N., Mélade, J., Pascalis, H., Aherfi, S., Dellagi, K., Charrel, RN., Carrat, F. & de Lamballerie, X. (2014). Influenza C virus high seroprevalence rates observed in 3 different population groups. J. Infect. Aug;69(2):182-9. doi: 10.1016/j.jinf.2014.03.016. Epub 2014 Apr 4.

### Faisant partie du sujet de thèse : publication acceptée

 Wilkinson, D.A., Mélade, J., Dietrich, M., Ramasindrazana, B., Soarimalala, V., Lagadec, E., le Minter, G., Tortosa, P., Heraud, J.M., de Lamballerie, X., Goodman, S.M., Dellagi, K. & Pascalis, H. (2014). Highly diverse morbillivirus-related paramyxoviruses in wild fauna of the southwestern Indian Ocean Islands: evidence of exchange between introduced and endemic small mammals. J. Virol. Aug;88(15):8268-77. doi: 10.1128/JVI.01211-14. Epub 2014 May 14.

### Faisant partie du sujet de thèse : publications soumises

- 3. **Mélade, J.**, McCulloch, S., Ramasindrazana, B., Lagadec, E., Turpin, M., Pascalis, H., Goodman, S.M., Markotter, W. & Dellagi, K. Serological evidence of diverse circulation of lyssaviruses among bats on southwestern Indian Ocean islands (en revision).
- 4. **Mélade**, **J**., Wieseke, N., Ramazindrazana, B., Flores, O., Lagadec, E., Gommard, Y., Goodman, S.M., Dellagi, K., & Pascalis, H. An eco-epidemiological study of Morbilli-related paramyxovirus infection in Madagascar bats reveals host-switching as the dominant macro-evolutionary mechanism (en revision).

### Faisant partie du sujet de thèse : publication en cours de soumission

5. **Mélade, J.**, Müller, M., Ramasindrazana, B., Lagadec, E., Turpin, M., Pascalis, H., Goodman, S.M., Drosten, C. & Dellagi, K. Serological evidence of widespread circulation of single stranded RNA viruses in wild fauna from the South Western Indian Ocean islands (en preparation).

# En rapport avec le sujet de thèse : publication soumise

6. Ghawar, W., Pascalis, H., Bettaie, J., Wilkinson, D., **Mélade, J**., El Gharbi, A., Snoussi, M.A., Laouini, D., Ben Salah, A. & Dellagi, K. Insight into the global evolution of Rodentia associated Morbillirelated Paramyxoviruses reveals the role of Rattus as a worldwide spreader (en revision).

## Annexe 8. Formations, congrès

- Congrès international « Bats, small mammals and infectious agents » (2013, St Denis, La Réunion). Communication orale.
- 2. Congrès international « **African Small Mammal Symposium** » (2015, Mantasoa, Madagascar). Communication orale.
- 3. Journées scientifiques de la « **Fédération Environnement, Biodiversité et Santé** » (2014, Saint-Denis La Réunion). Communication orale.
- 4. Colloque **FEDER POCT Biodiversité** (2015, Saint-Denis La Réunion). Communication orale.
- 5. Stage doctoral de 3 semaines du 6 Septembre 2014 au 27 Septembre 2014 au Centre Hospitalier Universitaire de Bonn en Allemagne dans le laboratoire de Virologie du Pr Christian Drosten.
- 6. Stage doctoral de 5 mois à l'Université de Pretoria en Afrique du Sud dans le laboratoire de virologie du Pr Wanda Markotter.

# Résumé

La faune sauvage a été depuis longtemps incriminée dans la survenue de zoonoses et joue le rôle de réservoir d'agents pathogènes (virus *Nipah*, *Hendra*, *Ebola*, *Hantaan* etc.) pour l'homme. Les îles tropicales et subtropicales du Sud-Ouest de l'Océan Indien (SOOI) constituent l'une des 34 régions reconnues comme « *hotspot* » de biodiversité au niveau mondial. Elles sont caractérisées par un très fort endémisme de la faune sauvage surtout sur l'Ile de Madagascar. Le caractère multi-insulaire de la région du SOOI, la diversité de ses biotopes et ses disparités biogéographiques et humaines offrent un champ d'investigation unique pour explorer « *in natura* » la dynamique évolutive des agents infectieux et les relations hôtes-virus.

Nos travaux de recherche ont porté sur deux modèles de virus à ARN de polarité négative, les paramyxovirus et les lyssavirus. Le premier modèle viral nous a permis d'aborder les questions relatives à la dynamique de transmission virale au sein de communauté d'hôtes, plus particulièrement, les chauves-souris et les petits mammifères terrestres de Madagascar et d'identifier les facteurs agissant sur cette dynamique de transmission et de diversification virale, en particulier les facteurs bio-écologiques associés à leurs hôtes. Le second modèle viral, les lyssavirus, nous a permis de décrire sur l'ensemble des îles du SOOI, la circulation virale dans ce système multi-insulaire diversifié, au sein des chauves-souris dont la plupart des espèces sont endémiques à cette région.

Dans l'ensemble, nos investigations ont permis de mettre en évidence des échanges viraux (« host-switch ») importants entre chauves-souris, petits mammifères terrestres endémiques de Madagascar et les rongeurs introduits, le rôle de ces mammifères en tant que réservoir viral majeur et souligner le rôle disséminateur de Rattus rattus. Par ailleurs, nous avons pu identifier ce phénomène de « host-switch » comme étant le mécanisme macro-évolutif prépondérant et l'importance des facteurs biotiques et abiotiques à l'origine de la dynamique de transmission et de la diversification virale observée chez les paramyxovirus de chauves-souris de Madagascar.

**Mots-clés** : faune sauvage, zoonoses, SOOI, biodiversité, lyssavirus, paramyxovirus, diversité virale, réservoir viral, disséminateur viral, « *host-switch* » facteurs écologique