



HAL
open science

Variabilité intraspécifique de la résistance à l'embolie et trade-offs fonctionnels chez le tournesol (*Helianthus annuus* L.) et le pin maritime (*Pinus pinaster* Ait.)

Hafiz Bashir Ahmad

► To cite this version:

Hafiz Bashir Ahmad. Variabilité intraspécifique de la résistance à l'embolie et trade-offs fonctionnels chez le tournesol (*Helianthus annuus* L.) et le pin maritime (*Pinus pinaster* Ait.). Sylviculture, foresterie. Université de Bordeaux, 2017. Français. NNT : 2017BORD0747 . tel-01661596

HAL Id: tel-01661596

<https://theses.hal.science/tel-01661596>

Submitted on 12 Dec 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THESE

Pour obtenir le grade de

DOCTEUR DE L'UNIVERSITE DE BORDEAUX

Spécialité: Ecologie évolutive, fonctionnelle et des communautés

Ecole doctorale: Sciences et Environnements

Intraspecific variability in embolism resistance and functional trade-offs in sunflower (*Helianthus annuus* L.) and Maritime pine (*Pinus pinaster* Ait.)

Variabilité intraspécifique de la résistance à l'embolie et trade-offs fonctionnels chez le tournesol (*Helianthus annuus* L.) et le pin maritime (*Pinus pinaster* Ait.)

Hafiz Bashir AHMAD

Directeur: Sylvain DELZON (DR, INRA Bordeaux)

Co-directeur: Annabel PORTÉ (CR, INRA Bordeaux)

Soutenue le 14 Novembre 2017

Membres du jury:

M. Christophe PLOMION	(DR, INRA Bordeaux)	Président
M. Hervé COCHARD	(DR, INRA Clermont-Ferrand)	Rapporteur
M. Philippe ROZENBERG	(DR, INRA Orléans)	Rapporteur
M. Nicolas MARTIN-StPAUL	(CR, INRA Avignon)	Examinateur
Mme. Annabel PORTÉ	(CR, INRA Bordeaux)	Co-directeur de thèse
M. Sylvain DELZON	(DR, INRA Bordeaux)	Directeur de thèse

Title: Intraspecific variability in embolism resistance and functional trade-offs in sunflower (*Helianthus annuus* L.) and Maritime pine (*Pinus pinaster* Ait.)

Summary

As a consequence of current global climate change, forest and crop ecosystems are decreasing in productivity worldwide and are under increased threat of widespread drought-induced mortality due to hydraulic failure caused by xylem embolism. The range of embolism resistance across species has been intensively studied but little attention has been paid to intraspecific variability. This thesis aimed therefore at improving our understanding of the intraspecific variation in embolism resistance in one crop species (*Helianthus annuus* L.) and one conifer species (*Pinus pinaster*). In the first place, vulnerability to embolism was determined across four sunflower accessions with significant variations in P_{50} (pressure inducing 50% loss of hydraulic conductivity) ranging between -2.67 and -3.22 MPa. A trade-off between hydraulic safety and growth was observed but not between hydraulic safety and efficiency. Only few anatomical traits, such as vessel density and vessel lumen area, were found related to embolism resistance. Secondly, we phenotyped 477 *P. pinaster* full-sib F2 genotypes corresponding to a three-generation inbred pedigree obtained by the self-pollination of an inter “Corsica × Landes” provenance hybrid, for embolism resistance and wood density. The variability for P_{50} within species was remarkably low ($CV_P=5.6\%$). We also found a lack of trade-off between hydraulic safety and wood density. The phenotypic variance explained by QTLs (PEV) detected for P_{50} accounted for 23.6% of the phenotypic variation but no QTL collocated between P_{50} and wood density, suggesting that embolism resistant genotypes in this family are not dependent on denser wood. A multi-trait analysis of leaf and xylem functioning was performed through seasonal measurements of photosystem II fluorescence and stem hydraulic traits on a full-sib family of an inter “Morocco × Landes” provenance hybrid of *P. pinaster*; it revealed that with decreasing water potential, the efficiency of both photosystem II and stomatal conductance decreased which resulted in rapid decrease in assimilation rate. Xylem resistance to embolism was found having no relationship with leaf functioning traits. The absence of trade-off between these traits makes it possible to consider selection for both efficient and resistant genotypes.

Key words: embolism, drought, intraspecific variability, xylem, vessel anatomy, QTL, wood density

Titre: Variabilité intraspécifique de la résistance à l'embolie et trade-offs fonctionnels chez le tournesol (*Helianthus annuus* L.) et le pin maritime (*Pinus pinaster* Ait.)

Résumé

Plusieurs études ont montré que la productivité des forêts et des cultures décroît sur l'ensemble de la planète en réponse au changement climatique, et que les risques de mortalité liée à la sécheresse s'accroissent. Ces mortalités massives ont été observées dans tous les écosystèmes forestiers et seraient dues à un dysfonctionnement hydraulique des plantes, causé par de l'embolie xylémiène. Le spectre interspécifique de résistance à l'embolie a été largement étudiée, mais peu d'attention a été accordée à la variabilité intra-spécifique de ce trait. C'est pourquoi, l'objectif de ce travail de thèse était d'améliorer notre compréhension de la variabilité intra-spécifique de la résistance à l'embolie chez une espèce cultivée (*Helianthus annuus* L.) et chez un conifère (*Pinus pinaster*). Nous avons déterminé la vulnérabilité à l'embolie de quatre variétés de tournesol, présentant des différences significatives de P_{50} (pression induisant 50% de perte de conductivité hydraulique), allant de -2.67 à -3.22 MPa. Un compromis entre sécurité hydraulique et croissance a été observé mais pas entre la sécurité hydraulique et la capacité de transport de l'eau du xylème. Seuls quelques caractères anatomiques, tels que la densité des vaisseaux et la surface du lumen, présentent une corrélation avec la résistance à l'embolie. Dans un second temps, nous avons phénotypé pour la résistance à l'embolie et la densité du bois 477 génotypes de *P. pinaster*, issus d'une F2 de pleins-frères correspondant à la 3^{ème} génération consanguine obtenue par auto-fécondation d'un hybride inter-provenance "Corse × Landes. La variabilité de la P_{50} au sein de l'espèce était extrêmement faible ($CV_P=5.6\%$). D'autre part, nous avons mis en évidence l'absence de compromis entre sécurité hydraulique et densité du bois. La part de variance du caractère P_{50} expliquée par QTLs (PEV) correspondait à 23.6% de la variance phénotypique totale mais aucun QTL commun n'a pu être détecté entre P_{50} et densité du bois, suggérant qu'au sein de cette famille, la plus forte résistance à l'embolie de certains génotypes n'est pas liée à un bois plus dense. Une analyse multi-caractère du fonctionnement de la feuille et du xylème a été réalisée via l'étude de la dynamique saisonnière de la fluorescence du photosystème II et des traits hydrauliques de la tige, sur une famille de pleins-frères d'un hybride inter-provenance "Maroc × Landes" de *P. pinaster*; en réponse à une diminution du potentiel hydrique, l'efficacité du photosystème II et la conductance stomatique ont tous deux diminué, induisant une décroissance rapide du taux d'assimilation. La résistance à l'embolie du xylème n'a pas montré de relation significative avec les traits foliaires. L'absence de compromis entre ces caractères permet donc d'envisager la sélection génétique de génotypes à la fois efficaces et résistants.

Mots-Clefs: embolie, sécheresse, variabilité intraspécifique, xylème, anatomie des vaisseaux, QTL, densité du bois

BIOGECO

[Biodiversité, gènes et communautés, UMR 1202, 33610 Pessac, France]

Table of contents

Summary.....	i
Résumé.....	ii
Table of contents.....	iii
Introduction.....	1
Drought-induced plant mortality and crop productivity decline.....	2
Water transport in vascular plants (The cohesion-tension theory).....	4
Cavitation and embolism formation in Plants.....	5
Resistance to embolism formation.....	6
Assessment of xylem vulnerability to drought induced embolism.....	7
Interspecific variability in embolism resistance.....	11
Intraspecific variability in embolism resistance.....	14
Intraspecific variability in embolism resistance in crops.....	16
Objectives of the thesis.....	16
Part I: Intraspecific variability in embolism resistance in sunflower crop.....	20
Chapter 1: Intraspecific variation in embolism resistance and stem anatomy across four sunflower (<i>Helianthus annuus</i> L.) accessions. (Accepted in <i>Physiologia Plantarum</i>).....	21
Supplementary information.....	51
Part II: Intraspecific variability in embolism resistance and its relation with genetic architecture and photosynthetic traits in Maritime pine.....	53
Chapter 2: Dissecting the genetic architecture of resistance in Maritime pine reveals that hydraulic safety is not linked to wood density. (in preparation).....	54
Supplementary information.....	76
Chapter 3: Intraspecific variability of the physiological response to water stress in a full-sib Maritime pine family. (in preparation).....	86
Discussion.....	109
Vulnerability curves assessment in long vesselled species.....	110
Intraspecific variability in embolism resistance in sunflower and maritime pine.....	111
Trade-offs with hydraulic traits.....	115
Implications and future perspectives.....	119
References.	122

Introduction

Introduction

Drought-induced plant mortality and crop productivity decline

Ecosystems across the globe are facing continuous threats by anthropogenic climate change, resulting natural populations to adapt, migrate or extinct (Walther et al., 2002). Shifting precipitation patterns, rising temperatures, heat waves combined with severe and frequent drought spells, triggered by climate change during the last few decades, may increase plant mortality (Allen et al., 2010; Bentz et al., 2010; Elderd et al., 2010; O' Grady and Mitchell, 2015;). The impacts of this changing climate have already been well recognized in ecosystems across the world, ranging from coral-reef dieback due to ocean acidification and temperature rise (Hoegh-Guldberg et al., 2007), extreme climatic events affecting many terrestrial ecosystems (Allen et al., 2010; Zhao and Running, 2010) to severely impacting human population globally, through crop failure, storm damage, extensive flooding and causing serious health issues by the spread of infectious fatal diseases (Patz et al., 2005). Existing climate models predict further decreases in soil moisture contents in the top 10 cm layer (Dai, 2013) and widespread increases in the intensity and duration of periodic water stress having severe consequences on plant growth and distribution worldwide during the 21st century (Van Mantgem et al., 2009; Anderegg et al., 2012; Engelbrecht, 2012; IPCC, 2014; McDowell et al., 2015).

According to FAO (2006) 30% of the world's land surface is under forests, including woodlands and savannas. Innumerable tangible and intangible benefits, around the globe, are derived from forests, ranging from the provision of essential goods and services such as timber (Payn *et al.*, 2015) and watershed protection/conservation to recreational, aesthetic, and spiritual amity. Understanding the mechanism underlying as well as predicting the consequences of these climatic changes in ecosystems and forecasting their impacts on forests and other vegetation types, is therefore, of particular importance and emerging as one of the striking challenges for scientists focusing on global change impacts (Boisvenue and Running, 2006; Bonan, 2008). The common implicated causal factor of forest die-offs and tree mortality in recent decades in southern Europe (Breda et al., 2006; Bigler et al., 2006), temperate forests of Western North America (van Mantgem et al., 2009), boreal forests of Canada (Peng et al., 2011) and extensive death of several tree species across different forest types over an area of about 10 million hectares (Raffa et al., 2008) is elevated temperatures and/or water stress. This indicates that the forests at a global scale across all eco-

regions are under greater risk and respond increasingly to drought and ongoing warming and drying (Allen et al., 2010; 2015).

In the face of climate change, similar to forest research, studies on predicting responses to non-lethal water stress levels in crops have also been focused for better understanding of breeding programs and water management (Blum, 2005; Tardieu, 2012) in order to obtain varieties with improved drought tolerance (Xoconostle-Cásarez et al., 2010). A pronounced decline in crop net primary productivity (NPP) was observed in 2003 throughout Europe following strong heatwaves (northern Italy, France) and drought (Ukraine, Romania). For instance, the greatest drop in maize NPP (-36%) was found in the Italian Po valley, where extremely high temperatures prevailed (Ciais et al., 2005). A continental scale analysis for 1901–2012 conducted by Trnka et al. (2016) also highlighted that barley and wheat yields were affected by climate variability in Central and Western Europe. The yield decline was more pronounced in Central Europe than in Western Europe due to comparatively drier conditions and variation in annual temperatures to a greater range. Pinke and Lövei, (2017) recently showed that increasing temperatures over the 1981-2010 period have negatively impacted the yields of barley, wheat, maize and rye in Hungary. Specifically, a 1°C rise in temperature reduced the yield of all four cereals by 9.6 to 14.8% in this Eastern European country that is situated on the western border of the Eurasian Steppe Belt. Overall, these studies confirm the high vulnerability of European agricultural regions to ongoing climate warming. Similarly, 32 to 39% of maize, rice, wheat and soybean yields are vulnerable to climate variability worldwide (Ray et al. 2015), which translates into large fluctuations in global crop production.

Direct and indirect effects of climate change (Wheeler and von Braun, 2013) associated with soil degradation/erosion (Pimentel et al., 1995), heatwave and drought (Ciais et al., 2005) and shrinking freshwater reserves (Tester & Langridge, 2010; Trnka et al., 2011) along with some other socio-economic changes such as decreasing fertilization in Europe (Peltonen-Sainio and Jauhiainen, 2014) are the main drivers of the recent decreases in crop productivity recorded in some regions. This causes serious concerns about global food security (Godfray et al., 2010) due to increase in per capita food consumption, which is predicted to further increase by ca. 70% by 2050 (FAO, 2009; Mansharamani, 2015) by rapidly growing human population (Mitchell, 2008). Many regions worldwide are recognized being affected from more frequent, prolonged, and severe

drought episodes (Stocker et al., 2013). The subsequent impact of transformation in climatic regimes on the biosphere is a long-term process, small data sets, based on regional observations, allow only limited conclusions (IPCC, 2013). Hence, in-depth analyses of regional patterns of climate change would be required to complement usefully in global analyses for better understanding of geographical variation of all these factors and to improve our knowledge of plants' response to drought and other abiotic stresses associated with climate change. It is, indeed, also required for improved modelling of future plant distributions, to predict areas highly vulnerable for plant mortality and to plan ahead by selection of species and/or varieties to assist in breeding programs for agriculture as well as forestry sector.

Water transport in vascular plants (The cohesion-tension theory)

The ability of plants to transport water through xylem is one of the most important aspects in plant-water relations. Plant growth depends on a number of morphological and physiological traits as well as on environmental factors like precipitation, temperature and nutrient availability (Wright and Westoby 1999). Efficiency of nutrient uptake (Cornelissen et al. 1997) and water status (Major and Johnsen 1999; Hsiao and Xu 2000) are important physiological factors which are known to influence plant growth. During the process of photosynthesis 200 to 400 molecules of water are needed to fixate one molecule of CO₂. Generally about 90% of the water absorbed from soil is transmitted to the atmosphere during this process (Zimmermann, 1983; Kramer and Boyer, 1995; Cruziat et al., 2001; Chavarria and Pessoa Dos Santos, 2012). Trees are the unique organisms capable of transporting huge amounts of water from the soil to the leaves at the top of the crown covering long distances of more than 100 meters (Ryan et al., 2006). Water moves through the entire plant body and evaporates into the atmosphere reaching the surface of the leaf through microscopic pores called stomata, which also allow the entrance of carbonic gas required essentially for photosynthesis. The phenomenon is called transpiration. To replace the water lost by transpiration, continuous flow of water is necessary to insure the growth and survival of plants.

The cohesion-tension theory is generally accepted mechanism for long-distance passive water-transport (Dixon and Joly, 1894; Sperry et al., 1996; Wei et al., 1999; Tyree and Zimmermann, 2002; Angeles et al., 2004). Evaporation of water from surface of leaves, a process called transpiration, is the driving force for water transport. Water molecules stick together due to cohesive forces (hydrogen bonding) and are gradually pulled up in the form of water-column

through the entire plant by the surface-tension of each air/sap meniscus in the leaves creating a water-potential gradient which becomes more negative following the soil-plant water continuum. A water potential is the potential energy of water per unit volume relative to pure water in a reference condition which is 0 MPa. Plant water potentials are negative (from 0 to -10 MPa), whereas water potential in the atmosphere is even more negative. This difference in water potentials creates a pressure gradient that drives the evaporation of water from leaves into the atmosphere (Zimmermann, 1983). Water in a plant is in a metastable thermodynamic state and remains in liquid phase even under negative pressure due to the substantial cohesive forces among the water molecules. Therefore, according to the cohesion-tension theory, as water evaporates via stomata, it decreases the leaf water potential, enabling water to move from stems to leaves thus lowering the water potential in stems and consequently causing water movement from roots to stems and from soil to roots.

Cavitation and embolism formation in Plants

The efficient water transport system enables plants to extract water from dry soils, and move it to long distances above the ground against the gravity pull, it depends entirely on maintaining an intact vascular network filled with water essentially in a metastable liquid state and hence, under a permanent risk of breakdown of column which disrupts the water flow (Tyree and Sperry, 1989). Indeed, during drought, the pressure becomes more negative and tension in the plant increases which means that air-bubbles can be drawn into conductive xylem elements from neighboring air-filled spaces or cells, according to the “air-seeding hypothesis” (Cochard et al., 2009), leading to partial blockage of water flow through the conduits, reducing the ability of plants to transport water from soil to leaf. This phenomenon, also called cavitation, has been described repeatedly over the last decades but clear understanding is still to be achieved (Schenk et al., 2015). Trees can protect themselves from this situation by reducing transpiration through closing their stomata and prevent the spread of embolism. However, during long and/or intense drought spells, stomatal control is not sufficient to regulate the gradient pressure and tension may reach such a highly negative value that the small initial gas-bubble spreads to fill the entire cell, which is then air-filled, or embolized, and therefore no longer remain functional, thereby reducing the plant’s overall capacity to transport water to its aerial organs (Sperry and Tyree, 1988; Cochard et al., 2006). The number of cavitation events increases with increasing evaporative demand at the leaf

level which results in lowering (more negative) the soil water-potential, reducing the hydraulic conductance drastically (Sperry and Sullivan, 1992). Hydraulic failure is one of the major causes of plant mortality due to xylem embolism under severe drought events (Choat et al., 2012). A conclusive evidence of a direct relation between plant death and dramatic failure of water transport has been highlighted in conifers (Brodribb and Cochard, 2009; Brodribb et al., 2010) as well as in angiosperms (Barigah et al., 2013; Urli et al., 2013) demonstrating that the species tolerance to water stress and their ecological distribution depends, at least partly, on the efficiency of their hydraulic system and their ability to withstand embolism (Maherali et al., 2004; Cochard et al., 2008; Anderegg et al., 2015). In a recent meta-analysis on mortality rates across 475 species from 33 studies around the globe, it was found that species-specific mortality rates in a given drought were associated with plant hydraulic traits. It showed that across all species, including angiosperms and gymnosperms, P_{50} is the trait that better explained mortality in trees, supporting the hypothesis that hydraulic traits hold key mechanisms determining tree death and highlight that physiological traits can improve prediction of tree mortality during climatic extremes (Anderegg et al., 2016).

Resistance to embolism formation

Air leakage from embolized to functional conduits is the main reason of xylem cavitation due to the presence of tiny inter-conduit openings called pits (Crombie et al., 1985; Cochard et al., 1992; Jarbeau et al., 1995; Tyree and Zimmermann, 2002). It was thought that in both angiosperms and gymnosperms, when the pressure in the xylem becomes negative enough, capillary rupture of an air-sap meniscus in a pit pair (air-seeding) results in xylem cavitation, enabling air to spread from a non-functional (air-filled, embolized) conduit to an adjacent functional (sap filled) conduit (Cochard et al., 2009; Delzon et al., 2010). Porosity of the xylem conduit wall would therefore influence the resistance to cavitation. Thus, variations in xylem anatomical traits among species are related to cavitation resistance in angiosperms (Sperry and Hacke, 2004; Jansen et al., 2009; Lens et al., 2011) and conifers (Hacke et al., 2004; Domec et al., 2006; Delzon et al., 2010; Pittermann et al., 2010, Bouche et al. 2014). In angiosperms, a porous membrane of micro fibrils is located inside the pit chamber in inter-vessel pits. Embolism may spread by capillary rupture of an air-sap meniscus depending on the size of the largest pores in the pit membranes (Christman et al., 2009). However, it has been suggested by some authors that embolism formation might not be caused by immediate expansion of an air bubble as described by the air-seeding hypothesis due to

the presence of nanobubbles in the sap of angiosperms (Schenk et al., 2015). Instead, these authors suggest that due to gas supersaturation of xylem sap or decreasing pressure, stable nanobubbles might become unstable and could lead to embolism formation. However, the presence of nanobubbles in conifer sap is not confirmed and so far, the air-seeding hypothesis related to xylem anatomy is still privileged. In bordered pit pairs of conifers, pit membranes are composed of two distinct regions: a torus located in the center serving as a valve to isolate the air-filled tracheids from the rest of the system (Bailey, 1916) and a porous margo region which surrounds the torus acting as a barrier to air bubbles and prevents the spread of embolism throughout the xylem. In a well-watered condition, the torus-margo structure is in a neutral position within the pit chamber and water flows through the large pores of the margo. Whereas under water stress conditions, the negative pressure in the xylem deflects the torus against the pit aperture. This valve effect creates a seal that prevents the spread of air into the xylem depending on the width of the torus relative to the pit aperture (Bailey, 1913; Hacke et al., 2004; Domec et al., 2006; Delzon et al., 2010). This seems to confer increased resistance to drought-induced cavitation in Gymnosperms, which are generally more resistant than Angiosperms (Maherali et al., 2004; Choat et al., 2012). Two hypotheses have been proposed for the spread of cavitation in this sophisticated “airtight” bordered pit: i) when the torus and the inner wall of the pit membrane are not completely sealed, air bubbles pass through pores at the edge of the torus (seal capillary seeding hypothesis, Cochard et al., 2009; Delzon et al., 2010; Pittermann et al., 2010) and ii) the torus structure is not fully airtight enabling the air bubble pass through tiny pores (torus capillary seeding hypothesis, Jansen et al., 2012). However, the later hypothesis was not confirmed by Bouche et al. (2014) who proposed that embolism was most likely to occur by seal capillary-seeding by overlapping the torus on pit aperture, whereas torus capillary-seeding could provide an additional mechanism in Pinaceae.

Assessment of xylem vulnerability to drought induced embolism

Cavitation is a fundamental phenomenon of plant-water relations and its multidimensional links with xylem anatomy and ecophysiological implications are important aspect to understand. Indeed, xylem vulnerability to cavitation is one of the major issue driving forest productivity and drought resistance in plants (Anderegg et al. 2012, Choat et al. 2012). With the increased understanding of plant water-transport, a variety of techniques have been developed to induce and measure the

extent of cavitation in order to estimate xylem vulnerability to drought induced cavitation (VCs, Cochard et al., 2013).

Embolism induction:

Bench dehydration method is the most simple and natural way of inducing embolism slowly in a whole intact plant or in cut branches by letting them dehydrate *in situ* or on a bench under ambient light conditions. Xylem pressure is assessed with a pressure chamber on non-transpiring covered leaves or with stem psychrometers (Tyree et al., 1992; Bréda et al., 1993).

Air pressurization method induces cavitation by forcing the air into a xylem segment inserted into a standard Scholander pressure chamber or a double ended pressure chamber, which allows the increase of air pressure, while the water flow through the sample is also measured at the same time (Crombie et al., 1985b; Sperry and Tyree, 1990; Cochard et al., 1992; Choat et al., 2005).

Centrifugation method exposes xylem to highly negative pressure by applying centrifugal forces (Pockman et al., 1995). The principle is to reduce xylem pressure in the middle part of a short branch segment by spinning it in a centrifuge.

Embolism Measurement:

1. *Acoustic emissions* (AEs), from audible “clicks” to ultrasound emissions, produced during drought stress are recorded automatically by a sensor clamped to a branch, leaf or trunk, even under field conditions, while the plant is subjected to natural or induced water stress (Tyree et al., 1984; Tyree and Sperry, 1989a).

2. *Observations of xylem water content* can be made with different methodologies to make direct or indirect observations of xylem water content such as magnetic resonance imaging technology (Holbrook et al., 2001) or cryo-scanning electron microscopy (Canny, 1997; Cochard et al., 2004). Another appropriate method for visualizing water or air in xylem conduits with relatively high spatial resolution is the high-resolution X-ray computed tomography (Fromm et al., 2001; Brodersen et al., 2012; Torres-Ruiz et al., 2015). However, this method is yet not being used for the routine study of xylem embolism in plants.

3. *Hydraulic detection of embolism* involve directly monitoring decrease in water-flow as stress is imposed to a sample to induce negative pressure in the xylem analogous to that experienced by

the plant during drought. Decrease in xylem hydraulic efficiency (specific conductance, k_s) caused by cavitation events is measured. The degree of embolism in a xylem segment can be quantified as the percentage loss of hydraulic conductance (PLC), which can be determined gravimetrically (Sperry et al. 1988) or with a Xylem apparatus. The principle of these techniques is to measure the relative decrease in the efficiency of xylem to transport water caused by the presence of air bubbles which considerably increase the resistance to water flow through the conduits.

Cavitron technique is the *in situ* flow centrifuge technique described by Cochard, (2002a) and Cochard et al. (2005) that enables the construction of a vulnerability curve in a very short time of less than half an hour. Samples are placed in a custom-built rotor with both ends in reservoirs filled with water. Small holes are made in the reservoirs at two distances from the rotation axis in order to generate a pressure gradient (ΔP , MPa) across the sample and to induce water flow through the sample. The hydraulic conductivity can then be determined directly during centrifugation and an entire vulnerability curve is constructed, thus allowing large scale investigations of interspecific, intraspecific, genetic and phenotypic variability of cavitation resistance in Gymnosperm as well as in Angiosperms (Delzon et al., 2010; Lamy et al., 2011; Wortemann et al., 2011).

By visualizing the percent loss of xylem conductance (PLC) as a function of xylem pressure (MPa), one obtains a vulnerability curve (VC, generally a sigmoid), from which the cavitation resistance of a sample is estimated, notably from the inflexion point, i.e., the pressure at which 50% of total xylem hydraulic conductance is lost (P_{50} , in Mega Pascal), and the slope of the curve (S , in % MPa⁻¹). A sigmoid function (Pammenter and Vander Willigen 1998) is fitted to the VC using the following equation:

$$PLC = \frac{100}{\left[1 + \exp\left(\frac{S}{25} \times (\Psi - P_{50})\right)\right]}$$

where P_{50} (MPa) is the xylem pressure inducing 50% loss of conductance and S (% MPa⁻¹) is the slope of the vulnerability curve at the inflexion point (Figure 1).

Although there is general agreement for the use of centrifuge technique to generate the vulnerability curves for short conduits bearing species (e.g. conifers), the application of both the static and *in the situ* flow centrifuge methods for long vessel bearing species remains a subject of debate, as they are potentially prone to artefact (Torres-Ruiz et al., 2014, 2017; Hacke et al., 2015,

Choat et al., 2016). The high losses of hydraulic conductivity (K_s) induced by the centrifuge technique at relatively mild xylem pressures for




Figure: 1 Example of *Pinus pinaster* vulnerability curve carried out with the cavitron technique showing the percentage loss of xylem conductance as a function of xylem pressure (MPa). The main hydraulic parameter P_{50} (dashed lines) fitted using the Pammenter and Vander Willigen model (1998).

species with long vessels results in the production of exponential vulnerability curves (the so-called ‘r-shaped’ curves). It is considered that these curves are methodological artefact resulting due to the presence of open vessels, i.e. vessels having no end within the stem segment (Choat et al., 2010; Cochard et al., 2013; Wheeler et al., 2013; Torres-Ruiz et al., 2014; Rockwell et al., 2014; Pivovarov et al., 2016). However, these curves have been taken by some other authors to be a reliable estimate of embolism resistance in these species, i.e. the significant embolism within

moderate ranges of water deficit (Sperry et al., 2012; Hacke et al., 2015). These contradictory views generate a confusion reaching conclusive results in the field of plant hydraulics and impedes the progress in understanding the role of hydraulics in plant function (Delzon & Cochard, 2014). A lot of work has been done to clarify whether there is an open-vessel artefact in using the standard centrifuge technique, but even working with the same plant species, contradictions in results are still there. For example, olive (*Olea europaea*) has been observed to produce contrasting vulnerability curves and resistances to embolism according to the methods being used for assessing the degree of embolism. Although it has been observed with high-resolution computed tomography that olive stems are not embolized until a certain threshold in xylem pressure is reached (approximately -3.0 MPa) and that vulnerability curves are sigmoidal (s-shaped curve) for this species (Torres- Ruiz et al., 2014), significant decreases in K_s at relatively high xylem water potentials have been reported by (Hacke et al., 2015), suggesting high embolism vulnerability with r-shaped curve in this species. In a recent study Torres-Ruiz et al. (2017) demonstrated the unreliability of the standard centrifuge technique for the measurement of resistance to embolism in xylem samples with some proportion of open vessels, regardless of the rotor design used. They observed significant decrease in hydraulic conductance at a less negative xylem pressure than the minimum pressure the plant material had experienced previously, indicating that embolism resistance in long vessels bearing speciss, is not as low as previously suggested.

Interspecific variability in embolism resistance

During water stress, hydraulic failure in trees has been reported to be one of the primary causes of forest decline (Anderegg et al., 2015, 2016). The negative pressure inside the water conducting cells get intensified due to drought, making the liquid xylem sap more vulnerable to air-entry and hence embolized (Lens et al., 2013). High levels of embolisms may ultimately lead to desiccation and plant death (Barigah et al., 2013; Urli et al., 2013). Plant resistance to embolism is, therefore, considered a key-parameter in determining the drought tolerance of plants and is estimated on the basis of so-called vulnerability curves (VCs), from which the P_{50} (i.e. the xylem pressure causing 50% loss of hydraulic conductivity) can be estimated (Cochard et al., 2013). P_{50} values are, therefore, good proxies for drought stress tolerance in plants which varies widely among species (Maherali et al., 2004) and have been reported for hundreds of angiosperms and gymnosperms (Delzon et al., 2010; Choat et al., 2012), with vulnerable species rapidly reducing

hydraulic conductance at moderate negative pressure (-2.0 to -0.5 MPa), whereas some species can resist high xylem negative pressure (< -19 MPa) before embolism starts to occur (Larter et al., 2015). This shows a huge range of P_{50} at interspecific level. However, a global convergence in tree vulnerability to drought across ecosystems was found in a study comparing the severity of drought experienced by trees (i.e. the lowest water-potentials observed seasonally) to levels of drought inducing death by hydraulic failure (Choat et al., 2012), indicating that wet and dry forest are equally prone to lethal levels of drought due to limited hydraulic safety margin. Martin-StPaul et al. (2017) reported the existence of an absolute limit by which stomata close to avoid rapid death under drought events. They showed that the water potential triggering stomatal closure and the xylem pressure causing embolism formation were equal for only a small number of species, and the difference between these two traits increased continuously with increasing embolism resistance, indicating the absence of functional coordination between stomatal closure and the onset of embolism formation.

Previous work on hydraulic traits highlights considerable variation in resistance to cavitation among coniferous species from contrasting environments, which seems to be linked to their ecology, with species from xeric regions being more resistant than those from mesic areas (Brodribb and Hill, 1999; Maherali et al., 2004; Choat et al., 2005, 2012; Vinya et al., 2013). It shows that the evolution of cavitation resistance in conifers enables them to adapt in arid environmental conditions. However, little is known about the relationship between cavitation resistance and xylem anatomy in conifers. It could be suggested that the variability of cavitation resistance and species distribution may be explained on the basis of structure of bordered pits.

In recent studies it has been revealed that cavitation events occur at air-sap interfaces within pores between embolized and functional conduits (Choat et al., 2008; Lens et al., 2013). A valve-like structure called torus, present in bordered pits of conifers, is deflected to block the pit aperture, protecting functional tracheids from air entry (Bailey, 1916). Though it provides conifers with increased protection from embolism compared to angiosperms (Maherali et al., 2004; Pittermann et al., 2005; Choat et al., 2012), embolism still occurs due to incomplete sealing of the pit aperture by the torus (Bouche et al., 2014). Embolism resistance is required to balance (xylem safety) with optimizing rates of water transport (xylem efficiency) to the photosynthetic organs of the plants (Tyree & Zimmermann, 2002). In competitive settings, the xylem safety gains are

obvious (i.e. increased survival during drought), whereas increased carbon allocation to leaves relative to sapwood area is allowed by xylem efficiency, resulting in rapid growth and optimum photosynthetic rate (Santiago et al., 2004; Poorter et al., 2010). In drought-tolerant species, the efficiency of wider conductive elements to transport water but their weakness under the mechanical stress imposed by negative xylem pressure, creates an indirect link between safety and efficiency (Sperry et al., 2008). Pit morphology, notably the pit membrane, is influenced by thicker cell walls, thereby increasing xylem embolism resistance (Tyree & Sperry, 1989b; Li et al., 2016). Species ecology and phylogeny strongly determine the final compromise between safety and efficiency. For example, species growing in comparatively wetter environments are more likely to maximize efficiency over safety (Sperry et al., 2006; Choat et al., 2012). At a higher ecological and taxonomic scale, only a few or no evidence is available of a universal and strong relationship between safety and efficiency (Gleason et al., 2016).

In the context of evolutionary biogeography of conifers, pit structure in 15 species of Cupressaceae revealed the hydraulic tradeoffs associated with cavitation resistance at the pit level, and the way they vary within the structural components of the inter tracheid pit. Cavitation resistance was found strongly correlated with the ratio of the torus to pit aperture diameter but did not vary systematically with the porosity of margo. Moreover, it was found that pit conductivity was restricted by pit aperture in more drought-resistant taxa, whereas in species that are more vulnerable to cavitation, pit conductivity was more likely to be regulated by increased margo resistance (Pittermann et al., 2010). The morphological and physiological differences between modern-day members of the Cupressaceae brought by Cenozoic climatic change described by Pittermann et al. (2012) showed that drought-resistant crown clades - the Cupressoid and Callitroid clades - most likely evolved from drought-intolerant Mesozoic ancestors, and that the pattern was consistent with the shifts in post-Eocene paleoclimates. Analyses based on phylogenetic methods suggest that the ancestors of the Cupressaceae conifers were dependent upon humid habitats and the current drought-resistant physiology evolved along with increasing habitat aridity from the Oligocene onward, largely by their capacity to adapt to drought. It was evidenced that within the Cupressaceae, the evolution of drought-resistant xylem was coupled with increased carbon allocation in xylem conduit walls and reduced xylem transport efficiency, and at leaf level, with reduced photosynthetic capacity (Pittermann et al., 2012).

In a recent study based on a time-calibrated phylogeny, Larter et al. (2017) found a wide range in embolism resistance across the *Callitris* clade (P_{50} : -3.8 to 18.8 MPa) which was significantly correlated with water scarcity and tracheid diameter. It was further evidenced that the evolution of extreme xylem embolism resistance and diversification into xeric environments accompanied the radiation of this clade and was driven by increasing aridity of the Australian continent during the last 30 million years.

Intraspecific variability in embolism resistance

While the interspecific differences in functional traits have been investigated by a vast majority of research, variation at intraspecific level is less understood, particularly for hydraulic traits, and is important for understanding and modeling the impacts of climate change on vegetation. The degree of variation of intraspecific traits has major consequences for species sensitivity to climatic shifts and required migration rates to track these shifts. Hydraulic characteristics variability within a single species could have a major impact on population demographics and responses to extreme climatic changes, such as ability of a population to withstand a drought, as well as spatial patterns and thresholds in drought-induced mortality (Martinez-Vilalta et al., 2009; L. D. L. Anderegg et al., 2013).

Intraspecific studies showed no variability of cavitation resistance in branches among populations belonging to different environments, arguing that within a species, this trait is highly confined to the branch level (Sáenz-Romero et al., 2013; Anderegg et al., 2014; Lamy et al., 2014). A comparative study of vulnerability to cavitation between the different organs within a single plant, has been studied to understand drought resistance at the whole-plant level (Bouche et al., 2015). Stomatal conductance and photosynthesis essentially depend on efficient water transport from roots to the leaves. Thus, the way different plant organs cope with cavitation risks has an impact on plant growth and survival. The hydraulic segmentation hypothesis was first proposed by Zimmermann, (1983) suggesting that distal plant organs would be more subject to cavitation events because of the decline in water potential from proximal to distal organs. Later on, this hypothesis was slightly modified by Tyree and Ewers (1991) showing that distal tissues were more vulnerable to cavitation than proximal tissues to prevent cavitation events in the main stem axis. However, significant variability in resistance to cavitation has been obtained between intra-plant studies for a given species and organs (Sperry and Ikeda, 1997; Martínez-Vilalta et al., 2002;

Cochard et al., 2004; Domec et al., 2006; Dalla-Salda et al., 2009; Charra-Vaskou et al., 2012; McCulloh et al., 2014), which might be linked to the different methods applied to measure cavitation resistance across organs.

Cavitation resistance at intra-specific level shows a lack of scaling with environment even for species distributed very widely. For example, climatic aridity had no significant effect on P_{50} whereas other traits such as branch leaf-to-sapwood area ratio were plastic over the wide range of scots pine (*P. sylvestris*) (Martínez-Vilalta et al., 2009). Similarly, Lamy et al., (2011) found less genetic differentiation for cavitation resistance (Q_{st}) than expected under genetic drift (neutral markers, F_{st}), while comparing populations from whole climatic range of maritime pine (*P. pinaster*), with the authors concluding that the trait is likely canalized, suggesting genetic constraints on cavitation resistance variability in this species.

Variation in xylem resistance to embolism (P_{50}) and xylem specific hydraulic conductivity (K_s) among *P. pinaster* populations and families in another study (Corcuera et al., 2011) were found to be mainly controlled by phenotypic plasticity, while wood density was found to be controlled genetically and was not under the influence of environment. Rosner *et al.* (2017) used wood density as a functional trait to highlight its variability within a conifer species, Norway spruce (*Picea abies*) and found that P_{50} of trunk wood of different cambial age from mature and young trees was strongly related to wood density.

There is scarcity of literature on genetic determinism of cavitation resistance, although studies on some species such as Arabidopsis (Tixier et al., 2013), and search for QTLs (Quantative Trait Loci – Lamy, 2012) are being pursued to follow in the future. As cavitation resistance is related to the xylem anatomy, especially at the inter-tracheid pit level, genetic architecture of cell and cell-wall development is probably linked to it, and is, therefore, likely under strong genetic regulation, consequently growth, water-transport efficiency and mechanical support may be affected by random changes. Same gene could be implicated in determining P_{50} and other traits (pleiotropy), and a gene cluster could also interact to establish the same phenotype (polygenic trait) - such complex genetic networks necessarily limit the impact of a single mutation on a single gene, therefore reducing the probability of variation on which natural (and artificial) selection can work. However, it is important to mention that recent results from a widespread angiosperm species,

Fagus sylvatica, are opposite to this trend (Schuldt et al., 2015), showing within-species variation for P_{50} is higher in Angiosperms than in Gymnosperms (Anderegg, 2015).

Intraspecific variability in embolism resistance in crops

Vulnerability of woody species (Angiosperm and gymnosperm) to embolism have been investigated in many studies, but there are only a few studies focusing on crop xylem hydraulics (Sperry et al., 2003). Lens et al. (2016) recently studied hydraulic failure in 26 species of herbaceous angiosperms (mainly European grasses, Poaceae) and found a wide range in P_{50} (the xylem pressure inducing a 50% loss of hydraulic conductivity) from -0.5 to -7.5 MPa, showing that they were generally more resistant to embolism than previously assumed. It was further revealed that embolism formation were not routinely experienced by herb as they had the ability to withstand high negative water potentials without embolism formation during drought stress, but the intraspecific variability of embolism resistance is yet to be investigated in herbs.

The limited number of studies have been performed on the relation between hydraulic failure and the agronomic performance of different crops such as sugarcane (Neufeld et al., 1992), maize (Tyree et al., 1986; Cochard, 2002b; Li et al., 2009), soybean (Sperry, 2000), rice (Stiller et al., 2003) and sunflower (Stiller and Sperry, 2002; Trifilo et al., 2003a, 2003b), but there is lack of information on intraspecific variability in embolism resistance and other hydraulic traits. In the face of enhanced drought risk and increasing drought-affected area globally with the rising global temperature associated with Climatic variability (IPCC, 2014), negative yield trends of important cereal crops, even in high-technology, high-yielding areas across the world, have been recorded (Kang et al., 2009; Ray et al., 2012; van Wart et al., 2013; Martín-Retortillo & Pinilla, 2015). Knowledge of the role of embolism formation in crops and its variability at intraspecific level is, thus essential, if we are to understand the susceptibility of the water transport pathway of crop plants to drought and to cope with the threats posed by changing climatic conditions, which may be crucial for selecting drought resistant, high yielding genotypes to meet the future food demands of increasing population.

Objectives of the thesis

The main objective of this thesis was to investigate the intraspecific variability in embolism resistance and other hydraulic traits in combination with xylem anatomy, and genetics. The work

was categorized in two parts. The first part deals with intraspecific variability in hydraulic traits in sunflower, which is an important oilseed crop in both Europe and South Asia. In Pakistan, the local production of edible oil meets only 32% of the country's domestic requirement while the remaining 68% demand is met through import involving huge expenditure, which accounts for almost 10% of the country's total import bill (Hussain, 2000; Joyo et al., 2016). 75 to 87% area of Pakistan already lies in extremely arid to semi-arid zones and only less than 10% is in humid regions, which are vulnerable to drought under the current climate change scenario, showing a significant increasing trend in maximum annual temperature by $0.014\text{ }^{\circ}\text{C year}^{-1}$ and $0.018\text{ }^{\circ}\text{C year}^{-1}$ in extremely arid and humid regions, respectively (Haider and Adnan, 2014; Adnan et al., 2017). Recent studies show that the sunflower seed yield could reduce by up to 15% by the 2020s and may further decrease up to 25% by the 2050s (Nasim et al., 2016). Characterization of the hydraulic traits of sunflower is, therefore, of utmost important for agricultural scientists, agronomists and hydrologists to plan the growing and breeding programs of the crop in accordance with the regional climate so that varieties/accessions with improved drought tolerance may be selected. The second part aims at finding the intraspecific variability in hydraulic traits and their link with genetics and other physiological traits in Maritime pine, which is the biggest coniferous forest species in Europe and presents a high ecological and economic importance in the Western Mediterranean basin. In France this species covers 6% of the total forest land (88% of the private forest area; CNPPF-IDF, 2015), 5% of the standing volume and more than 26% of the annual wood production (IGN 2016). In Southwestern France, forest productivities are known to be more negatively affected than in Northern France, in response to increase in air and soil drought frequency and intensity because of a more pronounced shift in seasonal rainfall from summer to winter (Loustau et al., 2005). It is, therefore, important to investigate the genetic basis of hydraulic traits and the correlations between key traits that determines the survival of maritime pine during severe periods of drought if such characteristics are to be incorporated into breeding programs.

Part I: Intraspecific variability in embolism resistance in sunflower crop

The main goals of this study were to investigate the embolism resistance of sunflower (*Helianthus annuus* L.), its intraspecific variability among accessions, and its relationships with xylem anatomy. This provided new insights into our understanding of the mechanism of embolism resistance in sunflower and highlighted the potential trade-offs resulting from an increase in

embolism resistance. To properly define a hydraulic strategy, it is essential to account both for the growth/drought resistance trade-off and the hydraulic safety/efficiency trade-off in xylem conduits.

Chapter 1: Intraspecific variation in embolism resistance and stem anatomy across four sunflower (*Helianthus annuus* L.) accessions. (*Accepted in Physiologia Plantarum*)

The major question to be answered was how much sunflower (*Helianthus annuus* L.) accessions, growing under the same environmental conditions, vary in their ability to resist drought-induced embolism? Four sunflower accessions were investigated to assess the vulnerability to embolism using the *in situ* flow centrifuge method for the first time, using a 42 cm rotor to prevent open-vessel artifacts. The potential trade-off between growth traits and embolism resistance and the relationship between embolism resistance and xylem anatomical traits was also determined.

Part II: Intraspecific variability in embolism resistance and its relation with genetic architecture and photosynthetic traits in Maritime pine.

Maintaining the productivity under reduced water availability and survival under severe drought and/or heatwaves are two major challenges, regarding the tree adaptation to water deficit in the face of climate change. Maritime pine (*Pinus pinaster* Ait.) is known to have low intra-specific variability both between *in situ* natural populations and provenances growing in common gardens (Lamy et al. 2011, 2014) but provenance hybrids, that would have increased variability, have never been tested so far (see Plomion et al., 2016 for a review). In view of recent predictions of future climatic scenarios (IPCC, 2014), the results obtained will make it possible to estimate basic genetic parameters for embolism resistance and to evaluate breeding opportunities.

Chapter 2: Dissecting the genetic architecture of resistance in Maritime pine reveals that hydraulic safety is not linked to wood density. (*in preparation*)

Due to the inaccessibility of high throughput phenotyping techniques for large populations until recently, insights into the genetic architecture, i.e., the number of chromosomes, chromosomal location and effect of quantitative trait loci (QTL) on vulnerability to embolism and xylem hydraulic conductivity are still less known. In the context of improving our understanding and in continuation of work already done on the genetic determinisms and evolutionary history of

embolism resistance in Maritime pine, this study aimed at finding the relationship between embolism related traits (hydraulic safety) and wood density related traits by dissecting the genetic architecture of this species. To improve our understanding of the maintenance and evolution of genetic variation in the wild, recent studies showed that evolutionary forces have shaped phenotypic variability for these traits towards a low variability between natural populations. We used hybrids (F2 genotypes) that display more variations as they are potential catalysts of phenotypic evolution and hence diversification both in plants and animals. Moreover, hybridization can be used for transgressive segregation to develop qualitatively or quantitatively novel phenotypes in addition to get specific adaptive traits through lateral gene transfer and introgression.

Chapter 3: Intraspecific variability of the physiological response to water stress in a full-sib Maritime pine family. (*in preparation*)

The objective of this study was to investigate the intra-specific variability within a Maritime pine tree family in response to water availability by performing a multi-trait analyses of leaf and xylem functioning through seasonal measurements of photosystem II fluorescence, gas exchange and stem hydraulic properties. This will be the first step towards selecting the genotypes having the ability to sustain changes in water availability by resisting to drought, while maintaining the same growth rates during pre and post water stress phases.

**Part I: Intraspecific variability in embolism resistance in
sunflower crop**

Chapter 1: Intraspecific variation in embolism resistance and stem anatomy across four sunflower (*Helianthus annuus* L.) accessions. (Accepted in *Physiologia Plantarum*)

1
2
3
4 1 **Intraspecific variation in embolism resistance and stem anatomy across four**
5
6 2 **sunflower (*Helianthus annuus* L.) accessions**
7
8 3

9 4 Hafiz B. Ahmad^a, Frederic Lens^b, Gaelle Capdeville^a, Régis Burlett^a, Laurent J. Lamarque^a and
10 5 Sylvain Delzon^{a*}
11
12 6

13
14 7 ^aBIOGECO, INRA, Univ. Bordeaux, F-33610 Cestas, France

15 8 ^bNaturalis Biodiversity Center, Leiden University, P.O. Box 9517, 2300 RA Leiden, the Netherlands
16
17 9

18
19 10 *Corresponding author, email: sylvain.delzon@u-bordeaux.fr
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

11 Abstract

12 Drought-induced xylem embolism is a key process closely related to plant mortality during extreme
13 drought events. However, this process has been little investigated in crop species to date, despite the
14 observed decline of crop productivity under extreme drought conditions. Interspecific variation in
15 hydraulic traits has frequently been reported, but less is known about intraspecific variation in crops. We
16 assessed the intraspecific variability of embolism resistance in four sunflower (*Helianthus annuus* L)
17 accessions grown in well-watered conditions. Vulnerability to embolism was determined by the *in situ*
18 flow centrifuge method (cavitron), and possible trade-offs between xylem safety, xylem efficiency and
19 growth were assessed. The relationship between stem anatomy and hydraulic traits was also investigated.
20 Mean P_{50} was -3 MPa, but considerable variation was observed between accessions, with values ranging
21 between -2.67 and -3.22 MPa. Embolism resistance was negatively related to growth and positively
22 related to xylem-specific hydraulic conductivity. There is, therefore, a trade-off between hydraulic safety
23 and growth but not between hydraulic safety and efficiency. Finally, we found that a few anatomical traits,
24 such as vessel density and the area of the vessel lumen relative to that of the secondary xylem, were
25 related to embolism resistance, whereas stem tissue lignification was not. Further investigations are now
26 required to investigate the link between the observed variability of embolism resistance and yield, to
27 facilitate the identification of breeding strategies to improve yields in an increasingly arid world.

29 Abbreviations

30 PLC, percentage loss of hydraulic conductance; VCs, vulnerability curves.

31 Introduction

32 Crop productivity and biodiversity are currently undergoing major transformations due to climate change,
33 with increases in global temperature and atmospheric CO₂ concentration and changes in land use
34 (Vitousek et al. 1997, Parmesan and Yohe 2003). Global surface temperature increased by about 0.8°C
35 from 1861 to 2005, and further increases are predicted for the future (IPCC 2007, Dai 2013). The
36 predicted increases in temperature and the prevalence of extreme climatic events (Sterl et al. 2008, Wigley
37 2009) have been accompanied by increases in precipitation during the winter and fall and decreases in
38 precipitation during the summer, resulting in an intensification of the summer drought period in Western
39 Europe (Beniston et al. 2007, van Oldenborgh et al. 2009). These changes will undoubtedly have a major
40 impact on crop production (Boyer 1982, Hussain et al. 2012), with adverse effects on all stages of plant
41 growth and development (Jensen and Mogenson 1984), particularly for summer crops, such as maize,
42 soybean, sugar beet and sunflower. Ciais *et al.* (2005) reported a pronounced decrease in net primary
43 productivity (NPP) for maize (*Zea mays* L.) in agricultural regions of Europe affected by the 2003
44 heatwave and drought, with a record decrease of 36% in Italy.

45 World food security is mostly under threat from drought (Somerville and Briscoe 2001, Farooq et al.
46 2009), which affects crop plants immediately after their germination, resulting in poor seedling
47 establishment (Harris et al. 2002, Kaya et al. 2006) and, ultimately, low yields (Jaleel et al. 2007), due to
48 low rates of absorption of photosynthetically active radiation, poor radiation-use efficiency, and a low
49 harvest index (Earl and Davis 2003). Crop sensitivity to water deficit varies with phenological stage and
50 is, thus, crop-dependent. For example, water deficit has been shown to affect vegetative growth in
51 soybean, flowering and boll formation in cotton and the vegetative and yield formation stages of
52 sunflower and sugar beet (Kirda 2002). Critically low levels of vegetative growth and poor plant
53 development due to water deficit have also been reported in rice (Tripathy et al. 2000, Manickavelu et al.
54 2006). Many studies have demonstrated effects of drought on crop phenology and gas exchanges, but little
55 is known about the breakdown of the water transport due to embolism under drought conditions, even
56 though this process has been shown to be the major cause of drought-related death in perennial plants
57 (Urli et al. 2013, Anderegg et al. 2015).

58 The water transport system of plants consists of a complicated network of xylem conduits through
59 which water moves under negative pressure (tension), due to the pull of transpiration at the leaves,
60 transmitted via a continuous column of water down to the roots, where the resulting decrease in pressure
61 leads to the absorption of water from the soil (Tyree & Zimmermann 2002, Wheeler & Stroock 2008,
62 Trifilo et al. 2014). Water columns in plants must consist entirely of liquid and be free of air bubbles,
63 despite being under negative pressure. This metastable liquid state is prone to cavitation, a phase change
64 from liquid water to water vapor, resulting in the formation of gas bubbles (air embolism) that prevent

1
2
3 65 water from flowing through xylem conduits, thereby reducing the hydraulic conductivity of the xylem
4 66 from the soil to the foliage, where water is required for the maintenance of optimal cell hydration levels
5 67 (Tyree and Sperry 1989, Salleo et al. 2000, Cochard 2006). During prolonged and severe droughts, the
6 68 tension of the xylem sap increases the probability of embolism formation in the xylem, reaching
7 69 potentially lethal levels causing desiccation and mortality (Davis et al. 2002, Brodribb and Cochard 2009,
8 70 Hoffmann et al. 2011, Choat 2013, Urli et al. 2013).

9
10
11 71 Water stress-induced xylem embolism is one of the major causes of plant mortality during extreme
12 72 drought events (Anderegg et al. 2015). Many studies have focused on the vulnerability of woody species
13 73 to embolism, but there have been far fewer investigations of crop xylem hydraulics (Sperry et al. 2003).
14 74 Nevertheless, the limited number of studies performed to date on sunflower (Stiller and Sperry 2002), rice
15 75 (Stiller et al. 2003), maize (Tyree et al. 1986, Cochard 2002a, Li et al. 2009), sugarcane (Neufeld et al.
16 76 1992) and soybean (Sperry 2000) have indicated that hydraulic failure also has a major effect on crops,
17 77 within the physiological range of xylem pressure. Knowledge of the role of embolism formation in crops
18 78 is thus essential if we are to understand the susceptibility of the water transport pathway of plants to
19 79 drought.

20
21 80 There have been few assessments of vulnerability to embolism in herbaceous plants (Lens et al. 2016).
22 81 Lens et al. (2016) showed that herbaceous plants are generally more resistant to embolism than previously
23 82 thought and that they do not routinely experience xylem embolism. Major differences in embolism
24 83 resistance between herbaceous species have been reported, based on estimates from vulnerability curves
25 84 (VCs), on which P_{50} — the xylem pressure inducing a 50% loss of hydraulic conductivity — ranged from
26 85 -0.5 to -7.5 MPa, but nothing is known about the intraspecific variability of embolism resistance.

27
28 86 We focus here on sunflower (*Helianthus annuus*) a major high-yield oilseed crop (Lawal et al. 2011,
29 87 Bera et al. 2014) cultivated over a broad geographical area with diverse environmental conditions
30 88 worldwide (Liu and Baird 2003, Lopez-Valdez et al. 2011). Sunflower is generally considered to be a
31 89 drought-resistant crop (Unger 1990, Connor and Hall 1997). However, it has been reported to be affected
32 90 by extreme and frequent periods of water stress in southern Europe, where it is cultivated in low-rainfall
33 91 areas in which the soil is shallow and irrigation facilities are not available (Casadebaig et al. 2008). Water
34 92 deficit affecting the vegetative and flowering stages strongly has been reported to cause a 29% decrease in
35 93 yield (Velue and Palanisami 2001). The susceptibility of sunflower-producing areas to drought may
36 94 increase considerably in the near future, if the predicted climate changes occur (Dufresne et al. 2006).

37
38 95 The objective of this study was to investigate the variability of embolism resistance between sunflower
39 96 (*Helianthus annuus*) accessions grown in the same environmental conditions. We assessed vulnerability to
40 97 embolism in four sunflower accessions by the *in situ* flow centrifuge method, using a 42 cm rotor to
41 98 prevent open-vessel artifacts (Cochard et al. 2013, Pivovarovoff et al. 2016). We then investigated (i) the

99 potential trade-off between growth traits and embolism resistance and (ii) the link between embolism
100 resistance and xylem anatomical traits.

101

102 **Materials and methods**

103 **Plant material and experimental design**

104 The study was performed on four sunflower (*Helianthus annuus*) accessions: a rustic accession capable of
105 growing on shallow soil (Melody), two early-sown accessions (ES_Biba and ES_Ethic) and one late-sown
106 accession (LG_5660). Seeds for all four accessions were provided by free of charge by the *Centre*
107 *Technique Interprofessionnel des Oléagineux Métropolitains* (CETIOM), France. Plants were grown in a
108 randomized complete block design (RCBD) with four blocks, including six plants per accession per block,
109 giving a total of 96 plants ((6*4)*4). Seeds were sown on March 14, 2014, in pots filled with Peltracom
110 substrate containing peat, clay, plant fibers, vulcanic stones, sand and compost, with 14-16-18 kg m⁻³ NPK
111 fertilizer. The pots were placed in a greenhouse under full light, in non-limiting growth conditions, at the
112 University of Bordeaux, France. Three seeds were initially sown in each pot at the start of the experiment.
113 After germination (about 10 to 12 days after sowing), when the seedlings had reached a height of about 10
114 to 15 cm, we removed two of the seedlings, leaving only one healthy seedling per pot. An automatic
115 irrigation system fitted with an electronic water timer (Dual logic, CLABER, Italy) was used to irrigate
116 the pots. It was set to irrigate all the pots simultaneously, for 10 minutes every 12 hours, to ensure that all
117 the plants remained well-watered and free of water stress and embolism throughout the growing period.

118

119 **Sample preparation for the assessment of embolism resistance**

120 Plants were harvested about 12-15 weeks after sowing (i.e. between June 19 and July 9, 2014), when they
121 had reached a mean height of 0.85 m. Plant height (H, m) and stem diameter (D mm) were measured with
122 a tape measure and an electronic Vernier scale, respectively, and the plants were then cut for hydraulic
123 measurements. Five to seven plants with intact flowers from the different accessions were selected at
124 random each day and cut 1 cm above the soil at 8:00 am. All the leaves were removed from the plants just
125 before cutting, to reduce loss of water due to transpiration. The plants were immediately wrapped in wet
126 cotton cloth and placed in a plastic bag to minimize dehydration during their transport to the laboratory.
127 They were then cut, under water, to a standard length of 42 cm, and both ends were trimmed with a fresh
128 razor blade. On average, we were able to take measurements for 16 stems per accession by the end of the
129 experiment.

130

131 **Xylem vulnerability curves**

132 Xylem vulnerability to embolism was assessed with the *in situ* flow-centrifuge technique (Cavitron), in

1
2
3 133 which the percentage loss of hydraulic conductivity relative to xylem under negative pressure is
4 134 determined (Cochard 2002b, Cochard et al. 2005). All measurements were performed at the high-
5 135 throughput phenotyping platform for hydraulic traits (CaviPlace, University of Bordeaux, Talence,
6 136 France). Centrifugal force was used to establish negative pressure in the xylem and to provoke water
7 137 stress-induced cavitation, in a 42 cm-wide custom-built honeycomb aluminum rotor (DGMeca, Gradignan,
8 138 France) mounted on a temperature-controlled high-speed centrifuge (J6-MI, Beckman Coulter, USA). The
9 139 ends of the sample were placed in 25 mm OD polycarbonate centrifuge tubes (38 mL, Beckman Coulter,
10 140 USA) with holes located 42 and 14 mm from the extremities for the upstream and downstream reservoirs,
11 141 respectively. Samples were then secured in a slit across the center of the rotor, with the lid screwed down
12 142 tightly to hold the sample in place. A solution of 10 mM KCl and 1 mM CaCl₂ in ultrapure deionized
13 143 water was used as the reference ionic solution. The rotor was first spun at low xylem pressure ($P_x = -0.8$
14 144 MPa), corresponding to a rotation speed of 2087 rpm (revolutions per minute). The rotation speed of the
15 145 centrifuge was then gradually increased by -0.3 or -0.5 MPa, to expose samples to lower xylem pressures.
16 146 Rotor velocity was monitored with a 10 rpm-resolution electronic tachymeter (A2108-LSR 232, Compact
17 147 Inst, Bolton, UK) and xylem pressure was adjusted to about ± 0.02 MPa. Hydraulic conductances (K_i , m²
18 148 MPa⁻¹ s⁻¹) were determined at every rotation by measuring the displacement speed of the air-water
19 149 meniscus from the upstream to the downstream extremity of the sample, according to the equations of
20 150 Wang et al. (2014). These measurements were performed with a calibrated CCD camera (Scout sca640,
21 151 Basler, Germany) coupled to custom-written software (Cavisoft version 4.0, BIOGECO, University of
22 152 Bordeaux). Hydraulic conductances were measured three times per rotation. The mean values were used
23 153 to determine the percentage loss of hydraulic conductance (PLC) at each pressure, as follows:

$$154 \quad PLC = 100 \left(1 - \frac{K_i}{K_{max}} \right)$$

155 , where K_{max} is the maximum hydraulic conductance measured at low speed, i.e. at very high xylem
156 pressure. Vulnerability curves (VCs), corresponding to the percentage loss of xylem conductance as a
157 function of xylem pressure (MPa), were determined for each sample as follows (Pammenter and Vander
158 Willigen 1998):

$$159 \quad PLC = \frac{100}{\left[1 + \exp\left(\frac{s}{25} (\psi - P_{50}) \right) \right]}$$

160 , where P_{50} (MPa) is the xylem pressure inducing a 50% loss of conductance and S (% MPa⁻¹) is the slope
161 of the vulnerability curve at the inflexion point. The xylem-specific hydraulic conductivity (K_s , m² MPa⁻¹
162 s⁻¹) was calculated by dividing the maximum hydraulic conductivity measured at low speed (K_{max}) by the
163 xylem area of the sample. The xylem pressures at which 12 and 88% conductivity were lost (P_{12} and P_{88} ,
164 MPa, respectively) were calculated as follows:

$$165 \quad P_{12} = 2/(s/25) + P_{50}$$

166 and

$$P_{88} = -2/(s/25) + P_{50}$$

168 The mean embolism vulnerability values were calculated from the data for 15 to 17 samples per accession.

169

170 **Anatomical observations**

171 Anatomical observations were carried out on the samples used for hydraulic measurements. Three stems
 172 per accession were selected at random. A 2.5-3.0 cm-long segment was cut from the central portion of
 173 each sample, and the 12 segments obtained in this way were stored in jars filled with 60% ethanol. The
 174 samples were then taken to the Naturalis Biodiversity Center (Leiden, the Netherlands) for sectioning.
 175 Three to four transverse stem sections, each about 20-25 μm thick, were cut from each sample with a
 176 sledge microtome (Reichert, Germany), for light microscopy (DM2500 microscope, Leica, Germany). The
 177 sections were prepared according to the standardized protocol described by Lens et al. (2005). Briefly,
 178 sections were treated with household bleach for 1 minute and rinsed at least three times with distilled
 179 water. They were then stained with safranin-Alcian blue (consisting of two parts 1% safranin in 50%
 180 ethanol and one part 1% Alcian blue in H_2O) for 15 seconds, and then subjected to dehydration by
 181 successive steps of at least one minute in 50%, 70% and 96% ethanol. Sections were finally treated with a
 182 1:1 mixture of ethanol 96%-Histoclear before complete immersion in Histoclear. The sections were
 183 mounted in Euparal mounting medium and dried in an oven at 60°C for at least three weeks. The slides
 184 were then scanned with a Hamamatsu NANOZOOMER 2.0HT (Bordeaux Imaging Center, University of
 185 Bordeaux, France). Images of complete cross sections were taken at $\times 10$ magnification and analyzed with
 186 Adobe Photoshop CS2 (Version 9.0, Adobe Systems Inc., San Jose, CA, USA) and ImageJ (Version
 187 1.44p) software, using the particle analysis function.

188 For all subsequent calculations, the complete stem cross section with pith and bark was analyzed. The
 189 parameters measured included: total stem cross section area (A_{stem} , μm^2); total xylem area (A_{Tx} , μm^2);
 190 primary xylem area (A_{Px} , μm^2) and secondary xylem area (A_{Sx} , μm^2); pith area (A_{pith} , μm^2); proportion of
 191 pith area per unit stem area ($P_{p s}$); area of the cellular part of the stem (A_{cp} , μm^2) calculated by subtracting
 192 A_{pith} from A_{stem} ; lignified area (A_{lig} , μm^2) calculated by adding total xylem area (A_{Tx}) and fiber cap area
 193 (A_{fcap} , μm^2); proportion of lignified area per stem area ($P_{lig s}$) obtained dividing A_{lig} by A_{stem} ; proportion of
 194 lignified area relative to the cellular part of the stem ($P_{lig cp}$) calculated by dividing A_{lig} by A_{cp} ; fiber cell
 195 wall area (A_{fcw} , μm^2) in the secondary xylem measured by subtracting fiber lumen area (A_{flumen} , μm^2) from
 196 fiber cell area (A_{fcell} , μm^2); proportion of cell wall per fiber cell ($P_{cw f}$) obtained by dividing (A_{fcw}) by
 197 (A_{fcell}); total fiber wall area in the lignified area ($A_{fcw in A_{lig}}$, μm^2) calculated by multiplying $P_{cw f}$ by A_{lig} ;
 198 proportion of fiber wall in the lignified area per stem area ($P_{fcw in lig s}$) measured by dividing $A_{fcw in A_{lig}}$ by
 199 A_{stem} . We also calculated the following parameters for both primary and secondary xylem: vessel density

1
2
3 200 (VD); vessel lumen area (A_v); cumulative vessel lumen area (A_{cv}); relative vessel lumen area (A_{rv}) obtained
4
5 201 by dividing cumulative vessel lumen area by the corresponding xylem area; thickness-to-span ratio of
6
7 202 vessels (TD^{-1}) obtained by dividing double intervessel wall thickness (T_{vw}) by the maximum diameter of
8
9 203 the vessel (D_{max}); equivalent circle diameter (D) and hydraulically weighted vessel diameter (D_h),
10
11 204 respectively, calculated as $D = (4A/\pi)^{1/2}$ where A is vessel cross sectional surface area (μm^2), and $D_h = \sum$
12
13 205 $D^5/\sum D^4$ (Scholz et al. 2013). A list of all the measured traits, their symbols and units is provided in Table
14
15 206 1.

16 207

17 208 **Statistical analyses**

18 209 The differences in hydraulic traits (P_{50} , P_{12} , P_{88} , S , K_s) between accessions were assessed by one-way
19
20 210 analysis of variance (ANOVA). Correlations between variables were evaluated by calculating the Pearson
21
22 211 correlation coefficient (r), and the relationship between variables was considered to be significant if $P =$
23
24 212 < 0.05 . Statistical analyses of the data were performed with SAS software (version 9.4, SAS Institute,
25
26 213 Cary, NC, USA).

27 214

28 215 **Results**

29 216 **Differentiation between accessions**

30
31 217 Stem diameter did not differ significantly between the four sunflower accessions ($F = 1.69$, $P = 0.1781$)
32
33 218 but significant differences in height were observed: Melody and LG_5660 were significantly taller than
34
35 219 ES_Ethic and ES_Biba ($F = 20.96$, $P = < 0.0001$; Table 2). The vulnerability curves of the four accessions
36
37 220 followed a similar sigmoidal shape (Fig. 1). Embolism resistance (P_{50}) differed significantly between
38
39 221 accessions ($F = 45.59$, $P = < 0.0001$), with LG_5660 and ES_Ethic the most vulnerable and the most
40
41 222 resistant accession to embolism, respectively (Table 2, Fig. 2). ES_Ethic also differed significantly from
42
43 223 the other accessions in terms of P_{12} ($F = 6.27$, $P = 0.0009$), whereas mean P_{88} , which differed significantly
44
45 224 between accessions ($F = 33.95$, $P = < 0.0001$), was lowest for Melody and highest for LG_5660,
46
47 225 respectively (Table 2). We also found significant differences in S ($F = 6.66$, $P = 0.0006$) and K_s ($F = 3.15$,
48
49 226 $P = 0.0313$; Table 2) between accessions.

50 227

51 228 **Correlation between hydraulic, growth and anatomical traits**

52
53 229 The Pearson correlation analysis revealed several relationships between growth, hydraulic and anatomical
54
55 230 traits. Height was positively correlated with P_{50} ($r = 0.42$; $P = 0.0027$; Table 3, Fig. 3A) and P_{12} ($r = 0.54$;
56
57 231 $P = < 0.0001$), indicating that the vulnerability of the xylem to embolism increased with height. Height
58
59 232 was negatively associated with S ($r = -0.35$; $P = 0.0128$), indicating that embolism occurred more rapidly
60
233 in faster growing individuals. It was also negatively associated with xylem-specific hydraulic conductivity

234 (K_s) ($r = -0.56$; $P = <0.0001$; Fig. 3B), indicating a lower xylem efficiency in faster growing individuals.
 235 A negative correlation was also observed between P_{50} and K_s ($r = -0.30$; $P = 0.0174$; Fig. 3C), suggesting
 236 a lack of trade-off between xylem safety and efficiency. No correlation was found between stem diameter
 237 (D, mm) and hydraulic traits, except for K_s ($r = -0.47$; $P = 0.0007$; Table 3).

238 Hydraulic traits were significantly correlated with seven of the 34 anatomical traits measured (Tables 1
 239 and S1). P_{12} and P_{50} were strongly and negatively correlated with VD_{Tx} and VD_{Sx} (Table 3, Fig. 4A, B),
 240 indicating that stems with a higher vessel density in the total xylem area and the secondary xylem area are
 241 more resistant to both the entry of air into the xylem and a substantial loss of conductance. P_{50} was also
 242 negatively correlated with A_{rvSx} ($r = -0.60$; $P = 0.0485$; Table 3, Fig. 4C), whereas P_{12} was negatively
 243 correlated with inter-vessel double-wall thickness in the primary xylem (T_{vwPx} , $r = -0.60$; $P = 0.0490$) and
 244 the thickness-to-span ratio of vessels in the primary xylem (TD^J_{Px} , $r = -0.63$; $P = 0.0385$; Table 3). P_{50}
 245 was not related to the lignified area (A_{lig}) or to the proportion of lignified area per unit stem area ($P_{lig s}$;
 246 Table SA, Fig. 5). No correlation was detected between P_{88} and anatomical traits (Table 3 and S1).
 247 Xylem-specific hydraulic conductivity (K_s) was negatively correlated with A_{vPx} ($r = -0.63$; $P = 0.0380$) and
 248 D_{Px} ($r = -0.64$; $P = 0.0319$; Table 3) but this variable was not correlated with any other anatomical trait
 249 (Table S1).

250

251 Discussion

252 The mean P_{50} value of -2.99 ± 0.15 MPa found here is similar to that reported by Stiller and Sperry (2002)
 253 for well-watered *Helianthus annuus* (P_{50} , -3.0 ± 0.1 MPa). Sunflower is, thus, moderately vulnerable to
 254 embolism relative to other herbaceous and woody species, which have P_{50} values ranging from -0.5 to -7.5
 255 MPa (Lens et al. 2016) and from -0.5 to -18.8 MPa (Delzon et al. 2010, Choat et al. 2012, Bouche et al.
 256 2014, Larter et al. 2015), respectively. We found significant intraspecific differences in vulnerability to
 257 xylem embolism in sunflower, with the accessions at the two extremes of the scale differing in P_{50} by
 258 about 0.55 MPa. Vulnerability to embolism (P_{50}) was also positively related to growth and negatively
 259 related to xylem-specific hydraulic conductivity, highlighting a trade-off between embolism resistance and
 260 growth, but not between xylem safety and efficiency. Finally, we found that various anatomical traits,
 261 such as vessel density, were related to embolism resistance, whereas the degree of stem tissue lignification
 262 was not.

263

264 Intraspecific variability of embolism resistance

265 Despite the critical role of xylem embolism resistance in plant survival during drought events, only one
 266 previous study has reported variation in this trait (from -0.8 to -3 MPa) in sunflower (Stiller and Sperry
 267 2002). However, the authors used a single genotype subjected to drought and rewatering cycles. Our study

1
2
3 268 is thus the first to investigate intraspecific variation in xylem embolism resistance in sunflower. We found
4
5 269 a 0.55 MPa difference in P_{50} between the most resistant sunflower accession, the early-sown ES_Ethic,
6
7 270 and the most vulnerable accession, the late-sown LG_5660. This finding highlights the possibility of
8
9 271 selecting specific sunflower accessions on the basis of their greater resistance to xylem embolism, and
10
11 272 therefore to drought, opening up opportunities for the development of new varieties better adapted to the
12
13 273 drier environmental conditions of the future. Studies investigating the intraspecific variability of embolism
14
15 274 resistance in other crops have yielded contrasting results. Neufeld et al. (1992) and Li et al. (2009)
16
17 275 highlighted genetic differences in P_{50} in sugarcane clones (from -0.83 to -1.36 MPa) and maize hybrid
18
19 276 stems (-1.56 to -1.78 MPa), respectively. By contrast, Cochard (2002b) and Stiller et al. (2003) found no
20
21 277 such differences for maize hybrids, and for comparisons of upland and lowland rice varieties, respectively.
22
23 278 Mixed results for the intraspecific variation of xylem embolism resistance have also been reported for
24
25 279 woody plants. Moderate to low levels of intraspecific variation have been reported for P_{50} (-2.21 ± 0.19 to
26
27 280 -2.97 ± 0.12 MPa) in poplar (*Populus* sp) demes (Hajek et al. 2014), whereas no significant differences in
28
29 281 P_{50} were observed in European beech (*Fagus sylvatica* L.) populations (Hajek et al. 2016). Similarly
30
31 282 diverse observations have been reported for conifers (Lamy et al. 2011, Sáenz-Romero et al. 2013, Lamy
32
33 283 et al. 2014). These findings suggest that the resistance to embolism may often be linked to uniform
34
35 284 evolutionary selection and canalization (Lamy et al. 2011, 2014).

32 286 **Trade-off between growth traits and embolism resistance**

34 287 It has often been suggested that increases in resistance to xylem embolism are achieved at the expense of
35
36 288 slower plant growth, due to conflicts in the allocation of carbon to the construction of denser wood with
37
38 289 thicker cell walls (Hacke et al. 2001) or the construction of foliar and axial tissues to increase canopy
39
40 290 carbon gains and growth rate (Wikberg and Ögren, 2004, Ducrey et al. 2008). Our findings suggest that
41
42 291 height is a key factor governing embolism resistance in sunflower accessions. We found that shorter plants
43
44 292 had greater embolism resistance. Conflicting results have been published concerning the possible
45
46 293 existence of such a trade-off between P_{50} and growth-related traits. Cochard et al. (2007) found a close
47
48 294 relationship between xylem vulnerability and productivity in poplar and willow clones. However, Fichot
49
50 295 et al. (2010), for instance, observed that embolism-resistant genotypes of poplar grew more rapidly than
51
52 296 vulnerable genotypes. Similarly, Sterck et al. (2012) found that embolism resistance had a positive effect
53
54 297 on branch growth in Scots pine, whereas Hajek et al. (2014) found no relationship between vulnerability
55
56 298 to embolism and growth rate in poplar demes. Several recent studies have also failed to detect a trade-off
57
58 299 between vulnerability to embolism and growth-related traits (Guet et al. 2015, Hajek et al. 2016). The
59
60 300 relationship between embolism resistance and growth therefore remains a matter of debate.

301 Xylem-specific hydraulic conductivity (K_s) was negatively correlated with both height and stem

1
2
3 302 diameter, consistent with a trade-off between hydraulic efficiency and growth. However, no direct effect
4
5 303 of xylem specific hydraulic conductivity was observed on growth in Scots pine (Sterck et al. 2012). By
6
7 304 contrast, Hajek et al. (2014) found a positive relationship between K_s and growth rate in poplar, suggesting
8
9 305 that water conductance capacity is a useful growth-determining factor. Similarly, Schuldt et al. (2015)
10
11 306 reported a significant positive relationship between K_s and growth in European beech, showing that fast-
12
13 307 growing branches had a more efficient hydraulic system than slower growing branches. This conflicting
14
15 308 results suggest that there is still a lack of consensus concerning possible trade-offs between K_s and growth
16
17 309 traits.
18

19
20 310
21 311 **Relationship between hydraulic traits and anatomy**
22
23 312 Vessel density in the total xylem (VD_{Tx}) and in the secondary xylem (VD_{Sx}) strongly influenced both the
24
25 313 point of air entry during embolism formation (P_{12}) and the xylem pressure inducing a 50% loss of
26
27 314 conductance (P_{50}). These results indicate that embolism resistance in sunflower is increased by the
28
29 315 production of more vessels per unit xylem area. This finding is also supported by the close relationship
30
31 316 between P_{50} and vessel lumen area relative to secondary xylem area (A_{rvSx}). A similar relationship between
32
33 317 vessel density (VD) and P_{12} was reported by Schuldt et al. (2015) for European beech; however, Hajek et
34
35 318 al. (2014) found no close relationship between P_{50} and relative vessel lumen area in poplar.

36
37 319 We found no relationship between P_{50} and greater stem tissue lignification. The development of
38
39 320 embolism-resistant stems does not therefore involve tissue lignification. Several studies have reported a
40
41 321 link between greater embolism resistance and higher levels of lignification in herbaceous (Lens et al.
42
43 322 2013, Tixier et al. 2013, Lens et al. 2016) as (Awad et al. 2012) woody plants, but an increase in
44
45 323 lignification is not always required to achieve higher levels of embolism resistance (Watkins et al. 2010,
46
47 324 Pittermann et al. 2011).

48
49 325 Strong correlations between P_{12} and both intervessel double-wall thickness in primary xylem (T_{vwPx})
50
51 326 and the thickness-to-span ratio of vessels in the primary xylem (TD^I_{Px}) suggest that these two traits are
52
53 327 important for the onset of embolism formation. Greater wall thickness and thickness-to-span ratios result
54
55 328 in a lower xylem air entry pressure (i.e. resistant accessions have thicker tracheid walls relative to lumen
56
57 329 area). This association between increasing cavitation resistance and increasing thickness-to-span ratio has
58
59 330 also been reported in conifers (Bouche *et al.* 2014; Hacke et al. 2001) and in *Acer* species (Chave et al.
60
331 2009, Lens et al. 2011). A higher thickness-to-span ratio is thought to strengthen the vessel walls against
332 implosion, higher embolism resistance being associated with a lower negative sap pressure.

333 The negative relationship between K_s and P_{50} observed here shows that there is no trade-off between
334 xylem-specific hydraulic conductivity and embolism resistance. Plants with higher embolism resistance
335 also transport water more efficiently. This finding contrasts with that reported by Lens et al. (2011), who

1
2
3 336 found that higher levels of embolism resistance were strongly associated with lower stem-specific (K_{Sa})
4 and xylem-specific (K_{Xa}) conductivities. Ideally, plants should be able to maintain both the efficient
5 337 conductivity and safety of the hydraulic system. However, this is not the case in natural conditions, and
6 338 little or no support for a safety-efficiency trade-off has been obtained across species (Gleason et al. 2015).
7 339 A few studies have evaluated this trade-off at the intraspecific level, and found either no support for the
8 340 existence of a trade-off (Martínez-Vilalta et al. 2009, Schuldt et al. 2015) or, as here, an association
9 341 between greater conductivity and lower embolism resistance (Corcuera et al. 2011). The water-conducting
10 342 efficiency of vessels depends on vessel diameter, with wider vessels more efficient than narrower ones
11 343 (Sperry et al. 2006). Our finding that K_s is negatively correlated with vessel diameter in primary xylem
12 344 (D_{Px}) and vessel lumen area in the primary xylem (A_{vPx}) is, therefore, surprising. It may reflect the
13 345 limitation of water flow through vascular tissue mostly by the pit resistivity of the conduit end walls
14 346 (about 56% according to Sperry et al. 2006). Indeed, increases in lumen conductivity are not necessarily
15 347 associated with increases in total conduit conductivity. However, further studies are required to determine
16 348 how pits can be efficient for water transport, leading to low values of pit resistivity, whilst also limiting
17 349 air-seeding under high xylem tension. This contrasts with the findings of Lens et al. (2011) and Hajek et
18 350 al. (2014), who found a positive relationship between K_s and relative vessel lumen area and vessel
19 351 diameter, indicating that hydraulic conductivity was determined by vessel size. This view is supported by
20 352 other studies carried out on woody species (Zwieniecki et al. 2001, Sperry et al. 2005, Sperry et al. 2008,
21 353 Schuldt et al. 2015).
22 354
23 355

356 **Conclusion**

357 Our findings demonstrate the existence, in sunflower, of intraspecific variation in resistance to drought-
358 induced xylem embolism. It may, therefore, be possible to select drought-resistant accessions/genotypes,
359 which will be crucial for future farming, particularly in areas prone to drought. There was no trade-off
360 between hydraulic efficiency (xylem-specific hydraulic conductivity) and xylem safety (embolism
361 resistance), but we did find trade-offs between height and hydraulic safety and height and hydraulic
362 efficiency. Future studies should investigate (i) the variability of embolism resistance across a wider range
363 of accessions and (ii) the possible existence of a trade-off between embolism resistance and yield
364 potential. Indeed, high yield potential, which is the main target of most crop breeding programs, may not
365 be compatible with higher embolism resistance.
366

367 **Author contributions**

368 S.D. and H.B.A. designed the study. H.B.A. and G.C. performed the greenhouse experiment. G.C. and
369 R.B. provided assistance for hydraulic measurements. S.D., F.L. and H.B.A. carried out the anatomical

1
2
3 370 observations. S.D. performed the statistical analyses. H.B.A. and L.J.L. wrote the first version of the
4 371 manuscript, which was reviewed and revised by all the authors.

5
6 372

7 8 373 **Acknowledgments**

9
10 374 This study was carried out with financial support from the Cluster of Excellence COTE (ANR-10-LABX-
11 375 45, within the Water Stress and Vivaldi projects) and the 'Investments for the Future' program (ANR-10-
12 376 EQPX-16, XYLOFOREST) of the French National Agency for Research. We would like to thank all the
13 377 contributors from SIGDU-University of Bordeaux for their assistance with the greenhouse experiment.
14 378 L.J.L. holds a postdoctoral fellowship from IdEx Bordeaux.

15 379

16 380 **References**

- 17
18
19
20
21 381 Anderegg WRL, Flint A, Huang CY, Flint L, Berry JA, Davis FW, Sperry JS Field CB (2015) Tree
22 382 mortality predicted from drought-induced vascular damage. *Nat Geosci* 8: 367-371
- 23 383 Awad H, Herbette S, Brunel N, Tixier A, Pilate G, Cochard H, Badel E (2012) No trade-off between
24 384 hydraulic and mechanical properties in several transgenic poplars modified for lignins metabolism.
25 385 *Environ Exp Bot* 77: 85-195
- 26 386 Beniston M, Stephenson DB, Christensen OB, Ferro CAT, Frei C, Goyette S, Halsnaes K, Holt T, Jylha
27 387 K, Koffi B (2007) Future extreme events in European climate: an exploration of regional climate
28 388 model projections. *Climatic Change* 81: 71-95
- 29 389 Bera AK, Pramanik K, Mandal B (2014) Response of biofertilizers and homo-brassinolide on growth,
30 390 yield and oil content of sunflower (*Helianthus annuus L.*). *Afr J Agr Res* 9: 3494-3503
- 31 391 Bouche PS, Larter M, Domec JC, Burlett R, Gasson P, Jansen S, Delzon S (2014) A broad survey of
32 392 hydraulic and mechanical safety in the xylem of conifers. *J Exp Bot* 65: 4419-4431
- 33 393 Boyer JS (1982) Plant Productivity and Environment. *Sci* 218: 443-448
- 34 394 Brodribb TJ, Cochard H (2009) Hydraulic failure defines the recovery and point of death in water-stressed
35 395 conifers. *Plant Physiol* 149: 575-584
- 36 396 Casadebaig P, Debaeke P, Lecoœur J (2008) Thresholds for leaf expansion and transpiration response to
37 397 soil water deficit in a range of sunflower genotypes. *Eur J Agron* 28: 646-654
- 38 398 Chave J, Coomes D, Jansen S, Lewis SL, Swenson NG, Zanne AE (2009) Towards a worldwide wood
39 399 economics spectrum. *Ecol Lett* 12: 351-366
- 40 400 Choat B, Jansen S, Brodribb TJ, Cochard H, Delzon S, Bhaskar R, Bucci SJ, Feild TS, Gleason SM,
41 401 Hacke UG, Jacobsen AL (2012) Global convergence in the vulnerability of forests to drought. *Nature*,
42 402 491(7426), 752-755
- 43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 403 Choat B (2013) Predicting thresholds of drought-induced mortality in woody plant species. *Tree Physiol*
4 33: 669-671
5 404
6 405 Ciais P, Reichstein M, Viovy N, Granier A, Ogée J, Allard V, Aubinet M, Buchmann N, Bernhofer C,
7 Carrara A, Chevallier F (2005) Europe-wide reduction in primary productivity caused by the heat and
8 406 drought in 2003. *Nature* 437: 529-533
9 407
10 408 Cochard H (2002a) Xylem embolism and drought-induced stomatal closure in maize. *Planta* 215: 466-471
11 409 Cochard H (2002b) A technique for measuring xylem hydraulic conductance under high negative
12 410 pressures. *Plant Cell Environ* 25: 815-819
13 411 Cochard H, Damour G, Bodet C, Tharwat I, Poirier M, Améglio T (2005) Evaluation of a new centrifuge
14 412 technique for rapid generation of xylem vulnerability curves. *Physiol Plantarum* 124: 410-418
15 413 Cochard H (2006) Cavitation in trees. *Comptes Rendus Physique* 7: 1018-1026
16 414 Cochard H, Casella E, Mencuccini M (2007) Xylem vulnerability to cavitation varies among poplar and
17 415 willow clones and correlates with yield. *Tree Physiol* 27: 1761-1767
18 416 Cochard H, Badel E, Herbette S, Delzon S, Choat B, Jansen S (2013) Methods for measuring plant
19 417 vulnerability to cavitation: a critical review. *J Exp Bot* 64: 4779-4791
20 418 Connor DJ, Hall AJ (1997) Sunflower physiology. In: Schneiter AA (ed) *Sunflower Science and*
21 419 *Technology*. The American Society of Agronomy, Madison, WI, USA, pp 113-182
22 420 Corcuera L, Cochard H, Gil-Pelegrin E, Notivol E (2011) Phenotypic plasticity in mesic populations of
23 421 *Pinus pinaster* improves resistance to xylem embolism (P_{50}) under severe drought. *Trees* 25: 1033-
24 422 1042
25 423 Dai A (2013) Increasing drought under global warming in observations and models. *Nat Clim Change* 3:
26 424 52-58
27 425 Davis SD, Ewers FW, Sperry JS, Portwood KA, Crocker MC, Adams GC (2002) Shoot dieback during
28 426 prolonged drought in *Ceanothus* (Rhamnaceae) chaparral of California: a possible case of hydraulic
29 427 failure. *Am J Bot* 89: 820-828
30 428 Delzon S, Douthe C, Sala A, Cochard H (2010) Mechanism of water-stress induced cavitation in conifers:
31 429 bordered pit structure and function support the hypothesis of seal capillary-seeding. *Plant Cell Environ*
32 430 33: 2101-2111
33 431 Ducrey M, Huc R, Ladjal M, Guehl J (2008) Variability in growth, carbon isotope composition, leaf gas
34 432 exchange and hydraulic traits in the eastern Mediterranean cedars *Cedrus libani* and *C. brevifolia*. *Tree*
35 433 *Physiol* 28: 689-701
36 434 Dufresne JL, Salas y Mélia D, Denvil S, Tyteca S, Arzel O, Bony S, Braconnot P, Brockmann P, Cadule
37 435 P, Caubel A, Chauvin F (2006) Simulation du climat récent et future par les modèles du CNRM et de
38 436 l'IPSL. *La Météorologie* 55: 45-59
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 437 Earl HJ, Davis RF (2003) Effect of drought stress on leaf and whole canopy radiation use efficiency and
4 438 yield of maize. *Agron J* 95: 688-696
- 5
6 439 Farooq M, Wahid A, Kobayashi N, Fujita D, Basra SMA (2009) Plant drought stress: effects, mechanisms
7 440 and management. *Agron Sustain Dev* 29: 185-212
- 8
9 441 Fichot R, Barigah TS, Chamaillard S, Le Thiec D, Laurans F, Cochard H, Brignolas F (2010) Common
10 442 trade-offs between xylem resistance to cavitation and other physiological traits do not hold among
11 443 unrelated *Populus deltoids* × *Populus nigra* hybrids. *Plant Cell Environ* 33: 1553-1568
- 12
13 444 Gleason SM, Westoby M, Jansen S, Choat B, Hacke UG, Pratt RB, Bhaskar R, Brodribb TJ, Bucci SJ,
14 445 Cao KF Cochard H (2015) Weak trade-off between xylem safety and xylem-specific hydraulic
15 446 efficiency across the world's woody plant species. *New Phytol* 209: 123-136
- 16
17 447 Guet J, Fichot R, Lédée C, Laurans F, Cochard H, Delzon S, Bastien C and Brignolas F (2015) Stem
18 448 xylem resistance to cavitation is related to xylem structure but not to growth and water-use efficiency
19 449 at the within-population level in *Populus nigra* L. *J Exp Bot* 66: 4643-4652
- 20
21 450 Hacke UG, Sperry JS, Pockman WT, Davis SD, McCulloh KA (2001) Trends in wood density and
22 451 structure are linked to prevention of xylem implosion by negative pressure. *Oecologia* 126: 457-461
- 23
24 452 Hajek P, Leuschner C, Hertel D, Delzon S, Schuldt B (2014) Trade-offs between xylem hydraulic
25 453 properties, wood anatomy and yield in *Populus*. *Tree Physiol* 34: 744-756
- 26
27 454 Hajek P, Kurjak D, von Wühlisch G, Delzon S, Schuldt B (2016) Intraspecific variation in wood
28 455 anatomical, hydraulic and foliar traits in ten European beech provenances differing in growth yield.
29 456 *Front Plant Sci* 7: 791
- 30
31 457 Harris D, Tripathi RS, Joshi A (2002) 'On-farm' seed priming to improve crop establishment and yield in
32 458 dry direct-seeded rice. In: Pandey S, Mortimer M, Wade L, Tuong TP, Lopes K, Hardy B (eds)
33 459 Proceedings of the international workshop on direct seeding in Asian rice systems: research strategies
34 460 and opportunities. International Research Institute, Manila, Philippines, pp 231-240
- 35
36 461 Hoffmann WA, Marchin RM, Abit P, Lau OL (2011) Hydraulic failure and tree dieback are associated
37 462 with high wood density in a temperate forest under extreme drought. *Glob Change Biol* 17: 2731-2742
- 38
39 463 Hussain S, Ali A, Ibrahim M, Saleem MF, Alias MA, Bukhsh HA (2012) Exogenous application of
40 464 abscisic acid for drought tolerance in sunflower (*Helianthus annuus* L.): a review. *J Anim Plant Sci* 22:
41 465 806-826
- 42
43 466 IPCC (2007) In: Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL
44 467 (eds) Climate change 2007: the physical science basis. Contribution of Working Group I to the fourth
45 468 assessment report of the Intergovernmental Panel on Climate Change. Cambridge University Press,
46 469 Cambridge, UK & New York, NY, USA, pp 1009.

- 1
2
3 470 Jaleel CA, Manivannan P, Kishorekumar A, Sankar B, Gopi R, Somasundaram R, Panneerselvam R
4
5 471 (2007) Alterations in osmoregulation, antioxidant enzymes and indole alkaloid levels in *Catharanthus*
6
7 472 *roseus* exposed to water deficit. *Colloid Surface B* 59: 150-157
- 8 473 Jensen HE, Mogensen VO (1984) Yield and nutrient content of spring wheat subjected to water stress at
9
10 474 various growth stages. *Acta Agr Scand* 34: 527-533
- 11 475 Kaya MD, Okçub G, Ataka M, Çıkılıç Y, Kolsarıcıa Ö (2006) Seed treatments to overcome salt and
12
13 476 drought stress during germination in sunflower (*Helianthus annuus* L.). *Eur J Agron* 24: 291-295
- 14 477 Kirda C (2002) Deficit irrigation scheduling based on plant growth stages showing water stress tolerance.
15
16 478 FAO, Deficit Irrigation Practices, Water Reports 22: 3-10
- 17
18 479 Lamy JB, Bouffier L, Burlett R, Plomion C, Cochard H, Delzon S (2011) Uniform selection as a primary
19
20 480 force reducing population genetic differentiation of cavitation resistance across a species range. *PLoS*
21
22 481 *One* 6: e23476
- 23 482 Lamy JB, Delzon S, Bouche PS, Alia R, Vendramin GG, Cochard H, Plomion C (2014) Limited genetic
24
25 483 variability and phenotypic plasticity detected for cavitation resistance in a Mediterranean pine. *New*
26
27 484 *Phytol* 201: 874-886
- 28 485 Larter M, Brodrribb TJ, Pfautsch S, Burlett R, Cochard H, Delzon S (2015) Extreme aridity pushes trees to
29
30 486 their physical limits. *Plant Physiol* 168: 804-807
- 31 487 Lawal BA, Obigbesan GO, Akanbi WB, Kolawole GO (2011) Effect of planting time on sunflower
32
33 488 (*Helianthus annuus* L.) productivity in Ibadan, Nigeria. *Afr J Agr Res* 6: 3049-3054
- 34 489 Lens F, Dressler S, Jansen S, van Evelghem L, Smets E (2005) Relationships within balsaminoid Ericales:
35
36 490 a wood anatomical approach. *Am J Bot* 92: 941-953
- 37 491 Lens F, Sperry JS, Christman MA, Choat B, Rabaey D, Jansen S (2011) Testing hypotheses that link
38
39 492 wood anatomy to cavitation resistance and hydraulic conductivity in the genus *Acer*. *New Phytol* 190:
40
41 493 709-723
- 42 494 Lens F, Tixier A, Cochard H, Sperry JS, Jansen S, Herbette S (2013) Embolism resistance as a key
43
44 495 mechanism to understand adaptive plant strategies. *Curr Opin Plant Biol* 16: 287-292
- 45 496 Lens F, Picon-Cochard C, Delmas CE, Signarbieux C, Buttler A, Cochard H, Jansen S, Chauvin T, Doria
46
47 497 LC, del Arco M, Delzon S (2016) Herbaceous angiosperms are not more vulnerable to drought-induced
48
49 498 embolism than angiosperm trees. *Plant Physiol* 00829
- 50 499 Li Y, Sperry JS, Shao M (2009) Hydraulic conductance and vulnerability to cavitation in corn (*Zea mays*
51
52 500 L.) hybrids of differing drought resistance. *Environ Exp Bot* 66: 341-346
- 53 501 Liu X, Baird WM (2003) Differential expression of genes regulated in response to drought or salinity
54
55 502 stress in sunflower. *Crop Sci* 43: 678-687
- 56
57
58
59
60

- 1
2
3 503 Lopez-Valdez F, Fernández-Luqueño F, Ceballos-Ramírez JM, Marsch R, Olalde-Portugal V, Dendooven
4 L (2011) A strain of *Bacillus subtilis* stimulates sunflower growth (*Helianthus annuus* L.) temporarily.
5 504
6 Sci Hort 128: 499-505
7 505
- 8 506 Martínez-Vilalta J, Cochard H, Mencuccini M, Sterck F, Herrero A, Korhonen JFJ, Llorens P, Nikinmaa
9 E, Nole A, Poyatos R, Ripullone F (2009) Hydraulic adjustment of Scots pine across Europe. New
10 507
11 Phytol 184: 353-364
12 508
- 13 509 Manickavelu A, Nadarajan N, Ganesh SK, Gnanamalar RP, Babu RC (2006) Drought tolerance in rice:
14 510
15 morphological and molecular genetic consideration. Plant Growth Regul 50: 121-138
- 16 511 Neufeld HS, Grantz DA, Meinzer FC, Goldstein G, Crisosto GM, Crisosto C (1992) Genotypic variability
17 512
18 in vulnerability of leaf xylem to cavitation in water-stressed and well-irrigated sugarcane. Plant Physiol
19 513
20 100: 1020-1028
- 21 514 Pammenter NV, Vander Willigen C (1998) A mathematical and statistical analysis of the curves
22 515
23 illustrating vulnerability of xylem to cavitation. Tree Physiol 18: 589-593
- 24 516 Parmesan C, Yohe G (2003) A globally coherent fingerprint of climate change impacts across natural
25 517
26 systems. Nature 421: 37-42
- 27 518 Pittermann J, Limm E, Rico C, Christman MA (2011) Structure–function constraints of tracheid-based
28 519
29 xylem: a comparison of conifers and ferns. New Phytol 192: 449-461
- 30 520 Pivovarov AL, Burlett R, Lavigne B, Cochard H, Santiago LS, Delzon S (2016) Testing the ‘microbubble
31 521
32 effect’ using the Cavitron technique to measure xylem water extraction curves. AoB Plants 8: 011
- 33 522 Salleo S, Nardini A, Pitt F, Gullo MAL (2000) Xylem cavitation and hydraulic control of stomatal
34 523
35 conductance in laurel (*Laurus nobilis* L.). Plant Cell Environ 23: 71-79
- 36 524 Sáenz-Romero C, Lamy JB, Loya-Rebollar E, Plaza-Aguilar A, Burlett R, Lobit P, Delzon S (2013)
37 525
38 Genetic variation of drought-induced cavitation resistance among *Pinus hartwegii* populations from an
39 526
40 altitudinal gradient. Acta Physiol Plant 35: 2905-2913
- 41 527 Scholz A, Klepsch M, Karimi Z, Jansen S (2013) How to quantify conduits in wood? Front Plant Sci 4: 56
- 42 528 Schuldt B, Knutzen F, Delzon S, Jansen S, Müller-Haubold H, Burlett R, Clough Y, Leuschner C (2015)
43 529
44 How adaptable is the hydraulic system of European beech in the face of climate change-related
45 530
46 precipitation reduction? New Phytol 210: 443-458
- 47 531 Somerville C, Briscoe J (2001) Genetic engineering and water. Science 292: 2217-2217
- 48 532 Sperry JS (2000) Hydraulic constraints on plant gas exchange. Agr Forest Meteorol 104: 13-23
- 49 533 Sperry JS, Stiller V, Hacke UG (2003) Xylem hydraulics and the soil–plant–atmosphere continuum.
50 534
51 Agron J 95: 1362-1370
- 52 535 Sperry JS, Hacke UG, Wheeler JK (2005) Comparative analysis of end wall resistivity in xylem conduits.
53 536
54 Plant Cell Environ 28: 456-465
- 55
56
57
58
59
60

- 1
2
3 537 Sperry JS, Hacke UG, Pittermann J (2006) Size and function in conifer tracheids and angiosperm vessels.
4 538 Am J Bot 93: 1490-1500
- 5
6 539 Sperry JS, Meinzer FC, McCULLOH KA (2008) Safety and efficiency conflicts in hydraulic architecture:
7 540 scaling from tissues to trees. Plant Cell Environ 31: 632-645
- 8
9 541 Sterck FJ, Martínez-Vilalta J, Mencuccini M, Cochard H, Gerrits P, Zweifel R, Herrero A, Korhonen JF,
10 542 Llorens P, Nikinmaa E, Nole A (2012) Understanding trait interactions and their impacts on growth in
11 543 Scots pine branches across Europe. Funct Ecol 26: 541-549
- 12
13 544 Sterl A, Severijns C, Dijkstra H, Hazeleger W, van Oldenborgh GJ, van den Broeke M, Burgers G, van
14 545 den Hurk B, van Leeuwen PJ, van Velthoven P (2008) When can we expect extremely high surface
15 546 temperatures? Geophys Res Lett 35: 1-5
- 16
17 547 Stiller V, Sperry JS (2002) Cavitation fatigue and its reversal in sunflower (*Helianthus annuus* L.). J Exp
18 548 Bot 53: 1155-1161
- 19
20 549 Stiller V, Lafitte HR, Sperry JS (2003) Hydraulic properties of rice and the response of gas exchange to
21 550 water stress. Plant Physiol 132: 1698-1706
- 22
23 551 Tixier A, Cochard H, Badel E, Dusotoit-Coucaud A, Jansen S, Herbette S (2013) *Arabidopsis thaliana* as
24 552 a model species for xylem hydraulics: does size matter? J Exp Bot 64: 2295-2305
- 25
26 553 Trifilò P, Raimondo F, Lo Gullo MA, Barbera PM, Salleo S, Nardini A (2014) Relax and refill: xylem
27 554 rehydration prior to hydraulic measurements favours embolism repair in stems and generates
28 555 artificially low PLC values. Plant Cell Environ 37: 2491-2499
- 29
30 556 Tripathy JN, Zhang J, Robin S, Nguyen TT, Nguyen HT (2000) QTLs for cell-membrane stability mapped
31 557 in rice (*Oryza sativa* L.) under drought stress. Theor Appl Genet 100: 1197-1202
- 32
33 558 Tyree MT, Fiscus EL, Wullschlegel SD, Dixon MA (1986) Detection of xylem cavitation in corn under
34 559 field conditions. Plant Physiol 82: 597-599
- 35
36 560 Tyree MT, Sperry JS (1989) Vulnerability of xylem to cavitation and embolism. Annu Rev Plant Biol 40:
37 561 19-36
- 38
39 562 Tyree MT, Zimmermann MH (2002) Hydraulic architecture of whole plants and plant performance. In:
40 563 Xylem structure and the ascent of sap. 2nd Edn. Springer Berlin Heidelberg, pp 175-214
- 41
42 564 Urli M, Porté AJ, Cochard H, Guengant Y, Burlett R, Delzon S (2013) Xylem embolism threshold for
43 565 catastrophic hydraulic failure in angiosperm trees. Tree physiol 33: 672-683
- 44
45 566 Unger PW (1990) Sunflower. In: Stewart BA, Nielsen DR (eds) Irrigation of Agricultural Crops.
46 567 Agronomy Monograph No. 30. Madison, WI, USA, pp 775-794
- 47
48 568 van Oldenborgh GJ, Drijfhout S, van Ulden A, Haarsma R, Sterl A, Severijns C, Hazeleger W, Dijkstra H
49 569 (2009) Western Europe is warming much faster than expected. Clim Past 5: 1-12
- 50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 570 Velue G, Palanisami K (2001) Impact of moisture stress and ameliorants on growth and yield of
4 sunflower. Madras Agr J 88: 660-665
5 571
6 572 Vitousek PM, Mooney HA, Lubchenco J, Melillo JM (1997). Human domination of Earth's ecosystems.
7 Science 277: 494-499
8 573
9 574 Wang Y, Burrell R, Feng F, Tyree MT (2014) Improved precision of hydraulic conductance measurements
10 using a Cochard rotor in two different centrifuges. J Plant Hydraul 1: 007
11 575
12 576 Watkins JE, Holbrook NM, Zwieniecki MA (2010) Hydraulic properties of fern sporophytes:
13 consequences for ecological and evolutionary diversification. Am J Bot 97: 2007-2019
14 577
15 578 Wheeler TD, Stroock AD (2008) The transpiration of water at negative pressures in a synthetic tree.
16 Nature 455: 208-212
17 579
18 580 Wikberg J, Ögren E (2004) Interrelationships between water use and growth traits in biomass-producing
19 willows. Trees 18: 70-76
20 581
21 582 Wigley TML (2009) The effect of changing climate on the frequency of absolute extreme events.
22 Climatic Change 97: 67-76
23 583
24 584 Zwieniecki MA, Melcher PJ, Holbrook NM (2001) Hydraulic properties of individual xylem vessels of
25 *Fraxinus americana*. J Exp Bot 52: 257-264
26 585
27
28
29
30

31 **Supporting Information**

32 Additional Supporting Information may be found in the online version of this article:

33 **Table S1.** Correlations between all the hydraulic and anatomical traits studied.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

590 **Table S1.** Non-significant ($P > 0.05$) correlations between anatomical and hydraulic traits. r values are Pearson correlation coefficients. See Table
 591 1 for abbreviations.

592

Anatomical traits	Hydraulic traits									
	P_{12}		P_{50}		P_{88}		S		K_s	
	r	P	r	P	r	P	r	P	r	P
A_{stem} (μm^2)	0.43	0.1884	0.26	0.4453	-0.09	0.7936	-0.43	0.1855	-0.33	0.3152
A_{Tx} (μm^2)	0.51	0.1103	0.31	0.3513	-0.09	0.7768	-0.49	0.1199	-0.27	0.4213
A_{Px} (μm^2)	0.30	0.3638	0.16	0.6404	-0.09	0.7764	-0.26	0.4424	-0.23	0.4931
A_{Sx} (μm^2)	0.51	0.1080	0.31	0.3441	-0.09	0.7843	-0.50	0.1143	-0.26	0.4309
A_{pith} (μm^2)	0.41	0.2054	0.25	0.4435	-0.07	0.8323	-0.41	0.2083	-0.38	0.2515
P_{ps}	-0.01	0.9809	0.05	0.8691	0.09	0.7906	0.02	0.9420	-0.34	0.2964
A_{cp} (μm^2)	0.41	0.2045	0.22	0.4992	-0.11	0.7334	-0.43	0.1894	-0.21	0.5423
A_{lig} (μm^2)	0.52	0.1041	0.32	0.3305	-0.08	0.8006	-0.50	0.1162	-0.25	0.4481
P_{ligs}	0.29	0.3878	0.22	0.5156	0.01	0.9841	-0.22	0.5037	0.26	0.4340
P_{ligcp}	0.46	0.1580	0.49	0.1210	0.22	0.5027	-0.31	0.3578	-0.29	0.3914
A_{fcap} (μm^2)	0.37	0.2571	0.36	0.2723	0.12	0.7176	-0.31	0.3481	0.11	0.7517
A_{fcell} (μm^2)	0.46	0.1493	0.34	0.2977	-0.00	0.9990	-0.34	0.3008	-0.16	0.6383
A_{flumen} (μm^2)	0.47	0.1470	0.15	0.6613	-0.29	0.3901	-0.46	0.1488	0.00	0.9874
A_{fcw} (μm^2)	0.32	0.3343	0.39	0.2276	0.23	0.5005	-0.14	0.6763	-0.24	0.4723
P_{cwf}	-0.12	0.7331	0.19	0.5769	0.40	0.2217	0.25	0.4498	-0.17	0.6185
A_{fcw} in A_{lig} (μm^2)	0.43	0.1854	0.42	0.1929	0.15	0.6574	-0.32	0.3349	-0.30	0.3649
P_{fw} in $ligs$	0.07	0.8452	0.27	0.4226	0.31	0.3383	0.08	0.8111	0.01	0.9679
VD_{Px} (n mm^{-2})	-0.57	0.0637	-0.49	0.1207	-0.09	0.7737	0.32	0.3434	0.02	0.9526
A_{vSx} (μm^2)	0.38	0.2477	0.31	0.3496	0.04	0.9019	-0.23	0.4902	-0.49	0.1186
A_{cvPx} (μm^2)	0.11	0.7380	-0.01	0.9719	-0.14	0.6793	-0.15	0.6519	-0.42	0.1931
A_{cvSx} (μm^2)	0.22	0.5000	-0.09	0.7900	-0.38	0.2508	-0.36	0.2741	-0.33	0.3143
A_{rvPx} (μm^2)	-0.13	0.7021	-0.21	0.5255	-0.17	0.6146	-0.01	0.9662	-0.56	0.0732
T_{vwSx} (μm)	-0.25	0.4556	-0.31	0.3458	-0.18	0.5851	-0.03	0.9202	-0.23	0.4892
TD_{Sx}^{-1}	-0.09	0.7845	-0.01	0.9732	0.08	0.8053	0.00	0.9903	-0.11	0.7509
D_{Sx} (μm)	0.42	0.1966	0.35	0.2809	0.06	0.8532	-0.25	0.4595	-0.48	0.1330
D_{hPx} (μm)	0.02	0.9493	-0.13	0.6960	-0.21	0.5212	-0.09	0.7986	-0.45	0.1666
D_{hSx} (μm)	0.10	0.7626	0.04	0.8980	-0.05	0.8891	-0.07	0.8451	-0.44	0.1759

593 **Table 1.** List of the traits studied, including their units and descriptions

Traits (units)	Description
Growth	
H (m)	Height
D (mm)	Stem diameter
Hydraulics	
P_{12} (MPa)	Xylem pressure inducing a 12% loss of hydraulic conductance
P_{50} (MPa)	Xylem pressure inducing a 50% loss of hydraulic conductance
P_{88} (MPa)	Xylem pressure inducing a 88% loss of hydraulic conductance
S (% MPa ⁻¹)	Slope of the vulnerability curve at the inflexion point
K_s (m ² MPa ⁻¹ s ⁻¹)	Xylem specific hydraulic conductivity
Anatomy	
A_{stem} (μm ²)	Stem cross-section area
A_{Tx} (μm ²)	Total xylem area
A_{Px} (μm ²)	Primary xylem area
A_{Sx} (μm ²)	Secondary xylem area
A_{pith} (μm ²)	Pith area
A_{cp} (μm ²)	Area of the cellular part of stem
A_{lig} (μm ²)	Lignified area
A_{fcap} (μm ²)	Fiber cap area (sum of the areas of all fiber caps in the stem cross-section)
A_{fcell} (μm ²)	Fiber cell area
A_{flumen} (μm ²)	Fiber lumen area
A_{fcw} (μm ²)	Fiber cell wall area
$A_{fcw \text{ in } A_{lig}}$ (μm ²)	Total fiber wall area in the lignified area
A_{vPx} (μm ²)	Vessel lumen area in the primary xylem
A_{vSx} (μm ²)	Vessel lumen area in the secondary xylem
A_{cvPx} (μm ²)	Cumulative vessel lumen area in the primary xylem
A_{cvSx} (μm ²)	Cumulative vessel lumen area in the secondary xylem
A_{rvPx}	Vessel lumen area relative to primary xylem area
A_{rvSx}	Vessel lumen area relative to secondary xylem area

1		
2		
3	P_{ps}	Pith area as a proportion of stem area
4		
5	$P_{lig s}$	Lignified area as a proportion of stem area
6		
7	$P_{lig cp}$	Lignified area as a proportion of the area of the cellular part of the stem
8		
9	P_{cwf}	Cell wall area as a proportion of fiber cell area
10		
11	$P_{fw in lig s}$	Fiber wall area in the lignified area as a proportion of stem area
12	VD_{Px} (n mm ⁻²)	Vessel density in the primary xylem
13		
14	VD_{Sx} (n mm ⁻²)	Vessel density in the secondary xylem
15		
16	VD_{Tx} (n mm ⁻²)	Vessel density in the total xylem
17		
18	T_{vwPx} (μm)	Inter-vessel double-wall thickness in the primary xylem
19		
20	T_{vwSx} (μm)	Inter-vessel double-wall thickness in the secondary xylem
21		
22	TD^{-1}_{Px}	Thickness-to-span ratio of vessels in the primary xylem
23		
24	TD^{-1}_{Sx}	Thickness-to-span ratio of vessels in the secondary xylem
25		
26	D_{Px} (μm)	Equivalent circle diameter of vessels in the primary xylem
27		
28	D_{Sx} (μm)	Equivalent circle diameter of vessels in the secondary xylem
29		
30	D_{hPx} (μm)	Hydraulically weighted vessel diameter in the primary xylem
31	D_{hSx} (μm)	Hydraulically weighted vessel diameter in the secondary xylem
32		

594

595 **Table 2.** Mean values (\pm SE) of traits related to growth and hydraulic properties for four sunflower
 596 accessions. Letters in bold indicate significant statistical differences between accessions ($P < 0.05$).
 597 Sampling sizes are indicated in brackets.

598

Traits	Sunflower accessions			
	Melody	LG_5660	ES_Ethic	ES_Biba
Growth				
H	0.95 \pm 0.07 a (16)	0.91 \pm 0.07 a (14)	0.78 \pm 0.06 b (20)	0.78 \pm 0.11 b (18)
D	8.75 \pm 1.18 a (16)	8.68 \pm 1.24 a (14)	8.02 \pm 0.92 a (20)	8.57 \pm 1.12 a (18)
Hydraulics				
P_{12}	-2.19 \pm 0.23 a (16)	-2.18 \pm 0.27 a (17)	-2.57 \pm 0.33 b (16)	-2.29 \pm 0.31 a (15)
P_{50}	-3.09 \pm 0.12 b (16)	-2.67 \pm 0.16 a (17)	-3.22 \pm 0.13 c (16)	-3.01 \pm 0.18 b (15)
P_{88}	-3.99 \pm 0.27 c (16)	-3.14 \pm 0.15 a (17)	-3.86 \pm 0.27 bc (16)	-3.73 \pm 0.30 b (15)
S	59.78 \pm 17.88 c (16)	120.48 \pm 52.91 a (17)	95.10 \pm 48.49 ab (16)	78.78 \pm 30.78 bc (15)
K_s	1.6 $\times 10^{-4} \pm 0.9 \times 10^{-4}$ ab (16)	1.4 $\times 10^{-4} \pm 0.8 \times 10^{-4}$ b (17)	2.4 $\times 10^{-4} \pm 0.9 \times 10^{-4}$ a (16)	1.7 $\times 10^{-4} \pm 0.7 \times 10^{-4}$ ab (15)

599

Table 3. Relationships between hydraulic, growth and anatomical traits. The values shown are the Pearson correlation coefficients. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. See Table 1 for trait descriptions.

Hydraulic traits	Growth traits		Anatomical traits						
	H	D	VD_{Tx}	VD_{Sx}	A_{rvSx}	T_{vwPx}	TD^{-1}_{Px}	A_{vPx}	D_{Px}
P_{12}	0.54***	0.15	-0.76**	-0.77**	-0.34	-0.60*	-0.63*	0.37	0.41
P_{50}	0.42**	0.08	0.85***	0.92***	-0.60*	-0.31	-0.43	0.21	0.27
P_{88}	0.09	-0.02	-0.41	-0.51	-0.51	0.21	0.06	-0.09	-0.06
S	-0.35*	-0.20	0.44	0.41	0.10	0.56	0.45	-0.31	-0.32
K_s	-0.56***	-0.47***	0.19	0.25	-0.18	-0.26	-0.09	-0.63*	-0.64*

Figure legends

Fig. 1. Vulnerability curves (VCs) for individuals of the four sunflower accessions studied, for which xylem embolism was induced by *in situ* flow centrifugation according to the Cavitron technique. $n = 16$, 17, 16 and 15 for Melody, LG_5660, ES_Ethic and ES_Biba, respectively. VCs are expressed as the percentage loss of hydraulic conductivity (PLC) as a function of xylem pressure.

Fig. 2. Mean vulnerability curves (VCs) (\pm SE) for each of the four sunflower accessions studied. $n = 16$, 17, 16 and 15 for Melody, LG_5660, ES_Ethic and ES_Biba, respectively. VCs were generated by the *in situ* flow centrifugation (Cavitron) technique and are expressed as the percentage loss of hydraulic conductivity (PLC) as a function of xylem pressure.

Fig. 3. Relationships between height (H, m), xylem embolism resistance (P_{50} , MPa) and xylem specific hydraulic conductivity (K_s , $m^2 MPa^{-1} s^{-1}$). $n = 49$, 63 and 49 for panels A, B and C, respectively.

Fig. 4. Relationship between vessel density in secondary xylem (VD_{Sx} ; A), vessel density in total xylem (VD_{Tx} ; B), vessel lumen area relative to secondary xylem area (A_{rvSx} ; C) and xylem embolism resistance (P_{50} , MPa). $n = 11$.

Fig. 5. Transverse sections of (A, C) the most resistant (ES_Ethic) and (B, D) the most vulnerable (LG_5660) sunflower accessions. Plants were grown in pots filled with Peltracom substrate placed in a greenhouse under full light and non-limiting growing conditions. They were sectioned with a sledge microtome ((Reichert, Germany). (A, C) Sections cut from the middle of the plant stem. (B, D) Overview

1
2
3 625 of mature stems highlighting the similarity in lignified area (A_{lig}) and the difference in vessel density in the
4
5 626 secondary xylem (VD_{Sx}) (black arrows).
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review




Fig. 1. Vulnerability curves (VCs) for individuals of the four sunflower accessions studied, for which xylem embolism was induced by in situ flow centrifugation according to the Cavitron technique. $n = 16, 17, 16$ and 15 for Melody, LG_5660, ES_Ethic and ES_Biba, respectively. VCs are expressed as the percentage loss of hydraulic conductivity (PLC) as a function of xylem pressure.

304x203mm (96 x 96 DPI)

review




Fig. 2. Mean vulnerability curves (VCs) (\pm SE) for each of the four sunflower accessions studied. $n = 16, 17, 16$ and 15 for Melody, LG_5660, ES_Ethic and ES_Biba, respectively. VCs were generated by the in situ flow centrifugation (Cavitrion) technique and are expressed as the percentage loss of hydraulic conductivity (PLC) as a function of xylem pressure.

317x211mm (96 x 96 DPI)

review




Fig. 3. Relationships between height (H , m), xylem embolism resistance (P_{50} , MPa) and xylem specific hydraulic conductivity (K_s , m² MPa⁻¹ s⁻¹). $n = 49, 63$ and 49 for panels A, B and C, respectively.

338x190mm (96 x 96 DPI)

Review




Fig. 4. Relationship between vessel density in secondary xylem (VDS_x ; A), vessel density in total xylem (VDT_x ; B), vessel lumen area relative to secondary xylem area (A_{vSx} ; C) and xylem embolism resistance (P_{50} , MPa). $n = 11$.

338x190mm (96 x 96 DPI)

Review




Fig. 5. Transverse sections of (A, C) the most resistant (ES_Ethic) and (B, D) the most vulnerable (LG_5660) sunflower accessions. Plants were grown in pots filled with Peltracom substrate placed in a greenhouse under full light and non-limiting growing conditions. They were sectioned with a sledge microtome ((Reichert, Germany). (A, C) Sections cut from the middle of the plant stem. (B, D) Overview of mature stems highlighting the similarity in lignified area (Alig) and the difference in vessel density in the secondary xylem (VDSx) (black arrows).

338x190mm (96 x 96 DPI)

Review

Table: S1 Non-significant ($P > 0.05$) correlations between anatomical and hydraulic traits. r values are Pearson correlation coefficients. See Table 1 for abbreviations.

Anatomical traits	Hydraulic traits									
	P_{12}		P_{50}		P_{88}		S		K_s	
	r	P	r	P	r	P	r	P	r	P
A_{stem} (μm^2)	0.43	0.1884	0.26	0.4453	-0.09	0.7936	-0.43	0.1855	-0.33	0.3152
A_{Tx} (μm^2)	0.51	0.1103	0.31	0.3513	-0.09	0.7768	-0.49	0.1199	-0.27	0.4213
A_{Px} (μm^2)	0.30	0.3638	0.16	0.6404	-0.09	0.7764	-0.26	0.4424	-0.23	0.4931
A_{Sx} (μm^2)	0.51	0.1080	0.31	0.3441	-0.09	0.7843	-0.50	0.1143	-0.26	0.4309
A_{pith} (μm^2)	0.41	0.2054	0.25	0.4435	-0.07	0.8323	-0.41	0.2083	-0.38	0.2515
P_{ps}	-0.01	0.9809	0.05	0.8691	0.09	0.7906	0.02	0.9420	-0.34	0.2964
A_{cp} (μm^2)	0.41	0.2045	0.22	0.4992	-0.11	0.7334	-0.43	0.1894	-0.21	0.5423
A_{lig} (μm^2)	0.52	0.1041	0.32	0.3305	-0.08	0.8006	-0.50	0.1162	-0.25	0.4481
P_{ligs}	0.29	0.3878	0.22	0.5156	0.01	0.9841	-0.22	0.5037	0.26	0.4340
P_{ligcp}	0.46	0.1580	0.49	0.1210	0.22	0.5027	-0.31	0.3578	-0.29	0.3914
A_{fcap} (μm^2)	0.37	0.2571	0.36	0.2723	0.12	0.7176	-0.31	0.3481	0.11	0.7517
A_{fcell} (μm^2)	0.46	0.1493	0.34	0.2977	-0.00	0.9990	-0.34	0.3008	-0.16	0.6383
A_{flumen} (μm^2)	0.47	0.1470	0.15	0.6613	-0.29	0.3901	-0.46	0.1488	0.00	0.9874
A_{fcw} (μm^2)	0.32	0.3343	0.39	0.2276	0.23	0.5005	-0.14	0.6763	-0.24	0.4723
P_{cwf}	-0.12	0.7331	0.19	0.5769	0.40	0.2217	0.25	0.4498	-0.17	0.6185
A_{fcw} in A_{lig} (μm^2)	0.43	0.1854	0.42	0.1929	0.15	0.6574	-0.32	0.3349	-0.30	0.3649
P_{fw} in $ligs$	0.07	0.8452	0.27	0.4226	0.31	0.3383	0.08	0.8111	0.01	0.9679
VD_{Px} (n mm^{-2})	-0.57	0.0637	-0.49	0.1207	-0.09	0.7737	0.32	0.3434	0.02	0.9526
A_{vSx} (μm^2)	0.38	0.2477	0.31	0.3496	0.04	0.9019	-0.23	0.4902	-0.49	0.1186
A_{cvPx} (μm^2)	0.11	0.7380	-0.01	0.9719	-0.14	0.6793	-0.15	0.6519	-0.42	0.1931
A_{cvSx} (μm^2)	0.22	0.5000	-0.09	0.7900	-0.38	0.2508	-0.36	0.2741	-0.33	0.3143
A_{rvPx} (μm^2)	-0.13	0.7021	-0.21	0.5255	-0.17	0.6146	-0.01	0.9662	-0.56	0.0732
T_{vwSx} (μm)	-0.25	0.4556	-0.31	0.3458	-0.18	0.5851	-0.03	0.9202	-0.23	0.4892
TD^{-1}_{Sx}	-0.09	0.7845	-0.01	0.9732	0.08	0.8053	0.00	0.9903	-0.11	0.7509
D_{Sx} (μm)	0.42	0.1966	0.35	0.2809	0.06	0.8532	-0.25	0.4595	-0.48	0.1330
D_{hPx} (μm)	0.02	0.9493	-0.13	0.6960	-0.21	0.5212	-0.09	0.7986	-0.45	0.1666
D_{hSx} (μm)	0.10	0.7626	0.04	0.8980	-0.05	0.8891	-0.07	0.8451	-0.44	0.1759

Part II: Intraspecific variability in embolism resistance and its relation with genetic architecture and photosynthetic traits in Maritime pine.

Chapter 2: Dissecting the genetic architecture of resistance in Maritime pine reveals that hydraulic safety is not linked to wood density. *(in preparation)*

Dissecting the genetic architecture of embolism resistance in maritime pine (*Pinus pinaster* Ait.) reveals that hydraulic safety is not linked to wood density. (*in preparation*)

Abstract

Hydraulic failure due to vascular embolism is likely the main cause of drought-induced forest mortality reported worldwide. The vulnerability to embolism has therefore been intensively investigated across species during the last decades, but we still know little about the intraspecific variability of this trait and its genetic determinism. To provide better insight about the genetic determinism of resistance to embolism, we dissected here its genetic architecture in concert with that of wood density in a large-size three-generation inbred pedigree of maritime pine (*Pinus pinaster* Ait.). We carried out the largest effort of phenotyping both hydraulic and wood density in 477 maritime pine genotypes. The variability of embolism resistance was remarkably low compared to that of wood density and radial growth. No significant correlations between embolism resistance and wood density were found. In addition, the comparison of quantitative trait loci (QTL) locations between embolism resistance and wood density related traits failed to detect co-locating genomic regions. The lack of phenotypic correlation and physical coincidence of QTLs in this family strongly suggest the absence of causality between these two traits in this species.

Introduction

Extreme droughts are predicted to be more intense, more frequent, and longer in the 21st Century (Meehl and Tebaldi 2004; Bahn et al., 2014; Lindner et al., 2014), placing forest systems in a more and more vulnerable situation. In the light of these events, the European heat wave of 2003 was a drastic demonstration of the extent of impacts we need to expect more often in the future (Ciais et al., 2005). Forest ecosystems have already been altered by climate change, leading to decreased growth (Piao et al., 2011; Zhao et al., 2010) and species distribution shift (Delzon et al., 2013), and to widespread mortality of trees (Allen et al., 2010; Michaelian et al., 2011). This global drought-induced forest mortality could be explained by the high vulnerability of trees to

drought (Choat et al., 2012). With their long life-spans, trees cannot adapt rapidly to environmental changes, and most forest tree species will have to cope with these changes within a single generation (Lindner et al., 2010; Kremer et al., 2014). To better predict the resilience of forests to extreme droughts, a deeper understanding is needed about the key physiological traits underlying drought tolerance, their variations across and within species and their genetic determinisms (Plomion et al., 2016).

Xylem sap being transported under negative pressure in the vascular system of trees, during drought stress, sap pressures become excessively negative inside vessels and air embolism can form, responsible of hydraulic failure. Several studies demonstrated experimentally that the vulnerability to embolism is related to the survival of tree species in drought conditions (Brodribb and Cochard, 2009; Brodribb et al., 2010; Urli et al., 2013). In addition, the breakdown of the vascular system through embolism was found to be involved in multiple mass mortality events observed *in situ* under drought conditions (Anderegg et al., 2015, 2016). This mechanistic trait is therefore a crucial adaptive trait for trees to cope with severe drought (Larter et al., 2017). At the interspecific level, resistance to embolism has been described as highly variable across a wide range of species, species from xeric environments being more resistant to cavitation than species from mesic environments (Maherali et al., 2004; Cochard et al., 2008; Choat et al., 2012; Bouche et al., 2014). The incorporation of phylogenetic information also showed that embolism-related traits are under natural selection (Maherali et al., 2004; Pittermann et al., 2012; Larter et al., 2017). Conversely, little is known about the level and structure of intraspecific variability of embolism resistance, as well as its relationships with species autoecology. Based on common garden experiments (e.g. Lamy et al., 2011 for *Pinus pinaster* and Kavanagh et al., 1999 for *Douglas-fir ponderosa*) it was shown that intra-specific genetic variation in embolism resistance did not reveal significant differences between populations which leads Lamy et al. (2014) to suggest that this trait was under strong canalization, at least in *Pinus pinaster* where the large sample size of their study allowed such inference.

At the cellular level, vulnerability to cavitation has been shown to be intimately linked to the anatomy of xylem cells (Hacke et al., 2001), and in conifers, the mechanism of drought-induced cavitation largely depends on pit membrane properties (Delzon et al., 2010; Pittermann et al., 2010; Bouche et al., 2014). However, high throughput measurement of pit membrane anatomy is not feasible yet. At the interspecific level, Hacke et al. (2001) associated greater wall thickness per

unit diameter in conifers (12 species) and angiosperms (36 species) with greater wood density and proposed to use wood density as good proxy of resistance for embolism. Delzon et al. (2010) studied 40 conifers and also reported a significant relationship between wood density and resistance to embolism but returns the density of the wood to a secondary role because all species that are in the higher density range are not all resistant to embolism. Such a relationship between wood density and embolism resistance was tested at the intraspecific level in maritime pine (Lamy et al., 2012) and in Douglas-fir (Dalla-Salda et al., 2009). Whereas Douglas-fir trees with higher wood density exhibited higher probability to survive drought than trees with a lower density, no relationships with embolism resistance was reported in maritime pine. By contrast, Rosner et al. (2017) found that trunk wood density was strongly related to resistance to embolism across cambial age in Norway spruce.

If we are beginning to better understand the genetics of embolism resistance (reviewed by Plomion et al., 2016 for *Pinus pinaster*), insights into the genetic architecture (number, chromosomal location and effect of quantitative trait loci, QTL) of vulnerability to embolism are still lacking, owing to the fact that high throughput phenotyping techniques of large populations (Liebhard et al., 2003) were inaccessible until recently. To our knowledge, a single study published by Lauri et al. (2011) failed to detect any QTL for embolism resistance in a *Malus domestica* full-sib mapping population of 123 trees.

In this context, the objectives of the present study were two-fold: i/ dissect the genetic architecture of embolism related traits (hydraulic safety) and wood density related traits, and (ii) test the relationship between wood density and xylem safety at the phenotypic and QTL level. To this end we phenotyped an already genotyped (Bartholomé et al., 2016) large-size three-generation inbred pedigree of *Pinus pinaster*. This species presents a high ecological and economic importance in the Western Mediterranean basin where it spans over 4M ha. The most recent climatic scenarios (IPCC, 2014) predict a strong modification of summer precipitation and temperature in this region by the end of this century. It is therefore important to know about the genetic architecture of traits that determines maritime pine survival during severe period of droughts if such characteristics are to be incorporated into breeding programs.

Material and methods

Plant material, genotyping and linkage map construction

The mapping population used in this study consisted of a three-generation inbred pedigree (F2) obtained by the self-pollination of an inter “Landes x Corsica” provenance hybrid (accession H12). Seeds were planted in a nursery (June 1998) and transplanted into the field (March 1999). Genotyping and phenotypic data were obtained for 477 F2 trees. The genotyping of this population is detailed in Bartholomé et al. (2016). A total of 248 SNP markers were mapped along the 12 linkage groups of the linkage map established for accession H12. Overall, 530 branches (i.e. assessment of some trees were replicated) were used to phenotype the F2s for cavitation resistance and wood density.

Phenotyping

Cavitation resistance related traits

For each F2, two branches fully exposed to the sun were collected in the early morning from November 2012 to April 2013. Forty cm branches of less than 1 cm diameter were chosen. All needles were immediately removed, and then the branches were labeled and placed in black bags, wrapped in wet paper towels. They were kept wet and cool (3°C) until measurement within three weeks after collection. Prior to embolism resistance measurement, branches were cut under water to a standard length of 27 cm, and bark was removed with a razor blade.

Xylem vulnerability to embolism was assessed with the CAVITRON, a centrifuge technique following the procedure described by Cochard (Cochard, 2002; Cochard et al., 2005) at the high-throughput phenotyping platform for hydraulic traits (CavitPlace, University of Bordeaux, Talence, France; http://sylvain-delzon.com/?page_id=536). Centrifugal force was used to establish negative pressure in the xylem and to provoke water stress-induced embolism, using a custom-built honeycomb rotor (Precis 2000, Bordeaux, France) mounted on a high-speed centrifuge (Sorvall RC5, USA). This technique enables measurement of the hydraulic conductance of a branch under negative pressure. Xylem pressure (P) was first set to a reference pressure (-0.5 MPa) and the maximal hydraulic conductivity (K) was determined by measuring the flux of a reference ionic solution (10 mM KCl and 1 mM CaCl₂ in deionized water) through the sample. The centrifugation speed was then set to a higher value for 3 min to expose the sample at a more negative pressure. Conductances were measured four times for each step, and the average was used to compute the percent loss of xylem conductance (PLC in %). The procedure was repeated for at least eight pressure steps with a -0.5 MPa step increment until PLC reached at least 90%. Rotor velocity was monitored with a 10 rpm resolution electronic tachymeter and xylem pressure was

adjusted to about -0.02 MPa. We used Cavi_soft software (version2.0, BIOGECO, University of Bordeaux) for conductance measurements.

The percent loss of xylem conductivity as a function of xylem pressure (MPa) represents the sample's vulnerability curve (VC). A sigmoid function (Pammenter and van der Willigen, 1998) was fitted to the VC from each sample using the following equation:

$$PLC = \frac{100}{\left[1 + \exp\left(\frac{S}{25} \times (P - P_{50})\right) \right]}$$

where P_{50} (MPa) is the xylem pressure inducing 50% loss of hydraulic conductivity and S (% MPa⁻¹) is the slope of the vulnerability curve at the inflexion point. The xylem pressure inducing 12 and 88% of PLC (P_{12} , P_{88} , MPa) was calculated from the equation of Domec and Gartner (2001):

$$P_{12} = P_{50} + \frac{50}{S}; P_{88} = P_{50} - \frac{50}{S}$$

Finally, the traits listed in Table 1 were available for quantitative trait loci (QTL) analysis.

Wood density and radial growth

All branches used for embolism resistance measurement were cut into 2cm longitudinal pieces and microdensity profile was obtained using X-ray densitometry (Polge, 1966). Two orthogonal (longest and shortest axes,) radial density profiles were obtained by analyzing the scanned images with WinDENDRO software (Guay et al., 1992). Data from these two sections were averaged and the mean used as a new variable. Ring limits (Fig. S1) were determined automatically, checked manually and then corrected with this software. For each genotype, a series of traits (including early and late wood densities and ring widths, see Table 1) was obtained for all the rings and then averaged for statistical and QTL analyses. When replicates (i.e. several branches per genotype) were available, the average was used for subsequent analysis.

Statistical analysis and QTL mapping

Classical descriptive statistics were generated using R (R core team, 2013) for the studied traits measured on the 477 F2 trees, i.e. mean, standard deviation, and coefficient of variation. A correlation matrix was obtained using Pearson correlation coefficients (R package *Hmisc*) to build a network (R package *Qgraph*) showing the relationship between variables. For QTL detection we used two software: (i) the multiple QTL mapping procedure implemented in the R package *qtl* (Broman et al., 2003) (function *stepwiseqtl* with the imputation method and a maximum of five QTLs for forward selection). In total, 1000 permutations were performed for each trait with the function *scantwo*, to estimate the type I error rate at genome level. A threshold error rate of 5% was used to define significant QTLs. The 95% Bayesian credible interval for each QTL was calculated using the function *bayesint* with default parameter (Manichaikul et al., 2006). The effects of the QTL as well as the percentage of the explained phenotypic variance were also calculated. Additive (a) and dominance (d) effects of a QTL were also calculated; (ii) the Multiple Interval Mapping (MIM) procedure (Jansen and Stam, 1994) implemented in MultiQTLV2.6 (Haifa, Israel, 2005; [<http://www.multiqtl.com/>]) following the approach described in Marguerit et al. (2014). Since both methods provided similar outcomes we only report here the result from the R package.

Results and discussion

Genotyping (248 SNPs mapped on the 12 linkage groups of the genetic map of accession H12) and phenotyping data (see material and method section) are provided in Table S1 for the 477 offspring of the F2 mapping population. The average, variance and distribution of each trait are presented in Table S2 and Fig. S2.

Hydraulic safety is not linked with wood density

Phenotypic variability and correlations among traits

All vulnerability curves for each genotype follow a sigmoid shape with the air-entry (P_{12}) around -3.15 ± 0.27 MPa as illustrated in Fig. 1A. The mean and standard deviation of the main variable P_{50} is -3.70 ± 0.21 MPa, i.e. similar to what was found in an earlier study (Lamy et al., 2011). The total amount of variability observed in our study is much larger than the variability observed across ten different Maritime pine ecotypes (Lamy et al., 2014) (Fig. 1B). However, as already observed for this species, the coefficient of variation for P_{50} is remarkably low ($CV_P=5.6\%$

against 10.6% for mean wood density, 14.5% for mean ring width and 26.8% for mean ring area). The slope of the vulnerability curve (S) was relatively steep (99% MPa^{-1}) compared to those reported for conifer species in general but in the range of values observed for pine species (Delzon et al., 2010). As slope values $>50\%$ MPa^{-1} indicate a very fast rate of embolism, maritime pine can be considered as a vulnerable species to runaway embolism. However, the coefficient of variation of S is high ($\text{CV}_P=29\%$) showing the large variability between genotypes. The phenotypic correlations among variables (Table S3) show that the four variables of hydraulic safety (P_{50} , P_{12} , P_{88}) are positively correlated to each other. In particular there is a strong correlation between P_{50} and P_{12} ($r=0.80$) or P_{88} ($r=0.77$). Besides, there is no significant correlation between P_{50} and $Slope$. CV_P for mean wood density was around 10.5%, which agrees with a previous investigation with different Maritime pine ecotypes (Lamy et al., 2012). Both early- and late-wood densities were on the same range and highly correlated ($r=0.8$) to each other. CV_P for mean ring area was 14% and that for mean ring width was twice higher. Both traits were highly correlated ($r=0.97$). However, wood density- and growth-related traits segregated almost independently in two different clusters (Fig. 2 and Table S3) suggesting the absence of significant trade-off between carbon allocated to cell wall thickening (density) on the one hand and cell division and expansion (growth) on the other hand.

Lack of relationship between embolism resistance and wood density

No significant correlations between embolism resistance (P_{12} , P_{50} , P_{88} , S) and wood density (EWD_av, LWD_av, WD_av) were found in the present study, suggesting independence between these traits (Fig. 2 and Table S3). This result suggests that in this family, the more embolism resistance genotypes did not have the denser wood. This finding that has already been observed at the intraspecific level (Lamy et al., 2012) contradicts the main theory suggesting that dense wood is supposed to be less vulnerable due to a smaller lumen, higher conduit wall thickness and reduced endangerment through cell wall implosion (Hacke et al., 2001; Jacobsen et al., 2005). A mechanical constraint on tracheid anatomy was the premise of this theory as embolism resistant species generally experienced more negative xylem water potential and require to build tracheids with a higher thickness to span ratio to resist mechanical stresses. Indeed, Bouche et al. (2014) reported a significant trade-off between hydraulic and mechanical safety across conifer species with a strong link between increasing embolism resistance and increasing thickness to span ratio of tracheids. In addition, wood density is well known to correlate with the thickness to span ratio

of conifers (Pitterman et al., 2006). However, a direct link between wood density and embolism resistance at both the inter- and intra-specific levels is still not convincingly confirmed. In conifers, those relationships were found across species (Hacke et al., 2001; Hacke and Jansen, 2009) and within a given species (Rosner et al., 2014; Rosner, 2017) but several studies reported lack of such correlations (Martínez-Vilalta et al., 2009; Corcuera et al., 2011; Lamy et al., 2012; Gleason et al., 2016). On a physiological perspective, the lack of link between wood density and embolism resistance in maritime pine is not surprising given the mechanistic role of the pit membrane in preventing embolism formation (Cochard et al., 2009; Delzon et al., 2010; Pittermann et al., 2010). Indeed, the best trait explaining the variability in embolism resistance across species is the ratio of torus to pit aperture diameter, so-called torus overlap (Bouche et al., 2014). More insights into the morphological, physiological or developmental mechanisms underlying the observed variability in cavitation resistance could then follow an approach consisting in finely studying the two extremes of the phenotypic distribution as reported for oak for water-use efficiency (Roussel et al., 2009a, b).

QTLs for cavitation resistance and wood density do not coincide

Following the mapping procedure described in Bartholomé et al. (2016), the linkage map of accession H12 included 248 SNP markers on 12 linkage groups (Table S1), corresponding to the haploid number chromosomes of the maritime pine genome. The length of the map was 1,754cM (146cM/linkage group on average) resulting into a density of 1 SNP every 7 cM, a density that is considered appropriate for QTL mapping.

No study has yet identified consistent QTLs for xylem water transport safety because it requires a high-throughput phenotyping method in a dedicated facility to measure a sufficient number of genotypes and reach a satisfactory statistical detection power. In this study we took advantage of the Cavitplace, a platform for hydraulic trait measurement at the University of Bordeaux (Talence, France) to screen the highest number ever of full-sibs with a single family, in order to dissect the genetic architecture of these traits and provide novel information regarding its genetic architecture and positional coincidence with wood density QTLs. As a result, highly significant QTLs were detected for all tested traits with high statistical support.

Considering the variability in the number of rings between the F2s (ranging from 2 to 5), we first analyzed the effect of this variable on embolism resistance traits. We only considered the

87, 328, 61 genotypes respectively with 2, 3 and 4 rings, because only one genotype presented 5 rings. Analysis of variance shows that the number of rings does not affect P_{12} and P_{50} (P -values > 0.05), while it barely affects P_{88} (P -value=0.0217) and had a significant impact ($P= 0.0018$) on S (Table S4). Therefore, while the full dataset (477 genotypes) was kept for QTL analysis for all the traits, we also performed the analysis for the 328 genotypes with only 3 rings for S . Since the results did not differ between both analyses we only report here the results for the whole dataset. For P_{50} , five QTLs were detected on five different linkage groups (LG1, LG4, LG6, LG9, and LG10, Fig. 3). The phenotypic variance explained by each QTL (PEV) ranged from 3% to 6.5%, altogether accounting for 23% of the phenotypic variation (Table S5). Noteworthy, the same QTLs were detected with the restricted dataset based on three rings (data not shown). For P_{88} , four of these five QTLs were detected on the same chromosomal location of LG1, LG4, LG9 and LG10, and an additional one was detected in LG7. For both traits the strongest QTL (up to 10.3% of PEV) coincided on LG4. Two QTLs on LG6 and LG8 were detected for P_{12} . The former overlapped the confidence interval (CI) of the P_{50} -QTL on the same chromosome. Finally, a single significant QTL was found in LG4 for S with CI overlapping that of P_{50} and P_{88} .

For mean wood density three QTLs were detected on LG1, LG2 and LG8 (Fig. 3). The overall PEV amounted to 10.3% (Table S5). The QTL on LG1 was detected for both early wood (EW) and late wood (LW) on the same position, while that of LG2 and LG8 were found for EW only. These results suggest common genetic factor controlling basic wood properties whatever the type of wood considered, as well as specific molecular mechanisms differentiating EW vs. LW. Thus, none of the QTL for cavitation resistance coincided with that of wood density.

Conclusion

Comparison of QTL locations between embolism resistance and wood density related traits failed to detect co-locating regions. This result is in agreement with the absence of phenotypic correlations between both traits in this family. Whether this result would still hold true in other genetic backgrounds will have to be investigated, but the lack of phenotypic and genetic correlations across families of five ecotypes (Lamy et al., 2012) strongly support the lack of causality between these two traits in this species. Non-competing physiological mechanisms for wood density and cavitation resistance is interesting from a breeding perspective, as it leaves the possibility of selecting genotypes with high wood density and resistant to severe drought. This

pattern should however be assessed at the trunk level where embolism resistance was found to be related to wood density, as reported by Rosner et al. (2017) in Norway spruce trunkwood at different cambial age.

Table 1: List of traits measured in this study.

Measured with the Cavitron	Abbreviation
Xylem pressure inducing 50% loss of hydraulic conductance (MPa) (<i>hydraulic safety = drought tolerance</i>)	<i>P₅₀</i>
Xylem pressure inducing 12% loss of hydraulic conductance (MPa) (<i>air-entry</i>)	<i>P₁₂</i>
Xylem pressure inducing 88% loss of hydraulic conductance (MPa)	<i>P₈₈</i>
Slope of the vulnerability curve (<i>hydraulic safety = embolism rates</i>)	<i>Slope</i>
Measured using X-ray microdensitometry	
Late wood density (kg.m ⁻³)	<i>LWD_{av}</i>
Late wood ring surface (mm ²)	<i>LWarea_{av}</i>
Late wood ring width (mm)	<i>LWwidth_{av}</i>
Late wood percent (%)	<i>LWpercent_{av}</i>
Early wood density (kg.m ⁻³)	<i>EWD_{av}</i>
Early wood ring surface (mm ²)	<i>EWarea_{av}</i>
Early wood ring width (mm)	<i>EWwidth_{av}</i>
Mean wood density (kg.m ⁻³)	<i>WD_{av}</i>
Mean ring surface (mm ²)	<i>area_{av}</i>
Mean ring width (mm)	<i>width_{av}</i>
Mean heterogeneity=Average of LWD - Average of EWD (kg.m ⁻³)	<i>het_{av}</i>
Measured with a caliper	
Branch diameter (mm)	<i>branch_{diam}</i>

Figure captions:

Figure 1: Vulnerability curves of the F2 trees (**A**, red line: fitted mean, grey dark: fitted standard error, light grey: individual raw data) together with vulnerability curves obtained from a range of maritime pine ecotypes (**B**, Cazorla, Landais, Porto-vecchio, Tamjout correspond to four natural populations of the Saint Alban common garden described in Harfouche and Kremer, (2000) and, Bayubas de Abajo, Coca, Erdeven, Mimizan, Oria, San-Cipriano, Tamrabta are seven natural populations from the common garden described in Lamy et al. 2011).

Figure 2: Network of Pearson correlation coefficients between the studied traits. Each node represents a variable and each edge a correlation. The width of the edge indicates the strength of the correlation. Only significant coefficients (P -values < 0.05) are shown here. The negative correlations are in gray and the positive correlations in red.

Figure 3: QTL map for cavitation and wood density related traits. QTLs are shown along the 12 linkage groups (1 to 12) for the traits listed in Table 1. The confidence interval of each QTL is indicated in orange with a color gradient ranging from the peak to the end of the confidence interval.

Supplementary information

Table S1: Genotypic and phenotypic data.

Table S2: Descriptive statistics for the traits listed in table 1.

Table S3: Significant Pearson correlation coefficients for cavitation (P_{50} , P_{88} , P_{12} , *Slope*), branch diameter (*branch_diam*), wood density (LWD_{av} , EWD_{av} , het_{av} , WD_{av}) and radial growth ($LWarea_{av}$, $LWwidth_{av}$, $LWpercent_{av}$, $EWarea_{av}$, $EWwidth_{av}$, $area_{av}$) related traits in the F2 mapping population (P -value <0.05).

Table S4: A. Analysis of variance (cavitation related traits as dependent variable and number of rings as independent variable). **B.** Descriptive statistics for groups of sample with two rings, three rings or four rings.

Table S5: QTL results from the analysis of the F2 mapping population for the traits listed in table 1. Position (cM) of QTLs on linkage groups (LG) with flanking SNP markers listed in table S1, \log_{10} likelihood ratio scores (LOD), genome wide P -value, 95% Bayesian credible intervals with marker interval, % variance explained by each QTL (PEV), and total PEV, genotypic class means and standard errors, additive-dominance effects are presented.

Figure S1: X-ray image of the radial section of a maritime pine branch. The dark grey color of early wood is related to its lower density compare to late wood indicated by a narrower whiter area.

Figure S2: Distribution of the studied traits listed in table 1.

References

- Allen CD, Macalady AK, Chenchouni H, Bachelet D, McDowell N, Vennetier M, Gonzalez P et al (2010) A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology Management* 259:660-684.
- Anderegg WR, Klein T, Bartlett M, Sack L, Pellegrini AF, Choat B, Jansen S (2016) Meta-analysis reveals that hydraulic traits explain cross-species patterns of drought-induced tree mortality across the globe. *Proceedings of the National Academy of Sciences* 113:5024-5029.
- Anderegg WR, Flint A, Huang CY, Flint L, Berry JA, Davis FW, Field CB et al (2015) Tree mortality predicted from drought-induced vascular damage. *Nat Geo* 8:367-371.
- Bahn M, Reichstein M, Dukes JS, Smith MD, McDowell NG (2014) Climate–biosphere interactions in a more extreme world. *New Phytology* 202 :356-359.
- Bartholomé J, van Heerwaarden J, Bink MCAM, Chancerel E, Boury C, Lesur I, Isik F, Bouffier L, Plomion C (2016) Linkage and association mapping for two major traits of the maritime pine breeding program: tree height and stem straightness. *PlosOne* 11:e0165323.
- Bouche PS, Larter M, Domec JC, Burlett R, Gasson P, Jansen S, Delzon S (2014) A broad survey of hydraulic and mechanical safety in the xylem of conifers. *Journal of Experimental Botany* 65:4419-4431.
- Brodribb TJ, Bowman DJ, Nichols S, Delzon S, Burlett R (2010) Xylem function and growth rate interact to determine recovery rates after exposure to extreme water deficit. *New Phytology* 188:533-542.
- Brodribb TJ, Cochard H (2009) Hydraulic failure defines the recovery and point of death in water-stressed conifers. *Plant physiology* 149:575-584.
- Broman KW, Wu H, Sen Ś, Churchill GA (2003) R/qtl: QTL mapping in experimental crosses. *Bioinformatics* 19: 889–90.
- Choat B, Jansen S, Brodribb TJ, Cochard H, Delzon S, Bhaskar R, Jacobsen AL et al (2012). Global convergence in the vulnerability of forests to drought. *Nature* 491:752-755.
- Ciais P, Reichstein M, Viovy N, Granier A, Ogee J, Allard V, Aubinet M, Buchmann N, Bernhofer C, Carrara A et al (2005) Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature* 437:529-533.
- Cochard H, Hölttä T, Herbette S, Delzon S, Mencuccini M (2009) New insights into the mechanisms of water-stress-induced cavitation in conifers. *Plant Physiology* 151:949-954.
- Cochard H, Barigah ST, Kleinhentz M, Eshel A (2008) Is xylem cavitation resistance a relevant criterion for screening drought resistance among *Prunus* species?. *Journal of plant physiology* 165: 976-982.

- Cochard H, Damour G, Bodet C, Tharwat I, Poirier M, Améglio T (2005) Evaluation of a new centrifuge technique for rapid generation of xylem vulnerability curves. *Physiologia Plantarum* 124:410-418.
- Cochard H (2002) A technique for measuring xylem hydraulic conductance under high negative pressures. *Plant, Cell & Environment* 25:815-819.
- Corcuera L, Cochard H, Gil-Pelegrin E, Notivol E (2011) Phenotypic plasticity in mesic populations of *Pinus pinaster* improves resistance to xylem embolism (P_{50}) under severe drought. *Trees* 25:1033-1042.
- Dalla-Salda G, Martinez-Meier A, Cochard H, Rozenberg P (2009) Variation of wood density and hydraulic properties of Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) clones related to a heat and drought wave in France. *Forest Ecology and Management* 257:182-189.
- Delzon S, Urli M, Samalens JC, Lamy JB, Lischke H, Sin F, Porté AJ et al (2013) Field evidence of colonisation by Holm oak, at the northern margin of its distribution range, during the Anthropocene period. *PloS one* 8:e80443.
- Delzon S, Douthe C, Sala A, Cochard H (2010) Mechanism of water-stress induced cavitation in conifers: bordered pit structure and function support the hypothesis of seal capillary-seeding. *Plant, Cell & Environment* 33:2101-2111.
- Domec JC, Gartner BL (2001) Cavitation and water storage capacity in bole xylem segments of mature and young Douglas-fir trees. *Trees-Structure and Function* 15:204-214.
- Gleason SM, Westoby M, Jansen S, Choat B, Hacke UG, Pratt RB, Bhaskar R, Brodribb TJ, Bucci SJ, Cao KF, Cochard H, Delzon S, Domec J-C, Cao KF, Feild TS, Jacobsen AL, Jacobsen AL, Lens F, Maherali H, Martínez-Vilalta J, Mayr S, McCulloh KA, Mencuccini M, Mitchell PJ, Morris H, Nardini A, Pittermann J, Plavcová L, Schreiber SG, Sperry JS, Wright IJ, Zanne AE. (2016) Weak tradeoff between xylem safety and xylem-specific hydraulic efficiency across the world's woody plant species. *New Phytologist* 209:123-136.
- Guay R, Gagnon R, Morin H (1992) A new automatic and interactive tree ring measurement system based on a line scan camera. *Forestry Chronicle* 68:138-141
- Hacke UG, Sperry JS, Pockman WT, Davis SD, McCulloh KA (2001) Trends in wood density and structure are linked to prevention of xylem implosion by negative pressure. *Oecologia* 126: 457-461.
- Hacke UG, Jansen S (2009) Embolism resistance of three boreal conifer species varies with pit structure. *New Phytologist* 182:675-686.
- Hacke U, Sperry J (2001) Functional and ecological xylem anatomy. *Perspectives in Plant Ecology, Evolution and Systematics* 4:97-115.
- Harfouche A, Kremer A (2000) Provenance hybridization in a diallel mating scheme of maritime pine (*Pinus pinaster*). I. Means and variance components. *Canadian journal of forest research* 30: 1-9.

- IPCC Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, Chatterjee M, Ebi KL, Estrada YO, Genova RC, Girma B, Kissel ES, Levy AN, MacCracken S, Mastrandrea PR, White LL (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1132 pp.
- Jacobsen AL, Ewers FW., Pratt RB, Paddock III WA, Davis SD (2005) Do Xylem Fibers Affect Vessel Cavitation Resistance? *Plant Physiology* 139:546-556.
- Jansen RC, Van Ooijen JW, Stam P, Lister C, Dean C (1995) Genotype-by-environment interaction in genetic mapping of multiple quantitative trait loci. *Theoretical and Applied Genetics*.91:33-37.
- Kremer A, Potts BM, Delzon S (2014) Genetic divergence in forest trees: understanding the consequences of climate change. *Functional Ecology* 28:22-36.
- Lamy JB, Delzon S, Bouche P, Alia R, Vendramin GG, Cochard H, Plomion C (2014) Limited genetic variability and phenotypic plasticity for cavitation resistance in a Mediterranean pine. *New Phytology* 201:874-86.
- Lamy JB, Lagane F, Plomion C, Cochard H, Delzon S (2012) Micro-evolutionary patterns of juvenile wood density in a pine species. *Plant Ecology* 213:1781-1792.
- Lamy JB, Bouffier L, Burlett R, Plomion C, Cochard H, Delzon S (2011) Uniform selection as the primary evolutionary force of cavitation resistance across a species range. *PlosOne* 6: e23476.
- Kavanagh KL, Bond BJ, Aitken SN, Gartner BL, Knowe S (1999) Shoot and root vulnerability to xylem cavitation in four populations of Douglas-fir seedlings. *Tree Physiology* 19:31-37.
- Larter M, Pfautsch S, Domec JC, Trueba S, Nagalingum N, Delzon S (2017) Aridity drove the evolution of extreme embolism resistance and the radiation of conifer genus *Callitris*. *New Phytologist* 215: 97-112.
- Lauri PE, Gorza O, Cochard H, Martinez S, Celton J, Ripetti V, Lartaud M, Bry X, Trottier C, Costes E (2011) Genetic determinism of anatomical and hydraulic traits within an apple progeny. *Plant, Cell & Environment*, 34:1276-1290.
- Liebhart R, Kellerhals M, Pfammatter W, Jertmini M, Gessler C (2003) Mapping quantitative physiological traits in apple (*Malus× domestica* Borkh.). *Plant molecular biology* 52:511-526.
- Lindner M, Fitzgerald JB, Zimmermann NE, Reyer C, Delzon S, van der Maaten E, Suckow F et al (2014) Climate change and European forests: what do we know, what are the uncertainties, and what are the implications for forest management?. *Journal of environmental management* 146:69-83.

- Lindner M, Maroschek M, Netherer S, Kremer A, Barbati A, Garcia-Gonzalo J, Lexer MJ et al (2010) Climate change impacts, adaptive capacity, and vulnerability of European forest ecosystems. *Forest ecology and management* 259:698-709.
- Maherali H, Pockman WT, Jackson RB (2004) Adaptive variation in the vulnerability of woody plants to xylem cavitation. *Ecology* 85: 2184-2199.
- Manichaikul A, Dupuis J, Sen S, Broman KW (2006) Poor Performance of Bootstrap Confidence Intervals for the Location of a Quantitative Trait Locus. *Genetics* 174:481-9.
- Marguerit E, Bouffier L, Chancerel E, Costa P, Lagane F, Guehl JM et al (2014) The genetics of water-use efficiency and its relation to growth in maritime pine. *Journal of Experimental Botany* 65:4757-68.
- Martínez-Vilalta J, Cochard H, Mencuccini M, Sterck F, Herrero A, Korhonen JF, Llorens P, Nikinmaa E, Nolè A, Poyatos R, Ripullone F, Sass-Klaassen U, Zweifel R (2009) Hydraulic adjustment of Scots pine across Europe. *New Phytologist* 184:353-364.
- Meehl GA, Tebaldi C (2004) More intense, more frequent, and longer lasting heat waves in the 21st century. *Science* 305:994-997.
- Michaelian M, Hogg EH, Hall RJ, Arsenault E (2011) Massive mortality of aspen following severe drought along the southern edge of the Canadian boreal forest. *Global Change Biology* 17:2084-2094.
- Pammenter NV, Van der Willigen C (1998) A mathematical and statistical analysis of the curves illustrating vulnerability of xylem to cavitation. *Tree physiology* 18: 589-593.
- Piao SL, Wang XH, Ciais P, Zhu B, Wang T, Liu J (2011) Changes in satellite-derived vegetation growth trend in temperate and boreal Eurasia from 1982 to 2006. *Global Change Biology* 17:3228-3239.
- Pittermann J, Stuart SA, Dawson TE, Moreau A (2012) Cenozoic climate change shaped the evolutionary ecophysiology of the Cupressaceae conifers. *Proceedings of the National Academy of Sciences* 109:9647-9652.
- Pittermann J, Choat B, Jansen S, Stuart SA, Lynn L, Dawson TE (2010) The relationships between xylem safety and hydraulic efficiency in the Cupressaceae: the evolution of pit membrane form and function. *Plant Physiology* 153:1919-1931.
- Pittermann J, Sperry JS, Wheeler JK, Hacke UG, Sikkema EH (2006) Mechanical reinforcement of tracheids compromises the hydraulic efficiency of conifer xylem. *Plant, Cell & Environment* 29:1618-1628.
- Plomion C, Bartholomé J, Bouffier L, Brendel O, Cochard H, de Miguel M, Delzon S, Gion JM, González-Martínez SC, Guehl JM, Lagravelle H, Le Provost G, Marguerit E, Porté A (2016) The genetic basis of adaptation to water deficit in trees through the examination of water use efficiency and cavitation resistance: maritime pine as a case study. *Journal of Plant Hydraulics* 3:e-008.

- Polge H (1966) Etablissement des courbes de variation de la densité du bois par exploration densitométrique de radiographie d'échantillons prélevés à la tarière sur des arbres vivants. *Annals of Forest Science* 23:1-206.
- R Core Team (2013) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Rosner S (2017) Wood density as a proxy for vulnerability to cavitation: Size matters. *Journal of Plant Hydraulics* 4:e001.
- Rosner S, Světlík J, Andreassen K, Børja I, Dalsgaard L, Evans R, Karlsson B, Tollefsrud MM, Solberg S. 2014. Wood density as a screening trait for drought sensitivity in Norway spruce. *Canadian Journal of Forest Research* 44:154-161.
- Roussel M, Dreyer E, Montpied P, Le-Provost G, Guehl J-M, Brendel O. 2009a. The diversity of ^{13}C isotope discrimination in a *Quercus robur* full-sib family is associated with differences in intrinsic water use efficiency, transpiration efficiency, and stomatal conductance. *Journal of Experimental Botany* 60:2419-2431.
- Roussel M, Le Thiec D, Montpied P, Ningre N, Guehl J-M, Brendel O. 2009b. Diversity of water use efficiency among *Quercus robur* genotypes: contribution of related leaf traits. *Annals of Forest Science* 66:408.
- Urli M, Porté AJ, Cochard H, Guengant Y, Burrett R, Delzon S (2013) Xylem embolism threshold for catastrophic hydraulic failure in angiosperm trees. *Tree physiology* 33:672-683.
- Zhao M, Running SW (2010) Drought-induced reduction in global terrestrial net primary production from 2000 through 2009. *Science* 329:940-943.

Figure 1: Vulnerability curves of the F2 trees (**A**, red line: fitted mean, grey dark: fitted standard error, light grey: individual raw data) together with vulnerability curves obtained from a range of maritime pine ecotypes (**B**, Cazorla, Landais, Porto-vecchio, Tamjout correspond to four natural populations of the Saint Alban common garden described in Harfouche and Kremer, (2000) and, Bayubas de Abajo, Coca, Erdeven, Mimizan, Oria, San-Cipriano, Tamrabta are seven natural populations from the common garden described in Lamy et al. (2011)).





Figure 2: Network of Pearson correlation coefficients between the studied traits. Each node represents a variable and each edge a correlation. The width of the edge indicates the strength of the correlation. Only significant coefficients (P -values < 0.05) are shown here. The negative correlations are in gray and the positive correlations in red.





Figure 3: QTL map for cavitation and wood density related traits. QTLs are shown along the 12 linkage groups (1 to 12) for the traits listed in Table 1. The confidence interval of each QTL is indicated in orange with a color gradient ranging from the peak to the end of the confidence interval.



Supplementary information

Table S1: Genotypic and phenotypic data. (Data could be provided on demand)

Table S2: Descriptive statistics for the traits listed in table 1.

Traits	P50	P12	P88	slope	branch_diam	LWD_av	LWarea_av	LWwidth_av	LWpercent_av	EWD_av	EWarea_av	EWwidth_av	het_av	WD_av	area_av	width_av
nbr.val	465	465	465	465	465	466	466	466	466	466	466	466	466	466	466	466
nbr.na	12	12	12	12	12	11	11	11	11	11	11	11	11	11	11	11
min	-4.251	-3.771	-5.28	38.428	4.88	0.284	2.919	0.815	19.075	0.216	0.811	0.429	0.007	0.261	2.662	0.859
max	-3.178	-2.148	-3.515	234.635	9.32	0.549	19.783	2.509	85.75	0.444	17.196	2.261	0.154	0.483	14.415	2.095
range	1.073	1.623	1.765	196.207	4.44	0.265	16.864	1.694	66.675	0.228	16.385	1.832	0.147	0.222	11.753	1.236
median	-3.701	-3.181	-4.23	96.505	7.58	0.397	8.393	1.588	48.247	0.314	7.4145	1.47	0.083	0.356	8.12	1.549
mean	-3.699933	-3.147105	-4.25277	98.972955	7.5146882	0.3972597	8.6373519	1.6048798	48.80014807	0.3144635	7.6402768	1.4719742	0.0827961	0.3558455	8.1388391	1.5384292
SE.mean	0.0096854	0.012523	0.0122293	1.3437006	0.0417026	0.0019785	0.1332118	0.0130624	0.510883698	0.0016614	0.1379791	0.0139867	0.0010808	0.0017456	0.1011144	0.0103446
CI.mean.0.95	0.0190327	0.0246087	0.0240317	2.6404924	0.0819493	0.0038879	0.2617717	0.0256686	1.003926684	0.0032648	0.2711398	0.027485	0.0021239	0.0034302	0.1986977	0.020328
var	0.0436201	0.0729233	0.0695435	839.57209	0.8086831	0.0018241	8.2693533	0.0795114	121.6270032	0.0012863	8.8718139	0.0911629	0.0005444	0.0014199	4.7644366	0.0498672
std.dev	0.2088543	0.2700432	0.263711	28.97537	0.8992681	0.04271	2.8756483	0.2819777	11.02846332	0.0358651	2.978559	0.3019319	0.0233316	0.0376815	2.182759	0.2233096
coef.var	-0.056448	-0.085807	-0.062009	0.2927605	0.1196681	0.1075116	0.3329317	0.1757002	0.225992415	0.1140517	0.3898496	0.2051204	0.2817961	0.105893	0.2681905	0.1451543
skewness	-0.090702	0.564199	-0.407894	0.9213723	-0.261898	0.1014238	0.4106431	0.0115472	0.221617939	-0.018864	0.3878128	-0.223004	-0.095484	0.0119548	0.0885417	-0.230671
skew.2SE	-0.400526	2.4914237	-1.801204	4.068651	-1.156505	0.4483516	1.8152787	0.0510452	0.979678794	-0.08339	1.7143558	-0.985805	-0.422096	0.0528472	0.3914054	-1.019698
kurtosis	-0.351155	0.2292819	0.355305	1.6349223	-0.5745	0.2743173	-0.002732	-0.240316	0.020341513	0.2689728	-0.095225	0.1482016	0.1797559	0.349072	-0.156857	0.0597896
kurt.2SE	-0.776966	0.5073095	0.7861486	3.6174321	-1.271139	0.6076003	-0.00605	-0.532289	0.045055527	0.5957626	-0.210919	0.3282597	0.3981512	0.7731787	-0.347431	0.1324313
normtest.W	0.9959744	0.9783594	0.9895499	0.9567244	0.9856567	0.9940065	0.9851837	0.9969886	0.995750882	0.9938627	0.9873573	0.9924261	0.9963251	0.9916166	0.9972077	0.9947861
normtest.p	0.2843538	2.12E-06	0.0021481	1.92E-10	0.0001506	0.0632142	0.0001091	0.548108	0.240680416	0.0564674	0.0004552	0.0183482	0.3608435	0.0098387	0.6186437	0.1161574

Table S3: Significant Pearson correlation coefficients for cavitation (P₅₀, P₈₈, P₁₂, *Slope*), branch diameter (branch_diam), wood density (LWD_av, EWD_av, het_av, WD_av) and radial growth (LWarea_av, LWwidth_av, LWpercent_av, EWarea_av, EWwidth_av, area_av) related traits in the F2 mapping population (*P*-value <0.05).

	P50	P12	P88	slope	branch_diam	LWD	LWarea	LWwidth	LWpercent	EWD	EWarea	EWwidth	het	WD	area	width
P50																
P12	0.8															
P88	0.77	0.29														
slope		-0.53	0.58													
branch_diam	0.33	0.31	0.23													
LWD					-0.1											
LWarea	0.13	0.13			0.41	0.11										
LWwidth	0.12	0.12			0.4	0.12	0.99									
LWpercent					-0.09	0.27	0.43	0.45								
EWD					-0.16	0.8			0.13							
EWarea	0.12	0.16		-0.1	0.43	-0.11	0.13	0.12	-0.67	-0.1						
EWwidth	0.13	0.16		-0.09	0.43		0.17	0.16	-0.68		0.97					
het						0.55	0.26	0.29	0.26							
WD					-0.14	0.96			0.21	0.93	-0.11		0.3			
area	0.18	0.2	0.11		0.57		0.72	0.71	-0.15	-0.11	0.75	0.75	0.14			
width	0.17	0.19	0.1	-0.09	0.55		0.74	0.74	-0.16	-0.11	0.71	0.76	0.19		0.97	

Table S 4: A. Analysis of variance (cavitation related traits as dependent variable and number of rings as independent variable). **B.** Descriptive statistics for groups of sample with two rings, three rings or four rings.

A

Source P ₁₂	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	9	0.31261047	0.03473450	0.47	0.8970
Error	467	34.77562535	0.07446601		
Corrected Total	476	35.08823583			

Source P ₅₀	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	9	0.31957841	0.03550871	0.81	0.6069
Error	467	20.46539659	0.04382312		
Corrected Total	476	20.78497500			

Source P ₈₈	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	9	1.35033274	0.15003697	2.19	0.0217
Error	467	32.01024837	0.06854443		
Corrected Total	476	33.36058112			

Source S _{LOPE}	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	9	22147.2872	2460.8097	2.98	0.0018
Error	467	385726.7682	825.9674		
Corrected Total	476	407874.0554			

B

Traits	Mean	Std Deviation	Min	Max	N
P ₁₂ (2 rings)	-3.05	0.29	-3.60	-2.13	87
P ₁₂ (3 rings)	-3.21	0.27	-3.77	-2.15	318
P ₁₂ (4 rings)	-3.16	0.26	-3.64	-2.33	61
P ₅₀ (2 rings)	-3.70	0.21	-4.18	-3.26	87
P ₅₀ (3 rings)	-3.64	0.21	-4.25	-3.18	328
P ₅₀ (4 rings)	-3.72	0.20	-4.24	-3.23	61
P ₈₈ (2 rings)	-4.36	0.26	-5.28	-3.79	87
P ₈₈ (3 rings)	-4.08	0.26	-5.10	-3.52	318
P ₈₈ (4 rings)	-4.27	0.27	-4.99	-3.78	61
S _{LOPE} (2 rings)	85.85	21.38	38.30	138.35	87
S _{LOPE} (3 rings)	141.62	76.02	81.37	241.95	328
S _{LOPE} (4 rings)	98.9	28.85	49.69	197.51	61

Table S 5: QTL results from the analysis of the F2 mapping population for the traits listed in table 1. Position (cM) of QTLs on linkage groups (LG) with flanking SNP markers listed in table S1, log10 likelihood ratio scores (LOD), genome wide *P*-value, 95% Bayesian credible intervals with marker interval, % variance explained by each QTL (PEV), and total PEV, genotypic class means and standard errors, additive-dominance effects are presented.

	Trait	n	LG	Position	LOD	BCI	PEV	PEV tot	Add effect	Dom effect
Cavitation	P50	465	1	73	6.97	61 - 94	4.25	23.05	0.06044	0.03084
	P50	465	4	43	9.04	17 - 54.38	6.46		0.07314	0.02841
	P50	465	6	55	6.04	46 - 63	4.07		-0.06252	0.01396
	P50	465	9	100.69	4.07	88 - 104	3.05	7.19	-0.04139	-0.04669
	P50	465	10	148.29	7.15	140 - 158	4.72		-0.05809	-0.02975
	P12	465	6	49	5.48	45 - 60	4.84	26.13	-0.08438	-0.02451
	P12	465	8	154.57	3.49	134 - 157.85	2.51		0.06085	-0.00139
	P88	465	1	56	8.4	51 - 67	5.58		0.07673	0.06126
	P88	465	4	25	14.08	2 - 52	10.3		0.12273	0.03421
	P88	465	7	28	4.67	23 - 46	3.26		0.06004	0.02766
	P88	465	9	100.69	4.3	87 - 105	2.72		-0.04918	-0.05549
	P88	465	10	148	6.56	138 - 157	4.7		-0.08043	-0.03969
slope	465	4	4	4.77	0 - 32	4.34	9.08926	0.32		
Wood density	LWD_av	466	1	126	4.86	121 - 142	3.56	11.75	-0.01098	-0.00651
	EWD_av	466	1	139	5.18	123 - 145	4.85		-0.01152	0.00046
	EWD_av	466	2	26.32	4.41	18 - 40	3.93		0.01003	-0.00439
	EWD_av	466	8	97.56	4.01	90 - 121	3.34	10.33	0.00897	0.00382
	EWwidth_av	466	12	109.69	3.84	100 - 116	2.95		0.06914	-0.04371
	WD_av	466	1	139	5.72	123 - 143	4.33		-0.01121	-0.00074
	WD_av	466	2	26	4.2	11 - 40	3.34		0.00976	-0.00391
	WD_av	466	8	97.56	3.48	88 - 126	2.85		0.00873	0.00534

Figure S 1: X-ray image of the radial section of a maritime pine branch. The dark grey color of early wood is related to its lower density compare to late wood indicated by a narrower whiter area.










Figure S 2: Distribution of the studied traits listed in table 1.




branch_diam




LWD




LWarea




LWwidth




LWpercent




EWD







EWarea



EWwidth





Chapter 3:

Intra-specific variability of the physiological response to water stress in a full-sib Maritime pine family (*in preparation*)

Intra-specific variability of the physiological response to water stress in a full-sib Maritime pine family

This work was performed in the framework of the ANR project WUEtree - Understanding the genetic and environmental determinisms of water use efficiency to improve sustainability of forest tree plantations (Project ID: ANR-13-ADAP-0012) - as a collaborative project with Marina de Miguel Vega.

Abstract

In the context of climate change and increasing water stress due to drought events, tree breeders are considering new selection criteria to dispose of genotypes that can resist droughts and maintain productivity under changing environmental conditions. Prior to such a program, there is an urgent need to assess the intra-specific variability of functional traits related to survival and growth. We performed a multi-trait analysis of leaf and xylem functioning including seasonal measurements of photosystem II fluorescence, gas exchange and stem hydraulic traits on a full-sib family of provenance hybrid of maritime pine (*Pinus pinaster* Ait.) established in Cestas (France). Xylem vulnerability to embolism was assessed on 480 individuals of inter “Morocco x Landes” and “Corsica x Landes” families and leaf functioning traits were performed on a subsample consisting of 108 individuals from the “Morocco x Landes” family. The mean P_{50} values varied from -3.2 MPa to -4.2 MPa and were in average of -3.6 and -3.8 MPa for “Corsica x Landes” and “Morocco x Landes” families, respectively. The efficiency of both photosystem II and stomatal conductance declined with increasing seasonal water stress, resulting in a rapid decrease in assimilation rate. We found no evidence of trade-off between xylem resistance to embolism (P_{50}) and leaf functioning traits indicating the possibility to consider selection for both efficient and resistant genotypes in future.

Introduction

Ongoing climate change is rapid compared to past climate change questioning the ability of long-lived organisms such as trees to adapt to the changes. For example, the rate of warming increased from 0.097 ° C per decade in the early 20th century (1901-1950) compared to 0.254 ° C between 1979 and 2012 (Hartmann et al. 2013). Watering regimes are also expected to increase in intensity and number of drought events (Meel and Tabaldi 2004). Maintaining species in the new environment will be possible provided populations have adaptive capacities to withstand growing constraints (Davis and Shaw 2001). To assess the adaptive capacity of trees to climate change or to select trees that are better suited to new conditions, the phenotypic and genetic diversity of many individuals should be assessed at inter- and intra-specific levels.

The study of the intra-specific variability of ecophysiological traits is less developed on forest trees (Olivas-García et al 2000, Gutierrez et al., 2002, Bigras 2005) compared to fruit trees (Petridis et al., 2012, Pettigrew and Turley 1998, Sofo et al 2009, Viljevac et al., 2013). Forest studies have focused on morphological features (SLA, leaf shape) and growth (e.g. Martinez Meier et al., 2008). However, in order to understand the variability of responses in a changing environment, to identify or select suitable genotypes or populations that are resistant to increasing stresses and / or capable of maintaining efficient functioning in their new environment, a detailed analysis of the functioning of forest trees is paramount.

In the analysis of the resistance to water stress, recent developments have demonstrated that xylem vulnerability to cavitation is a good indicator of the resistance of species to drought (Brodribb et al. 2010, Choat et al. 2012, Urli et al. 2013). At the intra-specific level, first analyses evaluating phenotypic and genotypic variability of cavitation resistance (Lamy et al., 2012, 2014) suggested that it was possible to identify individuals more resistant to water stress. Apart from resistance to drought and cavitation, water and carbon plant functioning in response to drought is of interest to investigate in order to select genotypes that can maintain some production functions under changing environments. Reduction of carbon assimilation in response to water stress is resulting from either stomatal closure or biochemical limitations. Both limitations are changing with the magnitude and duration of water stress, the biochemical limitation inducing 50% loss of carbon assimilation under severe stress (Grassi and Magnani 2005). The maintenance of photosynthetic capacities in response to water stress was identified as an adaptive character by comparing two *Nothofagus* species (Piper and al. 2007). Moreover, at the intra-specific level, families were shown to have distinct responses to drought, maintaining their leaf functioning (photosynthesis, stomatal opening) in the event of water stress, as shown by Bigras (2005) on spruce (*Picea glauca*) or Gilbert et al. (2011) on soybean. A recent study made it possible to distinguish poplar clones by differentiating their photosynthetic capacities (Guo et al., 2010). However, there also appears to be a trade-off between resistance to water stress (cavitation, defoliation) and foliar functional capacities (photosynthesis, water use efficiency, growth) at the inter-specific level (Fu et al., 2012) or intra-specific level (Bigras 2005): the most resistant species or families also demonstrated the most limited growth.

Maritime pine (*Pinus pinaster* Ait.) is a Mediterranean tree species, whose range is expanding from Morocco in the South to the Aquitania region - France in the North and East, to Greece in

the West. Its different provenances are presenting variability in some characters such as more drought resistant in Spain and Morocco, more productive in the Landes region, higher stem straightness in Corsica (Harfouche and Kremer 2000). In France, the species represent 6% of its forested area, 5% of its standing volume and more than 26% of its annual wood production (IGN 2016). A breeding program was initiated in the 60s to improve volume production and stem rectitude (Bouffier et al. 2009). As a consequence, in 2010, 25% of Maritime pine plantations in the Landes de Gascogne forests were corresponding to one of the two selected varieties. In the context of climate change and increasing water stress, breeding is considering new selection criteria related to the plant response to water availability and other strategies such as provenance crossings. Prior to such a program, it is compulsory to analyse the intra-specific variability of functional traits related to water and carbon functioning.

The objective of this study was to investigate the intra-specific variability within a Maritime pine tree family in response to water availability by performing a multi-trait analysis of leaf and xylem functioning through seasonal measurements of photosystem II fluorescence, gas exchange and stem hydraulic properties. Our objective was to evaluate the level of variability of these traits, as well as identifying potential trade-offs between these characters. This will be the first step towards selecting genotypes having the ability to sustain changes in water availability by resisting to drought, while maintaining the same growth rates during pre or post water stress phases.

Materials and Methods

Site and sampling

Analyses were performed on adult *Pinus pinaster* in the field, to avoid working on potted seedling in limited soil environment. The Abiogen trial (Figure 1) is a full-sib family essay planted in 2007 in Cestas (France) as a repeated block design: each block contains full-sibs from either Morocco x Landes, Corsica x Landes or Corsica x Morocco families (see Lagraulet 2015 for more details). The study was performed in spring (April, May) and summer (July) of 2015 and 2016 on two blocks of the Morocco x Landes family, corresponding to 108 trees (except for gas exchanges in summer 2015, performed on 40 individuals).

During each campaign, we sampled and measured 10 to 12 trees per day. Leaf physiological measurements were performed on 1-year-old needles selected on the southern branch of the 2- or 3-year-old whorl. Before sunrise, 4 pairs of needles were collected per tree, put into tubes closed

with a cap, where wet paper placed inside the tube maintained saturated air humidity to avoid plant transpiration.

Lab measurements

Immediately arriving at the lab, one pair of needles was used for predawn water potential measurements using a Scholander pressure bomb. The second pair was used to perform dark adapted fluorescence measurements using a PAM 2100 (Heinz Walz GmbH, Effeltrich, Allemagne) to get maximum F_m and minimum F_0 fluorescence values. The third and fourth pairs were cut under water at the base and put into water, for further gas exchanges and fluorescence measurements. After at least one hour under lamps, fluorescence of light adapted needles was measured using the PAM 2100 to get maximum and minimum levels of fluorescence of light adapted needles, F_m' and F_0' . Using these measurements, fluorescence parameters of maximum efficiency of photosystem II (F_v/F_m), actual efficiency of photosystem II (yield), electron transport rate ETR and non-photochemical quenching (NPQ) were calculated according to equations and definitions of Roháček (2002).

Gas exchanges were performed with a Walz GFS 3000 (Heinz Walz GmbH, Effeltrich, Allemagne) to get steady-state values of Assimilation (A , $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), transpiration (E , $\text{mmol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) and stomatal conductance (g_s , $\text{mmol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$). Conditions within the measuring chamber were air CO_2 concentration of 400 ppm, air temperature of 20°C and relative humidity of 70% to get a constant air VPD, PAR light intensity of 1800 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$. Water use efficiency and instantaneous water use efficiency were calculated as the ratio of A to E or g_s , respectively.

After gas exchange measurements, needles were measured in diameter / length using an electronic caliper (10-2mm) and a ruler (mm) to estimate their leaf area, then placed in an oven to reach constant dry weight (g) to get SLA estimations.

A total of 480 individuals were sampled for xylem hydraulic measurements during two campaigns, first from October 2014 to February 2015 and second from January 2016 to March 2016. In order to measure the same number of rings in each sample, branches were collected on the 2012 whorl to further measured 2 year-old growth units (2013 during the 1st campaign, 2014 during the second campaign). For each individual, a branch fully exposed to the sun with a basal diameter less than 1 cm was collected early in the morning. All needles were immediately removed, branches were labeled and placed in black bags, wrapped in wet paper towels to avoid

transpiration and dessication. They were then kept wet and cool (3°C) until measurement within maximum three weeks after field sampling. Prior to embolism resistance measurement, branches were cut under water to a standard length of 27 cm, and bark was removed with a razor blade.

Xylem vulnerability to embolism was assessed using the CAVITRON, a centrifuge technique following the procedure described by Cochard (Cochard 2002; Cochard et al. 2005) at the high-throughput phenotyping platform for hydraulic traits (CavitPlace, University of Bordeaux, Talence, France). Centrifugal force was used to establish negative pressure in the xylem and to provoke water stress-induced embolism, using a custom-built honeycomb rotor (Sam Precis 2000, Bordeaux, France) mounted on a high-speed centrifuge (Sorvall RC5, USA). This technique enables measurement of the hydraulic conductance of a branch under negative pressure. Xylem pressure (P) was first set to a reference pressure (-0.5 MPa) and maximal conductance (Ks) was determined by measuring the flux of a reference ionic solution (10 mM KCl and 1mM CaCl₂ in deionized water) through the sample. The centrifugation speed was then set to a higher value for 3 min to expose the sample to a more negative pressure. Conductances were measured three times for each step, and the average value was used to compute the percent loss of xylem conductance (PLC in %). The procedure was repeated for at least eight pressure steps with a -0.3 to -0.5 MPa step increment until PLC reached at least 90%. Rotor velocity was monitored with a 10 rpm resolution electronic tachymeter and xylem pressure was adjusted to about -0.02 MPa. We used the Cavi_soft software (version 4.0, for the first campaign and version 5.0 for the second campaign, BIOGECO, University of Bordeaux) for conductance measurements.

Data and statistical analyses

Hydraulic measurements were analysed using vulnerability curves (VC), the percent loss of xylem conductivity plotted as a function of xylem pressure (MPa). A sigmoid function (Pammenter and Van der Willigen, 1998) was fitted to the VC from each sample using the following equation:

$$PLC = \frac{100}{\left[1 + \exp\left(\frac{S}{25} \times (P - P_{50})\right) \right]} \quad (1)$$

where P_{50} (MPa) is the xylem pressure inducing 50% loss of hydraulic conductivity and S (% MPa⁻¹) is the slope of the vulnerability curve at the inflexion point.

The xylem-specific hydraulic conductivity (K_s , m² MPa⁻¹ s⁻¹) was calculated by dividing the maximum hydraulic conductivity measured at low speed by the sapwood area of the sample.

Correlation between ecophysiological traits were performed on the 108 individuals of the Morocco x Landes family that were used for both stem hydraulic measurements and leaf physiological measurements (40 individuals only, for gas exchanges in July 2015). Analyses were performed using the SAS Proc CORR (SAS 9.4, SAS Institute Inc., Cary, NC, USA).

Results

Both years were corresponding to two different water availability conditions (Figure 2): in spring 2015 and 2016, predawn leaf water potential remained high (close to 0) and drop to more negative values during the summer, especially in 2015 where it reached -1.5 MPa. As a result of decreasing water potential, leaf functioning decreased, whatever the trait. Assimilation rate decreased rapidly (Figure 3A) as a consequence of both a decrease in the efficiency of photosystem II (Figure 3B) and in stomatal conductance (Figure 3C). The decrease in efficiency of photosystem II was marked on light adapted needles as the actual yield of light efficiency decrease from 30 to 10%. The potential efficiency of photosystem II, F_v/F_m , slightly decreased with water potential indicating that the photosynthesis apparatus was partly damaged in response to water stress. The decrease in both assimilation and stomatal conductance resulted in a decrease in water use efficiency, in response to water stress (Figure 3D), as assimilation rate presented a steeper decrease in response to decreasing water potential.

All physiological leaf traits presented a large variability between genotypes (Figure 3, Table 1). Traits related to water functioning (E , g_s , WUE, WUE_i) were more variable under water-stressed conditions in July 2015, compared to spring 2015 or 2016, whereas traits related to photosynthesis (ETR , yield, A) were less variable under stressed conditions.

Vulnerability curves are all s-shaped (Figure 4) and P_{50} values vary from -3.2 MPa to -4.2MPa. Figure 5 shows that more than 70% of the genotypes has a P_{50} value ranging in between -3.5 and -3.9MPa. The average P_{50} values were -3.6 and -3.8MPa for “Corsica x Landes” and “Morocco x Landes” families, respectively. The difference between families is highly significant

($p < 0.0001$). There was no trade-off between hydraulic safety (P_{50}) and hydraulic efficiency (K_s) for both Corsica x Landes ($P = 0.836$) and Morocco x Landes ($P = 0.1867$) families (Figure 7).

Overall, there was no correlation between xylem resistance to cavitation quantified using P_{50} and the leaf functioning traits we measured in the Morocco x Landes family (Table 2), neither under stressful nor under non-stressful water conditions. The only significant correlations were observed under the non-stressful conditions of April 2015: a weak positive relationship was observed with photosynthetic variables (ETR, Yield) and a negative one with water use efficiency (WUE, WUEi). However, these relationships were not significant through similar water conditions in 2016, nor under stressful conditions (July 2015-16). In particular, it is worth noticing that there is no significant and consistent trade-off (negative relationship) between plant functioning and plant resistance. However, there was a significant linear relationship between stem hydraulic conductivity and the actual efficiency of photosystem II under non stressful conditions in spring 2015: plants with the highest water conductivity also presented the highest light use efficiency (Figure 6).

Discussion - conclusion

Through the analysis of two years of physiological measurements under well-watered and water-stressed conditions, we investigated the variability of leaf traits related to transpiration and assimilation in a full-sib Maritime pine family. This variability was also analysed according to P_{50} values measured on stem xylem, without evidencing proof of any trade-off between plant functioning and plant resistance to cavitation.

Our results showed that decreasing water potential resulted in rapid decrease in assimilation rate due to decrease in efficiency of photosystem II and in stomatal conductance within maritime pine. Water deficit-induced decrease in photosynthetic productivity was also observed by López et al (2009) who studied the responses to water deficit in seedlings of *Pinus canariensis* from five provenances under controlled conditions. Transpiration control for ensuring nutrients supply and regulating leaf temperature, and maintaining CO_2 entry of into the leaf, are the main functions of stomata. Photosynthesis is primarily restrained by stomatal closure in response to water deficit. (Flexas and Medrano 2002), affecting plant growth and temperature regulation under drought conditions. The magnitude of the variability of the leaf physiological traits within the Landes x Morocco maritime pine family was of the same magnitude than the one observed for similar traits

within a Spanish maritime pine full-sib family (Oria x Pontevedra, de Miguel et al. 2011). The decrease in maximum efficiency of PSII with water-stress was minimum compared to the one observed in provenances of *Pinus canariensis* submitted to drought (from 0.77 to 0.4, Lopez et al. 2009), or in *Pinus halepensis* (from 0.82 to 0.6, Prieto et al. 2009); this would indicate that the photosynthetic apparatus of *P. pinaster* was structurally less affected by the water stress compared to the one of *P. canariensis* or *P. halepensis*.

Under non stressful conditions, there was a significant relationship between stem hydraulic conductivity and the actual efficiency of PSII; this is showing that the efficiency of leaf functioning traits is strongly affected by the supply capacity of the hydraulic system. Our results are in agreement with Brodribb and Field, (2000) who also found a strong relationship between hydraulic supply of water (hydraulic conductivity) to leaves and maximum photosynthetic capacity (quantum yield) in a community of rainforest comprising of conifers and angiosperms suggesting that the maximum photosynthetic rate of leaves was constrained by their vascular supply. Similar results showing significant positive correlations between leaf-specific hydraulic conductivity of upper branches and maximum rates of net CO₂ assimilation per unit leaf area and stomatal conductance (gs) were also found across 20 species of canopy trees (Santiago et al, 2004). The variability within the Maritime pine family was of the same magnitude than the one observed at the inter-specific level both on coniferous and broadleaved trees (Brodribb and Field 2000). Our results indicate that such a relationship exists not only at interspecific but also at intraspecific level.

On the contrary, during the stress, hydraulic conductivity was not linked to leaf physiology, and overall the resistance to cavitation of the stem was not correlated to functional leaf traits. Since both Pine species present a similar vulnerability to cavitation as assessed by their P50 value, -3.9 for *P. pinaster* (Bouche et al. 2016) and -3.9 / - 4 MPa for *P. canariensis* (Lopez et al. 2013) under temperate climate, this suggests that there is no relationship between the xylem vulnerability and the leaf functioning. The absence of trade-off between resistance and functioning and the existence of a large variability in the traits would leave open the possibilities for a breeding program, to select efficient and resistant genotypes of Maritime pine. To go further into these considerations, investigations could however focus on different genetic material, and should consider the genetic determinisms of this variability, through association genetics studies.

Table 1: Mean (SD) of leaf physiological traits per campaign

Campaign	Ψ_I	Fv/Fm	ETR	Yield	NPQ	E	gs	A	WUE	WUEi
April 2015	-0.29 (0.15)	0.85 (0.01)	206.78 (57.81)	0.28 (0.08)	1.35 (0.48)	0.44 (0.08)	61.41 (11.41)	9.38 (1.35)	21.6 (2.24)	0.15 (0.02)
July 2015	-1.55 (0.15)	0.84 (0.02)	111.05 (38.28)	0.18 (0.05)	4.53 (3.04)	0.14 (0.11)	16.7 (13.21)	1.36 (0.71)	17.27 (18.76)	0.14 (0.15)
	()	()	()	()	()	()	()	()	()	()
May 2016	-0.21 (0.07)	0.87 (0.01)	145.64 (42.56)	0.22 (0.07)	1.97 (0.48)	0.24 (0.04)	40.96 (8.31)	6.61 (0.88)	28 (5.21)	0.17 (0.03)
July 2016	-0.5 (0.14)	0.86 (0.01)	153.14 (39.98)	0.22 (0.05)	2.12 (0.48)	0.29 (0.05)	37.59 (6.45)	4.03 (0.85)	13.82 (2.66)	0.11 (0.02)

Ψ_I predawn leaf water potential (MPa), Fv/Fm maximum efficiency of PSII, ETR electron transport rate, Yield actual efficiency of PSII, NPQ non photochemical quenching, E leaf transpiration rate ($\text{mmol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), gs stomatal conductance ($\text{mmol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), A assimilation rate ($\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), WUE water use efficiency ($\mu\text{mol}\cdot\text{mmol}^{-1}$), WUEi intrinsic water use efficiency ($\mu\text{mol}\cdot\text{mmol}^{-1}$).

Table 2: Correlation between P50 and leaf physiological traits.

Campaign	Colonne1	Fv/Fm	ETR	Yield	NPQ	E	gs	A	WUE	WUEi
April 2015	Pearson r	0.07	0.20	0.22	-0.08	0.18	0.16	0.03	-0.28	-0.25
	p-value	0.50	0.04	0.02	0.44	0.06	0.10	0.76	0.00	0.01
July 2015	Pearson r	-0.18	-0.04	-0.04	0.00	0.26	0.26	0.05	-0.28	-0.28
	p-value	0.07	0.67	0.67	0.98	0.10	0.11	0.73	0.08	0.08
	n	108.00	107.00	107.00	107.00	40.00	40.00	44.00	40.00	40.00
May 2016	Pearson r	-0.08	-0.11	-0.12	-0.10	0.01	-0.03	0.00	0.02	0.05
	p-value	0.43	0.27	0.23	0.29	0.92	0.79	0.99	0.87	0.64
July 2016	Pearson r	-0.06	0.05	0.01	-0.02	-0.09	-0.14	-0.23	-0.16	-0.13
	p-value	0.56	0.62	0.93	0.84	0.35	0.14	0.02	0.09	0.20

Fv/Fm maximum efficiency of PSII, ETR electron transport rate, Yield actual efficiency of PSII, NPQ non photochemical quenching, E leaf transpiration rate ($\text{mmol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), gs stomatal conductance ($\text{mmol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), A assimilation rate ($\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), WUE water use efficiency ($\mu\text{mol}\cdot\text{mmol}^{-1}$), WUEi intrinsic water use efficiency ($\mu\text{mol}\cdot\text{mmol}^{-1}$). Except when indicated otherwise (n), number of data used for the analyses was 108. Significant correlations ($p < 0.05$) are indicated in bold characters.

Figure captions:

Figure 1: Abiogen full-sib family essay. Each block contains individuals (dots) of a full-sib family. Measurements were performed on two blocks of the Landes x Morocco family (bold letters).

Figure 2: Predawn leaf water potential (average + standard deviation) during each campaign (April and July 2015, May and July 2016)

Figure 3: Leaf functioning as a function of predawn leaf water potential: (A) assimilation (A, $\mu\text{mol.m}^{-2}.\text{s}^{-1}$). (B) Potential efficiency of photosystem II (Fv/Fm, blue dots) and actual efficiency of PSII (Yield, red dots). (C) Stomatal conductance (gs, $\text{mmol.m}^{-2}.\text{s}^{-1}$). (D) Intrinsic water use efficiency (WUEi, $\mu\text{mol.mmol}^{-1}$).

Figure 4: Vulnerability curves obtained on 480 individuals from the two studied families growing in common garden. Each color represents a different individual.

Figure 5: Frequency distribution of P_{50} (pressure inducing 50 % loss of hydraulic conductance) in both “Corsica x Landes” and “Morocco x Landes” families of maritime pine.

Figure 6: Actual efficiency of photosystem II for light adapted needles (Yield) as a function of xylem specific hydraulic conductivity (K_s , $\text{kg.m}^{-1}.\text{s}^{-1}.\text{MPa}^{-1}$).

Figure 7: Relationship between hydraulic safety (P_{50} , MPa) and hydraulic efficiency (K_s , $\text{m}^2 \text{MPa}^{-1} \text{s}^{-1}$) across Corsica x Landes (blue) and Morocco x Landes (orange). N = 240 for each Corsica x Landes and Morocco x Landes families.

References

- Bigras, F.J. 2005. Photosynthetic response of white spruce families to drought stress. *New Forests*, 29(2), 135-148.
- Bouche, P. S., Delzon, S., Choat, B., Badel, E., Brodribb, T. J., Burlett, R., Cochard, H., Charra-Vaskou, K., Lavigne, B., Li, S., Mayr, S., Morris, H., Torres Ruiz, J.M., Zufferey, V., & Jansen, S. (2016). Are needles of *Pinus pinaster* more vulnerable to xylem embolism than branches? New insights from X-ray computed tomography. *Plant, cell & environment*, 39(4), 860-870.
- Bouffier, L., Raffin, A. & Kremer, A. (2009). La gestion de la variabilité génétique dans le programme d'amélioration du pin maritime (*Pinus pinaster* Ait.). *Rev. For. Fr.* LXI – 4: 369-388.
- Brodribb, T. J., Bowman, D. J., Nichols, S., Delzon, S., & Burlett, R. (2010). Xylem function and growth rate interact to determine recovery rates after exposure to extreme water deficit. *New Phytologist*, 188(2), 533-542.
- Choat B, S Jansen, TJ Brodribb, H Cochard, S Delzon, R Bhaskar, SJ Bucci, TS Feild, SM Gleason, UG Hacke, AL Jacobsen, F Lens, H Maherali, J Martinez-Vilalta, S Mayr, M Mencuccini, PJ Mitchell, A Nardini, J Pittermann, RB Pratt, JS Sperry, M Westoby, IJ Wright & AE Zanne. (2012). Global convergence in the vulnerability of forests to drought. *Nature* 491: 752-756
- Cochard, H. (2002). A technique for measuring xylem hydraulic conductance under high negative pressures. *Plant, Cell & Environment*, 25(6), 815-819.
- Cochard, H., Damour, G., Bodet, C., Tharwat, I., Poirier, M., & Améglio, T. (2005). Evaluation of a new centrifuge technique for rapid generation of xylem vulnerability curves. *Physiologia Plantarum*, 124(4), 410-418.
- Davis, M.B. & Shaw, R.G. (2001). Range shifts and adaptive responses to Quaternary climate change. *Science*. 292:673-679.
- de Miguel, M., Sánchez-Gómez, D., Cervera, M. T., & Aranda, I. (2011). Functional and genetic characterization of gas exchange and intrinsic water use efficiency in a full-sib family of *Pinus pinaster* Ait. in response to drought. *Tree physiology*, 32(1), 94-103.
- Flexas, J. & Medrano, H. (2002). Drought-inhibition of photosynthesis in C-3 plants: stomatal and non-stomatal limitations revisited. *Ann. Bot.*, 89, 183–189.
- Fu, P. L., Jiang, Y. J., Wang, A. Y., Brodribb, T. J., Zhang, J. L., Zhu, S. D., & Cao, K. F. (2012). Stem hydraulic traits and leaf water-stress tolerance are co-ordinated with the leaf phenology of angiosperm trees in an Asian tropical dry karst forest. *Annals of botany*, 110(1), 189-199.

- Grassi, G. & Magnani, F. (2005). Stomatal, mesophyll conductance and biochemical limitations to photosynthesis as affected by drought and leaf ontogeny in ash and oak trees. *Plant, Cell and Environment* 8, 834–849
- Guo, X. Y., Zhang, X. S., & Huang, Z. Y. (2010). Drought tolerance in three hybrid poplar clones submitted to different watering regimes. *Journal of Plant Ecology*, rtq007.
- Gutierrez, L., Casares, A., Sánchez-Tamés, R., & Majada, J. (2002). Early growth, biomass allocation and physiology in three *Eucalyptus nitens* populations under different water regimes. *Forestry*, 75(2), 139-148.
- Harfouche, A. & Kremer, A. (2000). Provenance hybridization in a diallel mating scheme of maritime pine (*Pinus pinaster*). I. Means and variance components. *Canadian journal of forest research*, 30(1), 1-9.
- Hartmann, Klein Tank, Rusticucci, Alexander, Brönnimann, Charabi, Dentener, Dlugokencky, Easterling, Kaplan, Soden, Thorne, Wild and Zhai. (2013). Observations: Atmosphere and Surface. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. pp 159-254.
- IGN. (2016). Le mémento, inventaire forestier, la forêt en chiffres et en cartes. St Mandé, IGN. 17 pp.
- Lagraulet, H. (2015). Plasticité phénotypique et architecture génétique de la croissance et de la densité du bois du pin maritime (*Pinus pinaster* Ait.) (Doctoral dissertation, Université de Bordeaux).
- Lamy, J. B., Delzon, S., Bouche, P. S., Alia, R., Vendramin, G. G., Cochard, H., & Plomion, C. (2014). Limited genetic variability and phenotypic plasticity detected for cavitation resistance in a Mediterranean pine. *New Phytologist*, 201(3), 874-886.
- Lamy, J. B., Plomion, C., Kremer, A., & Delzon, S. (2012). $QST < FST$ as a signature of canalization. *Molecular Ecology*, 21(23), 5646-5655.
- López, R., Rodríguez-Calcerrada, J., & Gil, L. (2009). Physiological and morphological response to water deficit in seedlings of five provenances of *Pinus canariensis*: potential to detect variation in drought-tolerance. *Trees*, 23(3), 509-519.
- López, R., López de Heredia, U., Collada, C., Cano, F. J., Emerson, B. C., Cochard, H., & Gil, L. (2013). Vulnerability to cavitation, hydraulic efficiency, growth and survival in an insular pine (*Pinus canariensis*). *Annals of botany*, 111(6), 1167-1179.

- Martinez Meier, A. G., Sanchez, L., Dalla Salda, G., Pastorino, M. J., Gautry, J. Y., Gallo, L. A., & Rozenberg, P. (2008). Genetic control of the tree-ring response of Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) to the 2003 drought and heat-wave in France. *Annals of forest science*, 65(1), 102-102.
- McAdam, S. A., & Brodribb, T. J. (2015). The Evolution of Mechanisms Driving the Stomatal Response to Vapor Pressure Deficit. *Plant Phys* 167 (3): 833-843
- Meehl, G.A. & Tebaldi, C. (2004). More intense, more frequent, and longer lasting heat waves in the 21st century. *Science*. 305:994-997.
- Olivas-García, J. M., Cregg, B. M., & Hennessey, T. C. (2000). Genotypic variation in carbon isotope discrimination and gas exchange of ponderosa pine seedlings under two levels of water stress. *Canadian journal of forest research*, 30(10), 1581-1590.
- Pettigrew, W. T., & Turley, R. B. (1998). Variation in photosynthetic components among photosynthetically diverse cotton genotypes. *Photosynthesis research*, 56(1), 15-25.
- Petridis, A., Therios, I., Samouris, G., Koundouras, S., & Giannakoula, A. (2012). Effect of water deficit on leaf phenolic composition, gas exchange, oxidative damage and antioxidant activity of four Greek olive (*Olea europaea* L.) cultivars. *Plant physiology and biochemistry*, 60, 1-11.
- Piper, F.I., Corcuera, L.J., Alberdi, M. & Lusk, C. (2007). Differential photosynthetic and survival responses to soil drought in two evergreen *Nothofagus* species. 2007. *Ann. For. Sci.* 64 : 447–452
- Prieto, P., Penuelas, J., Llusia, J., Asensio, D., & Estiarte, M. (2009). Effects of long-term experimental night-time warming and drought on photosynthesis, Fv/Fm and stomatal conductance in the dominant species of a Mediterranean shrubland. *Acta Physiologiae Plantarum*, 31(4), 729-739.
- Roháček, K. (2002). Chlorophyll fluorescence parameters: the definitions, photosynthetic meaning, and mutual relationships. *Photosynthetica*, 40(1), 13-29.
- Santiago, L. S., Goldstein, G., Meinzer, F. C., Fisher, J. B., Machado, K., Woodruff, D., & Jones, T. (2004). Leaf photosynthetic traits scale with hydraulic conductivity and wood density in Panamanian forest canopy trees. *Oecologia*, 140(4), 543-550.
- Sofa, A., Dichio, B., Montanaro, G., & Xiloyannis, C. (2009). Photosynthetic performance and light response of two olive cultivars under different water and light regimes. *Photosynthetica* 47 (4): 602-608
- Urli, M., Porté, A. J., Cochard, H., Guengant, Y., Burrell, R., & Delzon, S. (2013). Xylem embolism threshold for catastrophic hydraulic failure in angiosperm trees. *Tree physiology*, 33(7), 672-683.

Viljevac, M., Dugalić, K., Mihaljević, I., Šimić, D., Sudar, R., Jurković, Z., & Lepeduš, H. (2013). Chlorophyll content, photosynthetic efficiency and genetic markers in two sour cherry (*Prunus cerasus* L.) genotypes under drought stress *Acta Bot. Croat.* 72 (2), 221–235.

Figure 1:

Abiogen full-sib family essay. Each block contains individuals (dots) of a full-sib family. Measurements were performed on two blocks of the Landes x Morocco family (bold letters).




Figure 2:

Predawn leaf water potential (average + standard deviation) during each campaign (April and July 2015, May and July 2016).




Figure 3:

Leaf functioning as a function of predawn leaf water potential: (A) assimilation (A , $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$). (B) Potential efficiency of photosystem II (F_v/F_m , blue dots) and actual efficiency of PSII (Yield, red dots). (C) Stomatal conductance (g_s , $\text{mmol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$). (D) Intrinsic water use efficiency (WUE_i , $\mu\text{mol}\cdot\text{mmol}^{-1}$).




Figure 4:

Vulnerability curves obtained on 480 individuals of inter “Morocco x Landes” and “Corsica x Landes” families of maritime pine (*Pinus pinaster* Ait.) established in Cestas (France). Each color represents a different individual.




Figure 5:

Frequency distribution of P_{50} (pressure inducing 50 % loss of hydraulic conductance) in both “Corsica x Landes” and “Morocco x Landes” families of maritime pine (*Pinus pinaster* Ait.).




Figure 6:

Actual efficiency of photosystem II for light adapted needles (Yield) as a function of xylem specific hydraulic conductivity (Ks, kg.m-1.s-1.MPa-1).





Figure 7: Relationship between hydraulic safety (P_{50} , MPa) and hydraulic efficiency (K_s , $\text{m}^2 \text{MPa}^{-1} \text{s}^{-1}$) across Corsica x Landes (blue) and Morocco x Landes (orange). $N = 240$ for each Corsica x Landes and Morocco x Landes families.



Discussion

Discussion

The aim of this project was to investigate the intraspecific variability in embolism resistance in annual crop and conifer tree species and to find the relationship between hydraulic traits and traits related to wood density, xylem anatomy, genetics and leaf physiology.

In Part I, we provide a novel examination of variation in vulnerability to drought in sunflower. Few studies have considered the possibility of using embolism resistance to find more drought resistant crop varieties. We therefore investigated and determined vulnerability to embolism across four sunflower accessions growing under same environmental conditions with significant variations in P_{50} (pressure inducing 50% loss of hydraulic conductivity) ranging between -2.67 and -3.22 MPa. Moreover, the study of interactions between hydraulic traits, growth traits and xylem anatomical traits highlighted the potential trade-offs between hydraulic safety and growth but not between hydraulic safety and efficiency. The thorough tests of correlation between anato-morphological features and vulnerability provides a complete picture of the key factors interacting to determine P_{50} . We indeed found that several anatomical traits, such as vessel density and vessel lumen area correlated with embolism resistance.

In part II we provide new insight into the genetic determinism of resistance to embolism at intraspecific level in a large-sized three-generation inbred pedigree of *Pinus pinaster*. We dissected the genetic architecture of embolism related traits (hydraulic safety) and wood density related traits, and tested the relationship between wood density and xylem safety at the phenotypic and QTL level. Mean P_{50} value was found to be around -3.70 ± 0.21 MPa in both chapter 2 (Corsica x Landes) and chapter 3 (Corsica x Landes and Morocco x Landes families) with considerably low variability for P_{50} ($CV_P=5.6\%$). There were a lack of relationship between hydraulic safety and hydraulic efficiency, wood density and leaf functional traits. The results also explained that embolism resistant genotypes in this family are not dependent on denser wood as no QTL co-located between P_{50} and wood density.

Vulnerability curves assessment in long vesselled species

For the construction of vulnerability curves (VCs) in sunflowers, from which the embolism resistance of a sample is estimated, notably from the inflexion point, i.e., the pressure at which 50% of total xylem hydraulic conductance is lost (P_{50} , in Mega Pascal), and the slope of the curve

(S , in % MPa^{-1}), we use 42 cm rotor diameter. This large-diameter rotor was developed for species having long vessels in order to avoid the so called ‘open vessel’ artifact that may favor exponential (r-shaped) vulnerability curves. This has been demonstrated several times in woody species with long vessels (Cochard et al., 2013; Martin-StPaul et al., 2014; Torres-Ruiz et al., 2014, 2017; Cochard et al., 2015; Choat et al., 2016). Prior to the main experiment, we collected six sunflower stems to test whether we were able to accurately assess vulnerability curves with the standard cavitron (rotor diameter of 27 cm). All VCs obtained with this standard rotor were r-shaped while all VCs obtained with the 42 cm large rotor were s-shaped (see Figure. 1). Then, we performed maximum vessel lengths measurements on six stems by injecting air at 2 bars and cutting the apical end of the water-immersed stem section until the air bubbles emerged. This procedure allowed us to find that the sunflower stems have a maximum vessel length of 23 cm and therefore confirm that the VCs cannot be assessed with the standard cavitron. Indeed a significant proportion of open-cut vessels surpass the center of the plant segment or even permeate through its whole length when using a 27 cm large rotor.

Embolism resistance is typically assessed by decreasing xylem pressures and determining the corresponding losses in hydraulic conductivity (K_s). However, by using centrifuge technique, it is not possible to determine with certainty whether this loss in K_s reflects patterns of embolism that occur *in situ* or whether it comes from an artefact due to the presence of open vessels. The issue of this open-vessel artefact may be avoided and the process of embolism formation be evaluated with greater reliability by the use of alternative techniques that enables the direct observation of xylem embolism in a non-invasive, unbiased manner, such as MRI (Holbrook et al., 2001; Kaufmann et al., 2009; Choat et al., 2010) and X-ray micro tomography (Cochard et al., 2015; Torres-Ruiz et al., 2015, 2016; Bouche et al., 2016; Brodribb et al., 2016; Choat et al., 2016; Torres-Ruiz et al., 2017).

Intraspecific variability in embolism resistance in sunflower and maritime pine

Mean value of embolism resistance (P_{50}) for four sunflower accessions measured was -2.99 ± 0.15 MPa. There was a significant intraspecific variability in resistance to embolism among four accessions ranging from -2.67 to -3.22 MPa, between the most vulnerable and the most resistant accessions, respectively.




Figure: 1. Vulnerability curves (VCs) of six sunflower stems, for which xylem embolism was induced by in situ flow centrifugation according to the Cavitron technique. Three stems were measured with the standard rotor design (27cm, top panel) and three with the medium size rotor (42cm, bottom panel). VCs are expressed as the percentage loss of hydraulic conductivity (PLC) as a function of xylem pressure. This figure shows the discrepancy between the two rotor sizes: all VCs were r-shaped with the 27cm diameter rotor and s-shaped with the 42 cm diameter rotor, leading to a difference of 2MPa between them. The r-shaped curves (top panel) result from an open vessel artifact (Torres-Ruiz et al., 2014, 2017).

The difference in P_{50} values between the most resistant sunflower accession, the early-sown ES_Ethic, and the most vulnerable accession, the late-sown LG_5660 was found to be 0.55 MPa. Apart from showing significant intraspecific differences in vulnerability to embolism, these findings may also help in terms of developing selection and breeding strategies in future for drought prone areas on the basis of the greater resistance of specific sunflower accessions to xylem embolism. In comparison to many other crop species, sunflower is better able to resist drought events and is often cultivated in arid regions, however, like many other oil seed crops, sunflower is also sensitive to water stress, especially at the germination stage (Ahmad et al., 2009). Stiller and Sperry, (2002) reported variation in P_{50} (from -0.8 to -3 MPa) in a single sunflower genotype subjected to drought and re-watering cycles with mean P_{50} value -3.0 ± 0.1 MPa under well-watered conditions, which is quite similar to one found in our study. Despite the critical role of xylem resistance to embolism in plant survival during drought events (Anderegg et al. 2015), this trait is less studied in crops (Tyree et al., 1986; Sperry et al., 2003) and hence in sunflower. However, in some studies, decrease in leaf water potential in various sunflower lines has been observed due to water deficit (Ashraf and O'Leary, 1996). Leaf water potential in sunflower normally ranges between - 0.48 and - 1.74 MPa under different agro-climatic conditions (Prasad et al., 1985; Rachidi et al., 1993), however, under reduced water supply, it can further drop as below as - 3.0 MPa (Wise et al., 1990). This species seems therefore to have a null safety margin and works close to its hydraulic failure limit as already reported for trees by Choat et al. (2012). Moreover, it has been reported that dwarf sunflower lines are more resistant to drought than tall lines, showing a smaller decrease in leaf osmotic potential in response to drought stress (Angadi and Entz, 2002a). Maintenance of turgor plays a significant role in stress tolerance of plants due to its role in stomatal regulation, and hence photosynthesis (Ludlow et al., 1985). It has been shown that sunflower cultivars with higher turgor potential under water stress resulted in less decrease in yield (Angadi and Entz, 2002b). However, Ashraf and O'Leary, (1996) demonstrated that stress tolerance was not associated with leaf turgor potential. Indeed, the importance of stomata in understanding the overall hydraulic and leaf functioning performance cannot be under estimated as the closure of stomata is the primary limitation to photosynthesis in response to water deficit (Flexas & Medrano 2002) and play a key role in plant drought resistance. The timing of stomatal closure and xylem embolism during plant dehydration, as well as the fate of embolized organs, is a topic of hot debate in current years. While some favor the view that complete stomatal closure

precedes the onset of xylem embolism, others argue that the two processes occur simultaneously and are accompanied with daily xylem refilling. In a recent meta-analysis related to stomatal response to drought and embolism resistance (Martin-StPaul et al., 2017), it was shown that the range of variation of water potential at which stomata close is much smaller when it is considered in the light of the full range of embolism resistance, indicating the absence of functional coordination between stomatal closure and the onset of embolism formation. Thanks to the development of a new optical technique (Brodrribb et al., 2016), Hochberg et al. (2017) observed the complete stomatal closure before the appearance of embolism in the leaves of grapevines (*Vitis vinifera* L.) by several days. They further reported that the basal leaves once embolized, were shed, and prevented further water loss, thereby protecting the hydraulic integrity of younger leaves and the stem. I propose here to use this technique to test whether sunflowers also close their stomata before the onset of cavitation and also to quantify the leaf hydraulic safety margin between accessions (difference between the water potential inducing 90% stomatal closure and the potential leading to 12% of embolism in leaf veins).

The mean P_{50} value measured for 477 *Pinus pinaster* F2 genotypes “Corsica x Landes” provenance hybrid, growing in common garden, sampled between November 2012 and April 2013 was -3.70 MPa (Chapter 2) and are very similar to the mean P_{50} value for “Corsica x Landes” and “Morocco x Landes” families comprising a total of 480 genotypes sampled and measured during two campaigns, from October 2014 to February 2015 and from January 2016 to March 2016, which was 3.67 MPa (Chapter 3). The mean P_{50} values measured here are similar to what was found in an earlier study (Lamy et al. 2011 (P_{50} , -3.93 MPa), 2014 (P_{50} , 3.7 MPa).

Considerably low value ($CV_P=5.6\%$) of coefficient of variation for P_{50} for “Corsica x Landes” and more than 70% of the genotypes for “Corsica x Landes” and “Morocco x Landes” families with P_{50} value ranging between -3.5 and -3.9 MPa shows that the intraspecific variability in embolism resistance in maritime pine is very low, which shows they could be more similar for this trait than expected under genetic drift as previously shown by Lamy et al., (2011, 2014). This evolutionary stasis could be due to uniform selection (Lamy et al., 2011), with all populations selected towards an optimum value, which is, therefore, not under climate control. It is worth noting that some species show phenotypic plasticity for embolism resistance, with a slight trend of increasing P_{50} with increasing aridity in natural populations of *P. canariensis* (López et al., 2013) and in *Fagus*

sylvatica (Aranda et al., 2014; Schuldt et al., 2015). Similarly López et al. (2016) found that the intraspecific variability of hydraulic traits was largely due to phenotypic plasticity in six populations of Canary Island pine (*Pinus canariensis*). Vulnerability to embolism showed a plastic response in the driest provenance trial located in the ecological limit of the species. This trait was significantly correlated with annual precipitation, temperature oscillation and drought period at the origin of the populations, indicating that vulnerability to cavitation was directly related to survival in the dry and the xeric provenance trials, showing its importance in determining drought resistance. Contrasting results for variation of xylem embolism resistance at intraspecific level, as for example, moderate to low levels of intraspecific variation (P_{50} , -2.21 to -2.97 MPa) in poplar (*Populus* sp) (Hajek et al., 2014), and no significant variability in P_{50} in European beech (*Fagus sylvatica* L.) (Hajek et al., 2016) emphasized the need to further investigate this trait at broader spectrum. At broader evolutionary levels there is significant variability in P_{50} (Maherali et al., 2004; Pittermann et al., 2012; Bouche et al., 2014). On the other hand, at intraspecific level there is generally low variability (Delzon et al., 2010) and little genetic differentiation for xylem embolism resistance (Lamy et al., 2011; Sáenz-Romero et al., 2013).

Trade-offs with hydraulic traits

Safety/growth and safety/efficiency trade-offs

In sunflower we found a trade-off between P_{50} and growth-related traits i.e. height but not between hydraulic safety and xylem efficiency (Chapter 1), showing that shorter plants had greater embolism resistance. Previous studies show conflicting results regarding the possible existence of such a trade-off between P_{50} and growth-related traits. Positive relationship between embolism resistance and growth related traits was reported in poplar (Fichot et al., 2010) and Scot pine (Sterck et al., 2012). However, Cochard et al. (2007) found a close relationship between xylem vulnerability and productivity in poplar and willow clones. Several recent studies found no evidence of a trade-off between vulnerability to embolism and growth-related traits (Hajek et al., 2014; Guet et al., 2015; Hajek et al. 2016). In maritime pine family, we did not find any relationship of embolism resistance with wood density (Chapter 2) and with leaf functioning traits (Chapter 3), indicating that embolism resistant genotypes are independent on denser wood. It has been reported that xylem embolism resistance increases at the cost of slower growth as plant either has to allocate more carbon for thicker cell walls to construct denser wood (Hacke et al. 2001) or

to construct foliar and axial tissues for increased canopy carbon gains and growth rate (Wikberg and Ogren, 2004, Ducrey et al. 2008). Increased survival is the evident advantage of xylem safety during drought events while increased carbon allocation to leaves relative to sapwood area is achieved by xylem efficiency which allows rapid growth rate and optimizing rates of photosynthesis in competitive environment (Santiago et al., 2004; Poorter et al., 2010). Thus plants need a balanced state between embolism resistance (xylem safety) and water transport (xylem efficiency) to their photosynthetic organs for better performance (Tyree & Zimmermann, 2002; Hacke et al., 2006). As plants get advantages from a water transport system that is both efficient as well as safe from embolism at the same time, variability in both efficiency and safety are expected to reflect ecological and evolutionary differences among species (Sperry, 2003; Baas et al., 2004). Indeed, it is generally believed that there is a trade-off between xylem safety and hydraulic efficiency (Lens et al., 2011). However, in our study we found no trade-off between these two traits at intraspecific level in both sunflower and maritime pine, which are in agreement with those found by Gleason et al. (2015) who showed no or weak trade-off between safety and efficiency at interspecific level across a wide range of angiosperm and gymnosperm species. It may be due to reason that in gymnosperms, safety seems to arise largely by the amount of overlap between torus relative to the size of pit aperture having minimum influence on efficiency (Delzon et al., 2010; Pittermann et al., 2010; Bouche et al., 2014). In angiosperms, mechanical stability is largely decoupled from wood density, vessel lumen fraction, and therefore from hydraulic efficiency, resultantly a strong tradeoff between hydraulic efficiency and mechanical safety across angiosperms could not be expected.

Relationship between hydraulic traits and xylem anatomy

We found that both the point of air entry during embolism formation (P_{12}) and the xylem pressure inducing a 50% loss of conductance (P_{50}) were strongly influenced by the vessel density in secondary xylem (VD_{Sx}) and total xylem (VD_{Tx}) (Chapter 1). These results indicate that higher production of vessels per unit xylem area increases the embolism resistance in sunflower. Increased embolism resistance by the production of more number of vessels has also been reported by Schuldt et al. (2015) for European beech. They found that this relationship was stronger at air entry point (P_{12}) as we found in case of sunflower. However, in poplar no such close relationship between vessel density and embolism resistance was found (Hajek et al., 2014). Another xylem

anatomical trait “vessel grouping index” found closely related to vessel density (Lens et al., 2011) has also been reported to closely related with xylem embolism at the point of air entry (P_{12}) Schuldt et al. (2015). These results in a way support the assumption put forwarded by Carlquist, (1966) that in species adapted to water-limited environments having a xylem tissue composed of nonconductive fibers and no vasicentric tracheids, vessels should group to a larger degree to maintain the conductive flow path by pathway redundancy in case if the largest vessel of the group embolized.

We also found strong correlations between P_{12} and both inter vessel double-wall thickness in primary xylem (T_{vwPx}) and the thickness-to-span ratio of vessels in the primary xylem (TD^{-1}_{Px}), which suggest that these two traits are important to directly prevent embolism formation or to protect the vascular system from higher tensions. A higher thickness-to-span ratio has been reported to strengthen the vessel walls against implosion and the relationship between increased embolism resistances with increasing thickness-to-span ratio has also been reported in conifers (Hacke et al., 2001; Bouche et al. 2014). Previous studies (Pittermann et al., 2005; Sperry et al., 2006) show that species resistant to cavitation have thicker tracheid walls relative to lumen area. Whereas Pittermann et al. (2006b) and Sperry et al. (2006) explained that variation in the thickness to span ratio was determined by lumen diameter and not by the cell wall thickness. Bouche et al. (2014) showed that embolism resistant species had thicker tracheid walls with only minor reduction in lumen diameter (conduit size) and minimum impact on hydraulic conductivity. According to Maherali et al. (2004) and Choat et al. (2012) higher embolism resistance is associated with lower negative sap pressure, tracheids with a higher thickness to span ratio are required to resist mechanical stresses. From a functional point of view, thicker walls relative to lumen area do not improve drought-induced embolism resistance as this phenomenon occurs at the bordered pit level. Embolism resistance might, therefore, be indirectly linked to thickness to span ratio.

The relationship between higher levels of lignification and greater embolism resistance has been reported in woody plants (Awad et al., 2012) as well as in herbaceous plants (Lens et al., 2013; Tixier et al., 2013; Lens et al., 2016). However, on the contrary we found no relationship between P_{50} and greater stem tissue lignification, indicating that the development of embolism-resistant stems does not involve lignification of tissues. Our results are supported by previous

studies showing that achievement of higher levels of embolism resistance is not always dependent on increased lignification of the xylem tissues (Watkins et al., 2010; Pittermann et al., 2011).

Relationship between xylem specific hydraulic conductivity and leaf physiological traits

We found a significant linear relationship between stem hydraulic conductivity and the actual efficiency of photosystem II under non stressful conditions. It was found that plants with the highest water conductivity also presented the highest light use efficiency resulting in the highest photosynthetic productivity (Chapter 3). The significant relationship between stem hydraulic conductivity and the actual efficiency of Photosystem II under non stressful conditions show that the efficiency of leaf functioning traits is strongly affected by the capacity of the hydraulic system to transport water. Similar results were also found by Brodribb and Field, (2000) demonstrating a strong relationship between hydraulic supply of water (hydraulic conductivity) to leaves and maximum photosynthetic capacity (quantum yield) in a community of rainforest comprising of conifers and angiosperms suggesting that the maximum photosynthetic rate of leaves was constrained by their vascular supply. This study illustrates that the photosynthetic capacity of leaves is related to the hydraulic conductance of the supporting xylem tissue under optimal conditions. Similar results showing significant positive correlations between leaf-specific hydraulic conductivity of upper branches and maximum rates of net CO₂ assimilation per unit leaf area and stomatal conductance (gs) were also found across 20 species of canopy trees (Santiago et al., 2004). The variability within the Maritime pine family was of the same magnitude as the one observed at the inter-specific level both on coniferous and broadleaved trees (Brodribb and Field 2000). Our results indicate that such a relationship exists not only at interspecific but also at intraspecific level. This relationship between hydraulic efficiency and photosynthetic productivity is understandable. Loss of water through transpiration is monitored by opening and closing of stomata, which is governed by the turgor of the guard-cells. During drought, plants can restrict loss of water by regulating overall stomatal conductance at the cost of reduced photosynthesis during drought. If stomata are kept fully open during drought, leaf and xylem water potentials (ψ_L and ψ_X , respectively) would decrease rapidly to a point where permanent damage to photosynthetic apparatus can occur through embolism. Plants avoid this area of dangerous water-potential by closing stomata, thereby stopping water-flow to the leaves, consequently photosynthesis is reduced drastically by restricting carbon entry. The relationship between hydraulic conductivity and leaf

functioning traits under stressful conditions have been studied extensively (Brodrribb et al., 2003; Domec et al., 2008) however, this relationship is required to be investigated under non-stressful conditions. As previously discussed that no relationship between the xylem vulnerability and the leaf functioning was found in maritime pine which is also supported by the similar P_{50} values -3.9 MPa for *P. pinaster* (Bouche et al. 2016) and -3.9 / - 4 MPa for *P. canariensis* (Lopez et al., 2013) under temperate climate. This absence of trade-off between resistance to embolism and leaf functioning traits and the existence of a high variability in the physiological traits would pave a way to open the possibilities for a breeding program and to select efficient and resistant genotypes of Maritime pine by focusing on different genetic material and should consider the genetic determinisms of this variability through integration of genetics studies.

Implications and future perspectives

The results presented in this thesis greatly improved our understandings of the intraspecific variability in hydraulic traits and their association with other traits like genetics, xylem anatomy and leaf functioning. Having knowledge in advance regarding the plant species which are under threat and are most vulnerable to drought-induced mortality, we can make efforts to prevent major impacts of future droughts and heatwaves on the performance of plant functioning. As climate disasters are the main reason for fluctuations in crop yields and occurrence of drought is one of the most widespread climate disasters affecting agricultural productivity (UNDP 2004; Dilley et al., 2005; Helmer and Hilhorst, 2006) and since many conifers are planted or ecologically dominant notably in the Northern Hemisphere, this knowledge is vital for managing forest and crop ecosystems in the context of climate change.

The intraspecific variation in resistance to drought-induced xylem embolism in sunflower explained in chapter 1 may be helpful to select drought-resistant accessions/genotypes, which will be of vital importance for future farming, particularly to take decisions for selection of suitable genotypes to grow in areas prone to drought. The study highlighted the potential tradeoffs between height and hydraulic safety and height and hydraulic efficiency and emphasized the need to intensively investigate the variability of embolism resistance across a wider range of accessions. Future implications regarding sunflower could envisage to investigate

- (i) The possible existence of a trade-off between embolism resistance and yield potential. High yield potential is, indeed, the main objective of the most crop breeding programs, which may not be compatible with higher embolism resistance.
- (ii) The relationship between embolism resistance and pit membrane thickness as higher embolism resistance has been attributed to inter-conduit pitting in *Zea mays* (Li et al., 2009).
- (iii) Leaf hydraulic vulnerability to embolism using the new optical technique described by Brodribb et al. (2016). It is a new image processing technique that detects dynamic embolism by recording rapid changes in the transmission of light through the veins of leaf samples, thus enabling embolisms to be filmed and mapped during the development of leaf water stress. Plant photosynthetic performance can be well quantified on the basis of leaf hydraulic performance, which is a sensitive indicator of photosynthetic productivity (Sack and Scoffoni, 2013). Moreover, during water stress the vulnerability of leaves to hydraulic dysfunction has been demonstrated to provide valuable information regarding the timing of plant death, and the mechanisms of adaptation to increasing aridity (Blackman et al., 2010; Brodribb et al., 2014). Sunflower leaves are, therefore, required to investigate whether they are more vulnerable to embolism than stem and to test whether sunflower leaves close their stomata before the onset of embolism formation.

In maritime pine intraspecific variability is low (Lamy et al., 2011) and the comparison of QTL locations between embolism resistance and wood density related traits presented in chapter 2 failed to detect co-locating regions. Whether these results would still hold true, the lack of phenotypic and genetic correlations across families of five ecotypes (Lamy et al., 2012) strongly support the lack of causality between these two traits in this species. Finally, even if we detected strong QTL, the gain for future breeding program could be small or limited due to the low variability in P_{50} observed in maritime pine. Alternatively, we could find some interspecific hybrid to make a cross with *P. pinaster*. The best potential candidate for this would be *Pinus brutia* with P_{50} about -4.7 MPa. With its strong drought tolerance and control of apical dominance, this species is considered very important in the low/mid rainfall zone as its drought tolerance exceeds that of *P. pinaster*. It has already been tested as *P. pinaster* x *P. brutia* hybrid trial in 2005, east of York, Australia to

improve drought tolerance and apical dominance of Maritime pine with survival rate of 88% at the age of one year (Butcher, 2007).

References

- Adnan, S., Ullah, K., Gao, S., Khosa, A. H., & Wang, Z. (2017). Shifting of agro-climatic zones, their drought vulnerability, and precipitation and temperature trends in Pakistan. *International Journal of Climatology*, 37 (Suppl.1): 529–543.
- Ahmad, M. I., Ali, A., Khan, A., Jamro, S. A., Sher, A., & Shafeeq-ur-Rahman, A. R. (2017). OILCROP-SUN Model for Nitrogen Management of Diverse Sunflower (*Helianthus annuus* L.) Hybrids Production under Agro-Climatic Conditions of Sargodha, Pakistan. *American Journal of Plant Sciences*, 8(03), 412-427.
- Ahmad, S., Ahmad, R., Ashraf, M. Y., Ashraf, M., & Waraich, E. A. (2009). Sunflower (*Helianthus annuus* L.) response to drought stress at germination and seedling growth stages. *Pak. J. Bot*, 41(2), 647-654.
- Alder, N. N., Pockman, W. T., Sperry, J. S., & Nuismer, S. (1997). Use of centrifugal force in the study of xylem cavitation. *Journal of Experimental Botany*, 48(3), 665-674.
- Allen, C.D., Breshears, D.D., McDowell, N.G. (2015). On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* 6: art129.
- Allen, C.D., Macalady, A.K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger, T., Rigling, A., Breshears, D.D., Hogg, E.T. and Gonzalez, P. (2010). A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest ecology and management*, 259(4), 660-684.
- Anderegg, L.D.L., Anderegg, W.R., Berry, J.A. (2013). Not all droughts are created equal: translating meteorological drought into woody plant mortality. *Tree Physiology* 33: 672-683.
- Anderegg, W. R., Berry, J. A., & Field, C. B. (2012a). Linking definitions, mechanisms, and modeling of drought-induced tree death. *Trends in plant science*, 17(12), 693-700.
- Anderegg, W. R., Klein, T., Bartlett, M., Sack, L., Pellegrini, A. F., Choat, B., & Jansen, S. (2016). Meta-analysis reveals that hydraulic traits explain cross-species patterns of drought-induced tree mortality across the globe. *Proceedings of the National Academy of Sciences*, 113(18), 5024-5029.
- Anderegg, W.R. (2014). Spatial and temporal variation in plant hydraulic traits and their relevance for climate change impacts on vegetation. *New phytologist* 205: 1008-1041.
- Anderegg, W.R., Berry, J.A., Smith, D.D., Sperry, J.S., Anderegg, L.D.L., Field, C.B. (2012b). The roles of hydraulic and carbon stress in a widespread climate- induced forest die-off. *Proceedings of the National Academy of Sciences USA* 109: 233–237

- Anderegg, W.R., Flint, A., Huang, C.Y., Flint, L., Berry, J.A., Davis, F.W., Sperry, J.S. and Field, C.B. (2015). Tree mortality predicted from drought-induced vascular damage. *Nature Geoscience*, 8(5), 367-371.
- Angadi, S. V., & Entz, M. H. (2002a). Water relations of standard height and dwarf sunflower cultivars. *Crop science*, 42(1), 152-159.
- Angadi, S. V., & Entz, M. H. (2002b). Agronomic performance of different stature sunflower cultivars under different levels of interplant competition. *Canadian journal of plant science*, 82(1), 43-52.
- Angeles G, Bond B, Boyer JS, Brodribb TJ, Brooks JR, Burns MJ, Cavender-Bares J, Clearwater M, Cochard H, Comstock J, et al (2004) The Cohesion-Tension Theory. *New Phytologist* 163: 451–452
- Aranda, I., Cano, F.J., Gascó, A., Cochard, H., Nardini, A., Mancha, J.A., López, R. and Sánchez-Gómez, D. (2014). Variation in photosynthetic performance and hydraulic architecture across European beech (*Fagus sylvatica* L.) populations supports the case for local adaptation to water stress. *Tree physiology*, 35(1), 34-46.
- Ashraf, M., & O'Leary, J. W. (1996). Effect of drought stress on growth, water relations, and gas exchange of two lines of sunflower differing in degree of salt tolerance. *International Journal of Plant Sciences*, 157(6), 729-732.
- Awad, H., Herbette, S., Brunel, N., Tixier, A., Pilate, G., Cochard, H., & Badel, E. (2012). No trade-off between hydraulic and mechanical properties in several transgenic poplars modified for lignins metabolism. *Environmental and Experimental Botany*, 77, 185-195.
- Bailey, I.W. (1913). The preservative treatment of wood. II. The structure of the pit membranes in the tracheids of conifers and their relation to the penetration of gases, liquids and finely divided solids into green and seasoned wood. *Forestry Quarterly* 11: 12-20.
- Bailey, I.W. (1916). The Structure of the Bordered Pits of Conifers and Its Bearing Upon the Tension Hypothesis of the Ascent of Sap in Plants. *Bot Gaz* 62: 133.
- Barigah TS, Bonhomme M, Lopez D, Traore A, Douris M, Venisse JS, Cochard H, Badel E. (2013). Modulation of bud survival in *Populus nigra* sprouts in response to water stress-induced embolism. *Tree Physiology* 33: 261-274.
- Bentz, B.J., Régnière, J., Fettig, C.J., Hansen, E.M., Hayes, J.L., Hicke, J.A., Kelsey, R.G., Negrón, J.F. and Seybold, S.J. (2010). Climate change and bark beetles of the western United States and Canada: direct and indirect effects. *BioScience*, 60(8), 602-613.
- Bigler, C., Bräker, O. U., Bugmann, H., Dobbertin, M., & Rigling, A. (2006). Drought as an inciting mortality factor in Scots pine stands of the Valais, Switzerland. *Ecosystems*, 9(3), 330-343.

- Blackman, C. J., Brodribb, T. J., & Jordan, G. J. (2010). Leaf hydraulic vulnerability is related to conduit dimensions and drought resistance across a diverse range of woody angiosperms. *New Phytologist*, 188(4), 1113-1123.
- Blum, A. (2005). Drought resistance, water-use efficiency, and yield potential—are they compatible, dissonant, or mutually exclusive? *Australian Journal of Agricultural Research*, 56(11), 1159-1168.
- Boisvenue, C., & Running, S. W. (2006). Impacts of climate change on natural forest productivity—evidence since the middle of the 20th century. *Global Change Biology*, 12(5), 862-882.
- Bonan, G. B. (2008). Forests and climate change: forcings, feedbacks, and the climate benefits of forests. *Science*, 320(5882), 1444-1449.
- Bouche PS, Delzon S, Choat B, Badel E, Brodribb TJ, Burlett R, Cochard H, Charra-Vaskou K, Lavigne B, Li S, et al (2016). Are needles of *Pinus pinaster* more vulnerable to xylem embolism than branches? New insights from X-ray computed tomography. *Plant, cell & environment*, 39(4), 860-870.
- Bouche, P. S., Larter, M., Domec, J. C., Burlett, R., Gasson, P., Jansen, S., & Delzon, S. (2014). A broad survey of hydraulic and mechanical safety in the xylem of conifers. *Journal of Experimental Botany*, 65(15), 4419-4431.
- Bouche, PS. (2015). Cavitation Resistance and the Functional Role of Bordered Pits in Xylem of Conifers: From Inter-specific to Within Tree Variability (Doctoral dissertation, Universität Ulm).
- Bréda N, Cochard H, Dreyer E, Granier A. (1993). Field comparison of transpiration, stomatal conductance and vulnerability to cavitation of *Quercus petraea* and *Quercus robur* under water stress. *Annales des Sciences Forestières* 50, 571–582.
- Bréda, N., Huc, R., Granier, A., & Dreyer, E. (2006). Temperate forest trees and stands under severe drought: a review of ecophysiological responses, adaptation processes and long-term consequences. *Annals of Forest Science*, 63(6), 625-644.
- Brodersen CR, McElrone AJ, Choat B, Matthews MA, Shackel KA. (2010). The dynamics of embolism repair in xylem: in vivo visualizations using high-resolution computed tomography. *Plant Physiology* 154: 1088–1095.
- Brodribb T, Hill RS (1999). The importance of xylem constraints in the distribution of conifer species. *New Phytol* 143: 365–372
- Brodribb TJ, Bowman DJMS, Nichols S, Delzon S, Burlett R. (2010). Xylem function and growth rate interact to determine recovery rates after exposure to extreme water deficit. *New Phytologist* 188: 533-542.
- Brodribb, T. J., & Cochard, H. (2009). Hydraulic failure defines the recovery and point of death in water-stressed conifers. *Plant physiology*, 149(1), 575-584.

- Brodribb, T. J., & Feild, T. S. (2000). Stem hydraulic supply is linked to leaf photosynthetic capacity: evidence from New Caledonian and Tasmanian rainforests. *Plant, Cell & Environment*, 23(12), 1381-1388.
- Brodribb, T. J., Holbrook, N. M., Edwards, E. J., & Gutierrez, M. V. (2003). Relations between stomatal closure, leaf turgor and xylem vulnerability in eight tropical dry forest trees. *Plant, Cell & Environment*, 26(3), 443-450.
- Brodribb, T. J., McAdam, S. A., Jordan, G. J., & Martins, S. C. (2014). Conifer species adapt to low-rainfall climates by following one of two divergent pathways. *Proceedings of the National Academy of Sciences*, 111(40), 14489-14493.
- Brodribb, T. J., Skelton, R. P., McAdam, S. A., Bienaimé, D., Lucani, C. J., & Marmottant, P. (2016). Visual quantification of embolism reveals leaf vulnerability to hydraulic failure. *New Phytologist*, 209(4), 1403-1409.
- Butcher, T. B. (2007). Achievements in forest tree genetic improvement in Australia and New Zealand 7: Maritime pine and Brutian pine tree improvement programs in Western Australia. *Australian forestry*, 70(3), 141-151.
- Canny MJ. (1997). Vessel contents during transpiration: embolisms and refilling. *American Journal of Botany* 84: 1223–1230.
- Carlquist, S. (1966). Wood anatomy of Compositae: a summary, with comments on factors controlling wood evolution. *Aliso: A Journal of Systematic and Evolutionary Botany*, 6(2), 25-44.
- Carnicer, J., Coll, M., Ninyerola, M., Pons, X., Sanchez, G., & Penuelas, J. (2011). Widespread crown condition decline, food web disruption, and amplified tree mortality with increased climate change-type drought. *Proceedings of the National Academy of Sciences*, 108(4), 1474-1478.
- Charra-Vaskou K, Badel E, Burlett R, Cochard H, Delzon S, Mayr S. (2012). Hydraulic efficiency and safety of vascular and non-vascular components in *Pinus pinaster* leaves. *Tree Physiology* 32: 1161-1170.
- Chavarria G and Pessoa dos Santos H. (2012). *Plant Water Relations: Absorption, Transport and Control Mechanisms. Advances in Selected Plant Physiology Aspects*, Dr. Giuseppe Montanaro (Ed.), ISBN: 978-953-51-0557-2, InTech.
- Choat, B., Badel, E., Burlett, R., Delzon, S., Cochard, H., Jansen, S. (2016). Noninvasive Measurement of Vulnerability to Drought-Induced Embolism by X-Ray Microtomography. *Plant Physiol* 170: 273–282
- Choat, B., Cobb, A. R., & Jansen, S. (2008). Structure and function of bordered pits: new discoveries and impacts on whole-plant hydraulic function. *New phytologist*, 177(3), 608-626.

- Choat, B., Drayton, W. M., Brodersen, C., Matthews, M. A., Shackel, K. A., Wada, H., & McElrone, A. J. (2010). Measurement of vulnerability to water stress-induced cavitation in grapevine: a comparison of four techniques applied to a long-vesseled species. *Plant, Cell & Environment*, 33(9), 1502-1512.
- Choat, B., Jansen, S., Brodribb, T. J., Cochard, H., Delzon, S., Bhaskar, R., Bucci, S. J., Field, T. S., Gleason, S. M., Hacke, U. G., et al. (2012). Global convergence in the vulnerability of forests to drought. *Nature* 491: 752–755.
- Choat, B., Lahr, E., Melcher, P. J., Zwieniecki, M. A., Holbrook, N. M. (2005). The spatial pattern of air seeding thresholds in mature sugar maple trees. *Plant, Cell & Environment* 28: 1082–1089.
- Christman, M.A., Sperry, J.S., Adler, F. R. (2009). Testing the ‘rare pit’ hypothesis for xylem cavitation resistance in three species of *Acer*. *New Phytologist* 182: 664-674.
- Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogée, J., Allard, V., Aubinet, M., Buchmann, N., Bernhofer, C., Carrara, A. and Chevallier, F. (2005). Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature*, 437(7058), 529-533.
- CNPPF-IDF, (2015). Centre national professionnel de la propriété forestière - Institut pour le développement forestier. Private Forest Property in France – Key Figures. 2015.
- Cochard H, Badel E, Herbette S, Delzon S, Choat B, Jansen S. (2013). Methods for measuring plant vulnerability to cavitation: a critical review. *Journal of Experimental Botany* 64: 4779-4791.
- Cochard H, Barigah ST, Kleinhentz M, Eshel A. (2008). Is xylem cavitation resistance a relevant criterion for screening drought resistance amongst *Prunus* species? *Journal of Plant Physiology* 165: 976-982.
- Cochard H, Cruizat P, Tyree MT. (1992). Use of positive pressures to establish vulnerability curves. Further support for the air-seeding hypothesis and implications for pressure-volume analysis. *Plant Physiology* 100: 205–209.
- Cochard H, Damour G, Bodet C, Tharwat I, Poirier M, Améglio T. (2005). Evaluation of a new centrifuge technique for rapid generation of xylem vulnerability curves. *Physiologia Plantarum* 124: 410–418.
- Cochard H, Delzon S, Badel E (2015). X-ray microtomography (micro-CT): a reference technology for high-resolution quantification of xylem embolism in trees. *Plant Cell Environ* 38: 201–206
- Cochard H, Froux F, Mayr S, Coutand C. (2004). Xylem wall collapse in water-stressed pine needles *Plant Physiology* 134: 401-408.
- Cochard H, Holtta, Herbette S, Delzon S, Mencuccini M. (2009). New Insights into the Mechanisms of Water-Stress Induced Cavitation in Conifers. *Plant Physiology* 151: 949-954.
- Cochard H. (2006). Cavitation in trees. *Comptes Rendus Physique* 7: 1018–1126.

- Cochard, H. (2002a). A technique for measuring xylem hydraulic conductance under high negative pressures. *Plant, Cell & Environment* 25: 815–819.
- Cochard, H. (2002b). Xylem embolism and drought-induced stomatal closure in maize. *Planta*, 215(3), 466-471.
- Corcuera, L., Cochard, H., Gil-Pelegrin, E., & Notivol, E. (2011). Phenotypic plasticity in mesic populations of *Pinus pinaster* improves resistance to xylem embolism (P_{50}) under severe drought. *Trees*, 25(6), 1033-1042.
- Cornelissen, J. H. C., & Thompson, K. (1997). Functional leaf attributes predict litter decomposition rate in herbaceous plants. *The New Phytologist*, 135(1), 109-114.
- Crombie DS, Hipkins MF, Milburn JA. (1985a). Gas penetration of pit membranes in the xylem of *Rhododendron* as the cause of acoustically detectable sap cavitation. *Australian Journal of Plant Physiology* 12: 445-53.
- Crombie DS, Milburn JA, Hipkins MF. (1985b). Maximum sustainable xylem sap tensions in *Rhododendron* and other species. *Planta* 163 : 27-33.
- Cruziat P, Améglio T, Cochard H. (2001). La cavitation: un mécanisme perturbant la circulation de l'eau chez les végétaux. *Mécanique et Industries*. 2: 289-298.
- Dai, A. (2013). Increasing drought under global warming in observations and models. *Nature Climate Change*, 3(1), 52-58.
- Dalla-Salda G, Martinez-Meier A, Cochard H, Rozenberg P. (2009). Variation of wood density and hydraulic properties of Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) clones related to a heat and drought wave in France. *Forest Ecology and Management* 257: 182-189.
- Davis SD, Ewers FW, Sperry JS, Portwood KA, Crocker MC, Adams GC. (2002). Shoot dieback during prolonged drought in *Ceanothus* (Rhamnaceae) chaparral of California: a possible case of hydraulic failure. *American Journal of Botany* 89: 820-828.
- Delzon S, Douthe C, Sala A, Cochard H. (2010). Mechanism of water-stress induced cavitation in conifers: bordered pit structure and function support the hypothesis of seal capillary-seeding. *Plant Cell and Environment* 33: 2101–2111.
- Delzon, S., & Cochard, H. (2014). Recent advances in tree hydraulics highlight the ecological significance of the hydraulic safety margin. *New Phytologist*, 203(2), 355-358.
- Dilley M, Chen RS, Deichmann U, Lerner-Lam AL, Arnold M (2005) Natural disaster hotspots: a global risk analysis. Synthesis report. International Bank for Reconstruction and Development, Washington, DC. Available at <http://sedac.ciesin.columbia.edu/hazards/hotspots/synthesisreport.pdf>
- Dixon HH and Joly J. (1895). On the ascent of sap. *Philosophical Transactions of the Royal Society of London B* 186: 563–576.

- Domec JC, Lachenbruch B, Meinzer FC. (2006). Bordered pit structure and function determine spatial patterns of air-seeding thresholds in xylem of Douglas-fir (*Pseudotsuga menziesii*; Pinaceae) trees. *American Journal of Botany* 93: 1588–1600.
- Domec, J. C., Lachenbruch, B., Meinzer, F. C., Woodruff, D. R., Warren, J. M., & McCulloh, K. A. (2008). Maximum height in a conifer is associated with conflicting requirements for xylem design. *Proceedings of the National Academy of Sciences*, 105(33), 12069-12074.
- Ducrey, M., Huc, R., Ladjal, M., & Guehl, J. (2008). Variability in growth, carbon isotope composition, leaf gas exchange and hydraulic traits in the eastern Mediterranean cedars *Cedrus libani* and *C. brevifolia*. *Tree Physiology-Victoria*, 28(5), 689-701.
- Elder, B. D., Rehill, B. J., Haynes, K. J., & Dwyer, G. (2013). Induced plant defenses, host–pathogen interactions, and forest insect outbreaks. *Proceedings of the National Academy of Sciences*, 110(37), 14978-14983.
- FAO (2006). Global forest resources assessment 2005-progress towards sustainable forest management. FAO Forestry Paper No. 147. Rome.
- Fichot, R., Barigah, T. S., Chamaillard, S., Le Thiec, D., Laurans, F., Cochard, H., & Brignolas, F. (2010). Common trade-offs between xylem resistance to cavitation and other physiological traits do not hold among unrelated *Populus deltoides* × *Populus nigra* hybrids. *Plant, Cell & Environment*, 33(9), 1553-1568.
- Flexas, J., & Medrano, H. (2002). Drought-inhibition of photosynthesis in C3 plants: stomatal and non-stomatal limitations revisited. *Annals of botany*, 89(2), 183-189.
- Fromm JH, Sautter I, Matthies D, Kremer J, Schumacher P, Ganter C. (2001). Xylem water content and wood density in spruce and oak trees detected by high-resolution computed tomography. *Plant Physiology* 127: 416–425.
- Gleason, S.M., Westoby, M., Jansen, S., Choat, B., Hacke, U.G., Pratt, R.B., Bhaskar, R., Brodribb, T.J., Bucci, S.J., Cao, K.F. and Cochard, H. (2015). Weak tradeoff between xylem safety and xylem-specific hydraulic efficiency across the world's woody plant species. *New Phytologist*, 209(1), 123-136.
- Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., Pretty, J., Robinson, S., Thomas, S.M. & Toulmin, C. (2010). Food security: the challenge of feeding 9 billion people. *Science*, 327(5967), 812-818.
- GOP, (2015). Cotton: Pakistan Economic Survey 2013-14, Finance division, Economic Advisor's wing, Islamabad, pp. 19.
- Guadagno, C.R., Ewers, B.E., Speckman, H.N., Aston, T.L., Huhn, B.J., DeVore, S.B., Ladwig, J.T., Strawn, R.N. and Weinig, C. (2017). Dead or alive? Using membrane failure and chlorophyll fluorescence to predict mortality from drought. *Plant Physiology*, pp-00581.
- Guet, J., Fichot, R., Lédée, C., Laurans, F., Cochard, H., Delzon, S., Bastien, C. and Brignolas, F. (2015). Stem xylem resistance to cavitation is related to xylem structure but not to growth

- and water-use efficiency at the within-population level in *Populus nigra* L. *Journal of experimental botany*, 66(15), 4643-4652.
- Hacke, U. G., Sperry, J. S., Pittermann, J. (2004). Analysis of circular bordered pit function-II. Gymnosperm tracheids with torus–margo pit membranes. *American Journal of Botany* 91: 386-400.
- Hacke, U. G., Sperry, J. S., Pockman, W. T., Davis, S. D., & McCulloh, K. A. (2001). Trends in wood density and structure are linked to prevention of xylem implosion by negative pressure. *Oecologia*, 126(4), 457-461.
- Hacke, U. G., Sperry, J. S., Wheeler, J. K., & Castro, L. (2006). Scaling of angiosperm xylem structure with safety and efficiency. *Tree physiology*, 26(6), 689-701.
- Hacke, U. G., Venturas, M. D., MacKinnon, E. D., Jacobsen, A. L., Sperry, J. S., & Pratt, R. B. (2015). The standard centrifuge method accurately measures vulnerability curves of long-vesselled olive stems. *New Phytologist*, 205(1), 116-127.
- Haider, S., & Adnan, S. (2014). Classification and assessment of aridity over Pakistan provinces (1960-2009). *International Journal of Environment*, 3(4), 24-35.
- Hajek, P., Kurjak, D., von Wühlisch, G., Delzon, S., & Schuldt, B. (2016). Intraspecific variation in wood anatomical, hydraulic and foliar traits in ten European beech provenances differing in growth yield. *Frontiers in Plant Science*, 7, 791.
- Hajek, P., Leuschner, C., Hertel, D., Delzon, S., & Schuldt, B. (2014). Trade-offs between xylem hydraulic properties, wood anatomy and yield in *Populus*. *Tree Physiol*, 34(7), 744-756.
- Helmer, M., & Hilhorst, D. (2006). Natural disasters and climate change. *Disasters*, 30(1), 1-4.
- Hochberg, U., Windt, C. W., Ponomarenko, A., Zhang, Y. J., Gersony, J., Rockwell, F. E., & Holbrook, N. M. (2017). Stomatal closure, basal leaf embolism and shedding protect the hydraulic integrity of grape stems. *Plant Physiology*, pp-01816.
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K, et al (2007). Coral Reefs Under Rapid Climate Change and Ocean Acidification. *Science* (80-) 318: 1737–1742
- Holbrook NM, Ahrens ET, Burns MJ, Zwieniecki MA. (2001). In vivo observation of cavitation and embolism repair using magnetic resonance imaging. *Plant Physiology* 126: 27–31.
- Hsiao, T. C., & Xu, L. K. (2000). Sensitivity of growth of roots versus leaves to water stress: biophysical analysis and relation to water transport. *Journal of experimental botany*, 51(350), 1595-1616.
- Hussain, S. S. (2000). Patronage of Non-traditional Oilseed Crops. *Pakistan J. Agri. Econ*, 4.
- IGN. (2016). Le mémento, inventaire forestier, la forêt en chiffres et en cartes. St Mandé, IGN. 17 pp.

- IPCC (2014). *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Field CB, VR Barros, DJ Dokken, KJ Mach, MD Mastrandrea, TE Bilir, M Chatterjee, KL Ebi, YO Estrada, RC Genova, B Girma, ES Kissel, AN Levy, S MacCracken, PR Mastrandrea, LL White (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1132 pp
- Iqbal, N., Ashraf, M., & Ashraf, M. Y. (2008). Glycinebetaine, an osmolyte of interest to improve water stress tolerance in sunflower (*Helianthus annuus* L.): water relations and yield. *South African Journal of Botany*, 74(2), 274-281.
- Jansen S, Choat B, Pletsers A. (2009). Morphological variation of intervessel pit membranes and implications to xylem function in angiosperms. *American Journal of Botany* 96: 409-419.
- Jansen S, Lamy JB, Burlett R, Cochard H, Gasson P, Delzon S. (2012). Plasmodesmatal pores in the torus of bordered pit membranes affect cavitation resistance of conifer xylem. *Plant, Cell & Environment* 35: 1109-1120.
- Jarbeau J, Ewers F, Davis S. (1995). The mechanism of water-stressed induced embolism in two species of chaparral shrubs. *Plant, Cell & Environment* 18: 189-196.
- Joyo, M. A., Ram, N., Koondhar, M. A., & Abbasi, F. (2016). Impact Analysis: Farmer's Knowledge and Economics of Sunflower Production in Golarchi District of Badin Sindh Province of Pakistan.
- Kang, Y., Khan, S., & Ma, X. (2009). Climate change impacts on crop yield, crop water productivity and food security – A review. *Progress in Natural Science*, 19, 1665–1674.
- Kaufmann, I., Schulze-Till, T., Schneider, H. U., Zimmermann, U., Jakob, P., & Wegner, L. H. (2009). Functional repair of embolized vessels in maize roots after temporal drought stress, as demonstrated by magnetic resonance imaging. *New Phytologist*, 184(1), 245-256.
- Kramer, P. and Boyer, J. (1995). *Water relations of plants and soils*. Academic Press, San Diego
- Kursar, T. A., Engelbrecht, B. M., Burke, A., Tyree, M. T., El Omari, B., & Giraldo, J. P. (2009). Tolerance to low leaf water status of tropical tree seedlings is related to drought performance and distribution. *Functional Ecology*, 23(1), 93-102.
- Lamy J-B, Bouffier L, Burlett R, Plomion C, Cochard H, Delzon S. (2011). Uniform Selection as a Primary Force Reducing Population Genetic Differentiation of Cavitation Resistance across a Species Range. *PLOS ONE* 6(8): e23476.
- Lamy J-B, Delzon S, Bouche P, Alia R, Vendramin GG, Cochard H, Plomion C. (2014). Limited genetic variability and phenotypic plasticity detected for cavitation resistance in a Mediterranean pine. *New Phytologist* 201: 874-888.
- Lamy J-B, Plomion C, Kremer A, Delzon S (2012). QST is less than FST As a signature of canalization. *Mol Ecol* 21: 5646–5655

- Larter, M., Brodribb, T. J., Pfautsch, S., Burlett, R., Cochard, H., & Delzon, S. (2015). Extreme aridity pushes trees to their physical limits. *Plant physiology*, 168(3), 804-807.
- Larter, M., Pfautsch, S., Domec, J. C., Trueba, S., Nagalingum, N., & Delzon, S. (2017). Aridity drove the evolution of extreme embolism resistance and the radiation of conifer genus *Callitris*. *New Phytologist*. DOI: 10.1111/nph.14545
- Lens F, Sperry JS, Christman M, Choat B, Rabaey D, Jansen S. (2011). Testing hypotheses that link wood anatomy to cavitation resistance and hydraulic conductivity in the genus *Acer*. *New Phytologist* 190: 709-723.
- Lens, F., Picon-Cochard, C., Delmas, C.E., Signarbieux, C., Buttler, A., Cochard, H., Jansen, S., Chauvin, T., Doria, L.C., del Arco, M. and Delzon, S. (2016). Herbaceous angiosperms are not more vulnerable to drought-induced embolism than angiosperm trees. *Plant physiology*, pp-00829.
- Lens, F., Tixier, A., Cochard, H., Sperry, J. S., Jansen, S., & Herbette, S. (2013). Embolism resistance as a key mechanism to understand adaptive plant strategies. *Current Opinion in Plant Biology*, 16(3), 287-292.
- Li, S., Lens, F., Espino, S., Karimi, Z., Klepsch, M., Schenk, H. J., Schmitt M, Schuldt B., & Jansen, S. (2016). Intervessel pit membrane thickness as a key determinant of embolism resistance in angiosperm xylem. *Iawa Journal*, 37(2), 152-171.
- Li, Y., Sperry, J. S., & Shao, M. (2009). Hydraulic conductance and vulnerability to cavitation in corn (*Zea mays* L.) hybrids of differing drought resistance. *Environmental and Experimental Botany*, 66(2), 341-346.
- Lichtenthaler, H. K. (1996). Vegetation stress: an introduction to the stress concept in plants. *Journal of plant physiology*, 148(1-2), 4-14.
- López, R., Cano, F. J., Choat, B., Cochard, H., & Gil, L. (2016). Plasticity in vulnerability to cavitation of *Pinus canariensis* occurs only at the driest end of an aridity gradient. *Frontiers in plant science*, 7.
- López, R., López de Heredia, U., Collada, C., Cano, F. J., Emerson, B. C., Cochard, H., & Gil, L. (2013). Vulnerability to cavitation, hydraulic efficiency, growth and survival in an insular pine (*Pinus canariensis*). *Annals of botany*, 111(6), 1167-1179.
- Loustau, D., Bosc, A., Colin, A., Ogée, J., Davi, H., François, C., Dufrêne, E., Déqué, M., Cloppet, E., Arrouays, D. and Le Bas, C. (2005). Modeling climate change effects on the potential production of French plains forests at the sub-regional level. *Tree Physiology*, 25(7), 813-823.
- Ludlow, M. M., Fisher, M. J., & Wilson, J. R. (1985). Stomatal adjustment to water deficits in three tropical grasses and a tropical legume grown in controlled conditions and in the field. *Functional Plant Biology*, 12(2), 131-149.

- Maherali H, Pockman WT. (2004). Adaptive variation in the vulnerability of woody plants to xylem cavitation. *Ecology* 85: 2184-2199.
- Major, J. E., & Johnsen, K. H. (1999). Shoot water relations of mature black spruce families displaying a genotype× environment interaction in growth rate. II. Temporal trends and response to varying soil water conditions. *Tree physiology*, 19(6), 375-382.
- Martínez-Vilalta J, Cochard H, Mencuccini M, Sterck F, Herrero a., Korhonen JFJ, Llorens P, Nikinmaa E, Nolè a., Poyatos R, et al (2009). Hydraulic adjustment of Scots pine across Europe. *New Phytol* 184: 353–364
- Martínez-Vilalta J, Prat E, Oliveras I, Piñol J. (2002). Xylem hydraulic properties of roots and stems of nine Mediterranean woody species. *Oecologia* 133: 19-29.
- Martín-Retortillo, M., & Pinilla, V. (2015). On the causes of economic growth in Europe: Why did agricultural labour productivity not converge between 1950 and 2005? *Cliometrica*, 9, 359–396.
- Martin-StPaul, N. K., Longepierre, D., Huc, R., Delzon, S., Burlett, R., Joffre, R., ... & Cochard, H. (2014). How reliable are methods to assess xylem vulnerability to cavitation? The issue of ‘open vessel’ artifact in oaks. *Tree Physiology*, 34(8), 894-905.
- Martin-StPaul, N., Delzon, S., & Cochard, H. (2017). Plant resistance to drought depends on timely stomatal closure. *Ecology Letters*. DOI: 10.1111/ele.12851
- McCulloh KA, Johnson DM, Meinzer FC, Woodruff DR. (2014). The dynamic pipeline: hydraulic capacitance and xylem hydraulic safety in four tall conifer species. *Plant, Cell & Environment* 37: 1171-1183.
- McDowell, N., Pockman, W.T., Allen, C.D., Breshears, D.D., Cobb, N., Kolb, T., Plaut, J., Sperry, J., West, A., Williams, D.G. and Yezpez, E.A. (2008). Mechanisms of plant survival and mortality during drought: why do some plants survive while others succumb to drought? *New phytologist*, 178(4), 719-739.
- McDowell, N.G., Williams, A.P., Xu, C., Pockman, W.T., Dickman, L.T., Sevanto, S., Pangle, R., Limousin, J., Plaut, J., Mackay, D.S. and Ogee, J. (2016). Multi-scale predictions of massive conifer mortality due to chronic temperature rise. *Nature Climate Change*, 6(3), 295-300.
- Mitchell, D. (2008). A note on rising food prices (Vol. 4682). Washington, DC: World Bank.
- Nasim, W., Belhouchette, H., Ahmad, A., Habib-ur-Rahman, M., Jabran, K., Ullah, K., Fahad, S., Shakeel, M. and Hoogenboom, G. (2016). Modelling climate change impacts and adaptation strategies for sunflower in Pakistan. *Outlook on AGRICULTURE*, 45(1), 39-45.
- Neufeld, H. S., Grantz, D. A., Meinzer, F. C., Goldstein, G., Crisosto, G. M., & Crisosto, C. (1992). Genotypic variability in vulnerability of leaf xylem to cavitation in water-stressed and well-irrigated sugarcane. *Plant Physiology*, 100(2), 1020-1028.

- O'Grady, A. P., & Mitchell, P. J. (2015). Looking forward, looking back: capturing drought in flagrante delicto and uncovering its broader consequences for forest ecosystems. *Tree physiology*, 35(8), 803-805.
- Pammenter NW, Vander Willigen C. (1998). A mathematical and statistical analysis of the curves illustrating vulnerability to cavitation. *TreePhysiology* 18: 589-593.
- Patz JA, Campbell-Lendrum D, Holloway T, Foley JA (2005). Impact of regional climate change on human health. *Nature* 438: 310–317
- Payn, T., Carnus, J. M., Freer-Smith, P., Kimberley, M., Kollert, W., Liu, S., Orazio, C., Rodriguez, L., Silva, L.N. and Wingfield, M.J. (2015). Changes in planted forests and future global implications. *Forest Ecology and Management*, 352, 57-67.
- Peltonen-Sainio, P., & Jauhiainen, L. (2014). Lessons from the past in weather variability: sowing to ripening dynamics and yield penalties for northern agriculture from 1970 to 2012. *Regional environmental change*, 14(4), 1505-1516.
- Peng, C., Ma, Z., Lei, X., Zhu, Q., Chen, H., Wang, W., Liu, S., Li, W., Fang, X. and Zhou, X. (2011). A drought-induced pervasive increase in tree mortality across Canada's boreal forests. *Nature climate change*, 1: 467–471
- Pimentel, D., Harvey, C., Resosudarmo, P., Sinclair, K., Kurz, D., McNair, M., Crist, S., Shpritz, L., Fitton, L., Saffouri, R. and Blair, R. (1995). Environmental and economic costs of soil erosion and conservation benefits. *Science-AAAS-Weekly Paper Edition*, 267(5201), 1117-1122.
- Pinke, Z. and Lövei, G.L. (2017). Increasing temperature cuts back crop yields in Hungary over the last 90 years. *Global Change Biology*. DOI: 10.1111/gcb.13808
- Pittermann, J., Limm, E., Rico, C., & Christman, M. A. (2011). Structure–function constraints of tracheid-based xylem: a comparison of conifers and ferns. *New Phytologist*, 192(2), 449-461.
- Pittermann, J., Sperry, J. S., Hacke, U. G., Wheeler, J. K., & Sikkema, E. H. (2005). Torus-margo pits help conifers compete with angiosperms. *Science*, 310(5756), 1924-1924.
- Pittermann, J., Sperry, J. S., Wheeler, J. K., Hacke, U. G., & Sikkema, E. H. (2006). Mechanical reinforcement of tracheids compromises the hydraulic efficiency of conifer xylem. *Plant, Cell & Environment*, 29(8), 1618-1628.
- Pittermann, J., Stuart, S. A., Dawson, T. E., & Moreau, A. (2012). Cenozoic climate change shaped the evolutionary ecophysiology of the Cupressaceae conifers. *Proceedings of the National Academy of Sciences*, 109(24), 9647-9652.
- Pittermann, J., Choat, B., Jansen, S., Stuart, S., Lynn, L., Dawson, T.E. (2010). The relationships between xylem safety and hydraulic efficiency in the Cupressaceae: the evolution of pit membrane form and function. *Plant Physiology* 153: 1919-1931.

- Pivovarovff, A. L., Burlett, R., Lavigne, B., Cochard, H., Santiago, L. S., & Delzon, S. (2016). Testing the ‘microbubble effect’ using the Cavitron technique to measure xylem water extraction curves. *AoB Plants*, 8. plw011.
- Pockman WT, Sperry JS, O’Leary JW. (1995). Sustained and significant negative water pressure in xylem. *Nature* 378: 715-716.
- Poorter, L., McDonald, I., Alarcón, A., Fichtler, E., Licona, J.C., Peña-Claros, M., Sterck, F., Villegas, Z. and Sass-Klaassen, U. (2010). The importance of wood traits and hydraulic conductance for the performance and life history strategies of 42 rainforest tree species. *New phytologist*, 185(2), 481-492.
- Prasad, T. G., Sashidhar, V. R., Chari, M., Rao, S. R., Devendra, R., & Kumar, M. U. (1985). Regulation of water loss under moisture stress in sunflower genotypes: stomatal sensitivity in relation to stomatal frequency, diffusive resistances and transpiration rate, at different canopy positions. *The Journal of Agricultural Science*, 105(3), 673-678.
- Rachidi, F., Kirkham, M. B., Stone, L. R., & Kanemasu, E. T. (1993). Soil water depletion by sunflower and sorghum under rainfed conditions. *Agricultural Water Management*, 24(1), 49-62.
- Raffa, K. F., Aukema, B. H., Bentz, B. J., Carroll, A. L., Hicke, J. A., Turner, M. G., & Romme, W. H. (2008). Cross-scale drivers of natural disturbances prone to anthropogenic amplification: the dynamics of bark beetle eruptions. *AIBS Bulletin*, 58(6), 501-517.
- Ray, D. K., Gerber, J. S., MacDonald, G. K., & West, P. C. (2015). Climate variation explains a third of global crop yield variability. *Nature communications*, 6, 5989.
- Ray, D. K., Ramankutty, N., Mueller, N. D., West, P. C., & Foley, J. A. (2012). Recent patterns of crop yield growth and stagnation. *Nature communications*, 3, 1293.
- Reddy, A. R., Chaitanya, K. V., & Vivekanandan, M. (2004). Drought-induced responses of photosynthesis and antioxidant metabolism in higher plants. *Journal of plant physiology*, 161(11), 1189-1202.
- Rockwell FE, Wheeler JK, Holbrook NM (2014). Cavitation and Its Discontents: Opportunities for Resolving Current Controversies. *Plant Physiol* 164: 1649–1660.
- Rosner, S. (2017). Wood density as a proxy for vulnerability to cavitation: Size matters. *Journal of Plant Hydraulics*, 4, 001.
- Ryan MG, Phillips N, Bond BJ. (2006). The hydraulic limitation hypothesis revisited. *Plant, Cell & Environment* 29: 367–381.
- Sack, L., & Scoffoni, C. (2013). Leaf venation: structure, function, development, evolution, ecology and applications in the past, present and future. *New Phytologist*, 198(4), 983-1000.
- Sáenz-Romero C, Lamy J-B, Loya-Rebollar E, Plaza-Aguilar A, Burlett R, Lobit P, Delzon S (2013). Genetic variation of drought-induced cavitation resistance among *Pinus hartwegii* populations from an altitudinal gradient. *Acta Physiol Plant* 35: 2905–2913

- Sáenz-Romero, C., Larter, M., González-Muñoz, N., Wehenkel, C., Blanco-García, A., Castellanos-Acuña, D., Burlett, R. and Delzon, S. (2017). Mexican conifers differ in their capacity to face climate change. *Journal of Plant Hydraulics*, 4, 003.
- Santiago, L. S., Goldstein, G., Meinzer, F. C., Fisher, J. B., Machado, K., Woodruff, D., & Jones, T. (2004). Leaf photosynthetic traits scale with hydraulic conductivity and wood density in Panamanian forest canopy trees. *Oecologia*, 140(4), 543-550.
- Santiago, L. S., Goldstein, G., Meinzer, F. C., Fisher, J. B., Machado, K., Woodruff, D., & Jones, T. (2004). Leaf photosynthetic traits scale with hydraulic conductivity and wood density in Panamanian forest canopy trees. *Oecologia*, 140(4), 543-550.
- Schenk HJ, Steppe K, Jansen S. (2015). Nanobubbles: a new paradigm for air-seeding in xylem. *Trends in Plant Science* 20: 200-205.
- Schuldt, B., Knutzen, F., Delzon, S., Jansen, S., Müller-Haubold, H., Burlett, R., R., Clough, Y. and Leuschner, C. (2016). How adaptable is the hydraulic system of European beech in the face of climate change-related precipitation reduction? *New Phytologist*, 210(2), 443-458.
- Sperry JS and Ikeda T. (1997). Xylem cavitation in roots and stems of Douglas-fir and white fir. *Tree Physiology* 17: 275-280.
- Sperry JS, Donnelly JR, Tyree MT. (1988). A method for measuring hydraulic conductivity and embolism in xylem. *Plant, Cell & Environment* 11: 35–40.
- Sperry JS, Hacke UG. (2004). Analysis of circular bordered pit function. I. Angiosperm vessels with homogenous pit membranes. *American Journal of Botany* 91: 369-385.
- Sperry JS, Saliendra NZ, Pockman WT, Cochard H, Cruiziat P, Davis SD, Ewers FW, Tyree MT. (1996). New evidence for large negative xylem pressures and their measurement by the pressure chamber method. *Plant, Cell & Environment* 19: 427-436.
- Sperry JS, Sullivan JE (1992). Xylem embolism in response to freeze-thaw cycles and water stress in ring-porous, diffuse-porous, and conifer species. *Plant Physiol* 100: 605–613.
- Sperry JS, Tyree MT (1988). Mechanism of water stress-induced xylem embolism. *Plant Physiol* 88: 581-7.
- Sperry JS, Tyree MT. (1990). Water-stress-induced xylem embolism in three species of conifers. *Plant, Cell & Environment* 13: 427–436.
- Sperry, J. S. (2000). Hydraulic constraints on plant gas exchange. *Agricultural and forest meteorology*, 104(1), 13-23.
- Sperry, J. S., Christman, M. A., Torres-Ruiz, J. M., Taneda, H., & Smith, D. D. (2012). Vulnerability curves by centrifugation: is there an open vessel artefact, and are ‘r’ shaped curves necessarily invalid?. *Plant, Cell & Environment*, 35(3), 601-610.
- Sperry, J. S., Hacke, U. G., & Pittermann, J. (2006). Size and function in conifer tracheids and angiosperm vessels. *American journal of botany*, 93(10), 1490-1500.

- Sperry, J. S., Meinzer, F. C., & McCULLOH, K. A. (2008). Safety and efficiency conflicts in hydraulic architecture: scaling from tissues to trees. *Plant, Cell & Environment*, 31(5), 632-645.
- Sperry, J. S., Stiller, V., & Hacke, U. G. (2003). Xylem hydraulics and the soil–plant–atmosphere continuum. *Agronomy Journal*, 95(6), 1362-1370.
- Sterck, F. J., Martínez-Vilalta, J., Mencuccini, M., Cochard, H., Gerrits, P., Zweifel, R., ... & Nole, A. (2012). Understanding trait interactions and their impacts on growth in Scots pine branches across Europe. *Functional Ecology*, 26(2), 541-549.
- Stiller, V., & Sperry, J. S. (2002). Cavitation fatigue and its reversal in sunflower (*Helianthus annuus* L.). *Journal of experimental botany*, 53(371), 1155-1161.
- Stiller, V., Lafitte, H. R., & Sperry, J. S. (2003). Hydraulic properties of rice and the response of gas exchange to water stress. *Plant Physiology*, 132(3), 1698-1706.
- Stocker TF, Qin D, Plattner G-K, Tignor MM, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM (2013). IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA
- Tardieu, F. (2011). Any trait or trait-related allele can confer drought tolerance: just design the right drought scenario. *Journal of experimental botany*, 63(1), 25-31.
- Tester, M., & Langridge, P. (2010). Breeding technologies to increase crop production in a changing world. *Science*, 327(5967), 818-822.
- Tixier A, Cochard H, Badel E, Dusotoit-Coucaud A, Jansen S, Herbette S (2013). *Arabidopsis thaliana* as a model species for xylem hydraulics: Does size matter? *J Exp Bot* 64: 2295–2305
- Torres-Ruiz JM, Cochard H, Mayr S, Beikircher B, Diaz-Espejo A, Rodriguez-Dominguez CM, Badel E, Fernández JE (2014). Vulnerability to cavitation in *Olea europaea* current-year shoots: further evidence of an open-vessel artifact associated with centrifuge and air-injection techniques. *Physiol Plant* 152: 465–474.
- Torres-Ruiz JM, Jansen S, Choat B, McElrone A, Cochard H, Brodribb TJ, Badel E, Burlett R, Bouche PS, Brodersen C, Li S, Morris H, Delzon S. (2015). Direct X-ray microtomography observation confirms the induction of embolism upon xylem cutting under tension. *Plant Physiology* 167: 40-43.
- Torres-Ruiz, J. M., Cochard, H., Mencuccini, M., Delzon, S., & Badel, E. (2016). Direct observation and modelling of embolism spread between xylem conduits: a case study in Scots pine. *Plant, cell & environment*, 39(12), 2774-2785.
- Torres-Ruiz, J.M., Cochard, H., Choat, B., Jansen, S., López, R., Tomášková, I., Padilla-Díaz, C.M., Badel, E., Burlett, R., King, A. and Lenoir, N. (2017). Xylem resistance to embolism:

- presenting a simple diagnostic test for the open vessel artefact. *New Phytologist*, 215(1), 489-499.
- Trifilò, P., Nardini, A., Lo Gullo, M. A., & Salleo, S. (2003). Vein cavitation and stomatal behaviour of sunflower (*Helianthus annuus*) leaves under water limitation. *Physiologia Plantarum*, 119(3), 409-417.
- Trnka, M., Olesen, J. E., Kersebaum, K. C., Rötter, R. P., Brázdil, R., Eitzinger, J., ... & Balek, J. (2016). Changing regional weather-crop yield relationships across Europe between 1901 and 2012. *Climate Research* 70: 195-214
- Trnka, M., Olesen, J.E., Kersebaum, K.C., Skjelvåg, A.O., Eitzinger, J., Seguin, B., PELTONEN-SAINIO, P., Rötter, R., Iglesias, A.N.A., Orlandini, S. and Dubrovský, M. (2011). Agroclimatic conditions in Europe under climate change. *Global Change Biology*, 17(7), 2298-2318.
- Tyree MT and Ewers F. 1991. The hydraulic architecture of trees and other woody plants. *New Phytologist* 119: 345-360.
- Tyree MT, Alexander J, Machado JL. (1992). Loss of hydraulic conductivity due to water stress in intact juveniles of *Quercus rubra* and *Populus deltoides*. *Tree Physiology* 10: 411-415.
- Tyree MT, Dixon MA, Tyree EL, Johnson R. (1984). Ultrasonic acoustic emissions from the sapwood of cedar and hemlock. An examination of three hypotheses regarding cavitations. *Plant Physiology* 75: 988-992.
- Tyree MT, Sperry JS. (1989a). Characterization and propagation of acoustic emission signals in woody plants: towards an improved acoustic emission counter. *Plant, Cell & Environment* 12: 371-382.
- Tyree MT, Zimmermann MH (2002). *Xylem Structure and the Ascent of Sap*. Springer Science & Business Media
- Tyree, M. T., & Sperry, J. S. (1989b). Vulnerability of xylem to cavitation and embolism. *Annual review of plant biology*, 40(1), 19-36.
- Tyree, M. T., Fiscus, E. L., Wullschlegel, S. D., & Dixon, M. A. (1986). Detection of xylem cavitation in corn under field conditions. *Plant Physiology*, 82(2), 597-599.
- UNDP (United Nations Development Programme) (2004) *Reducing disaster risk: a challenge for development. A global report*. UNDP, New York. Available at www.undp.org/cpr/whats_new/rdr_english.pdf
- Urli M, Porté AJ, Cochard H, Guengant Y, Burlett R, Delzon S. (2013). Xylem embolism threshold for catastrophic hydraulic failure in angiosperm trees. *Tree Physiology* 33: 672-683.
- Van Mantgem, P.J., Stephenson, N.L., Byrne, J.C., Daniels, L.D., Franklin, J.F., Fulé, P.Z., Harmon, M.E., Larson, A.J., Smith, J.M., Taylor, A.H. and Veblen, T.T. (2009). Widespread increase of tree mortality rates in the western United States. *Science*, 323(5913), 521-524.

- Van Wart, J., Kersebaum, K. C., Peng, S., Milner, M., Cassman, K. G. (2013). Estimating crop yield potential at regional to national scales. *Field Crops Research*, 143, 34–43.
- Vinya R, Malhi Y, Fisher JB, Brown N, Brodribb TJ, Aragoa LE. (2013). Xylem cavitation vulnerability influences tree species' habitat preferences in miombo woodlands. *Oecologia* 113: 711-720.
- Walther G-R, Post E, Convey P, Menzel A, Parmesan C, Beebee TJ, Fromentin J-M, Hoegh-Guldberg O, Bairlein F (2002). Ecological responses to recent climate change. *Nature* 416: 389–395
- Wang R, Zhang L, Zhang S, Cai J, Tyree MT (2014). Water relations of *Robinia pseudoacacia* L.: do vessels cavitate and refill diurnally or are R-shaped curves invalid in *Robinia*? *Plant Cell Environ.* doi: 10.1111/pce.12315
- Watkins, J. E., Holbrook, N. M., & Zwieniecki, M. A. (2010). Hydraulic properties of fern sporophytes: consequences for ecological and evolutionary diversification. *American Journal of Botany*, 97(12), 2007-2019.
- Wei C, Tyree MT, Steudle E. (1999). Direct measurement of xylem pressure in leaves of intact maize plants: a test of the cohesion-tension theory taking hydraulic architecture into consideration. *Plant Physiology* 121: 1191-1205.
- Wheeler, J. K., Huggett, B. A., Tofte, A. N., Rockwell, F. E., & Holbrook, N. M. (2013). Cutting xylem under tension or supersaturated with gas can generate PLC and the appearance of rapid recovery from embolism. *Plant, Cell & Environment*, 36(11), 1938-1949.
- Wheeler, T., & Von Braun, J. (2013). Climate change impacts on global food security. *Science*, 341(6145), 508-513.
- Wikberg, J., & Ögren, E. (2004). Interrelationships between water use and growth traits in biomass-producing willows. *Trees*, 18(1), 70-76.
- Wise, R. R., Frederick, J. R., Alm, D. M., Kramer, D. M., Hesketh, J. D., Crofts, A. R., & Ort, D. R. (1990). Investigation of the limitations to photosynthesis induced by leaf water deficit in field-grown sunflower (*Helianthus annuus* L.). *Plant, Cell & Environment*, 13(9), 923-931.
- Wortemann R, Herbette S, Barigah TS, Fumanal B, Alia R, Ducouso A, Gomory D, Roedel-Drevet P, Cochard H. (2011). Genotypic variability and phenotypic plasticity of cavitation resistance in *Fagus sylvatica* L. across Europe. *Tree Physiology* 31: 1175-1182.
- Wright, I. J., & Westoby, M. (1999). Differences in seedling growth behaviour among species: trait correlations across species, and trait shifts along nutrient compared to rainfall gradients. *Journal of Ecology*, 87(1), 85-97.
- Xoconostle-Cázares, B., Ramirez-Ortega, F. A., Flores-Elenes, L., & Ruiz-Medrano, R. (2010). Drought tolerance in crop plants. *Am J Plant Physiol*, 5(5), 1-16.
- Zhao M, Running SW (2010). Drought-Induced Reduction in Global Terrestrial Net Primary Production from 2000 Through 2009. *Science* (80-) 329: 940–943.

Zimmermann MH. (1983). Xylem Structure and the Ascent of Sap. Springer Verlag, Berlin