



# Eigenvalues and eigenvectors of large matrices under random perturbations

Alkéos Michaël

## ► To cite this version:

Alkéos Michaël. Eigenvalues and eigenvectors of large matrices under random perturbations. Functional Analysis [math.FA]. Université Sorbonne Paris Cité, 2018. English. NNT : 2018USPCB214 . tel-02468213

**HAL Id: tel-02468213**

**<https://theses.hal.science/tel-02468213>**

Submitted on 5 Feb 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



UNIVERSITÉ  
**PARIS**  
**DESCARTES**

**U-PC**  
Université Sorbonne  
Paris Cité

**UNIVERSITÉ PARIS DESCARTES**

Laboratoire MAP5 UMR CNRS 8145

École doctorale 386 : Sciences Mathématiques de Paris Centre

**THÈSE**

Pour obtenir le grade de

**DOCTEUR EN MATHÉMATIQUES**

*Spécialité : Mathématiques appliquées*

Présentée par

**ALKÉOS MICHAÏL**

---

# Eigenvalues and eigenvectors of large matrices under random perturbations

---

Soutenue publiquement le 6 juillet 2018 devant un jury composé de

Florent BENAYCH-GEORGES	Université Paris Descartes	Directeur de thèse
Djalil CHAFAÏ	Université Paris-Dauphine	Examineur
Nathanaël ENRIQUEZ	Université Paris Sud	Examineur
Jamal NAJIM	Université de Marne La Vallée	Rapporteur
Sean O'ROURKE	University of Colorado Boulder	Rapporteur



---

## ABSTRACT

---

The present thesis is devoted to the study of the effect of a perturbation on the spectrum of a Hermitian matrix by a random matrix with small operator norm and whose entries in the eigenvector basis of the first one were independent, centered and with a variance profile. This is carried out through perturbative expansions of various types of spectral laws of the considered perturbed large matrices. First, we demonstrate different perturbative expansions of the empirical spectral measure in the cases of the perturbative regime and the semi-perturbative regime and highlight well known heuristic patterns in Physics, as the transition between semi-perturbative and perturbative regimes. Secondly, we provide a thorough study of the semi-perturbative regime and prove the new fact that this regime could be decomposed into infinitely many sub-regimes. Finally, we prove, through a perturbative expansion of spectral measures associated to the state defined by a given vector, a perturbative expansion of the coordinates of the eigenvectors of the perturbed matrices.

---

## RÉSUMÉ

---

La présente thèse est consacrée à l'étude de l'effet d'une perturbation sur le spectre d'une matrice hermitienne perturbée par une matrice aléatoire de petite norme opérateur et dont les entrées dans la base propre de la première matrice sont indépendantes, centrées et possèdent un profil de variance. Ceci est réalisé au travers de développements perturbatifs de divers types des lois spectrales des grandes matrices perturbées considérées. Dans un premier temps, nous démontrons différents développements perturbatifs de la mesure spectrale empirique dans les cas du régime perturbatif et du régime semi-perturbatif et mettons en évidence des modèles heuristiques bien connus en physique, comme la transition entre les régimes semi-perturbatifs et perturbatifs. Dans un deuxième temps, nous proposons une étude approfondie du régime semi-perturbatif et prouvons le fait nouveau que ce régime peut être décomposé en un nombre infini de sous-régimes. Enfin, nous démontrons, au travers d'un développement perturbatif des mesures spectrales associées à un vecteur donné, un développement perturbatif des coordonnées des vecteurs propres des matrices perturbées que nous considérons.



---

---

# Remerciements

Ma gratitude s'adresse en tout premier lieu à mon directeur de thèse, Florent Benaych-Georges, pour m'avoir encadré tout au long de ces dernières années et m'avoir ainsi permis de réaliser l'un de mes vœux les plus chers : m'initier à la recherche en mathématiques. La générosité avec laquelle il m'a partagé son savoir, son intégrité et sa vision des mathématiques ont été une réelle source d'inspiration.

Je tiens également à remercier Nathanaël Enriquez avec qui j'ai eu le plaisir de travailler durant de nombreuses journées. Je lui suis reconnaissant pour l'excellent exemple qu'il a su me prodiguer en tant que mathématicien mais aussi, plus généralement, en tant que philosophe.

J'adresse aussi mes plus sincères remerciements à Jamal Najim, Djalil Chafaï et Sean O'Rourke qui ont accepté d'effectuer le long travail qui consiste à rapporter une thèse et me font l'honneur de constituer aujourd'hui, avec Florent Benaych-Georges et Nathanaël Enriquez, mon jury de thèse.

Je garderai un excellent souvenir de l'accueil auquel j'ai eu droit au sein du laboratoire MAP5 au cours de ces années d'études. A ce titre, je voudrais exprimer ma profonde reconnaissance à Annie Raoult et Fabienne Comte, directrices successives du laboratoire, ainsi qu'à Christine Graffigne, directrice de l'UFR de mathématiques de l'Université Paris Descartes, pour l'aide et le soin particulier qu'elles portent aux doctorants. Durant ce doctorat j'ai eu le plaisir de m'enrichir aussi bien sur le plan mathématique que culturel aux côtés de nombreux chercheurs actuels et anciens du MAP5, je leur adresse à tous mes remerciements et plus particulièrement à Flora Alarcon, Andrés Almansa, Avner Bar-Hen, Etienne Birmelé, Olivier Bouaziz, Charles Bouveyron, Maya de Buhan, Thierry Cabanal-Duvillard, Antoine Chambaz, Manon Defosseux, Julie Delon, Sylvain Durand, Céline Duval, Nathalie Eisenbaum, Anne Estrade, Jean-Claude Fort, Bruno Galerne, Valentine Genon-Catalot, Joan Glaunès, Bérénice Grec, Catherine Huber, Georges Koepfler, Raphaël Lachièze-Rey, Rachid Lounes, Éric Luçon, Sébastien Martin, Nicolas Meunier, Lionel Moisan, Vittorio Perduca, Christophe Pouzat, Edoardo Provenzi, Ellen Saada, Camille Sutour et Bernard Ycart. Je remercie également Christophe Castellani, Marie-Hélène Gbaguidi, Augustin Hangat, Valérie Hoarau Carpentier, Sandrine Fallu, Azedine Mani, Arnaud Meunier, Maureen Némorin, Max Paisley, Thierry Raedersdorff et Isabelle Valero pour leur gentillesse et leur aide informatique et administrative.

---

Je souhaite exprimer de surcroît toute ma reconnaissance aux doctorants et post-doctorants anciens et actuels du MAP5 pour leur amitié et l'émulation intellectuelle dont j'ai bénéficié à leur contact. Je remercie ainsi Rémy Abergel, Morgan André, Andréa Bondesan, Thomas Browne, Khaled Chacouche, Ismaël Chouraqi, Ronan Costaouec, Alessandro Cucchi, Valentin de Bortoli, Christophe Denis, Anne-Claire Egloff, Christelle Etchegaray, Noura Faraj, Julie Fournier, Mélina Gallopin, Kévin Kuoch, Vivien Goepp, Ronan Herry, Antoine Houdart, Allan Jérôlon, Maud Kerebel, Claire Launay, Arthur Leclaire, Arthur Leroy, Gwennaëlle Mabon, Anne-Sophie Macé, Pierre-Alexandre Mattei, Juliana Pegoraro, Léo Planche, Thomas Picchetti, Angelina Roche, Mariano Rodriguez, Alasdair Newson, Samuel Ronsin, Maurizia Rossi, Pierre Roussillon, Alexandre Saint-Dizier, Matias Tassano, Vincent Vidal et Fabien Vergnet et exprime ma reconnaissance particulière à Charlotte Dion, Warith Harchaoui et Jean Rochet pour l'aide précieuse qu'ils ont su m'apporter si amicalement au cours de ces dernières années.

Tout au long de cette thèse de doctorat j'ai eu le plaisir de m'instruire en assistant aux exposés mensuels du groupement de recherche dénommé *Matrices et graphes aléatoires* organisé notamment par Florent Benaych-Georges, Mireille Capitaine, Mylène Maïda et Jamal Najim à l'Institut Henri Poincaré. Pour cette raison je remercie ces derniers ainsi que Camille Male, qui a initié ce projet, et les chercheurs avec qui j'ai beaucoup partagé au cours de ces réunions ; Fanny Augeri, Marwa Banna, Raphaël Butez, Sandrine Dallaporta, Laure Dumaz, Maxime Février, Benjamin Groux, Adrien Hardy, Thomas Leblé, Nathan Noiry, Laurent Ménard, Sandrine Péché et Pierre Youssef.

L'accès à une thèse de doctorat ne peut s'effectuer qu'en conséquence d'une formation solide et à ce titre je souhaite exprimer ma plus profonde gratitude à l'ensemble de mes professeurs qui ont su me faire évoluer tant sur le plan scolaire que sur le plan humain et plus particulièrement à Olivier Beuvelet, Laurent Delobel, Jacques Féjoz, Olivier Gingolani, Alain Grumbach, Irina Kourkova, Janine Laroche, Nicolas Lerner, Bernard Ott, Cyrille Rimboeuf, Catherine Roche et Martine Querette pour le parcours d'excellence sur lequel ils m'ont sagement guidé.

J'adresse ma vive affection à mes amis qui partagent mon quotidien et me donnent le bonheur qui me permet d'entreprendre et d'achever gaiement des projets tels que la présente thèse. Je remercie notamment Bernard Barral, Marie-Christine Boulanger, Mikael Beatriz, Antoine Brault, Richard de Cabrol de Moute, Michel Cherix, Julie Coniglio, Louise Coniglio, Thomas Crassous, Jacques de Fontenailles, Katy Frade, Jefferson Gandelin-Paris, Olivier Guitton, Pierre Holleaux, Gaël Lagarde, Alexandre Lejoncour, Hadrien de March, Arnaud Meurin, François Petit, Emmanuelle Robert et Karine Scellier pour l'influence positive qu'ils ont eu sur ma personne et conséquemment mon travail de recherche.

Je tiens à exprimer ma très grande reconnaissance à Nadia Coniglio et Bruno Coniglio, ainsi qu'au reste de ma belle-famille, pour leur soutien précieux tout au long de mon parcours étudiant et leur constante bienveillance. Je remercie chaleureusement mon amie et compagne de vie, Pauline Coniglio, qui par sa grandeur d'âme et sa munificente bienfaisance contribue à mon bonheur quotidien.

---

Bien que mes remerciements seront toujours en deçà de la dette véritablement due, je souhaite remercier mon père, Vassilios Michaïl, ainsi que, à titre posthume, ma mère, Artémis Michaïl, pour s'être employés avec grande industrie à m'insuffler les aspirations et les idéaux en l'absence desquels il n'est de sens.

Enfin, mes dernières pensées vont à mon pays, la France, qui de ma plus tendre jeunesse à aujourd'hui m'a fait bénéficier de tous les soins dont un citoyen pourrait rêver. Cette thèse, entièrement financée par le ministère de la recherche, n'en est que l'une des conséquences et lui en dédier le résultat ne saurait épuiser ma dette.

---

# Contents

<b>Abstract</b>	<b>3</b>
<b>Remerciements</b>	<b>5</b>
<b>1 Introduction</b>	<b>13</b>
1.1 Universality: from square matrices to circular distributions . . . . .	15
1.1.1 Wigner's semi-circle law . . . . .	15
1.1.2 Girko's circle law . . . . .	17
1.1.3 Marchenko-Pastur's quarter circle law . . . . .	18
1.1.4 Single Ring Theorem . . . . .	20
1.2 Eigenvectors in random matrix theory . . . . .	22
1.2.1 Spectral measure . . . . .	22
1.2.2 Anderson's transition . . . . .	23
1.3 Perturbation theory . . . . .	24
1.3.1 The problem of Weyl . . . . .	24
1.3.2 Eigenvalues of a perturbed matrix: bulk and outliers . . . . .	25
1.3.3 Eigenvectors of a perturbed matrix . . . . .	27
1.4 Purpose of this thesis . . . . .	29
1.4.1 Organization of the thesis . . . . .	29
<b>2 Useful tools and theorems</b>	<b>31</b>
2.1 Stieltjes transform . . . . .	31
2.2 Hellfer-Sjöstrand formula . . . . .	32
2.3 CLT extension lemma . . . . .	33

2.4	A functional density lemma . . . . .	34
<b>3</b>	<b>Empirical spectral distribution of a matrix under perturbation</b>	<b>37</b>
3.1	Introduction . . . . .	38
3.2	Main result . . . . .	40
3.2.1	Definition of the model and assumptions . . . . .	40
3.2.2	Main result . . . . .	42
3.3	Examples . . . . .	43
3.3.1	Uniform measure perturbation by a band matrix . . . . .	43
3.3.2	Triangular pulse perturbation by a Wigner matrix . . . . .	44
3.3.3	Parabolic pulse perturbation by a Wigner matrix . . . . .	44
3.4	Relation to free probability theory . . . . .	46
3.5	Strategy of the proof . . . . .	48
3.6	Stieltjes transforms convergence . . . . .	48
3.6.1	Two statements . . . . .	49
3.6.2	Proof of Proposition 21 . . . . .	50
3.6.3	Proof of Proposition 22 . . . . .	55
3.6.4	A local type convergence result . . . . .	57
3.6.5	Possible extensions to larger $\varepsilon_n$ . . . . .	58
3.7	Convergence in probability in the semi-perturbative regime . . . . .	59
3.8	Convergence in distribution towards the Gaussian variable $Z_\phi$ . . . . .	61
3.8.1	Coincidence of the expectation of $\mu_n^\varepsilon$ with its deterministic approximation . . . . .	61
3.8.2	Domination of the variance of $\mu_n^\varepsilon$ . . . . .	62
3.8.3	Proof of the convergences in distribution of Theorem 19 . . . . .	63
<b>4</b>	<b>Expansions of the empirical spectral measure of a perturbed matrix</b>	<b>65</b>
4.1	Introduction . . . . .	66
4.2	Main results . . . . .	67
4.2.1	Definition of the model and assumptions . . . . .	67
4.2.2	Convergence in distribution of $\mu_n^\varepsilon$ . . . . .	69
4.2.3	A local type convergence result . . . . .	70

---

4.3	Examples and simulations . . . . .	71
4.3.1	Explanation of the first three deterministic terms of the expansion of $\mu_n^\varepsilon$ . . . . .	71
4.3.2	Numerical simulations . . . . .	73
	Parabolic pulse perturbation by a Wigner matrix . . . . .	73
	Uniform measure perturbation by different band matrices . . . . .	74
4.4	Strategy of the proof . . . . .	74
4.5	Proofs . . . . .	75
4.5.1	Statements . . . . .	75
4.5.2	How the deterministic terms $C_k(z)$ appear . . . . .	76
4.5.3	Asymptotic behavior of the terms resulting from the expansion of the resolvent of $D_n^\varepsilon$ . . . . .	78
4.5.4	Proof of Theorem 28 . . . . .	83
4.5.5	Proof of the local type convergence result . . . . .	84
4.6	A possible extension to other classes of test functions . . . . .	85
<b>5</b>	<b>Eigenvectors of a matrix under random perturbation</b>	<b>87</b>
5.1	Introduction . . . . .	87
5.2	Main result . . . . .	89
5.3	Proof (of Theorem 32) . . . . .	91
5.3.1	Stieltjes transform . . . . .	92
5.3.2	From Stieltjes transform to $\mathcal{C}^5$ functions . . . . .	95
5.4	Consequence for the eigenvectors . . . . .	96
	<b>List of Figures</b>	<b>98</b>
	<b>Bibliography</b>	<b>99</b>

---



## CONTENTS

---

# Chapter 1

## Introduction

### Contents

---

<b>1.1</b>	<b>Universality: from square matrices to circular distributions .</b>	<b>15</b>
1.1.1	Wigner’s semi-circle law . . . . .	15
1.1.2	Girko’s circle law . . . . .	17
1.1.3	Marchenko-Pastur’s quarter circle law . . . . .	18
1.1.4	Single Ring Theorem . . . . .	20
<b>1.2</b>	<b>Eigenvectors in random matrix theory . . . . .</b>	<b>22</b>
1.2.1	Spectral measure . . . . .	22
1.2.2	Anderson’s transition . . . . .	23
<b>1.3</b>	<b>Perturbation theory . . . . .</b>	<b>24</b>
1.3.1	The problem of Weyl . . . . .	24
1.3.2	Eigenvalues of a perturbed matrix: bulk and outliers . . . . .	25
1.3.3	Eigenvectors of a perturbed matrix . . . . .	27
<b>1.4</b>	<b>Purpose of this thesis . . . . .</b>	<b>29</b>
1.4.1	Organization of the thesis . . . . .	29

---

Numerous mathematical problems naturally lead us to fill a matrix with random coefficients and to study its spectral properties. Thus, in the 1920s, the statistician Wishart initiated the random matrix theory by studying in [Wis28] the first ever model of a random matrix: a random covariance matrix. Given  $n$  centered i.i.d. random vectors,  $X_1, \dots, X_n$  in  $\mathbb{R}^p$  modeling a sample of a given population, assume that we want to estimate the covariance matrix  $\Sigma$  of their common distribution. We know from the law of large numbers that the random matrix  $S_n = \frac{1}{n}(X_1, \dots, X_n)(X_1, \dots, X_n)^T$  is a good estimator of  $\Sigma$ , as long as  $p$  remains small compared to  $n$ . But one of the reasons we may be interested in the

spectrum of these matrices,  $S_n$ , now identified as *Wishart matrices*, is that in the case where  $p$  has an order close to that of  $n$ , to obtain satisfying information about  $\Sigma$ , we need to use other statistical methods, such as a principal component analysis, which involve the largest eigenvalues of  $S_n$  and their associated eigenvectors.

However, it was not until the 1950s that a concrete mathematical theory of the spectrum of random matrices emerged through the work [Wig55] of Nobel Prize winner Eugene Wigner. In the context of his physical studies of heavy nuclei, the behavior of a nucleus is determined by a Hamiltonian operator. Wigner's idea was then to approximate this Hamiltonian operator by a Hermitian random matrix of high dimension. The interest for the spectrum of these matrices followed the postulate that the spacings between the lines in the spectrum of a heavy atom nucleus should resemble the spacings between their eigenvalues. Indeed, during this last study Wigner then conjectured a phenomenon of universality, namely that the asymptotic spectral properties of random matrices are described by universal laws.

This latter conjecture, which will be illustrated in Section 1.1, as well as the work of Dyson and Mehta, marked the beginning of the interest in random matrices as pure mathematical objects. Indeed, many important tools aiding the analysis of the properties of the spectrum of random matrices were then developed, for example, the method of orthogonal polynomials by Mehta and Gaudin in [MG60]. In addition, two years later in [Dys62], Dyson classified Hamiltonians into several categories based on their symmetry properties, implying the existence of three major sets of matrices: the Gaussian orthogonal ensemble (GOE), the Gaussian unitary ensemble (GUE) and the Gaussian symplectic ensemble (GSE). Those three ensembles of Gaussian random matrices, which are respectively real symmetric, complex Hermitian, and quaternionic self-dual, rapidly became fundamental in various areas of theoretical physics, mesoscopic physics, disordered electron systems, and in the field of quantum chaos.

Until the late 1980s, research on the properties of random matrices in the mathematical community was apparently not as intensive as it was in theoretical physics. Among the few purely mathematical articles of the two decades after the 1960s, we can cite in particular three works that greatly influenced the theory of random matrices: the work [MP67] of Marcenko and Pastur describing the spectrum of random covariance matrices, the work [FK81] of Furedi and Komlos on the distribution of eigenvalues in ensembles of random matrices with independent entries and finally Montgomery's work [Mon73], which provided an important link between the theory of random matrices and deep-rooted problems of number theory. Indeed, he revealed a certain similarity between the eigenvalues of matrices of the GUE and the zeros of the Riemann zeta function.

The situation changed considerably after the 1990s, thanks, in particular, to the works [Voi91, DVDN92] of Voiculescu *et al*, who developed the free probability theory. Since then, random matrices have been increasingly studied. Today, random matrix theory is a very large subject with applications in many disciplines of natural science, engineering and finance.

In the following sections of this introduction, we present some fundamental results of random matrix theory that will help us progressively introduce the main ideas and tools that we will use in this thesis, as well as presenting the historical context in which this thesis is ingrained. Finally, the last subsection presents the precise subject of this thesis.

## 1.1 Universality: from square matrices to circular distributions

The first, and perhaps the most fundamental, mathematical object used to study the eigenvalues of a random matrix was introduced by Eugen Wigner and corresponds to the following probability measure, commonly known as *empirical spectral measure*.

**Definition 1** (Empirical spectral measure). For any  $n \times n$  square matrix  $A$ , the probability measure,  $\mu_n$ , which puts equal mass on each eigenvalue of  $A$  is called the empirical spectral measure (or distribution) of  $A$ .

For example, in the specific case where we consider an  $n \times n$  Hermitian (or real symmetric) matrix  $A$ , as the eigenvalues  $\lambda_1, \dots, \lambda_n$  of  $A$  are all real, its empirical spectral measure is a real probability measure that could be expressed as

$$\mu_n = \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_i}.$$

Moreover, one can note that in this last case for any Borel set  $B \subseteq \mathbb{R}$ ,

$$\mu_n(B) = \frac{1}{n} \#\{\lambda_i \in B \mid i \in \llbracket 1, n \rrbracket\}$$

is the proportion of eigenvalues of  $A$  contained in the set  $B$ . For this reason, many studies have investigated the empirical spectral measure of various models of random matrices in order to study the distribution of their eigenvalues. As we will see in the next fundamental examples presented in this introductory chapter, many studies have focused on the limiting spectrum of an  $n \times n$  matrix  $A_n$  when its dimension,  $n$ , tends to infinity.

### 1.1.1 Wigner's semi-circle law

The matrix model initially introduced by Eugen Wigner in 1955 was a specific model of a random Hermitian matrix. He introduced it to study the eigenvalues of such a matrix when its dimension tends to infinity in order to analyze by approximation a self-adjoint operator that interested him in his work in the field of physics of heavy nuclei. This model is given as follows.

**Definition 2** (Wigner matrix). A Wigner matrix is a random Hermitian matrix,  $W_n = (W_n(i, j))_{i,j=1}^n$  such that the entries above its diagonal  $\{W_n(i, j), 1 \leq i < j \leq n\}$  are real or complex valued i.i.d. random variables with zero mean and unit variance, and the diagonal entries  $\{W_n(i, i), 1 \leq i \leq n\}$  are i.i.d. centered real-valued random variables with bounded mean and variance and are independent from the upper-triangular entries.

**Remark 1.** This definition may vary; rather than considering that the upper-triangular entries have a second moment equal to 1, other authors consider that the entries have just a bounded moment of order 2, or bounded moments of any order.

By studying the eigenvalues of this last model, Wigner demonstrated the following theorem which can be considered as the starting point of random matrix theory.

**Theorem 3** (Wigner's semi-circle law). *The empirical spectral measure,  $\mu_{\frac{1}{\sqrt{n}}W_n}$ , of an  $n \times n$  normalized Wigner matrix,  $\frac{1}{\sqrt{n}}W_n$ , converges almost surely as  $n$  tends to infinity to the Wigner semi-circle distribution with parameter 2:*

$$\mu_{\frac{1}{\sqrt{n}}W_n} \xrightarrow[n \rightarrow \infty]{a.s.} \mu_{sc}$$

with

$$\mu_{sc} := \frac{1}{2\pi} \sqrt{4 - x^2} \mathbb{1}_{x \in [-2, 2]} dx.$$

In other words, the eigenvalues of a normalized large Wigner matrix will be approximately distributed in a semi-circle of radius 2. This phenomenon is illustrated in Figure 1.1, where we can observe the convergence to the semi-circle distribution of the eigenvalues as the dimension,  $n$ , grows.

The reason we have to normalize the matrix  $W_n$  by  $\frac{1}{\sqrt{n}}$  comes from the fact that we want its spectrum to be bounded. Indeed, the magnitude of the eigenvalues of  $W_n$  is of order  $O(\sqrt{n})$ . Various proofs of this order of magnitude can be found in the literature. We can cite for example the so-called *Bai-Yin Theorem* which was proved in [BY86] and states that, in the case where the entries of  $W_n$  have finite fourth moment, almost surely:

$$\lim_{n \rightarrow \infty} \frac{\|W_n\|_{\text{op}}}{\sqrt{n}} = 2.$$

Another noteworthy fact of Wigner's theorem is that since the eigenvalues of a random matrix are random variables, the involved empirical spectral measure is a random measure. It is therefore a convergence theorem of a random probability measure to a deterministic probability measure.

Finally, as for the classical central limit theorem, this last result shows a limiting distribution which does not depend on the initial laws of the entries of the considered matrix. In random matrix theory this phenomenon is called *universality*. In the next subsections, we present three other historically important results of universality for different classes of random matrices.




Figure 1.1: **Illustration of the Wigner's semi-circle law.** The red curve represents the density  $x \mapsto \frac{1}{2\pi} \sqrt{4 - x^2} \mathbb{1}_{x \in [-2, 2]}$  of the semi-circle distribution with parameter 2 and the blue histogram is that of the eigenvalues of an  $n \times n$  Gaussian Wigner Matrix  $\frac{1}{\sqrt{n}} W_n$  for different values of  $n$ .

### 1.1.2 Girko's circle law

In 1965 the French mathematical physicist, Jean Ginibre, noticed and demonstrated in [Gin65] that the limiting spectral distribution of an  $n \times n$  random matrix which lacks any condition of symmetry and whose entries are all i.i.d. centered Gaussian random variables, all with variance equal to  $1/n$ , is the uniform distribution over the unit disc.

Since this study, the ensemble of square matrices with Gaussian distribution of entries is called the *Ginibre ensemble* and has been very successful in describing various physical phenomena.

Despite the fact that, in 1984, Vyacheslav Girko introduced in [Gir85], an approach that allowed for the establishment of the circular law for more general distributions, it was not until 45 years after the initial result of Ginibre that, in 2010, Tao and Vu proved in [TV10] the universality of the so-called *Girko's circle law* under the minimum assumptions set out below.

**Theorem 4** (Girko's circle law). *Let  $(G_n)_{n \in \mathbb{N}^*}$  be a sequence of  $n \times n$  matrix ensembles whose entries are i.i.d. copies of a complex random variable with mean 0 and variance 1. Then, the empirical spectral measure of  $\frac{1}{\sqrt{n}}G_n$  converges almost surely to the uniform measure on the unit disk as  $n$  tends to infinity:*

$$\mu_{\frac{1}{\sqrt{n}}G_n} \xrightarrow[n \rightarrow \infty]{a.s.} \mu_{disk}$$

with

$$\mu_{disk} := \frac{1}{\pi} \mathbb{1}_{\{x^2+y^2 \leq 1\}}(x, y) \, dx dy$$

The eigenvalues, which are in this case complex random variables, will be approximately distributed in a disk of the complex plane of radius 1. This phenomenon is illustrated in Figure 1.2, where we can observe the convergence to the uniform distribution on the unit disk of the eigenvalues as the dimension  $n$  grows.




Figure 1.2: **Illustration of the Girko's circle law.** The blue dots represent the eigenvalues in the complex plane of a real Ginibre matrix  $\frac{1}{\sqrt{n}}G_n$  for different values of  $n$ .

### 1.1.3 Marchenko-Pastur's quarter circle law

In 1967, two years after the first result of Jean Ginibre on the eigenvalues of Gaussian matrices, the two Ukrainian mathematicians Vladimir Marchenko and Leonid Pastur proved in [MP67] that the limiting distribution of the singular values of a large rectangular random matrix is a probability distribution, hence named the *Marchenko-Pastur distribution*.

We recall that the singular values of a matrix  $M$  are the eigenvalues of the matrix  $\sqrt{MM^*}$ . Thus, in the case where  $M$  is a rectangular matrix and therefore does not have eigenvalues, its singular values are helpful to compute its pseudoinverse, to determine its rank, its range, its operator norm, etc.

We present here a simpler, reformulated version, of the initial Marchenko-Pastur theorem which illustrates the convergence of the empirical spectral measure to the quarter circle distribution.

**Theorem 5** (Marchenko-Pastur's quarter circle law). *If  $X_n$  is an  $n \times n$  random matrix such that its entries are independent, centered and have variance 1, then the empirical spectral distribution,  $\mu_{\frac{1}{\sqrt{n}}\sqrt{MM^*}}$ , of the  $n \times n$  random matrix  $\frac{1}{\sqrt{n}}\sqrt{MM^*}$  converges to the Marchenko-Pastur's quarter circle distribution:*

$$\mu_{\frac{1}{\sqrt{n}}\sqrt{MM^*}} \xrightarrow[n \rightarrow \infty]{a.s.} \mu_{MP}$$

with

$$\mu_{MP} := \frac{1}{\pi} \sqrt{4 - x^2} \mathbb{1}_{[0,2]}(x) \, dx$$




Figure 1.3: **Illustration of the Marchenko-Pastur's quarter circle law.** The red curve represents the density  $x \mapsto \frac{1}{\pi} \sqrt{4 - x^2} \mathbb{1}_{[0,2]}(x)$  of the quarter circle distribution and the blue histogram is that of the singular values of an  $n \times n$  random matrix such that its entries are Gaussian random variables which are independent, centered and with unit variance, for different values of  $n$ .



### 1.1.4 Single Ring Theorem

A more recent result of universality that we can cite in this introduction is that of Guionnet, Krishnapur and Zeitouni. In 2011, they demonstrated, in an article entitled *Single Ring Theorem*, a new universality result for another class of matrices. For a fixed  $n$ -tuple  $(s_1, \dots, s_n)$  of non-negative reals, these matrices could be defined as the matrices which are uniformly distributed among the matrices with singular values  $(s_1, \dots, s_n)$ . More explicitly, any matrix of this class can be written under the form,

$$A_n = U_n D_n V_n$$

for  $U_n$  and  $V_n$  two independent Haar distributed random matrices, that is to say two matrices uniformly distributed among the ensemble of unitary matrices, and  $D_n = \text{diag}(s_1, \dots, s_n)$  a non-negative diagonal matrix independent of  $U_n$  and  $V_n$ . This result [GKZ11], was initially partially demonstrated by the physicists Feinberg and Zee in [FZ97], and its assumptions have since been amended. Specifically, in [RV14], Rudelson and Vershynin showed that a hard-to-check assumption on the smallest singular value of  $(A_n - z.I_n)$  was unnecessary and in [BD<sup>+</sup>13], Basak and Dembo weakened the hypotheses so that the empirical spectral measure of  $D_n$  allows for the existence of some atoms. We state here a shortened version with optimal assumptions of the *Single Ring Theorem* borrowed from the very recent work [BES16] of Bao, Erdős and Schnelli which applies, like the first paper [BG<sup>+</sup>17] of Benaych-Georges, to study the *Local Single Ring Theorem*, that is to say the *Single Ring Theorem* within the bulk regime.

**Theorem 6** (Single Ring Theorem). *Let  $(A_n)_{n \in \mathbb{N}^*}$  be a sequence of  $n \times n$  matrices defined as previously. If the empirical spectral measure,  $\mu_{D_n}$ , of  $D_n$  converges weakly to a measure  $\mu$  whose support contains more than one point and the sequence of matrices  $(D_n)_{n \in \mathbb{N}^*}$  is uniformly bounded, then the empirical spectral measure of  $A_n$  converges in probability to a deterministic probability measure  $\nu$  which possesses a radially-symmetric density,  $\rho(z)$ , with respect to the Lebesgue measure on  $\mathbb{C}$  depending only on  $\mu$  and supported on the single ring  $\{z \in \mathbb{C}, r \leq |z| \leq R\}$  for*

$$r = \frac{1}{\sqrt{\int x^{-2} \mu(dx)}} \quad \text{and} \quad R = \sqrt{\int x^2 \mu(dx)}$$

To put it another way, the distinctive property of matrices  $(A_n)$  is that their eigenvalues tend to spread over a single annulus centered in the origin. For example, if the eigenvalues of the matrix  $D_n$  are distributed such that the empirical spectral measure of  $D_n$  tends to the uniform measure on an interval  $[a, b]$ , then the eigenvalues of  $A_n$  will spread as  $n$  tends to infinity over the single ring  $\{z \in \mathbb{C}, r = \sqrt{ab} \leq |z| \leq R = \sqrt{\frac{a^2 + ab + b^2}{3}}\}$ . We illustrate this phenomenon in Figure 1.4 in the case where  $\mu$  is the uniform measure on the interval  $[1, 6]$ .




Figure 1.4: **Illustration of the Single Ring Theorem.** The blue dots represent the eigenvalues in the complex plane of the matrix  $A_n = U_n D_n V_n$ , for different values of  $n$  and for  $U_n$  and  $V_n$  two  $n \times n$  Haar distributed random matrices and  $D_n$  an  $n \times n$  random diagonal matrix such that its empirical spectral measure converges to the uniform measure on  $[1, 6]$ . The smaller circle has radius  $r = \sqrt{6} \approx 2.45$  and the bigger one has radius  $R = \sqrt{\frac{43}{3}} \approx 3.78$ .

Furthermore, one can note that if the measure  $\mu$  verifies  $\sqrt{\int x^{-2} \mu(dx)} = \infty$ , then  $r = 0$  and the support of the limiting measure  $\nu$  is a disk, leading to the previously presented Ginibre case. Moreover, in the case where  $\mu$  is a Dirac measure,  $r = R$  so that the support of  $\nu$  is simply a circle. These relations are illustrated in Figure 1.5.




Figure 1.5: **Relation of the Single Ring Theorem to other results.** In the three figures above, the blue dots represent the eigenvalues in the complex plane respectively from left to right of a real Ginibre matrix  $\frac{1}{\sqrt{n}} G_n$ , of a matrix  $A_n = U_n D_n V_n$  such that  $\mu$  is the uniform measure on  $[0, \sqrt{3}]$  and of a matrix  $A_n = U_n D_n V_n$  such that  $\mu = \delta_1$ .

## 1.2 Eigenvectors in random matrix theory

Although the earliest foundational works of random matrix theory were concerned with eigenvalues, many studies were subsequently focused on eigenvectors. Indeed, many problems, particularly related to graph theory, naturally lead to questions about the eigenvectors of random matrices. We can mention for example problems related to spectral clustering or even the famous PageRank algorithm used by Google to sort its search results. Nevertheless, the eigenvectors literature is currently narrower than the eigenvalues one. Nevertheless, the literature concerning eigenvectors is currently scarce compared to that relating to eigenvalues.

This could be explained in particular by the fact that many ensembles of random matrices, such as those of the GOE or the GUE, are rotationally invariant in law which makes the study of their eigenvectors pointless as they are uniformly distributed on the unit sphere.

### 1.2.1 Spectral measure

In this section dedicated to eigenvectors, we introduce one of the tools that will be later addressed in this thesis: the spectral measure.

**Definition 7** (Spectral measure over a vector). For an  $n \times n$  hermitian matrix,  $H_n$ , with eigenvalues  $\lambda_1, \dots, \lambda_n$  and associated eigenvectors  $\mathbf{u}_1, \dots, \mathbf{u}_n$ , we define the spectral measure of  $H_n$  over a vector  $\mathbf{v}$ ,  $\mu_{H_n, \mathbf{v}}$ , by

$$\mu_{H, \mathbf{v}} = \sum_{j=1}^n |\mathbf{v} \cdot \mathbf{u}_j|^2 \delta_{\lambda_j}$$

In contrast with the empirical spectral measure, the spectral measure over a vector  $\mathbf{v}$  does not weigh all the eigenvalues of a matrix  $H_n$  equally; the weighting depends here on the eigenvectors of  $H_n$ . Thus, the spectral measure gives information on the eigenvector basis of  $H_n$ . Moreover, this measure is often applied over specific vectors as vectors  $(\mathbf{e}_i)_{i=1}^n$  of the canonical base of  $\mathbb{R}^n$ , as in this case it satisfies, for any function  $\phi$  defined on the spectrum of  $H_n$ , the identity

$$(\phi(H_n))_{i,i} = \sum_{j=1}^n |\mathbf{e}_i \cdot \mathbf{u}_j|^2 \phi(\lambda_j) = \int \phi(x) \, d\mu_{H_n, \mathbf{e}_i}$$

which follows directly from the spectral theorem.

For this reason as well as others, this measure is often used in studies on the eigenvectors of random matrices. For example, many estimates on eigenvectors of a random matrix,  $H_n$ , are obtained from its resolvent matrix,  $(H_n - zI_n)^{-1}$  and using the previous formula with  $\phi : x \mapsto \frac{1}{x-z}$  could be helpful in the sense that the entries of the resolvent matrix are easy to compute.

### 1.2.2 Anderson's transition

One of the sizeable current issues concerning the eigenvectors of random matrices is the so-called Anderson's transition. It concerns matrices called *band matrices*, that is to say Hermitian matrices whose entries are null beyond a band located around their diagonal, and which represent physical systems with local interactions. The question is then to know to which bandwidth,  $W$ , the spectral properties remain local, that is to say close to those of a diagonal matrix with independent diagonal entries. Roughly speaking, we say that an eigenvector is localized if it is essentially carried by few of its coordinates (see e.g. [Cha10]).

Although this topic is not addressed in this thesis, it may be interesting to give some fundamental results, because as we will later see we will approach it by studying the spectrum of a diagonal matrix summed with a random band matrix.

Despite the fact that this has not yet been formally demonstrated, a phase transition between the localized regime and the delocalized regime of the eigenvectors seems to occur when the bandwidth of an  $n \times n$  random matrix is of order  $\sqrt{n}$ . Another way of formulating this latter conjecture could be the following: the average localization length of the eigenvectors of an  $n \times n$  band matrix, of bandwidth  $W$ , is of the order of  $W^2$ . Figure 1.6 illustrates this transition for a  $4000 \times 4000$  band matrix, as  $\sqrt{4000} \approx 63$ .

In [Sch09], Jeffrey Schenker shows that the localization length of the eigenvectors of a band random matrix whose bandwidth is equal to  $W$ , is lower than  $W^8$ . Similarly, in 2012, Erdős, Yau and Yin demonstrated, in [EYY12a] that, under certain assumptions, the localization length of the eigenvectors of a band matrix is greater than  $W$ . We can also mention the work [EKYY13] of Erdős, Knowles, Yau and Yin in which they prove that the eigenvectors of an  $n \times n$  random band matrix are delocalized when the width of the strip  $W$  is of order greater than  $n^{4/5}$ . More generally, all the results currently in existence regarding this subject are in agreement with the conjecture previously mentioned.




Figure 1.6: **Illustration of Anderson's transition.** For an  $n \times n$  random band matrix whose entries are standard Gaussian variables, we have represented one of its eigenvectors for a bandwidth equal to  $W = 30$  (left),  $63$  (center) and  $70$  (right). Here  $n = 4000$ .

## 1.3 Perturbation theory

A natural and central question, in Mathematics and Physics, is to understand how the spectral properties of an operator are altered when the operator is subject to a small perturbation. This question is at the center of *Perturbation Theory* and was more precisely introduced by Rayleigh and Schrödinger as follows.

Let us consider a Hamiltonian operator,  $H$ , whose eigenvectors and eigenvalues are known and a second operator,  $X$ , whose entries are small compared to those of  $H$ . Thus, if we look at the operator  $H + X$ , the effect of  $X$  is to modify the spectrum of  $H$ . This can model, physically speaking, the perturbation by a weak magnetic or electric field represented by  $X$  of an atomic nucleus represented by  $H$ . The purpose of Perturbation Theory is then to give an approximate expression of the eigenvalues and eigenvectors of  $H + X$  in terms of those of  $H$ . We refer the reader to Kato's book [Kat13] for a thorough account on this subject which has been studied in many different contexts.

### 1.3.1 The problem of Weyl

In 1912, the German mathematician, theoretical physicist and philosopher Hermann Weyl stated in [Wey12] the following problem: if  $A$  and  $B$  are two Hermitian matrices, how are the eigenvalues of the matrix  $A + B$  related to those of  $A$  and  $B$  ?

This problem, which is more general than that of a perturbation, because here no hypothesis is made on the "smallness" of the perturbing matrix, currently does not have so many answers. For instance, by using the notation  $\lambda_1(M) \leq \dots \leq \lambda_n(M)$  to denote the eigenvalues of an  $n \times n$  Hermitian matrix  $M$ , it is easy to see, from linearity of the trace, that

$$\mathrm{Tr}(A + B) = \mathrm{Tr}(A) + \mathrm{Tr}(B) \iff \sum_{i=1}^n \lambda_i(A + B) = \sum_{i=1}^n \lambda_i(A) + \sum_{i=1}^n \lambda_i(B),$$

or that if  $A$  and  $B$  commute, so that  $A$  and  $B$  are diagonalizable in a same eigenbasis, then for any  $i \in \llbracket 1, n \rrbracket$ ,

$$\lambda_i(A + B) = \lambda_i(A) + \lambda_i(B),$$

but overall precise answers under broad hypotheses are scarce. Nevertheless, Weyl's answer to this problem has remained one of the most satisfactory thus far.

**Theorem 8** (Weyl's inequality). *Let  $A$  and  $B$  be two  $n \times n$  Hermitian matrices, then for any  $i \in \llbracket 1, n \rrbracket$ ,  $i \in \llbracket 0, n - i \rrbracket$ ,*

$$\lambda_i(A + B) \leq \lambda_{i+j}(A) + \lambda_{n-j}(B)$$

*Moreover, if  $A$  and  $B$  have no common eigenvector, then every inequality is strict.*

From this first theorem many other inequalities have been deduced, for instance,

**Corollary 9.** *Let  $A$  and  $B$  be two  $n \times n$  Hermitian matrices, then for any  $i \in \llbracket 1, n \rrbracket$ ,*

$$\lambda_i(A) + \lambda_1(B) \leq \lambda_i(A + B) \leq \lambda_i(A) + \lambda_n(B)$$

or even,

**Corollary 10.** *Let  $A$  and  $B$  be two  $n \times n$  Hermitian matrices, then for any  $(i, j) \in \llbracket 1, n \rrbracket^2$  such that  $i + j - 1 \leq n$ ,*

$$\lambda_{i+j-1}(A + B) \leq \lambda_i(A) + \lambda_j(B)$$

All these inequalities, known today as Weyl's inequalities, are very useful in estimating perturbations in the spectrum of Hermitian matrices. In fact, in the case of a perturbation,  $H + P$ , if we have a bound on the perturbing matrix,  $P$ , in the sense that we know that its operator norm satisfies  $\|P\|_{\text{op}} \leq \varepsilon$ , then it follows, by definition of the operator norm, that all the eigenvalues of  $P$  are bounded in absolute value by  $\varepsilon$ . Applying Weyl's inequalities, it follows that the spectra of  $H + P$  and  $H$  are close in the sense that for all  $i \in \llbracket 1, n \rrbracket$ ,

$$|\lambda_i(H + P) - \lambda_i(H)| \leq \varepsilon.$$

In other words, these inequalities guarantee that the spectrum of a Hermitian matrix is stable with respect to small perturbations, and are therefore a good starting point for any study of a perturbed system.

### 1.3.2 Eigenvalues of a perturbed matrix: bulk and outliers

Questions about perturbed matrices are as diverse (global, local, bulk, edge...) as questions about non perturbed matrices in random matrix theory. Namely, if we consider a deterministic  $n \times n$  Hermitian matrix,  $H_n$ , perturbed by a random matrix  $X_n$ , as various new asymptotic behaviors of the spectrum of  $H_n + X_n$  can appear, we can study, for instance, questions relating to the extreme eigenvalues of  $H_n + X_n$  or the typical spacings between two consecutive eigenvalues of  $H_n + X_n$ .

We can distinguish two categories of perturbations; those which are small in terms of rank and those which are small in terms of operator norm.

Those of the first category are often referred to as *low rank perturbations*. They consist in perturbing a matrix  $H_n$  by a matrix  $X_n$  whose rank is a fixed integer which does not vary with  $n$  or by a matrix  $X_n$  whose rank is small compared to its dimension (as for example  $\log(n)$ ). Weyl's inequalities guarantee that such perturbations do not influence the global statistics of the eigenvalues as the dimension  $n$  tends to infinity. Thus, the empirical spectral measures of  $H_n + X_n$  and  $H_n$  have the same asymptotics and are governed by the universal law associated to the matrix ensemble to which  $H_n$  belongs. However, at a




Figure 1.7: **Illustration of a finite rank perturbation.** In both figures, the red curve represents the density  $x \mapsto \frac{1}{2\pi}\sqrt{4-x^2} \mathbb{1}_{[-2,2]}(x)$  of the semi circle distribution. The blue histogram of the left figure is that of the eigenvalues of an  $n \times n$  Wigner matrix,  $\frac{1}{\sqrt{n}}W_n$ , whose entries are standard Gaussian variables. The blue histogram of the right figure is that of the eigenvalues of the same Wigner matrix but perturbed this time by a random diagonal matrix of rank 3, two outliers represented here by red dots are exhibited out of the bulk spectrum.

microscopic scale, the behavior of individual eigenvalues may change dramatically under such a perturbation. Indeed, the perturbed matrix may exhibit *outliers*, that is to say eigenvalues detached from the bulk spectrum.

Regarding the outliers of a matrix perturbed by a low rank matrix, there is a phase transition called the *BBP phase transition*, from the names of the authors Baik, Ben Arous and P      who have highlighted it in [BAP<sup>+</sup>05] in the case of covariance matrices, after the initial work [Joh01] of Johnstone. The general principle of this transition is that if the amplitude of the perturbation remains below a certain threshold, the largest eigenvalues of the perturbed system do not move significantly, whereas beyond this threshold, they are at a macroscopic distance from the bulk spectrum. Studies on other ensembles of matrices, such as the Wigner ensemble, have shown similar phase transitions. We can cite for example the pioneering work [FK81] of F  redi and Koml  s about the outliers of perturbed Wigner matrices, and the works [BGN11] or [BGGM<sup>+</sup>11] of Benaych-Georges, Guionnet, Maida and Nadakuditi that have generalized this transition to larger ensembles.

In this thesis we are interested in perturbations belonging to the second category: those which are small in terms of operator norm. This second kind of perturbation has, unlike the first, a tendency to macroscopically perturb the bulk spectrum of the considered matrix, and also has a phase transition phenomenon.

More precisely, in the typical case where the spacing between two consecutive eigenvalues of the matrix,  $H_n$ , is of order  $n^{-1}$  the phase transition takes place at  $\|X_n\|_{\text{op}} \sim n^{-1}$ . The regime  $\|X_n\|_{\text{op}} \ll n^{-1}$  is called the *perturbative regime* and the regime  $n^{-1} \ll \|X_n\|_{\text{op}} \ll 1$  is often called the *semi-perturbative regime*, as the more the operator norm of a perturbation approaches the order 1, the more the considered regime begins to resemble that of a simple

matrix addition. We illustrate this phase transition in Figure 1.8, where the distinction between the spectra of  $H_n$  and  $H_n + X_n$  begins to be observable effectively as soon as the operator norm of the perturbative matrix,  $X_n$ , begins to be of order greater than  $n^{-1}$ .




Figure 1.8: **Illustration of the phase transition phenomenon.** The blue histogram is that of the eigenvalues of the diagonal matrix  $H_n = \text{diag}(\frac{1}{n}, \frac{2}{n}, \dots, \frac{n}{n})$  and the beige histogram is that of the eigenvalues of  $H_n + X_n$  where  $X_n$  is a Gaussian Wigner Matrix. Here  $\|X_n\|_{\text{op}} \sim n^{-1.2}$  (top left),  $n^{-1}$  (top right),  $n^{-0.5}$  (bottom left) and  $n^{-0.2}$  (bottom right).

In general, and as it is very well explained in [Fer00], the study of the perturbative regime mainly concerns applications in quantum mechanics, while the semi-perturbative regime deals with applications in the context of covariance matrices, as in [LP11], or applications to finance, as in [AB12].

### 1.3.3 Eigenvectors of a perturbed matrix

Currently, the existing results about the spectral properties of perturbed systems mostly concern the eigenvalues, and we have very little knowledge about eigenvectors. We can distinguish two major themes among the works focused on the study of the eigenvectors of a perturbed Hermitian matrix; those that provide bounds on the deviations of these eigenvectors under perturbation, as [OVW16, OVW17, vSW17, Zho17], and those, as [AB12, AB14, ABB14, Ben17], that provide explicit perturbative expansions.



For example, in [AB12], Allez and Bouchaud investigate the stability under a small perturbation of the subspace spanned by some consecutive eigenvectors of a generic symmetric matrix, and in [AB14] they consider the perturbation of a matricial stochastic process and they study the effect of this perturbation on the dynamics of a given eigenvector. In more recent work [Zho17], Zhong gives bounds of deviations of the eigenvector associated to the maximal eigenvalue of the perturbed matrix.

Let us suppose for exemple  $H_n$  to be diagonal. In this sense, it is possible to understand the current problem of perturbation as that of a perturbation of the canonical basis. Figure 1.9 illustrates this interpretation of the problem.




Figure 1.9: **Effect of a perturbation on an eigenvector.** For a diagonal matrix  $H_n = \text{diag}(\frac{1}{n}, \frac{2}{n}, \dots, \frac{n}{n})$  and a Gaussian Wigner matrix,  $X_n$ , which operator norm is of order  $n^{-0.4}$ , one can visualize in the left figure the  $\frac{n}{2}$ -th eigenvector of the original matrix,  $H_n$ , and in the right figure the  $\frac{n}{2}$ -th eigenvector of the perturbed matrix,  $H_n + X_n$ . Here  $n = 1000$ .

Major technical difficulties remain to approximate precisely the eigenvectors of a perturbed matrix. However, it is possible to find in the recent literature, some advances on this specific subject. We can cite for example the paper [Ben17] of Benigni, in which he shows that in the case of a perturbation of a diagonal matrix,  $H_n$ , by a Wigner matrix,  $X_n$ , with operator norm of order  $n^{-1/2} \ll \varepsilon \ll 1$ , we have for  $m$  specifically selected eigenvectors,  $(\mathbf{u}_{k_i})_{i=1}^m$ , of  $H_n + X_n$  and any unit vector  $\mathbf{q}$ , the convergence in moments of the quantities  $(\langle \mathbf{u}_{k_i}, \mathbf{q} \rangle)_{i=1}^m$  to those of a Gaussian vector if  $X_n$  is symmetric or to those of a complex Gaussian vector if  $X_n$  is Hermitian. By choosing, in particular, a unit vector  $\mathbf{q}$  of the canonical basis this result could then specify an average on some coordinates of an eigenvector of the perturbed matrix,  $H_n + X_n$ .

## 1.4 Purpose of this thesis

The present thesis is devoted to the study of the sensitivity of the eigenvalues and the eigenvectors of a given operator under small perturbations, and more precisely to the understanding of the effect of a perturbation on the spectrum of a Hermitian matrix by a random matrix with small operator norm and whose entries in the eigenvector basis of the first one were independent, centered and with a variance profile.

Throughout this thesis we will consider the following model:

We perturb an  $n \times n$  Hermitian matrix,  $D_n$ , by a random Hermitian matrix with small operator norm. Moreover, as we can diagonalize  $D_n$  up to a change of basis, we suppose it is diagonal and consider that the entries of the perturbative matrix,  $X_n$ , are independent in the eigenvector basis of  $D_n$  with a variance profile<sup>1</sup>. Furthermore, to better describe the magnitude of the perturbations, we suppose that the operator norm of  $X_n$  is of order 1, introduce a real sequence  $\varepsilon_n$  that tends to zero, and consider the perturbed matrix

$$D_n^\varepsilon := D_n + \varepsilon_n X_n$$

for different rates of convergence of  $\varepsilon_n$  to zero. In the following we prove that, depending on the order of magnitude of the perturbation, different regimes appear.

### 1.4.1 Organization of the thesis

This thesis is organized as follows:

- In Chapter 2, we introduce some useful tools and results which will be used throughout the thesis.
- In Chapter 3, we are interested in the empirical spectral distribution  $\mu_n^\varepsilon$  of the perturbed matrix,  $D_n^\varepsilon = D_n + \varepsilon_n X_n$ , in the regime where the matrix size  $n$  tends to infinity and  $\varepsilon_n$  tends to 0. We shall prove the following perturbative expansion of  $\mu_n^\varepsilon$ ,

$$\begin{aligned} \mu_n^\varepsilon &\approx \mu_n + \frac{\varepsilon_n}{n} dZ && \text{if } \varepsilon_n \ll n^{-1} \\ \mu_n^\varepsilon &\approx \mu_n + \frac{\varepsilon_n}{n} (cdF + dZ) && \text{if } \varepsilon_n \sim \frac{c}{n} \\ \mu_n^\varepsilon &\approx \mu_n + \varepsilon_n^2 dF && \text{if } n^{-1} \ll \varepsilon_n \ll 1 \end{aligned}$$

for  $\mu_n$  the empirical spectral measure of  $D_n$ ,  $dZ$  a Gaussian random linear form related to the so-called one-dimensional Gaussian free field, and  $dF$  a deterministic

---

<sup>1</sup>Note that if the perturbing matrix belongs to the GOE or GUE, then its law is invariant under this change of basis

linear form related to free probability theory. We will prove, moreover, that if  $n^{-1} \ll \varepsilon_n \ll n^{-1/3}$ , then the last of those three convergences can be refined as follows:

$$\mu_n^\varepsilon \approx \mu_n + \varepsilon_n^2 dF + \frac{\varepsilon_n}{n} dZ.$$

- The aim of Chapter 4 is to provide a thorough study of the semi-perturbative regime  $n^{-1} \ll \varepsilon_n \ll 1$ . We will see that this regime could be decomposed into infinitely many sub-regimes and that for all positive integer  $p$ , when  $n^{-1/(2p-1)} \ll \varepsilon_n \ll n^{-1/(2p+1)}$ , the perturbative expansion of  $\mu_n^\varepsilon$  could be decomposed in the following manner,

$$\mu_n^\varepsilon \approx \mu_n + \frac{\varepsilon_n}{n} dZ + \varepsilon_n^2 dC_2 + \varepsilon_n^4 dC_4 + \cdots + \varepsilon_n^{2p} dC_{2p}$$

for some deterministic linear forms ( $dC_{2k}$ ). In this chapter we will explain how we should reinforce the hypothesis of our model each time we choose a slower rate of convergence to zero for  $\varepsilon_n$  and how the deterministic terms appear then.

- Finally, in Chapter 5, we will provide a perturbative expansion of the coordinates of the eigenvectors of the perturbed matrix  $D_n^\varepsilon$ . This will be done through a perturbative expansion of spectral measures associated to the state defined by a given vector.

Chapters 3 and 5 correspond respectively to the publications [BGEM17] and [BGEM18] done in collaboration with Florent Benaych-Georges and Nathanaël Enriquez. Chapter 4 corresponds to work still undergoing study and unpublished as of yet. Some of the concepts mentioned in this introduction have not been removed from these chapters so that they can be read and understood independently of the rest of this thesis.

# Chapter 2

## Useful tools and theorems

### Contents

2.1	Stieltjes transform . . . . .	31
2.2	Hellfer-Sjöstrand formula . . . . .	32
2.3	CLT extension lemma . . . . .	33
2.4	A functional density lemma . . . . .	34

The tools and results we use along this thesis, namely the Stieltjes transform, the Hellfer-Sjöstrand formula, the CLT extension lemma of Shcherbina and Tirozzi and a functional density lemma with its proof, can be found by the reader in the following sections.

### 2.1 Stieltjes transform

In this section we present one of the fundamental tools of random matrix theory called Stieltjes transform and which is very useful for proving the convergence of a sequence of measures.

**Definition 11** (Stieltjes transform). Given a probability measure  $\mu$  on  $\mathbb{R}$  we define, for any  $z$  outside of its support, its Stieltjes transform by

$$s_\mu(z) := \int_{\mathbb{R}} \frac{1}{x - z} d\mu(x)$$

One of the main advantages of the Stieltjes transform is that, as it is invertible, we can use the following theorem to study the convergence of a sequence of measures.

**Theorem 12.** *If  $(\mu_n)_{n \in \mathbb{N}}$  is a sequence of measures on  $\mathbb{R}$  and  $\mu$  another measure on  $\mathbb{R}$ , then  $\mu_n$  converges almost surely to  $\mu$  as  $n$  tends to infinity if and only if  $s_{\mu_n}(z)$  converges almost surely as  $n$  tends to infinity to  $s_\mu(z)$  for any  $z \in \mathbb{C} \setminus \mathbb{R}$ .*

Moreover, in the special case of the study of the empirical spectral measure,  $\mu_n = \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_i}$  of a matrix  $M_n$ , this theorem allows us to resume our study to the convergence of the normalized trace of its resolvent matrix,

$$s_{\mu_n}(z) = \frac{1}{n} \sum_{i=1}^n \frac{1}{\lambda_i - z} = \frac{1}{n} \text{Tr} (M_n - zI_n)^{-1}.$$

Finally, we will use in the next three chapters the following identity,

**Lemma 13** (Resolvent expansion). *For two hermitian matrices  $A$  and  $B$ , and for any  $z$  that is not in the spectrum of  $A$ ,  $B$  and  $A + B$ , we have*

$$\frac{1}{z - (A + B)} = \frac{1}{z - A} + \frac{1}{z - A} B \frac{1}{z - (A + B)}.$$

## 2.2 Helffer-Sjöstrand formula

The Helffer-Sjöstrand formula expresses a regular function  $\phi$  on  $\mathbb{R}$  as an integral against functions  $x \mapsto \frac{1}{x-z}$  and is very useful to extend to any  $\mathcal{C}^p$  compactly supported function on  $\mathbb{R}$  a convergence which has been previously proven for test functions of the type  $x \mapsto \frac{1}{x-z}$ . In the next chapters, this formula will allow us to extend the convergence results initially demonstrated for the resolvent of the Hermitian matrix  $D_n^\varepsilon$  to its empirical spectral measure against regular functions.

**Proposition 14** (Helffer-Sjöstrand formula). *Let  $n \in \mathbb{N}$  and  $\phi \in \mathcal{C}^{p+1}(\mathbb{R})$ . We define the almost analytic extension of  $\phi$  of degree  $p$  through*

$$\tilde{\phi}_p(x + iy) := \sum_{k=0}^p \frac{1}{k!} (iy)^k \phi^{(k)}(x).$$

*Let  $\chi \in \mathcal{C}_c^\infty(\mathbb{C}; [0, 1])$  be a smooth cutoff function. Then for any  $\lambda \in \mathbb{R}$  satisfying  $\chi(\lambda) = 1$  we have*

$$\phi(\lambda) = \frac{1}{\pi} \int_{\mathbb{C}} \frac{\bar{\partial}(\tilde{\phi}_p(z) \chi(z))}{\lambda - z} d^2 z,$$

*where  $d^2 z$  denotes the Lebesgue measure on  $\mathbb{C}$  and  $\bar{\partial} := \frac{1}{2}(\partial_x + i\partial_y)$  is the antiholomorphic derivative.*

The proof of this formula can be found, e.g. in [BGK16].

## 2.3 CLT extension lemma

The following CLT extension lemma is borrowed from the paper of Shcherbina and Tirozzi [ST10]. We state here the version that can be found in the Appendix of [BGGM14].

**Lemma 15.** *Let  $(\mathcal{L}, \|\cdot\|)$  be a normed space with a dense subspace  $\mathcal{L}_1$  and, for each  $n \geq 1$ ,  $(N_n(\phi))_{\phi \in \mathcal{L}}$  a collection of real random variables such that:*

- *for each  $n$ ,  $\phi \mapsto N_n(\phi)$  is linear,*
- *for each  $n$  and each  $\phi \in \mathcal{L}$ ,  $\mathbb{E}[N_n(\phi)] = 0$ ,*
- *there is a constant  $C$  such that for each  $n$  and each  $\phi \in \mathcal{L}$ ,  $\text{Var}(N_n(\phi)) \leq C\|\phi\|^2$ ,*
- *there is a quadratic form  $V : \mathcal{L}_1 \rightarrow \mathbb{R}_+$  such that for any  $\phi \in \mathcal{L}_1$ , we have the convergence in distribution  $N_n(\phi) \xrightarrow[n \rightarrow \infty]{} \mathcal{N}(0, V(\phi))$ .*

*Then  $V$  is continuous on  $\mathcal{L}_1$ , can (uniquely) be continuously extended to  $\mathcal{L}$  and for any  $\phi \in \mathcal{L}$ , we have the convergence in distribution  $N_n(\phi) \xrightarrow[n \rightarrow \infty]{} \mathcal{N}(0, V(\phi))$ .*

One of the assumptions of previous the lemma concerns a variance domination. The next proposition provides a tool in order to check it. Let us first remind the definition of the Sobolev space  $\mathcal{H}_s$ .

For  $\phi \in L^1(\mathbb{R}, dx)$ , we define

$$\widehat{\phi}(k) := \int e^{ikx} \phi(x) dx \quad (k \in \mathbb{R})$$

and, for  $s > 0$ ,

$$\|\phi\|_{\mathcal{H}_s} := \|k \mapsto (1 + 2|k|)^s \widehat{\phi}(k)\|_{L^2}.$$

We define the Sobolev space  $\mathcal{H}_s$  as the set of functions with finite  $\|\cdot\|_{\mathcal{H}_s}$  norm. Let us now state Proposition 2 of the paper [ST10] of Shcherbina and Tirozzi.

**Proposition 16.** *For any  $s > 0$ , there is a constant  $C = C(s)$  such that for any  $n$ , any  $n \times n$  Hermitian random matrix  $M$ , and any  $\phi \in \mathcal{H}_s$ , we have*

$$\text{Var}(\text{Tr } \phi(M)) \leq C \|\phi\|_{\mathcal{H}_s}^2 \int_{y=0}^{\infty} y^{2s-1} e^{-y} \int_{x \in \mathbb{R}} \text{Var}(\text{Tr}((x + iy - M)^{-1})) dx dy.$$

## 2.4 A functional density lemma

We did not find Lemma 18 in the literature, so we provide its proof. Subsequently, the latter will be used with the CLT extension Lemma 15. Recall that for any  $z \in \mathbb{C} \setminus \mathbb{R}$ ,

$$\varphi_z(x) = \frac{1}{z - x}.$$

**Lemma 17.** *For any  $z \in \mathbb{C} \setminus \mathbb{R}$ , we have, in the  $L^2$  sense,*

$$\widehat{\varphi_z} = (t \mapsto -\operatorname{sgn}(\Im z) 2\pi i \mathbb{1}_{\Im(z)t > 0} e^{itz}) \quad (2.1)$$

and  $\varphi_z$  belongs to each  $\mathcal{H}_s$  for any  $s \in \mathbb{R}$ .

**Proof.** It is well known that if  $\Re z > 0$ , then  $\frac{1}{z} = \int_{t=0}^{+\infty} e^{-tz} dt$ .

Let  $z = E + i\eta$ ,  $E \in \mathbb{R}$ ,  $\eta > 0$ . For any  $\xi \in \mathbb{R}$ , we have

$$\varphi_z(\xi) = \frac{-i}{i(\xi - z)} = -i \int_{t=0}^{+\infty} e^{-it(\xi - z)} dt = -i \int_{t=0}^{+\infty} e^{-it\xi} e^{itz} dt.$$

We deduce (2.1) for  $\Im z > 0$ .

The general result can be deduced by complex conjugation.  $\square$

**Lemma 18.** *Let  $\mathcal{L}_1$  denote the linear span of the functions  $\varphi_z(x) := \frac{1}{z-x}$ , for  $z \in \mathbb{C} \setminus \mathbb{R}$ . Then the space  $\mathcal{L}_1$  is dense in  $\mathcal{H}_s$  for any  $s \in \mathbb{R}$ .*

**Proof.** We know, by Lemma 17, that  $\mathcal{L}_1 \subset \mathcal{H}_s$ . Recall first the definition of the Poisson kernel, for  $E \in \mathbb{R}$  and  $\eta > 0$ ,

$$P_\eta(E) = \frac{1}{\pi} \frac{\eta}{E^2 + \eta^2} = \frac{1}{2i\pi} (\varphi_{i\eta}(E) - \varphi_{-i\eta}(E))$$

and that, by Lemma 17,

$$\widehat{P_\eta}(t) = e^{-\eta|t|}.$$

Hence for any  $f \in \mathcal{H}_s$ , we have

$$\|f - P_\eta * f\|_{\mathcal{H}_s}^2 = \int (1 + 2|x|)^{2s} |\widehat{f}(x)|^2 (1 - e^{-\eta|x|})^2 dx,$$

so that, by dominated convergence,  $P_\eta * f \rightarrow f$  in  $\mathcal{H}_s$  as  $\eta \rightarrow 0$ .

To prove Lemma 18, it suffices to prove that any smooth compactly supported function can be approximated, in  $\mathcal{H}_s$ , by functions of  $\mathcal{L}_1$ . So let  $f$  be a smooth compactly supported function. By what precedes, it suffices to prove that for any fixed  $\eta > 0$ ,  $P_\eta * f$  can be approximated, in  $\mathcal{H}_s$ , by functions of  $\mathcal{L}_1$ . For  $x \in \mathbb{R}$ ,

$$\begin{aligned} P_\eta * f(x) &= \frac{1}{\pi} \int f(t) \frac{\eta}{\eta^2 + (x-t)^2} dt \\ &= -\frac{1}{\pi} \int f(t) \Im(\varphi_{t+i\eta}(x)) dt \\ &= \frac{1}{2\pi i} \int f(t) (\varphi_{t-i\eta}(x) - \varphi_{t+i\eta}(x)) dt. \end{aligned}$$

Without loss of generality, one can suppose that the support of  $f$  is contained in  $[0, 1]$ . Then, for any  $n \geq 1$ ,

$$P_\eta * f(x) = \frac{1}{2n\pi i} \sum_{k=1}^n f\left(\frac{k}{n}\right) \left( \varphi_{\frac{k}{n}-i\eta}(x) - \varphi_{\frac{k}{n}+i\eta}(x) \right) + R_n(x) \quad (2.2)$$

where for  $[t]_n := \lceil nt \rceil / n$ ,

$$R_n(x) = \frac{1}{2\pi i} \int f(t) (\varphi_{t-i\eta}(x) - \varphi_{t+i\eta}(x)) - f([t]_n) (\varphi_{[t]_n-i\eta}(x) - \varphi_{[t]_n+i\eta}(x)) dt.$$

The error term  $R_n(x)$  rewrites

$$\begin{aligned} R_n(x) &= \frac{1}{2\pi i} \int (f(t) - f([t]_n)) (\varphi_{t-i\eta} - \varphi_{t+i\eta})(x) dt \\ &\quad + \frac{1}{2\pi i} \int f([t]_n) (\varphi_{t-i\eta} - \varphi_{[t]_n-i\eta} + \varphi_{t+i\eta} - \varphi_{[t]_n+i\eta})(x) dt. \end{aligned}$$

Now, note that for any  $t \in \mathbb{R}$  and  $\eta \in \mathbb{R} \setminus \{0\}$ , we have by Lemma 17,

$$\widehat{\varphi_{t+i\eta}} = (x \mapsto -\operatorname{sgn}(\eta) 2\pi i \mathbb{1}_{\eta x > 0} e^{ixz}),$$

so that when, for example,  $\eta > 0$ , for any  $t \in \mathbb{R}$ ,

$$\|\varphi_{t+i\eta}\|_{\mathcal{H}_s}^2 = 4\pi^2 \int_0^\infty (1+2|x|)^{2s} e^{-2\eta x} dx$$

does not depend on  $t$  and for any  $t, t' \in \mathbb{R}$ ,

$$\begin{aligned} \|\varphi_{t+i\eta} - \varphi_{t'+i\eta}\|_{\mathcal{H}_s}^2 &= 4\pi^2 \int_0^\infty (1+2|x|)^{2s} |e^{itx} - e^{it'x}|^2 e^{-2\eta x} dx \\ &= 4\pi^2 \int_0^\infty (1+2|x|)^{2s} |e^{i(t-t')x} - 1|^2 e^{-2\eta x} dx \end{aligned}$$

depends only on  $t' - t$  and tends to zero (by dominated convergence) when  $t' - t \rightarrow 0$ .

We deduce that  $\|R_n\|_{\mathcal{H}_s} \rightarrow 0$  as  $n \rightarrow \infty$ , which closes the proof, by (2.2).  $\square$





# Chapter 3

## Empirical spectral distribution of a matrix under perturbation

### Contents

---

<b>3.1</b>	<b>Introduction</b>	<b>38</b>
<b>3.2</b>	<b>Main result</b>	<b>40</b>
3.2.1	Definition of the model and assumptions	40
3.2.2	Main result	42
<b>3.3</b>	<b>Examples</b>	<b>43</b>
3.3.1	Uniform measure perturbation by a band matrix	43
3.3.2	Triangular pulse perturbation by a Wigner matrix	44
3.3.3	Parabolic pulse perturbation by a Wigner matrix	44
<b>3.4</b>	<b>Relation to free probability theory</b>	<b>46</b>
<b>3.5</b>	<b>Strategy of the proof</b>	<b>48</b>
<b>3.6</b>	<b>Stieltjes transforms convergence</b>	<b>48</b>
3.6.1	Two statements	49
3.6.2	Proof of Proposition 21	50
3.6.3	Proof of Proposition 22	55
3.6.4	A local type convergence result	57
3.6.5	Possible extensions to larger $\varepsilon_n$	58
<b>3.7</b>	<b>Convergence in probability in the semi-perturbative regime</b>	<b>59</b>
<b>3.8</b>	<b>Convergence in distribution towards the Gaussian variable <math>Z_\phi</math></b>	<b>61</b>
3.8.1	Coincidence of the expectation of $\mu_n^\varepsilon$ with its deterministic approximation	61
3.8.2	Domination of the variance of $\mu_n^\varepsilon$	62

## 3.1 Introduction

In this chapter, we provide a perturbative expansion for the empirical spectral distribution of a Hermitian matrix with large size perturbed by a random matrix with small operator norm whose entries in the eigenvector basis of the first one are independent with a variance profile. More explicitly, let  $D_n$  be an  $n \times n$  Hermitian matrix, that, up to a change of basis, we suppose diagonal<sup>1</sup>. We denote by  $\mu_n$  the empirical spectral distribution of  $D_n$ . This matrix is additively perturbed by a random Hermitian matrix  $\varepsilon_n X_n$  whose entries are chosen at random independently and scaled so that the operator norm of  $X_n$  has order one. We are interested in the empirical spectral distribution  $\mu_n^\varepsilon$  of

$$D_n^\varepsilon := D_n + \varepsilon_n X_n$$

in the regime where the matrix size  $n$  tends to infinity and  $\varepsilon_n$  tends to 0. We shall prove that, depending on the order of magnitude of the perturbation, several regimes can appear. We suppose that  $\mu_n$  converges to a limiting measure  $\rho(\lambda)d\lambda$  and that the variance profile of the entries of  $X_n$  has a macroscopic limit  $\sigma_d$  on the diagonal and  $\sigma$  elsewhere. We then prove that there is a deterministic function  $F$  and a Gaussian random linear form  $dZ$  on the space of  $\mathcal{C}^6$  functions on  $\mathbb{R}$ , both depending only on the limit parameters of the model  $\rho, \sigma$  and  $\sigma_d$  such that if one defines the distribution  $dF : \phi \mapsto -\int \phi'(s)F(s)ds$ , then, for large  $n$ :

$$\mu_n^\varepsilon \approx \mu_n + \frac{\varepsilon_n}{n} dZ \quad \text{if } \varepsilon_n \ll n^{-1} \quad (3.1)$$

$$\mu_n^\varepsilon \approx \mu_n + \frac{\varepsilon_n}{n} (cdF + dZ) \quad \text{if } \varepsilon_n \sim \frac{c}{n} \quad (3.2)$$

$$\mu_n^\varepsilon \approx \mu_n + \varepsilon_n^2 dF \quad \text{if } n^{-1} \ll \varepsilon_n \ll 1 \quad (3.3)$$

and if, moreover,  $n^{-1} \ll \varepsilon_n \ll n^{-1/3}$ , then convergence (3.3) can be refined as follows:

$$\mu_n^\varepsilon \approx \mu_n + \varepsilon_n^2 dF + \frac{\varepsilon_n}{n} dZ. \quad (3.4)$$

In Section 3.3 several figures show a very good matching of random simulations with these theoretical results. The definitions of the function  $F$  and of the process  $Z$  are given below in (3.6) and (3.7). In many cases, the linear form  $dF$  can be interpreted as the integration with respect to the signed measure  $F'(x)dx$ . The function  $F$  is related to free probability theory, as explained in Section 3.4 below, whereas the linear form  $dZ$  is related to the so-called one-dimensional Gaussian free field defined, for instance, at [Dub09, Sect. 4.2].

---

<sup>1</sup>If the perturbing matrix belongs to the GOE or GUE, then its law is invariant under this change of basis, hence our results in fact apply to any self-adjoint matrix  $D_n$ .

If the variance profile of  $X_n$  is constant, then it is precisely the Laplacian of the Gaussian free field, defined in the sense of distributions.

The transition at  $\varepsilon_n \sim n^{-1}$  is the well-known transition, in quantum mechanics, where the *perturbative regime* ends. Indeed, one can distinguish the two following regimes:

- The regime  $\varepsilon_n \ll n^{-1}$ , called the *perturbative regime* (see [Fer00]): the size of the perturbation (i.e. its operator norm) is much smaller than the typical spacing between two consecutive eigenvalues (level spacing), which is of order  $n^{-1}$  in our setting.
- The regime  $n^{-1} \ll \varepsilon_n \ll 1$ , sometimes called the *semi-perturbative regime*, where the size of the perturbation is not small compared to the level spacing. This regime concerns many applications [LP11, AB12] in the context of covariance matrices and applications to finance.

A surprising fact discovered during this study is that the semi-perturbative regime  $n^{-1} \ll \varepsilon_n \ll 1$  decomposes into infinitely many sub-regimes. In the case  $n^{-1} \ll \varepsilon_n \ll n^{-1/3}$ , the expansion of  $\mu_n^\varepsilon - \mu_n$  contains a single deterministic term before the random term  $\frac{\varepsilon_n}{n} dZ$ . In the case  $n^{-1/3} \ll \varepsilon_n \ll n^{-1/5}$ , the expansion of  $\mu_n^\varepsilon - \mu_n$  contains two of them. More generally, for all positive integer  $p$ , when  $n^{-1/(2p-1)} \ll \varepsilon_n \ll n^{-1/(2p+1)}$ , the expansion contains  $p$  of them. For computational complexity reasons, the only case we state explicitly is the first one. We refer the reader to Section 3.6.5 for a discussion around this point.

In the papers [WW95, AB12, AB14, BABP16, ABB14], Wilkinson, Walker, Allez, Bouchaud *et al* have investigated some problems related to this one. Some of these works were motivated by the estimation of a matrix out of the observation of its noisy version. The present work differs from these ones mainly by the facts that firstly, we are interested in the perturbations of the *global* empirical distribution of the eigenvalues and not of a single one, and secondly, we push our expansion up to the random term, which does not appear in these papers. Besides, the noises they consider have constant variance profiles (either a Wigner-Dyson noise in the four first cited papers or a rotationally invariant noise in the fifth one). The transition at  $\varepsilon_n \sim n^{-1}$  between the perturbative and the semi-perturbative regimes is already present in these texts. They also consider the transition between the perturbative regime  $\varepsilon_n \ll 1$  and the *non perturbative* regime  $\varepsilon_n \asymp 1$ . As explained above, we exhibit the existence of an infinity of sub-regimes in this transition and focus on  $\varepsilon_n \ll 1$  for the first order of the expansion and to  $\varepsilon_n \ll n^{-1/3}$  for the second (and last) order. The study of other sub-regimes is postponed to forthcoming papers.

This chapter is organized as follows. Results, examples and comments are given in Sections 3.2 to 3.4, while the rest of the chapter is devoted to the proofs, except for Section 3.6.5, where we discuss the sub-regimes mentioned above.

**Notations.** For  $a_n, b_n$  some real sequences,  $a_n \ll b_n$  (resp.  $a_n \sim b_n$ ) means that  $a_n/b_n$  tends to 0 (resp. to 1). Also,  $\xrightarrow{P}$  and  $\xrightarrow{\text{dist.}}$  stand respectively for convergence in probability and convergence in distribution for all finite marginals.

## 3.2 Main result

### 3.2.1 Definition of the model and assumptions

For all positive integer  $n$ , we consider a real diagonal matrix  $D_n = \text{diag}(\lambda_n(1), \dots, \lambda_n(n))$ , as well as a Hermitian random matrix

$$X_n = \frac{1}{\sqrt{n}} [x_{i,j}^n]_{1 \leq i,j \leq n}$$

and a positive number  $\varepsilon_n$ . The normalizing factor  $n^{-1/2}$  and our hypotheses below ensure that the operator norm of  $X_n$  is of order one. We then define, for all  $n$ ,

$$D_n^\varepsilon := D_n + \varepsilon_n X_n.$$

We now introduce the probability measures  $\mu_n$  and  $\mu_n^\varepsilon$  as the respective uniform distributions on the eigenvalues (with multiplicity) of  $D_n$  and  $D_n^\varepsilon$ . Our aim is to give a perturbative expansion of  $\mu_n^\varepsilon$  around  $\mu_n$ .

We make the following hypotheses:

- (a) the entries  $x_{i,j}^n$  of  $\sqrt{n}X_n$  are independent (up to symmetry) random variables, centered, with variance denoted by  $\sigma_n^2(i, j)$ , such that  $\mathbb{E}|x_{i,j}^n|^8$  is bounded uniformly on  $n, i, j$ ,
- (b) there are  $f, \sigma_d, \sigma$  real functions defined respectively on  $[0, 1]$ ,  $[0, 1]$  and  $[0, 1]^2$  such that, for each  $x \in [0, 1]$ ,

$$\lambda_n(\lfloor nx \rfloor) \xrightarrow{n \rightarrow \infty} f(x) \quad \text{and} \quad \sigma_n^2(\lfloor nx \rfloor, \lfloor nx \rfloor) \xrightarrow{n \rightarrow \infty} \sigma_d(x)^2$$

and for each  $x \neq y \in [0, 1]$ ,

$$\sigma_n^2(\lfloor nx \rfloor, \lfloor ny \rfloor) \xrightarrow{n \rightarrow \infty} \sigma^2(x, y).$$

We make the following hypothesis about the rate of convergence:

$$\eta_n := \max\{n\varepsilon_n, 1\} \times \sup_{1 \leq i \neq j \leq n} (|\sigma_n^2(i, j) - \sigma^2(i/n, j/n)| + |\lambda_n(i) - f(i/n)|) \xrightarrow{n \rightarrow \infty} 0.$$

Let us now make some assumptions on the limiting functions  $\sigma$  and  $f$ :

- (c) the function  $f$  is bounded and the push-forward of the uniform measure on  $[0, 1]$  by the function  $f$  has a density  $\rho$  with respect to the Lebesgue measure on  $\mathbb{R}$  and a compact support denoted by  $\mathcal{S}$ ,

- (d) the variance of the entries of  $X_n$  essentially depends on the eigenspaces of  $D_n$ , namely, there exists a symmetric function  $\tau(\cdot, \cdot)$  on  $\mathbb{R}^2$  such that for all  $x \neq y$ ,  $\sigma^2(x, y) = \tau(f(x), f(y))$ ,
- (e) the following regularity property holds: there exist  $\eta_0 > 0, \alpha > 0$  and  $C < \infty$  such that for almost all  $s \in \mathbb{R}$ , for all  $t \in [s - \eta_0, s + \eta_0]$ ,  $|\tau(s, t)\rho(t) - \tau(s, s)\rho(s)| \leq C|t - s|^\alpha$ .

We add a last assumption which strengthens assumption (c) and makes it possible to include the case where the set of eigenvalues of  $D_n$  contains some outliers:

- (f) there is a real compact set  $\tilde{\mathcal{S}}$  such that

$$\max_{1 \leq i \leq n} \text{dist}(\lambda_n(i), \tilde{\mathcal{S}}) \xrightarrow{n \rightarrow \infty} 0.$$

**Remark 1** (About the hypothesis that  $D_n$  is diagonal). (i) If the perturbing matrix  $X_n$  belongs to the GOE (resp. to the GUE), then its law is invariant under conjugation by any orthogonal (resp. unitary) matrix. It follows that in this case, our results apply to any real symmetric (resp. Hermitian) matrix  $D_n$  with eigenvalues  $\lambda_n(i)$  satisfying the above hypotheses.

(ii) As explained after Proposition 20 below, we conjecture that when the variance profile of  $X_n$  is constant, for  $\varepsilon_n \gg n^{-1}$ , we do not need the hypothesis that  $D_n$  is diagonal neither. However, if the perturbing matrix does not have a constant variance profile, then for a non-diagonal  $D_n$  and  $\varepsilon \gg n^{-1}$ , the spectrum of  $D_n^\varepsilon$  should depend heavily on the relation between the eigenvectors of  $D_n$  and the variance profile, which implies that our results should not remain true.

(iii) At last, it is easy to see that the random process  $(Z_\phi)$  introduced at (3.7) satisfies, for any test function  $\phi$ ,

$$\frac{1}{\varepsilon_n} \sum_{i=1}^n \left( \phi(\lambda_n(i) + \frac{\varepsilon_n}{\sqrt{n}} x_{ii}) - \phi(\lambda_n(i)) \right) \xrightarrow[n \rightarrow \infty]{\text{dist.}} Z_\phi.$$

Thus, regardless to the variance profile, the convergence of (3.8) rewrites, informally,

$$\mu_n^\varepsilon = \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_n(i) + (\varepsilon_n/\sqrt{n})x_{ii}} + o(\varepsilon_n/n). \quad (3.5)$$

A so simple expression, up to a  $o(\varepsilon_n/n)$  error, of the empirical spectral distribution of  $D_n^\varepsilon$ , with some independent translations  $\frac{\varepsilon_n}{\sqrt{n}}x_{ii}$ , should not remain true without the hypothesis that  $D_n$  is diagonal or that the distribution of  $X_n$  is invariant under conjugation.

### 3.2.2 Main result

Recall that the *Hilbert transform*, denoted by  $H[u]$ , of a function  $u$ , is the function

$$H[u](s) := \text{p. v.} \int_{t \in \mathbb{R}} \frac{u(t)}{s - t} dt$$

and define the function

$$F(s) = -\rho(s)H[\tau(s, \cdot)\rho(\cdot)](s). \quad (3.6)$$

Note that, by assumptions (c) and (e),  $F$  is well defined and supported by  $\mathcal{S}$ . Besides, for any  $\phi$  supported on an interval where  $F$  is  $\mathcal{C}^1$ ,

$$-\int \phi'(s)F(s)ds = \int \phi(s)dF(s),$$

where  $dF(s)$  denotes the measure  $F'(s)ds$ .

We also introduce the centered Gaussian field,  $(Z_\phi)_{\phi \in \mathcal{C}^6}$ , indexed by the set of  $\mathcal{C}^6$  complex functions on  $\mathbb{R}$ , with covariance defined by

$$\mathbb{E}Z_\phi Z_\psi = \int_0^1 \sigma_d(t)^2 \phi'(f(t))\psi'(f(t))dt \quad \text{and} \quad \overline{Z_\psi} = Z_{\overline{\psi}}. \quad (3.7)$$

Note that the process  $(Z_\phi)_{\phi \in \mathcal{C}^6}$  can be represented, for  $(B_t)$  is the standard one-dimensional Brownian motion, as

$$Z_\phi = \int_0^1 \sigma_d(t)\phi'(f(t))dB_t.$$

**Theorem 19.** *For all compactly supported  $\mathcal{C}^6$  function  $\phi$  on  $\mathbb{R}$ , the following convergences hold:*

- **Perturbative regime:** if  $\varepsilon_n \ll n^{-1}$ , then,

$$n\varepsilon_n^{-1}(\mu_n^\varepsilon - \mu_n)(\phi) \xrightarrow[n \rightarrow \infty]{\text{dist.}} Z_\phi. \quad (3.8)$$

- **Critical regime:** if  $\varepsilon_n \sim c/n$ , with  $c$  constant, then,

$$n\varepsilon_n^{-1}(\mu_n^\varepsilon - \mu_n)(\phi) \xrightarrow[n \rightarrow \infty]{\text{dist.}} -c \int \phi'(s)F(s)ds + Z_\phi. \quad (3.9)$$

- **Semi-perturbative regime:** if  $n^{-1} \ll \varepsilon_n \ll 1$ , then,

$$\varepsilon_n^{-2}(\mu_n^\varepsilon - \mu_n)(\phi) \xrightarrow[n \rightarrow \infty]{P} - \int \phi'(s)F(s)ds, \quad (3.10)$$

and if, moreover,  $n^{-1} \ll \varepsilon_n \ll n^{-1/3}$ , then,

$$n\varepsilon_n^{-1} \left( (\mu_n^\varepsilon - \mu_n)(\phi) + \varepsilon_n^2 \int \phi'(s)F(s)ds \right) \xrightarrow[n \rightarrow \infty]{\text{dist.}} Z_\phi. \quad (3.11)$$

**Remark 2** (Sub-regimes for  $n^{-1/3} \ll \varepsilon_n \ll 1$ ). In the semi-perturbative regime, the reason why we provide an expansion up to a random term, only for  $\varepsilon_n \ll n^{-1/3}$ , is that the study of the regime  $n^{-1/3} \ll \varepsilon_n \ll 1$  up to such a precision, requires further terms in the expansion of the resolvent of  $D_n^\varepsilon$  that make appear, beside  $dF$ , additional deterministic terms of smaller order, which are much larger than the probabilistic term containing  $Z_\phi$ . The computation becomes rather intricate without any clear recursive formula. As we will see in Section 3.6.5, there are infinitely many regimes. Precisely, for any positive integer  $p$ , when  $n^{-1/(2p-1)} \ll \varepsilon_n \ll n^{-1/(2p+1)}$ , there are  $p$  deterministic terms in the expansion before the term in  $Z_\phi$ .

**Remark 3** (Local law). The approximation

$$\mu_n^\varepsilon(I) \approx \mu_n(I) + \varepsilon_n^2 \int_I dF$$

of (3.10) should stay true even for intervals  $I$  with size tending to 0 as the dimension  $n$  grows, as long as the size of  $I$  stays much larger than the right-hand side term of (3.30), as can be seen from Proposition 23.

**Remark 4.** The second part of Hypothesis (b), concerning the speed of convergence of the profile of the spectrum of  $D_n$  as well as of the variance of its perturbation, is needed in order to express the expansion of  $\mu_n^\varepsilon - \mu_n$  in terms of limit parameters of the model  $\sigma$  and  $\rho$ . We can remove this hypothesis and get analogous expansions where the terms  $dF$  and  $dZ$  are replaced by their discrete counterparts  $dF_n$  and  $dZ_n$ , defined thanks to the “finite  $n$ ” empirical versions of the limit parameters  $\sigma$  and  $\rho$ .

## 3.3 Examples

### 3.3.1 Uniform measure perturbation by a band matrix

Here, we consider the case where  $f(x) = x$ ,  $\sigma_d(x) \equiv m$  and  $\sigma(x, y) = \mathbb{1}_{|y-x| \leq \ell}$ , for some constants  $m \geq 0$  and  $\ell \in [0, 1]$  (the relative width of the band). In this case,  $\tau(\cdot, \cdot) = \sigma(\cdot, \cdot)^2$ , hence


$$F(s) = \mathbb{1}_{(0,1)}(s) \text{ p. v. } \int_t \frac{\tau(s, t)}{s - t} dt = -\mathbb{1}_{(0,1)}(s) \log \frac{\ell \wedge (1 - s)}{\ell \wedge s} \quad (3.12)$$

and  $(Z_\phi)_{\phi \in \mathcal{C}^6}$  is the centered complex Gaussian process with covariance defined by

$$\mathbb{E} Z_\phi \overline{Z_\psi} = m^2 \int_0^1 \phi'(t) \overline{\psi'(t)} dt \quad \text{and} \quad \overline{Z_\psi} = Z_{\overline{\psi}}.$$

Theorem 19 is then illustrated by Figure 3.1, where we plotted the cumulative distribution functions.





**Figure 3.1: Deforming the uniform distribution by a band matrix.** Cumulative distribution function of  $\varepsilon_n^{-2}(\mu_n^\varepsilon - \mu_n)$  (in blue) and function  $F(\cdot)$  of (3.12) (in red). The non smoothness of the blue curves results of the noise term  $Z_\phi$  in Theorem 19. Each graphic is realized thanks to one single matrix (no averaging) perturbed by a real Gaussian band matrix.

### 3.3.2 Triangular pulse perturbation by a Wigner matrix

Here, we consider the case where  $\rho(x) = (1 - |x|)\mathbb{1}_{[-1,1]}(x)$ ,  $\sigma_d \equiv m$ , for some real constant  $m$ , and  $\sigma \equiv 1$  (what follows can be adapted to the case  $\sigma(x, y) = \mathbb{1}_{|y-x| \leq \ell}$ , with a bit longer formulas). In this case, thanks to the formula (9.6) of  $H[\rho(\cdot)]$  given p. 509 of [Kin09], we get

$$F(s) = (1 - |s|)\mathbb{1}_{[-1,1]}(s) \{ (1 - s) \log(1 - s) - (1 + s) \log(1 + s) + 2s \log |s| \}. \quad (3.13)$$

and the covariance of  $(Z_\phi)_{\phi \in \mathcal{C}^6}$  is given by

$$\mathbb{E} Z_\phi \overline{Z_\psi} = m^2 \int_{-1}^1 (1 - |t|) \phi'(t) \overline{\psi'(t)} dt \quad \text{and} \quad \overline{Z_\psi} = Z_{\overline{\psi}}.$$

Theorem 19 is then illustrated by Figure 3.2 in the case where  $\varepsilon_n \gg n^{-1/2}$ . In Figure 3.2, we implicitly use some test functions of the type  $\phi(x) = \mathbb{1}_{x \in I}$  for some intervals  $I$ . These functions are not  $\mathcal{C}^6$ , and one can easily see that for  $\varepsilon_n \ll n^{-1/2}$ , Theorem 19 cannot work for such functions. However, considering imaginary parts of Stieltjes transforms, i.e. test functions

$$\phi(x) = \frac{1}{\pi} \frac{\eta}{(x - E)^2 + \eta^2} \quad (E \in \mathbb{R}, \eta > 0)$$

gives a perfect matching between the predictions from Theorem 19 and numerical simulations, also for  $\varepsilon_n \ll n^{-1/2}$  (see Figure 3.3, where we use Proposition 22 and (3.17) to compute the theoretical limit).

### 3.3.3 Parabolic pulse perturbation by a Wigner matrix

Here, we consider the case where  $\rho(x) = \frac{3}{4}(1 - x^2)\mathbb{1}_{[-1,1]}(x)$ ,  $\sigma_d \equiv m$ , for some real constant  $m$ , and  $\sigma \equiv 1$  (again, this can be adapted to the case  $\sigma(x, y) = \mathbb{1}_{|y-x| \leq \ell}$ ). Theorem 19 is




Figure 3.2: **Triangular pulse perturbation by a Wigner matrix: density and cumulative distribution function.** *Top left:* Cumulative distribution function of  $\varepsilon_n^{-2}(\mu_n^\varepsilon - \mu_n)$  (in blue) and function  $F(\cdot)$  of (3.13) (in red). *Top right and bottom:* Density  $\rho$  (red dashed line), histogram of the eigenvalues of  $D_n^\varepsilon$  (in black) and theoretical density  $\rho + \varepsilon_n^2 F'(s)$  of the eigenvalues of  $D_n^\varepsilon$  as predicted by Theorem 19 (in blue). Here,  $n = 10^4$  and  $\varepsilon_n = n^{-\alpha}$ , with  $\alpha = 0.25$  (up left),  $\alpha = 0.4$  (up right),  $0.25$  (bottom left) and  $0.1$  (bottom right).




Figure 3.3: **Triangular pulse perturbation by a Wigner matrix: Stieltjes transform.** Imaginary part of the Stieltjes transform of  $\varepsilon_n^{-2}(\mu_n^\varepsilon - \mu_n)$  (in blue) and of the measure  $dF$  (in red) at  $z = E + i$  as a function of the real part  $E$  for different values of  $\varepsilon_n$ . Here,  $n = 10^4$  and  $\varepsilon_n = n^{-\alpha}$ , with  $\alpha = 0.2, 0.5$  and  $0.8$  (from left to right).

then illustrated by Figure 3.4. In this case, thanks to the formula (9.10) of  $H[\rho(\cdot)]$  given p. 509 of [Kin09], we get

$$F(s) = -\frac{9}{16}(1-s^2)\mathbb{1}_{[-1,1]}(s) \left\{ 2s - (1-s^2) \ln \left| \frac{s-1}{s+1} \right| \right\} \quad (3.14)$$

and the covariance of  $(Z_\phi)_{\phi \in \mathcal{C}^6}$  is given by

$$\mathbb{E} Z_\phi \overline{Z_\psi} = \frac{3m^2}{4} \int_{-1}^1 (1-t^2) \phi'(t) \overline{\psi'(t)} dt \quad \text{and} \quad \overline{Z_\psi} = Z_{\overline{\psi}}.$$




Figure 3.4: **Parabolic pulse perturbation by a Wigner matrix.** *Top left:* Cumulative distribution function of  $\varepsilon_n^{-2}(\mu_n^\varepsilon - \mu_n)$  (in blue) and function  $F(\cdot)$  of (3.14) (in red). *Top right and bottom:* Density  $\rho$  (red dashed line), histogram of the eigenvalues of  $D_n^\varepsilon$  (in black) and theoretical density  $\rho + \varepsilon_n^2 F'(s)$  of the eigenvalues of  $D_n^\varepsilon$  as predicted by Theorem 19 (in blue). Here,  $n = 10^4$  and  $\varepsilon_n = n^{-\alpha}$ , with  $\alpha = 0.25$  (up left),  $\alpha = 0.4$  (up right),  $0.2$  (bottom left) and  $0.18$  (bottom right).

### 3.4 Relation to free probability theory

Let us now explain how this work is related to free probability theory. If, instead of letting  $\varepsilon_n$  tend to zero, one considers the model

$$D_n^t := D_n + \sqrt{t} X_n$$

for a fixed  $t > 0$ , then, by [CG93a, CG93b, Shl96, AZ06], the empirical eigenvalue distribution of  $D_n^t$  has a limit as  $n \rightarrow \infty$ , that we shall denote here by  $\mu_t$ . The law  $\mu_t$  can

be interpreted as the law of the sum of two elements in a non-commutative probability space which are free with an amalgamation over a certain sub-algebra (see [Shl96] for more details). The following proposition relates the function  $F$  from (3.6) to the first order expansion of  $\mu_t$  around  $t = 0$ .

**Proposition 20.** *For any  $z \in \mathbb{C} \setminus \mathbb{R}$ , we have*

$$\frac{\partial}{\partial t} \Big|_{t=0} \int \frac{d\mu_t(\lambda)}{z - \lambda} = - \int \frac{F(\lambda)}{(z - \lambda)^2} d\lambda = - \int F(\lambda) \frac{\partial}{\partial \lambda} \left( \frac{1}{z - \lambda} \right) d\lambda.$$

This is related to the fact that in Equations (3.1)–(3.4), for  $\varepsilon_n$  large enough, the term  $\varepsilon_n^2 dF$  is the leading term.

In the particular case where  $X_n$  is a Wigner matrix,  $\mu_t$  is the free convolution of the measure  $\rho(\lambda)d\lambda$  with a semicircle distribution and admits a density  $\rho_t$ , by [Bia97b, Cor. 2]. Then, Theorem 19 makes it possible to formally recover the *free Fokker-Planck equation with null potential*:

$$\begin{cases} \frac{\partial}{\partial t} \rho_t(s) + \frac{\partial}{\partial s} \{ \rho_t(s) H[\rho_t](s) \} = 0, \\ \rho_0(s) = \rho(s), \end{cases}$$

where  $H[\rho_t]$  denotes the Hilbert transform of  $\rho_t$ . This equation is also called *McKean-Vlasov* (or *Fokker-Planck*) *equation with logarithmic interaction* (see [Bia97a, BS98, BS01]).

Note also that when  $X_n$  is a Wigner matrix, the hypothesis that  $D_n$  is diagonal is not required to have the convergence of the empirical eigenvalue distribution of  $D_n^t$  to  $\mu_t$  as  $n \rightarrow \infty$ . This suggests that, even for non diagonal  $D_n$ , the convergence of (3.10) still holds when  $X_n$  is a Wigner matrix.

**Proof of Proposition 20.** By [Shl96, Th. 4.3], we have

$$\int \frac{d\mu_t(\lambda)}{z - \lambda} = \int_{x=0}^1 C_t(x, z) dx, \quad (3.15)$$

where  $C_t(x, z)$  is bounded by  $|\Im z|^{-1}$  and satisfies the fixed-point equation

$$C_t(x, z) = \frac{1}{z - f(x) - t \int_{y=0}^1 \sigma^2(x, y) C_t(y, z) dy}.$$

Hence as  $t \rightarrow 0$ ,  $C_t(x, z) \rightarrow \frac{1}{z - f(x)}$  uniformly in  $x$ . Thus

$$\begin{aligned} C_t(x, z) - \frac{1}{z - f(x)} &= \frac{t \int_{y=0}^1 \sigma^2(x, y) C_t(y, z) dy}{(z - f(x) - t \int_{y=0}^1 \sigma^2(x, y) C_t(y, z) dy)(z - f(x))} \\ &= t \frac{1}{(z - f(x))^2} \int_{y=0}^1 \sigma^2(x, y) C_t(y, z) dy + o(t) \\ &= t \frac{1}{(z - f(x))^2} \int_{y=0}^1 \frac{\sigma^2(x, y)}{z - f(y)} dy + o(t) \end{aligned}$$

where each  $o(t)$  is uniform in  $x \in [0, 1]$ . Then, by (3.15), we deduce that

$$\frac{\partial}{\partial t} \Big|_{t=0} \int \frac{d\mu_t(\lambda)}{z - \lambda} = \int_{(x,y) \in [0,1]^2} \frac{\sigma^2(x,y)}{(z - f(x))^2(z - f(y))} dx dy.$$

The right-hand side term of the previous equation is precisely the number  $B(z)$  introduced at (3.17) below. Then, one concludes using Proposition 22 from Section 3.6.1.  $\square$

### 3.5 Strategy of the proof

We shall first prove the convergence results of Theorem 19 for test functions  $\phi$  of the form  $\varphi_z(x) := \frac{1}{z-x}$ . This is done in Section 3.6 by writing an expansion of the resolvent of  $D_n^\varepsilon$ .

Once we have proved that the convergences hold for the resolvent of  $D_n^\varepsilon$ , we can extend them to the larger class of compactly supported  $\mathcal{C}^6$  functions on  $\mathbb{R}$ .

In Section 3.7, we use the Helffer-Sjöstrand formula to extend the convergence in probability in the semi-perturbative regime (3.10) to the case of compactly supported  $\mathcal{C}^6$  functions on  $\mathbb{R}$ .

In Section 3.8, the convergences in distribution (3.8), (3.9) and (3.11) are proved in two steps. The overall strategy is to apply an extension lemma of Shcherbina and Tirozzi which states that a CLT that applies to a sequence of *centered* random linear forms on some space can be extended, by density, to a larger space, as long as the variance of the image of these random linear forms by a function  $\phi$  of the larger space is uniformly bounded by the norm of  $\phi$ . Therefore, our task is twofold. We need first to prove that the sequences of variables involved in the convergences (3.8), (3.9) and (3.11) can be replaced by their *centered* counterparts  $n\varepsilon_n^{-1}(\mu_n^\varepsilon(\phi) - \mathbb{E}[\mu_n^\varepsilon(\phi)])$  (i.e. they differ by  $o(1)$ ). In a second step, we dominate the variance of these latter variables, in order to apply the extension lemma which is precisely stated in Chapter 2 as Lemma 15.

### 3.6 Stieltjes transforms convergence

As announced in the previous section, we start with the proof of Theorem 19 in the special case of test functions of the type  $\varphi_z := \frac{1}{z-x}$ . We decompose it into two propositions. Their statement and proof are the purpose of the three following subsections. The two last subsections 3.6.4 and 3.6.5 are devoted respectively to a local type convergence result and to a discussion about the possibility of an extension of the expansion result to a wider range of rate of convergence of  $\varepsilon_n$ , namely beyond  $n^{-1/3}$ .

### 3.6.1 Two statements

Let denote, for  $z \in \mathbb{C} \setminus \mathbb{R}$ ,

$$Z(z) := Z_{\varphi_z} \quad \text{for} \quad \varphi_z(x) := \frac{1}{z - x} \quad (3.16)$$

where  $(Z_\phi)_{\phi \in \mathcal{C}^6}$  is the Gaussian field with covariance defined by (3.7). We also introduce, for  $z \in \mathbb{C} \setminus \mathbb{R}$ ,

$$B(z) := \int_{(s,t) \in [0,1]^2} \frac{\sigma^2(s,t)}{(z - f(s))^2(z - f(t))} ds dt \quad (3.17)$$

and

$$\Delta G_n(z) := (\mu_n^\varepsilon - \mu_n)(\varphi_z) = \frac{1}{n} \text{Tr} \frac{1}{z - D_n^\varepsilon} - \frac{1}{n} \text{Tr} \frac{1}{z - D_n}. \quad (3.18)$$

**Proposition 21.** *Under Hypotheses (a), (b), (f),*

- if  $\varepsilon_n \ll n^{-1}$ , then for all  $z \in \mathbb{C} \setminus \mathbb{R}$ ,

$$n\varepsilon_n^{-1} \Delta G_n(z) \xrightarrow[n \rightarrow \infty]{\text{dist.}} Z(z) \quad (3.19)$$

- if  $\varepsilon_n \sim c/n$ , with  $c$  constant, then for all  $z \in \mathbb{C} \setminus \mathbb{R}$

$$n\varepsilon_n^{-1} \Delta G_n(z) \xrightarrow[n \rightarrow \infty]{\text{dist.}} cB(z) + Z(z), \quad (3.20)$$

- if  $n^{-1} \ll \varepsilon_n \ll n^{-1/3}$ , then for all  $z \in \mathbb{C} \setminus \mathbb{R}$

$$n\varepsilon_n^{-1} (\Delta G_n(z) - \varepsilon_n^2 B(z)) \xrightarrow[n \rightarrow \infty]{\text{dist.}} Z(z). \quad (3.21)$$

- if  $n^{-1} \ll \varepsilon_n \ll 1$ , then for all  $z \in \mathbb{C} \setminus \mathbb{R}$ ,

$$\varepsilon_n^{-2} \Delta G_n(z) - B(z) \xrightarrow[n \rightarrow \infty]{P} 0. \quad (3.22)$$

**Remark.** Note that (3.20) is merely an extension of (3.21) in the critical regime.

The following statement expresses  $B(z)$  as the image of a  $\varphi_z$  by a linear form. So, in the expansion of the previous proposition, both quantities  $Z(z)$  and  $B(z)$  depend linearly on  $\varphi_z$ . Note that as  $F$  vanishes at  $\pm\infty$ , Proposition 22 does not contradicts the fact that as  $|z|$  gets large,  $B(z) = O(|z|^{-3})$ .

**Proposition 22.** *Under Hypotheses (c), (d), (e), for any  $z \in \mathbb{C} \setminus \mathcal{S}$ , for  $F$  defined by (3.6),*

$$B(z) = - \int \frac{F(s)}{(z - s)^2} ds = - \int \varphi'_z(s) F(s) ds.$$

### 3.6.2 Proof of Proposition 21

The proof is based on a perturbative expansion of the resolvent  $\frac{1}{n} \text{Tr} \frac{1}{z - D_n^\varepsilon}$ . To make notations lighter, we shall sometimes suppress the subscripts and superscripts  $n$ , so that  $D_n^\varepsilon$ ,  $D_n$ ,  $X_n$  and  $x_{i,j}^n$  will be respectively denoted by  $D^\varepsilon$ ,  $D$ ,  $X$  and  $x_{i,j}$ . Let us fix  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ . We can deduce from the expansion of the resolvent of  $D^\varepsilon$ :

$$\Delta G_n(z) = A_n(z) + B_n(z) + C_n(z) + R_n^\varepsilon(z),$$

with

$$\begin{aligned} A_n(z) &:= \frac{\varepsilon_n}{n} \text{Tr} \frac{1}{z - D} X \frac{1}{z - D} = \frac{\varepsilon_n}{n} \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{x_{i,i}}{(z - \lambda_n(i))^2} \\ B_n(z) &:= \frac{\varepsilon_n^2}{n} \text{Tr} \frac{1}{z - D} X \frac{1}{z - D} X \frac{1}{z - D} = \frac{\varepsilon_n^2}{n^2} \sum_{i,j} \frac{|x_{i,j}|^2}{(z - \lambda_n(i))^2 (z - \lambda_n(j))} \\ C_n(z) &:= \frac{\varepsilon_n^3}{n} \text{Tr} \frac{1}{z - D} X \frac{1}{z - D} X \frac{1}{z - D} X \frac{1}{z - D} \\ &= \frac{\varepsilon_n^3}{n^{5/2}} \sum_{i,j,k=1}^n \frac{x_{i,j} x_{j,k} x_{k,i}}{(z - \lambda_n(i))^2 (z - \lambda_n(j)) (z - \lambda_n(k))} \\ R_n^\varepsilon(z) &:= \frac{\varepsilon_n^4}{n} \text{Tr} \frac{1}{z - D} X \frac{1}{z - D} X \frac{1}{z - D} X \frac{1}{z - D} X \frac{1}{z - D^\varepsilon}. \end{aligned}$$

The purpose of the four following claims is to describe the asymptotic behavior of each of these four terms.

**Claim 1.** *The finite dimension marginals of the centered process*

$$(n\varepsilon_n^{-1} A_n(z))_{z \in \mathbb{C} \setminus \tilde{\mathcal{S}}}$$

*converge in distribution to those of the centered Gaussian process  $(Z(z))_{z \in \mathbb{C} \setminus \tilde{\mathcal{S}}}$ . Besides, there is  $C > 0$  such that for any  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ ,*

$$\mathbb{E}[|n\varepsilon_n^{-1} A_n(z)|^2] \leq \frac{C}{\text{dist}(z, \tilde{\mathcal{S}})^4}. \quad (3.23)$$

**Proof.** Estimate (3.23) follows from

$$\mathbb{E}[|A_n(z)|^2] = \frac{\varepsilon_n^2}{n^3} \sum_{i=1}^n \frac{\mathbb{E}[|x_{i,i}|^2]}{|z - \lambda_n(i)|^4} \leq \frac{\varepsilon_n^2}{n^3} \sum_{i=1}^n \frac{\sigma_n^2(i, i)}{\text{dist}(z, \tilde{\mathcal{S}})^4}$$

and from the existence of a uniform upper bound for  $\sigma_n^2(i, i)$  which comes from Hypothesis (a) which stipulates that the 8-th moments of the entries  $x_{i,j}$  are uniformly bounded.

We turn now to the proof of the convergence in distribution of  $n\varepsilon_n^{-1}A_n(z)$  which actually does not depend on the sequence  $(\varepsilon_n)$ . For all  $\alpha_1, \beta_1, \dots, \alpha_p, \beta_p \in \mathbb{C}$  and for all  $z_1, \dots, z_p \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ ,

$$\sum_{i=1}^p \alpha_i (n\varepsilon_n^{-1}A_n(z_i)) + \beta_i \overline{(n\varepsilon_n^{-1}A_n(z_i))} = \frac{1}{\sqrt{n}} \sum_{j=1}^n x_{j,j} \left( \sum_{i=1}^p \xi_n(i, j) \right)$$

$$\text{for } \xi_n(i, j) = \frac{\alpha_i}{(z_i - \lambda_n(j))^2} + \frac{\beta_i}{(\bar{z}_i - \lambda_n(j))^2}.$$

On one hand, by dominated convergence, the covariance matrix of the above two dimensional random vector converges.

On the other hand,  $\mathbb{E}|x_{i,j}|^4$  is uniformly bounded in  $i, j$  and  $n$ , by Hypothesis (a). Moreover, for  $n$  large enough, for all  $i, j$ ,

$$|\xi_n(i, j)| \leq 2 \max_{1 \leq i \leq p} (|\alpha_i| + |\beta_i|) \times \left( \min_{1 \leq i \leq p} \text{dist}(z_i, \mathcal{S}) \right)^{-1}.$$

Hence, the conditions of Lindeberg Central Limit Theorem are satisfied and the finite dimension marginals of the process  $(n\varepsilon_n^{-1}A_n(z))_{z \in \mathbb{C} \setminus \tilde{\mathcal{S}}}$  converge in distribution to those of the centered Gaussian process  $(Z_z)_{z \in \mathbb{C} \setminus \tilde{\mathcal{S}}}$  defined by its covariance structure

$$\begin{aligned} \mathbb{E} \left( Z(z) \overline{Z(z')} \right) &= \lim_{n \rightarrow \infty} \mathbb{E} \left[ (n\varepsilon_n^{-1}A_n(z)) \cdot \overline{(n\varepsilon_n^{-1}A_n(z'))} \right] \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i,j=1}^n \frac{\mathbb{E}[x_{i,i} \overline{x_{j,j}}]}{(z - \lambda_n(i))^2 (\bar{z}' - \lambda_n(j))^2} \\ &= \int_0^1 \frac{\sigma_d(t)^2}{(z - f(t))^2 (\bar{z}' - f(t))^2} dt \end{aligned}$$

and by the fact that  $\overline{Z(z)} = Z(\bar{z})$  which comes from  $\overline{A_n(z)} = A_n(\bar{z})$ . □

**Claim 2.** *There is a constant  $C$  such that, for  $\eta_n$  as in Hypotheses (b),*

- if  $\varepsilon_n \ll n^{-1}$ , then

$$\mathbb{E}[|n\varepsilon_n^{-1}B_n(z)|^2] \leq \frac{C(n\varepsilon_n)^2}{\text{dist}(z, \tilde{\mathcal{S}})^6} + \frac{C\eta_n^2}{\text{dist}(z, \tilde{\mathcal{S}})^8},$$

- if  $\varepsilon_n \sim c/n$  or if  $n^{-1} \ll \varepsilon_n \ll 1$ , then

$$\mathbb{E}[|n\varepsilon_n^{-1}(B_n(z) - \varepsilon_n^2 B(z))|^2] \leq \frac{C\varepsilon_n^2}{\text{dist}(z, \tilde{\mathcal{S}})^6} + \frac{C\eta_n^2}{\text{dist}(z, \tilde{\mathcal{S}})^8}.$$

**Proof.** Remind that,

$$B_n(z) = \frac{\varepsilon_n^2}{n^2} \sum_{i,j} \frac{|x_{i,j}|^2}{(z - \lambda_n(i))^2 (z - \lambda_n(j))}.$$



Introduce the variable  $b_n^\circ(z)$  obtained by centering the variable  $n\varepsilon_n^{-2}B_n(z)$ :

$$b_n^\circ(z) := n\varepsilon_n^{-2}(B_n(z) - \mathbb{E}B_n(z)) = \frac{1}{n} \sum_{i,j} \frac{|x_{i,j}|^2 - \sigma_n^2(i,j)}{(z - \lambda_n(i))^2(z - \lambda_n(j))}$$

and the defect variable

$$\begin{aligned} \delta_n(z) &:= \varepsilon_n^{-2} \mathbb{E}B_n(z) - B(z) \\ &= \frac{1}{n^2} \sum_{i,j} \frac{\sigma_n^2(i,j)}{(z - \lambda_n(i))^2(z - \lambda_n(j))} - \int_{(s,t) \in [0,1]^2} \frac{\sigma^2(s,t)}{(z - f(s))^2(z - f(t))} \mathrm{d}s \mathrm{d}t. \end{aligned}$$

In the two regimes  $\varepsilon_n \ll n^{-1}$  and  $\varepsilon_n \geq c/n$ , we want to dominate the  $L^2$  norms respectively of

$$n\varepsilon_n^{-1}B_n(z) = \varepsilon_n b_n^\circ(z) + n\varepsilon_n(\delta_n(z) + B(z)) \quad \text{and} \quad n\varepsilon_n^{-1}(B_n(z) - \varepsilon_n^2 B(z)) = \varepsilon_n b_n^\circ + n\varepsilon_n \delta_n(z).$$

For this purpose, we successively dominate  $b_n^\circ$ ,  $\delta_n(z)$  and  $B(z)$ .

Using the independence of the  $x_{i,j}$ 's, the fact that they are bounded in  $L^4$  and the fact that  $z$  stays at a macroscopic distance of the  $\lambda_n(i)$ 's, we can write for all  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$

$$\begin{aligned} \mathbb{E}[|b_n^\circ(z)|^2] &= \frac{1}{n^2} \mathrm{Var} \left( \sum_{i \leq j} (x_{i,j}^2 + \mathbb{1}_{i \neq j} \overline{x_{i,j}}^2) \frac{1}{(z - \lambda_n(i))^2(z - \lambda_n(j))} \right) \\ &= \frac{1}{n^2} \sum_{i \leq j} \mathrm{Var} \left( (x_{i,j}^2 + \mathbb{1}_{i \neq j} \overline{x_{i,j}}^2) \frac{1}{(z - \lambda_n(i))^2(z - \lambda_n(j))} \right) \\ &\leq C \mathrm{dist}(z, \tilde{\mathcal{S}})^{-6}. \end{aligned} \tag{3.24}$$

Now, the term  $\delta_n(z)$  rewrites

$$\begin{aligned} \delta_n(z) &= O(n^{-1}) \\ &+ \int_{(s,t) \in [0,1]^2} \mathbb{1}_{[ns] \neq [nt]} \left( \frac{\sigma_n^2([ns], [nt])}{(z - \lambda_n([ns]))^2(z - \lambda_n([nt]))} - \frac{\sigma^2(s,t)}{(z - f(s))^2(z - f(t))} \right) \mathrm{d}s \mathrm{d}t. \end{aligned}$$

Since, for  $M_\sigma := \sup_{0 \leq x \neq y \leq 1} \sigma(x,y)^2$  and for any fixed  $z \notin \tilde{\mathcal{S}}$ , the function

$$\psi_z : (s, \lambda, \lambda') \in [0, M_\sigma + 1] \times \{x \in \mathbb{R}; \mathrm{dist}(x, \tilde{\mathcal{S}}) \leq \mathrm{dist}(z, \tilde{\mathcal{S}})/2\}^2 \mapsto \frac{s}{(z - \lambda)^2(z - \lambda')}$$

is  $C \mathrm{dist}(z, \tilde{\mathcal{S}})^{-4}$ -Lipschitz, for  $C$  a universal constant, by Hypothesis (b),

$$\delta_n(z) = O(n^{-1}) + \frac{O(\eta_n)}{\max\{n\varepsilon_n, 1\} \mathrm{dist}(z, \tilde{\mathcal{S}})^4}. \tag{3.25}$$

Finally, the expression of  $B(z)$  given in (3.17) implies,

$$B(z) \leq \frac{C}{\text{dist}(z, \tilde{\mathcal{S}})^3} \quad (3.26)$$

Collecting estimations (3.24), (3.25) and (3.26), we conclude.  $\square$

**Claim 3.** *There is a constant  $C$  such that for any  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ ,*

$$\mathbb{E}[|n\varepsilon_n^{-1}C_n(z)|^2] \leq \frac{C\varepsilon_n^4}{\text{dist}(z, \tilde{\mathcal{S}})^8}.$$

**Proof.** We start by writing for all  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$

$$\begin{aligned} \mathbb{E}[|n\varepsilon_n^{-1}C_n(z)|^2] &= \frac{\varepsilon_n^4}{n^3} \mathbb{E} \left[ \left| \sum_{i,j,k=1}^n \frac{x_{i,j} x_{j,k} x_{k,i}}{(z - \lambda_n(i))^2 (z - \lambda_n(j)) (z - \lambda_n(k))} \right|^2 \right] \\ &= \frac{\varepsilon_n^4}{n^3} \sum_{i,j,k,l,m,p=1}^n \frac{\mathbb{E}(x_{i,j} x_{j,k} x_{k,i} \overline{x_{l,m} x_{m,p} x_{p,l}})}{(z - \lambda_n(i))^2 (z - \lambda_n(j)) (z - \lambda_n(k)) (\bar{z} - \lambda_n(l))^2 (\bar{z} - \lambda_n(m)) (\bar{z} - \lambda_n(p))}. \end{aligned}$$

Generically, the set of "edges"  $\{(l, m), (m, p), (p, l)\}$  must be equal to the set  $\{(i, j), (j, k), (k, i)\}$  in order to get a non zero term. Therefore, the complexity of the previous sum is  $O(n^3)$ . Note that other non zero terms involving third or fourth moments are much less numerous. Hence,

$$\mathbb{E}[|n\varepsilon_n^{-1}C_n(z)|^2] \leq \frac{\varepsilon_n^4}{n^3} \times \frac{O(n^3)}{\text{dist}(z, \tilde{\mathcal{S}})^8} \leq \frac{C\varepsilon_n^4}{\text{dist}(z, \tilde{\mathcal{S}})^8}$$

$\square$

**Claim 4.** *There is a constant  $C$  such that for any  $z \in \mathbb{C} \setminus \mathbb{R}$ ,*

$$\mathbb{E}[|n\varepsilon_n^{-1}R_n^\varepsilon(z)|^2] \leq \frac{O(n^2\varepsilon_n^6)}{|\Im(z)|^2 \text{dist}(z, \tilde{\mathcal{S}})^8}.$$

**Proof.** Remind that,

$$R_n^\varepsilon(z) := \frac{\varepsilon_n^4}{n} \text{Tr} \frac{1}{z - D} X \frac{1}{z - D} X \frac{1}{z - D} X \frac{1}{z - D} X \frac{1}{z - D^\varepsilon}.$$

Hence,

$$\begin{aligned}
 \mathbb{E}[|n\varepsilon_n^{-1}R_n^\varepsilon(z)|^2] &\leq \varepsilon_n^6 \mathbb{E} \left[ \left| \text{Tr} \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D^\varepsilon} \right|^2 \right] \\
 &\leq \varepsilon_n^6 \mathbb{E} \left[ \text{Tr} \left| \left( \frac{1}{z-D} X \right)^4 \right|^2 \times \text{Tr} \left| \frac{1}{z-D^\varepsilon} \right|^2 \right] \\
 &\leq \varepsilon_n^6 \mathbb{E} \left[ \text{Tr} \left( \left( \frac{1}{z-D} X \right)^4 \left( \overline{\frac{1}{z-D} X} \right)^4 \right) \frac{n}{|\text{Im}(z)|^2} \right] \\
 &\leq \frac{n\varepsilon_n^6}{|\text{Im}(z)|^2} \mathbb{E} \left[ \text{Tr} \left( \left( \frac{1}{z-D} X \right)^4 \left( \overline{\frac{1}{z-D} X} \right)^4 \right) \right] \\
 &\leq \frac{n\varepsilon_n^6}{|\text{Im}(z)|^2} \frac{O(n^5)}{n^4 \text{dist}(z, \tilde{\mathcal{S}})^8} \leq \frac{O(n^2\varepsilon_n^6)}{|\text{Im}(z)|^2 \text{dist}(z, \tilde{\mathcal{S}})^8}.
 \end{aligned}$$

The inequality of the last line takes into account that

- the  $L^8$  norm of the entries of  $\sqrt{n}X$  is uniformly bounded
- the norm of the entries of  $X$  is of order  $n^{-1/2}$
- the norm of the coefficients of  $(z-D)^{-1}$  is smaller than  $\text{dist}(z, \tilde{\mathcal{S}})^{-1}$
- the complexity of the sum defining the trace is of order  $O(n^5)$  since its non-null terms are encoded by four edges trees which have therefore five vertices.

□

We gather now the results of the previous claims.

For any rate of convergence of  $\varepsilon_n$ , Claim 1 proves that the process  $n\varepsilon_n^{-1}A_n(z)$  converges in distribution to the centered Gaussian variable  $Z(z)$ . Moreover,

- if  $\varepsilon_n \ll n^{-1}$ , then as Claims 2, 3 and 4 imply that the processes  $n\varepsilon_n^{-1}B_n(z)$ ,  $n\varepsilon_n^{-1}C_n(z)$  and  $n\varepsilon_n^{-1}R_n^\varepsilon(z)$  converge to 0 in probability, we can conclude, by Slutsky's theorem, that for any  $z \in \mathbb{C} \setminus \mathbb{R}$ :

$$n\varepsilon_n^{-1}\Delta G_n(z) \xrightarrow[n \rightarrow \infty]{\text{dist}} Z(z)$$

- if  $\varepsilon_n \sim \frac{c}{n}$ , then, as Claims 2, 3 and 4 imply that the processes  $n\varepsilon_n^{-1}B_n(z)$ ,  $n\varepsilon_n^{-1}C_n(z)$  and  $n\varepsilon_n^{-1}R_n^\varepsilon(z)$  converge respectively to  $cB(z)$ , 0 and 0 in probability, we can conclude, by Slutsky's theorem, that for any  $z \in \mathbb{C} \setminus \mathbb{R}$ :

$$n\varepsilon_n^{-1}\Delta G_n(z) \xrightarrow[n \rightarrow \infty]{\text{dist}} Z(z) + cB(z)$$

- if  $n^{-1} \ll \varepsilon_n \ll n^{-1/3}$ , then, as Claims 2, 3 and 4 imply that the three processes  $n\varepsilon_n^{-1}(B_n(z) - \varepsilon_n^2 B(z))$ ,  $n\varepsilon_n^{-1}C_n(z)$  and  $n\varepsilon_n^{-1}R_n^\varepsilon(z)$  converge to 0 in probability, we can conclude, by Slutsky's theorem, that for any  $z \in \mathbb{C} \setminus \mathbb{R}$ :

$$n\varepsilon_n^{-1} (\Delta G_n(z) - \varepsilon_n^2 B(z)) \xrightarrow[n \rightarrow \infty]{\text{dist.}} Z(z)$$

Regarding the convergence in probability (3.22), in the case  $n^{-1} \ll \varepsilon_n \ll 1$ , Claims 1, 2, 3 and 4 imply that the processes  $\varepsilon_n^{-2}A_n(z)$ ,  $\varepsilon_n^{-2}B_n(z) - B(z)$ ,  $\varepsilon_n^{-2}C_n(z)$  and  $\varepsilon_n^{-2}R_n^\varepsilon(z)$  converge to 0.

This finishes the proof of the convergences of Proposition 21.  $\square$

### 3.6.3 Proof of Proposition 22

Recall that

$$B(z) = \int_{(s,t) \in [0,1]^2} \frac{\sigma^2(s,t)}{(z - f(s))^2(z - f(t))} ds dt.$$

Recall that  $\rho$  is the density of the push-forward of the uniform measure on  $[0,1]$  by the map  $f$ .

Let  $\tau$  be as in Hypothesis (d). We have

$$B(z) = \int_{\mathbb{R}^2} \frac{\tau(s,t) \rho(s) \rho(t)}{(z - s)^2 (z - t)} ds dt.$$

By a partial fraction decomposition we have for all  $a \neq b$

$$\frac{1}{(z - a)^2(z - b)} = \frac{1}{(b - a)^2} \left( \frac{1}{z - b} - \frac{1}{z - a} - \frac{b - a}{(z - a)^2} \right).$$

Thus, as the Lebesgue measure of the set  $\{(y_1, y_2) \in [0,1]^2 ; y_1 = y_2\}$  is null, we have

$$B(z) = \int_{\mathbb{R}^2} \frac{\tau(s,t) \rho(s) \rho(t)}{(t - s)^2} \left( \frac{1}{z - t} - \frac{1}{z - s} - \frac{t - s}{(z - s)^2} \right) ds dt.$$

Moreover, for  $\varphi_z$  the function  $\varphi_z : x \mapsto \frac{1}{z - x}$ , we obtain

$$B(z) = \int_{\mathbb{R}^2} \frac{\tau(s,t) \rho(s) \rho(t)}{(t - s)^2} (\varphi_z(t) - \varphi_z(s) - (t - s)\varphi'_z(s)) ds dt.$$

Now, we want to prove that  $B(z) = - \int_{\mathbb{R}^2} \frac{\tau(s,t) \rho(s) \rho(t)}{t - s} \varphi'_z(s) ds dt$ .

To do this, we will use a symmetry argument: in fact both terms in  $\varphi_z(t)$  and  $\varphi_z(s)$  neutralize each other, and it remains only to prove, that we did not remove  $\infty$  to  $\infty$  and that the remaining term has the desired form.

Let us define

$$B^\eta(z) := \int_{|s-t|>\eta} \frac{\tau(s,t) \rho(s) \rho(t)}{(t-s)^2} (\varphi_z(t) - \varphi_z(s) - (t-s)\varphi'_z(s)) ds dt.$$

By the Taylor-Lagrange inequality we obtain:

$$\left| \frac{\tau(s,t) \rho(s) \rho(t)}{(t-s)^2} (\varphi_z(t) - \varphi_z(s) - (t-s)\varphi'_z(s)) \right| \leq \frac{\rho(s) \rho(t) \|\tau(\cdot, \cdot)\|_{L^\infty} \|\varphi''_z\|_{L^\infty}}{2}.$$

So that, since  $\rho$  is a density, by dominated convergence, we have

$$\lim_{\eta \rightarrow 0} B^\eta(z) = B(z).$$

Moreover, by symmetry, for any  $\eta$ ,

$$B^\eta(z) = \int_{|s-t|>\eta} \frac{\tau(s,t) \rho(s) \rho(t)}{t-s} (-\varphi'_z(s)) ds dt.$$

So

$$\begin{aligned} B(z) &= \lim_{\eta \rightarrow 0} \int_{|s-t|>\eta} \frac{\tau(s,t) \rho(s) \rho(t)}{t-s} (-\varphi'_z(s)) dt ds \\ &= - \lim_{\eta \rightarrow 0} \int_{s \in \mathbb{R}} F_\eta(s) \varphi'_z(s) ds \end{aligned} \tag{3.27}$$

where for  $\eta > 0$  and  $s \in \mathbb{R}$ , we define

$$F_\eta(s) := \rho(s) \int_{t \in \mathbb{R} \setminus [s-\eta, s+\eta]} \frac{\tau(s,t) \rho(t)}{t-s} dt.$$

Note that that by definition of the function  $F$  given at (3.6), for any  $s$ , we have

$$F(s) = \lim_{\eta \rightarrow 0} F_\eta(s). \tag{3.28}$$

Thus by (3.27) and (3.28), to conclude the proof of Proposition 22, by dominated convergence, one needs only to state that  $F_\eta$  is dominated, uniformly in  $\eta$ , by an integrable function. This follows from the following computation.

Note first that by symmetry, we have

$$F_\eta(s) = \rho(s) \int_{t \in \mathbb{R} \setminus [s-\eta, s+\eta]} \frac{\tau(s,t) \rho(t) - \tau(s,s) \rho(s)}{t-s} dt. \tag{3.29}$$

Let  $M > 0$  such that the support of the function  $\rho$  is contained in  $[-M, M]$ . Then, for  $\eta_0, \alpha, C$  as in Hypothesis (e), using the expression of  $F_\eta(s)$  given at (3.29), we have

$$\begin{aligned} |F_\eta(s)| &\leq 2C\rho(s) \int_{t=s}^{s+\eta_0} |t-s|^{\alpha-1} dt + \int_{t \in [s-2M, s-\eta_0] \cup [s+\eta_0, s+2M]} \left| \frac{\tau(s, t)\rho(s)\rho(t)}{t-s} \right| dt \\ &\leq \frac{2C\rho(s)}{\alpha} \eta_0^\alpha + \frac{1}{\eta_0} \int_{t \in \mathbb{R}} |\tau(s, t)\rho(s)\rho(t)| dt \\ &\leq \frac{2C\rho(s)}{\alpha} \eta_0^\alpha + \frac{\|\tau(\cdot, \cdot)\|_{L^\infty}}{\eta_0} \rho(s). \end{aligned}$$

□

### 3.6.4 A local type convergence result

One can precise the convergence (3.22) by replacing the complex variable  $z$  by a complex sequence  $(z_n)$  which converges slowly enough to the real axis. This convergence won't be used in the sequel. As it is discussed in [BGK16], this type of result is a first step towards a local result for the empirical distribution.

**Proposition 23.** *Under Hypotheses (a), (b), (f), if  $n^{-1} \ll \varepsilon_n \ll 1$ , then for any nonreal complex sequence  $(z_n)$ , such that*

$$\Im(z_n) \gg \max \left\{ (n\varepsilon_n)^{-1/2}, \left( \frac{\eta_n}{n\varepsilon_n} \right)^{1/4}, \varepsilon_n^{2/5} \right\} \quad (3.30)$$

the following convergence holds

$$\varepsilon_n^{-2} \Delta G_n(z_n) - B(z_n) \xrightarrow[n \rightarrow \infty]{P} 0.$$

**Remark.** In the classical case where  $\frac{\eta_n}{n\varepsilon_n} = \sup_{i \neq j} (|\sigma_n^2(i, j) - \sigma^2(i/n, j/n)| + |\lambda_n(i) - f(i/n)|)$

is of order  $\frac{1}{n}$ , the above assumption boils down to  $\Im(z_n) \gg \max \left\{ (n\varepsilon_n)^{-1/2}, \varepsilon_n^{2/5} \right\}$ .

**Proof.** Assume  $n^{-1} \ll \varepsilon_n \ll 1$ . One can directly obtain, for all non-real complex sequences  $(z_n)$ , that

- by Claim 1, if  $\text{dist}(z_n, \tilde{\mathcal{S}}) \gg (n\varepsilon_n)^{-1/2}$ , then

$$\mathbb{E}[|\varepsilon_n^{-2} A_n(z_n)|^2] \leq \frac{C}{(n\varepsilon_n)^2 \text{dist}(z_n, \tilde{\mathcal{S}})^4} \xrightarrow[n \rightarrow \infty]{} 0,$$

- by Claim 2, if  $\text{dist}(z_n, \tilde{\mathcal{S}}) \gg \max \left\{ n^{-1/3}, (\eta_n/(n\varepsilon_n))^{1/4} \right\}$ , then

$$\mathbb{E}[|\varepsilon_n^{-2} B_n(z_n) - B(z_n)|^2] \leq \frac{C}{n^2 \text{dist}(z_n, \tilde{\mathcal{S}})^6} + \frac{C\eta_n^2}{(n\varepsilon_n)^2 \text{dist}(z_n, \tilde{\mathcal{S}})^8} \xrightarrow[n \rightarrow \infty]{} 0,$$

- by Claim 3, if  $\text{dist}(z_n, \tilde{\mathcal{S}}) \gg (\varepsilon_n/n)^{1/4}$ , then

$$\mathbb{E}[|\varepsilon_n^{-2} C_n(z_n)|^2] \leq \frac{C\varepsilon_n^2}{n^2 \text{dist}(z_n, \tilde{\mathcal{S}})^8} \xrightarrow{n \rightarrow \infty} 0,$$

- by Claim 4, if  $|\Im(z_n)| \text{dist}(z_n, \tilde{\mathcal{S}})^4 \gg \varepsilon_n^2$ , then

$$\mathbb{E}[|\varepsilon_n^{-2} R_n^\varepsilon(z_n)|^2] \leq \frac{O(\varepsilon_n^4)}{|\Im(z_n)|^2 \text{dist}(z_n, \tilde{\mathcal{S}})^8} \xrightarrow{n \rightarrow \infty} 0.$$

Therefore, when

$$\text{dist}(z_n, \tilde{\mathcal{S}}) \gg \max \left\{ (n\varepsilon_n)^{-1/2}, n^{-1/3}, \left( \frac{\eta_n}{n\varepsilon_n} \right)^{1/4}, \left( \frac{\varepsilon_n}{n} \right)^{1/4} \right\} \text{ and } |\Im(z_n)| \text{dist}(z_n, \tilde{\mathcal{S}})^4 \gg \varepsilon_n^2,$$

the four processes,  $\varepsilon_n^{-2} A_n(z_n)$ ,  $\varepsilon_n^{-2} B_n(z_n) - B(z_n)$ ,  $\varepsilon_n^{-2} C_n(z_n)$  and  $\varepsilon_n^{-2} R_n^\varepsilon(z_n)$  converge to 0 in probability. Since  $\text{dist}(z_n, \tilde{\mathcal{S}}) \geq \Im(z_n)$ , the above condition is implied by

$$\Im(z_n) \gg \max \left\{ (n\varepsilon_n)^{-1/2}, n^{-1/3}, \left( \frac{\eta_n}{n\varepsilon_n} \right)^{1/4}, \left( \frac{\varepsilon_n}{n} \right)^{1/4}, \varepsilon_n^{2/5} \right\}.$$

Observing finally that the two terms  $n^{-1/3}$  and  $(\frac{\varepsilon_n}{n})^{1/4}$  are dominated by the maximum of the three other ones, we conclude the proof.  $\square$

### 3.6.5 Possible extensions to larger $\varepsilon_n$

The convergence in distribution result of Theorem 19 is valid for  $\varepsilon_n \ll n^{-1/3}$  but fails above  $n^{-1/3}$ . Let us consider, for example, the case where  $n^{-1/3} \ll \varepsilon_n \ll n^{-1/5}$ . In this case, the contribution of the first term  $A_n(z)$  in the expansion of  $\Delta G_n(z)$  which yields the random limiting quantity, is dominated *not only* by the term  $B_n(z)$  as it used to be previously. It is also dominated by a further and smaller term  $D_n(z)$  of the expansion

$$\Delta G_n(z) = A_n(z) + B_n(z) + C_n(z) + D_n(z) + E_n(z) + R_n^\varepsilon,$$

with:

$$\begin{aligned} A_n(z) &:= \frac{\varepsilon_n}{n} \text{Tr} \frac{1}{z-D} X \frac{1}{z-D} \\ &\vdots \\ E_n(z) &:= \frac{\varepsilon_n^5}{n} \text{Tr} \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} \\ R_n^\varepsilon(z) &:= \frac{\varepsilon_n^6}{n} \text{Tr} \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} \end{aligned}$$

In this case, the random term  $Z(z)$  is still produced by  $A_n(z)$  and has an order of magnitude of  $\varepsilon_n/n$ . Meanwhile, the term  $D_n(z)$  writes

$$D_n(z) := \frac{\varepsilon_n^4}{n^3} \sum_{i,j,k,l=1}^n \frac{x_{i,j} x_{j,k} x_{k,l} x_{l,i}}{(z - \lambda_n(i))^2 (z - \lambda_n(j)) (z - \lambda_n(k)) (z - \lambda_n(l))}.$$

All the indices satisfying  $j = l$  contribute to the previous sum, since they produce a term in  $|x_{i,l}|^2 |x_{k,l}|^2$ . Their cardinality is of order  $n^3$ . Therefore, the term  $D_n(z)$  is of order  $\varepsilon_n^4$  which prevails on the order  $\varepsilon_n/n$  of  $A_n(z)$ , as soon as  $\varepsilon_n \gg n^{-1/3}$ . One can also observe that the odd terms  $C_n(z)$  and  $E_n(z)$  in the expansion are negligible with respect to  $A_n(z)$  due to the fact that the entries  $x_{i,j}$  are centered. One can then state an analogous result to Proposition 21, but the deterministic limiting term  $D(z)$  arising from  $D_n(z)$  does not find a nice expression as the image of  $\varphi_z$  by a linear form as it was the case for  $B(z)$  in Proposition 22. Therefore we did not state an extension of Theorem 19.

More generally, for all positive integer  $p$ , when  $n^{-1/(2p-1)} \ll \varepsilon_n \ll n^{-1/(2p+1)}$ , the expansion will contain  $p$  deterministic terms, produced by the even variables,  $B_n(z)$ ,  $D_n(z)$ ,  $F_n(z)$ ,  $H_n(z)$  ... All the other odd terms,  $C_n(z)$ ,  $E_n(z)$ ,  $G_n(z)$  ... being negligible due to the centering of the entries. The limits of the even terms  $B_n(z)$ ,  $D_n(z)$ ,  $F_n(z)$ ,  $H_n(z)$  ... can be expressed thanks to operator-valued free probability theory, using the results of [Shl96] (namely, Th. 4.1), but expressing these limits as the images of  $\varphi_z$  by linear forms is a quite involved combinatorial problem that we did not solve yet.

### 3.7 Convergence in probability in the semi-perturbative regime

Our goal now is to extend the convergence in probability result (3.22) of Proposition 21, proved for test functions  $\varphi_z(x) := \frac{1}{z-x}$ , to any  $\mathcal{C}^6$  and compactly supported function on  $\mathbb{R}$ . We do it in the following lemma by using the Helffer-Sjöstrand formula which is stated in Proposition 14 of Chapter 2.

**Lemma 24.** *If  $n^{-1} \ll \varepsilon_n \ll 1$ , then, for any compactly supported  $\mathcal{C}^6$  function  $\phi$  on  $\mathbb{R}$ ,*

$$\varepsilon_n^{-2}(\mu_n^\varepsilon - \mu_n)(\phi) \xrightarrow[n \rightarrow \infty]{P} - \int \phi'(s) F(s) \, ds.$$

**Proof.** Let us introduce the Banach space  $\mathcal{C}_{b,b}^1$  of bounded  $\mathcal{C}^1$  functions on  $\mathbb{R}$  with bounded derivative, endowed with the norm  $\|\phi\|_{\mathcal{C}_{b,b}^1} := \|\phi\|_\infty + \|\phi'\|_\infty$ .

On this space, let us define the random continuous linear form

$$\Pi_n(\phi) := \varepsilon_n^{-2}(\mu_n^\varepsilon - \mu_n)(\phi) + \int \phi'(s) F(s) \, ds.$$



Convergence (3.22) of Proposition 21 can now be formulated as

$$\forall z \in \mathbb{C} \setminus \mathbb{R}, \quad \Pi_n(\varphi_z) \xrightarrow[n \rightarrow \infty]{P} 0.$$

Actually, we can be more precise by adding the upper bounds of Claims 1, 2, 3 and 4, and obtain, uniformly in  $z$ ,

$$\begin{aligned} \mathbb{E}[|\Pi_n(\varphi_z)|^2] &= \mathbb{E}[|\varepsilon_n^{-2} \Delta G_n(z) - B(z)|^2] \\ &\leq \frac{(n\varepsilon_n)^{-2}}{\min \left( \text{dist}(z, \tilde{\mathcal{S}})^4, \text{dist}(z, \tilde{\mathcal{S}})^8, |\Im(z)|^2 \text{dist}(z, \tilde{\mathcal{S}})^8 \right)}. \end{aligned} \quad (3.31)$$

Now, let  $\phi$  be a compactly supported  $\mathcal{C}^6$  function on  $\mathbb{R}$  and let us introduce the almost analytic extension of degree 5 of  $\phi$  defined by

$$\forall z = x + iy \in \mathbb{C}, \quad \tilde{\phi}_5(z) := \sum_{k=0}^5 \frac{1}{k!} (iy)^k \phi^{(k)}(x).$$

An elementary computation gives, by successive cancellations, that

$$\bar{\partial} \tilde{\phi}_5(z) = \frac{1}{2} (\partial_x + i\partial_y) \tilde{\phi}_5(x + iy) = \frac{1}{2 \times 5!} (iy)^5 \phi^{(6)}(x). \quad (3.32)$$

Furthermore, by Helffer-Sjöstrand formula (Proposition 14), for  $\chi \in \mathcal{C}_c^\infty(\mathbb{C}; [0, 1])$  a smooth cutoff function with value one on the support of  $\phi$ ,

$$\phi(\cdot) = -\frac{1}{\pi} \int_{\mathbb{C}} \frac{\bar{\partial}(\tilde{\phi}_5(z)\chi(z))}{y^5} y^5 \varphi_z(\cdot) d^2 z$$

where  $d^2 z$  denotes the Lebesgue measure on  $\mathbb{C}$ .

Note that by (3.32),  $z \mapsto \mathbb{1}_{y \neq 0} \frac{\bar{\partial}(\tilde{\phi}_5(z)\chi(z))}{y^5}$  is a continuous compactly supported function and that  $z \in \mathbb{C} \mapsto \mathbb{1}_{y \neq 0} y^5 \varphi_z \in \mathcal{C}_{b,b}^1$  is continuous, hence,

$$\Pi_n(\phi) = \frac{1}{\pi} \int_{\mathbb{C}} \frac{\bar{\partial}(\tilde{\phi}_5(z)\chi(z))}{y^5} y^5 \Pi_n(\varphi_z) d^2 z.$$

Therefore,

$$\begin{aligned} \mathbb{E}(|\Pi_n(\phi)|^2) &= \mathbb{E} \left( \left| \frac{1}{\pi} \int_{\mathbb{C}} \frac{\bar{\partial}(\tilde{\phi}_5(z)\chi(z))}{y^5} y^5 \Pi_n(\varphi_z) d^2 z \right|^2 \right) \\ &\leq \mathbb{E} \left( \frac{1}{\pi^2} \int_{\mathbb{C}} \left| \frac{\bar{\partial}(\tilde{\phi}_5(z)\chi(z))}{y^5} y^5 \Pi_n(\varphi_z) \right|^2 d^2 z \right) \\ &= \frac{1}{\pi^2} \int_{\mathbb{C}} \left| \frac{\bar{\partial}(\tilde{\phi}_5(z)\chi(z))}{y^5} \right|^2 y^{10} \mathbb{E}(|\Pi_n(\varphi_z)|^2) d^2 z. \end{aligned}$$

Since the function  $\left| \frac{\bar{\partial}(\tilde{\phi}_5(z)\chi(z))}{y^5} \right|^2$  is continuous and compactly supported and that, by (3.31), for  $n^{-1} \ll \varepsilon_n \ll 1$ , uniformly in  $z$ ,

$$y^{10} \mathbb{E} (|\Pi_n(\varphi_z)|^2) \leq y^{10} \frac{o(1)}{\min(y^4, y^{10})} \xrightarrow{n \rightarrow \infty} 0.$$

Thus, for any compactly supported  $\mathcal{C}^6$  function on  $\mathbb{R}$ ,

$$\mathbb{E} (|\Pi_n(\phi)|^2) \leq \frac{1}{\pi^2} \int_{\mathbb{C}} \left| \frac{\bar{\partial}(\tilde{\phi}_5(z)\chi(z))}{y^5} \right|^2 y^{10} \mathbb{E} (|\Pi_n(\varphi_z)|^2) d^2z \xrightarrow{n \rightarrow \infty} 0$$

which implies that  $\Pi_n(\phi)$  converges to 0 in probability.  $\square$

### 3.8 Convergence in distribution towards the Gaussian variable $Z_\phi$

The purpose of this section is to extend the convergences in distribution of Proposition 21, from test functions of the type  $\varphi_z := \frac{1}{z-x}$ , to compactly supported  $\mathcal{C}^6$  functions on  $\mathbb{R}$ . To do so, we will use an extension lemma of Shcherbina and Tirozzi, stated as Lemma 15 in Chapter 2, which concerns the convergence of a sequence of *centered* random fields with uniformly bounded variance. Hence, we need to show first that our non centered random sequence is not far from being centered, which is done in subsection 3.8.1 by using again the Helffer-Sjöstrand formula (14). In subsection 3.8.2, we dominate the variance of this centered random field thanks to another result of Shcherbina and Tirozzi stated in Proposition 16 of Chapter 2. Subsection 3.8.3 collects the preceding results to conclude the proof.

#### 3.8.1 Coincidence of the expectation of $\mu_n^\varepsilon$ with its deterministic approximation

The asymptotic coincidence of the expectation of  $\mu_n^\varepsilon$  with its deterministic approximation is the content of next lemma:

**Lemma 25.** *Let us define, for  $\phi$  a  $\mathcal{C}^1$  function on  $\mathbb{R}$ ,*

$$\Lambda_n(\phi) := \begin{cases} n\varepsilon_n^{-1} (\mathbb{E}[\mu_n^\varepsilon(\phi)] - \mu_n(\phi)) & \text{if } \varepsilon_n \ll n^{-1}, \\ n\varepsilon_n^{-1} (\mathbb{E}[\mu_n^\varepsilon(\phi)] - \mu_n(\phi) + \varepsilon_n^2 \int \phi'(s)F(s)ds) & \text{if } \varepsilon_n \sim c/n \text{ or } n^{-1} \ll \varepsilon_n \ll n^{-1/3}. \end{cases}$$

*Then, as  $n \rightarrow \infty$ , for any compactly supported  $\mathcal{C}^6$  function  $\phi$  or any  $\phi$  of the type  $\varphi_z(x) = \frac{1}{z-x}$ ,  $z \in \mathbb{C} \setminus \mathbb{R}$ , we have*

$$\Lambda_n(\phi) \xrightarrow{n \rightarrow \infty} 0.$$

**Proof.** First note that, as the variables  $x_{i,j}$  are centered,  $\mathbb{E}[A_n(z)] = 0$ . Moreover, by adding the renormalized upper bounds of Claims 2, 3 and 4 one can directly obtain the two following inequalities for any  $z \in \mathbb{C} \setminus \mathbb{R}$ :

- If  $\varepsilon_n \ll n^{-1}$ , then

$$\begin{aligned}
 |\Lambda_n(\varphi_z)| &= n\varepsilon_n^{-1} |\mathbb{E}[\Delta G_n(z)]| \\
 &\leq n\varepsilon_n^{-1} (|\mathbb{E}[A_n(z)]| + \mathbb{E}[|B_n(z)|] + \mathbb{E}[|C_n(z)|] + \mathbb{E}[|R_n^\varepsilon(z)|]) \\
 &\leq \frac{C(n\varepsilon_n + \eta_n)}{\min \left\{ \text{dist}(z, \tilde{\mathcal{S}})^3, \text{dist}(z, \tilde{\mathcal{S}})^4, |\Im(z)| \text{dist}(z, \tilde{\mathcal{S}})^4 \right\}} \xrightarrow{n \rightarrow \infty} 0.
 \end{aligned}$$

- If  $\varepsilon_n \sim c/n$  or  $n^{-1} \ll \varepsilon_n \ll n^{-1/3}$ , then

$$\begin{aligned}
 |\Lambda_n(\varphi_z)| &= n\varepsilon_n^{-1} |\mathbb{E}[\Delta G_n(z) - \varepsilon_n^2 B(z)]| \\
 &\leq n\varepsilon_n^{-1} (|\mathbb{E}[A_n(z)]| + \mathbb{E}[|B_n(z) - \varepsilon_n^2 B(z)|] + \mathbb{E}[|C_n(z)|] + \mathbb{E}[|R_n^\varepsilon(z)|]) \\
 &\leq \frac{C(\varepsilon_n + \eta_n + n\varepsilon_n^3)}{\min \left\{ \text{dist}(z, \tilde{\mathcal{S}})^3, \text{dist}(z, \tilde{\mathcal{S}})^4, |\Im(z)| \text{dist}(z, \tilde{\mathcal{S}})^4 \right\}} \xrightarrow{n \rightarrow \infty} 0.
 \end{aligned}$$

Hence, in all cases,  $\Lambda_n(\varphi_z) \xrightarrow{n \rightarrow \infty} 0$ .

The extension of this result to compactly supported  $\mathcal{C}^6$  test functions on  $\mathbb{R}$  goes the same way as for  $\Pi_n$  in the proof of Lemma 24.  $\square$

### 3.8.2 Domination of the variance of $\mu_n^\varepsilon$

The second ingredient goes through a domination of the variance of  $\mu_n^\varepsilon(\phi)$ :

**Lemma 26.** *Let  $s > 5$ . There is a constant  $C$  such that for each  $n$  and each  $\phi \in \mathcal{H}_s$ ,*

$$\text{Var}(n\varepsilon_n^{-1} \mu_n^\varepsilon(\phi)) \leq C \|\phi\|_{\mathcal{H}_s}^2.$$

**Proof.** By Proposition 16, it suffices to prove that

$$\int_{y=0}^{\infty} y^{2s-1} e^{-y} \int_{x \in \mathbb{R}} \text{Var}(\varepsilon_n^{-1} \text{Tr}((x + iy - D_n^\varepsilon)^{-1})) dx dy$$

are bounded independently of  $n$ .

Note that for  $\Delta G_n(z)$  defined in (3.18),

$$\text{Var}(\varepsilon_n^{-1} \text{Tr}((z - D_n^\varepsilon)^{-1})) = n^2 \varepsilon_n^{-2} \text{Var}(\Delta G_n(z)).$$

Moreover, the sum of the inequalities of Claims 1, 2, 3 and 4 yields

$$\text{Var}(n\varepsilon_n^{-1} \Delta G_n(z)) \leq \frac{C}{\text{dist}(z, \tilde{\mathcal{S}})^4} + \frac{C}{|\Im(z)|^2 \text{dist}(z, \tilde{\mathcal{S}})^8}.$$

Let  $M > 0$  such that  $\tilde{\mathcal{S}} \subset [-M, M]$ . Then

$$\text{dist}(z, \tilde{\mathcal{S}}) \geq \begin{cases} y & \text{if } |x| \leq M, \\ \sqrt{y^2 + (|x| - M)^2} & \text{if } |x| > M. \end{cases}$$

Thus  $\text{dist}(z, \tilde{\mathcal{S}}) \geq y$  if  $|x| \leq M$  and, for  $|x| > M$ ,

$$\frac{1}{\text{dist}(z, \tilde{\mathcal{S}})} \leq \frac{y^{-1}}{\sqrt{1 + ((|x| - M)/y)^2}}$$

and for any  $y > 0$ ,

$$\begin{aligned} \int_{x \in \mathbb{R}} \text{Var}(n\varepsilon_n^{-1} \Delta G_n(x + iy)) dx &\leq 2CM(y^{-10} + y^{-4}) + 2C \int_0^{+\infty} \frac{y^{-4}}{(1 + (\frac{x}{y})^2)^2} + \frac{y^{-10}}{(1 + (\frac{x}{y})^2)^4} dx \\ &\leq 2CM(y^{-10} + y^{-4}) + C \left( \frac{\pi}{2} y^{-3} + \frac{5\pi}{16} y^{-9} \right) \\ &\leq k (y^{-10} + y^{-3}), \end{aligned}$$

for a suitable constant  $k$ .

We deduce that, as soon as  $2s - 10 > 0$ , i.e.  $s > 5$ ,

$$\int_{y=0}^{\infty} y^{2s-1} e^{-y} \int_{x \in \mathbb{R}} \text{Var}(\varepsilon_n^{-1} \text{Tr}((x + iy - D_n^\varepsilon)^{-1})) dx dy \leq k \int_0^{\infty} y^{2s-1} e^{-y} (y^{-10} + y^{-3}) dy < \infty.$$

□

### 3.8.3 Proof of the convergences in distribution of Theorem 19

Since we have proved in Lemma 25 that for all compactly supported  $\mathcal{C}^6$  function  $\phi$ , the deterministic term  $\mu_n(\phi)$  could be replaced by  $\mathbb{E}[\mu_n^\varepsilon(\phi)]$ , we only have to prove, that for all  $\phi \in \mathcal{C}^6$ ,

$$n\varepsilon_n^{-1}(\mu_n^\varepsilon(\phi) - \mathbb{E}[\mu_n^\varepsilon(\phi)]) \xrightarrow[n \rightarrow \infty]{\text{dist.}} Z_\phi.$$

For the time being, we know this result to be valid for functions  $\phi$  belonging to the space  $\mathcal{L}_1$ , defined as the linear span of the family of functions  $\varphi_z(x) := \frac{1}{z-x}$ ,  $z \in \mathbb{C} \setminus \mathbb{R}$ .

By applying Lemma 15 to the centered field  $\mu_n^\varepsilon - \mathbb{E}[\mu_n^\varepsilon]$ , we are going to extend the result from the space  $\mathcal{L}_1$  to the Sobolev space  $(\mathcal{H}_s, \|\cdot\|_{\mathcal{H}_s})$  with  $s \in (5, 6)$ . Note that, since  $s < 6$ , this latter space contains the space of  $\mathcal{C}^6$  compactly supported functions (see [Hör03, Sec. 7.9]).

It remains to check the two hypotheses of Lemma 15. First, the subspace  $\mathcal{L}_1$  is dense in every space  $(\mathcal{H}_s, \|\cdot\|_{\mathcal{H}_s})$ . This is the content of Lemma 18 of Chapter 2. Second, by Lemma 26, since  $s > 5$ ,  $\text{Var}(n\varepsilon_n^{-1}\mu_n^\varepsilon(\phi)) \leq C\|\phi\|_{\mathcal{H}_s}^2$  for a certain constant  $C$ .

This concludes the proof.

# Chapter 4

## Expansions of the empirical spectral measure of a perturbed matrix

### Contents

---

<b>4.1</b>	<b>Introduction . . . . .</b>	<b>66</b>
<b>4.2</b>	<b>Main results . . . . .</b>	<b>67</b>
4.2.1	Definition of the model and assumptions . . . . .	67
4.2.2	Convergence in distribution of $\mu_n^\varepsilon$ . . . . .	69
4.2.3	A local type convergence result . . . . .	70
<b>4.3</b>	<b>Examples and simulations . . . . .</b>	<b>71</b>
4.3.1	Explanation of the first three deterministic terms of the expansion of $\mu_n^\varepsilon$ . . . . .	71
4.3.2	Numerical simulations . . . . .	73
<b>4.4</b>	<b>Strategy of the proof . . . . .</b>	<b>74</b>
<b>4.5</b>	<b>Proofs . . . . .</b>	<b>75</b>
4.5.1	Statements . . . . .	75
4.5.2	How the deterministic terms $C_k(z)$ appear . . . . .	76
4.5.3	Asymptotic behavior of the terms resulting from the expansion of the resolvent of $D_n^\varepsilon$ . . . . .	78
4.5.4	Proof of Theorem 28 . . . . .	83
4.5.5	Proof of the local type convergence result . . . . .	84
<b>4.6</b>	<b>A possible extension to other classes of test functions . . . . .</b>	<b>85</b>

---

## 4.1 Introduction

In the present chapter, we study this problem in the special case of Hermitian matrices. Namely, we perturb an  $n \times n$  Hermitian matrix,  $D_n$ , by a random Hermitian matrix with small operator norm. Moreover, as up to a change of basis, we can diagonalize  $D_n$ , we suppose it diagonal and consider that the entries of the perturbative matrix,  $X_n$ , are independent in the eigenvector basis of  $D_n$  with a variance profile<sup>1</sup>. Furthermore, to better describe the magnitude of the perturbations, we suppose that the operator norm of  $X_n$  is of order 1 and introduce a real sequence  $\varepsilon_n$  that tends to zero, so as to interest us in the empirical spectral distribution,  $\mu_n^\varepsilon$ , of the perturbed matrix

$$D_n^\varepsilon := D_n + \varepsilon_n X_n$$

for different rates of convergence of  $\varepsilon_n$  to zero. In the following we prove that, depending on the order of magnitude of the perturbation, an infinity of different regimes appear.

Previously, in [BGEM17] (Chapter 3), we have explicated the three first regimes, respectively called *perturbative*, *critical* and *semi-perturbative regimes*, and appear under the three following expressions:

$$\begin{aligned} \mu_n^\varepsilon &\approx \mu_n + \frac{\varepsilon_n}{n} dZ && \text{if } \varepsilon_n \ll n^{-1} \\ \mu_n^\varepsilon &\approx \mu_n + \frac{\varepsilon_n}{n} (cdC_2 + dZ) && \text{if } \varepsilon_n \sim \frac{c}{n} \\ \mu_n^\varepsilon &\approx \mu_n + \varepsilon_n^2 dC_2 && \text{if } n^{-1} \ll \varepsilon_n \ll 1 \end{aligned}$$

where,  $\mu_n$  is the empirical spectral measure of  $D_n$ ,  $dC_2$  a deterministic linear form depending of the limit parameters of the off-diagonal entries of  $X_n$  and  $dZ$  a random linear form on the space of  $\mathcal{C}^1$  functions on  $\mathbb{R}$  depending only on the limit parameters of the diagonal entries of  $X_n$ . This transition at  $\varepsilon_n \sim n^{-1}$  is the well-known transition, in quantum mechanics, where the *perturbative regime* ends. A surprising fact discovered during this first study was that the semi-perturbative regime  $n^{-1} \ll \varepsilon_n \ll 1$ , which concerns many applications in the context of covariance matrices in finance (see, for example, [AB12] or [ABB14]), decomposes into infinitely many sub-regimes. In the case  $n^{-1} \ll \varepsilon_n \ll n^{-1/3}$ , the expansion of  $\mu_n^\varepsilon - \mu_n$  contains a single deterministic term before the random term  $\frac{\varepsilon_n}{n} dZ$ . In the case  $n^{-1/3} \ll \varepsilon_n \ll n^{-1/5}$ , the expansion of  $\mu_n^\varepsilon - \mu_n$  contains two of them. And more generally, for all positive integer  $p$ , when  $n^{-1/(2p-1)} \ll \varepsilon_n \ll n^{-1/(2p+1)}$ , the expansion contains  $p$  of them:

$$\mu_n^\varepsilon \approx \mu_n + \frac{\varepsilon_n}{n} dZ + \varepsilon_n^2 dC_2 + \varepsilon_n^4 dC_4 + \cdots + \varepsilon_n^{2p} dC_{2p}.$$

In this case, each linear form  $dC_{2k}$  may concern up to the  $k$  first derivatives of the test function  $\phi$  on which we evaluate the measure  $\mu_n^\varepsilon$ .

---

<sup>1</sup>Note that if the perturbing matrix belongs to the GOE or GUE, then its law is invariant under this change of basis

Thereby, the aim of this chapter is, now, to provide a thorough study of those sub-regimes of the semi-perturbative regime, explaining how we should reinforce the hypothesis of our model each time we choose a slower rate of convergence to zero for  $\varepsilon_n$  and how the deterministic terms appear then.

The perturbative expansion of the empirical spectral measure is, in this paper, done for real test-functions of the type  $\varphi_z(x) = \frac{1}{z-x}$ , for a non-real complex number  $z$ . This class of test functions, which plays a central role in random matrix theory, is very useful because it can be used to extend easily a convergence in law to larger classes of test functions, as it is discussed in Section 4.6.

In this case, the deterministic terms  $(C_k)$ , which come from the linear forms  $(dC_k)$ , of this expansion are related to some combinatorial objects as, for example, rooted planar trees and the random linear form  $dZ$  is related to the so-called one-dimensional Gaussian free field defined, for instance at [Dub09, Sect. 4.2] (if the variance profile of  $X_n$  is flat, then it is precisely the Laplacian of the Gaussian free field, defined in the sense of distributions).

This chapter is organized as follows; the main results, comments and examples are given in Sections 4.2 and 4.3. Sections 4.4 and 4.5 are devoted to the proofs. And finally, Section 4.6 is a discussion about possible extensions of the main results to larger classes of test functions.

**Notations.** For  $a_n, b_n$  some real sequences,  $a_n \ll b_n$  (resp.  $a_n \sim b_n$ ) means that  $a_n/b_n$  tends to 0 (resp. to 1). Also,  $\xrightarrow{P}$  and  $\xrightarrow{\text{dist.}}$  stand respectively for convergence in probability and convergence in distribution for all finite marginals.

## 4.2 Main results

### 4.2.1 Definition of the model and assumptions

For all positive integer  $n$ , we consider a real diagonal matrix  $D_n = \text{diag}(\lambda_n(1), \dots, \lambda_n(n))$ , as well as a Hermitian random matrix

$$X_n = \frac{1}{\sqrt{n}} [x_{i,j}^n]_{1 \leq i,j \leq n}$$

and a positive number  $\varepsilon_n$ . The normalizing factor  $n^{-1/2}$  and our hypotheses below ensure that the operator norm of  $X_n$  is of order one. We then define, for all  $n$ ,

$$D_n^\varepsilon := D_n + \varepsilon_n X_n.$$

We now introduce the probability measures  $\mu_n$  and  $\mu_n^\varepsilon$  as the respective uniform distributions on the eigenvalues (with multiplicity) of  $D_n$  and  $D_n^\varepsilon$ . Our aim is to explicit all the different perturbative expansions of  $\mu_n^\varepsilon$  around  $\mu_n$  when  $n^{-1} \ll \varepsilon_n \ll 1$ .

We make the following general hypotheses:



- (a) the entries  $x_{i,j}^n$  of  $\sqrt{n}X_n$  are independent (up to symmetry) random variables, centered, with variance denoted by  $\sigma_n^2(i, j)$ ,
- (b) there are  $f, \sigma_d, \sigma$  real functions defined respectively on  $[0, 1]$ ,  $[0, 1]$  and  $[0, 1]^2$  such that, for each  $x \in [0, 1]$ ,

$$\lambda_n(\lfloor nx \rfloor) \xrightarrow[n \rightarrow \infty]{} f(x) \quad \text{and} \quad \sigma_n^2(\lfloor nx \rfloor, \lfloor nx \rfloor) \xrightarrow[n \rightarrow \infty]{} \sigma_d(x)^2$$

and for each  $x \neq y \in [0, 1]$ ,

$$\sigma_n^2(\lfloor nx \rfloor, \lfloor ny \rfloor) \xrightarrow[n \rightarrow \infty]{} \sigma^2(x, y).$$

We make the following hypothesis about the rate of convergence:

$$\eta_n := \max\{n\varepsilon_n, 1\} \times \sup_{1 \leq i \neq j \leq n} (|\sigma_n^2(i, j) - \sigma^2(i/n, j/n)| + |\lambda_n(i) - f(i/n)|) \xrightarrow[n \rightarrow \infty]{} 0.$$

Let us now make a last general assumption which makes it possible to include the case where the set of eigenvalues of  $D_n$  contains some outliers:

- (c) there is a real compact set  $\tilde{\mathcal{S}}$  such that

$$\max_{1 \leq i \leq n} \text{dist}(\lambda_n(i), \tilde{\mathcal{S}}) \xrightarrow[n \rightarrow \infty]{} 0.$$

Furthermore, we add a last particular hypothesis which depends on the rate of convergence of  $\varepsilon_n$ . In the case  $n^{-1/(2p-1)} \ll \varepsilon_n \ll n^{-1/(2p+1)}$  :

- (d) the moment  $\mathbb{E}|x_{i,j}^n|^{4p+4}$  is bounded uniformly on  $n, i, j$

Finally, we must introduce the last notion of rooted planar tree that is used in the two main results.

**Definition 27** (Dyck path). A *Dyck path* of length  $k \in 2\mathbb{N}$ , also called *non-negative Bernoulli walk* or *Bernoulli excursion*, is a function  $\gamma : \llbracket 0, k \rrbracket \mapsto \mathbb{N}$  satisfying

$$\gamma(0) = \gamma(k) = 0 \quad \text{and} \quad \forall i \in \llbracket 1, k-1 \rrbracket, \quad |\gamma(i+1) - \gamma(i)| = 1$$

**Remark 5.** The set of Dyck paths of length  $k \in 2\mathbb{N}$  is in bijection with several other sets of combinatorial objects, as the set of rooted planar trees with  $k/2$  edges, that is to say the set of trees which are embedded in the plane and are given one distinguished vertex called the root. One can find, for example, a proof of this well known result in the Subsection 1.1.2 of [Gui08]. Namely, the bijection consists in considering a clockwise path around the tree which begins and finishes at its root. This way, each time the walk around the tree meets a vertex for the first time then the Dyck path rises by one unit and each time the walk around the tree meets an already visited vertex the Dyck path descends from one unit, as it can be seen in Figure 4.1.




Figure 4.1: A Dyck path and its corresponding rooted planar tree (root is white)

**Notation.** In the following we formulate the main results in terms of rooted planar trees rather than in terms of Dyck paths. In order to be able to precisely designate the different vertices and edges of these rooted planar trees we labeled them in the following way: for a rooted planar tree with  $k + 1$  vertices, the root is labeled 1 and as the other vertices could be arbitrarily labeled from 2 to  $k + 1$  (only the structure of the tree matters), we choose the so-called *left-to-right depth-first labelling with source the root*, that is to say we label the vertices in the order that the clockwise walk around the tree beginning at the root, induced by the associated Dyck path, discover them. This arbitrary labeling could be observed in Figure 4.2.




Figure 4.2: Three different labeled planar rooted trees

Moreover, we denote  $\mathfrak{T}(k)$  the set of the rooted planar trees with  $k$  edges. Thereby, for any tree  $T \in \mathfrak{T}(k)$ , we denote by  $\mathcal{E}_T$  the set of its (un-oriented) edges and by  $v_\ell^T$  the degree of the vertex  $\ell$ .

#### 4.2.2 Convergence in distribution of $\mu_n^\varepsilon$

Let introduce, for  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ , a centered Gaussian process,  $(Z(z))_{z \in \mathbb{C} \setminus \tilde{\mathcal{S}}}$ , with distribution defined by its covariance structure

$$\mathbb{E}(Z(z)Z(z')) = \int_0^1 \frac{\sigma_d(t)^2}{(z - f(t))^2 (z' - f(t))^2} dt$$

and by the fact that  $\overline{Z(z)} = Z(\bar{z})$ .

Note that the field  $(Z(z))_{z \in \mathbb{C} \setminus \tilde{\mathcal{S}}}$  can be represented as

$$Z(z) = \int_0^1 \frac{\sigma_d(t)}{(z - f(t))^2} dB_t$$

where  $(B_t)$  is the standard one-dimensional Brownian motion.

We also introduce, for  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ ,

$$\Delta G_n(z) := (\mu_n^\varepsilon - \mu_n)(\varphi_z) = \frac{1}{n} \text{Tr} \frac{1}{z - D_n^\varepsilon} - \frac{1}{n} \text{Tr} \frac{1}{z - D_n}.$$

**Theorem 28.** *Let  $p \in \mathbb{N}^*$ . Under the semi-perturbative regime  $n^{-1/(2p-1)} \ll \varepsilon_n \ll n^{-1/(2p+1)}$  and under Hypotheses (a), (b), (c), (d), the following convergence in distribution holds,*

$$n\varepsilon_n^{-1} \left( (\mu_n^\varepsilon - \mu_n)(\varphi_z) - \sum_{\substack{k=2 \\ k \text{ even}}}^{2p} \varepsilon_n^k C_k(z) \right) \xrightarrow[n \rightarrow \infty]{\text{dist.}} Z(z)$$

where the terms  $(C_k(z))$  are deterministic functions of  $z$ , defined by

$$C_k(z) = \sum_{T \in \mathfrak{T}(\frac{k}{2})} \int_{[0,1]^{\frac{k}{2}+1}} \frac{\prod_{\{i,j\} \in \mathcal{E}_T} \sigma^2(x_i, x_j)}{(z - f(x_1)) \prod_{i=1}^{k/2+1} (z - f(x_i))^{v_i^T}} dx_1 \dots dx_{\frac{k}{2}+1} \quad (4.1)$$

**Remark 6.** The second part of Hypothesis (b), concerning the speed of convergence of the profile of the spectrum of  $D_n$  as well as of the variance of its perturbation, is needed in order to express the expansion of  $\mu_n^\varepsilon - \mu_n$  in terms of limit parameters of the model  $\sigma$  and  $\rho$ . We can remove this hypothesis and get analogous expansions where the terms  $C_k(z)$  and  $dZ$  are replaced by their discrete counterparts  $C_k^{(n)}$  and  $dZ_n$ , defined thanks to the "finite  $n$ " empirical versions of the limit parameters  $\sigma$  and  $\rho$ .

### 4.2.3 A local type convergence result

One can precise the convergence of Theorem 28 by replacing the complex variable  $z$  by a complex sequence  $(z_n)$  which converges slowly enough to the real axis. As it is discussed in [BGK16], this type of result is a first step towards a local result for the empirical distribution.

**Theorem 29.** *Under Hypotheses (a), (b), (c), (d), and if  $n^{-1} \ll \varepsilon_n \ll 1$ , then for any nonreal complex sequence  $(z_n)$ , such that*

$$\Im(z_n) \gg \max \left\{ (n\varepsilon_n^{p-1})^{-1/2}, \left( \frac{\eta_n}{n\varepsilon_n^{p-1}} \right)^{1/4}, \varepsilon_n^{\frac{2}{p+3}} \right\}.$$

and any even  $p \geq 2$ , the following convergence in probability holds

$$\varepsilon_n^{-p}(\mu_n^\varepsilon - \mu_n)(\varphi_{z_n}) - \sum_{\substack{k=0 \\ k \text{ even}}}^{p-2} \varepsilon_n^{-k} C_{p-k}(z_n) \xrightarrow[n \rightarrow \infty]{P} 0,$$

**Remark.** In the classical case where  $\frac{\eta_n}{n\varepsilon_n} = \sup_{i \neq j} (|\sigma_n^2(i, j) - \sigma^2(\frac{i}{n}, \frac{j}{n})| + |\lambda_n(i) - f(\frac{i}{n})|)$  is of order  $\frac{1}{n}$ , the above assumption boils down to  $\Im(z_n) \gg \max \left\{ (n\varepsilon_n^{p-1})^{-1/2}, \varepsilon_n^{\frac{2}{p+3}} \right\}$ .

## 4.3 Examples and simulations

To illustrate Theorem 28 we explicit, in Subsection 4.3.1, the terms of the expansion of  $\mu_n^\varepsilon$  until the semi-perturbative regime where  $\varepsilon_n \sim n^{-1/7}$  and give some simulations based on them in Subsection 4.3.2.

### 4.3.1 Explanation of the first three deterministic terms of the expansion of $\mu_n^\varepsilon$

In accordance with Theorem 28, the three first sub-regimes,  $n^{-1} \ll \varepsilon_n \ll n^{-1/3}$ ,  $n^{-1/3} \ll \varepsilon_n \ll n^{-1/5}$  and  $n^{-1/5} \ll \varepsilon_n \ll n^{-1/7}$ , of the semi-perturbative regime, involve the three first deterministic terms,  $C_2(z)$ ,  $C_4(z)$  and  $C_6(z)$ , that we explicit in this subsection in order to re-use them in the next subsection which is dedicated to simulations.


The term first deterministic term  $C_2(z)$  is linked to the only planar rooted tree with a single double-edge of the set  $\mathfrak{T}(1)$ ,



Thus, one can deduce directly from formula (4.1) that

$$C_2(z) := \int_{(x_1, x_2) \in [0, 1]^2} \frac{\sigma^2(x_1, x_2)}{(z - f(x_1))^2 (z - f(x_2))} dx_1 dx_2.$$


Similarly, as  $C_4(z)$  is linked to the set,  $\mathfrak{T}(2)$  containing the 2 planar rooted trees with 2 double-edges,



we deduce from formula (4.1) that,

$$C_4(z) = \int_{[0,1]^3} \frac{\sigma^2(x_1, x_2) \sigma^2(x_2, x_3)}{(z - x_1)^2 (z - x_2)^2 (z - x_3)} dx_1 dx_2 dx_3 \\ + \int_{[0,1]^3} \frac{\sigma^2(x_1, x_2) \sigma^2(x_1, x_3)}{(z - x_1)^3 (z - x_2) (z - x_3)} dx_1 dx_2 dx_3.$$

And finally, as  $C_6(z)$  is linked to the set  $\mathfrak{T}(3)$  containing the 5 planar rooted trees with 4 double-edges,



one can deduce from formula (4.1) and by symmetry between two of those trees that,

$$C_6(z) = \int_{[0,1]^4} \frac{\sigma^2(x_1, x_2) \sigma^2(x_2, x_3) \sigma^2(x_3, x_4)}{(z - f(x_1))^2 (z - f(x_2))^2 (z - f(x_3))^2 (z - f(x_4))} dx_1 dx_2 dx_3 dx_4 \\ + \int_{[0,1]^4} \frac{\sigma^2(x_1, x_2) \sigma^2(x_2, x_3) \sigma^2(x_2, x_4)}{(z - f(x_1))^2 (z - f(x_2))^3 (z - f(x_3)) (z - f(x_4))} dx_1 dx_2 dx_3 dx_4 \\ + 2 \int_{[0,1]^4} \frac{\sigma^2(x_1, x_2) \sigma^2(x_1, x_3) \sigma^2(x_3, x_4)}{(z - f(x_1))^3 (z - f(x_2)) (z - f(x_3))^2 (z - f(x_4))} dx_1 dx_2 dx_3 dx_4 \\ + \int_{[0,1]^4} \frac{\sigma^2(x_1, x_2) \sigma^2(x_1, x_3) \sigma^2(x_1, x_4)}{(z - f(x_1))^4 (z - f(x_2)) (z - f(x_3)) (z - f(x_4))} dx_1 dx_2 dx_3 dx_4$$

### 4.3.2 Numerical simulations

#### Parabolic pulse perturbation by a Wigner matrix

We consider the case where

$\rho(x) = \frac{3}{4}(1 - x^2)\mathbb{1}_{[-1,1]}(x)$ ,  $\sigma_d \equiv m$ , for some real constant  $m$ , and  $\sigma \equiv 1$  (this can be adapted to the case  $\sigma(x, y) = \mathbb{1}_{|y-x| \leq \ell}$ ). Theorem 28 is then illustrated by Figure 4.3, where we have considered, for visual reasons, the imaginary parts of the studied Stieltjes transforms,

$$\frac{1}{\pi} \Im(\mu_n^\varepsilon(\varphi_z) - \mu_n(\varphi_z)),$$

which boils down to replace, in Theorem 28, the function  $\varphi_z$  by the function

$$\phi(x) = \frac{1}{\pi} \frac{\eta}{(x - E)^2 + \eta^2} \quad (E \in \mathbb{R}, \eta > 0).$$




Figure 4.3: **Parabolic pulse perturbation by a Wigner matrix.** For different regimes of  $n^{-1/(2p-1)} \ll \varepsilon_n \ll n^{-1/(2p+1)}$  (from left to right and from top to bottom  $\varepsilon_n = n^{-\alpha}$  for  $\alpha = 0.5, 0.25, 0.17$  and  $0.143$ ): imaginary part of the term  $(\mu_n^\varepsilon - \mu_n)(\varphi_z) - \sum_{k=1}^{p-1} \varepsilon_n^k C_{2k}(z)$  (in blue) and imaginary part of the last deterministic term of the expansion  $C_{2p}(z)$  (in red) at  $z = E + i$  as a function of the real part  $E$ . In the four cases,  $n = 10^4$ .

As it can be seen on Figure 4.3, the predictions from Theorem 28 give a perfect matching with the numerical simulations for any  $n^{-1/7} \ll \varepsilon_n \ll 1$ .

### Uniform measure perturbation by different band matrices

The deterministic terms,  $C_k(z)$ , of the perturbative expansion we provide are functions of the non-diagonal entries of the perturbative matrix  $X_n$ . Thus if  $X_n$  is a band matrix then a natural question might be that of the precision of the theoretical model, stated in Theorem 28, as a function of the width,  $\ell$ , of the band. Indeed, the thinner the band, the fewer non-diagonal elements are taken into account in the terms  $(C_k(z))$ .

To study this question, we consider, here, the case where  $f(x) = x$ ,  $\sigma_d(x) \equiv m$  and  $\sigma(x, y) = \mathbb{1}_{|y-x| \leq \ell}$ , for some constants  $m \geq 0$  and  $\ell \in [0, 1]$ , which is the relative width of the band.

Theorem 28 is then illustrated by Figure 4.4 where we plotted once again the imaginary parts of the Stieltjes transforms, for different band matrices.

Figure 4.4 shows that the predictions from Theorem 28 give a perfect matching with the numerical simulations and, finally, that the width of the band does not influence significantly the accuracy of the result.

## 4.4 Strategy of the proof

The proof of Theorems 28 and 29 is based on an iteration of the resolvent formula, which states that for two hermitian matrices  $A$  and  $B$ , and for any  $z$  that is not in the spectrum of  $A$ ,  $B$  and  $A + B$  the following equality holds;

$$\frac{1}{z - (A + B)} = \frac{1}{z - A} B \frac{1}{z - (A + B)}.$$

This formula permits to subdivide  $(\mu_n^\varepsilon - \mu_n)(\varphi_z)$  into several terms that either converge in distribution to  $Z(z)$ , either converge in probability to the previously cited deterministic terms, or are negligible in probability compared to the other ones. This is the purpose of Subsection 4.5.3. Note that, the deterministic terms,  $C_k(z)$ , are to be related to planar rooted trees and Catalan numbers. The way that they are obtained is explicited in Subsection 4.5.2.

Once, those convergences are proved, then, thanks to Slutsky's theorem one can deduce both Theorems 28 and 29 as it is done respectively in Subsections 4.5.4 and 4.5.5.




Figure 4.4: **Uniform measure perturbation by a Gaussian band matrix.** Imaginary part of the Stieltjes transform of  $(\mu_n^\varepsilon - \mu_n)(\phi) - \varepsilon_n^2 C_2(z)$  (in blue) and of the imaginary part of the last deterministic term  $\varepsilon_n^4 C_4(z)$  (in red) at  $z = E + i$  as a function of the real part  $E$  for different widths of the band of the perturbative matrix  $X_n$ . Here,  $n = 10^4$ ,  $\varepsilon_n = n^{-\frac{1}{4}}$  and the band's width take the following values, from left to right and from top to bottom,  $\ell = 0.25$ ,  $\ell = 0.5$ ,  $\ell = 0.75$  and  $\ell = 1$ .

## 4.5 Proofs

### 4.5.1 Statements

Moreover, for  $p \in \mathbb{N}^*$ , let suppose that  $n^{-1/(2p-1)} \ll \varepsilon_n \ll n^{-1/(2p+1)}$ . One can deduce from the expansion of the resolvent of  $D_n^\varepsilon$  that,

$$\Delta G_n(z) = A_1^{(n)}(z) + A_2^{(n)}(z) + A_3^{(n)}(z) + \cdots + A_{2p+1}^{(n)}(z) + A_{2p+2}^\varepsilon(z), \quad (4.2)$$



for

$$\begin{aligned}
 A_1^{(n)}(z) &:= \frac{\varepsilon_n}{n} \operatorname{Tr} \frac{1}{z-D} X \frac{1}{z-D} = \frac{\varepsilon_n}{n} \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{x_{i,i}}{(z-\lambda_n(i))^2} \\
 A_2^{(n)}(z) &:= \frac{\varepsilon_n^2}{n} \operatorname{Tr} \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} = \frac{\varepsilon_n^2}{n^2} \sum_{i,j} \frac{|x_{i,j}|^2}{(z-\lambda_n(i))^2 (z-\lambda_n(j))} \\
 &\vdots \\
 A_k^{(n)}(z) &:= \frac{\varepsilon_n^k}{n} \operatorname{Tr} \underbrace{\frac{1}{z-D} X \frac{1}{z-D} X \dots \frac{1}{z-D} X}_{k \text{ times } \frac{1}{z-D} X} \frac{1}{z-D} \\
 &= \frac{\varepsilon_n^k}{n^{\frac{k+2}{2}}} \sum_{i_1, \dots, i_k=1}^n \frac{x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{k-1}, i_k} x_{i_k, i_1}}{(z-\lambda_n(i_1))^2 (z-\lambda_n(i_2)) \dots (z-\lambda_n(i_k))} \\
 &\vdots \\
 A_{2p+1}^{(n)}(z) &:= \frac{\varepsilon_n^{2p+1}}{n} \operatorname{Tr} \underbrace{\frac{1}{z-D} X \frac{1}{z-D} X \dots \frac{1}{z-D} X}_{(2p+1) \text{ times } \frac{1}{z-D} X} \frac{1}{z-D} \\
 A_{2p+2}^\varepsilon(z) &:= \frac{\varepsilon_n^{2p+2}}{n} \operatorname{Tr} \underbrace{\frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} X \frac{1}{z-D} X}_{(2p+2) \text{ times } \frac{1}{z-D} X} \frac{1}{z-D^\varepsilon}.
 \end{aligned}$$

**Notations.** To make notations lighter, we shall sometimes suppress the subscripts and superscripts  $n$ , so that  $D_n^\varepsilon$ ,  $D_n$ ,  $A_k^{(n)}$ ,  $X_n$  and  $x_{i,j}^n$  will be respectively denoted by  $D^\varepsilon$ ,  $D$ ,  $A_k$ ,  $X$  and  $x_{i,j}$ .

### 4.5.2 How the deterministic terms $C_k(z)$ appear

In order to lighten the proof of Theorems 28 and 29, the way the deterministic terms  $C_k(z)$  are obtained is explicated in the following lemma.

**Lemma 30.** *For any  $k \in \mathbb{N}$  and any  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ ,*

$$\lim_{n \rightarrow \infty} \varepsilon_n^{-k} \mathbb{E}[A_k(z)] = \begin{cases} 0 & \text{if } k \text{ is odd} \\ C_k(z) & \text{if } k \text{ is even} \end{cases}.$$

**Remark 7.** The limits of the even terms  $A_2(z)$ ,  $A_4(z)$ ,  $A_6(z)$ ,  $\dots$  could also be expressed thanks to operator-valued free probability theory, using the result [Shl96, Th. 4.1].

To prove Lemma 30, we shall provide first a lemma of graph theory which provides a link between the cardinality of the set of the vertices,  $V$ , and the cardinality of the set of the edges,  $E$ , of a graph  $G$ .

**Lemma 31.** *For any connected graph,  $G = (V, E)$ , the following inequality holds,*

$$|V| \leq |E| + 1.$$

*Moreover, the equality holds if and only if  $G$  is a tree.*

**Proof.** The proof is performed by induction over  $|V|$ .

If the graph is composed of only one vertex then the property is true.

Assume now that  $|V| \geq 1$ . Let's choose a vertex  $v$  of  $V$  and list  $e_1, \dots, e_l$  the  $l$  edges of  $G$  that contain  $v$ . One can split the graph  $G$  into  $m$  connected graphs  $G_1, \dots, G_m$  such that each of the edges  $e_1, \dots, e_l$  is contained in only one of those  $m$  graphs. Note that,  $m \leq l$ . By denoting, for  $i \in \{1, \dots, m\}$ ,  $G_i := (V_i, E_i)$ , by induction hypothesis,  $|V_i| \leq |E_i| + 1$  and thus

$$|V| = 1 + \sum_{i=1}^m |V_i| = 1 + \sum_{i=1}^m (|E_i| + 1) = 1 + m + \sum_{i=1}^m |E_i| = 1 + m + |E| - l \leq 1 + |E|$$

In addition, for the case of equality, note that if  $|V| = |E| + 1$  then we must have, for all  $i$ ,  $|V_i| = |E_i| + 1$  and  $m = l$ . But, if there is a loop in  $G$ , then one can find a vertex  $v$  such that  $m < l$ . So, by contradiction, the case of equality is verified only and only if  $G$  is a tree.  $\square$

**Proof.** [of Lemma 4.1] Note that for any  $k \in \mathbb{N}$  and any  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ ,

$$\varepsilon_n^{-k} \mathbb{E}[A_k(z)] = \frac{1}{n^{\frac{k}{2}+1}} \sum_{i_1, \dots, i_k=1}^n \frac{\mathbb{E}[x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{k-1}, i_k} x_{i_k, i_1}]}{(z - \lambda_n(i_1))^2 (z - \lambda_n(i_2)) \dots (z - \lambda_n(i_k))}$$

- If  $k$  is odd, as the variables  $(x_{ij})$  are centered, in order to get a non zero term in  $\mathbb{E}[x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{k-1}, i_k} x_{i_k, i_1}]$ , we need to consider all the couplings such that no first order moment appear. In other words, one should consider all the sets of edges  $\{(i_1, i_2), \dots, (i_{k-1}, i_k), (i_k, i_1)\}$  verifying that every edge is equal at least to an other one. Thus, as we need to consider graphs with at most  $\lfloor \frac{k}{2} \rfloor$  different edges, by Lemma 31, we know that those graphs have at most  $\lfloor \frac{k}{2} \rfloor + 1$  vertices  $i_1, \dots, i_{\lfloor \frac{k}{2} \rfloor + 1}$ .

Thus, since the indices  $i_1, i_2, \dots$  vary from 1 to  $n$ , there at most  $n^{\lfloor \frac{k}{2} \rfloor + 1} = n^{\frac{k+1}{2}}$  indices contributing in the previous sum and finally,

$$\varepsilon_n^{-k} \mathbb{E}[A_k(z)] = \frac{1}{n^{\frac{k}{2}+1}} \underbrace{\sum_{i_1, \dots, i_k=1}^n \frac{\mathbb{E}[x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{k-1}, i_k} x_{i_k, i_1}]}{(z - \lambda_n(i_1))^2 (z - \lambda_n(i_2)) \dots (z - \lambda_n(i_k))}}_{O\left(n^{\frac{k+1}{2}}\right)} \xrightarrow{n \rightarrow \infty} 0$$

- If  $k$  is even, as the variables  $(x_{ij})$  are centered, we need once again to consider all the sets of edges  $\{(i_1, i_2), \dots, (i_{k-1}, i_k), (i_k, i_1)\}$  that are, at least, equal two by two, in order

to get a non zero term in  $\mathbb{E} [x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{k-1}, i_k} x_{i_k, i_1}]$ . Namely, we have to consider all the graphs that have at most  $\frac{k}{2}$  edges which are at least double.

From Lemma 31 one can deduce that among all these graphs, those with a triple-edge or more have at most  $\frac{k}{2} - 1$  edges and so at most  $\frac{k}{2}$  vertices. So the indices  $i_1, i_2, \dots$  which are associated to those last graphs contribute at most for  $O(n^{k/2})$  in the previous sum.

Moreover, for the remaining case of graphs having exactly  $\frac{k}{2}$  double-edges, note that a cyclic path of  $k$  edges,  $\{i_1, i_2\}, \{i_2, i_3\}, \dots, \{i_{k-1}, i_k\}, \{i_k, i_1\}$ , such that each edge is repeated exactly twice, that is to say such that

$$\forall \ell \in \llbracket 1, k-1 \rrbracket, \exists ! \ell' \in \llbracket 1, k-1 \rrbracket, \ell \neq \ell', \quad \{i_\ell, i_{\ell+1}\} = \{i_{\ell'}, i_{\ell'+1}\}$$

corresponds to a unique Dyck path, and so, to its associated rooted planar tree. Indeed, as a Dyck path of length  $k$  is non-negative and return at time  $k$  to 0, all the edges of its associated rooted planar tree form a cyclic path of  $k/2$  edges, from the root to the root, which are visited exactly twice (see Remark 5). Thus, as the cyclic order of the edges,  $\{(i_1, i_2), \dots, (i_{k-1}, i_k), (i_k, i_1)\}$ , induces a unique orientation to those trees which make them planar with  $\frac{k}{2}$  edges and with root  $i_1$  (the only vertex which is "visited" twice), one can deduce from Lemma 31 that they are constituted by  $\frac{k}{2} + 1$  vertices. Thereby, the indices  $i_1, i_2, \dots$  which contribute the most to the previous sum, that is to say for  $O(n^{\frac{k}{2}+1})$ , are exactly those that "reconstitute" the planar rooted trees of the set  $\mathfrak{T}(\frac{k}{2})$ .

Therefore, as the number of rooted planar trees with  $\frac{k}{2}$  edges is equal to the  $(\frac{k}{2} + 1)$ -th Catalan number, that is to say to  $\frac{2}{k+2} \binom{k+2}{\frac{k}{2}+1}$ , one can conclude that the sum

$$\sum_{i_1, \dots, i_k=1}^n \frac{\mathbb{E} [x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{k-1}, i_k} x_{i_k, i_1}]}{(z - \lambda_n(i_1))^2 (z - \lambda_n(i_2)) \dots (z - \lambda_n(i_k))}$$

could be decomposed into  $\frac{2}{k+2} \binom{k+2}{\frac{k}{2}+1}$  sums related to rooted planar trees and which complexity is  $O(n^{\frac{k}{2}+1})$ , and other sums which complexity is at most  $O(n^{\frac{k}{2}})$ . In other words,

$$\varepsilon_n^{-k} \mathbb{E}[A_k(z)] = \frac{1}{n^{\frac{k}{2}+1}} \sum_{i_1, \dots, i_{\frac{k}{2}+1}=1}^n \left( \sum_{T \in \mathfrak{T}(\frac{k}{2})} \frac{\prod_{(i,j) \in \mathcal{E}_T} \mathbb{E}[x_i, x_j]}{(z - \lambda_n(x_1)) \prod_{i=1}^{k/2+1} (z - \lambda_n(x_i))^{v_i^T}} + O(n^{\frac{k}{2}}) \right)$$

Thus, by recognition of Riemann sums, as  $n$  tends to infinity, the term  $\varepsilon_n^{-k} \mathbb{E}[A_k(z)]$  tends to the term  $C_k(z)$  defined in (4.1).  $\square$

### 4.5.3 Asymptotic behavior of the terms resulting from the expansion of the resolvent of $D_n^\varepsilon$ .

The purpose of the four following claims is to describe the asymptotic behavior of each of the terms  $(A_k(z))$ . Namely, we prove in Claim 5 that the first term,  $A_1(z)$ , of the expansion converges in distribution to a Gaussian variable, in Claim 6 that all the even terms

$A_2(z), A_4(z), A_6(z) \dots$  of the expansion converge in probability respectively to the deterministic terms  $C_2(z), C_4(z), C_6(z) \dots$ , in Claim 7 that all the even terms  $A_3(z), A_5(z), A_7(z) \dots$  converge in probability to zero and in Claim 8 that the last remainder term,  $A_{2p+2}^\varepsilon(z)$  converge in probability to zero and is negligible compared to the other terms.

**Claim 5.** *The finite dimension marginals of the centered process*

$$(n\varepsilon_n^{-1}A_1(z))_{z \in \mathbb{C} \setminus \tilde{\mathcal{S}}}$$

converge in distribution to those of the centered Gaussian process  $(Z(z))_{z \in \mathbb{C} \setminus \tilde{\mathcal{S}}}$ . Besides, there is  $C > 0$  such that for any  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ ,

$$\mathbb{E}[|n\varepsilon_n^{-1}A_1(z)|^2] \leq \frac{C}{\text{dist}(z, \tilde{\mathcal{S}})^4}. \quad (4.3)$$

**Proof.** Estimate (4.3) follows from

$$\mathbb{E}[|n\varepsilon_n^{-1}A_1(z)|^2] = \frac{1}{n} \sum_{i=1}^n \frac{\mathbb{E}[|x_{i,i}|^2]}{|z - \lambda_n(i)|^4} \leq \frac{1}{n} \sum_{i=1}^n \frac{\sigma_n^2(i, i)}{\text{dist}(z, \tilde{\mathcal{S}})^4}$$

and from the existence of a uniform upper bound for  $\sigma_n^2(i, i)$  which comes from Hypothesis (f) which stipulates that the  $(4p+4)$ -th moments of the entries  $x_{i,j}$  are uniformly bounded.

We turn now to the proof of the convergence in distribution of  $n\varepsilon_n^{-1}A_1(z)$  which actually does not depend on the sequence  $(\varepsilon_n)$ . For all  $\alpha_1, \beta_1, \dots, \alpha_p, \beta_p \in \mathbb{C}$  and for all  $z_1, \dots, z_p \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ ,

$$\sum_{i=1}^p \alpha_i (n\varepsilon_n^{-1}A_1(z_i)) + \beta_i \overline{(n\varepsilon_n^{-1}A_1(z_i))} = \frac{1}{\sqrt{n}} \sum_{j=1}^n x_{j,j} \left( \sum_{i=1}^p \xi_n(i, j) \right)$$

$$\text{for } \xi_n(i, j) = \frac{\alpha_i}{(z_i - \lambda_n(j))^2} + \frac{\beta_i}{(\overline{z_i} - \lambda_n(j))^2}.$$

On one hand, by dominated convergence, the covariance matrix of the above two dimensional random vector converges.

On the other hand,  $\mathbb{E}|x_{i,j}|^4$  is uniformly bounded in  $i, j$  and  $n$ , by Hypothesis (f). Moreover, for  $n$  large enough, for all  $i, j$ ,

$$|\xi_n(i, j)| \leq 2 \max_{1 \leq i \leq p} (|\alpha_i| + |\beta_i|) \times \left( \min_{1 \leq i \leq p} \text{dist}(z_i, \mathcal{S}) \right)^{-1}.$$

Hence, the conditions of Lindeberg Central Limit Theorem are satisfied and the finite dimension marginals of the process  $(n\varepsilon_n^{-1}A_1(z))_{z \in \mathbb{C} \setminus \tilde{\mathcal{S}}}$  converge in distribution to those of

the centered Gaussian process  $(Z_z)_{z \in \mathbb{C} \setminus \tilde{\mathcal{S}}}$  defined by its covariance structure

$$\begin{aligned} \mathbb{E} \left( Z(z) \overline{Z(z')} \right) &= \lim_{n \rightarrow \infty} \mathbb{E} \left[ \left( n \varepsilon_n^{-1} A_1(z) \right) \cdot \left( n \varepsilon_n^{-1} \overline{A_1(z')} \right) \right] \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i,j=1}^n \frac{\mathbb{E} [x_{i,i} \overline{x_{j,j}}]}{(z - \lambda_n(i))^2 (\overline{z'} - \lambda_n(j))^2} \\ &= \int_0^1 \frac{\sigma_d(t)^2}{(z - f(t))^2 (\overline{z'} - f(t))^2} dt \end{aligned}$$

and by the fact that  $\overline{Z(z)} = Z(\overline{z})$  which comes from  $\overline{A_1(z)} = A_1(\overline{z})$ .  $\square$

**Claim 6.** For any even  $k \in \{2, \dots, 2p+1\}$ , there is a constant  $C$  such that for any  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ ,

$$\mathbb{E} \left[ \left| n \varepsilon_n^{-1} (A_k(z) - \varepsilon_n^k C_k(z)) \right|^2 \right] \leq \frac{C \varepsilon_n^{2k-2}}{\text{dist}(z, \tilde{\mathcal{S}})^{2k+2}} + \frac{C \eta_n^2 \varepsilon_n^{2k-4}}{\text{dist}(z, \tilde{\mathcal{S}})^{2k+4}}$$

**Proof.** Introduce the variable  $A_k^o(z)$  obtained by centering the variable  $n \varepsilon_n^{-k} A_k(z)$ :

$$\begin{aligned} A_k^o(z) &:= n \varepsilon_n^{-k} (A_k(z) - \mathbb{E} A_k(z)) \\ &= \frac{1}{n^{\frac{k}{2}}} \sum_{i_1, \dots, i_k}^n \frac{x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{k-1}, i_k} x_{i_k, i_1} - \mathbb{E} [x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{k-1}, i_k} x_{i_k, i_1}]}{(z - \lambda_n(i_1))^2 (z - \lambda_n(i_2)) \dots (z - \lambda_n(i_k))} \end{aligned}$$

and the defect variable

$$\begin{aligned} \delta_n(z) &:= \varepsilon_n^{-k} \mathbb{E} A_k(z) - C_k(z) \\ &= \frac{1}{n^{\frac{k+2}{2}}} \sum_{i_1, \dots, i_k=1}^n \frac{\mathbb{E} [x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{k-1}, i_k} x_{i_k, i_1}]}{(z - \lambda_n(i_1))^2 (z - \lambda_n(i_2)) \dots (z - \lambda_n(i_k))} \\ &\quad - \lim_{n \rightarrow \infty} \frac{1}{n^{\frac{k+2}{2}}} \sum_{i_1, \dots, i_k=1}^n \frac{\mathbb{E} [x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{k-1}, i_k} x_{i_k, i_1}]}{(z - \lambda_n(i_1))^2 (z - \lambda_n(i_2)) \dots (z - \lambda_n(i_k))} \end{aligned}$$

we want to dominate the  $L^2$  norm of  $n \varepsilon_n^{-1} (A_k(z) - \varepsilon_n^k C_k(z)) = \varepsilon_n^{k-1} A_k^o(z) + n \varepsilon_n^{k-1} \delta_n(z)$ . For this purpose, we successively dominate  $A_k^o(z)$ ,  $\delta_n(z)$  and  $C_k(z)$ .

Using the independence of the  $x_{i,j}$ 's, the fact that they are bounded in  $L^{4p+4}$  and the fact that  $z$  stays at a macroscopic distance of the  $\lambda_n(i)$ 's, we can write for all  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$

$$\begin{aligned} \mathbb{E}[|A_k^o(z)|^2] &\leq \mathbb{E}[|n \varepsilon_n^{-k} A_k(z)|^2] \\ &\leq \mathbb{E} \left[ \left| \frac{1}{n^{\frac{k}{2}}} \sum_{i_1, \dots, i_k=1}^n \frac{x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{k-1}, i_k} x_{i_k, i_1}}{(z - \lambda_n(i_1))^2 (z - \lambda_n(i_2)) \dots (z - \lambda_n(i_k))} \right|^2 \right] \\ &\leq \frac{1}{n^k} \sum_{i_1, \dots, i_{2k}=1}^n \frac{\mathbb{E} [x_{i_1, i_2} \dots x_{i_k, i_{k+1}} x_{i_{k+1}, i_{k+2}} \dots x_{i_{2k-1}, i_{2k}}]}{\text{dist}(z, \tilde{\mathcal{S}})^{2k+2}} \end{aligned}$$

Generically, the set of "edges"  $\{(i_1, i_2), (i_2, i_3), \dots, (i_{k-1}, i_k)\}$  must be equal to the set  $\{(i_{k+1}, i_{k+2}), \dots, (i_{2k-1}, i_{2k})\}$  in order to get a non zero term. Therefore the complexity of the previous sum is  $O(n^k)$ . Note that other non zero terms involving third or fourth moments are much less numerous. Hence,

$$\mathbb{E}[|A_k^o(z)|^2] \leq \frac{O(n^k)}{n^k \text{dist}(z, \tilde{\mathcal{S}})^{2k+2}} \leq \frac{C}{\text{dist}(z, \tilde{\mathcal{S}})^{2k+2}}. \quad (4.4)$$

Now, note that the expression of the finite and deterministic term of  $C_k(z)$ , defined in (4.1), implies that,

$$C_k(z) \leq \frac{C}{\text{dist}(z, \tilde{\mathcal{S}})^{k+1}}. \quad (4.5)$$

Finally, the term  $\delta_n(z)$  rewrites

$$\begin{aligned} \delta_n(z) = & O(n^{-1}) + \sum_{T \in \mathfrak{T}(\frac{k}{2})} \int_{[0,1]^{\frac{k}{2}+1}} \mathbb{1}_{\mathfrak{B}} \left( \frac{\prod_{\{i,j\} \in \mathcal{E}_T} \sigma^2(\lfloor nx_i \rfloor, \lfloor nx_j \rfloor)}{(z - \lambda_n(\lfloor nx_1 \rfloor)) \prod_{i=1}^{k/2+1} (z - \lambda_n(\lfloor nx_i \rfloor))^{v_i^T}} \right. \\ & \left. - \frac{\prod_{\{i,j\} \in \mathcal{E}_T} \sigma^2(x_i, x_j)}{(z - f(x_1)) \prod_{i=1}^{k/2+1} (z - f(x_i))^{v_i^T}} dx_2 \dots dx_{\frac{k}{2}+1} \right) dx_1 \dots dx_{\frac{k}{2}+1}, \end{aligned}$$

where  $\mathfrak{B} := \left\{ (x_1, \dots, x_{\frac{k}{2}+1}) \in [0, 1]^{\frac{k}{2}+1} ; \forall i \neq j, \lfloor nx_i \rfloor \neq \lfloor nx_j \rfloor \right\}$ .

Since, for  $M_\sigma := \sup_{0 \leq x \neq y \leq 1} \sigma(x, y)^2$  and for any fixed  $z \notin \tilde{\mathcal{S}}$ , the function

$$\psi_z(s, \lambda_1, \dots, \lambda_{\frac{k}{2}+1}) \mapsto \frac{s}{(z - \lambda_1) \prod_{i=1}^{k/2+1} (z - \lambda_i)^{v_i^T}}$$

defined for  $(s, \lambda_1, \dots, \lambda_{\frac{k}{2}+1}) \in [0, M_\sigma + 1] \times \{x \in \mathbb{R} ; \text{dist}(x, \tilde{\mathcal{S}}) \leq \text{dist}(z, \tilde{\mathcal{S}})/2\}^{\frac{k}{2}+1}$  is  $C \text{dist}(z, \tilde{\mathcal{S}})^{-(k+2)}$ -Lipschitz, for  $C$  a universal constant, by Hypothesis (b),

$$\delta_n(z) = O(n^{-1}) + \frac{O(\eta_n)}{\max\{n\varepsilon_n, 1\} \text{dist}(z, \tilde{\mathcal{S}})^{k+2}}. \quad (4.6)$$

Collecting estimations (4.4), (4.5) and (4.6) we conclude.  $\square$

**Claim 7.** *For any odd  $k \in \{3, \dots, 2p+1\}$  there is a constant  $C$  such that for any  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ ,*

$$\mathbb{E}[|n\varepsilon_n^{-1} A_k(z)|^2] \leq \frac{C\varepsilon_n^{2k-2}}{\text{dist}(z, \tilde{\mathcal{S}})^{2k+2}}.$$

**Proof.** We start by writing for all  $z \in \mathbb{C} \setminus \tilde{\mathcal{S}}$ ,

$$\begin{aligned} \mathbb{E}[|n\varepsilon_n^{-1}A_k(z)|^2] &= \frac{\varepsilon_n^{2k-2}}{n^k} \mathbb{E} \left[ \left| \sum_{i_1, \dots, i_k=1}^n \frac{x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{k-1}, i_k} x_{i_k, i_1}}{(z - \lambda_n(i_1))^2 (z - \lambda_n(i_2)) \dots (z - \lambda_n(i_k))} \right|^2 \right] \\ &= \sum_{i_1, \dots, i_{2k}=1}^n \frac{\varepsilon_n^{2k-2} \mathbb{E} [x_{i_1, i_2} x_{i_2, i_3} \dots x_{i_{2k-1}, i_{2k}}]}{n^k (z - \lambda_n(i_1))^2 (z - \lambda_n(i_2)) \dots (z - \lambda_n(i_k)) (\bar{z} - \lambda_n(i_{k+1}))^2 (\bar{z} - \lambda_n(i_{k+2})) \dots (\bar{z} - \lambda_n(i_{2k}))}. \end{aligned}$$

Generically, the set of "edges"  $\{(i_1, i_2), (i_2, i_3), \dots, (i_{k-1}, i_k)\}$  must be equal to the set  $\{(i_{k+1}, i_{k+2}), \dots, (i_{2k-1}, i_{2k})\}$  in order to get a non zero term. Therefore, as  $k$  is odd, the complexity of the previous sum is  $O(n^k)$ . Note that other non zero terms involving third or fourth moments are much less numerous. Hence,

$$\mathbb{E}[|n\varepsilon_n^{-1}A_k(z)|^2] \leq \frac{\varepsilon_n^{2k-2}}{n^k} \times \frac{O(n^k)}{\text{dist}(z, \tilde{\mathcal{S}})^{2k+2}} \leq \frac{C\varepsilon_n^{2k-2}}{\text{dist}(z, \tilde{\mathcal{S}})^{2k+2}}$$

□

**Claim 8.** For any  $z \in \mathbb{C} \setminus \mathbb{R}$ ,

$$\mathbb{E}[|n\varepsilon_n^{-1}A_{2p+2}^\varepsilon(z)|^2] \leq \frac{O(n^2\varepsilon_n^{4p+2})}{|\text{Im}(z)|^2 \text{dist}(z, \tilde{\mathcal{S}})^{4p+4}}.$$

**Proof.** Remind that,

$$A_{2p+2}^\varepsilon(z) := \frac{\varepsilon_n^{2p+2}}{n} \text{Tr} \underbrace{\frac{1}{z-D} X \dots \frac{1}{z-D} X}_{(2p+2) \text{ times } \frac{1}{z-D} X} \frac{1}{z-D^\varepsilon}$$

Hence,

$$\begin{aligned} \mathbb{E}[|n\varepsilon_n^{-1}A_{2p+2}^\varepsilon(z)|^2] &\leq \varepsilon_n^{4p+2} \mathbb{E} \left[ \left| \text{Tr} \underbrace{\frac{1}{z-D} X \dots \frac{1}{z-D} X}_{(2p+2) \text{ times } \frac{1}{z-D} X} \frac{1}{z-D^\varepsilon} \right|^2 \right] \\ &\leq \varepsilon_n^{4p+2} \mathbb{E} \left[ \text{Tr} \left| \left( \frac{1}{z-D} X \right)^{2p+2} \right|^2 \times \text{Tr} \left| \frac{1}{z-D^\varepsilon} \right|^2 \right] \\ &\leq \varepsilon_n^{4p+2} \mathbb{E} \left[ \text{Tr} \left( \left( \frac{1}{z-D} X \right)^{2p+2} \left( \overline{\frac{1}{z-D} X} \right)^{2p+2} \right) \frac{n}{|\text{Im}(z)|^2} \right] \\ &\leq \frac{n\varepsilon_n^{4p+2}}{|\text{Im}(z)|^2} \mathbb{E} \left[ \text{Tr} \left( \left( \frac{1}{z-D} X \right)^{2p+2} \left( \overline{\frac{1}{z-D} X} \right)^{2p+2} \right) \right] \\ &\leq \frac{n\varepsilon_n^{4p+2}}{|\text{Im}(z)|^2} \frac{O(n^{(2p+3)})}{n^{2p+2} \text{dist}(z, \tilde{\mathcal{S}})^{4p+4}} \leq \frac{O(n^2\varepsilon_n^{4p+2})}{|\text{Im}(z)|^2 \text{dist}(z, \tilde{\mathcal{S}})^{4p+4}}. \end{aligned}$$

The inequality of the last line takes into account that

- the  $L^{4p+4}$  norm of the entries of  $\sqrt{n}X$  is uniformly bounded
- the norm of the entries of  $X$  is of order  $n^{-1/2}$
- the norm of the coefficients of  $(z - D)^{-1}$  is smaller than  $\text{dist}(z, \tilde{\mathcal{S}})^{-1}$
- the complexity of the sum defining the trace is of order  $O(n^{2p+3})$  since its non-null terms are encoded by  $(2p+2)$  edges trees which have therefore  $(2p+3)$  vertices.

Moreover, note that, as we have assumed that  $n^{-1/(2p-1)} \ll \varepsilon_n \ll n^{-1/(2p+1)}$ , the term  $n^2 \varepsilon_n^{4p+2}$  is  $o(1)$ .  $\square$

#### 4.5.4 Proof of Theorem 28

The proof of Theorem 28 is based on the perturbative expansion (4.2) of the resolvent  $\frac{1}{n} \text{Tr} \frac{1}{z - D_n^\varepsilon}$  and the four previous Claims.

**Proof.** We gather now the results of the previous claims. If  $n^{-1/(2p-1)} \ll \varepsilon_n \ll n^{-1/(2p+1)}$ , then, for any  $z \in \mathbb{C} \setminus \mathbb{R}$ , we need to make the following expansion until the order  $(2p+2)$ ,

$$n\varepsilon_n^{-1} \Delta G_n(z) = n\varepsilon_n^{-1} (A_1(z) + A_2(z) + A_3(z) + \cdots + A_{2p+1}(z) + A_{2p+2}^\varepsilon(z))$$

and

- Claim 5 proves that the process  $n\varepsilon_n^{-1} A_1(z)$  converges in distribution to the centered Gaussian variable  $Z(z)$ ,
- Claim 6 proves that for any even  $k \in \{2, \dots, 2p\}$  each term  $n\varepsilon_n^{-1} A_k(z)$  converges in probability to the deterministic term  $C_k(z)$ ,
- Claim 7 proves that for any odd  $k \in \{3, \dots, 2p+1\}$  each term  $n\varepsilon_n^{-1} A_k(z)$  converges in probability to zero,
- and Claim 8 proves that the remainder term  $n\varepsilon_n^{-1} A_{2p+2}^\varepsilon(z)$  converges in probability to zero.

Thus, by Slutsky's theorem, for any  $z \in \mathbb{C} \setminus \mathbb{R}$ :

$$n\varepsilon_n^{-1} (\Delta G_n(z) - \varepsilon_n^2 C_2(z) - \varepsilon_n^4 C_4(z) \cdots - \varepsilon_n^{2p} C_{2p}(z)) \xrightarrow[n \rightarrow \infty]{\text{dist.}} Z(z)$$

This finishes the proof of Theorem 28.  $\square$



### 4.5.5 Proof of the local type convergence result

Thanks to the perturbative expansion (4.2) of the resolvent  $\frac{1}{n} \text{Tr} \frac{1}{z - D_n^\varepsilon}$  and Claim 5, Claim 6, Claim 7 and Claim 8 we can also proceed to the proof of Theorem 29.

**Proof.** Assume  $n^{-1} \ll \varepsilon_n \ll 1$  and note that  $\varepsilon_n^{-p} \Delta G_n(z_n) - \sum_{\substack{k=0 \\ k \text{ even}}}^{p-2} \varepsilon_n^k C_{p-k}(z_n)$  rewrites

$$\varepsilon_n^{-p} (A_1(z_n)) + \varepsilon_n^{-p} (A_2(z_n) - \varepsilon_n^2 C_2(z_n)) + \varepsilon_n^{-p} A_3(z_n) + \varepsilon_n^{-p} (A_4(z_n) - \varepsilon_n^4 C_4(z_n)) + \cdots + \varepsilon_n^{-p} A_{p+2}^\varepsilon(z_n)$$

One can directly obtain, for all non-real complex sequences  $(z_n)$ , that

- by Claim 5, if  $\text{dist}(z_n, \tilde{\mathcal{S}}) \gg (n\varepsilon_n^{p-1})^{-1/2}$ , then

$$\mathbb{E}[|\varepsilon_n^{-p} A_1(z_n)|^2] \leq \frac{C}{(n\varepsilon_n^{p-1})^2 \text{dist}(z_n, \tilde{\mathcal{S}})^4} \xrightarrow{n \rightarrow \infty} 0,$$

- by Claim 6, for any even  $k \in \{2, p\}$ , if  $\text{dist}(z_n, \tilde{\mathcal{S}}) \gg \max \left\{ \left( \frac{\varepsilon_n^{2-p}}{n} \right)^{1/3}, (\eta_n / (n\varepsilon_n^{p-1}))^{1/4} \right\}$ , then

$$\mathbb{E} \left[ |\varepsilon_n^{-p} (A_k(z_n) - \varepsilon_n^k C_k(z_n))|^2 \right] \leq \frac{C\varepsilon_n^{2(k-p)}}{n^2 \text{dist}(z_n, \tilde{\mathcal{S}})^{2k+2}} + \frac{C\eta_n^2 \varepsilon_n^{-2p}}{n^2 \text{dist}(z_n, \tilde{\mathcal{S}})^{2k+4}} \xrightarrow{n \rightarrow \infty} 0$$

- by Claim 7, for any odd  $k \in \{3, p+1\}$ , if  $\text{dist}(z_n, \tilde{\mathcal{S}}) \gg \left( \frac{\varepsilon_n^{3-p}}{n} \right)^{1/4}$ , then

$$\mathbb{E}[|\varepsilon_n^{-p} C_k(z_n)|^2] \leq \frac{C\varepsilon_n^{2(k-p)}}{n^2 \text{dist}(z_n, \tilde{\mathcal{S}})^{2k+2}} \xrightarrow{n \rightarrow \infty} 0,$$

- by Claim 8, if  $|\Im(z_n)| \text{dist}(z_n, \tilde{\mathcal{S}})^{p+2} \gg \varepsilon_n^2$ , then

$$\mathbb{E}[|\varepsilon_n^{-p} A_{p+2}^\varepsilon(z_n)|^2] \leq \frac{O(\varepsilon_n^4)}{|\Im(z_n)|^2 \text{dist}(z_n, \tilde{\mathcal{S}})^{2p+4}} \xrightarrow{n \rightarrow \infty} 0.$$

Therefore, when

$$\text{dist}(z_n, \tilde{\mathcal{S}}) \gg \max \left\{ (n\varepsilon_n^{p-1})^{-1/2}, \left( \frac{\varepsilon_n^{2-p}}{n} \right)^{1/3}, \left( \frac{\eta_n}{n\varepsilon_n^{p-1}} \right)^{1/4}, \left( \frac{\varepsilon_n^{3-p}}{n} \right)^{1/4} \right\}$$

and

$$|\Im(z_n)| \text{dist}(z_n, \tilde{\mathcal{S}})^{p+2} \gg \varepsilon_n^2,$$

all the processes that constitute the process  $\varepsilon_n^{-p} \Delta G_n(z_n) - \sum_{\substack{k=0 \\ k \text{ even}}}^{p-2} \varepsilon_n^{-k} C_{p-k}(z)$  converge to 0 in probability. Since  $\text{dist}(z_n, \tilde{\mathcal{S}}) \geq \Im(z_n)$ , the above condition is implied by

$$\Im(z_n) \gg \max \left\{ (n\varepsilon_n^{p-1})^{-1/2}, \left( \frac{\varepsilon_n^{2-p}}{n} \right)^{1/3}, \left( \frac{\eta_n}{n\varepsilon_n^{p-1}} \right)^{1/4}, \left( \frac{\varepsilon_n^{3-p}}{n} \right)^{1/4}, \varepsilon_n^{\frac{2}{p+3}} \right\}.$$

Observing finally that the two terms  $\left( \frac{\varepsilon_n^{2-p}}{n} \right)^{1/3}$  and  $\left( \frac{\varepsilon_n^{3-p}}{n} \right)^{1/4}$  are dominated by the maximum of the three other ones, we conclude the proof.  $\square$

## 4.6 A possible extension to other classes of test functions

Theorems 28 and 29 could be extended from test functions of the type  $\varphi_z(x) := \frac{1}{z-x}$  to larger classes of test functions. To make this extension possible it is necessary to succeed in expressing the terms  $C_k(z)$ , defined in (4.1), as linear expressions of the function  $\varphi_z(x)$ . Thereby, the expansions of  $(\mu_n^\varepsilon - \mu_n)(\varphi_z)$  that we have previously considered, would be expressed entirely as linear forms of  $\varphi_z$ , which will allow to extend this convergence, by density of the linear span of the functions  $\varphi_z(x)$  in the fractional Sobolev space  $\mathcal{H}_s$ , for any  $s > 0$ , to a larger class of regular functions through a CLT extension theorem. For example, one could use the CLT extension lemma of Scherbina and Tirozzi which could be found in [ST10].

This extension have been done in [BGEM17], for all the regimes where  $\varepsilon_n \ll n^{-1/3}$ , that is to say for all the regimes that involve at most one deterministic term, which comes from  $C_2(z)$ . In fact, in this paper the expression  $C_2(z)$  have been expressed under the following form which is linear in  $\varphi_z(x)$ :

$$C_2(z) = \int_{[0,1]^2} \frac{\sigma^2(x_1, x_2)}{(z - f(x_1))^2 (z - f(x_2))} dx_1 dx_2 = - \int_{\mathbb{R}} \varphi'_z(s) F(s) ds$$

where  $F(s) = \int_{\mathbb{R}} \frac{\tau(s,t)\rho(s)\rho(r)}{s-t} dt$  for  $\tau$  a function such that  $\tau(s, t) = \sigma^2(f^{-1}(s), f^{-1}(t))$  and  $\rho$  the density of the push-forward of the uniform measure on  $[0, 1]$  by the function  $f$ .

But from the second deterministic term  $C_4(z)$  it becomes much more difficult to give a linear expression in  $\varphi_z(x)$  of  $C_k(z)$  without making too restrictive assumptions on the diagonal entries of the matrix  $X_n$ .

If one succeeds in overcoming this technical limitation on the diagonal entries of  $X_n$ , then by a similar approach that in [BGEM17] a result, as for example the following that in addition to using the above-mentioned CLT extension lemma uses the Helffer-Sjöstrand formula (see [HS89]) and the fact that  $\mathcal{C}^{2p+4}(\mathbb{R}) \subset \mathcal{H}_s$  for  $s < 2p + 4$  (see [Hör03]), could be proved:

Let  $p \in \mathbb{N}^*$ . Under the semi-perturbative regime, if  $n^{-1/(2p-1)} \ll \varepsilon_n \ll n^{-1/(2p+1)}$  then there exist  $p$  linear deterministic functions,  $D_1, \dots, D_p$ , defined on the space of compactly supported  $\mathcal{C}^{2p+4}$  functions on  $\mathbb{R}$  such that for any compactly supported  $\mathcal{C}^{2p+4}$  function,  $\phi$ , on  $\mathbb{R}$ , the following convergence holds :

$$n\varepsilon_n^{-1} ((\mu_n^\varepsilon - \mu_n)(\phi) + D_1(\phi) + \dots + D_p(\phi)) \xrightarrow[n \rightarrow \infty]{\text{dist.}} Z_\phi.$$

where  $(Z_\phi)_{\phi \in \mathcal{C}^1}$  is the centered Gaussian field indexed by the set of  $\mathcal{C}^1$  complex functions on  $\mathbb{R}$ , with covariance structure defined by

$$\mathbb{E}Z_\phi Z_\psi = \int_0^1 \sigma_d(t)^2 \phi'(f(t))\psi'(f(t))dt \quad \text{and} \quad \overline{Z_\psi} = Z_{\overline{\psi}}$$

and which can be represented as

$$Z_\phi = \int_0^1 \sigma_d(t) \phi'(f(t)) dB_t$$

where  $(B_t)$  is the standard one-dimensional Brownian motion.

Note that the computation of the deterministic functions  $D_1, \dots, D_p$  is rather intricate and without any clear recursive formula. Indeed, each term  $D_k$  comes from the deterministic term  $C_{2k}(z)$  after some computations which have no clear recurrence relation such that partial fraction decompositions. Also, according to this link between  $D_k$  and  $C_{2k}(z)$ , all the deterministic terms  $D_1(\phi), \dots, D_p(\phi)$  have a much larger order than the probabilistic term containing  $Z_\phi$  and, more precisely, for any  $k \in \{2, \dots, p-1\}$  the deterministic term  $D_k(\phi)$  has a smaller order than  $D_{k-1}(\phi)$  and a larger than  $D_{k+1}(\phi)$ . Namely,  $D_k(\phi)$  is of order  $(\varepsilon_n)^k$ .

# Chapter 5

## Eigenvectors of a matrix under random perturbation

### Contents

5.1	Introduction . . . . .	87
5.2	Main result . . . . .	89
5.3	Proof (of Theorem 32) . . . . .	91
5.3.1	Stieltjes transform . . . . .	92
5.3.2	From Stieltjes transform to $\mathcal{C}^5$ functions . . . . .	95
5.4	Consequence for the eigenvectors . . . . .	96

### 5.1 Introduction

This last chapter is devoted to the study of the sensitivity of the eigenvectors of a given operator under small perturbations. In the previous chapters we studied the effect of a perturbation on the spectrum of a Hermitian matrix by a random matrix with small operator norm and whose entries in the eigenvector basis of the first one were independent, centered, with a variance profile. We provided a perturbative expansion of the empirical spectral distribution, but did not consider the deformation of the eigenvectors basis with respect to the canonical basis. In the present paper, to complete this first study, we deal with the spectral measure of our matrix associated to the state defined by a given vector.

To define this measure, let us introduce some notations. We consider a real diagonal matrix  $D_n = \text{diag}(\lambda_1, \dots, \lambda_n)$  (the eigenvalue  $\lambda_i$  implicitly depends on  $n$ ), as well as a Hermitian random matrix

$$X_n = \frac{1}{\sqrt{n}} [x_{i,j}^n]_{1 \leq i,j \leq n}$$

such that the  $x_{ij}$  are independent (up to the symmetry), centered, with a variance profile. The normalizing factor  $n^{-1/2}$  and our hypotheses below ensure that the operator norm of  $X_n$  is of order one. We then define, for  $\varepsilon > 0$ ,

$$D_n^\varepsilon := D_n + \varepsilon X_n.$$

In contrast with [BGEM17], where we studied the empirical spectral measure  $\mu_n^\varepsilon$  of the matrix  $D_n^\varepsilon$ , we consider here the spectral measure  $\mu_{n, \mathbf{e}_i}^\varepsilon$  of  $D_n^\varepsilon$  over a vector  $\mathbf{e}_i$  of the canonical basis, defined through an eigenvector basis  $(\mathbf{u}_j^\varepsilon)_{j \in \{1, \dots, n\}}$  of  $D_n^\varepsilon$  and the related eigenvalues  $(\lambda_j^\varepsilon)_{j \in \{1, \dots, n\}}$  by

$$\mu_{n, \mathbf{e}_i}^\varepsilon := \sum_{j=1}^n |\langle \mathbf{u}_j^\varepsilon, \mathbf{e}_i \rangle|^2 \delta_{\lambda_j^\varepsilon}.$$

The interest of these measures is that they give information on the eigenvector basis of  $D_n^\varepsilon$ , while being tractable since they satisfy, for any test function  $\varphi$ , the key identity

$$\int \varphi(x) d\mu_{n, \mathbf{e}_i}^\varepsilon(x) = \sum_{j=1}^n |\langle \mathbf{u}_j^\varepsilon, \mathbf{e}_i \rangle|^2 \varphi(\lambda_j^\varepsilon) = (\varphi(D_n^\varepsilon))_{i,i}. \quad (5.1)$$

Our main result, Theorem 32, gives a perturbative expansion of  $\mu_{n, \mathbf{e}_i}^\varepsilon$ . More precisely, using a resolvent expansion and the Helffer-Sjöstrand formula, we give an asymptotic expansion of

$$\int_{\mathbb{R}} \varphi(t) d\mu_{n, \mathbf{e}_i}^\varepsilon(t)$$

for any  $\mathcal{C}^5$  test function  $\varphi$ . From that, we deduce, at Equation (5.17), a heuristic first order estimation of the overlaps  $|\langle \mathbf{u}_j^\varepsilon, \mathbf{e}_i \rangle|^2$  between the eigenvectors  $\mathbf{u}_j^\varepsilon$  of  $D_n^\varepsilon$  and the ones  $\mathbf{e}_i$  of  $D_n$ .

Some other works, on models closed to our one or contained in it, are devoted to the sensitivity to perturbations of the eigenvectors. Some of them, as [OVW16, OVW17, vSW17, Zho17], provide bounds on the deviations of these eigenvectors under perturbation, while some other, as [AB12, AB14, ABB14, Ben17], provide explicit perturbative expansions. As we explained in the previous paragraph, this is what we do here, and it allows us to recover (see (5.17)) the fact that the overlaps  $|\langle \mathbf{u}_j^\varepsilon, \mathbf{e}_i \rangle|^2$  have order  $\varepsilon^2(\lambda_j - \lambda_i)^2/n$ . This cannot be proved for all  $i, j$  but only on average over more or less large windows. This is why estimates of this flavor have already appeared in various papers, as in [AB12, AB14, ABB14, Ben17] under various forms (for example, one of the differences between our result and [Ben17, Coro. 1.6] is that we average on  $j$  and not on  $i$ , as explained in Remark 5). In addition to the fact that it only relies on short and elementary computations, one of the interests of this paper is also to consider rather general perturbations, since we do not suppose that all entries of  $X_n$  have the same variance nor that they are Gaussian. Another interest is to provide, with the functional  $\Xi_s(\varphi)$  from (5.5) and (5.9), an expression for the first order expansion of the measure  $\mu_{n, \mathbf{e}_i}^\varepsilon$  from (5.1), which, up to our knowledge, did not appear so far.

The paper is organized as follows. Results and comments are given in Section 5.2, proofs in Section 5.3, while Section 5.4 is devoted to the heuristic derivation of the consequences of Theorem 32, and to some simulations which corroborate our predictions.

**Notation.** For  $u = u_n$ ,  $v = v_n$  some sequences,  $u \ll v$  means that  $u_n/v_n$  tends to 0.

## 5.2 Main result

We consider a real diagonal matrix  $D_n = \text{diag}(\lambda_1, \dots, \lambda_n)$  (the eigenvalue  $\lambda_i$  implicitly depend on  $n$ ), as well as a Hermitian random matrix

$$X_n = \frac{1}{\sqrt{n}} [x_{i,j}^n]_{1 \leq i,j \leq n}$$

and define, for  $\varepsilon > 0$ ,

$$D_n^\varepsilon := D_n + \varepsilon X_n.$$

We make the following hypotheses:

- (a) the entries  $x_{i,j}^n$  of  $\sqrt{n}X_n$  are independent (up to symmetry) random variables, centered, with variance denoted by  $\sigma_n^2(i, j)$ , such that  $\mathbb{E}|x_{i,j}^n|^6$  is bounded uniformly on  $n, i, j$ ,
- (b) there are two bounded real functions,  $f$  and  $\sigma$ , defined respectively on  $[0, 1]$  and  $[0, 1]^2$  such that, denoting  $\lambda_i$  by  $\lambda_{n,i}$  to emphasize the implicit dependence in  $n$ , the error bound

$$\eta_n := \sup_{x \in [0,1]} |\lambda_{n,[nx]} - f(x)| + \sup_{(x,y) \in [0,1]^2} |\sigma_n^2([nx], [ny]) - \sigma^2(x, y)| \quad (5.2)$$

satisfies

$$\eta_n \xrightarrow{n \rightarrow \infty} 0.$$

Let us now make some assumptions on the limiting functions  $\sigma$  and  $f$ :

- (c) the push-forward of the uniform measure on  $[0, 1]$  by the function  $f$  has a density  $\rho$  with respect to the Lebesgue measure on  $\mathbb{R}$  and a compact support denoted by  $\mathcal{S}$ ,
- (d) the variances of the entries of  $X_n$  essentially depend on the eigenspaces of  $D_n$ , namely, there exists a symmetric function  $\tau(\cdot, \cdot)$  on  $\mathbb{R}^2$  such that for all  $x \neq y$ ,  $\sigma^2(x, y) = \tau(f(x), f(y))$ .

Let  $\mu_{n,\mathbf{e}_i}^\varepsilon$  denote the probability measure defined, for any test function  $\varphi$ , by

$$\int \varphi(t) d\mu_{n,\mathbf{e}_i}^\varepsilon(t) := (\varphi(D_n^\varepsilon))_{ii}. \quad (5.3)$$

One can equivalently define  $\mu_{n,\mathbf{e}_i}^\varepsilon$  by

$$\mu_{n,\mathbf{e}_i}^\varepsilon := \sum_{j=1}^n |\langle \mathbf{u}_j^\varepsilon, \mathbf{e}_i \rangle|^2 \delta_{\lambda_j^\varepsilon}, \quad (5.4)$$

where  $\mathbf{e}_i$  denotes the  $i$ -th vector of the canonical basis, the  $\lambda_j^\varepsilon$ 's denote the eigenvalues of  $D_n^\varepsilon$  and the  $\mathbf{u}_j^\varepsilon$ 's denote the associated eigenvectors.

We now introduce a functional which is central in the statement of our result. This functional admits another expression, given in Proposition 33 below.

Let, for  $s \in \mathbb{R}$  and  $\varphi : \mathbb{R} \rightarrow \mathbb{C}$  a  $\mathcal{C}^2$  function,

$$\Xi_s(\varphi) := \int_{\mathbb{R}} \tau(s, t) \rho(t) \frac{(\varphi(t) - \varphi(s) - (t-s)\varphi'(s))}{(t-s)^2} dt \quad (5.5)$$

**Theorem 32.** *Let us suppose that  $\varepsilon \ll 1$ . Let  $\varphi : \mathbb{R} \rightarrow \mathbb{C}$  be a compactly supported  $\mathcal{C}^5$  function. For  $x \in [0, 1]$ , set  $i = i(n, x) = \lfloor nx \rfloor$ . Then we have*

$$\int_{\mathbb{R}} \varphi(t) d\mu_{n,\mathbf{e}_i}^\varepsilon(t) = \varphi\left(\lambda_i + \frac{\varepsilon}{\sqrt{n}} x_{ii}\right) + \varepsilon^2 \Xi_{f(x)}(\varphi) + O_{L^2}\left(\varepsilon^2 \|\varphi^{(5)}\|_\infty (\varepsilon + n^{-1/2} + \eta_n)\right)$$

for  $\eta_n$  as in (5.2).

**Remark 2** (Leading order transition). Note that for any  $\mathcal{C}^1$  test function  $\varphi$ ,

$$\varphi\left(\lambda_i + \frac{\varepsilon}{\sqrt{n}} x_{ii}\right) = \varphi(\lambda_i) + \frac{\varepsilon}{\sqrt{n}} x_{ii} \varphi'(\lambda_i) + O_{L^2}\left(\frac{\varepsilon^2}{n} \|\varphi''\|_\infty\right).$$

Thus the previous theorem allows to expand the measure  $\mu_{n,\mathbf{e}_i}^\varepsilon$  around  $\delta_{\lambda_i}$  as follows. With the notations and the hypothesis of the theorem,

$$\begin{aligned} \int \varphi(t) d\mu_{n,\mathbf{e}_i}^\varepsilon(t) &= \varphi(\lambda_i) + \frac{\varepsilon}{\sqrt{n}} x_{ii} \varphi'(\lambda_i) + \varepsilon^2 \Xi_s(\varphi) \\ &\quad + O_{L^2}\left(\varepsilon^2 \|\varphi^{(5)}\|_\infty (\varepsilon + n^{-1/2} + \eta_n) + \frac{\varepsilon^2}{n} \|\varphi''\|_\infty\right). \end{aligned} \quad (5.6)$$

As far as the leading term of the expansion of  $\mu_{n,\mathbf{e}_i}^\varepsilon - \delta_{\lambda_i}$  is concerned, we deduce the following transition: for  $\eta_n$  small enough, if  $\varepsilon \ll n^{-1/2}$ , then the leading term is  $\frac{\varepsilon}{\sqrt{n}} x_{ii} \varphi'(\lambda_i)$ , whereas if  $n^{-1/2} \ll \varepsilon \ll 1$ , the leading term is  $\varepsilon^2 \Xi_s(\varphi)$ .

**Remark 3.** Strikingly, the image of a function  $\varphi$  by the operator  $\Xi_{f(x)}$  is not changed if one adds an affine function to  $\varphi$ . This can be understood because the measure  $\mu_{n, \mathbf{e}_i}^\varepsilon - \delta_{\lambda_i}$  is of null mass and with first moment of order  $o(\varepsilon^2)$  since by (5.1),

$$\int_{\mathbb{R}} x \, d(\mu_{n, \mathbf{e}_i}^\varepsilon - \delta_{\lambda_i}) = (D_n^\varepsilon)_{ii} - \lambda_i = \frac{\varepsilon}{\sqrt{n}} x_{ii} = o(\varepsilon^2).$$

Note that when both  $\varphi(f(x))$  and  $\varphi'(f(x))$  are null, the function  $\Xi_{f(x)}(\varphi)$  boils down to the integral

$$\Xi_{f(x)}(\varphi) = \int_{\mathbb{R}} \tau(f(x), t) \rho(t) \frac{\varphi(t)}{(t - f(x))^2} dt. \quad (5.7)$$

We will use this fact in Section 5.4 for test functions  $\varphi$  whose support does not contain  $f(x)$ .

**Proposition 33.** *Let us define, for any  $s \in \mathbb{R}$ , the function  $\zeta_s$  defined on  $\mathbb{R}$  by*

$$\zeta_s(y) := \int_1^{+\infty} \frac{r-1}{r^2} \tau(s, s+r(y-s)) \rho(s+r(y-s)) dr. \quad (5.8)$$

*Then for any  $\mathcal{C}^2$  function  $\varphi$  and any  $s \in \mathbb{R}$ , the functional  $\Xi_s$  defined at (5.5) rewrites*

$$\Xi_s(\varphi) = \int_{\mathbb{R}} \varphi''(y) \zeta_s(y) dy. \quad (5.9)$$

**Proof.** Taylor formula yields

$$\varphi(t) - \varphi(s) - (t-s)\varphi'(s) = \int_s^t \varphi''(x)(t-x) dx = (t-s)^2 \int_{u=0}^1 \varphi''(s+u(t-s))(1-u) du.$$

Hence,

$$\Xi_s(\varphi) = \int_{t \in \mathbb{R}} \int_{u=0}^1 \varphi''(s+u(t-s))(1-u) du \, \tau(s, t) \rho(t) dt$$

We now perform the change of variable  $(r, y) = \Psi_s(u, t)$  with

$$\Psi_s : (u, t) \in (0, 1) \times \mathbb{R} \mapsto (r, y) = \left( \frac{1}{u}, u(t-s) + s \right) \in (1, \infty) \times \mathbb{R}$$

which gives the result □

## 5.3 Proof (of Theorem 32)

The proof is divided into two parts. We shall first prove a convergence result for test functions  $\varphi$  of the type  $\varphi_z := \frac{1}{z-x}$ . This is the purpose of Subsection 5.3.1. It will be obtained by writing an expansion of the resolvent of  $D_n^\varepsilon$ .

Once we have proved that such a convergence holds for the resolvent of  $D_n^\varepsilon$ , we will be able to extend it to the class of compactly supported  $\mathcal{C}^5$  functions on  $\mathbb{R}$ , by using the Helffer-Sjöstrand formula (see [HS89] or [BGK16]) which expresses a regular function  $\varphi$  on  $\mathbb{R}$  as an integral against functions  $\varphi_z$  of the previous type. This is done in Subsection 5.3.2.



### 5.3.1 Stieltjes transform

Let us introduce the Banach space  $\mathcal{C}_b^2$  of bounded  $\mathcal{C}^2$  functions on  $\mathbb{R}$  with bounded first and second derivatives, endowed with the norm  $\|\varphi\|_{\mathcal{C}_b^2} := \|\varphi\|_\infty + \|\varphi'\|_\infty + \|\varphi''\|_\infty$ .

On this space, let us define, for  $x \in [0, 1]$  and  $i = \lfloor nx \rfloor$ , the random continuous linear form

$$\Pi_n(\varphi) := \varepsilon^{-2} \left( \int \varphi(t) d\mu_{n, \mathbf{e}_i}^\varepsilon(t) - \varphi(\lambda_i + \frac{\varepsilon}{\sqrt{n}} x_{ii}) \right) - \Xi_{f(x)}(\varphi).$$

**Lemma 34.** *Uniformly in  $z \in \mathbb{C} \setminus \mathbb{R}$ ,*

$$\mathbb{E}[|\Pi_n(\varphi_z)|^2] = O\left(\left(\eta_n^2 + \frac{1}{n}\right) |\Im z|^{-6}\right) + O(\varepsilon^2 |\Im z|^{-8}). \quad (5.10)$$

**Remark 4.** This result implies that  $\forall z \in \mathbb{C} \setminus \mathbb{R}$ ,  $\Pi_n(\varphi_z) \xrightarrow[n \rightarrow \infty]{P} 0$ .

Let us prove the above lemma. We denote, for short,  $x_{i,j}^n$  by  $x_{ij}$  and introduce the diagonal matrix

$$\tilde{D}_n^\varepsilon := \text{diag} \left( \left( \tilde{\lambda}_n^\varepsilon(i) := \lambda_i + \frac{\varepsilon}{\sqrt{n}} x_{ii} \right)_{i=1, \dots, n} \right) \quad (5.11)$$

which is the diagonal part of the matrix  $D_n^\varepsilon$ . Note that with this notation and by using identity (5.1), the quantity we are interested in can be written:

$$\Pi_n(\varphi_z) = \varepsilon^{-2} \left( (z - D_n^\varepsilon)^{-1} - (z - \tilde{D}_n^\varepsilon)^{-1} \right)_{ii} - \Xi_{f(x)}(z). \quad (5.12)$$

To deal with this quantity we introduce the null diagonal matrix

$$\tilde{X}_n := \varepsilon^{-1} (D_n^\varepsilon - \tilde{D}_n^\varepsilon) = X_n - n^{-1/2} \text{diag}((x_{ii})_{i=1, \dots, n})$$

obtained by vanishing the diagonal of the matrix  $X$ .

A perturbative expansion of the resolvent of  $D_n^\varepsilon = \tilde{D}_n^\varepsilon + \varepsilon \tilde{X}_n$  yields

$$\begin{aligned} (z - D_n^\varepsilon)^{-1} - (z - \tilde{D}_n^\varepsilon)^{-1} &= \varepsilon (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - \tilde{D}_n^\varepsilon)^{-1} \\ &\quad + \varepsilon^2 (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - \tilde{D}_n^\varepsilon)^{-1} \\ &\quad + \varepsilon^3 (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - D_n^\varepsilon)^{-1}. \end{aligned} \quad (5.13)$$

We now want to analyze the corresponding expansion of  $\left( (z - D_n^\varepsilon)^{-1} - (z - \tilde{D}_n^\varepsilon)^{-1} \right)_{ii}$ .

**Claim 9.** *For all  $i \in \llbracket 1, n \rrbracket$ ,  $\left( (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - \tilde{D}_n^\varepsilon)^{-1} \right)_{ii} = 0$ .*

**Proof.** This comes from the fact that the matrix  $\tilde{X}_n$  has a null diagonal.  $\square$

**Claim 10.** *If, for all  $i \in \llbracket 1, n \rrbracket$ , we denote*

$$B_n(z, i) := \left( (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - \tilde{D}_n^\varepsilon)^{-1} \right)_{ii},$$

*then, for all  $x \in [0, 1]$ ,  $B_n(z, \lfloor nx \rfloor) - \Xi_{f(x)}(\varphi_z) = O_{L_2} \left( \left( \eta_n + \frac{1}{\sqrt{n}} \right) |\Im z|^{-3} \right)$ .*

**Proof.** With the notations of (5.11), the term  $B_n(z, i)$  writes

$$B_n(z, i) = \frac{1}{n} \sum_j \frac{|x_{ij}|^2}{\left( z - \tilde{\lambda}_n^\varepsilon(i) \right)^2 \left( z - \tilde{\lambda}_n^\varepsilon(j) \right)},$$

and, for  $x \in [0, 1]$ ,

$$\begin{aligned} \Xi_{f(x)}(\varphi_z) &= \int_{t \in \mathbb{R}} \frac{\tau(f(x), t) \rho(t)}{(t - f(x))^2} \left( \frac{1}{z - t} - \frac{1}{z - f(x)} + \frac{t - f(x)}{(z - f(x))^2} \right) dt \\ &= \int_{t \in \mathbb{R}} \frac{\tau(f(x), t)}{(z - f(x))^2 (z - t)} \rho(t) dt \end{aligned} \quad (5.14)$$

$$= \int_{y \in [0, 1]} \frac{\sigma(x, y)^2}{(z - f(x))^2 (z - f(y))} dy. \quad (5.15)$$

The difference of these quantities writes,

$$\begin{aligned} &B_n(z, \lfloor nx \rfloor) - \Xi_{f(x)}(\varphi_z) \\ &= \frac{1}{n} \sum_{j=1}^n \frac{|x_{\lfloor nx \rfloor, j}|^2 - \sigma_n^2 \left( \frac{\lfloor nx \rfloor}{n}, \frac{j}{n} \right)}{\left( z - \tilde{\lambda}_n^\varepsilon(\lfloor nx \rfloor) \right)^2 \left( z - \tilde{\lambda}_n^\varepsilon(j) \right)} \\ &\quad + \frac{1}{n} \sum_{j=1}^n \frac{\sigma_n^2 \left( \frac{\lfloor nx \rfloor}{n}, \frac{j}{n} \right)}{\left( z - \tilde{\lambda}_n^\varepsilon(\lfloor nx \rfloor) \right)^2 \left( z - \tilde{\lambda}_n^\varepsilon(j) \right)} - \frac{\sigma_n^2 \left( \frac{\lfloor nx \rfloor}{n}, \frac{j}{n} \right)}{(z - \lambda_{\lfloor nx \rfloor})^2 (z - \lambda_j)} \\ &\quad + \int_{y \in [0, 1]} \frac{\sigma_n^2 \left( \frac{\lfloor nx \rfloor}{n}, \frac{\lfloor ny \rfloor}{n} \right)}{(z - \lambda_{\lfloor nx \rfloor}^2)(z - \lambda_{\lfloor ny \rfloor})} dy - \int_{y \in [0, 1]} \frac{\sigma(x, y)^2}{(z - f(x))^2 (z - f(y))} dy. \end{aligned}$$

The  $L_2$  norm of the first line of the right hand side of the previous equality writes

$$\left\| \frac{1}{n} \sum_{j=1}^n \frac{|x_{ij}|^2 - \sigma_n(i/n, j/n)^2}{(z - \lambda_i)^2 (z - \lambda_j)} \right\|_{L_2} = \frac{1}{n} \left( \sum_{j=1}^n \frac{\mathbb{E}[|x_{ij}|^2 - \sigma_n(i/n, j/n)^2]^2}{|z - \lambda_i|^4 |z - \lambda_j|^2} \right)^{\frac{1}{2}} = O \left( \frac{1}{\sqrt{n} |\Im z|^3} \right),$$

the  $L_1$  norm of the second line is bounded, for  $C > 0$ , by

$$\frac{C}{n} \sum_{j=1}^n \mathbb{E} \left[ \frac{\varepsilon}{\sqrt{n}} \frac{|x_{\lfloor nx \rfloor, \lfloor nx \rfloor} + |x_{jj}|}{|\mathfrak{Im}(z)|^3} \right] = O \left( \frac{\varepsilon}{\sqrt{n} |\mathfrak{Im}(z)|^3} \right)$$

and, finally, from assumption (b), the third line is  $O(\eta_n |\mathfrak{Im} z|^{-3})$ .  $\square$

**Claim 11.** For all  $i \in \llbracket 1, n \rrbracket$ ,

$$\left( (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X} (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X} (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X} (z - D_n^\varepsilon)^{-1} \right)_{ii} = O_{L^2}(|\mathfrak{Im} z|^{-4})$$

**Proof.** By taking into account that the  $L^6$  norm of the entries of  $\sqrt{n}X$  is finite, and that the norms of the coefficients of  $(z - \tilde{D}_n^\varepsilon)^{-1}$  and of  $(z - D_n^\varepsilon)^{-1}$  are smaller than  $|\mathfrak{Im}(z)|^{-1}$ , we deduce that

$$\begin{aligned} & \mathbb{E} \left[ \left| \left( (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - D_n^\varepsilon)^{-1} \right)_{ii} \right|^2 \right]^{\frac{1}{2}} \\ &= \mathbb{E} \left[ \left| \sum_{j,k,l=1}^n ((z - \tilde{D}_n^\varepsilon)^{-1})_{i,i} (\tilde{X}_n)_{i,j} ((z - \tilde{D}_n^\varepsilon)^{-1})_{j,j} (\tilde{X}_n)_{j,k} ((z - \tilde{D}_n^\varepsilon)^{-1})_{k,k} (\tilde{X}_n)_{k,l} ((z - D_n^\varepsilon)^{-1})_{l,i} \right|^2 \right]^{\frac{1}{2}} \\ &\leq \frac{1}{|\mathfrak{Im}(z)|^4} \mathbb{E} \left[ \left| \sum_{j,k,l=1}^n (\tilde{X}_n)_{i,j} (\tilde{X}_n)_{j,k} (\tilde{X}_n)_{k,l} \right|^2 \right]^{\frac{1}{2}} \\ &\leq \frac{1}{|\mathfrak{Im}(z)|^4 n^{3/2}} \mathbb{E} \left[ \left| \sum_{j,k,l=1}^n x_{i,j} x_{j,k} x_{k,l} \right|^2 \right]^{\frac{1}{2}} \\ &\leq \frac{1}{|\mathfrak{Im}(z)|^4 n^{3/2}} \mathbb{E} \left[ \sum_{j,k,l,m,p,q=1}^n x_{i,j} x_{j,k} x_{k,l} \overline{x_{i,m}} \overline{x_{m,p}} \overline{x_{p,q}} \right]^{\frac{1}{2}}. \end{aligned}$$

Since the entries  $(x_{i,j})$  are independent and centered, the set of “edges”  $\{(i, m), (m, p), (p, q)\}$  must be equal to the set  $\{(i, j), (j, k), (k, l)\}$  in order to get a non zero term. Therefore, the complexity of the previous sum is  $O(n^3)$ . Note that other non zero terms involving third or fourth moments are much less numerous.

Therefore,

$$\mathbb{E} \left[ \left| \left( (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - \tilde{D}_n^\varepsilon)^{-1} \tilde{X}_n (z - D_n^\varepsilon)^{-1} \right)_{ii} \right|^2 \right]^{\frac{1}{2}} \leq \frac{1}{|\mathfrak{Im}(z)|^4}$$

$\square$

Gathering Formulas (5.12), (5.13) and Claims 1, 2 and 3, we prove Lemma 34.

### 5.3.2 From Stieltjes transform to $\mathcal{C}^5$ functions

Now, let  $\varphi$  be a  $\mathcal{C}^5$  function on  $\mathbb{R}$  with bounded fifth derivative and let us introduce the almost analytic extension of degree 5 of  $\varphi$  defined by

$$\forall z = x + iy \in \mathbb{C}, \quad \tilde{\varphi}_4(z) := \sum_{k=0}^4 \frac{1}{k!} (iy)^k \varphi^{(k)}(x).$$

An elementary computation gives, by successive cancellations, that

$$\bar{\partial} \tilde{\varphi}_4(z) = \frac{1}{2} (\partial_x + i\partial_y) \tilde{\varphi}_4(x + iy) = \frac{1}{2 \times 4!} (iy)^4 \varphi^{(5)}(x). \quad (5.16)$$

Furthermore, by Helffer-Sjöstrand formula [BGE17, Propo. 9], for  $\chi \in \mathcal{C}_c^\infty(\mathbb{C}; [0, 1])$  a smooth cutoff function with value one on the support of  $\varphi$ ,

$$\varphi(\cdot) = -\frac{1}{\pi} \int_{\mathbb{C}} \frac{\bar{\partial}(\tilde{\varphi}_4(z)\chi(z))}{y^4} y^4 \varphi_z(\cdot) d^2z$$

where  $d^2z$  denotes the Lebesgue measure on  $\mathbb{C}$ .

Note that by (5.16),  $z \mapsto \mathbb{1}_{y \neq 0} \frac{\bar{\partial}(\tilde{\varphi}_4(z)\chi(z))}{y^4}$  is a continuous compactly supported function and that  $z \in \mathbb{C} \mapsto \mathbb{1}_{y \neq 0} y^4 \varphi_z \in \mathcal{C}_b^1$  is continuous, hence,

$$\Pi_n(\varphi) = \frac{1}{\pi} \int_{\mathbb{C}} \frac{\bar{\partial}(\tilde{\varphi}_4(z)\chi(z))}{y^4} y^4 \Pi_n(\varphi_z) d^2z.$$

Therefore, using the Cauchy-Schwarz inequality and the fact that  $\chi$  has compact support at the second step, for a certain constant  $C$ , we have

$$\begin{aligned} \mathbb{E} (|\Pi_n(\varphi)|^2) &= \mathbb{E} \left( \left| \frac{1}{\pi} \int_{\mathbb{C}} \frac{\bar{\partial}(\tilde{\varphi}_4(z)\chi(z))}{y^4} y^4 \Pi_n(\varphi_z) d^2z \right|^2 \right) \\ &\leq C \mathbb{E} \left( \int_{\mathbb{C}} \left| \frac{\bar{\partial}(\tilde{\varphi}_4(z)\chi(z))}{y^4} y^4 \Pi_n(\varphi_z) \right|^2 d^2z \right) \\ &= C \int_{\mathbb{C}} \left| \frac{\bar{\partial}(\tilde{\varphi}_4(z)\chi(z))}{y^4} \right|^2 y^8 \mathbb{E} (|\Pi_n(\varphi_z)|^2) d^2z. \end{aligned}$$

By (5.16), the function  $\left| \frac{\bar{\partial}(\tilde{\varphi}_5(z)\chi(z))}{y^5} \right|^2$  is continuous and compactly supported and bounded by  $C \|\varphi^{(5)}\|_\infty^2$  for some constant  $C$ . Besides, by Lemma 34, uniformly in  $z$ ,

$$y^8 \mathbb{E} (|\Pi_n(\varphi_z)|^2) = O \left( (1 + y^2) \left( \eta_n^2 + \frac{1}{n} + \varepsilon^2 \right) \right).$$

We deduce that

$$\mathbb{E} (|\Pi_n(\varphi)|^2) \leq C \int_{\mathbb{C}} \left| \frac{\bar{\partial}(\tilde{\varphi}_4(z)\chi(z))}{y^4} \right|^2 y^8 \mathbb{E} (|\Pi_n(\varphi_z)|^2) d^2z = O \left( \|\varphi^{(5)}\|_\infty^2 \left( \eta_n^2 + \frac{1}{n} + \varepsilon^2 \right) \right),$$

which closes the proof of Theorem 32.

## 5.4 Consequence for the eigenvectors

In this section we want to present some concrete applications of our main result and some numerical simulations that comfort well our developments. Therefore, some of the following considerations will be at a heuristic level.

Let  $x_0 \in [0, 1]$  and  $t \in f([0, 1])$ , with  $t \neq f(x_0)$ . Let us assume that the function  $f$  is invertible (which significantly lessens the generality of the model).

If we denote by  $\mu_n^\varepsilon$  the empirical spectral measure of  $D_n^\varepsilon$ , namely

$$\mu_n^\varepsilon := \frac{1}{n} \sum_{j=1}^n \delta_{\lambda_j^\varepsilon},$$

the definition of  $\mu_{n, \mathbf{e}_i}^\varepsilon$  implies, for  $t \neq f(x_0)$ ,

$$\mu_{n, \mathbf{e}_i}^\varepsilon(dt) \approx n |\langle \mathbf{u}_{[nf^{-1}(t)]}^\varepsilon, \mathbf{e}_{[nx_0]} \rangle|^2 \mu_n^\varepsilon(dt).$$

In addition, Formulas (5.7) and (5.6) give, for  $t \neq f(x_0)$ , the following expansion

$$\mu_{n, \mathbf{e}_i}^\varepsilon \approx \varepsilon^2 \frac{\tau(f(x_0), t) \rho(t)}{(t - f(x_0))^2} dt.$$

Keeping in mind the convergence of  $\mu_n^\varepsilon(t)$  towards the measure  $\rho(t)dt$ , we conclude that we have the asymptotic equality, in the sense of distributions, between functions of  $t$ ,

$$\varepsilon^{-2} n |\langle \mathbf{u}_{[nf^{-1}(t)]}^\varepsilon, \mathbf{e}_{[nx_0]} \rangle|^2 \approx \frac{\tau(f(x_0), t)}{(t - f(x_0))^2}. \quad (5.17)$$

We present now two simulations (displayed in Figures 5.1 and 5.2) which show a good matching with this theoretical prediction. First we consider the case where the deterministic matrix  $D_n$  is perturbed by a Gaussian Wigner matrix,  $X_n$ . More precisely, we take for  $D_n$  the diagonal matrix with  $\frac{i}{n}$  as  $i^{th}$  entry, so that  $f(x) = x$  and the density  $\rho$  is equal  $x \mapsto \mathbb{1}_{[0,1]}(x)$ . The entries of the perturbing matrix  $X_n$  are all Gaussian and independent with variance one. Then, we consider the case where the same matrix  $D_n$  is perturbed by a band matrix. In other words, we consider now that  $\sigma(x, y) = \mathbb{1}_{|x-y| \leq \ell}$ , where  $\ell \in [0, 1]$  is the relative width of the band. Note that in this second example, even though there is absolutely no deterministic reason why  $\langle \mathbf{u}_{[ny]}^\varepsilon, \mathbf{e}_{[nx]} \rangle$  would vanish when  $|y - x| > \ell$ , we see that at first order, it is actually almost zero (Figure 5.2). This is related to the question of the localization of the eigenvectors of band random matrices (see e.g. [Cha10, EKYY13, EK11b, EK11a, EYY12b, EYY12a, FM91, Sch09]).




Figure 5.1: **Uniform measure perturbation by a Wigner matrix.** The red curve represents (part of) the function  $t \in [0, 1] \mapsto \varepsilon^{-1} \sqrt{n} |\langle \mathbf{u}_{[nf^{-1}(t)]}^\varepsilon, \mathbf{e}_{[nx_0]} \rangle|$ , which results from the numeric computation of the eigenvectors of  $D_n^\varepsilon$  and the blue curve represents our theoretical prediction  $t \mapsto |t - x_0|^{-1}$ . As (5.17) is an equality between distributions, we have smoothed both curves thanks to a simple moving average with window length  $\sqrt{n}$ . Here  $n = 10^4$ ,  $\varepsilon = n^{-0.7}$  and  $x_0 = \frac{1}{2}$ .




Figure 5.2: **Uniform measure perturbation by a Gaussian band matrix.** The red curve represents (part of) the function  $t \in [0, 1] \mapsto \varepsilon^{-1} \sqrt{n} |\langle \mathbf{u}_{[nf^{-1}(t)]}^\varepsilon, \mathbf{e}_{[nx_0]} \rangle|$ , which results from the numeric computation of the eigenvectors of  $D_n^\varepsilon$  and the blue curve represents our theoretical prediction  $t \mapsto \mathbb{1}_{|t - x_0| \leq \ell} |t - x_0|^{-1}$ . As (5.17) is an equality between distributions, we have smoothed both curves thanks to a simple moving average with window length  $\sqrt{n}$ . Here  $n = 10^4$ ,  $\ell = 0.1$ ,  $\varepsilon = n^{-0.7}$  and  $x_0 = \frac{1}{2}$ .

**Remark 5.** It could be interesting to compare (5.17) with the recent work [Ben17] of Benigni. Corollary 1.6 of [Ben17] precisely implies the same estimate as the one we give in (5.17). More precisely, it says that if the distribution of the eigenvalues of  $D_n$  is regular enough (in our case, these conditions can be expressed as a bound on our error term  $\eta_n$

of (5.2)), if  $X_n$  is a Wigner matrix (in our case, it amounts to suppose that  $\tau \equiv 1$ ), if  $\varepsilon$  satisfies, roughly,  $n^{-1} \ll \varepsilon \ll 1$  and if  $\gamma_{j,\varepsilon}$  denotes the quantile of the density  $\rho^\varepsilon$  of the free convolution of the spectral distribution of  $D_n$  with a semicircle law with variance  $\varepsilon^2$ , i.e.

$$\int_{-\infty}^{\gamma_{j,\varepsilon}} \rho^\varepsilon(x) dx = \frac{j}{n},$$

then the approximation

$$\varepsilon^{-2} n |\langle \mathbf{u}_{j_0}^\varepsilon, \mathbf{e}_i \rangle|^2 \approx \frac{1}{(\lambda_i - \gamma_{j_0,\varepsilon})^2 + O(\varepsilon^4)},$$

holds when empirical means on  $i$  in the neighborhood of a fixed  $j_0$  have been taken. We recover the first order approximation from (5.17). Note however that here, empirical means are taken on  $i$ , whereas in our result, with these notations, they are taken on  $j_0$ .

# List of Figures

1.1	Illustration of the Wigner's semi-circle law . . . . .	17
1.2	Illustration of the Girko's circle law . . . . .	18
1.3	Illustration of the Marchenko-Pastur's quarter circle law . . . . .	19
1.4	Illustration of the Single Ring Theorem . . . . .	21
1.5	Relation of the Single Ring Theorem to other results . . . . .	21
1.6	Illustration of Anderson's transition . . . . .	23
1.7	Illustration of a finite rank perturbation . . . . .	26
1.8	Illustration of the phase transition phenomenon . . . . .	27
1.9	Effect of a perturbation on an eigenvector . . . . .	28
3.1	Deforming the uniform distribution by a band matrix . . . . .	44
3.2	Triangular pulse perturbation by a Wigner matrix: density and cumulative distribution function . . . . .	45
3.3	Triangular pulse perturbation by a Wigner matrix: Stieltjes transform . . .	45
3.4	Parabolic pulse perturbation by a Wigner matrix . . . . .	46
4.1	A Dyck path and its corresponding rooted planar tree . . . . .	69
4.2	Three different labeled planar rooted trees . . . . .	69
4.3	Parabolic pulse perturbation by a Wigner matrix . . . . .	73
4.4	Uniform measure perturbation by a Gaussian band matrix . . . . .	75
5.1	Uniform measure perturbation by a Wigner matrix . . . . .	97
5.2	Uniform measure perturbation by a Gaussian band matrix . . . . .	97



## LIST OF FIGURES

---

# Bibliography

- [AB12] Romain Allez and Jean-Philippe Bouchaud. Eigenvector dynamics: general theory and some applications. *Physical Review E*, 86(4):046202, 2012. (cited on pages 27, 28, 39, 66, and 88.)
- [AB14] Romain Allez and Jean-Philippe Bouchaud. Eigenvector dynamics under free addition. *Random Matrices: Theory and Applications*, 3(03):1450010, 2014. (cited on pages 27, 28, 39, and 88.)
- [ABB14] Romain Allez, Joël Bun, and Jean-Philippe Bouchaud. The eigenvectors of gaussian matrices with an external source. *arXiv preprint arXiv:1412.7108*, 2014. (cited on pages 27, 39, 66, and 88.)
- [AZ06] Greg W Anderson and Ofer Zeitouni. A clt for a band matrix model. *Probability Theory and Related Fields*, 134(2):283–338, 2006. (cited on page 46.)
- [BABP16] Joël Bun, Romain Allez, Jean-Philippe Bouchaud, and Marc Potters. Rotational invariant estimator for general noisy matrices. *IEEE Transactions on Information Theory*, 62(12):7475–7490, 2016. (cited on page 39.)
- [BAP<sup>+</sup>05] Jinho Baik, Gérard Ben Arous, Sandrine Péché, et al. Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices. *The Annals of Probability*, 33(5):1643–1697, 2005. (cited on page 26.)
- [BD<sup>+</sup>13] Anirban Basak, Amir Dembo, et al. Limiting spectral distribution of sum of unitary and orthogonal matrices. *Electronic Communications in Probability*, 18, 2013. (cited on page 20.)
- [Ben17] Lucas Benigni. Eigenvectors distribution and quantum unique ergodicity for deformed wigner matrices. *arXiv preprint arXiv:1711.07103*, 2017. (cited on pages 27, 28, 88, and 97.)
- [BES16] Zhigang Bao, László Erdős, and Kevin Schnelli. Local single ring theorem on optimal scale. *arXiv preprint arXiv:1612.05920*, 2016. (cited on page 20.)
- [BG<sup>+</sup>17] Florent Benaych-Georges et al. Local single ring theorem. *The Annals of Probability*, 45(6A):3850–3885, 2017. (cited on page 20.)

- [BGEM17] Florent Benaych-Georges, Nathanaël Enriquez, and Alkéos Michaël. Empirical spectral distribution of a matrix under perturbation. *Journal of Theoretical Probability*, pages 1–32, 2017. (cited on pages 30, 66, 85, 88, and 95.)
- [BGEM18] Florent Benaych-Georges, Nathanaël Enriquez, and Alkéos Michaël. Eigenvectors of a matrix under random perturbation. *arXiv preprint arXiv:1801.10512*, 2018. (cited on page 30.)
- [BGGM<sup>+</sup>11] Florent Benaych-Georges, Alice Guionnet, Mylène Maida, et al. Fluctuations of the extreme eigenvalues of finite rank deformations of random matrices. *Electronic Journal of Probability*, 16:1621–1662, 2011. (cited on page 26.)
- [BGGM14] Florent Benaych-Georges, Alice Guionnet, and Camille Male. Central limit theorems for linear statistics of heavy tailed random matrices. *Communications in Mathematical Physics*, 329(2):641–686, 2014. (cited on page 33.)
- [BGK16] Florent Benaych-Georges and Antti Knowles. Lectures on the local semicircle law for wigner matrices. *arXiv preprint arXiv:1601.04055*, To appear in *SMF series Panoramas et Synthèses*, 2016. (cited on pages 32, 57, 70, and 91.)
- [BGN11] Florent Benaych-Georges and Raj Rao Nadakuditi. The eigenvalues and eigenvectors of finite, low rank perturbations of large random matrices. *Advances in Mathematics*, 227(1):494–521, 2011. (cited on page 26.)
- [Bia97a] Philippe Biane. Free brownian motion, free stochastic calculus and random matrices, in free probability theory. *Fields Inst. Commun.*, 12:1–19, 1997. (cited on page 47.)
- [Bia97b] Philippe Biane. On the free convolution with a semi-circular distribution. *Indiana University Mathematics Journal*, pages 705–718, 1997. (cited on page 47.)
- [BS98] Philippe Biane and Roland Speicher. Stochastic calculus with respect to free brownian motion and analysis on wigner space. *Probability theory and related fields*, 112(3):373–409, 1998. (cited on page 47.)
- [BS01] Philippe Biane and Roland Speicher. Free diffusions, free entropy and free fisher information. In *Annales de l’Institut Henri Poincaré (B) Probability and Statistics*, volume 37, pages 581–606. Elsevier, 2001. (cited on page 47.)
- [BY86] ZD Bai and YQ Yin. Limiting behavior of the norm of products of random matrices and two problems of geman-hwang. *Probability theory and related fields*, 73(4):555–569, 1986. (cited on page 16.)
- [CG93a] G Casati and V Girko. Generalized wigner law for band random matrices. *Random Operators and Stochastic Equations*, 1(3):279–286, 1993. (cited on page 46.)

- [CG93b] G Casati and V Girko. Wigner’s semicircle law for band random matrices. *Random Operators and Stochastic Equations*, 1(1):15–22, 1993. (cited on page 46.)
- [Cha10] Djalil Chafaï. Anderson localization, blog note, url: <http://djalil.chafai.net/blog/2010/09/07/anderson-localization/>, 2010. (cited on pages 23 and 96.)
- [Dub09] Julien Dubédat. Sle and the free field: partition functions and couplings. *Journal of the American Mathematical Society*, 22(4):995–1054, 2009. (cited on pages 38 and 67.)
- [DVDN92] DV-Voiculescu, KJ Dykema, and Alexandru Nica. *Free Random Variables: A Noncommutative Probability Approach to Free Products with Applications to Random Matrices, Operator Algebras, and Harmonic Analysis on Free Groups*. American Mathematical Society, 1992. (cited on page 14.)
- [Dys62] Freeman J. Dyson. The threefold way. algebraic structure of symmetry groups and ensembles in quantum mechanics. *Journal of Mathematical Physics*, 3(6):1199–1215, 1962. (cited on page 14.)
- [EK11a] László Erdős and Antti Knowles. Quantum diffusion and delocalization for band matrices with general distribution. In *Annales Henri Poincaré*, volume 12, page 1227. Springer, 2011. (cited on page 96.)
- [EK11b] László Erdős and Antti Knowles. Quantum diffusion and eigenfunction delocalization in a random band matrix model. *Communications in Mathematical Physics*, 303(2):509–554, 2011. (cited on page 96.)
- [EKYY13] László Erdős, Antti Knowles, Horng-Tzer Yau, and Jun Yin. Delocalization and diffusion profile for random band matrices. *Communications in Mathematical Physics*, 323(1):367–416, 2013. (cited on pages 23 and 96.)
- [EYY12a] László Erdős, Horng-Tzer Yau, and Jun Yin. Bulk universality for generalized wigner matrices. *Probability Theory and Related Fields*, 154(1-2):341–407, 2012. (cited on pages 23 and 96.)
- [EYY12b] László Erdős, Horng-Tzer Yau, and Jun Yin. Rigidity of eigenvalues of generalized wigner matrices. *Advances in Mathematics*, 229(3):1435–1515, 2012. (cited on page 96.)
- [Fer00] Francisco M Fernández. *Introduction to perturbation theory in quantum mechanics*. CRC press, 2000. (cited on pages 27 and 39.)
- [FK81] Zoltán Füredi and János Komlós. The eigenvalues of random symmetric matrices. *Combinatorica*, 1(3):233–241, 1981. (cited on pages 14 and 26.)

## BIBLIOGRAPHY

---

- [FM91] Yan V Fyodorov and Alexander D Mirlin. Scaling properties of localization in random band matrices: a  $\sigma$ -model approach. *Physical review letters*, 67(18):2405, 1991. (cited on page 96.)
- [FZ97] Joshua Feinberg and A Zee. Non-gaussian non-hermitian random matrix theory: phase transition and addition formalism. *Nuclear Physics B*, 501(3):643–669, 1997. (cited on page 20.)
- [Gin65] Jean Ginibre. Statistical ensembles of complex, quaternion, and real matrices. *Journal of Mathematical Physics*, 6(3):440–449, 1965. (cited on page 17.)
- [Gir85] Vyacheslav L Girko. Circular law. *Theory of Probability & Its Applications*, 29(4):694–706, 1985. (cited on page 17.)
- [GKZ11] Alice Guionnet, Manjunath Krishnapur, and Ofer Zeitouni. The single ring theorem. *Annals of Mathematics*, 174(2):1189–1217, 2011. (cited on page 20.)
- [Gui08] Alice Guionnet. Large random matrices: Lectures on macroscopic asymptotics, 2008. (cited on page 68.)
- [Hör03] Lars Hörmander. The analysis of linear partial differential operators. i. distribution theory and fourier analysis. reprint of the second (1990) edition, 2003. (cited on pages 64 and 85.)
- [HS89] Bernard Helffer and J Sjöstrand. Equation de schrödinger avec champ magnétique et équation de harper. In *Schrödinger operators*, pages 118–197. Springer, 1989. (cited on pages 85 and 91.)
- [Joh01] Iain M. Johnstone. On the distribution of the largest eigenvalue in principal components analysis. *Ann. Statist.*, 29(2):295–327, 2001. (cited on page 26.)
- [Kat13] Tosio Kato. *Perturbation theory for linear operators*, volume 132. Springer Science & Business Media, 2013. (cited on page 24.)
- [Kin09] Frederick W. King. *Hilbert Transforms*, volume 2 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, 2009. (cited on pages 44 and 45.)
- [LP11] Olivier Ledoit and Sandrine Pécché. Eigenvectors of some large sample covariance matrix ensembles. *Probability Theory and Related Fields*, 151(1-2):233–264, 2011. (cited on pages 27 and 39.)
- [MG60] M.L. Mehta and M. Gaudin. On the density of eigenvalues of a random matrix. *Nuclear Physics*, 18:420 – 427, 1960. (cited on page 14.)
- [Mon73] Hugh L Montgomery. The pair correlation of zeros of the zeta function. In *Proc. Symp. Pure Math*, volume 24, pages 181–193, 1973. (cited on page 14.)

- [MP67] Vladimir A Marčenko and Leonid Andreevich Pastur. Distribution of eigenvalues for some sets of random matrices. *Mathematics of the USSR-Sbornik*, 1(4):457, 1967. (cited on pages 14 and 18.)
- [OVW16] Sean O’Rourke, Van Vu, and Ke Wang. Eigenvectors of random matrices: a survey. *Journal of Combinatorial Theory, Series A*, 144:361–442, 2016. (cited on pages 27 and 88.)
- [OVW17] Sean O’Rourke, Van Vu, and Ke Wang. Random perturbation of low rank matrices: Improving classical bounds. *Linear Algebra and its Applications*, 2017. (cited on pages 27 and 88.)
- [RV14] Mark Rudelson and Roman Vershynin. Invertibility of random matrices: unitary and orthogonal perturbations. *Journal of the American Mathematical Society*, 27(2):293–338, 2014. (cited on page 20.)
- [Sch09] Jeffrey Schenker. Eigenvector localization for random band matrices with power law band width. *Communications in Mathematical Physics*, 290(3):1065–1097, 2009. (cited on pages 23 and 96.)
- [Shl96] Dimitri Shlyakhtenko. Random gaussian band matrices and freeness with amalgamation. *International Mathematics Research Notices*, 1996(20):1013–1025, 1996. (cited on pages 46, 47, 59, and 76.)
- [ST10] Mariya Shcherbina and Brunello Tirozzi. Central limit theorem for fluctuations of linear eigenvalue statistics of large random graphs. *Journal of Mathematical Physics*, 51(2):023523, 2010. (cited on pages 33 and 85.)
- [TV10] Terence Tao and Van Vu. Random matrices: Universality of esds and the circular law. *The Annals of Probability*, pages 2023–2065, 2010. (cited on page 17.)
- [Voi91] Dan Voiculescu. Limit laws for random matrices and free products. *Inventiones mathematicae*, 104(1):201–220, 1991. (cited on page 14.)
- [vSW17] Per von Soosten and Simone Warzel. Non-ergodic delocalization in the rosenzweig-porter model. *arXiv preprint arXiv:1709.10313*, 2017. (cited on pages 27 and 88.)
- [Wey12] H. Weyl. Das asymptotische verteilungsgesetz der eigenwerte linearer partieller differentialgleichungen (mit einer anwendung auf die theorie der hohlraumstrahlung). *Mathematische Annalen*, 71:441–479, 1912. (cited on page 24.)
- [Wig55] Eugene P. Wigner. Characteristic vectors of bordered matrices with infinite dimensions. *Annals of Mathematics*, 62(3):548–564, 1955. (cited on page 14.)
-

## BIBLIOGRAPHY

---

- [Wis28] John Wishart. The generalised product moment distribution in samples from a normal multivariate population. *Biometrika*, 20(1/2):32–52, 1928. (cited on page 13.)
- [WW95] Michael Wilkinson and Paul N Walker. A brownian motion model for the parameter dependence of matrix elements. *Journal of Physics A: Mathematical and General*, 28(21):6143, 1995. (cited on page 39.)
- [Zho17] Yiqiao Zhong. Eigenvector under random perturbation: A nonasymptotic rayleigh-schrödinger theory. *arXiv preprint arXiv:1702.00139*, 2017. (cited on pages 27, 28, and 88.)

---

## ABSTRACT

---

The present thesis is devoted to the study of the effect of a perturbation on the spectrum of a Hermitian matrix by a random matrix with small operator norm and whose entries in the eigenvector basis of the first one were independent, centered and with a variance profile. This is carried out through perturbative expansions of various types of spectral laws of the considered perturbed large matrices. First, we demonstrate different perturbative expansions of the empirical spectral measure in the cases of the perturbative regime and the semi-perturbative regime and highlight well known heuristic patterns in Physics, as the transition between semi-perturbative and perturbative regimes. Secondly, we provide a thorough study of the semi-perturbative regime and prove the new fact that this regime could be decomposed into infinitely many sub-regimes. Finally, we prove, through a perturbative expansion of spectral measures associated to the state defined by a given vector, a perturbative expansion of the coordinates of the eigenvectors of the perturbed matrices.

---

## RÉSUMÉ

---

La présente thèse est consacrée à l'étude de l'effet d'une perturbation sur le spectre d'une matrice hermitienne perturbée par une matrice aléatoire de petite norme opérateur et dont les entrées dans la base propre de la première matrice sont indépendantes, centrées et possèdent un profil de variance. Ceci est réalisé au travers de développements perturbatifs de divers types des lois spectrales des grandes matrices perturbées considérées. Dans un premier temps, nous démontrons différents développements perturbatifs de la mesure spectrale empirique dans les cas du régime perturbatif et du régime semi-perturbatif et mettons en évidence des modèles heuristiques bien connus en physique, comme la transition entre les régimes semi-perturbatifs et perturbatifs. Dans un deuxième temps, nous proposons une étude approfondie du régime semi-perturbatif et prouvons le fait nouveau que ce régime peut être décomposé en un nombre infini de sous-régimes. Enfin, nous démontrons, au travers d'un développement perturbatif des mesures spectrales associées à un vecteur donné, un développement perturbatif des coordonnées des vecteurs propres des matrices perturbées que nous considérons.