

Patient safety in medical humanitarian action: medical error prevention and management

Jean-Marc Biquet

► To cite this version:

Jean-Marc Biquet. Patient safety in medical humanitarian action : medical error prevention and management. Santé publique et épidémiologie. Université de Lyon; Université de Genève, 2020. English. NNT : 2020LYSE1038 . tel-02884986

HAL Id: tel-02884986 https://theses.hal.science/tel-02884986

Submitted on 30 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



N°d'ordre NNT : 2020LYSE1038

THESE de DOCTORAT DE L'UNIVERSITE DE LYON

opérée au sein de l'Université Claude Bernard Lyon 1

Ecole Doctorale N° EA7425 - QUALITÉ SÉCURITÉ PERFORMANCE EN SANTÉ (HESPER)

Spécialité de doctorat :

Discipline : Science de la Santé

Soutenue publiquement le 20/02/2020, par :

Jean-Marc Biquet

PATIENT SAFETY IN

MEDICAL HUMANITARIAN ACTION.

Medical error prevention and management

Devant le jury composé de :

André, Flahaut. Professeur des Universités. Université de Genève. Président Migeot, Virginie. Professeure des Universités - Praticien Hospitalier Université de Poitiers. Rapporteur François, Patrice. Professeure des Universités - Praticien Hospitalier Université de Grenoble. Rapporteur Sprumont, Dominique. Professeur des Universités. Université de Neuchâtel. Rapporteur Calmy, Alexandra Professeur Associé, Université de Genève. Examinatrice Cyrille, Colin. Professeur des Universités - Praticien Hospitalier Université de Lyon. Examinateur Philippe, Michel. Professeure des Universités - Praticien Hospitalier Université de Lyon. Co-directeur de thèse Blanchet Karl. Professeur des Universités. Université de Genève. Co-Directeur de thèse Doris Schopper. Professeure retraitée. Université de Genève. Invitée

Acknowledgments

Thanks are due, first of all, to the members of my doctoral committee for their patience and support.

I am particularly grateful to those who, from the sidelines, have been an unwavering help and to those who had to bear with me as I invested my time and energy in this confounded undertaking.

The two oversight institutions, the University of Geneva's Institute of Global Health and Claude Bernard University, Lyon 1, both deserve special mention for having believed in my project and for having accepted me within their doctoral program.

I should also like to pay tribute to all humanitarian workers who, relentlessly, in spite of the difficulties and risks that their work involves, try to bring effective aid to their contemporaries. I hope that this modest labour on my part will be able to contribute to making them even more effective... Keywords: Medical error, Patient safety, Humanitarian Medicine, Safety culture, safety risks

Mots-clés : Erreur médicale, Sécurité du patient, Médecine Humanitaire, Culture de la sécurité, Risque de Sécurité

1. List of Publications

Published articles

Article 1

<u>Title:</u> 'A Call for the Application of Patient Safety Culture in Medical Humanitarian Action: A Literature Review'

Authors:

Jean-Marc Biquet, MSc, Institute of Global health, University of Geneva; Claude Bernard University, Lyon 1, HESPER EA 7425, F-69008 Lyon, France

Doris Schopper, PhD, MD, Medical Faculty, University of Geneva

Dominique Sprumont, PhD, Institute of Health Law, University of Neuchâtel

Philippe Michel, PhD, MD, Hospices Civils de Lyon, DOQRU, Lyon, F-69003, France ; Claude Bernard Université, Lyon 1, HESPER EA 7425, F-69008 Lyon, France

This article was submitted to and published in the Journal of Patient Safety in March 2020.

Article 2

<u>Title: '</u>Knowledge, Attitudes and Expectations of Medical Staff toward Medical Error Management Policies in Humanitarian Medicine: A Qualitative Study'

Authors:

Jean-Marc Biquet, MSc, Institute of Global health, University of Geneva; Claude Bernard University, Lyon 1, HESPER EA 7425, F-69008 Lyon, France

Doris Schopper, PhD, MD, Medical Faculty, University of Geneva

Dominique Sprumont, PhD, Institute of Health Law, University of Neuchâtel

Philippe Michel, PhD, MD, Hospices Civils de Lyon, DOQRU, Lyon, F-69003, France ; Claude Bernard Université, Lyon 1, HESPER EA 7425, F-69008 Lyon, France This article was submitted to and accepted for publication in the Journal of Patient Safety with minor modifications January 2020.

Oral Presentation:

Title: 'Medical Humanitarian Action and Patient Safety: The End of an Exception?'

Author:

Jean-Marc Biquet, jean-marc.biquet@unige.ch

This presentation (see Annex 1) was made at the Geneva Health Forum on 19 April 2016 in a parallel event organized by J.-M. Biquet and titled 'Quality of Care after Sudden Onset Disasters: Good Will Is Not Enough'.

Under Preparation

<u>Symposium</u>

Title: 'Patient safety: Management of medical errors in Medical Humanitarian Action'

Foreseen for Autumn 2020, the symposium aims at opening the debate and sharing the experiences in the medical humanitarian sector on patient safety, with a focus on medical error management. Expected partners for the symposium: CERAH, WHO, MSF, ICRC

Training

Title: 'Patient safety: Management of medical errors in Medical Humanitarian Action'

Thematic short Course organised (spring-summer 2020) by CERAH in partnership with MSF and ICRC aiming at training the medical manager of medical humanitarian action on management of medical errors in the field

2. Abstract in English

Patient safety has been recognized for some 20 years as one of the essential elements of healthcare quality¹ and has become an integral part of healthcare systems – at least in OCDE countries. It encompasses regulations, tools and strategies that affect all sectors of medicine. Today, research and implementation in the area of patient safety pertain above all to healthcare systems in the most developed countries whereas two thirds of estimated safety incidents occur in low- or mid-income countries.²

An exploratory phase aiming at developing the research strategy was made through discussions during informal meetings with international patient security specialists as well as with actors in the humanitarian sector. It confirmed that patient safety, per se, and the detection and management of medical errors have not yet been translated into the humanitarian assistance sector in a structured and adapted way. In order to understand the reasons for this gap in relation to criteria of healthcare quality as defined in the healthcare systems of countries where the headquarters of major international medical organisations are based, this research aimed to understand what the current status and perspectives of patient safety in medical humanitarian action are. An initial phase explored developments in the knowledge of safety and risk management that have dealt with the phenomenon of error and have developed theories and models regarding risk management at the origin of patient safety. Then, the current state of knowledge and the main developments in patient safety and especially medical error management were explored, before moving on to current limits and the stakes of implementation. This research on Google Search (Keywords: Patient safety and risk management) was followed by a rapid literature review on PubMed (Keywords: Patient safety, medical error, adverse event, review, systematic review, since 2013). 104 articles on the 245 references found, met the criteria and were analysed. In total a database of more than 1,200 articles has been established. In parallel, the characteristics of medical action as carried out by medical humanitarian organisations were analysed through a systematic literature review on PubMed and Embase databases. The searched keywords were: Patient safety, medical error, humanitarian, since 2000. 39 articles on the 308 references retrieved, met the inclusion criteria. This part of the research was the subject of a first article accepted by the "journal of Patient Safety" (in print).

The lessons learned in patient safety practises in OCDE healthcare systems show the necessary conjunction of action at three levels: at the macro level, regulations, the determination of objectives and a clear legal framework; at the micro level, personnel engaged in healthcare and its management to be sensitised, trained and directly involved; and at the meso level, healthcare institutions to be catalysts of patients safety owing to their commitment and leadership. It is at this level that the tools, processes and a favourable environment are created, institutions being responsible for establishing error-detection systems and a participatory management framework for correction and learning.

The second phase of our research was centred on semi-directive discussions with medical and paramedical personnel active within six medical humanitarian organisations. The selected organisations respected the following criteria: International humanitarian organisations based in an OECD country; active throughout the world with national and international staff; having a medical department implementing the organisation's medical policies; and whose medical action was, in terms of budget, the major area of activities (with the exception of ICRC, historically a medical organisation having diversified its portfolio of activities). A first interview was done with the medical director or the responsible of the quality of care of those organisations. Then, 39 interviews were done by Skype or face to face with medical and paramedical staff of those organisations responding to the following criteria: international staff with minimum 2 years of experience in the humanitarian sector. 36 interviews were analysed, the others being of too poor quality of recording to be transcribed effectively. The selected interviews integrally transcribed were coded and analysed by themes and sub-themes (Atlas ti ©). The objective was to understand the knowledge, attitudes and practises of these personnel with regards to patient safety and medical error management as well as their

expectations regarding the needs, as they perceived them, to integrate the preoccupation with patient safety and risk management into institutional priorities within their sector. The results of this research were reported in a second article presented and accepted with minor modifications by the "Journal of Patient Safety".

It appeared clearly that, while there may not yet be a structured approach in the sector regarding patient safety and, specifically, medical error management, this clearly corresponds to an expectation on the part of the humanitarian personnel interviewed. The reasons cited to explain this lack fell into roughly two groups. First, the specificity of medical humanitarian action such as the mind-set and mode of operation oriented to urgency (tendency to move from one priority to another depending on the level of urgency) as well as national and international personnel encompassing highly varied medical cultures. To this one might add the absence of a regulatory system in the sector as well as external pressures pushing toward investments in safer healthcare provision, and, finally, a great diversity of activities and intervention contexts as well as in the level of control of the humanitarian organisations regarding the care dispensed. Second, there are factors common to those found in OCDE country healthcare systems such as the absence of the ingredients of a safety culture: leadership; a learning, rather than a blaming, culture; appropriate support of personnel regarding sensitisation, training and development of tools for reporting and analysing medical error as well as clear guidance for the management of consequences.

Regarding the expectation of the persons interviewed, while they all acknowledged their role in the implementation of safety for their patients, they clearly identified the responsibility of their organisations to invest in the means of translating into practice the desire for a better awareness of patient safety. To achieve this, they call for collaboration among organisations to share experiences and to develop tools of implementation.

This research, to our knowledge the first of its kind, demonstrates the eagerness of the medical and paramedical staff engaged in humanitarian action to commit to an internal cultural revolution towards a safer healthcare provision, even in precarious situations.

Catching up the delays in adopting adapted patient safety and medical error management policies would reinforce the accountability to the vulnerable populations assisted by these organisations and save more lives, the essence of humanitarian purpose.

3. Abstract in French

La sécurité des patients est reconnue depuis une vingtaine comme un des éléments essentiels de la qualité des soins¹et est devenu une partie intégrante des systèmes de santé, tout au moins dans les pays de l'OCDE. Elle se déclinée en règlementations, outils et stratégies qui touchent tous les secteurs de la médecine. Aujourd'hui les recherches et applications de la sécurité des patients concernent surtout les systèmes de santé des pays les plus développés alors même que deux-tiers des incidents de sécurité estimés se produisent dans les pays à revenu faible ou moyen².

Une phase exploratoire visant à élaborer la stratégie de recherche a été menée au travers d'entretiens non-structurés avec d'experts internationaux sur la sécurité des patients et de responsables du secteur humanitaire. Cette phase a permis de confirmer que la sécurité du patient et la détection et gestion des erreurs médicales n'ont pas encore eu de traduction structurée, adaptée au secteur de l'assistance médicale humanitaire. Afin d'essayer de comprendre les raisons de ce décalage par rapport aux critères de qualité de l'offre de soins tels que définis dans les systèmes de santé de pays dont sont pourtant originaires la plupart des organisations médicales humanitaires, cette recherche vise à comprendre quels sont le statut actuel et les perspectives de la sécurité des patients dans l'action médicale humanitaire. Une première phase s'est penchée sur les développements dans les sciences de la sécurité et de la gestion des risques ayant étudié le phénomène de l'erreur et ayant développé des théories et modèles sur la gestion du risque à l'origine de la sécurité du patient. La recherche a ensuite abordé l'état de connaissance actuelle et les principaux développements en matière de sécurité des patients, et de la gestion des erreurs médicales en particulier, avant de s'attarder aux limites et enjeux actuels de sa mise en œuvre. Cette recherche sur Google Search (Mots-clés : Patient safety and risk management) a été suivie d'une revue de littérature rapide sur PubMed (Mots-clés : Patient safety, Medical error, Adverse event, Review, Systematic review since 2013). 104 articles sur les 245 articles trouvés répondaient aux critères. Au total, une base de données de plus de 1.200 articles a ainsi été constituée. En parallèle a été mené une documentation et analyse des caractéristiques de l'action médicale telle que déployée par les organisations médicales humanitaires à travers une revue systématique des articles sur les bases de donnes PubMed et Embase. Les mots-clés utilisés étaient : Humanitarian, patient safety, since 2000. 39 articles sur les 308 références trouvées répondaient aux critères. Cette partie de la recherche a fait l'objet d'un premier article accepté par la revue « Journal of Patient Safety (en attente d'impression).

Les leçons apprises dans les pratiques en matière de sécurité des patients dans les systèmes de santé de l'OCDE, montre la nécessaire conjonction d'actions à trois niveaux : au niveau macro, des règlementations, déterminations d'objectifs et un cadre légal clair ; au niveau micro, du personnel impliqué dans les soins et leur gestion qui soit sensibilisé, formé et impliqué ; et au niveau méso des institutions de soins, catalyseurs de la sécurité des patients grâce à l'engagement et au leadership. C'est à ce niveau que des outils, des processus et un environnement favorable sont initiés, car ces institutions ont la responsabilité d'établir des systèmes de détection d'erreur et un cadre de gestion participatif pour la correction et l'apprentissage.

La deuxième phase de la recherche s'est centrée sur des entretiens semi-directifs avec du personnel médical et paramédical actifs au sein de 6 organisations médicales humanitaires pour connaître l'état actuel des développements en matière de sécurité du patient et de la gestion des erreurs médicales. Les organisations sélectionnées répondaient aux critères suivants : Organisations humanitaires internationales basées dans un pays de l'OCDE ; actives partout dans le monde avec du personnel national et international, ayant un département médical déterminant leurs propres politiques médicales ; Et dont l'action médicale était, en termes budgétaires, l'activité principale (à l'exception du CICR, organisation médicale à la base dont l'activité s'est diversifiée dans le temps). Un premier

entretien a été mené avec les responsables des départements médicaux ou le responsable de la qualité des soins de ces 6 organisations. Ensuite, 39 entretiens ont été menés par Skype ou en face à face avec du personnel médical ou paramédical qui correspondaient aux critères suivants : personnel international avec 2 ans d'expérience minimum dans le secteur humanitaire. 36 entretiens ont été retenus pour analyse, les autres étant de trop mauvaise qualité d'enregistrement pour être exploitables. Les entretiens retenus, entièrement retranscrits ont été codés et analysés par thèmes et sous-thèmes (Atlas ti©).

L'objectif était de comprendre les connaissances, attitudes et pratiques de ce personnel dans ces domaines et entendre les attentes de ce même personnel en ce qui concerne les besoins pour intégrer la préoccupation de la sécurité du patient et la gestion des risques dans les priorités institutionnelles de leur secteur. Les résultats de cette analyse ont fait l'objet d'un deuxième article présenté et accepté, moyennant modifications mineures, par la revue « Journal of Patient Safety ».

Il apparait clairement que s'il n'existe actuellement pas encore dans le secteur d'approche structurée de la question de la sécurité du patient et plus spécifiquement de la gestion des erreurs médicales, cela répond clairement à une attente de la part du personnel humanitaire interviewé. Les raisons invoquées pour expliquer ce manque sont de deux ordres. Il y a d'abord celles en lien avec les spécificités de l'action médicale humanitaire telles que la mentalité et le mode d'action tourné vers l'urgence (tendance à passer d'une priorité à l'autre selon le degré d'urgence), un personnel national et international de culture médicale très variée. A cela s'ajoutent l'absence d'un système de régulation dans ce secteur d'activité tout comme de pressions extérieures poussant à investir dans la sécurité du patient et enfin une grande diversité des activités et des contextes d'intervention ainsi que dans le niveau de contrôle des organisations humanitaires sur les soins prodigués. D'autre part, il y a des facteurs communs à ceux que l'on a pu retrouver dans les systèmes de santé des pays de l'OCDE comme l'absence des ingrédients principaux de la culture de la sécurité : le leadership ; la culture de l'apprentissage et non du blâme ; un soutien approprié du personnel en matière de sensibilisation, formation et développement d'outils pour le rapportage et l'analyse des erreurs médicales ; ainsi qu'une guidance claire pour la gestion de leurs conséquences.

En matière d'attentes des personnes interviewées, si elles reconnaissent leur rôle dans la mise en œuvre de la sécurité de leurs patients, elles identifient clairement la responsabilité de leurs organisations pour un investissement clair dans les moyens de traduire dans les faits leur désir d'une meilleure prise en compte de la sécurité des patients. Pour ce faire, elles appellent de leurs vœux une collaboration entre organisations pour un partage d'expériences et le développement d'outils et d'approche.

Cette recherche, la première du genre selon nos informations, identifie la motivation du personnel médical et paramédical du secteur humanitaire à s'engager à mener une véritable révolution culturelle pour rendre l'offre de soins plus sûre, même dans des situations précaires. Rattraper le retard dans l'adoption de politiques en matière de sécurité du patient et de la gestion des erreurs médicales renforcerait la « redevabilité » de ces organisations envers les populations vulnérables qu'elles assistent et permettrait de sauver plus de vies, l'essence même de l'action humanitaire.

4. Table of Content

1.	List of Publications	3
Published articles		3
U	nder Preparation	4
2.	Abstract in English	5
3.	Abstract in French	8
4.	Table of Content	. 12
5.	General Introduction	. 13
6. A	Articles	. 19
Methodological contributions		. 19
	Phase 1: Patient Safety and Patient Safety in Medical Humanitarian Action: The State of Art	the 19
	Phase 2: Knowledge, Attitudes and Practices as Well as Expectations of Medical Humanitarian Actors Regarding Patient Safety	. 21
A	rticle 1	. 23
A	rticle 2	. 44
7. Conclusions and Perspectives		. 64
Annex 1: Oral Presentation during the Geneva Health Forum, April 2016		. 71
Annex 2: Patient Safety: State of the Art		. 77
	Developments in Safety Science	. 77
	1.1 James Reason's "Swiss Cheese Model"	. 77
	1.2 Rasmussen's Migration of Acceptable Boundaries	. 80
	1.3 High Reliability Organisations	. 81
	1.4 Organisational Approaches to Risk	. 82
Transposition of Safety Theories to Healthcare: Towards Patient Safety		. 85
	2.1 Late Investment on Healthcare Safety	. 85
	2.2 Patient Safety as an Attribute of Healthcare Systems	. 89
	2.3 Ingredients of Patient Safety	. 89
	2.4 Patient Safety: Some Still Problematic Issues	. 99
Annex 3: Humanitarian Medical Action and Patient Safety		104
	Definition of Medical Humanitarian Action and Links with Patient Safety	104
Ann	ex 4: Glossary	111
9. B	9. Bibliography17	

5. General Introduction

Patient safety is a concept that has only recently come fully into its own in the medical care sector. The recognition of error, as a subject of systemic concern, has been a slow process despite many individual calls³ and legendary initiatives, such as the development of standards of care by Florence Nightingale.⁴ The publication of the United States Institute of Medicine (IOM) report in 2000, To Err is Human: Building a Safer healthcare System (1999),⁵mentioning 44,000 to 98,000 Americans dying every year from medical errors, sparked a massive investment in error prevention and management. The awareness of numerous cases of incidents during medical care, with severe consequences for the patient, and their aftermath in terms of image, legal, social and psychological effect,ⁱ have forced healthcare systems and institutions – in the countries of the Organisation for Economic Co-operation and Development (OECD) at least – to establish comprehensive mitigation and prevention strategies. This came very late in comparison to other high-risk sectors (aviation, nuclear industry...) and can be explained by various factors: the lack of visibility of the consequences of medical errors (in the worst-case scenario, one patient may die at a time, hiding the extent of the problem); the very idea of doctors making mistakes or errors creates a cultural dissonance⁶ (medical doctors today still enjoy high social status; this is embedded not only in medical professionals' minds but also in the belief of the patients and, more widely, the belief of the general population); the complexity of medical care (involving different disciplines, practised in different locations for the same patient, with ever more complex medical cases); the litigation risks for the staff involved in a medical error and their institutions if it is disclosed.

i See for instance the 2006 Eurobarometer survey on medical errors available on https://ec.europa.eu/commfrontoffice/publicopinion/archives/ebs/ebs_241_en.pdf ; accessed 12 September 2019

The development of these new policies and practices to ensure safer medical care provision and complementing the already existing actions (pharmacovigilance, standards of care...) is known under the broad concept of patient safety.

Patient safety can be defined as "the identification, analysis and management of patientrelated risks and incidents, to make patient care safer and minimize harm to patients".⁷ It builds on the involvement of key actors at three levels: systemic, individual and institutional.

At the level of the healthcare system and its governing bodies, what is required is an environment conducive to the implementation of a political will, under legal and regulatory conditions, for individuals as institutions within the system to speak on, manage and work on preventing medical errors in a structured, responsible and safe manner. By developing recommendations and evaluating health services in terms of quality, the pertinent authorities can develop standards as well as global and national plans of action in the area of patient safety plus tools and guidelines to assist medical institutions and professionals in implementing measures for patient safety. They also have the capacity to nurture the links among the elements of the system, notably to influence universities and schools to develop the necessary training of individual medical staff in the concern, methods and management of patient safety.⁸ National authorities play a major role in influencing society's perception of its relation to healthcare expectations and limits⁹ while promoting actions.ⁱⁱ

At their level, the individual medical staff and managers of healthcare services are key actors in patient safety, not only because they are potentially involved in, or witnesses of, medical error commission, but because they must be fully engaged in the detection, analysis and management of medical errors and of their consequences. This supposes that they have been made thoroughly aware of the error-prone nature of their professional activities as well as understanding that error is human, meaning that there is not

ⁱⁱ For instance a call from the French Ministry of Health available on http://social-sante.gouv.fr/grandsdossiers/signalement-sante-gouv-fr/ Accessed on 18/3/17

necessarily negligence behind a deviation from what is generally considered the standard action to follow in a specific situation. These are requisites for them accepting to disclose detected errors then analyse them in search of lessons to be learned in order to avoid recurrence.

At the medical institution level, the creation of a safe, supportive learning environment is crucial to involving the staff in proper management of medical errors and their consequences and improving medical practices. Creating a no-blame culture and providing adapted support to staff when establishing reporting systems of medical errors are necessary conditions that a committed leadership must provide in order for errors to become a source of learning and subsequent improvement.

Medical error is now better understood,¹⁰ is recognized as an important cause of harm (third cause of death in the U.S.A.¹¹), and strategies, policies and tools have been developed at medical-institution and health-system levels to minimize the risks of error and to manage the occurring cases.¹² The consequences of medical error are impressive. For instance, a study sponsored by the Society for Actuaries for the U.S.A. alone estimated the cost of medical error, for 2008, at \$19.5 billion, of which 80% is additional medical costs: prescriptions, in- and outpatient costs, ancillary services.¹³ The human and economic costs for society are coupled with the legal, financial and reputational consequences for the involved medical staff and medical institutions.

Twenty years of research, experience-sharing and investments have allowed the establishment of patient safety as a discipline in itself, and as an attribute of healthcare systems.¹⁴ Patient safety concept and strategies have now percolated – in OECD countries at least – into all specialties of medicine and all care-giving settings: from ambulatory to home-based care, from general practice to secondary and tertiary level of care.¹⁵ Reports highlight improvements in safety of healthcare provision since the beginning of the 2000s. (For instance, from 2010 to 2014, the overall rate of hospital-acquired conditions declined by 17% in the U.S.A. Improvements were observed in about 60% of safety measures monitored by AHRQ.¹⁶ In Great Britain, a systematic analysis of underlying risk factors lead

to actions enabling a diminution of 20 to 30% of the falls of patient in hospitals¹⁷). However, considerable frustrations are emerging over what has (not) been achieved so far. The awareness that patients must be recognized as agents of their health has not yet translated into an acceptable role for them in error detection and management and its prevention. One of the most difficult tasks of patient safety specialists is to find appropriate measurement strategies to assess where priorities must be assigned in error prevention and to quantify the benefits of proposed measures. Healthcare systems experiencing ever greater budgetary constraints, along with pressures exerted on medical staff to increase productivity, do not favour investments in patient safety efforts. The movement towards patient safety of the last two decades, which complements continuous improvements of healthcare in the last centuries, will remain an endless battle owing to the evolution of medicine and the human nature of its practice. Technological evolution affects patient safety like a double-edged sword. On the one hand, it creates new opportunities and capacities to act on safety risks; on the other, it increases the complexity of medical practice, thereby increasing the risk of making errors. More research is needed to find appropriate measures to avoid and manage the risks. Presently, patient safety mainly targets hospital settings. There is still much effort needed in looking at the specificities of other areas of healthcare provision such as primary care.¹⁸ Today, research and patient safety initiatives are to be found mainly in developed countries despite estimations that two thirds of medical errors occur in low- or middle-income countries.²

It took a long time for patient safety to become a recognized necessary attribute of quality care provision and to be translated into policies, methods and practices. While there are still considerable challenges and much research is still necessary to overcoming their assessed limits, the attention to safety corresponds to the ethical, moral, legal, economic and social obligations of the medical professions and the management of healthcare services.

There is one area of medical practice that seems totally unexplored: medical humanitarian action. A Humanitarian medicine can be defined as "a set of medical or public health practices whose sole intent is to selflessly accommodate and address the tension created between compelling health needs and the ongoing deprivation of resources in a given population or community."¹⁹

An initial review of the literature and interviews with humanitarian aid actors strongly suggest that there is presently no global strategic thinking nor organized practice in the area of patient safety and specifically in medical error prevention and management. For instance, it was only in the 2018 version of the Sphere standards, a reference document in the humanitarian sector identifying norms to be observed in humanitarian response²⁰ and having specific chapters on health-related activities, that patient safety was mentioned several times and a specific paragraph devoted to adverse event management and prevention.

While, a priori, we can understand the difficulties imposed by the precariousness of humanitarian contexts (scarcity of resources, insecurity, competition among priority needs...), there are at least three sets of reasons calling for a real investment in patient safety: 1. ethical responsibilities: medical ethics as much as humanitarian ethics²¹ (calling for the non-malfeasance to patients and/or beneficiaries of aid); 2. legal duties (entering into a contractual relation, the medical staff is expected to deliver a service which is not detrimental to the patient); and 3. practical realities (patients' complaints regarding alleged sub-standard care, although poorly documented, are reported²² and deserve proper management; media start to relay stories of alleged medical error in the sector;ⁱⁱⁱ medical humanitarian action is implemented in more and more developed healthcare

iii See for instance two cases related to vaccination campaigns in South Sudan and in Syria. Available on https://www.bbc.com/news/world-africa-40135814 https://www.bbc.com/news/world-africa-40135814 and https://www.newsweek.com/15-syria-children-dead-after-measles-vaccination-271525 https://www.newsweek.com/15-syria-children-dead-after-measles-vaccination-271525. Accessed on 15 September 2019.

systems as in the Middle East or in transit or destination countries for migrants where the requirements in terms of quality by authorities and populations are high.

Although there are no reasons why patient safety could not, and should not, be addressed to improve the quality of care delivered in medical humanitarian action, it is reasonable to suppose that a simple transfer of strategies and experiences in patient safety from developed countries' healthcare systems may be difficult due to the specificities of humanitarian action.

Within the framework of this research, we sought to address the following questions.

Primary Question:

What are the present status and perspectives of patient safety, and medical error management in particular, in medical humanitarian action?

Specific Questions

- What are the characteristics, challenges and limits of patient safety strategies, policies and practices implemented in health systems of OECD countries?
- What are the specificities of medical humanitarian action affecting the possible adoption of OECD documented policies and practices of patient safety?
- What are the knowledge, attitude and expectations of medical and paramedical staff active in humanitarian organisations, regarding patient safety in general and medical error management in particular?

We began by exploring the evolution in safety and risk management science pertaining to medical error which gave rise to the theories and models of risk management at the origin of the patient safety discipline. We explored the genesis and the main developments of patient safety to describe the present limits and challenges of patient safety implementation (Annex 2). To limit the scope, this research focused on the experiences and practices of patient safety in hospitals. Confronted with a vast array of concepts and

debates on the appropriate terminology to use in the discipline of patient safety, we developed a glossary of the main terms related to patient safety used in the literature (Annex 3). We then documented and analysed the specificities of medical action deployed in humanitarian contexts (Annex 4) and how this may impact patient safety. This is the focus of the first published article.

The second part of the research went on to document the current knowledge, attitudes and practices in medical humanitarian organisations concerning patient safety and, in particular, error management and prevention and to explore the expectations of staff regarding further investments in this area. For this, we interviewed medical and paramedical staff active in international humanitarian organisations. The results were presented in the second article.

6. Articles

Methodological contributions

The research was done in two steps, each requiring a distinct methodology.

Phase 1: Patient Safety and Patient Safety in Medical Humanitarian Action: The State of the Art

Three strategies were developed in parallel for the literature review: 1. An exploratory literature review on patient safety; 2. Followed by more systematic literature review (rapid literature review) on reviews and systematic reviews articles on the same topic: and 3. a systematic literature review on medical humanitarian action and patient safety.

General Literature Overview on Patient Safety

The doctoral candidate (J.-M. Biquet) conducted a Google search of relevant sources related to patient safety research and practices presenting and analysing present strategies, policies, experiences and challenges on patient safety: evolution of risk management, experiences in other economic sectors, analysis of specific elements leading to medical errors, practices developed to manage them and avoid their recurrence. This search was guided by advice from specialists of patient safety and professionals in

humanitarian action consulted in the preliminary phase of the research. In this context, J.-M. Biquet followed an MOOC on "Leading Healthcare Quality and Safety" developed by George Washington University in the United States.

Rapid Literature Review on Selected Criteria on Patient Safety

Objective: to have a state-of-the-art appraisal of knowledge from present strategies, policies and practices in healthcare systems.

Method: a rapid literature review of all articles in PubMed related to patient safety, covering "reviews" or "systematic reviews" in the last five years and with "review" in the title; papers using the concepts of patient safety and medical errors or adverse events, written in the last five years, were collected and analysed.

For the selection process, two reviewers (J.-M. Biquet and P. Michel) independently screened the titles and abstracts yielded by the search against the inclusion criteria to identify relevant abstracts. Full reports were obtained for all the titles that appeared to meet the inclusion criteria even in cases where there were disagreements between the reviewers. The same two reviewers read the full-text articles obtained and selected those that met the inclusion criteria, categorizing the articles. Thereupon followed a qualitative narrative synthesis by J.-M. Biquet, who created lists of articles based on themes (authors, titles, main messages).

Systematic Literature Review on Medical Humanitarian Action and Patient Safety

Objective: to have a state-of-the-art appraisal of patient safety in medical humanitarian action.

Method: a systematic review of all articles on Embase and PubMed published since 1/1/2000, in French or English, mentioning the concepts of patient safety and humanitarian medicine. This part was managed by J.-M. Biquet. The selected articles were categorized by J.-M. Biquet according to their characteristics: call for patient safety improvements; ethical dimension of medical humanitarian action; training or management of staff; tools related to quality of care.

There was also a database search strategy to find studies on patient safety in medical humanitarian action:

Articles were searched for the following terms:

PubMed search key: (((((("patient safety"[MeSH Terms] OR ("patient"[All Fields] AND "safety"[All Fields]) OR "patient safety"[All Fields]) AND (humanitarian[All Fields] AND ("medicine"[MeSH Terms] OR "medicine"[All Fields]))) OR (medical[All Fields] AND humanitarian[All Fields] AND ("Practice (Birm)"[Journal] OR "practice"[All Fields]))) AND (English[lang] OR French[lang]))) AND ("2000/01/01"[PDat] : "2018/10/30"[PDat])) Sort by: PublicationDate Filters: Publication date from 2000/01/01 to 2018/09/31

Embase search key: ('patient safety'/exp OR 'patient safety') AND ('humanitarian'/exp OR humanitarian)

The three strategies combined led to the creation of a database of more than 1,200 articles.

Phase 2: Knowledge, Attitudes and Practices as Well as Expectations of Medical Humanitarian Actors Regarding Patient Safety

We approached eight international medical humanitarian organisations (six responded positively) in order to know the present status of patient safety in terms of strategy and policy and to be able to interview some of their active international medical and paramedical staff. The selection criteria for the organisations were: having a medical department enacting the organisation's medical policies; based in Organisation for Economic Co-operation and Development (OECD) countries but active throughout the world; working with national and international staff; medical action being the major area of activity in terms of budget (with the exception of ICRC, historically a medical organisation, having diversified its portfolio of activities)

In order to learn qualitative research methods, J.-M. Biquet took a course at the University of Geneva. The lecturer, Prof. Claudine Burton-Jeangros, agreed to review the methodology used for the research and to comment on an early draft of the article.

J.-M. Biquet selected the organisations to be approached after discussion with, and approval by, the doctoral committee. He also designed and tested the interview guide. The final version of the interview guide was drafted with D. Schopper. The selected organisations respected the following criteria: International humanitarian organisations based in an OECD country; active throughout the world with national and international staff; having a medical department implementing the organisation's medical policies; and whose medical action was, in terms of budget, the major area of activities (with the exception of ICRC, historically a medical organisation having diversified its portfolio of activities).

In addition to interviews with six medical directors or those responsible for quality of care and three legal advisors (the others declining the interview for confidentiality reasons), 39 interviews were done with medical and paramedical staff of those organisations. The interviewees respected to the following criteria: international staff with minimum 2 years of experience in the humanitarian sector. 36 interviews were analysed, the others being of too poor quality of recording to be transcribed effectively. The selected interviews integrally transcribed were coded and analysed by themes and sub-themes (Atlas ti ©).

All interviews were done by J.-M. Biquet. He did the transcription of interviews using Express Scribe[©] and the analysis with the data analysis software Atlas TI[©] (coding) to extract the trends and the main elements. The preliminary analysis results were discussed with the doctoral committee.

The writing of the two articles was carried out by J.-M. Biquet. All drafts were reviewed and commented by the members of the doctoral committee before their final approval for submission.

A Call for the Application of Patient Safety Culture in Medical Humanitarian Action: A Literature Review

Jean-Marc Biquet, MSc; Doris Schopper, PhD, MD; Dominique Sprumont, PhD; Philippe Michel, PhD, MD

Abstract

<u>Objectives:</u> The aim of the study was to assess lessons learned on patient safety in OECD countries and assess if they are applied or can be applied to humanitarian medicine.

<u>Methods</u>: 1. A 2013-2018 rapid literature review of reviews and systematic review of articles (PubMed database) on "patient safety" and "medical error" to look for lessons learned regarding patient safety in OECD countries; 2. a rapid literature review (PubMed and Embase databases) on "humanitarian medicine" and "patient safety", from their creation to 2018, to find any articles related to patient safety in humanitarian medicine. In both reviews, articles specifically related to one device, disease or medical act were excluded. These reviews were complemented by a Google search.

<u>Results:</u> 1. Of the 245 references retrieved, 104 met the inclusion criteria. 2. Of 308 references, 39 respected the inclusion criteria. In OECD countries, patient safety comprises correlated measures taken at three levels. The micro level focuses on individual staff involved in healthcare provision or management; the meso level focuses on medical institutions; the macro level focuses on national healthcare systems. Only one reference mentioned the implementation of a medical error reporting and analysis system in medical humanitarian organisation.

<u>Conclusions</u>: Adopting strategies and a culture of safety will require adapting to and addressing the variety of intervention contexts and responding first to the fears and

expectations of humanitarian staff. Medical humanitarian organisations, in the absence of an overarching authority for the sector, have a major responsibility in the development of a general patient safety policy applicable in all their operations.

<u>Keywords</u>: patient safety, safety culture, humanitarian medicine, medical error, literature review

Introduction:

Medical humanitarian organisationsiv are used to taking risks and to deploying ingenuity to provide medical care to victims of crises. Besides trying to overcome the obstacles to access to care, these organisations have long been concerned with the improvement of the quality of their services such as professionalization of staff, development of tools and guidelines, creation of joint platforms to analyse and share lessons learned. Unlike what is taking place in healthcare systems in Organisation for Economic Co-operation and Development (OECD) countries, most of them being of high-income, there is one domain, "the dark side of quality"¹ or the prevention and management of medical errors, which seems presently not fully explored and addressed by humanitarian aid providers. Human action is prone to error. Over the last two decades, the recognition that this also holds true for medical staff has sparked initiatives to analyse incidents where errors were committed and use them in learning processes. However, only when researchers aggregated what was seen, until then, as a series of individual cases of serious errors did the safety issues in healthcare appear so strikingly. The IOM's 2000 report To Err is Human² was a wake-up call for the healthcare sector. Error or adverse event, "an event that results in unintended harm to the patient by an act of commission or omission rather than by the underlying disease or condition of the patient",³ is now recognized as a major cause of harm (the third cause of death in the USA).⁴ It carries a heavy reputational and

^{iv} Medical humanitarian organisations can be international organisations or non-governmental organisations providing medical services in humanitarian crisis. There are, for instance, more than 700 of them collaborating in coordination mechanisms for the health response to crises. See: https://www.who.int/health-cluster/partners/current-partners/en/ 3

financial cost for healthcare systems. In Europe, 15% of hospital expenditure is related to treating consequences of safety incidents,⁵ and for society as a whole, the economic loss by medical error represents up to 23 million DALYs (Disability-Adjusted Life Year) per year in the world.⁶ Ensuring a safer provision of healthcare falls under the umbrella of patient safety, to wit "the absence of preventable harm to a patient during the process of healthcare and reduction of risk of unnecessary harm associated with healthcare to an acceptable minimum. An acceptable minimum refers to the collective notions of given current knowledge, resources available and the context in which care was delivered weighed against the risk of non-treatment or other treatment."7 Considered an integral component of healthcare systems and institutions, it is an important subset of the quality of care.⁸ We wanted to assess if humanitarian medicine, defined as "a set of medical or public health practices whose sole intent is to selflessly accommodate and address the tension created between compelling health needs and the ongoing deprivation of resources in a given population or community",⁹ has integrated patient safety into delivery of patient care. After an assessment of the current knowledge, lessons learned and discussions on medical error prevention and management in OECD countries in hospital institutions, this article explores the integration of patient safety principles into the care delivered by aid organizations. Finally, suggestions are provided for a path forward in developing a stronger patient safety environment for the delivery of medical care by humanitarian organisations.

Methodology

Data Sources

We used the rapid literature review method¹⁰ to find lessons learned regarding patient safety strategies and practices. The review was conducted in September 2018 and included studies published between 2013 and September 2018 in the PubMed bibliographic database for the lessons learned from patient safety. In parallel, we did a rapid literature review on PubMed and Embase databases to find research related to patient safety in medical humanitarian action. To complement these rapid literature

reviews, we searched Google for articles and textbooks related to lessons learned in patient safety as in quality of care in humanitarian action.

Search Strategy, Study/Paper Selection; and Data Analysis for the First Review of "Lessons Learned Regarding Patient Safety"

The inclusion criteria were peer-reviewed reviews and systematic review articles evaluating methods or tools aiming at improving the safety of care. Excluded were articles related to a specific pathology, medical act or medical material or concerning ambulatory or primary care services. The reference lists of selected articles were manually searched for additional citations. The search terms in PubMed were (Box 1): "patient safety" and "medical error" and a list of MESH terms in order to be as exhaustive as possible; filters were: review, systematic reviews, having "review" in the title. For the selection process, two reviewers (J.-M. Biquet and P. Michel), specialists in global health and patient safety, independently screened the titles and abstracts yielded by the search against the inclusion criteria in order to identify relevant abstracts. Articles with a title or abstract that did not meet one or more review criteria were rejected. Full reports were obtained for all the titles that appeared to meet the inclusion criteria even in cases where there were disagreements between the reviewers. The same two reviewers read the full-text articles obtained and selected those that met the inclusion criteria, categorizing the articles according to the following study characteristics (Fig1): articles related to: evaluation of tools; methods of patient safety; training or working conditions of staff; measurement or costs of medical errors; terminology used in patient safety; experiences of patient safety in non-OECD countries; safe management of drugs; disclosure of medical error; health information technology; safety culture. Thereupon followed a qualitative narrative synthesis. We created lists of articles per themes (authors, titles, main messages).

Search Strategy, Study/Paper Selection; and Data Analysis for the First Review of "Patient Safety in Medical Humanitarian Action"

The search key words (Box 2) were: "patient safety" and "humanitarian". The selection criteria were articles explicitly calling for or describing aspects of patient safety or quality 26

of care in medical humanitarian action. Excluded were articles not directly linked to humanitarian action or the quality of clinical practices or articles related to a specific disease or device (Fig.2). The selected articles were categorized according to their characteristics: call for patient safety improvements; ethical dimension of medical humanitarian action; training or management of staff; tools related to quality of care.

Results

The database search on patient safety and secondary healthcare (Fig. 1) identified 245 articles, of which 141, related to a specific disease, medication or medical act or to the use of highly sophisticated technology, were excluded. Also excluded were articles unrelated to patient safety in hospital care. The remaining 104 underwent a full text review. For medical humanitarian action, 308 articles were identified, of which 39 met the inclusion criteria and subsequently underwent a full text review (Fig. 2).

Lessons Learned Regarding Patient Safety Policies, Strategies and Practices

In OECD countries, at least and according to literature, patient safety efforts comprise correlated partially dependent measures at three levels. The micro level initiative focuses on individual staff involved in healthcare provision or management; the meso level, medical institutions; and the macro level, the national healthcare systems. At the macro level, laws and regulations influence (through incentives and disincentives) the reduction of medical errors and the improvement of safety by identifying patient safety obligations¹¹ and establishing legal and regulatory management of occurring cases. Accreditation bodies, such as The Joint Commission (TJC) in the U.S., determine accreditation requirements for healthcare professionals and institutions. Ministries of health, as professional organisations, are also empowered to reinforce links between different elements of the system, notably by motivating medical schools to develop training on patient safety methods and management,¹² and to leverage society's perception of healthcare risks and limitations.¹³ At the meso level, healthcare organizations, the cornerstone of healthcare provision, are the link between micro and macro levels, the enablers of patient safety through commitment and leadership,¹⁴ the crossroads of 27

frequently conflicting imperatives: quality of care vs efficiency. It is at their level that tools, processes and favourable environments are initiated, as they have the responsibility to establish error detection systems and a participatory management framework for remediation and learning. At the micro level, medical staff must be sensitized to the risk of error; must understand that involvement in an error does not equal negligence or incompetence; must learn to implement preventive and corrective actions. Their active participation in the management of errors is a necessity supposing favourable working conditions.^{15,16}

Accomplishments in Error Detection and Safety Culture

The complexity of patients' conditions (multi-morbidities), the importance of team work in patient care management,¹⁷ and extra-medical elements like stress, administrative burden or logistical aspects are some of the factors leading to errors.¹⁸ The analysis of these factors requires a multi-disciplinary approach.¹⁹ Root cause analytical grids can spot contributing factors²⁰ for which reactive measures can then be conceived such as sensitization strategies, staff training, protocol revision, work environment adaptation and improvement of communication. Depending on medical institutions' priorities, extensive root cause analysis may be limited to the most serious incidents. Classification methods assist in identifying points of vulnerability or failure in care processes.²¹ A proper internal feedback mechanism on lessons learned from errors along with selected strategies for safety improvement are then key to preventing staff from perceiving such reporting and analysis as just another administrative burden.²²

Error identification is structured through different data collection methods,²³ each with its advantages and limitations. Voluntary reporting requires a serious commitment from staff and may lead to under-reporting. Direct observation of clinical procedures contributes to detecting system errors but also requires resources. Trigger tools are less labour-intensive, but their validity has been questioned. Mortality-morbidity conferences require clear and complete patient files, clear case definitions and teamwork abilities.²⁴ Other sources of data such as patient complaints²⁵ complement these methods. Generally,

medical institutions use a combination of methods to reach the fullest vision of reality and to respond with preventive actions.

Beyond tools to collect and analyse data, collective investment in a safer provision of healthcare is underpinned by a common safety culture. This concept is defined as "an integrated pattern of individual and organizational behaviour, based upon shared beliefs and values that continuously seeks to minimize patient harm".²⁶ Within healthcare institutions, it translates into participatory mechanisms of reporting and analysis of errors (as opposed to the blame culture) and an accepted openness to system latencies.²⁷ In contrast to a traditional "blame-shame" culture, where those involved in a medical error may be sanctioned for what is considered culpable weakness, it is incumbent upon institutions to create a climate in which individual staff members understand that errors can be learned from and that they can expect management's support²⁸ and guidance. To keep this focus among competing imperatives implies genuinely strong and visionary leadership.²⁹

Unexplored Territory and Limitations

Patient safety, as a discipline with its own modalities and culture, is in constant evolution. In addition to improvements like hygiene,³⁰ increased reliability of medical devices and updated standards of care, patient safety recognizes that humans are prone to error, and thus becomes a continuous source of learning to improve medical practice. Beyond the development of techniques and processes to detect and act on care-related risks,³¹ specialists see leadership, human resource management and comprehensive therapeutic pathways as the challenges for breakthroughs in patient safety⁻³²

While the literature highlights undeniable improvements in patient safety since the beginning of the 2000s, much remains to be done.³³ Technology can provide a means of preventing errors, but there is tension between the promise of technology's contribution to decreasing human error and the documented evidence of increased risk of incidents arising from its ever greater complexity.³⁴ The slow integration of safety into healthcare planning strategies is linked to the complexity of healthcare practice. Not only does this 29

practice encompass a variety of professions, but it also involves large teams interacting at different levels of intensity, along different hierarchical lines and possibly performing in different locations.³⁵ Moreover, medicine is expected to develop a response to increasingly complex medical problems like co-morbidities and evolution of diseases,³⁶ making the identification of a problem's causal factors more difficult and requiring a systems approach. There are still many bridges needed between medical and legal professionals to agree on a regulatory environment providing a fair balance between, on the one hand, responsibility and compensation mechanisms³⁷ and, on the other, measures favouring patient safety, particularly the anonymity of error reporting and the facilitation of disclosure to patients.³⁸ Even proper evaluation of patient safety actions tends to produce patchy results and requires further investment in the search for adapted measurement methods.³⁹ The role of the patients in patient safety is still a matter of research.⁴⁰ Their involvement in patient safety efforts seems pertinent and beneficial⁴¹ beyond their right to play a significant role in the management of their own health. Currently, health budgets in OECD countries do not sufficiently address the need for research and investment in patient safety, notably in primary care.⁴² In low and middle income countries, patient safety remains embryonic.^{43,44} Admittedly, there are still discussions on optimal approaches to improve safety of care.⁴⁵

Patient Safety in Medical Humanitarian Action: A Humanitarian Exception?

Regarding medical care in humanitarian contexts, be they conflicts, natural disasters or significant deficiencies in access to health services for population groups, the initial literature review as well as discussions with humanitarian practitioners show that there is as yet no structured strategic thinking on patient safety. Only one initiative of patient safety policy implementation was identified,⁴⁶ although several articles call for such development.^{47,48,49}

A priori, it could be considered that, since humanitarian contexts are intrinsically precarious owing to a scarcity of resources, to insecurity and to competing needs, simply succeeding in providing healthcare services is a considerable achievement. Yet there are

at least four reasons to call for an increase in the safety of care provided by medical humanitarian organizations. First, ethical responsibilities, based on both medical ethics and humanitarian values,⁵⁰ require respect of the principle of non-maleficence in providing care to patients. Second, medical staff have the legal obligation to act with respect for the rights and integrity of their patients. Third, complaints of patients against humanitarian organizations for alleged substandard care are already a reality. (For instance, Médecins Sans Frontières' legal department received at least 15 complaints within two years following the earthquake in Haiti (Interview MSF in January 2013), and medical errors have indeed been reported.v). Fourth, inclusion of patient safety initiatives would complement the present efforts of medical humanitarians to improve the overall quality of the services they provide to the neediest populations. Striving for safer healthcare provision is not a luxury,⁵¹ even in the context of humanitarian healthcare delivery.

The Characteristics of Humanitarian Medicine

Medical humanitarian action embodies the values of solidarity and humanity and is mostly carried out in countries whose health systems are failing their populations at a specific time or in areas hit by a conflict or a natural disaster. By providing vital services in circumstances generally associated with disasters, both natural and man-made, it evokes images of heroism and urgency. Neither literature nor practice provides clarity on when medical humanitarian action is supposed to begin or end. This leads to medical humanitarian projects having multiple forms such as obvious emergencies, targeted structural support, assistance to substantially deficient healthcare systems, long-term projects, fighting the spread of a specific disease or medical assistance to a neglected

^v Example: In 2014 a vaccination campaign in Syria resulted in the death of 18 children. The vaccinators, who couldn't read English, mistakenly used an anaesthetic product instead of sterile water as a diluent for the vaccine. Newsweek. September 2014. Available on http://www.newsweek.com/15-syria-children-dead-after-measles-vaccination-271525. Accessed on 13 January 2018.

population within a stable context. Aid organizations apply their own criteria in selecting the needs they will respond to and the means to be deployed.

The humanitarian sector is by and large not structured or regulated; therefore each organization operates and defines its own mandate independently.⁵² There is no overarching governing body overseeing action, and no set of practical rules beyond what is locally imposed by each country's laws and regulations, while regulations and social norms differ from one region of the world to another.

In addition to the precariousness of the working environment, the lack of availability of qualified staff as well as the frequent lack of referral possibilities for complex cases are some of the factors pushing humanitarian aid medical volunteers to constantly take risks by pushing the limits of what can be done with scarce resources.⁵³ The aid provided is often a compromise between what should and what can be done, leading to demanding working environments for medical personnel, characterized notably by stress, long working days and different medical cultures among staff.⁵⁴ Having to compromise is also a source of moral distress for staff who are not always provided with the necessary institutional support.⁵⁵

On top of the intrinsic complexity of medical care, motivational factors to invest in patient safety are not as advanced in the humanitarian sector as in OECD countries. Patients' rights are rarely well defined or even broadly understood, much less subject to oversight. Monitoring and counter-power structures lack organisation (patient associations, judicial mechanisms). The very conditions of vulnerability, amplified by crisis, do not allow communities to hold accountable those who have volunteered to respond to their needs, and local health authorities may have limited capacities to oversee the care provided. Medical humanitarian organizations are used to monitoring and evaluating their activities mainly in quantitative terms such as number of people accessing their services, encouraged to do so by their donors, to whom they are primarily accountable. Measuring the qualitative aspects of the care is left mostly to the initiative of individual aid workers and the good will of organizations.

Discussion

Humanitarian action is based on high moral values. The commitment to "do good" goes together with the desire to "do it well".⁵⁶ The internal push for quality improvements is a consequence of the professionalization of the aid sector reinforced by the values of their committed medical volunteers (Do No Harm, quality initiatives...). This leads to a multiplication of efforts to constantly revise procedures without, until now, a clear focus on patient safety. The awareness of the perfectible quality is already present.⁵⁷ The adoption of adapted patient safety strategies could reinforce the convergence of values and practices and systematize the increase of safer care. The specific conditions under which humanitarian medical interventions take place can be perceived as being in contradiction to the basic requirements of patient safety. Yet, this does not preclude the possibility of creating a working environment that limits the stress of the staff while providing the safest care for patients, in other words avoiding the addition of another layer of stress to a situation that is already very stressful for both patients and caregivers. The absence of an overarching authority and the multiplicity of contexts at the macro level, associated with the great diversity of medical cultures of the ever-changing staff (international and local) at the micro level, put the responsibility to develop an adapted model of patient safety for medical humanitarian action first and foremost on the shoulders of the meso level, the organizations themselves. As in OECD countries, the sparking of an internal discussion on the human nature of (humanitarian) medical action and the inherent associated and multifactorial safety risks seems a necessary first step.⁵⁸ It should aim at creating adherence to an institutional investment in the specific dimension of quality of care and should empower individuals to engage. It should involve an assessment of internal fears but also expectations, as well as proposals from individual medical and managerial staff to integrate an adapted response to each of them.

At the institutional level, the initiation of a collective action on risk prevention and management at project sites requires two correlated steps: a clear declaration of commitment from organizational leaders, setting out principles and goals to achieve, and

the provision of structured direct support empowering front-line medical and paramedical staff with training, tools and procedures. At a later stage, when internal policies with their associated procedures are established, aid organizations will be in a better position to lay out to the external stakeholders (authorities, communities and donors) the inherent safety risks of adverse events in humanitarian contexts.

The methodology of rapid literature review has an inherent limitation: a limited number of accessed databases. The research team tried to complement the method by searches on Google Scholar and by meeting specialists of medical humanitarian action and of patient safety in OECD countries. The very limited number of articles focusing on patient safety in medical humanitarian action forced the research team to include articles that mentioned patient safety without elaborating what it exactly meant and how to implement it.

Conclusions

The use of a carbon copy of patient safety strategies and tools implemented in OECD countries does not seem appropriate given the range of operational realities in medical humanitarian aid.

The diversity of intervention circumstances and activities, such as the level of control over medical care provision and evolution of the project from acute emergency care to long term care, indicates that a proper assessment of each situation as well as local regulations, practices and beliefs would be useful in medical error management. These assessment criteria could be used to adapt current strategies and tools for practical implementation of overall patient safety at the operational level, depending on the acuteness of the emergency and availability of qualified medical staff. Reviewing medical care facilities, personnel and care delivery processes of current aid projects with an eye to patient safety, while applying lessons from what is presently implemented in OECD countries, should make it possible to provide a new framework for safer humanitarian medical aid delivery. Going beyond the challenge of offering access to healthcare in sometimes extreme conditions while working on safer delivery presents volunteers and their organizations 34
with a new challenge, based on a universal conviction that safety is part of what patients deserve regardless of the circumstances of delivery. This is a call for the application of patient safety culture to humanitarian medical aid.

References

1. Vincent C. Risk, safety, and the dark side of quality. BMJ 1997;314: 1775–1776.

2. Kohn LK, Corrigan J, Donaldson M (eds). Institute of Medicine, Committee on Quality of Health Care in America, To Err Is Human: Building a Safer Health System. Washington, DC: National Academies Press, 2000.

3. IOM Patient Safety: Achieving a New Standard for Care. Available at: http://www.nap.edu/catalog/10863.html p.327. Accessed December 19, 2018.

4. Makary M, Michael D. Medical error—the third leading cause of death in the US. *BMJ* 2016;3:i2139.

5. World Health Organization, Patient Safety: Making Health Care Safer (Geneva: World Health Organization, 2017), Available at: <u>http://www.who.int/patientsafety/publications/patient-safety-making-health-care-safer/en/</u>. Accessed December 19, 2018.

6. Jha A, Larizgoitia I, Audera-Lopez C, et al. The global burden of unsafe medical care: analytic modelling of observational studies. *BMJ Qual Saf* 2013;22:809–815.

7. World Health Organization web site 2018. Available at: http://www.who.int/patientsafety/en/. Accessed December 18, 2018.

8. Committee on Quality of Health Care in America, Institute of Medicine, Crossing the Quality Chasm: A New Health System for the 21st century, National Academy of Sciences, 2001. Available at: <u>http://www</u>. nationalacademies.org/hmd/Reports/2001/Crossing-the-Quality-Chasm-A-New-Health-System-for-the-21st-Century.aspx. Accessed December 19, 2018.

9. Calain P. In search of the "new informal legitimacy" of Medecins Sans Frontieres. *Public Health Ethics* 2012;5:56–66.

10. Khangura S, Konnyu K, Cushman R, et al. Evidence summaries: the evolution of a rapid review approach. *Syst Rev* 2012;1:10.

11. Baume C. Gestion des risques cliniques et responsabilité medico-hospitalière. Neuchâtel: Institut du droit de la santé de l'Université de Neuchatel. 2015.

12. Abbas M, Quince T, Wood D, et al. Attitudes of medical students to medical leadership and management: a systematic review to inform curriculum development. *BMC Med Educ* 2011;11:93.

13. Quintard B, Roberts T, Nitaro L, et al. Acceptability of health care–related risks: a literature review. *J Patient Saf* 2016;12:1–10.

14. Montgomery A. The relationship between leadership and physician well-being: a scoping review. *J Healthc Leadersh* Vol. 2016;8:71–80.

15. Mao X, Jia P, Zhang L, et al. An evaluation of the effects of human factors and ergonomics on health care and patient safety practices: a systematic review. *PLoS One* 2015;10:e0129948.

16. Gates M, Wingert A, Featherstone R, et al. Impact of fatigue and insufficient sleep on physician and patient outcomes: a systematic review. *BMJ Open* 2018;8:e021967.

17. Epstein N. Multidisciplinary in-hospital teams improve patient outcomes: a review. *Surg Neurol Int* 2014;5:S295–S303.

18. Gaffney T, Hatcher B, Milligan R. Nurses' role in medical error recovery: an integrative review. *J Clin Nurs* 2016;25:906–917.

19. Lawton R, McEachan R, Giles S, et al. Development of an evidence-based framework of factors contributing to patient safety incidents in hospital settings: a systematic review. *BMJ Qual Saf* 2012;21:369–380.

20. Vincent C, Taylor-Adams S, Chapman EJ, et al. How to investigate and analyse clinical incidents: clinical risk unit and association of litigation and risk management protocol. *BMJ* 2000;320:777–781.

21. World Health Organization. The conceptual framework for the International Classification for Patient Safety (Version 1.1). Final Technical Report. Geneva: Switzerland; 2009. Available at: https://www.who.int/patientsafety/taxonomy/icps_full_report.pdf. Accessed December 19, 2018.

22. Brunsveld-Reinders A, Arbous MS, De Vos R, et al. Incident and error reporting systems in intensive care: a systematic review of the literature. *International J Qual Health Care* 2016;28:2–13.

23. Michel P. Strengths and Weaknesses of Available Methods for Assessing the Nature and Scale of Harm Caused by the Health System: Literature Review. Geneva: WHO; 2000. Available at: http://www.who.int/patientsafety/research/P_Michel_Report_Final_version.pdf. Accessed December 19, 2018.

24. Giesbrecht V, Au S. Morbidity and mortality conferences: a narrative review of strategies to prioritize quality improvement. *Jt Comm J Qual Patient Saf* 2016;42:516–527.

25. Christiaans-Dingelhoff I, Smits M, Zwaan L, et al. To what extent are adverse events found in patient records reported by patients and healthcare professionals via complaints, claims and incident reports? *BMC Health Serv Res* 2011;11:49.

26. Halligan M, Zecevic A. Safety culture in healthcare: a review of concepts, dimensions, measures and progress. *BMJ Qual Saf* 2011;20:338–343.

27. Pettker C, Thung S, Raab C, et al. A comprehensive obstetrics patient safety program improves safety climate and culture. *Am J Obstet Gynecol* 2011;204:216.e1–216.e6.

28. Seys D, Scott S, Wu A, et al. Supporting involved health care professionals (second victims) following an adverse health event: a literature review. *Int J Nurs Stud* 2013;50:678–687.

29. Tsai T, Jha A, Gawande A, et al. Hospital board and management practices are strongly related to hospital performance on clinical quality metrics. *Health Aff* (Millwood) 2015;34:1304–1311.

30. Zingg W, Holmes A, Dettenkofer M, et al. Hospital organisation, management, and structure for prevention of health-care-associated infection: a systematic review and expert consensus. *Lancet Infect Dis* 2015;15:212–224.

31. Zegers M, Hesselink G, Geense W, et al. Evidence-based interventions to reduce adverse events in hospitals: a systematic review of systematic reviews. *BMJ Open* 2016;6:e012555.

32. Leape L, Berwick D, Clancy C, et al. Transforming healthcare: a safety imperative. *Qual Saf Health Care* 2009;18:424–428.

33. Wachter R. Patient safety at ten: unmistakable progress, troubling gaps. *Health Affairs* 2010;29:165–173.

34. Salahuddin L, Ismail Z. Classification of antecedents towards safety use of health information technology: a systematic review. *Int J Med Inform* 2015;84:877–891.

35. Kapur N, Parand A, Soukup T, et al. Aviation and healthcare: a comparative review with implications for patient safety. *JRSM Open* 2016;7.

36. Vincent C, Amalberti R. Safer Healthcare: Strategies for the Real World. Cham: Springer International Publishing; 2016. Available at: http://link.springer.com/10.1007/978-3-319-25559-0. Accessed December 19, 2018.

37. Runciman W, Merry A, Tito F. Error, blame, and the law in health care—an antipodean perspective. *Ann Intern Med* 2003;138:974–979.

38. Guillod O. Medical error disclosure and patient safety: legal aspects. *J Public Health Res* 2013;2:e31.

39. Hanskamp-Sebregts M, Zegers M, Vincent C, et al. Measurement of patient safety: a systematic review of the reliability and validity of adverse event detection with record review. *BMJ Open* 2016;6:e011078.

40. Longtin Y, Sax H, Leape L, et al. Patient participation: current knowledge and applicability to patient safety. *Mayo Clin Proc* 2010;85:53–62.

41. Doyle C, Lennox L, Bell D. A systematic review of evidence on the links between patient experience and clinical safety and effectiveness. *BMJ Open* 2013;3:e001570.

42. Ricci-Cabello I, Gonçalves D, Rojas-García A, et al. Measuring experiences and outcomes of patient safety in primary care: a systematic review of available instruments. *Fam Pract* 2015;32:106–119.

43. Bates D, Larizgoitia I, Prasopa-Plaizier N, et al. Global priorities for patient safety research. *BMJ* 2009;338:b1775.

44. Nejad S, Allegranzi B, Syed S, et al. Health-care-associated infection in Africa: a systematic review. *Bull World Health Organ* 2011;89:757–765.

45. Slawomirski L, Auraaen A, Klazinga N. The Economics of Patient Safety. Paris: OECD Publishing; 2017. Available at: http://www.oecd.org/els/health-systems/The-economics-of-patient-safety-March-2017.pdf. Accessed December 19, 2018.

46. Shanks L, Bil K, Fernhout J. Learning without borders: a review of the implementation of medical error reporting in Médecins Sans Frontières. *PLoS One* 2015;10:e0137158.

47. Djalali A, Ingrassia P, Corte F, et al. Identifying deficiencies in national and foreign medical team responses through expert opinion surveys: implications for education and training. *Prehosp Disaster Med* 2014;29:364–368.

48. Merchant A, Ingrassia P, Della Corte F, et al. Evaluating progress in the global surgical crisis: contrasting access to emergency and essential surgery and safe anesthesia around the world. *World J Surg* 2015;39:2630–2635.

49. Dohlman L. Providing anesthesia in resource-limited settings. *Curr Opin Anaesthesiol* 2017;30:496–500.

50. Slim H. Humanitarian Ethics: A Guide to the Morality of Aid in War and Disaster. London: Hurst & Co.; 2015.

51. Patient safety is not a luxury. *The Lancet* 2016;387:19.

52. Wessells M. Do no harm: challenges in organizing psychosocial support to displaced people in emergency settings. Refuge. 2008;25.

53. Hurst S, Mezger N, Mauron A. Allocating resources in humanitarian medicine. *Public Health Ethics* 2009;89–99.

54. Hunt M. Establishing moral bearings: ethics and expatriate health care professionals in humanitarian work. *Disasters* 2011;35:606–622.

55. Hunt M, Schwartz L, Sinding C, et al. The ethics of engaged presence: a framework for health professionals in humanitarian assistance and development work: the ethics of engaged presence. *Dev World Bioeth* 2014;14:47–55.

56. Beck M, Brauman R. Médecins Sans Frontières et la qualité médicale. Paris : CRASH/Fondation Médecins Sans Frontières. 2017 Available at: https://www.msfcrash.org/fr/publications/medecine-et-sante-publique/medecins-sans-frontieres-et-la-qualitemedicale. Accessed December 23, 2018.

57. Reason J, Carthey J, De Leval M. Diagnosing "vulnerable system syndrome": an essential prerequisite to effective risk management. *Qual Health Care* 2001;10:ii21–ii25.

Box 1: Database search strategy to find studies on patient safety policies, strategies and practices

Titles and abstracts of articles published in English were searched for the following terms:

((("patient safety"[MeSH Terms] OR Patient Safety[Title/Abstract] OR (("patients"[MeSH Terms] OR "patients"[All Fields] OR "patient"[All Fields]) AND Safeties[Title/Abstract])) AND ("medical errors"[MeSH Terms] OR (medical error[Title/Abstract] OR medical errors[Title/Abstract]) OR (medical mistake[Title/Abstract] OR medical mistakes[Title/Abstract]) OR wrong procedure errors[Title/Abstract] OR (wrong site surgeries[Title/Abstract] OR wrong site surgery[Title/Abstract]) OR (critical[All Fields]) OR (medical incident[Title/Abstract] OR medical errors[Title/Abstract]) OR (Critical[All Fields] AND (medical incident[Title/Abstract] OR medical incidents[Title/Abstract])) OR

medical critical incidents[Title/Abstract] OR (wrong patient surgeries[Title/Abstract] OR wrong patient surgery[Title/Abstract]) OR Never Event[Title/Abstract] OR Never Events[Title/Abstract] OR (diagnostic error[Title/Abstract] OR diagnostic errors[Title/Abstract]) OR Misdiagnosis[Title/Abstract] OR Misdiagnoses[Title/Abstract] OR (false negative reaction[Title/Abstract] OR false negative reactions[Title/Abstract]) OR (false positive reaction[Title/Abstract] OR false positive reactions[Title/Abstract]) OR (observer variation[Title/Abstract] OR observer variations[Title/Abstract]) OR Observer Bias[Title/Abstract] OR (interobserver variation[Title/Abstract] OR interobserver variations[Title/Abstract]) OR (inter observer variation[Title/Abstract] OR inter observer variations[Title/Abstract]) OR (inter observer variabilities[Title/Abstract] OR inter observer variability[Title/Abstract]) OR (intraobserver variation[Title/Abstract] OR intraobserver variations[Title/Abstract]) OR (medication error[Title/Abstract] OR medication errors[Title/Abstract]) OR Medication Error[Title/Abstract] OR drug use errors[Title/Abstract] OR (inappropriate prescribing[Title/Abstract] OR inappropriate prescribings[Title/Abstract]) OR (inappropriate prescription[Title/Abstract] OR inappropriate prescription,[Title/Abstract] OR inappropriate prescriptions[Title/Abstract]) OR over prescribing[Title/Abstract] OR (medication reconciliation[Title/Abstract] OR medication reconciliations[Title/Abstract] OR medication reconciliationv[Title/Abstract]) OR (Near[All Fields] AND Miss[All Fields] AND Healthcare[Title/Abstract]) OR (("delivery of healthcare"[MeSH Terms] OR ("delivery"[All Fields] AND "health"[All Fields] AND "care"[All Fields]) OR "delivery of healthcare"[All Fields] OR "healthcare"[All Fields]) AND Near Miss[Title/Abstract]) OR (("delivery of healthcare"[MeSH Terms] OR ("delivery"[All Fields] AND "health"[All Fields] AND "care"[All Fields]) OR "delivery of healthcare"[All Fields] OR "healthcare"[All Fields]) AND Near Misses[Title/Abstract]) OR (("delivery of healthcare"[MeSH Terms] OR ("delivery"[All Fields] AND "health"[All Fields] AND "care"[All Fields]) OR "delivery of healthcare"[All Fields] OR "healthcare"[All Fields]) AND Close Call[Title/Abstract]) OR (("delivery of healthcare"[MeSH Terms] OR ("delivery"[All Fields] AND "health"[All Fields] AND "care"[All Fields]) OR "delivery of healthcare"[All Fields] OR "healthcare"[All Fields]) AND Close Calls[Title/Abstract]) OR (radiotherapy setup error[Title/Abstract] OR radiotherapy setup errors[Title/Abstract]) OR event[Title/Abstract] OR adverse events[Title/Abstract] (adverse OR adverse events, [Title/Abstract]))) AND ((Review[ptyp] OR systematic[sb]) AND "2013/11/10"[PDat] : "2018/11/08"[PDat])) AND review[Title] AND ("2013/11/10"[PDat] : "2018/11/08"[PDat]



Flowchart showing the selection of publications on patient safety and medical error or adverse event, 2013-2018

Box 2: Database search strategy to find studies on patient safety in medical humanitarian action

Articles were searched for the following terms:

PubMed search key: (((((("patient safety"[MeSH Terms] OR ("patient"[All Fields] AND "safety"[All Fields]) OR "patient safety"[All Fields]) AND (humanitarian[All Fields] AND ("medicine"[MeSH Terms] OR "medicine"[All Fields]))) OR (medical[All Fields] AND humanitarian[All Fields] AND ("Practice (Birm)"[Journal] OR "practice"[All Fields]))) AND (English[Iang] OR French[Iang])))) AND ("2000/01/01"[PDat] : "2018/10/30"[PDat])) Sort by: PublicationDate Filters: Publication date from 2000/01/01 to 2018/09/31

Embase search key: ('patient safety'/exp OR 'patient safety') AND ('humanitarian'/exp OR humanitarian)



Flowchart showing the selection of publications on patient safety and humanitarian medicine, 2000-2018

Article 2

Submitted to and accepted for publication in the Journal of Patient Safety, January 2020.

Knowledge, Attitudes and Expectations of Medical Staff towards Medical Error Management Policies in Humanitarian Medicine: A Qualitative Study

Jean-Marc Biquet, MSc; Doris Schopper, PhD, MD; Dominique Sprumont, PhD; Philippe Michel, PhD, MD

Abstract

Background Patient safety, a major component of quality of care, is now an attribute of health care systems in developed countries at least. While there is ever more research on this subject in developed countries, humanitarian medicine, mainly implemented in resource-poor countries, has yet to structure its own set of policies and strategies on patient safety and the management of medical errors.

Objectives We assessed the knowledge, attitudes and expectations of medical humanitarian staff regarding the development of policies and strategies related to patient safety and medical error management in medical humanitarian action.

Methods We conducted 36 semi-structured interviews with international medical and paramedical staff active in six medical humanitarian organisations after having interviewed the medical directors or the person in charge of quality of care and the legal advisors. Interviews were transcribed verbatim and subjected to a thematic analysis.

Results The interviews confirmed the current absence of clear investments in dealing with safety risks in the selected medical humanitarian organisations. The difficulties experienced by medical staff in reporting medical errors such as blame culture, lack of training, absence of leadership committed on patient safety are non-specific. Other arguments are related to the specific conditions of humanitarian settings: coexistence of different medical culture; absence of international or local regulations or external pressures; great diversity of activities and contexts.

Conclusions Interviewed staff expressed high expectations of receiving guidance from their organisations and support to adopt clear patient safety and medical error management policies adapted to their complex operational and clinical realities.

Keywords: Medical error, Patient safety, Humanitarian Medicine, Safety culture, safety risks

Introduction

Patient safety is defined as "the absence of preventable harm to a patient during the process of health care and reduction of risk of unnecessary harm associated with health care to an acceptable minimum. An acceptable minimum refers to the collective notions of given current knowledge, resources available and the context in which care was delivered weighed against the risk of non-treatment or other treatment."¹. The management of medical errors, "a failure to carry out a planned action as intended or application of an incorrect plan"², is an essential part of patient safety³. Management of medical errors is most often done with a retrospective approach based on the implementation of an error reporting system.⁴ In the last 20 years patient safety has become a priority in healthcare systems of countries of the Organisation for Economic Co-operation and Development countries (OECD) and beyond. vi

One sector of medical activities, the health care provided by humanitarian aid organisations^{vii}, seems not to have been influenced yet by this groundswell. A recent literature review⁵ found only one initiative of humanitarian policy on medical error reporting and management⁶ and some articles calling for such development^{7,8,9}. There is no internationally agreed definition of humanitarian action, but it commonly refers to the

^{vi} OECD is regularly assessing the state of patient safety in healthcare systems of their country members and contributes to the exchange of experiences. See https://www.oecd.org/health/patient-safety.htm Accessed on December 10, 2019, More and more articles can be found in literature on initiatives developed in low- or middle-income countries.

vii Medical humanitarian organisations can be international organisations or non-governmental organisations providing medical services in humanitarian crisis. There is for instance more than 700 collaborating in coordination mechanisms for the health response to crises. See: https://www.who.int/health-cluster/partners/current-partners/en/ Accessed on December 12th, 2019. 45

assistance provided by professional aid organisations to save life and alleviate suffering in time of crisis¹⁰. Concerning specifically Humanitarian medicine, we selected the following definition: "a set of medical or public health practices whose sole intent is to selflessly accommodate and address the tension created between compelling health needs and the ongoing deprivation of resources in a given population or community"¹¹. It is performed by committed medical and paramedical staff assisted by support staff of medical humanitarian organisations which are numerous and decide independently the scope of their action and where to intervene. Contexts and the range of humanitarian interventions vary, from war-torn contexts, zones affected by natural disasters, to offering care to impoverished populations in stable contexts. The scarcity of means, the very diverse types of aid projects as much as shortage in quantity and quality of staff available to run them seem to be constant features of (medical) humanitarian action.

Besides trying to overcome the obstacles to access to care¹², humanitarian organisations have long been concerned with the improvement of the quality of their services through a process of professionalization, edition of guidelines, adoption of the WHO surgical checklist¹³, quality drugs procurement policies^{viii}. One domain though seems not yet to be addressed in a structured way: the management of medical errors.

In order to understand this gap, this research focussed on assessing the knowledge, attitude and expectations of medical and paramedical staff active in humanitarian organisations, regarding patient safety in general and medical error management in particular.

Method

We employed qualitative methods to capture humanitarian medical or paramedical staff perceptions of risk management; barriers they think explain the presently limited developments of medical error management strategies; and their expectations for the future.

viii See for instance the guidelines for the procurement of medical supplies aid organisations funded by the European commission must abide to. Available on https://www.dgecho-partnershelpdesk.eu/actions_implementation/procurement_in_humanitarian_aid/medical_supplies. Accessed on November 16th, 2019

Sampling

Eight organisations, a convenient choice based on previous contacts, were purposefully chosen to reach a relevant representation of the diversity of the sector (Anglo-Saxon and Latin organisations; emergency oriented or long-term philosophy). The selection criteria were having a medical department enacting the organisation's medical policies; based in Organisation for Economic Co-operation and Development (OECD) countries but active throughout the world; working with national and international staff; medical action being the major area of activity in terms of budget (with the exception of ICRC, historically a medical organisation, having diversified its portfolio of activities). From the eight organisations contacted, six responded positively (the seventh never responded; the medical director of the eighth declined, judging it impossible to convince his leadership to invest in patient safety). These six organisations were: the Swiss and Belgian sections of Médecins Sans Frontières (MSF); Médecins du Monde-France (MDM); Save the Children-UK (SC); International Rescue Committee–USA (IRC); International Committee of the Red Cross (ICRC).

After the signature of a research agreement, we interviewed the director of the medical department and/or the person in charge of quality of care to learn about the status of patient safety efforts related to medical error management. The medical directors or the human resource departments provided contacts of some active medical and paramedical staff corresponding to the selection criteria: active international medical or paramedical staff with a minimum of two years' experience in humanitarian action. These criteria were chosen in order to maximize the chances of meeting interviewees knowing their organisations' policies and who have carried out several missions. The recruitment of volunteers was done by the main researcher through snowballing until saturation, with a minimum set of five per organisation.

Participants were contacted by an invitation e-mail letter and provided an information sheet on the project (confidentiality, anonymization, purpose, use of the data) and a consent form.

Data Collection and Analysis

An interview guide of semi-open questions (Annex 1) was developed by the research team and pilot-tested on three humanitarian medical staff to assess acceptability and feasibility. The final guide was designed in English and French by two researchers^{ix}. All interviews were conducted by one Researcher. The interviews, face-to-face or via Skype and lasting from 45 to 120 minutes, were carried out from April 2017 to January 2018 without any offer of compensation. Interviewees were either still on mission, transiting between two field assignments, or based at the Headquarter of their organisation after several missions. The same researcher transcribed and coded verbatim each interview (Atlas Ti 8); carried out thematic analysis of the transcripts; and proceeded to the data extraction, analysis and review. Data were tabulated by themes and subthemes, then summarised and shared among the research team for discussion. Trends and illustrative quotations were compiled.

Results

1) Demographic Profiles of the Participants

Among the 48 participants contacted, 39 responded positively (81%). The analysis comprised thirty-six interviews, three were abandoned owing to the poor quality of the recording and concomitant impossibility of producing an intelligible transcription. The 23 women and 13 men interviewed (Table 1) had an average field experience in medical humanitarian action of 7 years and 4 months (from 2 to 23 years). Nineteen were Europeans, eleven Africans, five came from the Americas and one from Oceania. Sixteen of them had worked for more than one humanitarian organisation, and several continued to do clinical work in their country of origin between missions. They considered themselves to have been middle or senior managers in their last field assignment.

2) Institutional Policies and Strategies Related to Patient Safety

 ^{ix} The research protocol was presented in April 2017, for approval, to the Geneva "Commission cantonale d'éthique de la recherche" (CCER). The Commission exempted this protocol from ethics review.
 48

Currently, none of the six organisations has a fully operational structured institutional policy on patient safety and on medical error management. Sensitization level and actions on patient safety differ depending on the organisation: one was implementing a pilot project of quality measurement through score cards and a clinical incident reporting system in three missions. Another had a serious adverse-event reporting system sparsely used and promoted. A third had started an internal reflection on the quality of care in its medical services and ways to improve it. A fourth was considering implementing a reporting system for any major incident (medical or other). The last two were not yet developing an institutional approach to patient safety. While they were convinced that medical errors were sometimes committed in their organisations' work, medical directors and those responsible for quality in the six organisations reported that they were informed of such occurrences either rarely or only long afterwards There were no noticeable differences in the answers provided by the interviewees would they work presently at Headquarters or at the field level or according to their professional background, sex or origin.

3) Knowledge of the Concepts of Patient Safety and Medical Error

Most interviewees had already heard of the concept of patient safety, and more than half of the interviewed staff had already encountered patient safety policies in their professional life, outside of their humanitarian experiences (in medical institutions they worked for or during their studies). They confirmed a current absence of clear policies on patient safety and medical error management in their organisations while acknowledging efforts undertaken institutionally in quest of better quality of care. All had seen, participated in or at least heard about medical errors occurring in humanitarian projects. All considered that to err is human, hence anybody can err, even if there was some confusion in differentiating between medical error, intended deviation from a protocol and negligence. The problem I think is that we use the term "error" when we mean somebody did something bad or wrong, and I think the term there we want to focus on is negligence. – Medical doctor, 5 years' experience in 2 organisations When asked what the causes of medical errors were, the most spontaneously mentioned factors were: work overload, distraction, lack of training/supervision, miscommunication or a combination of these factors.

Medical errors seemed barely discussed within the organisations.

That is what I find sad in humanitarian action... It must be such a compelling situation [in order for us] to be able to question ourselves. Even the daily minor errors that we all make and will make again, those... they do not lead superiors to question themselves. – Nurse X, 3 years' experience

4) Prevention and Management of Medical Errors

The interviewees' experience, regardless of the organisation, was that detected cases of medical error were rarely well managed (analysis, search of explanatory factors,

recommendations from lessons learned). In the end, it depended on the team leader.

This is on case by case. There is in fact nothing generalized, standard or which has been approved. This is done according to who is the boss. – Nurse, 15 years' experience in 3 organisations

Information of an error occurrence might be forwarded to the upper hierarchical level

(coordination and/or headquarters) but usually triggered no feedback/support. This

created frustrations, because, according to them, either the case was immediately

hidden, or no proper analysis was done, and a "culprit" eventually sanctioned.

Most of the time it is discussed internally in the team that is working there, to understand what happened and how to deal with it and, if you have a really experienced team leader, even to find solutions for it, how to learn from it. It is the optimal case... Most of the time... you sit in the evening and talk about it... it shouldn't happen, what a stupid guy or a stupid nurse. – Paramedic, 8 years' experience in 3 organisations

According to interviewees, patients and their relatives were rarely informed.

I think, in my experience, in this case, the information given to the family is just: "Ben, we are sorry, despite all our treatments, the patient didn't survive." We don't enter into details because it would expose the medical staff and would expose the organisation. – Midwife, 15 years' experience

Good practices related to error detection or prevention were mentioned as happening regularly across organisations, although depending on the team initiatives: mortality and morbidity reviews, meeting of the entire medical team after an incident, or maternal death audits.

5) Factors Explaining the Absence of Comprehensive Medical Error Management Policies

When asked why medical humanitarian action had up to now escaped the almost 20year-old structured investments in medical error management, the main explanations spontaneously given were: the emergency mode (among so many priorities, patient safety gets subsumed, while the emergency mentality incites to focus exclusively on what is most pressing); fear of negative perception by colleagues or hierarchy (blame culture); lack of resources (to develop processes); lack of pressure to do it (staff and local populations unaware of patients' rights).

Cultural differences were said to play an important role: variety of staff without a common medical culture; communication difficulties between professional categories; different perceptions of responsibilities. The risks of consequences in case of disclosure of errors, even only internally, were acknowledged but discussed only if probed. Humanitarian action's emergency mode, by virtue of either the nature of the intervention or the organisation's internal culture, was frequently mentioned and considered an excuse not to include medical error management as a priority, yet, according to interviewees, it is out of line with the evolution of projects developed by humanitarian organisations, which runs from simple substitution in emergency to support of institutional health care in a stable context. The instability of the contexts regularly affecting the provision of medical services was seen as an inherent part of any humanitarian action and in essence a starting point for action, not a reason to prevent change. Interviewed staff fully recognized the challenge of implementing proper strategies and procedures related to medical errors management in their organisations. The vast variety of contexts, circumstances and work conditions plays a significant role. The varying level of control the organisations have over care provided in their projects 51

was perceived as a clear difficulty in determining responsibilities and in enforcing potential policies related to medical error management: the organisation may have direct control over the care provided by its own staff; the project can be managed through a mix of health ministry and organisation staff; the project can consist of a distant support to the local health ministry by the provision of necessary inputs (drugs, medical material...).

The organisational structure was seen as a hindrance to the development of medical error management policy: support to field teams at the medical department level is organised in silos and does not favour investments in transversal issues.

"Organisations should be pushed to do more, and I think if organisations don't do it, it is not because of a lack of interest but because, for example, of the way [the organisation] is constructed ... you have people focused on paediatric care or specific diseases. And then safety of patient is a very transversal issue" – Midwife, 3 years' experience

6) Expectations with regard to a policy on medical error management

Interviewees clearly saw the responsibility of making care safe at the level of both the individuals and the institution employing them. The latter should provide the means, working conditions and guidance enabling a practical enforcement of safety. The state was mentioned as possibly co-responsible by defining policies and framework. There was unanimity on the need, if not the urgency, to develop clear medical error management policies adapted to medical humanitarian action.

Yes, I think it is very important. We should be held accountable for this because it is not because you are humanitarian that you are exempted from all of these things that we do back in our home country. – Nurse Y, 3 years' experience

The main quoted expectations regarding such a policy referred to procedures, tools and guidance to clarify how to do it (prevention and management of errors), and explanations and objectives to explain why to do it (awareness of safety risks and their consequences). The need for sensitization and training arose with the need of a framework clarifying responsibilities.

The medical errors observed by interviewees did not lead to any institutional or legal consequences. When evoked by the interviewer, the idea of negotiating immunity for

the staff in case of medical error, in acknowledgement of all the difficulties in implementing proper medical services in humanitarian settings, was rejected by all interviewees.

Two thirds of the interviewees considered it more appropriate for each humanitarian organisation to develop its own medical error management policy and strategy, adapted to its specific operations. The other third favoured a broad initiative from the World Health Organisation. In any case, interviewees thought that there must be enough flexibility to enable project sites to adapt the general framework to local specificities. Many interviewees favoured collaboration among humanitarian organisations through sharing of experiences.

Me, I think on this that we should have experience sharing. It would be super-important, between organisations, that we learn from each other... I think that if we, at our level, try to work alone, it will take a lot of time. – Midwife, 23 years' experience

The general proposals of interviewees to improve medical error management in medical humanitarian action reinforced the above mentioned expectations such as more collaborative experience-sharing among actors, sensitization of donors to secure necessary funding for this, more training and sensitization of staff and management and better interaction with patients as well as with local health ministries.

"I think we have been going around the edges a lot, with all our efforts, around quality improvement, quality care... And, you know, I am not aware of how to start it. But that there just needs to be a more thoughtful focus on the full patient safety pieces out tackling it out of support or supervision and systems. – Nurse, 18 years' experience

Discussion

This study is the first to explore some of the reasons why patient safety and medical

error management in particular are not yet a priority for medical humanitarian

organisations, despite patient safety being now recognized as a discipline in itself¹⁴ and

medical error management promoted by the World Health Organisation¹⁵.

The difficulties experienced by humanitarian medical staff in reporting and managing medical errors correspond to what is experienced in OECD countries: blame culture¹⁶, 53

lack of training,¹⁷ absence of leadership committed to patient safety^{18,19}. Several studies done in OECD countries point to the reluctance among medical staff to admit and disclose errors, preventing them from detecting, analysing and conceiving means of avoiding error reoccurrences^{20,21}. To overcome this barrier, studies have highlighted elements to focus on: strong organisational and leadership support for safety; sensitisation about, and training on, error-reporting systems; a shared vision and blame-free culture.²²

However, other difficulties mentioned by the interviewees, related to the specific conditions of humanitarian settings, add further complexity. These include the emergency mode and mentality pushing the organisation to shift from one priority to the next leaving the issues considered less urgent for later, the coexistence of; different medical cultures among the staff as well as the assisted populations; the absence of regulations or external pressures to invest in medical error management; the great diversity of activities and contexts of intervention with varying levels of control on the care provided.

All these difficulties, taken in isolation or together, were nevertheless not considered by interviewed staff as sufficient in themselves to justify an absence of structured investments in patient safety in general and medical error management in particular by the humanitarian sector. While acknowledging their part of responsibility in error prevention and management, they clearly emphasized the responsibility of their organisation in providing them support and guidance as reported in other researches.²³ The field experience of the interviewees made them realistic regarding the time and effort needed to change the internal mentalities concerning medical errors. Also, to become an area of collective action, it must become, as in OECD countries since 2000, a subject of discussion.²⁴

Concerning the external factors as the absence of local regulations and of local recognition of patients' rights or the routine lack of automatic patient involvement in the care processes, there was a clear perception that it is incumbent on the organisations to

actively engage in an overall change relating to their duties and accountability in the domain of medical errors prevention and management.

To engage in change, the humanitarian sector has a range of assets enabling quick progress. A significant part of the interviewed international staff has already a good knowledge of patient safety. The high staff turnover amongst organisations may also favour cross fertilization on this subject. And, far from starting from scratch, the sector can benefit from the lessons learned in implementing patient safety and medical error management policies in OECD countries.

Strengths and limitations

The strengths of this study include its multicentre design, incorporating the perspectives of diverse organisations, levels of responsibility and types of professionals. The validity and transferability of our work is supported by the consistency of our results across the interviews.

The results represent the opinions of those interviewed and cannot be generalized, although the recurring experiences described, and opinions expressed tend to show a common desire for investment in patient safety and medical error management strategies. Questions raised, notably on interviewees' experiences with medical errors, are sensitive and may induce answers influenced by a social desirability bias. The review of interview transcripts and analysis was done by one researcher only even if the results were discussed with the research team.

Conclusions

Medical humanitarian organisations, while doing highly commendable work, still suffer from a lack of consistent incorporation of medical error management into their efforts to improve the quality of care. Reasons evoked by the interviewees can be divided in two categories: those which are commonly found in healthcare institutions in OECD countries, namely blame culture, absence of leadership committed to patient safety, lack of training and the ones related to the specificities of medical humanitarian action. If the latter add complexity, interviewed staff consider they are not sufficient to prevent a real investment by medical aid organisations in adapted patient safety and medical error management strategies.

This research demonstrates the aspiration of humanitarian health care workers to invest in what can be considered an internal cultural revolution with regards to patient safety. Interviewees mostly perceived this to be the responsibility of their organisations and linked it to the duty of organisations claiming their commitment to be accountable to the beneficiaries of their actions.

The expressed call by interviewees for more experience-sharing among organisations suggests considerable scope for institutions like the World Health Organisation to convene global reflections in this area.

More research is needed to know how to apply lessons learned from OECD countries in humanitarian settings and to further understand how existing barriers can be overcome, given that the field is ripe for change. Our findings suggest that the medical and paramedical staff committed to humanitarian values are keen to participate in institutional developments in the domain of patient safety and medical error management that will help to save more lives and alleviate suffering, the ultimate purposes of humanitarian action.

References

¹ World Health Organization website 2018. http://www.who.int/patientsafety/en/. Accessed December 18, 2018.

² World Health Organization. The Conceptual Framework for the International Classification for Patient Safety (Version 1.1). Final Technical Report. Geneva: Switzerland; 2009. Available at: https://www.who.int/patientsafety/taxonomy/icps_full_report.pdf. Accessed December 19, 2018

³ Macchi, L, Pietikäinen E, Reiman T, et al. Patient Safety Management. Available Models and Systems, VTT Working Papers 1459-7683. Available on: https://www.vtt.fi/inf/pdf/workingpapers/2011/W169.pdf. Accessed December 9, 2019.

⁴ World Alliance for patient safety. WHO Draft Guidelines for Adverse Event Reporting and Learning Systems. From information to action. World Health Organisation. Geneva;2005 Available at https://apps.who.int/iris/bitstream/handle/10665/69797/WHO-EIP-SPO-QPS-05.3eng.pdf?sequence=1&isAllowed=y. Accessed November 18, 2019.

⁵ Biquet J, Schopper D, Sprumont D, et al. A call for the application of patient safety culture in medical humanitarian action. A literature review, *J Patient Saf* in press (8/2019)

⁶ Shanks L, Bil K, Fernhout J. Learning without borders: a review of the implementation of medical error reporting in Médecins Sans Frontières. *PLoS One* 2015;10:e0137158.

⁷ Djalali A, Ingrassia P, Corte F, et al. Identifying deficiencies in national and foreign medical team responses through expert opinion surveys: implications for education and training. *Prehosp Disaster Med* 2014;29:364–368.

⁸ Merchant A, Hendel S, Shockley R, et al. Evaluating Progress in the Global Surgical Crisis: Contrasting Access to Emergency and Essential Surgery and Safe Anesthesia Around the World. *World J Surg* 2015; 39:2630–2635.

⁹ Dohlman L. Providing anesthesia in resource-limited settings. Curr Opin Anaesthesiol. 2017;30:496–500. ¹⁰ Rysaback-Smith H. History and Principles of Humanitarian Action. *Turk J Emerg Med* 2016;15(Suppl 1):5-7.

¹¹ Calain P. In search of the "new informal legitimacy" of Medecins Sans Frontieres. *Public Health Ethics* 2012;5:56–66.

12 Brauman, R. La Médecine Humanitaire. Paris: Presses universitaires de France, 2009.

¹³ de Jager E, McKenna C, Bartlett L, et al. Postoperative Adverse Events Inconsistently Improved by the World Health Organization Surgical Safety Checklist: A Systematic Literature Review of 25 Studies. *World J Surg* 2016;40:1842–58.

¹⁴ Emanuel L, Berwick D, Conway j, *et al.* What Exactly Is Patient Safety. In: Henriksen K, Battles JB, Keyes MA, et al. Eds. Advances in patient safety: new directions and alternative approaches. Rockville, MD: Agency for Healthcare Research and Quality 2008;1:1–18.

¹⁵ World Health Organization, Patient Safety: Making Health Care Safer (Geneva: World Health Organization, 2017). Available at: http://www.who.int/patientsafety/publications/patient-safety-making-health-care-safer/en/ Accessed December 19, 2018

¹⁶ Wachter R, and Pronovost P. Balancing "No Blame" with Accountability in Patient Safety. *N Engl J Med* 2009;361:1401–1406.

17 Liston B, Tartaglia K, Evans D, et al. Handoff Practices in Undergraduate Medical Education *J Gen Intern* 2015:29:765–69.

¹⁸ Morain S, Kass N, and Grossmann C, What Allows a Health Care System to Become a Learning Health Care System: Results from Interviews with Health System Leaders. Learn Health Syst 2017;1: e10015.

¹⁹ Castel E, Ginsburg L, Zaheer S, *et al.* Understanding Nurses' and Physicians' Fear of Repercussions for Reporting Errors: Clinician Characteristics, Organization Demographics, or Leadership Factors?. *BMC Health Serv Res* 2015;15:326.

²⁰ Moumtzoglou A. Factors That Prevent Physicians Reporting Adverse Events. *Int J Qual Health Care* 2010;23:51-58.

²¹Health Quality Ontario and others, Patient Safety Learning Systems: A Systematic Review and Qualitative Synthesis, *Ont Health Technol Assess Ser* 2017;17:1.

²² Vrbnjak D, Denieffe S, O'Gorman C, *et al.* Barriers to Reporting Medication Errors and near Misses among Nurses: A Systematic Review. *Int J Nurs Stud* 2016;63:162–78.

²³ Hunt M. Establishing Moral Bearings: Ethics and Expatriate Health Care Professionals in Humanitarian Work. *Disasters* 2011;35:606–22.

²⁴ Ramanujam R, Goodman P. The Challenge of Collective Learning from Event Analysis. *Safety Science* 2011;49:83–89.

<u>Interviewed</u> <u>staff</u>	Nurse	Medical practitioner	Midwife	Surgeon / Anesthetist	Paramedic	Total
MSF-CH	3		2	2		7
MSF-B	2	3	1			6
MDM	4	1	1			6
SC	3	2				5
IRC	2	3			1	6
ICRC	4			1	1	6
Total:	18	9	4	3	2	36

Table 1: Professional background of interviewees

Annex 1: Interview Guide

Interview guide for in-depth interviews

FRONT-LINE STAFF

Date interview	Mode	Place	of		Date	of	Date	of	Identificati
(dd/mmm/yy)	interview :	the		□ Information sheet	transc	ripti	coding		on code of
	□ F2F	intervie	ew:	received:	on		(dd/m	mm	the
					(dd/m	mm	/yy)		interview:
					/уу)				
	🗆 Skype			□ Consent form signed					
				and sent					

	Main Question	Probing Questions (probing questions to help
		interviewers in case of limited or no answer)
Ther	ne 1: Experience of Patient Safe	ty (PS) & Medical Error (ME)
1.1	What is your experience of patient safety in MHA or not?	 a. Have you already heard about this concept? b. Are you familiar with policies implementing PS? c. Is PS an individual, institutional or national responsibility?
1.2	What is, according to you, a ME?	 a. Do you make a difference between ME and adverse event (AE)? b. What are the causes or contributing factors of ME? c. Do you consider that any medical staff does/can do a ME?
1.3	Have you witnessed ME during your humanitarian experience? (Description)	 a. How was it handled? b. Was the management of the case responding to what you considered as appropriate? c. How would you have expected the case to be managed?

Provision of definition and explanations on the concepts of patient safety and medical error

Ther	ne 2: Management of ME in the	ir organisation
2.1	Does your organisation have	
	a policy or strategy on PS?	

2.2	lf yes,	please	provide	a.	How have you been informed?
	informatic	on on this I	PS policy	b.	Do you consider it useful to manage and prevent medical errors?
				c.	What are the missing and the strong points of the policy?
				d.	Which prevention and management procedures exist?
				e.	Is ME a subject of discussion in your organisation?
				f.	
2.2	If no, do yo	ou think it	would be	Wo	uld it be interesting to have such policy? Why
	interesting	g to have a	PS policy		
	to manage	e and prev	ent ME in		
	your orgai	nisation?			

In general terms, what are the needs in terms of the determination and adoption of a PS strategy in MHA?

Then	ne 3: Factors pushing for the ad	optio	on of a PS strategy in MHA
3.1	Is PS pertinent for medical	a.	If yes, what should be developed
	practice in humanitarian settings?	b.	If no, why?
3.2	According to you, which	С.	Ethics (medical and humanitarian)
	factors or considerations	d.	Legal
	should make PS an important		
	concern for aid organisations	e.	Accountability to patients
	providing medical care?	f.	Practical (stated policy)

1	
1	

In global terms, what are the difficulties or challenges in the determination and adoption of a PS strategy in MHA?

Then	ne 4: Factors making the adopti	on of a PS strategy in MHA difficult/not feasible
4.1	Which elements can explain	List local, institutional, international factors
	the absence of a PS policy in your organisation and in	Are the following factors important?
	general in medical	a. Absence of resources at their disposal
	humanitarian organisations?	b. Absence of legal framework on the responsibility of the medical staff in MHA
		c. A blame culture
		d. Absence of willingness from the organisation and/or staff and/or management
		e. No procedures of error reporting and analysis
		f. Precariousness of the medical activities
		g. Consequences in case of disclosure of ME to patient
		h. Absence of compensation framework
		i. Never thought about it
		j. Security risk
		k. Fear of defensive medicine
		I. Too much differences between staff (MoH, organisation, partners)

	Any other?

In global terms, what are the needed conditions for the determination and implementation of a PS strategy in MHA?

Then	ne 5: Necessary ingredients to	allow a	definition & implementation of a PS strategy in
MHA	l l		
5.1	According to you, which (pre-	Sugges	tions:
) conditions should be met to	a.	Legal framework (immunity, partial immunity)
	adapted to MHA?	b.	Training of staff
		C.	Sensitisation of the management
		d.	Sensitisation of staff
		e.	Proper procedures, tools and means
		f.	Absence of blame culture
		g.	A clear institutional policy framing responsibilities and security of the staff
		h.	
5.2	Who is responsible for	a.	Each organisation
	creating these favourable conditions?	b.	Each medical project
		c.	Host state
		d.	WHO
		e.	

5.3	What would you propose to	
	enhance patient safety in	
	Medical Humanitarian	
	action?	
Cond	lusion	
6	Before to conclude, have you	
6	Before to conclude, have you anything you would like to	
6	Before to conclude, have you anything you would like to add?	

Closure of the interview: thanks, and reminder of the confidentiality of the interview.

7. Conclusions and Perspectives

It took a long time, in OECD countries, for patient safety to become a recognized and necessary component of quality care provision and to be translated into policies, methods and practices. If patient safety is in constant evolution and faces many challenges, it responds to the deontological, moral, legal, economic and social obligations of the medical professions and the management of healthcare services. We saw that the example of successful strategies of error prevention in other high-risk industries, exemplified by the aviation industry, ²³ was not sufficient in itself to have the medical profession and institutions immediately embrace the recognition of errors and search for structured strategies to prevent them²⁴. A series of ingredients are considered as essential for a comprehensive investment in patient safety: A strong organisational and leadership support for safety; sensitisation about, and training on, error-reporting systems; a shared vision and blame-free culture.²⁵

Through interviews with medical and para-medical staff active in the humanitarian sector, we found that medical humanitarian action is confronted with the same initial difficulties as health care institutions in OECD countries: the effect of errors committed is rarely visible and therefore not yet a subject of discussion and research; and among medical staff employed by aid actors, as the resident medical staff, in some contexts at least, are highly considered by the communities they work in and their doings are not questioned..

We saw, through the literature review, that patient safety, gradually built up to become a major domain of investments in many countries, notably the high-income countries, is based on the conjunction of actions at three interdependent levels: the macro level, authorities who set the laws and regulations framing the rights and duties of the profession and institutions exercising clinical practices ; the meso level, the medical institutions, enablers of patient safety through training, implementation of tools, commitment and leadership; and the micro level, individuals directly or indirectly

participating in the patient care, whose active involvement is the cornerstone of the prevention and adequate management of medical errors and their consequences.

As in other high-risk industries, an adequate working climate is considered a key ingredient in gaining the active participation of all involved actors in the improvement of safety in healthcare provision. The safety culture²⁶ creates the environment within which the staff involved in clinical care dare to overcome their potential reluctance to speak about incidents, to participate voluntarily in its analysis and learn its lessons.

Beyond the implementation of adequate tools and strategies to detect, report and analyse incidents so as to produce clear recommendations to avoid their recurrence, at medical institutions but also medical departments, units or teams, this safety culture supposes: the clear commitment of the leadership to its enforcement; an active fight against the traditional blame culture; a proper support of the medical profession to not only learn how to methodically analyse incidents but to cope with the difficult experience of having participated in the commission of a perhaps harmful error and being in a position to disclose it appropriately to the patients and/or their relatives.

Interviews with medical and paramedical staff active in the humanitarian field have demonstrated two important lessons confirmed by the medical directors and/or those responsible for quality of care in the organisations involved in the study: 1. there is as yet no strategic thinking about patient safety translating into policies and investments, although there is a clear preoccupation with improving the quality of the care provided or supported by the organisations; 2. there is a willingness on the part of the staff to be trained and supported to implement patient safety as a priority of their organisation.

Otherwise, the explanations provided by the medical and paramedical staff interviewed on the current absence of structured strategies of prevention and management of medical errors can be divided into two categories: 1. the specificities of medical humanitarian action and 2. the absence of the main ingredients of a safety culture (leadership, no-blame culture, appropriate support of the staff in sensitization, training and guiding tools to report, analyse and disclose medical errors). For the latter point, the inspiration of how 65 the healthcare systems and institutions in OECD countries were able to develop and nurture this safety culture seems relevant. Collaborations, or at least consultations, with specialists in patient safety could allow medical aid organisations to build up their appropriate strategies for the development of an internal safety culture.

Related to the specificities of medical humanitarian action, interviewees point additional difficulties to circumvent in order to conceptualize structured and overall strategies of patient safety adapted to their operational realities: the emergency mode of aid organisations (the need to focus on the most pressing among the various other priorities); the highly varied types of projects and work conditions; the countries of intervention having limited resources and being also frequently characterized by social and political instability; the great variety of staff with different medical cultures and a considerable turnover; the limited duration of humanitarian projects (from some months to several years); the absence of regulatory bodies able to develop, impose and monitor patient safety policies in a sector with a multitude of independent aid organisations; and the different levels of control over the care provided to the patients (from projects totally managed by the staff of the humanitarian organisation up to an external support from the ministry of health structure and staff).

The role of medical aid organisations in developing patient safety strategies and medical error management tends to be reinforced by the absence of an overarching regulatory body for the sector coupled with regulatory deficits notably in the establishment of healthcare standards²⁷ in countries of intervention where medical errors are estimated to be very high.^{28,2} Additionally, there is a great variety of medical cultures among the staff of medical aid organisations. Embarking in the definition of adapted patient safety policies requires a clear commitment and investment decided at the higher level of the organisation so that that priority becomes an institutional one. Entrusted with ensuring that the quality of care provided through their projects is accountable to their donors as much as to the beneficiaries of their action, medical humanitarian organisations have the

power and the leverage to conceive, at least for their own actions, policies and strategies ensuring the safety of medical practices they implement and support.

While stating their ambitions in terms of patient safety, a specific subset of the quality of care they intend to ensure, they have the responsibility to set up appropriate support to the front-line medical and paramedical staff. This will come to pass by developing internal mechanisms and tools first, to sensitize the staff regarding the phenomenon of medical errors; second, to train the staff and particularly the medical managers to detect and manage consequences of occurring cases, to report cases in an appropriate manner through appropriate channels; and third to draw lessons from occurring cases to be reinjected into the entire organisation and its project recommendations for implementation. The organisations' leadership has a crucial role in this process, as much in the allocation of resources and means to develop and implement the decided policies as in determining the appropriate support mechanisms for the staff involved in the error making.

The great variety of medical contexts tends to require a high degree of autonomy to allow the project's medical managers to determine how the guiding principles in the organisation's policy can be translated into locally adapted objectives. This supposes that the initial assessment has equipped the field teams with a proper knowledge of local regulations, habits and capacities to, at least, respect local legal requirements and set-up locally realistic plans to improve the patient safety while preserving the security of the staff and the organisation. This tends to reinforce the necessity to invest in sensitisation and training of staff to develop the necessary competence.

In order to have humanitarian staff starting to offer medical services, these organisations must first mobilize financial and human resources, negotiate access in contexts of intervention, ensure the acceptance of their presence and activities by all stakeholders, and more – all challenges demanding substantial energy and capacities before the first medical intervention. Moving into the development and implementation of adapted patient strategies will require dedicated staff as well as extra resources to effect what

could be seen as a real cultural revolution. The active support of donors will therefore be necessary to secure the resource allocations for: the development of internal guidelines; adapted sensitization and training material; training of staff for the analysis of medical error and proposal of recommendations; training of staff on how to disclose the error to the family and/or relatives if appropriate; securing an appropriate arbitrage mechanism to discuss problematic cases; etc.

Initiating discussion of medical error even internally could pose a serious risk for the reputation of the organisation and its staff. It will mean opening a Pandora's box of complex questions. Who is responsible in the event of an error severely affecting a patient – the staff involved or the organisation they work for? When, how and to whom should the detected error be disclosed while respecting ethical precepts and ensuring the security of the staff member(s)? What compensation for the harmed patients is to envisage and what equitable criteria are to be applied across contexts to determine a fair level of compensation? What should or should not be reported to national medical and judicial authorities?

More research and reflection are needed to provide acceptable answers to these various crucial questions.

The expressed desire, by the staff interviewed in the course of this research, of collaboration among aid organisations to develop adapted patient safety and medical error management strategies can be a valuable avenue to pursue in the search for adequate solutions. Such collaboration and experience sharing can easily lead to the development of standardized terminology, sensitization and training tools inspired by the existing training schemes and curricula used in healthcare systems of OECD countries but based on traditional environments and realities encountered in medical humanitarian projects. Adequate options in response to the above questions could be developed under the auspices of the World Health Organisation, which is already providing guidance and standards on patient safety for healthcare systems.

If presently a non-problem (as long as cases of error are not known, disclosed and publicised) absent from cost evaluation, pursuing the definition of a patient safety strategy for humanitarian organisations will represent an important investment in time and energy. It supposes the development of qualitative objectives as well as the usual quantitative ones (number of hospitalisations, of consultations, of vaccinations...) which are naturally scrutinized, be it by the donors wanting to know the result of their investments or by the organisations themselves whose objectives are mainly and legitimately determined by public health considerations.

Medical humanitarian action embodies values of solidarity and humanity by being deployed in places where health systems fail, and by providing vital services in circumstances generally associated with catastrophe: war, natural disaster or huge epidemics like Ebola or cholera. Indeed, it is still associated with heroism.^x This reputation is nevertheless tarnished by various concomitant phenomena: the constant attempts at politicising aid, by which states try to hijack the benefit of the aid deployed to cover the absence of other options at their disposal; an increased resistance from the population affected by crisis, notably owing to false promises or expectations;²⁹ media scandals as "Oxfamgate",^{xi} which create the impression of a gap between public discourse on values and principles and perceived reality. Daring to accept the existence of medical error and to openly discuss may create, if not adequately handled, the risk of increasing distrust.

To tackle a hidden but real phenomenon is a responsibility of organisations that put accountability and quality at the forefront of their identity.^{xii} Confronted with a constant

^{*} Ex.: World Food Program: "What Makes a Humanitarian Hero" available on: <u>https://medium.com/human-development-project/what-makes-a-humanitarian-hero-440202edded8</u> <u>Accessed on October 16</u>, 2019

^{xi} See for instance https://www.thetimes.co.uk/article/the-times-view-on-the-oxfam-scandal-abuse-of-power-kf8w6x2gw accessed 28 August 2019

xⁱⁱ Accountability to beneficiaries is one of the most shared values among aid organisations and is mentioned as a priority in the various sectoral initiatives to develop standards of good practice. For instance, The Code of Conduct for the International Red Cross and Red Crescent Movement and Non-Governmental Organizations (NGOs) in Disaster Relief adopted in 1994 (See https://www.icrc.org/en/doc/assets/files/publications/icrc-002-1067.pdf); The Common Humanitarian 69

need of adaptation,³⁰ medical humanitarian aid organisations have everything to gain by integrating patient safety into their efforts and trying to devise by themselves what is adapted to their realities while ethically and practically sound. It may help to avoid as much as possible the judicialization of occurring cases that occurred in OECD countries,³¹ without having secured mechanisms of prevention and mitigation.

Opening the discussion with national health systems will also open the door to collaboration, to the prevention of unnecessary tensions. This is another area needing further investments. Legal advisors of medical humanitarian organisations may have great added value in understanding various local regulations and practices and in drafting of agreements with local health ministries regarding medical error management, prevention and dispute settlement.^{32,33}

There is no doubt that the incorporation of patient safety through structured strategies will be a long and demanding process. Making a qualitative step from "acting for access to care for vulnerable populations" to "acting for access to quality care for patients belonging to vulnerable populations" will involve numerous challenges as soon as specific proposals and their practical application gain momentum. It must necessarily begin by daring to open the discussion within the sector, one of the ambitions of this research. There is no reason why the "specialness" of the aid system, to paraphrase one of the interviewees, doing a fantastic job, should evade all the rules and the safety procedures expected in quality care provision.

Standards adopted in 2015 (See CHS alliance https://www.chsalliance.org/); The Sphere standards (last edition in 2018: https://spherestandards.org/wp-content/uploads/Sphere-Handbook-2018-EN.pdf).
Annex 1: Oral Presentation during the Geneva Health Forum, April 2016

Medical Humanitarian Action and Patient Safety: The End of an Exception?

Humanitarian actors embody the values of solidarity and humanity by going to those countries whose health systems fail their populations, and by providing vital services. Humanitarian medical staff accept the risks and the difficult working environment that these missions present because they believe it is necessary to respond to suffering wherever it occurs.

Historically the humanitarian sector has grown organically, in response to the demands of emergencies and human suffering. As a result, there is no single, uniform set of norms and standards. However, as the sector has become increasingly institutionalized, there has been a constant push towards professionalization, improved standards, and quality assurance.

There is no doubt – as I have personally witnessed in MSF – that humanitarian medical staff do an admirable job, bearing witness to much more than mere good intentions. But strange though it may seem, "patient safety"^{xiii} has not yet penetrated the aid sector vocabulary and policies, neither as a discipline nor as a central pillar in the prevention of harm to the patient in medical care delivery. A Google search for the terms "humanitarian action" and "patient safety" results in only two articles that mention both concepts.

^{xiii} "We define patient safety as a discipline in the healthcare professions that applies safety science methods toward the goal of achieving a trustworthy system of healthcare delivery. We also define patient safety as an attribute of healthcare systems that minimizes the incidence and impact of adverse events and maximizes recovery from such events." Emanuel L, Berwick D, Conway j, *et al.* What Exactly Is Patient Safety. In: Henriksen K, Battles JB, Keyes MA, et al. Eds. Advances in patient safety: new directions and alternative approaches. Rockville, MD: Agency for Healthcare Research and Quality 2008;1:1–18.

My aim here is to explore how it came to be that humanitarian medical practice has been an exception to the principle of patient safety, a principle that has otherwise been integrated into almost all areas of medicine. I shall also look at the forces that are pushing humanitarian medical providers to ending this exception.

Why Such an Exception?

While there is little consensus about the definition of "humanitarian medicine" or "medical humanitarian action", this sort of medical action is generally associated with catastrophic situations: war, natural disaster, immense epidemics like Ebola or cholera. Medical humanitarian action evokes an image of heroism and urgent response in dangerous situations.

Dr Philippe Calain, an aid actor himself, offered this definition for the concept of humanitarian medicine: "a set of medical or public health practices whose sole intent is to selflessly accommodate and address the tension created between compelling health needs and the ongoing deprivation of resources in a given population or community". ^{xiv} This definition is interesting in that it does not associate the action with acute emergency. It corresponds to this reality: medical humanitarian action is multiform and consists of a range of medical situations from emergencies to support to a healthcare system; from long-term projects to tackle the spread of a disease to providing basic medical assistance to a vulnerable population in a totally stable context.

Taking MSF as the archetypal medical humanitarian actor, we can see that almost half of its projects play out in stable contexts,^{xv} far from active conflicts or sudden disasters. MSF projects are selected for their relevance to the organization's priorities as much as for the medical needs uncovered during situation assessments. The final decision to undertake a

x^{iv} Calain P. In search of the "new informal legitimacy" of Medecins Sans Frontieres. Public Health Ethics 2012;5:56–66.

^{xv} See http://cdn.msf.org/sites/msf.org/files/msf_international_activity_report_2014_en.pdf Accessed on April 12th 2016.

project as well as its objectives and timeframe, are determined by the organisation, even if the process involves negotiation with the local medical authorities and other stakeholders. It is the humanitarian organization itself that decides which resources will be allocated to a project. This decision has implications for other projects, as any allocation of energy or of other means to one project reduces the resources available for another projects. The project performance standards – including standards of care – are frequently left for the organization itself to decide and monitor on its own.

In the vast majority of countries where humanitarian medical aid is deployed, little attention is paid to the concept and principle of patient safety, in terms of frameworks, clear procedures, or policies. It is hardly surprising that the medical humanitarian sector, which has emerged from such operational environments, has been slow to embrace patient safety as a common discipline.

We can delineate three main challenges for the creation of a rigorous patient safety culture in medical humanitarian action.

The Contexts of Intervention

In many contexts, there are more pressing concerns – such as the immediate security of both the patients and the humanitarian staff – that take precedence over considerations of patient safety. Humanitarian medical personnel often work in precarious situations with limited resources and under less than ideal working conditions. It is common to work without any real reference system, with the humanitarian staff solely responsible for the resources mobilized and sent by their organization. Medical cultures differ depending on the context, with widely varying written norms (when they exist at all), numerous ways of defining and perceiving quality of care, and differing levels of expectations (norms/standards). Societies frequently conceive of the medical world's relationship with patients in a paternalistic way: with doctors holding knowledge and power and patients expected to docilely accept whatever treatment is dispensed. In such a context, there is little pressure on the medical providers to focus closely on quality of care. The availability

of suitable qualified local medical staff also varies widely depending on the situation: this too affects the quality of care provided.

Organizational Characteristics

Humanitarian organizations employ heterogeneous, cosmopolitan medical staff, whether recruited locally or internationally. The effect of this is that many different medical cultures are represented within any single organisation. Projects are often planned and implemented extemporaneously, a drawback compounded by short-term engagement of staff members, resulting in considerable turnover both within individual projects and in the organization overall. The relationship with local health systems varies from one situation to another. Some projects use an independent structure created by the humanitarian organization; some are done in full collaboration with, or under the control of, the local health ministry; others are developed using health ministry resources but under full responsibility and supervision of the humanitarian organization.

In these differing contexts, the official frame of responsibilities is often unclear or undetermined. If a general agreement with the local authorities does exist, it often neglects to spell out who holds responsibility for what, and, more importantly, what the consequences might be in the event of a serious breach of care.

The General Context of Humanitarian Medical Action

The humanitarian sector, by and large, is neither structured nor regulated. Each organization operates independently, with no overarching governing body or common sets of rules. There is little published research or documentation on medical humanitarian activities and patient safety. A search on PubMed and Embase, for example, turns up not a single result linking the terms 'patient safety' and 'humanitarian medical aid'.

Potential for the Emergence of Patient Safety Culture in the Medical Humanitarian Sector

While there have been impediments to the development of a rigorous patient safety culture in the medical humanitarian aid sector, we can nonetheless identify some promising developments in that direction. Broadly speaking, there are two main reasons for such an evolution.

Humanitarian Organizations Are Aiming to Become More Accountable to Their "Beneficiaries"

The ethical and quality-assurance framework of many medical humanitarian organizations is rooted in the standards required by the Western world. Both within organizations and across the sector, there is a clear push for professionalization of the aid sector in general as well as for the development of high-quality aid, emphasising the well-being of the patient.

"Do no harm" is a central value to humanitarian action in general, and it reflects perfectly the culture of patient safety in medical ethics. These concepts are inextricably tied to notions of accountability to beneficiaries of humanitarian action, a major subject of discussion across the sector.

A General Trend Toward a Culture of Patient Safety All over the World.

There is an increasing number of examples of medical institutions in low- or middleincome countries implementing measures and tools to improve patient safety. Even if these are isolated instances, they nonetheless reflect a trend towards improved standards in medical care. There is a general demand from societies for more accountability from the humanitarian sector as humanitarian action is viewed with a critical eye from increasingly discerning beneficiaries. Even remotely situated and poor patients have begun to express their concerns and questions about the care they receive from humanitarian medical providers. In those places where expectations of medical standards are high, such as the Middle East, or in contexts where dissatisfied patients are likely to seek judicial intervention, like Haiti (with its proximity to the U.S. culture of litigation), the resulting pressure draws attention to the risks of patients' dissatisfaction and is a strong external factor impelling to better quality of care.

"Do No Harm" as an Operational Value

The scientific literature demonstrates that patient safety is preceded by the development of a more general culture of safety.xvi In settings that cater to individuals who demand a high level of quality, it is crucial that the institutional medical environment be sensitized to patient safety concerns, and that staff work as a team to address these concerns. Increasingly, patient safety is enshrined in health policies in high-income countries, but there is still fear, resistance, and difficulty in shifting from a culture of blame and hiding problems to a culture of learning from one's mistakes. In order for the humanitarian sector to undergo such a paradigm shift, humanitarian health institutions must cultivate a climate favourable to patient safety. Only then will they be able to make this difficult qualitative step.

Medical humanitarian personnel are at the forefront of the response to the pressing needs of victims of crises and emergencies. The development of a patient safety culture, including policies, tools and means, will allow the sector to innovate and increase the quality of its care, and deliver it in a responsible manner. As the Lancet, in a recent editorial, put it: "Patient safety needs to be integrated into the foundation of quality care – safety is not a special programme."xvii

The first step, therefore, is for the humanitarian medical sector to end its exception from the patient-safety groundswell.

^{xvi} "An integrated pattern of individual and organizational behavior, based upon shared beliefs and values, that continuously seeks to minimize patient harm which may result from the processes of care delivery" Aspden, Philip, Institute of Medicine (U.S.), and Committee on Data Standards for Patient Safety. Patient Safety Achieving a New Standard for Care. Washington, D.C.: National Academies Press, 2004. http://search.ebscohost.com/login.aspx?direct=true&scope=site&db=nlebk&db=nlabk&AN=112058.

^{xvii} Patient safety is not a luxury. The Lancet 2016;387:19.

Annex 2: Patient Safety: State of the Art

Developments in Safety Science

Any human activity contains some risk, defined as "the probability that an incident will occur".34 While risk management practice is ancient (noted as far back as the Renaissance period35), the analysis of risks and accidents as a subject of science had to wait until the 1970s to see a flowering of research.36 Similarly, the fallibility of humans has long been recognized (ERRARE HVMANVM EST, attributed variously to the Roman philosophers Seneca and Saint Augustine). Error is a normal cognitive process.³⁷ But, regarding the notion of risk, it is only in the second part of the twentieth century that theories and models of error processes were developed, allowing organized and structured actions of prevention and mitigation.

1.1 James Reason's "Swiss Cheese Model"

The human factor, defined as "the study of the interrelationships between humans, the tools they use, and the environment in which they live and work",³⁸ is one major factor behind accidents (among others such as material defect, environmental variations etc.).

The aviation sector pioneered in research to understand accidents^{xviii} and invested hugely to understand the human factor. Until serious investigations and research were undertaken, accidents were imputed to the pilot.

Scientific analysis and observation enabled researchers to understand that plane accidents, 70% to 80% of which are attributable at least partly to human error,³⁹ are mostly the end result of a series of causes, with the unsafe acts by the crew only the last.⁴⁰ Consequently, prevention measures could be imagined at various levels, which have had a significant effect on the number of accidents and fatalities since the 1970s. Air travel is now one of the safest forms of transportation.^{xix}

 ^{xviii} A specialized agency of the United Nations, the International Civil Aviation Organisation (ICAO) was even created in 1944 to promote the safe and orderly development of international civil aviation throughout the world. See http://www.icao.int/Pages/default.aspx Accessed on September 17, 2018.
^{xix} According to a report of Allianz Insurance company, although the air traffic has constantly grown, in 1959 the chance to have a fatal accident was 1 every 25'000 departure (USA & Canada) and in 2015 is 1 on 77

James Reason, a psychologist, is one of those researchers whose work on human error remains an important reference in the area of safety.⁴¹ With his observations and reflections on possible causes explaining the genesis of errors committed by front-line workers, he initiated a real paradigm shift in the understanding of human error. Human errors are not necessarily the result of a lack of ability on the part of front-line workers; they may also be the culmination of a series of prior errors (at the management level, in the maintenance processes etc.)

Starting by clarifying the separation between intentional and non-intentional actions,⁴² he made the distinction between what he designated as active errors and latent errors. Active errors are those whose effects are in direct relation to the occurrence of the adverse event and are committed by the front-line actor. Latent errors are those whose contributing factors are lying dormant within the system. Such errors are committed by people who are likely unaware that they are participating in a process that will result in an accident.

In a production process, various human errors, even very small, can contribute (contributing factors), each at its own level, to the occurrence of the accident. However, each error alone is not necessarily sufficient, by itself, to cause the accident.

Reason offers also a differentiation between various types of human error: slips (errors made under the influence of fatigue, stress or distraction), lapses (memory error) and mistakes ("deficiency or failure in the judgemental and/or inferential processes involved in the selection of an objective or in the specification of the means to achieve it, irrespective whether the actions directed by this decision-scheme run according to plan").⁴³ Slips and lapses are skill-based errors, while mistakes can be rules-based errors (bad application of the rules) or knowledge-based errors (the person has no idea what to do without clear rules, nor how to react to a specific situation).⁴⁴

²⁹ million departures (USA & EU), 100 time less than bicycle accident risk. http://www.skybrary.aero/bookshelf/books/3297.pdf. Accessed on July 12, 2016

The understanding of this differentiation allows the determination and implementation of specific actions to avoid future errors of the same kind: intelligent decision support systems (checklists), training, clearer written procedures for maintenance, environmental interface design, self-knowledge about error types and mechanisms.⁴⁵ One of the consequences of Reason's approach to error has been, in some high-risk industries, to deviate from the usual search for a guilty person to blame. It offers new options on where to act in the quest to prevent errors:⁴⁶ beyond interventions to enhance the abilities of front-line workers, actions upstream in the complex interaction of humans, machines and systems.⁴⁷

James Reason illustrated his reflections on human error in the famous "Swiss-Cheese Model" (Figure 1), a milestone in safety science.



Figure 1. James Reason: The Swiss Cheese model48

Unawareness and/or unacceptance of errors as being rooted in system processes leads to the "vulnerable system syndrome" making an organisation more error-prone.⁴⁹ Organisations suffering from this syndrome share three main characteristics: the impulsion to search for a guilty party to blame in the event of a problem; the nonrecognition that latent errors are points of vulnerability generated by the system; and a culture of organisation and management focused on productivity without caring for safety.⁵⁰

In Reason's model, accidents are explained by a linear succession of actions or a chain of various elements. There is an accident when factor A + factor B + factor C etc. are activated or because factor A causes factor B which causes factor C..., leading to the accident. This linearity is one of the main sources of criticism of the model,⁵¹ seen as an over-simplification of reality. Despite this criticism, the Swiss Cheese Model remains an essential landmark in safety science and its applications.⁵²

1.2 Rasmussen's Migration of Acceptable Boundaries

The main limit of the linear model proposed by Reason is that it does not sufficiently consider the complexity of human-decision mechanisms. The decision-making process is always a compromise among competing needs, be they from within the organisation (performance, work relations...) or from the outside (legislation, media attention...).⁵³ According to Jens Rasmussen, an engineer working for the nuclear industry, those needs have the potential to act as various sorts of opposing pressures⁵⁴ on the worker (workload, efficiency...).

In Rasmussen's model, the starting point is not the accident but the organisation and the way it is structured and organized,⁵⁵ which create the potential to generate or prevent errors.

Each part of the organisation has its own logic, different from that of other parts, yet produces outcomes influencing the performance of the other parts. The organisation management will have to create the necessary arbitrage among logics in order to limit the 80

potential negative influences that each part of the organisation can have on the others.⁵⁶ This means that decision-makers must analyse the dynamic of each part of a system and the interaction(s) among the parts to arrive at the decision having the least likelihood of erroneous outcome.⁵⁷ In constant evolution, the interaction(s) among the parts of the system can lead to migration of the risks. This then implies constant monitoring of processes to adapt functional safety measures to the evolving nature and place of risks.⁵⁸

Safety is never a given. It is the result of being constantly alert to any evolution and adapting accordingly. In this vision, the intention of the model is not to achieve a zero fault/error organisation but to make it reach an acceptable (low) level of risk. This level of acceptance varies depending on circumstances, if there were just one accident that attracted attention, the tendency would be to act forcefully to keep the risk to a minimum. After a while, attention, and therefore the investment of energy in safety, would tend to diminish, while the risk tolerance increased.⁵⁹

This approach innovates by introducing the notion of competing influences explaining the variations in human decisions that could lead to taking risk up to the commission of error. In a dynamic society, safety is built up at various levels affecting each other in a hierarchical way: government will impose rules and regulations; companies will have their decisions influenced by market evolution; managers will attribute more or less importance and means to safety; staff will have access or not to training and guidance...⁵⁶ It also shifts from the notion of failure or error to the idea of variability of behaviour and human decisions to adapt to the situation.⁵⁹

1.3 High Reliability Organisations

Psycho-sociologists, observing complex organisations, developed the concept of high reliability organisations (HROs). Such organisations or industries, despite the elevated level of complexity of their tasks and their exposure to risks, perform very well, with a very limited number of accidents (nuclear power plants, airplanes etc.).

High reliability organisations have common characteristics regarding their relation to risk.⁶⁰ First, they focus on detecting vulnerabilities (working a priori on risks) in 81

organisational structures or processes: detection of "near misses", those events or situations that could have resulted in an adverse event but did not, either by chance or by timely intervention⁶¹ through notably probabilistic risk assessment.⁶² Thus, they do not limit their attention to accidents that happened (working a posteriori on occurred errors), as these are quite rare.⁶³ Second, they develop multiple redundant protective barriers at recognized vulnerable points of procedures to avoid accidents (technological or human redundant checks). Third, they have created a real culture within the organisation in which everyone shares the same constant concern for safety and respects and trusts each other's competences beyond the hierarchical structure (collective mindfulness).⁶⁴

Theorists of the HRO model have come up with this important notion of an organisational culture that creates the cement among the parts of the organisation in pursuit of a single goal: a safe organisation.⁶⁵ This cementing culture is never a given and must be constantly nurtured.⁶⁶

Such industries take for granted that reality never completely matches what is described and promoted in guidelines. Standardization of processes and procedures is important, but, to respond to all real situations, they must demonstrate the flexibility (even allowing for violations of rules)⁶⁷ necessary to cope with the unexpected by accepting adjustments to what is advocated in guidelines (resilience).⁶⁸ From the point of view of human error, this model of high reliability organisation has shifted theory to a comprehensive and systemic approach:⁴⁸ there is a constant search for the optimization of the interaction between humans and technologies in the best possible design.⁶⁹

Unlike Reason's and Rasmussen's, the high reliability organisation model focuses on risks and looks at successful processes, instead of failures, to extract a set of ingredients favouring safety and explaining the successes of the organisation.⁵¹

1.4 Organisational Approaches to Risk

Drawing on various theories, Amalberti and Vincent¹² categorize organisations according to their relation to risk, in three groups.

1. Ultra-resilient organisations: risks are known and managed. Owing to a careful analysis of previous accidents and their contributing factors, the organisation works at managing the risks and limiting its exposure to them. The resilience of the organisation is based on the teamwork of people sharing the same values and concerns about safety and capable of flexibility and adaptive judgement regarding the most appropriate behaviour (i.e. oil companies, chemical industry).

2. Ultra-adaptive organisations (HRO): risk is a recognized, important and integral part of the activity. The staff must be able to adapt to any circumstance and, even if accidents are frequent owing to high-risk exposure, the staff have the autonomy to make the decision perceived as the most adapted to each situation. There is a limited number of standards and procedures, but also constant sensitization and training to upgrade individual skills, based on the best decision-making processes when confronted with risks. (i.e. fishing industry, combat troops).

3. Ultra-Safe organisations: reliance mainly on the avoidance of risks. They develop their working procedures by standardizing and automatizing them as much as possible. This allows the staff to be interchangeable⁶⁷ (i.e. nuclear industry, civil aviation).

In figure 2, sectors of activities are categorized according to their level of exposure and the adapted style of risk management.^{xx} Those on the left side rely on their capacity to adapt to risk in order to remain performing. Activities or industries on the right side avoid as much as possible the risks.

^{xx} ASA= American Anaesthesiology Association patient status (level 1 = normal healthy patient; level 5 = moribund patient who may not survive without the operation 83





<u>Figure 2:</u> Extract from Amalberti & al. "Average rate per exposure of catastrophes and associated deaths in various industries and human activities "⁷⁰

Safety science has evolved over time in function of various disciplines (psychology, engineering, psycho-sociology, law) and changes in focus: from individual human focus to a systems approach (from Reason to Rasmussen theories), from failures to risks (from Reason & Rasmussen approaches to high reliability organisation theories). This evolution allows a progressively greater overall understanding of risks, their nature and their causes. Their translation into safety and risk-management models requires adaptation to context as much as to the sort of activity. Looking at the complexity and diversity of medical activities, we can imagine that one model of risk management will not fit all sorts of activities, while all of them provide an interesting angle that suits, partially at least, the practice of medicine.⁷¹

Transposition of Safety Theories to Healthcare: Towards Patient Safety 2.1 Late Investment on Healthcare Safety

The recognition of error, as a subject of systemic concern in healthcare, took a long time despite many individual calls³ and legendary initiatives, such as the development of standards of care by Florence Nightingale.⁴ It was first met with scepticism⁷² and is still today greeted with some reticence by the medical professions, as demonstrated by some surveys.⁷³

The term "patient safety" was used for the first time in 1985 in the United States by anaesthesiologists who created the Anaesthesia Patient Safety Foundation (APSF).⁷⁴ But it was the publication of the U.S. Institute of Medicine (IOM) report To Err is Human: Building a Safer healthcare System in 1999⁵ that sparked the interest in "medical errors" and consequently a huge investment in their prevention and management.

How can this interest in safety in healthcare activities, latter day compared to other sectors, be explained? And why not a simple replication of what had been implemented so successfully in the aviation industry?

Four main explanations can be given: 1. the limited impact, in terms of number of affected people, of each individual error in healthcare; 2. the intrinsic characteristics of the medical ethos; 3. the complexity of healthcare provision; and 4. the legal environment creating fears of litigation if errors are disclosed.

A. The Impact of the Error

First, there is a major difference in the scope of errors or accidents occurring in healthcare delivery compared to other high-risk industries. When a plane crashes, there are many victims, while in healthcare, only one patient at a time is harmed.⁷⁵ This lack of "critical mass" visibility has a natural suppressive effect on collective awareness of the extent of the problem.

In parallel, in some countries, there are quite limited external pressures for the disclosure of the errors and for a search for explanations and remedial action. (The USA is an exception.)

It is only when researchers aggregated what was up to then seen as a series of individual cases of error that the limitations of safety in healthcare appeared striking and triggered a general wake-up call: the IOM report of 1999 spoke of between 44,000 to 98,000 Americans dying 85 every year in the United States from medical errors, while up to 1 million patients endured adverse events every year.⁵ In the U.K., a report in 2000, mentioned the figures of 400 deaths and of 10,000 patients enduring serious adverse event per year.⁷⁶ In Australia, according to research in 1985, there were 10,000 deaths each year and up to 50,000 permanent disabilities⁷⁷ due to medical errors.^{xxi}

Parallel to and beyond the assessment of the problem's extent and its impact on the patient, evaluation studies exploring the indirect consequences for society are now important incentives for investments in safety. A study sponsored by the Society for Actuaries estimated the cost of medical error for the United States, for 2008, at \$19.5 billion, of which 80% is additional medical costs: prescriptions, in- and outpatient costs, ancillary services.¹² In the U.K., clinical negligence claims alone cost an estimated at £400 million in 1998-1999.⁷⁸

The increase of mortality rates and loss of productivity (missed workdays) adds \$2.5 billion. In terms of quality adjusted life years (QALYs), the economic loss of deceased patients due to medical error represents up to \$1 trillion per year in the US.⁷⁹ Extrapolating to the entire planet with 421 million hospitalisations per year, there would be up to 42,7 million adverse events causing 23 million DALYs (Disability-adjusted life years) loss per year (two thirds occurring in low- or middle-income countries).²

B. Medical Ethos

A second explanation for the late investment in safety by the medical professions is related to the medical ethos that places the medical doctor in a dominant⁸⁰ position of perceived excellence, reinforced by the asymmetry of access to information, even to the total detriment of the patient.⁸¹ The simple idea of doctors making errors creates a cultural dissonance.⁶ This is embedded not only in medical professionals' minds but also in the beliefs of the patients and, more widely, in the beliefs general population.

xxi According to studies done in comparable healthcare systems, between 700 and 1,700 patients die every year in Switzerland, due to medical error. See http://www.patientensicherheit.ch/fr/th-mes/S-curit--despatients.html Accessed on September 9th, 2016

Further, the medical professions still sometimes consider medicine as craftsmanship needing total autonomy to attain excellence.⁸² Even today, medical culture favours a person-centred approach to human error linked to two illusions: the punishment illusion, according to which the person who makes an error, if punished, will not make it again; and the perfection illusion, claiming that error-free performance is attainable – if one only tries hard enough.⁸³

C. A Complex Human Activity

A third explanation for the slow progress in integrating safety into healthcare design strategy is linked to the complexity of healthcare practice. It not only encompasses different professions, it is often the result of large teams having interactions of different intensity, along different hierarchical lines and even performed in different locations.⁸⁴ Moreover, medicine is expected to develop a response to more and more complex medical problems (co-morbidities, evolution of diseases etc.),¹²rendering the detection and isolation of the causal factors of a problem more difficult.

The rapid evolution of technologies⁸⁵ to assist provision of care imposes a constant reshuffling of the procedures and human-machine interactions and uses ever more varied and varying technological means in its response. Individual staff involved in the medical management of a patient are submitted to constant pressure of using the latest technologies aiming at improving the care provided, which limit the possibilities to develop automatisms gained by experience.

Therefore, a simple transfer of practices or formulae from one organisation or commercial/industrial sector to another may not be suitable even though the pattern or the processes leading to an accident are essentially the same.^{86,75}

D. A Non-Incentive Environment

Healthcare systems and services are costly and under constant pressure to increase the productivity of the diverse services (staff reduction per service, payment per act...).

The risk of litigation associated with the disclosure or the discovery of an error is a strong disincentive at individual and institution levels. Unlike the airline and nuclear industries which are under a regimen of strict liability, where the company, not the individual worker, is 87

responsible for the potential damages in a lot of countries, the medical staff can be considered as guilty of any mistake done in the medical treatment. Even if there is a differentiation between private practice and practice done in public institution (for the latter the medical institution will be the one accountable). In some countries, there are also special funds for medical hazard (Sweden, France, New Zealand), the fear of long and painful legal procedures remains important and maintains expensive malpractice insurance policies.

Despite this general environment, a multitude of research initiatives on error in healthcare appeared after the shock created by the publication of the IOM report, To Err is Human. Scientific medical publications quickly adapted to this new area of research.^{xxii} Medical error is now better understood,¹⁰ is recognized as an important cause of harm (third cause of death in the United States¹¹), and strategies, policies and tools are developed at medical-institution and healthcare-system levels to minimize the risks of error and to manage the occurring cases.¹²

An alignment of pressures, both external (from society) and internal (within healthcare systems), has allowed the development of patient safety policies at institutional and national levels:⁸⁷ adoption of a legal framework; development of national policies with accreditation bodies using patient safety as one of their main criteria; public opinion and media pressures; educational strategies; and voluntary patient safety policies developed inside medical institutions.

^{xxii} In PubMed, the research key "patient safety" for publications in the 60s gives 38 items, 2,452 for the 80s, 10,310 for the 90s, 35,677 for the 00s and 54,071 for the 10s (up to end June 2016)

2.2 Patient Safety as an Attribute of Healthcare Systems

Although it is in hospital care that safety risks are mostly documented, structured and evaluated, patient safety is now an integral part of overall healthcare system concerns and responsibilities.

The concept of patient safety itself has evolved with time, depending on the chosen focus and the evolution of its integration into healthcare design. From a state of "absence of injury",⁵ it has progressively expanded to a process: "The avoidance, prevention, and amelioration of adverse outcomes or injuries stemming from the processes of healthcare"⁸⁸, a frequently used definition of patient safety. It is now considered an attribute of healthcare systems or institutions.¹³

This is intrinsically linked to the concept of quality of care of which it is an important subset (other aspects are hospitality, catering etc., services peripheral to the care itself⁸⁹).

The United Kingdom's NHS, for instance, has integrated patient safety into the definition of quality of care: "The quality of the [health] services is measured by looking at patient safety, the effectiveness of treatments that patients receive, and patient feedback about the care provided."⁹⁰

In the United States, the Institute of Medicine identifies six dimensions of healthcare quality: healthcare must be safe, effective, patient-centred, timely, efficient and equitable.⁹¹

The responsibility of the state⁹² and the level of demand in terms of quality and therefore patient safety are built into policies and strategies affecting healthcare provision at system, institution and individual levels and are so many conditions to delivering healthcare (HAS certification in France).⁹³

2.3 Ingredients of Patient Safety

As already touched upon in the previous chapters, patient safety represents a global change of thinking for some aspects of care delivery. In this chapter, we shall discuss the various elements allowing a full and optimal development of patient safety for the benefit of the overall quality of healthcare, the satisfaction of the care deliverer and the patients benefiting from care services, while contributing to an overall response of efficient healthcare delivery.

Patient safety is developed and implemented at three levels. It is applied, at the micro-level, by individuals/teams who are sensitized, trained and aware of their responsibilities. It is facilitated and supported through strategies developed by institutions, the meso level. And it is framed, at macro-level, by states' laws and policies integrating the expectations, priorities and representations of society.⁹⁴

Although totally interconnected and interdependent, these three distinct levels of responsibilities, development and ingredients will be discussed separately.

2.3.1 At the Systemic Level A. General and Legal Environment

Laws and regulations have an influence (incentives and disincentives) on the reduction of medical errors and the improvement of safety by framing patient safety obligations⁹⁵ of medical staff and institutions and by determining legal and regulatory management of occurring cases.⁹⁶

They first clarify generally the responsibilities of healthcare systems and institutions toward society, notably taking into account demands in terms of risk management⁹⁷ and the nature of contractual links existing between the patient and the consulted medical professional and institution (obligation of means not of results,⁹⁸ implying the respect of certain conditions)^{99.} They also allow identification of who must pay for patient safety and the consequences of an error, a thorny problem.¹⁰⁰

To concretely assist staff to meet ethical requirements of patients' right to information regarding what happened to them, legislation can promote the disclosure¹⁰¹ of errors: In some contexts,¹⁰² (U.S.A.,^{xxiii} U.K., Australia, Scandinavian countries) "apology or sorry laws" provide

^{xxiii} For instance, in the U.S.A., 36 states have such laws. See www.sorryworks.net/apology-laws-cms-143 Accessed on July 2, 2016

some sort of protection to the medical staff so that apologies expressed cannot be used as proofs of malpractice in a court case.¹⁰³

There are still many bridges needed between medical and legal professionals to agree on the legal and regulatory environment providing a fairer and optimal balance between, on the one hand, the search for responsibility and compensation mechanisms and, on the other, the search for measures to favour patient safety and prevention of error.¹⁰⁴

By allowing the anonymity of error reporting, as in Australia, regulations may assist in increasing the number of reported cases.¹⁰⁵

In the same vein of creating a favourable environment for working on reported errors, the determination of responsibility for compensation for alleged harm may have at least carry moral weight in influencing medical staff and medical institutions to set up a proper error reporting system.¹⁰⁶ (New Zealand has adopted a no-fault compensation system to manage complaints of patients of suspected malpractice. While it has affected the number of court cases, there is no tangible evidence that it assists in improving overall patient safety.)¹⁰⁷

Other alternative and complementary systems of conflict resolution to litigation procedures, like mediation,¹⁰⁸ can reduce safety-related costs for health systems in terms of liability costs such as excessive precautions (defensive medicine)¹⁰⁹ and, therefore, may indirectly incentivize reporting of errors.

B. Determination of Rules and Overall Objectives

The state body in charge of the regulation of healthcare systems determines the requirements imposed on health professionals and institutions to be accredited and certified. By developing recommendations and evaluation of health services in terms of quality, the authorities can develop general and national plans of action in the area of patient safety as well as tools and guidelines to assist medical institutions and professionals to actively implement measures for patient safety.

As safety is a moving target¹² owing to the constant evolution of technology and environment and to the difficulty of maintaining efforts underway, including the sensitization of the entire staff (project fatigue), safety programs and projects must constantly be updated and reframed. In healthcare institutions that have not developed maturity concerning the preparation and implementation of safety culture,⁶⁴ an overall and centralised vision assists in selecting the priorities in project development around patient safety.

It is mainly at the overall level that the cost of medical errors is perceptible. Working from claims data for 2008 in the U.S.A., the American Society of Actuaries estimated the cost of medical errors to be \$19.5 billion (\$17 billion for direct cost of care, drugs and hospitalization; \$1.4 billion following deaths due to medical errors, and \$1.1 billion from losses in productivity of affected patients).¹³

C. Interconnection among Elements of the System

By having an overall understanding of trends, strengths and weaknesses of the healthcare system and its components, the national authorities in charge have the capacity to nurture the links between the elements of the system, notably to influence universities and schools to develop the necessary training of individual medical staff to the concern, methods and management of patient safety⁸ and to influence the perception of society on its relation to healthcare expectations and limits⁹ while promoting actions,¹¹⁰

The sensitization and training regarding professional staff on safety matters as on human and systemic contributing factors to errors are conditions allowing them to actively contribute to safety efforts.⁸ Despite an increase in training methods (individual or team training) and curricula (computer-based¹¹¹ or in situ¹¹² simulation...) and in documented evidence of the effect of training,¹¹³ there still seems to be much progress to be made: full integration of patient safety matters into medical students' training curricula¹¹⁴ through provision of enough time and space for it.¹¹⁵

2.3.2 At individual Level

It is at the individual level that health staff contribute to medical errors in as much as they are key to detecting, analysing and reacting to those committed. Their active participation is a requisite for the success of any patient safety efforts.

A. Understanding of Human Nature and Humans' Error-Prone Activity

The active involvement of management staff supposes their belief in the human nature of their thus error-prone work. Through training and/or campaigns of sensitisation, medical staff and other professionals involved in the care system must be made to understand the nature of the problem, its magnitude as well as be convinced of their necessary collaboration in minimizing the risks and manage its consequences.

B. Convinced Regarding the Non-Blame Culture

Beyond the fear of sanctions coming from the hierarchy, staff involved in a medical error express their fear of sanctions from their peers.¹¹⁶ Proper policies, but also demonstration that the medical institution is committed to managing reported errors in a non-judgmental manner, are requisites for staff support of a collective investment in patient safety.

C. Sensitized Regarding the Possibility of Learning from Error

To have the staff willing to report errors and/or to detect points of vulnerability in healthcare procedures, they must be convinced that such reporting will not only assist in managing cases in the best interests of the patients, the institution and its staff, but also that it will effectively assist learning important lessons that could be useful in other services or circumstances.¹¹⁷

D. Encouraged/Trained to Team Discussion and Work on Errors

Studies have demonstrated that communication (or absence thereof) among involved staff is among the main factors in coming to terms with errors.¹¹⁸

As mentioned earlier, the management of patients involves ever more medical specialisation. For hospitalized patients, teamwork is the basis of care management.

Attention to medical error detection and management implies coordination and communication between specialists and team members.¹¹⁹

2.3.3 At the Institution Level

Researchers still insist on the need to support the staff (with training as much as with psychological support)120 and to develop patient-centred care¹²¹ through an effective organisational culture avoiding "tribalism and lack of engagement".¹²²

As central fixtures in healthcare provision, medical institutions are the link between individual and healthcare system levels, the enablers of patient safety as much as the actors at the crossroads of conflicting imperatives: the quality of care provided and the search for efficiency.

It is at this level that the most mentioned, but the most blurred, characteristic of patient safety is developed: safety culture. This concept is abundantly touched upon in the literature and can be defined as "the product of individual and group values, attitudes, perceptions, competencies and patterns of behaviour that determine the commitment to, and the style and proficiency of, an organisation's health and safety management".¹²³ Safety culture implies a comprehensive investment in the goal of safety: it is less an individual characteristic than a collective one of the health institution at management as well as at unit levels.¹²⁴ Safety culture is implemented within healthcare institutions in mechanisms of reporting and analysis of errors and near miss events, a diminution of the blame culture, a real openness to system latencies etc.¹²⁵

A. No Blame but Just Culture

The direct link between the harmed patient and the medical professionals who oversee his/her medical management creates a sense of guilt and shame that leads to suffering also for the involved staff. They may adopt a defensive approach leading to hiding the facts.¹²⁶

As illustrated by high reliability organisation theorists,⁶⁵ working on safety in a complex organisation means the creation of a safe environment for the staff.¹²⁷ The latter not only must be able to report cases without fear of blame but must be actively encouraged to do so.

Such a safe environment is an essential ingredient of what researchers consider a requisite for the organisation to act in favour of safe healthcare. From a traditional "blame-shame" culture by which actors of a medical error may be sanctioned for what is considered their weakness or a "name-blame-train" culture considering that the actors in healthcare must be retaught what proper attitudes are or action to take, institutions have the responsibility to create a climate within which individual staff and managers absorb system thinking enabling learning from errors. A badly perceived effort in patient safety has two combining effects on error reporting and analysis: a persistent fear to speak to colleagues, a fortiori to patients, of problems occurring in the medical management of the patients,¹²⁸ and suspicion that the search for systemic causes to problems means disempowerment.¹²⁹ It eventually sparks controversies regarding who is responsible¹³⁰ while in fact it should allow a necessary clarification of the accountability of both organisations (systems) and individuals. Improved coordination has to be imagined¹³¹ involving all actors: "when many are in charge, no one is".¹³²

Safety culture must remain a "just culture", a concept responding to the necessity to also consider acts of recklessness for which punishment must be conceived.¹³³

That an entire sector starts to speak of its error, while used to profiling its excellence, is a real paradigm shift.¹³⁴

B. Error Detection System (Detection, Analysis, Feedback)

Applying lessons learned from safety science, medical institutions developed systems of detection and analysis of medical errors; policies and tools of prevention; and assistance mechanisms for the management of the error consequences.

The documentation of errors is developed through different data collection methodologies,¹³⁵ all having advantages and limits, such as voluntary reporting (needing a real commitment from the staff in reporting cases), direct observation of clinical procedures (resource demanding), mortality-morbidity conferences (requiring clear and comprehensive patient files and a clear case definition),¹³⁶ trigger tools¹³⁷ (less labour-intensive but with hindsight bias risk).¹³⁸ Eventually, other sources of data are used to complement these methods: informal and formal patient complaints and medico-legal claims of patients or relatives¹³⁹ as well as questionnaires

sent to clinical practitioners.¹⁴⁰ Often medical institutions use a combination of these methods to get the most complete view of reality.¹⁴¹

Inspired by Reason's research on human error, root causes analytical grids are developed looking at contributing factors.¹⁴² Extended to apply also to system errors,¹⁴³, more comprehensive grids clarify the various contributory factors, such as clusters of root causes (e.g., London Protocol¹⁴⁴ promoted by HAS in France;¹⁴⁵ Yorkshire contributory factors framework¹¹⁸, used mainly for training purposes).

The root cause analysis requires time and some experience.¹⁴⁶ Depending on the priorities and policies of the medical institution, its use is limited to the most serious error cases, often named "sentinel events" (cases having a dramatic consequence for the patient and which trigger an immediate investigation¹⁴⁷).

The more medical errors and" near-misses" reported, the more there is a need to apply a method to determine where to invest energy in a complete root cause analysis and in the development of corrective or preventative measures:¹⁴⁸ classification methods assist in the determination of points of vulnerability or failure in the care process to decide where corrective/preventive actions are needed.¹⁴⁹ A scoring system¹⁵⁰ can be applied to the reported errors or the risks assessed independently of error occurrences.¹⁵¹ Interdisciplinary meetings involving the different decision-making layers of the institution can select its priorities to develop action plans.¹⁵²

Various methods are then promoted to develop action plans maximizing the chance of implementing successful and sustainable improvements in care processes and material used.¹⁵³

The choice of wording to name the sort of incident to be reported is important and subject to debate.¹⁵⁴ Behind the search for the most relevant or precise terms, the choice of concepts is frequently guided by the purpose sought by its user.¹⁵⁵ It is not anecdotal, for at least 3 reasons.

First, for the scope of what is reported: literature on safety and many authors on patient safety use mainly the term "error" referring to the sole human nature of the cause of the problem that has had or could have had negative consequences on the health of the patient.⁸⁹An error is "a generic term to encompass all those occasions in which a planned sequence of mental or physical activities fails to achieve its intended outcome".¹⁵⁶ As specified in the IOM report To err is Human, only a fraction of adverse events is caused by human error.⁵ The choice of terms and their definition can mean the underreporting of cases as sources of learning.¹⁵⁷The extension of reflection on accident causation to beyond the human factor, to embrace system failures in the scope of what must be addressed, introduces a more inclusive term: incident. Eventually to limit the scope of data collection, the concept of "adverse event" is chosen: "an undesired patient outcome that may or may not be the result of an error".¹⁵⁸

A second reason to be careful in the choice of wording is negative connotations associated with terms: "error" carries the notion of guilt¹⁵⁹ and may contribute to limiting the neutrality of the event's analysis and increasing the reluctance to report cases. Finally, the choice of terms and associated definitions, if too narrow, can facilitate or hinder the comparison of collected safety data shared among medical institutions.¹⁶⁰ It also has an important effect on the classification of reported incidents used to determine priorities in the development of corrective actions.¹⁶¹

Concerted efforts led by the WHO¹⁶² brought some standardization of terms, but it nevertheless does not prevent varied interpretations.

C. Support of the Involved Staff

The buy-in of medical staff will be seriously influenced by the way their colleagues, the institution they work for and, more generally, the "system" treat those involved in error reporting¹⁶³ (mostly those involved in error commission). Participating in such an event can be profoundly distressing and also has consequences for future professional capacities. The recognition of the "second victim" phenomenon¹⁶⁴ (if not third victim when considering the institution itself)¹⁶⁵ and the need to provide a proper support is not yet widely recognized¹⁶⁶ or even taught to medical professionals.¹⁶⁷

On the same level, the medical staff having to disclose to patients or their relatives the occurrence of a medical error and its consequences need proper training and a clear procedure designed by the institution to limit the harm to both patients and staff.¹⁶⁸

D. Individual and Organisational Learning

The implementation of a reporting system for errors and near misses may become useful and used if its purpose is clearly stated and shared within the organisation.¹⁶⁹ If not, it risks being perceived as just another administrative burden.¹⁷⁰ With the imperative involvement of staff trained and dedicated to the analysis of reported cases,¹⁷¹ a proper internal retro-information system on lessons learned and strategies chosen to avoid recurrences¹⁷² must be set up: the documentation of errors must be a reporting AND learning system.¹⁷³

It is an organisational responsibility to implement such a feedback loop.¹⁷⁴ It allows the realisation in due time that recommendations lead to implementation¹⁷⁵of system improvements, becoming by itself a "winning staff hearts and minds" strategy feeding a virtuous circle of commitment to safety¹⁷⁶ that must be constantly nurtured. The link with universities and research centres provides opportunities to implement patient safety concerns in all the curricula of the training of present and future staff.¹⁷⁷

The lessons to be taken from error occurrences and analysis imply participation not only of all the front-line staff, such as medical and paramedical specialists that may have different perspectives on the same processes¹⁷⁸, but also of the non-medical staff whose involvement¹⁷⁹ offers eventually a more comprehensive vision of what safety improvements can be implemented and how.¹⁸⁰

E. Leadership

Facing possibly conflicting imperatives of productivity and patient safety implies that managers of medical institutions demonstrate a genuine commitment by providing clear plans and means to achieve safety, while offering a safe environment to discuss and act on the risk of errors.¹⁸¹ Interviews with health system leaders can identify the elements necessary for teaching healthcare systems. Beyond the elements presented here, they must insist on the necessity of strong and visionary leadership as an overall structure enabling all the others.¹⁸²

Reports highlight serious improvements in safety of healthcare since the beginning of the 2000s. (From 2010 to 2014, the overall rate of hospital-acquired conditions declined by 17% in the U.S.A. Improvements were observed in about 60% of safety measures monitored by AHRQ.¹⁸³) But many frustrations are emerging regarding what has (not) been achieved so far.¹⁸⁴ While some authors mention the necessity of allowing time for change, citing aviation industry, which needed 30 years to achieve a real reduction in accidents,¹⁸⁵ numerous studies and analyses present reasons for the slow progress.¹⁸⁶

2.4 Patient Safety: Some Still Problematic Issues

A. The Role of the Patients

The role patients can play in safety improvements in healthcare is still a matter of debate and research in medical circles,¹⁸⁷ despite their undeniable right to be informed of risks and to be full actors in all matter regarding their own health.¹⁸⁸

Patients' perception of the quality of healthcare they receive, as well as of the human – and therefore error-prone – nature of medical acts, has an influence on the extent to which society trusts¹⁸⁹ the healthcare system and the medical profession.¹⁹⁰

There exist many possible organised roles for patients, depending on country or medical institution: the constitution of patients' organisations may be favoured by the national government, as in France,¹⁹¹ where they assist in the development of a fair representation of rights and duties of patients as much as in a real understanding of challenges related to healthcare provision.¹⁹² Programs like the WHO's "Patients for Patients Safety" (PFPS), established in 2005,¹⁹³ aim at empowering patients and making them full actors of their health.

Patients' point of view regarding their own experiences is also a good indicator of the quality and safety of healthcare as it is not limited by professional and organisational boundaries.¹⁹⁴ Patients reported that experience surveys can deliver an interesting picture of the safety of the medical institution¹⁹⁵ and be a source of information on errors that might not have been detected by the medical professionals,¹⁹⁶, especially on preventable medical errors.¹⁹⁷ 99 Therefore, beyond moral considerations, it seems interesting to get patients involved in safety for its multiple potential effects: patient's satisfaction and well-being;¹⁹⁸ their understanding of the risks and limits of medical practice;¹⁹⁹ reputational aspects²⁰⁰ in addition to value added from safety improvements. Such multi-layered potential has to be willingly and strategically organized.²⁰¹

The quality of communication among patients and medical staff and institutions when a serious incident occurs seems also to have an effect on the willingness of patient to complain through costly judicial processes.²⁰² If both professionals and patients seem to favour the disclosure of adverse events,²⁰³ this needs a proper environment and minimal training for the involved staff.²⁰⁴

B. Measurement Strategies

One of the biggest challenges, when working at safety improvements, is developing the ability to measure safety and safety improvements with the proper tools.²⁰⁵ Various measurement methods have been developed over the last decades, all having their limits and advantages.²⁰⁶ Proper evaluations of actions in patient safety tend to show patchy results²⁰⁷ requiring investment in adapted measurement methods even if they are based on some level of interpretation.²⁰⁸ Assessment of safety measurement methods of other industries is again a source of inspiration²⁰⁹ but cannot be transferred blindly. The complexity of healthcare and care services led a prominent charity in the U.K. to conclude a report by showing directions but no definite answers on how to measure the five main dimensions of safety in healthcare (measures of harm, of reliability of the health structure, of the capacity to monitor safety, of the ability to anticipate problems and be prepared, and of the capacity to integrate and learn from safety information).²¹⁰ It remains an open challenge requiring innovation,²¹¹, notably to analyse the cost-benefit of investing in patient safety.²¹²

C. Potential Threats to Patient Safety Investments

The financial pressure on health systems and institutions and the related effect on productivity of healthcare staff do not play in favour of attention to the safety aspect of healthcare, despite the indirect costs for society. The efficiency of methods and strategies adopted to tackle the problem of recurrent errors is also subject to measurement and leads to different appreciations depending on the chosen criteria.²¹³

D. Innovation and Safety

Technology, in the form of assistance to diagnosis, monitoring of patients or data processing, as well as related tools, is a promising source of assistance for the front-line medical staff. But much research must still be done to validate the added value and/or real efficacy of implemented new technological tools.²¹⁴

There is something of a paradox in the literature between the expected positive effect of technology on the decrease of human error and the documented evidence of increased complexity and thus the risk of incidents that it induces.²¹⁵

E. Lack of Research

If the optimization of technology use must be considered,²¹⁶ present health budgets cannot meet the challenge²¹⁷ and will not address the need of research in patient safety.²¹⁸

Moreover, even if quality of care is considered worldwide crucially important,²¹⁹ the investment and attention to patient safety differs from country to country.²²⁰ It is still embryonic in the low- and middle-income countries. While it is known that it must be adapted to suit various health system specificities,²²¹, globalization and the absence of borders for health-related problems require a global and worldwide investment in patient safety.

Most research on patient safety is done in the most developed countries, although there are more and more documented initiatives in poor countries.²²² As culture has an influence on the perception of patient safety as much as relations among team members, more research is necessary to design adapted safety strategies.²²³ Training curricula and tools must also be adapted to the different contexts.²²⁴ Some worldwide projects sponsored by the international community are ongoing²²⁵ (e.g. WHO initiatives like the "Clean your hands"²²⁶ campaign or "Surgical Safety Checklist"²²⁷ promotion). In the area of assistance to developing countries, some donors, such as USAID, are engaged in patient safety promotion²²⁸ and calls for action are emerging.²²⁹

F. Missing Territories

Research and strategies of patient safety are mainly targeting hospital settings. There is still much effort needed in looking at the specificities of other areas of healthcare provision, such as in primary care.¹⁸

Already mentioned is the minimal documentation on patient safety in emergent and developing countries. There is another area of medical practice which seems totally unexplored: medical humanitarian action. Frequently deployed in poor countries by organisations coming from OECD countries, medical humanitarian action has specific characteristics that must be explored to explain this exception and envisage an adapted safety strategy.

Patient safety is now, in OECD countries at least, part of the normal concerns of patients, medical and paramedical practitioners as it is of healthcare institutions and systems. If there is still considerable resistance and discussions on its costs and on the ways to improve current practices, it is more how to achieve safety which is questioned than the very fact that better safety is needed.

The movement toward patient safety of the last two decades, which complements continuous improvements of healthcare in the last centuries, will remain an endless battle owing to the evolution of medicine and the human nature of its practice. The last push related to the input of risk management science was started by the recognition of numerous errors made. It is facilitated by the stability of the work conditions of medical staff: the organisation of care, the constitution of medical teams and the regulations organizing the management of patients is "secured". With those conditions in place, the individual medical and management staff are invited to accept the fallibility of their actions and to revisit them; to detect what may have gone wrong at individual and team levels; and to learn through the establishment of new recommendations for improvement. Patient safety is built on two interlinked investments: the tools and design to work on error management and prevention, and the culture of safety

allowing all components of care management (medical and management staff) to bypass the question of responsibility and work on corrective measures.

Annex 3: Humanitarian Medical Action and Patient Safety

Definition of Medical Humanitarian Action and Links with Patient Safety

A tentative definition of humanitarian medicine or humanitarian medical action could be "the medical activities, allying clinical and public health medicine, practised in situations of humanitarian crisis.²³⁰ It is done selflessly¹⁹ with the sole intent to apply the right of access to healthcare for patients who would otherwise not access it."²³¹

While the label "humanitarian action" is frequently associated with any initiatives aiming at providing help to needy people, in the frame of this research, we focus on medical humanitarian action developed by permanent and structured humanitarian actors (NGOs, international organisations) implementing medical care themselves in keeping with the above definition.

Compared to medicine developed in national healthcare systems, humanitarian medical action has two major differences: It is carried out frequently in highly precarious situations, and it is often implemented in parallel to the existing healthcare system offering services for only a part of the population (the one having geographical, financial and authorized access). It is implemented by actors whose sole interest is to alleviate suffering and to restore the dignity of victims of human or natural disasters or of people who cannot access the existing healthcare services.

Humanitarian action is foremost a struggle to overcome all obstacles to the delivery of aid responding to unattended needs. The aid provided often, therefore, remains a compromise between what should and what could be done.²³²

But patient safety, as conceived and developed in OECD countries' healthcare systems, has not yet "crystallised" as an area of documented initiatives for medical humanitarian actors. There is, for instance, very limited documentation of experiences, practices or questions related to patient safety in humanitarian contexts. A literature review done on PubMed and Embase gives links to only one article addressing directly reporting of medical errors and their management.²² Other articles mainly mention the need of more investment in safety and quality.

Some explanations of this limited focus on patient safety are related to the specificities of this form of medical practice.

Characteristics of Medical Humanitarian Action

We can distinguish a series of characteristics specific to this sort of medical response to patients' needs: the philosophy of humanitarian action, the contexts of medical humanitarian interventions and the characteristics of the medical humanitarian actors themselves.

A. The Philosophy of Medical Humanitarian Action

Medical humanitarian action embodies the values of solidarity and humanity by being developed in those countries whose health systems fail their populations, and by providing vital services in circumstances generally associated with catastrophic situations: war, natural disaster or massive epidemics like Ebola or cholera. It evokes an image of heroism²³³ and urgent response in precarious situations. It refers first to the intention behind the action and remains open to what it is in terms of activities and how they are developed.

But goodwill is not enough to implement good intentions. Past experiences have shown the limits of the action itself and of those of the quality of what is offered to beneficiaries of medical assistance, as in Haiti after the 2010 earthquake, when only 10% of the humanitarian surgical teams deployed were considered competent and prepared to face the needs.²³⁴ Evaluations of responses to major crises of the 1990s and 2000s regularly pointed the finger at the weaknesses of humanitarian action.²³⁵ This has generated an important investment in research of improved quality of action,²³⁶ notably professionalization of staff,²³⁷ development of tools, guidelines,²³⁸ specific training^{xxiv} and creation of joint platforms to analyse and share

 ^{xxiv} Ex. Master in Humanitarian Aid to CERAH Genève http: //www.cerahgeneve.ch/formations.html (accessed 06.06.2014); Master in Humanitarian Assistance of the Tuft University (USA): http: // fic .tufts.edu / education / 105

experiences and lessons learned. ^{xxv} Improvement of humanitarian action, in terms of effectiveness, accountability and quality is a core theme of reflection for aid organisations. For instance, the medical humanitarian organisation Médecins Sans Frontières considers that "its primary responsibility is to improve the quality, relevance and extent of [its] own assistance... Obtaining quality clinical results while maintaining respect for the patient must be the major criteria used to evaluate the progress of our medical practice..."²³⁹

When facing substantial unaddressed medical needs and knowing that the care offered by their organisation is the sole option for the patients, humanitarian actors may decide what is acceptable in terms of quality of care in a debatable way.²⁴⁰ This leads, if not well managed, to a risk of some sort of "charity mentality" through which "good intentions directed towards disadvantaged populations can lead to the misconception that in resource-poor environments, any healthcare is good healthcare, regardless of the quality of services".²⁴¹

B. The Nature of Medical Humanitarian Action

Unaddressed health needs can be found everywhere and be of all sorts. Looking only at the direct provision of medical care by humanitarian organisation staff, medical action is multiform, and consists of a range of medical situations ranging from emergencies to the support to a health system; from long-term projects to tackle the spread of one specific disease to providing medical assistance for a neglected population within a totally stable context.

In a crisis endured by a population, the medical needs responded to are selected by the aid organisation in accordance with their own criteria. The addressed needs can be acute for an important group of population, or not. There is no clarity, either in the literature or in practice on when the action should start and when it should end. The means deployed to tackle the

maha / (accessed 06.06.2014), the NOHA Network of the European Union http://www.nohanet.org/en/ (accessed on 06/06/2014) or Master organized by Deakin

^{xxv} Ex. Creation of ALNAP in 1997 (Active Learning Network for Accountability and Performance in Humanitarian Action) http://www.alnap.org/about/french Accessed on May 23, 2016.
selected objectives are left to the discretion of the aid organisation and are the result of a conscious choice determined by organisation's criteria and the limits imposed by the context. Medical aid, by essence, is a political act: the refusal of the fate of absence of access to healthcare. It is therefore frequently implemented in more or less frontal opposition to, or in rivalry with, local actors and authorities.

C. Contexts of Intervention

Medical humanitarian projects are developed in contexts where the local capacities of healthcare service provision, as other public services, are, at least momentarily, disrupted by events (natural disaster, conflict, societal tensions...) or where society or its rulers exclude some of its members from access to healthcare. These projects are implemented in addition, or as a substitute, to care services provided locally, either in parallel to the healthcare system services or integrated into them with a significant level of autonomy of decision-making regarding the sort of services and how they will be offered.

Frequently, but not always, as demonstrated by humanitarian action for migrants in Europe, the contexts of action are characterised by insecurity and social and political instability.

If medical aid actors must be registered and recognized by local authorities and must abide to the local laws and regulations, notably in terms of quality of care, the clarity of those regulations, as well as that of the mechanisms to implement them, may be limited and differ from one context to another. The events at the origin of the humanitarian action may have disrupted the capacity to ensure standards of care considered locally acceptable.

The vulnerability of the patients coupled eventually with the absence of knowledge of patients' rights owing to the absence of a culture of counter-power, limits the existence of external incentive for the optimisation of the quality of care.

D. Means of Action

The precariousness of the working environment arising from logistical constraints, the limited available and qualified staff, added to the absence of reference possibilities for complex cases, are as many factors explaining the need to constantly push the limits of what can be done to respond to acute needs with limited and therefore less than ideal means.²⁴²

The volatility of contexts of intervention eventually imposes frequent adaptation of the project focus (shift of priority for organisations aiming at responding to the "most urgent" needs of the "most vulnerable"), design (according to access to resources and means) and duration (having an effect on what can be achieved and which services to offer).

The cultural differences also effect the possibility of approaching every situation in the same way²⁴³ for the delivery of care and the determination of the desired quality of relationship between the medical aid actor and the community.²⁴⁴

These specific characteristics of medical humanitarian action impose a constant need to adapt the action and develop innovative solutions to respond in a responsible manner to the needs it wants to address. This is a challenge for which at least some actors recognize that a lot is still to be done: "There is no doubt that we have ignored or failed in various medical issues over time, including a lack of attention to the information given to patients, to consideration of their concerns and choices, to the management of pain, and to the prescription of the most appropriate medicines. We must question our acceptance of this status quo and try to address what we are neglecting today."²⁴⁵

While the search for "zero fault" can become a major obstacle to the risk-taking inherent in crisis settings, the quality of action and the preparation to act according to acceptable standards of care are more and more questioned.

E. Nature of Actors

The humanitarian sector, by and large, is not structured and regulated. Each organisation operates and defines its own mandate independently, with no overarching governing body evaluating what is done and no common set of rules. In the quasi-absence of governance and any institutionalized professional body within the humanitarian sector,²⁴⁶ efforts in search of improvements are made on a voluntary basis. The applied norms are very often only non-binding principles, standards and recommendations. Certification mechanisms^{xxvi} that could

^{xxvi} Like the services of a newly created entity HQAI (for Humanitarian Quality Assurance Initiative) created in 2015. See <u>https://hqai.org/</u> Accessed on November 19, 2019

ensure a minimum of compliance with rules based on "soft law" are not yet sufficiently recognized as legitimate and credible. The process itself remains highly controversial.²⁴⁷

The conditions of work, as well as the limits to locally available and mobilized resources, create a unique set of features difficult to cope with for the medical personnel:²⁴⁸ stress, long working hours and days, limited available technology, small number of staff of various medical culture origins etc. While these difficulties are often experienced with a positive outlook,²⁴⁹ they are also a source of moral distress²⁵⁰ for staff not provided with the necessary capabilities and support.²⁵¹There is a high turnover of personnel in such organisations,²⁵², affecting in turn the capacity to sustain the quality of services.

From this quick panorama, some lessons and consequences in terms of patient safety can be drawn.

First, humanitarian actors are frequently left with the responsibility of determining what is good, notably in terms of quality of care, without other references than the ones they choose.²⁵³ This implies also a great responsibility on the part of the organisations regarding the means deployed to reach an acceptable level of quality of care.

Second, the care provided in such contexts is often constrained by other elements than what medical ethics guidance traditionally responds to: conflicts with the local authorities and/or communities on the action, its intentions and the way it is implemented or additional constraints like insecurity, scarcity of resources, total disorganisation of state functions...²⁵⁴

Third, the diversity of the local health capacities and expectations in terms of standards of care reinforces the difficulty for aid organisation to have standardized approaches and policies²⁵⁵for all contexts of action. There cannot be a "one size fits all" solution, and a blind copying of strategies and practices implemented in OECD healthcare system may not be appropriate.

The present lack of ownership of patient safety by the medical aid sector seems therefore to be related to at least three factors: the specificities of medical humanitarian action, the absence of motivational factors coming from the external environment (media or patient 109 pressure, applied regulatory frame of medical practice...) and the lack of proper research on which lessons learned from policies and practices of patient safety adopted in developed countries' healthcare systems could be adopted or adapted to its specific work environment.

~
\subseteq
σ
S
S
0
Ū
4
\times
еX
nex
nnex

the standard of $ig ullet$ Une faute grave réelle ou potentielle dans le	or harm caused standard des soins prodigués à un patient ou e	erformance of a patients ou des dommages causés à un patien	essionals des patients à la suite de la prestation d'un se	de santé et / ou de professionnels de santé	travaillant en son sein	• Assistance, protection et plaidoyer mis en	sponse to œuvre de manière impartiale en réponse aux	political besoins humains résultant de situations d'urge	politiques complexes et de catastrophes	naturelles.	Sauver des vies, soulager la souffrance et le	math of Jack Maintien de la dignité humaine pendant et à l	suite de crises.	• Les objectifs de l'action humanitaire sont de	human dignity	préserver la dignité humaine pendant et aprè	t and strengthen	uch situations.	l by the préparation à la survenue de ces situations.	
An actual or potential serious lapse in t	care provided to a patient or patients o	to a patient or patients through the pe	health service and/or healthcare profe	working within it. (WHO)		Assistance, protection and advocacy a	undertaken on an impartial basis in res	human needs resulting from complex _f	emergencies and natural hazards. ²⁵⁸		Saving lives, alleviating suffering and \mathfrak{m}	human dignity during and in the aftern	crises. ²⁵⁹	The objectives of humanitarian action	lives, alleviate suffering and maintain h	during and in the aftermath of man-mi	onstural disasters. as well as to prevent	preparedness for the occurrence of su	Humanitarian action should be guided	
Accident médical (Medical Mishap)						Action humanitaire/ Assistance humanitaire	(Humanitarian action/Humanitarian assistance						-			-				

humanitarian principles of humanity, meaning the	L'action humanitaire doit être guidée par les
centrality of saving human lives and alleviating	principes humanitaires d'humanité, ce qui signifie
suffering wherever it is found; impartiality, meaning	la centralité de sauver des vies humaines et de
the implementation of actions solely on the basis of	soulager la souffrance partout où elle se trouve;
need, without discrimination between or within	l'impartialité, ce qui signifie la mise en œuvre des
affected populations; neutrality, meaning that	actions uniquement sur la base des besoins, sans
humanitarian action must not favour any side in an	discrimination entre ou au sein des populations
armed conflict or other dispute where such action is	touchées; la neutralité, ce qui signifie que l'action
carried out; and independence, meaning the	humanitaire ne doit favoriser aucune partie à un
autonomy of humanitarian objectives from the	conflit armé ou autre conflit où une telle action
political, economic, military or other objectives that	est effectuée; et l'indépendance, ce qui signifie
any actor may hold with regard to areas where	l'autonomie des objectifs humanitaires des
humanitarian action is being implemented.	objectifs politiques, économiques, militaires ou
Humanitarian action includes the protection of	autres qu'un acteur pourrait avoir dans les zones
civilians and those no longer taking part in hostilities,	où l'action humanitaire est mise en œuvre.
and the provision of food, water and sanitation,	L'action humanitaire englobe la protection des
shelter, health services and other items of assistance,	civils et qui ne prennent plus part aux hostilités, et
undertaken for the benefit of affected people and to	la fourniture de nourriture, d'eau et
facilitate the return to normal lives and livelihoods. 260	d'assainissement, des abris, des services de santé
	et d'autres biens d'assistance, entrepris pour le
	bénéfice de personnes affectées et de faciliter

<u><u><u></u><u></u><u></u><u></u><u></u><u></u><u></u><u></u><u></u><u></u><u></u><u></u><u></u><u></u><u></u><u></u><u></u><u></u></u></u>	lumanitarian assistance is aid to an affected opulation that seeks, as its primary purpose, to save ves and alleviate suffering of a crisis-affected opulation. Humanitarian assistance must be rovided in accordance with the basic humanitarian rinciples of humanity, impartiality and neutrality. ²⁶¹	leur retour à la vie normale et aux moyens de subsistance. • L'aide humanitaire est une aide à une population touchée qui vise, comme objectif principal, à sauver des vies et alléger les souffrances d'une population touchée par la crise. L'aide humanitaire doit être fournie conformément aux principes humanitaires fondamentaux d'humanité, d'impartialité et de neutralité.
st culture) A an an mu	i just culture reconciles professional accountability nd the need to create a safe environment to report nedication errors; seeks to balance the need to learn rom mistakes and the need to take disciplinary ction. ¹³³	 Une culture juste réconcilie la responsabilité professionnelle et la nécessité de créer un environnement sûr pour signaler les erreurs de médication ; cherche à équilibrer la nécessité d'apprendre de ses erreurs et la nécessité de prendre des mesures disciplinaires

The phrase "just culture" was popularized in the	• L'expression « culture juste » a été popularisée
patient safety lexicon by a report (Marx D.) that	dans le lexique de la sécurité des patients par un
outlined principles for achieving a culture in which	rapport (Marx D.) qui établit les principes
frontline personnel feel comfortable disclosing	d'atteindre une culture dans laquelle le personnel
errors—including their own—while maintaining	de première ligne se sente confortable de
professional accountability. In summary, a just culture	divulguer les erreurs, y compris les-leurs-tout en
recognizes that competent professionals make	conservant la responsabilité professionnelle. En
mistakes and acknowledges that even competent	résumé, une culture juste reconnaît que les
professionals will develop unhealthy norms	professionnels compétents font des erreurs et
(shortcuts, "routine rule violations"), but has zero	reconnaît que même les professionnels
tolerance for reckless behaviour. (AHRQ)	compétents peuvent développer des actions
	malsaines (raccourcis, "des violations des règles
	de routine"), mais a une tolérance zéro pour
	comportement imprudent.
A culture in which frontline personnel are comfortable	 Une culture dans laquelle le personnel de
disclosing errors, including their own, while	première ligne se sentent à l'aise de divulguer les
maintaining professional accountability, recognizing	erreurs, y compris les-leurs, tout en conservant la
individual practitioners should not be held	responsabilité professionnelle, reconnaissant que
accountable for system failings over which they have	des praticiens individuels ne doivent pas être
no control, yet does not tolerate conscious disregard	tenus pour responsables des défaillances du
of clear risks to patients or gross misconduct. ²⁶²	système sur lesquels ils n'ont pas de contrôle,

mais ne tolère pas le mépris conscient des risques évidents pour les patients ou les fautes graves.		 Un modèle intégré du comportement individuel et organisationnel, basé sur des croyances et des valeurs partagées, qui cherche constamment à minimiser les dommages au patient qui peuvent résulter de processus de prestation de soins. 	 [Une culture qui présente ce qui suit] cinq attributs de haut niveau que [les professionnels de soins de santé] visent à opérationnaliser à travers la mise en œuvre de systèmes de gestion de la sécurité fortes. (1) Une culture où tous les
	-a culture de sécurité des soins est une production collective. Elle désigne les valeurs et les croyances communes à un groupe en matière de sécurité, et qui constituent un cadre commun aux comportements (HAS)	An integrated pattern of individual and organizational behaviour, based upon shared beliefs and values, that continuously seeks to minimize patient harm which may result from the processes of care delivery. (Aspden, CoE)	A culture that exhibits the following] five high-level attributes that [healthcare professionals] strive to operationalize through the implementation of strong safety management systems. (1) A culture where all workers (including front-line staff, physicians, and
	Culture de la Sécurité (Safety Culture)		

sécurité d'une organisation.	
compétence de gestion de la santé et de la	management. ²⁶³
déterminent l'engagement, et le style et la	proficiency of, an organization's health and safety
compétences et des modes de comportement qui	determine the commitment to, and the style and
collectives, des attitudes, des perceptions, des	competencies, and patterns of behaviour that
sécurité est le produit des valeurs individuelles et	individual and group values, attitudes, perceptions
 La culture de la sécurité d'une organisation de la 	The safety culture of an organization is the product of
sécurité efficaces.	
la responsabilité de maintenir des systèmes de	
fournit des ressources appropriées, la structure, e	
organisationnel d'accidents. (5) [Une culture qui]	Networks)
culture qui] prévoit l'apprentissage	safety systems. (Forum of End Stage Renal Disease
la résolution des problèmes de sécurité. (4) [Une	structure, and accountability to maintain effective
récompense l'identification, la communication et	(5) [A culture that] provides appropriate resources,
opérationnels. (3) [Une culture qui] encourage et	provides for organizational learning from accidents.
au-dessus des objectifs financiers et	resolution of safety issues. (4) [A culture that]
visiteurs. (2) [Une culture qui] priorise la sécurité	and rewards the identification, communication, and
mêmes, de leurs collègues, des patients et des	and operational goals. (3) [A culture that] encourages
acceptent la responsabilité ou la sécurité d'eux-	(2) [A culture that] prioritizes safety above financial
ligne, les médecins et les administrateurs)	themselves, their co-workers, patients, and visitors.
travailleurs (y compris le personnel de première	administrators) accept responsibility or the safety of

Organizations with effective safety cu	ultures share a	 Les organisations avec des cultures de sécurité
constant commitment to safety as a t	top-level priority,	efficaces partagent un engagement constant de la
which permeates the entire organizat	tion. Noted	sécurité comme une priorité de haut niveau, qui
components include (1) acknowledgm	nent of the high-	imprègne l'ensemble de l'organisation. Cela
risk, error-prone nature of an organiz	cation's activities,	implique notamment les composants suivants (1)
(2) a blame-free environment where i	individuals are	la reconnaissance du risque élevé générés par les
able to report errors or close calls wit	thout	activités sujettes à erreurs d'une organisation, (2)
punishment, (3) an expectation of col	llaboration	un environnement sans blâme où les individus
across ranks to seek solutions to vuln	ierabilities, and	sont en mesure de signaler des erreurs ou des
(4) a willingness on the part of the org	ganization to	échappées belles sans être punis, (3) une attente
direct resources to address safety cor	ncerns. ²⁶⁴	de collaboration à travers les niveaux pour
		chercher des solutions aux vulnérabilités, et (4)
		une volonté de la part de l'organisation pour
		orienter des ressources pour répondre aux
		préoccupations de sécurité.
Culture of Safety: The result of an org	ganizational	 Culture de la Sécurité : Le résultat d'un
commitment to safety permeating all	l levels from	engagement de l'organisation à la sécurité
frontline personnel to executive man	agement.	imprègne tous les niveaux, du personnel de

	Features of a culture of safety include	première ligne à la haute direction. Les
	acknowledgment of the high-risk, error prone nature	caractéristiques d'une culture de la sécurité
	of an organization's activities, a just environment	comprennent la reconnaissance du risque élevé,
	where individuals are able to report errors and near	les activités par nature sujettes à erreurs d'une
	misses without fear of reprimand or punishment, an	organisation, un environnement juste là où les
	expectation of collaboration across ranks to seek	individus sont capables de rapporter des erreurs
	solutions to vulnerabilities and a willingness on the	et des accidents évités de justesse, sans crainte de
	part of the organization to direct resources for	réprimande ou punition, une attente de la
	addressing safety concerns. (CPS)	collaboration à travers les niveaux pour chercher
		des solutions aux vulnérabilités et une volonté de
		la part de l'organisation de diriger les ressources
		pour répondre aux préoccupations de sécurité.
	Safety culture refers to the beliefs, values and	 La culture de sûreté se réfère à des croyances,
	attitudes of patient safety shared by all members of	des valeurs et des attitudes de la sécurité du
	the organisation. These shared values are reflected in	patient partagées par tous les membres de
	the day to day operations of the organization. ²⁶⁵	l'organisation. Ces valeurs communes sont
		reflétées dans les opérations quotidiennes de
		l'organisation.
Défaillance active (Active Failure)	Actions or processes during the provision of direct	 Actions ou processus au cours de la prestation
	patient care that fail to achieve their expected aims,	de soins directs aux patients qui ne parviennent
	for example, errors of omission or commission. While	pas à atteindre leurs objectifs attendus, par
119		

	some active failures may contribute to patient injury,	exemple, les erreurs d'omission ou de
	not all do. (Wade, 2002; Davies,2003) (COE)	commission. Alors que certaines défaillances
		actives peuvent contribuer à blesser le patient,
		toutes ne le font pas.
	Active failures are sometimes referred to as errors at	
	the sharp end, figuratively referring to a scalpel. In other words, errors at the sharp end are noticed first	 Les deraillances actives sont parfois appelees erreurs à « l'extrême pointe », au sens figuré se
	because they are committed by the person closest to the nationt (AHRO)	rererant a un scalpel. En d'autres termes, les erreurs à l'extrême pointe sont remarquées
		d'abord parce qu'elles sont commises par la
		personne la plus proche du patient.
	An error that is precipitated by the commission of	 Une erreur qui est précipité par la commission
	errors and violations. These are difficult to anticipate	d'erreurs et violations. Celles-ci sont difficiles à
	and have an immediate adverse impact on safety by	anticiper et ont un impact négatif immédiat sur la
	breaching, bypassing, disabling existing defences.	sécurité en violant, en contournant ou en
	(Khoja)	désactivant les défenses existantes.
Dommage ou Maladie latrogène (latrogenic	Any undesirable condition in a patient occurring as	 Toute condition indésirable survenant chez un
injury) / ou Nosocomiale (nosocomial injury)	the result of treatment by a physician (or other health	patient à la suite d'un traitement par un médecin

a santé) relatives à	e résultant d'une	u d'un autre élément		ecevant des soins ou	tion de soins de santé.	u causées par un	c medecin »), y olontaires ou inutiles	t de tout aspect de la	/ compris les	s de commission ou		fessionnelles des	nt, des professionnels	oliqué à des troubles	
(ou autre professionnel de l	une maladie ou une blessur	procédure, de la thérapie, c	des soins.	Une infection acquise en	des services dans l'organisa	Blessures provenant de or	médecin (iatros, grec pour < compris des dommages inv	ou des souffrances résultan	gestion des soins de santé,	problèmes découlant d'acte	d'omission.	 résultant des activités pro 	médecins, ou, plus largeme	de la santé. Initialement apl	
professional)Pertaining to an illness or injury	resulting from a procedure, therapy, or other element	of care. (COE)		An infection acquired while receiving care or services	in the healthcare organization. (Aspden)	Injury originating from or caused by a physician	(iatros, Greek for "physician"), including unintended	or unnecessary harm or suffering arising from any aspect of healthcare management including	broblems arising from acts of commission or omission.	(Aspden)		Resulting from the professional activities of physicians. or. more broadly. from the activities of	health professionals. Originally applied to disorders	induced in the patient by autosuggestion based on	
															121

	a physician's examination, manner, or discussion, the	induits chez le patient par l'autosuggestion basée
	term is currently applied to any undesirable	sur l'examen, la manière, ou la discussion d'un
	condition in a patient occurring as a result of	médecin, le terme est actuellement appliqué à
	treatment by a physician (or other health	toute situation indésirable chez un patient
	professional), especially to infections acquired by the	survenue à la suite d'un traitement par un
	patient during the course of treatment. ²⁶⁶	médecin (ou un autre professionnel de la santé),
		en particulier aux infections acquises par le
		patient au cours du traitement.
	Pertaining to an illness or injury resulting	 Relatif à une maladie ou un dommage résultant
	trom a procedure, therapy, or other element of care. (Khoja)	d'une procédure, thérapie, ou autre élément des
		soins.
Erreur (error)	A generic term to encompass all those occasions in	 Un terme générique pour englober toutes ces
	which a planned sequence of mental or physical	occasions dans lesquelles une séquence planifiée
	activities fails to achieve its intended outcome. ¹⁴⁹	d'activités mentales ou physiques ne parvient pas
		à atteindre son résultat escompté.
	The failure of a planned action to be completed as	 L'échec d'une action planifiée à s'achever
	intended or use of a wrong, inappropriate, or	comme prévu ou l'utilisation d'un plan mauvais,
	incorrect plan to achieve an aim ¹⁵⁷	inapproprié ou incorrect pour atteindre un
		objectif.

 Déviation dans un processus de soins qui peut, ou non, causer un préjudice aux patients 	 Un acte de commission ou d'omission qui a causé ou contribué à la survenue d'une blessure 		
Deviation in a process of care that may or may not cause harm to patients ²⁶⁷	An act of commission or omission that caused, or contributed to the cause of, the unintended injury ²⁶⁸	-'erreur découle d'une omission ou d'un acte non ntentionnel, mais ne relève pas de l'infraction d'une « règle » ²⁶⁹	 'exécution non conforme d'un acte prévu ou 'application d'un plan incorrect, peut se produire orsque, pendant la planification ou l'exécution, on fait quelque chose alors qu'il ne fallait pas le faire (erreur par commission) ou lorsqu'on ne fait pas quelque chose alors qu'il fallait le faire (erreur par

 L'échec d'une action à se finaliser comme prévu L'échec d'une action à se finaliser comme prévu (par exemple, erreur d'exécution), et l'utilisation d'un mauvais plan pour atteindre un but (cà-d. erreur de planification) (Institute of Medicine, 2000). Il comprend également l'échec d'une action imprévue qui aurait dû être finalisée (omission). 	
Une erreur a été définie comme étant « une action planifiée qui n'a pas été exécutée comme prévu (à savoir, erreur d'exécution) ou l'utilisation d'un mauvais plan pour atteindre un objectif (à savoir, erreur de planification) ». ¹⁴⁷ erreur de planification) ». ¹⁴⁷ The failure of a planned action to be completed as intended (i.e., error of execution), and the use of a wrong plan to achieve an aim (i.e., error of planning) (Institute of Medicine, 2000). It also includes failure of an unplanned action that should have been completed (omission). ²⁷¹	Une erreur est définie comme l'exécution non conforme d'un acte prévu ou de l'application d'un plan non correct" *erreur par commission : quand pendant la planification ou l'exécution d'une action on fait quelque chose qu'il ne fallait pas faire

	* erreur par omission : lorsque l'on	
	ne fait pas quelque chose que l'on aurait	
	dû faire ¹⁵⁶	
Erreur (Mistake)	In some contexts, errors are dichotomized as slips or	 Dans certains contextes, les erreurs sont
_	mistakes, based on the cognitive psychology of task-	divisées en erreurs d'attention ou méprises, selon
	oriented behaviour. Mistakes reflect failures during	la psychologie cognitive du comportement
	attentional behaviours—behaviour that requires	attentionnel. Les méprises reflètent des échecs
	conscious thought, analysis, and planning, as in active	lors de comportements attentionnels-
	problem solving. Rather than lapses in concentration	comportement qui exige la pensée, l'analyse et la
	(as with slips), mistakes typically involve insufficient	planification conscientes, comme dans la
	knowledge, failure to correctly interpret available	résolution active de problèmes. Plutôt que de
	information, or application of the wrong cognitive	manque de concentration (comme avec les
	heuristic or rule. (AHRQ)	erreurs d'attention), les méprises impliquent
		généralement des connaissances insuffisantes,
		une insuffisance d'interprétation correcte des
		informations disponibles, ou l'application d'une
		mauvaise règle ou heuristique cognitive.
	Deficiency or failure in the judgmental and/or	 La carence ou l'échec dans les processus de
	inferential processes involved in the selection of an	jugement et / ou déductif impliqués lors de la
-	objective or in the specification of the means to	sélection d'un objectif ou dans la spécification des

achieve it, irrespective whether or not t	the actions	moyens pour y parvenir, indépendamment de
directed by this decision- scheme run a	ccording to	savoir si ou non les actions réalisées par ce
plan (Reason, 1990, p.9) ; errors of con	scious though	schéma décisionnel se développent
including rule based errors that occur d	uring problem	conformément au plan (Reason, 1990 , p.9);
solving when a wrong rule is chosen, ar	id knowledge-	erreurs de pensée consciente, y compris des
based errors that arise because of lack	of knowledge	erreurs de règles qui se produisent pendant la
or misinpretation of the problem (Leap	e, 1994). ²⁷²	résolution de problème quand une règle erronée
		est choisie, et erreurs basées sur la connaissance
		qui se posent en raison du manque de
		connaissances ou de mésinterprétation du
		problème (Leape,1994).
A commission or an omission with pote	ntially	 Une commission ou une omission avec
negative consequences for the patient	that would	potentiellement des conséquences négatives pour
have been judged wrong by skilled and		le patient qui aurait été jugée mauvaise par des
knowledgeable peers at the time it occi	ırred,	pairs qualifiés et bien informés au moment où elle
independent of whether there were an	y negative	a eu lieu, indépendamment de savoir s'il y avait
consequences. This definition excludes	the natural	des conséquences négatives. Cette définition
history of disease that does not respon	d to treatment	exclut le développement naturel de la maladie qui
and the foreseeable complications of a	correctly	ne répond pas au traitement et les complications
performed procedure, as well as cases i	n which there	prévisibles d'une procédure correctement
		appliquée, ainsi que les cas dans lesquels il y a un

	is a reasonable disagreement over whether a mistake	désaccord raisonnable quant à savoir si une
	occurred. ²⁷³	méprise est survenue.
	An error that occurs when choosing inappropriate goals through conscious deliberation. The actions may	 Une erreur qui se produit lors du choix d'obiectifs inappropriés par décision consciente.
	conform exactly to the plan, but the plan is	Les actions peuvent être exactement conformes
	inadequate to achieve its indented outcome. (Khoja)	au plan, mais le plan est inadapté pour atteindre
		son résultat escompté.
Erreur active (Active Error)	An error that occurs at the level of the frontline	• Une erreur qui se produit au niveau de
	operator and whose effects are felt almost	l'opérateur de première ligne et dont les effets se
	immediately (Kohn)	font sentir presque immédiatement.
	An error associated with the performance of the	 Une erreur associée à la performance de
	'front-line' operator of a complex system and whose	l'opérateur de la 'ligne de front' d'un système
	effects are felt almost immediately. (Reason, 1990,	complexe et dont les effets se font sentir presque
	p.173) (CoE)	immédiatement. (Reason, 1990, p.173)

Erreur d'attention// Erreur de mémoire	Slips refer to failures of schematic behaviours, or	 Les erreurs d'attention se réfèrent à des
(Slip/lapses)	lapses in concentration (e.g., overlooking a step in a	défaillances de comportements schématiques, ou
	routine task due to a lapse in memory, an experienced	à un manque de concentration (par exemple,
	surgeon nicking an adjacent organ during an	omettre une étape dans une tâche de routine en
	operation due to a momentary lapse in concentration)	raison d'une défaillance de la mémoire, un
	Slips occur in the face of competing sensory or	chirurgien expérimenté entaille un organe
	emotional distractions, fatigue, and stress. Reducing	adjacent lors d'une opération en raison d'une
	the risk of slips requires attention to the designs of	défaillance momentanée de la concentration) Les
	protocols, devices, and work environments—using	erreurs d'attention se produisent quand confronté
	checklists so key steps will not be omitted, reducing	à des distractions sensorielles ou émotionnelles
	fatigue among personnel (or shifting high-risk work	concurrentes, à la fatigue et le stress. Réduire les
	away from personnel who have been working	risques d' erreur d'attention nécessite l'attention
	extended hours), removing unnecessary variation in	sur la conception des protocoles, des
	the design of key devices, eliminating distractions	équipements et de l'environnement de travail en
	(e.g., phones) from areas where work requires intense	utilisant des listes de contrôle-donc les étapes clés
	concentration, and other redesign strategies. Slips can	ne seront pas omises, en réduisant la fatigue chez
	be contrasted with mistakes, which are failures that	le personnel (ou en évitant le travail à risque élevé
	occur in attentional behaviour such as active problem	au personnel qui a travaillé de longues heures), en
	solving. (AHRQ)	supprimant la variation inutile dans la conception
		des équipements clés, en éliminant les
		distractions (par exemple, les téléphones) des
		zones où le travail exige une concentration
		intense, et d'autres stratégies de restructuration.

	Les erreurs d'attention peuvent être comparées à
	des méprises, qui sont les échecs qui se
	produisent dans le comportement attentionnel
	tels que la résolution active de problèmes.
Errors which result from some failure in the execution	 Erreurs qui résultent de quelque défaillance
and/or storage stage of an action sequence, ()	dans l'exécution et / ou l'étape de stockage d'une
largely involving failures of memory, that do not	séquence d'action, () impliquant largement les
necessarily manifest themselves in actual behaviour	défaillances de la mémoire, qui ne se manifestent
and may be only apparent to the person who	pas nécessairement dans le comportement réel et
experience them. (Reason, 1990, p.9); internal events	peut être qu'apparente à la personne qui les
[that] generally involve failures of memory. (Reason,	éprouvent. (Reason, 1990, p.9) ; événements
1997, p.71) (COE)	internes [que] impliquent généralement des
	défaillances de mémoire. (Reason, 1997, p.71)
A slip or lapse occurs when the action conducted is	 Une erreur d'attention ou de concentration se
not what was intended. It is an error of execution. The	produit lorsque l'action menée n'est pas ce qui
difference between a slip and a lapse is that a slip is	était prévu. C'est une erreur d'exécution. La
observable, and a lapse is not. ²⁷⁴	différence entre une erreur d'attention et une
	erreur de concentration est que la première est
	observable, l'autre ne l'est pas.

 Erreurs qui se produisent lorsque les individus fonctionnent en mode automatique. 	 Un échange d'informations fausses ou manquantes ou une mauvaise interprétation ou un malentendu 	 Erreur due au manqué de compétences ou connaissances. 	 Erreur liée à un raisonnement vicié. 	 Echec conscient d'adhérer aux procédures ou règlements.
Errors that occur when individuals are functioning in the automatic mode. (Khoja)	Missing or wrong information exchange or misinterpretation or misunderstanding. ²⁷⁵	Error due to lack of knowledge or skill. (Forum of End Stage Renal Disease Networks)	Une erreur commise par suite d'une défaillance cognitive dans l'évaluation de l'information, l'élaboration des plans et la détermination des conséquences probables d'une intervention prévue (CHUM) Error related to flawed reasoning ²⁷⁶	Conscious failure to adhere to procedures or regulation. (Forum of End Stage Renal Disease Networks)
	Erreur de Communication (error of communication)	Erreur de compétence/maîtrise (Error of proficiency)	Erreur de jugement (error of judgment)	Erreur de non-respect des règles (Error of violation)

	A deliberate – but not necessarily reprehensible –	 Un écart délibéré - mais pas nécessairement
	deviation from those practices deemed necessary (by	condamnable - par rapport à ces pratiques jugées
	designers, managers and regulatory agencies) to	nécessaires (par les concepteurs, les gestionnaires
	maintain the safe operation of a potentially hazardous	et les organismes de réglementation) afin de
	system. ⁷	maintenir la sécurité de fonctionnement d'un
		système potentiellement dangereux.
	Action de transgresser de mauvaise foi une loi, une	
	règle de soin, une obligation, un engagement, un	
	devoir au cours de la prestation de soins et de	
	services dans le domaine de la santé et/ou des	
	services sociaux. (CHUM)	
	Appreciated by the individual as being required by	 Apprécié par l'individu comme étant requis par
	regulation, or necessary or advisable to achieve an	la réglementation, ou nécessaire ou souhaitable
	appropriate objective while maintaining safety and	pour atteindre un objectif approprié tout en
	the ongoing operation of a device or system.	maintenant la sécurité et l'exploitation continue
	(Runciman, 2003)	d'un dispositif ou d'un système.
Erreur de planning (Error of planning)	The original intended action is not correct. (Kohn)	 L'action originellement voulue n'est pas correcte.

Erreur de procedure (Error of procedure)	Procedures were followed with the wrong	•Les procédures sont suivies d'une mauvaise mise
	execution. ²⁷⁷	en œuvre.
Erreur d'exécution (error of execution)	A correct action that does not proceed as intended. (Kohn)	 Une action correcte qui ne se déroule pas comme prévu
Erreur latente (Latent error// System error)	(Ou défaillance latente) : Un défaut dans la conception, l'organisation, la formation ou l'entretien d'un système qui engendre des erreurs et dont les effets sont généralement retardés. (EU)	
	(or latent conditions), refer to less apparent failures of organization or design that contributed to the occurrence of errors or allowed them to cause harm to patients. (AHRQ)	 (ou conditions latentes), se rapporte aux échecs moins apparents de l'organisation ou de la conception, qui ont contribué à l'émergence d'erreurs ou leur a permis de causer un préjudice aux patients.
	Latent failure/Latent error: errors in the design, organization, training, or maintenance that lead to operator errors. They may lie dormant in the system for lengthy periods of time. (Kohn)	 Echec latent / erreur latente : erreurs dans la conception, l'organisation, la formation, ou l'entretien qui mène à des erreurs de l'opérateur. Ils peuvent rester dormants dans le système pendant de longues périodes de temps.

	The delayed consequences of technical design or	 Les effets à long terme de la conception
	organizational issues and decisions. Also referred to	technique ou des questions d'organisation et des
	as latent errors. An error that is not the result of an	décisions. Aussi appelé erreurs latentes. Une
	individual's actions, but the predictable outcome of a	erreur qui n'est pas le résultat de l'action d'un
	series of actions and factors that comprise a	individu, mais le résultat prévisible d'une série
	diagnostic or treatment process. (Khoja)	d'actions et de facteurs que comprennent un
		processus de diagnostic ou de traitement.
Erreur liée au médicament/Erreur	Any preventable event that may cause or lead to	 Tout événement évitable qui peut avoir causé
Médicamenteuse (Medication error // Drug	inappropriate medication use or patient harm while	ou entraîné une utilisation de médicaments
Adverse Event // Medication-related	the medication is in the control of the healthcare	inappropriée ou nuire au patient alors que le
Adverse Events//Administration Error))	professional, patient, or consumer. ²⁷⁷	traitement est sous le contrôle du professionnel
		de santé, du patient ou du consommateur.
	A deviation from the prescriber's handwritten or	 Un écart par rapport à l'ordonnance
	turned medication order or from the order that the	médicamenteuse manuscrite ou dactylographiée
		du médecin ou de l'ordonnance que le médecin a
	prescriber has entered into the computer system.	entré dans le système informatique. Les erreurs
	Medication errors are typically viewed as related to	μο πόdication cont πόνότα]οπιοτί του είδος
	administration of a medication, but they can also	מב ווופטרמנוסוו צסוור צפוופומופווופוור נסווצומפופפא
	include errors in ordering or delivering medication. ²⁷⁸	comme liées à l'administration d'un médicament,
		mais elles peuvent également inclure des erreurs

dans la commande ou l'administration d'un médicament. • Tout événement évitable qui peut causer l'utilisation inappropriée de médicaments ou de compromettre la sécurité des patients			 Une erreur dans les processus de commande, de transcription, de distribution, d'administration ou de surveillance des médicaments, quel qu'en soit le résultat (blessure pour le patient)
Any preventable event that may cause inappropriate nedication use or jeopardize patient safety. (Aspden)	Jne erreur qui survient dans une ou plusieurs étapes lu processus d'utilisation d'un médicament : orescription, transcription, préparation, idministration ou monitorage. (CHUM)	Jne erreur commise dans le processus comprenant la prescription jusqu'à l'administration d'un médicament CHUM)	An error in the processes of ordering, transcribing, dispensing, administering, or monitoring medications, rrespective of the outcome (i.e., injury to the oatient). ²⁷⁹
			, I

A failure of some kind in the process of medication	 L'échec de quelque nature que ce soit dans le
administration. ²⁸⁰	processus d'administration de médication.
A discrepancy between what a physician orders and	•Un décalage entre ce que le médecin prescrit et
what is reported to occur. Types of medication errors	ce qui est rapporté comme s'étant produit. Les
include omission, unauthorized drug, extra dose,	types d'erreurs de médication comprennent
wrong dose, wrong dosage form, wrong rate,	l'omission, le médicament non autorisé, la dose
deteriorated drug, wrong administration technique,	supplémentaire, la mauvaise dose, la mauvaise
and wrong time. An omission medication error is the	forme de dosage, la mauvaise mesure, le
failure to give an ordered dose; a refused dose is not	médicament détérioré, la mauvaise technique
counted as an error if the nurse responsible for	d'administration, et le mauvais timing. Une erreur
administering the dose tried but failed to persuade	d'omission de médication est l'incapacité à
the patient to take it. Doses withheld according to	donner la dose prescrite ; une dose refusée n'est
written policies, such as for x-ray procedures, are not	pas comptée comme une erreur si l'infirmière
counted as omission errors. An unauthorized drug	responsable de l'administration de la dose a
medication error is the administration of a dose of	essayé mais n'a pas réussi à persuader le patient
medication not authorized to be given to that patient.	de la prendre. Les doses non-administrées en
Instances of "brand or therapeutic substitution" are	fonction des politiques écrites, telles que des
counted as unauthorized medication errors only when	procédures de radiographie, ne sont pas
prohibited by organization policy. A wrong dose	comptées comme des erreurs d'omission. Une
medication error occurs when a patient receives an	erreur d'administration de médicament non
amount of medicine that is greater or less than the	autorisé est l'administration d'une dose de
_	

	amount ordered; the range of allowable deviation is	médicament non autorisé à ce patient. Les cas de
*	based on each organization's definition. (Aspden)	« marque ou la substitution de marque ou de
		traitement » sont comptés comme des erreurs de
		médication non autorisée seulement
		lorsqu'interdit par la politique de l'organisation.
		Une erreur de mauvais dosage de médication se
		produit quand un patient reçoit une quantité de
		médicament qui est supérieure ou inférieure à la
		quantité prescrite ; le niveau de déviation permis
		est basé sur la définition de chaque organisation.
*	Any preventable event (i.e., professional practice,	
5	drug products, procedures, systems, prescribing,	•Tout événement évitable (cà-d., pratique
5	order communication, product	professionnelle, produits pharmaceutiques,
_	labelling/packaging/nomenclature, compounding,	procédures, systèmes, prescription,
5	dispensing, distribution, administration, education,	communication de la prescription, étiquetage /
<u>_</u>	monitoring and use) that may cause or lead to	emballages / nomenclature des produits,
	inappropriate medication use or patient harm while	composition, préparation, distribution,
ť	the medication is in the control of the healthcare	administration, l'éducation, la surveillance et
<u></u>	professional, patient, or consumer. ²⁸¹	utilisation) qui peut causer ou entraîner
		l'utilisation inappropriée de médicaments ou de
		nuire au patient pendant le traitement est sous le

		contrôle d'un professionnel de la santé, du patient
		ou du consommateur.
	A deviation from an interpretable written prescription or medication order, including written modification of the prescription made by a pharmacist following contact with the prescriber or in compliance with the pharmacy policy [or] any deviation from professional or regulatory references, or guidelines affecting dispensing procedures. (CoE)	 Un écart par rapport à une prescription écrite interprétable ou d'une ordonnance de médicaments, y compris la modification écrite de l'ordonnance faite par un pharmacien ayant pris contact avec le prescripteur ou en conformité avec la politique de la pharmacie [ou] toute déviation de références professionnelles ou réglementaires, ou des lignes directrices touchant les procédures de distribution.
Erreur par omission (error of omission)	Erreur par omission : lorsque l'on ne fait pas quelque chose que l'on aurait dû faire (Brami) Failing to provide the patient with a medical intervention from which the patient would have likely benefited (Aspden)	 Ne pas fournir au patient une intervention médicale dont le patient aurait probablement bénéficié.

	The failure to administer an ordered dose to a	 L'incapacité à administrer une dose prescrite à
	resident by the time the next dose is due, assuming	un patient au moment où la dose suivante est
	there has been no prescribing error. Exceptions would	due, en supposant qu'il n'y ait pas eu d'erreur de
	include a resident's refusal to take the medication and	prescription. Des exceptions pourront inclure le
	failure to administer the dose because of recognized	refus d'un patient de prendre le médicament et
	contraindications. (Khoja)	l'échec à administrer la dose en raison de contre-
		indications reconnues.
Erreur par Commission (error of commission)	Quand, pendant la planification ou l'exécution d'une	
	action, I'on fait quelque chose qu'il ne fallait pas faire	
	(Brami)	
	An error that occurs as a result of an action taken ²⁸²	 Une erreur qui se produit à la suite d'une mesure prise.
Événements critiques (Critical Event)	En particulier, les incidents graves, indicateurs	
	potentiels d'un dysfonctionnement grave du système,	
	qui pourraient engendrer un préjudice grave pour le	
	patient voire son décès et entraîner une perte de	
	confiance des citoyens à l'égard des services de santé.	
	(EU)	
Evènement Indésirable (Adverse event)	Adverse events are untoward incidents, therapeutic	 Les événements indésirables sont les incidents
	misadventures, iatrogenic injuries or other adverse	fâcheux, mésaventures thérapeutiques, blessures

occurrences directly associated with care or services	iatrogènes ou autres événements indésirables
provided within the jurisdiction of a medical center,	directement associés aux soins ou aux services
outpatient clinic or other facility. Adverse events may	fournis au sein de la juridiction d'un centre
result from acts of commission or omission (e.g.,	médical, clinique externe ou autre établissement.
administration of the wrong medication, failure to	Les événements indésirables peuvent résulter
make a timely diagnosis or institute the appropriate	d'actes commis ou omis (par exemple,
therapeutic intervention, adverse reactions or	l'administration du mauvais médicament,
negative outcomes of treatment, etc.) ⁶¹	omission de faire un diagnostic en temps
	opportun ou d'instaurer l'intervention
	thérapeutique appropriée, les effets indésirables
	ou les résultats négatifs du traitement, etc.)
Il s'agit d'un événement non désiré et imprévisible,	
susceptible de causer un préjudice. C'est la nature du	
préjudice (conséquence) qui détermine la	
classification de l'événement et sa gravité. (CHUM)	
An injury that was caused by medical management or	 Une blessure qui a été causée par la direction ou
complication instead of the underlying disease and	complication médicale la place de la maladie sous-
that resulted in prolonged hospitalization or disability	jacente et qui a abouti à une hospitalisation
at the time of discharge from medical care, or both $^{\rm 283}$	

		או טוטווצבב טע מוו וומווטורמע מע וווטווובוור עב ומ זטו גוב
		des soins médicaux, ou les deux.
	An undesired patient outcome that may or may not oe the result of an error. ²⁶⁹	 Un résultat indésirable pour le patient qui peut ou pas être le résultat d'une erreur.
	An event or omission arising during clinical care and ausing physical or psychological injury to a batient. ^{284,285}	 Un événement ou omission survenant pendant les soins cliniques et causant des blessures physiques ou psychologiques à un patient.
<u>م</u>	A negative consequence of care that results in Inintended injury or illness which may or may not	 Une conséquence négative des soins qui se
<u></u>	lave been preventable. ²⁸⁶	traduit par une blessure ou une maladie imprévue qui peut ou peut ne pas avoir été évitable.
<u> </u>	ivent means a discrete, auditable, and clearly defined occurrence. Adverse describes a consequence of care hat results in an undesired outcome. It does not iddress preventability. ²⁸⁸	 Evènement signifie une occurrence discrète, vérifiable et clairement définie. Indésirable décrit la conséquence de soins qui se traduit par un résultat indésirable. Il ne traite pas l'viabilité.

 Une blessure qui a été causée par la gestion médicale et qui résulte en une invalidité mesurable. 	 Une blessure causée par la gestion médicale (plutôt que par la maladie sous-jacente) qui prolonge l'hospitalisation, produit un handicap au moment de la décharge, ou les deux ; Les El sont causés par des complications du médicament, des infections de plaies et des complications techniques, et celles qui sont dues la négligence [provoquée par] des mésaventures diagnostiques, des mésaventures thérapeutiques, et les événements qui se produisent dans la salle d'urgence. 	 Un événement indésirable survenant dans le cadre de soins médicaux qui produit un changement mesurable de l'état du patient.
An injury that was caused by medical management and that results in measurable disability. ²⁸⁷	An injury caused by medical management (rather than by the underlying disease) which prolongs hospitalization, produces a disability at the time of discharge, or both; AEs are caused by drug complications, wound infections, and technical complications, and those due to negligence [caused by] diagnostic mishaps, therapeutic mishaps, and events occurring in the emergency room. (Segen)	An undesirable event occurring in the course of medical care that produces a measurable change in patient status. ²⁸⁸

An event that results in unintended harm to the patient by an act of commission or omission rather	 Un événement qui donne lieu à des dommages involontaires au patient par un acte de 																		
than by the underlying disease or condition of the patient. (Aspden)	commission ou d'omission plutôt que par la maladie sous-jacente ou l'état du patient.																		
An unintended injury caused by medical management rather than by a disease process. ²⁸⁹	 Une blessure involontaire causée par le traitement médical plutôt que par un processus de la maladie. 																		
Any injury caused by medical care (AHRQ)	•Toute blessure causée par les soins médicaux.																		
An incident in which harm resulted to a person receiving healthcare ²⁹⁰	 Un incident dont résultent des dommages pour la personne recevant des soins. 																		
Situation qui s'écarte de procédures ou de résultats escomptés dans une situation habituelle et qui est ou																			
qui serait potentiellement source de dommages. Il existe plusieurs types d'événements indésirables : les dysfonctionnements (non-conformité, anomalie,																			
ents, les événements sentinelles, les	oresque accidents, les accidents ²⁹¹	t shall mean an event that occurs in • Un événement indésirable désigne un	health professional activity, including evénement aui se produit dans le cadre d	vity or in connection with supply of I'activité professionnelle de la santé, y co	about medicines. Adverse events l'activité préhospitalière ou en relation av	and unknown events and errors that de et l'information sur les médicaments.	y the patient's disease, and which evénements indésirables comprennent d	ul or could have been harmful had événements et des erreurs connus et inco	oided beforehand or for other he sont pas causés par la maladie du pati	occur. ²⁹² qui sont soit nuisibles ou soit auraient pu	nuisibles s'ils ont pu être évités au préala	qui ne sont pas survenus pour d'autres ra	ads to or prolongs a hospitalization, • [Un événement] qui conduit à une	causes death, or is associated with hospitalisation ou la prolonge, contribue	enital anomaly. ²⁹³ provoque la mort, ou qui est associée à u	ou une anomalie congénitale.	e Un événement qui entraîne la mort ou l	of bodily function lasting more than d'une partie du corps ou d'invalidité ou d	de fonction corporelle qui dure plus de se
---------------------------------------	---	---	--	---	--	---	--	--	---	--	--	---	--	---	--	------------------------------	--	---	--
défaut), les incid	précurseurs, les	An adverse even	connection with	pre-hospital acti	and information	comprise known	are not caused b	either are harmf	they not been av	reasons did not			Evènement indésirable grave (Serious Adverse [An event] that l	Event) contributes to o	cancer or a cong		An event that re	disability or loss	

	seven days or still present at the time of discharge	ou encore présent au moment de la sortie d'un
	from an inpatient healthcare facility or, when	établissement de soins de santé en milieu
	referring to other than an adverse event, an event	hospitalier ou, en se référant à autre qu'un
	whose occurrence is grave. (NQF)	événement indésirable, un événement dont la
		survenance est grave.
	Unexpected occurrences involving death or serious	
	physical or psychological injury. Any adverse event	 Evénements inattendus ayant entraîné la mort
	(AE) that is fatal, life-threatening, or permanently	ou une blessure physique ou psychologique grave.
	Disabling, or that results in new or prolonged	Tout événement indésirable (El) qui est mortelle,
	hospitalization. (Khoja)	met la vie en danger, ou handicape de façon, ou
		qui résulte en une hospitalisation nouvelle ou
		prolongée.
Evènement Indésirable Grave Evitable (Never	Never Events are serious, largely preventable patient	 Les évènements indésirables graves évitables
Event)	safety incidents that should not occur if the available	sont des incidents de sécurité graves, en grande
	preventative measures have been implemented. ²⁸⁶	partie évitables, qui ne se seraient pas produits si
		les mesures préventives disponibles avaient été
		mises en œuvre.
Evènement Indésirable Sans Préjudice (No-	When an error does not result in an adverse event for	 Quand une erreur ne se mue pas en évènement
Harm Event)	the patient and the absence of injury is owed to	indésirable pour le patient et que l'absence de
	chance. This differs from a near miss, in which injury is	dommage est dû à la chance. C'est différent des
	absent because the error was "caught." ^{294.}	

		« presque incidents » pour lesqueis il n'y a pas de préjudice puisque l'erreur a été évitée.
Incident (Incident // Occurence) Synonyme d'évènement indésirable	Une action ou une situation qui n'entraîne pas de conséquences sur l'état de santé ou le bien-être d'un usager, du personnel, d'un professionnel concerné ou d'un tiers, mais dont le résultat est inhabituel et qui en d'autres occasions, pourrait entraîner des conséquences. (CHUM) An event that, under slightly different circumstances,	 un événement qui, lors de circonstances légèrement différentes, aurait pu être un
Incident Potentiel (No Harm Event)	Une grave erreur ou un grave incident qui peut éventuellement causer un incident, mais ne le fait pas en raison du hasard ou parce que l'erreur est constatée à temps (aussi appelé « quasi-incident »). (EU)	
Infraction / Violation (Violation) Médecine Humanitaire (Humanitarian Medicine)	Une infraction implique le non-respect délibéré d'une procédure, d'une norme ou d'une règle. (WHO) Médecine qui s'adresse à des populations marginalisées, éprouvées par une crise ou privées	

d'accès aux soins. Celle qui s'exerce sans autre	
objectif que de se rendre utile. ²³¹	
A set of medical or public health practices whose sole	 Un ensemble de pratiques médicales ou de
intent is to selflessly accommodate and address the	santé publique dont le but unique est d'accueillir
tension created between compelling health needs and	et de répondre, de manière désintéressée, à la
the ongoing deprivation of resources in a given	tension créée entre les besoins de santé
population or community. ¹⁹	incontestables et la privation continue d'accès aux
	ressources pour une population ou une
	communauté donnée.
While all medical intervention to reduce a person's	
sickness and suffering is in essence humanitarian,	 Alors que toute intervention médicale pour
Humanitarian Medicine goes beyond the usual	réduire la maladie et la souffrance d'une personne
therapeutic act and promotes, provides, teaches,	est par essence humanitaire, la médecine
supports, and delivers people's health as a human	humanitaire va au-delà de l'acte thérapeutique
right, in conformity with the ethics of Hippocratic	habituelle et favorise, fournit, enseigne, soutient,
teaching, the principles of the World Health	et délivre la santé des gens en tant que droit de
Organization, the Charter of the United Nations, the	l'homme, en conformité avec l'éthique
Universal Declaration of Human Rights, the Red Cross	l'enseignement d'Hippocrate, les principes de
Conventions and other covenants and practices that	l'Organisation mondiale de la Santé, de la Charte
ensure the most humane and best possible level of	des Nations Unies, la Déclaration universelle des

care, without any discrimination or consideration of	droits de l'homme, les conventions de la Croix-
material gain. ²⁹⁵	Rouge et d'autres conventions et pratiques qui
	garantissent le niveau le plus humain et le
	meilleur de soins possible, sans aucune
	discrimination ou considération de gain matériel.
Humanitarian medicine is made up of a wide range of	 La médecine humanitaire est constituée d'un
practices with few obvious connections between	large éventail de pratiques avec quelques
them. Battlefield medicine and surgery, rural	connexions évidentes entre elles. La médecine et
dispensaries in remote areas, campaigns to raise	chirurgie du champ de bataille, les dispensaires
awareness about health problems in poor countries,	ruraux dans les zones reculées, des campagnes de
emergency teams in disaster situations, vaccination	sensibilisation sur les problèmes de santé dans les
campaigns, health education, help for marginalised	pays pauvres, les équipes d'urgence dans les
groups in affluent countries and public health advice	situations de catastrophe, les campagnes de
are just some examples of actions that fall within the	vaccination, l'éducation à la santé, l'aide aux
scope of "humanitarian medicine" when they are	groupes marginalisés dans les pays riches et les
carried out by organisations and in circumstances that	conseils de la santé publique ne sont que
can be classified as "humanitarian". The field is	quelques exemples de actions qui entrent dans le
defined not by a particular set of techniques but by	champ d'application de la «médecine
the setting in which the action takes place and the	humanitaire» quand ils sont effectués par des
stated aim of those involved. This may appear	organisations et dans des circonstances qui
inadequate as a definition insofar as it combines some	peuvent être classés comme «humanitaire». Le

very different areas under a single heading, both from	champ est défini non pas par un ensemble
the point of view of the context in which the action	particulier de techniques, mais par le contexte
takes place and the status of the organisations	dans lequel se déroule l'action et l'objectif déclaré
involved. Heterogeneity, however, is a defining	de ceux qui y sont impliqués. Cela peut paraître
characteristic of humanitarian medical practice. ²⁹⁶	insuffisant comme une définition dans la mesure
	où il combine des domaines très différents sous
	une seule rubrique, tant du point de vue du
	contexte dans lequel se déroule l'action et le
	statut des organisations impliquées.
	L'hétérogénéité, cependant, est une
	caractéristique déterminante de la pratique
	médicale humanitaire.
La médecine humanitaire se définit comme « la	
médecine pour ceux qui n'y ont pas accès » ²³²	
La médecine humanitaire n'est pas une pratique à la	
marge de la biomédecine et de la santé publique,	
mais une tentative de répondre aux attentes de	
populations qui en sont privées alors que leur poids	
démographique est parfois considérable. 297	

Mésaventure médicale (Medical	Personal injury that is caused by medical error or	• Dommage causé par une erreur médicale ou une
Misadventure)	medical mishap. Medical misadventure required the	mésaventure médicale. Une mésaventure
	injury suffered by the claimant to be caused by fault	médicale nécessite que le préjudice subi par le
	on the part of the health professional providing	plaignant soit causé par une faute de la part du
Terme juridique en Nouvelle Zélande jusqu'en	treatment (medical error), or for the injury resulting	professionnel de santé qui fournit le traitement
2005. Remplacé depuis par « Blessure de	from properly given treatment leading to an adverse	(erreur médicale), ou soit le préjudice résultant du
traitement » (Treatment Injury)	consequence to be both rare and severe (medical	traitement dûment donné menant à une
	mishap). ²⁹⁸	conséquence négative étant à la fois rare et grave
		(accident médical).
Négligence/Faute (Medical negligence)	An accident or unintentional act, as in an occupation- related "homicide by misadventure"; in medicine, the term has become an elegant euphemism for a therapeutic error, as in a surgical misadventure in which the wrong leg was amputated. (Segen) An incident causing harm, damage or loss is the result of doing something wrong or failing to provide a reasonable level of care in a circumstance in which one has a duty of care. ²⁹⁹	 Un accident ou acte involontaire, comme un « homicide accidentel » lié à la profession ; en médecine, le terme est devenu un euphémisme élégant pour une erreur thérapeutique, comme dans une mésaventure chirurgicale dans laquelle la mauvaise jambe a été amputée. Un incident, causant des lésions, dommages ou pertes, est le résultat du fait de faire quelque chose de mal ou de ne pas fournir un niveau de soins raisonnable dans une situation où l'on a une

Manque de soin, de vigilance. Manque de precaution.	
(CHUM)	
In law, negligence refers to a breach of a duty of care	 En droit la négligence se réfère à un
owed by one person or entity to another, which if	manutement à rue obligation de soins dû nar rue
breached, results in foreseeable and compensable	personne ou entité à une autre, qui, si manguée.
harm to that other, or to some interest of his or	entraîne des dommages prévisibles et
hers.T132 FPT More narrowly, negligence can refer to	indemnisables à cette autre, ou à un intérêt des
one element of the cause of action for negligence, i.e.	siens. Plus étroitement, la négligence peut se
whether the defendant's conduct met the standard of	référer à un élément de la cause de l'action pour
care. 10/	négligence, à savoir si la conduite du défendeur a
	respecté la norme de soins.
The [British] law of medical negligence operates on	 Le droit [britannique] de la négligence médicale
two principles: that the patient must agree to	fonctionne sur deux principes : le patient doit
treatment and that treatment must be carried out	accepter le traitement et le traitement doit être
with proper skill by the doctors involved. But it holds	effectué avec la compétence voulue des médecins
doctors and other healthcare professionals liable only	impliqués. Mais il tient les médecins et les autres
for that subset of iatrogenic injury that occurs when	professionnels de santé pour responsables de ce
there is a breach of the duty to use reasonable care	sous-ensemble de dommage iatrogène qui se
and, as a consequence, the patient experiences an	produit quand il y a une violation de l'obligation

injury In principle, adverse outcomes consistent	de soins raisonnable et qu'en conséquence, le
with "normal" risk must be borne by the patient. ³⁰⁰	patient encourt un dommage En principe, les
	résultantes défavorables cohérentes avec un
	risque « normal » doivent être supportées par le
	patient.
Care provided failed to meet the standard of care	 Les soins fournis ne répondent pas à la norme
reasonably expected of an average practitioner	de soins raisonnablement attendue d'un praticien
qualified to care for the patient in question. (COE)	moyen qualifié pour soigner le patient en
	question.
The failure (usually on the part of a physician or other	 L'échec (habituellement de la part d'un médecin
healthcare professional) to exercise ordinary,	ou d'un autre professionnel de la santé) d'exercer
reasonable, usual, or expected care, prudence, or skill	des soins raisonnables, ordinaires, habituels ou
(that would usually or customarily be exercised by	attendus, de la prudence, ou de compétences (ce
other reputable physicians treating similar patients) in	qui serait normalement ou habituellement fait par
the performance of a legally recognized duty,	d'autres médecins réputés qui traitent des
resulting in foreseeable harm, injury; or loss to	patients similaires) dans l'exécution d'une
another; negligence may be an act of omission (i.e.,	obligation légalement reconnue, ce qui entraîne
unintentional) or commission (i.e., intentional),	un préjudice prévisible, un dommage; ou une
characterized by inattention, recklessness,	perte à un autre; la négligence peut être un acte

inadvertence, thoughtlessness, or wantonness; in	d'omission (cà-d. non-intentionnelle) ou de la
healthcare, negligence implies a substandard	commission (cà-d. intentionnelle), caractérisée
deviation from the "standard of medical practice" that	par l'inattention, l'imprudence, l'inattention,
would be exercised by a similarly trained professional	l'insouciance ou la sauvagerie; dans les soins de
under similar circumstances. (Segen)	santé, la négligence implique une piètre déviation
	des «standards de la pratique médicale» qui
	seraient appliqués par un professionnel qualifié de
	manière équivalente dans des circonstances
	similaires.
Care that fell below the standard expected of	 Soins tombé en dessous de la norme attendue
physicians in their community. Care provided failed to	de médecins dans leur communauté. Les soins
meet the standard of care reasonably expected of an	fournis n'atteignent pas la norme de soins
average practitioner qualified to care for the patient	raisonnablement attendue d'un praticien moyen,
in question. The failure of an individual to exercise the	qualifié pour soigner le patient en question.
ordinary degree of care that would be expected from	L'échec d'un individu à exercer le degré de soins
a reasonable and prudent person acting in the same	ordinaire qui serait attendu d'une personne
situation, under the same or similar circumstances.	raisonnable et prudente agissant dans la même
This term is generally synonymous with malpractice. A	situation ou dans des circonstances similaires. Ce
legal conclusion that is reached when it has been	terme est généralement synonyme de faute
determined that: (1) the defendant owed a duty of	professionnelle. Conclusion juridique atteinte
care to the plaintiff; (2) the defendant breached the	quand il a été déterminé que : (1) le défendeur

	duty of care; (3) the plaintiff was injured. as a result of	avait une obligation de diligence envers le
	the breach of the duty of care; and (4) legally	demandeur ; (2) le défendeur transgresse le
	cognizable damages resulted from the injury. (Khoja)	devoir de soins dus au plaignant ; (3) le plaignant a
		eu des dommages à la suite de la violation de
		l'obligation de soins ; et (4) un préjudice est
		reconnu par la loi à la suite des dommages.
Presque incident // Quasi incident //	An event or situation that did not produce patient	 Un événement ou une situation qui n'a pas
Echappée belle (Near miss // Close call // No	injury, but only because of chance. This good fortune	produit de blessure au patient, mais seulement
harm incident)	might reflect robustness of the patient (e.g., a patient	parce que dû au hasard. Cette bonne fortune
	with penicillin allergy receives penicillin, but has no	pourrait refléter la robustesse du patient (par
	reaction) or a fortuitous, timely intervention (e.g., a	exemple, un patient atteint d'allergie à la
	nurse happens to realize that a physician wrote an	pénicilline reçoit la pénicilline, mais n'a pas de
	order in the wrong chart). This definition is identical	réaction) ou d'une intervention fortuite rapide
	to that for close call. (AHRQ)	(par exemple, une infirmière arrive à se rendre
		compte que le médecin a écrit un ordre dans le
		mauvais tableau). Cette définition est identique à
		celle « d'échappée belle ».
	An error of commission or omission that could have	 Une erreur de commission ou d'omission qui
	harmed the patient, but serious harm did not occur as	aurait pu nuire au patient, mais le préjudice grave
		ne se produit par chance.

	Un « quasi-incident » est une erreur ou un accident grave qui peut éventuellement causer un incident, mais qui ne le cause pas par hasard ou parce que l'erreur a été constatée à temps. (EU)	
	Un incident qui n'a pas provoqué d'atteinte (WHO)	
	An event or situation that could have resulted in an adverse event but did not, either by chance or through timely intervention. (VA)	 Un événement ou une situation qui aurait pu conduire à un événement indésirable, mais n'a pas, par hasard ou à la suite d'une intervention en temps opportun.
Qualité des Soins	Degree to which health services for individuals and populations increase the likelihood of desired health outcomes and are consistent with current professional knowledge. (Kohn)	 Degré auquel les services de santé pour les individus et les populations augmentent la probabilité de résultats souhaités en santé et qui sont conformes aux connaissances professionnelles actuelles
	Quality of care is the kind of care which is expected to maximize an inclusive measure of patient welfare, after one has taken account of the balance of	 La qualité des soins est le type de soins qui devrait maximiser une mesure inclusive du bien- être patient, après que l'on a tenu compte de

expected gains and losses that attend the process of	l'équilibre des gains et des pertes attendues qui
care in all its parts. ³⁰¹	participent au processus de soins dans toutes ses
	parties.
Healthcare that uses the available and annronriate	 Les soins de santé qui utilisent les ressources
resources in an efficient way to equitably contribute	disponibles et appropriées d'une manière
to the improvement of the health of the populations	efficiente pour contribuer équitablement à
and patients. This implies that provision of care is	l'amélioration de la santé des populations et des
consistent with current professional knowledge.	patients. Cela implique que la fourniture de soins
focuses on the needs and goals of individuals. their	est conforme aux connaissances professionnelles
families and communities prevents and avoid harm	en cours, se concentre sur les besoins et les
related to care and involves nersons/natients as key	objectifs des individus, de leurs familles et des
narthers in the nroress of care ³⁰²	communautés, prévient et évite les dommages
	liés aux soins, et implique les personnes / patients
	en tant que partenaires-clés dans le processus de
	soins.
Quality of care is the level of attainment of health	 La qualité des soins est le niveau de réalisation
systems' intrinsic goals for health improvement and	des objectifs intrinsèques des systèmes de santé
responsiveness to legitimate expectations of the	pour améliorer la santé et la réactivité aux
population. ³⁰³	attentes légitimes de la population.

The degree of conformity with accepted principles	 Le degré de conformité avec les principes et les
and practices (standards), the degree of fitness for the	pratiques acceptées (standards), le degré
patient's needs, and the degree of attainment of	d'adaptation aux besoins du patient, et le degré
achievable outcomes (results), consonant with the	de réalisation des objectifs réalisables (résultats),
appropriate allocation or use of resources. The phrase	conforme avec l'attribution ou l'utilisation des
quality of care carries the concept that quality is not	ressources appropriées. La phrase « qualité des
equivalent to "more" or "higher technology" or higher	soins » soutient l'idée que la qualité n'est pas
cost. The degree of conformity with standards focuses	équivalente à « plus » ou « plus haute technologie
on the provider's performance, while the degree of	» ou à un coût plus élevé. Le degré de conformité
fitness for the patient's needs indicates that the	avec les standards se concentre sur la
patient may present conditions that override strict	performance du prestataire de soins, tandis que le
conformity with otherwise prescribed procedures.281	degré d'adéquation aux besoins du patient
	indique que le patient peut présenter des
	conditions qui outrepassent le strict respect des
	procédures prescrites autrement.
Healthcare that is effective, safe and responds to the	 Les soins de santé qui sont efficaces, sûrs et
needs and preference of patients. Other dimensions	répondent aux besoins et aux préférences des
of quality of care, such as efficiency, access and equity	patients. D'autres dimensions de la qualité des
	soins, tels que l'efficacité, l'accès et l'équité sont

g considérés comme faisant partie d'un débat plus	large et sont traitées dans d'autres enceintes.		 [La qualité est] le degré par lequel les services de santé aux individus et aux populations augmentent la probabilité de résultats souhaités en santé et sont conformes aux connaissances professionnelles en cours. L'Institute of Médecir a identifié six dimensions de la qualité des soins de santé. Ceux-ci affirment que les soins de sant doivent être : sûrs, efficaces, centrés sur le patient, rapides, efficients et équitables.
are seen as being part of a wider debate and are bein	addressed in other fora. ³⁰⁴	La qualité des soins de santé signifie la prestation des meilleurs soins qui soient et l'atteinte des meilleurs résultats possibles chaque fois qu'une personne a recours au système de soins de santé ou utilise ses services. Il s'agit essentiellement de faire le meilleur travail possible en fonction des ressources disponibles. ³⁰⁵	[quality is] the degree to which health services for individuals and populations increase the likelihood of desired health outcomes and are consistent with current professional knowledge. The Institute of Medicine has identified six dimensions of healthcare quality. These state that healthcare must be: safe, effective, patient-centred, timely, efficient and equitable. (IOM)

	Quality of care is defined by the proportion of	 La qualité des soins est définie par la proportion
	potential health gain actually delivered by a	de gain potentiel en matière de santé
	healthcare organization for its set of patients. It	effectivement délivré par un organisme de soins
	reflects the gap between what can be achieved and	de santé pour l'ensemble de ses patients. Il reflète
	what actually happens. When the gap is small, quality	l'écart entre ce qui peut être accompli et ce qui se
	is achieved, if gap large, bad quality. 15 (Vincent)	passe réellement. Lorsque l'écart est faible, la
		qualité est atteinte, s'il y a un grand écart, elle est
		de mauvaise qualité.
	La qualité a un périmètre plus large que la sécurité car	
	touche à d'autres aspects que ce qui est directement	
	lié au soin comme l'accueil, l'hôtellerie Cela étant	
	chaque action dans la démarche qualité est à la fois	
	plus locale et procédurale : chaque action porte sur	
	une cause supposée d'un incident et corrige le	
	problème par la mise en place d'une nouvelle	
	procédure spécifique. Or souvent les causes sont	
	multiples (Brami)	
Sécurité du Patient (Patient Safety)	The avoidance, prevention, and amelioration of	 L'évitement, la prévention et l'amélioration des
	adverse outcomes or injuries stemming from the	résultats ou des dommages indésirables découlant
	processes of healthcare. These events include	des processus de soins de santé. Ces événements

ccidents." Safety comprennent des "erreurs", "dévi of the components of "accidents" de sécurité. La sécurit	in a person, device, or l'interaction des composants du syst	/ depends on learning réside pas dans une personne, disposi	interactions of the département. Améliorer la sécurité dé a subset of healthcare l'apprentissage de comment la sécurit	des interactions entre composants. La	patients est un sous-ensemble de la q soins de santé.	ry; ensuring patient nent of operational inimize the likelihood of hood of intercepting et ³⁰⁶) et ³⁰⁶) tet ³⁰⁶) hood of intercepting et ³⁰⁶) et ³⁰⁶)	luals and organizations • Les actions menées par des individuents from being harmed organisations pour éviter aux bénéficients et le la
"errors," "deviations," and "ac emerges from the interaction	the system; it does not reside	department. Improving safety	how safety emerges from the components. Patient safety is	quality. ⁸⁸		Freedom from accidental injuu safety involves the establishm systems and processes that m errors and maximize the likelil them when they occur. (Kohn	Actions undertaken by individ to protect healthcare recipien

Freedom from accidental injuries during the course of	 L'absence de dommages accidentels au cours de
medical care; activities to avoid, prevent, or correct	soins médicaux ; activités pour éviter, prévenir ou
adverse outcomes which may result from the delivery	corriger les effets indésirables qui peuvent
of healthcare. (CoE)	résulter de la prestation des soins de santé.
The identification, analysis and management of	 L'identification, l'analyse et la gestion des
patient-related risks and incidents, in order to make	risques et incidents liés au patient, afin de rendre
patient care safer and minimize harm to patients.	les soins aux patients plus sûrs et de minimiser les
(CoE)	dommages pour les patients.
The reduction and mitigation of unsafe acts within the	 La réduction et la réduction des actes dangereux
health-care system, as well as through the use of best	au sein du système de soins de santé, ainsi que
practices shown to lead to optimal patient outcomes.	l'utilisation de pratiques exemplaires qui donnent
(Davies)	des résultats optimaux aux patients.
The prevention and mitigation of harm to patients.	 La prévention et la réduction des risques pour
(NQF)	les patients.
La sécurité du patient est définie comme l'absence,	
pour un patient, d'atteinte inutile ou potentielle	
associée aux soins de santé. (WHO)	

 La prévention des dommages causés par les erreurs et les omissions. 	 La sécurité des patients se réfère à l'absence de dommages accidentels ou évitables générés par les soins médicaux. Ainsi, des pratiques ou des interventions qui améliorent la sécurité des patients sont celles qui réduisent la survenue d'événements indésirables évitables. 	 Le processus par lequel une organisation rend les soins aux patients plus sûrs. Cela devrait inclure une évaluation des risques, l'identification et la gestion des risques liés aux patients, l'analyse et le reporting des incidents, et la capacité d'apprendre de et de suivre les incidents et de mettre en œuvre des solutions pour minimiser le risque de leur reproduction. Le terme « sécurité du patient » remplace les termes « risque clinique », « risque de non-clinique » et la « santé et la sécurité des patients ».
The prevention of harm caused by errors of commission and omission. (Aspden, IOM)	Patient safety refers to freedom from accidental or preventable injuries produced by medical care. Thus, practices or interventions that improve patient safety are those that reduce the occurrence of preventable adverse events. (AHRQ)	The process by which an organisation makes patient care safer. This should involve risk assessment, the identification and management of patient-related risks, the reporting and analysis of incidents, and the capacity to learn from and follow-up on incidents and implement solutions to minimise the risk of them recurring. The term 'patient safety' is replacing 'clinical risk', 'non-clinical risk' and the 'health and safety of patients'. (NHS)

La sécurité du patient se définit comme la réduction	
de tout risque de préjudice évitable subi par le patient	
(HAS)	
	 L'évitement, la prévention et l'amélioration des
Avoidance, prevention and amelioration of adverse outcomes or iniuries stemming from the process of	résultats ou dommages indésirables découlant du
healthcare. (Vincent)	processus des soins de santé.
Patient safety is the sustained, proactive process of	
identifying, avoiding and rapidly resolving errors,	• La sécurité du patient est le processus proactif
omissions, mishaps and miscommunications that	visant à identifier, éviter et résoudre rapidement
could affect a patient's healing, health or well-being	les erreurs, les omissions, les accidents et les
at any point, at any time, in any care setting. (Khoja)	malentendus qui pourraient affecter la guérison,
	la santé ou le bien-être d'un patient, à tout
	moment, en tout temps et dans tous les
	structures de soins.
A discipline in the healthcare sector that applies	 Une discipline dans le secteur des soins de santé
safety science methods toward the goal of achieving a	qui applique les méthodes scientifiques de
trustworthy system of healthcare delivery. Patient	sécurité en vue de parvenir à un système de soins
safety is also an attribute of healthcare systems; it	fiable. La sécurité du patient est aussi un attribut

	minimizes the incidence and impact of, and maximizes	des systèmes de soins de santé ; il minimise
	recovery from, adverse events. ¹⁴	l'incidence et l'impact des événements
		indésirables et en maximise la récupération.
Situation dangereuse/Risque (Hazard)	A situation or event that introduces or increases the	 Une situation ou un événement qui introduit ou
	probability of an adverse event arising from a danger	augmente la probabilité d'un événement
	or peril, or that increases the extent of an adverse	indésirable découlant d'un danger ou péril, ou qui
	event. (JCHAO)	augmente l'étendue d'un événement indésirable.
	The potential source of harm (e.g., a hazard can be an	 La source potentielle de dommages (par
	error in the system itself or a misuse of the system). 307	exemple, un risque peut être une erreur dans le
		système lui-même ou le détournement du
		système).
	Le risque est la probabilité qu'un événement causant	
	ou pouvant causer des dommages se produise.	
	Lorsque cet événement se produit, on dit alors que le	
	risque s'est réalisé. (CHUM)	
		 Toute menace à la sécurité, par exemple
	Any threat to safety, e.g. unsafe practices, conduct,	pratiques, conduite, équipement, étiquettes,
	equipment. labels. names. (WHO. 2005)	noms dangereux.

	A set of circumstances or a situation that could harm a person's interests, such as their health or welfare. ³⁰⁸	 Un ensemble de circonstances ou une situation qui pourrait nuire aux intérêts d'une personne, comme leur santé ou leur bien-être.
	Anything that can cause harm. (NQF)	 Tout ce qui peut causer des dommages.
	Le risque est la probabilité de survenue d'un incident (WHO Delphi)	
	Both active and latent failures exist that create a hazard increasing the risk of harm (Forum of End Stage Renal Disease Networks)	 Il existe à la fois des défaillances actives et latentes qui créent un danger d'augmentation du risque de préjudice.
	The chance of something happening that will have an impact on individuals and/or organisations. It is measured in terms of likelihood and consequences. (NHS)	 La chance de quelque chose qui se passe d'avoir un impact sur les individus et / ou organisations. Elle est mesurée en termes de probabilité et de conséquences.
Standard de Soins (Standard of Care)	Les principes et les pratiques reconnues par les professionnels de la santé pour un état de santé	

donné dans des circonstances données. Les standards	
de soins sont des consensus d'experts basés sur des	
recherches spécifiques. (CHUM)	
	 Un ensemble de caractéristiques ou de
A set of characteristics or quantities that describes	quantités qui décrit les caractéristiques d'un
features of a product, process, service, interface, or	produit, processus, service, interface, ou matériel.
material. The description can take many forms, such	La description peut prendre de nombreuses
as the definition of terms, specification of design and	formes. telles aue la définition des termes. la
construction, detailing of procedures, or performance	spécification de la conception et de la
criteria against which a product, process, and other	construction le détail des procédures ou des
factors can be measured (National Research Council,	contraction, to define and proceeding of activity
1995). (Aspden)	כוונבובא מב הבווסוווומווכב המו ובאלמבוא מוו הוסממוני
	un processus, ainsi que d'autres facteurs peuvent
	être mesurés. (National Research Council, 1995).
A level of competence in performing medical tasks	 Un niveau de compétence dans l'exécution des
that is accepted as reasonable and reflective of a	tâches médicales qui est accepté comme
skilled and diligent healthcare provider, which obliges	raisonnable et reflétant un prestataire de soins de
a physician to confine his practice of medicine only to	santé qualifié et diligent, qui oblige un médecin de
those areas of his expertise; such standards may be	limiter sa pratique de la médecine uniquement
delineated by a hospital's medical staff bylaws or the	aux domaines de sa compétence ; ces standards
standards published by a specialty college. (Segen)	peuvent être délimitées par les statuts du

disregarded geographical considerations, holding that	cas impliquant une spécialisation, cependant,
in the practice of a board-certified medical or surgical	certains tribunaux ne tiennent pas compte des
specialty, the standard should be that of a reasonable	considérations géographiques, estimant que dans
specialist practicing medicine or surgery in the same	la pratique d'une spécialité médicale ou
special field. If a physician's conduct falls below the	chirurgicale certifiée par un Conseil, la norme doit
standard of care, he or she may be liable for any	être celle d'un spécialiste raisonnable pratiquant
injuries or damages resulting from that conduct.	la médecine ou la chirurgie dans le même
(Aspden)	domaine de spécialité. Si la conduite d'un médecin
	tombe en dessous des standards de soins, il ou
	elle peut être tenu responsable des blessures ou
	dommages résultant de ce comportement.
The principles and practices which have been accepted by a health-care profession as expected to be applied for a patient under ordinary circumstances. (Davies)	 Les principes et les pratiques qui ont été acceptées par une profession de soins de santé comme étant à appliquer pour un patient dans des circonstances ordinaires.
That standard of care is as found in a policy, or clinical guideline, or in common practice — a set of steps that would be followed or an outcome that would be expected. The standard of care may vary by	 Cette norme de soins est, comme on le trouve dans une politique ou une directive clinique, ou dans la pratique courante, un ensemble d'étapes qui seraient suivies ou un résultat qui serait

0	community (eg, due to resource constraints). When	attendu. La norme de soins peut varier selon la
t	he term is used in the clinical setting, the standard of	communauté (par exemple, en raison de
0	are is generally felt not to vary by specialty or level of	contraintes en termes de ressources). Lorsque le
t	raining. In other words, the standard of care for a	terme est utilisé dans le cadre clinique, la norme
0	ondition may well be defined in terms of the	de soins est généralement estimée ne devant pas
<u> </u>	tandard expected of a specialist, in which case a	varier selon la spécialité et le niveau de formation.
	eneralist (or trainee) would be expected to deliver	En d'autres termes, la norme de soins pour un
t	he same care or make a timely referral to the	état de santé peut ainsi être défini en termes de la
	ppropriate specialist (or supervisor, in the case of a	norme attendue de la part d'un spécialiste, auquel
t	rainee). Standard of care is also a term of art in	cas il est attendu d'un généraliste (ou d'un
	nalpractice law, and its definition varies from	stagiaire) de fournir le même soin ou de référer le
Ĺ	urisdiction to jurisdiction. When used in this legal	patient en temps opportun au spécialiste
<u>v</u>	ense, often the standard of care is specific to a given	appropriée (ou superviseur, dans le cas d'un
<u>v</u>	pecialty; it is often defined as the care expected of a	stagiaire). La norme de soins est aussi un terme en
_	easonable practitioner with similar training practicing	droit de la responsabilité, et sa définition varie
	n the same location under the same circumstances.	d'une juridiction à l'autre. Lorsqu'il est utilisé dans
)	Khoja)	ce sens juridique, souvent la norme de soins est
		spécifique à une spécialité donnée ; elle est
		souvent définie comme le soin attendu de la part
		d'un praticien raisonnable, avec une formation
		pratiquent au même endroit et dans les mêmes
		circonstances.

9. Bibliography

2 Jha A, Larizgoitia I, Audera-Lopez C, et al. The global burden of unsafe medical care: analytic modelling of observational studies. *BMJ Qual Saf* 2013;22:809–815.

3 McIntyre N, Popper K, The critical attitude in medicine: the need for a new ethics. *BMJ* 1983;287:1919-1923.

4 Woodward S, Review: Patient Safety: A Core Value of Nursing - so Why Is Achieving It so Difficult? *J Res Nurs* 2011;16:224–225.

5 Kohn LK, Corrigan J and Donaldson M(eds). Institute of Medicine, Committee on Quality of healthcare in America, To Err is Human: Building a Safer Health System. Washington, D.C.: National Academies Press, 2000.

6 Galbraith R, Holtman M, and Clyman S. Use of Assessment to Reinforce Patient Safety as a Habit. *Qual Saf Health Care* 2006;15(Suppl I):i30–i33.

7 Committee of Experts (CoE) on Management of Safety and Quality in Healthcare of the Council of Europe in "Glossary of terms related to patient and medication safety", 2005. Available on http://www.bvs.org.ar/pdf/seguridadpaciente.pdf Accessed on July 16th, 2015

8 Nendaz M, et A Perrier. « Diagnostic Errors and Flaws in Clinical Reasoning: Mechanisms and Prevention in Practice ». *Swiss Med Wkly* 2012;142:w13706.

9 Quintard B, Roberts T, Nitaro L, et al. Acceptability of health care–related risks: a literature review.

J Patient Saf 2016;12:1–10.

10 Galam, E. Dédramatiser et Travailler Nos Erreurs. Rev Prat Med Gen 2005;19:1-10.

11 Makary M, Michael D. Medical error—the third leading cause of death in the US. *BMJ* 2016;3:i2139.

12 Vincent C, Amalberti R. Safer Healthcare. Cham: Springer International Publishing, 2016. Available on http://link.springer.com/10.1007/978-3-319-25559-0. Accessed on June 10th 2016.

13 Shreve j, Van Den Bos J, Gray T, et al. The Economic Measurement of Medical Errors. Milliman Inc, 2010. Available on:

http://stage.build.gedigitalmedia.com/sites/default/files/data_set/Edward%20Lee/research-econmeasurement.pdf. Accessed on July 10, 2016

¹ Committee on Quality of Health Care in America, Institute of Medicine, Crossing the Quality Chasm: A New Health System for the 21st century, National Academy of Sciences, 2001. Available at: http://www.nationalacademies.org/hmd/Reports/2001/Crossing-the-Quality-Chasm-A-New-Health-System-for-the-21st-Century.aspx Accessed December 19, 2018.

14 Emanuel L, Berwick D, Conway j, et al. What Exactly Is Patient Safety. In: Henriksen K, Battles JB, Keyes MA, et al. Eds. Advances in patient safety: new directions and alternative approaches. Rockville, MD: Agency for Healthcare Research and Quality 2008;1:1–18.

15 Vincent, C. Patient Safety. Wiley-Blackwell, BMJ Books, (1ère edition: 2006), Oxford, 2010. pp. 416

16 AHRQ "2015 National Healthcare Quality and Disparities Report", 2015 Available on http://www.ahrq.gov/sites/default/files/wysiwyg/research/findings/nhqrdr/nhqdr15/2015nhqdr.pdf Accessed on July, 28, 2016

17 Rob Morris R, O'Riordan S. Prevention of Falls in Hospital, *Clinical Medicine* 2017;17:360–62.

18 Spencer R, Campbell S. Tools for primary care patient safety: a narrative review. *BMC Fam Pract* 2014;15:166.

19 Calain P. In search of the "new informal legitimacy" of Medecins Sans Frontieres. Public Health Ethics

2012;5:56-66.

20 Sphere Project. The Sphere Handbook: Humanitarian Charter and Minimum Standards in Humanitarian Response. Geneva: Sphere Project, ed. 2018. Available on: https://spherestandards.org/handbook-2018/ Accessed on September 14, 2019.

21 Slim H, Humanitarian Ethics. A guide to the morality of aid in war and disaster. London: Hurst & Co edition, 2015. Pages 111-121

22 Shanks L, Bil K, Fernhout J. Learning without borders: a review of the implementation of medical error reporting in Médecins Sans Frontières. *PLoS One* 2015;10:e0137158.

23 Helmreich R. On Error Management: Lessons from Aviation. BMJ 2000;320:781–5.

24 N. Kapur et al., 'Aviation and Healthcare: A Comparative Review with Implications for Patient Safety', JRSM Open 7, no. 1 (4 January 2016), https://doi.org/10.1177/2054270415616548.

25 Vrbnjak D, Denieffe S, O'Gorman C, et al. Barriers to Reporting Medication Errors and near Misses among Nurses: A Systematic Review. Int J Nurs Stud 2016;63:162–78.

26 M. Halligan and A. Zecevic, 'Safety Culture in Healthcare: A Review of Concepts, Dimensions, Measures and Progress', BMJ Quality & Safety 20, no. 4 (1 April 2011): 338–43, https://doi.org/10.1136/bmjqs.2010.040964.

27 Nathan Cortez, 'A medical malpractice model for developing countries?' 4 (n.d.): 25.

28 R. M. Wilson et al., 'Patient Safety in Developing Countries: Retrospective Estimation of Scale and Nature of Harm to Patients in Hospital', BMJ 344, no. mar13 3 (13 March 2012): e832–e832, https://doi.org/10.1136/bmj.e832.

29 Biquet Jean-Marc, 'Haiti: Between Emergency and Reconstruction: An inadequate response', Revue internationale de politique de développement, no. 4.3 (30 January 2014), https://doi.org/10.4000/poldev.1600. 30 Clarke Paul and Ben Ramalingam, 'Organisational Change in the Humanitarian Sector', n.d., 63.

31 OECD, *Medical Malpractice: Prevention, Insurance and Coverage Options*, Policy Issues in Insurance (OECD, 2006), https://doi.org/10.1787/9789264029057-en.OECD, Medical Malpractice: Prevention, Insurance and Coverage Options, Policy Issues in Insurance (OECD, 2006), https://doi.org/10.1787/9789264029057-en.

32 Elgafi S. Medical Liability in Humanitarian Missions The Journal of Humanitarian Assistance November 11, 2014 Available on https://sites.tufts.edu/jha/archives/2111 Accessed on October 17, 2019.

33 Ranjan Rajesh, et al., 'Epidemiology of Violence against Medical Practitioners in a Developing Country (2006-2017)', Journal of Health Research and Reviews 5, no. 3 (2018): 153, https://doi.org/10.4103/jhrr.jhrr_84_17.

34 WHO Patient, World Health Organisation, ICPS: Report on the Results of the Web-Based Modified Delphi Survey of the International Classification for Patient Safety Overview 8 juin 2007. Available on: http://www.who.int/patientsafety/taxonomy/ps_modified_delphi_survey.pdf Consulted on: 15/06/2015

35 Bernstein, Peter L. 'The New Religion of Risk Management'. Harvard Business Review 74, no. 2 (1996): 47.

36 Aven, Terje. 'Risk Assessment and Risk Management: Review of Recent Advances on Their Foundation'. European Journal of Operational Research 253, no. 1 (August 2016): 1–13. doi:10.1016/j.ejor.2015.12.023.

37 Esterman, M., M. D. Rosenberg, and S. K. Noonan. 'Intrinsic Fluctuations in Sustained Attention and Distractor Processing'. Journal of Neuroscience 34, no. 5 (29 January 2014): 1724–30. doi:10.1523/JNEUROSCI.2658-13.2014.

38 Weinger BM., Pantiskas C., Wiklund M., Carstensen P "Incorporating human factors into the design of medical devices" JAMA, Nov. 1998 – Vol 280, n°17 p. 1484

39 Shappell, S. & Wiegmann D."The Human Factors Analysis and Classification System–HFACS" Office of Aviation Medicine Washington, 2000. Available on https://www.nifc.gov/fireInfo/fireInfo_documents/humanfactors_classAnly.pdf Accessed on June 12, 2016

40 Wiegmann, Douglas A., and Scott A. Shappell. 'Human Error Perspectives in Aviation'. The International Journal of Aviation Psychology 11, no. 4 (2001): 341–57.

41 Reason J. "Human error" Cambridge University press, Cambridge 1990 pp 302

42 Idem p. 5-8.

43 Idem p.9

44 Idem p.53

45-Jdem p.234-248

46 Reason, J. 'Human Error: Models and Management'. BMJ 320, no. 7237 (2000): 768–70.

47 Helmreich L. On Error Management: Lessons from Aviation. BMJ 2000;320:781-5.

48 Reason J. "Human error" p. 202.

49 Reason J, Carthey J, De Leval M. Diagnosing "vulnerable system syndrome": an essential

prerequisite to effective risk management. *Qual Health Care* 2001;10:ii21–ii25.

50 Espin S. Persistence of Unsafe Practice in Everyday Work: An Exploration of Organisational and Psychological Factors Constraining Safety in the Operating Room. *Qual Saf Health Care* 2006;15:165–170.

51 Hollnagel E, Wears R, Braithwaite J. From Safety-I to Safety-II: A White Paper. 2015. Accessible on: https://www.england.nhs.uk/signuptosafety/wp-content/uploads/

sites/16/2015/10/safety-1-safety-2-whte-papr.pdf Accessed on Sept 9, 2019.

52 Reason J. Hollnagel E, Paries J. Revisiting the "Swiss Cheese" model of accident. EUROCONTROL Experimental Centre, EEC-SAS, Bretigny-sur-Orge, France, EEC Note N° 13/06, octobre 2006

53 Le Coze J, Salvi J, Gaston D. Complexity and Multi (Inter or Trans)-Disciplinary Sciences: Which Job for Engineers in Risk Management? *Journal of Risk Research* 2006;5:569–82.

54 Rasmussen J. Risk management in a dynamic society: A modeling problem. *Safety Science* 1997;27:183-213, 1997. available on http://sunnyday.mit.edu/16.863/rasmussen-safetyscience.pdf Accessed on June 20, 2016

55 Rasmussen J. The Role of Error in Organizing Behaviour. *Qual Saf Health Care* 2003;12:377–385.

56 Cook R. 'Going Solid': A Model of System Dynamics and Consequences for Patient Safety. *Qual Saf Health Care* 2005;14:130–134.

57 Le Coze J. New Models for New Times. An Anti-Dualist Move. Safety Science 2013;59:200–218.

58 Kirwan, Barry. 'Incident Reduction and Risk Migration'. Safety Science 2011;49: 11–20.

59 Agence Nationale d'Acréditation et d'Evaluation en Santé (ANAES). Principes méthodologiques pour la gestion des risques en établissement de santé. Paris, 2003 Page 12. Available on : http://www.hosmat.eu/assurqualite/gestion_risques.pdf. Accessed on July 3, 2016

60 Sutcliffe K. High Reliability Organisations (HROs). *Best Pract Res Clin Anaesthesiol* 2011;25: 133–44.

61 VA National Center for Patient Safety. Glossary of Patient Safety Terms. Available on: http://www.patientsafety.va.gov/professionals/publications/glossary.asp Accessed on: June 18, 2015.

62 Wreathall J. Assessing Risk: The Role of Probabilistic Risk Assessment (PRA) in Patient Safety Improvement. *Qual Saf Health Care* 2004;13:206–12.

633 ip L, Farmer B. High Reliability Organisations—Medication Safety. J Med Toxicol 2015;11: 257–61.

64 Chassin M, Loeb J. High-Reliability healthcare: Getting There from Here. *Milbank Quarterly* 2013;91(3): 459–90.

65 Weick K. Organisational culture as a source of high reliability. *California management review* 1987;29(2):112-127. Available on http://www.itn.liu.se/mit/education/courses/tnfl05-risk-och-olycksanalys/vecka-48/1.305709/Weick1987.pdf Accessed June 16th, 2016

66 Roberts K. A Case of the Birth and Death of a High Reliability Healthcare Organisation. *Qual Saf Health Care* 2005;14(3): 216–20.

67 Amalberti R, Vincent C, Auroy Y, et al. Violations and Migrations in healthcare: A Framework for Understanding and Management. *Qual Saf Health Care* 2006;15 (suppl1):i66–71.

68 Bergström J, van Winsen R, Henriqson E. n the Rationale of Resilience in the Domain of Safety: A Literature Review. *Reliability Engineering & System Safety* 2015;141: 131–41.

69 Kleiner B, Hettinger L, DeJoy D, et al. Sociotechnical Attributes of Safe and Unsafe Work Systems. *Ergonomics* 2015;58:635–49.

70 Idem p. 758

71 Vincent C, Amalberti R. page 34

72 Vincent, C. Research Into Medical Accidents: A Case Of Negligence? BMJ 1989;299: 1150-153.

73 Smith K, Harris K, Potters L, et al. Physician Attitudes and Practices Related to Voluntary Error and near-Miss Reporting. *J Oncol Pract* 2014;10:e350–57.

74 See http://www.apsf.org/about_safety.php Accessed on 01/07/2016

75 Reason J. Safety in the Operating theatre–Part 2: Human Error and Organisational Failure. *Qual Saf Health Care* 2005;14:56–60.

76 Great Britain, ed. An Organisation with a Memory: Report of an Expert Group on Learning from Adverse Events in the NHS. London, 2000.

77 latrogenic Injury in Australia. Australian Patient Safety Foundation, 2001. p.15. Available on http://www.apsf.net.au/dbfiles/latrogenic_Injury.pdf Accessed on 05/07/2016

78 Wakefield A, Attree M, Braidman I, et al. Patient Safety: Do Nursing and Medical Curricula Address This Theme?. *Nurse Education Today* 2005;25:333–340.

79 Andel C, Davidow S, Hollander M, et al. The Economics of healthcare Quality and Medical Errors. *J Health Care Finance* 2012;39:1-12.

80 Måseide P. Possibly Abusive, Often Benign, and Always Necessary. On Power and Domination in Medical Practice. *Sociology of Health & Illness* 1991;13:545–61.

81 Barondess J. Care of the Medical Ethos, with Some Comments on Research: Reflections after the Holocaust. *Perspectives in Biology and Medicine* 2000;43:308–24.

82 Amalberti R, Auroy Y, Berwick D, et al. Five System Barriers to Achieving Ultrasafe healthcare. *Ann Intern Med* 2005;142:756–64.

83 Carthey J. Understanding Safety in Healthcare: The System Evolution, Erosion and Enhancement Model. *Journal of Public Health Research* 2013;2:144-149..

84 Kapur N, Parand A, Soukup T, et al. Aviation and Healthcare: A Comparative Review with Implications for Patient Safety. *JRSM Open* 2016;7:1-11.

85 Wears R, Berg M. Computer Technology and Clinical Work: Still Waiting for Godot. *Jama* 2005;293:1261–63.

86 Hughes C, Travaglia J, Braithwaite J. Bad Stars or Guiding Lights? Learning from Disasters to Improve Patient Safety. *Qual Saf Health Care* 2010;19(4): 332–36.

87 Small S, Barach P. Patient Safety and Health Policy: A History and Review. *Hematol Oncol Clin North Am* 2002;16(6):1463–82.

88 Cooper J, Gaba D, Liang B et al. The National Patient Safety Foundation Agenda for Research and Development in Patient Safety. *MedGenMed*. 2000;2(3):E38. Mentioned in: Safety, WHO Patient, World Health Organisation, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: http://apps.who.int/iris/handle/10665/70882 Accessed on 18/6/2015

89 Brami J, Amalberti R. La sécurité du patient en médecine générale. Paris : Ed. Springer, 2010.

90 See NHS Website:

http://www.nhs.uk/aboutNHSChoices/professionals/healthandcareprofessionals/quality-accounts/Pages/about-quality-accounts.aspx Accessed on 05/07/2016

91 Institute of Medicine (IOM). Crossing the Quality Chasm: A New Health System for the 21st Century. Washington, D.C: National Academy Press; 2001.

92 Patient safety is not a luxury. The Lancet 2016;387:19.

93 See Haute Autorité de la Santé (HAS) http://www.hassante.fr/portail/jcms/fc_1249603/fr/accreditation-certification Accessed on 05/07/2016.

94 Ballard K. Patient Safety: A Shared Responsibility. Online J Issues Nurs 2003;8(3): 1-10.

95 Baume C. "Gestion des risques cliniques et responsabilité medico-hospitalière" Neuchâtel : Institut du droit de la santé de l'Université de Neuchatel, 2015.

96 OECD Medical Malpractice. Prevention, insurance and coverage options. *Policy issues in insurance*. 2006;11. Available on http://www.oecd.org/daf/fin/insurance/medicalmalpractice-preventioninsuranceandcoverageoptions.htm Accessed on August 10, 2014.

97 Baume C. pages 80-131.

98 Baume C. pages 13-14.

99 Scolan V, Telmon N, Rouge J, et al. Responsabilité médicale et infections nosocomiales. *Rev Med Interne* 2000;2:1-7.

100 Lindenauer P. Strengthening the Business Case for Patient Safety. AHRQ, Patient Safety Network Perspectives on Safety. 2013. Available on

https://psnet.ahrq.gov/perspectives/perspective/142/strengthening-the-business-case-for-patient-safety# Accessed August 1, 2016.

101 Guillod O. Medical Error Disclosure and Patient Safety: Legal Aspects. *Journal of Public Health Research* 2013;2:182-185.

102 Baume C. Pages 185-190

103 Mastroianni A, Mello M, Sommer S, et al. The Flaws In State 'Apology' And 'Disclosure' Laws Dilute Their Intended Impact On Malpractice Suits. *Health Affairs* 2010;29:1611–19.

104 Dekker S. The Criminalization of Human Error in Aviation and Healthcare: A Review. *Safety Science* 2011;49(2):121–27.

105 Runciman B, Merry A, McCall Smith A. Improving patients' safety by gathering information. Anonymous reporting has an important role. *BMJ* 2001;323:298.

106 Runciman B, Merry A, Tito F. Error, Blame, and the Law in healthcare—an Antipodean Perspective. *Ann Intern Med* 2003;138:974–79.

107 Gilmour J. Patient Safety, Medical Error and Tort Law: An International Comparison. Final Report 2006. Available on :

http://digitalcommons.osgoode.yorku.ca/cgi/viewcontent.cgi?article=1041&context=reports. Accessed on /3/17.

108 Liebman C, Hyman C. A Mediation Skills Model to Manage Disclosure of Errors and Adverse Events to Patients. *Health Affairs* 2004;23(4):22–32.

109 Asher E, Greenberg-Dotan S, Halevy J, et al. Defensive Medicine in Israel–a Nationwide Survey. *PloS One* 2012;7(8):e42613.

110 For instance a call from the French Ministry of Health available on http://socialsante.gouv.fr/grands-dossiers/signalement-sante-gouv-fr/ Accessed on 18/3/17.

111 Karakuş A, Duran L, Yavu Y, et al. Computer-Based Simulation Training in Emergency Medicine Designed in the Light of Malpractice Cases. *BMC Med Educ* 2014;14(1):155.

112 Wheeler D, Geis G, Mack E, et al. High-Reliability Emergency Response Teams in the Hospital: Improving Quality and Safety Using in Situ Simulation Training. *BMJ Qual Saf* 2013;22(6):507–14.

113 Gillespie B, Chaboyer W, Murray P. Enhancing Communication in Surgery Through Team Training Interventions: A Systematic Literature Review. *AORN J* 2010;92(6):642–57.

114 Wolff M, Macias C, Garcia E, et al. Patient Safety Training in Pediatric Emergency Medicine: A National Survey of Program Directors." Edited by Bema Bonsu. *Acad Emerg Med* 2014;21(7):835–38.

115 Tregunno D., Ginsburg L, Clarke B, et al. Integrating Patient Safety into Health Professionals' Curricula: A Qualitative Study of Medical, Nursing and Pharmacy Faculty Perspectives. *BMJ Qual Saf* 2014;23(3):257–64.

116 Okuyama A, Wagner C, Bijnen B. Speaking up for patient safety by hospital-based healthcare professionals: a literature review. *BMC Health Serv Res* 2014;14(1):1.

117 Moumtzoglou A. Factors That Prevent Physicians Reporting Adverse Events. *Int J Health Care Qual Assur* 2010;23(1):51-58.

118 Lawton R, McEachan R, Giles S, et al. Development of an evidence-based framework of factors contributing to patient safety incidents in hospital settings: a systematic review. *BMJ Qual Saf* 2012;21:369–380.

119 Baines R, de Bruijne M, Langelaan M, et al. What are the safety risks for patients undergoing treatment by multiple specialties: a retrospective patient record review study. *BMC Health Serv Res* 2013;13(1):1.

120 Royal College of Physicians (RCP). Work and Wellbeing in the NHS: Why Workforce Health Matters to Patient Care", 2015. Available on http://www.rcpsych.ac.uk/pdf/RCP-%20WorkWellbeingNHS.pdf accessed on July 31, 2016.

121 Renedo A, Marston C. Developing Patient-Centred Care: An Ethnographic Study of Patient Perceptions and Influence on Quality Improvement. *BMC Health Serv Res* 2015;15(1):1-11.

122 Dixon-Woods M, McNicol S, Martin M. Ten challenges in improving quality in healthcare: lessons from the Health Foundation's programme evaluations and relevant literature. *BMJ Qual Saf* 2012; 21:876–884. p 181.

123 Taken back by WHO Patient Safety pages. Available on

http://www.who.int/patientsafety/research/methods_measures/human_factors/intro_tools/en/ Accessed on July 20, 2016, this definition is developed by the Great Britain, Health and Safety Executive, and Human Engineering Limited. A Review of Safety Culture and Safety Climate Literature for the Development of the Safety Culture Inspection Toolkit. Sudbury: HSE Books, 2005.

124 Leroy H, Dierynck B, Anseel F, et al. Behavioral Integrity for Safety, Priority of Safety, Psychological Safety, and Patient Safety: A Team-Level Study. *Journal of Applied Psychology* 2012;97(6):1273.

125 Pettker C., Thung S, Raab C, et al. A Comprehensive Obstetrics Patient Safety Program Improves Safety Climate and Culture. *Am J Obstet Gynecol* 2011;204(3):216.e1–216.e6..

126 Ullstrom S, Sachs M, Hansson J et al. Suffering in Silence: A Qualitative Study of Second Victims of Adverse Events. *BMJ Qual Saf* 2014;23(4):325–31.

127 Castel E., Ginsburg L, Zaheer S, et al. Understanding Nurses' and Physicians' Fear of Repercussions for Reporting Errors: Clinician Characteristics, Organisation Demographics, or Leadership Factors? *BMC Health Serv Res* 2015;15:326.

128 Bowman C, Neeman N, Sehgal L. Enculturation of Unsafe Attitudes and Behaviors: Student Perceptions of Safety Culture. *Acad Med* 2013;88(6):802–810.

129 Fabri P, Zayas-Castro J. Human Error, Not Communication and Systems, Underlies Surgical Complications. *Surgery* 2008;144(4):557–65.

130 Dekker S, Leveson N.. 'The Systems Approach to Medicine: Controversy and Misconceptions'. *BMJ Qual Saf* 2015;24(1):7–9.

131 Craig C, Eby D, Whittington J. Care Coordination Model: Better Care at Lower Cost for People with Multiple Health and Social Needs. IHI Innovation Series white paper. Cambridge, Massachusetts: Institute for Healthcare Improvement; 2011. Available on www.IHI.org Accessed on July 31, 2016

132 National Advisory Group on the Safety of Patients in England. A Promise to Learn—A Commitment to Act: Improving the Safety of Patients in England", 2013. Available on: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/226703/Berwick_R eport.pdf. Accessed July 31, 2016

133 Marx D. Patient Safety and the "Just Culture": A Primer for healthcare Executives. New York, NY: Columbia University; 2001. Available at:

http://www.safer.healthcare.ucla.edu/safer/archive/ahrq/FinalPrimerDoc.pdf Accessed on 27/6/2015

134 Schwappach D, Gehring K. Silence That Can Be Dangerous: A Vignette Study to Assess Healthcare Professionals' Likelihood of Speaking up about Safety Concerns. *PLoS ONE* 2014;9(8):e104720.

135 Michel P. Strengths and weaknesses of available methods for assessing the nature and scale of harm caused by the health system: literature review. Report of 2004 commissioned by WHO. Available on http://www.who.int/patientsafety/research/P_Michel_Report_Final_version.pdf?ua=1. Accessed on 15/3/2017.

136 Fraser J. The Morbidity and Mortality Meeting: Time for a Different Approach?. *Arch Dis Child* 2016;101:4–8.

137 Classen D, Resar R, Griffin F, et al. "Global Trigger Tool" Shows That Adverse Events In Hospitals May Be Ten Times Greater Than Previously Measured. *Health Affairs* 2011;30(4):581–89.

138 Henriksen K, Kaplan K. Hindsight Bias, Outcome Knowledge and Adaptive Learning. *Qual Saf Health Care* 2003;12(Suppl II):ii46–ii50.

139 Christiaans-Dingelhoff I, Smits M, Zwaan L, et al. To What Extent Are Adverse Events Found in Patient Records Reported by Patients and Healthcare Professionals via Complaints, Claims and Incident Reports? *BMC Health Serv Res* 2011;11:49.

140 Van Beuzekom M, Boer F, Akerboom S, et al. Patient safety in the operating room: an intervention study on latent risk factors. *BMC Surg* 2012;12:10.

141 Lipczak H, Knudsen J, Nissen A. Safety Hazards in Cancer Care: Findings Using Three Different Methods. *BMJ Qual Saf* 2011;20(12):1052–56.
142 Stein J, Heiss K. The Swiss Cheese Model of Adverse Event occurrence—Closing the Holes. *Semin Pediatr Surg* 2015;24(6):278–82.

143 Waterson P. A Critical Review of the Systems Approach within Patient Safety Research. *Ergonomics* 2009;52(10):1185–95.

144 Taylor-Adams S, Vincent C. Systems Analysis of Clinical Incidents: The London Protocol. *Clinical Risk* 2004;10(6):211–20.

145 http://www.has-sante.fr/portail/jcms/c_1215806/fr/grille-alarm-un-outil-pour-structurer-l-analyse-des-causes

146 Hettinger A, Rollin F, Hegde S, et al. An Evidence-Based Toolkit for the Development of Effective and Sustainable Root Cause Analysis System Safety Solutions. *J Healthc Risk Manag* 2013;33(2):11–20.

147 EU. Les systèmes de signalement capables de tirer des enseignements des défaillances concernant les incidents relatifs à la sécurité des patients en Europe. Rapport du sous-groupe chargé du signalement du groupe de travail sur la sécurité des patients et la qualité des soins (GTSPQS) de la Commission européenne, 2014. Available on : https://eur-lex.europa.eu/legal-content/FR/TXT/PDF/?uri=CELEX:52014DC0371&from=FR. Accessed on 21/06/2015.

148 Kaplan H, Fastman B. Organisation of Event Reporting Data for Sense Making and System Improvement. *Qual Saf Health Care* 2003;12(Suppl II):ii68–ii72.

149 Safety, WHO Patient, World Health Organisation, and others. 2010. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009'. Available on: http://apps.who.int/iris/handle/10665/70882. Accessed on: 21/06/2015.

150 Pham J, EColantuoni E, Dominici F, et al. The Harm Susceptibility Model: A Method to Prioritise Risks Identified in Patient Safety Reporting Systems. *BMJ Qual Saf* 2010;19(5):440–45.

151 See guideline promoted by NHS "Healthcare risk assessment made easy" available on http://www.nrls.npsa.nhs.uk/resources/?entryid45=59825&q=0%C2%ACrisk%C2%AC&p=3 Accessed on July 12, 2016.

152 Frankel A. Improving Patient Safety across a Large Integrated healthcare Delivery System'. *Int J Qual Health Care* 2003; 15(Suppl. 1):i31–i40.

153 Taylor M, McNicholas C, NicolayC, et al. Systematic Review of the Application of the Plan–do– study–act Method to Improve Quality in Healthcare. *BMJ Qual Saf* 2014;23(4):290–98.

154 Dovey S, Phillips R. What Should We Report to Medical Error Reporting Systems? *Qual Saf Health Care* 2004;13:322–323.

155 Schubert C, Winslow G, Montgomery S. Defining Failure: The Language, Meaning and Ethics of Medical Error. Int J Humanit Soc Sc 2012;2(22):30-42.

156 Reason J. Understanding adverse events: The human factor. In: Vincent CA, ed. Clinical Risk Management: Enhancing Patient Safety. London: BMJ Publications, 2001, pp. 9-30. Mentioned by: 179

Committee of Experts on Management of Safety and Quality in healthcare (SP-SQS) http://www.bvs.org.ar/pdf/seguridadpaciente.pdf . Accessed on: 16/06/2015

157 Tamuz M. Defining and Classifying Medical Error: Lessons for Patient Safety Reporting Systems. *Qual Saf healthcare* 2004;13:13–20.

158 Thomas E, Brennan T. Errors and adverse events in medicine: An overview. In: Vincent C, ed. Clinical Risk Management: Enhancing Patient Safety. London: BMJ Publishing, 2001, pp. 31–43

159 Schmitt E. Qu'entend-on par erreur médicale ? *Revue Pratiques* 2012 ;59. Available on : https://pratiques.fr/Les-soignants-et-l-aptitude. Accessed on July 20, 2016

160 Pintor-Mármol A, Baena M, Fajardo P, et al. Terms Used in Patient Safety Related to Medication: A Literature Review: terminology of medication patient safety. *Pharmacoepidemiol Drug Saf* 2012;21(8):799-809.

161 Morimoto, T. Adverse Drug Events and Medication Errors: Detection and Classification Methods. *Qual Saf healthcare* 2004;13(4):306–14.

162 WHO 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. http://apps.who.int/iris/handle/10665/70882 accessed on July 21, 2016

163 Jones, A. The Role of Employee Whistleblowing and Raising Concerns in an Organisational Learning Culture – Elusive and Laudable? Comment on "Cultures of Silence and Cultures of Voice: The Role of Whistleblowing in Healthcare Organisations". *Int J Health Policy Manag* 2015;5:67–69.

164 Seys, D, Scott S, Wu A, et al. Supporting Involved healthcare Professionals (second Victims) Following an Adverse Health Event: A Literature Review. Int J Nurs Stud 2013;50(5):678–87.

165 Mira J, Lorenzo , Carrillo I, et al. Interventions in Health Organisations to Reduce the Impact of Adverse Events in Second and Third Victims. *BMC Health Serv Res* 2015;15:341.

166 Cabilan C, Kynoch K. The Experiences of and Support for Nurses as Second Victims of Adverse Nursing Errors: A Qualitative Systematic Review Protocol. *JBI Database System Rev Implement Rep* 2015;13(9):63.

167 Pronovost P, Joseph Bienvenu O. From Shame to Guilt to Love. JAMA 2015;314(23):2507-8.

168 Birks Y, Harrison R, Bosanquet K, et al. An Exploration of the Implementation of Open Disclosure of Adverse Events in the UK: A Scoping Review and Qualitative Exploration. *Health Services and Delivery Research* 2014;2(20):1-196.

169 Pfeiffer Y, Briner M, Wehner T, et al. Motivational Antecedents of Incident Reporting: Evidence from a Survey of Nurses and Physicians'. *Swiss Med Wkly*, 2013;12.

170 Brunsveld-Reinders A,. Sesmu Arbous M, De Vos R, et al. Incident and Error Reporting Systems in Intensive Care: A Systematic Review of the Literature. *Int J Qual Health Care* 2015;1–12.

171 Vincent C. Analysis of clinical incidents: a window on the system not a search for root causes Q_{W} and Q_{W} and Q

172 Wangler M, Fujikawa R, Hestbaek L, et al. Creating European Guidelines for Chiropractic Incident Reporting and Learning Systems (CIRLS): Relevance and Structure. *Chiropr Man Therap* 2011;19:1.

173 Vincent C. Patient Safety. P. 75.

174 Tamuz M, Franchois E, Thomas R. What's Past Is Prologue: Organisational Learning from a Serious Patient Injury. *Safety Science* 2011;49:75–82.

175 Wrigstad J, Bergström J, Gustafson P. Mind the Gap between Recommendation and Implementation—principles and Lessons in the Aftermath of Incident investigations: A Semi-Quantitative and Qualitative Study of Factors Leading to the Successful Implementation of Recommendations. BMJ Open 2014;4(5):e005326.

176 Wachter J, Yorio P. A System of Safety Management Practices and Worker Engagement for Reducing and Preventing Accidents: An Empirical and Theoretical Investigation. *Accident Analysis & Prevention* 2014;68:117–30.

177 Pronovost P, Mathews S, Chute C, et al. Creating a Purpose-Driven Learning and Improving Health System: The Johns Hopkins Medicine Quality and Safety Experience. *Learn Health Sys* 2017;1:e10018.

178 Verbeek-Van Noord I, Wagner C, Van Dyck, et al. Is Culture Associated with Patient Safety in the Emergency Department? A Study of Staff Perspective'. International Journal for Quality in Health Care 2014;26(1):64–70.

179 Travaglia J, Nugus P, Greenfield D, et al. Visualising Differences in Professionals' Perspectives on Quality and Safety. *BMJ Qual Saf* 2012;21(9):778–83.

180 Smits M Wagner C, Spreeuwenberg P, et al. Measuring Patient Safety Culture: An Assessment of the Clustering of Responses at Unit Level and Hospital Level. *Qual Saf healthcare* 2009;18(4):292–96.

181 Jeffs L, Abramovich L, Hayes C, et al. Implementing an Interprofessional Patient Safety Learning Initiative: Insights from Participants, Project Leads and Steering Committee Members. *BMJ Qual Saf* 2013;22:923-930.

182 Morain S, Kass N, Grossmann C. What Allows a healthcare System to Become a Learning healthcare System: Results from Interviews with Health System Leaders. *Learn Health Sys* 2017;1:e10015.

183 AHRQ. 2015 National Healthcare Quality and Disparities Report. 2015 Available on http://www.ahrq.gov/sites/default/files/wysiwyg/research/findings/nhqrdr/nhqdr15/2015nhqdr.pdf Accessed on July, 28, 2016

184 Wachter R. Patient Safety At Ten: Unmistakable Progress, Troubling Gaps. *Health Affairs* 2010;29:165–73.

185 Shah R, Brereton J, Roberson D. Quality in Otolaryngology: Where We Have Been and Where We Are Going. *Otolaryngology–Head and Neck Surgery* 2014;150(5):779–784.
186 Pronovost P, Holzmueller C, Ennen C, et al. Overview of Progress in Patient Safety. *Am J Obstet Gynecol* 2011;204:5–10.

187 Berger Z, Flickinger T, Pfoh E, et al. Promoting Engagement by Patients and Families to Reduce Adverse Events in Acute Care Settings: A Systematic Review. BMJ Qual Saf 2013;23:548–555.

188 Thurman A. Institutional Responses to Medical Mistakes: Ethical and Legal Perspectives. *Kennedy Inst Ethics J* 2001;11(2):147–56.

189 Rowe R, Calnan M. Trust relations in healthcare—the new agenda. *Eur J Public Health* 2005; 16(1):3–4. Available from: http://eurpub.oxfordjournals.org/cgi/doi/10.1093/eurpub/ckl003 Accessed on August 10th, 2016

190 Schwappach D. Frequency of and Predictors for Patient-Reported Medical and Medication Errors in Switzerland. *Swiss Med Wkly* 2011;141:w13262.

191 In France, The state proceeds to the accreditation of such association active in the defence of patient's rights (Art. R1114-1 du code de la santé publique)

192 Evin C. Les droits des malades. *Pouvoirs* 1999:15–30.

193 See WHO programme available on : http://www.who.int/patientsafety/patients_for_patient/programme/en/ Accessed on 18/3/2017

194 O'Hara J, Lawton R. At a Crossroads? Key Challenges and Future Opportunities for Patient Involvement in Patient Safety. *BMJ Qual Saf* 2016;25(8):565–68.

195 Doyle C, Lennox L, Bell D. A Systematic Review of Evidence on the Links between Patient Experience and Clinical Safety and Effectiveness. *BMJ Open* 2013;3: e001570.

196 Ward J, Armitage G. Can Patients Report Patient Safety Incidents in a Hospital Setting? A Systematic Review. *BMJ Qual Saf* 2012;21(8):685–99.

197 Khan A, Furtak S, Melvin P, et al. Parent-Reported Errors and Adverse Events in Hospitalized Children. *JAMA Pediatr* 2016;170(4):e154608.

198 Jani B, Blane D, Mercer S. The Role of Empathy in Therapy and the Physician-Patient Relationship. *Forsch Komplementmed* 2012;19:252–257.

199 Hotvedt R, Førde O. Doctors Are to Blame for Perceived Medical Adverse Events. A Cross Sectional Population Study. The Tromsø Study. *BMC Health Serv Res* 2013;13:1.

200 Weingart S, Zhu J, Chiappetta L, et al. Hospitalized Patients' Participation and Its Impact on Quality of Care and Patient Safety. *Int J Qual Health Care* 2011;23(3):269–77.

201 Davis R, Sevdalis N, Vincent C. Patient Involvement in Patient Safety: How Willing Are Patients to Participate? *BMJ Qual Saf* 2011;20:108–14.

202 Vincent C, Phillips A, Young M. Why do people sue doctors? A study of patients and relatives taking legal action *The Lancet* 1994;343(8913):1609-1613.

203 O'Connor E, Coates H, Yardley I, et al. Disclosure of Patient Safety Incidents: A Comprehensive Review. *Int J Qual Health Care* 2010;22(5):371–79.

182

204 Wells R, Kaptchuk J. To Tell the Truth, the Whole Truth, May Do Patients Harm: The Problem of the Nocebo Effect for Informed Consent. *Am J Bioeth* 2012;12(3):22–29.

205 Cassel C, Conway P, Delblanco S, et al. Getting more performance from performance measurement. *N Engl J Med*, 2014;371(23):2145-2147.

206 Thomas E, Petersen L. Measuring errors and adverse events in healthcare. *J Gen Intern Med* 2003;18:61-67.

207 Dixon-Woods M, McNicol S, Martin M. Ten challenges in improving quality in healthcare: lessons from the Health Foundation's programme evaluations and relevant literature *BMJ Qual Saf* 2012;21:876–884.

208 Jha A, Pronovost P. Toward a Safer healthcare System: The Critical Need to Improve Measurement. *JAMA* 2016;315(17):1831–32.

209 Vincent C, Burnett S, Carthey J. The Measurement and Monitoring of Safety: Drawing Together Academic Evidence and Practical Experience to Produce a Framework for Safety Measurement and Monitoring. BMJ Qual Saf 2014;23:670–677.

210 Idem

211 McGlynn E, McDonald K, Cassel C. Measurement Is Essential for Improving Diagnosis and Reducing Diagnostic Error: A Report From the Institute of Medicine. *JAMA* 2015;314(23):2501-2502.

212 De Rezende B, Or Z, Com-Ruelle L, et al. Economic evaluation in patient safety: a literature review of methods. *BMJ qual saf* 2012;21:457e465.

213 Slawomirski L, Auraaen A, Klazinga N. The economics of patient safety. Paris: OECD Publishing, 2017. http://www.oecd.org/els/health-systems/The-economics-of-patient-safety-March-2017.pdf.

214 Ranji S, Rennke S, Wachter R. Computerised Provider Order Entry Combined with Clinical Decision Support Systems to Improve Medication Safety: A Narrative Review. *BMJ Qual Saf* 2014;23(9):773-80.

215 Salahuddin L, Ismail Z. Classification of Antecedents towards Safety Use of Health Information Technology: A Systematic Review. *Int J Med Inform* 2015;84(11):877-91..

216 National Patient Safety Foundation. Free from Harm: Accelerating Patient Safety Improvement Fifteen Years after To Err Is Human. 2015: 1-159 Available on: <u>http://www.ihi.org/resources/Pages/Publications/Free-from-Harm-Accelerating-Patient-Safety-</u> <u>Improvement.aspx</u> accessed on: July 2016.

217 See OECD Health Statistics 2015, website: http://www.oecd.org/health/slow-growth-in-health-spending-but-europe-lags-behind.htm Accessed on July 2016

218 Vincent C, Batalden P, Davidoff F. Multidisciplinary Centres for Safety and Quality Improvement: Learning from Climate Change Science. *BMJ Qual Saf* 2011;20(Suppl 1):i73–78..

219 Leatherman S, Donaldson L, Eisenberg J. International Collaboration: Harnessing Differences to Meet Common Needs in Improving Quality of Care. *Quality in Health Care* 2000;9:143–145.

220 Adhikari N. Patient Safety without Borders: Measuring the Global Burden of Adverse Events. *BMJ Qual Saf* 2013;22(10):798–801.

221 Fujita S, Seto K, Ito S, et al. The Characteristics of Patient Safety Culture in Japan, Taiwan and the United States. *BMC Health Serv Res* 2013;13:1.

222 Siddiqi S, Elasady R, Khorshidi I, FortuneT, et al. Friendly Hospital Initiative: From Evidence to Action in Seven Developing Country Hospitals. *Int J Qual Health Care* 2012;24(2):144–51.

223 Raghunathan K. Checklists, Safety, My Culture and Me. BMJ Qual Saf 2012;21(7):617-20.

224 Dondorp A, Iyer S, Schultz J. Critical Care in Resource-Restricted Settings. *JAMA* 2016;315(8):753–754.

225 Rutter P, Syed S, Storr J, et al. Development of an Evaluation Framework for African–European Hospital Patient Safety Partnerships. BMJ Qual Saf 2014;23:332–337.

226 http://www.who.int/gpsc/5may/en/ Accessed on July 10, 2017

227 http://www.who.int/patientsafety/safesurgery/ss_checklist/en/ Accessed on July 10, 2017

228 See https://www.usaidassist.org/topics/patient-safety Accessed on July 10, 2017

229 Salzburg Global Seminar How to Make healthcare Better in Low and Middle Income Countries, A Call to Action. 2012. Available on:

http://www.salzburgglobal.org/fileadmin/user_upload/Salzburg_Statement-Better_care_for_all_every_time.PDF Accessed on July 20, 2016

230 Brauman R, La médecine humanitaire. Paris : Presse Universitaire de France, 2009.

231 Brauman R. Médecins sans Frontières, quarante ans après. Entretien avec le Docteur Rony Brauman, ancien Président de « Médecins sans Frontières ». *Médecine de l'Homme, Revue du Centre Catholique des Médecins Français*, 2012 ;7. Available on: https://www.msf.fr/actualites/interviewdu-dr-rony-brauman-msf-40-ans-d-aventure-humaine. Accessed on: June 28,2015.

232 Anderson M. You Save My Life Today, But for What Tomorrow? Some Moral Dilemmas of Humanitarian Aid, in Jonathan Moore (ed.), Hard Choices: Moral Dilemmas in Humanitarian Intervention. Rowman and Littlefield, 1998:137-56.

233 Ex.: World Food Program : "What Makes a Humanitarian Hero" available on: https://hdp.press/what-makes-a-humanitarian-hero-440202edded8#.9obx8s6c3

234 Herard P, Boillot F. Quality Orthopaedic Care in Sudden-Onset Disasters: Suggestions from Médecins Sans Frontières-France. *International Orthopaedics (SICOT)* 201;40:435–438.

235 Humbert N. Comment être une ONG humanitaire non toxique ? in Le Temps, Genève. 14 octobre 2013

236 Biquet, J. « Accountability Towards Beneficiaries »ou De la responsabilité des humanitaires envers leurs populations-cibles . Mémoire en vue de l'obtention du Master 2 Politiques et Stratégies de l'Action Publique Internationale. Option : OI/ONG. 2009 184 237 Eberwein W. Le Paradoxe Humanitaire? Normes et Pratiques. Cultures et Conflits 2005 ;15–37.

238 Brauman R, Tanguy J. The MSF Experience. 1998 available on http://www.doctorswithoutborders.org/msf-experience Accessed on March 23, 2015.

239 MSF « La Mancha Agreement » – 25 June 2006, Athens. Page 3. Available on : http://association.msf.org/sites/default/files/rst_library_item/La%20Mancha%20Agreement%20EN. pdf. Consulted on August 10th, 2016

240 Montandon D, Quinodoz P, Pittet D. La chirurgie plastique humanitaire en question. *Ann Chir Plast Esthet* 2004;49(3):314–19.

241 Berry N. Did We Do Good? NGOs, Conflicts of Interest and the Evaluation of Short-Term Medical Missions in Sololá, Guatemala. *Social Science & Medicine* 2014;120:344–51..

242 Hurst S, Mezger N, Mauron A. Allocating Resources in Humanitarian Medicine'. *Public Health Ethics* 2009;89–99.

243 Quadrell, S, Colt H, Garcia C. L'éthique Humanitaire et La Notion de Justice. *Rev can santé publique* 2011;102(3):210-14.

244 Schwartz L, Hunt M, Sinding M, et al. Models for Humanitarian healthcare Ethics. *Public Health Ethics* 2012;5:81–90.

245 MSF « La Mancha Agreement » page 2. Available on : http://association.msf.org/sites/default/files/rst_library_item/La%20Mancha%20Agreement%20EN. pdf. Consulted on August 10th, 2016

246 Barnett M. Humanitarian Governance. Annu. Rev. Political Sci. 2013;16:379-98.

247 Eberwein W. Certification des ONG humanitaires, à quoi bon ? Grotius International,1 novembre 2013. Available on: https://grotius.fr/certification-des-ong-humanitaires-a-quoi-bon/#.Xe-bNvzjLIU Accessed on July 20, 2016

248 Hunt M. Establishing Moral Bearings: Ethics and Expatriate healthcare Professionals in Humanitarian Work. *Disasters* 2011;35():606–22.

249 Ager A, Iacovou M. The Co-Construction of Medical Humanitarianism: Analysis of Personal, Organisationally Condoned Narratives from an Agency Website. *Social Science & Medicine* 2014;120: 430–38.

250 Bjerneld M, Lindmark G, Diskett P, et al. Perceptions of Work in Humanitarian Assistance: Interviews with Returning Swedish Health Professionals. *Disaster Management & Response* 2004;2(4)101–108.

251 Djalali A, Ingrassia P, Della Corte F, et al. Identifying Deficiencies in National and Foreign Medical Team Responses Through Expert Opinion Surveys: Implications for Education and Training. *Prehosp Disaster Med* 2014;29(4):364–68.

252 Loquercio D, Hammersley M, Emmens B. Understanding and Addressing Staff Turnover in Humanitarian Agencies. London: Overseas Development Institute, 2006.

253 Chu K, Trelles M, Ford P. Quality of Care in Humanitarian Surgery. *World J Surg* 2011;35(6):1169–72.

254 Gordon S. The Military Physician and Contested Medical Humanitarianism: A Dueling Identity? *Social Science & Medicine* 2014;120:421–29.

255 LeBrun, D, Chackungal S, Chao T, et al. Prioritizing Essential Surgery and Safe Anesthesia for the Post-2015 Development Agenda: Operative Capacities of 78 District Hospitals in 7 Low- and Middle-Income Countries. *Surgery* 2014;155(3):365–73.

256 Segen J. Current Med Talk: A Dictionary of Medical Terms, Slang & Jargon. Stanford, CT: Appleton and Lange, 1995. Mentioned in Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on:http://apps.who.int/iris/handle/10665/70882 Accessed on June 18, 2015.

257 Senders J. Medical devices, medical errors, and medical accidents. In: Bogner MS, ed. Human Error in Medicine. Hillsdale, NJ: Lawrence Erlbaum Associates, 1994. Mentioned in Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: http://apps.who.int/iris/handle/10665/70882 Accessed on June 18, 2015.

258 ALNAP Annual Review 2003 Humanitarian Action Available in: http://www.livestockemergency.net/userfiles/file/common-standards/ALNAP-2003.pdf Accessed on June 25, 2015.

259 OECD. Towards Better Humanitarian Donorship. 12 Lessons from DAC Peer Reviews. 2012:1-56. Available on: http://www.oecd.org/dac/peer-reviews/12lessons.pdf. Accessed on June 25, 2015.

260 GHD. Good Humanitarian Donorship. Principles and good practice of humanitarian donorship. 2003. Available on: <u>http://www.mofa.go.jp/mofaj/gaiko/jindo/pdfs/pgphd.pdf</u> Accessed on June 26, 2015.

261 UNOCHA. Guidelines on the Use of Military and Civil Defence Assets to Support United Nations Humanitarian Activities in Complex Emergencies. 2003. Available on :

http://www.unicef.org/pathtraining/Documents/Session%204%20Humanitarian%20Principles/Participant%20Manual/4.4%20Military%20and%20Civil%20Defense%20Assets%20Guidelines.doc. Accessed on June 26, 2015.

262 Center For Patient Safety (CPS). Patient safety organization terminology and acronyms Available on: <u>http://www.centerforpatientsafety.org/wp-content/uploads/2011/07/CPS-Glossary.pdf</u> Accessed on June 25,2015.

263 ACSNI Study Group on Human Factors. Organising for Safety (Third Report to Health and Safety Commission. ACSNI Study Group on Human Factors, Advisory Committee on the Safety of Nuclear Installations). London: Health and Safety Commission, 1993. Mentioned in Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification 186 for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: <u>http://apps.who.int/iris/handle/10665/70882</u> Accessed on June 18, 2015.

264 Pizzi L, Goldfarb N, Nash D. Procedures for obtaining informed consent. In: Making healthcare Safer: A Critical Analysis of Patient Safety Practices. Evidence Report/Technology Assessment, Number 43. Rockville, MD: Agency for Healthcare Research and Quality, 2001. Mentioned in Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on:http://apps.who.int/iris/handle/10665/70882 Accessed on June 18, 2015.

265 Schein E. Organisational Culture and Leadership . 2nd edn. San Francisco, CA: Jossey-Bass, 1992 Cité dans Vlayen, Annemie, Johan Hellings, Neree Claes, Hilde Peleman, and Ward Schrooten. 'A Nationwide Hospital Survey on Patient Safety Culture in Belgian Hospitals: Setting Priorities at the Launch of a 5-Year Patient Safety Plan'. *BMJ Qual Saf* 2012;21(9):760–67.

266 Khoja T, Kamel A, Rawaf S. Council of Health Ministers of G.C.C. States, and Executive Board. Glossary of patient safety. Riyadh: Executive Board of the Health Ministers' Council for Cooperation Council States, 2010.

267 Thomas E, Brennan T. Errors and adverse events in medicine: An overview. In: Vincent C, ed. Clinical Risk Management: Enhancing Patient Safety. London: BMJ Publishing, 2001, pp. 31–43

268 AHRQ Accessible on : http://psnet.ahrq.gov/glossary.aspx?indexLetter=E Accessed on 16/06/2015

269 CHUM. Glossaire canadien sur la prestation sécuritaire des soins et services au patient. Centre hospitalier de l'Université de Montréal, Direction de la gestion de l'information et de la qualité-performance, 2004. http://site.ebrary.com/id/10222210.

270 WHO Patient, World Health Organization, CISP: Rapport sur les résultats de l'enquête Delphi modifiée par Internet sur la Classification internationale pour la sécurité des patients Introduction 8 juin 2007. Available on :

http://www.who.int/patientsafety/taxonomy/ps_modified_delphi_survey_fr.pdf Accessed on : June 15, 2015.

271 Aspden, Philip, Institute of Medicine (U.S.), and Committee on Data Standards for Patient Safety. Patient Safety Achieving a New Standard for Care. Washington, D.C.: National Academies Press, 2004. http://site.ebrary.com/id/10057003.

272 National Steering Committee on Patient Safety (Canada). Building a Safer System: A National Integrated Strategy for Improving Patient Safety in Canadian healthcare. [Ottawa]: National Steering Committee on Patient Safety, 2002.

273 Wu A, Cavanaugh T, McPhee S, et al. To tell the truth: Ethical and practical issues in disclosing medical mistakes to patients. *J Gen Intern Med* 1997;12:770–775. Mentioned by Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: http://apps.who.int/iris/handle/10665/70882 Accessed on June 16, 2015. 187

274 WHO "WHO Draft Guidelines for Adverse Event Reporting and Learning Systems" 2005 http://www.who.int/patientsafety/events/05/Reporting_Guidelines.pdf Accessed on June, 20,2015.

275 Forum of End Stage Renal Disease Networks, National Patient Safety Foundation, Renal Physicians Association, Renal Physicians Association. National ESRD Patient Safety Initiative: Phase II Report. Chicago: National Patient Safety Foundation, 2001 Mentioned by Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: htp://apps.who.int/iris/handle/10665/70882 Accessed on June 16, 2015.

276 Couch N, Tilney N, Rayner A, et al. The high cost of low-frequency events: The anatomy and economics of surgical mishaps. *N Engl J Med* 1981;304:634-637.

277 Cousins D. Developing a uniform reporting system for preventable adverse drug events. Clin Therap 1998;20(suppl C):C45–C59. Mentioned by Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: http://apps.who.int/iris/handle/10665/70882 Accessed on 18June 18, 2015.

278 Cohen M, Kilo C. High-alert medications: Safeguarding against errors. In: Cohen MR, ed. Medication Errors. Washington, DC: American Pharmaceutical Association, 1999. Mentioned by Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: http://apps.who.int/iris/handle/10665/70882. Accessed on June 18, 2015.

279 Kaushal R, Bates D. Computerized physician order entry (CPOE) with clinical decision support systems (CDSSs). In: Making healthcare Safer: A Critical Analysis of Patient Safety Practices. Evidence Report/Technology Assessment, Number 43. Rockville, MD: Agency for Healthcare Research and Quality, 2001. Mentioned by Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: http://apps.who.int/iris/handle/10665/70882 Accessed on June 18. 2015.

280 Slee V, Slee D, Schmidt H. healthcare Terms, 3d ed. St. Paul, MN: Tringa Press, 1996. . Mentioned by Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: http://apps.who.int/iris/handle/10665/70882. Accessed on June 18, 2015.

281 Institute for Safe Medication Practices Canada, Canadian Medication Incident Reporting and Prevention System. Definitions of Terms. www.ismp-canada.org/definitions.htm. Accessed 14 February 2006. Mentioned by Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: <u>http://apps.who.int/iris/handle/10665/70882</u> <u>Accessed on June 18</u>, 2015.

282 Joint Commission on Accreditation of Healthcare Organizations, editor. Lexicon: Dictionary of healthcare Terms, Organizations, and Acronyms. 2nd ed. Oakbrook Terrace: Joint Commission on Aggreditation of Healthcare Organizations; 1998.

283 Brennan T, Leape L, Laird N, et al. Incidence of adverse events and negligence in hospitalized patients: Results of the Harvard Medical Practice Study. *N Engl J Med* 1991;324:370–376.

284 NHS. "Building a safer NHS for patients" 2002. Available on: http://www.bvsde.paho.org/bvsacd/cd65/medicationsafety.pdf Accessed on 18/6/2015

285 IOM Patient Safety: Achieving a New Standard for Care. Available on : http://www.nap.edu/catalog/10863.html Accessed on June 18, 2015.

286 NQF. National Quality Forum. Serious Reportable Events in Healthcare: A Consensus Report. Washington, DC: National Quality Forum, 2002. Available on : http://www.nothingleftbehind.org/uploads/NQF_SRE_s_2011.pdf. Accessed on 24/06/2015

287 Quality Interagency Coordination Task Force. Doing What Counts for Patient Safety: Federal Actions to Reduce Medical Errors and Their Impact. Washington, DC: Quality Interagency Coordination Task Force, 2000. Mentioned by Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on:http://apps.who.int/iris/handle/10665/70882 Accessed on June 18, 2015.

288 Boxwala A, Dierks M, Keenan M, et al. Organization and Representation of Patient Safety Data: Current Status and Issues around Generalizability and Scalability. *J Am Med Inform Assoc* 2004;11:468-478

289 Michel P, Quenon JL, de Sarasqueta AM, et al. Comparison of three methods for estimating rates of adverse events and rates of preventable adverse events in acute care hospitals. *BMJ*, 2004;.328(7433):1-5.

290 NSQHS. National Safety and Quality Health Service Standards", 2012. Available on: http://www.safetyandquality.gov.au/wp-content/uploads/2011/09/NSQHS-Standards-Sept-2012.pdf Accessed on June 18, 2015.

291 HAS. Ce qu'il faut savoir sur l'évaluation de la démarche qualité et gestion des risques au bloc opératoire. 2015. Accessible on: http://www.has-sante.fr/portail/jcms/c_2025542/fr/ce-qu-il-faut-savoir-sur-l-evaluation-de-la-demarche-qualite-et-gestion-des-risques-au-bloc-operatoire Accessed on June 25, 2015.

292 Danish healthcare Act (no. 913 of 13/07/2010). Available on: http://www.patientsikkerhed.dk/in-english/act-on-patient-safety.aspx. Accessed on 20/06/2015

293 Rogers A, Israel E, Smith C, et al. Physician knowledge, attitudes, and behavior related to reporting adverse drug events. *Arch Intern Med* 1988;148:1596–1600. Mentioned by Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: http://apps.who.int/iris/handle/10665/70882 Accessed on June 18, 2015.

294 Wald H, Shojania K. Incident reporting. In: Making healthcare Safer: A Critical Analysis of Patient Safety Practices. Evidence Report/Technology Assessment, Number 43. Rockville, MD: Agency for Healthcare Research and Quality, 2001. Mentioned by Safety, WHO Patient, World Health 189

Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: http://apps.who.int/iris/handle/10665/70882 Accessed on June 18, 2015.

295 IAHM. International Association for Humanitarian Medicine Brock Chisholm Website. Available on: http://www.iahm.org/eng/home_medicinaumanitaria.htm Accessed on June 28, 2015.

296 Brauman R."Humanitarian Medicine. 1999. Available on : http://www.msfcrash.org/drive/1433-humanitarian-medicine.pdf Accessed on June 28, 2015.

297 Bradol J. Soigner la santé Agir à tout prix ? Négociations humanitaires : l'expérience de Médecins Sans Frontières. Sous la direction de Claire Magone, Michaël Neuman et Fabrice Weissman. 2001 Available on: http://msf-crash.org/livres/agir-a-tout-prix/soigner-la-sante. Accessed on 29June 29, 2015.

298 NZ. New Zealand Supreme Court considers a case of medical misadventure, unexpected pregnancy, personal injury and the Accident Compensation Act. Available on : <u>http://www.minterellison.co.nz/New_Zealand_Supreme_Court_considers_a_case_of_medical_misa_dventure_unexpected_pregnancy_personal_injury_and_the_Accident_Compensation_Act_August_2_012/ Accessed on June 25, 2015.</u>

299 Runciman B, Merry A, Walton M. Safety and Ethics in Healthcare: A Guide to Getting it Right Hampshire: Ashgate Publishing Limited, 2007.

300 Mulcahy L. Medication of medical negligence actions: An option for the future? In: Rosenthal MM, Mulcahy L, Lloyd-Bostock S, eds. Medical Mishaps: Pieces of the Puzzle. Buckingham, UK: Open University Press, 1999, pp. 154–167 Mentioned in Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on:

http://apps.who.int/iris/handle/10665/70882 Accessed on June 18, 2015.

301 Donabedian A. The Definition of Quality and Approaches to Its Assessment. Vol 1. Explorations in Quality Assessment and Monitoring. Health Administration Press; 1980. Mentioned by EXPH. Expert Panel on Effective Ways of Investigating in Health "Future EU Agenda on quality of healthcare with a special emphasis on patient safety" 9 October 2014.

http://ec.europa.eu/health/expert_panel/opinions/docs/006_safety_quality_of_care_en.pdf Accessed_on 26June 26, 2015.

302 EXPH. Expert Panel on Effective Ways of Investigating in Health "Future EU Agenda on quality of healthcare with a special emphasis on patient safety" 9 October 2014. Available on: http://ec.europa.eu/health/expert_panel/opinions/docs/006_safety_quality_of_care_en.pdf Accessed on June 26, 2015.

303 WHO. 2000. Mentioned by EXPH. Expert Panel on Effective Ways of Investigating in Health "Future EU Agenda on quality of healthcare with a special emphasis on patient safety" 9 October 2014. <u>http://ec.europa.eu/health/expert_panel/opinions/docs/006_safety_quality_of_care_en.pdf</u> Accessed on 26June 26, 2015.

304 European Commission 2010. Mentioned by: EXPH. Expert Panel on Effective Ways of Investigating in Health. Future EU Agenda on quality of healthcare with a special emphasis on patient safety. 9 October 2014. Available on:

http://ec.europa.eu/health/expert_panel/opinions/docs/006_safety_quality_of_care_en.pdf Accessed on June 26, 2015.

305 Website of Canadian government. Accessible on: http://www.hc-sc.gc.ca/hcs-sss/qual/index-fra.php Accessed on 18/6/2015

306 Institut canadien pour la sécurité des patients. Available on: http://www.patientsafetyinstitute.ca/English/toolsResources/PatientSafetyIncidentManagementToo lkit/Pages/Glossary.aspx Accessed on: 20/06/2015

307 Voges U. Minimisation of risk in medical systems by system design for safety. In: Vincent C, de Mol B, eds. Safety in Medicine. Amsterdam: Pergamon, 2000. Mentioned by: Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: http://apps.who.int/iris/handle/10665/70882 Accessed on June 18, 2015.

308 Davies J, Hebert P, Hoffman C, editors. The Canadian Patient Safety Dictionary. Calgary: Royal College of Physicians and Surgeons of Canada and Health Canada. 2003. Mentioned by: Safety, WHO Patient, World Health Organization, and others. 'Conceptual Framework for the International Classification for Patient Safety Version 1.1: Final Technical Report January 2009', 2010. Available on: http://apps.who.int/iris/handle/10665/70882 Accessed on June 18, 2015.