



# Investigation of the neural circuits mediating instrumental active avoidance and non instrumental contextual avoidance

Juliette Viellard

## ► To cite this version:

Juliette Viellard. Investigation of the neural circuits mediating instrumental active avoidance and non instrumental contextual avoidance. Neuroscience. Université de Bordeaux; Universidade de São Paulo (Brésil), 2019. English. NNT : 2019BORD0456 . tel-03217639

**HAL Id: tel-03217639**

**<https://theses.hal.science/tel-03217639>**

Submitted on 5 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THÈSE EN COTUTELLE PRÉSENTÉE

POUR OBTENIR LE GRADE DE

**DOCTEUR DE**

**L'UNIVERSITÉ DE BORDEAUX**

**ET DE L'UNIVERSITÉ DE SÃO PAULO**

ÉCOLE DOCTORALE SCIENCE DE LA VIE ET DE LA SANTE (SVS)

ÉCOLE DOCTORALE DE LA BIOLOGIE DES SYSTEMES

SPÉCIALITÉ NEUROSCIENCES

Par **Juliette VIELLARD**

**ETUDES DES CIRCUITS NEURONAUX ORGANISANT L'EVITEMENT ACTIF  
INSTRUMENTAL ET L'EVITEMENT CONTEXTUEL NON INSTRUMENTAL**

*INVESTIGATION OF THE NEURAL CIRCUITS MEDIATING INSTRUMENTAL ACTIVE  
AVOIDANCE AND NON INSTRUMENTAL CONTEXTUAL AVOIDANCE*

Sous la direction de Cyril HERRY  
et de Newton CANTERAS

Soutenue le 20 décembre 2019

**Membres du jury :**

M. Yann HUMEAU, Directeur de recherche	UBX (CNRS)	Président
M. Antonio Padua de Carobrez, Professeur	UFSC	Rapporteur
Mme Deborah SUCHECKI, Professeure associée	EPM/UNIFESP	Rapporteur
Mme Caroline BLANCHARD, Professeure Emérite	Université de Hawaii	Examineur
M. Cyril Herry, Directeur de recherche	UBX (INSERM)	Directeur de Thèse
M. Newton Canteras, Professeur	USP	Directeur de Thèse



## ACKNOWLEDGMENT

These last four years I did my Phd in collaboration with the University of Bordeaux, in the lab of Doctor Cyril Herry at the Magendie institute, and with the University of São Paulo, in the lab of Professor Newton Canteras. As my thesis is successfully coming to an end, I would like to thank, along with my two thesis directors, and the committee members of my thesis, and all of the people who have helped, and support me during these four years.

First of all Thank you Newton and Cyril, for giving me the opportunity to work in both of your labs, where I had the chance to learn high quality science and evolve a lot in my scientific life. It also meant a lot to me that you agreed to go along in this “cotutelle” adventure with me, that, not without a bit of work, was a successful collaboration. Newton, I have been honored to work with you. Working in your lab helped me becoming independent in my research project and transmitted me the excitement of discussing burning questions of today’s science. I thank you for your infinite support and your wise advice. Cyril, it was a great chance for me to work with you, thank you for transmitting your expertise and giving me the will to always expect the best of myself. I really thank you both for the complementarity of your mentorship, and I hope that our collaboration is just starting.

I Thank my thesis committee members for agreeing to participate to the evaluation of my work. Thank you Dr. Deborah Suchecki for agreeing to be the President of my Jury, I thank you Prof. Caroline Blanchard, Prof. Antonio de Padua Carobrez, and Dr. Yann Humeau for examining my work.

I would like to thank Stéphane Olié, Head of François Magendie Institute, and Prof. Dr. Claudimara Ferini Pacicco Lotfifor, Head of the department of Anatomy of the Institute of Biomedical Science, for hosting me and giving me the opportunity to pursue my work in great conditions.

I would like to thank the Fundação de Amparo à Pesquisa do Estado de S. Paulo (FAPESP) for funding my thesis since 2016 (n° 2016/08640-7, 2017/05082-6).

A special thank you to *Marilia Oliveira* and *Stéphanie Danaux*, for your great help in the finalisation of the cotutelle contract.

I would like to show my special gratitude to *Suzana Khoder*, who was my lab partner in Bordeaux, and with whom I shared half of my thesis. Thank you Suzana for welcoming me so openly in your project, and for being such a great mentor. I learned a lot by your side. I also think that our collaboration couldn't have been better, and that our complementarity made our strength. With your energy, your sense of organization, your search for perfection, and your great openness for discussion, I couldn't expect a better partnership for my thesis.

Thank you Yann and Ha Rang for your help and collaboration in the project. Your expertise of *in-vitro* patch-clamp recordings gave added a great complement to our study.

From my two labs, I am very grateful for the diversity of great people that I have worked with and I wish you all a lot of success in your careers and life. I would like to give a special thank you to Amanda and Delphine who welcomed me and guided me in the labs. I would like to show my gratitude to the team members that showed me their expertise and helped me during my project, *Suzana, Simone, Miguel X., Natxo, Paula, Fernando M. and Ricardo B.* Thank you other team members for your support and for our great moments together. In Bordeaux *Cyril D., Mario, Daniel, Thomas, Maiena, Ha Rang, Kibong, Nancy, Yohan.* In São Paulo, *Kelvia, Miguel R., Ricardo M., Helen, Fernando R., Gabriela, Tatiane, João, Alisson, Erika, Carol, Leonardo, Luan,*

The life of a Phd student wouldn't be possible without friends around, as I was in Bordeaux, São Paulo or other places, you guys were there to support me and cheer me up. So thank you *Camille, Mathilde, Vladimir, Frank, Julia, Diane, Josi, Geoffroy, Afonso* and others I may forget. And of course I cannot forget my flatmates who followed the adventure from the first row, whether you liked it or not... So Victor, Aurélien, Pierre Yves, Marylou, Luka, Clément, Marine, Annabelle, thank you for your support, in all of your own beautiful ways.

Finally I want to show my gratitude to my dear family. This project made me travel back and forth between France and Brazil during four years, and your support was always present and strong. The kilometers did not weaken the strength you gave me. So thank you my Parents, thank you Antoine, Marie, Annabelle, and Nastia. I also have a special thanks for my uncle *Prof. Guillaume Bécard*, thank you for mentoring me since my first attempts to start a Phd. I thank you for your advice. I also want to thank you *Victor* (and you too *Yvan* and *Eva* who arrived

later), my “Brazilian” cousins who made my stay in Brazil feel even more like home. Your support was precious.

## Résumé

Les mammifères, y compris les rongeurs, présentent un large éventail de comportements défensifs actifs, tels que l'évitement, ou passif, tel que l'immobilisation (le freezing), à des fins d'adaptation et de survie. La réponse d'évitement est une réaction apprise dans laquelle un individu prend le contrôle de situations dangereuses pour faire face aux menaces. Une forme d'évitement qui a été étudiée est l'évitement actif signalé, où les individus sont entraînés à éviter une situation et fuient en réponse à un signal précédemment associé à un stimulus aversif. Il a été souligné que le Cortex préfrontal dorso-médian (CPFdm) joue un rôle important dans l'encodage de l'acquisition et de l'expression du freezing ainsi que dans les réponses d'évitement. Cependant, sa contribution à l'acquisition et à l'expression de comportements d'évitement n'est pas claire, et les circuits neuronaux du CPFdm qui gèrent ensemble les stratégies d'adaptation actives et passives, restent à découvrir. Pour répondre à cette question, nous avons développé un nouveau paradigme comportemental dans lequel une souris a la possibilité de se figer ou d'éviter un stimulus aversif en fonction des contingences contextuelles. Premièrement, nous avons étudié le rôle de la voie entre le CPFdm et la matière grise periaqueducale (PAG) dans l'évitement actif signalé, et sa relation avec le freezing. Nos résultats indiquent que (i) le CPFdm et le dl/IPAG sont activés lors du comportement d'évitement, (ii) et que l'inhibition optogénétique de cette voie bloque l'acquisition de l'évitement conditionné. Une forme d'évitement non instrumental est également étudiée, dans laquelle l'individu apprend à éviter l'environnement aversif en utilisant uniquement des indices contextuels et affichant des comportements d'évaluation des risques à l'encontre de l'environnement dangereux. Il a été précédemment démontré que dans cette situation, un circuit septohippocampale-hypothalamique-tronc cérébral est spécialement activé. Cette analyse a aussi révélé que le noyau dorsal pré-mamillaire (PMD) devait être impliqué de manière critique dans l'évitement passif contextuel. Nous avons analysé l'influence de la modulation du PMD et de ses projections au niveau de ses cibles principales, sur le processus d'expression et de reconsolidation de l'évitement passif contextuel. Nos résultats ont montré qu'une (i) voie septohippocampale-hypothalamique-tronc cérébral spécifique était impliquée dans notre paradigme d'évitement passif. (ii) De plus, l'inhibition du PMD lors d'une exposition au contexte aversif altère à la fois l'expression des comportements d'évitement et la reconsolidation de la mémoire. (iii) Enfin l'inhibition au niveau des terminaux du PMD altère l'expression et la reconsolidation de la mémoire dans le dlPAG et dans l'AMv. Les expériences de ce projet ont été menées à bien à l'aide d'analyse Fos, d'inhibition pharmacogénétique et optogénétiques.

**Mots clés:** Evitement actif. Evitement Passif. Cortex préfrontal. Hypothalamus. Optogénétique

## Resumo

Mamíferos, incluindo roedores, mostram uma ampla gama de comportamentos defensivos como forma de lidar ativamente, como comportamentos de esquiva, ou passivamente, como comportamento de congelamento (“freezing”). A resposta de esquiva é uma resposta aprendida na qual um indivíduo assume o controle em situações perigosas para lidar com ameaças. Uma forma de esquiva investigada é a esquiva ativa sinalizada, na qual os indivíduos são treinados a se esquivar de um estímulo aversivo fazendo uma tarefa aprendida em resposta a apresentação de uma pista previamente associada ao estímulo aversivo. Foi evidenciado que o CPFdm desempenha um papel importante na codificação da aquisição e expressão de congelamento, bem como nas respostas de esquiva. No entanto, sua contribuição para a aquisição e expressão do comportamento de esquiva não é clara e os circuitos neurais do processamento do CPFdm ainda não foram descobertos. Para resolver essa questão, desenvolvemos um novo paradigma comportamental, no qual o camundongo tem a possibilidade de “freeze” passivamente ao estímulo aversivo ou evitá-lo ativamente em função de contingências contextuais. Nessa primeira parte do projeto, estudamos o papel da via entre o CPFdm e a PAG na esquiva ativa sinalizada e sua relação com o congelamento. Nossos resultados indicam que (i) o CPFdm e a dl/IPAG são ativadas durante o comportamento de esquiva, (ii) a inibição optogenética dessa via bloqueou a aquisição de esquiva condicionada mas não alterou a resposta de congelamento. Uma forma não instrumental de esquiva, foi investigada, onde o indivíduo aprende a evitar o ambiente aversivo usando apenas pistas contextuais e exibindo comportamentos de avaliação de risco em relação ao ambiente aversivo. Nesta situação foi demonstrado que uma via específica septohipocampo-hipotalâmico-tronco encefálico está envolvida. Esta análise revelou que o núcleo pré-mamilar dorsal (PMD) deve estar criticamente envolvido na expressão de esquiva passiva. Nos analisamos como a manipulação do PMD e suas projeções para seus principais alvos influencia os processos de expressão e re-consolidação da esquiva passiva contextual. Nossos resultados mostraram que (i) uma via específica septohipocampo-hipotalâmico-tronco encefálico está envolvida em nosso paradigma de esquiva passiva. (ii) O silenciamento do PMD durante a exposição ao contexto prejudica tanto a expressão de esquiva quanto a reconsolidação de memória e que (ii) a inibição no nível terminal prejudica a expressão e a reconsolidação de memória tanto em dlPAG quanto em AMv. investigamos essas questões com a análise imunoquímica de Fos, manipulações de circuitos neurais usando técnicas optogenéticas e fármacogenéticas.

**Palavras-chave:** Esquiva ativa. Esquiva passiva. Cortex préfrontal. Hipotalamo. Optogenética

## Abstract

Mammals, including rodents show a broad range of defensive behaviors as a mean of coping actively, such as avoidance behaviors, or passively such as freezing behavior. The avoidance response is a learned response in which an individual takes control in dangerous situations to deal with threats. One form of avoidance that has been investigated is the signaled active avoidance, where individuals are trained to avoid an environment, and escape in response to a cue previously associated with an aversive stimulus. It has been emphasized that the dmPFC plays an important role in encoding freezing acquisition and expression as well as active avoidance responses. However the neural circuits of the dmPFC processing the expression and acquisition of both active and passive coping strategies are yet to be discovered. To adress this question, we developed a novel behavioral paradigm in which a mouse has the possibility to either passively freeze to an aversive stimulus or to actively avoid it as a function of contextual contingencies. We first investigated the role of the pathway between the dmPFC and PAG in signaled active avoidance, and its relation with freezing. Our results indicate that (i) dmPFC and dl/IPAG sub-regions are activated during avoidance behavior, (ii) and that the optogenetic inhibition of this pathway blocked the acquisition of active avoidance. A non-instrumental form of avoidance is also investigated where the individual learns to avoid the aversive environment using contextual clues only, and displaying risk assessment behaviors toward the fearful environment. It has been previously shown that in this situation, a circuit involving the septohippocampal-hypothalamic-brainstem pathway is involved. It also revealed that the dorsal premammillary nucleus (PMD) must be critically involved in contextual passive avoidance. We analysed how the manipulation of the PMD and its projections to its main targets influences the expression and reconsolidation processes of contextual passive avoidance. Our results showed that (i) a specific septohippocampal-hypothalamic-brainstem pathway is involved in our passive avoidance paradigm. (ii) Silencing the PMD during context exposure impairs both avoidance expression and memory reconsolidation and that (iii) the inhibition at terminal level impairs the expression and memory reconsolidation in both dlPAG and AMv. Both parts of the project assessed these questions using Fos immunocytochemistry analysis, manipulations of neural circuits using optogenetic, and pharmacogenetic techniques.

**Keywords:** Active avoidance. Passive avoidance. Prefrontal cortex. Hypothalamus. Optogenetics.

## Résumé de la thèse

Tout au long du règne animal, les individus présentent un répertoire de comportements défensifs liés à leurs besoins de survie, selon leur environnement. En effet, les animaux adoptent des stratégies défensives pour se protéger et/ou protéger leurs congénères contre des menaces externes. Les mammifères, y compris les rongeurs, présentent un large éventail de comportements défensifs. Ces comportements peuvent être exprimés de manière active, tels que des comportements d'évitement, ou passive, tel que le comportement d'immobilisation communément appelé « freezing ». Lorsque l'individu adapte son comportement à des fins de survie, il sélectionne une stratégie défensive appropriée en tenant compte de ses coûts énergétiques, de la menace, et du contexte dans lequel elle se manifeste (Hofmann et Hay, 2018). C'est pourquoi lorsque le danger est évitable, des comportements défensifs actifs tels que l'évitement et la fuite sont adoptés (Ramirez et al., 2015; Blanchard et Blanchard, 1969). L'évitement est une réaction apprise dans laquelle un individu prend le contrôle de situations dangereuses en s'éloignant de la menace potentielle.

Une forme d'évitement qui est couramment modélisée en laboratoire est l'évitement actif signalé. Il s'agit d'une simulation où les individus sont entraînés à éviter une situation, en s'échappant suite à un signal précédemment associé à un stimulus aversif (Moscarello and LeDoux, 2013). Ce modèle d'évitement actif sera le sujet de la première partie de cette thèse. Le projet a été développé en collaboration avec Dr. Suzana Khoder, dans l'équipe de Dr Herry à l'INSERM, Bordeaux. L'exposé des résultats de cette première partie est le détail de ma contribution.

Il existe aussi une autre forme d'évitement exposé dans la littérature, qui est un modèle non instrumental, dans lequel l'individu apprend à éviter un contexte précédemment associé à un stimulus aversif en utilisant uniquement des indices contextuels. Dans cette situation, l'individu adopte des comportements d'évaluation des risques à l'encontre de l'environnement dangereux, et un refus d'approcher le contexte aversif. Ce sujet sera exploré dans la deuxième partie de la thèse, impliquant des expériences mises en œuvre dans l'équipe du Prof. Canteras, à l'université de São Paulo, au Brésil.

L'évitement actif signalé est un phénomène impliquant des fonctions cognitives complexes. Il a été souligné que le Cortex préfrontal dorso-médian (CPFdm) joue un rôle important dans l'encodage de l'acquisition et de l'expression de freezing ainsi que dans les

réponses d'évitement. Cependant, son implication dans l'acquisition et l'expression des comportements d'évitements n'est pas claire, et les circuits neuronaux du CPFdm qui gèrent ensemble les stratégies d'adaptation active et passive, restent à découvrir. Pour répondre à cette question, nous avons développé un nouveau paradigme comportemental dans lequel une souris a la possibilité de se figer passivement à un stimulus aversif ou de l'éviter activement, ceci dépendant des contingences contextuelles. Dans la première partie de ce projet, nous avons étudié, à l'aide d'analyse immunohistochimique de Fos, d'enregistrements électrophysiologiques et de manipulations de circuits neuronaux utilisant des techniques d'optogénétiques, le rôle de la voie entre le cortex préfrontal dorso-médian (et la matière grise périaqueducule (PAG) dans l'évitement actif signalé, et sa relation avec le freezing.

Dans un deuxième temps l'intérêt s'est porté sur l'évitement passif non instrumental. L'utilisation d'un nouveau paradigme donne la possibilité à l'animal d'entrer ou non dans un environnement familier associé à un événement aversif (un choc). D'après une étude similaire effectuée dans notre laboratoire, chez le rat (Viellard et al., 2016), et une étude approfondie de la littérature de peur innée contextuelle, il était d'attrait de se pencher sur un circuit septohippocampal-hypothalamo-tronc cérébral. Grâce à ces données, il a été révélé que le noyau prémammillaire dorsal (PMD) devait être impliqué de manière critique dans l'expression de l'évitement contextuel et la reconsolidation de la mémoire aversive. En utilisant des outils immunohistochimique, d'inhibition pharmacogénétique, d'optogénétique, nous avons analysé l'influence de la modulation du PMD et de ses principales cibles (à savoir, le PAG dorso-latéral et le Thalamus antéro-ventral (AMv)), sur le processus d'expression et de reconsolidation de l'évitement passif contextuel.

### **Résultat partie I : Rôle de la voie préfrontale – tronc cérébrale médiant le comportement d'évitement actif instrumental.**

Nos différentes études ont prouvé que le nouveau paradigme permettait aux animaux de répondre de manière passive (freezing), ou active (évitement) suivant le même stimulus conditionné. Ce paradigme, par la complexité de sa tâche, révèle une hétérogénéité chez les individus dans l'apprentissage de l'évitement, créant un groupe de bons esquivants et un groupe de mauvais esquivants. Par ailleurs cette hétérogénéité n'est pas retrouvée au niveau du taux d'expression de freezing pendant l'exercice. Une étude de nage forcée (FST), test utilisé en général pour vérifier l'efficacité des antidépresseurs, a prouvé que ces différences n'étaient

pas la résultante du paradigme d'évitement en lui-même, mais était inhérente aux individus. En effet, le test, utilisé à la fin des sessions d'apprentissage, n'a pas mis en valeur deux groupes distincts suivant le critère du temps d'immobilité dans l'eau. De plus, d'après une première étude d'immunohistochimie révélant la protéine C-Fos, marqueur de l'activité neuronale, nous avons pu remarquer que le CPFdm et les sous-régions dorsolatérales et latérales de la matière grise périaqueducale (dl/IPAG) sont préférentiellement activées lors du comportement d'évitement. En concordance avec les résultats de la thèse de Dr. Suzana Khoder démontrant qu'une sous-population de neurones du CPFdm vers le dl/IPAG code l'évitement actif, mais pas le comportement de freezing, il a été vérifié que, l'activation ou l'inhibition optogénétique de cette voie favorisait et bloquait l'acquisition de l'évitement conditionné, et n'avait pas d'effet sur le freezing. En effet, après les sessions d'acquisition de l'exercice, les souris n'ayant pas appris l'évitement (les mauvais esquivants) ont été stimulées optogénétiquement pendant deux sessions. Afin de mesurer l'effet de cette stimulation sur la voie CPFdm-dl/IPAG, deux sessions supplémentaires sans stimulations ont été nécessaires. Les souris ont ainsi amélioré leur performance d'évitement pendant et après la stimulation de la voie, mais le taux de freezing n'a pas été affecté. En ce qui concerne les souris ayant appris la tâche (les bons esquivants), aucun effet n'a été mis en évidence lorsque qu'on a inhibé la voie, suivant le même protocole expérimental. En revanche, une expérience plus probante a été l'inhibition optogénétique de la voie PFCdm-dl/IPAG à chaque session d'apprentissage. Les résultats indiquent que par cette inhibition quotidienne, aucune des souris n'a été capable d'apprendre l'évitement. A la suite de ces observations, il a pu être confirmé, et décrit dans la thèse de Dr. Khoder, que l'apprentissage de l'évitement était associé au développement de la plasticité des neurones du CPFdm au niveau de leurs synapses dans le dl/IPAG.

### **Résultat partie II : Rôle des voies hypothalamiques médiant l'évitement passif non instrumental.**

A la suite de cette première partie nous avons étudié l'influence des circuits hypothalamiques dans une situation d'évitement passif non instrumentale. Suivant une étude précédemment faite chez le rat, les animaux répondent de différentes manières en fonction des conditions dans lesquelles ils sont réexposés à un environnement devenu aversif. Il a été intéressant de constater que lorsque l'animal a la possibilité d'éviter le contexte aversif, il exprime une palette de comportements d'évaluation du risque comme des comportements d'approche et de scannage de l'environnement. Lorsque l'animal est confiné dans son contexte

aversif, il exprime majoritairement du freezing. Utilisant une technique de révélation de la protéine C-Fos, il a été mis en évidence une voie septohippocampale-hypothalamique-tronc cérébrale spécifique, impliquée dans la situation d'évitement passif, c'est-à-dire lorsque l'individu est libre d'accéder ou non au contexte aversif. Cette voie n'est par ailleurs pas impliquée lorsque l'animal est confiné dans un contexte devenu aversif. Afin d'en savoir davantage sur l'aspect fonctionnel du PMD, l'usage d'outils pharmacogénétiques a permis d'inhiber temporairement le PMD lors de l'exposition au contexte aversif après conditionnement. Cela a montré la diminution de l'expression des comportements d'évitement ainsi que la latence de l'animal à entrer dans la cage de conditionnement. De plus l'animal a été réexposé au même contexte le jour suivant, montrant les mêmes faibles réactions au contexte aversif. L'hypothèse a donc été émise que l'inhibition du PMD altérerait non seulement l'expression des comportements d'évitement, mais aussi la reconsolidation de la mémoire liée à ce contexte aversif. Enfin, par l'usage d'outils optogénétiques l'expérience a été répétée suivant le même protocole expérimental. L'inhibition au niveau des terminaux du PMD, qui projettent dans le dIPAG et l'AMv, altère de la même manière l'expression des comportements d'évitement et la reconsolidation de la mémoire.

## Conclusion

A la fin de ce projet il a pu être démontré dans un premier temps, qu'un circuit du cortex préfrontal dorsomédian projetant vers la partie dorsolatérale du PAG, est spécifiquement impliqué dans la réponse d'évitement actif ; et dans un second temps, l'évitement passif est contrôlé en partie par un noyau de l'hypothalamus postérieur, le PMD, et ce, de manière équivalente pour ses deux sites de projections principales, l'AMv et le dIPAG. Nos résultats révèlent une dynamique neuronale très intéressante entre des réponses impliquant des fonctions cognitives plus complexes, comme l'apprentissage de l'évitement actif signalé dépendant d'une voie spécifique du cortex préfrontal vers le dIPAG. Des réponses reposant sur un apprentissage contextuel plus simple telles que l'évitement passif, en revanche, dépendent de sites hypothalamiques, qui par ailleurs, sont également impliqués de manière critique dans les comportements défensifs innés liés aux menaces prédatrices et sociales.

## Table of content

Introduction.....	16
1) Active and Passive avoidance.....	16
a/ Fear defensive strategies .....	16
b/ Paradigms of avoidance learning.....	18
2) Summary of the structures involved in conditioned active avoidance.....	21
a/ structures involved in signalled active avoidance.....	21
b/Hypothesis and interests of the study .....	23
3) Summary of the structures involved in conditioned passive avoidance.....	26
a/ structures involved in passive avoidance.....	26
b/Hypothesis and interests of the study .....	27
4) Anatomy and connectivity of our structures of interest.....	27
a/ The dorso-medial Prefrontal Cortex .....	27
b/ The dorsal Premammillary nucleus of the Hypothalamus.....	30
<i>Aims of the thesis</i> .....	32
I/ Role of the prefrontal-brainstem pathway in mediating avoidance behavior .....	33
Materials and methods .....	33
Results .....	40
II/ Hypothalamic pathways of shock-based Passive avoidance .....	60
Materials and methods .....	60
Results .....	66
Discussion.....	85
First Part .....	85
Second Part.....	89
References.....	97

## List of Abbreviations

AAV	Adeno-associated virus
ACC	Anterior cingulate cortex
AHN	Anterio hypothalamic nucleus
AMPA	$\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid
AMv	Ventral part of the anteromedial nucleus of the thalamus
ArchT	Archeorodhopsin
BA	Basal nucleus of amygdala
BLA	Basolateral nucleus of amygdala
CAV	Canine adenovirus
CCK	Cholecystokinin
CeA	Central nucleus of amygdala
CeL	Centrolateral nucleus of amygdala
CeM	Centromedial nucleus of amygdala
CS	Conditioned stimulus
CS-	Control stimulus (not paired with shock)
CS+	Conditioned stimulus (paired with shock)
dIPAG	Dorsolateral periaqueductal gray
dmPAG	Dorsomedial periaqueductal gray
dmPFC	Dorsomedial prefrontal cortex
DREADD	Designer receptors exclusively activated by designer drugs
GABA	Gamma-aminobutyric acid
GFP	Green Fluorescent Protein
hM4Di	Gi-coupled human M4 muscarinic DREADD receptor
IL	Infra limbic
IN	Interneuron
LA	Lateral nucleus of amygdala
LH	Lateral hypothalamus
LHAjd	Lateral hypothalamic area juxtadorsomedial region
IPAG	Lateral periaqueductal gray
MD	Medial segment of the Thalamus

MDTB	Mouse Defense Test Battery
mPFC	Medial prefrontal cortex
Nac	Nucleus accumbens
NacC	Nucleus accumbens core
NacS	Nucleus accumbens shell
NMDA	N-Methyl- d-aspartic acid
NPY	Neuropeptide Y
PAG	Periaqueductal gray
PCA	Principal Component Analysis
PFC	Prefrontal cortex
PL	Prelimbic cortex
PMD	Dorsal Premammillary nucleus
PN	Pyramidal neurons
PPN	Putative pyramidal neurons
PTSD	Post-traumatic stress disorder
PV	Parvalbumin
RA	Risk assessment
SigA	Signalled active avoidance
SST	Somatostatin
SUBv	Ventral Subiculum
UR	Unconditioned response
US	Unconditioned stimulus
vHPC	Ventral hippocampus
VIP	Vaso active intestinal peptide
vIPAG	Ventrolateral periaqueductal gray
vIPFC	Ventrolateral prefrontal cortex
VMH	Ventral part of the medial hypothalamus
vmPFC	Ventromedial prefrontal cortex
VTa	Ventral tegmental area

## Introduction

### 1) Active and Passive avoidance

#### a/ Fear defensive strategies

Depending on the environment, animals present a repertoire of defensive behaviors related to their survival needs. Indeed, animals adopt defensive strategies to protect themselves and/or their conspecifics against environmental dangers. Moreover, when the danger is escapable, more active defensive behaviors such as avoidance, escape and flight are adopted (Blanchard and Blanchard, 1969; Ramirez et al., 2015). Adaptation includes selecting the appropriate defensive strategy taking into account its costs, the threat, and the context in which it occurs (Hofmann and Hay, 2018). As mentioned above, avoidance is one defensive strategy adopted when an individual is exposed to harm but has the possibility to put distance with the threat. However, under certain circumstances, for instance inescapable situation, individuals eventually adopt other defensive strategies, like freezing (LeDoux, 2012). Promising fields of research have been explored to study emotional coping strategies, and a large variety of paradigms have been developed in order to disentangle the circuits recruited in defenses responses of an individual to fear. The most complete study about defensive behaviors in rodents had been carried out by the Blanchards. The idea was to predict which defensive behavior would be selected depending on the different contextual and stimuli changes. An example of this grouping of tasks is the Mouse Defense Test Battery MDTB (Blanchard et al., 2003a; Blanchard, 2017). In these studies, numerous defensive responses in rodents exposed to threatful situations have been observed: flight, hiding, freezing, attack and risk assessment. An example of MDTB tests is a long oval runaway permitting to quantify escape behavior that can be modified and transformed to an unescapable arena to study the switch to freezing strategy. Indeed, freezing and avoidance has been one of the most studied defensive behaviors. Regarding freezing, some studies describe it as being a passive tonic immobilization (Blanchard and Blanchard, 1972; LeDoux, 2000), but other researchers argue that freezing is an active preparation state during which the organism gets ready to flight, avoid or fight (Gladwin et al., 2016). It is why this passive response is

interesting to be compared with active behaviors like avoidance, in terms of brain circuits and behavior selection.

During threatening escapable situation, individuals usually demonstrate predictable goal directed behaviors. Indeed, there are two main categories of motor responses learned under negative reinforcement: escape behavior and risk assessment/avoidance behaviors. Escape behavior is a motor action performed by the animal to terminate an ongoing aversive stimulus. This behavior is negatively reinforced by the elimination of the unpleasant stimulus. For instance, a rat will flee the room if receiving a shock on an electrified floor. Fleeing to stop the shock is an escape behavior. One characteristic of flight behavior, in more naturalistic situations, is that the initiation of the movement is very sensitive to the distance separating the animal from the potential threat. For example, the rat needs to be quite close to the predator to elicit a flight response, it is the concept of threat imminence (Kim et al., 2013). In learned tasks, escape behavior is converted into avoidance behavior by giving a signal before the aversive stimulus starts. In this case, avoidance represents complex motor actions learned by repetitive trials of conditioning paradigms (Moscarello and LeDoux, 2013). In innate situations, or open threatful environments, avoidance behavior would be the action to not approximate the localized threat, by scanning the environment with flat back approaches and oriented stretched postures (Blanchard et al., 2003b; Dielenberg et al., 2001). Risk assessment behaviors (RA) are expressed as mentioned before in natural conditions (Blanchard et al., 2003a), as for example the exposure to a cat or cat odor (Blanchard et al., 2005; Osada et al., 2013). RA also correspond to the animal scanning of the environment to detect routes of possible hiding or escape (Ellard and Eller, 2009). In a recent study, these behaviors are proposed to be a good model to compare anxious behaviors in human (Blanchard and Meyza, 2019).

These active defensive strategies are encountered in various escapable situations that can be modeled using different types of experimental avoidance paradigms.

## b/ Paradigms of avoidance learning

Most of the avoidance paradigms encountered in the literature are based on Pavlovian conditioning. It is the way to associate an aversive unconditioned stimulus (US), to a neutral conditioned stimulus (CS) which can be either an acute signal (Morgan and LeDoux, 1995), or the context itself (Baldi et al., 2004).

The association of the CS with the aversive US creates a new fearful memory and is considered as conditioned learning (Pavlov, 2010). The acquisition of a conditioned memory goes with the consolidation of the CS-US association. It refers to a memory that was temporary and labile, and enters into a more stable state (Lechner et al., 1999; Squire et al., 2015). This process is strengthened by an other stage of consolidation during sleep (Rasch et al., 2007). The memory formation processes occur primarily in the Hippocampus, and the memory storage is gradually distributed to cortical regions. Its dependence of the hippocampus then decreases. This is the process of systems consolidation of memory (Kitamura et al., 2017). Depending on the retrieval conditions, the conditioned memory will enter again in a labile state and can undergo either reconsolidation or extinction, maintaining or inhibiting the expression of the original fearful memory. These opposite memory processes recruiting specific signalling pathways are essential to re-stabilize the memory, in the case of reconsolidation, or inhibit the expression of the fearful memory, in the case of extinction (Merlo et al., 2015). When an individual has been confronted to an aversive situation, the right balance of the modulation of these two processes is crucial for the creation of a new safe environment (Lee et al., 2006). In the case of a persisting reconsolidation, poorly balanced by extinction of the fearful memory, it leads to pathological states like Post Traumatic Stress Disorder (PTSD) (Shin and Liberzon, 2010).

The dynamic of fearful memories is then important to untangle, regarding the memory stage, as well as the specific behaviors triggered by this memory. In the literature, the conditioning paradigms to study these phenomenon are diverse, and the use of specific US are of different natures. The most commonly used ones are mild electrical shocks both in humans (Löw et al., 2015) and rodents (Bravo-Rivera et al., 2015; LeDoux et al., 2017), however other types of US can be encountered, like airpuffs

(Moriarty et al., 2012), aversive odors (Osada et al., 2013), or predators (live or robots) (Blanchard et al., 2005; Kim et al., 2015). In the literature, avoidance paradigms are usually divided in two types of study: the active and the passive avoidance studies. The next paragraphs will describe the different paradigms encountered in each type of study, introducing our choice of paradigm to study instrumental active avoidance, and non instrumental passive avoidance.

#### *Instrumental Active avoidance paradigm*

This project will first focus on the strategy of active avoidance, which consists on taking action to prevent harm. It is often studied using one-way or two-way active avoidance paradigms. In one-way active avoidance paradigms, only one of the two chambers of a shuttle-box is aversive (Gebhardt et al., 2013) and associated with a shock presentation. In two-way active avoidance, both chambers can be aversive, therefore the behaviors expressed are less context dependent as compared to one-way avoidance paradigms. Two-way active avoidance paradigms can be either signalled by a stimulus such as a tone or a light, or unsignalled (Servatius, 2016). In unsignalled (or Sidman) avoidance conditioning, the individual receives an aversive stimulus at fixed intervals, without any warning signal. In order to reset the timer to zero and cancel the shocks, a shuttle to the other side is required. However, unsignalled active-avoidance is very difficult to acquire in rodents, which is why signaled two-way active avoidance is preferred in our case. The two-way signaled avoidance (SigA) is a more complex paradigm that involves two forms of conditioning, the Pavlovian and the instrumental, which produce conflicting behavioral responses, and must be reconciled to ensure that the individual responds adequately in order to avoid the aversive stimulus. Associative, or Pavlovian learning, is a simple and fundamental form of memory formation (Pavlov, 2010), where as described before, and individual associate and aversive unconditioned stimulus (US), to a neutral conditioned stimulus (CS). Instrumental, or operant conditioning, initiated by the behaviorist Skinner (Skinner, 1938), is the association of an action that will lead to a specific outcome when a motivational event is repeatedly displayed. This motivation, or reinforcement, to perform an action can be either positive or negative. the two-way sigA is and experiment that requires a shuttle-box separated

into two compartments by a door or a hurdle. The animal learns to cross during the warning signal to anticipate the delivery of the unconditioned stimulus (US) (Ramirez et al., 2015). Therefore, the two-way SigA is based on what is called the two factor theory proposed by Mowrer (Mowrer, 1947) as the task reconciles the two principles of Pavlovian and instrumental learning.

In our case, we chose the two-way signal active avoidance paradigm, with the difference that the contextual contingencies demonstrate either escapable (opening of the door between the two compartments), or unescapable (the door stays closed) situations. The rationales of this choice will be described later in the introduction.

#### *Non instrumental Passive avoidance paradigm*

Passive avoidance, also labelled inhibitory avoidance refers to abstaining from entering a likely to be aversive environment (i.e. entering a footshock compartment). It is important to note that passive avoidance does not mean passive coping behaviors. Interestingly, while assessing the environment and integrating aversive cues the individual expresses a range of risk assessment behaviors (RA), that are likely to be opposite to the freezing state (Blanchard et al., 2003a). Passive avoidance studies are of a strong importance for different reasons, as for instance investigating the neural circuits underlying the learning of “what to not do”. It is described in the step-down inhibitory avoidance paradigm, like deciding to step or not on an electrified platform where the animal had previously received a shock (Canto-de-Souza and Mattioli, 2016). The paradigm of contextual passive avoidance is also commonly used in innate threat exposure. The animal is usually exposed to a predator (a cat, an aggressive conspecific, or a snake) in a known environment. The next day, the animal is exposed to the same environment and has the possibility through a corridor to enter or not the predator cage (Gross and Canteras, 2012). Passive avoidance can also be implemented by using a two-compartment behavior apparatus, with a shock grid floor, the animal will receive a shock in the preferred compartment. The latency to enter the shock compartment again will be measured (Ambrogio Lorenzini et al., 1999; LeDoux et al., 2017). The passive avoidance tasks are interesting paradigms, as the acquisition is very rapid and hard to extinct, even

with the lack of negative reinforcement. Passive avoidance gives also the possibility to vary the nature of the threat (shock, predator exposure like a snake or a cat).

In our case, we used a novel paradigm previously implanted in our lab in rats (Viellard et al., 2016). we used an experimental apparatus developed for our experiments of fear conditioning to social and predatory threats as described above. In this case, the animal enters a shock-grid cage where it receives a series of shocks and is exposed to the whole apparatus (safe cage, corridor and grid cage) the next day, where the fear responses are measured.

## 2) Summary of the structures involved in conditioned active avoidance.

### a/ structures involved in signalled active avoidance

As described before, SigA paradigm involves complex mechanisms of conditioning learning and strategy coping. According to the Two-factors theory, in early-training phases, active avoidance learning depends on Pavlovian associative processes and lead to increased fear, expressed in terms of freezing. In a second step, avoidance responses are developed depending on instrumental associative processes to ultimately reduce the negative state generated by the CS presentation (LeDoux et al., 2017; Mowrer, 1947). In several avoidance studies both freezing and avoidance are quantified allowing to assess the effect of lesions on both freezing and avoidance behaviors in the same paradigm. So far the literature emphasize a strong role of the medial Prefrontal Cortex (mPFC) in coping strategy selection. It is a nucleus involved in higher processes, regulating a broad range of brain functions related to attention, executive control or working memory (Euston et al., 2012; Smith and Lane, 2016). It is also broadly investigated for its role in the regulation of emotional behavior as it is well known that the dysfunction of the mPFC is related to psychiatric conditions such as post-traumatic stress disorder (PTSD) (Shin and Liberzon, 2010). In the case of active avoidance, it is thought that lesions of the mPFC (ACC, IL and PL) disrupt the acquisition but not the expression of goal-directed behaviors pre-training (Gabriel et al., 1991). Lesions of the IL (the ventromedial PFC) region increased freezing

expression and disrupted two-way active avoidance learning (Moscarello and LeDoux, 2013). Furthermore, according to Moscarello and colleagues, the expression of passive freezing behavior and active avoidance are inversely correlated and depends on a balance of activity between the IL and the amygdala. However the role of the PL (the dorsomedial PFC) is not confirmed in the acquisition and expression of active avoidance (Moscarello and LeDoux, 2013). In contrast, other studies using different active avoidance paradigm show a role of the PL, and the ACC in the acquisition of avoidance learning (Bravo-Rivera et al., 2014). That is why for the time being, the dorso-ventral axis of the mPFC depending on conditions, does not have a clean frontier in terms of role in acquisition and expression of avoidance. One of the many targets of the mPFC is the Ventral Striatum, a particular region of this structure that seems to be involved in active avoidance would be the Nucleus Accumbens (Nac). Even though it has been widely studied in reward and appetitive reinforcement, some studies emphasized its role in acquisition of avoidance learning (Bravo-Rivera et al., 2014) and discrimination of the aversive CS with neutral tones (Oleson et al., 2012). There seems to be a complex implication of the core of the Nucleus accumbens core (NacC) and the shell (NacS) that are respectively, involved in the acquisition and the expression of active avoidance (Moscarello and LeDoux, 2013). However the role of the NacC is still unclear as contradictory studies have been published refuting its role in acquisition (Corbit et al., 2001; Ramirez et al., 2015). Furthermore, the Amygdala is indeed a structure broadly studied for its role in classical fear conditioning and freezing expression (Herry et al., 2006; Maren, 2001). Is also a candidate for active avoidance, but working in an opposite manner as it does for freezing. Indeed amygdala nuclei are reported to participate differentially in avoidance acquisition. First, the LA is shown to be crucial for the acquisition of both freezing and avoidance behaviors (Amorapanth et al., 2000). The BLA and LA but not the CeA impaired the acquisition of Sidman active avoidance behavior in rodents (Lázaro-Muñoz et al., 2010). Even if the LA is an important site for storing the CS-US association, there are probably other circuits regulating that same function as the lesion of the latter impairs early sessions of active avoidance learning, but not the late ones. Moreover, lesioning the CeA nucleus blocks freezing responses and can facilitate avoidance behavior learning in bad performers (Lázaro-Muñoz et al., 2010). In conclusion the amygdala seems to be crucial in short term

avoidance expression but probably relays the information to other systems for long term memory, which could be involving the ventral hippocampus. Indeed a study demonstrated that the ventral hippocampus contributes to the two-way sigA learning (Wang et al., 2015). Another structure that could be part of a putative pathway for avoidance processing, is the periaqueductal gray matter. The first evidence of the involvement of the PAG in mediating defensive behavior was carried out by Bandler and colleagues. The injection of excitatory Amino acid in different parts of the PAG shed light on the different roles of its columns (Keay and Bandler, 2001). The dorsal PAG is a key structure for flight responses and other active behaviors like aggression (Motta et al., 2017). Whereas the ventral columns are inducing more passive behaviors like freezing (Carrive, 1993; Kim et al., 2013). However electrical stimulations of the dorsal PAG of different intensity induce first freezing then flight responses (Vianna et al., 2001). These works point out the dual role of the dorsal PAG on active and passive behaviors, and the complexity of the PAG columns communication. Likewise more recent studies on a communication circuit between the ventrolateral and the dorsolateral PAG showed that the activation of the dIPAG glutamatergic projections to the vIPAG blocks freezing and promotes active defensive behavior expression (Tovote et al., 2016). The dmPFC and Lateral Hypothalamus are potential candidates to mediate this circuit, as they both projects on the dIPAG (Halladay and Blair, 2015).

To summarize, various structures of the brain have a role in either active avoidance or freezing, but specific studies show a clear role of the mPFC, the amygdalar nuclei and the PAG in monitoring active avoidance system as well as freezing expression.

## b/Hypothesis and interests of the study

The literature suggests that, among other structures, the interaction between the Amygdala, the mPFC and the PAG are key structures for driving adapted fear behaviors. Yet it is still unclear if freezing and active avoidance rely on the same, or different circuits. And the structure involved in processing avoidance behavior and the contribution of distinct prefrontal circuits to both freezing and avoidance responses are largely unknown. Our interest is to understand which projection of the dmPFC is a key switch between

avoidance and freezing. The role of the amygdalar nuclei, as described above is major in these two behaviors, however they do not seem to have the same dynamic while processing them. That is why our attention focused on the dIPAG, considering the fact that the structures hosts neural processes implicated in both behaviors. To further investigate the role of dmPFC circuits in encoding passive and active fear coping strategies, in the laboratory of Cyril Herry, I worked in collaboration with Suzana Khoder who had developed a novel behavioral paradigm in which a mouse has the possibility to either passively freeze to an aversive stimulus, or to actively avoid it, depending on the contextual contingencies. Using this behavioral paradigm we investigated whether the same circuits mediate freezing and avoidance behaviors or if distinct neuronal circuits were involved. **To address this question, a combination of behavioral, neuronal tracing, immunochemistry, single-unit, patch-clamp recordings and optogenetic techniques were used to study the role of the dmPFC to dIPAG pathway in both active avoidance and freezing acquisition and expression. As Dr. Suzana Khoder published her thesis last year, I will briefly explain in this introduction a part of the conclusions of her thesis, and develop with more details my contribution to the work in the “Result” section of part I.**

*After validating the behavior paradigm, it was demonstrated that the active avoidance learning paradigm using a two way shuttle box showed variability of learning between the two groups. The good avoiders, who would discriminate the task and learn to avoid at the tone onset (CS) avoiding the shock, by shuttling to the other compartment, in the open door situation. When the door remained closed during the CS paired to a shock (CS+), the good avoiders also learned to freeze and discriminate with the unpaired CS (CS-). The bad avoiders were unable to learn the task after six days of training, and would freeze more in the closed door situation. However they were able to discriminate between the CS+ and the CS-.*

*Using in vivo electrophysiological recordings, the results showed that the dmPFC of Good avoiders indicated that most avoidance-inhibited dmPFC PPNs (putative pyramidal neurons) are modulated by both freezing and avoidance, while most avoidance-activated dmPFC PPNs are modulated exclusively by avoidance behavior. Moreover, it has also been demonstrated that changes in firing activity of avoidance-activated dmPFC*

*neurons is not an effect of an increase in locomotion during avoidance and likely reflects associative learning. Furthermore the antidromic stimulations data clearly indicated that the subpopulation of dmPFC PPNs neurons exhibiting an increased activity during avoidance learning (avoidance-activated / freezing non responsive cells) project to the dlPAG.*

*It was then interesting to take advantage of the fact that a subgroup of animals could not learn the avoidance task. In this view the PL to dlPAG pathway was activated using optogenetic tools. The data pointed out that optogenetic stimulation of dmPFC-dl/IPAG projections progressively promotes learning of avoidance behavior. Once again it has also been proved that the 10Hz optogenetic stimulation of the pathway is not a locomotor effect. Thus, supporting the electrophysiological results, the activation of dmPFC neurons projecting to the dl/IPAG did not affect conditioned freezing behavior.*

*To reinforce these data, it was also demonstrated using in vitro whole cell recordings by measuring the AMPA/NMDA receptor ratio, that the switch of Bad avoiders into Good avoiders upon the optogenetic stimulation of the dmPFC dl/IPAG pathway is associated with the development of synaptic potentiation at dmPFC inputs onto dl/IPAG cells. (Khoder, 2018)*

In order to give a stronger insight on these differences between the two groups, my contribution to the work was to investigate the activation pattern of the mPFC and the PAG, in good and bad avoiders. Also, it is still unclear, with the optogenetic activation of the pathway only, to understand if the cells projecting from the dmPFC to the dlPAG are involved in the expression of avoidance and/or its acquisition. These questions will then be assessed using Fos immunochemistry analysis, inhibitory optogenetic strategies, and other behavioral tests.

### 3) Summary of the structures involved in conditioned passive avoidance.

#### a/ structures involved in passive avoidance

Passive avoidance is a non instrumental form of avoidance and has been studied through lesion studies, using inhibitory avoidance tasks. For instance, the literature indicates that the mPFC is a potential structure to mediate passive avoidance. Indeed, studies of the mPFC demonstrate that lesions of the PL in rats in the step-through passive avoidance paradigm impaired fear memory whereas a stimulation of the region improves it (Canto-de-Souza and Mattioli, 2016). The ventral hippocampus and lateral septum also have an important role in the encoding of association of contextual cues (Gross and Canteras, 2012). In fact, the septo-hippocampal system has been proposed to play a pivotal role in anxiety in response to conflicted situations, by interrupting ongoing behavior and increasing the level of arousal and attention to enhance gathering information (Gray and McNaughton, 2003). It is also known that anxiogenic-like state is provoked by selective stimulation of BLA to ventral hippocampus projections. And the stimulation of the amygdala has been shown to disrupt inhibitory passive avoidance (Gold et al., 1973). Notably, the striatum has been demonstrated to play a role, but as opposed to active avoidance, fear retrieval following conditioning is disrupted by the nucleus accumbens shell, but not core, region inactivation (Piantadosi et al., 2018). It is important to note that most of the data on passive avoidance were collected using step-through or step-down inhibitory avoidance paradigms. The difference with the paradigm used in this study is the proximity with the threatening location. The presence of a corridor imposes a distance between the safe cage and the conditioning cage, and this situation may need the recruitment of particular neural circuits for the encoding and retrieval of fearful contextual cues. So far, according to a fos study in rats using this experimental apparatus on shock based passive avoidance, the circuit recruited during fear retrieval seems to involve a specific septo/hippocampal-hypothalamic-brainstem pathway, namely the ventral subiculum, lateral septum, the juxtadorsomedial part of the lateral hypothalamus (LHAjd), the dorsal premammillary nucleus (PMD) and the dorsal and lateral parts of the

PAG (dl/IPAG) (Viellard et al., 2016). Notably, the PMD occupies a pivotal role in this circuit, and its hodological relationships will be discussed below.

## b/Hypothesis and interests of the study

According to recent work of our lab, and the literature on innate fear learning, we are understanding that shock-based contextual passive avoidance is mediated in part by a septo/hippocampal-hypothalamic-brainstem pathway. We are hypothesising that a key structure of this pathway is the PMD. The PMD is largely influenced by septo-hippocampal processing (Comoli et al., 2000) and, on the efferent side, sends projections to both the AMv (likely to be involved in encoding fear memory) and the dorsal PAG, which is known to participate in the expression of active and passive defensive behaviors. After validating the activation of this septo/hippocampal-hypothalamic-brainstem pathway in mouse using a Fos immunochemistry analysis, we will focus on the PMD and its projection sites using a combination of behavioral, pharmacogenetic and optogenetic approaches, to evaluate their roles on the expression of inhibitory avoidance and fear memory reconsolidation process.

## 4) Anatomy and connectivity of our structures of interest

This next paragraph will give a rapid insight on the anatomy of our two structures of interest: the dmPFC and the PMD.

### a/ The dorso-medial Prefrontal Cortex

The rodent mPFC can be divided into four distinct areas which are, descending from the most dorsal region, the medial precentral cortex (PrCm), the Anterior Cingulate Cortex (ACC) (dorsal and ventral part), the Prelimbic Cortex and the Infralimbic Cortex. Our interest will focus on the dorsal part of the medial cortex, which are the PL and the ACC. The ACC areas regulates various motor behavior, whereas the PL will regulates emotional, mnemonic, and cognitive processes (Heidbreder and Groenewegen, 2003). The cortex has a paralleled laminar organization divided in 6 layers that are called from the most superficial to the deepest: the molecular layer, which is poorly dense in neurons;


the external granular layer; the external pyramidal layer; The layer 4, or granular layer, (this layer is not present in rodents); the internal pyramidal layer, that is composed of sparse and large Pyramidal neurons (PNs) vertically oriented; and lastly the polymorph layer, that contains various neuronal types without specific organization. These six layers have their own organization in terms of connectivity. For instance, layers 2/3 support the cortico-cortical connections, layers 1 and 4 receive thalamic inputs and layers 5 and 6 are respectively the main sources of thalamic and subthalamic projections (Harris and Shepherd, 2015).

The cortex is composed of two main classes of neurons: the glutamatergic pyramidal neurons (PNs) and the GABAergic interneurons (INs) that represent respectively 80% and 20% of the cortical neurons. PNs used glutamate as a neurotransmitter and are located in all six cortical layers, except layer 1. As opposed to PNs, the vast majority of INs do not leave the cortex and are restricted to a local environment (Spruston, 2008). That is why our projection studies will focus on PNs. Finally, INs can be characterized at neurochemical level, indeed a numerous variety of peptides is encountered in interneurons, that give them their neuronal subtypes (i.e. PV, CR, CB, SST, VIP, CCK, NYP). Because of their morphological, electrophysiological and molecular diversity, INs are believed to differentially sculpt cortical activity. (Hubel and Wiesel, 1962).

*Inputs:* The dmPFC, our structure of interest, receives its major inputs from the medial segment of the thalamus (MD), it also projects back through a descending pathway to the MD (Groenewegen, 1988). The whole PFC, including the PL receives also massive inputs from the BLA. The paralimbic cortex sends reciprocal projections back to the PL. Another important input is coming from the vHPC (CA1 region and subiculum) and terminates in all layers of the PL, with sparse inputs from the dorsal hippocampus.

*outputs:* The ACC and PL project mostly to the BA, as opposed to the IL for example, that will send projections to the CeA and LA (Hoover and Vertes, 2007). The mPFC shares reciprocal connectivity with the ventral tegmental area (VTA), the basal ganglia (Groenewegen, 1988), and the dorsal and lateral regions of the PAG (Gabbott et al., 2005). It also projects to the hypothalamus, like the PMD (Comoli et al., 2000). The PL

also project internally to the ventral ACC and the IL region sending outputs preferentially to the PrCm and dorsal ACC (Hoover and Vertes, 2007).



**Figure 1** Schematic sagittal sections summarizing the main efferents projections of the PL in rats Sections are modified from the rat atlas of Paxinos and Franklin (Paxinos and Franklin, 2008)

Abbreviations: AA, anterior area of amygdala; AHN, anterior nucleus of hypothalamus; Al,d,v, agranular insular cortex, dorsal, ventral divisions; AM, anteromedial nucleus of thalamus; AON, anterior olfactory nucleus; BMA, basomedial nucleus of amygdala; C, cerebellum; CEM, central medial nucleus of thalamus; CLA, claustrum; COA, cortical nucleus of amygdala; C-P, caudate/putamen; DBh, nucleus of the diagonal band, horizontal limb; DMH, dorsomedial nucleus of hypothalamus; DR, dorsal raphe nucleus; EN, endopiriform nucleus; IAM, interanteromedial nucleus of thalamus; IC, inferior colliculus; IMD, intermediodorsal nucleus of thalamus; IP, interpeduncular nucleus; LHy, lateral hypothalamic area; LPO, lateral preoptic area; LS, lateral septal nucleus; MEA, medial nucleus of amygdala; MO, medial orbital cortex; MPO, medial preoptic area; MR, median raphe nucleus; N7, facial nucleus; OT, olfactory tubercle; PBI, parabrachial nucleus, medial and lateral divisions; PFC, perifornical region of hypothalamus; PN, nucleus of pons; PRC, Reunions nucleus; RE, perirhinal cortex; RH, rhomboid nucleus of thalamus; SI, substantia innominata; SLN, supralemniscal nucleus (B9); SUM, supramammillary nucleus; TTd, taenia tecta, dorsal part; VLO, ventral lateral orbital cortex; VO, ventral orbital cortex. Reprinted from Vertes (2004).

## b/ The dorsal Premammillary nucleus of the Hypothalamus

The PMD is a small dense structure of the posterior ventral hypothalamus with anatomical and neuronal properties that are poorly understood. It is part of the mammillary complex but has unique projections and functions compare to the other mammillary nuclei (Canteras and Swanson, 1992). It is known to be a structure mostly glutamatergic.

*inputs:* The PMD receives a dense input from the ventral tegmental nucleus , and unlike other mammillary nuclei, the PMD does not receive a direct input from subicular regions of the hippocampal formation but instead it receives a massive input from the anterior hypothalamic nucleus and the juxtadorsomedial part of the lateral hypothalamic area (LHAjd) (Comoli et al., 2000; Hahn and Swanson, 2012).The anterior hypothalamic nucleus integrates and transmits (either directly or indirectly) information from the prefrontal cortex, amygdala, hippocampus, and septal region, whereas the LHAjd receives massive inputs from the subiculum and lateral septum (Comoli et al., 2000; Hahn and Swanson, 2012). Moreover, more sparse inputs to the PMD arise from the ventromedial hypothalamus (VHM)and prelimbic cortex PL (Canteras and Swanson, 1992; Comoli et al., 2000).

*Outputs:* ascending branch of the PMD projection ends massively in the ventral part of the anteromedial nucleus of the thalamus (AMv) and anterior hypothalamic nucleus (AHN); this branch also provides moderate inputs to rostral parts of the zona incerta, the nucleus reuniens, and to perifornical areas of the lateral hypothalamic area. The descending branch of the PMD projection courses to and through the posterior hypothalamic nucleus and end densely in the dorsolateral part, but also the medial and lateral parts of the periaqueductal gray; as well as in the deep and intermediate gray layers of the superior colliculus, and caudal parts of the midbrain reticular formation (including the cuneiform nucleus) (Canteras and Swanson, 1992).




FIG. 4. Major input/output relations of the PMd, medial mamillary nucleus (MM), and reticular part of substantia nigra (SNr). Hypothetical pathway from the periaqueductal gray (PAG) to the spinal cord (SP) may be direct or indirect. AHN, anterior hypothalamic nucleus (n.); AM, anteromedial n. thalamus; AMY, amygdala; CER, cerebellum; CG, cingulate gyrus; CUN, cuneiform n.; MO, motor cortex; MRN, mesencephalic reticular n.; PAG, periaqueductal gray; PFC, prefrontal cortex; SBC, subicular complex; SC, superior colliculus; SP, spinal cord; SUBv, ventral subiculum; TRN, tegmental reticular n.; VM, ventral medial n. thalamus; VTN, ventral tegmental n.

Figure 2 . Scheme of the inputs and outputs related to the PMD (Canteras and Swanson, 1992)

## *Aims of the thesis*

### **First part: Role of the prefrontal-brainstem pathway in mediating avoidance behavior**

1. Establish the behavioral paradigm for SigAA, and determine the criteria for the good and bad avoiders;
2. Examine the fos activation pattern of the PAG, mPFC and amygdala
3. Inhibit the dmPFC-dIPAG pathway during avoidance using optogenetic to untangle whether the dmPFC-dIPAG pathway is involved in the expression and/or acquisition of avoidance behavior.

### **Second part: Hypothalamic pathways of shock-based Passive avoidance**

1. Validate the passive avoidance paradigm and the Fos activation pattern of the septohippocampal-hypothalamic-brainstem pathway
2. Investigation of the role of the PMD in the expression and reconsolidation processes of contextual passive avoidance.
3. Inhibition of the projections from the PMD to its main target (the dorsal PAG and ventral part of the anteromedial thalamic nucleus) and study their influence on the expression and reconsolidation processes of contextual passive avoidance.

#### **Note :**

The results section is divided in two main parts, and each result part is preceded by its own material and methods section. The first part is devoted to the investigation of role of the prefrontal-brainstem pathway in mediating avoidance behavior, a work carried out in Dr. Herry's lab, at the university of Bordeaux. This part of the result section is my contribution to the work of Dr. Suzana Khoder's, that has been published in her thesis in november 2018. The manuscript of an article is under finalization before submission to a high profile journal where Suzana and I will be first co-author of the work. The second part of the results is a second project carried out in the university of São Paulo, in the lab of Prof. Newton Canteras during which we investigated Hypothalamic pathways of shock-based Passive avoidance.

# I/ Role of the prefrontal-brainstem pathway in mediating avoidance behavior

## Materials and methods

### I. Animals

Animals SigAA paradigm: Male C57BL6/J mice (3 months old; Janvier) weighing 30-35 g at the time of surgery, were group-housed upon arrival in a 22°C colony room, on a 12-hours light/dark cycle (lights on at 7 a.m.) and were provided with food and water ad libitum. Mice were housed individually 3 days before surgeries and manipulated daily. Animals' experiments were carried out in accordance with standard ethical guidelines (European Communities Directive 86/60-EEC).

### II. Experimental protocol

The apparatus is composed of two identical square compartments of 25cmx25cm. It is separated by a descending door linked to the Imetronic software. The sound and the shock are also monitored by the Imetronic software.

## Behavioral Paradigm




Figure 3 **Behavioral paradigm.** In the door opened trials (DO), 3 types of behavioral readouts were scored: **error trials** during which animals stayed during the whole CS<sup>+</sup> and the US delivery; **escape trials** during which mice crossed to the opposite compartment of a shuttle-box during the US and **avoid trials** during which animals crossed during the CS<sup>+</sup> and avoided the US. In door-closed trials (DC), **freezing** behavior was assessed during the sound presentation.

Avoidance learning protocol:

**Habituation to context** (Figure 4, first block) : On day 1, the mice were placed in the shuttle box and habituated to walk freely between the two compartments for 15 minutes.

**Habituation to door and tones** (Figure 4, second block) : On day 2, the mice were placed in the apparatus and were habituated to the protocol events of the acquisition task, without receiving a shock.

The events were the followings : door openings and different tones (7.5KHz pips for CS+, white noise for CS-). The different trials were the followings:

- CS+/CS- door opened (DO) : A sound was presented for 23.1 seconds (7.5kHz pips for CS+, white noise for CS-) with the door closed. The door opened and the sound continued for 8.9 seconds which was the time for the animal to shuttle to the other compartment. The door closed again either at the 8.9 seconds, or at the moment the animal passed the door.
- CS+/CS- door closed (DO) : A sound was presented for 33 seconds (7.5KHz pips for CS+, white noise for CS-).

The animals were left 3 minutes freely exploring the environment at the beginning and at the end of each session. All trial – CS+DO, CS-DO, CS+DC, CS-DC – were presented in a shuffled manner 9 times each. An inter trial interval (ITI) with a value from 15 to 40 seconds was used with the door opened between each trial. The number of shuttles for each type of trials (CS+DO or CS-DO) was counted, in order to monitor the effect of the sound type on the shuttling rate.

**Acquisition** (Figure 4, third block) : During day 3 to 8, the mice were placed in the apparatus to performed a 6-day training to acquire the task.

- CS-/CS+ door opened : A sound was presented for 23.1 seconds (7.5 kHz pips for CS+, white noise for CS-) with the door closed. The door opened and the sound continued for 8.9 seconds which was the time for the animal to shuttle to the other compartment. For the CS+ trials, if the mice had not shuttled during the

door opened time laps, a shock was triggered (4 sec, 0.6mA), with the door left opened. The door closed again either after the 4 second shock, or at the moment the animal passed the door. If the animal shuttles during the 8.9 seconds, it is an avoid trial, if the animal shuttles during the shock, it is an escape trial, if the animal does not shuttle, it is an error trial (Figure 3).

- CS-/CS+ door closed : A sound was presented for 33 seconds 7.5kHz pips for CS+, white noise for CS-). For the CS+DC trials, the 33 second sound is followed by a 4 second shock. The door stayed closed during the whole trial.

All trials – CS+DO, CS-DO, CS+DC, CS-DC – were presented in a shuffled manner 15 times each. An inter trial interval (ITI) with a value from 15 to 40 seconds was used with the door opened between each trial. The following outcomes were counted: number of error trials, escape trials, avoid trials, the freezing level, latencies (for avoid and escape trials), discrimination index (difference of avoid trials between the CS+DO and the CS-DO).

At the end of the acquisition sessions (day 8), the animals were divided into two groups:

- The good avoiders : with an avoid trial level superior or equal to 5 and a discrimination index superior or equal to 20%.
- The bad avoiders : with an avoid trial level inferior to 5 and a discrimination index inferior to 20%.




Figure 4 **Behavioral protocol** On day 1, mice were habituated during 15 min to the shuttle-box. On day 2, animals were habituated to the presentation of two sounds that were played in two contextual conditions : (i) door-closed trials (DC) during which the sound was played for 33 s , and (ii) door-opened trials (DO) during which 23.1 s following the sound's onset the door was slid-down (DO) and slid-up again 8.8 sec after. 9 trials of each of the DO and DC types of trials were played for both CS<sup>-</sup> and CS<sup>+</sup> and the session lasted about 45 min. From day 3 to day 8, animals underwent 6 training sessions lasting each about 1h20min and during which the same type of trials than during day 2 were played except that the number of trials was increased to 15, and that CS<sup>+</sup> trials were followed by a 4 s shock in the DC condition. At day 8, animals were categorized into **Good** or **Bad avoiders** based on their behavioral avoidance scores.

### III. Forced-swim test (FST)

Following avoidance training, **Good** and **Bad avoiders** underwent a FST session during which each mouse was individually placed in a cylindrical tank (50 cm height and 20 cm width) filled with clean tap water ( $24 \pm 1$  °C). Mice were forced to swim during 6 minutes. The first two minutes were considered as an acclimatization time and during the last 4 minutes, Behavioral Observation Research Interactive Software (BORIS) was used to process the recorded behavioral video allowing to score the total duration of immobility. Mice were scored to show immobility when they floated without struggling and making only those movements necessary to keep their heads above the water: namely moving only their hind paws but not front paws. In the end of the FST, mice were carefully dried before being returned to their home cages.

### IV. Histological processing of Fos immunohistochemistry

Following the 6th session of training mice were divided into three groups and underwent a last 7th behavioral session. The first control group received, at a 7th behavioral session, only 15 CS- trials. The second and the third groups, which were respectively classified at session 6 as Bad and Good avoiders underwent a 7th behavioral session during which they received 15 trials of CS+ presentations without footshocks. A forth group of naïve mice was also used as a control. Ninety minutes after the 6th behavioral session (not for the naïve mice), The animals were perfused after being deeply anesthetized with sodium pentobarbital (40 mg/kg, i.p.) and perfused transcardially with a solution of 4 % paraformaldehyde in 0.1 M phosphate buffer (pH 7.4). The brains were removed and left overnight in a solution of 20% sucrose in 0.1 M phosphate buffer at 4 °C. The brains were then frozen, and 5 series of 40- $\mu$ m-thick sections in the frontal plane were cut using a sliding microtome. One series of sections was processed for immunohistochemistry. The sections were incubated with anti-Fos antiserum raised in rabbit (Ab-5; Calbiochem) at a dilution of 1:20,000. The primary antiserum was detected using a variation of the avidin–biotin complex system. In brief, the sections were incubated for 90 min at room temperature in a solution of biotinylated goat anti-rabbit

IgG (Vector Laboratories) and then placed in a mixed avidin–biotin horseradish peroxidase (HRP) complex solution (ABC Elite Kit; Vector Laboratories) for 90 min. The black-blue peroxidase complex was visualized after a 5 min exposure to a chromogen solution containing 0.02% 3,3′ diaminobenzidine tetrahydrochloride (DAB; Sigma) with 0.3% nickel ammonium sulfate in 0.05 M Tris buffer (pH 7.6), followed by a 20 min incubation in a chromogen solution containing hydrogen peroxide (1:3000). The reaction was stopped using potassium phosphate-buffered saline (KPBS; pH 7.4). The sections were mounted on gelatin-coated slides, dehydrated and cover slipped using DPX mounting media (Sigma). An adjacent series of sections was stained with thionin (Nissl stain) to serve as a reference series for cytoarchitectonic purposes. Images of the selected brain regions (PFC, PAG, amygdala) were generated using a Nikon Eclipse 80i (10 x magnification, Nikon Corporation, Chiyoda-Ku, Tokyo-To, Japan) microscope equipped with a Nikon digital camera (DXM1200F, Nikon Corporation). To quantify the density of the Fos-labeled cells, we first delineated the borders of the selected brain regions by referring to the reference (Nissl-stained) sections and the mouse brain atlas (Franklin and Paxinos, 2008). Then, the Fos-labeled cells were counted. Only darkly labeled oval nuclei that fell within the borders of a region of interest were counted. The density of Fos labeling was determined by dividing the number of Fos-immunoreactive cells by the area of the region of interest. Both the cell counting and area measurements were performed with the aid of a computer program (Image-Pro Plus, version 4.5.1; Media Cybernetics, Silver Spring, MD, USA). Cell densities were obtained on both sides of the brain and were averaged for each mouse.

## **V. Viral injections and optogenetics**

For specific optogenetic manipulation of the dmPFC-dl/IPAG pathway during behavior we used C57BL6/J wild-type mice. Animals were bilaterally injected with glass pipettes (tip diameter 10-20  $\mu$ m) connected to a picospritzer (Parker Hannifin Corporation;  $\sim$  0.2  $\mu$ L per hemisphere) with a cocktail of Cav2-Cre, HSV-Cre retrograde virus and AAV-hSyn-mCherry in the dl/IPAG at the following coordinates relative to bregma:  $-4.50$  mm AP;  $\pm$  0.5 ML;  $-1.45$  DV from dura. The same animals received also an injection of AAV9-FLEXArchT-GFP, or AAV5-FLEX-GFP (UNC Vector Core Facility) in the dmPFC at the

following coordinates relative to bregma: +1.8 mm AP;  $\pm$  0.40 ML; -1.3 DV from dura. Following 4 weeks of recovery from injections, mice were implanted with optic fibers in the two hemisphere at the following coordinates relative to bregma: +1.8 mm AP;  $\pm$  0.55 mm ML; -1.15 mm DV from dura; lowered at an angle of 10°.

#### *Optical stimulation*

A laser generating a continuous yellow light at 593 nm or a blue light at 473nm (DPSSL lasers) was connected to a 200  $\mu$ m diameter optic fibre patch cable (Plexon) and calibrated to produce a fibre tip irradiance of approximately 16-18 mW. Computer software (Imetronic) controlled the timing and duration of laser pulses. Laser stimulations were delivered to the dmPFC to transiently activate or inactivate pyramidal projection of the dmPFC neurons to the dIPAG.

## **VI. Statistical analysis**

Analyses were performed with Matlab, Graphpad Prism and Statview. For all datasets normality was tested using the Shapiro-Wilk normality test ( $\alpha < 0.05$ ) to determine whether parametric or non-parametric analyses were required. Parametric analyses included t. tests and one- and two-way repeated-measures ANOVA followed by Bonferroni's, or Fisher's PLSD (for behavior only) multiple comparison post-hoc test if a significant main effect or interaction was observed. For parametric data, correlation analyses were made using Pearson's correlation. If datasets did not meet normality assumptions non-parametric analyses were used (mainly non-parametric Mann-Whitney test). If significance was observed, these non-parametric analyses were followed by Dunn's multiple comparison post hoc tests. For non-parametric data, correlation analyses were made using Spearman's correlation. Apart from t. tests, the asterisks in the figures represent the P-values of post hoc tests corresponding to the following values \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$  based on mean  $\pm$  S.E.M.

## Results

### **I. Behavior**


In order to investigate the neural circuits underlying freezing and avoidance responses, a novel paradigm was developed. It would create the possibility to condition the animal with the same CS to both an open door situation (DO) where the animal has the possibility to escape/avoid a shock, and a closed door situation (DC) where the animal would only have the choice to freeze waiting for the shock.

The results showed that this behavioral paradigm generated a difference of learning profile within the group of trained mice (Figure 5. **a, b**). The animals were then categorized in two groups, the bad avoiders and the good avoiders. The bad avoiders (55%) showed a deficit in learning performances as they couldn't reach 5/15 avoidances after the 6<sup>th</sup> day of training. The good avoider group (33%) (Figure 6d) was selected following two parameters. First, they had to reach 5/15 avoidances at the end of the 6th session, second, they had to discriminate between the CS+ and the CS-, according to the avoidance discrimination index (Figure 6a). Notably, 13% of the animals (Figure 6d.), successfully avoided during the CS+ but also during the CS-. These animals were called the generalizers and were excluded from the analysis as they had a very low discrimination index (Figure 6b,d). Another indicator of learning was the latency to avoid. The good learners learned to avoid the CS+ faster than the CS- (Figure 5d). Interestingly, the bad avoiders did not learn to avoid, however, the escape latency decreases across the sessions (Figure 5c.). Importantly, the escape latency of the good avoiders also decreased (Figure 5, c.), but was not different from the bad avoiders. It indicates that the good and the bad avoiders had the same escape kinetics.

Regarding freezing expression, the Bad avoiders froze significantly more than the Good avoiders during CS+ presentation in DC condition. However they had a significant difference of freezing level between the CS+ and the CS- (Figure 5e). It indicates that they would discriminate between the two CSs. During the DO condition they showed no difference of freezing before and after the door opening (Figure 5f).


For the Good avoiders, they also discriminated the CS+ and CS- in terms of freezing level during the DC condition (Figure 5**g**). And as they would learn to change their strategy during DO condition and avoid, as expected, they would have a significant reduced freezing level after the opening of the door (Figure 5**h**.).

It should be noted that freezing discrimination was rapidly acquired in both groups, as the difference is clear from the first day of training. However the avoidance acquisition was a longer process, as the good avoiders would learn to avoid and discriminate between the CSs at late training stage (Figure 5**b**., Figure 6**a**.).



**Figure 5 Behavioral characterization of Good and Bad avoiders.** **a.** Trial counts (shuttle CS<sup>-</sup>, avoid CS<sup>+</sup>, escape, errors) across 6 training sessions in **Bad avoiders** ( $n = 22$ ) (two-way repeated measures ANOVA; group:  $F_{(3,420)} = 104.5$ ,  $p < 0.0001$ , training session:  $F_{(5,420)} = 0.50$ ,  $p = 0.77$ , group  $\times$  training session:  $F_{(15,420)} = 3.98$ ,  $p < 0.0001$ ). **b.** Trial counts (shuttle CS<sup>-</sup>, avoid CS<sup>+</sup>, escape, errors) across 6 training sessions in **Good avoiders** ( $n = 13$ ) (two-way repeated measures ANOVA; group:  $F_{(3,240)} = 28.43$ ,  $p < 0.0001$ , training session:  $F_{(5,240)} = 1.13$ ,  $p = 0.34$ , group  $\times$  training session:  $F_{(15,240)} = 8.83$ ,  $p < 0.0001$ , and  $p < 0.01$ ). **c.** Escape latency (s) for both **Good** and **Bad avoiders** across 6 training sessions (two-way repeated measures ANOVA; group:  $F_{(1,184)} = 0.54$ ,  $p = 0.45$ , training session:  $F_{(5,184)} = 2.55$ ,  $p = 0.02$ , group  $\times$  training session:  $F_{(5,184)} = 0.82$ ,  $p = 0.53$ ). **d.** Avoid latencies during DO trials for both CS<sup>+</sup> and CS<sup>-</sup> trials in **Good avoiders** ( $n = 13$ ) (two-way

repeated measures ANOVA; group:  $F_{(1,114)} = 1.56, p = 0.21$ , training session:  $F_{(5,114)} = 7.03, p < 0.0001$ , group x training session:  $F_{(5,114)} = 2.45, p = 0.03, * p < 0.05$ ). **e.** Averaged freezing behavior in **Bad avoiders** ( $n = 22$ ) across training sessions for both  $CS^+$  and  $CS^-$  at door-closed trials (two-way repeated measures ANOVA; group:  $F_{(1,210)} = 30.80, p < 0.0001$ , training session:  $F_{(5,210)} = 1.96, p = 0.08$ , group x training session:  $F_{(5,210)} = 0.76, p = 0.57$ ). **f.** Averaged freezing behavior in **Bad avoiders** across training before and after door opening (8.8s pre-DO and post-DO) (two-way repeated measures ANOVA; group:  $F_{(1,210)} = 2.05, p = 0.15$ , training session:  $F_{(5,210)} = 3.68, p = 0.003$ , group x training session:  $F_{(5,210)} = 0.34, p = 0.88$ ). **g.** Averaged freezing behavior in **Good avoiders** ( $n = 13$ ) across training sessions for both  $CS^+$  and  $CS^-$  at door-closed trials (two-way repeated measures ANOVA; group:  $F_{(1,120)} = 17.82, p = 0.0003$ , training session:  $F_{(5,120)} = 1.53, p = 0.18$ , group x training session:  $F_{(5,120)} = 1.04, p = 0.39$ ). **h.** Averaged freezing behavior in **Good avoiders** across training before and after door opening (8.8 s pre-DO and post-DO) (two-way repeated measures ANOVA; group:  $F_{(1,120)} = 23.67, p < 0.0001$ , training session:  $F_{(5,120)} = 0.47, p = 0.79$ , group x training session:  $F_{(5,120)} = 0.37, p = 0.86$ ).



**Figure 6** A subset of Good avoiders generalized between CS<sup>-</sup> and CS<sup>+</sup> during avoidance

**a.** Avoidance discrimination index calculated as following  $((\text{Avoidance counts CS}^+) - (\text{avoidance counts CS}^-)) / ((\text{Avoidance counts CS}^+) + (\text{Avoidance counts CS}^- + 1))$  for **Good avoiders** discriminators (paired t. test:  $t = 2.81$ ,  $p = 0.015$ ) and generalizers in panel **b.** (paired t.test:  $t = 0.57$ ,  $p = 0.59$ ) at first and sixth training sessions. The dashed line at 20% represents the cut-off that we consider to classify mice as generalizers versus discriminators. **c.** Trial counts (shuttle CS<sup>-</sup>, avoid CS<sup>+</sup>, escape, errors) across 6 training sessions in generalizers ( $n = 5$ ) (two-way repeated measures ANOVA; group:  $F_{(3, 80)} = 17.90$ ,  $p < 0.0001$ , training session:  $F_{(5, 80)} = 2.30$ ,  $p = 0.052$ , group x training session:  $F_{(15, 80)} = 6.96$ ,  $p < 0.0001$ ). **d.** Pie-chart representative of avoidance-based profiles for the 40 mice tested. **Bad avoiders** represent 55 % ( $n = 22$ ), **Good avoiders** discriminators 33 % ( $n = 13$ ) and generalizers 12 % ( $n = 5$ ). **e.** Averaged freezing behavior in generalizers ( $n = 5$ ) across training sessions for both CS<sup>+</sup> and CS<sup>-</sup> at DC trials. (Two-way repeated measures ANOVA; group:  $F_{(1, 40)} = 5.36$ ,  $p = 0.04$ , training session:  $F_{(5, 40)} = 1.24$ ,  $p = 0.30$ , group x training session:  $F_{(5, 40)} = 1.24$ ,  $p = 0.30$ ).

$_{40}) = 0.47, p = 0.78$ ). f. Averaged freezing behavior in generalizers across training before and after door opening (8.8 s pre-DO and post-DO) (two-way repeated measures ANOVA; group:  $F_{(1, 40)} = 6.05, p = 0.03$ , training session:  $F_{(5, 40)} = 2.80, p = 0.02$ , group x training session:  $F_{(5, 40)} = 0.40, p = 0.83$ ).


*In summary, we developed a novel behavioral paradigm allowing a mouse to acquire and perform discriminative passive (freezing) and active (avoidance) behavior to a single conditioned stimulus depending on contextual contingencies. The kinetics of acquisition of both behaviors were dissimilar; discriminative freezing being acquired very rapidly as compared to a progressive acquisition of discriminative avoidance. In terms of our behavioral paradigm, two categories of mice were identified:*

- Bad avoiders: acquired discriminative freezing very rapidly but did not acquire discriminative avoidance.*
- Good avoiders: acquired discriminative freezing early during training and discriminative avoidance progressively with training.*

## II. Forced Swim Test

Our behavioral paradigm resulted in Good and Bad avoidance learners. Although this may simply represent a phenotypic trait, we were concerned that a repetitive footshock experience in DC trials might promote learned helplessness behaviors, which could be manifested by a lack of avoidance responses, like in the Bad avoiders group. Therefore we conducted a standard test that would model learned helplessness: the forced swim test (FST) (Figure 7a.).

The results of this test illustrate the fact that good and bad avoiders show no difference of swimming/immobility strategy during the force swim test (Figure 7c.). They show the same amount of immobility and there is not correlation with their performance during the 6<sup>th</sup> session of the avoidance training (Figure 7d). Interestingly, If we take into account two animals that were excluded from the analysis (a generalizer, and an animal that received every shock without escaping), there will be a difference between the two groups (Figure 7b.). In fact the former spent the entire time swimming without stopping and the latter stayed immobile in the water and was rescued from drowning. These two animals are representing a very small percentage of our study, and the FST was a good indicator to exclude them.



**Figure 7 Bad and Good avoiders do not differ in the forced swim test**

**a.** Mice undergoing the avoidance behavioral protocol for 8 days (see Figure 4) were classified into **Good** and **Bad avoiders** and were exposed one day later to the forced swim test (FST) during 6 minutes. **b.** Time spent immobile during the 4 last minutes of the FST for **Good** and **Bad avoiders**. Filled red circle represent a generalizer and empty red circle represent a mouse with a learned helplessness profile which spent all the 4 minutes immobile that was about to drown at the end of the session (unpaired t. test:  $t = 2.19$ ,  $p = 0.04$ ). **c.** Time spent immobile during the 4 last minutes of the FST for **Good** and **Bad avoiders** excluding the learned helplessness profile mouse (unpaired t. test:  $t = 1.47$ ,  $p = 0.16$ ). **d.** Correlation between the number of avoidance to the CS<sup>+</sup> and the time spent immobile during the 4 last minutes of the FST (Spearman correlation  $r = -0.34$ ,  $p = 0.18$ ). Filled circles concern the **Good avoiders** and empty circles concern the **Bad avoiders**. The horizontal dashed lines represent the lower and upper limits of immobility time range of a control group of mice exposed to the same FST protocol (Kara, et al., 2014; Kara, et al., 2016). The vertical dashed red line represents the threshold separating **Bad** and **Good avoiders**.

*The disparity of performance of the animals is not reflected in the forced swim test. The fact that repeated shocks during training did not make them change strategies of coping during the forced swim test, can exclude the fact that Bad avoiders present learned helplessness.*

*So far the disparity in the behavioral profiles observed (Good versus Bad learners) is an interesting phenomenon as it allows investigating the underlying neuronal mechanisms and to perform loss and gain of function optogenetic experiments.*

### III. C-fos Immunoreactivity

After determining the behavioral dynamic of our paradigm, we studied the fos pattern of different regions of interest right after the end of the training sessions. Our interest in this analysis was to see the difference of Fos staining pattern between the bad and the good avoiders. We modified the protocol by separating the animals in four groups, and testing them only to one specific trial contingency (see **Forced-swim test**).

The results showed an upregulation of the fos protein in the dmPFC in the good avoiders group (**Table 1, Figure 8a., Figure 9top panel**). This group was different from the Bad avoiders and the controls. When looking at the activation pattern of the vmPFC, the results showed no difference between the groups (**Table 1, Figure 8b.**). The Good avoiders also showed a difference with the control groups in the dlPAGc (**Table 1., Figure 8d., Figure 9 bottom Panel**). However the group was not different from the Bad avoiders. Other structures like the the dm,l, vIPAG and the BLA showed no differences in fos staining between the groups (**Table 1, Figure 8c, e, f , g, h**).

Structures	Atlas level	Groups				F(3,18) ; p
		Home Cage (n=6)	CS- (n=6)	Bad Avoiders (n=5)	Good Avoiders (n=5)	
dmPFC		539,95 ± 49,41	576,7 ± 56,04	353,03 ± 53,41	913,28 ± 127,11 °,*,**	8,98 ; 0,0007
vmPFC	1,94	294,35 ± 14,69	372,59 ± 6,51	116,56 ± 9,46	439,34 ± 28,27	2,37 ; 0,1043
BLA ant	-1,2	269,85 ± 38,57	278,48 ± 58,14	345,23 ± 64,74	391,21 ± 97,05	2,41 ; 0,1005
BLA post	-1,82	122,71 ± 43,5	142,07 ± 41,9	153,73 ± 42,62	208,06 ± 36,04	2,83 ; 0,0675
BA post		144,54 ± 47,55	157,64 ± 43,76	194,68 ± 38,15	225,99 ± 85,77	2,69 ; 0,0771
LA post		94,64 ± 25,43	124,77 ± 22,1	85,93 ± 47,29	181,56 ± 60,51	2,48 ; 0,0938
PAGdmr	-4,48	171,29 ± 33,11	153,23 ± 39,19	244,17 ± 38,55	285,49 ± 35,47	2,47 ; 0,0948
PAGdlr		102,43 ± 21,38	76,95 ± 13,94	93,71 ± 22,33	110,78 ± 12,56	0,65 ; 0,5934
PAGlr		314,34 ± 57,58	231,19 ± 27,28	286,33 ± 60,07	290,73 ± 42,93	0,58 ; 0,6384
PAGvlr		245,8 ± 67,75	196,19 ± 78,52	196,75 ± 50,51	234,01 ± 173,35	0,39 ; 0,7649
PAGdmc	-4,84	56,81 ± 7,1	64,43 ± 15,88	129,59 ± 33,08	138,6 ± 40,64	2,8 ; 0,0698
PAGdlc		48,34 ± 17,74	25,45 ± 15,34	49,16 ± 21,59	101,88 ± 31,14 °	3,92 ; 0,0258
PAGlc		253,34 ± 54,01	244,58 ± 41,23	220,87 ± 48,18	338,34 ± 46,63	1,05 ; 0,396
PAGvlc		149,12 ± 37,16	127,39 ± 21,5	123,61 ± 22,64	203,71 ± 35,81	1,38 ; 0,2808


Bonferroni/Dunn Post-Hoc test: Comparisons of all groups to Good Avoiders group are not significant unless the corresponding p-value is less than 0,0083.

° Different from Home Cage (HC)


\* Different from CS-

\*\* Different from Bad Avoiders

**Table 1 C-fos immunostaining in Bad and Good Avoiders.** C-fos immunoreactivity cell counts/mm<sup>2</sup> in the prefrontal cortex, amygdala and PAG of Bad and Good avoiders following 6 avoidance training sessions. dmPFC includes ACC and PL, vmPFC includes IL. **Ant**: anterior, **post**: posterior, **r**: rostral and **c**: caudal.



**Figure 8 C-fos is expressed in the dmPFC and dIPAG of Good avoiders.** Quantification of c-fos expression in home cage controls (HC), mice exposed to CS<sup>-</sup>, **Bad** and **Good avoiders** exposed to CS<sup>+</sup> in the dmPFC (a), vmPFC (b), caudal dmPAG (dmPAGc) (c), caudal dIPAG (dIPAGc) (d), caudal IPAG (IPAGc) (e), caudal vIPAG (vIPAGc) (f), BLA ant (g) and BLA post (h).



**Figure 9 C-fos immunostaining in Bad and Good avoiders.** Representative examples of c-fos staining in the prefrontal cortex (top panels) and the PAG (bottom panels) of a **Bad avoider** (left column) compared to a **Good avoider** (right column)

*The present Fos study, established on our behavioral paradigm provided important information about subregional activations at the level of two structures. the dmPFC and the PAG. As expected, it suggests that it is rather the dmPFC, (and not the vmPFC) and the dlPAG which are activated during avoidance learning. Our data describe that the lack of learning in Bad avoiders is indeed correlated to a lower activation of the dmPFC. These data strengthen the optogenetic results on the activation of the dmPFC to dlPAG pathway promoting avoidance in Bad avoiders.*

#### IV. Optogenetic manipulation


After validating the differences of fos activation between the Good and the Bad avoiders in the dIPAG and the dmPFC, we wanted to inhibit the dmPFC to dIPAG projections in order to understand its influence in the acquisition and expression of avoidance learning.

As we had previously optically activated the dmPFC neurons projecting to the dIPAG showing an improvement in the Bad avoiders performances after the training sessions, we wanted to see whether the inhibition of the pathway would impair the avoidance performances of the good avoiders, and make them bad avoiders.

According to our results, the optogenetic stimulation of the good avoiders the day after the 6<sup>th</sup> session had no effect on the avoidance level of the animals, and there was no difference with the GFP control group (Figure 10b.). This result let us think that inhibiting the dmPFC-dIPAG pathway did not affect animals that had already learnt the task. Regarding the latency to avoid, the good avoiders did not show differences in their latency to avoid after the stimulation (Figure 10d.). The discrimination between the CS+ and the CS- was also not impaired (Figure 10e.).


Interestingly, the inhibition of the pathway did not induce differences with the control groups in terms of freezing expression during the DC trials (Figure 10a, b.). Furthermore, in neither the shock nor the avoidance trials the animals reduced or increased their freezing level (Figure 10c, d.).

*To conclude, the inhibition of the dmPFC-dIPAG pathway does not alter the performance of the good learners after they have learned the task. These results could indicate that the dmPFC to dIPAG pathway is not involved in the expression of active avoidance.*



**Figure 10 Optogenetic inhibition of the dmPFC-dl/IPAG pathway does not impair avoidance expression**

**a. Left panels:** Photomicrographs of the fiber implantation and infection of the dmPFC to dlPAG pyramidal neurons in the PL with a (left) GFP and (right) ArchT cre dependent virus; **right panel**: photomicrograph of the retrograde cav-cre virus in the dl/IPAG **b.** CS<sup>+</sup> avoid counts at pre-stimulation, second stimulation and second post-stimulation sessions in two groups of **Good avoiders** infected with ArchT or GFP. (Two-way repeated measures ANOVA; group:  $F_{(1, 22)} = 2.50$ ,  $p = 0.14$ , training session:  $F_{(2, 22)} = 0.74$ ,  $p = 0.48$ , group x training session:  $F_{(2, 22)} = 0.28$ ,  $p = 0.75$ ). **c.** CS<sup>-</sup> shuttle counts at pre-stimulation, second stimulation and second post-stimulation sessions in two groups of **Good avoiders**: ArchT and GFP groups. (Two-way repeated measures ANOVA; group:  $F_{(1, 22)} = 3.01$ ,  $p = 0.11$ , training session:  $F_{(2, 22)} = 1.56$ ,  $p = 0.23$ , group x training session:  $F_{(2, 22)} = 1.08$ ,  $p = 0.35$ ). **c.** Mean avoidance latency to CS<sup>+</sup> trials in two groups of **Good avoiders** expressing ArchT or GFP during the second stimulation session, the pre-stimulation and the post-stimulation sessions (two-way repeated measures ANOVA; group:  $F_{(1, 33)} = 1.13$ ,  $p = 0.29$ , training session:  $F_{(2, 33)} = 0.07$ ,  $p = 0.92$ , group x training session:  $F_{(2, 33)} = 0.24$ ,  $p = 0.78$ ). **d.** Avoidance discrimination index in ArchT **Good avoiders** mice during the pre-stimulation session and the second stimulation session (paired t-test:  $t = 0.63$ ,  $p = 0.55$ ).




**Figure 11 Optogenetic inhibition of the dmPFC-dl/IPAG pathway does not impair freezing expression**

CS<sup>+</sup>-evoked freezing (across 15 trials) **(a)**, and CS<sup>-</sup>-evoked freezing (across 15 trials) **(b)** at pre-stimulation, second stimulation and second post-stimulation sessions in two groups of good avoiders: ArchT and GFP groups (panel **a**: two-way repeated measures ANOVA; group:  $F_{(1, 22)} = 0.10$ ,  $p = 0.75$ , training session:  $F_{(2, 22)} = 1.66$ ,  $p = 0.21$ , group x training session:  $F_{(2, 22)} = 4.12$ ,  $p = 0.03$ , and panel **b**: two-way repeated measures ANOVA; group:  $F_{(1, 26)} = 0.47$ ,  $p = 0.50$ , training session:  $F_{(2, 26)} = 0.91$ ,  $p = 0.41$ , group x training session:  $F_{(2, 26)} = 0.35$ ,  $p = 0.70$ ). CS<sup>+</sup>-evoked freezing during (i) the interval between the door opening and the shock delivery (shock trials) **(c)** and (ii) the interval between the door opening and avoidance response (avoid trials) **(d)** at pre-stimulation, second stimulation and second post-stimulation sessions in ArchT and GFP control mice (panel **c**, two-way repeated measures ANOVA; group:  $F_{(1, 32)} = 2.15$ ,  $p = 0.15$ , training session:  $F_{(2, 32)} = 2.07$ ,  $p = 0.14$ , group x training session:  $F_{(2, 32)} = 4.24$ ,  $p = 0.02$ , and panel **d**: two-way repeated measures ANOVA; group:  $F_{(1, 33)} = 3.37$ ,  $p = 0.07$ , training session:  $F_{(2, 33)} = 0.39$ ,  $p = 0.67$ , group x training session:  $F_{(2, 33)} = 0.04$ ,  $p = 0.95$ ). e. CS<sup>+</sup>-evoked freezing during the interval between the DO and the shock delivery (post-DO) and the same interval of time (namely 8.8 seconds) preceding the door opening (pre-DO) at pre-stimulation, second stimulation and second post-stimulation sessions in **Good avoiders** (two-way repeated measures ANOVA; group:  $F_{(1, 28)} = 0.001$ ,  $p = 0.96$ , training session:  $F_{(2, 28)} = 0.05$ ,  $p = 0.94$ , group x training session:  $F_{(2, 28)} = 0.34$ ,  $p = 0.70$ ).

Our next step was to focus our interest on the influence of the inhibition of the dmPFC-dIPAG pathway at early learning stage. As inhibiting the pathway had no effect on avoidance after training, it ruled out the role of this pathway in the expression of active avoidance, however, an inhibition during training will give us more information about its involvement in learning. We then inhibited the pathway in DO condition during the six days of training. To do so we simplified the paradigm by exposing the animals to the opened door condition only (DO+, DO-), as this classical active avoidance paradigm is known to generate a bigger proportion of good avoiders (Figure 12a.).

When inhibiting the pathway at early learning stage, the stimulated group was not able to avoid during the CS+ at the end of the 6<sup>th</sup> day, whereas the control group reached significantly higher avoidance rates (Figure 12c, d.). Interestingly, when left with no stimulation during six more days of training, the ArchT group stayed with bad performances, incapable to learn the task, while the control group kept improving (Figure 12c, d.). Regarding the freezing expression level, there is no significative difference between the stimulated sessions and the non-stimulated sessions (Figure 12e, f. ). However the ArchT group displayed significantly more freezing than the control group. This result cannot be explained by the stimulation but more likely by the fact that the ArchT group avoided less, ergo froze more, as shown with the bad learners (Figure 5).

*In conclusion, the inhibition of the dmPFC to dIPAG pathway at early training stage abolished avoidance learning but did not affect conditioned freezing behavior.*



**Figure 12 Inhibition of the dmPFC-dl/IPAG pathway abolished avoidance learning**

**a.** Adapted behavioral protocol. On Day 1, mice were habituated during 15 min to the shuttle-box. On Day 2, animals were habituated to the presentation of the CS<sup>-</sup> and CS<sup>+</sup> during opened condition (DO) only. After 23.1 seconds following the sound's onset, the door was slid-down (DO) and slid-up again 8.8 seconds after. 9 trials of CS<sup>-</sup> and CS<sup>+</sup> were played. From Day 3 to Day 8, animals underwent 6 training during which the same type of trials were played except that the number of trials was increased to 15 CS<sup>+</sup> followed by a 4 s shock if the animal did not escape or avoid, and the yellow laser was turned on continuously for 9 seconds following door opening. From day 9 till 14 animals underwent the same training sessions except that no laser stimulation was delivered. **b.** Avoidance-based profiling after 6 sessions of training with no laser stimulation (non-stimulated group). **c.** CS<sup>+</sup> avoidance counts (two-way repeated measures ANOVA; group:  $F_{(1, 209)} = 10.06$ ,  $p = 0.005$ , training session:  $F_{(11, 209)} = 7.62$ ,  $p < 0.0001$ , group x training session:  $F_{(11, 209)} = 2.42$ ,  $p = 0.007$ ). **d.** CS<sup>-</sup> shuttles counts (two-way repeated measures ANOVA; group:  $F_{(1, 209)} = 7.72$ ,  $p = 0.01$ , training session:  $F_{(11, 209)} = 3.91$ ,  $p < 0.0001$ , group x training session:  $F_{(11, 209)} = 1.38$ ,  $p = 0.18$ ). **e, f.** Averaged freezing behavior during pre-door opening CS<sup>+</sup> trials (**e**: two-way repeated measures ANOVA; group:  $F_{(1, 209)} = 4.19$ ,  $p = 0.05$ , training session:  $F_{(11, 209)} = 4.27$ ,  $p < 0.0001$ , group x training session:  $F_{(11, 209)} = 0.61$ ,  $p = 0.81$ ), and CS<sup>-</sup> trials (**f**: two-way repeated measures ANOVA; group:  $F_{(1, 209)} = 0.35$ ,  $p = 0.55$ , training session:  $F_{(11, 209)} = 2.72$ ,  $p = 0.002$ , group x training session:  $F_{(11, 209)} = 1.22$ ,  $p = 0.27$ ).

*In summary, the optogenetic inhibition of dmPFC-dl/IPAG does not impair avoidance behavior in already good avoiders, but abolished the capacity to learn the task when the animals are repeatedly inhibited at early stage.*

*Altogether with the fos data, and the data on the optogenetic activation of the pathway in bad avoiders promoting active avoidance, we can propose that the dmPFC to dIPAG pathway promotes active avoidance but does not seem to be involved in its expression, nor in freezing. It is also supported with the electrophysiological data, showing a specific activation of the dmPFC projecting cells during avoidance.*

## II/ Hypothalamic pathways of shock-based Passive avoidance

### Materials and methods

#### I. Animals


*FOS expression experiments:* Male C57BL6/J mice (3 months old) weighing 30-35 g were used in these experiments. They were housed the same way as previously described for the SigAA experiments.

*Pharmacogenetic, optogenetic and electrophysiological experiments:* CCK-Cre transgenic male mice (3-month-old, Jackson Laboratories) were used in these experiments. These animals express the Cre protein under control of the cholecystokinin promoter (CCK). Interestingly, this animal is ideal for manipulating the PMD, since this nucleus differs from neighboring structures by the abundant expression of CCK. They were housed the same way as above. We aimed at inactivating the PMD using pharmacogenetic tools. The PMD is a very small hypothalamic site, and the use of conventional forms of pharmacological inhibition would spread to other neighboring areas, rendering very difficult to ascertain the specific roles of the PMD in passive contextual avoidance. To circumscribe, as much as possible, the inactivation to the PMD, we took advantage to the fact that PMD cells present a characteristic expression of CCK (cholecystokinin) peptides, differing from the neighboring structures.

#### II. Experimental protocols

##### A/ Compared conditions Paradigm

The experimental apparatus consists of a safe cage (25x25x25 cm) with a door, connected to a 30cm corridor with access to a second cage (conditioning cage), with the same dimensions as the safe cage. The whole experimental apparatus is made of acrylic. The conditioning cage has a floor with a grid composed of steel bars spaced 7 mm and connected to a current generator (Insight).



**Figure 13** Behavioral protocol for the study of Fos expression in the contextual non-instrumental avoidance.

For three days, each animal was habituated to this experimental apparatus, where the animals were placed in the safe cage, allowing the animal to explore the entire experimental apparatus for 10 minutes. On the 4<sup>th</sup> day, the animals were placed in the apparatus, and when they entered the conditioning cage the door was shut, after two minutes of habituation they received five footshocks (0.6mA, for 1 second, following a random triggering pattern) and stayed in this box for an additional 2 minutes. The control groups had exactly the same treatment without receiving footshocks (not conditioned). The next day, four groups of animals were tested, namely:


Group 1 (**Free+**) : Animals that received shocks on day 4 were placed in the safe cage (Figure 13 panel “whole apparatus”) with an access to the whole apparatus, including the conditioning cage and were observed for 8 minutes.

Group 2 (**Confined+**): Animals that received shocks on day 4 were confined in the conditioning cage for 8 minutes (Figure 13 panel “confined”).

Group 3 (**Free-**) : unconditioned naive animals that were placed in the safe cage with access to the conditioning cage, and were observed for 8 minutes (Figure 13 panel “whole apparatus”).

Group 4 (**Confined-**): unconditioned naive animals that were confined in the conditioning cage for 8 minutes (Figure 13 panel “confined”).

#### B/ Passive avoidance Paradigm: PMD study



**Figure 14** Behavioral protocol for the study of the role of the PMD in the contextual non-instrumental avoidance.

For the experiments of the PMD modulations, the animals were conditioned the same way as in the other paradigm (see “*compared conditions paradigm*”). On **day 5**, the animals were placed in the safe cage with access to the conditioning cage and were

observed for 8 minutes. On **Day 6**, the different groups were exposed again to the same context for 8 minutes.

Pharmacogenetic experiments: on **day 5** the animals injected with an inhibitory DREADD virus (HM4D (Gi)) were administered a dose of 300µl of CNO (Clozapine *N*-oxide) solution, 30min before the context exposure. To measure the putative interoceptive stimulus effect of clozapine, the control group was also administered CNO but were injected with a non DREADD GFP virus (. On **day 6**, none of the groups received treatment. (Figure 19)

Optogenetic experiments: on day 5, both groups (Halorodopsin and GFP) were continuously stimulated with yellow light (wavelength of 589nm) for 8 minutes, when entering the conditioning apparatus. On day 6, the animals were not stimulated with light (


Figure 24,

Figure 26).

Generalisation experiment: The animals encountered the same procedure as in the “*passive avoidance paradigm*” but were exposed prior to the conditioning apparatus, to a neutral open field (60cmx40cmx20cm) for 5 minutes. They were exposed during the three days of habituation, and the two days after the conditioning day (Figure 22).

### III. Behavioral analysis

Animals were filmed and data analyzed later by a trained observer using the ethological analysis software The Observer (Noldus). The counted behaviors were the ones below (see Figure 15) The behavioral analysis was done during the last day of habituation, the conditioning phase and the next day in the test of contextual responses. The analysis involved spatiotemporal (for contextual responses) and behavioral (for conditioning and contextual responses) measurements. Spatiotemporal measurements are related to the time (in seconds) the animal spent in the safe cage, corridor, or conditioning box. Behavioral data were processed in terms of duration (8 min per session).



**Figure 15 Behavioral features analysed during experiments** The following behavioral responses were categorized: "**Freezing**" - the animal remained completely immobile in a freezing state except for the breathing movements; "**Crouch sniff**" - the animal remained still, with arched back, making movements with the head smelling and scanning the environment; "**Stretch Attend Posture**" - the animal extended the head and part of the body forward, kept the tail elevated, but did move; "**Stretch Approach**" - the animal retained the same anterior posture, but moved forward; "**Up right position**" - the animal was standing with its hind and front legs extended and supported on the walls of the apparatus; "**Locomotion**" - the animal moved more than 1cm with arched back;

#### IV. Histological processing of Fos immunohistochemistry

90 minutes following the exposure to the aversive context, the animals were anesthetized and perfused following the same technique as the first part. (see *Histological processing of Fos immunohistochemistry Part I*).

#### V. surgeries and optogenetics

*Viral injection surgery:* Following the same procedures described in the SigAA paradigm, we made bilateral injection in the PMD (coordinates: -2.43 mm antero posterior,  $\pm$  0.3 medio lateral, -5.4 ventral to the dura), where cre-dependent virus to express DREADD was injected in CCK-IRES-CRE mice.

*Optical fiber implantation surgery:* Three weeks after viral infection, using the same surgical and stereotaxic procedures previously described, cranial holes were drilled at the following stereotaxic coordinates  $\pm$  2.3 mm anterior-posterior from the bregma and  $\pm$  0.3 mm medial-lateral. For the inhibition of PMD projections, bilateral optic fibers were

implanted either in the dIPAG (-4.2 mm anterior-posterior from the bregma and + 0.5 (8°angle) mm medial-lateral, -1.35 mm ventral to the dura-mater); or in the Anterior thalamus (-0.7 mm anterior-posterior from the bregma and +/- 1.4 (10°angle) mm medial-lateral, -3.35 mm ventral to the dura-mater). After surgery, mice were allowed to recover for at least 7 days. They were handled daily during the three days preceding the behavioral tests to habituate them to the connection procedure.

*Pharmacogenetics inhibition:* To circumscribe the DREADD expression in the PMD we used CCK-Cre transgenic male mice (Jackson Laboratories), which express the Cre protein under control of the cholecystokinin promoter (CCK), and applied CRE-dependent virus for the expression of HM4D (Gi) in the PMD to be responsive to the inactivation with CNO.

*Optical stimulation (see material and method part I):* Terminals were inhibited by a continuous yellow light at 593 nm using a laser (DPSSL lasers) that was connected to a 200 µm diameter optic fibre patch cable (Thorlab) and calibrated to produce a fibre tip irradiance of approximately 16-18 mW. The animals were illuminated by yellow light while entering the apparatus to immediately block any possible contextual cue gathering.


## VI. Statistical analysis

(see Statistical analysis in part I)

## Results

### I. Behavior analysis of the non-instrumental contextual avoidance paradigm.

In this part of the project we developed a paradigm where the animals, after being conditioned to a shock would be placed, depending on the group, in different conditions for fear retrieval. As shown in **Figure 16b**, the **Free-** group explored the entire apparatus and spent  $217 \pm 23.1$  s in the footshock chamber. In contrast, the **Free+** group did not enter the footshock chamber and stayed  $467 \pm 22.9$  s in the safe cage and  $28 \pm 4.9$  s in the corridor (**Figure 16b**). Looking at the behaviors, the **Free+** group (**Figure 16a**) showed a small amount of freezing but showed a lot of risk assessment (i.e., “crouch-sniff” and “stretch postures”) (**Figure 16a**). Moreover, during the test period, the **Confined-** group explored fearlessly the conditioning chamber (**Figure 16a**). The **Confined+** group, on the contrary, expressed freezing behavior and risk assessment during the exposure to the shock chamber (**Figure 16a, c**).




**Figure 16.** Behavior expression strategies differ depending on the fear recall exposure conditions. **a.** Pie charts of the behaviors counted during context exposure of: **Confined+** (top left panel), **Free+** (top right panel), **Confined-** (bottom left panel) and **Free-** (bottom right panel) groups. **b.** Comparison of the spatiotemporal measurements between **Free+** and **Free-** there is a difference in group\*spatiotemporal distribution interaction: two-way ANOVA  $F_{(1,8)}=48.423$   $p>0.0001$ . **c.** Comparison of freezing expression level among the groups: ANOVA  $F_{(3,16)}=13.909$   $p<0.0001$ ; Fisher PLSD post hoc: \*\*\* $p<0.001$  different from all groups.


*These results indicate that specific behaviors are expressed during fear retrieval according to different exposure situations. Even though the animals have been conditioned the same way, the one in a free condition will express more risk assessment behaviors, and will not enter the conditioning cage, whereas the animals in a closed situation will express more freezing.*

## II. Analysis of Fos expression in the non-instrumental contextual avoidance paradigm.

After understanding the behavioral pattern of the defense responses toward different exposure conditions, we analysed the Fos upregulation of the structures of our interest. The selected areas for evaluation of the Fos protein expression were based on a previous study (Viellard et al. 2016), where during passive contextual avoidance we showed, in rats, the mobilization of the circuit formed by the ventral Subiculum (SUBv), lateral septum (LS), juxta-dorsomedial part of the lateral hypothalamus (LHAjd) and the dorsal preammillary (PMD). In addition, we investigated the expression of Fos in the dorsomedial, dorsolateral, lateral and ventrolateral sectors of the PAG. Our results show an upregulation of fos expression for the **Free+** group in the SUBv, the LS, the LHAjd, the PMD, the dmPAG, and the lPAG different from all of the other groups (Figure 17a, b, Figure 18). The Fos expression of the vIPAG is different between the **Free+** and the **Confined+** group but the **Free+** group is not different from its control (**Free-**). The **Free+** group also differs in Fos expression in the dIPAG from both control groups (**Free-**, **Confined-**), but not from the **Confined+** group. (Figure 17b.)



**Figure 17** Density of Fos-labeled cells in selected brain regions during exposure to the context. The structures selected showed an up-regulation in Fos expression in the **Free+** group compared to the **Confined+** group and controls. **a.** The **Free+** group showed Fos upregulation in septohippocampal structures: **LS**: Lateral Septum, **SUBv**: ventral part of the Subiculum; and in hypothalamic structures: **LHAJD**: lateral hypothalamic area juxtadorsomedial region, **PMD**: dorsal premammillary nucleus. **b.** The **Free+** group showed Fos upregulation in brainstem structures **PAGdm**: periaqueductal gray, dorsomedial part; **PAGdl**: periaqueductal gray, dorsolateral part; **PAGl**: periaqueductal gray, lateral part; **PAGvl**: periaqueductal gray, ventrolateral part. Bonferroni Post Hoc test, after a Group effect on the ANOVA ( $p < 0.0083$ ). When differs significantly from conditioned groups, \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ ). When differs significantly from Control groups, ° $p < 0.05$ ; °° $p < 0.01$ ; °°° $p < 0.001$ .




**Figure 18** Photomicrographs of frontal Fos-stained sections illustrating the distribution of Fos-labeled cells in the LHAjd (a), PMD (b), PAG (c) and SUBv (d), comparing the **Free+** (left column) and the **Confined+** (right column) groups. Abbreviations: **PaDC** : paraventricular hypothalamic nucleus, dorsal cap; **mfb** : medial forebrain bundle; **opt** : olivary pretectal nucleus; **LH**: lateral hypothalamic area; **Gem**: gemini hypothalamic nucleus; **PH**: posterior hypothalamic area; **SuMM**: supramammillary nucleus, medial part; **PMD**: premammillary nucleus, dorsal part; **PMV**: premammillary nucleus, ventral part; **DRD**: dorsal raphe nucleus, dorsal part; **PaS**: parasubiculum ; **Ect**: ectorhinal cortex; **PRh**: perirhinal cortex; **LEnt**: lateral entorhinal cortex **3V**: third ventricle; **f** : fornix; **LHAjd**–lateral hypothalamic area juxtadorsomedial region; **dIPAG** – periaqueductal gray, dorsolateral part; **dmPAG** - periaqueductal gray, dorsomedial part; **lPAG** periaqueductal gray, lateral part; **vlPAG** – periaqueductal gray, ventrolateral part; **PMD** - dorsal premammillary nucleus; **S**: subiculum.


*The overall data illustrate the idea that the fear conditioned animals which were able to avoid the conditioning chamber presented increased Fos expression in the circuit formed by the ventral subiculum, the juxtadorsomedial lateral hypothalamic area, the dorsal premammillary nucleus and the lateral and dorsal parts of the periaqueductal gray. Anatomical and functional data suggest that this septo/hippocampal-hypothalamic-brainstem circuit should be putatively involved in mediating contextual avoidance.*

### III. Roles of the PMD on the non-instrumental contextual avoidance: Pharmacogenetic manipulation

After validating the recruitment of a specific pathway in our passive avoidance paradigm, we targeted the PMD, a nucleus of the pathway that, according to our hypothesis, is a key structure of the circuit. That is why we aimed at inactivating the PMD using pharmacogenetic tools. To do so we applied a CRE-dependent inhibitory DREADD virus (HM4D (Gi)) in the PMD in order to inactivate temporarily the structure with CNO during context exposure (Figure 19).



**Figure 19** The inhibition of the PMD has an effect on the expression and the reconsolidation of passive avoidance **a**. Injection sites and viral strategy. Injection of a cre dependent virus expressing the DREADD inhibitory promoter. **b**. Photograph of the fluorescence of the DREADD infected cells of the PMD. The virus will recombine in the CCK positive cells of the PMD that express cre-recombinase in CCK-CRE-IRES mouse line. **c**. Behavioral protocol to test the PMD's role on the passive contextual avoidance. On **day 5** both groups (DREADD and GFP) were injected with CNO (300  $\mu$ L intraperitoneal) 30 minutes before exposure to the context. On **day 6**, animals were placed again in the apparatus, without pharmacological treatment.




**Figure 20** The inhibition of the PMD has an effect on the expression and the reconsolidation of passive avoidance. Behavioral and spatiotemporal analysis in the contextual avoidance apparatus during habituation (**a**), first exposure to context (**b**) and re-exposure (**c**). **a.** No behavior difference between the groups during **habituation**: group\*behavior (two-way ANOVA  $F_{(1,6)}=0.864$   $p = 0.5259$ ); and no time difference of time spent in the conditioning cage (unpaired t-test  $p=0.263$ ). **b.** Behavior\*group interaction difference during **context exposure** (two-way ANOVA  $F_{(1,6)}=12,406$   $p < 0.001$ ); group effect on **exploration**  $p < 0.01$ , **crouch sniff**  $p < 0.05$ , and **stretch postures**  $p < 0.05$ ; and the **conditioning cage** (t-test  $p=0.0168$ ). **c.** During **context re-exposure** there is also a difference between groups for the behavioral measurements (two-way ANOVA  $F_{(1,6)}= 16.557$   $p < 0.001$ ); group effect on **exploration**, **crouch sniff** and **stretch postures**  $p < 0.05$ ). The DREADD animals entered the **conditioning cage** (t-test  $p=0.0369$ ).

On the last day of habituation (day 3), the behavior analysis show no difference between the DREADD and control group, as well as the time spent in the conditioning cage (Figure 20a.). On the context exposure day, there are significant differences between the DREADD and control groups in the behavioral expression and the time spent in the conditioning cage (Figure 20b.). In fact, the DREADD group drastically decreased risk assessment behaviors, like crouch sniffing and stretch postures, they also entered the conditioning cage, showing a loss of fearful recall of the conditioning day. Interestingly, during context re-exposure, we also found significant differences between the groups for the behavioral measurements, as well as the spatiotemporal measurements (Figure 20c.). Again, there was a significant decrease in risk assessment behaviors like stretch postures and crouch sniffing (Figure 20c. **left panel**). Yet again, while the control group wouldn't enter the conditioning cage, the DREADD group entered fearlessly the space, suggesting a impairment in memory reconsolidation of the context cues (Figure 20c. **right panel**).

#### IV. Behavioral and virus controls

##### Virus control

As described earlier in the ("Experimental protocols" section), all cre dependent viruses that we injected during our experiments infected CCK positive cells in CCK-IRES-CRE transgenic animals. In the posterior hypothalamus, these CCK positive cells are present exclusively in the PMD, and the rest of the mammillary bodies (ML, MM, MMn). Unfortunately, the injections targeting the PMD would also infect partially the mammillary bodies. According to the literature, cytotoxic lesions of the mammillary bodies as no effect on fear expression and recall during cat exposure in rats (Cezario et al., 2008). Acknowledging this information, we analysed the activation pattern of this region during passive avoidance, to make sure that the nuclei had no importance in our study as well.




**Figure 21** The mammillar nuclei present no activity during passive avoidance. **a.** cell density counting (cell/mm<sup>2</sup>) of fos immunostaining in the mammillar nuclei (ML, MM). There is no difference of Fos staining between the **Free+** and the **Free-** groups : t-test  $p=0.9647$ . **b.** Photomicrograph of a fos immunostaining of the mammillar region in a free+ animal (**left panel**), and its control (**right panel**). Abbreviations: : **MMn**: median mammillary nucleus, **ML**: medial mammillary nucleus, lateral part, **MM**: medial mammillary nucleus, medial part; **pm**: mammillary peduncle; **SuM**: supramammillary nucleus; **VTM**: ventral tuberomammillary nucleus.

When observing the Fos expression in the ML and the MM in the **Free+** and the **Free-** groups, there is only a very weak Fos-positive-cell density in the mammillar region (~32cell/mm<sup>2</sup>). Furthermore, the animals placed in free condition during context exposure (**Free+**) showed no difference of cell activation with the control (**Free-**) (Figure 21). These data comfort us in our hypothesis that the ML and MM are not implicated in passive avoidance.

#### Behavioral control

We then wanted to test whether the conditioning protocol would create generalisation. In order to do so we tested the risk of generalisation using a discrimination protocol. The animals would be presented to a neutral Open Field before entering the conditioning apparatus. We then observed the ratio of fearful versus fearless behaviors along the different sessions, in the two contexts (Figure 22a., see Experimental protocols).

Our results illustrate that the behavior of the animals during the Open field exposure is not altered after the conditioning day, as there is no difference of ratio between the three days (Figure 22b.). Regarding the conditioning apparatus exposure, as expected, the post conditioning ratio of fearful vs. fearless behaviors drastically increases (Figure 22c.). These data suggest that the animals discriminate between the two contexts, and that only the conditioning apparatus becomes aversive after the conditioning day.




**Figure 22** During Open field exposure, the fearful vs fearless behaviors ratio is not altered after the conditioning day, but the Conditioning apparatus exposure is. **a.** Contextual discrimination protocol. The animals were exposed to an open field 5 min before exposition to the conditioning apparatus during habituation days, exposure and re-exposure. **b.** Effect of the conditioning day on the ratio of fearful behaviors versus fearless behaviors, in the open field,  $n=7$ , ANOVA ( $F_{(2,18)}=0.263$   $p < 0.77$ ). **c.** Effect of the conditioning day on the ratio of fearful behaviors versus fearless behaviors, in the conditioning apparatus,  $n=7$ , ANOVA ( $F_{(2,18)}=28.069$   $p < 0.001$ ); Fisher'PLSD post hoc test (cont1 vs. cont2)  $p=0.1843$ ; (Hab vs. cont1)  $p<0.001$ ; (Hab vs. cont1)  $p<0.001$ .

*Overall, after showing that the viral infection of the mammillar bodies does not disturb the behavior response as it is not involved in passive avoidance, and that the test does not create generalisation, it can be proposed that inactivation of PMD affects contextual passive avoidance. It also greatly weakens the process of reconsolidation of the aversive context.*

## V. role of the PMD projections in both the expression and reconsolidation of passive avoidance


Our next goal was to understand the role the terminal fields of the PMD, which are the dlPAG and AMv (Figure 23). These two structures are densely targeted by the PMD, and using optogenetic terminal inhibition we are going to examine their potential roles in the expression and reconsolidation of contextual avoidance.



**Figure 23** Sagittal views of the PMD projections in the ventral part of the anterior thalamus and the dorsal part of the PAG. **left panel**, projections of virus infected cells in the PMD and Mammillary bodies. **Right panel**, green coloration representation of the PMD projections only to the PAG (left direction), and the AMv, (right direction).

### a. Optogenetic Manipulation on dIPAG terminals

We first inhibited the terminals of the PMD in the dIPAG using Halorodpsin virus inhibition during the context exposure (Figure 24).



**Figure 24 Optogenetic dPAG terminal inhibition strategy** **a.** Photomicrographs of the PMD's terminal fields in the dIPAG . **b.** scheme of the viral and optogenetic inhibition strategy. CCK-IRES-Cre transgenic male mouse that received CRE dependent virus for the expression of Halorodopsin and the reporter YFP (rAAV9/CAG-Flex-eNpHR-YFP) in the PMD. Optical stimulation of the PMD terminals in the dIPAG. **c.** Behavioral protocol of the role of the PMD to dIPAG projections in passive avoidance. on **day 5**, both groups (Halorodopsin and GFP) were continuously stimulated with yellow light (wavelength of 589nm) during 8 minutes, while entering the conditioning apparatus. On **day 6**, the animals were re exposed to the same context but were not optically stimulated.

On the third day of habituation, the two groups presented no differences of behaviors, nor spatiotemporal distribution (Figure 25a.). However, the day of exposure to the aversive context (**day 5**), the Halo group significantly reduced risk assessment behaviors expression compared to the control GFP group. In fact, the inhibited animals expressed more exploration behavior, and less crouch sniffing and stretch postures (Figure 25b.left panel). They also entered the conditioning cage, with an average of 50 sec during the 8min of exposure (Figure 25b.right panel), whereas the control group never entered the cage.





Figure 25 Behavioral and spatiotemporal analysis in the contextual avoidance apparatus during habituation, first exposure to context and re-exposure. **a. left panel** There is no difference during habituation behavior\*group interaction two-way ANOVA  $F_{(1,6)}=2.087$ ,  $p=0.0699$ . **right panel** There is no difference of time spent in the conditioning cage between the two groups. **b. left panel** the two groups show behavior differences during **context exposure**. Group\*behavior interaction two-way ANOVA  $F_{(1,6)}=19.141$ ,  $p<0.0001$ . T-test group difference in **exploration** ( $p=0.0002$ ); **crouch** ( $p=0.0034$ ); **stretch postures** ( $p=0.0052$ ). **right panel** the Halo group entered the conditioning cage, but not the GFP group. (t-test,  $p=0.0227$ ) **c. left panel** the two groups show behavior differences during **context re-exposure**. Group\*behavior interaction two-way ANOVA  $F_{(1,6)}=5.635$ ,  $p<0.0001$ . t-test group differences in **exploration** ( $p=0.0242$ ); **crouch** ( $p=0.0159$ ). **right panel** Halo group entered the conditioning cage, but not the GFP group (t-test  $p=0.0038$ ).

On the 6<sup>th</sup> day, the animals were exposed again in the same aversive context, they also decreased in risk assessment responses. They expressed more exploration, and less crouch than the control group (Figure 25c.**left panel**). However, the stretch postures expression is not different from the controls (Figure 25c.**left panel**). They also entered the conditioning cage on the second day, which was still not the case for the control group (Figure 25c.**right panel**).


## VI. Optogenetic Manipulation on Thalamic terminals

In a second part, we inhibited the terminals of the PMD in the AMv using Halorodopsin virus inhibition during the context exposure (

Figure 26).



**Figure 26 Optogenetic Thalamus terminal inhibition strategy.** **strategy a.** Photomicrographs of the PMD's terminal fields in the AMv. **b.** scheme of the viral and optogenetic inhibition strategy. CCK-IRES-Cre transgenic male mouse that received CRE dependent virus for the expression of Halorodopsin and the reporter YFP (rAAV9/CAG-Flex-eNpHR-YFP) in the PMD. Optical stimulation of the PMD terminals in the AMv. **c.** Behavioral protocol of the role of the PMD to AMv projections in passive avoidance. on **day 5**, both groups (Halorodopsin and GFP) were continuously stimulated with yellow light (wavelength of 589nm) during 8 minutes, while entering the conditioning apparatus. On **day 6**, the animals were re exposed to the same context but were not optically stimulated.



**Figure 27 Behavioral and spatiotemporal analysis in the contextual avoidance apparatus during habituation, first exposure to context and re-exposure. a. left panel** There is no difference during habituation behavior\*group interaction two-way ANOVA  $F_{(1,6)}=0.351$ ,  $p=0.9052$ . **right panel** There is no difference of time spent in the conditioning cage between the two groups. **b. left panel** the two groups show behavior differences during **context exposure**. Group\*behavior interaction two-way ANOVA  $F_{(1,6)}=7.525$ ,  $p<0.0001$ . t-test group difference in **exploration** ( $p=0.0121$ ); **crouch** ( $p=0.0391$ ); **upright** ( $p=0.0337$ ). **right panel** the Halo group entered the conditioning cage, but not the GFP group. (t-test,  $p=0.0643$ ) **c. left panel** the two groups show behavior differences during **context re-exposure**. Group\*behavior interaction two-way ANOVA  $F_{(1,6)}=14.681$ ,  $p<0.0001$ . t-test group differences in **exploration** ( $p=0.0039$ ); **crouch** ( $p=0.0003$ ); **upright** ( $p=0.0115$ ). **right panel** Halo group entered the conditioning cage, but not the GFP group (t-test  $p=0.0270$ ).

On the third day of habituation, the two groups present no differences of behaviors, nor spatiotemporal distribution (**Figure 27a.**). The results showed a difference in behaviors between the **Halo** group and its **control** during context exposure. The Halo group expressed more exploration and upright behaviors, and less crouch sniffing compared to the control (**Figure 27b. left panel**). Interestingly the amount of stretch postures expression does not differ with the controls. The Halo group also entered the conditioning cage, with an average of 50 seconds during the 8 minutes of exposure whereas the control group did not enter the compartment (**Figure 27b. right panel**). These results suggest that the inhibition of the AMv terminals partially impairs passive avoidance behaviors during context exposure.

On the re-exposure day, the Halo group also decreased in risk assessment responses compared to the control group (**Figure 27c. left panel**). They expressed more exploration and upright behaviors, and less crouch sniffing than the control group. However, the stretch postures expression is not different from the controls. When re-exposed to the aversive context, the Halo group also entered the conditioning cage, which was still not the case of the control group (**Figure 27c. right panel**). According to these data the yellow light stimulation also impacted the re-exposure day, suggesting an effect on memory reconsolidation.

*projections of the PMD to AMv and dIPAG are both involved in expression and reconsolidation of passive avoidance. However, even if the comparison with their controls shows a tendency of the dIPAG inhibition to disrupt more the expression of the behaviors, and the inhibition of the AMv has more effect on the re-exposure day which indicates a more important role in memory reconsolidation, the two groups do not differ between them (data not shown). Knowing the latter, no conclusion can be made about their specific implication differences in passive avoidance expression and reconsolidation.*

## Discussion

### First Part

**A novel behavioral paradigm to study avoidance and freezing:** In the first part of the project we focused our attention on the projection of the dmPFC to the dIPAG, and its role in avoidance and freezing acquisition and expression. To that extent we developed a novel behavioral paradigm of active avoidance during which a single CS was associated with two conditioned behavioral outcomes (freezing versus avoidance) depending on the state of the door in between compartments. Freezing behavior was evident to acquire for all the mice tested in our paradigm. All mice froze significantly more to the CS+ (between 40% to 60 % in DC condition) compared to the CS- and, discriminated between the two CSs. Interestingly, the freezing level evoked by the CS- was relatively high in all mice. The fact that mice cannot predict whether the door will open or stay close increases the attentional processes and promotes immobility. It could be an explanation of their high freezing level during CS-. A second potential explanation could be linked to the random trial presentation. Even though random presentation of different trial types makes the learning more complex, it also potentially enhances attentional processes and prevents the development of habitual avoidance learning (Dickinson, 1985; Wood and Neal, 2007). Our objective being to study goal-directed avoidance learning and not habitual avoidance we opted for presenting the trials in an intermingled manner.

Interestingly, the second behavioral outcome (avoidance) was not learned by all mice. Indeed, we categorized mice based on (i) avoidance scores and (ii) discrimination between CS- and CS+ trials. This categorization led to Bad avoiders (mice that did not learn to avoid), Good avoiders (mice that learned discriminative avoidance) and generalizers (Good avoiders that learned to shuttle/avoid to the other compartment each time the door got opened regardless of the CS). In terms of freezing, Bad avoiders, Good avoiders and generalizers also differ at two levels even though all three groups discriminate between CS- and CS+. During DC condition, Bad avoiders present the highest freezing levels to CS+ (mean~55-60%) followed by discriminators (mean~45-50%) and generalizers which exhibited very low freezing levels to CS+ presentations (mean~35-40%). Therefore, it seems that the DC condition allows to categorize animals in terms of

freezing levels. During DO trials, at door opening Bad avoiders continue to freeze at high levels post-DO whereas Good avoiders and generalizers present a drop in their freezing levels since they switch to an active defensive strategy.

This heterogeneity in acquiring active defensive strategies have been already reported in active avoidance studies (Galatzer-Levy et al., 2014) and is of a relative importance from a clinical point of view because it transduces the heterogeneity of response of humans facing traumatic events. Regarding the proportions of the different behavioral profiles, both the original paradigm and the simplified adapted paradigm (only DO condition) resulted in ~35% of Good avoiders discriminators which acquired discriminative avoidance behavior. For connected animals (optic fibers, electrodes cables), around 45 to 55% of the population were classified as Bad avoiders and the rest as Generalizers. In all the experiments, generalizers were not further considered.

Regarding the FST, the results showed that there was no difference between the two groups of animals. This test was important in this study to measure the impact of repetitive shocks on the animals. As the FST induces learned helplessness (Yankelevitch-Yahav et al., 2015), it is important to note that the bad avoiders did not change their strategy, in this test, even though failing to learn the avoidance task. However it does not give us insight on the background of the animals, and why such a great number of them cannot learn the task. Even though the two way avoidance shuttle box paradigm does not impact the emotional state of the bad avoiders, extreme behaviors (the non avoider and the generalizer) are however reflected in the FST, as the non-avoider almost drowned, and the generalizer did not stop swimming. We can then advance, that abnormal stress level in a small portion of the animals impacted their performance of the two-way SigA paradigm. Some studies also suggest that the FST reveal a certain heterogeneity in immobility levels after chronic and acute stress experiments ((Suvrathan et al., 2010). To validate this hypothesis, an interesting experiment would be to administer an anxiolytic drug before each SigA training day in order to see if the proportion of good and bad avoiders will be different. It can then also be interesting to study how afferent hypothalamic projections, like the PVH, known to be involved in stress (Xu et al., 2019), would indirectly impact the dmPFC to dIPAG pathway.

**The active avoidance learning preferentially activates the dmPFC and the cdlPAG:** The immediate-early gene c-fos study we performed revealed a clear significant upregulation of c-fos in Good avoiders as compared to Bad avoiders and controls in the caudal dmPFC (ACC, PL). Our results are in concordance with several studies in rodents using a platform-mediated avoidance paradigm (Bravo-Rivera et al., 2015) or lever-press avoidance paradigm (Beck et al., 2014) demonstrating that PL drives avoidance behavior acquisition/expression. Our results are also consistent with clinical results indicating that in healthy humans avoidance is linked to an increased reactivity of the anterior ACC and the dmPFC (Schlund et al., 2015). Based on our fos results, we also identified a structure considered to regulate the defensive output behavioral responses, namely the PAG and more specifically the dIPAG. Even though the dIPAG show a clear upregulation in the animals performing the avoidance task, this structure did not show a significative difference between the good and the bad avoiders. It is however important to note that no direct correlation between the dIPAG Fos upregulation and freezing was found (data not shown). As mentioned before, the dIPAG is involved in both passive and active defensive responses (Vianna et al., 2001). Knowing that can explain the fact that the dIPAG could have been recruited by the dmPFC in the case of the good avoiders performing avoidance; and by other direct inputs as the hypothalamus, or indirect inputs as the Amygdala via the vIPAG, in the case of bad avoiders with a higher freezing level (Halladay and Blair, 2015; Rozeske et al., 2018; Tovote et al., 2016).

**The caudal dmPFC promotes active avoidance :** Using the conclusions brought by our in vivo electrophysiological, antidromic and optogenetic data, we can strongly suggest that avoidance behavior is driven by an activation of a subpopulation of dmPFC PNs (Khoder, 2018) which opposes the results of a recent paper (Diehl et al., 2018) suggesting that avoidance is rather associated with an inhibition of dmPFC activity. We think that those discrepancies are linked to the differences in the rostro-caudal axis of manipulation/recordings at the dmPFC level. Indeed our recordings in the dmPFC and optogenetic manipulations are made in the caudal dmPFC (A.P. < +2.1) whereas in the platform-mediated paradigm (Diehl, et al., 2018) the results concern the rostral dmPFC (A.P. > 2.1). Furthermore, a pilot study from our lab tends to show that inhibiting the

rostral dmPFC to dlPAG pathway promotes avoidance learning, whereas the caudal pathway abolishes it. The opposing roles in avoidance learning played by the rostral dmPFC and caudal dmPFC rise an important question, being to determine which structure is critically involved in the selection of the behavioral response during avoidance. Does the selection of avoidance behavior depend on the rostral vs caudal dmPFC local connectivity or is the selection made at downstream structures like the dl/IPAG? It also reopens the question about where the behavioral switch between freezing and avoidance is made. We demonstrated here that the caudal PL was specifically involved in avoidance learning, and not freezing. However, as our recordings in the cdmPFC infer, there are cells activated in both freezing and avoidance (Khoder, 2018). rostral dmPFC to dlPAG is yet to be studied to understand its role in this behavioral switch. Additional experiments will be required to specifically address this question.

**The dmPFC to dlPAG pathway is necessary for promoting avoidance behavior but not freezing:** The modulation of the dmPFC to dlPAG pathway was made in two steps. Firstly we activated the pathway in the bad avoiders to see the evolution of their performance in the two way SigA paradigm. Interestingly, the two days of post training stimulation made the **bad avoiders** improve their avoidance performances, and the animals discriminate better the CS+ and the CS-. However the freezing level in DC condition was unchanged (Khoder, 2018). The fact that the animals kept improving after the stimulation days, showed us that the stimulation of the pathway couldn't promote avoidance expression only. This hypothesis was clarified with the inhibition of the pathway in the **good avoiders**. In fact early training inhibition but not post training inhibition impaired their capacity to acquire avoidance. These results validated the fact that the dmPFC pathway is sufficient and necessary to promote avoidance behavior. However its role in freezing is not present. These results refute the fact that freezing and avoidance are mediated by the same caudal dmPFC to dlPAG pathway, and go along with other works emphasizing the involvement of the dmPFC in platform-mediated avoidance but not freezing behavior (Bravo-Rivera et al., 2014; Diehl et al., 2018). However as opposed to these works, our electrophysiological data showed that a proportion cdmPFC neurons encode for freezing only. It is then more likely that freezing and avoidance are driven in

the dmPFC by independent subsets of neuron populations. A possible candidate mediating freezing expression alone would be the dmPFC to BLA pathway, as it is proposed in the literature (Courtin et al., 2014).

## Second Part

**Behaviors involved in the passive avoidance paradigm:** In the second part, using the paradigm based on our previous work on rats, we were able to reproduce the results in mice (Viellard et al., 2016). The two groups of animals presented clear behavioral differences in terms of contextual fear responses. Following the day of the conditioning, the group kept enclosed in the in the shock chamber, spent close to 25% of the time frozen, and 51% of the time immobile in a crouched back posture, sniffing the environment (Blanchard and Blanchard, 1969). In contrast, animals placed in the home cage with open access to the conditioning chamber presented only a minimal amount of freezing (1%), and spent most of the time of the test risk assessing the environment with either crouch sniff (61%) or doing stretch postures (21%). This group of animals did not enter the conditioning chamber and, in addition to the fear responses, also actively explore the home box and corridor close to 17% of the time. The groups that did not receive shocks actively explored their environment fearlessly in both conditions. As reported before, the different conditions in which the animal recalls a threatening environment, affects the responses of the animal (Viellard et al., 2016). The present findings of this shock-based passive avoidance paradigm can be compared with previous studies from our lab using the same apparatus and experimental design for either cat exposure or social defeat. As we have just found for footshock conditioning, animals conditioned with either predator threat or social defeat, presented a similar form of contextual fear responses (i.e., risk assessment). And when placed in the home cage with access to the compartment associated with either predator threat or social defeat, they largely avoided this chamber. The amplitude of the fear response is difficult to compare as the impact of the different threats (i.e., footshock, aggressive conspecific and predator) on the animal is not measurable. However, compared to other shock-based passive avoidance paradigms, this one leaves a stronger conditioning as the animal entirely avoid the conditioning chamber. In our case, on the conditioning day, the animal

received a series of shocks enclosed in the conditioning chamber, whereas in the step-down inhibitory avoidance another form of shock-based passive avoidance paradigm, the animal has the possibility to escape after the first shock (Ambrogio Lorenzini et al., 1999). Furthermore the long term pre-exposure habituation (three days) is known to influence the conditioning process, as it has been shown that context pre-exposure facilitates and strengthen the learning of context-shock association (Fanselow, 1980; Rudy, 2009). One could argue that a strong conditioning as the one in our paradigm could lead to generalization. However, the results in the open field showed the behavioral ratio between fearful and fearless behaviors does not change after the conditioning day, whereas this ratio greatly increases in the conditioning apparatus. Thus suggesting that the animals differentiate the aversive and neutral contexts. The experiment was also set using mild shocks of 0.6mA, which are unlikely to create generalization (Baldi et al., 2004). Compared to animals tested enclosed in conditioning cage, the present paradigm (using a shock as a controllable threat) yields the expression of a larger range of risk assessment behaviors, which are good candidates for modeling anxiety behaviors in humans (Blanchard and Meyza, 2019).

**Septo-hippocampal–hypothalamic-brainstem circuit putatively involved in inhibitory avoidance: comparison to other threats and conditions:** The present results are also in line with our previous results, in rats (Viellard et al., 2016), showing that the fear conditioned animals, which were able to avoid the conditioning chamber, presented increased Fos expression in a circuit formed by the subiculum, the lateral septum, the juxtadorsomedial part of lateral hypothalamic area (LHAjd), the dorsal premammillary nucleus and the lateral and dorsal parts of the periaqueductal gray. Anatomical and functional data suggest that this septo/hippocampal-hypothalamic-brainstem circuit should be putatively involved in mediating contextual avoidance. Interestingly, social defeat to an aggressive conspecific, exposure to a snake predator and restraint stress also up-regulate Fos expression in this same circuit. Notably, in response to all these threats, animals displayed a significant increase in Fos expression in the juxtadorsomedial region of the lateral hypothalamic area (LHAjd) (Motta and Canteras, 2015). The LHAjd conveys information to the dorsal premammillary nucleus from the septo-hippocampal system

(Hahn and Swanson, 2012). The septo-hippocampal system has been proposed to play a pivotal role in anxiety in response to conflict situations, by interrupting ongoing behavior and increasing the level of arousal and attention to enhance gathering information (Gray and McNaughton, 2003). In fact, the hippocampus may work as a context analyzer providing a spatial mapping of the environment derived from two sets of information: one based on the external environment and the other based on self-motion (Burgess et al., 2002). Of relevance to the present study, the hippocampus contains a special kind of cell, the boundary vector cell (BVC), which codes for environmental boundaries (irrespective of their sensory nature (Stewart et al., 2014). Interestingly, the distribution of the BVCs and the cells that project to the LHAjd seem to overlap, at least partially, in the subiculum (Hahn and Swanson, 2012). The concept of an environmental boundary is somewhat abstract and represents an effective obstacle to locomotion that does not necessarily involve physical prevention of movement (Stewart et al., 2014). Considering the evidences, all these forms of threats (i.e., physical constraint, exposure to an aggressive conspecific or a snake predator, and the avoidance of a threatening chamber) set clear environmental boundaries, constraining the animals either physically (by the restraining apparatus) or behaviorally (conspecific aggressor, snake predator, and the threatening chamber). Therefore, it would be interesting to investigate whether the avoidance of the threatening chamber would work as an environmental boundary signaled by BVC cells. As previously mentioned, on the efferent side, the LHAjd projects densely to the dorsal premammillary nucleus (PMD), in addition to the dorsomedial and lateral parts of the periaqueductal gray (PAGdm,l) (Hahn and Swanson, 2012), all of which have been shown to present a significant Fos increase in response to passive avoidance, as well as to a social aggressor and snake threat (Faturi et al., 2013; Motta et al., 2009). The present results gives further support to the idea that there are interesting commonalities among restraint stress, social defeat, snake threat and passive contextual avoidance, suggesting a septo-hippocampal–hypothalamic-brainstem path likely to respond to the environmental boundary restriction that may act as common stressor component for all these types of stress.

**PMD influences both inhibitory avoidance and memory reconsolidation:** The PMD has a pivotal role in the septo/hippocampal-hypothalamic-brainstem circuit putatively involved in mediating contextual avoidance. On the afferent side, the PMD integrates hippocampal information likely related to signaling environmental boundaries, and on the efferent side, the nucleus projects to the periaqueductal gray, which is critically involved in the expression of avoidance responses (Motta et al., 2017). The present results indicate that pharmacogenetic inhibition of the PMD resulted in a general decrease in risk assessment behaviors. Thus, the CNO-injected animals expressing hM4D receptor in the PMD spent around 150 seconds risk assessing the shock-related context of the context exposure, in comparison to close to 300 seconds for the control group. Moreover, the group in which the PMD was inhibited spend about 130 seconds exploring the conditioning cage whereas the control group did not enter this cage. Notably, in the CCK CRE line used in this experiment, apart from the PMD, the expression of the hM4D receptor spread to a certain degree over the mammillary bodies, which also contain CCK cell bodies. However, CNO-injected animals containing the hM4D receptor only in the mammillary bodies did not reduce risk assessment and did not enter the shock-related chamber during the day after the conditioning. Moreover, our Fos analysis showed no involvement of the mammillary bodies in passive avoidance. In line with the present results, pharmacological inactivation of the PMD, but not of the nearby mammillary nuclei, was able to significantly reduce the contextual conditioned responses to predatory threats (Cezario et al., 2008). As in the present case, in this experiment, animals were tested in a similar apparatus with a home cage linked to a corridor and the threatening chamber, and muscimol injection in the PMD, on the day after cat exposure, drastically reduced risk assessment responses and the animal entered the threatening chamber (Cezario et al., 2008).

On the day following PMD inhibition, we found a decrease in the inhibitory avoidance in the animals re-exposed to the threatening context. Thus, suggesting that PMD inhibition influenced memory processes related to fear reconsolidation during exposure to the threatening environment. Accordingly, the group of animals expressing hM4D in the PMD that received CNO during the first day of exposure to the shock-related environment, when re-tested the following day in the same context, presented decreased risk assessment responses and spent significantly higher amount of time in the

conditioning compartment. In line with the present results, pharmacological blockade of either beta-adrenoceptor or NMDA receptor in the PMD, but not in the adjacent mammillary bodies, immediately before the conditioning session, reduced the defensive response to the cat odor and also, 24 hours later, to the cat-odor related environment (Canteras et al., 2008; Do Monte et al., 2008). The PMD's role in fear memory may be viewed as either an impairment in fear memory processing or the result of decreased emotional component of the aversive event during the learning stage. In favor of the view that the decrease of emotional component during the learning stage does not necessarily influence fear memory, (de Andrade Rufino et al., 2019) found that ventral periaqueductal gray lesions resulting in clear decrease of innate defensive responses to a predator did not affect anti-predatory contextual fear learning.

Overall, our results indicate that the PMD influences both the expression of inhibitory avoidance and the memory reconsolidation processes during exposure to the shock-related context.

#### **How the PMD's targets influence passive avoidance and fear memory reconsolidation :**

The functional role of the PMD appears to depend on its branched pathway to the periaqueductal gray (PAG) and the ventral part of the anteromedial thalamic nucleus (AMv) (Canteras and Swanson, 1992). Therefore, we examined how the PMD projections to the PAG and AMv influences defensive responses during the contextual avoidance and the memory reconsolidation process during exposure to the shock-related context. To this end, we induced halorodopsin expression in the PMD cells, and silenced the PMD's terminals in either the PAG or the AMv. At first, we were expecting that silencing the projections to the PAG would influence passive avoidance during exposure to the shock-related context, whereas inactivation of the projections to the AMv would disrupt the memory reconsolidation process. However, inactivation of PMD terminals in the PAG or in the AMv had similar effects. Thus, optogenetic inhibition of PMD projection to the PAG or to the AMv during exposure to the shock-related context resulted in decreased risk assessment responses and increase the time spent in the conditioning chamber. Moreover, compared to the control group, animals that received optogenetic inhibition of PMD's terminals in the PAG or in the AMv, when re-tested 24 hours late in the same

context, presented a reduction in risk assessment responses and spent significantly higher amount of time in the conditioning chamber. Therefore, silencing the PMD's projections to the PAG or the AMv interfere with both the expression of defensive responses during contextual avoidance and the memory reconsolidation process. At this point, we need to understand how the PMD's targets could influence both the expression of defensive responses during the inhibitory avoidance and the memory reconsolidation processing.

Previous studies have shown that pharmacological inactivation of the AMv disrupts the acquisition of contextual memory to predatory threats (de Lima et al., 2017). The AMv role on memory processing seems to depend on its projection to a cortical network (formed by the prelimbic, anterior cingulate, visual associative and ventral retrosplenial areas), which influences fear memory and has access to key elements involved in memory processing, such as the basolateral amygdala and the hippocampal formation (de Lima et al., 2019).

In the PAG, particularly its dorsal part has been shown to support fear learning. Of relevance, the dorsal PAG seems critical for the acquisition of contextual fear memory to predator threats (de Andrade Rufino et al., 2019; Souza and Carobrez, 2016). Moreover, several studies using classical fear conditioning to sound-, light- or odor-conditioned stimuli (CS) have shown that electrical, chemical or optogenetic stimulation of the dorsal PAG may work as a useful US to support associative learning (Deng et al., 2016; Di Scala et al., 1987; Di Scala and Sandner, 1989; Kim et al., 2013; Kincheski et al., 2012). The dorsal PAG provides a number of parallel thalamic paths likely to influence fear learning. Thus, the dorsal PAG provides direct inputs to the nucleus reuniens, the central lateral nucleus, the lateral dorsal nucleus, the supragenicolate nucleus, and the parvicellular subparafascicular nucleus (Kincheski et al., 2012). The nucleus reuniens represents the main thalamic source of projections to the hippocampal formation (Vertes et al., 2006); the central lateral nucleus and the lateral dorsal nucleus project to cortical areas involved in the cortical circuit mentioned above that influences fear learning (i.e., the anterior cingulate and retrosplenial areas) (Furlong et al., 2010; van Groen and Wyss, 1992); and the supragenicolate and the parvicellular subparafascicular nuclei project

densely to the lateral amygdalar nucleus (Linke et al., 2000). However, further studies are needed to address how these dorsal PAG-thalamic pathways may influence fear learning. During exposure to environments previously associated with a threat, such as a predator, an aggressive conspecific or, as in the present case, a footshock, the threat is more ambiguous and evokes risk assessment responses, including a careful scanning of the environment in the crouched position (crouch sniffing) and attempts to approach the threatening stimulus by stretching the body (stretch postures) (Faturi et al., 2013; Ribeiro-Barbosa et al., 2005; Viellard et al., 2016). Previous studies using cytotoxic lesions and pharmacological inactivation have shown that the dorsal PAG appears to exert critical control on risk assessment responses (Faturi et al., 2013; Pobbe et al., 2011). In agreement with this idea, the present results showed that optogenetic inhibition of the PMD's projection to the PAG, which is putatively a glutamatergic projection, decreased risk assessment response during exposure to the shock-related context. Risk assessment responses are very complex, and it is not clear how the dorsal PAG influences these responses. Nevertheless, ascending dorsal PAG projections to prosencephalic targets have been proposed to influence risk assessment behaviors (Motta et al., 2017).

One of our most puzzling results was the drastic reduction of risk assessment in response to the optogenetic inhibition of the PMD projection to the AMv in animals exposed to the shock-related context. Recent results from our lab indicate that the AMv-related cortical network may influence the expression of contextual fear responses. In this way, we have found that optogenetic inhibition of anterior cingulate projection to the dorsal PAG significantly reduced risk assessment responses during exposure to context previously related to a predator. Therefore, the PMD would influence the expression of inhibitory avoidance during exposure to shock-related context through its direct projection to the PAG and the through the projection to the AMv, which may ultimately impact on the anterior cingulate area – dorsal PAG pathway.

To help understand better the nature and the specific role of the cells projecting to the AMV and the dIPAG, further experiments will be done using electrophysiological recordings of PMD cells and try to correlate their firing rate with specific behavioral responses. Concerning the afferent pathway of the PMD, it has been previously noted how important the hippocampus is in passive avoidance and fear learning, and future

studies will aim to investigate whether the avoidance of the threatening chamber would work as an environmental boundary perhaps signaled by BVC cells.

## References

- Ambrogi Lorenzini, C.G., Baldi, E., Bucherelli, C., Sacchetti, B., Tassoni, G., 1999. Neural topography and chronology of memory consolidation: a review of functional inactivation findings. *Neurobiol. Learn. Mem.* 71, 1–18. <https://doi.org/10.1006/nlme.1998.3865>
- Amorapanth, P., LeDoux, J.E., Nader, K., 2000. Different lateral amygdala outputs mediate reactions and actions elicited by a fear-arousing stimulus. *Nat. Neurosci.* 3, 74–79. <https://doi.org/10.1038/71145>
- Baldi, E., Lorenzini, C.A., Bucherelli, C., 2004. Footshock intensity and generalization in contextual and auditory-cued fear conditioning in the rat. *Neurobiol. Learn. Mem.* 81, 162–166. <https://doi.org/10.1016/j.nlm.2004.02.004>
- Beck, K.D., Jiao, X., Smith, I.M., Myers, C.E., Pang, K.C.H., Servatius, R.J., 2014. ITI-Signals and Prelimbic Cortex Facilitate Avoidance Acquisition and Reduce Avoidance Latencies, Respectively, in Male WKY Rats. *Front. Behav. Neurosci.* 8, 403. <https://doi.org/10.3389/fnbeh.2014.00403>
- Blanchard, D.C., 2017. Translating dynamic defense patterns from rodents to people. *Neurosci. Biobehav. Rev., Translational Neuroscience & Mental Disorders: bridging the gap between animal models and the human condition* 76, 22–28. <https://doi.org/10.1016/j.neubiorev.2016.11.001>
- Blanchard, D.C., Blanchard, R.J., 1972. Innate and conditioned reactions to threat in rats with amygdaloid lesions. *J. Comp. Physiol. Psychol.* 81, 281–290. <https://doi.org/10.1037/h0033521>
- Blanchard, D.C., Canteras, N.S., Markham, C.M., Pentkowski, N.S., Blanchard, R.J., 2005. Lesions of structures showing FOS expression to cat presentation: Effects on responsivity to a Cat, Cat odor, and nonpredator threat. *Neurosci. Biobehav. Rev., Defensive Behavior* 29, 1243–1253. <https://doi.org/10.1016/j.neubiorev.2005.04.019>
- Blanchard, D.C., Griebel, G., Blanchard, R.J., 2003a. The Mouse Defense Test Battery: pharmacological and behavioral assays for anxiety and panic. *Eur. J. Pharmacol., Animal Models of Anxiety Disorders* 463, 97–116. [https://doi.org/10.1016/S0014-2999\(03\)01276-7](https://doi.org/10.1016/S0014-2999(03)01276-7)
- Blanchard, D.C., Li, C.I., Hubbard, D., Markham, C.M., Yang, M., Takahashi, L.K., Blanchard, R.J., 2003b. Dorsal premammillary nucleus differentially modulates defensive behaviors induced by different threat stimuli in rats. *Neurosci. Lett.* 345, 145–148. [https://doi.org/10.1016/S0304-3940\(03\)00415-4](https://doi.org/10.1016/S0304-3940(03)00415-4)

- Blanchard, D.C., Meyza, K., 2019. Risk assessment and serotonin: Animal models and human psychopathologies. *Behav. Brain Res.* 357–358, 9–17.  
<https://doi.org/10.1016/j.bbr.2017.07.008>
- Blanchard, R.J., Blanchard, D.C., 1969. Passive and active reactions to fear-eliciting stimuli. *J. Comp. Physiol. Psychol.* 68, 129–135.  
<https://doi.org/10.1037/h0027676>
- Bravo-Rivera, C., Roman-Ortiz, C., Brignoni-Perez, E., Sotres-Bayon, F., Quirk, G.J., 2014. Neural Structures Mediating Expression and Extinction of Platform-Mediated Avoidance. *J. Neurosci.* 34, 9736–9742.  
<https://doi.org/10.1523/JNEUROSCI.0191-14.2014>
- Bravo-Rivera, C., Roman-Ortiz, C., Montesinos-Cartagena, M., Quirk, G.J., 2015. Persistent active avoidance correlates with activity in prelimbic cortex and ventral striatum. *Front. Behav. Neurosci.* 9.  
<https://doi.org/10.3389/fnbeh.2015.00184>
- Burgess, N., Maguire, E.A., O'Keefe, J., 2002. The human hippocampus and spatial and episodic memory. *Neuron* 35, 625–641. [https://doi.org/10.1016/s0896-6273\(02\)00830-9](https://doi.org/10.1016/s0896-6273(02)00830-9)
- Canteras, N.S., Kroon, J.A.V., Do-Monte, F.H.M., Pavesi, E., Carobrez, A.P., 2008. Sensing danger through the olfactory system: The role of the hypothalamic dorsal premammillary nucleus. *Neurosci. Biobehav. Rev.* 32, 1228–1235.  
<https://doi.org/10.1016/j.neubiorev.2008.05.009>
- Canteras, N.S., Swanson, L.W., 1992. The dorsal premammillary nucleus: an unusual component of the mammillary body. *Proc. Natl. Acad. Sci. U. S. A.* 89, 10089–10093.
- Canto-de-Souza, L., Mattioli, R., 2016. The consolidation of inhibitory avoidance memory in mice depends on the intensity of the aversive stimulus: The involvement of the amygdala, dorsal hippocampus and medial prefrontal cortex. *Neurobiol. Learn. Mem.* 130, 44–51.  
<https://doi.org/10.1016/j.nlm.2016.01.012>
- Carrive, P., 1993. The periaqueductal gray and defensive behavior: functional representation and neuronal organization. *Behav. Brain Res.* 58, 27–47.
- Cezario, A.F., Ribeiro-Barbosa, E.R., Baldo, M.V.C., Canteras, N.S., 2008. Hypothalamic sites responding to predator threats - the role of the dorsal premammillary nucleus in unconditioned and conditioned antipredatory defensive behavior. *Eur. J. Neurosci.* 28, 1003–1015. <https://doi.org/10.1111/j.1460-9568.2008.06392.x>

- Comoli, E., Ribeiro-Barbosa, E.R., Canteras, N.S., 2000. Afferent connections of the dorsal premammillary nucleus. *J. Comp. Neurol.* 423, 83–98.  
[https://doi.org/10.1002/1096-9861\(20000717\)423:1<83::aid-cne7>3.0.co;2-3](https://doi.org/10.1002/1096-9861(20000717)423:1<83::aid-cne7>3.0.co;2-3)
- Corbit, L.H., Muir, J.L., Balleine, B.W., 2001. The role of the nucleus accumbens in instrumental conditioning: Evidence of a functional dissociation between accumbens core and shell. *J. Neurosci. Off. J. Soc. Neurosci.* 21, 3251–3260.
- Courtin, J., Chaudun, F., Rozeske, R.R., Karalis, N., Gonzalez-Campo, C., Wurtz, H., Abdi, A., Baufreton, J., Bienvenu, T.C.M., Herry, C., 2014. Prefrontal parvalbumin interneurons shape neuronal activity to drive fear expression. *Nature* 505, 92–96. <https://doi.org/10.1038/nature12755>
- de Andrade Rufino, R., Mota-Ortiz, S.R., De Lima, M.A.X., Baldo, M.V.C., Canteras, N.S., 2019. The rostradorsal periaqueductal gray influences both innate fear responses and acquisition of fear memory in animals exposed to a live predator. *Brain Struct. Funct.* 224, 1537–1551. <https://doi.org/10.1007/s00429-019-01852-6>
- de Lima, M.A.X., Baldo, M.V.C., Canteras, N.S., 2019. Revealing a Cortical Circuit Responsive to Predatory Threats and Mediating Contextual Fear Memory. *Cereb. Cortex* 29, 3074–3090. <https://doi.org/10.1093/cercor/bhy173>
- de Lima, M.A.X., Baldo, M.V.C., Canteras, N.S., 2017. A role for the anteromedial thalamic nucleus in the acquisition of contextual fear memory to predatory threats. *Brain Struct. Funct.* 222, 113–129. <https://doi.org/10.1007/s00429-016-1204-2>
- Deng, H., Xiao, X., Wang, Z., 2016. Periaqueductal Gray Neuronal Activities Underlie Different Aspects of Defensive Behaviors. *J. Neurosci.* 36, 7580–7588.  
<https://doi.org/10.1523/JNEUROSCI.4425-15.2016>
- Di Scala, G., Mana, M.J., Jacobs, W.J., Phillips, A.G., 1987. Evidence of Pavlovian conditioned fear following electrical stimulation of the periaqueductal grey in the rat. *Physiol. Behav.* 40, 55–63. [https://doi.org/10.1016/0031-9384\(87\)90185-5](https://doi.org/10.1016/0031-9384(87)90185-5)
- Di Scala, G., Sandner, G., 1989. Conditioned place aversion produced by microinjections of semicarbazide into the periaqueductal gray of the rat. *Brain Res.* 483, 91–97. [https://doi.org/10.1016/0006-8993\(89\)90038-3](https://doi.org/10.1016/0006-8993(89)90038-3)
- Dickinson, A., 1985. Actions and habits: the development of behavioural autonomy. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 308, 67–78.  
<https://doi.org/10.1098/rstb.1985.0010>
- Diehl, M.M., Bravo-Rivera, C., Rodriguez-Romaguera, J., Pagan-Rivera, P.A., Burgos-Robles, A., Roman-Ortiz, C., Quirk, G.J., 2018. Active avoidance requires

- inhibitory signaling in the rodent prelimbic prefrontal cortex. *eLife* 7, e34657.  
<https://doi.org/10.7554/eLife.34657>
- Dielenberg, R.A., Hunt, G.E., McGregor, I.S., 2001. 'When a rat smells a cat': the distribution of Fos immunoreactivity in rat brain following exposure to a predatory odor. *Neuroscience* 104, 1085–1097. [https://doi.org/10.1016/S0306-4522\(01\)00150-6](https://doi.org/10.1016/S0306-4522(01)00150-6)
- Do Monte, F.H.M., Canteras, N.S., Fernandes, D., Assreuy, J., Carobrez, A.P., 2008. New Perspectives on -Adrenergic Mediation of Innate and Learned Fear Responses to Predator Odor. *J. Neurosci.* 28, 13296–13302.  
<https://doi.org/10.1523/JNEUROSCI.2843-08.2008>
- Ellard, C.G., Eller, M.C., 2009. Spatial cognition in the gerbil: computing optimal escape routes from visual threats. *Anim. Cogn.* 12, 333–345.  
<https://doi.org/10.1007/s10071-008-0193-9>
- Euston, D.R., Gruber, A.J., McNaughton, B.L., 2012. The role of medial prefrontal cortex in memory and decision making. *Neuron* 76, 1057–1070.  
<https://doi.org/10.1016/j.neuron.2012.12.002>
- Fanselow, M.S., 1980. Conditional and unconditional components of post-shock freezing. *Pavlov. J. Biol. Sci. Off. J. Pavlov.* 15, 177–182.  
<https://doi.org/10.1007/BF03001163>
- Faturi, C.B., Jr, M.J.R., Baldo, M.V.C., Canteras, N.S., 2013. Functional mapping of the circuits involved in the expression of contextual fear responses in socially defeated animals. *Brain Struct. Funct.* 219, 931–946.  
<https://doi.org/10.1007/s00429-013-0544-4>
- Franklin, K.B.J., Paxinos, G., 2008. The mouse brain in stereotaxic coordinates, Compact 3. ed. ed. Elsevier, Academic Press, Amsterdam.
- Furlong, T.M., Cole, S., Hamlin, A.S., McNally, G.P., 2010. The role of prefrontal cortex in predictive fear learning. *Behav. Neurosci.* 124, 574–586.  
<https://doi.org/10.1037/a0020739>
- Gabbott, P.L.A., Warner, T.A., Jays, P.R.L., Salway, P., Busby, S.J., 2005. Prefrontal cortex in the rat: Projections to subcortical autonomic, motor, and limbic centers. *J. Comp. Neurol.* 492, 145–177. <https://doi.org/10.1002/cne.20738>
- Gabriel, M., Kubota, Y., Sparenborg, S., Straube, K., Vogt, B.A., 1991. Effects of cingulate cortical lesions on avoidance learning and training-induced unit activity in rabbits. *Exp. Brain Res.* 86, 585–600.  
<https://doi.org/10.1007/bf00230532>
- Galatzer-Levy, I.R., Moscarello, J., Blessing, E.M., Klein, J., Cain, C.K., LeDoux, J.E., 2014. Heterogeneity in signaled active avoidance learning: substantive and

- methodological relevance of diversity in instrumental defensive responses to threat cues. *Front. Syst. Neurosci.* 8. <https://doi.org/10.3389/fnsys.2014.00179>
- Gebhardt, N., Bär, K.-J., Boettger, M.K., Grecksch, G., Keilhoff, G., Reichart, R., Becker, A., 2013. Vagus nerve stimulation ameliorated deficits in one-way active avoidance learning and stimulated hippocampal neurogenesis in bullectomized rats. *Brain Stimulat.* 6, 78–83. <https://doi.org/10.1016/j.brs.2012.01.009>
- Gladwin, T.E., Hashemi, M.M., van Ast, V., Roelofs, K., 2016. Ready and waiting: Freezing as active action preparation under threat. *Neurosci. Lett.* 619, 182–188. <https://doi.org/10.1016/j.neulet.2016.03.027>
- Gold, P.E., Macri, J., McGaugh, J.L., 1973. Retrograde amnesia produced by subseizure amygdala stimulation. *Behav. Biol.* 9, 671–680. [https://doi.org/10.1016/s0091-6773\(73\)80128-2](https://doi.org/10.1016/s0091-6773(73)80128-2)
- Gray, J.A., McNaughton, N., 2003. *The Neuropsychology of Anxiety*. Oxford University Press. <https://doi.org/10.1093/acprof:oso/9780198522713.001.0001>
- Groenewegen, H.J., 1988. Organization of the afferent connections of the mediodorsal thalamic nucleus in the rat, related to the mediodorsal-prefrontal topography. *Neuroscience* 24, 379–431. [https://doi.org/10.1016/0306-4522\(88\)90339-9](https://doi.org/10.1016/0306-4522(88)90339-9)
- Gross, C.T., Canteras, N.S., 2012. The many paths to fear. *Nat. Rev. Neurosci.* 13, 651–658. <https://doi.org/10.1038/nrn3301>
- Hahn, J.D., Swanson, L.W., 2012. Connections of the Lateral Hypothalamic Area Juxtadorsomedial Region in the Male Rat. *J. Comp. Neurol.* 520, 1831–1890. <https://doi.org/10.1002/cne.23064>
- Halladay, L.R., Blair, H.T., 2015. Distinct ensembles of medial prefrontal cortex neurons are activated by threatening stimuli that elicit excitation vs. inhibition of movement. *J. Neurophysiol.* 114, 793–807. <https://doi.org/10.1152/jn.00656.2014>
- Harris, K.D., Shepherd, G.M.G., 2015. The neocortical circuit: themes and variations. *Nat. Neurosci.* 18, 170–181. <https://doi.org/10.1038/nn.3917>
- Heidbreder, C.A., Groenewegen, H.J., 2003. The medial prefrontal cortex in the rat: evidence for a dorso-ventral distinction based upon functional and anatomical characteristics. *Neurosci. Biobehav. Rev.* 27, 555–579. <https://doi.org/10.1016/j.neubiorev.2003.09.003>
- Herry, C., Trifilieff, P., Micheau, J., Lüthi, A., Mons, N., 2006. Extinction of auditory fear conditioning requires MAPK/ERK activation in the basolateral amygdala. *Eur. J. Neurosci.* 24, 261–269. <https://doi.org/10.1111/j.1460-9568.2006.04893.x>

- Hofmann, S.G., Hay, A.C., 2018. Rethinking Avoidance: Toward a Balanced Approach to Avoidance in Treating Anxiety Disorders. *J. Anxiety Disord.* 55, 14–21.  
<https://doi.org/10.1016/j.janxdis.2018.03.004>
- Hoover, W.B., Vertes, R.P., 2007. Anatomical analysis of afferent projections to the medial prefrontal cortex in the rat. *Brain Struct. Funct.* 212, 149–179.  
<https://doi.org/10.1007/s00429-007-0150-4>
- Hubel, D.H., Wiesel, T.N., 1962. Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. *J. Physiol.* 160, 106–154.  
<https://doi.org/10.1113/jphysiol.1962.sp006837>
- Keay, K.A., Bandler, R., 2001. Parallel circuits mediating distinct emotional coping reactions to different types of stress. *Neurosci. Biobehav. Rev.* 25, 669–678.  
[https://doi.org/10.1016/S0149-7634\(01\)00049-5](https://doi.org/10.1016/S0149-7634(01)00049-5)
- Khoder, S., 2018. Role of the prefrontal-brainstem pathway in mediating avoidance behavior [WWW Document]. URL  
<http://www.sudoc.abes.fr/xslt/DB=2.1//SRCH?IKT=12&TRM=234768711>  
 (accessed 11.7.19).
- Kim, E.J., Horovitz, O., Pellman, B.A., Tan, L.M., Li, Q., Richter-Levin, G., Kim, J.J., 2013. Dorsal periaqueductal gray-amygdala pathway conveys both innate and learned fear responses in rats. *Proc. Natl. Acad. Sci. U. S. A.* 110, 14795–14800.  
<https://doi.org/10.1073/pnas.1310845110>
- Kim, E.J., Park, M., Kong, M.-S., Park, S.G., Cho, J., Kim, J.J., 2015. Alterations of hippocampal place cells in foraging rats facing a “predatory” threat. *Curr. Biol. CB* 25, 1362–1367. <https://doi.org/10.1016/j.cub.2015.03.048>
- Kincheski, G.C., Mota-Ortiz, S.R., Pavesi, E., Canteras, N.S., Carobrez, A.P., 2012. The dorsolateral periaqueductal gray and its role in mediating fear learning to life threatening events. *PloS One* 7, e50361.  
<https://doi.org/10.1371/journal.pone.0050361>
- Kitamura, T., Ogawa, S.K., Roy, D.S., Okuyama, T., Morrissey, M.D., Smith, L.M., Redondo, R.L., Tonegawa, S., 2017. Engrams and circuits crucial for systems consolidation of a memory. *Science* 356, 73–78.  
<https://doi.org/10.1126/science.aam6808>
- Lázaro-Muñoz, G., LeDoux, J.E., Cain, C.K., 2010. Sidman Instrumental Avoidance Initially Depends on Lateral and Basal Amygdala and is Constrained by Central Amygdala-mediated Pavlovian Processes. *Biol. Psychiatry* 67, 1120–1127.  
<https://doi.org/10.1016/j.biopsych.2009.12.002>
- Lechner, H.A., Squire, L.R., Byrne, J.H., 1999. 100 Years of Consolidation—Remembering Müller and Pilzecker. *Learn. Mem.* 6, 77–87.  
<https://doi.org/10.1101/lm.6.2.77>

- LeDoux, J., 2012. Rethinking the Emotional Brain. *Neuron* 73, 653–676.  
<https://doi.org/10.1016/j.neuron.2012.02.004>
- LeDoux, J.E., 2000. Emotion Circuits in the Brain. *Annu. Rev. Neurosci.* 23, 155–184.  
<https://doi.org/10.1146/annurev.neuro.23.1.155>
- LeDoux, J.E., Moscarello, J., Sears, R., Campese, V., 2017. The birth, death and resurrection of avoidance: a reconceptualization of a troubled paradigm. *Mol. Psychiatry* 22, 24–36. <https://doi.org/10.1038/mp.2016.166>
- Lee, J.L.C., Milton, A.L., Everitt, B.J., 2006. Reconsolidation and Extinction of Conditioned Fear: Inhibition and Potentiation. *J. Neurosci.* 26, 10051–10056.  
<https://doi.org/10.1523/JNEUROSCI.2466-06.2006>
- Linke, R., Braune, G., Schwegler, H., 2000. Differential projection of the posterior paralaminar thalamic nuclei to the amygdaloid complex in the rat. *Exp. Brain Res.* 134, 520–532. <https://doi.org/10.1007/s002210000475>
- Löw, A., Weymar, M., Hamm, A.O., 2015. When Threat Is Near, Get Out of Here: Dynamics of Defensive Behavior During Freezing and Active Avoidance. *Psychol. Sci.* 26, 1706–1716. <https://doi.org/10.1177/0956797615597332>
- Maren, S., 2001. Neurobiology of Pavlovian Fear Conditioning. *Annu. Rev. Neurosci.* 24, 897–931. <https://doi.org/10.1146/annurev.neuro.24.1.897>
- Merlo, E., Bekinschtein, P., Jonkman, S., Medina, J.H., 2015. Molecular Mechanisms of Memory Consolidation, Reconsolidation, and Persistence. *Neural Plast.* 2015, 1–2. <https://doi.org/10.1155/2015/687175>
- Morgan, M.A., LeDoux, J.E., 1995. Differential contribution of dorsal and ventral medial prefrontal cortex to the acquisition and extinction of conditioned fear in rats. *Behav. Neurosci.* 109, 681–688. <https://doi.org/10.1037//0735-7044.109.4.681>
- Moriarty, O., Roche, M., McGuire, B.E., Finn, D.P., 2012. Validation of an air-puff passive-avoidance paradigm for assessment of aversive learning and memory in rat models of chronic pain. *J. Neurosci. Methods* 204, 1–8.  
<https://doi.org/10.1016/j.jneumeth.2011.10.024>
- Moscarello, J.M., LeDoux, J.E., 2013. Active Avoidance Learning Requires Prefrontal Suppression of Amygdala-Mediated Defensive Reactions. *J. Neurosci.* 33, 3815–3823. <https://doi.org/10.1523/JNEUROSCI.2596-12.2013>
- Motta, S.C., Canteras, N.S., 2015. Restraint stress and social defeat: What they have in common. *Physiol. Behav.* 146, 105–110.  
<https://doi.org/10.1016/j.physbeh.2015.03.017>
- Motta, S.C., Carobrez, A.P., Canteras, N.S., 2017. The periaqueductal gray and primal emotional processing critical to influence complex defensive responses, fear

- learning and reward seeking. *Neurosci. Biobehav. Rev., Translational Neuroscience & Mental Disorders: bridging the gap between animal models and the human condition* 76, 39–47.  
<https://doi.org/10.1016/j.neubiorev.2016.10.012>
- Motta, S.C., Goto, M., Gouveia, F.V., Baldo, M.V.C., Canteras, N.S., Swanson, L.W., 2009. Dissecting the brain's fear system reveals the hypothalamus is critical for responding in subordinate conspecific intruders. *Proc. Natl. Acad. Sci.* 106, 4870–4875. <https://doi.org/10.1073/pnas.0900939106>
- Mowrer, O.H., 1947. On the dual nature of learning—a re-interpretation of “conditioning” and “problem-solving.” *Harv. Educ. Rev.* 17, 102–148.
- Oleson, E.B., Gentry, R.N., Chioma, V.C., Cheer, J.F., 2012. Subsecond dopamine release in the nucleus accumbens predicts conditioned punishment and its successful avoidance. *J. Neurosci. Off. J. Soc. Neurosci.* 32, 14804–14808.  
<https://doi.org/10.1523/JNEUROSCI.3087-12.2012>
- Osada, K., Kurihara, K., Izumi, H., Kashiwayanagi, M., 2013. Pyrazine analogues are active components of wolf urine that induce avoidance and freezing behaviours in mice. *PloS One* 8, e61753. <https://doi.org/10.1371/journal.pone.0061753>
- Pavlov, I.P., 2010. Conditioned reflexes: An investigation of the physiological activity of the cerebral cortex. *Ann. Neurosci.* 17. <https://doi.org/10.5214/ans.0972-7531.1017309>
- Piantadosi, P.T., Yeates, D.C.M., Floresco, S.B., 2018. Cooperative and dissociable involvement of the nucleus accumbens core and shell in the promotion and inhibition of actions during active and inhibitory avoidance. *Neuropharmacology* 138, 57–71.  
<https://doi.org/10.1016/j.neuropharm.2018.05.028>
- Pobbe, R.L.H., Zangrossi, H., Blanchard, D.C., Blanchard, R.J., 2011. Involvement of dorsal raphe nucleus and dorsal periaqueductal gray 5-HT receptors in the modulation of mouse defensive behaviors. *Eur. Neuropsychopharmacol. J. Eur. Coll. Neuropsychopharmacol.* 21, 306–315.  
<https://doi.org/10.1016/j.euroneuro.2010.05.004>
- Ramirez, F., Moscarello, J.M., LeDoux, J.E., Sears, R.M., 2015. Active Avoidance Requires a Serial Basal Amygdala to Nucleus Accumbens Shell Circuit. *J. Neurosci.* 35, 3470–3477. <https://doi.org/10.1523/JNEUROSCI.1331-14.2015>
- Rasch, B., Buchel, C., Gais, S., Born, J., 2007. Odor Cues During Slow-Wave Sleep Prompt Declarative Memory Consolidation. *Science* 315, 1426–1429.  
<https://doi.org/10.1126/science.1138581>
- Ribeiro-Barbosa, E.R., Canteras, N.S., Cezário, A.F., Blanchard, R.J., Blanchard, D.C., 2005. An alternative experimental procedure for studying predator-related

- defensive responses. *Neurosci. Biobehav. Rev.* 29, 1255–1263.  
<https://doi.org/10.1016/j.neubiorev.2005.04.006>
- Rozeske, R.R., Jercog, D., Karalis, N., Chaudun, F., Khoder, S., Girard, D., Winke, N., Herry, C., 2018. Prefrontal-Periaqueductal Gray-Projecting Neurons Mediate Context Fear Discrimination. *Neuron* 97, 898–910.e6.  
<https://doi.org/10.1016/j.neuron.2017.12.044>
- Rudy, J.W., 2009. Context representations, context functions, and the parahippocampal-hippocampal system. *Learn. Mem. Cold Spring Harb. N* 16, 573–585. <https://doi.org/10.1101/lm.1494409>
- Schlund, M.W., Brewer, A.T., Richman, D.M., Magee, S.K., Dymond, S., 2015. Not so bad: avoidance and aversive discounting modulate threat appraisal in anterior cingulate and medial prefrontal cortex. *Front. Behav. Neurosci.* 9, 142.  
<https://doi.org/10.3389/fnbeh.2015.00142>
- Servatius, R.J., 2016. Editorial: Avoidance: From Basic Science to Psychopathology. *Front. Behav. Neurosci.* 10. <https://doi.org/10.3389/fnbeh.2016.00015>
- Shin, L.M., Liberzon, I., 2010. The neurocircuitry of fear, stress, and anxiety disorders. *Neuropsychopharmacol. Off. Publ. Am. Coll. Neuropsychopharmacol.* 35, 169–191. <https://doi.org/10.1038/npp.2009.83>
- Skinner, B.F., 1938. *The behavior of organisms: an experimental analysis*, *The behavior of organisms: an experimental analysis*. Appleton-Century, Oxford, England.
- Smith, R., Lane, R.D., 2016. Unconscious emotion: A cognitive neuroscientific perspective. *Neurosci. Biobehav. Rev.* 69, 216–238.  
<https://doi.org/10.1016/j.neubiorev.2016.08.013>
- Souza, R.R., Carobrez, A.P., 2016. Acquisition and expression of fear memories are distinctly modulated along the dorsolateral periaqueductal gray axis of rats exposed to predator odor. *Behav. Brain Res.* 315, 160–167.  
<https://doi.org/10.1016/j.bbr.2016.08.021>
- Spruston, N., 2008. Pyramidal neurons: dendritic structure and synaptic integration. *Nat. Rev. Neurosci.* 9, 206–221. <https://doi.org/10.1038/nrn2286>
- Squire, L.R., Genzel, L., Wixted, J.T., Morris, R.G., 2015. Memory Consolidation. *Cold Spring Harb. Perspect. Biol.* 7, a021766.  
<https://doi.org/10.1101/cshperspect.a021766>
- Stewart, S., Jeewajee, A., Wills, T.J., Burgess, N., Lever, C., 2014. Boundary coding in the rat subiculum. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 369, 20120514.  
<https://doi.org/10.1098/rstb.2012.0514>

- Suvrathan, A., Tomar, A., Chattarji, S., 2010. Effects of chronic and acute stress on rat behaviour in the forced-swim test. *Stress Amst. Neth.* 13, 533–540. <https://doi.org/10.3109/10253890.2010.489978>
- Tovote, P., Esposito, M.S., Botta, P., Chaudun, F., Fadok, J.P., Markovic, M., Wolff, S.B.E., Ramakrishnan, C., Fenno, L., Deisseroth, K., Herry, C., Arber, S., Lüthi, A., 2016. Midbrain circuits for defensive behaviour. *Nature* 534, 206–212. <https://doi.org/10.1038/nature17996>
- van Groen, T., Wyss, J.M., 1992. Connections of the retrosplenial dysgranular cortex in the rat. *J. Comp. Neurol.* 315, 200–216. <https://doi.org/10.1002/cne.903150207>
- Vertes, R.P., Hoover, W.B., Valle, A.C.D., Sherman, A., Rodriguez, J.J., 2006. Efferent projections of reuniens and rhomboid nuclei of the thalamus in the rat. *J. Comp. Neurol.* 499, 768–796. <https://doi.org/10.1002/cne.21135>
- Vianna, D.M., Graeff, F.G., Brandão, M.L., Landeira-Fernandez, J., 2001. Defensive freezing evoked by electrical stimulation of the periaqueductal gray: comparison between dorsolateral and ventrolateral regions. *Neuroreport* 12, 4109–4112. <https://doi.org/10.1097/00001756-200112210-00049>
- Viellard, J., Baldo, M.V.C., Canteras, N.S., 2016. Testing conditions in shock-based contextual fear conditioning influence both the behavioral responses and the activation of circuits potentially involved in contextual avoidance. *Behav. Brain Res.* 315, 123–129. <https://doi.org/10.1016/j.bbr.2016.08.033>
- Wang, J., Bast, T., Wang, Y.-C., Zhang, W.-N., 2015. Hippocampus and two-way active avoidance conditioning: Contrasting effects of cytotoxic lesion and temporary inactivation. *Hippocampus* 25, 1517–1531. <https://doi.org/10.1002/hipo.22471>
- Wood, W., Neal, D.T., 2007. A new look at habits and the habit-goal interface. *Psychol. Rev.* 114, 843–863. <https://doi.org/10.1037/0033-295X.114.4.843>
- Xu, Yuanzhong, Lu, Y., Cassidy, R.M., Mangieri, L.R., Zhu, C., Huang, X., Jiang, Z., Justice, N.J., Xu, Yong, Arenkiel, B.R., Tong, Q., 2019. Identification of a neurocircuit underlying regulation of feeding by stress-related emotional responses. *Nat. Commun.* 10, 3446. <https://doi.org/10.1038/s41467-019-11399-z>
- Yankelevitch-Yahav, R., Franko, M., Huly, A., Doron, R., 2015. The Forced Swim Test as a Model of Depressive-like Behavior. *J. Vis. Exp.* 52587. <https://doi.org/10.3791/52587>