

Le rôle des bactéries dans le filtrage du chlorométhane un gaz destructeur de la couche d'ozone : des souches modèles aux communautés microbiennes de sols forestiers

Pauline Chaignaud

▶ To cite this version:

Pauline Chaignaud. Le rôle des bactéries dans le filtrage du chlorométhane un gaz destructeur de la couche d'ozone : des souches modèles aux communautés microbiennes de sols forestiers. Bactériologie. Université de Strasbourg; Universität Bayreuth, 2016. Français. NNT : 2016STRAJ027 . tel-03336569

HAL Id: tel-03336569 https://theses.hal.science/tel-03336569

Submitted on 7 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





ÉCOLE DOCTORALE DES SCIENCES DE LA VIE ET DE LA SANTÉ

UMR 7156 Génétique Moléculaire Génomique, Microbiologie

THÈSE présentée par :

Pauline CHAIGNAUD

soutenue le : 29 Juin 2016

pour obtenir le grade de : Docteur de l'université de Strasbourg

Discipline : Sciences du vivant

Spécialité : Aspects moléculaires et cellulaires de la biologie

Le rôle des bactéries dans le filtrage du chlorométhane, un gaz destructeur de la couche d'ozone – des souches modèles

aux communautés microbiennes de sols forestiers

Bacteria as chloromethane sinks – from model strains to forest soil communities

THÈSE dirigée par :	Directrice de recherche, Université de Strasbourg, France	
Mme. BRINGEL Françoise	Docteur, Leibnizzentrum für Agrarlandschaftsforschung	
M. KOLB Steffen	(ZALF) und Universität Bayreuth, Allemagne	
RAPPORTEURS : Mme. LAUGA Béatrice M. HORN Marcus	Professeure, Université de Pau, France Docteur, Universität Bayreuth, Allemagne	

AUTRES MEMBRES DU JURY : M. POTIER Serge Mme. KNIEF Claudia

Professeur, Université de Strasbourg, France Professeure, Universität Bonn, Allemagne



École Doctorale des Sciences de la Vie et de la Santé S T R A S B O U R G







Ce travail de thèse est une cotutelle entre l'Université de Strasbourg et l'Université de Bayreuth. Il a été réalisé respectivement dans l'équipe AIME dirigée par le Professeur Stéphane Vuilleumier au sein du laboratoire de Génétique Moléculaire, Génomique et Microbiologie, ainsi qu'au sein du département d'écologie microbienne (EMIC) dirigé par le Professeur Harold Drake à Bayreuth.

Je tiens à remercier les membres de mon jury de thèse ; le Pr. Serge Potier, Dr. Marcus Horn, Pr. Béatrice Lauga et le Pr. Claudia Knief, d'avoir accepté d'évaluer mon travail.

Je remercie également mes 2 directeurs de thèse pour leur encadrement, le Dr. Françoise Bringel et le Dr. Steffen Kolb. Françoise, merci pour tes nombreuses remarques et critiques qui m'ont permis d'avancer et d'aller au bout de mon cursus universitaire. Steffen, merci pour ton accueil à Bayreuth et pour ton aide dans la réalisation de ce projet.

Merci aux membres de l'équipe AIME pour son soutien tout au long de ma thèse et d'autant plus ces derniers mois. Stéphane, merci de m'avoir fait confiance lors de mon stage de master, et montré que je pouvais me dépasser. Sans ce stage, je n'en serai pas là aujourd'hui. Cette expérience n'aurait pas été possible sans ton aide, ton écoute, et ton soutien à toute épreuve.

Un grand merci à Thierry pour m'avoir proposé de faire mon stage de master au sein de l'équipe. Merci pour ta disponibilité, tes conseils et les nombreuses corrections de mon mémoire. Tes blagues sur mon choix de partir en thèse vont me manquer !

Christelle, je ne te remercierai jamais assez pour tout surtout pour tes blagues et clichés sur Lille et le Nord... Je pourrai faire un chapitre complet de remerciements !

Merci à Yousra, ton grain de folie, tes chansons chantées en yaourt et tes prises de kung-fu au milieu de couloir me feront toujours rire !

Merci aux post-docs, et thésards rencontrés pendant toutes ces années dans l'équipe. Sabrina, ma compatriote du Nord, merci pour ton aide, ton écoute et ta bonne humeur. Merci de m'avoir permis de me remémorer des monuments de la chanson française injustement oubliés ; Larusso, Ophélie Winter.... Merci à Bruno, pour ta présence au quotidien (et le soir à table). Qui va te saouler, maintenant, avec des débats qui se terminent toujours en monologue ?! Merci pour ta disponibilité et ton aide. Ludo, merci pour ton aide et surtout ton sourire qui illuminait notre bureau tous les jours...

Merci également aux « anciens » thésards de l'équipe, Fahran et Omniae pour leurs conseils et Louis, Panyu, Elova (et ses Elovades...) ainsi qu'à Emmanuel.

Merci également à Jackson, Jean-Seb, Christian, David et Julien pour leur aide, soutien et les bières après le labo.

Merci à toute l'équipe EMIC pour son accueil et sa bienveillance. Un merci tout particulier à Mareen, pour son aide, ses conseils et sa disponibilité lors des expériences de SIP. Merci aussi à Adam et Sindy pour avoir pris le temps de me faire visiter la ville de Bayreuth.

Enfin, merci à mes proches pour leur présence ces dernières années, mes parents, Charlotte, Antoine, Caro, Brigitte, Lucile et Sophie sans vous je n'en serai pas là aujourd'hui.

Diese Diplomarbeit entstand mit der Unterstützung der Universität Straßburg und der Universität Bayreuth.

Sie wurde realisiert durch das AIME Team von dem Professor des Labors für molekulare Genetik, Genomik und Mikrobiologie Stephane Vuilleumier, sowie Professor Harold Drake, Leiter der Abteilung für mikrobielle Ökologie (EMIC) in Bayreuth.

Ich möchte den Mitgliedern meines Bewertungskomitees danken; Professor Serge Potier, Doktor Marcus Horn, Professor Béatrice Lauga und Professor Claudia Knief, die sich bereiterklärt haben meine Arbeit zu bewerten.

Ich danke auch meinen beiden Betreuern für ihre Führung, Doktor Françoise Bringel und Doktor Steffen Kolb.

Françoise, danke für die vielen Kommentare und die Kritik, die mich weitergebracht haben und es mir ermöglicht haben meinen Studiengang an der Universität zu beenden.

Steffen, vielen Dank mich in Bayreuth willkommen geheißen zu haben und für deine Hilfe an diesem Projekt.

Vielen Dank an die Mitglieder des AIME Teams für ihre Unterstützung während meiner Diplomarbeit und vor allem in den letzten Monaten.

Stéphane, danke für dein Vertrauen in mich während meiner Masterarbeit, das mir zeigte mich selbst zu übertreffen. Ohne dieses Referendariat wäre ich heute nicht hier. Diese Erfahrung wäre nicht möglich gewesen ohne deine Hilfe, dein aufmerksames Ohr und deine unermüdliche Unterstützung.

Ein großes Dankeschön an Thierry für die Einladung meinen Master innerhalb des Teams abzuleisten. Vielen Dank für deine Verfügbarkeit, deine Beratung und zahlreiches "auf die Sprünge helfen" meines Gedächtnisses. Deine Witze über meine Wahl eine Diplomarbeit zu schreiben werden mir fehlen.

Christelle, ich könnte dir niemals genug danken vor allem für deine Witze und Klischees über Lille und den Norden ... Ich könnte ein ganzes Kapitel mit Danksagungen vollschreiben!

Vielen Dank an Yousra, deine Verrücktheit, deine in Joghurt gesungen Lieder und deine Kung-Fu-Vorstellungen inmitten des Flurs, die mich immer zum Lachen bringen werden!

Vielen Dank an die "Post-Docs" und Doktoranden, die sich all die Jahre im Team trafen.

Sabrina, meine Kollegin des Nordens, vielen Dank für deine Hilfe, dein Ohr und deine gute Laune. Vielen Dank, mich an die zu Unrecht vergessenen, französischen Lieder erinnert zu haben ; Larusso, Ophélie Winter

Vielen Dank an Bruno für deine tägliche Anwesenheit (und Abends am Tisch). Wer wird dich jetzt mit Diskussionen nerven, die immer im Monolog enden? Vielen Dank für deine Verfügbarkeit und deine Hilfe.

Ludo, ich danke dir für deine Hilfe und vor allem dein Lächeln, das unser Büro jeden Tag erleuchtete.

Vielen Dank an das "alte" Team von Doktoranden, Farhan und Omniae für ihre Beratung und Louis, Panyu, EloVA (und seine Elovades ...), sowie Emmanuel.

Danke auch an Jackson, Jean-Seb, Christian, David und Julian für ihre Hilfe, Unterstützung und das Bier nach der Arbeit im Labor.

Vielen Dank an das EMIC Team für ihre Gastfreundschaft und Freundlichkeit.

Ein ganz besonderer Dank an Mareen, für ihre Hilfe, Beratung und Verfügbarkeit bei SIP-Experimenten. Vielen Dank ebenfalls an Adam und Sindy, die sich die Zeit genommen haben, mir die Stadt Bayreuth zu zeigen.

Abschließend danke Ich meiner Familie für ihre Anwesenheit in den letzten Jahren, meinen Eltern, Charlotte, Anthony, Caro, Brigitte, Lucile und Sophie. Ohne euch würde ich heute nicht hier stehen.

Table of contents

List of abbreviations	4
List of figures	6
List of tables	10
Chapter 1. Introduction (in English)	10
1. Chlorinated methanes	
2. Chloromethane	13
2.1. Sources of chloromethane	
2.1.1. Abiotic production	14
2.1.2. Biotic production	15
2.2. Chloromethane sinks	22
2.2.1. Abiotic degradation of chloromethane	22
2.2.2. Biotic degradation of chloromethane	22
2.3. Global chloromethane balance in soil	25
3. Methylotrophy	27
3.1. Methylotrophic pathways	28
3.2. Bacterial degradation of dichloromethane (CH ₂ Cl ₂)	
3.2.1. Dichloromethane utilization in anoxic conditions	
3.2.2. Dichloromethane utilization in oxic conditions	
3.3. Bacterial degradation of chloromethane	34
3.3.1. Degradation under anoxic conditions	34
3.3.2. Degradation in oxic conditions	34
4. The <i>Methylobacterium</i> model	40
4.1. <i>Methylobacterium</i> , a model for methylotrophy studies	41
4.2. Genetic tools available for <i>M. extorquens</i>	44
4.3. Global approaches in <i>M. extorquens</i>	44
4.3.1. Genomic studies in <i>M. extorquens</i>	44
4.3.2. Transcriptomic studies in <i>M. extorquens</i>	47
4.3.3. Proteomic studies in <i>M. extorquens</i>	47
4.3.4. Metabolomic studies in <i>M. extorquens</i>	48
5. PhD thesis aims and questions	49
Chapter 1. Introduction (en français)	55
1. Les méthanes chlorés	55
2. Le chlorométhane	58
2.1. Les sources de chlorométhane	59
2.1.1. Production abiotique	59
2.1.2. Production biotique	60
2.2. Les puits de chlorométhane	66
2.2.1. Dégradation abiotique du chlorométhane	66
2.2.2. Dégradation biotique du chlorométhane	67

Table of contents

2.3. Le budget global du chlorométhane du sol	70
3. La méthylotrophie	72
3.1. Les voies de la méthylotrophie	73
3.2. La dégradation bactérienne du dichlorométhane (CH ₂ Cl ₂)	78
3.2.1. Utilisation du dichlorométhane en condition anoxique	78
3.2.2. Utilisation du dichlorométhane en condition oxique	78
3.3. La dégradation bactérienne du chlorométhane	79
3.3.1. Dégradation du chlorométhane en condition anoxique	79
3.3.2. Dégradation du chlorométhane en condition oxique	80
4. Le modèle <i>Methylobacterium</i>	85
4.1. Methylobacterium, bactérie modèle pour l'étude de la méthylotrophie	86
4.2. Outils génétiques disponible chez <i>M. extorquens</i>	89
4.3. Approches globales chez <i>M. extorquens</i>	89
4.3.1. Etudes génomiques chez <i>M. extorquens</i>	89
4.3.2. Approches de transcriptomique chez <i>M. extorquens</i>	92
4.3.3. Etudes de protéomique chez <i>M. extorquens</i>	93
4.3.4. Etudes de métabolomique chez <i>M. extorquens</i>	94
5. Objectif et questions du projet de thèse	95
Chapter 2. Methodological section	101
1. Methods used to study <i>M. extorguens</i> model strains	
1.1. Aerobic methylotrophic growth	
1.2. RNA preparation	
1.2.1. DNA removal	
1.2.2. rRNA removal	
1.2.3. RNA quality control	
1.3. Construction of the directional cDNA bank	104
2. Methods used to study chloromethane-degrading bacterial communities in for	est soils
	106
2.1. Soil sampling and preincubation	107
2.2. Stable isotope probing (SIP): the [¹³ C]-chloromethane labeling step	107
2.3. DNA extraction from soil	109
2.4. Separation of isotopically-labeled DNA by isopycnic density gradient	109
2.5. Metagenomics of PCR-amplified markers	111
2.5.1. Design of new primers targeting environmental chloromethane dehalog	genase
CmuA-encoding gene	112
2.5.2. Design of new primers targeting environmental methanol dehydrogena	ses
MDH-encoding genes	
	116
3. Illumina sequencing	118
4. Bioinformatics	120
4.1. MicroScope platform	120

	4.2.	Transcriptome Analyses based on MAssive sequencing of RnAs: RNAseq datab	ase
			122
Cha	pter 3.	RNAseq study of chlorinated compound utilization in <i>M. extorquens</i>	.129
1.	Intr	oduction	129
2.	Arti	cle	133
3.	Sup	plemental data	185
	3.1.	Defining the core genome	185
	3.2.	RNAseq data of the variable genome only shared by dechlorinating strains	185
	3.3.	RNAseq data of plasmid pCMU01-borne gene with chromosomal homologs	189
	3.4.	RNAseq data of genes with differential sense/antisense transcript abundance	191
	3.5.	RNAseq data of formate dehydrogenase subunits encoding-genes	195
4.	Con	clusion	196
Cha	pter 4.	Ecotypes of microbial chloromethane utilizers in a forest soil	.199
1.	Intr	oduction	199
2.	Mat	erials and methods	203
	2.1.	Study site and sampling	203
	2.2.	Soil activation and microcosm set-up	203
	2.3.	Chemical analyses	204
	2.4.	[¹³ C]-CO ₂ analysis	205
	2.5.	Nucleic acid extraction and RNA removal	205
	2.6.	DNA fractionation	206
	2.7.	PCR amplification	206
	2.8.	DNA sequencing	209
	2.9.	Analysis of the 16S rRNA gene	209
	2.10.	Analysis of functional genes	210
	2.11.	Identification of labeled OTUs	210
	2.12.	Statistical and phylogenetic analyses	211
3.	Res	ults	211
	3.1.	Microcosm setup for detection of CH_3CI -utilizing methylotrophs in forest soil	211
	3.2.	Mineralization and assimilation of chloromethane in forest soil microcosms	213
	3.3.	Taxa associated with chloromethane utilization in forest soils	213
	3.4.	Diversity of cmu pathway chloromethane utilizers in forest soil microcosms	220
	3.5.	Diversity of methanol utilizers in forest soil microcosms	221
4.	Disc	cussion	224
	4.1.	New insights into CH ₃ Cl utilizers in forest soil	225
	4.2.	cmuA OTUs and metabolic diversity in the environment	228
	4.3.	Chloromethane-associated methylotrophy	229
5.	Con	clusion	229
6.	Sup	plemental materials	230
Cha	pter 5.	Conclusions and perspectives	.239

Table of contents

1. News insights in chloromethane utilization deduced from the study of the model strain
M. extorquens CM4240
1.1. In depth study of the bacterial adaptive response to chloromethane utilization240
1.2. Search of chloromethane-utilization-dependent regulatory elements241
2. Identification of chloromethane-utilizing ecotypes in forest soil microcosms246
3. Assessing the diversity of chloromethane-utilizing pathway in soil
4. Need for new functional biomarkers for evaluation of the microbial chloromethane sinks
Bibliography
Apendices
Preliminary version of the thesis summary validated by Université de Strasbourg (Long version in
French)
PhD thesis abstract and key words281
Resumé de la thèse et mots-clés280
Zusammenfassung dissertation und stichworte

List of abbreviations

аа	Amino acid	HOL	Harmless to ozone layer
ADH:	Alcohol dehydrogenase	IARC	International Agency for Research
ADN/DNA	Acide desoxyribonculéique		on Cancer
	/deoxyribonucleic acid	IS	Insertion sequence
ADP	Adenosine diphosphate	kb	Kilo base pair
ANAH	Adenine Nucleotide Alpha	KEGG	Kyoto Encyclopedia of Genes and
	Hydrolase		Genomes
ARN/RNA	Acide ribonucléique	log2fc	log2 fold-change
	/ribonucleic acid	M3	Mineral media for
ATP	Adenosine triphosphate		Methylobacterium
BLAST	Basic local alignment search tool	MADH	Methylamine dehydrogenase
bp	Base pair	MaGe	Magnifying Genomes
BSA	Bovine serum albumin	Mb	Mega base pair
cDNA	Complementary deoxyribonucleic	Mchl	Methylobacterium
	acid		chloromethanicum
CDS	Coding DNA sequence	MDH	Methanol dehydrogenase
CM	Chlorométhane	MEGA	Molecular Evolutionary Genetics
	/chloromethane		Analysis
Cmu	Chloromethane utilization	METDI	Methylobacterium extorquens
CoA	Coenzyme A		dichloromethane
COV/VOC	Composé organique volatile	MNNG	N-methyl-N'-nitrosoguanidine
	/volatile organic compound	mRNA	Messenger Ribonucleic acid
CsCl	Cesium chloride	MS	Methanesulfonate d'éthyle
СТАВ	Cetyl trimethylammonium bromide	NAD(H)	Nicotinamide adenine dinucleotide
DCM	Dichlorométhane	NADP (H)	Nicotinamide adenine dinucleotide
	/dichloromethane		phosphate
ddNTP	Dideoxynucleotide triphosphate	NCBI	National Center for Biotechnology
DEPC	Diethyl pyrocarbonate		Information
DESeq	Differential expression analysis for	nd	Not determined
	sequence count data	NGS	Next generation sequencing
DNase	Deoxyribonuclease	NJ	Neighbour-joining
dNTP	Deoxyribonucleotide triphosphate	NMDS	Non-metric multidimensional
DMB	Dimethylbenzimidazole		scaling
ECD	Electron capture detector	nt	Nucleotide
EDTA	Ethylenediaminetetraacetic acid	OD	Optical density
FDH	Formate dehydrogenase	OTU	Operational taxonomic unit
FDR	False discovery rate	PCR	Polymerase chain reaction
GC	Gas chromatography	PEG	Polyethylene glycol
GC-MS	Gas chromatography-mass	PHB	Poly-beta-hydroxybutyrate
	spectometry	PICRUST	Phylogenetic investigation of
GEI	Genomic islet		communities by reconstruction of
GST	Glutathion S-transferase		unobserved states
HEPES	4-(2-hydroxyethyl)-1-	PkGDB	Prokaryotic Genome DataBase
	piperazineethanesulfonic acid		

рММО	Particulate methanol	SAM	S-adenosyl L-methionine
	monooxygenase	SBS	Sequencing by synthesis
ppb	Part per billion	SIP	Stable isotope probing
PPi	Pyrophosphate inorganique	sMMO	Soluble methanol monooxygenase
ppt	Part per trillion	SMRT	Single molecule real time
Pptv	Part per trillion by volume	ssDNA	Single-stranded deoxyribonucleic
PQQ	Pyrroloquinoline quinone		acid
qPCR	Quantitative polymerase chain	STAMP	Statistical analysis of metagenomic
	reaction		profiles
RefSeq	Reference Sequence collection	TAMARA	Transcriptome analyses based on
RIN	RNA integrity number		massive sequencing of RNAs
RNase	Ribonuclease	TCA	Tricarboxylic acid cycle
RNAseq	Ribonucleic acid sequencing	TCD	Thermal conductivity detector
RPM	Rotation per minute	TCM	Tetrachloromethane
rRNA	Ribosomal ribonucleic acid	TE	Tris-EDTA buffer (10 mM Tris-HCl, 1
RT-qPCR	Reverse transcription-quantitative		mM disodium EDTA, pH 8.0)
	polymerase chain reaction	UV	Ultraviolet
RuBP	Ribulose biphosphate	WGS	Whole genome sequencing
RuMP	Ribulose monophosphate		
SAH	S-adenosyl L-homocystein		

List of figures

Chapter 1 (in English) Introduction

igure 1.1. Ozone production and degradation reactions in the atmosphere	13
-igure 1.2. Abiotic chloromethane production in soils	15
igure 1.3. Chloromethane production in Arabidopsis thaliana	16
-igure 1.4. Cross-section of forest soil top revealing horizons	17
igure 1.5. Chloromethane production by the <i>Phellinus pomaceus</i> fungus	18
igure 1.6. Volatile organic compounds emitted by soils	20
-igure 1.7. Chloromethane dosage using a bioreporter strain	21
igure 1.8. Chloromethane degradation reactions by fungi	25
-igure 1.9. Seasonal chloromethane fluxes in forest soil	26
igure 1.10. Scheme of different methylotrophic pathways in <i>M. extorquens</i>	30
-igure 1.11. Carbon assimilation in <i>M. extorquens</i>	31
igure 1.12. Dichloromethane degradation by the hydrolytic pathway of glutathione S-	
transferase	33
igure 1.13. Proposed mechanism of chloromethane oxidation in Methylomicrobium album E	3G8
	35
igure 1.14 The cmu pathway of Methylobacterium extorquens CM4	37
igure 1.15 The chloromethane utilization pCMU01 plasmid in <i>M. extorquens</i> CM4	38
igure 1.16. Comparison of <i>cmu</i> gene organization in bacteria harboring gene <i>cmuA</i>	40
igure 1.17. 2D gel picture of protein extracts of <i>M. extorquens</i> CM4 grown with chlorometh	ane
or with methanol	48
igure 1.18. Sinks and emissions of chloromethane and methanol in forest soil	50

Chapitre 1 (en français) Introduction

Figure 1.1. Réactions de formation et de destruction de l'ozone dans l'atmosphère58
Figure 1.2. Formation abiotique du chlorométhane dans les sols60
Figure 1.3. Production de chlorométhane chez Arabidopsis thaliana61
Figure 1.4. Organisation schématique du sol forestier62
Figure 1.5. Production de chlorométhane par le champignon Phellinus pomaceus63
Figure 1.6. Les composés organiques volatiles émis dans les sols65
Figure 1.7. Souche bio-rapportrice pour la détection du chlorométhane66
Figure 1.8. Réactions de dégradation du chlorométhane par les champignons70
Figure 1.9. Flux du chlorométhane dans un sol forestier en fonction des saisons71
Figure 1.10. Schéma synthétique des différentes voies de la méthylotrophie chez <i>M. extorquens</i>
75
Figure 1.11. Assimilation du carbone chez <i>M. extorquens</i> 76
Figure 1.12. Dégradation du dichlorométhane par la voie hydrolytique de la glutathion79
Figure 1.13. Mecanisme proposé d'oxydation du chlorométhane par Methylomicrobium album
BG881
Figure 1.14 Voie cmu chez Methylobacterium extorquens CM482

Figure 1.15. Plasmide pCMU01 d'utilisation du chlorométhane chez M. extorquens CM483
Figure 1.16. Comparaison de l'organisation des gènes cmu chez des bactéries contenant le gène
<i>cmuA</i> 85
Figure 1.17. Image d'un gel 2D des protéines extraites après croissance de <i>M. extorquens</i> CM4
en présence de chlorométhane ou de méthanol94
Figure 1.18. Emissions et consommations de chlorométhane et de méthanol dans les sols
forestiers96

Chapter 2 Methodological section

Figure 2.1. Electrophoregram of RNAs extracted from <i>M. extorquens</i> CM4 growing with	
chloromethane	103
Figure 2.2. Directional cDNA bank construction	105
Figure 2.3. Validation of cDNA bank constructions with the 2100 Bioanalyzer	106
Figure 2.4. Overview of the stable isotope probing experiment	108
Figure 2.5. Material required for ultracentrifugation and fractionation	111
Figure 2.6. Comparing two sets of primers targeting environmental chloromethane	
dehalogenase CmuA-encoding gene	115
Figure 2.7. PCR amplification of methanol dehydrogenase-like MxaF and XoxF encoding ge	nes
	116
Figure 2.8. Schematic representation of Illumina 'Genome Analyser' flow cell for high	
throughput sequencing	119
Figure 2.9. A screen capture of the "MicroScope" web platform	121
Figure 2.10. A screen capture of the "mapping overview" data available in "TAMARA" web	
platform	123
Figure 2.11. A screen capture of the "Raw read count" data available in TAMARA web plat	form
	124
Figure 2.12. A screen capture of the "Analysis" interface available in TAMARA web platform	n.125

Chapter 3 RNAseq study of chlorinated compound utilization in *M. extorquens*

Figure 3.1. Aerobic C ₁ dehalogenases	129
Figure 3.2. Adaptation to growth with chlorinated methanes as the sole carbon and energy	
source	132

Chapter 4 Ecotypes of microbial chloromethane utilizers in a forest soil

Figure 4.1. Targeted functional genes in methylotrophic utilization of chloromethane and	
methanol	202
Figure 4.2. Carbon mineralization in the SIP experiment	212
Figure 4.3. Clustering analysis of 16S rRNA gene amplicon sequence datasets from all	
microcosms	215
Figure 4.4. Taxonomic affiliation of labeled family-level 16S rRNA OTUs	217
Figure 4.5. Phylogenetic relationships between partial sequences of the 16S RNA gene of	
labeled OTUs and of reference chloromethane-utilizing strains	219

Figure 4.6. Affiliation and relative abundance of <i>cmuA</i> OTUs	220
Figure 4.7. Affiliation and relative abundance of MDH OTUs	222
Figure 4.8. Affiliation and relative abundance of <i>mch</i> OTUs	223

Chapter 5 Conclusions and perspectives

Figure 5.1. Overview of research avenues to assess the role of bacteria as chloromethane	
emission filters in forest soil	239
Figure 5.2. Central questions for future in lab experiments with model strains	242
Figure 5.3. Exploration of chloromethane-dependent regulation mechanisms in model strai	ins
	245

List of tables

Chapter 1 (in English) Introduction

Table 1.1. Chlorinated methane utilizations and characteristics	.11
Table 1.2. Chlorinated methanes in the troposphere	.12
Table 1.3. Estimation of chloromethane sources in the atmosphere	.14
Table 1.4. Estimation of atmospheric chloromethane sinks	.22
Table 1.5. Chloromethane-utilizing bacterial isolated from contrasting environments	.23
Table 1.6. Characteristics of the first Methylobacterium sequenced genomes	.43
Table 1.7. "Omics" studies in <i>M. extorquens</i> strains	.46

Chapitre 1 (en français) Introduction

Tableau 1.1. Propriétés et utilisations des méthanes chlorés	.56
Tableau 1.2. Les méthanes chlorés dans la troposphère	.57
Tableau 1.3. Estimation des sources de chlorométhane dans l'atmosphère	.59
Tableau 1.4. Estimation des puits de chlorométhane dans l'atmosphère	.66
Tableau 1.5. Souches bactérienne chlorométhane-dégradantes isolées de différents	
environnements	.68
Tableau 1.6. Caractéristiques des premières souches de Methylobacterium dont le génome a	
été séquencé	.88
Tableau 1.7. Etudes «omiques» réalisées dans des souches de <i>M. extorquens</i>	.92

Chapter 2 Methodological section

Table 2.1. PCR amplification program to validate DNA removal	102
Table 2.2. Set of primers available for <i>cmuA</i> amplification in the environment	113
Table 2.3. Primer sets available for <i>mxaF_xoxF</i> amplification for NGS analysis in the	
environment	117
Table 2.4. TAMARA project DehaloRNAseq: sample name reference	122

Chapter 3 RNAseq study of chlorinated compound utilization in *M. extorquens*

Table 3.1. RNAseq data of genes shared exclusively by <i>M. extorquens</i> CM4 and DM4	.186
Table 3.2. RNAseq data for pCMU01-borne genes with chromosomal homologs	.189
Table 3.3. RNAseq data of genes with differential sense/antisense transcript abundance	.193
Table 3.4. RNAseq data of formate dehydrogenase subunits encoding-genes	.195

Chapter 4 Ecotypes of microbial chloromethane utilizers in a forest soil

Table 4.1. Microcosm setup and one-carbon supplementation	204
Table 4.2. Biomarker gene primer sequence, PCR amplicon size and OTU occurrence	208
Table 4.3. Number of filtered amplicon sequences obtained from heavy DNA fractions	214
Table 4.4. Diversity indices for 16S rrnA amplicon sequences obtained from heavy and light	t DNA
fractions	216
Table 4.5. Dominant genera within labeled family-level [¹³ C] OTUs	218

Table S4.1. Summary of sequence data obtained in the SIP experiment	183
Table S4.2. Relative abundance (%) and labeled status of functional gene OTUs in heavy	DNA
fractions of [¹³ C]-labeled microcosms	184

Chapter 1 Introduction

Chapter 1. Introduction (in English)

This chapter begins with the general information on chlorinated methanes and chloromethane in particular. Chlorinated methane chemical properties and industrial use as well as their impact on human health and the environment will be described. The following paragraphs are dedicated to chloromethane, including its production (sources) and consumption (sinks), with a focus on the role of microorganisms. Knowledge on microbial degradation of chloromethane will be presented in the context of one carbon compound metabolism and the genetics of methylotrophic model strains. The last paragraph will explain the aims and objectives of my PhD thesis.

1. Chlorinated methanes

Chlorinated methanes are one-carbon organic compounds with one or more chlorine atoms. A carbon atom has four covalent bonds. Four chlorinated methanes are found: chloromethane (CH₃Cl also called methyl chloride), dichloromethane (CH₂Cl₂), chloroform (CHCl₃) and tetrachloromethane (CCl₄). Of those, only chloromethane is in a gaseous state at room temperature and one atmosphere pressure. Other chlorinated methanes are liquid and colorless (Huang *et al.*, 2014). These compounds have different physical and chemical properties and are, since the 20th century, produced and used in large quantities by the industry (Table 1.1). Chlorinated methanes are also produced naturally. Natural production of chloroform and dichloromethane are estimated at 0.56 and 0.25 Tg.year⁻¹, respectively (Cox *et al.*, 2003; Xiao, 2008). However, chloromethane is the most abundant chlorinated methane found in the atmosphere with 2.8 Tg.year⁻¹ (Ruecker *et al.*, 2014) (Table 1.2).

Compound	State	Water solubility (g.L ⁻¹)	Effect on health ^a	Pollutant ^b	Industrial utilization ^c
	colorless		group 3	no	silicone production,
chloromethane	gas	5.3			construction
	0				products
	colorless liquid	13.0	group 2B	yes	paint stripping,
dichloromethane					pharmaceuticals
					products
chloroform	colorless 8. liquid	8.0	group 2B	yes	solvents, chemicals
					synthesis
					pharmaceuticals
tetrachloromethane	colorless 0.8 liquid	0.8	group 2B	yes	products,
		0.0			cleaning products,
					agrochemicals

Table 1.1. Chlorinated methane utilizations and characteristics

^a Data from the International Agency Research against Cancer (IARC). Group 2B contains compounds probably carcinogenic to humans. Group 3 consists of compounds not classified such as carcinogenic for humans (Huang *et al.*, 2012).

^b Compounds recognized as pollutants by the United States, China. Dichloromethane and chloroform are also recognized as pollutant by the European Union (Huang *et al.*, 2014). ^c Data from Frascari *et al.*, 2015

Chlorinated methanes are present in soil, water and air (http://www.epa.gov). With the exception of chloromethane, they have been classified by the International Agency for Research on Cancer (IARC) as possibly carcinogenic, and are considered as pollutants in many countries (Table 1.1). The impact of chlorinated compounds on the environment has been extensively documented (Harper, 2000).

	Troposphere		Flow (Gg. year ⁻¹)		
Conc (nntv)		Half-life		Conc. (ppty)	
	•••••• (PP•••)	(year)		(PP)	
chloromethane	$550\pm30^{\text{a}}$	1.3	chloromethane	$550\pm30^{\text{a}}$	
dichloromethane	25 ± 5^{d}	0.38	dichloromethane	25 ± 5^{d}	
chloroform	15 ± 5^{c}	0.41	chloroform	15 ± 5^{c}	
tetrachloromethane	95 ± 5^{f}	26	tetrachloromethane	95 ± 5^{f}	

Table 1.2. Chlorinated methanes in the troposphere

From S. Roselli PhD thesis, 2009

^a Montzka and Fraser, 2003

^b McCulloch *et al.*, 2003

^c O'Doherty *et al.*, 2001

^d Schauffler, 2003

^e Kurylo and Rodriguez, 1998

^f Prinn *et al.*, 2000

^g UNEP, 2005

In the stratosphere, chlorinated methanes are activated by ultraviolet radiation. This process results in the formation of radical Cl-, which will contribute to the depletion of the ozone layer by catalyzing the formation of O_2 (Figure 1.1). It is estimated that one chlorine atom can destroy 100,000 ozone molecules (http://www.epa.gov/ozone/science/process.html). An example of this impact is the formation of a hole in the ozone layer over the Antarctic every spring (Clerbaux *et al.*, 2007).



Figure 1.1. Ozone production and degradation reactions in the atmosphere In the presence of chlorinated methanes (CHxClx), natural reactions of ozone (O₃) are disrupted. UVA between 315 nm and 400 nm and UVB radiation from 280 nm to 315 nm (Figure modified from http://www.cec.org).

Because of their impact on the environment, the use of halogenated compounds has been regulated by the Montreal Protocol in 1987, thus significantly reducing their use in the industry, up to a total cessation of production of compounds such as carbon tetrachloride in 1995. This resulted in the augmentation of the share of chloromethane, the chlorinated C₁, compound the most produced naturally in the environmental impact of chlorinated methanes. Indeed, from chlorinated methanes found in the troposphere, chloromethane is the most abundant (Table 1.2). Today it is estimated that 16 % of the degradation of the ozone layer is due to chlorinated compounds (Montzka *et al.*, 2011).

2. Chloromethane

The overall budget of chloromethane, or its concentration in the atmosphere, is determined by the sum of its production sources, subtracted from all sinks dissipation. The diversity of abiotic and biotic sources and sinks of chloromethane is detailed below. This budget remains uncertain mainly because of the unknown quantitative impact of bacteria on the biotic filtering of chloromethane emitted from terrestrials and other environments.

2.1. Sources of chloromethane

Total annual production is estimated at 2.8 Tg (Ruecker *et al.*, 2014; Sailaukhanuly *et al.*, 2014). As more than 5,000 halogenated substances, chloromethane is produced predominantly

naturally. The natural production of chloromethane based on abiotic and biotic sources (Table 1.3). Its low production of anthropogenic origin is due to the burning of fossil fuels such as coal, the combustion of municipal wastes, and other industrial activities (Table 1.3, McCulloch and Aucott, 1999).

Type	Estimation	Low/high values
туре	(Gg. year ⁻¹) ^a	(Gg. year ⁻¹) ^b
Plant senescence	1,800 ^d	30/2500
Forest fires	911	655/ 1,125
Tropical plants	910	820/ 8,200
Oceans	600	325/ 1,300
Salt marshes	170	65/ 440
Fungi	160	43/ 470
Combustion of fossil organic matters	105	5/ 205
Incineration	45	15/75
Wetlands	40	6-270 ^c
Industrial	10	10
Rice fields	5	2,4/ 4,9
Total sources	4,746	2,019/ 1,4378

 Table 1.3. Estimation of chloromethane sources in the atmosphere

^a Data from Keppler *et al.*, 2005

^b Data from Clerbaux *et al.*, 2007

^c Data from Montzka and Fraser, 2003

^d Data from Montzka *et al.*, 2011

2.1.1. Abiotic production

Forest fires are the largest source of abiotic chloromethane (Sailaukhanuly *et al.*, 2014). The produced chloromethane is released into the atmosphere, but is also found in the soil surface, where abiotic processes of the production of halogenated methanes have also been characterized (Keppler *et al.*, 2005). Abiotic production of chloromethane by redox reactions (Figure 1.2) and substitution reactions were demonstrated (Hamilton, 2003; Keppler *et al.*, 2005; Wishkerman *et al.*, 2008) although their share in budget chloromethane was not

assessed. The oceans represent the second source of abiotic chloromethane (Xiao, 2008; Hu et al., 2010).



Figure 1.2. Abiotic chloromethane production in soils

Halogenated alkyl formation by the reduction of Fe (III) with the organic material in the presence of chloride ions in soils (from Keppler *et al.*, 2003).

2.1.2. Biotic production

The main production of chloromethane is the result of biotic processes in plants (Hamilton, 2003; Saito *et al.*, 2008), fungi (Moore *et al.*, 2005), seaweed (Traunecker *et al.*, 1991) or salt marshes (Rhew *et al.*, 2003) (Table 1.3). Terrestrial ecosystems contribute significantly to this production, with more than 50% of emissions of chloromethane derived from them (Redeker and Kalin, 2012). In the environment, the presence of chloride ions is highly variable and may influence the emission rates of chloromethane. The first results of a possible correlation are mentioned in the literature and are described below, as well as the current knowledge of chloromethane production by plants and in soils. Also explained below is the emission rate of chloromethane as compared to other Volatile Organic Compound (VOC) and the development of a bio-reporter for its quantification.

Production by plants

The genetic mechanism of production of chloromethane was characterized in detail in the model plant *Arabidopsis thaliana* (Rhew *et al.*, 2003). This study shows the production of chloromethane but also bromomethane (CH₃Br) and iodomethane (CH₃I), by the action of a S-Adenosyl Methionine (SAM)-dependent methyltransferase, named HOL for "Harmless to Ozone Layer". The HOL gene is conserved in plants, suggesting that the production of chloromethane is a ubiquitous process of plants (Rhew *et al.*, 2003; Nagatoshi and Nakamura,

2007). Chloromethane is produced by the transfer of the methyl group of the SAM to a chloride ion (Cl⁻) (Figure 1.3). The SAM is involved in numerous synthesis reactions of lignin, a component of the plant cell wall (Campbell and Sederoff, 1996). Rhew *et al.*, 2003 has shown that the methyltransferase coded by the HOL gene catalyzes the methylation of chloride, and is dependent on SAM.



Figure 1.3. Chloromethane production in *Arabidopsis thaliana* (Figure obtained and modified from the thesis manuscript of Muhammad Farhan UI Haque, May 2013).

In fact, different studies have demonstrated the existence of plants that emit chloromethane. For example, of 187 plants studied in a subtropical region, 33 species belonging to different families emit chloromethane (Yokouchi *et al.*, 2007). Similarly, the study of chloromethane fluxes in a tropical forest showed that 25 out of 117 plants studied emit chloromethane (Saito *et al.*, 2008). The most important emissions were observed in fern species (*Osmunda banksiifolia, Cibotium balometz, Angiopteris palmiformis*), and in halophile plants (*Vitex rotundifolia, Vitex trifolia* and *Excoecaria agalloch*) with emission rates in the order of 1 µg of chloromethane.g⁻¹ of dry matter. h⁻¹. Nevertheless, the capacity of chloromethane production depends more on the plant species than on the family (Yokouchi *et al.*, 2007).

Production of chloromethane by the soil and decomposing vegetable matter
 Decomposing plants are the biggest source of chloromethane (Hamilton, 2003; Montzka *et al.*, 2011). Production of chloromethane is mostly found in the O horizon of soils and seems
 to be insignificant in lower horizons (Figure 1.4) (Redeker and Kalin, 2012).



Figure 1.4. Cross-section of forest soil top revealing horizons

The forest soil is divided into different horizons: O, A, E, B, C, and R. Horizon O is mostly composed of decomposing organic matter; A is equally composed of organic and mineral matter; E is a leached horizon, where soluble nutrients are no longer present due to precipitation or irrigation; B is a horizon rich in minerals from the leaching of horizons A and E; C is a non-consolidated underground horizon; finally, E is a rocky horizon, composed of rocks such as granite or limestone. Horizons O and A are biologically active and highly oxygenated, while the others are poor in organic matter and have lower microbial activity. Within the framework of my thesis project, an experiment on the forest soil was performed at the level of the O horizon.

(Figure modified from http://www.ctahr.hawaii.edu/mauisoil/a_profile.aspx)

Soil is an environment rich in chloride ions, and the bioavailability of these is essential to the formation of chloromethane. However, the chloride ion concentration in the soil is difficult to estimate due to the intervention of several processes and mechanisms such as the flow between the vegetation and soil, alterations of the rock, or the decomposition of organic matter (Bastviken *et al.*, 2007; Öberg, 2003). It has been observed that a part of the chloride ions is considered non-bioavailable because of its absorption by vegetation or microorganisms (Bastviken *et al.*, 2007). Chloride ions also occur in tree trunks and fungi that colonize the wood utilize them for the chloromethane synthesis (Watling and Harper, 1998). The amount of Cl⁻ in the wood of temperate forest species is estimated to vary between 2.4 and 123 mg.kg⁻ ¹ of dry matter (Watling and Harper, 1998).

Studies of fungi have demonstrated that members of the *Hymenochaetaceae* are able of reemitting a proportion of the chloride ions found in their substrates in the form of chloromethane (Harper and Kennedy, 1986). In fact, decomposing wood is colonized by fungi that, by means of a group of peroxidases, degrade lignin, a major component of wood, into SAM. The SAM and the Cl⁻ ions present in the medium forms chloromethane, a reaction mediated by a methyltransferase (Harper, 2000; Anke and Weber, 2006). Such chloromethane production mechanism has been studied and described in the fungus *Phellinus tuberculosus* (White, 1982; Wuosmaa and Hager, 1990); however, the most common example is that of *Phellinus pomaceus* (Figure 1.5), which is capable of re-emitting as chloromethane 90% of the chloride of its degraded substrates (Harper *et al.*, 1988). This chloromethane production occurs when a Cl⁻ ion is transferred to the methyl group of the SAM (Figure 1.5).



Figure 1.5. Chloromethane production by the *Phellinus pomaceus* **fungus** (Figure modified from Harper, 2000)

Chloromethane production by fungi associated to the root systems of forest soils exists, as well. Roots represent an important source of Volatile Organic Compounds (VOCs) (Lin *et al.*, 2007). In the study carried out by Redeker *et al.*, 2004, 9 ectomycorrhizal fungi isolates, among them *Cenococcum geophilum*, one of the most widespread and common species, (Trappe, 1977; Massicotte *et al.*, 1992), demonstrated their ability to produce halogenated methanes, chloromethane being one of them. They produce between 0.003 µg and 65 µg of

chloromethane per gram of dry matter (Redeker *et al.*, 2004). However, it is difficult to separately evaluate the role of the root system, since it is closely tied to the microbial system, which itself influences the flow of VOCs in the soil (Peñuelas *et al.*, 2014). In fact, 95% of plants' short roots are colonized by fungi which are located close to communities of mycorrhiza helper bacteria that aid fungi in the rhizosphere, notably during processes of mycorrhizal development (Frey-Klett *et al.*, 2007; Bonfante and Anca, 2009; Rigamonte *et al.*, 2010).

Chloromethane is emitted at lower rates than other VOCs

Plants re-emit a proportion of fixed as VOCs, methanol being among them (Vorholt, 2012; Bringel and Couée, 2015). These compounds are emitted into the atmosphere or into the soil, where they can be directly assimilated by microorganisms (Schade and Goldstein, 2001). Thus, the largest part of atmospheric methanol is linked to the production by plants (Jacob, 2005; Galbally and Kirstine, 2002). Other VOCs are produced by plants, among which there are aromatic compounds and halogenated compounds (Guenther *et al.*, 2012; Forczek *et al.*, 2015). The plant production of 4 chlorinated methanes has been described, even though chloromethane is the most produced, with a production of 0.8 to 8 Tg.year⁻¹ (Clerbaux *et al.*, 2007). This production is lower than that of methanol (50 - 132 Tg. year⁻¹) (Jacob, 2005; Forczek *et al.*, 2015).

In the case of forest soils, chloromethane is far from being the only VOC produced (Keppler *et al.*, 2005; Redeker and Kalin, 2012; Peñuelas *et al.*, 2014). Soil is a heterogeneous complex ecosystem, an assembly of complex microenvironments (Peñuelas *et al.*, 2014) with an huge microbial diversity. Bacterial and fungal diversity can be evaluated by the number of different species, respectively, 10⁵ and 10⁶ species per gram of soil (Peñuelas *et al.*, 2014). Numerous VOCs are emitted by these microorganisms (Figure 1.6). This production is directly related to the availability of substrates and of growth conditions (Kai *et al.*, 2010; Blom *et al.*, 2011). A large part of emitted VOCs comes from microbial degradation of plant litter (dead plant matter), of the decomposition of organic matter (Kögel-Knabner, 2002; Peñuelas *et al.*, 2014), as well as from root exudates.



Figure 1.6. Volatile organic compounds emitted by soils

Bacterial emissions are marked in red, and fungal emissions are marked in blue (modified from Peñuelas *et al.,* 2014). The complete inventory of these VOCs is listed in a database (http://bioinformatics.charite.de/mvoc/). The red arrow marks the category of VOCs in which chloromethane is found.

In general, bacteria produce more alkenes, alcohols, ketones, and terpenes, while fungi produce more alcohols, benzenoids, acids, and aldehydes (Figure 1.6). Chlorinated compounds are produced less, and this production in the soil is more important in fungi than in bacteria (Figure 1.6). The "chloride" group includes chloromethane (indicated with a red arrow).

Chloromethane dosage using a bioreporter strain

The quantification of the low amounts of chloromethane produced by plants and soils (in the order of ppt) is difficult using classical approaches of gas chromatography (GC) dosage that is not sensitive enough (Figure 1.7). In the laboratory in Strasbourg, with the goal of evaluating chloromethane emissions by plants, *Methylobacterium extorquens* CM4 was transformed in order to be able to emit fluorescence in proportion to the weak concentration of chloromethane (Farhan UI Haque *et al.*, 2013). This bioreporter strain is based on the presence of a plasmid that carries the *yfp* gene, which then encodes a fluorescent protein under the control of the promoter of gene *cmuA* (Figure 1.7). The bioreporter strain allows a sensitive and proportional detection of chloromethane concentrations, with a detection threshold of 2

pM. This bioreporter has been tested for the quantification of chloromethane emitted by the plants *A. thaliana* and *V. rotundifolia*, which were demonstrated to have emission rates of respectively 13 and 2800 ng.g of dry matter⁻¹. h⁻¹ (Farhan Ul Haque *et al.*, 2013).





(A) In the bioreporter strain, a plasmid that carries the *yfp* gene under the control of the promoter of the *cmuA* gene (encoding a protein with a methyltransferase domain and a corrinoid-binding domain, essential for chloromethane utilization in *M. extorquens* CM4 (Studer *et al.*, 2001, 2002)), which is induced in the presence of chloromethane, is present in *M. extorquens* CM4 (Farhan UI Haque *et al.*, 2013). The plasmid carries a gene resistant to kanamycin in order to maintain its presence during the propagation of the bioreporter strain. (B) The emission of fluorescence of the Yfp protein is specific to the presence of chloromethane. The threshold of detection with the bioreporter strain is marked in yellow, and in black with the GC dosage.

2.2. Chloromethane sinks

2.2.1. Abiotic degradation of chloromethane

Different abiotic sinks of chloromethane, which vary in their ability to eliminate chloromethane, have been identified (Tableau 1.4). Even though an abiotic degradation of VOCs has been observed in soils, abiotic degradation of chloromethane has not been described (Miller *et al.*, 2004; Insam and Seewald, 2010).

Sink type	Estimation	Low/high values		
зик туре	(Gg. year ⁻¹) ^a	(Gg. year ⁻¹) ^b		
Tropospheric reaction with OH	-3,180	-3800/ -4,100		
Stratospheric loss	-200	-100/ -300		
Reaction with Cl ⁻ in marine boundary layer	-370	-18/ -550		
Microbial degradation in soil	< -1,000	-100/ -1,600		
Cold ocean water loss	-75	-93/-145		
Total sinks	< -4,875	-4273/ -6,695		

Table 1.4. Estimation of atmospheric chloromethane sinks

^a Data from Keppler *et al.*, 2005

^b Data from Clerbaux *et al.,* 2007

2.2.2. Biotic degradation of chloromethane

Biotic degradation of chloromethane is mainly due to microorganisms activity, although its estimation is variable (from 180 to 1,600 Gg.an⁻¹) (Keppler *et al.*, 2005).

Degradation by bacteria in marine environments

Marine environments play a role in the chloromethane degradation. Bacterial strains able to utilize chloromethane have been isolated from marine environments, such as *Leisingera methylohalidovorans* MB2, the first class of *Rhodobacteracea* shown to be able to degrade chloromethane (Goodwin *et al.*, 1997; Schaefer; 2002, Tableau 1.5). Three other chloromethane-degrading strains of this family have been isolated from marine environments: *Roseovarius* sp. 179, *Roseovarius* sp. 217 and *Ruegeria* sp. 198 (Schäfer *et al.*, 2005; Tableau 1.5).

Table 1.5. Chloromethane-utilizing bacterial isolated from contrasting environments

Bacterial strain	Origin	Gram type	Metabolism	Trophic type	Presence of <i>cmuA</i> ^a	Reference
Acetobacterium dehalogenans MC	Activated sludge	Positive	Anaerobic	Homoacetogenic	nd	Traunecker <i>et al.,</i> 1991
Aminobacter ciceronei IMB1	Fumigated strawberry field	Negative	Aerobic	Facultative methylotroph	Yes	Hancock <i>et al.,</i> 1998
Aminobacter lissarensis CC495	Beech woodland soil	Negative	Aerobic	Facultative methylotroph	Yes	Coulter <i>et al.,</i> 1999
Hyphomicrobium sp. AT1	Phyllosphere	Negative	Aerobic	Facultative methylotroph	Yes	Nadalig <i>et al.,</i> 2011
Hyphomicrobium sp. AT2	Phyllosphere	Negative	Aerobic	Facultative methylotroph	Yes	Nadalig <i>et al.,</i> 2011
Hyphomicrobium sp. AT3	Phyllosphere	Negative	Aerobic	Facultative methylotroph	Yes	Nadalig <i>et al.,</i> 2011
Hyphomicrobium sp. AT4	Phyllosphere	Negative	Aerobic	Facultative methylotroph	Yes	Nadalig <i>et al.,</i> 2011
Hyphomicrobium sp. MC1	Industrial sewage plant	Negative	Aerobic	Facultative methylotroph	Yes	Hartmans <i>et al.,</i> 1986
Hyphomicrobium sp. MC2	Soil from a petrochemical factory	Negative	Aerobic	Facultative methylotroph	Yes	Doronina <i>et al.,</i> 1996
Leisingera methylohalidovorans MB2	Marine tide pool	Negative	Aerobic	Facultative methylotroph	No	Schaefer, 2002
Methylomicrobium album BG8	Fresh water	Negative	Aerobic	Obligatory methylotroph	nd	Han and Semrau, 2000
Methylobacterium extorquens CM4	Soil from a petrochemical factory	Negative	Aerobic	Facultative methylotroph	Yes	Doronina <i>et al</i> ., 1996
Pseudomonas aeruginosa NB1	Activated sludge	Negative	Anaerobic	Facultative methylotroph	nd	Freedman <i>et al.,</i> 2004
Roseovarius sp. strain 179	Coastal seawater	Negative	Aerobic	Facultative methylotroph	Yes	Schäfer <i>et al.,</i> 2005
Roseovarius sp. strain 198	Coastal seawater	Negative	Aerobic	Facultative methylotroph	Yes	Schäfer <i>et al.,</i> 2005
Roseovarius sp. strain 217	Seawater	Negative	Aerobic	Facultative methylotroph	Yes	Schäfer <i>et al.,</i> 2005

^a nd; not determined

Bacterial degradation in the phyllosphere

The aerial parts of plants constitute the phyllosphere, which is home for many microorganisms. The total leaf surface (bottom and upper side) corresponds to twice the surface of Earth, with a bacterial density of 10⁷.cm⁻² (Vorholt, 2012). Few studies have studied so far the role of phyllospheric microorganisms in the degradation of plant-emitted chloromethane. Some bacteria able to utilize chloromethane have been isolated from the surface of leaves of *A. thaliana* (Nadalig *et al.*, 2011). Nevertheless, the potential role of these epiphytic bacteria as biotic filters for chloromethane has so far not been taken into account.

Bacterial degradation in soils

Soil contains a large variety of compounds, such as halogenated compounds. Soil has been shown to be major sink for some of halogenated compounds, and for instance 70% of bromomethane (CH₃Br) is consumed by soils (Shorter *et al.*, 1995). Although chloromethane degradation in soils is misevaluated, it is estimated to exceed 1,000 Gg. year⁻¹ (Tableau 1.4). Chloromethane consumption in soil has been correlated to bacterial and fungal degradation mainly in the O horizon with was found to be very weak in lower horizons (Redeker and Kalin, 2012).

The exchange of chloromethane between the terrestrial ecosystems and the atmosphere are modulated by bacteria living in soil (Miller *et al.*, 2004; Borodina *et al.*, 2005; Keppler *et al.*, 2005; Clerbaux *et al.*, 2007; Schäfer *et al.*, 2007; Rhew *et al.*, 2010). A variety of bacteria affiliated to *Alpha*- or *Beta-proteobacteria* that are able to utilize chloromethane have isolated from forest soils, but their contribution to the chloromethane budget remains to be evaluated (Miller *et al.*, 2004; Borodina *et al.*, 2005).

Chloromethane consumption has also been demonstrated in fungi via reactions with produced secondary metabolites including more than 200 halogenated metabolites (Gribble, 2003) such as brominated, fluorinated, iodinated, and chlorinated compounds (Anke et Weber, 2006). Among chlorinated metabolites, chloromethane can be released during methylation reactions in fungi (Figure 1.8). Methylation is a common reaction of fungal metabolism (Anke and Weber, 2006).





Methylation in fungi involves SAM, as a donor of the methyl group, as found for chloromethane synthesis in plants. In fungi, the utilization of chloromethane is preferred over SAM, since the reaction needs less energy (Harper, 2000). When chloromethane serves as a methyl group donor to an hydroxyl (-OH) or carboxyl (-COOH) group, products such as veratryl alcohol or methyl benzoate are produced (Figure 1.8). Veratryl alcohol is a secondary metabolite involved in the synthesis of enzymes that are necessary for lignin degradation (Dekker *et al.*, 2001). Methyl benzoate is an olfactive compound synthesized to attract pollinators, and acts in inhibition of *A. thaliana* root growth (Horiuchi *et al.*, 2007). Chloromethane balance in fungi is difficult to determine since they are able to simultaneously produced and degraded chloromethane (Anke and Weber, 2006).

Although the role of soil as a chloromethane sink does not need to be further evidenced, uncertainties persist about global chloromethane fluxes in soils (see section below).

2.3. Global chloromethane balance in soil

Soil is a complex environment for which chloromethane sources and sinks are difficult to estimate. Besides its intrinsic heterogeneity (Figure 1.4), contrasting variations of physicochemical, climatic and biological parameters occur. For instance, a seasonal effect on chloromethane production has been described (Peñuelas *et al.*, 2014). Increased chloromethane emissions have been observed in summer (Llusia *et al.*, 2013; Oderbolz *et al.*, 2013), compared to in the falls where the leaf cover reduce emission of COVs such as chloromethane (Figure 1.9). Temperature also has an impact on chloromethane production.
The release of chloromethane by decomposing plants increases with higher temperatures (Hamilton, 2003).



Figure 1.9. Seasonal chloromethane fluxes in forest soil

Chloromethane emissions from a forest soil have been monitored during one year. JFM, January, February and March; AMJ, April, May and June, JAS, July, August and September; OND, October, November and December (from Redeker *et al.*, 2012). Error bars show standard errors.

Chloromethane production can oscillate between 30 and 60 % (Saito and Yokouchi, 2008). Estimation of chloromethane production is difficult since chloromethane fluxes are bidirectional (from the soil to the atmosphere and from the atmosphere to the soil), and forest soil microorganisms modulate these exchanges (Schäfer *et al.*, 2005). The estimated chloromethane sinks in soils varies from 256 Gg.year⁻¹ (Clerbaux *et al.*, 2007) to 1,000 Gg.year⁻¹ (Keppler *et al.*, 2005), which represents 6 up to 20% of the total tropospheric chloromethane sinks. Further studies are needed to better understand the global cycle of chloromethane in forest soils, which are both source and sink of chloromethane.

Quantification uncertainties of different chloromethane sources and sinks prevent accurate global balance calculations. The fact that some environments are sources and sinks, even makes it more complex to estimate. The estimation of chloromethane production and consumption can be improved using analytical methods basing on carbon and hydrogen isotope fractionation (Keppler *et al.*, 2005; Greule *et al.*, 2012; Nadalig *et al.*, 2013) and the development of three-dimensional model of terrestrial chloromethane fluxes (Xiao *et al.*, 2010). Studies of chloromethane degradation in forest soils have shown the presence of bacteria active on chloromethane degradation (Borodina *et al.*, 2005; Miller *et al.*, 2004). An unpublished study of a forest soil in Steigerwald in Germany (samples from this site were used

for stable isotope probing experiments described in chapter 4) demonstrated chloromethane degradation by bacteria in oxic conditions (Rüffer, 2013).

3. <u>Methylotrophy</u>

Microorganisms able to grow in chloromethane as sole carbon and energy source are methylotrophic microorganisms (Table 1.5). Methylotrophy is defined as the ability of an organism to utilize a compound with no C-C bonds as the sole source of carbon and energy. Substrates for methylotrohic growth include one-carbon compounds such as methane, (CH₄), methanol (CH₃OH), formaldehyde (HCHO), and methylamine (CH₃NH₂), as well as halogenated compounds such as chloromethane and dichloromethane, or multi-carbon compounds such as dimethylamine (C₂H₇N) (Chistoserdova, 2011; Muller *et al.*, 2011a).

Methylotrophic organisms are found ubiquitous in the air (DeLeon-Rodriguez *et al.*, 2013), in fresh water and marine environments (Neufeld *et al.*, 2008), in plant roots leaves (Nadalig *et al.*, 2011; Knief *et al.*, 2012), and in soils and marine sediments (Kolb, 2009a). These organisms belong to the three domains of living organisms: bacteria, archea (methanotrophic archea) (Antony *et al.*, 2012), and eukaryotes (yeasts) (Yurimoto, 2009). Yeasts of the genus *Pichia*, *Candida* and *Hansenula* are able to metabolize methanol via an alcohol oxidase present in the peroxisome cellular compartment (van der Klei *et al.*, 2006).

The methylotrophic metabolism has been the best studied in bacteria (Kolb, 2009; Chistoserdova, 2011; Peyraud *et al.*, 2011; Peyraud *et al.*, 2012). Since the discovery of bacteria able to utilize methane and methanol as the only source of carbon and energy (Loew, 1892; Käserer, 1906; Söhngen, 1906), a large variety of bacteria from different genus have been identified, including facultative methylotrophs (able to grow with C₁ and multi-carbon substrates such as succinate) (Whittenbury *et al.*, 1970; Lidstrom, 2006; Boden *et al.*, 2008; Hung *et al.*, 2011). Methylotrophic bacteria can be found among *Alpha, Beta* and *Gamma proteobacteria*, Gram negative and Gram positive bacteria, using aerobic or anaerobic metabolisms.

Methylotrophic bacteria are classified in two groups. The first group of methanotrophs is able to utilize methane as the sole source of carbon and energy, are often strict methylotrophs (*Methylomonas, Methylobacter, Methylosinus,* and *Methylocystis*). The second group of non-methanotrophic methylotrophs (unable to oxidize methane), are often facultative methylotrophs, like bacteria of the genus *Hyphomicrobium* or *Methylobacterium* (Chistoserdova and Lidstrom, 2013).

27

3.1. Methylotrophic pathways

Methylotrophy involved 3 major metabolic steps:

- Initial transformation of the methylotrophic compound into formaldehyde (HCHO) (not the case for chloromethane);

- Complete oxidation of the methylotrophic compound to CO₂ in order to produce energy;

- Carbon assimilation for biomass production via 3 alternative pathways: the ribulose biphosphate (RuBP) pathway that utilizes CO₂, the ribulose monophosphate (RuMP) pathway from formaldehyde, or the serine cycle from methylenetetrahydrofolate (CH₂=H₄F) and CO₂ (Chistoserdova, 2011).

For most methylotrophs, the first metabolic step is the oxidation or the hydrolysis of the carbon compound into formaldehyde. The enzymes involved in this step are specific to their substrates: the soluble methane monooxygenase (sMMO) or the particular methane monooxygenase (pMMO) for methane, the methylamine deshydrogenase (MADH, periplasmic) for methylamine, the methanol deshydrogenase (Mxa, periplasmic) for methanol, or a dichloromethane deshydrogenase for dichloromethane (DcmA, cytoplasmic). The mxaF gene has been used as a molecular marker of methanol oxidation (Neufeld et al., 2008). More widespread in the environment than the mxa genes, alternative methanol degradation systems exist such as the xox genes or the mdh gene (Kalyuzhnaya et al., 2008). Among these genes, *xoxF* codes for a subunit of a methanol dehydrogenase that shares more than 50% of amino acid identity with MxaF (Schmidt et al., 2010; Taubert et al., 2015). Several methanol oxidation pathways can be simultaneously present in an organism as found in M. extorquens (Figure 1.10, Skovran et al., 2011). Several methylamine oxidation pathways have also been found in *M. extorquens* strains (Vuilleumier et al., 2009; Gruffaz et al., 2014; Nayak et Marx, 2014). Halogenated compounds (chloromethane, dichloromethane) can be utilized by methylotrophic bacteria such as M. extorquens and Hyphomicrobium spp. (Gälli and Leisinger, 1985; La Roche and Leisinger, 1990; Vannelli et al., 1998; McAnulla et al., 2001). The dehalogenation of dichloromethane is catalyzed by a dichloromethane dehalogenase encoded by gene *dcmA*, and formaldehyde is produced.

Formaldehyde is transformed into formate by either a tetrahydrofolate-dependent (H₄F) pathway or a tetrahydromethanopterin (H₄MPT)-dependent pathway in *Methylobacterium*

28

strains. The first pathway is not the main one regarding the production of energy and involves inter-conversions between methylene-H₄F, methenyl-H₄F, formyl-H₄F and formate (Marison and Attwood, 1982). This metabolic pathway might play a major role by keeping adequate concentrations of the different H₄F-containing metabolites, which are essential intermediates for carbon assimilation via the serine cycle pathway (Marx *et al.*, 2005; Crowther *et al.*, 2008). In the second pathway, discovered in 1998, H₄MPT is condensed to formaldehyde, and successively transformed into methylene-H₄F, methenyl-H₄F, formyl-H₄F, and finally into formate (Chistoserdova et al., 1998). This H₄MPT-dependent pathway has been first described in the Archaea, and is more efficient in formaldehyde oxidation into CO2 than the H4Fdependent pathway, because its corresponding enzymes display high activities (Vorholt, 2002). Despite the fact that most methylotrophic substrates are oxidized into formaldehyde, this is the case for chloromethane. Chloromethane dehalogenation produces methylenetetrahydrofolate ($CH_2=H_4F$), which either enters the serine cycle pathway for biomass production or is oxidized into formate for energy production (Figure 1.10). As a matter of facts, M. extorquens CM4 converts directly chloromethane into methylene-tetrahydrofolate (CH₂=H₄F) without formaldehyde production thanks to the activity of CmuA (methyltransferase and corrinoid-binding domain-containing protein), CmuB (methylcobalamin: H₄F methyltransferase) and MetF2 (methylene-H₄F reductase) (Studer et al., 1999; Studer et al., 2001; Studer et al., 2002) (see section 1.3.4.2.).



Figure 1.10. Scheme of different methylotrophic pathways in *M. extorquens*

Oxidation state is shown on the Y axis. The reactions specific for chloromethane $(CH_{3}CI)$ and dichloromethane $(CH_{2}CI_{2})$ utilization as sole source of carbon and energy are shown in green and purple, respectively: CmuABC, chloromethane utilizing genes; MetF2, methylene-H₄F reductase; DcmA, dichloromethane dehalogenase. Enzymes for initial oxidation are in black: MDH, methanol dehydrogenase; MADH, methylamine dehydrogenase. Reactions leading to complete oxidation of carbon to energy are in orange: Fae, formaldehyde activation enzyme; MtdAB, methylene-tetrahydromethanopterin (H₄MPT) dehydrogenase; Mch, methenyl-H₄MPT cyclohydrolase; Fch, methylene-tetrahydrofolate (H₄F) cyclohydrolase; FtfL, formyl-H₄F ligase; FhcABCD, formyltransferase-hydrolase complex; Fdh, formate dehydrogenase. Reactions for carbon assimilation to biomass are in grey: FtfL, formyl-H₄F ligase; Fuchylene-H₄F cyclohydrolase; MtdA, methylene-H₄MPT dehydrogenase; RuMP, ribulose monophosphate cycle; RuBP, ribulose biphosphate cycle.

More than 86 genes involved in *M. extorquens* C₁ compound utilization have have been identified (Marx *et al.*, 2003). These genes are involved in the methanol oxidation methylamine, formaldehyde, and formate or in the serine-cycle pathway (Figure 1.11).





Entering C₁ compounds are circled in blue and main biosynthesis pathway outputs are indicated in green. Bacteria oxidize methanol (CH₃OH) or dichloromethane (CH₂Cl₂) to formaldehyde that is condensed with a tetrahydromethanopterin (H₄MPT), and further oxidized to formate (HCOOH). Formate is further converted to methylene-tetrahydrofolate (H₄F) via the serine cycle or transformed in CO₂. Chloromethane (CH₃Cl) is assimilated via the transfer of the methyl group to H₄F which is then assimilated via the serine cycle to produce biomass or oxidized to formate (HCOOH). Acetyl-CoA assimilation and glyoxylate regeneration proceed via the ethylmalonyl-CoA pathway. The tricarboxylic acid (TCA) cycle is mainly used for assimilation of multi-carbon compounds (Šmejkalová *et al.*, 2010). Circled numbers identify distinct reactions: 1, serine hydroxymethyl transferase; 2, serine-glyoxylate aminotransferase; 3, hydroxypyruvate reductase; 4, glycerate kinase; 5, enolase; 6,

phosphoenolpyruvate carboxylase; 7, malate dehydrogenase; 8, malate-CoA ligase (malate thiokinase); 9, L-malyl-CoA/ β -methylmalyl-CoA lyase; 10, β -cetothiolase; 11, acetoacetyl-CoA reductase; 12, crotonase (R- specific); 13, crotonyl-CoA carboxylase reductase; 14, ethylmalonyl-CoA/methylmalonyl-CoA epimerase; 15, ethylmalonyl-CoA mutase; 16, methylsuccinyl-CoA deshydrogenase; 17, mesaconyl-CoA hydratase; 18, propionyl-CoA carboxylase; 19, methylmalonyl-CoA mutase; 20, succinyl-CoA synthase; 21, succinate deshydrogenase; 22, fumarase; 23, citryl-CoA lyase; 24, aconitate hydratase; 25, NADP-dependent isocitrate dehydrogenase; 26, 2-oxoglutarate dehydrogenase complex; 27, methanol dehydrogenase; 28, dichloromethane dehalogenase; 29, formate dehydrogenase. For a given reaction, the number of parallel arrows correspond to the number of molecules involved in the reaction. PHB, polyhydroxybutyrate; Q, quinone (adapted from Šmejkalová *et al.*, 2010).

In *Methylobacterium*, for biomass production, formaldehyde incorporates the serine cycle pathway via formate and methylene-H₄F. The GlyA enzyme (serine hydroxymethyltransferase) catalyzes the reaction between methylene-H₄F and glycine to produce serine (Figure 1.11). Then, serine (C₃ compound) is successively transformed into hydroxypyruvate, glycerate, glycerate-2-phosphate and phosphoenolpyruvate. Phosphoenolpyruvate is subsequently converted into oxaloacetate, then malate by incorporation of CO₂. Malate (C₄ compound) coupled to co-enzymeA (CoA) is then cleaved in 2 compounds: glyoxylate allowing to close the cycle serine, and acetyl-CoA which is integrated in the ethylmalonyl-CoA pathway where a CO₂ molecule is incorporated (Erb *et al.*, 2008; Peyraud *et al.*, 2009; Šmejkalová *et al.*, 2010). This cycle generates hydroxybutyryl-CoA, a precursor of the polyhydroxybutyrate (PHB) synthesis and granule production for carbon and phosphate storage in the cell.

3.2. Bacterial degradation of dichloromethane (CH₂Cl₂)

Bacterial dichloromethane is better characterized in oxic than in anoxic conditions.

3.2.1. Dichloromethane utilization in anoxic conditions

Dichloromethane can be fermented by *Dehalobacterium formicoaceticum* (Yurimoto, 2009). This anaerobic bacterium has a cobalamin- and H₄F-dependent methyltransferase. Dichloromethane and H₄F are converted into methylene-tetrahydrofolate (CH₂=H₄F) and into inorganic chloride. Part of CH₂=H₄F is converted into formate, whereas another part of the CH₂=H₄F incorporates CO₂ to produce acetate (Mägli *et al.*, 1998).

3.2.2. Dichloromethane utilization in oxic conditions

The first characterized isolate able to utilize dichloromethane for growth, belongs to the genus *Methylopila* (Brunner *et al.*, 1980). Three aerobic dichloromethane degradation pathways are known. Dichloromethane utilization as carbon and energy sources is possible only in one of these pathways and involved the expression of a dichloromethane dehalogenase belonging to the glutathion-S-transferase (GST) family that is encoded by gene *dcmA* (Figure 1.12). The majority of strains able to utilize dichloromethane for their growth, harbor gene *dcmA* (Vuilleumier *et al.*, 2001; Muller *et al.*, 2011b). When detected, gene *dcmA* is highly conserved with up to 98-100% identity at the nucleotide level. The DcmA protein catalyzes the dechlorination of one mole of dichloromethane into two moles of protons (H⁺), two moles of chloride (Cl⁻) and one mole of formate. This reaction involved two genotoxic compounds, S-chloromethylglutathion and formaldehyde (Figure 1.12).



Figure 1.12. Dichloromethane degradation by the hydrolytic pathway of glutathione S-transferase

(adapted from Muller *et al.,* 2011). spont.: spontaneous, abiotic reaction; GSH, glutathione; SG, glutathionyl.

The aerobic bacterial degradation of dichloromethane has been best described in *M. extorquens* DM4, a strain able to utilize dichloromethane as unique source of carbon and

energy (Roche and Leisinger, 1991; Kayser *et al.*, 2002; Muller *et al.*, 2011a). Utilization genes are located on a catabolic transposon containing four *dcm* genes, framed by insertion sequences. Among *dcm* genes, *dcmA* encoded a dichloromethane dehalogenase, *dcmR* encodes a transcriptional factor, and genes *dcmB* and *dcmC* encode for proteins of unknown function. The *dcm* transposon is localized on a larger genomic island of 126 Kb (Vuilleumier *et al.*, 2009). Dichloromethane utilization in *M. extorquens* DM4 generates formaldehyde production, which is readily oxidized into formate via the H₄MPT-dependent pathway. Formate is reduced into CH₂=H₄F, which enters the serine cycle for biomass production, or is substrate to formate dehydrogenase (FDH) to produce CO₂ and energy (Figure 1.10).

3.3. Bacterial degradation of chloromethane

Bacteria affiliated to various genera and isolated from contrasting environments were described as able to degrade chloromethane (Table 1.5). Chloromethane can serve as carbon and energy source but in other cases, chloromethane is degraded and not assimilated as a carbon source in both oxic and anoxic conditions (Traunecker *et al.*, 1991; Han and Semrau, 2000; Freedman *et al.*, 2004).

3.3.1. Degradation under anoxic conditions

Chloromethane dehalogenation in anaerobic condition is catalyzed by corrinoid-containing enzyme which transfers the methyl group (CH₃) to H₄F. Products of this reaction are chloride ions and methyl-H₄F, which is then oxidized into formate and CO₂ or assimilated in the biomass via the acetyl-CoA pathway (Harper, 2000). *Acetobacterium dehalogenans* strain MC is able to utilize chloromethane in anaerobiosis (Table 1.5) (Traunecker *et al.*, 1991; Harper, 2000). This dehalogenation requires a corrinoid-dependent methyltransferase, which in unlike other corrinoid-dependent methyltransferases, maintains cobalt int the corrinoid cofactor using an ATPase dependent system (Studer *et al.*, 1999). *Pseudomonas aeruginosa* NB1 is also able to utilize chloromethane as unique source of carbon and energy in nitrate respiration conditions (Freedman *et al.*, 2004), by a dechlorination mechanism that remains unknown.

3.3.2. Degradation in oxic conditions

The first described strain able to utilize chloromethane as sole source of carbon and energy was *Hyphomicrobium* sp. MC1 (Hartmans *et al.*, 1986). Like this strain, the majority of the

chloromethane-degrading strains isolated so far utilize chloromethane under oxic conditions. They were isolated from various environments such as petrochemical factory soil, wastewater treatment plans, forest soils, freshwater and marine environments (Tableau 1.5). They are affiliated to the class of *Alphaproteobacteria* of the genera *Methylobacterium*, *Hyphomicrobium* and *Aminobacter*. As of this date, the *cmu* pathway is the best-characterized bacterial chloromethane-degradation pathway with a majority of the studies performed in *M. extorquens* CM4, but uncharacterized *cmuA*-independent pathway exist (see below).

Chloromethane degradation pathway in *Methylomicrobium album* BG8
The methanotroph *Methylomicrobium album* BG8 degrades chloromethane in oxic conditions, independently of the *cmu* pathway, only in presence of another carbon source such as methanol (Figure 1.13) (Han et Semrau, 2000).



Figure 1.13. Proposed mechanism of chloromethane oxidation in *Methylomicrobium album* BG8

Chloromethane can be oxidized by the particulate methane monooxygenase (pMMO) since *M. album* BG8 stops degrading chloromethane in presence of acetylene, a pMMO inhibitor. Methanol is oxidizes into formaldehyde (HCHO) by the methanol dehydrogenase (MDH). Part of the produced formaldehyde enters the C₁ metabolism to produce biomass and the rest is oxidized into CO₂ and reducing equivalents (2[H]) for energy production via the formaldehyde deshydrogenase (FDH) and the formate dehydrogenase (FDH). Compared to methanol, chloromethane degradation generates 2 reducing equivalents less (From Han and Semrau, 2000).

Chloromethane stimulates *M. album* BG8 growth with methanol, and when using [¹⁴C]-labeled chloromethane, it was demonstrated that the labeled carbon was assimilated into the biomass (38%) and released as CO₂ (50%). This strongly suggested that chloromethane can be used as carbon and energy source when methanol was present (Han and Semrau, 2000). When provided alone, chloromethane is unable to sustain growth and serve as the sole source of carbon. The authors suggested that *M. album* BG8 would not be able to generate enough reducing equivalents from chloromethane degradation to sustain growth with chloromethane. In presence of another carbon source, in larger quantities, the chloromethane utilization was observed (5 mM of methanol and 2.6 mM of chloromethane) (Han and Semrau, 2000).

– The *cmu* pathway

Combined biochemical and genetic studies in *M. extorquens* CM4 were used to identify the *cmu* (chloromethane utilization) pathway under oxic conditions (Vannelli *et al.*, 1998, 1999). *M. extorquens* CM4 metabolizes one mole of CO_2 and one mole of chloride (Vannelli *et al.*, 1998). Using random mutagenesis with the minitransposon *Tn*5, 9 mutants were detected with wild-type growth with methanol or methylamine, but impaired growth with chloromethane (Vannelli *et al.*, 1998). Analysis of the *Tn*5 insertion sites in mutants that lost the ability to grow with chloromethane, demonstrated the essential role of *cmuABC*, *purU* and *metF2* (Studer *et al.*, 2001, 2002). In addition, *cmuAB* genes were demonstrated to be essential for chloromethane dechlorination (Studer *et al.*, 2001b, 2002). CmuA and CmuB catalyze the methyl group transfer from chloromethane to H₄F to form methyl-H₄F (CH₃-H₄F) (Figure 1.14) (Studer *et al.*, 2001). CH₃-H₄F is oxidized either into formate and then CO₂ for energy production, or into methylene-H₄F for biomass production via the serine cycle (Vannelli *et al.*, 1999).



Figure 1.14 The *cmu* pathway of *Methylobacterium extorquens* CM4

The methyl group of chloromethane is transferred to tetrahydrofolate (H₄F) by *cmuA* and *cmuB*. The methylene-H₄F produced by the oxidation of methyl-H₄F enters the serine cycle to produce biomass and is also oxidized to CO₂ to produce energy. CmuA, protein with a methyltransferase domain and a corrinoid-binding domain (Studer *et al.*, 2001), CmuB, methyltransferase methylcobalamin: H₄F (Studer *et al.*, 1999); MetF2, methylene reductase; FoID, bifunctional enzyme methylene-H₄F dehydrogenase/ methenyl-H₄F cyclohydrolase; PurU, 10-formyl-H₄F hydrolase.

All the genes of *cmu* pathway in *M. extorquens* CM4 are plasmid pCMU01-borne in a region of 180 kb (Figure 1.15). The *cmu* genes are clustered in two groups 30 kb apart that displays an atypical GC content and is devoid of detectable mobile elements. Upstream of *purU* and *foID*, gene *foIC2* encoded a bi-functional folylpolyglutamate synthase / dihydrofolate synthase protein. This region contains also *paaE-like*, *fmdB* and *hutI*, encoding respectively a putative oxidoreductase, a putative transcriptional regulator and a putative imidazole hydrolase.





The 21 kb distant clusters 1 and 2 contain the essential genes for chloromethane utilization (*cmu*) known before the genome has been sequenced (Vannelli *et al.*, 1999; Studer *et al.*, 2002). The *cmu* region covers 40 kb (cluster 1 and 2). The S1, S2, S3 regions correspond to a region of 69 kb conserved between pCMU01 and the p1METDI plasmid of *M. extorquens* DM4. The S4 region shows a synteny with a duplicated region on *M. extorquens* DM4 chromosome. Circles represent (from the outside): 1, percentage of deviation of the GC content in a 1 kb window; 2, predicted CDS transcribed in the clockwise direction; 3, predicted CDS transcribed in the counterclockwise direction; 4, GC skew (G+C/G-C) in a 1 kb window; 5, transposable elements (in pink) and pseudogenes (in grey) (adapted from the PhD manuscript of Sandro Roselli, 2009).

This region contains 23 genes involved in biosynthesis and transport of cobalamin, those involved on H₄F biosynthesis, both cofactors are essential for chloromethane utilization via the *cmu* pathway (Figure 1.15). Some of these plasmid-borne genes have chloromsomal homologs (Roselli *et al.*, 2013). Genome comparative analysis of strains harboring *cmu* genes, demonstrated two types of *cmu* gene organization; one found in *M. extorquens* CM4 where

genes are localized in two clusters spaced by 30 kb and a second one, found in others bacteria, where *cmu* are contiguous (Figure 1.15). A high degree of similarity in *cmu* genes organization where found in chloromethane-degrading strains (Figure 1.15, Schäfer *et al.*, 2007), all affiliated to the class of *Alphaproteobacteria*. The genes order into the *cmu* cluster is highly conserved (Nadalig *et al.*, 2011). In *M. extorquens* CM4, *cmuB* and *cmuC* genes are part of the same transcriptional unit, unlike gene *cmuA*, which has its own promoter (Studer *et al.*, 2002). The genome of some chloromethane degrading-strains lack detectable *cmu* genes as in *Roseovarius* sp. 217 and *Leisingera methylohalidovorans* MB2 (Schäfer *et al.*, 2007). Chloromethane stable isotope analysis revealed different isotopic fractionation signatures for the carbon and hydrogen in *cmu*-containing strains and *L. methylohalidovorans* MB2 which not possess those genes (Nadalig *et al.*, 2014). The *cmu*-independent chloromethane degradation pathway in *L. methylohalidovorans* MB2, needs to be solved.



Figure 1.16. Comparison of *cmu* gene organization in bacteria harboring gene *cmuA*

Encoding genes are represented with arrows. Genes with an incomplete determined sequenced are represented by incomplete arrows. The gene reference colors are based on those from *M. extorquens* CM4 representation, where they are separated by 30 kb. Homologous genes are represented by the same color, the percentage of homology with proteins encoded by *M. extorquens* CM4 genes are written into arrows. The genetic representation is represented at the scale. To note that in the strain MB2, only one part of *cmuA*, corresponding to corrinoid cofactor binding-domain is present.

4. The Methylobacterium model

The genus *Methylobacterium* was first described in 1976 (Patt *et al.*, 1976), and now encompasses 20 species. This genus is affiliated to *Alphaproteobacteria*, of the order *Rhizobiales* and of the *Methylobacteriacaea* family as deduced from nucleotidic sequence analysis of 16S rRNA encoding genes (Lidstrom, 2006). They are facultative methylotrophs, strictly aerobic, mesophiles with an optimal temperature between 25 and 30°C, Gram

negative. They test positive with the catalase and oxidase assays. Cells often display poly- β hydroxybutyrate inclusions involved in energy and carbon storage. They display a pink coloration due to carotenoids (Korotkova and Lidstrom, 2001; Van Dien *et al.*, 2003). *Methylobacterium* strains have been isolated from contrasting environments either handmade (road borders, water tanks) or natural (soil, leaf surface) (Tableau 1.6) (Radajewski *et al.*, 2002; Nadalig *et al.*, 2011; Bai *et al.*, 2015). They metabolize C₁ compounds such as methanol and methylamine, and a few strains also degrade toxic chlorinated methanes such as dichloromethane or chloromethane (Dourado *et al.*, 2015). Methylotrophs have been among the first bacteria to be studied in the environment using molecular approaches (Holmes *et al.*, 1995; Kolb et Stacheter, 2013).

4.1. *Methylobacterium*, a model for methylotrophy studies

One of the most studied species of the genus Methylobacterium is M. extorquens which has been isolated from various environments such as plants, soil, waste waters or cloud water (Bai et al., 2015; Bringel et Couée, 2015; Nadalig et al., 2011; Ochsner et al., 2015; Temkiv et al., 2012). M. extorquens AM1 is the most studied strain which has been used along with Methylobacterium organophilum, as Methylobacterium reference strains, in plantmicroorganism interaction studies (Dourado et al., 2015). M. extorquens AM1 harbours five replicons and 174 insertion sequences, which complicates its utilization in molecular biology studies. Thus, strain AM1 has been replaced by *M. extorquens* PA1 whose genome consists of a unique chromosome and only 20 insertion sequences (Nayak et Marx, 2014). M. extorquens strains (AM1, BJ001, CM4, DM4, DSM 13060 and PA1) are able to utilize methanol, but only a few of those are able to utilize chlorinated methanes as a sole carbon and energy source (Vuilleumier et al., 2009; Marx et al., 2012). M. extorquens CM4 has been isolated from a Russian petrochemical industry soil, and is able to grow with chloromethane (Doronina et al., 1996). On the other hand, *M. extorquens* DM4 able to grow on dichloromethane, has been isolated from a chlorinated alkane-polluted Swiss soil (Gälli et Leisinger, 1985). M. extorquens genomes display high GC content of at least 68% with a large chromosome sized between 5.5 and 5.9 Mb (Tableau 1.6).

Unlike strains PA1 and DSM 13060, further *M. extorquens* strains harbour at least one plasmid, including *M. extorquens* strain AM1, which remarkably harbors of a megaplasmid (Ochsner *et al.*, 2015). At the end of December 2015, 24 additional *Methylobacterium* genomes have been

sequenced from strains isolated from plant phyllosphere and rhizosphere (Bai et al., 2015).

These recently sequenced genomes harbour no *cmuA* gene homologue.

Introduction

Parameters ^a	<i>M. extorquens</i> AM1	<i>M. extorquens</i> BJ001 ^b	M. extorquens CM4	<i>M. extorquens</i> DM4	<i>M. extorquens</i> DSM13060	<i>M. extorquens</i> PA1	<i>M. nodulans</i> ORS 2060	<i>M. radiotolerans</i> JCM 2831	<i>M.</i> sp. 4-46
Genome									
Size (Mb)	6,88	5,85	6,18	6,12	6,73	5,47	8,84	6,90	7,73
% GC	68,5	69,4	68,0	68,0	68,23	68,2	68,4	71,0	71,5
Number of replicon	5	3	3	3	1	1	8	9	3
Number of predicted CDS	6335	5945	6276	5706	7132	5153	10333	7356	8479
Genbank accession number	NC_012808	NC_010725	NC_011757	NC_012988.1	NZ_AGJK0000 0000.1	NC_010172	NC_011894	NC_010505	NC_010511
Chromosome									
Size (Mb)	5,51	5,8	5,77	5,94	6,73	5,47	7,78	6,1	7,70
% GC	68,7	69,4	68,2	68,1	68,23	68,2	68,9	71,5	71,6
Number of plasmids	4	2	2	2	0	0	7	8	2
Sequencing (date)	UW Genome Sequencing Center, (2003/2006)	JGI (2008)	JGI (2008)	Génoscope (2006)	JGI (2010)	JGI (2007)	JGI (2009)	JGI (2008)	JGI (2008)
Genome publication	Vuilleumier <i>et al.,</i> 2009	Marx <i>et al.,</i> 2012	Marx <i>et al.,</i> 2012	Vuilleumier <i>et al.,</i> 2009	Koskimäki <i>et al.,</i> 2015	Marx <i>et al.,</i> 2012	Marx <i>et al.,</i> 2012	Marx <i>et al.,</i> 2012	Marx <i>et al.,</i> 2012

Table 1.6. Characteristics of the first *Methylobacterium* sequenced genomes

^a Génoscope (http://www.cns.fr/agc/microscope/home/index.php) or Craig Venter institute (http://www.jcvi.org/)

^b Originally named Methylobacterium populi (Van Aken et al., 2004), but based on 16S rRNA identity, was affiliated to M extorquens (Marx et al., 2012)

4.2. Genetic tools available for *M. extorquens*

Different genetic tools are available for *M. extorquens* AM1. They include tools for random mutagenesis, allelic gene replacement for targeted gene deletion and insertion, but also for modulated gene expression (constitutive or inducible). A complete list of genetic tools for *M. extorquens* AM1 is detailed in a recent review (Ochsner *et al.*, 2015).

4.3. Global approaches in *M. extorquens*

Our knowledge of methylotrophy in bacteria belonging to the genus *Methylobacterium* and in particular in the species *M. extorquens* has increased considerably over last years with high-throughput *Next Generation Sequencing* (NGS) technologies, allowing the study of genes (genomics), their expression at the RNA level (transcriptomics), at the protein level (proteomics) as well as of cell metabolites (metabolomics).

4.3.1. Genomic studies in *M. extorquens*

In 1998, the first Methylobacterium genome sequence was obtained from strain AM1 (Chistoserdova et al., 2003). It has been completed and assembled only in 2009 when the genome of strain DM4 was available (Vuilleumier et al., 2009). Later, in 2012, the genome of strain *M. extorquens* CM4 was sequenced simultaneously with the genomes of other *M. extorquens* strains unable to degrade chlorinated methanes (BJ001; PA1) (Marx *et al.*, 2012). When I started my PhD thesis, the MaGe platform hosted by the Génoscope (Evry, France), offered easy annotation and comparative analysis of 6 *M. extorquens* genomes (AM1, BJ001, CM4, DM4, DSM 13060 and PA1), with characterized growth abilities for methanol and chlorinated methanes. Intra-species *M. extorquens* comparative genomic study has detected the presence of a high number of insertion sequences (IS) in the vicinity of gene clusters involved in methanol utilization, suggesting a role of these mobile elements in the evolution of phylogenetically closely-related genomes of strains AM1 and DM4 (Vuilleumier et al., 2009). Nevertheless, the history of the evolution of *M. extorquens* genome remains unclear. As a matter of facts, despite the fact that M. extorquens AM1 and PA1 have similar GC content (68.2% and 68.5%, respectively) and share a large number of common genes involved in methylotrophy (90 genes with at least 95% identity between their corresponding encoded proteins), a recent study highlighted differences in growth rates for a variety of C_1 compounds

such as methanol, methylamine or formaldehyde, which would suggest the implication of different mechanisms (Nayak et Marx, 2014).

Combined random mutagenesis and comparative proteomic studies have evidenced the growth with chlorinated methanes involved genes localized on genomic islands, in addition to housekeeping genes involved in stress response, and central metabolism (Michener *et al.*, 2014a; Muller *et al.*, 2011a; Roselli *et al.*, 2013). Chloromethane or dichloromethane utilization generates a cellular stress that requires specific adaptive responses (Kayser et Vuilleumier, 2001; Kayser *et al.*, 2002; Roselli *et al.*, 2013). To decipher those adaptive mechanisms, Michener *et al.* (2014) have tested the ability to utilize dichloromethane of naïve non-degrading strains. The plasmid-borne *dcmA* gene has been transferred to strain AM1 (unable to utilize dichloromethane for its growth) (Michener *et al.*, 2014b). Transconjugants displayed poor growth with dichloromethane and but had a detectable dehalogenation activity. The subsequent genome sequencing of adapted clones enabled the identification of mutation within gene *clcA* encoding an antiporter Cl⁻/H⁺. When a plasmid harbouring genes *dcmA* and the mutated *clcA* were introduced in *M. extorquens* PA1 (unable to utilize dichloromethane.

Complementary studies of chloromethane utilization and adaptation in naïve strains (unable to utilize chloromethane for their growth) have been performed (Michener *et al.*, submitted article). A plasmid harbouring genes of the *cmu* pathway (*folD*, *metF2*, *paaE-like*, *purU*) and of associated genes (*fmdB*, *hutl*), has been inserted by conjugation in *M. extorquens* AM1, PA1, BJ001, *M. radiotolerans* and *M. nodulans* strains. The resulting transconjugants acquired the ability to growth with chloromethane, although cultures were of low optical densities. When the mutated gene *clcA*, which has been demonstrated to increase cell fitness for growth with dichloromethane, was also present in the transconjugants, no improvement of the fitness was observed for cells grown with chloromethane. This suggest that the effectiveness of heterologous catabolism of chloromethane and dichloromethane are uncorrelated in *Methylobacterium* strains (Michener *et al.*, submitted article).

There remains a need for a better understanding of the mechanisms involved in methylotrophic growth in *M. extorquens* using complementary approaches such as transcriptomics, proteomics or metabolomics in addition to the studies already available (Tableau 1.7 completed from Ochsner *et al.*, 2015).

45

« omic » study	Reference	Description			
Genome	Vuilleumier <i>et al.,</i> 2009	AM1, DM4			
	Roselli <i>et al.,</i> 2013	CM4			
	Muller <i>et al.,</i> 2011	DM4			
	Doi at al 2015	Isolates from phyllosphere and rhizosphere			
	Bai <i>et al.,</i> 2015	(24 genomes)			
Modelling	Peyraud <i>et al.,</i> 2011	AM1			
Proteomic	Bosch <i>et al.,</i> 2008;	AM1, methanol and succinate comparison DM4, dichloromethane et methanol comparison			
	Laukel <i>et al.,</i> 2004				
	Muller <i>et al.,</i> 2011				
	Roselli <i>et al.,</i> 2013	CM4, chloromethane et methanol comparison			
	o	AM1, « Profiling » of metabolites in methanol			
	Guo et Liastrom, 2008	and succinate			
Transcriptomic	Olucha at al. 2007	AM1, methanol and succinate comparison			
	Okubo <i>et al.,</i> 2007	(microarray)			
	Francez-Charlot et al.,	AM1, phyR mutant, general stress regulation			
	2009	(microarray)			
Metabolomic	Kiefor et al 2008 2011	AM1, central metabolites concentration in			
	Kielel <i>et ul.</i> , 2008, 2011	methanol and succinate			
	Dermand et al. 2012	AM1, [¹³ C]-fluxomic, [¹³ C]-CH ₃ OH in co-			
	Peyradu <i>et ul.,</i> 2012	utilization with succinate			
	No	AM1, ¹³ C methanol, metabolite labeling during			
	rang <i>et ul.</i> , 2013	growth			
	Cui et al 2016	AM1 with overexpression of ethylmalonyl-CoA			
	Cui <i>et u</i> i., 2010	mutase in methanol			
	Possor at al 2016	AM1, monitoring over time of metabolite after			
	Neaser et ul., 2010	incorporating [¹³ C]-CH ₃ OH			

Table 1.7. "Omics" studies in *M. extorquens* strains

4.3.2. Transcriptomic studies in *M. extorquens*

Transcriptomics is a qualitative and quantitative study of the transcriptome, which encompasses in principle all RNAs produced by transcription (Wang *et al.*, 2009). In different growth conditions, the spectrum of transcripts and their abundance is modified. Thus, the quantitative inventory of transcripts enables a better understanding of the adaptive processes involved in response to growth in various conditions.

Today, few transcriptomic studies have been performed in *M. extorquens*, and when performed, it was using DNA chips (microarrays) in strain AM1. These studies demonstrated the role of genes involved in C₁ metabolism in comparative studies of cultures grown with methanol versus succinate (Okubo *et al.*, 2007; Francez-Charlot *et al.*, 2009), as well as the involvement in the stress response of the transcriptional gene regulator *phyR* (Francez-Charlot *et al.*, 2009). Sequencing and analysis of RNA (RNA sequencing, i.e. a global transcriptome analysis) has not been described so far for *M. extorquens*, and I provide in this doctoral work the first transcriptome study for strains CM4 and DM4.

4.3.3. Proteomic studies in *M. extorquens*

There are more published proteomic than transcriptomic studies in *M. extorquens* (Tableau 1.7, Gourion *et al.*, 2006; Muller *et al.*, 2011; Roselli *et al.*, 2013). In *M. extorquens* AM1, proteins involved in methylotrophy have been detected in cultures grown with methanol (the reference growth substrate for methylotrophy studies) compared to cultures grown with succinate ($C_4H_6O_4$) (Laukel *et al.*, 2004; Bosch *et al.*, 2008). Other studies were performed in *M. extorquens* strains able to degrade chlorinated methanes. In *M. extorquens* DM4, the protein content of cultures grow with dichloromethane compared to methanol was used to identify proteins involved in dichloromethane utilization (see chapter 3) (Muller *et al.*, 2011a). Similarly, *M. extorquens* AM1 colonization mechanisms of

A. thaliana phyllosphere has been better understood using proteomic approaches (Tableau 1.7, Knief *et al.*, 2010, 2012; Vorholt, 2012). Overexpression of the regulatory protein PhyR was employed to evidence its role in plant colonization (Gourion *et al.*, 2006).

Proteomic analysis in *M. extorquens* CM4 chloromethane utilization was used to demonstrated that 49 proteins more abundant in cultures grown with chloromethane compared tomethanol (Figure 1.17).



Figure 1.17. 2D gel picture of protein extracts of *M. extorquens* CM4 grown with chloromethane or with methanol

CmuA, CmuB and PurU framed in red, dark green and light green respectively, were identified by mass spectrometry and were more abundant with chloromethane (Sandro Roselli PhD thesis, 2009).

By combining genomic and proteomic approaches, genes involved in chloromethane utilization highlighted the central implication of plasmid pCMU01 in growth with chloromethane in *M. extorquens* CM4 (Roselli *et al.*, 2013).

4.3.4. Metabolomic studies in *M. extorquens*

Most of the metabolomic studies were performed with cultures of *M. extorquens* AM1 to better characterize the methylotrophic and central metabolism of carbon assimilation. In cultures grown with methanol compared tosuccinate, metabolites specifically associated to growth in methylotrophic or involved in glyoxylate regeneration such as β -hydroxybutyrate, methylsuccinyl-CoA or ethylmalonyl-CoA have been detected (Figure 1.11) (Guo et Lidstrom, 2008). Metabolites profiling were also performed in cultures of

M. extorquens AM1 grown with compounds in C₂ (ethylamine) and in C₄ (succinate) (Yang *et al.*, 2009). Similarly, *M. extorquens* AM1 grown with acetate (CH₃COO⁻) as the sole source of energy and carbon was used to demonstrated that the ethylmalonyl-CoA pathway replaced the isocitrate lyase pathway for the regeneration of glyoxylate, needed for carbon assimilation in biomass via the serine cycle (Schneider *et al.*, 2012, Figure 1.11). These results illustrate metabolic adjustments to growth with different carbon sources.

5. PhD thesis aims and questions

Since the Montreal protocol in 1987, limiting the utilization of chlorinated compounds, chloromethane is responsible for 16% of ozone layer destruction related to chlorinated compounds (Montzka et al., 2011). The global budget of chloromethane remains uncertain even if sources and sinks are identified (Tableau 1.4 and Tableau 1.5, (Montzka and Fraser, 2003; Clerbaux et al., 2007). The role of the soil, both source and sink of chloromethane, in this global budget is shown (Gribble, 2003; Harper et al., 2003; Harper, 2000; Miller et al., 2004). Otherwise uncertainties remain regarding the importance of the soil contribution in chloromethane exchanges with the atmosphere (Saito et al., 2008). One of those uncertainties comes from the bacterial capacity to play a role as a chloromethane emission filter, limiting chloromethane into the atmosphere. Otherwise, low chloromethane concentrations available in forest soil (1 pM. g⁻¹) (Harper et al., 2003) could not allow enough to support an efficient growth for chloromethane degrader communities, requiring other carbon and energy sources to grow. In fact, methanol concentration, estimated at 1 nM. g⁻¹, in forest soils, is higher than those of chloromethane (Figure 1.18, Stacheter and Kolb, 2013). In the environment, methylotrophic microorganisms that use chloromethane could also use methane and methanol. Methane and methanol emissions form terrestrial ecosystems are within same range (10¹² mol per year) (Kolb, 2009a). Methanol is a central compound of methylotrophic metabolism, as more than 83% of methylotrophic strains growing in aerobic conditions were isolated from soil and were able to grow using methanol (Kolb, 2009). It is also shown in the strain *M. extorquens* CM4, which is unable to use methane (Van Aken, 2004). One of the objectives of my PhD thesis was to characterize bacterial populations within forest soil that are able to use chloromethane, and to test the hypothesis of chloromethane and methanol co-utilizations. My PhD project was articulated around two major axes. First, an in depth study of chloromethane utilization in controlled laboratory conditions. The sequencing of transcripts (RNAseq), was done to identify genes differentially expressed in M. extorquens CM4 in methylotrophic growth conditions with chloromethane and methanol. The second part of my PhD work, based on a Stable Isotope Probing (SIP) approach (Neufeld et al., 2007), consisted of highlighting methylotrophic bacteria active in forest soil, and its ability to degrade chloromethane. In this case, incubations with chloromethane and/or methanol labelled with [¹³C], were created to detect bacteria able to assimilate those compounds, and to incorporate the [13C] into the biomass. Enriched [13C]-DNA ("Heavy" DNA), corresponding to the DNA of microorganisms assimilating the labelled-carbon source, was separated from the DNA not labelled ("Light" DNA constituted only by [¹²C]) by ultracentrifugation. Apart from the heavy DNA, the bacterial diversity is estimated by sequencing of PCR products of taxonomic and functional markers.

Three gene markers were used targeting:

- Aerobic degradation of the chloromethane (*cmuA* gene encoding a chloromethane methyltransferase belonging to the *cmu* pathway)
- Aerobic degradation of the methanol (*mxaF* and *xoxF* genes encoding a methanol dehydrogenase subunit)
- Central methylotrophy metabolism (*mch* gene) (Stacheter and Kolb, 2013).

The main goal of those combined approaches was:

To have a better understanding of chloromethane utilization in model strains cultivated in the laboratory, to identify new biomarkers of chloromethane degradation, and to detect more efficiently new biomarkers of chloromethane degradation in forest soils.

Identify new bacterial methylotrophy taxa, active in aerobic chloromethane consumption in sub-surface forest soil microcosms, and the study of their functions within the bio-capture of terrestrial chloromethane emissions.



Figure 1.18. Sinks and emissions of chloromethane and methanol in forest soil Chloromethane emissions data from Harper *et al.*, 2003. Methanol fluxes from Stacheter *et al.*, 2013.

Chapitre 1

Introduction

Chapter 1. Introduction (en français)

Ce premier chapitre débutera par la présentation des méthanes chlorés en général, et en particulier du chlorométhane, avec la description de leurs propriétés chimiques, leurs utilisations industrielles, et leurs impacts sur la santé et l'environnement. Suivront des paragraphes dédiés spécifiquement au chlorométhane, à ses sources de production et ses puits de consommation, notamment par les microorganismes. Les connaissances sur la dégradation microbienne du chlorométhane seront ensuite présentées dans le contexte du métabolisme des composés à un atome de carbone et la génétique de souches modèles méthylotrophes. Enfin, le dernier paragraphe posera les problématiques et les objectifs de mes travaux de thèse.

<u>1.</u> <u>Les méthanes chlorés</u>

Les méthanes chlorés sont des composés organiques ne comportant qu'un seul atome de carbone relié à un ou plusieurs atomes de chlore. Un atome de carbone possède quatre liaisons covalentes, il existe donc quatre méthanes chlorés, le chlorométhane (CH₃Cl), le dichlorométhane (CH₂Cl₂), le chloroforme (CHCl₃) et le tétrachlorométhane (CCl₄). A l'exception du chlorométhane qui se trouve à l'état de gaz à température ambiante, les autres méthanes chlorés sont liquides et incolores (Huang *et al.*, 2014). Ces composés ont des propriétés physico-chimiques différentes et sont, depuis le 20^e siècle, produits et utilisés en grande quantité dans l'industrie (Tableau 1.1). Les méthanes chlorés peuvent également être produits naturellement. Les productions naturelles de chloroforme et de dichlorométhane sont estimées respectivement à 0,56 et 0,25 Tg.an⁻¹ (Cox *et al.*, 2003; Xiao, 2008). Cependant, le chlorométhane est le méthane chloré le plus présent dans l'atmosphère avec 2,8 Tg.an⁻¹ (Ruecker *et al.*, 2014) (Tableau 1.2).

Composé	Apparence	Solubilité dans l'eau (g.L ⁻¹)	Effet sur la santé ^a	Polluant ^b	Utilisation dans l'industrie ^c
chlorométhane	gaz incolore	5,3	groupe 3	non	production de silicone, produits de construction
dichlorométhane	liquide incolore	13,0	groupe 2B	oui	décapant pour peinture, produits pharmaceutiques
chloroforme	liquide incolore	8,0	groupe 2B	oui	solvants, synthèse de produits chimiques
tétrachlorométhane	liquide incolore	0,8	groupe 2B	oui	produits pharmaceutiques, produits de nettoyage, produits agrochimiques

Tableau 1.1. Propriétés et	utilisations des	méthanes	chlorés
----------------------------	------------------	----------	---------

^a Données issues de l'Agence internationale de recherche contre le cancer (IARC). Le groupe 2B regroupe les composés probablement cancérigènes pour l'Homme. Le groupe 3 regroupe les composés non classifiables comme carcinogène pour l'Homme (Huang *et al.*, 2012). ^b Composés reconnus comme polluants par les Etats-Unis, la Chine et pour le dichlorométhane et le chloroforme reconnus également par l'Union Européenne (Huang *et al.*, 2014).

^c Données issues de Frascari *et al.,* 2015.

Les méthanes chlorés sont présents dans les sols, l'eau et l'air (http://www.epa.gov). A l'exception du chlorométhane, ils ont été classés par l'IARC (International Agence for Research on Cancer) comme étant potentiellement cancérigènes, et comme polluants par de nombreux pays (Tableau 1.1). L'impact des composés chlorés dans l'environnement fait qu'ils sont très étudiés (Harper, 2000).

	Troposphère		Flux (Gg. an ⁻¹)	
	Conc. (pptv)	Demi-vie (année)	Industrie	Naturel
chlorométhane	$550\pm30^{\text{a}}$	1,3	10 ^a	1 900ª
dichlorométhane	$25\pm5^{\text{d}}$	0,38	530 ^e	160 ^e
chloroforme	15 ± 5^{c}	0,41	60 ^b	595 ^b
tétrachlorométhane	$95\pm5^{\mathrm{f}}$	26	26 46 ^g	
D'après manuscrit de thèse de S. Roselli (2009)				

Tableau 1.2. Les méthanes chlorés dans la troposphère

D'après manuscrit de thèse de S. Roselli (^a Montzka et Fraser, 2003 ^b McCulloch *et al.*, 2003 ^c O'Doherty *et al.*, 2001 ^d Schauffler, 2003 ^e Kurylo et Rodriguez, 1998 ^f Prinn *et al.*, 2000 ^g UNEP, 2005

Dans la stratosphère, les méthanes chlorés sont activés par les ultra-violets. Il en résulte la formation de radicaux Cl⁻, qui vont contribuer à la déplétion de la couche d'ozone en catalysant la formation d'O₂ (Figure 1.1). Il est estimé qu'un atome de chlore peut détruire 100 000 molécules d'ozone (http://www.epa.gov/ozone/science/process.html), avec pour exemple de cet impact, la formation d'un trou dans la couche d'ozone au-dessus de l'Antarctique à chaque printemps (Clerbaux *et al.*, 2007).



Figure 1.1. Réactions de formation et de destruction de l'ozone dans l'atmosphère En présence de méthanes chlorés (CH_xCl_x), les réactions naturelles de l'ozone (O₃) sont perturbées. Rayons UVA entre 315 nm et 400 nm et rayons UVB entre 280 nm et 315 nm (Figure modifiée de http://www.cec.org)

En raison de leur impact sur l'environnement, l'utilisation des composés halogénés a été régulée par le protocole de Montréal en 1987, diminuant ainsi grandement leur utilisation dans l'industrie, allant jusqu'à un arrêt total de la production de composés comme le tétrachlorométhane en 1995. Ceci a eu pour conséquence d'augmenter la part du chlorométhane, le C₁ chloré le plus produit naturellement, dans l'impact environnemental des méthanes chlorés. En effet, parmi les méthanes chlorés retrouvés dans la troposphère, le chlorométhane est le plus abondant (Tableau 1.2). Aujourd'hui, on estime qu'il est responsable de 16 % de la dégradation de la couche d'ozone liée aux composés chlorés (Montzka *et al.*, 2011).

2. <u>Le chlorométhane</u>

Le budget global du chlorométhane, c'est-à-dire sa concentration dans l'atmosphère, est défini par la somme de ses sources de production, soustrait de l'ensemble des puits de dissipation. La diversité des sources et puits abiotiques et biotiques du chlorométhane est détaillée ci-dessous. Ce budget reste cependant incertain notamment du fait de la part des bactéries dans le filtrage du chlorométhane émis dans différents environnements.

2.1. Les sources de chlorométhane

La production totale annuelle est estimée à 2,8 Tg (Ruecker *et al.*, 2014; Sailaukhanuly *et al.*, 2014). Comme plus de 5 000 substances halogénées, le chlorométhane est produit majoritairement naturellement. La production naturelle de chlorométhane repose sur des sources abiotiques et biotiques (Tableau 1.3). Sa faible production d'origine anthropogénique est due à la combustion de matières fossiles comme le charbon, la combustion de déchets municipaux, ou à d'autres activités industrielles (Tableau 1.3, McCulloch et Aucott, 1999).

Origino	Estimation	Valeurs basses et		
Ongine	(Gg. an⁻¹) ^a	hautes (Gg. an ⁻¹) ^b		
Senescence végétale	1800 ^d	30/2500		
Feux de forêts	911	655/ 1125		
Plantes tropicales	910	820/ 8200		
Océans	600	325/ 1300		
Marais salants	170	65/ 440		
Champignons	160	43/ 470		
Combustion de matière organique	105	F / 20F		
fossile	105	5/ 205		
Incinération	45	15/ 75		
Zones humides	40	6-270 ^c		
Industrie	10	10		
Rizières	5	2,4/ 4,9		
Total des sources	4746	2019/ 14378		

Tableau 1.3. Estimation des sources de chlorométhane dans l'atmosphère

^a Données issues de (Keppler *et al.*, 2005)

^b Données issues de (Clerbaux *et al.,* 2007)

^c Données issues de (Montzka et Fraser, 2003)

^d Données issues de (Montzka *et al.,* 2011)

2.1.1. Production abiotique

Les feux de forêts constituent la plus grande source abiotique de chlorométhane (Sailaukhanuly *et al.*, 2014). Le chlorométhane produit est émis dans l'atmosphère mais également retrouvé à la surface des sols, où des processus abiotiques de production de méthanes halogénés ont également été caractérisés (Keppler *et al.*, 2005). Une production abiotique de chlorométhane par des réactions redox (Figure 1.2) et des réactions de substitution ont été démontrées (Hamilton, 2003; Keppler *et al.*, 2005; Wishkerman *et al.*,

2008) bien que leur part dans le bugdet du chlorométhane n'ait pas été évaluée. Les océans représentent la 2^e source abiotique de chlorométhane (Xiao, 2008; Hu *et al.*, 2010).



Figure 1.2. Formation abiotique du chlorométhane dans les sols

Formation des alkyls halogénés par la réduction du Fe (III) avec la matière organique en présence d'ions chlorure dans les sols (issue de Keppler *et al.,* 2003).

2.1.2. Production biotique

La production principale de chlorométhane résulte de processus biotiques chez les plantes (Hamilton, 2003; Saito *et al.*, 2008), les champignons (Moore *et al.*, 2005), les algues marines (Traunecker *et al.*, 1991) ou encore les marais salants (Rhew *et al.*, 2003) (Tableau 1.3). Les écosystèmes terrestres contribuent largement à cette production, avec plus de

50 % des émissions de chlorométhane qui en sont issues (Redeker et Kalin, 2012). Dans l'environnement, la présence d'ions chlorure est très variable et pourrait influencer les taux d'émissions de chlorométhane. Les premiers résultats d'une corrélation possible évoquée dans la littérature sont décrits ci-après ainsi que l'état des connaissances des productions du chlorométhane par les plantes d'une part et dans les sols d'autre part. On situera également les taux d'émissions du chlorométhane par rapport aux autres Composé Organiques Volatiles (COVs) et le développement d'un bio-rapporteur pour sa quantification.

Production par les plantes

Le mécanisme génétique de production du chlorométhane a été caractérisé en détails chez la plante modèle *Arabidopsis thaliana* (Rhew *et al.*, 2003). Cette étude montre la production de chlorométhane mais aussi de bromométhane (CH₃Br) et d'iodométhane (CH₃I), par l'action d'une méthyltransférase S-adénosylméthionine (SAM)-dépendante nommée *HOL* pour « *Harmless to Ozone Layer* ». Le gène *HOL* est conservé chez les plantes, laissant penser que la production de chlorométhane par les plantes est un processus répandu (Rhew *et al.*, 2003;

Nagatoshi et Nakamura, 2007). Le chlorométhane est produit par transfert du groupement méthyl du SAM à un ion chlorure (Cl⁻) (Figure 1.3). La SAM intervient dans de nombreuses réactions de synthèse de la lignine, un constituant de la paroi cellulaire végétale (Campbell et Sederoff, 1996). Rhew *et al.*, 2003 a démontré que la méthyltransférase codée par le gène *HOL* catalyse la méthylation du chlorure, dépendamment du SAM.





De fait, différentes études ont démontré l'existence de plantes émettrices de chlorométhane. Par exemple, dans une région subtropicale, sur 187 plantes étudiées, 33 espèces appartenant à des familles différentes émettent du chlorométhane (Yokouchi *et al.*, 2007). D'une manière similaire, l'étude des flux de chlorométhane en forêt tropicale a montré que 25 plantes sur les 117 étudiées émettent du chlorométhane (Saito *et al.*, 2008). Les émissions les plus importantes ont été observées chez des fougères (*Osmunda banksiifolia, Cibotium balometz, Angiopteris palmiformis*), et des plantes halophiles (*Vitex rotundifolia, Vitex trifolia* et *Excoecaria agalloch*) avec des taux d'émission de l'ordre de 1 µg de chlorométhane.g⁻¹ de matière sèche. h⁻¹. Toutefois, la capacité de production de chlorométhane dépend plus de l'espèce que de la famille à laquelle est affiliée la plante (Yokouchi *et al.*, 2007).

Production de chlorométhane par le sol et les matières végétales en décomposition
Les plantes en décomposition représentent la source majeure du chlorométhane
(Hamilton, 2003, Montzka *et al.*, 2011). La production de chlorométhane est surtout localisée
dans l'horizon « O » des sols et semble être quasi nulle dans les horizons inférieurs (Figure 1.4)
(Redeker et Kalin, 2012).

61



Figure 1.4. Organisation schématique du sol forestier

Le sol forestier type est organisé en différents horizons; O, A, E, B, C et R. L'horizon « O » est surtout constitué de matière organique en décomposition, l'horizon « A » constitué également de matière organique et de matière minérale, l'horizon « E » est un horizon lessivé, où les nutriments solubles ne sont plus présents en raison des précipitations ou de l'irrigation. L'horizon « B » est un horizon riche en minéraux issus du lessivage des horizons « A » et « E ». L'horizon « C » est un horizon de sous-sol, non consolidé. Enfin l'horizon R est un horizon rocheux, constitué de roche comme le granit ou encore du calcaire. Les horizons « O » et « A » sont biologiquement actifs et sont fortement oxygénées, tandis que les autres sont plus pauvres en matière organique et ont une activité microbienne plus faible. Dans le cadre de mon projet de thèse, un travail sur le sol forestier a été effectué au niveau de l'horizon « O ». (Figure modifiée de http://www.ctahr.hawaii.edu/mauisoil/a_profile.aspx)

Le sol est un environnement riche en ions chlorure, dont la biodisponibilité est essentielle à la formation du chlorométhane. Cependant, l'estimation de la concentration des ions chlorure dans les sols reste difficile car de nombreux mécanismes interviennent comme les flux entre la végétation et le sol, l'altération de la roche, ou encore la dégradation de la matière organique (Bastviken *et al.*, 2007; Öberg, 2003). Il est admis qu'une partie des ions chlorure est considérée comme non biodisponible du fait de son adsorption ou de son absorption par la végétation ou les microorganismes (Bastviken *et al.*, 2007). Les ions chlorure sont aussi contenus dans le bois et sont utilisés pour la synthèse de chlorométhane par les champignons les colonisant (Watling et Harper, 1998). On estime que le contenu en Cl⁻ dans le bois d'espèce de forêts tempérées varie entre 2,4 et 123 mg.kg⁻¹ de matière sèche (Watling et Harper, 1998).

Des études sur des champignons ont démontrées que des membres de la famille des *Hymenochaetaceae* sont capables de réémettre sous forme de chlorométhane une partie des ions chlorure présents dans leurs substrats (Harper et Kennedy, 1986). En effet, le bois mort en décomposition est colonisé par les champignons qui, via un ensemble de péroxydases, dégradent la lignine, un constituant majeur du bois, en SAM. La SAM, par interaction avec les ions Cl⁻ présents dans le milieu va, par l'intermédiaire d'une méthyltransférase, former du chlorométhane (Harper, 2000; Anke et Weber, 2006). Cette production a été étudiée et décrite chez le champignon *Phellinus tuberculosus* (White, 1982; Wuosmaa et Hager, 1990), cependant l'exemple le plus parlant est celui de *Phellinus pomaceus* (Figure 1.5) capable de réémettre 90 % des ions chlorure présents dans ses substrats sous forme de chlorométhane (Harper *et al.*, 1988). Cette production de chlorométhane se fait via transfert d'un ion Cl⁻ sur le groupement méthyl de la SAM (Figure 1.5).



Figure 1.5. Production de chlorométhane par le champignon *Phellinus pomaceus* (Figure modifiée de Harper, 2000)

Il existe également une production de chlorométhane par les champignons associés aux systèmes racinaires des sols forestiers. Les racines représentent une source importante de COVs (Lin *et al.*, 2007). Dans l'étude de Redeker *et al.*, 2004, 9 isolats de champignons ectomycorhiziens, dont *Cenococcum geophilum*, l'une des espèces les plus répandues et
communes (Trappe, 1977; Massicotte *et al.*, 1992), ont démontré leur capacité à produire des méthanes halogénés dont le chlorométhane. Ils produisent entre 0,003 µg et 65 µg de chlorométhane par gramme de matière sèche (Redeker *et al.*, 2004). Cependant, il reste difficile d'évaluer séparément le rôle du système racinaire, car il est étroitement lié au système microbien, lui-même influençant les flux de COVs dans les sols (Peñuelas *et al.*, 2014). En effet, plus de 95 % des racines courtes des plantes sont colonisées par des champignons, eux-mêmes localisés à proximité des communautés bactériennes aidant les champignons de la rhizosphère « *mycorrhiza helper bacteria* » notamment lors des processus de développement mycorhyzien (Frey-Klett *et al.*, 2007; Bonfante et Anca, 2009; Rigamonte *et al.*, 2010).

Le chlorométhane est faiblement émis par rapport à d'autres COVs

Les plantes réémettent une partie du carbone disponible sous forme de Composés Organiques Volatiles (COVs) dont le méthanol (Bringel et Couée, 2015; Vorholt, 2012). Ces composés sont émis dans l'atmosphère ou les sols, où ils peuvent ensuite être directement assimilés par les microorganismes (Schade et Goldstein, 2001). Ainsi, la majeure partie du méthanol atmosphérique serait liée à la production par les plantes (Jacob, 2005; Galbally et Kirstine, 2002). D'autres COVs sont produits par les plantes, dont des composés aromatiques mais également des composés halogénés (Guenther *et al.*, 2012; Forczek *et al.*, 2015). La production par les plantes chlorés a été décrites, bien que le chlorométhane soit le plus produit avec une production estimée de 0,8 à 8 Tg.an⁻¹ (Clerbaux *et al.*, 2007). Cette production reste inférieure à celle de méthanol (50-132 Tg. an⁻¹) (Jacob, 2005; Forczek *et al.*, 2015).

Au niveau de sols forrestiers, le chlorométhane est loin d'être le seul COV produit (Keppler *et al.*, 2005; Redeker et Kalin, 2012; Peñuelas *et al.*, 2014). Le sol est un milieu hétérogène, constitué de microenvironnements complexes (Peñuelas *et al.*, 2014) avec une diversité microbienne importante. La diversité bactérienne et des champignons est évaluée par le nombre d'espèces présentes, soit respectivement de 10⁵ et 10⁶ espèces par gramme de sol (Peñuelas *et al.*, 2014). De nombreux COVs sont émis par ces microorganismes (Figure 1.6). Cette production est directement liée à la disponibilité des substrats et des conditions de croissance (Kai *et al.*, 2010; Blom *et al.*, 2011). Une grande partie des COVs émis provient de la dégradation microbienne de la litière végétale (matière morte végétale), de la dégradation

de la matière organique (Kögel-Knabner, 2002; Peñuelas *et al.*, 2014) ainsi que des exsudats racinaires.



Figure 1.6. Les composés organiques volatiles émis dans les sols

Ceux émis par les bactéries sont représentés en rouge, et ceux émis par les champignons sont représentés en bleu (modifiée de Peñuelas *et al.,* 2014). L'ensemble de ces COVs est répertorié dans une base de données (http://bioinformatics.charite.de/mvoc/). La flèche rouge représente la catégorie de COVs dans laquelle se situe le chlorométhane.

De manière générale, les bactéries produisent plus d'alcènes, d'alcools, de cétones et de terpènes, tandis que les champignons produisent plus d'alcools, de benzénoïdes, d'acides et d'aldéhydes (Figure 1.6). Les composés chlorés sont moins produits, et cette production dans les sols est plus importante par les champignons que par les bactéries (Figure 1.6). Le chlorométhane produit est classé dans le groupe « chlorides » (indiqué par la flèche rouge).

Dosage du chlorométhane à l'aide d'une souche bio-rapportrice

La quantification de la très faible production de chlorométhane par les plantes et le sol (de l'ordre du ppt) est difficile par des approches classiques de dosage GC « *gas chromatography* » pas assez sensible (Figure 1.7). Au laboratoire à Strasbourg, dans le but d'évaluer les émissions de chlorométhane par les plantes, *Methylobacterium extorquens* CM4 a été transformée pour être capable d'émettre de la fluorescence, proportionnellement à la faible concentration de chlorométhane (Farhan UI Haque *et al.*, 2013). Cette souche bio-rapportrice est basée sur la présence d'un plasmide portant un gène *yfp*, codant une protèine fluorescente sous contrôle du promoteur du gène *cmuA* (Figure 1.7) et permet d'apporter une réponse sensible et proportionnelle à des concentrations de chlorométhane, avec un seuil de détection de 2 pM.

A l'aide de ce bio-rapporteur, la production de chlorométhane chez la plante *A. thaliana* et chez *V. rotundifolia* a pu être démontrée et quantifiée avec des taux d'émission respectifs de 13 et 2 800 ng.g de matière séche⁻¹. h⁻¹ (Farhan Ul Haque *et al.*, 2013).



Figure 1.7. Souche bio-rapportrice pour la détection du chlorométhane

(A) Dans la souche bio-rapportrice, un plasmide porteur du gène *yfp* placé sous le contrôle du promoteur du gène *cmuA* (codant une protéine avec un domaine méthyltransférase et un de liaison corrinoïde, esentielle à l'utilisation du chlorométhane chez *M. extorquens* CM4 (Studer *et al.*, 2001, 2002)), induit en présence de chlorométhane, est présent dans *M. extorquens* CM4 (Farhan Ul Haque *et al.*, 2013). Le plasmide porte un gène de résistance à la kanamycine pour maintenir la présence de ce plasmide lors de la propagation de la souche bio-rapportrice.
(B) L'émission de fluorescence de la protéine Yfp est spécifique à la présence de chlorométhane. En jaune, le seuil de détection avec la souche bio-rapportrice et en noir avec le dosage GC.

2.2. Les puits de chlorométhane

2.2.1. Dégradation abiotique du chlorométhane

Différents puits abiotiques de chlorométhane ont été identifiés et contribuent à des degrés plus ou moins importants à l'élimination du chlorométhane (Tableau 1.4). Bien qu'une dégradation abiotique de COVs ait été observée dans les sols, celle du chlorométhane n'a pu y être décrite (Miller *et al.*, 2004; Insam et Seewald, 2010).

Nature des puits Estimation (Gg. an ⁻¹) ^a	Valeurs basses et hautes (Gg. an ⁻¹) ^b
--	--

Tableau 1.4. Estimation des puits de chlorométhane dans l'atmosphère
--

Puits	
-------	--

Réaction avec les OH de la troposphère	-3 180	-3800/ -4100		
Perte dans la stratosphère	-200	-100/ -300		
Réaction avec les Cl ⁻ à la surface des	-270	-18/-550		
océans	-370	-18/ -330		
Dégradation microbienne dans le sol	< -1 000	-100/ -1600		
Perte dans les océans polaire	-75	-93/-145		
Total des puits	< -4 875	-4273/ -6695		

^a Données issues de (Keppler et al., 2005)

^b Données issues de (Clerbaux *et al.,* 2007)

2.2.2. Dégradation biotique du chlorométhane

La dégradation biotique du chlorométhane est principalement liée à l'activité microbienne même si son estimation reste très variable (entre 180 et 1600 Gg.an⁻¹) (Keppler *et al.*, 2005).

Dégradation bactérienne dans les environnements marins

Les environnements marins contribuent à la dégradation du chlorométhane (Tableau 1.4). Des bactéries issues d'environnements marins, capables d'utiliser le chlorométhane, ont été isolées. C'est le cas de *Leisingera methylohalidovorans* MB2, premier isolat des *Rhodobacteracea* à démontrer cette capacité (Goodwin *et al.*, 1997; Schaefer; 2002, Tableau 1.5). Trois autres souches appartenant à cette famille et dégradant le chlorométhane ont été isolées d'environnements marins; *Roseovarius* sp. 179, *Roseovarius* sp. 217 et *Ruegeria* sp. 198 (Schäfer *et al.*, 2005; Tableau 1.5).

Chapitre 1 (en français)

Tableau 1.5. Souches bactérienne chlorométhane-dégradantes isolées de différents environnements

Souche bactérienne	Origine	Gram	Métabolisme	Type trophique	Présence de <i>cmuA</i> ª	Références
Acetobacterium dehalogenans MC	Boues activées	positive	anaérobie	Homoacétogène	nd	Traunecker <i>et al</i> ., 1991
Aminobacter ciceronei IMB1	Champ de fraises fumigé	négative	aérobie	Méthylotrophe facultatif	oui	Hancock <i>et al.,</i> 1998
Aminobacter lissarensis CC495	Sol forestier	négative	aérobie	Méthylotrophe facultatif	oui	Coulter <i>et al.,</i> 1999
Hyphomicrobium sp. AT1	Phyllosphère	négative	aérobie	Méthylotrophe facultatif	oui	Nadalig <i>et al</i> ., 2011
Hyphomicrobium sp. AT2	Phyllosphère	négative	aérobie	Méthylotrophe facultatif	oui	Nadalig <i>et al</i> ., 2011
Hyphomicrobium sp. AT3	Phyllosphère	négative	aérobie	Méthylotrophe facultatif	oui	Nadalig <i>et al</i> ., 2011
Hyphomicrobium sp. AT4	Phyllosphère	négative	aérobie	Méthylotrophe facultatif	oui	Nadalig <i>et al</i> ., 2011
Hyphomicrobium sp. MC1	Station d'épuration industrielle	négative	aérobie	Méthylotrophe facultatif	oui	Hartmans et al., 1986
Hyphomicrobium sp. MC2	Sol d'usine pétrochimique	négative	aérobie	Méthylotrophe facultatif	oui	Doronina <i>et al</i> ., 1996
Leisingera methylohalidovorans MB2	Bassin de marée	négative	aérobie	Méthylotrophe facultatif	non	Schaefer, 2002
Methylomicrobium album BG8	Eau douce	négative	aérobie	Méthylotrophe obligatoire	nd	Han et Semrau, 2000
Methylobacterium extorquens CM4	Sol d'usine pétrochimique	négative	aérobie	Méthylotrophe facultatif	oui	Doronina <i>et al.,</i> 1996
Pseudomonas aeruginosa NB1	Boues activées	négative	anaérobie	Méthylotrophe facultatif	nd	Freedman <i>et al</i> ., 2004
Roseovarius sp. 179	Eau de mer côtière	négative	aérobie	Méthylotrophe facultatif	oui	Schäfer <i>et al.,</i> 2005
Roseovarius sp. 198	Eau de mer côtière	négative	aérobie	Méthylotrophe facultatif	oui	Schäfer <i>et al.,</i> 2005
Roseovarius sp. 217	Eau de mer	négative	aérobie	Méthylotrophe facultatif	oui	Schäfer <i>et al.,</i> 2005

^a nd; non determiné

Dégradation bactérienne dans la phyllosphère

La phyllosphère correspond à la partie aérienne des plantes et constitue un des habitats microbiens les plus importants. La surface totale des faces inférieures et supérieures des feuilles est égale à deux fois la surface terrestre avec une densité bactérienne de 10⁷.cm⁻² (Vorholt, 2012). Peu d'études sont disponibles sur le rôle des microorganismes de la phyllosphère comme puits de chlorométhane. Des bactéries capables de l'utiliser ont été isolées à la surface des feuilles de la plante *A. thaliana* (Nadalig *et al.*, 2011), mais le rôle potentiel de filtre de chlorométhane par les bactéries de la phyllosphère n'est pas pris en compte pour le moment.

Dégradation bactérienne dans les sols

Le sol contient de nombreux composés, dont certains halogènes, et constitue un puits majeur pour certains d'entre eux. Ainsi, 70 % du bromométhane (CH₃Br) est consommé dans les sols (Shorter *et al.*, 1995). La dégradation du chlorométhane dans les sols est mal évaluée mais est estimée comme étant supérieure à 1000 Gg. an⁻¹ (Tableau 1.4). Cette consommation dans les sols est liée à une dégradation bactérienne mais aussi fongique et serait majoritaire dans l'horizon « O » et est plus faible dans les horizons inférieurs (Redeker et Kalin, 2012).

Les échanges de chlorométhane entre les écosystèmes terrestres et l'atmosphère sont modulés par les bactéries présentes dans les sols forestiers (Miller *et al.*, 2004; Borodina *et al.*, 2005; Keppler *et al.*, 2005; Clerbaux *et al.*, 2007; Schäfer *et al.*, 2007; Rhew *et al.*, 2010). Différentes bactéries appartenant aux alpha et béta-protéobactéries et capables d'utiliser le chlorométhane, ont été isolées de sols forestiers mais leur contribution dans le budget du chlorométhane n'est pas clairement définie (Miller *et al.*, 2004; Borodina *et al.*, 2005).

Cette consommation de chlorométhane est également liée aux champignons. En effet, ces derniers produisent des métabolites secondaires, dont plus de 200 sont des métabolites halogénés (Gribble, 2003), incluant des métabolites bromés, fluorés, iodés ou chlorés (Anke et Weber, 2006). Ces métabolites chlorés, dont le chlorométhane, peuvent être éliminés lors de réactions de méthylation chez les champignons (Figure 1.8). La méthylation est un processus fréquent de la biosynthèse de métabolites par les champignons (Anke et Weber, 2006).



Figure 1.8. Réactions de dégradation du chlorométhane par les champignons (Modifiée de Anke et Weber, 2006)

Cette méthylation peut se faire chez les champignons via le transfert du groupement méthyl de la SAM comme dans la synthèse du chlorométhane chez les plantes, mais il semble que l'utilisation de chlorométhane soit privilégiée par rapport au SAM, car moins coûteuse d'un point de vue énergétique (Harper, 2000). Plus particulièrement le chlorométhane sert de donneur de groupement méthyle sur des fonctions hydroxy (OH) ou carboxy (COOH), permettant ainsi la synthèse de métabolites, comme l'alcool de vératryl ou le benzoate de méthyl (Figure 1.8). L'alcool de vératryl est un métabolite secondaire intervenant dans la synthèse d'enzymes nécessaires à la dégradation de la lignine (Dekker *et al.*, 2001). Le benzoate de méthyl est un composé olfactif synthétisé pour attirer les pollinisateurs ou encore jouant un rôle dans la réduction des racines chez *A. thaliana* (Horiuchi *et al.*, 2007). Cependant, le bilan net du chlorométhane chez les champignons (Anke et Weber, 2006). Bien que l'importance du sol comme puits du chlorométhane ne soit plus à démontrer, de nombreuses incertitudes persistent quant au bilan net de chlorométhane dans les sols (voir paragraphe suivant).

2.3. Le budget global du chlorométhane du sol

Le sol est un environnement complexe rendant difficile une estimation nette du bilan global des puits et sources de chlorométhane. En plus de la grande hétérogéneité intrinsèque du sol (Figure 1.4), le sol est soumis à la variation de nombreux paramètres aussi bien physicochimiques, climatiques que biologiques. Par exemple, un effet saisonnier sur la production de chlorométhane a été décrit (Peñuelas *et al.*, 2014). En effet, une augmentation des émissions de chlorométhane a été observée durant l'été (Llusia *et al.*, 2013; Oderbolz *et al.*, 2013), expliquée par le fait qu'en automne l'émission des COVs, dont le chlorométhane, vers l'atmosphère est atténuée par la chute des feuilles restant à la surface du sol (Figure 1.9). La température a également un impact sur la production du chlorométhane. La libération du chlorométhane par du matériel végétal en décomposition est plus importante lorsque la température augmente (Hamilton, 2003).



Figure 1.9. Flux du chlorométhane dans un sol forestier en fonction des saisons Mesure du chlorométhane émis par un sol forestier (µg.m⁻³. jour⁻¹), tout au long d'une année. JFM, janvier, février et mars; AMJ, avril, mai et juin, JAS, juillet, août et septembre; OND, octobre, novembre et décembre (issue de Redeker *et al.,* 2012). Les barres d'erreur représentent les erreurs standards.

Ainsi les estimations de la production de chlorométhane peuvent varier de 30 % à 60 % (Saito et Yokouchi, 2008). De plus, les flux de chlorométhane sont bidirectionnels avec une émission depuis le sol vers l'atmosphère et inversement de l'atmosphère vers le sol, avec une importance des microorganismes du sol forestier dans la modulation de ces échanges rendant difficile cette estimation (Schäfer *et al.*, 2005). En effet, les estimations du puits du chlorométhane dans le sol varient de 256 Gg.an⁻¹ (Clerbaux *et al.*, 2007) à plus de 1000 Gg.an⁻¹ (Keppler *et al.*, 2005) soit 6 % à plus de 20 % du puits troposphérique total du chlorométhane, d'où l'importance de mieux étudier et comprendre le cycle global du chlorométhane dans les sols forestiers à la fois source et puits de chlorométhane.

Les incertitudes sur les quantifications des différents puits et sources de chlorométhane ne permettent pas de fournir un budget global sûr. De plus, la double fonction de certains environnements, à la fois puits et sources, renforce ces incertitudes. Une amélioration des estimations de production et de consommation du chlorométhane passe par le développement de méthodes d'analyse performantes telle que l'isotopie du carbone et de l'hydrogène (Keppler *et al.*, 2005; Greule *et al.*, 2012; Nadalig *et al.*, 2013) et la proposition de

modèle tridimensionnel des flux terrestres de chlorométhane (Xiao *et al.,* 2010). Des études de la dégradation du chlorométhane dans des sols forestiers ont mis en évidence la présence de bactéries actives dans la dégradation du chlorométhane (Borodina *et al.,* 2005; Miller *et al.,* 2004). Une étude non publiée, réalisée au sein du sol forestier de Steigerwalt en Allemagne (site d'étude pour l'expérience de *Stable Isotope Probing* décrite au chapitre 4) démontre également une dégradation du chlorométhane par des bactéries en condition oxique (Rüffer, 2013).

3. La méthylotrophie

Les microorganismes capables de croître sur le chlorométhane comme source de carbone et d'énergie sont des microorganismes méthylotrophes (Tableau 1.5). La méthylotrophie est définie comme la capacité d'un microorganisme à utiliser un composé sans liaison carbonecarbone comme unique source de carbone et d'énergie. Les substrats de la méthylotrophie regroupent des composés à un seul atome de carbone comme le méthane (CH₄), le méthanol (CH₃OH), le formaldéhyde (HCHO), la méthylamine (CH₃NH₂), ainsi que des composés halogénés comme le chlorométhane et le dichlorométhane, et également des composés à plusieurs atomes de carbone comme la diméthylamine (C₂H₇N) (Chistoserdova, 2011; Muller et al., 2011a). Les organismes méthylotrophes sont retrouvés de façon ubiquitaire dans l'air (DeLeon-Rodriguez et al., 2013), dans les eaux douces et marines (Neufeld et al., 2008), dans les racines et les feuilles des plantes (Nadalig et al., 2011; Knief et al., 2012) ou encore dans le sol et les sédiments marins (Kolb, 2009a). Ces organismes sont retrouvés dans les trois domaines du vivant; les bactéries, les archées (dont des archées méthanotrophes, (Antony et al., 2012)) et les eucaryotes (dont des levures (Yurimoto, 2009)). Chez les eucaryotes, des levures du genre Pichia, Candida et Hansenula sont capables de métaboliser le méthanol via une alcool oxydase dans un compartiment cellulaire nommé peroxysome (van der Klei et al., 2006). Le métabolisme méthylotrophe est toutefois mieux étudié chez les bactéries (Kolb, 2009; Chistoserdova, 2011; Peyraud et al., 2011; Peyraud et al., 2012). Depuis la découverte des bactéries capables d'utiliser le méthane ou le méthanol comme unique source de carbone et d'énergie (Loew, 1892; käserer, 1906; Söhngen, 1906), de nombreux genres bactériens méthylotrophes ont été identifiés, dont certains sont méthylotrophes facultatifs, capables aussi de croître sur des substrats multicarbonés comme le succinate (Whittenbury et al., 1970; Lidstrom, 2006; Boden et al., 2008; Hung et al., 2011). Les bactéries méthylotrophes

appartiennent aux classes des alpha, béta et gamma protéobactéries, sont positifs ou négatifs à la coloration de Gram, avec des métabolismes aérobies ou anaérobies facultatifs.

Les bactéries méthylotrophes sont classées en 2 groupes; le premier regroupant les bactéries méthanotrophes, capables d'utiliser le méthane comme seule source de carbone et d'énergie. Elles sont souvent méthylotrophes strictes, comme *Methylomonas, Methylobacter, Methylosinus* et *Methylocystis*. Le 2^e groupe comprend celles qui sont incapables d'oxyder le méthane, appelées bactéries méthylotrophes non-méthanotrophes. Ces dernières sont souvent méthylotrophes facultatives, comme les bactéries du genre *Hyphomicrobium* ou *Methylobacterium* (Chistoserdova et Lidstrom, 2013).

3.1. Les voies de la méthylotrophie

La méthylotrophie comprend 3 étapes :

- la transformation initiale des composés en formaldéhyde (HCHO) (pas pour l'utilisation du chlorométhane) ;

- l'oxydation complète de la source de carbone en CO₂ pour la génération d'énergie;

- l'assimilation de carbone pour la production de biomasse qui peut se faire par 3 voies distinctes, la voie du ribulose biphosphate (RuBP) utilisant du CO_2 , la voie du ribulose monophosphate (RuMP) à partir de formaldéhyde ou par le cycle de la sérine à partir du méthylène-tetrahydrofolate (CH₂=H₄F) et de CO₂ (Chistoserdova, 2011).

Pour la majorité des souches méthylotrophes, la première étape du métabolisme est l'oxydation ou l'hydrolyse de la source de carbone et d'énergie en formaldéhyde. Les enzymes catalysant cette étape sont spécifiques au substrat : la méthane monooxygénase MMO soluble (sMMO) ou particulaire (pMMO) pour le méthane, la méthylamine déshydrogénase (MADH, périplasmique) pour la méthylamine, la méthanol déshydrogénase (Mxa, périplasmique) pour le méthanol, ou encore une dichlorométhane déshydrogénase pour le dichlorométhane (DcmA, cytoplasmique). Le gène *mxaF* est utilisé comme marqueur moléculaire de l'oxydation du méthanol (Neufeld *et al.*, 2008). Cependant, il existe des systèmes alternatifs pour la dégradation du méthanol, comme les gènes *xox* plus répandus dans l'environnement ou encore *mdh* (Kalyuzhnaya *et al.*, 2008). Parmi ces gènes, *xoxF* code une sous-unité d'une méthanol déshydrogénase qui présente plus de 50 % d'identité au niveau des acides aminés avec MxaF (Schmidt *et al.*, 2010; Taubert *et al.*, 2015). Plusieurs

systèmes d'oxydation du méthanol peuvent être présents simultanément au sein d'un organisme comme par exemple chez *M. extorquens* (Figure 1.10, Skovran *et al.*, 2011). Plusieurs systèmes d'oxydation pour l'utilisation de la méthylamine ont également été mis en évidence chez certaines souches de *M. extorquens* (Vuilleumier *et al.*, 2009; Gruffaz *et al.*, 2014; Nayak et Marx, 2014). Des composés halogénés sont également utilisés par les bactéries méthylotrophes comme chez *M. extorquens* et *Hyphomicrobium* où deux souches différentes peuvent utiliser soit le dichlorométhane soit le chlorométhane (Gälli et Leisinger, 1985; La Roche et Leisinger, 1990; Vannelli *et al.*, 1998; McAnulla *et al.*, 2001). La déshalogénation du dichlorométhane par la dichlorométhane déshalogenase codée par le gène *dcmA* conduit à la formation de formaldéhyde.

Chez Methylobacterium, deux voies sont actuellement caractérisées pour transformer le formaldéhyde en formiate : l'une est dépendante du tétrahydrofolate (H₄F) et l'autre utilise la tétrahydrométhanoptérine (H₄MPT). La première de ces deux voies n'est pas la voie majeure pour la génération d'énergie, et met en jeu des inter-conversions entre le méthylène-H₄F, le méthényl-H₄F, le formyl-H₄F et enfin le formiate (Marison et Attwood, 1982). Cette voie joue probablement un rôle important dans le maintien de concentrations suffisantes des différents intermédiaires métaboliques contenant du H₄F, nécessaires pour l'assimilation du carbone via le cycle de la sérine (Marx et al., 2005; Crowther et al., 2008). La seconde voie, découverte plus tardivement en 1998, utilise la H₄MPT qui, condensée au formaldéhyde, est successivement transformée en méthyléne-H₄MPT, méthényl-H₄MPT, formyl-H₄MPT, puis formiate (Chistoserdova et al., 1998). Cette voie, initialement décrite chez les Archaea, est plus efficace pour l'oxydation du formaldéhyde en CO₂ que la voie dépendante du H₄F grâce aux fortes activités de ses enzymes (Vorholt, 2002). Cependant, tous les substrats de la méthylotrophie ne sont pas oxydés en formaldéhyde; c'est le cas du chlorométhane. La déshalogénation du chlorométhane va mener à la formation de méthylène-tétrahydrofolate (CH₂=H₄F), qui va directement être assimilé dans la biomasse via le cycle de la sérine mais également oxydé en formiate pour fournir de l'énergie (Figure 1.10). Ainsi, M. extorquens CM4 convertit directement le chlorométhane en méthylène-tétrahydrofolate (CH₂=H₄F) sans passer par le formaldéhyde grâce aux gènes cmuA, cmuB, metF2 codant respectivement une protéine avec un domaine méthyltransférase et un de liaison corrinoïde, une méthyltransférase méthylcobalamine : H₄F et une méthylène-H₄F réductase (Studer et al., 1999 ; Studer et al., 2001; Studer et al., 2002) (voir section 1.3.4.2.).

74



Figure 1.10. Schéma synthétique des différentes voies de la méthylotrophie chez *M. extorquens*

Le degré d'oxydation des molécules est représenté par l'échelle de gauche. Les réactions spécifiques liées à l'utilisation du chlorométhane (CH₃Cl) et du dichlorométhane (CH₂Cl₂) comme unique source de carbone et d'énergie sont représentées respectivement en vert et en violet : CmuABC, « chloromethane utilizing genes », MetF2, méthylène- H4F réductase, DcmA, dichlorométhane déshalogénase. Les autres réactions spécifiques aux substrats utilisés sont en noir : MDH, méthanol déshydrogénase; MADH, méthylamine déshydrogénase. Les réactions conduisant à l'oxydation complète du carbone pour la production d'énergie sont en orange : Fae, enzyme d'activation du formaldéhyde; MtdAB, méthylènetétrahydrométhanoptérine déhydrogénase; Mch, méthenyl-tétrahydrométhanoptérine méthylène-tétrahydrofolate cyclohydrolase; Fch, cyclohydrolase; FtfL, formyltétrahydrofolate ligase; FhcABCD, complexe formyltransférase-hydrolase; Fdh, formiate déshydrogénase. Les réactions de l'assimilation du carbone pour produire de la biomasse sont en gris : FtfL, formyl-tétrahydrofolate ligase; Fch, méthylène-tétrahydrofolate cyclohydrolase; MtdA, méthylène-tétrahydrométhanoptérine déhydrogénase; RuMP, cycle du ribulose monophosphate; RuBP, cycle du ribulose biphosphate.

Plus de 86 gènes impliqués dans le métabolisme des composés en C₁ chez *M. extorquens* ont été identifiés (Marx *et al.*, 2003). Ces gènes sont impliqués dans l'oxydation du méthanol, de





Figure 1.11. Assimilation du carbone chez M. extorquens

Les molécules d'entrée sont encadrées en bleu. Les principales connexions vers d'autres voies métaboliques sont indiquées par des rectangles verts. Les bactéries transforment le méthanol (CH₃OH) ou le dichlorométhane (CH₂Cl₂) en formaldéhyde (HCHO) qui est condensé avec une molécule de tétrahydrométhanoptérine (H₄MPT), puis oxydé en formiate (HCOOH). Le formiate peut être transformé en CO₂, ou assimilé via le méthylène-tétrahydrofolate (H₄F) par le cycle de la sérine. L'assimilation du chlorométhane (CH₃Cl), se fait directement par transfert

du groupement méthyl sur le H₄F, qui est alors assimilé via le cycle de la sérine pour fournir de la biomasse, ou oxydé en formiate (HCOOH). L'assimilation de l'acétyl-CoA et la régénération du glyoxylate se font au travers de la voie de l'éthylmalonyl CoA. Le cycle TCA (cycle des acides tricarboxyliques) est quant à lui principalement utilisé pour l'assimilation des composés multicarbonés (Šmejkalová et al., 2010). Le nombre de flèches correspond au nombre de molécules impliquées dans la réaction. 1, sérine hydroxyméthyl transférase; 2, sérine-glyoxylate aminotransférase; 3, hydroxypyruvate réductase; 4, glycérate kinase; 5, énolase; 6, phosphoénolpyruvate carboxylase; 7, malate déshydrogénase; 8, malate-CoA ligase (malate thiokinase); 9, L-malyl-CoA/β-méthylmalyl-CoA lyase; 10, β-cétothiolase; 11, acétoacétyl-CoA réductase; 12, crotonase (R-spécifique); 13, crotonyl-CoA carboxylase réductase; 14, éthylmalonyl-CoA/méthylmalonyl-CoA épimerase; 15, éthylmalonyl-CoA mutase; 16, méthylsuccinyl-CoA déshydrogénase; 17, mésaconyl-CoA hydratase; 18, propionyl-CoA carboxylase; 19, méthylmalonyl-CoA mutase; 20, succinyl-CoA synthase; 21, succinate déshydrogénase; 22, fumarase; 23, citryl-CoA lyase; 24, aconitate hydratase; 25, NADP-dépendent isocitrate déshydrogenase; 26, complexe 2-oxoglutarate déshydrogénase; 27, méthanol déshydrogénase; 28, dichlorométhane déshalogénase; 29, formiate déshydrogenase. Pour une même réaction le nombre de flèche correspond au nombre de molécules impliquées dans cette réaction. PHB, polyhydroxybutyrate; Q, quinone (figure modifiée d'après Šmejkalová et al., 2010).

Chez Methylobacterium, une partie du formaldéhyde produit intègre le cycle de la sérine via le formiate et le méthylène-H₄F pour la production de biomasse. L'enzyme GlyA (sérine hydroxyméthyltransférase) permet la condensation du méthylène-H₄F avec la glycine pour produire la sérine (Figure 1.11). La sérine (composé à trois atomes de carbone) est ensuite transformée en hydroxypyruvate, glycérate, glycérate-2-phosphate et en phosphoénolpyruvate. Le phosphoénolpyruvate est converti en oxaloacétate puis en malate, composés à quatre atomes de carbone par incorporation de CO₂. Le malate, couplé au coenzymeA (CoA), est scindé en deux composés, le glyoxylate permettant de refermer le cycle de la sérine et l'acétyl-CoA qui est intégré dans la voie de l'éthylmalonyl-CoA où une molécule de CO₂ est incorporée (Erb et al., 2008; Peyraud et al., 2009; Šmejkalová et al., 2010). Ce cycle peut également produire de l'hydroxybutyryl-CoA, matière première de la synthèse de polyhydroxybutyrate (PHB), formant des granules de réserve en carbone et phosphate pour la cellule.

3.2. La dégradation bactérienne du dichlorométhane (CH₂Cl₂)

L'utilisation bactérienne du dichlorométhane est mieux connue en condition oxique qu'anoxique.

3.2.1. Utilisation du dichlorométhane en condition anoxique

Un processus de fermentation du dichlorométhane a été décrit chez la bactérie *Dehalobacterium formicoaceticum* (Mägli *et al.,* 1998). Le métabolisme anaérobie de cette bactérie fait intervenir une combinaison de méthyltransférases dépendantes de la cobalamine et du H₄F. Le dichlorométhane et le H₄F sont convertis en méthylène-tétrahydrofolate (CH₂=H₄F), et en chlorure inorganique. Une partie du CH₂=H₄F est transformée en formiate, tandis qu'une autre partie du CH₂=H₄F forme de l'acétate par incorporation de CO₂ présent dans le milieu (Mägli *et al.,* 1998).

3.2.2. Utilisation du dichlorométhane en condition oxique

La première souche capable d'utiliser le dichlorométhane comme substrat de croissance à avoir été isolée et caractérisé est une bactérie du genre *Methylopila* (Brunner *et al.*, 1980). Trois voies de dégradation du dichlorométhane en aérobie sont connues. Cependant, une seule de ces voies permet l'utilisation du dichlorométhane comme source de carbone et d'énergie. L'utilisation du dichlorométhane comme source de carbone et d'énergie est possible par l'expression d'une dichlorométhane déshalogénase appartenant à la famille des glutathion-S-transférase (GST), codée par le gène *dcmA* (Figure 1.12). Dans la majorité des cas, les souches capables d'utiliser le dichlorométhane comme substrat de croissance possèdent le gène *dcmA* (Vuilleumier *et al.*, 2001; Muller *et al.*, 2011b). De plus, ce gène est fortement conservé avec des identités au niveau nucléotidique comprises entre 98 % et 100 %. La protéine DcmA catalyse la réaction de déchloration d'une mole de dichlorométhane en deux moles de protons (H⁺), deux moles de chlorure (Cl⁻), et une mole de formiate, via la formation consécutive de S-chlorométhylglutathion et de formaldéhyde, deux composés génotoxiques (Figure 1.12).



Figure 1.12. Dégradation du dichlorométhane par la voie hydrolytique de la glutathion S-transférase

(Figure adaptée de Muller *et al.*, 2011). spont.: spontanée, réaction abiotique; GSH, glutathion; SG, glutathionyl.

La dégradation bactérienne aérobie du dichlorométhane a été largement décrite chez *M. extorquens* DM4 qui utilise le dichlorométhane comme unique source de carbone et d'énergie (Roche et Leisinger, 1991; Kayser *et al.*, 2002; Muller *et al.*, 2011a). Les gènes d'utilisation du dichlorométhane sont portés par un transposon catabolique possédant quatre gènes *dcm* encadrés par des séquences d'insertion. Ces gènes sont la dichlorométhane déshalogénase (*dcmA*), *dcmR* qui code un facteur de transcription, ainsi que deux gènes de fonction encore inconnue (*dcmB* et *dcmC*). Ce transposon est lui-même localisé sur un îlot génomique d'une taille de 126 kb (Vuilleumier *et al.*, 2009). L'utilisation du dichlorométhane par *M. extorquens* DM4 aboutit à la formation de formaldéhyde, qui est ensuite oxydé en formiate via la voie de la tétrahydrométhanoptérine (H₄MPT). Le formiate est soit réduit en CH₂=H₄F pour fournir de la biomasse via le cycle de la sérine, soit oxydé en CO₂ par une formiate déshydrogénase (FDH) pour produire de l'énergie (Figure 1.10).

3.3. La dégradation bactérienne du chlorométhane

Plusieurs genres bactériens, isolés de différents environnements, sont capables de dégrader le chlorométhane (Tableau 1.5). Le chlorométhane peut être utilisé comme source de carbone et d'énergie ou dégradé sans servir de source de carbone en conditions oxique ou anoxique (Traunecker *et al.*, 1991; Han et Semrau, 2000; Freedman *et al.*, 2004).

3.3.1. Dégradation du chlorométhane en condition anoxique

La déshalogénation du chlorométhane en anaérobie est catalysée par une enzyme qui transfère le groupement méthyl (CH₃) au H₄F via une protéine corrinoïde, produisant des ions chlorure et du méthyl-H₄F, qui est ensuite oxydé en formiate puis en CO₂ ou assimilé par la voie de l'acétyl CoA (Harper, 2000). La bactérie *Acetobacterium dehalogenans* MC utilise le chlorométhane en anaérobie (Tableau 1.5) (Traunecker *et al.*, 1991; Harper, 2000). Cette

déshalogénation est basée sur une méthyl transférase corrinoïde-dépendante qui contrairement à d'autres méthyl transférase corrinoïde-dépendantes, nécessite le maintien du cobalt dans la protéine corrinoïde par un système ATP-dépendant (Studer *et al.*, 1999). La souche *Pseudomonas aeruginosa* NB1, quant à elle, est capable d'utiliser le chlorométhane comme unique source de carbone et d'énergie en condition de réduction des nitrates (Freedman *et al.*, 2004). Cependant, le mécanisme de déchloration reste inconnu.

3.3.2. Dégradation du chlorométhane en condition oxique

La première souche décrite comme utilisant le chlorométhane comme unique source de carbone et d'énergie est *Hyphomicrobium* sp. MC1 (Hartmans *et al.*, 1986). Comme cette souche, la majorité des souches chlorométhanes-dégradantes isolées à ce jour sont des microorganismes se développant en condition oxique. Les souches ont été isolées d'environnements variés comme des sols d'usines pétrochimiques, des stations d'épuration, des sols forestiers ou des milieux aquatiques marins et dulçaquicoles (Tableau 1.5). Elles appartiennent à la classe des *Alphaproteobacteriacea* et aux genres *Methylobacterium*, *Hyphomicrobium* et *Aminobacter*. A ce jour, la dégradation du chlorométhane la mieux caractérisée est celle de *M. extorquens* CM4 selon la voie *cmu*, mais elle n'est pas la seule voie de dégradation (voir paragraphes ci-dessous).

Voie de dégradation du chlorométhane chez *Methylomicrobium album* BG8 A part la voie *cmu*, le chlorométhane peut aussi être dégradé en condition oxique par une bactérie méthanotrophe, *Methylomicrobium album* BG8, mais seulement en présence d'une autre source carbonée telle que le méthanol (Figure 1.13) (Han et Semrau, 2000). Medium



Figure 1.13. Mecanisme proposé d'oxydation du chlorométhane par *Methylomicrobium album* BG8

En présence d'acéthylène, inhibiteur de l'activité de la pMMO, la souche ne dégrade plus le chlorométhane, démontrant que le chlorométhane est oxydé par la méthane monooxygénase particulaire (pMMO). Le méthanol est oxydé en formadéhyde (HCHO) par la méthanol deshydrogénase (MDH). Le formaldéhyde rejoint le métabolisme en C₁ pour la synthèse de biomasse et pour la production d'énergie sous forme d'équivalents réducteurs (2[H]) en impliquant la formaldéhyde déshydrogénase (FADH) et la formiate déshydrogénase (FDH). On notera que par rapport au méthanol, le chlorométhane génère deux équivalents réducteurs en moins. (D'après Han et Semrau, 2000).

Dans une culture de *M. album* BG8 en présence de méthanol, le marquage au C¹⁴ du CH₃Cl a permis de démontrer que le chlorométhane stimule la croissance et l'assimilation du carbone dans la biomasse jusqu'à 38 % avec 50 % du chlorométhane transformé en CO₂ suggérant que le chlorométhane peut servir comme source de carbone et d'énergie (Han and Semrau, 2000). Cependant, comme le chlorométhane ne peut pas servir comme unique source de carbone, les auteurs de ce travail proposent que *M. album* BG8 ne puisse pas tirer assez d'équivalents réduits de la dégradation du chlorométhane pour permettre sa croissance sur le chlorométhane. En présence d'une autre source carbonée en quantité suffisante par rapport au chlorométhane, l'utilisation du chlorométhane est possible (2,6 mM de chlorométhane et 5 mM de méthanol) (Han et Semrau, 2000).

– La voie *cmu*

Des études biochimiques et génétiques chez la souche *M extorquens* CM4 ont permis d'identifier une voie d'utilisation du chlorométhane en condition oxique, appelée voie *cmu*

(chlorométhane utilisation) (Vannelli *et al.*, 1998, 1999) (Figure 1.14). *M extorquens* CM4 métabolise 1 mole de chlorométhane et 1,5 moles d'O₂ pour produire 1 mole de CO₂ et 1 mole de chlorure (Vannelli *et al.*, 1998). L'inactivation de gènes par mutagénèse aléatoire avec le minitransposon *Tn*5 a permis d'identifier 9 mutants dont la croissance était maintenue en méthanol ou en méthylamine, mais impossible sur chlorométhane (Vannelli *et al.*, 1998). L'étude de ces mutants a mis en évidence le caractère essentiel des gènes *cmuABC*, des méthyltransférases et *purU* et *metF2*, codant respectivement une méthylène-H₄F réductase et une formyl- H₄F, pour la croissance en présence de chlorométhane (Studer *et al.*, 2001, 2002). De plus, les gènes *cmuAB* sont essentiels pour la déchloration du chlorométhane (Studer *et al.*, 2002).



Figure 1.14 Voie cmu chez Methylobacterium extorquens CM4

Le groupement méthyl du chlorométhane est transféré au tétrahydrofolate (H₄F) par *cmuA* et *cmuB*. Le méthényl-tétrahydrofolate produit par oxydation du méthyle-tétrahydrofolate peut soit incorporer le cycle de la sérine pour produire de la biomasse, ou être finalement oxydé en CO₂ pour produire de l'énergie. CmuA, protéine avec un domaine méthyltransférase et un de liaison de groupe corrinoïde (Studer *et al.*, 2001), CmuB, méthyltransférase méthylcobalamine:H₄F (Studer *et al.*, 1999); MetF2, méthylène réductase; FoID, enzyme bifonctionnelle méthyléne-H₄F déshydrogénase/ méthényl-H₄F cyclohydrolase; PurU, 10-formyl-H₄F hydrolase.

Les protéines CmuA et CmuB vont catalyser le transfert du groupement méthyl du chlorométhane au H₄F pour former le méthyl-H₄F (CH₃-H₄F) (Studer *et al.*, 2001). Le CH₃-H₄F est ensuite oxydé en formiate et CO₂, avec une assimilation du carbone au niveau du méthylène-H₄F (Vannelli *et al.*, 1999). Chez *M. extorquens* CM4, l'ensemble des gènes de la voie *cmu* sont localisés sur le plasmide pCMU01 dans une région de 180 kb (Figure 1.15). Les gènes sont regroupés en 2 clusters distants de 30 kb, contenant un taux de GC atypique et dépourvus d'éléments mobiles. Directement en aval de *purU* et *folD* se trouve le gène *folC2*, codant une enzyme bifonctionnelle folylpolyglutamate synthase et dihydrofolate synthase. Cette région comporte aussi des gènes non caractérisés fonctionnellement : *paaE-like*, *fmdB* et *hutl*, codant respectivement une putative oxydoréductase, un régulateur transcriptionnel putatif, et une imidazole hydrolase putative.



Figure 1.15. Plasmide pCMU01 d'utilisation du chlorométhane chez *M. extorquens* **CM4** Les clusters 1 et 2 distants de 21 kb contiennent tous les gènes essentiels d'utilisation du chlorométhane (*cmu*) connues avant le séquençage du génome (Vannelli *et al.*, 1999; Studer *et al.*, 2002). L'ensemble de cette région couvre 40 kb de la région *cmu* du plasmide de 380 kb. Les zones S1, S2, S3 correspondent à une région de 69 kb conservée entre le plasmide pCMU01

et le plasmide p1METDI de *M. extorquens* DM4. La région S4 présente une synténie avec une région dupliquée du chromosome de la souche dichlorométhane-dégradante DM4. Les cercles représentent depuis l'extérieur : 1, pourcentage de déviation GC (GC fenêtre – GC moyen) dans une fenêtre de 1000 pb; 2, CDS prédits transcrits dans le sens des aiguilles d'une montre; 3, CDS prédits transcrits dans le sens contraire des aiguilles d'une montre; 4, GC skew (G+C/G-C) dans une fenêtre de 1000 pb; 5, éléments transposables (rose) et pseudogènes (gris) (Figure issue du manuscrit de thèse de Sandro Roselli, 2009)

Cette région contient également 23 gènes impliqués dans la biosynthèse et le transport de la cobalamine, ainsi que ceux nécessaires à la biosynthèse du tétrahydrofolate, deux cofacteurs essentiels à la voie *cmu* d'utilisation du chlorométhane (Figure 1.15). Une partie d'entre eux présente une copie chromosomique (Roselli *et al.*, 2013). Toutefois, l'analyse des génomes de bactéries possédant les gènes *cmu* a fait apparaître deux organisations génétiques : une retrouvée chez *M. extorquens* CM4, où les gènes sont distribués en deux clusters éloignés de 30 kb, et une deuxième organisation, présente chez les autres bactéries, où les gènes *cmu* sont situés sur le même cluster (Figure 1.15). Un degré important de conservation dans l'organisation des gènes *cmu* chez les bactéries appartiennent toute à la classe des *Alphaproteobacteria*. L'ordre des gènes dans le cluster *cmu* est fortement conservé (Nadalig *et al.*, 2011). Cependant chez *M. extorquens* CM4, les gènes *cmu* as fortement conservé (Nadalig *et al.*, 2011). Cependant chez *M. extorquens* CM4, les gènes *cmu* est fortement conservé (Studer *et al.*, 2002).

Des souches chlorométhane-dégradantes ne possédant pas les gènes *cmu* ont été isolées comme *Roseovarius* sp. 217 et *Leisingera methylohalidovorans* MB2 (Schäfer *et al.,* 2007). Des analyses en isotopes stables du chlorométhane ont montré que les fractionnements isotopiques du carbone et de l'hydrogène étaient différents pour les bactéries chlorométhane-dégradantes possédant les gènes *cmu* et *L. methylohalidovorans* MB2 ne les possédant pas (Nadalig *et al.,* 2014). La voie de dégradation du chlorométhane par *L. methylohalidovorans* MB2, indépendante des gènes *cmu*, reste à élucider.



Figure 1.16. Comparaison de l'organisation des gènes *cmu* chez des bactéries contenant le gène *cmuA*

Les flèches représentent les gènes codant des protéines. Les flèches incomplètes représentent la présence de gène dont la séquence n'est pas entièrement déterminée. La couleur de référence des gènes est celle indiquée chez *M. extorquens* CM4, où ces gènes sont retrouvés sur deux clusters séparés de 30 kb. Les gènes homologues sont représentés de la même couleur; Le pourcentage d'homologie avec les protéines codées par les gènes de *M. extorquens* CM4 est indiqué à l'intérieur des flèches. L'organisation génétique est représentée à l'échelle. Chez la souche MB2, on notera que seulement une partie du gène *cmuA* correspondant au domaine de liaison du co-facteur corrinoide est présente.

4. Le modèle Methylobacterium

Le genre *Methylobacterium* a été décrit pour la première fois par Patt, Cole et Hanson en 1976 (Patt *et al.*, 1976). Ce genre compte une vingtaine d'espèces appartenant, selon la séquence nucléotidique du gène codant l'ARNr 16S, aux Alpha-protéobactéries, à l'ordre des *Rhizobiales* et à la famille des *Methylobacteriacaea* (Lidstrom, 2006). Elles sont méthylotrophes

facultatives, aérobies strictes, mésophiles avec un optimal de température compris entre 25 et 30°C, de coloration Gram négative et elles montrent une réaction positive aux tests de catalase et d'oxydase. Les cellules possèdent souvent des inclusions de poly-β-hydroxybutyrate, réserves d'énergie et de carbone, et ont en général une coloration rose en raison de la présence de caroténoïdes (Korotkova et Lidstrom, 2001; Van Dien *et al.*, 2003). Les souches de *Methylobacterium* ont été isolées de lieux très divers, anthropisés (bords de route, réservoirs d'eau) mais aussi naturels (sol, surface de feuilles) (Tableau 1.6) (Radajewski *et al.*, 2002; Nadalig *et al.*, 2011; Bai *et al.*, 2015). Ces bactéries ont un rôle

environnemental important de par leur capacité à métaboliser des composés toxiques comme le méthanol, le dichlorométhane ou encore le chlorométhane (Dourado *et al.*, 2015). Les bactéries méthylotrophes ont été parmi les premières étudiées dans l'environnement à l'aide d'outils moléculaires (Holmes *et al.*, 1995; Kolb et Stacheter, 2013).

4.1. *Methylobacterium,* bactérie modèle pour l'étude de la méthylotrophie

Une des espèces les plus étudiées du genre Methylobacterium est M. extorquens retrouvée dans de nombreux environnements comme les plantes, le sol, les eaux usées ou les nuages (Bai et al., 2015; Bringel et Couée, 2015; Nadalig et al., 2011; Ochsner et al., 2015; Temkiv et al., 2012). M. extorquens AM1 est la souche la plus étudiée et a été considérée, avec Methylobacterium organophilum, comme le modèle d'étude du genre Methylobacterium, pour la méthylotrophie ou les intéractions plantes-microorganismes (Dourado et al., 2015). La présence de cinq réplicons et de 174 séquences d'insertion chez M. extorquens AM1 complique son utilisation en biologie moléculaire et elle est en voie d'être remplacée par M. extorquens PA1 qui ne comporte qu'un chromosome et seulement 20 séquences d'insertion (Nayak et Marx, 2014). L'ensemble des souches de M. extorquens (AM1, BJ001, CM4, DM4, DSM 13060 et PA1) sont capables d'utiliser le méthanol, mais seules M. extorquens CM4 et DM4 sont capables d'utiliser des substrats chlorés comme unique source de carbone et d'énergie (Vuilleumier et al., 2009; Marx et al., 2012). M. extorquens CM4, isolée d'un sol d'usine pétrochimique en Russie, est capable de croître avec du chlorométhane (Doronina et al., 1996), tandis que M. extorquens DM4, capable de croître sur dichlorométhane, a été isolée d'un sol pollué par des alcanes chlorés (Gälli et Leisinger, 1985). Les génomes des souches de *M. extorquens* sont caractérisés par un taux de GC supérieur à 68 %, avec un chromosome de taille importante (entre 5,5 et 5,9 Mb) (Tableau 1.6).

A l'exception de la souche PA1 et DSM 13060, les souches de *M. extorquens* possèdent un ou plusieurs plasmides, avec pour la souche *M. extorquens* AM1 la présence d'un mégaplasmide (Ochsner *et al.*, 2015). Fin décembre 2015, 24 génomes de *Methylobacterium* isolés de la phyllosphère et de racines de plantes ont été séquencées (Bai *et al.*, 2015), mais aucun gène similaire à *cmuA* n'a été détecté.

Chapitre 1 (en français)

Tableau 1.6. Caractéristiques des premières souches d	le Methylobacterium	dont le génome	a été séquencé
---	---------------------	----------------	----------------

Paramètres ^a	M. extorquens AM1	<i>M. extorquens</i> BJ001 ^b	<i>M. extorquens</i> CM4	M. extorquens DM4	<i>M. extorquens</i> DSM13060	<i>M.</i> extorquens PA1	<i>M. nodulans</i> ORS 2060	<i>M. radiotolerans</i> JCM 2831	<i>M.</i> sp. 4-46
Génome									
Taille (Mb)	6,88	5,85	6,18	6,12	6,73	5,47	8,84	6,90	7,73
Contenu en G+C (%)	68,5	69,4	68,0	68,0	68,23	68,2	68,4	71,0	71,5
Nombre de réplicons	5	3	3	3	1	1	8	9	3
Nombre de CDS prédites	6335	5945	6276	5706	7132	5153	10333	7356	8479
N° d'accession Genbank	NC_012808	NC_010725	NC_011757	NC_012988.1	NZ_AGJK00000000.1	NC_010172	NC_011894	NC_010505	NC_010511
Chromosome									
Taille (Mb)	5,51	5,8	5,77	5,94	6,73	5,47	7,78	6,1	7,70
Contenu en G+C (%)	68,7	69,4	68,2	68,1	68,23	68,2	68,9	71,5	71,6
Nombre de plasmides	4	2	2	2	0	0	7	8	2
Séquençage (date)	UW Genome Sequencing Center, (2003/2006)	JGI (2008)	JGI (2008)	Génoscope (2006)	JGI (2010)	JGI (2007)	JGI (2009)	JGI (2008)	JGI (2008)
Publication du génome	Vuilleumier <i>et al.,</i> 2009	Marx <i>et al.,</i> 2012	Marx <i>et al.,</i> 2012	Vuilleumier <i>et al.,</i> 2009	Koskimäki et al., 2015	Marx <i>et al.,</i> 2012	Marx <i>et al.,</i> 2012	Marx <i>et al.,</i> 2012	Marx <i>et al.,</i> 2012

^a Génoscope (http://www.cns.fr/agc/microscope/home/index.php) ou Craig Venter institute (http://www.jcvi.org/)

^b Originellement nommée *Methylobacterium populi* (Van Aken *et al.,* 2004), mais sur la base de l'identité de l'ARNr 16S a été affiliée à *M extorquens* (Marx *et al.,* 2012)

4.2. Outils génétiques disponible chez *M. extorquens*

De nombreux outils génétiques sont disponibles chez *M. extorquens* AM1. Ces outils génétiques sont utilisés dans des approches de mutagénèse aléatoire, de délétion ou d'insertion ciblée de gènes, ainsi que d'expression génique constitutive ou inductible. Une liste des outils génétiques disponibles chez *M. extorquens* AM1 a été mise à jour récemment dans une revue (Ochsner *et al.*, 2015).

4.3. Approches globales chez *M. extorquens*

Les connaissances portant sur le métabolisme méthylotrophe chez les bactéries du genre *Methylobacterium* et plus particulièrement de l'espèce *M. extorquens* ont considérablement augmenté ces dernières années grâce aux apports de séquençage nouvelle génération à hautdébit « *Next Generation Sequencing* » ou NGS, permettant l'étude des gènes (génomique), de leurs expressions (transcriptomique), ainsi que l'étude du protéome (protéomique) et des métabolites (métabolomique).

4.3.1. Etudes génomiques chez *M. extorquens*

Un projet de séquençage du génome complet de AM1 a été entrepris en 1998 (Chistoserdova *et al.*, 2003). L'assemblage du génome n'a été complété qu'en 2009 lorsque celui de la souche DM4 a été disponible (Vuilleumier *et al.*, 2009). Le génome de la souche *M. extorquens* CM4 a été disponible en même temps que le génome d'autres souches non déchlorantes de la même espèce (BJ001; PA1) (Marx *et al.*, 2012). Au début de ma thèse, la plateforme MaGe, mise en place par le Génoscope (Evry, France), permettait l'accès à l'annotation et aux analyses comparatives de six génomes de *M. extorquens* (AM1, BJ001, CM4, DM4, DSM 13060 et PA1) dont on connaissait bien le phénotype de croissance sur le méthanol et les méthanes chlorés. La première analyse génomique comparative de souches de *M. extorquens* a démontré la présence de nombreux éléments d'insertion (IS) ainsi qu'une organisation groupée de ces gènes, notamment ceux du métabolisme du méthanol, suggérant ainsi le rôle prédominant des IS dans l'évolution des génomes de souches phylogénétiquement proches comme AM1 et DM4 (Vuilleumier *et al.*, 2009a). Cependant, l'histoire évolutive des génomes de *M extorquens* AM1 et PA1 aient un taux de GC proche (respectivement 68,2 % et 68,5 %), et partagent plus de 90 gènes

impliqués dans la méthylotrophie avec au moins 95 % d'identité, une étude récente montre que ces souches présentent des taux de croissance différents avec des substrats en C₁, comme le méthanol, la méthylamine ou encore le formaldéhyde suggérant l'implication d'autres mécanismes (Nayak et Marx, 2014).

Des études de mutagénèse aléatoire et de protéomique comparative ont mis en évidence qu'en croissance sur un méthane chloré, M. extorquens requière la présence de gènes localisés sur des îlots génomiques, mais également des gènes de ménage impliqués dans la réponse au stress, ainsi que dans le métabolisme central (Michener et al., 2014a; Muller et al., 2011a; Roselli et al., 2013). L'utilisation de chlorométhane ou de dichlorométhane génère un stress cellulaire auquel la cellule doit s'adapter (Kayser et Vuilleumier, 2001; Kayser et al., 2002; Roselli et al., 2013). Pour comprendre ces adaptations, Michener et al., 2014 ont testé la capacité d'utilisation du dichlorométhane chez des souches non dichlorométhanedégradantes. Pour cela, le gène *dcmA*, inséré sur un plasmide, a été transféré dans la souche AM1 (incapable d'utiliser le dichlorométhane pour sa croissance) (Michener et al., 2014b). Les transconjugants obtenus ont une croissance difficile en présence de dichlorométhane, mais montrent une activité de déshalogénation. Le séquençage du génome de ces souches a permis l'identification de mutation dans le gène *clcA* codant un antiporteur Cl⁻/H⁺. L'introduction d'un plasmide contenant les gènes dcmA et clcA mutés dans la souche M. extorquens PA1 (incapable de croître en présence de dichlorométhane) a permis d'établir une croissance efficace en présence de dichlorométhane chez cette souche.

Des travaux complémentaires, portant sur l'adaptation à l'utilisation du chlorométhane par des souches non chlorométhane-dégradantes, ont été réalisés et font l'objet d'un article en cours de soumission (Michener *et al.*, soumis). Un plasmide contenant des gènes de la voie *cmu* (*folD*, *metF2*, *paaE-like*, *purU*), et des gènes associées (*fmdB*, *hutI*) ont été transféré dans des souches non chlorométhane-dégradantes, *M. extorquens* AM1, PA1, BJ001, *M. radiotolerans* et *M. nodulans*. Les transconjugants de chacune de ces souches montrent une croissance sur chlorométhane, bien que la croissance en présence de chlorométhane soit faible. Si le gène *clcA* muté, identifié comme bénéfique à l'adaptation à la croissance sur le dichlorométhane, est inséré dans le plasmide en plus des autres gènes du cluster *cmu*, cela n'améliore pas le fitness des transconjugants sur le chlorométhane. Ceci suggère que l'adaptation à la croissance sur le chlorométhane et le dichlorométhane impliquerait des réponses adaptatives distinctes (Michener *et al.*, submitted article).

Ainsi, malgré de nombreuses études sur la méthylotrophie, de nombreux points ne sont pas élucidés, d'où l'importance de l'utilisation d'approches diverses mais néanmoins complémentaires comme la transcriptomique, la protéomique ou la métabolomique pour comprendre l'ensemble des mécanismes impliqués dans la méthylotrophie chez *M. extorquens* (Tableau 1.7 complété à partir de Ochsner *et al.*, 2015).

Etude « omique »	Réfèrence	Description
Génome	Vuilleumier <i>et al.,</i> 2009	AM1, DM4
	Roselli <i>et al</i> ., 2013	CM4
	Muller <i>et al.,</i> 2011	DM4
	Bai <i>et al.,</i> 2015	Isolats de la phyllosphère et de la rhizosphère (24 génomes)
Modélisation	Peyraud <i>et al.,</i> 2011	AM1
Protéomique	Bosch <i>et al.,</i> 2008; Laukel <i>et al.,</i> 2004	AM1, comparaison méthanol et succinate
	Muller <i>et al.,</i> 2011	DM4, comparaison dichloroméhane et méthanol
	Roselli <i>et al.,</i> 2013	CM4, comparaison chlorométhane et méthanol
	Guo et Lidstrom, 2008	AM1, « Profiling » des métabolites en méthanol et succinate
Transcriptomique	Okubo <i>et al.,</i> 2007	AM1, comparaison méthanol et succinate (microarray)
	Francez-Charlot et al., 2009	AM1, mutant <i>phyR</i> de la régulation du stress général (microarray)
Métabolomique	Kiefer <i>et al.,</i> 2008, 2011	AM1, Concentration des métabolites centraux en méthanol et succinate
	Peyraud et al., 2012	AM1, [¹³ C]-fluxomique, [¹³ C]-CH₃OH en co-utilisation avec le succinate
	Yang et al., 2013	AM1, ¹³ C méthanol, marquage de métabolite au cours la croissance
	Cui <i>et al.,</i> 2016	AM1 avec surexpression d'éthylmalonyl-CoA mutase en méthanol
	Reaser <i>et al.,</i> 2016	AM1, suivi au cours du temps de métabolite après incorporation de [¹³ C]-CH₃OH

Tableau 1.7. Etudes «omiques» réalisées dans des souches de M. extorquens

4.3.2. Approches de transcriptomique chez *M. extorquens*

La transcriptomique est l'étude qualitative et quantitative du transcriptome, c'est-à-dire, l'ensemble des ARNs produits lors de la transcription (Wang *et al.*, 2009). En fonction des conditions de croissance, la cellule modifie le type de transcrits produits ainsi que leurs abondances. Ainsi l'étude de ces transcrits permet de mieux comprendre les adaptations mises en place par la cellule en réponse à diverses conditions de croissance.

Peu d'études de transcriptomique ont été réalisées chez *M. extorquens*. Toutes ont été réalisées chez AM1 et sont basées sur des approches de puces à ADN (microarray). Ces études ont mis en évidence des gènes intervenant dans le métabolisme des C₁, par comparaison des transcriptomes de AM1 en croissance sur du méthanol ou du succinate

(Okubo *et al.*, 2007; Francez-Charlot *et al.*, 2009), ou encore le rôle de *phyR*, un régulateur transcriptionnel de la réponse au stress (Francez-Charlot *et al.*, 2009). Le séquençage et

l'analyse des transcrits (RNA sequencing) réalisés dans ce projet de thèse constituent la première approche de ce genre chez *M. extorquens,* mais aussi la première étude des transcriptomes chez les souches CM4 et DM4.

4.3.3. Etudes de protéomique chez *M. extorquens*

Les études de protéomique chez M. extorquens sont plus nombreuses que celles de transcriptomique (Tableau 1.7, Gourion et al., 2006; Muller et al., 2011; Roselli et al., 2013). Des travaux sur la méthylotrophie, notamment chez *M. extorquens* AM1, ont mis en évidence les protéines induites lors de croissance en présence de méthanol (substrat de référence dans l'étude de la méthylotrophie), en comparaison avec des substrats comme le succinate (C₄H₆O₄) (Laukel et al., 2004; Bosch et al., 2008). Comme décrit précédemment, M. extorquens est capable de dégrader de nombreux composés dont les méthanes chlorés. Des études réalisées chez M. extorquens DM4, en croissance sur du méthanol versus du dichlorométhane, ont permis de mieux comprendre les mécanismes impliqués dans l'utilisation de ces composés (voir chapitre 3) (Muller et al., 2011a). De même, les mécanismes de colonisation de la phyllosphère de A. thaliana par M. extorquens AM1 sont mieux compris grâce à l'utilisation de ces techniques (Tableau 1.7, (Knief et al., 2010, 2012; Vorholt, 2012)). Ainsi, la surexpression de la protéine régulatrice PhyR, nécessaire à la colonisation de la plante a été mise en évidence (Gourion et al., 2006). Concernant l'utilisation de chlorométhane par M. extorquens CM4, 49 protéines plus exprimées en condition de croissance sur chlorométhane par rapport à du méthanol ont été identifiées (Figure 1.17).



Figure 1.17. Image d'un gel 2D des protéines extraites après croissance de *M. extorquens* CM4 en présence de chlorométhane ou de méthanol

Les protéines CmuA, CmuB et PurU, encadrées respectivement en rouge, vert foncé et vert clair, ont été identifiées par spectrométrie de masse et sont plus abondantes en condition chlorométhane (Thèse de Sandro Roselli, 2009).

Cette approche de protéomique combinée à la génomique a permis de mettre en évidence les gènes requis pour l'utilisation du chlorométhane par *M. extorquens* CM4 et également l'importance des gènes portés par le plasmide pCMU01 dans la croissance avec chlorométhane (Roselli *et al.*, 2013).

4.3.4. Etudes de métabolomique chez *M. extorquens*

L'ensemble des études de métabolomique réalisées chez *M. extorquens* AM1 avaient pour but de mieux caractériser le métabolisme méthylotrophe et le métabolisme central d'assimilation du carbone. La croissance sur du méthanol ou sur du succinate met en évidence la présence de métabolites spécifiquement liés à la croissance en condition méthylotrophe ou encore des métabolites intervenant dans la régénération du glyoxylate comme le β -hydroxybutyrate, le méthylsuccinyl-CoA ou encore l'éthylmalonyl-CoA (Figure 1.11) (Guo et Lidstrom, 2008). Des différences dans les métabolites produits ont été observées lors de la croissance de AM1 sur un composé en C₂ (l'éthylamine) et en C₄ (le succinate) (Yang *et al.*, 2009). De même, chez *M. extorquens* AM1 en croissance avec l'acétate (CH₃COO⁻) comme unique source de carbone et d'énergie, la voie de l'éthylmalonyl-CoA remplace la voie de l'isocitrate lyase pour régénérer le glyoxylate, nécessaire à l'assimilation du carbone dans la biomasse via le cycle de la sérine (Schneider *et al.,* 2012, Figure 1.11). Ces résultats ont permis de préciser comment la cellules modifie son métabolisme pour s'adapater aux sources de carbone disponible.

5. Objectif et questions du projet de thèse

Depuis le protocole de Montréal, en 1987, réduisant l'utilisation des composés chlorés, le chlorométhane est responsable de 16 % de la dégradation de la couche d'ozone liée aux composés chlorés (Montzka et al., 2011). Le budget atmosphérique du chlorométhane reste mal évalué même si de nombreuses sources et puits ont été identifiés (Tableau 1.3 etTableau 1.4, Clerbaux et Cunnold, 2007; Montzka et Fraser, 2003). Le rôle du sol, à la fois source et puits de chlorométhane, dans ce budget atmosphérique du chlorométhane n'est plus à démontrer (Harper, 2000; Harper et al., 2003; Gribble, 2003; Miller et al., 2004). Cependant de nombreuses incertitudes persistent quant à l'importance des sols forestiers dans les échanges de chlorométhane avec l'atmosphère (Saito et Yokouchi, 2008). Une des incertitudes est notamment liée à la capacité des bactéries du sol à jouer le rôle de filtre de chlorométhane limitant ainsi son relargage dans l'atmosphère. Toutefois, en théorie, les faibles concentrations de chlorométhane disponibles dans les sols (1 pM. g⁻¹) (Harper et al., 2003) ne pourraient pas à elles seules permettre une croissance suffisante pour le maintien de communautés chlorométhane-dégradantes, qui nécessiteraient l'utilisation d'autres sources de carbone et d'énergie bio-disponibles pour leur croissance. De fait, la concentration du méthanol, estimée à 1 nM.g⁻¹ dans les sols, est nettement supérieure à celle du chlorométhane (Figure 1.18, Stacheter et Kolb, 2013). Dans l'environnement, les microorganismes méthylotrophes utilisant le chlorométhane pourraient utiliser le méthane et le méthanol, en plus du chlorométhane. Les émissions de méthane et de méthanol à partir d'écosystèmes terrestres sont du même ordre de grandeur (10¹² mol par an) (Kolb, 2009a). Le méthanol est un composé central dans le métabolisme méthylotrophe et 83 % des souches méthylotrophes poussant en aérobiose qui ont été isolées du sol sont capables d'utiliser le méthanol (Kolb, 2009). C'est le cas de la souche de référence de la dégradation aérobie du chlorométhane, M. extorquens CM4, incapable d'utiliser le méthane (Van Aken, 2004). Un des objectifs de mon projet de thèse était de caractériser les populations bactériennes méthylotrophes de sols forestiers capables d'utiliser le chlorométhane et de tester l'hypothèse de co-utilisation du chlorométhane et du méthanol.



Figure 1.18. Emissions et consommations de chlorométhane et de méthanol dans les sols forestiers

Données pour les émissions de chlorométhane issues de Harper *et al.*, 2003. Celles pour les flux de méthanol sont issues de Stacheter *et al.*, 2013.

Mon projet s'est articulé autour de deux volets. Dans un premier temps, une étude approfondie de l'utilisation du chlorométhane en conditions contrôlées a été réalisée au laboratoire. Le séquençage de transcrits à haut-débit (RNAseq) visait à identifier l'ensemble des gènes différentiellement exprimés chez *M. extorquens* CM4 en condition de croissance méthylotrophe avec le chlorométhane et le méthanol. Le deuxième volet de mes travaux, basé sur une approche de type Stable Isotope Probing (SIP) (Neufeld *et al.*, 2007) a consisté à mettre en évidence les bactéries méthylotrophes chlorométhane-dégradantes actives dans des microcosmes de sols forestiers. Ainsi des incubations avec du chlorométhane et/ou du méthanol marqués avec du [¹³C] ont été réalisées afin de détecter les bactéries capables d'assimiler ces composés, et donc d'incorporer le [¹³C] dans leurs ADNs. L'ADN enrichi en [¹³C] (ADN « lourd »), correspondant à l'ADN des microorganismes ayant utilisé la source de carbone marquée, a été séparé de l'ADN non marqué (« léger » constitué de [¹²C]) par ultracentrifugation. A partir de l'ADN « lourd », la diversité bactérienne est estimée par séquençage de produits PCR d'un marqueur taxonomique (gène codant l'ARNr 16S) et de marqueurs fonctionnels. Trois gènes marqueurs ont été utilisés ciblant :

- la dégradation aérobie du chlorométhane (gène *cmuA* codant une chlorométhane méthyltransférase de la voie *cmu*);
- la dégradation aérobie du méthanol (gènes *mxaF* et *XoxF* codant une sous-unité de méthanol déshydrogénase) ;

96

- le métabolisme méthylotrophe central (gène mch) (Stacheter et Kolb, 2013).

Ces approches combinées avaient pour but :

 de mieux comprendre l'utilisation du chlorométhane chez une souche modèle cultivée au laboratoire pour éventuellement pouvoir identifier de nouveaux marqueurs de la dégradation du chlorométhane en vue de détecter plus efficacement les bactéries capables de dégrader le chlorométhane dans les sols de forêts tempérées ;

 d'identifier les taxa de bactéries méthylotrophes actifs dans l'utilisation aérobie du chlorométhane en microcosmes de sols forestiers de surface et l'étude de leur rôle dans la bio-capture des émissions de chlorométhane terrestres.

Chapitre 2

Methodological section
Chapter 2. Methodological section

1. Methods used to study *M. extorquens* model strains

During this thesis, eight cDNA banks were constructed using RNA extracted from *M. extorquens* CM4, the model strain for growth with chloromethane (2 biological independent replicates), and from *M. extorquens* DM4, the model strain for growth with dichloromethane (2 biological independent replicates). In addition, RNAs were extracted from both strains grown using a methylotrophic reference substrate, methanol (2 biological independent replicates).

1.1. Aerobic methylotrophic growth

Cultured of 220 mL, were performed into M3 medium defined for *Methylobacterium* with 10 mM of carbon (Muller *et al.*, 2011a). To prepare 1 liter of M3 medium, 1.9 g of KH₂PO₄, 6.39 g of Na₂HPO₄·2 H₂O, 0.1 g of MgSO₄·7 H₂O and 0.2 g de (NH₄)₂SO₄ are dissolved into 800 mL of milliQ water. After autoclaving, 1 mL of Ca(NO₃)₂·4 H₂O at 25 g.L⁻¹ is added with 1 mL of a trace solution. This solution contains H₂SO₄ concentrated (95 %) at 9.2 g.L⁻¹, Fe(II)SO₄·7 H₂O at 1.0 g.L⁻¹, Na₂MoO₄·H₂O at 25 g.L⁻¹, H₃BO₃ at 0.1 g.L⁻¹, CuCl₂·2 H₂O at 0.28 g.L⁻¹, ZnSO₄·7 H₂O at 0.53 g.L⁻¹, NH₄VO₃ at 0.1 g.L⁻¹, Co(NO₃)₂·6 H₂O à 0.25 g.L⁻¹, NiSO₄·7 H₂O at 0.1 g.L⁻¹. Growth conditions and total RNA extractions are described in the article "Contribution of the Core and Variable Genome to the Transcriptomes of two *M extorquens* Strains Grown with Chloromethane, Dichloromethane or Methanol" (see Chapter 3, section 3.2.).

1.2. RNA preparation

The total RNA extraction protocol is described in the Chapter 3 in the Materials and Methods section. No kit was utilized to the RNAs extraction, but a protocol with a phenol/chloroform step. These methods have some advantages in comparison with kits; no restriction in the total amount of RNA extracted, and no size selection. Once total RNA has been extracted, 2 additional steps before cDNA bank construction are needed: rRNA removal and RNA quality check.

1.2.1. DNA removal

A DNase treatment was used to remove DNA contaminants (Turbo DNAse Ambion Thermo Fischer Scientific). A total of 3 µL of Turbo DNAse was added (2U/µL) with 2 µL of RNAsin (Promega), 10 µL of TURBOTM buffer (10 X; Ambion) and 78µL of milli DEPC-treated water. The reactionnal mix is incubated two times, at 30 minutes at 37°C, with 2µL of TURBOTM DNase added between those two incubations. Evaluation of the efficiency of DNA-free RNA extracts was checked by PCR with three sets of primers targeting the 16S RNA-encoding gene *rrsA* and *cliA* present in both strains. Additional PCR targeted genes *cmuA* and *dcmA*, which are specific of strains CM4 and DM4 respectively, were also used (see article "Contribution of the Core and Variable Genome to the Transcriptomes of two *M. extorquens* Strains Grown with Chloromethane, Dichloromethane or Methanol" (see Chapter 3, section 3.2.). PCR conditions are summarized in Table 2.1.

Table 2.1. P	CR amplificatior	program to	validate DNA	removal
I GIALC ELT I I	en ampinication			10110101

Mix	Volume (μL)
H2O milliQ water	17.6
PCR buffer (Biolabs, 10X)	2.5
dNTP 20mM each	0.25
Forward primer (20µM)	1
Reverse primer (20µM)	1
Taq polymerase	0.1
DNA	1

PCR program					
	Pre-denaturation, 2 min at 95°C				
40 cycles	Denaturation, 30 sec at 94°C				
	Annealing, 45 sec at 60°C				
	Extension, 30 sec at 72°C				
Final extension, 10 min at 72°C					

1.2.2. rRNA removal

The abundance of ribosomal RNA (rRNA) accounts for 95-98 % of total RNA in bacteria. Thus, a step to efficiently remove rRNA is necessary to achieve optimal coverage of mRNA, good detection sensitivity and reliable results. The necessity to optimize efficient rRNA removal methods for RNA sequencing in other GC-rich bacteria has been described previously (Peano *et al.*, 2013). Two types of rRNA depletion approaches exist: the first one is based on a selective degradation of rRNAs by an enzyme recognizing specifically the 5' monophosphate extremity of rRNAs, whereas mRNAs are protected by their triphosphates extremities (Evguenieva-Hackenberg and Klug, 2011). The second method relies on subtractive hybridization with probes specifically designed to target rRNA conserved regions (van Dijk *et al.*, 2014). This second method was the most efficient in rRNA depletion of samples extracted from *M. extorquens* strains as tested at the Génoscope in Evry (experiments performed by

Béatrice Ségurens and Adriana Albertini). Extracted RNA samples were treated with the RiBoZero Magnetic kit Gram-negative bacteria (Tebu-Bio). Evaluation of the efficiency of RNA removal was checked by PCR with primers targeting the 16S RNA-encoding gene *rrsA* and *cliA* as a control. Despite the validation of rRNA removal by PCR and bioanalyzer profil, 14 up to 34 % of the RNAseq reads mapped to rRNA (Chapter 3 (section 3.2, Table 1). Thus, the RiboZero Magnetic Kit Gram-negative bacteria is not optimized to target *M. extorquens* rRNA (data confirmed by the Tebu-Bio company).

1.2.3. RNA quality control

Enzymes that degrade RNA, ribonucleases (RNases), are so ubiquitous (reagents, skin, dust) that working in an RNase-free environment can be challenging. The integrity of the extracted RNA was systematically checked using the Agilent biotechnology 2100 Bioanalyzer System. This system is based on classical gel electrophoresis adapted to a chip format. The RNA nano and RNA pico kits are used to detect RNA in the range of 5-500 ng.μL⁻¹ and 50-5000 pg.μL⁻¹ respectively. Each RNA sample is mixed to a fluorescent dye and loaded into a well connected to micro-channels for capillary electrophoresis. The dye binds to nucleic acids, then RNA-dye complex are detected by laser-induced fluorescence. Fluorescence is recorded for nucleic acid visualization. RNA fragments are separated based on their size and fragment size assignment is deduced from the electrophoregram of a co-migrating RNA size ladder (Figure 2.1).



Figure 2.1. Electrophoregram of RNAs extracted from *M. extorquens* CM4 growing with chloromethane

Peak area is assigned in arbitrary fluorescence units (FU). (A) RNA nano kit-based electrophoregram of RNA before rRNA depletion (RIN value of 6.7). In *M. extorquens* an additional band is observed. (B) RNA pico kit-based electrophoregram of RNA after the rRNA depletion (RIN value of 2.6).

Profiles generated on the Agilent 2100 Bioanalyzer System yield information on concentration, allow a visual inspection of RNA integrity, and generate ribosomal ratios (16S versus 23S rRNA). Two kits are available for RNA analysis: RNA nano and RNA pico kits, depending on their sensibility (5-500 ng.μL⁻¹ and 50-5000 pg.μL⁻¹⁰ respectively). The manufacturer Agilent Technologies has introduced the RNA Integrity Number (RIN). The RIN is an algorithm for assigning integrity values to RNA measurements: a value of 1 means that RNAs are totally degraded while a value of 10 means "perfect" RNAs integrity with no detected degradation. The RIN value is defined by a complex calculations fixed by the software using the ratio 16S/23S. Moreover, the RIN value should be use with caution. In some bacteria, RNAs can be naturally fragmented, as described in *M. extorquens* (Zahn *et al.*, 2000; Evguenieva-Hackenberg, 2005). This maturation will lead to the 23S fractionation, illustrating by the apparition a new peak between those for the 16S and 23S, as observed in Figure 2.1.

1.3. Construction of the directional cDNA bank

To construct the cDNA banks, Béatrice Segurens and Adriana Albertini performed preliminary tests at the Génoscope in Evry. Three protocols were tested basing on the TruSeq Small RNA, the TruSeq stranded mRNA, and a Génoscope "house" protocol. The TruSeq stranded mRNA LT kit (Illumina) was chosen. The directional bank relies on dUTP incorporation during the second strand synthesis. This will avoid the amplification of the second strand, keeping only information from the first strand cDNA. The addition of two different adapters, in 3' and 5' extremities will keep the strand orientation. Fifty to sixty ng of rRNA-depleted RNA were used to construct directional cDNA banks as described in Figure 2.2.



Figure 2.2. Directional cDNA bank construction

The TruSeq stranded mRNA kit (Illumina) was used. (1) RNAs were primed with random hexamers to synthetize the first cDNA strand by reverse transcription. (2) The complementary cDNA strand was synthetized with the incorporation of dUTP instead of dTTP, whereas the RNA template was degraded; (3) A single adenosine nucleotide (A) was added to the 3' ends of the blunt extremity of each cDNA fragment to prevent them from ligation with other cDNA fragments; (4) Ligation of multiple indexing adapters to both the 5' and 3' ends of the cDNA fragments; (5) Selectively enriched DNA fragments: PCR amplification and treatment with magnetic beads (AMP Xpure, Agencourt, Agilent) for PCR product size selection; (6) cDNAs with a size around 260 bp were hybridized to surface-bound primers in the flow cell. Then sequencing can start.

The cDNA bank constructions were validated if the final product had an expected size around 260 bp using the Agilent DNA 1000 kit. An example of an electrophoregram of cDNA with a DNA chip is shown (Figure 2.3). Validated cDNA banks were sent to the Génoscope to be sequenced in paired-end with the HiSeq2000.



Figure 2.3. Validation of cDNA bank constructions with the 2100 Bioanalyzer

2. Methods used to study chloromethane-degrading bacterial communities in forest soils

The environment is a complex system, with a high bacterial diversity, and a lot of different metabolisms. The majority of bacteria found in the environment is not cultivable in the laboratory (Pham and Kim, 2012) so that the identification in situ of a specific bacterial associated-metabolism may be challenging (Rappé and Giovannoni, 2003). Cultivationindependent methods have been developed to identify which microorganisms are carrying out a specific set of metabolic processes in the natural environment (Coyotzi et al., 2016). Among those methods, the stable isotope probing (SIP) method relies on the incorporation of a labeled substrate that is highly enriched in a stable isotope, such as [¹³C], in order to identify the active microorganisms by the selective recovery and analysis of isotope-enriched cellular components (Dumont et al., 2011). The stable isotope of carbon [¹³C] is found naturally although it only represents about 1.1 % of all natural carbon on Earth while carbon [¹²C] makes up to 98.9 %. The labeling of microorganisms with the [¹³C]-carbon is the direct consequence of the [¹³C]-labelled substrate consumption. The approach has been applied to lipids (Evershed et al., 2006), nucleic acids (Neufeld et al., 2007) and proteins (Jehmlich et al., 2008). The incorporation of heavy isotopes into newly synthesized nucleic acids has been used to characterize the diversity of active bacteria in soil, after separation of newly synthesized from existing rRNA and DNA in RNA-SIP and DNA-SIP experiments, respectively (Rettedal and Brözel, 2015). RNA-SIP targets transcriptionally active cells producing new ribosomes after addition of the label, whereas DNA-SIP focuses on DNA replicating, yielding an inventory of all dividing cells (Chen and Murrell, 2010). Since rRNA-SIP is more sensitive and does not require cell division to label RNAs, the RNA-SIP approach was envisaged in addition to the DNA-SIP to characterize active chloromethane-degraders in forest soils. Nevertheless, reverse transcribed-soil RNA, were used as template for PCR amplification of *cmuA*, *mch* and *mxaF/xoxF*, but no amplification was observed, that's why RNA were not further studied. Moreover, the DNA-SIP approach allowed subsequent metagenomic analysis.

2.1. Soil sampling and preincubation

The soil sampling was performed at a forest site that now serves as a model forest ecosystem for investigation of methylotrophic communities of temperate climate zone is the forest at Steigerwald (49°52'N, 10°28'E; sandy loam (soil type: Dystric Cambisol) with a pH of 4.6; mean annual temperature: 7.5°C; mean annual precipitation: 725 mm). The Steigerwald forest covers an area of 115,2 km² located between Würzburg and Nürnberg in Germany. And though the area is developed, there are no industrial activities nearby and access is limited. A total of 5 different samples were taken from the upper layer in an area of 20 m of diameter, in September 2014. Mycelium and leaves were removed. The soil samples were sieved through a 2 mm screen and 500 g of each sieved soil sample were mixed. Before the SIP experiment per se, a pre-incubation was performed during two weeks at 20°C in the dark in presence of 1 % of [¹²C]-CH₃Cl per 270 g in 1L Müller-Krempel flasks. The resulting activated soils were exposed to air to remove remaining chloromethane gas, and mixed together. The average water content was determined to be of 18 % by measurement of the weight difference before and after drying (ten days at 60°C) of 1.2 g of soil placed into an Eppendorf tube.

2.2. Stable isotope probing (SIP): the [¹³C]-chloromethane labeling step

Microcosms were set up using 100 g of the activated soil introduced into sterile 0.5 L Müller-Krempel flasks. Chloromethane and methanol were added alone or in combination in a total of 5 successive carbon pulses per condition. The next carbon pulse occurred when chloromethane concentration decreased below 10 % of the initial concentration. In parallel, each methanol pulse took place at the same time as the reference chloromethane pulse. To optimize carbon availability and homogenization without transforming the soil into mud, 1 mL of liquid phase (water or methanol) was added into each microcosm per pulse. Control conditions without added carbon sources were included in the study (Figure 2.4). Each pulse of chloromethane corresponded in the addition of 1 % of CH₃Cl. This resulted in a final concentration of 18 mM (volume flask of 0.5L; chloromethane solubility of 5 g. L⁻¹). Equivalent amounts of chloromethane and methanol were added per each pulse (see Table 4.1 in Chapter 4). Thus, for methanol, 1mL of a methanol stock solution at 216 nM was

added per pulse (70g of soil contains 12 mL of water) to obtain a final concentration of 18 mM.





During the course of the labeling experiment, four parameters were monitored:

- a) pH. CH₃Cl consumption or CO₂ reactions may produce protons and subsequently modify bacterial diversity and hinder the detection of the chloromethane-degrading communities. To be sure that pH values do not change during the experiment, the pH was checked on sieved soil, before the pre-incubation step, and after each carbon source pulse. The pH was measured by suspending 0.5 g of soil into 1 mL of water using a pH electrode (R422 InLab Semi-Micro; Mettler-Toledo).
- b) Chloromethane concentrations. Chloromethane concentrations were monitored using gas chromatography (GC) directly after CH₃Cl has been added to the microcosms. To correlate the peak area and CH₃Cl concentration, a calibration curve was done, by using 500 μL of standard CH₃Cl concentrations prepared at 10 %, 2 %, 1 %, 0.5 %, 0.25 %, 0.001 % and 0.005 % in serum flasks (correlation factor R² of 0.95).
- c) **Total CO₂ concentration.** The CO₂ was measured by using a thermal conductivity detector (TCD). For more details, see Chapter 4. As for CH₃Cl, a calibration curve was realized to correlate peak areas and CO₂ concentrations.

d) [¹³C]-CO₂. A carbon source can be assimilated into biomass, or reduce to produce energy leading to a CO₂ production. To estimate the part of carbon mineralized, the [¹³C]-CO₂ was measured by gas chromatography coupled with mass spectrometry (GC-MS).

2.3. DNA extraction from soil

The bacterial community DNA was extracted from soil after the third chloromethane addition pulse according to Griffith et al., (2000) with minor modifications. Elution buffer was prepared by mixing equal volume of potassium-phosphate buffer and hexadecyltrimethylammonium bromide (CTAB)/ NaCl solution. The potassium-phosphate buffer was prepared by mixing 39.30 g of K_2 HPO₄ and 1.96 g of KH₂PO₄ qsp 1000 mL of milliQ H₂O and adjusted at pH 8. The CTAB/ NaCl solution was prepared with 10 g of CTAB with 4.09 g qsp 100 mL of DEPC water and then heated at 65°C. After mixing potassium-phosphate buffer and CTAB/NaCl solution, the elution buffer was mixed and heated at 60°C. Then, 0.5 mL was added to 0.5 g of soil previously placed into a tube with sterile zirconium beads (0.5 g of those of 0.1 mm and 0.5 g of 0.5 mm diameters, Roth) before to be mixed using a vortex. A volume of 0.5 mL of phenol/chloroform/isoamyl alcohol (ratio 25:24:1) was added before to place the tube in a bead beater for 30 seconds at 5.5 m.sec⁻¹ and immediately placed on ice before to be centrifuged at 4°C at 13,000g during 5 minutes. The upper phase was transferred in another tube and mixed by inversion with chloroform/isoamylalcohol (24:1) and centrifuged at 4°C at 13,000 g during 5 min. The supernatant was transferred to another tube with 2-3 volume of precipitation buffer, mixed and incubated at room temperature for two hours. This buffer was prepared with 30 % PEG 3,000 at pH 7 in 0.1 M of 2-[4-(2-hydroxyethyl) piperazin-1-yl] ethane sulfonic acid (HEPES). The samples were centrifuged at 4°C during 10 min at 13,000 g then the pellet was washed with ethanol 70 % before to be centrifuged again, and dried at room temperature. The DNA pellet was suspended in 30 µL DEPC water and treated with RNAse A to remove contaminant RNAs. DNA was quantified by fluorescence with the Quant-iT™ PicoGreen[®] kit (Invitrogen).

2.4. Separation of isotopically-labeled DNA by isopycnic density gradient

The "heavy"-DNA is separated from the "light"-DNA by ultracentrifugation in a cesiumchloride (CsCl) gradient where DNA molecules migrate to the point where they have the same density as the gradient. The CsCl solution and gradient buffer were prepared as recommended (Neufeld *et al.*, 2007). To dissolve CsCl in milliQ water (1 g mL⁻¹), the solution was heated at 30°C. The density of the CsCl solution at 20°C absolutely needs to be between 1.88 and 1.89 g mL⁻¹. To prepare the gradient buffer, 3.75 g of KCl, 50 mL of Tris-HCl, 1 mL of 0.5 M EDTA were dissolved in a final volume to 500 mL of milliQ water. The CsCl solution and the gradient buffer were mixed in the ratio 4:1 to give an expected density of 1.725 g.mL⁻¹ at 20°C. This gradient solution (4.98 mL) was mixed to DNA (5 µg in 20 µL of RNA-free milliQ-water), and transfered into a column Quick-Seal (Beckman Coulter). A control (without added nucleic acids) was run in parallel to validate the density gradient after ultracentrifugation. Each tube was sealed and placed into the rotor Vti 65.2 (Beckman Coulter) for ultracentrifugation for 40 hours at 177,000 g at temperature 20°C (Figure 2.5). To avoid gradient destruction, a deceleration phase of 2 hours was programmed. After the centrifugation, the "heavy"-DNA and the "light"-DNA were located at the bottom or the top of the tube, respectively. Immediately after the rotor stopped, the isotopically-labeled and unlabeled nucleic acids present along the isopycnic density gradient were separated in 10 fractions per tube. According to the literature, the collected fractions 6-7 are expected to have a density of 1.725 g. mL⁻¹ and all other fractions a density comprised between 1.690 and 1.760 g.mL⁻¹ (Neufeld *et al.*, 2007). The density of the collected fractions was established from a standard curve established by weighting each fraction of the control tube after the temperature has been stabilized at 20°C in a water bath.



Figure 2.5. Material required for ultracentrifugation and fractionation

(A) Nucleic acids dissolved in the gradient solution are transferred to a microcentriguge tube (Tube, Quick-Seal, Beckman Coulter); (B) A total of 16 tubes can be placed vertically into the rotor Vti 65.2 (Beckman Coulter). Each tube is circumscribed in a rotor well with a screw-on adaptor; (C) Ultracentrifuge (LE-70, Beckman Coulter); (D) After centrifugation, each tube containing the gradient is fractionated to split heavy, middle and light fractions. Each tube is fixed and pierced carefully in the top and bottom with needles (30 x 0.63 mm, BD). Using a low-flow pump (BioRad Econo Pump), a solution of water containing-dye was injected into the top with a flux fixed at 450 μ L.min⁻¹. Drops are collected during one minute before to change the tube to collect the next fraction. This technique provides smooth delivery of displacement liquid, which minimizes disruption of the gradient and ensures a consistent volume in the collected fractions.

The collected nucleic acid fractions were precipitated for 2 hours at 4°C in presence of 1 μ L of glycogen (20 μ g. μ L⁻¹) and 2 volumes of polyethylene glycol PEG 6000 30 % / 1.6 M NaCl solution (150 g PEG 6000, 46.8 g of NaCl dissolved in total volume of 500 mL of milliQ water). Precipitated nucleic acids were harvested by centrifugation at 13,000 g at 20°C during 30 minutes, washed once with ethanol 70 %, dried at room temperature, and resuspended in 30 μ L of TE Buffer previously prepared to have a final concentration of 10 mM Tris-HCl (pH 8), and 1 mM EDTA (pH 8).

2.5. Metagenomics of PCR-amplified markers

To assess the diversity of chloromethane-degraders in forests soil, the *rrsA* gene was amplified in the different microcosms prepared. In addition, genes encoding enzymes associated with mono carbon metabolisms are suitable to detect methylotrophs in the environment (McDonald *et al.*, 2005). The genes *cmuA*, *mxaF/xoxF* and *mch*, encoding respectively a chloromethane deshalogenase, a methanol dehydrogenase subunit and methenyl cyclohydrolase were targeted.

The corresponding PCR products were sent for sequencing at LGC Genomics GmbH (Berlin, Germany). To reduce sequencing costs, PCR fragments were extended by small nucleotides sequences called barcode and specific to each sample. A total of 48 barcode sequences were used; 24 barcodes for the 16S PCR amplicons (eight microcosms, with three fractions by microcosm (heavy, middle and light)), and 24 for functional genes.

2.5.1. Design of new primers targeting environmental chloromethane dehalogenase CmuA-encoding gene

PCR amplification targeting the diversity of *cmuA* in samples from contrasting environments such as soil, marine ecosystems, and phyllosphere have been performed previously using primer targeting the end of the methyltransferase domain (Miller *et al.*, 2004; Borodina *et al.*, 2005; Nadalig *et al.*, 2011) (Table 2.2; box 1). These studies as well as sequencing of new chloromethane-degrading *cmuA* gene enriched the database allowing the design of new primers, which may be better in detecting *cmuA* genes in the environment for NGS and associated qPCR analysis.

Table 2.2. Set of primers available for *cmuA* amplification in the environment

Utilization	Product (bp)	Primer orientation, name, sequence 5'->3' (degeneracy rate) ^b	CM4 genome position (nt) ^c	GC (%)	Sequences with mismatch (number of aligned sequences) ^d	Reference
Box1 NGS	442	Forward, cmuA802F, TTCAACGGCGAYATGTATCCYGG (4)	336191-336213	52.2	Vibrio orientalis, Thermosediminibacter oceani, Desulfotomaculum alcoholivorax (32)	Miller <i>et al.,</i> 2004
		Reverse, cmuA1244R, TABTCCATDATGGCYTCGAC (18)	335790-335771	56.3	Marine methyl halide utilising bacteria, Vibrio orientalis, Thermosediminibacter oceani, Thermincola potens, Desulfomonile tiedjei, Desulfotomaculum alcoholivorax (113)	PhD thesis, Farhan Ul Haque
ox1 qPCR 1	166	Forward, cmuA802F, TTCAACGGCGAYATGTATCCYGG (4)	336191-336213	52.2	Vibrio orientalis, Thermosediminibacter oceani, Desulfotomaculum alcoholivorax (32)	Miller <i>et al.</i> , 2004
	100	Reverse, cmuA968R, CCRCCRTTRTAVCCVACYTC (166)	336066-336047	57.0	Vibrio orientalis (113)	Nadalig et al., 2011
Box2 NGS	422	Forward, cmuAf422 GARGTBGGITAYAAYGGHGG (24)	336046-336066	52.5	Vibrio orientalis (113)	This work
		Reverse, cmuAr422 TCRTTGCGCTCRTACATGTCICC (4)	335666-335644	52.2	Thermosediminibacter oceani, Vibrio orientalis, Desulfomonile tiedjei, uncultured bacterium (113)	This work
Box2 qPCR 1	101	Forward, cmuAf422 GARGTBGGITAYAAYGGHGG (24)	336046-336066	52.5	Vibrio orientalis (113)	This work
	121	Reverse, cmuA1070R AYRTCCATCGGRTAYTTRAT (32)	335964-335945	38.0	Vibrio orientalis (113)	This work
Box2 qPCR	144	Forward, cmuA1227F, CGARGCSATIATGGAITAYGA (8)	335767-335788	43.0	Vibrio orientalis, Thermincola potens (113)	This work
		Reverse, cmuAr422 TCRTTGCGCTCRTACATGTCICC (4)	335666-335644	52.2	Vibrio orientalis, Thermincola potens (113)	This work

^a Sets of primers target either box 1 or box 2 of *cmuA* genes, which are localized at

nucleotides 336191-335790 and 336191-335790, respectively for *M. extorquens* CM4 genome (see footnote ^c). NGS stands for Next Generation Sequencing (MiSeq from Illumina)

^b Number of sequences in degenerate primers synthesized as mixtures of different sequences with B=C/G/T, D=A/G/T, H=A/C/T, I=inosine, R=A/G, S=G/C, Y=C/T, V=A/C/G, W=A/T

^c MaGe database http://www.genoscope.cns.fr/agc/mage)

^d Divergent sequences include sequences with a 3'-terminal mismatch (in bold) or one mismatch at the second last position of the 3'-terminal position or at least three mismatches. Sequence accession n° Desti_5447 (*Desulfomonile tiedjei* DSM 6799, Uniprot); TherJR_0143 (*Thermincola potens* JR, Uniprot); Toce_1533 (*Thermosediminibacter oceani* DSM 16646, Uniprot); VIA_000821 (*Vibrio orientalis* ATCC33934, Uniprot); WP_027363948 (Desulfotomaculum alcoholivorax, GenBank); Gi/66474906 (Marine methyl halide utilizing bacteria); Gi/664795/1 (Uncultured bacteria). *Desulfotomaculum kuznetsovii* DSM 6115 was excluded from this study. A total of 113 *cmuA* sequences were retrieved from NCBI database after a BLAST search using *M. extorquens cmuA* (Mchl_5697) sequence, and subsequently aligned using clustalW-multialign, Mobyle (http://mobyle.pasteur.fr/cgi-bin/portal.py#forms::clustalw-multialign alignment). Most divergent sequences shared 48.9 % of aa identity. Among those sequences, uncultured bacterium isolated from *cmuA* sequences from strains detected in soil (Miller *et al.*, 2004; Borodina *et al.*, 2005), marine environment (Cox *et al.*, 2012) and also *cmuA* sequences from *Proteobacteria* isolates that belong to the class of:

- Alphaproteobacteria of the genus Methylobacterium (strain CM4), Hyphomicrobium (strains AT1, AT2, AT3, MC1), Aminobacter (strains lissarensis CC495, ciceronei IMB-1, TW3), Rhodobacteraceae (strains 179, 198);
- Deltaproteobacteria of the genus Desulfotomaculum (D. alcoholivorax, D. kuznetsovii) and Desulfomonile tiedjei;
- Gammaproteobacteria such as Vibrio orientalis.

Two *cmuA* sequences of *Clostridia* isolates were also added in this study (*Thermincola potens* JR and *Thermosediminibacter oceani*).

Based on the alignment, new sets of primers (box 2) were designed for NGS and qPCR primers (Table 2.2). The PCR program for Box 2 targeted primers starts with pre-denaturation of 10 min at 95°C, then 30 cycles of two steps: denaturation of 15 seconds at 94°C followed by an annealing/elongation of 60 seconds at 56°C. The program ends with a post-elongation of 5 minutes at 72°C. The program for the primer set cmuA802F/cmuA1244 (box 1) was performed as previously described (Miller et al., 2004). To validate those new primers, PCR amplifications were performed on three different forest soils, a native forest one, a [¹²C]-CH₃Cl pre-incubated forest soil, and one incubated with [¹³C]-CH₃Cl, and compared to the old primers (box 1). Amplifications were more specific with the new primers cmuAF422/cmuAR422 (box 2) than cmuA802F/cmuA1244R 1), with (box but also more efficient with primers cmuA802F/cmuA1244R ten time more sensitive than others (Figure 2.6). The new primer set is still unable to amplify the cmuA gene found in Vibrio orientalis (Table 2.2). This organism is unable to dechlorinate chloromethane as tested by Ludovic Besaury, a post-doctorate fellow in the laboratory (data not shown). To confirm that no Vibrio orientalis cmuA-like sequence is present into the studied forest soil, a PCR reaction with primer set specially designed to target *Vibrio orientalis cmuA* sequence was performed. No amplification was detected in the DNA from soil while the positive control with *Vibrio orientalis* DNA template was amplified.



Figure 2.6. Comparing two sets of primers targeting environmental chloromethane dehalogenase CmuA-encoding gene

Gene *cmuA* PCR amplification targets the region between the conserved domains (in grey color) IPR006360 and IPR006158 of the encoded protein CmuA. A) Position of the primer box 1 (Miller *et al.*, 2004) and primer box 2 (this study as detailed in Table 2.2). For each primer, its corresponding name, localization and orientation (arrow) is shown. Amplification DNA fragments are indicated with lines, in grey when designed for qPCR only. Number indicate product size in bp. B) PCR-amplified *cmuA* products using reference genomic DNA and environmental DNA. Amplifications were performed using 10 ng of DNA. Track 1-3, DNA from isolated chloromethane-degrading strains including the negative control *Leisingera methylohalidovorans* (track 1) (*cmuA* undetected, Schäfer *et al.*, 2004); *M. extorquens* CM4 (2) and *Hyphomicrobium* sp. MC1 (3) are the positive controls. Tracks 4-6, environmental DNA from a native forest soil (4), a [¹²C]-CH₃Cl pre-incubated forest soil (5), and a forest soil incubated with [¹³C]-CH₃Cl (6). C) Sensitivity of two sets of primers sets, in PCR-amplification of *cmuA* using DNA from isolated chloromethane-degrading strains including strains including the negative control *Leisingera methylohalidovorans* and a forest soil incubated with [¹³C]-CH₃Cl (6). C) Sensitivity of two sets of primers sets, in PCR-amplification

2.5.2. Design of new primers targeting environmental methanol dehydrogenases MDH-encoding genes

In soil, species belonging to *Alpha-*, *Beta-*, and *Gammaproteobacteria*, *Verrucomicrobia*, *Firmicutes*, and *Actinobacteria* were described as involved in methanol oxidation (Kolb, 2009b). A pyrroloquinoline quinone (PQQ)-dependent methanol dehydrogenase encoded by *mxaFl* is found in Gram negative bacteria including Proteobacteria (Lidstrom, 2006; Chistoserdova *et al.*, 2009). A similar gene, *xoxF*, encoding a (PQQ)-dependent methanol dehydrogenase, occurs in all known methylotroph genomes and has been described as essential for growth in the presence of methanol (Chistoserdova *et al.*, 2011, Schmidt *et al.*, 2010). It was described that *mxaF* and *xoxF* can be present simultaneously in genomes of methylotrophs (Skovran *et al.*, 2011b). But *xoxF* and not *mxaF* is expressed even in samples where both genes have been detected (Beck *et al.*, 2015). This prompted us to design new primers that target both *mxaF* and related *xoxF* gene (Figure 2.7).



Figure 2.7. PCR amplification of methanol dehydrogenase-like MxaF and XoxF encoding genes

The dotted line shows the PCR product only detected with the "old primer set".

A total of 120 *mxaF/xoxF* sequences, mostly found in sequenced genomes, were aligned using clustalW-multialign, Mobyle (http://mobyle.pasteur.fr/cgi-bin/portal.py#forms::clustalw-multialign alignment). Based on this alignment, primer 1555R previously used to target *mxaF* in soil (Stacheter *et al.*, 2013) would poorly detect *xoxF* (Table 2.3), whereas the newly designed primer sets target all the aligned *mxaF* and *xoxF* genes (Table 2.3).

Table 2.3. Primer sets available for *mxaF_xoxF* amplification for NGS analysis in the environment

Product (bp)	Orientation, name, sequence 5'->3' (number of expected sequence) ^a	CM4 genome position (nt) ^b	GC content (%)	Primer occurrence: sequences with mismatch (number of aligned sequences) ^c	Reference
549	Forward, mxaF_1003f GCGGCACCAACTGGGGCTCGT	4786435-786452	67.6	Hyhomicrobium denitrificans ATCC 51888, Hyphomicrobium sp. MC1, Hyphomicrobium nitrativorans NL23, Methylocella silvestris BL2, Methylobacterium extorquens PA1, Methylobacillus flagellatus KT, Methylobacterium nodulans strain ORS2060 (nd)	McDonald <i>et al.,</i> 1997
	Reverse, mxaF_1555r CATGAABGGCTCCCARTCCAT (6)	4785922-785902	59.5	nd	Neufeld <i>et al.,</i> 2007
551	Forward, Mdh1/2F GCGGIWS <u>C</u> AICTGGGGGYT (8)	4786432-786452	69.0	Methylocella silvestris BL2 (97)	This work
	Reverse, MdhR GAASGGYTCSYARTCCATGCA (32)	4785925-785905	55.5	(120)	This work
551	Forward, Mdh2/2F GCGGIWS <u>G</u> AICTGGGGYT (8)	4786432-786452	69.0	(23)	This work
	Reverse, MdhR GAASGGYTCSYARTCCATGCA (32)	4785925-785905	55.5	(120)	This work

^a Degenerate primers are synthesized as mixtures of different sequences with B=C/G/T, D=A/G/T, H=A/C/T, I=inosine, R=A/G, S=G/C, Y=C/T, V=A/C/G, W=A/T

^b MxaF position, MaGe database http://www.genoscope.cns.fr/agc/mage)

^c Divergent sequences include sequences with a 3'-terminal mismatch (in bold) or one mismatch at the second last position of the 3'-terminal position or at least three mismatches. Sequence accession n° Hden_1305/ Hden_1617/ Hden_2848 (*Hyhomicrobium denitrificans* ATCC 51888, Uniprot); HYPMCv2_0800/ HYPMCv2_3613 (*Hyphomicrobium* sp. MC1, Uniprot); Mext_0099 (*M. extorquens* PA1), Mfla_1451 (*Methylobacillus flagellatus* KT, Uniprot); Mnod_2344 (*Methylobacterium nodulans* strain ORS2060, Uniprot); Msil_2260 / Q8KMI4_METSI (*Methylocella silvestris* BL2, Uniprot); W911_13290 (*Hyphomicrobium nitrativorans* NL23, Uniprot).

3. <u>Illumina sequencing</u>

The Illumina technology relies on sequencing by synthesis (SBS) and occurs on a solid phase, called a flow-cell, where short DNA sequences are anchored (Figure 2.8). The DNA fragments are immobilized on the flow cell surface thanks to the ligation of a DNA sequence complementary to the fixed short sequences. A flow-cell has the dimension of a microscope slide (26 x 76 mm) with 8 lanes. It is designed to facilitate access to enzymes. Densities on the order of ten million single-molecule clusters per square centimeter are achieved and the tens of millions of clusters on the flow cell surface are sequenced in parallel. Solid-phase amplification creates up to 1,000 identical copies of each single template molecule in close proximity (diameter of one micron or less). The sequencing is based on the using of four fluorescent tagged reversible terminator-bound dNTP (A, T, C, G). During each sequencing cycle, a single labeled dNTP is added to the nucleic acid chain. After each dNTP incorporation, the fluorescent dye is imaged to identify the base added and then enzymatically cleaved to allow incorporation of the next labeled nucleotide. Since all four labeled dNTPs are present as single, separate molecules, natural competition minimizes incorporation bias. Base calls are made directly from signal intensity measurements during each cycle, which greatly reduces raw error rates compared to other next generation sequencing technologies. The end result is highly accurate base-by-base sequencing that eliminates sequence-context specific errors, enabling robust base calling across the genome, including repetitive sequence regions and within homopolymers. Fragments in a range size of 150-400 bp can be sequenced using the Illumina technology.



Figure 2.8. Schematic representation of Illumina 'Genome Analyser' flow cell for high throughput sequencing

(a) The flow cell contained many capillaries, where millions of DNA fragments are anchored for sequencing. (b) Cross section of flow cell showing second strand synthesis. Laser light activates base-specific emission color from modified nucleotides incorporated at the end of each cycle. (c) Imaging the surface of the flow cell after each cycle records the base added at each 'spot' of DNA. (d) Image analysis 'reads out' the sequence of incorporated second strand bases at each spot (from Mike Gilchrist and was first published in the 2010 *Mill Hill Essays*, National Institute for Medical Research, http://www.historyofnimr.org.uk/mill-hill-essays/essays-yearly-volumes/2010-2/bringing-it-all-back-home-next-generation-sequencing-technology-and-you/).

The data generated by HiSeq2000 and MiSeq Illumina platforms are similar, but differ in read lengths, scale and therefore support different applications. For large projects where time is less of an issue but cost per sequence is a major concern, the HiSeq platform allows massively parallel sequencing at the lowest cost. Comparable data can be generated on the MiSeq for smaller projects where it is important to process samples quickly, for example, in environmental microbial community assessment (Caporaso *et al.*, 2012; Schmidt *et al.*, 2013). The price per sequence decreases on Illumina HiSeq2000 and MiSeq platforms, democratizing the sequencing (Tringe and Hugenholtz, 2008). Multiplexing is also used to reduce sequencing

costs. The incorporation of a barcoded nucleotide sequence specific to each sample was performed for both the RNAseq and the SIP studies (see Chapters 3 and 4), which are specifically recognized by the sequencer (Figure 2.8, Shearer *et al.*, 2012).

4. Bioinformatics

4.1. MicroScope platform

The MicroScope platform was developed by Génoscope (Evry, France). It provides an access to easy-use-tools to study, annote, and perform comparative genomic analysis of bacterial genomes (Vallenet *et al.*, 2013). This database relies on three major components; (i) the PKGDB (Prokaryotic Genome DataBase), used to access, store and modify information on organisms, sequences and genomic objects, (ii) a panel of bioinformatic tools linked to the PKGDB database to provide results of syntactic and functional annotation pipeline as metabolic pathway tools (KEGG, MicroCyc) and analysis for each genome; (iii), a graphic interface, MaGe (Magnifying Genome) allowing manual annotations and comparative genomic analysis (Vannellet *et al.*, 2006 and 2009). The MaGe database interface combines graphical interfaces and the analysis of gene conservation (synteny), together with genome data sources and metabolic pathway tools to assist in data evaluation in order to manually assign the best possible annotation to a given gene product (Figure 2.9) (Vannellet *et al.*, 2006; Vannellet *et al.*, 2009). It offers useful tools to explore genome data according to the specifications and requests of the annotator. MicroScope can be used as an open or restricted access resource.



Figure 2.9. A screen capture of the "MicroScope" web platform

The screenshot was taken during analysis of the genome of *M. extorquens* CM4 in the "Genome Browser" window. Several tools (A) can be used to explore the genome of the selected (pivot) organism (B). (C) Genomic objects (predicted genes in the 6 frames) are displayed as rectangles, along with (D) genes in synteny in other organisms in the MicroScope database (PkGBD, http://www.genoscope.cns.fr/agc/microscope) and publically available genome sequences (Refseq).

4.2. Transcriptome Analyses based on MAssive sequencing of RnAs: RNAseq database

TAMARA is the acronym for "Transcriptome Analyses based on Massive sequencing of RnAs" included in the MicroScope platform, which was developed by the Génoscope team at Evry (France). TAMARA pipeline enables to preprocess raw sequencing reads, map reads on reference genomes implemented in the microscope platform, compute transcript coverage along genome and expression levels for annotated genes, and evaluates differential biological expression between tested samples (https://www.genoscope.cns.fr/agc/microscope/transcriptomic/NGSProjectRNAseq.php ?projType=RNAseq&wwwpkgdb=8da0d938cc9a2c3d80982c6a2e87f0cf). It is a userfriendly tool for RNAseq database analysis. This platform stores data from the DESeq pipeline. Of the different methods available to analyze differentially-expressed genes, DESeq is one of the most effective (Qin et al., 2016). TAMARA platform converts data from DESeq and make them accessible for users and connect them with data from annotated genomes available on the MicroScope database. This facilitates data treatment and their modifications, making the analysis easy to do (Table 2.4).

Strain	Group	Carbon source	Name of independent biological samples
	C1		CM4 C2-2
CN44	CI		CM4 C3-1
CIVI4	<u></u>		CM4-4
	0	СпзОп	CM4 M4-1
	נח		C3-2C
	DZ		C3-3B
DIVI4	00		DM4-3
	DU	CH3OH	DM4 M2-3

Table 2.4. TAMARA	project	DehaloRNAseq:	sample name	reference

The TAMARA interface gives an access to a large set of tools:

- the sequence data overview (Figure 2.10)
- the raw read count (Figure 2.11)
- a differential analysis (Figure 2.12)

DehaloRNAseg : RNAseg Projects **Mapping Overview Raw Read Count** Analysis IGV В С Base quality along reads (BBC_IOSW_3_1_C2FUBACXX_IND4) C3-2C SSAHA2 version 2.5.1 dir mRNAseq solexa Total read number 43101981 100 % 2156706 5.00 % Nb of unmapped reads 40945275 95.00 % Nb of reads mapped at least once 14962708 34.71 % Nb of reads mapped on rRNA 25367570 58.85 % Nb of reliable reads 24942666 57.87 % Nb of reads kept on chromosome METDI 0.86 % 372629 Nb of reads kept on plasmid p1METDI 0.12 % 52275 Nb of reads kept on plasmid p2METDI Total reads mapped on genomic objects (except rRNA) into 2 4 6 8 10121416182022242628303234363840 2 6466 6870 72 7476 7880 8284 8688 9092 9496 980 23580430 54.71 % Read Position chromosome METDI BBC_IOSW_3_1_C2FUBACXX_IND4 Total reads mapped on genomic objects (except rRNA) into plasmid 203689 0.47 % p1METDI 60 % Total reads mapped on genomic objects (except rRNA) into plasmid 0.11 % 49335 10 % 40 % 70 % 100 % D2METDI - 80 % 20.% 50 % Download forward / reverse wig file Good quality calls SSAHA2 version 2.5.1 dir mRNAseq solexa DM4-3 SSAHA2 version 2.5.1 dir mRNAseq solexa Calls of reasonable quality DM4_M2-3 SSAHA2 version 2.5.1 dir mRNAseq solexa CM4 C2-2 SSAHA2 version 2.5.1 dir mRNAseq solexa Calls of poor quality CM4_C3-1 SSAHA2 version 2.5.1 dir mRNAseq solexa CM4-4 SSAHA2 version 2.5.1 dir mRNAseq solexa 40 46 52 58 82 S CM4_M4-1 SSAHA2 version 2.5.1 dir mRNAseq solexa Read Position

TAMARA - Transcriptome Analyses based on MAssive sequencing of RnAs

Figure 2.10. A screen capture of the "mapping overview" data available in "TAMARA" web platform

The screenshot was taken during analysis of the genome of *M. extorquens* CM4 in the "Transcriptomics" window. (A) "Mapping overview" gives access to information of mapped sequence data overview. (B) Summary table with total number of reads, unmapped reads, mapped reads on *M. extorquens* CM4 chromosome or on its two plasmids. Reads mapped at least once correspond to the number of reads used for the RNAseq analysis. (C) The quality of calls decreases as the run progresses. In the diagram called "Base quality along reads", the purple boxes represent the inter-quartile rage (25-75 %), the upper and lower whiskers represent the 10 % and 90 % points, the red line shows the mean value.



Figure 2.11. A screen capture of the "Raw read count" data available in TAMARA web platform

The "Raw read count" interface gives access to normalized read numbers matched in a given genomic object. Results are accessible following a 5 steps process which is described below. (B) Choose one or several reference sequences. (C) Select at least one experiment and compute the associated read count number per genomic object. (D) It is possible to restrict the query to one or several given classes of genomic objects (CDS, fCDS, rRNA, tRNA, miscRNA or all). (E) Export functions. This section allows users to make all genes (or subsets of genes) available for other analysis tools. (F) Information about genes (label, name, product, begin and end, length and frame). (G) Sense and antisens read numbers.



Figure 2.12. A screen capture of the "Analysis" interface available in TAMARA web platform

The "Analysis" interface give access to the differential gene expressions. (B) Selection of conditions for the differential analysis. (C) FDR (false discovery rate) threshold. Data normalized using (D) 55 reference genes or (E) the complete set of CDS.

Chapitre 3

RNAseq study of chlorinated compound utilization in

M. extorquens

Chapter 3. RNAseq study of chlorinated compound utilization in *M. extorquens*

1. Introduction

Advanced sequencing technologies now provide the opportunity to improve our understanding of the structures and regulatory roles of functional elements involved in adaptation to chloromethane. This chapter describes the first chloromethane transcriptome. To better target the inventory of genes that are specifically expressed in growth with chloromethane, a differential RNAseq approach was performed with two *M. extorquens* reference strains cultivated in methylotrophic conditions. The chloromethane-degrading reference strain *M. extorquens* CM4 was grown with chloromethane versus methanol while the closely-related dichloromethane reference strain *M. extorquens* DM4 was grown with dichloromethane versus methanol. Chapter 3 consists with an introduction, a submitted article of the comparative dehalogenation transcriptomic study, additional data, and the Chapter ends with a brief conclusion.





Numerous families of dehalogenase enzymes are known, totalling more than 200 annotation types in sequence databases (David *et al.*, 2008). Diverse mechanistic types (hydrolytic or thiolytic, dehydrodehalogenative, involving methyl transfer; reductive, oxidative and/or respiratory; see e.g. (Janssen *et al.*, 2005) are known, and many representatives of each mechanistic type have been investigated in detail in the laboratory. Aerobic microbial pathways for utilization of chlorinated methanes have been characterized in the chloromethane and the dichloromethane-degrading strains *M. extorquens* CM4 and DM4, respectively. In *M. extorquens* CM4, two methyltransferases encoded by *cmuA* and *cmuB* are

essential for growth with chloromethane, and require two cofactors, a cobalt-containing corrinoid compound and tetrahydrofolate (H₄F) (Vannelli *et al.*, 1999). In *M. extorquens* DM4, the dichloromethane dehalogenase encoded by the *dcmA* gene is essential for dichloromethane utilization (Vuilleumier *et al.*, 2009b). DcmA belongs to the glutathione-S-transferase family and uses glutathione as a cofactor. For each C₁ dehalogenase, cofactors are highlighted in grey colour. In both cases, the product of aerobic chlorinated methane dehalogenation leads to intracellular HCl production. A major difference between chloromethane and dichloromethane utilization is that the carbon of the chloromethane is funnelled into a specific metabolic pathway lacking formaldehyde (Kohler-Staub and Leisinger, 1985; Studer *et al.*, 2002; Vannelli *et al.*, 1999).

The aim of the intraspecies comparative transcriptomics was to explore RNA regulatory elements with the identification of transcript abundance patterns specific of:

- The chloromethane response that involves a corrinoid compound and tetrahydrofolate (H₄F)-dependent methylotrophic pathway (only detected in the chloromethane-grown CM4 strain). The chloromethane utilization (cmu) pathway is initiated by expression of the chloromethane dehalogenase that differs from the dichloromethane dehalogenase (Figure 3.1). In the chloromethane-degrading strain, some atypical, surnumerary H₄F-dependent enzymes not found in other Methylobacterium strains have been detected (gene metF2; folC2; purU; folD for instance), but their function or regulation has not been elucidated (Roselli et al., 2013). Among those genes, some genes, e.g. metF2 and folC2 may have functional redundancy with the "core genome" gene copy. Moreover, redundancy of genes coding for transport and biosynthesis of the chloromethane dehalogenase corrinoid cofactor have also been detected in strain CM4. Among those, expression of *cob* genes adjacent to *cmu* genes may be required for growth with chloromethane but whether these genes are transcribed has not been previously tested (Roselli et al., 2013). By comparing the chloromethane transcriptome to the methanol transcriptome in strain CM4, we expected to reveal RNA dependent regulatory mechanisms specifically involved in expression of the *cmu* H₄F-dependent methylotrophic pathway, as opposed to genes involved in standard formaldehyde and the tetrahydromethanopterindependent methyltrophic pathways (see Chapter 1, Figure 1.11).
- The halogenated C₁ methylotrophy and dehalogenation-linked adaptation such as intracellular chloride and acid stress (common to the chloromethane-grown CM4 strain and the dichloromethane-grown DM4 strain).

The transcriptomic data are discussed in view of our model for bacterial adaptation to the transformation of chlorinated methanes (Figure 3.2). By comparing transcriptomics of cultures grown with methanol to those grown with chlorinated methanes, we identified core and variable genome genes with transcription patterns modulated by only chloromethane, dichloromethane and methanol, or shared by both chlorinated methanes. To our knowledge, this is the first study that has experimentally identified the carbon source-dependent core genome contribution in methylotrophic conditions of chlorinated compounds in natural isolates of two closely related strains of the species *M. extorquens*. Those results will lead to a better understanding of key enzymes involved in bacterial degradation of chlorinated methanes and the regulation of their expression will provide molecular ecological tools to evaluate biodegradation in ecosystems contaminated by such compounds.



Figure 3.2. Adaptation to growth with chlorinated methanes as the sole carbon and energy source

This figure proposes a model for adaptation to growth with chlorinated methanes and has been adapted from Muller and collaborators (2011). Previous genetic, comparative genomics, proteomics and random mutagenesis studies have demonstrated that cultures grown with these compounds elicit a complex adaptive response encoded by the core genome, which is shared by both dechlorinating and non-dechlorinating *Methylobacterium* strains. It requires the expression of dedicated essential dehalogenase enzymes encoded by strain-specific dehalogenation gene modules. The variable genome includes strain-specific pathways of chloromethane and dichloromethane that are initiated by the expression of dehalogenation modules that respectively encode the *cmu* genes located on the 380 kb pCMU01 plasmid in strain CM4 and the *dcm* genes nested within a genomic island in strain DM4 (Marx *et al*, 2012; Vuilleumier *et al.*, 2009). Both pathways produce intracellular HCI that lowers pH and increases inorganic chloride. Moreover, the microbial response to such pollutants can affect biological function at different levels (e.g. membrane structure, inactivation of key molecules, through genotoxic effects, and may involve metabolic adjustments) (Muller *et al.*, 2011a; Roselli *et al.*, 2013; Vuilleumier, 2002).

<u>2.</u> <u>Article</u>

Contribution of the Core and Variable Genome to the Transcriptomes of two
 Methylobacterium extorquens Strains Grown with Chloromethane, Dichloromethane or
 Methanol

- 4
- 5 Pauline Chaignaud^{1,2}, Marion Weiman³, Adriana Alberti³, Stéphane Vuilleumier¹, Steffen
 6 Kolb⁴,² Stéphane Cruveiller³, Françoise Bringel^{1*}
- 7
- 8 Affiliations:
- 9¹, CNRS Molecular Genetics, Genomics, Microbiology, Université de Strasbourg UMR 7156
- 10 UNISTRA CNRS, Strasbourg, France
- 11 ², Department of Ecological Microbiology, University of Bayreuth, Bayreuth, Germany
- 12 ³, CNRS-UMR 8030 and Commissariat à l'Energie Atomique et aux Energies Alternatives
- 13 CEA/DSV/IG/Genoscope LABGeM, Evry, France.
- 14 ⁴, Institute for Landscape Biogeochemistry Leibniz Centre for Agricultural Landscape
- 15 Research ZALF, Müncheberg, Germany
- 16 *Corresponding author: <u>francoise.bringel@unistra.fr</u>; Phone: + 33 (0)3 68 85 18 15; Fax: + 33
- 17 (0)3 68 85 20 28
- 18 Address: 28 rue Goethe, 67083 Strasbourg CEDEX, France
- 19
- 20 Running title: Transcriptomics of chlorinated C₁ compound utilization
- 21

22 ABSTRACT. The transcriptomes for the utilization of methanol, chloromethane or 23 dichloromethane as sole carbon and energy source were compared in two closely related 24 Methylobacterium extorquens strains. Both strains grow with methanol, but strain CM4 grows 25 on chloromethane while strain DM4 grows on dichloromethane. Both chlorinated methane 26 utilization pathways produce intracellular HCl that lowers pH and increases inorganic chloride. 27 Only 11 out of 4,620 common genes had higher transcript abundance in cultures grown with 28 chlorinated methanes compared to methanol. These genes included stress response and 29 related bioenergetics genes (*depP*, *hppA*). On the other hand, the variable genome (25 %) 30 which includes the dehalogenation modules that encode the *cmu* genes in strain CM4 and the 31 dcm genes in strain DM4 harbored the majority of the genes whose transcription was 32 enhanced by chloromethane on plasmid pCMU01 such as the <u>cmu</u> and coenzyme-associated 33 *cmu*-pathway genes for vitamin B₁₂ and for H₄F. In the variable genome of strain DM4, the 34 dcm genes nested within a genomic island (GEI) were enhanced in cultures grown with 35 dichloromethane, whereas central metabolism genes (ackA, adh-like, xfp, cytochromeassociated genes) located on redundant GEIs had increased transcript abundance in cultures 36 37 grown with methanol. A major finding is that core and variable genome-encoded genes 38 involved in redox status adjustments were identified for growth with different C₁ compounds. 39 Transcription responses to growth with chloromethane and dichloromethane were 40 predominantly uncorrelated.

41

42 **IMPORTANCE**. Methylotrophic microorganisms often differ in their ability to use one-carbon 43 compounds due to differential possession of gene modules. To grow with chlorinated onecarbon compounds, a methylotrophic microorganism must acquire a one-carbon 44 45 dehalogenation module that integrates the existing methylotrophic metabolic and regulatory 46 networks. This prompted us to investigate transcriptional adjustments during growth with 47 different one-carbon growth substrates in the genomic background of Methylobacterium 48 extorquens, the best-characterized microbial model for the study of aerobic methylotophy. M. 49 extorquens has a broad ability to utilize one-carbon compounds, including the reference C1 50 compound methanol. Using intraspecies comparative transcriptomics, we identified key 51 functions encoded by the core genome or by genomic islands that are necessary for growth with different C₁ compounds, toxic natural (chloromethane) and industrial (dichloromethane) 52 53 chlorinated compounds.

134

54 INTRODUCTION

One-carbon compounds such as methanol (CH₃OH), chloromethane (CM, CH₃Cl) and 55 dichloromethane (DCM, CH₂Cl₂) are found in both pristine and polluted environments. In 56 57 plants, methanol and chloromethane are both produced, although chloromethane emissions are at least twelve orders of magnitude lower than that of methanol (1,2). Chloromethane is 58 59 released from terrestrial environments as a gas subsequently found in the atmosphere, where it is the most important halogenated trace gas contributing to ozone destruction (2). 60 61 Dichloromethane, the most produced industrial worldwide-chlorinated solvent, is commonly 62 detected in polluted sites. A few methylotrophic microorganisms, including bacteria and yeast, are able to utilize chlorinated methanes (such as chloromethane and dichloromethane) 63 as their sole carbon and energy source. Methanol is utilized by a majority of methylotrophs, 64 65 while only a restricted number of methylotrophs are able to degrade chloromethane or dichloromethane (3). Thus, methanol is the reference reduced one-carbon compound. A long 66 67 history of biochemical and genetic utilization studies has set basis for genome-scale reconstruction of metabolic networks in methylotrophic bacteria (4,5). The genomes of 68 69 methylotrophic bacteria contain a wide variety of gene-encoding methylotrophic metabolic 70 pathway modules. These modules are unique, redundant, and sometimes interchangeable (6). 71 Such methylotrophic modules were divided into three groups: for primary one-carbon 72 oxidation, formaldehyde dissimilation, and assimilation pathways. In particular, 73 dehalogenation modules initiate halogenated methane catabolism (Fig. 1, green box for 74 chloromethane; purple box for dichloromethane).

75

76 FIG 1 Metabolism of chlorinated C₁ compounds chloromethane and dichloromethane and 77 of methanol in *M. extorquens* CM4 and DM4. Chloromethane dehalogenation and methyl 78 transfer to tetrahydrofolate (H_4F) metabolic steps are boxed in green. Dichloromethane 79 dehalogenation and methanol oxidation into formaldehyde are boxed in purple and blue, 80 respectively. Black boxes delimit common methylotrophic oxidation modules: H₄F and 81 tetrahydromethanopterin (H₄MPT)-dependent C₁ transfer reactions, and formate oxidation 82 for C₁ dissimulation. The serine-glyoxylate cycle for C₁ assimilation are schematized with two 83 connected circles. Reactions related to essential co-factors for chlorinated methane utilization 84 are highlighted with colored spots for H₄F (yellow), corrinoid cofactor (green), glutathione 85 (purple) and pyrroquinoline quinone (blue). Genes located on genomic islands (GEI) in 86 M. extorquens DM4 or on the plasmid pCMU01 in M. extorquens CM4 are shown in grey-87 shaded boxes. RNAseq data are summarized using color-coding for gene names when 88 transcripts were detected as more abundant on chloromethane (green), dichloromethane 89 (purple), both chloromethane and dichloromethane (dark red), or methanol (both 90 *M. extorquens* CM4 and DM4 (dark blue), or methanol in *M. extorquens* DM4 (sky blue).
105

91 Compounds involved in energy balance reactions are highlighted with names written in orange 92 when produced and red when consumed. The proposed figure is based on previous studies in 93 Methylobacterium or in other Proteobacteria: cytochromes bd quinol oxidase, cytochrome d 94 terminal oxidase, methanol dehydrogenase (Mxa, Mdh) and formate dehydrogenase (Fdh) 95 complexes (4,7); acetone carboxylase complex (8), the membrane-bound transhydrogenase 96 encoded by the *pnt* complex (9); the membrane-bound proton translocating pyrophosphatase 97 transporter HppA (10); the ClcA H⁺/Cl⁻ antiporter (11); the DctA multi-carbon compound/ H⁺ 98 symport-transporter (12,13); phbC encoding a poly-(R)-3-hydroxybutyrate (PHB) synthase 99 (14); and *phaABC* encoding a β-ketothiolase, an acetoacetyl-CoA reductase and poly-β-100 hydroxybutyrate synthase respectively, involved in PHB biosynthesis (15). PHB metabolism 101 has previously been shown to be linked to acetone production (16), to beta-oxidation by 3-102 hydroxy-butyryl CoA (17) and patatin-like phospholipase (18). Cbi, cobilamide; Cbl, cobalamin; 103 CoA, co-enzyme A; PEP, phosphoenol pyruvate; OXA, oxaloacetate; TCA, tricarboxylic acid 104 cycle.

106 The dynamics of acquisition, loss, cohabitation and regulation patterns of metabolic pathway-107 encoding modules are driven by the specific habitat's physico-chemical parameters and 108 genetic repertoire differences. To grow with "alternative" one-carbon compounds, a 109 methylotrophic microorganism must acquire a primary one-carbon oxidation or 110 dehalogenation module that integrates the existing methylotrophic metabolic and regulatory 111 networks. Thus, these bacteria need to adjust their overall genome expression during 112 "alternative" chlorinated compounds utilization driven by the expression of variable genome modules and fine-tuning of functional methylotrophic modules. 113

114 While methanol can be assimilated by most methylotrophs, growth with "alternative" one-115 carbon sources such as chlorinated methanes can be quite challenging (11,19). 116 Chloromethane and dichloromethane utilization starts with a dehalogenation step (Fig. 1), 117 which induces physiological stress due to the production of intracellular HCl, lower pH and 118 higher concentration of inorganic chloride, as well as formation of DNA adducts (19,20). 119 Growth with dichloromethane in experimental evolution studies showed that the horizontal 120 transfer of dehalogenase-encoding modules for dichloromethane dechlorination in naïve 121 methylotrophic bacteria led to post-transfer refinement (11). In addition, previous studies of 122 genome-wide random mutagenesis, differential proteomics studies have shown that 123 adaptation to growth with chlorinated methanes requires genes located on genomic islands 124 (GEI) or plasmids, as well as core genome-encoded housekeeping functions involved in stress 125 response and central metabolic tuning (11,21,22).

126 The most extensively studied methylotroph is the alpha-proteobacteria *M. extorquens*, a 127 species found in a wide variety of habitats including plants, soil, clouds, and wastewater 128 (23,24–27). The genome of five *M. extorquens* strains (AM1, BJ001, CM4, DM4 and PA1) has 129 been completely sequenced (28,29). Isolates of *M. extorquens* are able to grow with 130 methanol, but only *M. extorquens* strains CM4 and DM4 are able to utilize chlorinated 131 compounds as their sole source of carbon and energy. M. extorquens CM4, isolated from 132 petrochemical factory soil in Russia, grows on chloromethane (30) while M. extorquens DM4 133 isolated from contaminated industrial waste grows on dichloromethane (31). M. extorquens 134 genomes are characterized by a high GC content (68 %), large chromosome size (between 5.5 135 and 5.9 Mb) with plasmid presence, except for strain PA1 (23). In *M. extorquens* CM4, plasmid 136 pCMU01 (380 kb long) harbors two chloromethane dehalogenation modules that initiate the 137 chloromethane-utilization pathway (cmu) and require expression of the essential genes cmuA 138 and cmuB (22,32). The methyl group of chloromethane is transferred to the corrinoid 139 methyltransferase protein CmuA encoded by the *cmuA* gene and then to tetrahydrofolate 140 (H₄F) catalyzed by the methyltransferase CmuB (32). For each mole of chloromethane, one 141 mole of HCl and one of methyl-H₄F are produced. CmuAB protein activity requires two 142 essential cofactors, a corrinoid cofactor and H₄F. Thus, adjacent to the *cmuAB* genes it was 143 not surprising to find the corrinoid compound vitamin B₁₂-associated aerobic biosynthesis and 144 transport, and H₄F-dependent metabolism plasmid genes in addition to chromosomal copies 145 conserved in *M. extorquens* genomes (22,33). In *M. extorquens* DM4, the dichloromethane 146 utilization dcm genes are part of a genomic island of 126 kb located on the chromosome 147 where dcmA encodes a dichloromethane dehalogenase DcmA of the gluthathione S-148 transferase family (22,29). The dichloromethane dehalogenase, converts dichloromethane 149 into inorganic chloride and S-chloromethyl glutathion, which is genotoxic, unstable and 150 decomposes spontaneously into glutathione, inorganic chloride and formaldehyde (HCHO) 151 (19,34). For each mole of dichloromethane, two moles of HCl and one mole of formaldehyde 152 are produced.

The *cmu* and *dcm* genes are the only known chlorinated methane degradation modules and have best been characterized in the *M. extorquens* strains CM4 and DM4, respectively (35,36). Several methanol primary oxidation modules with either *mxaFl, xoxF* or *mdh2* genes have been identified and may coexist simultaneously in genomes of methylotrophs (37,38). The first one discovered is the *mxa* gene cluster in *M. extorquens* AM1 containing the structural *mxaFl* genes coding for the methanol dehydrogenase (MDH) subunits and its associated cyctochrome c accepting the electrons (7). While gene *mxaF* has been proposed as a gene 160 marker for detection of methanol oxidation in the environment (39), *xoxF*-related genes are 161 more widespread in metagenomic datasets of environmental samples. XoxF-related genes 162 encode a MDH (methanol dehydrogenase) which shares approximately 50 % amino acid 163 identity (aa Id) with MxaF, while the *mdh2* gene encodes a single-subunit MDH that belongs 164 to a class of pyrroloquinoline quinone (PQQ)-linked dehydrogenases (38,40). For each mole of 165 methanol, one mole of formaldehyde (HCHO) is produced by these dehydrogenases.

166 The major metabolic difference between chloromethane and dichloromethane utilization is 167 the carbon entry in the methylotrophic metabolism and the link to formate production (Fig. 1). 168 Chloromethane requires a reversal of the assimilatory H₄F pathway since methyl-H₄F must be 169 oxidized into formate to yield energy. The formate produced via the H₄F-depending pathway 170 essential for growth with chloromethane requires the bifunctional enzyme FoID (with 171 methylene-H₄F dehydrogenase and a methenyl-H₄F cyclo-hydrolase activities) associated with 172 *metF2* (coding a methyl-H₄F reductase) and *purU* (coding a formyl-H₄F hydrolase) (41). These 173 enzymes are part of the *cmu* pathway first characterized in *M. extorquens* CM4 (41). In other 174 *M. extorquens* strains and most methylotrophs, formaldehyde is the product of one-carbon 175 compound primary oxidization when grown with methanol (42). M. extorquens DM4 grown 176 with dichloromethane forms formaldehyde that is oxidized into formate via the 177 tetrahydrometanopterine (H₄MPT) pathway. To produce biomass via the serine cycle, formate 178 is reduced into methylene-tetrahydrofolate ($CH_2=H_4F$) (43) while to produce energy, formate 179 is oxidized to CO₂ by formate dehydrogenase (FDH). *M. extorquens* core genomes harbor four 180 characterized FDH (44,45).

181 In this study, we searched for phenotypic expression profiles of common or specific genes by 182 comparative transcriptomics of phylogenetically very-closely related strains after growth 183 under methylotrophic conditions with different one-carbon compounds. M. extorquens 184 strains CM4 and DM4, isolated from soil on the basis of their ability to grow with 185 chloromethane and dichloromethane respectively, acquired different chlorinated one-carbon 186 dehalogenation modules (22,29,46). In this study, the core and variable genome content was 187 analyzed using intraspecies comparative genomics of 5 sequenced genomes of *M. extorquens* 188 strains with respect to their capacity to degrade chlorinated methanes. By comparing 189 transcriptomics of cultures grown with methanol to those grown with chlorinated methanes, 190 we identified core and variable genome genes with transcription patterns modulated by only 191 methanol, chloromethane, dichloromethane or shared by both chlorinated methanes. To our 192 knowledge, this is the first study that has experimentally identified the carbon source-193 dependent core genome contribution in methylotrophic conditions of chlorinated compounds

194 in natural isolates of two closely related strains of the species *M. extorquens*.

RESULTS 195

196 Deep sequencing was performed on cDNA prepared from samples of partially rRNA-depleted 197 RNA extracts from methyltrophically-grown cultures with different one-carbon compounds 198 (Table 1). Four transcriptomes were obtained, two for each of the two *M. extorquens* strains 199 tested. For M. extorquens CM4, chloromethane or methanol was used as the sole source of 200 carbon and energy. For *M. extorquens*, DM4, dichloromethane or methanol was used. Both 201 *M. extorquens* CM4 and DM4 strains grew with a similar generation time (3.2 h) on methanol, 202 which served as the reference growth condition. Longer generation times were needed for 203 chloromethane (5.4 h) and even longer for dichloromethane (9.0 h), in agreement with earlier studies (21,22). Criteria for gene expression profiling included differential expression profiles 204 calculated as log2 fold-change (log2fc) of normalized read numbers in cultures grown with 205 206 either one of the two chlorinated methanes than with methanol, the reference non-207 chlorinated one carbon compound.

208

209 TABLE 1 Overview of *M. extorquens* CM4 and DM4 growth, genomic properties and 210 transcriptomic data

Strain	Genomeª	Carbon source (generation time in h)⁵	Total reads ^c	Mapped reads (%)	rRNA ^d (%)
		methanol	30,337,270	98.6	18.3
CM4	chromosome 5.8 Mb, GC % = 68.2 pCMU01: 380.2 kb, GC % = 66.3 p2MCHL: 22.6 kb, GC % = 63.9	(3.0 ± 0.2)	37,414,003	98.2	14.3
		chloromethane	53,790,411	96.3	18.6
		(5.4 ± 0.4)	36,665,352	97.5	25.4
		methanol	48,154,448	99.1	12.3
	chromosome 5.9 Mb, GC % = 68.1 p1METDI 141.5 kb, GC % = 65.3 p2METDI: 38.6 kb, GC % = 63.7	(3.4 ± 0.4)	38,757,418	97.9	23.0
UM4		dichloromethane	43,101,981	95.0	34.7
		(9.0 ± 0.7)	32,066,920	95.0	20.0

211 212 ^a GenBank accession n° for CM4, CP001298, CP001299, CP001300; for DM4, FP103042, FP103043, FP103044

^b Aerobic growth in M3 media in presence of 10 mM one-carbon compound provided as the sole source of carbon and energy

213 ^c High-throughput sequencing with HiSeq2000 from Illumina. Coverage 600X

- 214 ^d Percentage of total read number (mapped and unmapped)
- 215

216 Extensive gene synteny between strains CM4 and DM4. M. extorquens CM4 and DM4 217 genomes share a large majority of their genes (75 %, 4,620 CDS) when defining common genes 218 as encoding proteins with at least 80 % amino acid sequence identity on 80 % of their length. 219 Their gene organization along the genome is conserved overall in large synteny segments 220 (Fig. 2).

221

FIG 2 Overview of genomic islands and gene synteny in genomes of *M. extorquens* CM4 and DM4. Strand conservation (in purple) and strand inversion (in blue) lines representing a conserved group of at least 6 genes were drawn using the Conserved Synteny LinePlot on the MaGe plateform. One-carbon primary oxidation genomic islands (GEI) are described in Table 4: for dichloromethane (*dcm* genes on 126 kb genomic island (29), for methylamine (*mau* genes on 70 kb genomic island (29), and for chloromethane (*cmu* genes on the 380 kb plasmid pCMU01 (22), as well as other GEIs.

229

230 The genomes of three non-dechlorinating strains (AM1, BJ001, and PA1) affiliated to *M. extorquens* were then compared to those of the chloromethane-degrading strain CM4 and 231 232 the dichloromethane-degrading strain DM4 (28,29). The assembly of the gene content found 233 in five *M. extorquens* genomes resulted in a pan genome of 12,273 unique CDS for a total of 234 12.3 Mb. The core genome included 3,489-shared CDS (Fig. 3). The variable genome specific 235 to strain CM4 and potentially specific to growth with chloromethane included 1,512 CDS. The 236 variable genome specific to strain DM4 and potentially specific to growth with 237 dichloromethane was smaller (952 CDS). The variable genome shared only by dechlorinating 238 strains and potentially specific to growth with chlorinated methanes included 163 CDS. Of 239 those, 85 genes were detected on two plasmids (pCMU01 in CM4; p1METDI in DM4) with no 240 predicted function besides plasmid-related functions.

241

FIG 3 Core, variable and specific genome of five strains affiliated to the species
 Methylobacterium extorquens. Comparative analyses were performed using the fully
 sequenced genomes of *M. extorquens* strain AM1 (GenBank accession n°CP001510,
 CP001511, CP001512, CP

001513 and CP001514), strain BJ001 (CP001029, CP001030 and CP001031), strain CM4
(CP001298, CP001299 and CP001300), strain DM4 (FP103042, FP103043 and FP103044) and
strain PA1 (CP000908) (28,29). The pan genome consists 12,274 genes. Homologous genes
encode for proteins with at least 80 % of amino acid identity on 80 % of the CDS length.
Genome size of the common (core), variable (found in two up to four genomes) and specific
(only found in one strain) are indicated in the Venn diagram obtained with the MaGe platform
interface pan/core-genome.

253

254 Methanol-enhanced expression of a core genome C₁/C₂ dehydrogenase *mdh2*-associated 255 gene module. As expected, the overall transcription patterns are conserved in the two strain 256 cultures grown with methanol as the sole carbon and energy source (Fig. S1A). Among those 257 genes with unchanged transcript abundance patterns, we found genes encoding components 258 of the well-characterized pyrroloquinoline quinone (PQQ)–dependent methanol259 dehydrogenase MxaFI (27) (Fig. S2). A minority of the core genome-encoded genes showed significantly higher transcript abundance in cultures grown with methanol compared to 260 261 chlorinated methanes (9 genes group E in Table S2; $\log_2 fc \leq -2.1$, p-value <0.029). One of 262 these, a highly transcribed gene, encodes a C₄-dicarboxylate transport (Dct) protein that displays 77 % aa identity with the characterized Rhizobium DctA transporter (47). Although 263 264 Dct protein transport specificity has not been studied in *M. extorquens*, in other aerobic 265 bacteria, Dct transports succinate, malate, fumarate, aspartate and a pyrimidine precursor 266 orotate, from the periplasm across the inner membrane (12). We also found that a large 267 majority of the methanol-enhanced transcript abundant genes (7 out of 9) were part of a 268 single primary C₁/C₂ carbon compound oxidizing cluster. This cluster harbors the highly 269 transcribed PQQ-dependent methanol/ethanol dehydrogenase-encoding gene mdh2 which is 270 involved in methanol utilization in other methylotrophs (40).

271

272 Chlorinated methane-enhanced expression of core genome genes were involved in stress 273 response and bioenergetics. Utilization of either chloromethane or dichloromethane involves 274 exposure to intracellular acidification and production of chloride, so we decided to focus on 275 identifying a common adaptive response to the use of chlorinated methanes versus methanol. 276 To do so, we searched for core genome-encoded genes with similar transcription patterns in 277 cultures of strains CM4 and DM4. We found that 11 genes, representing 0.02 % of 278 *M. extorquens* genome, had higher transcript abundance in cultures grown with chlorinated 279 methanes compared to methanol (Fig. 4; Table S2 group B; log2fc from 2.0 up to 6.6).

280

281 FIG 4 Carbon source-dependent shared categories of gene transcript abundance between 282 M. extorquens CM4 and DM4 genomes. The Venn diagram includes a total of 4,620 genes 283 encoding CDS with at least 80 % aa identity on 80 % of the CDS length detected in both M. 284 extorquens CM4 and DM4 genomes. Categories were defined on the basis of the log2 fold-285 change (log2fc) values of RNAseq reads for *M. extorquens* CM4 grown with chloromethane 286 versus methanol, and for *M. extorquens* DM4 grown with dichloromethane versus methanol 287 as follows: (A) constitutive expression (log2fc between 2 and -2); higher transcript abundance 288 $(\log_2 fc > 2)$ in dichloromethane and chloromethane (B), in chloromethane only (C), in 289 dichloromethane only (D); higher transcript abundance (log2fc < -2) in methanol (F), in strain 290 CM4 only (except for the gene Mchl_1717/METDI2103 coding for a conserved protein of 291 unknown function); and in strain DM4 only (G). Details found in Table S2.

292 Two genes had predicted functions. First, gene *depP* encodes a putative periplasmic serine

293 protease (log2fc value of 6.6). Its corresponding *Escherichia coli* homolog HtrA (39 % aa Id) is

a central housekeeping molecular chaperon protein in the bacterial envelope, which under

295 stress including membrane damage, controls the folded state of proteins and the biosynthesis 296 of partially folded outer-membrane proteins (48). Second, gene hppA encodes a membrane-297 bound proton translocating pyrophosphatase. HppA in Carboxydothermus hydrogenoformans 298 (48 % aa Id) uses the energy of pyrophosphate hydrolysis as the driving force for proton 299 extrusion to build up a proton motive force (49). In Rhodospirullum rubrum (72 % aa Id), this 300 proton pump is involved in stress bioenergenetics (10). Membrane-bound proton 301 translocating pyrophosphatases confer resistance to salt stress (50). Thus, hppA may also play 302 a role in the adaptive response and bioenergetics of chloromethane and dichloromethane 303 aerobic utilization in *M. extorquens*. The stress-related core genome genes *depP* and *hppA* are 304 located on different genomic regions.

305

306 The chloromethane transcriptome. Among 137 genes detected as more abundant with 307 chloromethane than methanol, 40 genes have been previously associated with growth with 308 chloromethane (22) ($\log 2fc \ge 2$; Tables 2 and S3). These genes, classified in the chloromethane 309 utilization group, showed the highest chloromethane-dependent regulation levels (log2fc 310 values 7.7 to 10). These included essential genes for growth with chloromethane (cmuABC, 311 metF2, purU), genes previously found to be part of the cmu pathway (folD) (41), genes 312 systematically associated with *cmu* gene clusters in various chloromethane-degrading strains (fmdB, hutI, paaE-like) (22,51), and corrinoid cofactor riboswitch regulatory elements (52,53) 313 314 (Fig. 1; misc _RNA_1 see Table 2).

315

316	TABLE 2 M. extorquens CM4 transcripts differentially abundant in cultures grown with
317	chloromethane compared to methanol

				Occurr	RNAseq data		
	Label ^a	Name	Product	ence ^b	log2fc ^c	Adjusted p-value ^d	
Chlor	omethane utilization						
	Mchl_5694	hutl	putative imidazolonepropionase	pCMU01	7.7	< 0.001	
	Mchl_5696	-	putative regulatory protein FmdB	pCMU01	7.7	< 0.001	
	Mchl_5697	стиА	methyltransferase/corrinoid binding protein CmuA	pCMU01 ^f	8.3	< 0.001	
	Mchl_5698	cmuC2	putative methyltransferase, CmuC-like protein	pCMU01	8.4	< 0.001	
	Mchl_5699	purU	formyltetrahydrofolate hydrolase	pCMU01 ^f	10.0	< 0.001	
_	Mchl_5700	folD	bifunctional methylenetetrahydrofolate dehydrogenase/methenyltetrahydrofolate cyclohydrolase	pCMU01	8.6	< 0.001	
	Mchl_5717	-	putative oxidoreductase FAD/NAD(P)-binding domain protein, PaaE-like	pCMU01 ^e	8.1	< 0.001	
	Mchl_5726	metF2	methylenetetrahydrofolate reductase	pCMU01	9.6	< 0.001	
_	Mchl_5727	стиВ	methylcobalamin:H4folate methyltransferase CmuB	pCMU01 ^e	9.5	< 0.001	
	Mchl_5728	стиС	putative methyltransferase CmuC	pCMU01	9.1	< 0.001	

Coba	lamin and precursor	metabolisn	n and transport			
_	Mchl_1718	-	putative cobalt transporter, subunit CbtA	core	2.5	< 0.001
_	Mchl_1719	-	putative cobalt transporter subunit CbtB	core	2.7	< 0.001
-	Mchl_2855	-	putative TonB-dependent siderophore receptor	core	3.4	< 0.001
-	Mchl_5676	-	putative cobalamin outer membrane transporter (BtuB)	pCMU01 ^e	6.7	< 0.001
-	Mchl_5677	-	ABC transporter periplasmic binding component, putative vitamin B12 transporter (BtuF)	pCMU01 ^e	6.0	< 0.001
-	Mchl_5678	-	ABC transporter membrane component, putative vitamin B12 transporter subunit (BtuC)	pCMU01 ^e	5.3	< 0.001
_	Mchl_5679	-	ABC transporter ATP-binding component, putative vitamin B12 transport (BtuD)	pCMU01 ^e	5.0	< 0.001
-	Mchl_5681	-	putative P-loop containing nucleoside triphosphate hydrolase	pCMU01 ^e	4.8	< 0.001
-	Mchl_5682	-	putative TonB-dependent receptor	pCMU01 ^e	4.4	< 0.001
-	Mchl_5685	cobM2	precorrin-4 C(11)-methyltransferase	pCMU01 ^e	7.0	< 0.001
-	Mchl_5686	-	putative cobalamin biosynthesis protein CobE	pCMU01 ^e	4.8	< 0.001
-	Mchl_5687	cobL2	precorrin-6Y C(5,15)-methyltransferase (decarboxylating)	pCMU01 ^e	4.7	< 0.001
-	Mchl_5689	cobJ2	precorrin-3B C(17)-methyltransferase	pCMU01 ^e	5.5	< 0.001
-	Mchl_5690	cobl2	precorrin-2 C(20)-methyltransferase	pCMU01 ^e	5.7	< 0.001
-	Mchl_5691	cobH2	precorrin-8X methylmutase	pCMU01 ^{e,f}	5.9	< 0.001
-	Mchl_5692	-	putative cobalamin biosynthesis protein, cobaltochelatase subunit CobN-like domain	pCMU01 ^e	7.0	< 0.001
_	Mchl_5693	-	conserved exported protein of unknown function, CoxB-related protein	pCMU01 ^e	8.9	< 0.001
_	Mchl_5702	cobU2	nicotinate-nucleotide-dimethylbenzimidazole phosphoribosyltransferase	pCMU01	6.8	< 0.001
_	Mchl_5714	czcB	RND efflux transporter, putative membrane fusion protein	pCMU01 ^e	2.5	< 0.001
_	Mchl_5715	czcA2	RND divalent metal cation efflux transporter membrane component, cobalt-zinc-cadmium resistance protein	pCMU01 ^e	2.3	< 0.001
_	Mchl_5721	cobP2	bifunctional adenosylcobinamide kinase and adenosylcobinamide-phosphate guanylyltransferase	pCMU01	6.7	< 0.001
_	Mchl_5722	cobO2	cob(I)yrinic acid a,c-diamide adenosyltransferase	pCMU01	6.5	< 0.001
_	Mchl_5723	cobQ2	cobyric acid synthase	pCMU01	6.2	< 0.001
_	Mchl_5724	cobD2	cobalamin biosynthesis protein	pCMU01	2.3	< 0.001
_	Mchl_5729	cbiD	cobalamin biosynthesis protein, putative cobalt- precorrin-6A synthase	pCMU01	4.3	< 0.001
_	Mchl_5730	cobC2	L-threonine-O-3-phosphate decarboxylase domain	pCMU01	4.9	< 0.001
_	Mchl_5731	cobA	uroporphyrinogen-III C-methyltransferase	pCMU01	5.6	< 0.001
_	Mchl_5732	bluB2	5,6-dimethylbenzimidazole synthase (flavin destructase), putative cob(II)yrinic acid a,c-diamide reductase	pCMU01	5.2	< 0.001
_	misc_RNA_1	-	cobalamin riboswitch upstream of <i>btuB</i> (Mchl_5676)	pCMU01	5.9	< 0.001
	Rmisc_RNA_4	-	cobalamin riboswitch upstream of the uncharacterized gene (Mchl_1720)	core	3.1	< 0.001
Tet	rahydrofolate metab	olism				
_	Mchl_0812	gcvP	glycine cleavage complex protein P, PLP-dependent glycine dehydrogenase	core	4.6	< 0.001
_	Mchl_0813	gcvH	glycine cleavage complex protein H	core	4.5	< 0.001
_	Mchl_0814	gcvT	glycine cleavage complex protein T, aminomethyltransferase tetrahydrofolate-dependent	core ^f	3.0	< 0.001
_	Mchl_5701	folC2	putative folylpolyglutamate synthase and dihydrofolate synthase	pCMU01	7.8	< 0.001
	Rmisc_RNA_14 Rmisc_RNA_15	-	glycine riboswitch upstream of gcvT	core	2.2	< 0.001
C ₁ r	netabolism	1				
_	Mchl_1538	-	c-cytochrome c550 protein, putative PQQ-dependent methanol/ethanol oxidation system	core ^e	-3.9	< 0.001
_	Mchl_1540	mdh2	PQQ-dependent methanol/ethanol dehydrogenase	core ^e	-3.4	< 0.001
	Mchl_5683	gck2	glycerate kinase	pCMU01	5.1	< 0.001

MCv2_5148	pqqA	coenzyme pyrroloquinoline quinone (PQQ) biosynthesis protein A	core	2.2	< 0.001
MCv2_5149	pqqA	coenzyme PQQ biosynthesis protein A	core	2.1	< 0.001
entral metabolism					
Mchl_5057	-	putative seryl-tRNA synthetase	variable	2.2	< 0.001
Mchl_5058	acd	acyl-CoA dehydrogenase	variable	2.2	< 0.001
Mchl_5060	-	putative monooxygenase, small component, putative flavin: NADH reductase	variable	2.5	< 0.001
Mchl_5066	-	putative homoserine O-succinyltransferase (MetA)	variable	2.0	< 0.001
Mchl_5154	-	aldo/keto reductase family	core	2.1	< 0.001
Mchl_5521	acxC	acetone carboxylase gamma subunit	pCMU01	2.5	< 0.001
Mchl_5522	асхВ	acetone carboxylase alpha subunit	pCMU01	2.1	< 0.001
Mchl_5523	acxA	acetone carboxylase beta subunit, putative N-methylhydantoinase A	pCMU01	3.7	< 0.001
ergy					
Mchl_2986	pntAA	NAD(P) ⁺ transhydrogenase, subunit alpha part 1	core ^e	2.7	< 0.001
Mchl_2987	pntAB	NAD(P) ⁺ transhydrogenase, subunit alpha part 2	core	3.6	< 0.001
Mchl_2988	pntB	NAD(P) ⁺ transhydrogenase, subunit beta	core	3.4	< 0.001
Mchl_3408	hppa	H(*) translocating pyrophosphate synthase	core	2.4	< 0.001
aptation to stress and	l transport				
Mchl_2781	-	putative L,D-transpeptidase catalytic domain (YkuD)	core	2.5	< 0.001
Mchl_3002	-	putative manganese catalase	core ^f	2.1	< 0.001
Mchl_3283	dctA	C ₄ -dicarboxylate transport protein	core	-2.9	< 0.001
Mchl_4687	ada	fused DNA-binding transcriptional dual regulator, O6-methylguanine-DNA methyltransferase	core	5.6	< 0.001
Mchl_5016	ibpA	small heat shock protein	core	2.8	< 0.001
Mchl_5157	depP	periplasmic serine protease (DegP)	core	4.6	< 0.001
Mchl_5405	clcA2	putative H(*)/Cl(`) exchange transporter	pCMU01	5.5	< 0.001
Mchl_5500	-	transporter, major facilitator superfamily	pCMU01 ^e	4.4	< 0.001
Mchl_5718	-	putative membrane protein, putative transport protein	pCMU01	7.1	< 0.001
Mchl_5719	-	putative membrane protein, putative transport protein	pCMU01	7.7	< 0.001

318

319 ^b Genes labeled as core or variable are respectively found in all or at least 2 genomes of *M. extorquens* strains AM1, BJ001, CM4, DM4 and 320 PA1 (80 % of identity on at least 80 % of the protein length). Genes labeled as pCMU01 are located on the 380 kb plasmid and some have 321 chromosomal gene copies described previously (22).

322 ^c log2 fold-change of normalized read numbers in cultures grown with chloromethane compared to methanol.

323 ^d False discovery rate, only adjusted p-value under 0.1 were considered.

324 ^e Part of gene clusters: Mchl 1533-1540; Mchl 5676-5693 gene; Mchl 5714-5716; Mchl 5499-5500..

325 ^f Corresponding encoded protein found more abundant in cell protein extracts of chloromethane- versus methanol-grown cultures (22) 326

327 In addition, 30 genes were involved in the metabolism and transport of the corrinoid cofactor, 328 which is essential for CmuA dehalogenase and chloromethane utilization (32) (Table 2). Two 329 transport systems related to corrinoid cofactors were identified: cbtAB genes encoding a 330 putative cobalt transporter Cbt complex (log2fc around 2.6), and btu genes encoding the 331 corrinoid transporter Btu complex formed by 4 proteins (BtuB, BtuC, BtuD, BtuF) associated 332 with TonB (54) with corresponding genes all highly transcribed. A corrinoid cofactor riboswitch 333 (53) was also found with higher transcript abundance in cultures grown with chloromethane 334 compared to methanol upstream of gene *btuB* (misc_RNA_1, Table 2) (log2fc value of 5.9). 335 The gene *bluB2* had a high log2fc value of 5.2. It encodes for a protein highly similar to an 336 enzyme that cleaves reduced flavin mononucleotide (FMNH₂) in an oxygen-dependent 337 reaction to form erythrose 4-phosphate (E₄P) and dimethylbenzimidazole (DMB) (55). These 338 products of the enzymatic reaction catalyzed by BluB2 are precursors of the corrinoid and H₄F 339 cofactor biosynthesis, which are both essential cofactors for the *cmu* pathway (32). In the 340 metabolism of the coenzyme H₄F, gene *folC2* encodes a predicted bifunctional dihydrofolate 341 synthetase/H₄F synthase. Gene *folC2* displayed a tightly regulated chloromethane-dependent 342 transcript abundance (high log2fc value of 7.8). The gcvPHT gene cluster, also involved in 343 coenzyme H₄F metabolite interconversion, encodes the glycine cleavage complex which 344 catalyzes the H₄F transformation to methylene-H₄F, and its upstream predicted glycine 345 riboswitchs (53,56). The gcvPHT genes had transcripts with higher abundance with 346 chloromethane than methanol (Fig. 4, group C; Table 2).

347 Other genes not previously linked to growth on chloromethane had higher transcript 348 abundance in cultures grown with chloromethane compared to methanol. These included (i) 349 the *pnt* gene cluster encoding the transhydrogenase PntAA, PntAB and PntB subunits needed 350 in the transfer of reducing equivalent between NAD(H) and NADP(H) to the translocation of 351 protons across the membrane, as demonstrated in *M. extorquens* AM1 (57) (log2fc values \geq 352 2.7), (ii) two contiguous ppqA genes highly homologous to M. extorquens AM1 PqqA precursor 353 of a low molecular weight redox active cofactor of dehydrogenases (23); (iii) a gene encoding 354 an ubiquitous family of L,D-transpeptidase catalytic domain YkuD involved in cell wall 355 composition adjustments (58) (log2fc value of 2.5); (iv) ada encoding a bifunctional 356 transcriptional activator/DNA repair enzyme Ada in response to alkylation damage in DNA (51 357 % aa Id) (log2fc value of 5.6) (59); and (v) *ibpA* encoding a small heat shock protein (sHSP) 358 displaying 57 % aa identity with the characterized E. coli IbpA that binds to misfolded proteins 359 and protects enzymes from stressful oxidants (60). In conclusion, genes found with higher 360 transcript abundance in cultures grown with chloromethane were involved in chloromethane 361 dechlorination and associated corrinoid cofactor and H₄F-metabolism, as well as in 362 adaptation, energy and central metabolisms.

363

Plasmid pCMU01 harbors a higher proportion of genes with chloromethane-enhanced
 transcription than the chromosome of strain CM4. The 380 kb pCMU01 plasmid harbors 62
 genes out of the 137 genes detected with transcript reads more abundant in cultures grown

367 with chloromethane compared to methanol. Among these, unique genes with no 368 chromosomal homologs were involved in carbon assimilation of the methylotrophic *cmu* 369 pathway and related functions as described in the previous paragraph, and also in ATP-370 dependent carboxylation of C₂ compounds with the complete set of the acx genes highly similar to those characterized in aerobic bacterium Xanthobacter autotrophicus acetone 371 372 carboxylase components AcxABC (>81 % aa Id) (61). A total of 31 duplicated genes displayed 373 38 to 87 % aa identity with chromosomal homologs (22) and had higher transcript abundance 374 on chloromethane, unlike their corresponding chromosomal copy. A majority of the plasmid-375 borne homologs were involved in corrinoid cofactor metabolism such as the *cob* genes (Fig. 376 5A).

377

378 FIG 5 Involvement of plasmid pCMU01 in the transcriptional response to chloromethane 379 Genes encoded by plasmid pCMU01 (in grey) or by the chromosome (in black) were grouped 380 according to their log2fc values with log2fc \geq 2.0 and \leq -2.0 meaning reads more abundant on 381 chloromethane and methanol, respectively. (A) Comparison of pCMU01- and chromosome-382 encoded paralogs with indication of their involvement in cofactor metabolisms essential for 383 growth with chloromethane. (B) Ratio calculated by dividing the gene number in defined 384 log2fc range, by the total number of genes located on the chromosome or on the plasmid 385 multiplied by one hundred. *M. extorquens* CM4 chromosome and plasmid pCMU01 encode 386 for a total number of 6261 and 361 genes, respectively. 387

388 Other plasmid homologs have been previously associated with the *cmu* pathway (*metF2* and 389 purU) (32,35), the serine cycle for carbon assimilation (gck2), and for adaptation to the 390 chloride stress (clc2) (Fig. 1; Table 2). For instance, the highly transcribed glycerate kinase 391 encoding gck2 gene seems to be more tightly regulated by chloromethane (log2fc value of 392 5.1) than the characterized chromosomal copy (log2fc value of - 0.5) (42), although they 393 encode for closely related homologs with 85 % aa Id. Similarly, the H⁺/Cl⁻ antiporter encoded 394 by the plasmid *clcA* gene was highly transcribed on chloromethane (log2fc value of 5.5) unlike 395 the chromosomal copy (68 % aa Id) that is required for adaptation to chloride stress of 396 M. extorquens DM4 on dichloromethane (11). Among 98 plasmid-borne genes that had 397 chromosomal homologs in M. extorquens, 70 genes were constitutively expressed under the 398 tested conditions, including genes involved in one-carbon metabolism, such as the 399 phosphoserine aminotransferase-encoding serC2 gene and the PQQ-linked methanol 400 dehydrogenase large subunit-encoding xoxF2 gene (88 % aa identity with the corresponding 401 chromosome homolog). Altogether, the plasmid pCMU01 had a higher proportion of genes

that are differentially transcribed in cultures grown with chloromethane versus methanol than
the chromosomal homologues (Fig. 5B). This highlights the central role of plasmid pCMU01borne genes in chloromethane utilization.

405

406 The dichloromethane transcriptome. The transcripts of 59 genes were found to be more 407 abundant in cultures grown with dichloromethane compared to methanol (log2fc \geq 2) (Tables 408 3 and S4). These included all four *dcm* genes of the dichloromethane utilization module (21). 409 Genes *dcmA* and *dcmB* were among the most abundant transcripts (Fig. S2B). The majority of 410 the genes that had increased reads on dichloromethane belonged to the core genome of 411 *M. extorquens* (Fig 4., group D, 29 genes). Among these, the putative glutathione peroxidase 412 (METDI0190, log2fc value of 2.5) is connected to the activity of the DcmA enzyme that 413 catalyzes the glutathione-dependent hydrolysis of dichloromethane. The formate 414 dehydrogenase homologs (fdh4AB) had lower read counts in cultures grown with methanol 415 compared to dichloromethane. This concurs with a *fdh4A* mutant phenotype found in 416 *M. extorquens* AM1 that was impaired for methanol utilization with formate accumulation in 417 the medium (45), suggesting that the *fdh4A* homolog may play a role in methanol dissimilation in different *M. extorquens* strains. Sets of uncharacterized molybdenum cofactor-dependent 418 419 enzyme-encoding genes had increased transcript abundance in cultures grown with 420 dichloromethane compared to methanol. These include the molybdopterin oxidoreductase 421 (METDI2693, log2fc value of 3.8) and three components of a molybdenum-containing 422 oxidoreductase (METDI0091-METDI0093). Their role remains unknown, as is the case with a 423 majority of the core genome encoding genes with higher transcript abundance on 424 dichloromethane (Tables 3 and S4). We found in this study that within 126 kb *dcm* GEI (29) 425 only the four *dcm* genes had enhanced expression on dichloromethane.

- 426
- 427

428 **TABLE 3** *M. extorquens* **DM4** transcripts differentially abundant in cultures grown with 429 dichloromethane compared to methanol

Labelª		News	Draduat	Occurrence	Directional RNAseq data		
		Name	Product	Occurrence	log2fc ^c	Adjusted p- value ^d	
DC	M utilization						
	METDI2655	dcmR	transcriptional repressor of DCM dehalogenase	dcm GEI ^g	2.7	0.040	
	METDI2656	dcmA	dichloromethane dehalogenase (DCM dehalogenase)	dcm GEI ^g	4.3	< 0.001	
_	METDI2657	-	conserved protein of unknown function DcmB	dcm GEI	4.0	< 0.001	

METDI2658	-	conserved protein of unknown function DcmC	dcm GEI	3.5	< 0.001
C ₁ metabolism					•
METDI2873	fdh4B	formate dehydrogenase subunit B	core	-3.2	< 0.001
METDI2874	fdh4A	formate dehydrogenase subunit A	core	-4.0	< 0.001
METDI5146	mxaW	conserved hypothethical protein	core	2.1	0.002
Central metabolism	l				
METDI0091	-	oxidoreductase, 2Fe-2S subunit	core ^e	2.5	< 0.001
METDI0092	-	oxidoreductase, molybdenum cofactor binding subunit	core ^e	2.9	< 0.001
METDI0128	ino	inositol-3-phosphate synthase	core	2.1	0.003
METDI0129	-	putative dTDP-glucose 4,6-dehydratase	core	2.2	0.002
METDI0190	-	putative glutathione peroxidase	core	2.5	0.002
METDI0283	adh-like	putative alcohol dehydrogenase	GEI 160	-3.3	0.029
METDI0286	xfp	D-xylulose 5-phosphate/ D-fructose 6-phosphate phosphoketolase	GEI 160	-2.9	0.099
METDI2432	-	putative phosphotransferase	core	2.7	< 0.001
METDI2693	-	putative molybdopterin oxidoreductase	core	3.8	0.008
METDI3569	arcB	ornithine cyclodeaminase	core ^f	2.6	< 0.001
METDI4540	adh-like	putative alcohol dehydrogenase	GEI 197	-3.7	0.012
METDI4543	xfp	D-xylulose 5-phosphate/D-fructose 6-phosphate phosphoketolase	GEI 197	-3.1	0.085
METDI4670	-	putative monooxygenase with ATPase activity	core	2.0	0.003
Energy	1	1			1
METDI0301	-	cation-transporting ATPase (P-type)	GEI 160	-3.3	0.071
METDI2011	cydB	cytochrome d terminal oxidase, polypeptide subunit II	core ^e	-3.4	0.036
METDI3863	hppa	H ⁺ translocating pyrophosphate synthase	core ^f	3.5	0.020
METDI4554	-	cation-transporting ATPase (P-type)	GEI 197	-3.5	0.059
METDI4816	qxtB	cytochrome bd-quinol oxidase subunit II	core ^e	-2.7	0.061
Lipid-related metab	olism	1	1	I	
METDI0292	-	putative poly-beta-hydroxyalkanoate synthase (PhbC)	GEI 160	-3.1	0.085
METDI0293	fabl	enoyl-(acyl-carrier-protein) reductase (NADH)	GEI 160	-3.6	0.046
METDI0294	-	putative phosphate butyryltransferase	GEI 160	-3.6	0.045
METDI0295	-	putative acetate kinase (AckA)	GEI 160	-3.4	<0.001
METDI1770	ohyA	oleate hydratase	variable	-3.4	0.044
METDI4905	-	putative patatin-like phospholipase	core	2.5	< 0.001
Nitrogen-related m	etabolism				1
METDI1697	-	putative formate/nitrate transporter	variable	-3.4	<0.001
METDI1754	-	putative cysteine desulfurase (SufS domain)	variable	-2.2	0.094
METDI2867	glnK	nitrogen regulatory protein P-II	core ^e	-2.5	0.029
METDI3157	glnll	glutamine synthetase, type II	core	-3.2	0.084
METDI5532	nrtB	nitrate transport permease protein	core ^e	-2.4	0.001
METDI5533	nrtA	nitrate transporter component	core ^e	-2.7	<0.001
Transport and adap	tation		1		1
METDI0093	-	putative voltage-dependent anion channel	core ^e	2.7	0.001
METDI0288	сlpВ	protein disaggregation chaperone	GEI 160	-3.6	0.014
METDI1223	-	putative cobalt/nickel resistance NcrA-like major facilitator superfamily permease	GEI 270	2.1	0.009
METDI1224	-	putative cobalt/nickel resistance NcrB-like regulator	GEI 270	2.8	< 0.001
METDI3043	-	conserved protein of unknown function with 2 CBS domains	core	4.4	0.002
METDI3067	-	putative methyl-accepting chemotaxis sensory transducer	core	2.5	0.009

METDI3303	-	putative extracellular protein with a calcium ion binding domain	core	2.8	< 0.001
METDI3835	dctA	C ₄ -dicarboxylate transport protein	core	-3.2	0.020
METDI4470	-	putative L,D-transpeptidase catalytic domain (YkuD)	GEI 197	-2.3	<0.001
METDI4699	-	transcriptional regulator, AraC family	core ^e	4.3	< 0.001
METDI5068	-	putative transcriptional regulator	variable ^f	3.2	0.011
METDI5746	depP	periplasmic serine protease (DegP)	core	6.6	0.010
METDI5875	-	putative photosynthesis gene regulator (bchF-crtJ)	variable	2.5	0.040
METDI5891	-	putative endonuclease	core	3.2	< 0.001

430 ^a MaGe annotation (https://www.genoscope.cns.fr/agc/microscope). See complete list Table S4

431 ^b Genes labeled as core, variable or specific are respectively found in all, at least 2 genomes of *M. extorquens* strains AM1, BJ001, CM4, 432

DM4 and PA1 (80 % of identity on at least 80 % of the protein length), or only in the genome of strain DM4. 433

^c log2 fold-change of normalized read numbers in cultures grown with dichloromethane compared to methanol.

434 ^d False discovery rate, only adjusted p-value under 0.1 were considered.

435 ^e Part of gene clusters: METDI1978-1985; METDI2004-2012[:] METDI2867-2874; METDI4813-4816; METDI5532-5533; METDI0091-0093;

436 METDI4695-4699.

437 ¹Adjacent to a gene of unknown function with transcript abundance higher in dichloromethane versus methanol

438 ^m Corresponding encoded protein found more abundant in cell protein extracts of dichloromethane versus methanol-grown cultures (21). 439

440 Methanol-enhanced transcript abundance of gene clusters located on redundant genomic 441 islands in *M. extorquens* DM4. Most of the genes with higher transcript abundance in cultures 442 grown with methanol compared to dichloromethane (Tables 3 and S4, 121 genes) were 443 localized on three GEI with sizes ranging from 160 to 270 kb (Table 4, 76 genes). The 444 proportion of genes with differential transcript abundance represented 19 and 31 % of the 445 gene content of two partially redundant GEI of 160 and 197 kb, called GEI 160 and GEI 197, 446 respectively (Table 4). Amongst these, an *ackA*-like gene displayed the strongest differential 447 transcript abundance for methanol in this study (log2fc value of -5.0 for METDI4544; Table 448 S4). A comparative genetic map of the two redundant GEI regions revealed transcription 449 pattern similarities between homologous genes sharing at least 74 % aa identity between their 450 encoded proteins (Fig. S3). Differentially expressed genes have predicted functions involved 451 in: i) central metabolism (two copies of putative acetate kinase-encoding ackA-like genes, one 452 copy of an alcohol dehydrogenase-encoding adh-like gene, two copies of a D-xylulose 5-453 phosphate/D-fructose 6-phosphate phosphoketolase-encoding *xfp* gene), ii) energy 454 metabolism (two copies of a cation-transporting ATPase, fourteen copies of cytochrome c and 455 cytochrome c oxidase SU), iii) lipid metabolism (one copy of an enoyl-(acyl-carrier-protein) 456 reductase-encoding *fabl* gene and a putative poly-beta-hydroxyalkanoate synthase-encoding 457 phbC gene), iv) stress response (a copy of the heat shock protein encoding dnaJ gene and of a 458 protein disaggregation chaperone-encoding *clpB* gene), and v) uncharacterized functions (five 459 copies of genes coding for members of conserved proteins containing domain ANAH (adenine 460 nucleotide alpha hydrolase protein superfamily Interpro SSF52402) (Fig. S3; S4).

Among the differentially transcribed homologous genes more abundant in methanol, a few were localized both on the variable and the core genomes, such as genes coding a cationtransporting ATPase, a cytochrome c oxidase, a Xfp enzyme and two members of the ANAH domain containing protein superfamily (Table 4). In conclusion, unique and accessory duplicated genes located on GEI may be involved in central metabolism, including carbon, energy and lipid metabolisms and stress for *M. extorquens* DM4 grown with methanol.

TABLE 4 Genomic island (GEI)-encoded transcripts differentially abundant in cultures of *M. extorquens* strains CM4 or DM4 grown with chlorinated methanes compared to methanol

Genomic island (GEI) and plasmid features ^a					Higher rea (CDS I	ad counts ^c METDI)		
Name	Length (kb)	Start-End (CDS Label) ^b	Start-End (nt)	(left border) inside (right border)	Chlorinated methane	Methanol	Number of regulated-genes (ratio) & relevant functions (genes)	
Within M. e	extorquens	DM4						
dcm GEI	126	2551-2682	2459291- 2585910	(none) tRNA-int- mob-misc_RNA- SIGI-AH (none)	2655-2658	none	4 (3) DCM utilization (<i>dcmRABC)</i>	
GEI 160	160	0225-0425	232397- 392215	(tRNA)-int-SIGI- AH (int)	0344	0259>0386	35 (18) central metabolism (<i>ackA</i> -like; <i>adh</i> - like; <i>xfp</i>), energy (ATPase; cyt. c; cyt. c oxidase SU), lipid metabolism (<i>fabI</i> ; <i>phbC</i>), stress (<i>clpB</i> ; <i>DnaJ</i>),	
GEI 197	197	4320-4570	4260147- 4456880	(none) tRNA-int- mob-SIGI-AH (IS)	None	4470; 4503>4565	33 (13) central metabolism (<i>ackA</i> -like; <i>adh</i> - like; <i>xfp</i>), energy (ATPase; cyt. c; cyt. c oxidase SU), members of ANAH-like superfamily	
GEI 270	270	1158-1475	1065633- 1335359	(tRNA) int-mob- misc_RNA-SIGI- AH (tRNA)	1223-1224	1414>1446	8 (3) metabolism (uncharacterized oxidoredcutase)	
Within <i>M.</i> e	extorquens	CM4						
GEI 84	84	1228-1308	1250078- 1333760	SIGI-IVOM- Specific_Region	1258 ; 1287-1288	none	3 (4) transcriptional regulator	
GEI 56	56	1470-1536	1531700- 1587516	tRNA-int-SIGI- IVOM- Specific_Region	1525	1533>1536	5 (8) metabolism (NosX precursor)	
GEI 12	12	1537-1558	1597506- 1609551	(none)	None	1537>1540	4 (19) energy (cyt. c), metabolism (<i>mdh2</i>)	
GEI 107	107	4756-4848	5048681- 5155763	tRNA-int-SIGI- IVOM- Specific_Region	4776-4777; 4779	none	3 (3) proteins of unknown functions	
GEI 28	28	5040-5066	5394282- 5422511	(none)	5057-5058; 5060; 5066	none	4 (15) central metabolism (<i>metA</i> monooxygenase, tRNA synthetase) lipid metabolism (<i>acd</i>)	
pCMU01	380	5386-5736	0-380207	nd ^d	5499; 5676-5704; 5714-5717; 5721-5732	none	62 (18) chloromethane utilization (<i>cmu</i>), corrinoïd cofactor biosynthesis (<i>cob</i> , <i>czc, bluB2</i>), transposase	

472 ^a Basing on "Regions of Genomic Plasticity" finder (MaGe plateform) using *M. extorquens* CM4, DM4, PA1 and BJ001 genomes. Int, integrase;

473 mob, mobility determinant; AH, region rich in genes detected by Alien Hunter based on variable-length k-mers bias (Vernikos and Parkhill,

474 2006) and SIGI, region with biased codon usage (62). Plasmid pCMU01 detected so far only in *M. extorquens* CM4 (22).

475 ^b Genes are labelled as Mchl and METDI in *M. extorquens* CM4 and DM4, respectively.

476 ^c Chlorinated methanes (log2fc > 2), methanol (log2fc < -2) as defined in material and methods.

- 477 ^d Not determined
- 478

479 **DISCUSSION**

- 480 The ratio of carbon source-dependent transcript abundance of *M. extorquens* genes was
- 481 significantly lower for genes in the core genome than in the variable genome (Table 5).
- 482

TABLE 5 Contribution of core and variable genome genes to carbon source-dependent differential transcriptional expression

			Differential transcript abundance			
Strain	Genome	CDS number		Gene number ^a		
			Ratio (%)	Up	Down	
	core	3,489	1.6	45	11	
M. extorquens CM4	variable ^b (CM4 only)	2,783 1,512	3.4 2.0	92 <i>29</i>	2 1	
	core	3,489	1.9	31	34	
M. extorquens DM4	variable ^b (DM4 only)	2,112 <i>952</i>	5.4 <i>5.6</i>	28 10	87 43	

485 ^a Transcript abundance higher (log2fc > 2) or lower (log2fc < -2) in cultures grown with chlorinated methane compared
 486 to with methanol.

487 ^bVariable genome includes genes not found in all *M. extorquens* genomes. Italics refer to strain-specific genes.

488

489 Strain-specific genes were among the most transcribed genes, especially gene clusters 490 involved in chlorinated methane dehalogenation. The *dcm* genes were more expressed in 491 growth with dichloromethane compared to methanol: *dcmA* coding for the dichloromethane 492 dehalogenase DcmA essential for growth with dichloromethane, dcmR coding for a regulator 493 of its own gene and of *dcmA* transcription (34,46), *dcmB* coding for a protein of unknown 494 function previously demonstrated to be dichloromethane-induced (21), and the 495 uncharacterized dcmC. Similarly, the cmuAB genes coding for essential chloromethane 496 dehalogenation enzymes underwent massive induction in cultures grown with 497 chloromethane, as previously described (63). Adjacent to the *cmuAB* genes on plasmid 498 pCMU01, other cmu pathway-associated genes may also be massively induced (cmuC, cmuC2, 499 folD, hutI, metF2, paaE, purU) as suggested in this study (22,51). In addition, plasmid-borne 500 genes with chromosomal homologs had higher transcript abundance in cultures with 501 chloromethane compared to methanol, unlike the chromosomal copies. This includes 14 cob 502 genes of the *cmu* pathway-associated corrinoid coenzyme biosynthesis (22), as well as *gck2*, 503 which encodes for a putative glycerate kinase displaying 85 % of aa identity with the 504 chromosomal homolog (Mchl_2974) involved in carbon assimilation by the serine pathway 505 (log2fc value of 5.1) (Table 2; clcA homologs are discussed in the next paragraph). This 506 suggests that a yet uncharacterized chloromethane-dependent regulation mechanism favors 507 the expression of genes of the variable genome over homologous genes of the core genome. 508 In the chloromethane transcriptome, the *acx* genes of the acetone carboxylase complex had 509 higher transcripts in cultures grown with chloromethane compared to methanol (log2fc values 510 of 2.1 to 3.7; Table 2). The *acx* genes, part of the variable genome located on plasmid pCMU01 511 in *M. extorquens* CM4, were conserved in the few sequenced genomes of isolates that 512 degrade chloromethane by the cmu pathway (51). Acetone has been described as the 513 substrate that regulates expression of the acx genes in the acetone-degrading Helicobacter 514 pylori (8), but the compound that induces acx gene expression during growth with 515 chloromethane has yet to be identified.

516

517 The transcriptional response common to *M. extorquens* growing with chloromethane or 518 dichloromethane is limited to a few genes only. Bacteria that grow with chlorinated 519 methanes have to cope with intracellular production of chloride ions and protons that are 520 formed during the first catabolic reactions (Fig. 1). Thus, we expected to find higher expression 521 of housekeeping genes involved in chloride and proton associated stress and bioenergetics as 522 a response to growth with chloromethane or dichloromethane. We identified such genes in 523 this study but only 11 among 4,620 genes were shared between the chloromethane- and 524 dichloromethane-degrading strains (Fig. 4 group B; Table S2). Of these genes, depD and hppA 525 had characterized homologs in other bacteria and were highly expressed in cultures grown 526 with chlorinated methanes (Table S2, group B). The central housekeeping molecular chaperon 527 protein DepD (48) repairs misfolded proteins damaged by reactive oxygen produced during 528 aerobic chlorinated methane utilization (21,22). As a proton pump, the membrane-bound 529 proton translocating pyrophosphatase HppA most likely plays a role in proton extrusion and 530 stress bioenergenetics, as demonstrated for a highly similar protein characterized in 531 Rhodospirullum rubrum (72 % aa identity with M. extorquens homolog) (10). For chloride 532 extrusion, the H⁺/Cl⁻ exchange transporter ClcA was encoded by a constitutively highly 533 expressed core genome gene in *M. extorquens* DM4, which confirms previous RT-qPCR data 534 in wild-type dichloromethane-degrading isolates (11). A different situation occurred in 535 *M. extorquens* CM4 that harbors two *clcA* paralogs (69 % aa Id). The *clcA* chromosomal core 536 genome copy displayed moderate constitutive transcript abundance whereas the clcA2 537 plasmid-borne gene significantly increased in transcript abundance with chloromethane

538 (log2fc value of 5.5), suggesting that *clcA*-driven chloride extrusion is also required during 539 growth with chloromethane. The occurrence of *clcA* paralogs with specific expression patterns 540 suggests that growth with chloromethane in *M. extorquens* CM4 and growth with 541 dichloromethane in *M. extorquens* DM4 both generate an adaptive response to chloride stress that is similar but not identical. Previous studies have shown that when gene clcA is 542 543 transcribed from its native promoter present in a synthetic expression module containing 544 dcmA, it confers higher fitness to dichloromethane in Methylobacterium strains that had not 545 been previously exposed to dichloromethane (11). This is not the case when using a synthetic 546 expression module containing the *cmu* pathway for growth with chloromethane (64) (J. K. 547 Michener, S. Vuilleumier, F. Bringel, C. J. Marx, personal communications; submitted article). 548 In addition, in laboratory evolution experiments, the genetic modules for heterologous 549 utilization of chloromethane and dichloromethane have shown that phylogeny poorly predicts 550 the utility of a challenging horizontally transferred gene in *Methylobacterium* strains (65), 551 which would suggest that strain-specific rather than core genome-encoded genes play a 552 crucial role in adaptation to growth with chlorinated methanes.

553 In both *M. extorquens* strains, 9 genes with lower transcript abundance were identified in 554 cultures grown with chlorinated methanes compared to methanol (Fig. 4 group E; Table S2). 555 Among these, *dctA* encodes a (Na⁺/H⁺)-C₄-dicarboxylate transporter (Dct) component highly similar to characterized DtcA proteins (77 % and 63 % aa identity with Rhizobium meliloti and 556 557 E. coli DctA, respectively) (47). For each C₄-dicarboxylate molecule imported, the Dct 558 transporter imports two protons (12). Therefore, during growth with chlorinated methanes, 559 an additional increase in intracellular proton concentration would be detrimental and may 560 explain the higher dctA transcript levels in cultures grown with methanol compared to those 561 grown with chloromethane or dichloromethane. In a previous study in M. extorquens AM1, 562 higher dctA transcript levels were found in cultures grown with succinate compared to 563 methanol while other two core genome dctA copies were not differentially expressed (66). Of 564 the three DctA homologs found in *M. extorquens* genomes that share around 50 % aa Id, only 565 one *dctA* copy has been identified as differentially expressed in this study (Table S2). Thus, the 566 expression of the same *dctA* homolog was modulated by carbon source-dependent growth 567 conditions in the three *M. extorquens* strains. In *E. coli*, DctA is a component of the Dct 568 transporter, as well as a co-regulator of the membrane-bond carboxylate sensor kinase DctS

569 (67). Therefore the differentially transcribed *dctA* gene may also play a potential role in570 carboxylate compound sensing in *M. extorquens*.

571 This study did not find any differentially expressed genes amongst the 163 genes shared 572 exclusively between the genomes of the two chlorinated methane-degrading strains and 573 absent in other non-degrading closely-related *M. extorquens* strains (Fig. 3; data not shown). 574 Conversely, a larger number of genes only present in either *M. extorquens* CM4 (6 genes) or 575 M. extorquens DM4 (10 genes) were differentially expressed in cultures grown with the 576 corresponding chlorinated methane compared to methanol (Tables 2, 3, S3 and S4). Our study 577 found that core genome encoded housekeeping proteins were differentially expressed in 578 cultures grown with dichloromethane compared to methanol (two uncharacterized 579 oxidoreductase with molybdenum cofactors METDI0092 and METDI2693; a putative 580 glutathione peroxidase; an ornithine cyclodeaminase ArcB). Other genes were only detected 581 in cultures grown with chloromethane (components of the NAD(P)⁺ transhydrogenase Pnt and 582 of the glycine cleavage system Gcv; fused transcriptional regulator and O6-methylguanine-583 DNA methyltransferase Ada) (Fig. 4). We found that the transcriptional responses to growth 584 with chloromethane or dichloromethane are predominantly uncorrelated, except a restricted 585 number of core-encoded genes involved in stress and bioenergetic intracellular HCl 586 production (*clcA* paralogs, *depP*, *ibpA* and *hppA*).

587

588 Core genome and strain-specific transcription patterns of methanol dehydrogenase and 589 alternative genes for alcohol/aldehyde utilization. The *M. extorquens* core genome harbors 590 several sets of methanol dehydrogenase (MDH)-encoding genes including the canonical 591 MxaFI, the environmental-widespread XoxF-type, and a Mdh2-type MDH encoded gene 592 cluster closely related to alcohol dehydrogenase ADH. The mxaFI and xoxF genes were not 593 significantly more expressed during growth with methanol compared to chloromethane or 594 dichloromethane (core genome-encoded genes mxa and xox and pCMU01 plasmid-borne 595 gene xoxF2). The unchanged expression profile with xoxF was expected since transcription of 596 xoxF-type genes are often undetected in cultures grown in the laboratory under conditions 597 different from environmental samples (68). Earlier microarray mRNA quantification in 598 M. extorquens strain AM1 found that mxaFI transcript abundance in cultures grown with 599 methanol was higher than with succinate, suggesting that *mxaFI* genes were induced during 600 methylotophic growth, whereas no gene belonging to the *mdh2* gene cluster was more

601 abundant (66). In *M. extorquens* AM1, *mdh2* alone cannot sustain growth with methanol since 602 a mxaFI mutant was unable to utilize methanol (69). In this study, surprisingly, all 8 genes of 603 the *mdh2* gene cluster were more expressed in methanol-grown strains CM4 and DM4 (Fig. 4). 604 The Mdh2 protein typically has low affinity for methanol except in *Betaproteobacteria* species 605 (40). In *M. extorquens*, Mdh2 enzyme affinity for methanol and other alcohol or aldehyde 606 compounds still needs to be tested whereas MxaFI catalyzes the oxidation of methanol as well 607 as formaldehyde (69). In this study, very high normalized read numbers of mxaFI genes were 608 detected in all the tested methylotrophic cultures although formaldehyde is a byproduct of 609 methanol and dichloromethane degradation but not of chloromethane dehalogenation 610 catalyzed by CmuAB (Fig. 1) (34,41,69). The equilibrium between free formaldehyde and 611 methylene-tetrahydrofolate (CH2=H4F) lies overwhelmingly on the side of methylene-H4F 612 formation, so significant reverse flux from CH₂=H₄F to formaldehyde and then to formate 613 seems unlikely during chloromethane utilization (70). The high level of normalized reads of 614 mxaFI mRNA during growth with chloromethane (Fig. S2A) suggests a yet unknown role of 615 MxaFI during growth with chloromethane or a loss of regulation in strain CM4.

616 Two *adh*-like genes of the family of zinc-dependent alcohol dehydrogenases (ADH) encode 617 similar proteins in strain DM4 (METDI0283 and METDI4540 display 90 % aa Id). They catalyze 618 the NAD(P)(H)-dependent interconversion of alcohols into aldehydes or ketones (71) and may 619 be involved in methanol utilization or C₂ metabolism (Fig. 1). These two *adh* genes, like a large 620 majority of the 121 transcripts found to be more abundant in cultures grown with methanol 621 compared to dichloromethane, are encoded by the variable genome and in particular two 622 partially-redundant GEI (68 genes; GEI 160 and GEI 197, Fig. S3; Tables 3 and S4). This shared 623 GEI region is also found as a single copy on the genome of *M. extorquens* AM1 nested within 624 a 64 kb-long GEI (Fig. S3). However, no differential transcript abundance has been detected in 625 previous microarray mRNA quantification in cultures grown with methanol compared to 626 succinate as growth substrate (66). This would suggest a dichloromethane-dependent 627 repression of the GEI genes or alternatively, a strain-specific methanol-dependent regulation 628 process. The adh-like genes and adjacent genes located on two redundant GEI found in 629 M. extorquens DM4 may be involved in central metabolism readjustments during 630 methylotrophic growth with methanol compared to dichloromethane.

631

155

632 Redox status adjustment as a possible key adaptation to growth with different C1 633 compounds. Formate dehydrogenases are central methylotrophic modules for C₁ 634 dissimulation (Fig. 1) and are encoded by four core genome fdh gene clusters (44,45). In 635 previous studies, the protein Fhd2B was detected as more abundant in strain DM4 grown with 636 dichloromethane compared to methanol (21), while *fdh1* and *fdh2* genes had higher transcript 637 abundance in strain AM1 grown with methanol compared to succinate (66). In this study, the 638 only difference in transcript abundance for fdh4AB was detected in M. extorquens DM4 639 (Tables 3 and S4). Taken together, expression of different copies of NAD-dependent formate 640 dehydrogenase components would suggest that methanol versus dichloromethane utilization 641 may affect specific redox reactions. This hypothesis is confirmed by the observation that 642 several types of cytochrome oxidase-encoded gene clusters, which belong to the core and the 643 variable genome, had higher transcript counts in methanol-grown cultures (Fig. 1; Table S3, 644 for instance *cbaB*, *qxtB*, *cydB* genes for an ATP transporter and an adjacent cytochrome d 645 terminal oxidase (METDI2004-METDI2012). In particular, within the variable genome of M. 646 extorquens DM4, a large set of GEI-encoded genes involved in cytochrome electron transport 647 and the formation of H⁺ gradient had enhanced transcript abundance in cultures grown with 648 methanol, and thus may be involved in ATP production (Fig. 1).

649 Other energy-associated adjustments in *M. extorquens* DM4 grown with methanol compared 650 to dichloromethane include very high transcript reads of genes involved in lipid metabolism 651 (Fig. 1; Tables 3 and S4.). Poly-beta-hydroxyalkanoate (PHB) constitutes a carbon and energy 652 reserve material during growth with both multicarbon- and single-carbon substrates (72). It 653 can represent 2.3 % of the cell dry weight of *M. extorquens* growing with methanol (4). PHB 654 metabolism controls carbon and energy spillage and its effectiveness depends on nitrogen 655 availability (9,73). In this study we identified five genes of the core genome with higher 656 transcript abundance in cultures grown with methanol. These five genes encode nitrogen-657 related functions (regulatory protein (glnK), transporters (atmB; ntrAB; putative 658 formate/nitrate transporter METDI1697) and enzymes (glutamine synthase glnII; cysteine 659 desulfurase; Table 3). In summary, the differential RNAseq (methanol versus 660 dichloromethane) study revealed the differential expression of core and variable genome-661 encoded genes with functions associated with energy metabolism.

For each molecule of chloromethane transformed to formate by the *cmu* pathway, two molecules of NADPH are generated from $NADP^+$ in two successive reactions catalyzed by 664 MetF2 and FoID (Fig. 1). The corresponding genes metF2 and foID are two of the most 665 differentially expressed genes in the chloromethane transcriptome (Table 2). Thus, lowering the intracellular pool of NADP⁺ should limit growth with chloromethane unless a metabolic 666 667 reaction could regenerate NADP⁺. A good candidate for this is the membrane-bound 668 transhydrogenase encoded the pnt gene cluster, which was significantly more expressed on 669 chloromethane both at the level of transcription and protein (Table 2) (22). It catalyzes the 670 reversible reaction NADPH + NAD⁺ + H⁺_{in} <=> NADP⁺ + NADH + H⁺_{out} (74). This membrane-671 bound transhydrogenase plays a predominant role in laboratory evolution of *M. extorquens* 672 and E. coli deficient in NADPH production (49,66), which would be the case in cells utilizing 673 chloromethane with the cmuA pathway. In addition to NADP⁺ regeneration via the membrane-674 bound transhydrogenase, concomitant proton efflux would also lower internal cellular pH 675 occurring during chloromethane dehalogenation. Whether this core genome membrane-676 bound transhydrogenase is essential for NADP⁺ regeneration in cultures grown with 677 chloromethane still needs to be tested.

678

679 Finally, we observed that the difference in transcript abundance was the highest for genes of 680 the variable genome (*cmu* genes) during growth with chloromethane compared to methanol 681 (log2fc of 10 instead of 4 for *dcm* genes). On the other hand, similar transcript abundances 682 were detected for core genome-encoded genes when both strains grew with methanol 683 (Fig. S2). Thus, chloromethane-dependent transcription regulation seems very effective and 684 confirms earlier studies (63). Here, we observed a looser dichloromethane-dependent control 685 at the transcription level with high dcm transcript abundance even in the non-induced 686 condition (Table 3). At the protein level, nevertheless, no DcmA protein has been detected in 687 the methanol proteome while it accounts for about 25 % of total proteins in the 688 dichloromethane proteome (21). These observations strongly suggest that an effective post-689 transcriptional regulation of the variable genome of the *dcm* islet of *M. extorquens* DM4 may 690 take place. Future transcriptional start site determination will be conducted to characterize 691 core and variable genome transcription regulatory elements and mechanisms responsible for 692 growth with different C₁ compounds.

693

694 MATERIALS AND METHODS

695 Bacterial growth conditions and RNA purification. M. extorquens strains CM4 and DM4 grew 696 aerobically at 30°C on a rotary shaker (120 rpm) in 1.2-liter Erlenmeyer flasks closed with gas-697 tight screw caps with mininert valves (Supelco) in 220 mL Methylobacterium mineral medium 698 (M3) adapted from (75) (ZnSO₄ replaced ZnCl₂ and (NH₄)₂SO₄ was at final concentration 0.2 699 g.L⁻¹). The one-carbon compounds added at 10 mM final concentration were either methanol 700 (2.75 mL of a methanol filter-sterilized stock solution at 800 mM), dichloromethane (141 μ L 701 of dichloromethane 100 %) or chloromethane (40 mL of CH₃Cl 100 % assuming a Henry constant of 0.0106 m³.atm.-mol⁻¹ at 30°C as previously described) (76). The 220-mL cultures 702 703 were stopped during the mid-exponential phase ($OD_{600nm} \approx 0.15$) by the addition of 27.5 mL 704 of stabilization buffer. The stabilization buffer was prepared as follows: 5 mL of phenol was mixed with 5 mL of 1 M sodium acetate pH 5.5 and 90 mL of DEPC-treated water. After 705 706 centrifugation at 1800 g for 3 min, 5 mL of the inferior phase was mixed with 95 mL of ethanol 707 100 %. Cell suspensions were centrifuged at 5000 rpm for 5 min at 4°C and resuspended in 10 708 mL of TE containing 2 mg. mL⁻¹ of lysozyme (Euromedex). After an incubation 15 min at 37°C, 709 the suspension was centrifuged 10 min at 4°C, then the pellet was resuspended with 10 mL of 710 Trizol (Invitrogen), then 2.5 mL of chloroform was added. RNAs were precipitated with 711 isopropanol and washed with ethanol (70 %), then resuspended in DEPC-water and treated 712 with DNAse (TurboDNAse, Invitrogen). DNA depletion was checked by PCR on rrs (27f, 5'-713 GTTTGATCCTGGCTCAG-3' and 927r, 5'-CCGTCAATTCCTTTRAGTTT-3') and on specific genes: cli (cli1R, 5'-ATCGTCCTGATCGCACCT-3' and cli1F, 5'-GCCTCGAAACGTTGAACACC-3'), dcmA 714 5'-TCGATCTAGATAGCGCGGTCTAAGGGAC-3' 5'-715 (dcmAiR, and dcmAiF, 716 GACTTCTAGACGTGAGCCCGAATCAAC-3') or стиА (cmuA802F, 5'-717 TTCAACGGCGAYATGTATCCYGG-3' and cmuA1244R, 5'-TABTCCATDATGGCYTCGA-3'). RNA 718 quality was checked on a Bioanalyzer 2100 using RNA 6000 Nano ship (Agilent) and quantified 719 with Qubit RNA kit (Invitrogen). Then, rRNA were removed by treating 5 µg of total RNA with 720 Gram negative RiboZero Magnetic kit (Tebu-Bio) according to the manufacturer's protocol.

721

cDNA library preparation, sequencing and data normalization. Fifty to sixty ng of rRNAdepleted RNA were used to construct cDNA banks with the TruSeq stranded mRNA LT kit (Illumina). This construction started with the step "Purify and fragment mRNA" from Illumina protocol, by adding 13 μ L of the solution "Fragment, Prime, Finish Mix" to the 5 μ L of RNA containing 50-60 ng of RNA. Then the manufacture's protocol was followed. Bank 727 constructions were validated by Bioanalyzer with DNA 1000 kit (Agilent). Sequencing was 728 performed using HiSeq2000. Paired-end sequence mapping was used to limit mapping 729 artifacts and remove any ambiguity at mapping level between homologs. For data 730 normalization (Fig. S1), the log2fc value of normalized read numbers in chlorinated methanes versus methanol was normalized with a set of 55 reference genes listed in Table S1, and a cut-731 732 off value = 1. The reference genes were unique and constitutively expressed in the tested 733 conditions. Genes were defined as differentially abundant in cultures grown with chlorinated 734 methane compared to methanol when log2fc values were \geq 2.0 and differentially abundant in 735 cultures grown with methanol compared to chlorinated methanes when log2fc values were ≤ 736 -2.0 (77). Each condition was performed in duplicate; the number of reads was based on the average of read sense and antisense for each condition. Data were only considered significant 737 738 when the p-value was \leq 0.1. The complete RNAseq dataset is accessible online (https:// 739 www.genoscope.cns.fr/agc/microscope/transcriptomic/NGSProjectRNAseq.php?projType=R 740 NAseq).

741

742 Comparative genomics and phylogenetic analysis. The core genome of *M. extorquens* was 743 done using the sequenced genomes of five strains of the *M. extorquens* species (GenBank 744 accession n°): strain AM1 (CP001511, CP001512, CP001513 and CP001514); strain BJ001 (CP001029, CP001030 and CP001031); strain CM4 (CP001298, CP001299, CP001300); strain 745 746 DM4 (FP103042 and FP103043 and FP103044) and strain PA1 (CP000908) (28,29). The core 747 genome established via the MaGe platform was 748 (https://www.genoscope.cns.fr/agc/microscope/compgenomics/pancoreTool.php?). A gene 749 was defined as belonging to the core genome when it encodes a protein sharing at least 80 % 750 of identity on 80 % of CDS length in all the tested genomes. Gene was defined as belonging to 751 the core genome when it is shared by the 5 strains, or belonging to the variable genome when 752 it is shared at least by 2 strains and maximum by 4 strains. Sequences were aligned via ClustalX 753 (78). The phylogenetic tree was constructed using the maximum parsimony method with a 754 bootstrap of 500 replicates with MEGA4 software (79).

755

756 SUPPLEMENTAL MATERIAL

FIGURE S1 Comparison of genome wide gene transcription between *M. extorquens* strains
 CM4 and DM4 grown with methanol. The percentage of reads represents the number of
 reads for each gene divided by the cumulated number of reads of the 4,620 genes shared (at

least 80 % of identity on the protein level on 80 % of the CDS length) in a given strain. Name
of highly transcribed genes are indicated. (A) Normalization using 55 reference genes. (B)
Normalization with the standard method using the complete set of CDS.

763

FIGURE S2 Genes with highest differential transcription in cultures grown with chlorinated methanes compared to methanol. Black rectangles indicate values of log2fc (values higher than 2.0 or lower than -2.0 mean normalized reads more abundant in cultures grown with chlorinated methanes or with methanol, respectively). Grey rectangles represent normalized read numbers divided by gene length in kb. (A) RNAseq data of *M. extorquens* CM4 cultures grown with chloromethane compared to methanol. (B) RNAseq data of *M. extorquens* DM4 cultures grown with dichloromethane compared to methanol.

771

772 FIGURE S3 Transcription profiles and gene redundancy of key gene clusters more highly 773 transcribed in methanol- than dichloromethane-grown cultures of *M. extorquens* DM4. 774 Conserved stretches between two GEI-borne regions (Table 4) are delimited with grey 775 rectangles. Numbers indicate the percentage of amino acid identity between encoded METDI 776 protein labels defined in MicroScope 777 (https://www.genoscope.cns.fr/agc/microscope/mage/viewer. php?S id=240). Genes are 778 indicated as colored arrows: in red when higher normalized read numbers were detected in 779 cultures grown with methanol compared to dichloromethane (log2fc \leq -2), in blue when no 780 significant difference was observed ($-2 < \log_2 fc$ values >2), and in purple for the METDI0344-781 encoding gene detected with higher reads in dichloromethane (log2fc value of 2.2). 782 Interrupted GEI segments are indicated with zigzagged arrows. IS stands for insertion 783 sequences. The ANAH conserved domain is found in proteins of the "adenine nucleotide alpha 784 hydrolases-like superfamily" including N type ATP PPases, ATP sulphurylases, universal stress 785 response proteins and electron transfer flavoproteins 786 (http://supfam.cs.bris.ac.uk/SUPERFAMILY/cgi-bin/scop.cgi?sunid=52402).

787

788 FIGURE S4 Phylogenetic tree of proteins with adenine nucleotide alpha hydrolase (ANAH) 789 domains in *M. extorquens* strains CM4 and DM4. Alphaproteobacteria members of the 790 aligned ANAH family proteins. Protein containing only one ANAH domain (Interpro SSF52402) 791 are boxed. Alignments were performed using ClustalX (78) for proteins mean sizes between 792 270-280 aa. The phylogenetic tree was constructed using software MEGA (79) with the 793 neighbor joining method (Saitou and Nei, 1987). Color-coded protein names correspond to 794 genes with higher (in orange) or unchanged (in black) transcript abundance in *M. extorquens* 795 cells grown with methanol compared to chlorinated methanes. In parenthesis, protein-796 encoded gene occurrence in the variable genome (genomic island) or in *M. extorquens* shared 797 genes (core). Symbols indicate when an ANAH-encoding gene is located in the vicinity (less 798 than 2 kb with) of a gene coding for an ATPase (star), a cytochrome c (disk), an alcohol 799 dehydrogenase-like (ADH-like) (triangle) or *fixJL* (square) encoding a putative two component 800 low oxygen sensor histidine kinase. Sequence accession numbers for *M. extorquens* CM4: 801 Mchl_0097 (WP_012605265), Mchl_1555 (WP_015950242), Mchl_3807 (WP_015951826), 802 Mchl 3811 (WP 015951830), Mchl 3815 (WP 015951832), Mchl 4817 (WP 015952568); 803 for M. extorguens DM4: METDI0258 (WP 0127790266), METDI0272 (WP 012779075), 804 METDI0274 (WP 012779077), METDI0276 (WP 012779079), METDI0302 (WP 012779104), 805 METDI0320 (WP_012779122), METDI0323 (WP_012779125), METDI2004 (WP_015821999), 806 METDI2442 (WP_01582230), METDI4304 (WP_015823643), METDI4309 (WP_015823647),

METDI4314 (WP_015823651), METDI4521 (WP_012753369), METDI4535 (WP_015823830),
METDI4537 (WP_012753383), METDI4539 (WP_012753385), METDI4555 (WP_0158238333),
METDI4563 (WP_012753411), METDI4565 (WP_01275344); for *Bradyrhizobium japonicum*(WP_011085546), *Sinorhizobium meliloti* (WP_015242313) and *Xanthobacter autotrophicus*(WP_012115156).

- 813 TABLE S1 Set of 55 genes used for RNAseq data normalization
- 814
- 815TABLE S2 Groups of genes with similar transcription shared between *M. extorquens* strains816CM4 and DM4
- 817

TABLE S3 Differentially abundant transcripts in cultures of *M. extorquens* CM4 grown with
 chloromethane compared to methanol (Table 2 complementary list)

- 821TABLE S4 Differentially abundant transcripts in cultures of *M. extorquens* DM4 grown with822dichloromethane compared to methanol
- 823

824 ACKNOWLEDGMENTS

825 PhD grant to CP (funded by Région Alsace (France) and partially by DFG grant ko2912/5-1 826 (Germany); the German Academic Exchange Service (DAAD); University of Bayreuth) and 827 Franco-German mobility funds (Deutscher Akademischer Austausch Dienst (PROCOPE)) are 828 gratefully acknowledged. Thanks to Beatrice Segurens (Commissariat à l'Energie Atomique et 829 aux Energies Alternatives (CEA) CNRS-UMR8030, France) for RNA extraction protocol 830 optimization, and David A. C. Beck (University of Washington, Seattle, USA) and Anne Francez-831 Charlot (ETH Zurich, Swiss) for checking proposed RNAseq reference genes in unpublished M. 832 extorquens transcriptomic libraries. 833

834 **REFERENCES**

- 1. Galbally IE, Kirstine W. 2002. The production of methanol by flowering plants and the global
 cycle of methanol. J Atmospheric Chem 43: 195–229. 10.1023/A:1020684815474.
- 837 2. Clerbaux C, Cunnold DM, Anderson J. 2006. Halogenated long-lived compounds, scientific
- assessment of ozone depletion. Global Ozone Research and Monitoring Project 10 Report
- 839 No 50. World Meteorological Organization, Geneva.
- 840 3. Trotsenko YA, Doronina NV. 2003. The biology of methylobacteria capable of degrading
- 841 halomethanes. Microbiology **72**: 121–131. doi:10.1023/A:1023218326407

842	4. Peyraud R, Schneider K, Kiefer P, Massou S, Vorholt JA, Portais J-C. 2011. Genome-scale
843	reconstruction and system level investigation of the metabolic network of
844	Methylobacterium extorquens AM1. BMC Syst Biol 5 : 189. doi:10.1186/1752-0509-5-189
845	5. Šmejkalová H, Erb TJ, Fuchs G. 2010. Methanol assimilation in Methylobacterium
846	extorquens AM1: demonstration of all enzymes and their regulation. PLoS ONE 5: e13001.
847	doi:10.1371/journal.pone.0013001
848	6. Chistoserdova L. 2011. Modularity of methylotrophy, revisited: modularity of
849	methylotrophy, revisited. Environ Microbiol 13 : 2603–2622. doi:10.1111/j.1462-
850	2920.2011.02464.x
851	7. Schmidt S, Christen P, Kiefer P, Vorholt JA. 2010. Functional investigation of methanol
852	dehydrogenase-like protein XoxF in <i>Methylobacterium extorquens</i> AM1. Microbiology 156 :
853	2575–2586. doi:10.1099/mic.0.038570-0
854	8. Brahmachary P, Wang G, Benoit SL, Weinberg MV, Maier RJ, Hoover TR. 2008. The human
855	gastric pathogen Helicobacter pylori has a potential acetone carboxylase that enhances its
856	ability to colonize mice. BMC Microbiol 8: 14. doi:10.1186/1471-2180-8-14
857	9. Sauer U. 2003. The soluble and membrane-bound transhydrogenases UdhA and PntAB
858	have divergent functions in NADPH metabolism of <i>Escherichia coli</i> . J Biol Chem 279: 6613–
859	6619. doi:10.1074/jbc.M311657200
860	10. Lopez-Marques RL, Perez-Castineira JR, Losada M, Serrano A. 2004. Differential
861	regulation of soluble and membrane-bound inorganic pyrophosphatases in the
862	photosynthetic bacterium Rhodospirillum rubrum provides insights into pyrophosphate-

- 863
 based stress bioenergetics. J Bacteriol
 186:
 5418–5426.
 doi:10.1128/JB.186.16.5418

 864
 5426.2004
- Michener JK, Neves AAC, Vuilleumier S, Bringel F, Marx CJ. 2014. Effective use of a
 horizontally-transferred pathway for dichloromethane catabolism requires post-transfer
 refinement. eLife 3: e04279. doi:10.7554/eLife.04279.001
- I2. Janausch IG, Zientz E, Tran QH, Kröger A, Unden G. 2002. C₄-dicarboxylate carriers and
 sensors in bacteria. Biochim Biophys Acta BBA-Bioenerg 1553: 39–56. doi:10.1016/S00052728(01)00233-x
- 13. Davies SJ, Golby P, Omrani D, Broad SA, Harrington VL, Guest JR, et al. 1999. Inactivation
- and regulation of the aerobic C_4 -dicarboxylate transport *dctA* gene of *Escherichia coli*. J
- 873 Bacteriol **181**: 5624–5635. doi:10/1999; 181(18):5624-35.

- 14. Naik S, Venu Gopal SK, Somal P. 2008. Bioproduction of polyhydroxyalkanoates from
 bacteria: a metabolic approach. World J Microbiol Biotechnol 24: 2307–2314.
 doi:10.1007/s11274-008-9745-z
- 877 15. Korotkova N, Lidstrom ME. 2001. Connection between poly-hydroxybutyrate biosynthesis
 878 and hrowth on C₁ and C₂ compounds in the methylotroph *Methylobacterium extorquens*
- AM1. J Bacteriol **183**: 1038–1046. doi:10.1128/JB.183.3.1038-1046.2001
- 16. Thomson AW, O'Neill JG, Wilkinson JF. 1976. Acetone production by methylobacteria.
 Arch Microbiol 109: 243–246. doi:10.1007/BF00446635
- 882 17. Eggers J, Steinbuchel A. 2013. Poly-3-Hydroxybutyrate degradation in *Ralstonia eutropha* 883 H16 is mediated stereoselectively to (S)-3-hydroxybutyryl coenzyme A (CoA) via crotonyl-
- 884 CoA. J Bacteriol **195**: 3213–3223. doi:10.1128/JB.00358-13
- 18. Sznajder A, Jendrossek D. 2014. To be or not to be a poly-3-hydroxybutyrate (PHB)
 depolymerase: PhaZd1 (PhaZ6) and PhaZd2 (PhaZ7) of *Ralstonia eutropha*, highly active
 PHB depolymerases with no detectable role in mobilization of accumulated PHB. Appl
 Environ Microbiol 80: 4936–4946. doi:10.1128/AEM.01056-14
- 19. Kayser MF, Vuilleumier S. 2001. Dehalogenation of dichloromethane by dichloromethane
 dehalogenase/glutathione S-transferase leads to formation of DNA adducts. J Bacteriol
 183: 5209–5212. doi:10.1128/JB.183.17.5209-5212.2001
- 20. Torgonskaya ML, Doronina NV, Hourcade E, Trotsenko YA, Vuilleumier S. 2011. Chlorideassociated adaptive response in aerobic methylotrophic dichloromethane-utilising
 bacteria. J Basic Microbiol 51: 296–303. doi:10.1002/jobm.201000280
- 895 21. Muller EEL, Hourcade E, Louhichi-Jelail Y, Hammann P, Vuilleumier S, Bringel F. 2011.
 896 Functional genomics of dichloromethane utilization in *Methylobacterium extorquens* DM4:
 897 functional genomics of *Methylobacterium* dichloromethane utilization. Environ Microbiol
- 13: 2518–2535. doi:10.1111/j.1462-2920.2011.02524.x
- 899 22. Roselli S, Nadalig T, Vuilleumier S, Bringel F. 2013. The 380 kb pCMU01 plasmid encodes 900 chloromethane utilization genes and redundant genes for vitamin B₁₂- and 901 tetrahydrofolate-dependent chloromethane metabolism in Methylobacterium extorquens 902 CM4: а proteomic and bioinformatics study. PLoS ONE **8**: e56598. 903 doi:10.1371/journal.pone.0056598
 - 163

- 904 23. Ochsner AM, Sonntag F, Buchhaupt M, Schrader J, Vorholt JA. 2015. *Methylobacterium* 905 *extorquens*: methylotrophy and biotechnological applications. Appl Microbiol Biotechnol
- 906 **99**: 517–534. doi:10.1007/s00253-014-6240-3
- 907 24. Bringel F, Couée I. 2015. Pivotal roles of phyllosphere microorganisms at the interface
 908 between plant functioning and atmospheric trace gas dynamics. Front Microbiol 06.
 909 doi:10.3389/fmicb.2015.00486
- 910 25. Temkiv TŠ, Finster K, Hansen BM, Nielsen NW, Karlson UG. 2012. The microbial diversity
- 911 of a storm cloud as assessed by hailstones. FEMS Microbiol Ecol 81: 684–695.
 912 doi:10.1111/j.1574-6941.2012.01402.x
- 913 26. Kolb S. 2009. Aerobic methanol-oxidizing in soil. FEMS Microbiol Lett 300: 1–10.
 914 doi:10.1111/j.1574-6968.2009.01681.x
- 915 27. Mitsui R, Omori M, Kitazawa H, Tanaka M. 2005. Formaldehyde-limited cultivation of a
 916 newly isolated methylotrophic bacterium, *Methylobacterium* sp. MF1: enzymatic analysis
 917 related to C₁ metabolism. J Biosci Bioeng **99**: 18–22. doi:10.1263/jbb.99.18
- 918 28. Marx CJ, Bringel F, Chistoserdova L, Moulin L, Farhan Ul Haque M, Fleischman DE, et al.
- 919 2012. Complete genome sequences of six strains of the genus *Methylobacterium*. J
 920 Bacteriol **194**: 4746–4748. doi:10.1128/JB.01009-12
- 921 29. Vuilleumier S, Chistoserdova L, Lee M-C, Bringel F, Lajus A, Zhou Y, et al. 2009.
- 922 *Methylobacterium* genome sequences: a reference blueprint to investigate microbial 923 metabolism of C₁ compounds from natural and industrial sources. Ahmed N, editor. PLoS
- 924 ONE **4**: e5584. doi:10.1371/journal.pone.0005584
- 30. Doronina NV, Sokolov AP, Trotsenko YA. 1996. Isolation and initial characterization of
 aerobic chloromethane-utilizing bacteria. FEMS Microbiol Lett 142: 179–183.
 doi:10.1016/0378-1097(96)00262-5
- 31. Gälli R, Leisinger T. 1985. Specialized bacterial strains for the removal of dichloromethane
 from industrial waste. Conservation & Recycling 8: 91–100. 10.1016/0361-3658(85)900281
- 32. Studer A, Stupperich E, Vuilleumier S, Leisinger T. 2001. Chloromethane: tetrahydrofolate
 methyl transfer by two proteins from *Methylobacterium chloromethanicum* strain CM4.
- 933 Eur J Biochem **268**: 2931–2938. doi:10.1046/j.1432-1327.2001.02182.x
- 934 33. Nadalig T, Farhan Ul Haque M, Roselli S, Schaller H, Bringel F, Vuilleumier S. 2011.
 935 Detection and isolation of chloromethane-degrading bacteria from the *Arabidopsis*

- thaliana phyllosphere, and characterization of chloromethane utilization genes:
 chloromethane-degrading bacteria from *A. thaliana*. FEMS Microbiol Ecol **77**: 438–448.
 doi:10.1111/j.1574-6941.2011.01125.x
- 34. Leisinger T, Bader R, Hermann R, Schmid-Appert M, Vuilleumier S. 1994. Microbes,
 enzymes and genes involved in dichloromethane utilization. Biodegradation 5: 237–248.
 doi:10.1007/BF00696462
- 35. Studer A, McAnulla C, Buchele R, Leisinger T, Vuilleumier S. 2002. Chloromethaneinduced genes define a third C₁ utilization pathway in *Methylobacterium chloromethanicum* CM4. J Bacteriol 184: 3476–3484. doi:10.1128/JB.184.13.34763484.2002
- 36. Schmid-Appert M, Zoller K, Traber H, Vuilleumier S, Leisinger T. 1997. Association of
 newly discovered IS elements with the dichloromethane utilization genes of
 methylotrophic bacteria. Microbiology 143: 2557–2567. doi:10.1099/00221287-143-82557
- 37. Nakagawa T, Mitsui R, Tani A, Sasa K, Tashiro S, Iwama T, *et al.* 2012. A catalytic role of
 XoxF1 as La³⁺-dependent dethanol dehydrogenase in *Methylobacterium extorquens* strain
 AM1. Battista JR, editor. PLoS ONE **7**: e50480. doi:10.1371/journal.pone.0050480

38. Skovran E, Palmer AD, Rountree AM, Good NM, Lidstrom ME. 2011. XoxF is required for
expression of methanol dehydrogenase in *Methylobacterium extorquens* AM1. J Bacteriol
193: 6032–6038. doi:10.1128/JB.05367-11

- 39. Neufeld JD, Chen Y, Dumont MG, Murrell JC. 2008. Marine methylotrophs revealed by
 stable-isotope probing, multiple displacement amplification and metagenomics:
 metagenomics of active marine methyotrophs. Environ Microbiol 10: 1526–1535.
 doi:10.1111/j.1462-2920.2008.01568.x
- 40. Kalyuzhnaya MG, Hristova KR, Lidstrom ME, Chistoserdova L. 2008. Characterization of a
 novel methanol dehydrogenase in representatives of *Burkholderiales*: implications for
 environmental detection of methylotrophy and evidence for convergent evolution. J
 Bacteriol 190: 3817–3823. doi:10.1128/JB.00180-08
- 41. Vannelli T, Messmer M, Studer A, Vuilleumier S, Leisinger T. 1999. A corrinoid-dependent
 catabolic pathway for growth of a *Methylobacterium* strain with chloromethane. Proc Natl
 Acad Sci 96: 4615–4620. doi:10.1073/pnas.96.8.4615

967	42. Šmejkalová H, Erb TJ, Fuchs G. 2010. Methanol assimilation in Methylobacterium
968	extorquens AM1: demonstration of all enzymes and their regulation. PLoS ONE 5: e13001.
969	doi:10.1371/journal.pone.0013001
970	43. Marx CJ, Van Dien SJ, Lidstrom ME. 2005. Flux analysis uncovers key role of functional
971	redundancy in formaldehyde metabolism. PLoS Biol.3: e16.
972	doi:10.1371/journal.pbio.0030016
973	44. Chistoserdova L, Laukel M, Portais J-C, Vorholt JA, Lidstrom ME. 2004. Multiple formate
974	dehydrogenase enzymes in the facultative methylotroph Methylobacterium extorquens
975	AM1 are dispensable for growth on methanol. J Bacteriol 186 : 22–28.
976	doi:10.1128/JB.186.1.22-28.2004
977	45. Chistoserdova L, Crowther GJ, Vorholt JA, Skovran E, Portais J-C, Lidstrom ME. 2007.
978	Identification of a fourth formate dehydrogenase in Methylobacterium extorquens AM1
979	and confirmation of the essential role of formate oxidation in methylotrophy. J Bacteriol
980	189 : 9076–9081. doi:10.1128/JB.01229-07
981	46. Kayser MF, Ucurum Z, Vuilleumier S. 2002. Dichloromethane metabolism and C_1
982	utilization genes in <i>Methylobacterium</i> strains. Microbiology 148 : 1915–1922.
983	doi:10.1099/00221287-148-6-1915
984	47. Engelke T, Jording D, Kapp D, Pühler A. 1989. Identification and sequence analysis of the
985	<i>Rhizobium meliloti dctA</i> gene encoding the C ₄ -dicarboxylate carrier. J Bacteriol 171 : 5551–
986	5560. doi:10.1007/BF00284690
987	48. Ge X, Wang R, Ma J, Liu Y, Ezemaduka AN, Chen PR, et al. 2014. DegP primarily functions
988	as a protease for the biogenesis of β -barrel outer membrane proteins in the Gram-negative
989	bacterium <i>Escherichia coli</i> . FEBS J 281 : 1226–1240. doi:10.1111/febs.12701
990	49. Belogurov GA. 2002. A Lysine Substitute for K ⁺ . J Biol Chem 277: 49651–49654.
991	doi:10.1074/jbc.M210341200
992	50. Mirete S, Mora-Ruiz MR, Lamprecht-Grandío M, de Figueras CG, Rosselló-Móra R,
993	González-Pastor JE. 2015. Salt resistance genes revealed by functional metagenomics from
994	brines and moderate-salinity rhizosphere within a hypersaline environment. Front
995	Microbiol 6 . doi:10.3389/fmicb.2015.01121
996	51. Nadalig T, Greule M, Bringel F, Keppler F, Vuilleumier S. 2014. Probing the diversity of
997	chloromethane-degrading bacteria by comparative genomics and isotopic fractionation.
998	Front Microbiol 5 . doi:10.3389/fmicb.2014.00523

- 52. Trausch JJ, Ceres P, Reyes FE, Batey RT. 2011. The structure of a tetrahydrofolate-sensing
 riboswitch reveals two ligand binding sites in a single aptamer. Structure 19: 1413–1423.
 doi:10.1016/j.str.2011.06.019
- 1002 53. Vitreschak AG. 2003. Regulation of the vitamin B₁₂ metabolism and transport in bacteria
 1003 by a conserved RNA structural element. RNA 9: 1084–1097. doi:10.1261/rna.5710303
- 1004 54. Bassford PJ, Kadner RJ. 1977. Genetic analysis of components involved in vitamin B₁₂
 1005 uptake in *Escherichia coli*. J Bacteriol 132: 796–805. 10.1371/journal.pone.0062382
- 1006 55. Yu T-Y, Mok KC, Kennedy KJ, Valton J, Anderson KS, Walker GC, *et al.* 2012. Active site
 1007 residues critical for flavin binding and 5,6-dimethylbenzimidazole biosynthesis in the flavin
 1008 destructase enzyme BluB. Protein Sci 21: 839–849. doi:10.1002/pro.2068
- 56. Mandal M. 2004. A glycine-dependent riboswitch that uses cooperative binding to control
 gene expression. Science. 306: 275–279. doi:10.1126/science.1100829
- 1011 57. Chou H-H, Marx CJ, Sauer U. 2015. Transhydrogenase promotes the robustness and
 1012 evolvability of *Escherchia coli* deficient in NADPH production. PLoS Genet 11: e1005007–
 1013 e1005007. doi:10.1371/journal.pgen.1005007
- 1014 58. Bielnicki J, Devedjiev Y, Derewenda U, Dauter Z, Joachimiak A. 2005. Derewenda ZS.
 1015 Bascillus subtilis ykuD protein at 2.0 Å resolution: insights into the structure and function
- 1016 of a novel, ubiquitous family of bacterial enzymes. Proteins Struct Funct Bioinforma **62**:

1017 144–151. doi:10.1002/prot.20702

1018 59. Mc Carthy T V, Lindahl T. 1985. Methyl pbosphotnesters in alkylated DNA are repaired by
 1019 the Ada regulatory protein of *Escherichia coli*. Nucleic Acids Research 13: 2683–2698.

1020 60. Kitagawa M, Miyakawa M, Matsumura Y, Tsuchido T. 2002. Escherichia coli small heat

1021 shock proteins, IbpA and IbpB, protect enzymes from inactivation by heat and oxidants. Eur

```
1022 J Biochem 269: 2907–2917. doi:10.1046/j.1432-1033.2002.02958.x
```

- 1023 61. Sluis MK, Larsen RA, Krum JG, Anderson R, Metcalf WW, Ensign SA. 2002. Biochemical,
 1024 molecular, and genetic analyses of the acetone carboxylases from *Xanthobacter* 1025 *autotrophicus* strain Py2 and *Rhodobacter capsulatus* strain B10. J Bacteriol 184: 2969–
 1026 2977. doi:10.1128/JB.184.11.2969-2977.2002
- 1027 62. Waack S, Keller O, Asper R, Brodag T, Damm C, Fricke WF, *et al.* 2006. Score-based
 prediction of genomic islands in prokaryotic genomes using hidden Markov models. BMC
 Bioinformatics 7: 142.

- 1030 63. Farhan Ul Haque M, Nadalig T, Bringel F, Schaller H, Vuilleumier S. 2013. Fluorescence 1031 based bacterial bioreporter for specific detection of methyl halide emissions in the
 1032 environment. Appl Environ Microbiol. 79: 6561–6567. doi:10.1128/AEM.01738-13
- Michener, J.K., Vuilleumier, S., Bringel, F., Marx, C.J. submitted. Effectiveness of
 heterologous catabolism of chloromethane and dichloromethane are uncorrelated in
 Methylobacterium strain. Front Microbiol
- 1036 65. Michener JK, Vuilleumier S, Bringel F, Marx CJ. 2014. Phylogeny poorly predicts the utility
- 1037 of a challenging horizontally transferred gene in *Methylobacterium* strains. J Bacteriol 196:
 2101–2107. doi:10.1128/JB.00034-14
- 1039 66. Okubo Y, Skovran E, Guo X, Sivam D, Lidstrom ME. 2007. Implementation of microarrays
 1040 for *Methylobacterium extorquens* AM1. OMICS J Integr Biol 11: 325–340.
 1041 doi:10.1089/omi.2007.0027
- 1042 67. Steinmetz PA, Wörner S, Unden G. 2014. Differentiation of DctA and DcuS function in the
 1043 DctA/DcuS sensor complex of *E scherichia coli* : function of DctA as an activity switch and
 1044 of DcuS as the C₄ -dicarboxylate sensor: DctA and DcuS functional differentiation in
 1045 DctA/DcuS. Mol Microbiol 94: 218–229. doi:10.1111/mmi.12759
- 1046 68. Delmotte N, Knief C, Chaffron S, Innerebner G, Roschitzki B, Schlapbach R, *et al.* 2009.
 1047 Community proteogenomics reveals insights into the physiology of phyllosphere bacteria.
 1048 Proc Natl Acad Sci 106: 16428–16433. doi:10.1073/pnas.0905240106
- 1049 69. Nunn DN, Lidstrom ME. 1986. Phenotypic characterization of 10 methanol oxidation
 1050 mutant classes in *Methylobacterium* sp. strain AM1. J Bacteriol 166: 591–597. doi:00211051 9193/86/050591-07\$02.00/0
- 1052 70. Kallen RG, Jencks WP. 1966. The mechanism of the condensation of formaldehyde with
 1053 tetrahydrofolic acid. J Biol Chem 241: 5851–5863.
- 1054 71. Persson B, Hedlund J, Jörnvall H. 2008. Medium- and short-chain
 1055 dehydrogenase/reductase gene and protein families: The MDR superfamily. Cell Mol Life
 1056 Sci 65: 3879–3894. doi:10.1007/s00018-008-8587-z
- 1057 72. Escapa IF, García JL, Bühler B, Blank LM, Prieto MA. 2012. The polyhydroxyalkanoate
 1058 metabolism controls carbon and energy spillage in *Pseudomonas putida*. Environ Microbiol
- 1059 **14**: 1049–1063. doi:10.1111/j.1462-2920.2011.02684.x
- 1060 73. Hervas AB, Canosa I, Santero E. 2008. Transcriptome analysis of *Pseudomonas putida* in
 1061 response to nitrogen availability. J Bacteriol 190: 416–420. doi:10.1128/JB.01230-07

- 1062 74. Carroll SM, Marx CJ. 2013. Evolution after introduction of a novel metabolic pathway
 1063 consistently leads to restoration of wild-type physiology. Schneider D, editor. PLoS Genet
 1064 9: e1003427. doi:10.1371/journal.pgen.1003427
- 1065 75. Vannelli T, Studer A, Kertesz M, Leisinger T. 1998. Chloromethane metabolism by
 1066 *Methylobacterium* sp. strain CM4. Appl Environ Microbiol 64: 1933–1936. doi:05/1998;
 1067 64(5):1933-6.
- 1068 76. Chen F, Freedman DL, Falta RW, Murdoch LC. 2012. Henry's law constants of chlorinated
 1069 solvents at elevated temperatures. Chemosphere 86: 156–165.
 1070 doi:10.1016/j.chemosphere.2011.10.004
- 1071 77. Yang Y-X, Wang M-M, Yin Y-L, Onac E, Zhou G-F, Peng S, et al. 2015. RNA-seq analysis
- 1072 reveals the role of red light in resistance against *Pseudomonas syringae* pv. tomato DC3000
- 1073 in tomato plants. BMC Genomics **16**: 120. doi:10.1186/s12864-015-1228-7
- 1074 78. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, et al.
 1075 2007. Clustal W and Clustal X version 2.0. Bioinformatics 23: 2947–2948.
 1076 doi:10.1093/bioinformatics/btm404
- 1077 79. Tamura K, Dudley J, Nei M, Kumar S. 2007. MEGA4: Molecular Evolutionary Genetics
- 1078 Analysis (MEGA) software version 4.0. Mol Biol Evol **24**: 1596–1599

FIG 1 Metabolism of chlorinated C₁ compounds chloromethane and dichloromethane and of methanol in *M. extorquens* CM4 and DM4




FIG 2 Overview of genomic islands and gene synteny in genomes of *M. extorquens* CM4 and DM4

FIG 3 Core, variable and specific genome of five strains affiliated to the species *Methylobacterium extorquens*



FIG 4 Carbon source-dependent shared categories of gene transcript abundance between *M. extorquens* CM4 and DM4 genomes



FIG 5 Involvement of plasmid pCMU01 in the transcriptional response to chloromethane





FIGURE S1 Comparison of genome wide gene transcription between *M. extorquens* strains CM4 and DM4 grown with methanol

FIGURE S2 Genes with highest differential transcription in cultures grown with chlorinated methanes compared to methanol





FIGURE S3 Transcription profiles and gene redundancy of key gene clusters more highly transcribed in methanol- than dichloromethane-grown cultures of *M. extorquens* DM4



FIGURE S4 Phylogenetic tree of proteins with adenine nucleotide alpha hydrolase (ANAH) domains in *M. extorquens* strains CM4 and DM4



S1 Table. Set of 55 genes used for RNAseq data normalization

Lab	elª	Gene	Product
Strain CM4	Strain DM4		
Mchl_0002	METDI0001	dnaA	DNA replication initiator protein
Mchl_0220	METDI4779	-	putative phosphodiesterase
Mchl_0372	METDI0202	glnE	glutamate-ammonia-ligase adenylyltransferase
Mchl_0480	METDI0517	zwf	glucose-6-phosphate 1-dehydrogenase
Mchl_0641	METDI0609	-	conserved protein of unknown function
Mchl_0710	METDI0680	-	conserved protein of unknown function
Mchl_0713	METDI0683	qxtB	cytochrome bd ubiquinol oxidase, subunit II
Mchl_0917	METDI1100	prmC	protein-(glutamine-N5) methyltransferase, release factor-specific
Mchl_1568	METDI2018	-	putative CelB-like protein
Mchl_1668	METDI2051	-	conserved protein of unknown function precursor
Mchl_1859	METDI2245	trmE	tRNA modification GTPase trmE
Mchl_2123	METDI2469	lysA	diaminopimelate decarboxylase
Mchl_2201	METDI2686	-	conserved protein of unknown function
Mchl_2204	METDI2689	-	transglutaminase family protein cysteine peptidase
Mchl_2251	METDI2738	-	putative DNA topoisomerase I
Michl_2277	METDI2766	gmk 	guanylate kinase
Mchl_2458	METDI2953	adk	adenylate kinase
Mchl_2482	METDI2981	тар	methionine aminopeptidase
Mchl_2549	METDI3056	rnc	ribonuclease III
Mchl_2552	METDI3059	rsuA	16S rRNA pseudouridylate 516 synthase
Mchl_2581	METDI3088	ung	uracil-DNA glycosylase superfamily protein
Mchl_2691	METDI3197	-	ribonuclease BN family protein
Mchl_2750	METDI3257	-	putative decarboxylase, thiamine pyrophosphate requiring
Mchl_2817	METDI3327	thrB	homoserine kinase
Mchl_2824	METDI3334	recA	RecA protein
Mchl_3177	METDI3720	IpxC	UDP-3-O-acyl N-acetylglucosamine deacetylase
Mchl_3288	METDI3840	gatB	aspartyl/glutamyl-tRNA amidotransferase subunit B
Mchl_3478	METDI3942	-	ATP-dependent DNA helicase
Mchl_3479	METDI3943	-	conserved protein of unknown function
Mchl_3486	METDI3950	ribE	6,7-dimethyl-8-ribityllumazine synthase
Mchl_3504	METDI3967	mutY	A/G-specific adenine glycosylase
Mchl_3512	METDI3975	-	putative adenylate cyclase protein
Mchl_3524	METDI3989	rimi	ribosomal-protein-alanine N-acetyltransferase
Mchl_3541	METDI4005	murl	glutamate racemase
Mchl_3578	METDI4045	thrC	threonine synthase
Michl_3610	METDI4078	nhaA	sodium:proton antiporter
Michl_3635	METDI4104	руrВ	aspartate carbamoyitransferase
IVICHI_3693	METDI4186	-	conserved protein of unknown function
Nichi_3946	IVIETDI4648	JaaD	acyl-coa synthetase
4052		-	tonB-dependent siderophore receptor protein
Nichi_4185		prs	phosphoridosylpyrophosphate synthetase
Nichi_4253	METDI4871	gia da v D	tRNA uridine 5-carboxymethylaminomethyl modification enzyme
Nobl 4306	METDI4930	апав	replicative DNA nelicase DnaB
Nabl 4404	IVIETDI4968	- ftall	
		JISK	cell division protein
		-	conserved protein of unknown function
	METDIE 427	pijc	olongation factor Tu (FE-Tu)
Mchl 4014	METDI5427	luj	nutative ATP_independent RNA belicase
	METDIE602	dell	DNA ligaça NAD(*) dopondont
Mchl 5143	METDIS092	InvP	linid_A_disaccharide synthase
Mchl 5150	METDIS731	Vals	valine tRNA synthetase
Mchl 5242	METDI5748	nhnC	valine trive synthetase
Mchl 5245	METDI5053	hem	5-aminolevulinic acid synthase
	METDISOOT	nema	s-annihitevullille actu synthase
101011-2212		-	

^a MaGe annotation (https://www.genoscope.cns.fr/agc/microscope) of genes with constitutive transcript abundance (log2fc value between -0.9 and 1.2 of normalized read numbers in cultures grown with chlorinated methanes compared to methanol.

S2 Table. Groups of genes with similar transcription shared between *M. extorquens* strains CM4 and DM4

Group and label ^a		Name	Product	log2	!fc⁵	Adjusted p-value ^c	
CM4	DM4	ittaine		CM4	DM4	CM4	DM4
Group B: mor	e abundant on	both chl	orinated methanes				
Mchl_0651	METDI0620	-	conserved protein of unknown function	2.3	3.8	< 0.001	< 0.001
Mchl_1347	METDI1765	-	conserved protein of unknown function	2.1	3.3	< 0.001	0.004
Mchl_2116	METDI2461	-	conserved protein of unknown function	2.6	2.5	< 0.001	0.016
Mchl_2560	METDI3067	-	putative methyl-accepting chemotaxis sensory transducer	2.0	2.5	< 0.001	0.009
Mchl_2793	METDI3303	-	protein of unknown function	2.7	2.8	< 0.001	< 0.001
Mchl_3129	METDI3671	-	conserved protein of unknown function, DUF1328	2.5	2.7	< 0.001	< 0.001
Mchl_3408	METDI3863	hppa	H ⁺ translocating pyrophosphate synthase	2.4	3.5	< 0.001	0.019
Mchl_3885	METDI4584	-	protein of unknown function	2.2	2.5	< 0.001	0.006
Mchl_3919	METDI4618	-	conserved protein of unknown function	2.4	2.9	< 0.001	< 0.001
Mchl_4560	METDI5190	-	protein of unknown function	2.1	3.8	< 0.001	< 0.001
Mchl_5157	METDI5746	depP	periplasmic serine protease (DegP)	4.6	6.6	< 0.001	< 0.001
Group C: mor	e abundant on	chlorom	ethane				
Mchl_1416	METDI1841	-	putative membrane protein	2.2	1.4	< 0.001	0.071
Mchl_1525	METDI1969	-	conserved protein of unknown function	2.0	1.9	< 0.001	0.006
Mchl_2287	METDI2777	-	conserved protein of unknown function	2.2	1.9	< 0.001	0.083
Mchl_3132	METDI3674	-	protein of unknown function	2.2	1.9	< 0.001	0.021
Mchl_4484	METDI5109	-	protein of unknown function	4.2	1.7	< 0.001	0.059
Mchl_5057	METDI5648	-	putative seryl-tRNA synthetase	2.2	1.6	< 0.001	0.072
Mchl_5058	METDI5649	acd	acyl-CoA dehydrogenase	2.2	1.7	< 0.001	0.036
MCv2_5149	METDI2503	pqqA	coenzyme PQQ biosynthesis protein A	2.1	1.5	< 0.001	0.063
Group D: more	abundant on o	dichloron	nethane				
Mchl_0009	METDI0008	-	protein of unknown function	1.9	2.4	< 0.001	< 0.001
Mchl_0154	METDI0091	iorA	isoquinoline 1-oxidoreductase, alpha subunit	0.9	2.5	0.014	< 0.001
Mchl_0155	METDI0092	-	oxidoreductase, molybdenum cofactor binding subunit	1.5	2.9	< 0.001	< 0.001
Mchl_0156	METDI0093	-	voltage dependent anion channel protein	1.9	2.7	< 0.001	0.001
Mchl_0171	METDI0108	-	protein of unknown function	1.0	2.0	0.017	0.011
Mchl_0189	METDI0128	-	inositol-1-phosphate synthase	1.7	2.1	< 0.001	0.003
Mchl_0190	METDI0129	-	putative dTDP-glucose 4,6-dehydratase	1.8	2.2	< 0.001	0.002
Mchl_0194	METDI0133	-	putative glycosyl transferase	1.3	2.6	0.048	0.006
Mchl_0360	METDI0190	-	putative glutathione peroxidase	1.1	2.5	0.002	0.009
Mchl_1112	METDI1593	-	transposase of ISMex11, IS3 family	0.7	2.1	0.410	0.260
Mchl_1492	METDI1934	-	protein of unknown function	1.7	2.2	< 0.001	0.014
Mchl_1516	METDI1959	-	conserved protein of unknown function precursor	0.7	2.5	0.047	0.003
Mchl_1729	METDI2115	-	conserved protein of unknown function	0.1	3.0	0.920	< 0.001
Mchl_2091	METDI2432	-	putative aminoglycoside phosphotransferase	1.7	2.7	< 0.001	< 0.001
Mchl_2208	METDI2693	-	putative molybdopterin oxidoreductase	1.9	3.8	< 0.001	0.008
Mchl_2537	METDI3043	-	conserved protein of unknown function with 2 CBS domains	1.0	4.4	0.003	0.002
Mchl_2848	METDI3359	-	protein of unknown function	1.1	2.1	0.029	0.002
Mchl_2894	METDI3426	-	conserved protein of unknown function	1.3	2.1	< 0.001	0.006
Mchl_3030	METDI3569	arcB	ornithine cyclodeaminase	1.6	2.6	< 0.001	< 0.001
Mchl_3921	METDI4620	-	protein of unknown function	1.2	2.0	0.026	0.002
Mchl_3964	METDI4670	-	putative monooxygenase with ATPase activity	1.8	2.0	< 0.001	0.003
Mchl_4030	METDI4699	-	transcriptional regulator, AraC family	-0.6	3.2	< 0.001	< 0.001
Mchl_4335	METDI4959	-	conserved protein of unknown function	1.5	2.2	0.001	< 0.001
Mchl_4440	METDI5067	-	conserved protein of unknown function	1.9	4.3	0.098	0.002
Mchl_4491	METDI5117	-	protein of unknown function	1.2	3.5	0.007	< 0.001
Mchl_4727	METDI1224	-	putative NcrB-like regulator, DUF156	0.8	2.8	0.059	< 0.001
Mchl_4728	METDI1223	-	putative NcrA-like major facilitator superfamily permease	0.9	2.1	0.034	0.009
Mchl_5285	METDI5875	-	putative photosynthesis gene regulator (bchF-crtJ)	1.1	2.5	0.001	0.037
Mchl_5301	METDI5891	-	putative endonuclease	1.5	3.2	< 0.001	< 0.001
Group E: mor	e abundant on	methan	bl				
Mchl_1165	METDI1517	-	conserved membran protein of unknown function	-3.1	-2.3	< 0.001	< 0.001
Mchl_1533	METDI1978	-	conserved protein of unknown function	-2.9	-2.1	< 0.001	0.037
Mchl_1534	METDI1979	-	putative NosX precursor (nitrous oxide reductase)	-3.3	-3.2	< 0.001	0.013
Mchl_1535	METDI1980	-	putative regulatory protein NosR	-3.4	-2.6	< 0.001	0.019

Mchl_1536	METDI1981	-	conserved protein of unknown function	-3.7	-2.5	< 0.001	< 0.001
Mchl_1537	METDI1982	-	conserved exported protein of unknown function	-3.3	-3.3	< 0.001	< 0.001
Mchl_1538	METDI1983	-	cytochrome c550 protein, putative PQQ-dependent methanol/ethanol oxidation system	-3.9	-3.8	< 0.001	< 0.001
Mchl_1539	METDI1984	-	conserved protein of unknown function, periplasmic binding protein precursor, putative periplasmic binding protein-like II	-4.5	-3.8	< 0.001	< 0.001
Mchl_3283	METDI3835	dctA	C ₄ -dicarboxylate transport protein	-2.9	-3.2	< 0.001	0.029
Group F: mor	e abundant on	methan	ol in strain CM4 only				
Mchl_0844	METDI1024	-	conserved membran protein of unknown function	-2.0	-1.8	< 0.001	0.031
Group G: mo	re abundant on	methan	ol in strain DM4 only				
Mchl_0500	METDI0537	-	ABC transporter, membrane protein	1.6	-2.5	< 0.001	0.059
Mchl_0501	METDI0538	-	ABC transporter, permease	1.1	-2.0	0.015	0.073
Mchl_1243	METDI1697	-	putative formate/nitrate transporter	-1.6	-3.4	< 0.001	< 0.001
Mchl_1337	METDI1753	eshA	eshA	-1.1	-3.3	< 0.001	< 0.001
Mchl_1338	METDI1754	-	putative cysteine desulfurase (SufS domain)	-1.1	-2.2	< 0.001	0.094
Mchl_1341	METDI1761	-	conserved exported protein of unknown function	-1.4	-2.6	< 0.001	< 0.001
Mchl_1717	METDI2103	-	conserved protein of unknown function	2.5	-2.4	< 0.001	0.003
Mchl_2380	METDI2873	fdh4B	formate dehydrogenase subunit B	-1.2	-3.2	0.001	< 0.001
Mchl_2381	METDI2874	fdh4A	formate dehydrogenase subunit A	-0.7	-4.0	0.048	< 0.001
Mchl_2653	METDI3157	glnII	glutamine synthetase II	-0.8	-3.2	0.081	0.065
Mchl_3806	METDI4303	-	putative transcriptional regulator	1.5	-3.6	< 0.001	0.048
Mchl_3807	METDI4304	-	conserved protein with ANAH-like domain	1.1	-3.0	0.004	0.054
Mchl_3808	METDI4305	-	conserved protein of unknown function	0.8	-2.4	0.069	0.039
Mchl_3810	METDI4307	-	protein of unknown function	0.9	-2.7	0.028	0.034
Mchl_4141	METDI4814	-	protein of unknown function, putative exported protein	0.9	-4.0	0.036	0.001
Mchl_4142	METDI4816	qxtB	cytochrome bd-quinol oxidase subunit II	1.2	-2.7	0.008	0.061
Mchl_4650	METDI5284	-	putative membrane of unknown function	-1.6	-3.1	< 0.001	< 0.001
Mchl_4958	METDI5532	nrtB	nitrate transport permease protein	-1.1	-2.4	0.017	0.001
Mchl_4959	METDI5533	nrtA	nitrate transporter component	-1.1	-2.7	0.007	< 0.001

^a MaGe annotation (https://www.genoscope.cns.fr/agc/microscope). Group as defined in Fig 3.

^b log2fc of normalized read numbers in cultures grown with chlorinated methanes compared to methanol.

^c False discovery rate, only adjusted p-value under 0.1 were considered.

S3 Table. Differentially abundant transcripts in cultures of *M. extorquens* CM4 grown with chloromethane compared to methanol (Table 2 complementary list)

	Nomo	Droduct	Occurr-	Directional RNAseq data		
Laber	Name	Product	ence⁵	log2 fc°	Adjusted p-value ^d	
Mchl_0472	-	putative PYP-like sensor domain	core	4.8	< 0.001	
Mchl_1288	-	putative transcriptional regulator, AraC family	variable	2.2	< 0.001	
Mchl_1416	-	putative membrane protein	core	2.2	< 0.001	
Mchl_1525	-	protein of unknown function	variable	2.0	< 0.001	
Mchl_1533	-	protein of unknown function	core ^e	-2.9	< 0.001	
Mchl_1534	-	putative NosX precursor (nitrous oxide reductase)	core ^e	- 3.3	< 0.001	
Mchl_1535	-	conserved membrane protein, FMN-binding and 4Fe-4S binding domains	core ^e	-3.4	< 0.001	
Mchl_1536	-	conserved protein of unknown function, putative YVTN beta- propeller repeat family protein	core ^e	-3.7	< 0.001	
Mchl_1537	-	conserved exported protein of unknown function	core ^e	-3.3	< 0.001	
Mchl_1539	-	conserved protein of unknown function, putative periplasmic binding protein-like II	core ^e	-4.5	< 0.001	
Mchl_2560	-	putative exported sensory transducer protein	core	2.0	< 0.001	
Mchl_2920	-	putative exported sensory transducer protein	variable	-2.4		
Mchl_3129	-	conserved protein of unknown function, DUF1328	core	2.5	< 0.001	
Mchl_4776	-	conserved protein of unknown function, putative esterase	specific	2.1	< 0.001	
Mchl_4777	-	putative outer-membrane protein	variable	2.1	< 0.001	
Mchl 4779	-	conserved protein of unknown function	specific	2.4	< 0.001	
 Mchl_5060	-	putative monooxygenase, small component; putative flavin:NADH reductase	variable	2.5	< 0.001	
Mchl_5347	-	conserved protein of unknown function, DUF892	core	2.1	< 0.001	
Mchl_5465	-	transposase (fragment)	pCMU01	2.3	< 0.001	
Mchl_5499	-	response regulator in two-component regulatory system, sigma- 54 interaction domain	pCMU01	2.4	< 0.001	
Mchl_5505	-	integrase catalytic region	pCMU01	2.9	0.018	
Mchl_5526	-	transposase of ISMex11, IS3 family (ORF 2)	pCMU01	2.3	< 0.001	
Mchl 5581	-	transposase (fragment)	pCMU01	2.3	0.011	
	-	conserved protein of unknown function (fragment)	pCMU01	2.2	< 0.001	
Mchl 5663	-	conserved protein of unknown function	pCMU01	2.1	< 0.001	
	-	protein of unknown function	pCMU01	4.8	< 0.001	
	-	putative transposase (fragment)	pCMU01	4.2	< 0.001	
Mchl 5703	-	putative transposase (fragment)	pCMU01	2.4	0.015	
	-	putative transposon (fragment)	pCMU01	3.2	< 0.001	
	-	conserved protein of unknown function, CobS-like domain	pCMU01	7.1	< 0.001	
	-	tRNA-Thr	core	2.1	< 0.001	
Mchl R0005	-	tRNA-Glu	core	2.3	< 0.001	
Mchl R0006	-	tRNA-Ser	core	2.0	< 0.001	
Mchl R0057	-	tRNA-Thr	core	2.3	< 0.001	
Mchl R0078	-	tRNA-Asn	core	2.1	< 0.001	
Mchl_RncRNA 2297962R	-	SRP RNA, RNA component of signal recognition particule	core	2.1	< 0.001	
p1MCv2 0094	-	protein of unknown function	pCMU01	2.3	< 0.001	
p1MCv2 0095	-	conserved protein of unknown function	pCMU01	2.3	< 0.001	
p1MCv2 0379	-	protein of unknown function	pCMU01	2.4	< 0.001	
p1MCv2_0380	-	protein of unknown function	pCMU01	2.1	< 0.001	
		· · · · · · · · · · · · · · · · · · ·				

^a MaGe annotation (https://www.genoscope.cns.fr/agc/microscope).

^b Genes labeled as core, variable or specific are respectively found in all, at least 2 genomes of *M. extorquens* genomes of strains AM1, BJ001, CM4, DM4 and PA1 (80 % of identity on at least 80 % of the protein length), or only in the chromosome of strain CM4. Genes labeled as pCMU01 are located on the 380 kb plasmid (Roselli *et al.*, 2013).

^c log2 fold-change of normalized read numbers in cultures grown with chloromethane compared to methanol.

^d False discovery rate, only adjusted p-value under 0.1 were considered.

^e Methanol-more expressed gene cluster Mchl_1533 to Mchl_1540.

S4 Table. Differentially abundant transcripts in cultures of *M. extorquens* DM4 grown with dichloromethane compared to methanol

l a b a la	Nama	Durchust	Occurr-	Directional RNAseq data		
Label®	Name	Product	ence ^b	log2fc ^c	Adjusted p-value ^d	
List of genes mor	e abundant o	on dichloromethane				
METDI0008	-	protein of unknown function	core	2.4	< 0.001	
METDI0108	-	protein of unknown function	variable	2.0	0.012	
METDI0190	-	putative glutathione peroxidase	core	2.5	0.001	
METDI0620	-	conserved exported protein of unknown function	core	3.8	< 0.001	
METDI1593	-	transposase of ISMex11, IS3 family (ORF 2)	variable	2.1	0.009	
METDI1597	-	exported protein of unknown function	specific	2.3	0.001	
METDI1598	-	protein of unknown function	specific	2.1	0.001	
METDI1755	-	conserved protein of unknown function	specific	2.1	0.004	
METDI1765	-	protein of unknown function	core	3.3	0.004	
METDI1934	-	protein of unknown function	variable	2.2	0.014	
METDI1959	-	conserved protein of unknown function precursor	variable	2.5	0.036	
METDI2/12		nutative phosphotransforase	variable	3.0	< 0.001	
METDI2432		protein of unknown function	specific	2.7	0.016	
METDI2401		putative molybdonterin oxidoreductase	core	3.8	0.010	
METDI2033	-	conserved protein with 2 CBS domains	core	4.4	0.002	
METDI3067	-	putative methyl-accepting chemotaxis sensory transducer	core	2.5	0.009	
METDI3303	-	putative extracellular protein with a Ca binding domain	core	2.8	0.001	
METDI3359	-	protein of unknown function	core	2.1	0.002	
METDI3426	-	conserved protein of unknown function	core	2.1	0.006	
METDI4190	-	exported protein of unknown function	core	2.1	0.048	
METDI4320		putative alcohol dehydrogenase (fragment)	specific	2.6	0.064	
METDI4584	-	protein of unknown function	core	2.5	0.006	
METDI4608	-	conserved membrane protein of unknown function	variable	2.2	0.006	
METDI4618	-	conserved protein of unknown function	core	2.9	0.004	
METDI4620	-	protein of unknown function	variable	2.2	0.002	
METDI4670	-	putative monooxygenase with ATPase activity	core	2.0	0.004	
METDI4741	-	putative response regulator, (CheY-like protein)	core	2.0	0.009	
METDI4905	-	putative patatin-like phospholipase	core	2.5	< 0.001	
METDI4959	-	conserved protein of unknown function	core	2.2	< 0.001	
METDI5117	-	protein of unknown function	variable	3.5	< 0.001	
METDI5146	mxaW	conserved hypothethical protein	core	2.1	0.001	
METDI5190	-	protein of unknown function	variable	3.8	0.001	
METDI5265	-	protein of unknown function	variable	2.4	0.002	
METDI5746	depP	periplasmic serine protease	core	6.6	< 0.001	
METDI5875	-	putative photosynthesis gene regulator (<i>bcnF-crtJ</i>)	Variable	2.5	0.004	
IVIE I DIS891	-		core	3.2	< 0.001	
		METDI0091-METDI0093 gene cluster				
METDI0091	-	oxidoreductase, 2Fe-2S subunit	core	2.5	< 0.001	
METDI0092	-	oxidoreductase, molybdenum cofactor binding subunit	core	2.9	< 0.001	
METDI0093	-	putative voltage-dependent anion channel	core	2.7	0.001	
		METDI0128-METDI0133 gene cluster				
METDI0128	ino	inositol-3-phosphate synthase	core	2.1	0.003	
METDI0129	-	putative dTDP-glucose 4,6-dehydratase	core	2.2	0.002	
METDI0133	-	putative glycosyl transferase	variable	2.6	0.006	
		GEI 160				
METDI0344	-	conserved protein of unknown function	variable	2.2	0.012	
		GEI 270				
			1		1	
METDI1223	-	superfamily permease	variable	2.1	0.009	
METDI1224	-	putative cobalt/nickel resistance NcrB-like regulator	variable	2.8	< 0.001	
		dcm GEI				
METDI2655	dcmR	transcriptional repressor of DCM dehalogenase	specific ^f	2.7	0.046	
METDI2656	dcmA	dichloromethane dehalogenase (DCM dehalogenase)	specific ^f	4.3	< 0.001	
METDI2657	-	conserved protein of unknown function DcmB	specific	4.0	< 0.001	
METDI2658	-	conserved protein of unknown function DcmC	specific	3.5	< 0.001	
		METDI3569-METDI3571 gene cluster				

METDI3569	arcB	ornithine cyclodeaminase	core	2.6	< 0.001				
METDI3671	-	conserved membrane protein of unknown function	core	2.7	< 0.001				
		METDI3863-METDI3864 gene cluster							
	hana	Lit translagating a real asphate surthage	0010	2.5	0.010				
IVIE I DI3863	прри	transiocating pyrophosphate synthase	core	3.5	0.019				
IVIE I DI3804	-		specific	2.4	0.004				
	METDI4695-METDI4699 gene cluster								
	prol/	glycine/betaine/proline ABC transporter,	variable	2.2	0.001				
IVIE I D14095	ριον	ATP-binding component	Valiable	2.5	0.001				
METDI4696	opuAB	glycine/betaine/proline ABC transporter, membrane component	variable	2.0	0.016				
METDI4699	-	transcriptional regulator, AraC family	core	4.3	< 0.001				
		METDI5067-METDI5068 gene cluster							
METDI5067	_	conserved protein of unknown function	core	3.0	0.002				
METDI5068	_	nutative transcriptional regulator	variable ^e	3.0	0.002				
List of source man			Variable	5.2	0.001				
List of genes mor	e abundant d	on methanol	1						
METDI1517	-	conserved membrane protein of unknown function	core	-2.3	< 0.001				
METDI1697	-	putative formate/nitrate transporter	variable	-3.4	< 0.001				
METDI1842	-	conserved protein of unknown function	core	-2.1	0.005				
METDI2103	-	conserved protein of unknown function	core	-2.4	0.003				
METDI3157	glnII	glutamine synthetase, type II	core	-3.2	0.008				
METDI3835	dctA	C ₄ -dicarboxylate transport protein	core	-3.2	0.003				
METDI4261	-	conserved protein of unknown function	core	-2.6	0.001				
METDI5284	-	membrane protein of unknown function	core	-3.1	< 0.001				
p2METDI0004	-	conserved protein of unknown function	specific	-3.7	0.002				
p2METDI0005	-	protein of unknown function	specific	-3.1	0.005				
		METDI1978-METDI1985 gene cluster							
METDI1978	_	conserved protein of unknown function	core	-2.1	0.004				
		conserved protein of unknown function.							
METDI1979	-	NosX-related protein	core	-3.2	0.012				
		conserved membrane protein. FMN-binding and							
METDI1980	-	4Fe-4S binding domains	core	-2.6	0.002				
METDI1981	-	conserved exported protein of unknown function	core	-2.5	< 0.001				
METDI1982	-	conserved exported protein of unknown function	core	-3.3	< 0.001				
METDI1983	-	cvtochrome c550	core	-3.8	< 0.001				
METDI1984	-	conserved periplasmic protein of unknown function	core	-3.8	< 0.001				
METDI1985	mdh2	PQQ-dependent methanol/ethanol dehydrogenase	core	-1.9	0.301				
		GEI 160	1		I				
1457010055					0.004				
METDI0255	-	RND efflux transporter, MFP subunit	specific	-2.1	0.084				
METDI0260	-	glucokinase (C-terminal fragment)	variable	-4.3	< 0.001				
METDI0262	-	glucokinase (N-terminal fragment)	variable	-2.6	0.016				
METDI0263	-	protein of unknown function	specific	-2.7	0.003				
METDI0266	-	putative ion transport domain	variable	-3.1	< 0.001				
METDI0267	-	conserved protein of unknown function	variable	-2.8	0.007				
METDI0270	-	protein of unknown function	variable	-2.6	0.094				
METDI0275	-	putative cytochrome c	variable	-2.9	0.045				
METDI0283	adh-like	putative alcohol dehydrogenase	variable	-3.3	0.029				
METDI0285	-	protein of unknown function	specific	-3.3	0.048				
METDI0286	xfp	D-xylulose 5-phosphate/D-fructose 6-phosphate phosphoketolase	variable	-2.9	0.098				
METDI0288	сірВ	protein disaggregation chaperone	core	-3.6	0.014				
METDI0289	-	putative heat shock protein DnaJ	specific	-3.9	0.008				
METDI0290	-	DnaJ-associated protein of unknown function	specific	-4.0	0.001				
METDI0292	phbC	putative poly-beta-hydroxyalkanoate synthase	specific	-3.1	0.085				
METDI0293	fabl	enoyl-[acyl-carrier-protein] reductase (NADH)	specific	-3.6	0.046				
METDI0294	-	putative phosphate butyryltransferase	specific	-3.6	0.046				
METDI0295	-	putative acetate kinase (AckA)	specific	-3.4	< 0.001				
METDI0300	-	protein of unknown function	specific	-2.6	0.029				
METDI0301	-	cation-transporting A Pase	variable	-3.3	0.077				
METDI0306	-	protein of unknown function	specific	-2.7	< 0.001				
METDI0322	-	putative cytochrome c, class l	specific	-2.2	0.034				
METDI0329	-	KND efflux transporter, MFP subunit	specific	-3.0	0.062				
METDI0330	-	RND ettlux transporter, HME family, translocase subunit	variable	-3.0	< 0.001				
METDI0333	-	putative haloacid dehalogenase-like hydrolase	specific	-2.7	< 0.001				
METDI0334	-	exported protein of unknown function	specific	-2.2	0.017				
METDI0349	-	putative protein disulfide reductase/isomerase, putative	specific	-2.4	0.057				
		thioredoxin							
METDI0363	-	putative cytochrome c oxidase, subunit l	specific	-2.6	0.011				
METDI0364	-	putative cytochrome c oxidase, subunit II (CbaB)	specific	-3.4	< 0.001				
METD10365	-	conserved membane protein of unknown function	specific	-3.3	< 0.001				

METDI0366	_	nutative sytoshrome s	specific	-29	0.001				
METDIO267	-	putative cytochrome c	specific	-2.5	< 0.001				
	-	putative cytochionie c	specific	-2.7	< 0.001				
IVIE I DIU384	-		specific	-2.5	0.004				
METDI0386	-	histone deacetylase family protein	variable	-2.9	0.029				
		METDI0537-METDI0538 gene cluster ^e							
		ABC transporter, fused tandem ATPase							
METDI0537	-	and permease domains	core	-2.5	0.059				
METDI0538	-	ABC-2 transporter, permease	core	-2.0	0.073				
METDI1414	ardC	antirestriction protein (ArdC)	specific	-2.2	0.026				
METDI1416	-	transposase of ISMdi10, IS110 family	specific	-3.7	0.008				
METDI1418	-	exported protein of unknown function	specific	-3.5	0.059				
METDI1419	-	protein of unknown function	specific	-3.6	0.046				
METDI1420	-	protein of unknown function	specific	-2.9	0.084				
METDI1446	-	protein of unknown function	specific	-2.8	0.003				
		METDI1753-METDI1770 gene cluster							
				2.2	0.001				
MEIDI1/53	eshA	eshA	core	-3.3	< 0.001				
METDI1754	-	putative cysteine desulfurase (SufS domain)	variable	-2.2	0.094				
METDI1761	-	conserved exported protein of unknown function	variable	-2.6	< 0.001				
METDI1762	-	conserved protein of unknown function, DUF1236	variable	-2.1	0.007				
METDI1769	-	conserved protein of unknown function	core	-3.7	0.034				
METDI1770	ohyA	oleate hydratase	variable	-3.4	0.043				
		METDI2004-METDI2012 gene cluster ^e							
		concerned protein with ANALL like domain	0010	2.2	0.046				
	-		COTE	-5.5	0.040				
IVIET DI2005	-		core	-3.2	0.042				
METDI2007	-	ABC transporter, fused ATPase and permease domains (CydD-like)	core	-3.4	< 0.001				
METDI2009	-	ABC transporter related, fused ATPase and permease domains	core	-3.3	< 0.001				
		(CydD-like)	00.0	0.0					
METDI2011	cydB	cytochrome d terminal oxidase, subunit II	core	-3.4	0.036				
METDI2012	-	putative exported cyd operon protein	specific	-3.7	< 0.001				
METDI2867-METDI2874 gene cluster									
METDI2867	alnK	nitrogen regulatory protein P-II	core	-25	0.029				
	fdh AD	formate debudrogenase subunit P	core	2.5	< 0.025				
IVIET DI2873	JUN4B		core	-3.2	< 0.001				
IVIE I DI2874	Jun4A	Tormate denydrogenase subunit A	core	-4.0	< 0.001				
		METDI4303-METDI4313 gene cluster ^e							
METDI4303	-	putative transcriptional regulator, Crp/Fnr family	variable	-3.6	0.048				
METDI4304	-	conserved protein with ANAH-like domain	core	-3.1	0.054				
METDI4305	-	conserved protein of unknown function	variable	-2.4	0.008				
METDI4307	-	protein of unknown function	variable	-2.7	0.034				
MFTDI4308	-	protein of unknown function	variable	-3.4	< 0.001				
METDI4310	_	protein of unknown function	specific	-2.7	0.003				
	1		speeme	2.7	0.000				
	1	GEI 197	n						
METDI4470	-	putative L,D-transpeptidase catalytic domain (YkuD)	specific	-2.3	< 0.001				
METDI4503	-	putative cytochrome c oxidase, subunit I	specific	-2.4	0.008				
METDI4504	-	putative cytochrome c oxidase, subunit II	specific	-3.0	< 0.001				
METDI4505	-	conserved membrane protein of unknown function	specific	-2.5	0.034				
METDI4506	-	putative cytochrome c, putative exported protein	specific	-2.5	0.006				
METDI4507	-	putative cytochrome c	specific	-3.3	< 0.001				
METDI4514	-	putative transporter, SulP family	specific	-3.3	0.076				
METDI4515	-	histone deacetylase family protein	variable	-3.8	0.033				
METDI4517	-	protein of unknown function	specific	-2.5	0.007				
METDI4520	_	putative transcriptional regulator. Crn/Enr family	variablo	2.5	0.072				
	-	putative dialscriptional regulator, crp/rni family	variable	-3.2	0.072				
IVIET DI4522	-	putative glucokinase (C-terminal fragment)	specific	-5.5	< 0.001				
IVIE I DI4524	-		Variable	-2.0	0.035				
METDI4525	-	protein of unknown function	specific	-2.8	0.046				
IVIE I DI4526	-	conserved protein of unknown function	variable	-3.0	0.075				
METDI4527	-	putative biotin/lipoyl attachment domain	variable	-2.8	0.054				
METDI4528	-	putative ion transport domain	variable	-3.1	< 0.001				
METDI4530	-	conserved protein of unknown function	variable	-3.0	0.034				
METDI4531	-	putative manganese transporter	variable	-3.3	0.051				
METDI4536	-	protein of unknown function	variable	-2.4	0.011				
METDI4538	-	putative cytochrome c, class I	variable	-2.8	0.034				
METDI4540	adh-like	putative alcohol dehydrogenase	variable	-3.7	0.011				
METDI4543	xfp	D-xylulose 5-phosphate/D-fructose 6-phosphate phosphoketolase	variable	-3.1	0.085				
METDI4544		nutative acetate kinase (nartial)	variable	-5.0	< 0.001				
MFTDI4545	-	protein of unknown function	variable	-3.6	0 044				
	1		variable	5.0	0.044				

METDI4547	-	protein of unknown function	specific	-3.3	0.003
METDI4548	-	conserved protein of unknown function	variable	-3.2	0.083
METDI4552	-	protein of unknown function	variable	-3.5	0.036
METDI4553	-	exported protein of unknown function	variable	-2.9	0.003
METDI4554	-	putative cation-transporting ATPase (P-type)	variable	-3.5	0.059
METDI4555	-	conserved protein with ANAH-like domain	variable	-3.3	0.001
METDI4557	-	exported protein of unknown function	variable	-3.4	< 0.001
METDI4563	-	conserved protein with ANAH-like domain	variable	-3.3	0.051
METDI4564	-	putative cytochrome c, class I	variable	-2.6	0.003
METDI4565	-	conserved protein with ANAH-like domain	variable	-2.9	0.051
		METDI4813-METDI4816 gene cluster			
METDI4813	-	RND efflux transporter, MFP subunit	core	-3.6	< 0.001
METDI4814	-	exported protein of unknown function	variable	-4.0	0.001
METDI4815	-	protein of unknown function	variable	-2.7	0.003
METDI4816	qxtB	cytochrome bd-quinol oxidase subunit II	core	-2.7	0.061

METDI5532-METDI5533 gene cluster

METDI5532	nrtB	nitrate transport permease protein	core	-2.4	0.001
METDI5533	nrtA	nitrate transporter component	core	-2.7	< 0.001

^a MaGe annotation (https://www.genoscope.cns.fr/agc/microscope)

^b Genes labeled as core, variable or specific are respectively found in all, at least 2 genomes of *M. extorquens* genomes of strains AM1, BJ001,

CM4, DM4 and PA1 (80 % of identity on at least 80 % of the protein length), or only in the genome of strain DM4.

^c log2 fold-change of normalized read numbers in cultures grown with dichloromethane compared to methanol.

^d False discovery rate, only adjusted p-value under 0.1 were considered.

^e Adjacent to dichloromethane-enhanced gene encoding a protein of unknown function

^f Using 2D gel proteomics of cell extracts of dichloromethane- versus methanol-grown *M. extorquens* DM4, proteins detected as more abundant on dichloromethane (METDI2655 and METDI2656) and less abundant on methanol (METDI0320) (Muller *et al.*, 2011)

3. <u>Supplemental data</u>

3.1. Defining the core genome

M. extorquens CM4 and DM4 are the only known M. extorquens strains that have been described in the literature to be able to utilize chlorinated compounds. Conversely, all *M. extorquens* strains are able to grow with methanol as sole carbon and energy source (Doronina *et al.*, 1996; Vuilleumier *et al.*, 2009). Thus, the two dechlorinated methanes may share specific gene sets involved in adaptation to growth with either chloromethane or dichloromethane but not with methanol. Comparative genomics of assembled reference genome sequences of five *M. extorquens* strains AM1, BJ001, CM4, DM4 and PA1 (Vuilleumier et al., 2009; Marx et al., 2012) aimed to identify the genes of the core genome. The core genome is defined as the set of genes (orthologous genes) shared by all the strains of the same bacterial species. A set of 3,489-shared CDS were defined for genes encoding for proteins with at least 80 % of amino acid identity on 80 % of the CDS length. The two dechlorinating strains, the chloromethane-degrading strain CM4 and the dichloromethane-degrading strain DM4, share a total of 4,620 genes as found using the MaGe plateform "Phyloprofile Exploration" tool (https://www.genoscope.cns.fr/agc/microscope/home/). Among those, 99 % are encoding for proteins with at least 90 % of amino acid identity on 80 % of the CDS length (data not shown). Within the MaGe plateform, "Mchl" label refer to genetic element of strain CM4 basing on its historical first taxonomical name: Methylobacterium chloromethanicum (Doronina et al., 1996). Similarity, genes in the genome of strain DM4 are labelled as "METDI" for Methylobacterium extorquens dichloromethane-degrading strain. The correspondence of "Mchl"/"METDI" names (except for transposable elements) is listed in Table 3.1 (https://seafile.unistra.fr/f/a924b1b359/).

3.2. RNAseq data of the variable genome only shared by dechlorinating strains

The variable genome exclusively shared by the genomes of the dechlorinating strains included 163 genes (listed in Table 3.2) with 67 plasmid-borne genes with no predicted functions besides those related to plasmids (replication, integrase, transfert). They were located on two plasmids: pCMU01 in *M. extorquens* CM4 and p1METDI in *M. extorquens* DM4. The extent of relatedness between these plasmids is remarkable since 47 % of p1METDI-borne CDS have a homolog on plasmid pCMU01 in three synteny regions (39 genes; (Roselli, 2009)). A few

chromosomal genes had predicted function in central metabolism (oxidoreductase, ATPhydrolyzing enzyme, succinyl CoA synthetase), and stress response (beta-lactamase, esterase) (Table 3.2). Using intraspecies comparative RNAseq analysis, no gene shared only by *M. extorquens* CM4 and DM4 were detected as differentially abundant in cultures grown with chlorinated methanes compared to methanol.

				M. extorquens		M. extorquens M. extorquens	
CM4 label	DM4 label	Gene	Product	C	M4	D	M4
					Adjusted	log2fc	Adjusted
					p-value ^b		p-value
Mchl_0111	METDI4323	-	protein of unknown function	0.7	0.140	-0.9	0.530
Mchl_0185	METDI0124	-	protein of unknown function	1.2	0.014	1.4	1.000
Mchl_0225	METDI4784	-	protein of unknown function	1.3	<0.001	1.6	0.130
Mchl_0318	METDI4325	-	protein of unknown function	0.4	0.450	-0.7	0.700
Mchl_0688	METDI0657	-	fragment of putative flagellin (FlaA-like; FlaB-like)	-1.0	<0.001	-0.9	0.830
Mchl_0906	METDI1089	-	putative OsmB-like lipoprotein	0.4	0.340	1.0	0.770
Mchl_0969	METDI1157	-	phage integrase	0.1	0.920	0.8	0.580
Mchl_0970	METDI1158	-	phage integrase	-1.0	<0.001	-0.2	0.990
Mchl_0971	METDI1159	-	putative phage-associated protein	-0.8	0.047	-1.0	0.550
Mchl_0974	METDI1162	-	putative ferrichrome-iron receptor protein	-0.1	0.940	-0.3	0.970
Mchl_0975	METDI1163	-	putative exported esterase (IroE-like)	0.4	0.670	-0.2	1.000
Mchl_0976	METDI1164	-	protein of unknown function	0.2	0.730	0.8	0.590
Mchl_0977	METDI1165	-	protein of unknown function	-0.2	0.740	-0.6	0.840
Mchl_0979	METDI1168	-	putative resolvase/recombinase	0.5	0.280	0.5	0.900
Mchl_0980	METDI1170	-	putative resolvase/recombinase	0.5	0.200	0.5	0.900
Mchl_0982	METDI1174	-	putative phosphohydrolase/phosphoesterase	0.5	0.470	-0.1	1.000
Mchl_0983	METDI1175	-	putative transcriptional repressor	0.5	0.430	-1.3	0.160
Mchl_0984	METDI1180	-	protein of unknown function	-0.1	0.990	0.4	0.850
Mchl_0985	METDI1181	-	putative ATP-hydrolyzing enzyme	0.5	0.380	-0.1	1.000
Mchl_0988	METDI1314	-	fragment of protein of unknown function	0.3	0.720	0.4	0.860
Mchl_0990	METDI1315	-	protein of unknown function	0.1	0.980	0.4	0.890
Mchl_1005	METDI1336	-	protein of unknown function	0.6	0.480	1.4	0.240
Mchl 1022	METDI1354	-	protein of unknown function	0.9	0.011	1.2	0.770
Mchl 1037	METDI1369	-	protein of unknown function	1.1	< 0.001	0.7	0.910
	METDI1216	-	protein of unknown function	0.6	0.590	-0.3	0.970
	METDI1282	-	conserved exported protein of unknown function	0.4	0.620	0.9	0.570
_ Mchl 1136	METDI1461	opdE	transcription regulatory protein	0.1	0.960	-0.4	0.930
Mchl 1137	METDI1462	-	putative short-chain dehydrogenase/reductase SDR	0.1	1.000	-0.3	0.970
 Mchl 1138	METDI1463	-	conserved exported protein of unknown function	0.1	0.970	-0.1	1.000
Mchl 1141	METDI1467	-	putative short-chain dehydrogenase/reductase SDR	-0.4	0.690	-0.7	0.850
	METDI1468	-	putative transcriptional regulator. LysR family	-0.1	0.940	0.3	0.930
Mchl 1143	MFTDI1469	-	oxidoreductase. 2Ee-2S subunit	-0.2	0.810	0.7	0.610
Mchl 1144	METDI1470	-	oxidoreductase. FAD-binding subunit	0.1	0.960	0.8	0.500
Mchl 1145	MFTDI1471	-	oxidoreductase, molybdopterin-binding subunit	0.2	0.800	0.6	0.750
Mchl 1147	MFTDI1473	-	protein of unknown function	0.4	0.530	0.5	0.820
Mchl 1259	METDI1666	-	putative transcriptional regulator. LysR family	-0.1	0.900	-0.1	1.000
			ABC transporter, periplasmic protein: putative taurine	0.1	0.000	0.12	1.000
Mchl_1260	METDI1667	-	transporter	-0.6	0.280	-0.2	0.980
			ABC transporter permease component: putative				
Mchl_1261	METDI1668	-	tauring transporter	-0.5	0.520	-0.1	1.000
Mcbl 1262	METDI1660		APC transporter ATPace putative tauring transporter	0.6	0.420	0.2	0.060
Mcbl 1264	METDI1671	-	alutamyl-tRNA(Gln) amidotransforaça subunit A	-0.0	1 000	-0.1	1 000
Mcbl 1490	METDI1071		motal dependent phosphobudrolase Subuliit A	1 7	<1 000	0.1	1.000
Machi 1494		-	transcriptional regulator. Are C family	1./	<1.000	0.1	1.000
Nabl 1481		-	u anscriptional regulator, Arac family	-0.2	0.890	0.4	0.680
Nichi 1482		-		0.3	0.770	-0.9	0.030
Nichi_1484	IVIETDI1927	-	protein of unknown function	0.8	0.048	0.3	0.9940
IVICNI_1493	INIE I DI 1932	-	conserved protein of unknown function	0.1	0.830	0.3	0.930

Table 3.1. RNAsed	ı data of ge	enes shared	exclusively by	v M. ext	orauens CM4	and DM4

Mchl_2249	METDI2736	-	protein of unknown function	-0.3	0.730	0.2	0.970
Mchl_2370	METDI2862	-	protein of unknown function	0.5	0.760	-0.9	0.670
Mchl_2863	METDI3394	-	protein of unknown function	0.5	0.280	-0.1	1.000
Mchl_3241	METDI3788	-	conserved protein of unknown function	0.4	0.390	0.1	0.990
			putative sensory box-containing diguanylate		0.000	0.7	0.070
IVICNI_3333	METDI0326	-	cyclase/phosphodiesterase	0.4	0.390	0.7	0.870
Mchl 3438	METDI3901	-	conserved protein of unknown function	0.1	0.950	-0.3	0.960
Mchl 3466	METDI3928	-	putative addiction module antidote protein	-0.6	0.300	0.1	0.990
 Mchl 3840	METDI4356	-	phage-related integrase (fragment)	0.9	0.930	0.3	0.930
Mchl 3913	MFTDI4612	-	conserved protein of unknown function	0.5	0.510	-0.4	0.900
Mcbl 4141	METDI4814	-	protein of unknown function	0.9	0.096	-0.4	0.0001
Mcbl 4205	METDI0774	_	protein of unknown function	-0.1	0.000	0.5	0.0001
Mcbl 4205	METDI0775	sucD	suscing CoA synthetase, alpha subunit	0.1	0.120	1.7	0.000
Mcbl 4200		SUCC	succinyl CoA synthetase, apria subunit	0.5	0.130	1.7	0.022
Mchl 4207	METDI0777	Succ	nutativo succinato comialdobudo dobudrogonaco	0.0	0.320	1.5	0.030
Mahl 4208		-	putative succinate-semilaidenyde denydrogenase	0.0	0.290	1.4	0.120
Mchi_4209	METDI0778	-		0.5	0.010	1.9	0.008
Mchi_4211	METDI0781	-		-0.1	0.910	0.2	0.970
Wichi_4212	METDI0782	-	MFS transporter membrane protein	0.1	0.980	0.1	0.840
Mchl_4214	METDI0788	XSC	sulfoacetaldehyde acetyltransferase	-0.2	0.780	0.1	1.000
Mchl_4215	METDI0789	-	DNA-binding transcriptional repressor, IclR family	0.1	0.780	0.4	0.980
Mchl_4216	METDI0790	-	putative acetate kinase (AckA)	0.2	0.960	0.1	0.890
Mchl_4341	METDI4965	-	conserved protein of unknown function	0.7	0.092	0.1	1.000
Mchl_4342	METDI4966	-	conserved protein of unknown function	0.5	0.230	0.4	0.890
Mchl_4350	METDI4974	-	conserved protein of unknown function	0.6	0.200	0.7	0.670
Mchl_4444	METDI5071	-	protein of unknown function	-0.2	0.810	-0.5	0.840
Mchl_4550	METDI5180	-	conserved protein of unknown function	0.2	0.830	0.4	0.900
Mchl_4551	METDI5181	-	conserved protein of unknown function	0.1	0.930	0.5	0.900
Mchl_4659	METDI5293	-	conserved protein of unknown function	0.8	0.100	0.1	1.000
Mchl_4718	METDI5356	-	putative phage integrase	-0.1	0.840	0.4	0.880
Mchl_4719	METDI5357	-	putative transcriptional regulator	0.4	0.690	-0.4	0.960
Mchl_4720	METDI5358	-	putative metallo-beta-lactamase family protein	0.1	0.990	-0.5	0.910
Mchl_4721	METDI5359	-	conserved protein of unknown function	0.4	0.560	-0.4	0.890
Mchl_4722	METDI5360	-	protein of unknown function	0.6	0.260	0.6	0.800
Mchl_4723	METDI5361	-	protein of unknown function	0.4	0.670	0.6	0.780
Mchl_4729	METDI1222	-	protein of unknown function	0.5	0.270	1.4	0.120
Mchl_4742	METDI5367	-	protein of unknown function	1.4	< 0.001	0.1	0.950
Mchl 4743	METDI5371	-	protein of unknown function	0.3	0.810	-0.4	0.930
Mchl 4744	METDI5372	-	putative DNA helicase related protein	0.1	0.860	1.4	0.590
	METDI5375	-	putative ardC antirestriction protein	1.1	0.260	-1.3	0.690
	METDI2680	-	putative SAM-dependent methyltransferase domain	0.5	0.290	0.1	1.000
	METDI1912	-	protein of unknown function	0.6	0.280	1.1	0.370
	METDI0315/		P			1.1/	0.300/
Mchl_5447	METDI0325	-	transposase	1.2	0.001	0.9	0.460
Mchl 5456	METDI4486	-	protein of unknown function	0.7	0.190	-0.3	0.960
Mchl 5457	METDI4485	isnF	2C-methyl-D-erythritol 2 4-cyclodinhosphate synthase	-1 4	<0.001	-1 7	0.150
		1501	nutative nseudouridine synthase nutative 1-deoxy-D-		.01001		0.1200
Mchl_5458	METDI4484	-	xylulose 5-phosphate reductoisomerase	-1.6	<0.001	-1.8	0.260
Mcbl 5/159		adhA	alutamate debudrogenase (NAD(P)+) oxidoreductase	_1 3	<0.001	_1 /	0.650
Mchl 5461	METDI4405	guna		-1.9	<0.001	-1.1	0.000
Wichi_3401	WILT D14401	-	putative oxidoreductase	-1.0	<0.001	-1.1	0.780
Mchl_5546 ^c	p1METDI0092	- :	conserved protein of unknown function	1.4/1	0.056	0.2	0.920
			anatain of unlances for ation	.4	0.005	0.2	1 000
Nichi_5549°		-	protein of unknown function	1.4	0.085	0.2	1.000
IVICNI_5552°		-	conserved protein of unknown function	0.5	0.570	-0.1	1.000
IVICNI_5555°	p1WE1DI0102	-	conserved protein of unknown function	1.0	0.082	-0.3	0.940
Mchl_5558°	p1METDI0104	ardC	antirestriction protein ArdC	0.7	0.240	0.5	0.750
Mchl_5559°	p1METDI0105	-	conserved protein of unknown function	1.3	0.016	-0.6	0.091
Mchl_5564 ^c	p1METDI0110	-	conserved protein of unknown function	0.9	0.100	-0.4	0.970
Mchl_5567°	p1METDI0111	-	putative methylase/helicase	0.9	0.041	-0.5	0.940
Mchl_5568°	p1METDI0112	-	conserved protein of unknown function	1.0	0.045	1.1	0.220
Mchl_5569°	p1METDI0113	-	conserved protein of unknown function	1.1	0.052	-0.3	0.960
Mchl_5571 ^c	p1METDI0117	-	protein of unknown function	1.8	0.450	-0.2	1.000
Mchl_5572°	p1METDI0118	traG	conjugal transfer protein	-0.6	0.170	-0.1	1.000
Mchl_5573 ^c	p1METDI0119	traD	conjugal transfer protein	-0.8	0.048	-0.2	0.990

Mchi_5575	p1METDI0121	-	putative conjugal transfer protein (TraA)	0.5	0.370	-0.8	0.650
Mchi_5578	p1METDI0122	-	protein of unknown function	0.4	0.470	-0.1	1.000
Mchl_5579°	p1METDI0123	-	protein of unknown function	0.6	0.170	-0.1	1.000
Mchl_5591 ^c	p1METDI0125°	-	conserved membrane protein of unknown function	1.1	0.011	-0.4	1.000
Mchl_5592°	p1METDI0126 ^c	істВ	putative IcmB/DotO-related protein	0.7	0.082	-0.8	0.094
Mchl_5595°	p1METDI0129°	icmE	putative IcmE-related protein	0.6	0.210	-0.4	0.980
Mchl_5596 ^c	p1METDI0130 ^c	істК	putative IcmK-related protein	0.4	0.370	-1.1	0.620
Mchl_5597°	p1METDI0131 ^c	icmL	conserved protein of unknown function; putative IcmL	0.0	0.950	-1.1	0.780
Mchl_5600°	p1METDI0134 ^c	-	protein of unknown function	-0.0	1.000	-1.0	0.840
Mchl_5601 ^c	p1METDI0135 ^c	-	protein of unknown function	0.9	0.190	-0.3	0.990
Mchl_5602°	p1METDI0136 ^c	-	putative DotC-related protein	0.4	0.490	-0.3	1.000
Mchl_5603 ^c	p1METDI0137 ^c	-	putative Dot/Icm secretion system ATPase DotB	0.3	0.640	-0.2	1.000
Mchl_5604 ^c	p1METDI0138 ^c	-	conserved exported protein of unknown function	0.8	0.087	-0.2	0.990
Mchl_5605°	p1METDI0139 ^c	-	protein of unknown function	0.8	0.034	-0.9	0.610
Mchl_5606°	p1METDI0140 ^c	-	conserved protein of unknown function	0.9	0.009	-0.6	0.930
Mchl_5607°	p1METDI0141 ^c	-	protein of unknown function	1.1	0.021	-0.7	0.880
Mchl_5609	p1METDI0002 ^c	-	conserved protein of unknown function	1.0	0.004	-0.2	1.000
Mchl_5610 ^c	p1METDI0003 ^c	-	conserved exported protein of unknown function	1.2	< 0.001	-0.3	0.960
Mchl_5611 ^c	p1METDI0004 ^c	-	protein of unknown function	1.2	<0.001	-0.1	1.000
Mchl_5612 ^c	p1METDI0005 ^c	-	protein of unknown function	1.3	< 0.001	-0.4	0.960
Mchl 5613 ^c	p1METDI0006 ^c	-	protein of unknown function	1.6	< 0.001	-0.2	1.000
Mchl 5614 ^c	p1METDI0007 ^c	-	fragment of transposase related to IS481 family	1.6	< 0.001	0.3	0.940
_ Mchl 5615	p1METDI0008°	repA	plasmid partitioning protein RepA	0.4	0.380	0.4	0.120
Mchl 5616 ^c	p1METDI0009°	repB	plasmid partitioning protein RepB	0.5	0.200	0.5	0.960
	p1METDI0010 ^c	repC	replication protein C	0.4	0.340	0.4	1.000
Mchl 5620 ^c	p1METDI0111°	-	conserved protein of unknown function (fragment)	0.3	0.750	0.3	0.960
Mchl 5622	p1METDI0112	-	conserved protein of unknown function	1.2/1.3	1.000	1.0	0.710
Mchl 5644	p1METDI0053	-	conserved protein of unknown function. PilT domain	-0.3	0.620	-0.3	0.770
Mchl 5646	p1MFTDI0052	-	putative phage integrase	0.9	0.020	0.4	0.870
MCv2 0417	MFTDI0198	-	protein of unknown function	0.0	0.990	03	0.930
MCv2 0842	METDI0940	-	protein of unknown function	-0.5	0.380	-0.1	1 000
MCv2 1071	METDI1130	_	protein of unknown function	0.5	0.300	0.1	0.750
MCv2_1071	METDI1150	_	protein of unknown function	-0.1	0.710	_0.1	0.750
MCv2_1101	METDI1171	-	consonved protoin of unknown function	0.1	0.550	0.1	1 000
MCv2_1114	METDI1171		protoin of unknown function	1 1	0.000	0.1	0.020
MCv2_1113	METDI1172	-	protein of unknown function	0.5	0.624	0.3	0.550
MCv2_1120	METDI1220		protein of unknown function	0.5	0.020	0.5	0.540
MCv2_1140	METDI1220	-	protein of unknown function	-0.2	0.940	-0.0	0.750
MCv2_1145	METDI1333		protein of unknown function	0.2	0.370	1.1	0.000
MCv2 1217	METDI1202	-	protein of unknown function	0.3	0.780	0.4	0.430
MCv2_1216	METDI1203	-	protein of unknown function	0.2	0.800	0.4	0.900
MCv2_1510	METDI1474	-	concerved protein of unknown function	-0.5	0.790	-0.2	0.970
MCv2_1584	METDI1/9/	-	conserved protein of unknown function	1.5	0.005	-0.7	0.740
WCV2_1680	IVIE I DI 1888	-	protein of unknown function	0.8	0.350	0.2	0.950
MCv2_1703	METDI1921	-	putative phosphoglycerate/bisphosphoglycerate	0.2	0.750	-0.2	0.970
MC-2 1720	METDIADEO		mutase	1.1	0.010	0.2	0.000
MCV2_1739	METDI1950	-	protein of unknown function	1.1	0.018	-0.3	0.960
MCV2_1740	METDI1951	-	protein of unknown function	0.8	0.490	-0.1	1.000
MCV2_1741	METDI1952	-	conserved protein of unknown function	-0.1	1.000	-0.5	0.900
MCv2_1/51	METDI1961	-	conserved exported protein of unknown function	-0.3	0.740	-0.7	0.640
MCv2_2296	METDI2356	-	conserved protein of unknown function	0.7	0.320	0.5	0.840
MCv2_2310	METDI2369	-	protein of unknown function	0.3	0.760	-0.6	0.840
MCv2_2529	METDI2709	-	protein of unknown function	0.3	0.660	-0.5	0.810
MCv2_2712	METDI2882	-	protein of unknown function	1.3	0.060	0.8	0.580
MCv2_2972	METDI3136	-	protein of unknown function	0.5	0.460	0.5	0.820
MCv2_3197	METDI3358	-	protein of unknown function	0.3	0.720	0.2	0.950
MCv2_3231	METDI3407	-	conserved protein of unknown function	0.3	0.670	0.4	0.850
MCv2_3537	METDI3709	-	protein of unknown function	0.3	0.600	-0.8	0.580
MCv2_4420	METDI5044	-	protein of unknown function	0.1	0.980	0.8	0.590
MCv2_4813	METDI4912	-	protein of unknown function	-0.5	0.590	-0.3	0.940
MCv2_5307	METDI5397	-	protein of unknown function	0.2	0.900	-1.7	0.070
MCv2_5403	METDI4380	-	conserved protein of unknown function	1.0	0.046	0.1	1.000

- ^a log2 fold-change (log2fc) values ≥ 2 or ≤ -2 mean that transcript abundance is higher in cultures grown with chlorinated methanes compared to methanol, or higher with methanol compared to chlorinated methanes, respectively.
- ^b False discovery rate, only adjusted p-value under 0.1 were considered.
- ^c In synteny region between plasmid pCMU01 and plasmid p1METDI.

3.3. RNAseq data of plasmid pCMU01-borne gene with chromosomal homologs

Among plasmid-borne homologs, 28 out of 98 had transcript found as more abundant in cultures grown with chloromethane compared to methanol (Table 3.3; no homologs found as differentially less abundant). Among those, 24 were involved in biosynthesis of the essential chloromethane dehalogenase corrinoid compound (Studer *et al.*, 2001, 2002) (see section 3.2 article). Similarly, of the homologs encoding cobalt and cobalamin transporters, only copies located on plasmid pCMU01 has transcript with increased abundant in cultures grown with chloromethane compared to methanol, unlike core genome homologs (*czcA*, encoding a RND transporter or some transport proteins encoded by *tonB*, *exbB*/*exbD*, *tolQ*/*tolR* and *btuFCD*). This suggests that a yet uncharacterized chloromethane-dependent regulation mechanism favors the expression of genes of the variable genome over homologous genes of the core genome.

Homolog name & localization ^a		- Gono Broduct		aa Id	Plasmid pCMU01		Chromosome	
pCMU01	Chromosome	Gene	e Product (log2fc ^b	Adjusted p-value ^c	log2fc	Adjusted p-value
Mchl_5400	Mchl_2934	-	putative nitrile hydratase activator protein	77.3	1.3	0.050	-0.6	0.700
Mchl_5405	Mchl_3481	-	<pre>putative H(+)/Cl(-) exchange transporter ClcA</pre>	68.0	5.5	< 0.001	0.3	0.700
Mchl_5435	Mchl_1555	-	putative universal stress protein, UspA-like (fragment)	58.7	1.0	0.050	0.5	0.300
Mchl_5437	Mchl_4097	-	integrase family protein	58.4	-0.1	0.940	-0.1	0.960
Mchl_5444	Mchl_2145	xoxF2	putative PQQ-linked dehydrogenase of unknown specificity	88.2	0.9	0.019	0.5	0.240
Mchl_5445	Mchl_3593	-	sensor histidine kinase	40.6	0.9	0.047	-0.1	0.880
Mchl_5446	Mchl_3594	-	putative response regulator; CheY and HTH LuxR family	64.4	1.0	0.074	-0.2	0.890
Mchl_5455	Mchl_2361	-	putative cystathionine beta-synthase	82.1	1.0	0.006	-0.3	0.640
Mchl_5457	Mchl_3044	ispF	2C-methyl-D-erythritol 2,4-cyclodiphosphate synthase	94.5	-1.4	<0.001	0.1	0.990
Mchl_5460	Mchl_0670	serC2	phosphoserine aminotransferase	95.2	-1.2	<0.001	-0.6	0.150
Mchl_5461	Mchl_2219	-	putative oxidoreductase	52.7	-1.8	0.001	-0.2	0.740
Mchl_5462	Mchl_2317	hss2	homospermidine synthase	96.4	-1.4	0.001	-0.6	0.210
Mchl_5475	Mchl_0044	-	putative sensor protein (fragment)	45.7	0.8	0.091	1.4	< 0.001
	Mchl_0272			61.3			0.6	0.130
Mchl_5476	Mchl_0517	-	putative sensor protein (fragment)	46.2	1.2	0.009	1.0	0.034
	Mchl_1530			56.2			-0.1	1.000
Mchl_5478	Mchl_2189	-	conserved protein of unknown function	69.0	1.0	0.015	-0.7	0.110
Mchl_5491	Mchl_2220	shc2	squalene-hopene cyclase	56.7	-0.1	0.970	-0.5	0.430
Mchl_5495	Mchl_3651	-	RND efflux transporter, MFP subunit	47.8	0.9	0.270	-0.5	0.250
Mchl_5496	Mchl_1298 Mchl_3650	-	RND efflux transporter, HME family, translocase subunit	40.8 60.2	0.3	0.540	0.5 -0.7	0.320 0.140

Table 3.2. RNAsed	data for	oCMU01-borne	genes with	chromosomal	homologs

	Mchl_4139			40.3			0.4	0.630
	Mchl_3285			40.1			0.1	0.970
Mchl 5499	Mchl 1419	-	response regulator in two-component regulatory system	42.2	2.4	< 0.001	0.3	0.700
	Mchl 2980			40.2			0.2	0 750
Mahl 5502	Mchl 1050		concorried protein of unknown function	70.2	1 2	0 1 2 0	0.2	0.750
IVICIII_5502	Mahl 0550	-	conserved protein of unknown function	12.2	1.5	0.120	-0.5	0.470
Mchl 5504	IVICNI_0559	-	putative di-heme cytochrome c peroxidase (MauG-related)	41.3	1.0	0.021	0.3	0.650
-	Mchl_2999			40.3			-0.1	1.000
Mchl_5509	Mchl_4119	-	conserved exported protein of unknown function	45.4	0.9	0.041	0.4	0.550
Mchl_5512	Mchl_5137	-	putative molybdopterin molybdenumtransferase (MoeA)	42.3	0.7	0.350	0.2	0.790
		14.0	molybdate ABC transporter, periplasmic binding	64 0				0.050
Mcni_5514	IVICNI_4941	moaA2	component	61.2	0.6	0.140	0.1	0.950
Mchl 5515	Mchl 3926	modB2	molybdate ABC transporter, membrane component	75.4	0.4	0.490	-0.1	0.720
Mchl 5516	Mchl 3928	modC2	molyhdate ABC transporter. ATP-hinding component	55.7	0.6	0 210	0.1	0 520
			molybdenum-nterin-binding protein, nutative molybdate	0017	0.0	0.220	0.1	0.020
Mchl_5517	Mchl_2845	mop2	homoostasis factor	69.6	1.0	0.003	0.1	0.890
	M.L. 2274			44.0	1.0	.0.001	0.4	1 000
Wichi_5520	IVICNI_2371	-	conserved exported protein of unknown function	44.8	1.8	<0.001	0.1	1.000
	Mchl_2536			46.4			-1.9	<0.001
	Mchl_1417			40.7			-1.0	0.015
	Mchl_1055			43.5			0.4	0.740
Mchl_5526	Mchl_1112	-	transposase of ISMex11	100	2.3	<0.001	0.7	0.410
	Mchl_1119			64.3			0.1	0.910
Mchl 5528	Mchl 2715	-	putative diguanylate cyclase (fragment)	42.1	1.0	0.240	-0.9	0.920
-	 Mchl 2736			45.5			0.1	0.920
	Mchl 2255			46.3			0.4	0.490
	Mchl 0114			59.5			0.4	0.450
Mchl_5529		-	conserved protein of unknown function	36.5	1.4	0.003	0.1	0.980
	Michi_1956			/5./			-0.2	0.850
Mchl_5531	Mchl_0082	-	putative transcriptional regulator, Ros/MucR family	46.4	1.0	0.052	0.1	0.900
	Mchl_4811			49.4			-0.2	0.730
	Mchl_4927			53.4			0.3	0.620
Mchl_5542	Mchl_2091	-	putative aminoglycoside phosphotransferase	43.4	0.6	<0.001	-0.1	< 0.001
Mchl_5544	Mchl_2409	-	putative glycosyl transferase	61.6	1.4	0.038	0.3	0.730
	Mchl 0114			59.3			0.1	0.980
Mchl_5546		-	conserved protein of unknown function	75.7	1.3	0.056	-0.2	0.850
Mcbl 5556	Mchl 5373	_	nutative methyl-accenting chemotaxis protein	46.0	0.6	0 1 2 0	1.6	<0.001
<u>_</u> 5556	Mchl 0134			16.0	0.0	0.120	0.0	0.009
		andC		40.2	0.7	0.240	0.9	0.008
	Michi_4749	arac		40.8	0.7	0.240	1.0	0.260
MCNI_5566	IVICNI_0082	-	putative transcriptional regulator	40.8	1.0	0.001	0.1	0.900
	Mchl_4938			38.7			0.5	0.240
Mchl 5581	MCv2_4479	-	transposase (fragment)	90.9	23	0.011	-0.5	0.540
<u>.</u> 5561	Mchl_1117			77	2.5	0.011	1.6	0.038
Mahl FF03	Mabl 0521		class III alcohol dehydrogenase, S-(hydroxymethyl)	F0 /	1.0	<0.001	1 5	<0.001
WICHI_5562	WICHI_0551	-	glutathione dehydrogenase	56.4	1.0	<0.001	1.5	<0.001
	Mchl_2582			61.5			0.6	0.280
	Mchl 4105			59.2			0.5	0.380
Mchl 5618		-	protein of unknown function	69.4	2.2	< 0.001	0.1	0.850
Mchl 5623	Mchl 1137	-	dehydrogenase/reductase (fragment)	41.6	0.6	0 590	0.1	1 000
<u>5025</u>	Mchl 5004			46.7	0.0	0.550	0.1	0.800
Mahl E627	Mchl 4222		concorring protoin of unknown function	40.7	0.0	0 100	0.1	0.890
Wichi_5627	WICHI_4252	-		07.5	0.9	0.190	0.7	0.400
WICHI_5639	IVICNI_3405	-	conserved exported protein of unknown function	79.0	1.7	<0.001	1.7	<0.001
Mchl_5646	Mchl_4007	-	putative integrase	68.8	0.9	0.023	0.6	0.470
Mchl_5667	Mchl_3540	-	putative phosphohydrolase/phosphoesterase	52.9	1.3	<0.001	0.1	1.000
	Mchl_0982			42.0			0.5	0.470
	Mchl_1897			43.8			0.3	0.280
Mchl_5669	Mchl_0978	-	putative ATP-hydrolyzing enzyme	42.1	1.0	0.032	0.3	1.000
			ABC transporter periplasmic binding component, putative					
Mchl_5677	Mchl_3153	btuF	vitamin B12 transporter (BtuF)	61.1	6.0	<0.001	0.1	0.990
			ABC transporter membrane component putative vitamin					
Mchl_5678	Mchl_3152	btuC	R12 transporter, including component, putative vitalilli	73.5	5.3	< 0.001	-0.5	0.590
	Mahl 4450		ADC transporter ATD his diag service and the list in the	40.0			0.7	0.070
Mchl_5679	ivicni_4153	btuD	ABC transporter ATP-binding component, putative vitamin	40.0	5.0	< 0.001	-0.7	0.870
_	Mchl_3151		B12 transport (BtuD)	65.9			-0.1	0.970
Mchl_5680	Mchl_3150	-	protein of unknown function	66.3	4.8	< 0.001	0.1	1.000
Mchl_5682	Mchl_1004	-	putative TonB-dependent receptor	78.3	4.4	< 0.001	0.1	0.960
	Makl 2074	ack2	alveerate kinase	85.0	51	<0.001	-0.5	0.250

Mchl_5685	Mchl_1166	cobM2	precorrin-4 C(11)-methyltransferase	82.7	7.0	< 0.001	-0.2	0.720
Mchl_5686	Mchl_1162	cobE2	putative cobalamin biosynthesis protein CobE	64.5	4.8	<0.001	0.9	0.490
Mchl_5687	Mchl_1161	cobL2	precorrin-6Y C(5,15)-methyltransferase (decarboxylating)	76.1	4.7	< 0.001	-0.2	0.990
Mchl_5688	Mchl_5189	cobK2	precorrin-6A reductase	63.4	0.7	0.600	0.2	0.810
Mchl_5689	Mchl_1710	cobJ2	precorrin-3B C(17)-methyltransferase	79.1	5.5	< 0.001	0.4	0.620
Mchl_5690	Mchl_1711	cobl2	precorrin-2 C(20)-methyltransferase	80.7	5.7	0.015	0.5	0.320
Mchl_5691	Mchl_1712	cobH2	precorrin-8X methylmutase	89.2	5.9	< 0.001	0.5	0.370
	Mchl_2536		conserved exported protein of unknown function, CoxB-	40.5	0.0	-0.001	-1.9	<0.001
IVICII_5095	Mchl_4258	-	related protein	49.4	8.9	<0.001	-0.1	0.960
Mchl_5701	Mchl_4908	folC2	putative folylpolyglutamate synthase and dihydrofolate	46.3	7.8	<0.001	-0.2	0.720
			synthase					
Mchl_5702	Mchl_0872	cobU2	nhosnhorihosyltransferase	54.5	6.8	<0.001	-0.1	0.890
Mchl 5703	Mchl 3302	_	nutative transposase (fragment)	85.7	24	<0.001	-0.1	0 990
Mchl 5704	Mchl 4231	-	nutative transposon (fragment)	95.4	3.2	<0.001	-0.8	0.706
inem_s/e4	111011_1231		DNA-binding response regulator in two-component	55.1	5.2	40.001	0.0	0.700
Mchl 5713	Mchl 0376	czcR	regulatory system nutative activator of the heavy-metal	49.8	19	<0.001	-0.7	0 150
		02011	resistance system Czc	1010	2.0			01200
			RND divalent metal cation efflux transporter membrane					
Mchl_5715	Mchl_1072	czcA2	component	43.4	2.3	<0.001	0.2	0.670
			bifunctional adenosylcobalamin biosynthesis enzyme					
Mchl_5721	Mchl_1722	cobP2	(adenosylcobinamide kinase/adenosylcobinamide-	73.6	6.7	< 0.001	-0.3	0.630
			phosphate guanylyltransferase)					
Mchl_5722	Mchl_1723	cobO2	cob(I)yrinic acid a,c-diamide adenosyltransferase	83.5	6.5	<0.001	0.1	0.870
Mchl_5723	Mchl_5190	cobQ2	cobyric acid synthase	75.2	6.2	< 0.001	-0.4	0.460
Mchl_5724	Mchl_5191	cobD2	cobalamin biosynthesis protein	76.4	2.3	<0.001	0.3	0.710
Mchl_5730	Mchl_5192	cobC2	L-threonine-O-3-phosphate decarboxylase domain	51.2	5.0	<0.001	-0.1	0.960
	Mchl_0172			54.7			0.9	0.074
Mchl_5735	Mchl_4926	-	RNA polymerase ECF-type sigma factor	43.5	1.3	<0.001	-0.3	0.580
	Mchl_1650			41.4			-1.3	<0.001
p1MCv2_0095	Mchl_0651	-	conserved protein of unknown function	44.3	2.3	<0.001	2.3	< 0.001
p1MCv2_0107	Mchl_2145	-	XoxF, PQQ-linked dehydrogenase (fragment)	100	1.8	<0.001	0.5	0.240

^a Genes encoding for transposon-related elements with no differential transcript abundance in the studied conditions are not listed. Genes are homologs when they have 80 % of identity on at least 80 % of the protein length.

^b log2 fold-change (log2fc) values \geq 2 or \leq -2 mean that transcript abundance is higher in cultures grown with chlorinated methanes compared to methanol, or higher with methanol compared to chlorinated methanes, respectively.

^c False discovery rate, only adjusted p-value under 0.1 were considered.

3.4. RNAseq data of genes with differential sense/antisense transcript abundance

Directional RNAseq was performed so that the information about the DNA strand from which the RNA was transcribed can be traced. Knowing the direction of the transcription is crucial for resolving overlapping genetic features, detecting antisense transcription and assigning the sense strand for non-coding RNA (ncRNA). Antisense RNAs referred to as *cis*-encoded RNAs are transcribed opposite annotated genes and thus share extensive complementarity with the corresponding transcripts although the antisense RNA may have additional targets in *trans* (Brantl, 2007). In the submitted paper, mRNA quantification was studied with the sum of sense and antisense reads in order to detect sets of genes with differential transcript abundance). Among those, we searched for the occurrence of putative antisense transcription nested within predicted mRNA-encoding genes. We defined the "sense ratio" which was calculated by dividing the number of normalized sense reads by the total number of normalized total sense and antisense reads. A "sense ratio" < 0.5 may point out misannotated uncharacterized CDS nested within predicted CDS, or may result from transcription noise (Llorens-Rico *et al.*, 2016), read-through of adjacent genes, or involve regulation mechanisms (Georg and Hess, 2011; Chang *et al.*, 2015; Brophy and Voigt, 2016).

Three groups were defined on the basis of the "sense ratio" in cultures grown with chlorinated methanes compared to methanol (Table 3.4).

- Group A, the "sense ratio" is lower than 0.5 pointing out the prevalence of antisense transcription;
- Group B includes CDS for which the "sense ratio" is above 0.8 only in one growth condition, and at least half the value in the other tested growth condition. This suggests conditional antisense transcription in cultures grown with different onecarbon compounds;
- Group C includes CDS that do not fall in groups A and B. This may include genes with similar ratio of sense and antisense transcription.

Using these criteria, in *M. extorquens* CM4, 20 genes were detected of which 17 were plasmid pCMU01-borne including six genes of a uncharacterized highly transcribed gene cluster harbouring *cszA2* and *cobS*-like, seven mobile element-related pseudogenes, and gene *cobM2*. The CobM2 protein (69 % aa identity with the characterized *Pseudomonas denitrificans* precorrin-4 C(11)-methyltransferase (Debussche *et al.*, 1993) was found directly upstream of a mobile-element gene (Mchl_5684), which may be responsible, at least in part, of *cobM2* antisense transcription. In *M. extorquens* DM4, of the 4 highly transcribed genes in methanol that had a majority of antisense reads, two were members of the gene family of uncharacterized ANAH-like domain containing proteins localized on GEI 197 (METDI1426; METDI4555).

	c 0.01 m	Gene characteristics ^a		/ 411(15)		Directional	RNAsen dat	tab	
Sense/ —		Gene characteristics		log2fc		Polativo r	and count	Sonco	ratiod
antisense group	Gene	Product	Genomic occurrence	logzic	Adjusted p-value	CM or	CH ₃ 0H	CM or	CH ₃ OH
		M. extora	uens CM4			DCIVI		DCIVI	
Group A: Antis	ense transc	ription							
Mchl 5465	-	transposase (fragment)	pCMU01	2.3	< 0.001	1257	261	0.3	0.2
Mchl 5685	cobM2	precorrin-4 C(11)-methyltransferase	pCMU01	7.0	< 0.001	432	3	0.4	0.1
	-	conserved protein (fragment)	pCMU01	2.2	< 0.001	293	67	0.3	0.1
	-	putative transposase (fragment)	pCMU01	4.2	<0.001	2949	162	0.2	0.2
	-	putative transposase (fragment)	pCMU01	2.4	0.015	208	43	0.0	0.1
Group B: Carbo	on source-d	ependent transcription orientation							
Mchl 1416	-	putative membrane protein	core	2.2	< 0.001	226	50	0.9	0.5
Mchl 3283	dctA	C ₄ -dicarboxylate transport protein	core	-2.9	< 0.001	149	1120	0.4	0.8
	clcA2	putative H(*)/Cl() exchange transporter	pCMU01	5.5	< 0.001	3782	84	0.9	0.3
	acxA	acetone carboxylase beta subunit	pCMU01	3.7	<0.001	1770	133	0.8	0.3
Mchl 5716	-	conserved protein. CobS-like domain	pCMU01	7.1	< 0.001	539	125	0.5	1.0
Mchl 5718	-	putative membrane transport protein	pCMU01	7.1	< 0.001	79188	578	1.0	0.5
Mchl 5719	-	putative transport membrane protein	pCMU01	7.7	< 0.001	81772	385	1.0	0.3
Group C: Simila	ar ratio of s	ense and antisense transcription							
Mchl 5715	czcA2	RND divalent metal cation efflux transporter SU	pCMU01	2.3	< 0.001	204	42	0.5	0.3
Mchl 5663	-	conserved protein	pCMU01	2.1	< 0.001	2072	484	0.5	0.3
p1MCv2 0379	-	protein of unknown function	pCMU01	2.4	<0.001	1514	296	0.5	0.7
p1MCv2_0380	-	protein of unknown function	pCMU01	2.1	<0.001	1233	295	0.2	0.6
<u></u>		M. extora	uens DM4						
Group A: Antis	ense transc	ription							
METDI0260	-	glucokinase (fragment)	GEI160	-4 3	<0.001	14	292	0.1	0.1
METDI0364	-	nutative cytochrome c oxidase subunit (ChaB)	GEI 160	-3.4	<0.001	174	1791	0.1	0.1
METDI1414	ardC	antirestriction protein (ArdC)	GEI 270	-2.2	0.026	36	159	0.4	0.4
METDI1414	urue	nutative alycosyl transferase	GEI 270	-2.2	0.020	587	2867	0.2	0.3
METDI3864	_	protein of unknown function	specific	-2.5 2 A	0.150	116	84	0.2	0.2
METDI/261	_	conserved protein of unknown function	core	-2.4	0.003	2/18	1525	0.2	0.0
METDI4201		protein of unknown function	GEI 197	-2.0	<0.001	1330	5840	0.0	0.0
Group B: Carbo	n source-d	energent transcription orientation	011157	-3.4	\0.001	1333	5040	0.4	0.4
METDI0133	-	nutative glycosyl transferase	variable	2.6	0.005	3/	6	10	0.4
METDI0620	_	conserved protein of unknown function	core	2.0	<0.003	9568	681	1.0	0.4
METDI1755	-	conserved protein of unknown function	specific	2.0	0.001	446	104	0.9	0.4
METDI1762	-	conserved protein of unknown function	variable	-2.1	0.007	553	2370	0.5	0.9
METDI1842	_	conserved protein of unknown function	core	-2.1	0.007	82	343	0.5	0.5
METDI1978	-	conserved protein of unknown function	core	-2.1	0.005	31	130	0.2	0.7
METDI2432	_	nutative phosphotransferase	core	2.1	<0.000	846	129	0.4	0.0
METDI2432	fdh4B	formate dehydrogenase subunit B	core	-3.2	<0.001	112	1020	0.5	1.0
METDI3359	-	nrotein of unknown function	core	2.1	0.001	610	143	0.5	0.2
METDI4507	_	nutative cytochrome c	GEL 197	-3.3	<0.002	93	913	0.0	0.2
METDI5891	-	nutative endonuclease	core	3.5	<0.001	3714	408	0.9	0.5
Group C: Simila	ar ratio of s	ense and antisense transcription	6010	5.2	0.001	5714	400	0.5	0.4
METDIO344	-	conserved protein of unknown function	GEI 160	2.2	0.013	120	31	0.5	0.2
METDI0365	_	conserved protein of unknown function	GEI 160	_3.3	<0.013	70	802	0.0	0.2
METDI0366	_	nutative sytochrome c	GEI 160	-2 9	0.001	108	1/138	0.4	0.0
METDI0367	_	putative cytochrome c	GEI 160	-2.5	<0.001	1/8	035	0.0	0.4
METDI038/		protein of unknown function	GEI 160	-2.7	0.001	17/	977	0.0	0.5
METDI1598	_	protein of unknown function	variable	2.5	0.004	5/1	126	0.7	0.3
METDI2012	_	nutative exported cvd operan protein	variable	_2.1	<0.012	424	5591	0.5	0.3
METDI2012	_	conserved protein of unknown function	variable	-3.7	0.001	1090	5590	0.5	0.4
METDIA505	-	nutativo outochromo o ovideco CL		-2.4	<0.004	1009	1055	0.0	0.4
METDI4504	_	putative cytochrome c oxidase 50	GEI 197	-5.0	0.001	120	1055	0.5	0.4
METDI4500	-		GEI 197	-2.5	<0.000	208	2022	0.0	0.5
METDI4528	-	consonued protein with ANALL like domein	GEI 197	-3.1	0.001	308	2549	0.5	0.7
	-	nutativo outochromo o	GEI 197	-3.3	0.001	413	4105	0.5	0.4
METDI4504	-	protoin of unknown function	Ucriable	-2.0	0.003	476	106	0.5	0.0
METDI4620	- nrol/	glycine /betaine /proline APC transporter SU	variable	2.2	0.002	4/0	100	0.7	0.4
WIL 1 D14095	prov	Biyene, betaine, profine Abe transporter 50	variable	2.5	0.001	209	55	0.7	0.4

Table 3.3. RNAseq data of genes with differential sense/antisense transcript abundance

METDI4696	opuAB	glycine/betaine/proline ABC transporter SU	variable	2.0	1.000	15578	17755	0.5	0.3
METDI5532	nrtB	nitrate transport permease protein	core	-2.4	0.001	73	386	0.2	0.7

^a MaGe annotation (https://www.genoscope.cns.fr/agc/microscope). Genes labelled as "core" encode for proteins with 80 % aa identity on at least 80 % of the protein length that are shared between *M. extorquens* genomes. Genes labelled as "variable" encode for homologs shared at least between 2 genomes of strains AM1, BJ001, CM4, DM4 and PA1. GEI stands for genomic islands as detailed in Table 4 of the submitted article.

^b log2 fold-change (log2fc) values \geq 2 or \leq -2 mean that transcript abundance is higher in cultures grown with chlorinated methanes compared to methanol, or higher with methanol compared to chlorinated methanes, respectively.

^c Averaged normalized read numbers divided by gene length in kb in different methylotrophic growth conditions.

^d Number of normalized sense reads divided by total number of normalized sense and antisense reads.

^e Adjacent to the Mchl_5714 to Mchl_5716 gene cluster, gene encoding Mchl_5717 (log2fc of 8.1) displayed a majority of sense transcript reads.

3.5. RNAseq data of formate dehydrogenase subunits encoding-genes

Formate is central to methylotrophic growth (see Chapter 1, Figure 1.10): For carbon assimilation, *M. extorquens* reduced formate into methylene-H₄F (CH₂=H₄F) (Marx *et al.*, 2005) while for energy production, formate is oxidized to CO₂ by formate dehydrogenase (FDH). *M. extorquens* core genome harbour four characterized FDH (Chistoserdova *et al.*, 2004, 2007). Some FDH-encoding SU genes were detected to have one-carbon compound utilization-dependent transcript abundance variations (summarized in Table 3.4).

Formate	M. extorquens CM4					M. extorquens DM4				
(cellular localization) a	encoded 50	log2fcb	Adjusted	Relative rea	ad counts ^d	log2fc	Adjusted	Relative re	ad counts	
		IOBTIC	p-value ^c	СМ	CH₃OH	IOBTIC	p-value	DCM	CH₃OH	
Fdh1	<i>fdh</i> 1A, alpha	1.6	<0.001	9327	3164	2.3	0.150	17071	3401	methanol
tungsten-containing (cytoplasm)	<i>fdh</i> 1B, beta	1.5	<0.001	7155	2480	1.8	0.350	10588	2981	methanol
Fdh2	<i>fdh</i> 2A, alpha ^f	-0.7	0.085	2391	3891	0.9	0.800	8798	4767	methanol
NAD-dependent	fdh2B, beta	-0.7	0.059	1571	2260	0.5	0.890	4856	3349	methanol
molybdenum containing	fdh2C, gamma	-0.5	0.290	662	958	-0.4	0.960	970	1257	methanol
(cytoplasm)	<i>fdh</i> 2D, delta	-0.9	0.029	1505	2821	0.9	0.450	3409	1801	methanol
	<i>fdh</i> 3A, alpha	0.5	0.300	15087	1049 9	0.9	0.810	14985	8365	succinate
Fdh3 (periplasm)	<i>fdh</i> 3B, beta	0.7	0.069	15221	9360	0.8	0.830	13849	7864	succinate
	fdh3C, gamma	0.5	0.270	10983	7894	0.8	0.840	1127	6360	succinate
Fdh4	<i>fdh</i> 4A alpha	-0.7	0.048	1819	3041	-4.0	<0.001	140	2280	nd
(cytoplasm)	fdh4B, beta	-1.2	0.001	762	1779	-3.2	<0.001	112	1020	nd

Table 3.4. RNAseq data of formate dehydrogenase subunits encoding-genes

^a Data from Chistoserdova *et al.*, 2004, 2007; Peyraud *et al.*, 2011.

^b log2 fold-change (log2fc) values \geq 2 or \leq -2 mean that transcript abundance is higher in cultures grown with chlorinated methanes compared to methanol, or higher with methanol compared to chlorinated methanes, respectively.

^c False discovery rate, only an adjusted p-value under 0.1 was considered as significant.

^d Averaged normalized read numbers divided by gene length in kb in different methylotrophic growth conditions.

^e Microarray data from Okubo *et al.*, 2007. Transcript abundance higher in cultures grown with methanol, succinate or unchanged (nd).

^f Proteomic data from Muller *et al.*, 2011. Protein more abundant in cultures grown with dichloromethane compared to methanol.

<u>4.</u> <u>Conclusion</u>

The transcriptomes for the utilization of methanol, chloromethane or dichloromethane as sole carbon and energy source were compared in two *M. extorquens* strains. The core genome represents 75 % (4,620 CDS) of their shared genes. The variable genome includes strain-specific pathways of chloromethane and dichloromethane that are initiated by the expression of dehalogenation modules that respectively are encoded by the *cmu* genes located on the 380 kb pCMU01 plasmid in strain CM4 and the *dcm* genes nested within a genomic island (GEI) in strain DM4. The core genome had a lower proportion of differentially expressed genes than the variable genome. A few common genes had higher transcript abundance under the following growth conditions:

- (i) chlorinated methane, including stress response and related bioenergetics genes (*depP*, *hppA* encoding a membrane-bound proton translocating pyrophosphatase);
- (ii) chloromethane, including the membrane-bound transhydrogenase *pnt* gene cluster for NADP⁺ regeneration and the glycine cleavage complex *gcvPHT* gene cluster involved in C₁ carrier tetrahydrofolate (H₄F) metabolism;
- (iii) dichloromethane, including a putative glutathione peroxidase;
- (iv) methanol, including 8 genes of the PQQ-dependent C_1/C_2 dehydrogenase *mdh2*-associated gene module.

The variable genome of strain CM4 harbored the majority of the genes whose transcription was enhanced by chloromethane: carbon assimilation genes (*cmu*, *acxABC*, *gck2*), coenzymeassociated *cmu*-pathway genes for vitamin B₁₂ (14 *cob* genes, *btuBCDF*, *bluB2*) and for H₄F (*folC2*, *folD*, *metF2*, *purU*). In the variable genome of strain DM4, the *dcm* genes were more expressed genes in cultures grown with dichloromethane compared to methanol, whereas central metabolism genes (*ackA*, *adh*-like, *xfp*, cytochrome-associated genes) encoded by redundant GEIs had increased transcript abundance in cultures grown with methanol. The common transcriptional response for *M. extorquens* grown with chloromethane or dichloromethane was limited to a few core genome genes. This suggests that transcription responses to growth with chloromethane and dichloromethane are predominantly uncorrelated. Redox status adjustments are probably necessary for growth with different C₁ compounds as deduced from the transcription profiles of genes of the core and the variable genome.

Chapter 4 Ecotypes of microbial chloromethane utilizers in a forest soil

Chapter 4. Ecotypes of microbial chloromethane utilizers in a forest soil

<u>1.</u> <u>Introduction</u>

Chloromethane (CH₃Cl) is an ozone-destroying volatile organic compound (VOC), which is mainly produced by natural processes (Galbally and Kirstine 2002; Denman *et al.*, 2007; see Chapter 1). Its global emissions to the atmosphere are about 10 Gmol year⁻¹. Chloromethane is the most abundant halogenated carbon compound in the atmosphere (500-600 ppt) (Montzka *et al.*, 2003). Chloromethane production is mainly associated with biological activities of plants and white rot fungi, and with the degradation of plant organic matter in soils (Harper 2000; Keppler *et al.*, 2005; Schäfer *et al.*, 2007). Chloromethane affects ozone formation, and causes 16 % of chlorine-catalyzed ozone destruction (Harper *et al.*, 2003; Hines *et al.*, 1998; Redeker *et al.*, 2000; Rhew *et al.*, 2007).

Another biogenic, plant-derived VOC is methanol. Global methanol (CH₃OH) emissions are much larger than those of CH₃Cl. Methanol represents the second most abundant organic compound in the atmosphere (0.1–10 ppb) after methane (1800 ppb). Global emissions of CH₃OH are estimated at 5 Tg y⁻¹. The main source of atmospheric CH₃OH is also associated with plants (Galbally and Kirstine 2002; Jacob 2005; Wohlfahrt *et al.*, 2015), and specifically originates from methoxy groups of pectin and lignin upon production of plant biomass (Donnelly and Dagley, 1980; Schink and Zeikus, 1980; Fall and Benson, 1996; Warneke *et al.*, 1999).

A significant part of naturally produced CH₃OH does not reach the atmosphere but is dissolved in water and thus readily utilized as a growth substrate by many methylotrophic bacteria, in particular in terrestrial soil environments (Kolb and Stacheter, 2012). In the case of the more volatile CH₃Cl, a filtering effect mediated by CH₃Cl degrading methylotrophs is also likely, in particular in the aerial parts of plants (Nadalig *et al.*, 2011; Farhan UI Haque *et al.*, 2013; Bringel and Couée, 2015). CH₃Cl-utilizing bacteria have been isolated from soil of industrial sites (McDonald *et al.*, 2001) and from woodland and forest soils (Borodina *et al.*, 2005). Thus, microbial-driven reduction of CH₃Cl emissions from soils is in principle possible. However, this process remains to be characterized in detail, and its significance in soils has not yet been investigated. Soil is a physicochemically heterogeneous environment, and top soil is highly biologically active. The number of bacterial cells per gram of top soil (Ah horizon) likely exceeds 10^{11} , and soil microbial diversity has been estimated at between 10^5 and 10^6 species of gram of soil bacteria and fungi, respectively (Peñuelas *et al.*, 2014). Forest soil is a typical habitat of methylotrophic microorganisms capable of using C₁ compounds for growth. Most of them are facultative methylotrophic, and strictly aerobic (Kolb *et al.*, 2009). Their abundance was estimated in the range between 10^6 cells and $3 \cdot 10^8$ cells g.soil⁻¹ (Stacheter *et al.*, 2013). There is evidence that methylotrophic bacteria are highly active in soil, and that they are able to use a large set of substrates, including CH₃OH and also some halogenated compounds including CH₃Cl (Kolb, 2009; Stacheter *et al.*, 2013).

To date, the magnitude and kinetics of transformations involving CH₃Cl in forest soils remains a black box. CH₃Cl degradation in forest soil was estimated at around 137 Gg.y⁻¹ (Yoshida *et al.*, 2004). Biotic and abiotic processes occurring in forest soil are complex and subject to high temporal and spatial variability, and contribute to the difficulty in estimating the global budget of CH₃Cl (Redeker *et al.*, 2012). It has already been shown that exchange of CH₃Cl between forest ecosystems and the atmosphere are modulated by bacteria in forest soil (Schäfer *et al.*, 2007). A biochemical pathway for CH₃Cl utilization (the *cmu* pathway) has been characterized in detail in *M. extorquens* CM4 (Vannelli *et al.*, 1999; Roselli *et al.*, 2013; see Chapter 1 and Chapter 3). This pathway was subsequently reported for several CH₃Cl-degrading isolates (reviewed in Schäfer, 2007) including isolates from forest soil (Doronina *et al.*, 1996; Coulter *et al.*, 1999, McAnulla *et al.*, 2001; Miller *et al.*, 2004). There is increasing evidence that the *cmu* pathway is not the sole pathway for CH₃Cl utilization (Chapter 1, Section 1.3.4., and Nadalig *et al.*, 2014).

Different approaches including degradation kinetics, use of pure bacterial cultures and stable isotope tracer experiments with aerated soil have shown that aerobic methylotrophic microbes in soil are able to use CH₃Cl at low concentrations such as those encountered in the environment (Shorter *et al.*, 1995; Hines *et al.*, 1998; Yoshida *et al.*, 2006; Miller *et al.*, 2004). Environmental CH₃Cl concentrations are so low that it is likely that CH₃Cl-degrading methylotrophs also rely on some of the very many other carbon compounds present in forest soil as growth substrates. For example, *Methylomicrobium album* BG8, for which CH₃Cl cannot

serve as the sole growth substrate, displays enhanced growth with CH₃Cl in the presence of CH₃OH (Han and Semrau, 2003).

In this study, bacterial CH₃Cl utilization in Steigerwald forest soil (Germany) and the effect of CH_3OH as a methylotrophic co-substrate in CH_3Cl utilization were investigated. To identify metabolically active key players in this process, taxonomic and functional gene biomarkers were employed in combination with a stable isotope labeling experiment. Genes encoding subunits of key enzymes of C1 metabolism such as mxaF have been repeatedly used as biomarkers to detect methylotrophs in the environment (McDonald et al., 2008). Here, 4 key functional gene markers of methylotrophs (cmuA, mxaF, xoxF, and mch, Figure 4.1) were targeted by PCR in addition to the 16S rRNA gene. The gene *cmuA* encodes one of the two methyltransferases of the first reaction required for growth with CH_3Cl by the *cmu* pathway, and has already intensively been used as a biomarker for CH₃Cl-utilization (McAnulla et al., 2001a; Miller et al., 2004; Borodina et al., 2005; Schäfer et al., 2005; Nadalig et al., 2011). Genes mxaF and xoxF are both associated with the oxidation of CH₃OH for growth for a wide diversity of microorganisms. The gene mxaF encodes the catalytic subunit of pyrroloquinoline quinone (PQQ)-dependent CH₃OH dehydrogenase, and was proposed early on as a functional gene probe for methylotrophs (McDonald and Murrell, 1997). The xoxF gene, homologous to *mxaF*, occurs in the overwhelming majority of methylotroph genomes (Chistoserdova, 2011). Homologs of xoxF encode functional MDHs (Fitriyanto et al., 2011; Chu and Lidstrom 2016; Vu et al., 2016). The xoxF gene is present in a wide range of both methylotrophs and nonmethylotrophs from Alphaproteobacteria, Betaproteobacteria and Gammaproteobacteria (Taubert et al., 2015). The gene mch encodes a methenyl cyclohydrolase involved in the interconversion of key intermediates of methylotrophic metabolism, and has already been used to resolve methylotrophic community structure and diversity (Kalyuzhnaya et al., 2004, 2005; Stacheter et al., 2013).



Figure 4.1. Targeted functional genes in methylotrophic utilization of chloromethane and methanol

The 3 genes *cmuA*, *mxaF*/*xoxF* and *mch* investigated by amplicon sequencing are boxed. Cofactors of central methylotrophic metabolism tetrahydrofolate (H₄F), tetrahydromethanopterin (H₄MPT) and metabolic intermediates including formaldehyde (CH₂O) and formate (HCOOH) are also indicated. The CH₃Cl utilization (*cmu*) pathway is shown in green, with the *cmuA* gene essential for growth with CH₃Cl as sole source of carbon and energy in *M. extorquens* CM4 (Vannelli et al., 1999). Reactions of the cmu pathway (green arrows) are shown on the right with associated genes encoding CH₃Cl dehalogenase subunits corrinoid-dependent methyltransferase (cmuA) and methyl-H₄F transferase (cmuB), methylene-H₄F reductase (*metF2*), bifunctional methylene-H₄F (CH₂=H₄F) dehydrogenase / methenyl-H₄F (CH \equiv H₄F) cyclohydrolase (*folD*) and 10-formyl-H₄F hydrolase (*purU*). Reactions and genes for transformation of methanol (CH₃OH) for energy production and incorporation into biomass are shown in blue, and include genes for oxidation reactions methanol dehydrogenase (mxaF and xoxF subunits) and methylene-tetrahydromethanopterin dehydrogenases (mtdA and mtdB). Other genes involved in oxidation of CH₃OH in Methylobacterium include formaldehyde activation enzyme (fae), methenyl cyclohydrolase (mch), and the formyltransferase-hydrolase complex (fhcABCD). Formate dehydrogenase (fdh) is assumed to be used for the final oxidation step to CO₂ from formate originating from both CH₃OH and CH₃Cl. Genes involved in carbon assimilation from CH₃OH include formyltetrahydrofolate ligase (ftfL), methylene-tetrahydrofolate cyclohydrolase (fch), and methylene-tetrahydromethanopterin dehydrogenases (mtdA). In Methylobacterium, carbon assimilation into biomass occurs by the serine cycle and the ethylmalonyl-CoA pathway via methylene-H₄F (striped thick arrow). In other organisms, carbon assimilation can occur by other pathways via formaldehyde or CO_2 (grey and white thick arrows). In the performed SIP labeling experiment, assimilation into biomass of CO₂ originating from mineralization of labeled carbon was avoided (red cross) as described in Material and Methods.

In previous studies, the *cmuA* gene probe and DNA stable isotope probing (DNA SIP) with fully $[^{13}C]$ -CH₃Cl have been employed to target the diversity of CH₃Cl-utilizing bacteria in soil (Miller *et al.*, 2004; Borodina *et al*, 2005). The SIP method makes use of the fact that the DNA of active bacteria that are able to assimilate the provided $[^{13}C]$ -labeled carbon source will be enriched in $[^{13}C]$ -carbon in their DNA. This "heavy" DNA can be separated by isopycnic ultracentrifugation from DNA of bacteria that did not assimilate the labeled compound. Such labeled DNA from active microbial populations can be analyzed, either by direct sequencing or by PCR amplification of key genes of interest (Coyotzi *et al.*, 2016; see Chapter 2). Here, we explored the structural and functional diversity of CH₃Cl-utilizers in top soil samples of forest soil amended with CH₃Cl and CH₃OH, added alone or together as the $[^{13}C]$ -labeled form, or as the regular, $[^{12}C]$ -unlabeled form of the compound.

2. Materials and methods

2.1. Study site and sampling

Soil was sampled from the Steigerwald forest area, located between Würzburg and Nürnberg (49° 37' N, 10° 17' O). Despite extensive development in this area, only few anthropogenic activities and no industrial activities exist in the forest, which covers a total area of about 1152 km². Vegetation mainly consists of beech (*Fagus sylvatica*), with minor stocks of oak (*Quercus* sp.). Five samples were taken from the upper soil layer within a 20 m circle, mixed, homogeneized and sieved through a 2 mm screen.

2.2. Soil activation and microcosm set-up

Physico-chemical characterization of soils was performed as described in Chapter 2. Sieved soil was aliquoted in batches of 500 g in sealed flasks, and 1 % CH₃Cl (Sigma Aldrich) was added to the gas phase to activate soils. Flasks (1 L) were kept at 20°C, and concentration of CH₃Cl was followed by gas chromatography until all CH₃Cl was consumed (within 2 weeks). For each microcosm, 70 g activated soil (approx. 50 mL) was transferred to a 500 mL flask sealed by a Viton stopper. In total, 8 different microcosms were prepared in duplicate in this study, and amended with unlabeled or [¹³C]-CH₃Cl (Sigma Aldrich) and/or CH₃OH (Campro Scientific) (Table 4.1). Each microcosm received five carbon pulses (18 mM of carbon per pulse and

carbon source) over a period of 23 days (Figure 4.2). After each pulse, 5 aliquots of 1 g soil were sampled and immediately frozen at -80°C until further analysis.

Carbon source added	Total carbon added (mM) ^a	Gas phase supplementation ^b	Liquid phase supplementation ^c
[¹³ C]-CH ₃ Cl	18	[¹³ C]-CH ₃ Cl	H ₂ O
$[^{13}C]$ -CH ₃ Cl + CH ₃ OH	36	[¹³ C]-CH ₃ Cl	CH₃OH
$CH_3CI + [^{13}C]-CH_3OH$	36	CH₃Cl	[¹³ C]-CH₃OH
[¹³ C]-CH₃OH	18	-	[¹³ C]-CH₃OH
CH₃Cl	18	CH₃Cl	H ₂ O
CH ₃ Cl + CH ₃ OH	36	CH₃Cl	CH₃OH
CH₃OH	18	-	CH₃OH
Control	-	-	H ₂ O

^a Carbon added per pulse (5 pulses were performed by microcosm; Figure 4.2).

^b A volume of 6 mL CH₃Cl was added for each CH₃Cl pulse. In total, a gas phase volume of 56 mL (air with or without CH₃Cl) was added to 500 mL flasks at each pulse in order to maintain slight overpressure in the flasks.

 $^{\rm c}$ Either 1 mL of milliQ water or of a 216 mM CH_3OH stock solution was added to soil of each microcosm per pulse.

2.3. Chemical analyses

Microcosms were monitored by measuring CH₃Cl and CO₂ concentrations by gas chromatography (HP 5890 Series II, Agilent) using a Porapak Q 80/100 (Supelco, Bellefonte, Pe) column. Injection of a new carbon pulse was performed when remaining CH₃Cl was low in microcosms amended with a CH₃Cl pulse. The same time-point was chosen for microcosms amended with a CH₃OH pulse. For CH₃Cl analysis, 1 mL of the gas phase was analyzed with the gas chromatograph fitted with an Electron Capture Detector (ECD). The injector temperature was fixed at 250°C, and ECD temperature was fixed at 300°C with a flow rate of 20 mL.min⁻¹, and a helium - methane (95:5) mixture as the carrier gas. A calibration curve with CH₃Cl standards allowed to correlate peak areas and CH₃Cl concentration. For CO₂ analyses, a volume of 100 μl of the gas phase was injected at 150°C into the gas chromatograph fitted with a thermal conductivity detector (TCD), and eluted at 250°C with a flow at 15 mL.min⁻¹, using an argon - methane mixture (95:5) as the carrier gas.

Values of pH were routinely measured on sieved soil, before pre-incubation, and after every carbon pulse was consumed, using an InLab R422 pH electrode (InLab Semi-Micro; Mettler-Toledo, Gießen, Germany). Soil moisture content was determined by weighing the soil before and after drying (at 60°C for 10 days).

2.4. [¹³C]-CO₂ analysis

Analyses were performed at the UFZ - Helmholtz Centre for Environmental Research, Department of Molecular Systems Biology, Leipzig (Germany). GC-MS analysis was carried out using a Perkin–Elmer GC Clarus 600 system with an Rtx[®]-1 capillary column (60 m x 320 μ m), operated with an ionization energy of 70 eV. Mass spectra were taken from 14 to 70 Da. Helium was used as the carrier gas with a constant flow at 300 kPa, and 10 μ l gas samples (split ratio 10:1) were manually injected using gas-tight syringes. Each sample was measured five times. Total [¹²C]-CO₂ and [¹³C]-CO₂ were analyzed by extraction of masses 44 and 45 followed by peak integration. Peak areas were corrected for [¹²C]-CO₂ and [¹³C]-CO₂ indoor air values to calculate the 45/44 mass ratio of CO₂.

2.5. Nucleic acid extraction and RNA removal

Nucleic acids were extracted from soil after the third carbon pulse (chloromethane dissipation differed between each duplicate after the third pulse) was consumed in microcosms amended with CH₃Cl (Figure 4.2). DNA from both duplicates of each of the 8 microcosms was extracted according to the method described by Griffith *et al.* (2000), except that centrifugation was performed at 13'000 *g* and that NaCl was not added in the final precipitation step. Nucleic acids were resuspended in 30 µL DEPC-treated water and heat-treated for 3 min at 65°C. RNAs were removed by RNaseA treatment at 37°C for an hour, and DNA was precipitated with 21 µl isopropanol and 6 µl 5 M NaCl overnight at -20°C. After centrifugation at 14'000 *g* for 1 h at 4°C, the DNA pellet was washed with 70 % ethanol, air-dried at room temperature and resuspended in 30 µL DEPC-treated water. DNA was quantified by fluorescence using the Quant-iT^m PicoGreen[®] kit (Invitrogen).

2.6. DNA fractionation

A cesium chloride gradient solution was prepared as described by Neufeld et al., 2007). For each microcosm, a volume of 5 mL of this solution, containing 2 µg of RNA-free microcosm DNA in 20 µL of DEPC-treated water, was loaded in an ultracentrifugation tube (Quick-Seal, Beckman Coulter). Tubes were placed in a Vti 65.2 vertical rotor (Beckman Coulter) and centrifuged for 40 h at 177'000 g (LE-70 ultracentrifuge, Beckman Coulter) followed by 2 h of deceleration so as not to disturb the cesium chloride density gradient. Tubes were pierced at the bottom using a sterile 0.63 mm diameter needle, and a dye-containing water solution was injected from the top of the tube using a low-flow pump (BioRad Econo Pump), with a flow fixed at 450 µL.min⁻¹. Ten 450 µL fractions were collected for each tube for densities previously defined to be associated with heavy, middle and light DNA fractions (Neufeld et al., 2007). DNA from each fraction was precipitated as described (Neufeld et al., 2007), resuspended in 20 µL, and the defined 4 heavy, 3 middle and 3 light fractions were pooled together. Pooled heavy, middle and light DNA fractions of each microcosm were quantified with the Quant-iT™ PicoGreen[®] kit (Invitrogen). DNA concentrations ranged between 0.3 - 3.0 ng/µl for heavy fractions, 14.5 - 25.8 ng/µl for middle fractions and 18.5 -50.3 ng/µl for light fractions, respectively.

2.7. PCR amplification

PCR reactions of 16S rRNA gene, *mxaF/xoxF* and *mch* genes (Table 4.2) were carried out in 0.2 mL microcentrifuge tubes on a LabCycler thermal cycler (SensoQuest, Göttingen, Germany). The obtained PCR products were checked by agarose gel electrophoresis. The 50 μL reaction mix for amplification of the 16S rRNA gene consisted of 23.75 μL distilled water, 5 μL DNA template solution (between 1.5 - 15 ng), 2.5 μL of 10 μM degenerate primer stocks 341for and 785/805rev (Table 4.2) using Crystal Taq DNA polymerase (Biolab products GmbH, Lüneburg, Germany) as recommended by the supplier. After initial denaturation (95°C, 5 min), DNA amplification was achieved by 40 cycles of 1 min steps of denaturation at 95°C, annealing at 55°C, and extension at 72°C, followed by a final extension step of 5 min at 72°C. Amplification of the *cmuA* gene was performed as follows. A reaction mix of 5 μL of 10x buffer provided by the supplier (1x: Tris- HCl 10 mM, KCl 50 mM, MgCl₂ 1.5 mM, Triton 0.1 %, BSA 0.2 mg⁻ mL⁻¹, pH 9.0, MP Biomedicals) 0.5 μL dNTPs (20 mM), 3.25 μL forward primer cmuAF422 and 0.75 μL of the reverse primer cmuAR422 (Table 4.2) as 20 μM stocks, 5 μL DNA

template solution (between 1.5 - 15 ng), and 33.5 μ L of distilled water was prepared. After initial denaturation (95°C, 10 min), 2 μ L TAQ polymerase mix (prepared as 2 μ L of TAQ polymerase (5U/ μ L, MP biomedicals) added to 1 μ L of 10x buffer and 7 μ L H₂O) was added to the reaction mix. DNA amplification was then performed for 30 cycles consisting of 15 s denaturation at 95°C, annealing/extension for 30 s at 56°C, followed by a final extension step of 5 min at 72°C.
Gene	Function	PCR Primer	Sequence (5'-3') ^a	Amplicon size (bp)	SNP/ OTU [♭]	Total OTUs ^c	Labeled OTUs ^d	Reference
16S	ribosomal small subunit	341for	CCTACGGGNGGCWGCAG	464	0) 117	5	Muyzer <i>et al.,</i> 1988
rRNA		785/805rev	GACTACHVGGGTATCTAATCC	404	9			Herlemann <i>et al</i> ., 2011
cmuA	chloromethane methyltransferase	cmuAF422	GARGTBGGITAYAAYGGHGG	477	38	8	5	This study ^f
cinari		cmuAR422	TCRTTGCGCTCRTACATGTCICC	722	50			This study
mch	methylotrophy pathways	mch-2a	TGCCTCGGCTCKCAATATGCYGGBTGG	E40	100	o	4	Verbelt et al. 1000
men		mch-3	GCGTCGTTKGTKCKBCCCAT	549	100	0		vomoit et ui., 1999
		mdh1	GCGGIWS <u>C</u> AICTGGGGYT		39	6	6	
mxaF_ xoxF ^e	methanol dehydrogenase	mdh2	GCGGIWS <u>G</u> AICTGGGGYT	430				This study ^f
		mdhR	GAASGGYTCSYARTCCATGCA					

Table 4.2. Biomarker gene primer sequence, PCR amplicon size and OTU occurrence

^a Degenerate base mixtures: B (C,G,T), H (A,C,T), K (G,T), N (A,C,G,T), R (A,G), S (G,C), Y (C,T), V (A, C, T), W (A,T). Inosine (I) was used instead of the N mixture (Zheng *et al.*, 2008) in some cases.

^b Maximal Single Nucleotide Polymorphism (SNP) positions possible within an OTU.

^c Corresponding to the sum of OTUs detected in the 8 microcosms of the SIP experiment. Sequences were affiliated to the same OTU at cutoff values of 98 %, 90 %, and 80 % sequence identity at the nucleotide level for 16S rRNA gene, *cmuA* and *mxaF_xoxF*, and *mch* amplicons, respectively.

^d See Materials and Methods for the criteria applied to define OTUs as "labeled".

^e Primer pairs allow to amplify both *mxaF* and *xoxF* types of methanol dehydrogenase (MDH) subunits. Amplifications were performed with two different forward primers mdh1 and mdh2 in order to reduce primer degeneracy and improve PCR efficiency. Amplicons obtained with primer pairs mdh1/mdhR and mdh2/mdhR were pooled before sequencing.

^f See Chapter 2 for details.

Amplifications of *mxaF/xoxF* and *mch* genes were performed in 20 µL reactions using 8 µL of 2.5x 5 Prime Mastermix (MgCl₂ final concentration 1.5 mM), 1.5 µL 3 % BSA (Biolab products GmbH, Lüneburg, Germany), 1 µL each of forward and reverse primers (10 µM; Table 4.2 for primer sequences), 5 µL DNA template solution (between 1.5 - 15 ng), 0.5 µL distilled water, and 3 µL 5x TaqMaster PCR Enhancer (5 Prime GmbH, Hamburg, Germany). In the case of *mxaF/xoxF* genes, two different PCR reactions with two different forward primers mdh1 or mdh2 (Table 4.2) and the same reverse primer mdhR (Table 4.2 for primer sequences) were performed, yielding amplicons for both *mxaF* and *xoxF* types of methanol dehydrogenase genes. After initial denaturation (95°C, 5 min), DNA amplification was performed by 40 cycles of 1 min denaturation at 95°C, annealing for 1 min at 55°C, extension for 1 min at 72°C and a final extension step of 5 min at 72°C.

2.8. DNA sequencing

PCR products of mdh1/mdhR and mdh2/mdhR amplicons were mixed prior to sequencing. In total, 24 samples each of 16S rRNA gene, *cmuA*, *mch*, and *mxaF_xoxF* amplicons were sequenced using Illumina MiSEQ technology (see Chapter 2) by a commercial supplier (LGC Genomics GmbH). Briefly, a barcode oligonucleotide identifying sequence was ligated to each PCR product. Equimolar pools of all libraries were assembled, and the resulting combined library was analyzed using a standard MiSeq paired end (2×300 bp) flow cell and reagent cartridge.

2.9. Analysis of the 16S rRNA gene

Illumina reads were analyzed using mothur software package v.1.33.2 (Kozich et al., 2013) with the default parameters of the MiSeq standard operating protocol (http://www.mothur.org/wiki/MiSeq_SOP). Briefly, read pairs were assembled into contigs. Contigs shorter than 420 bp or longer than 460 bp were discarded. Sequences were preclustered in groups of sequences with up to 2 nucleotide differences. Chimera sequences predicted by UCHIME (Edgar et al., 2011) were removed. Remaining sequences were assigned using naïve Bayesian taxonomic classification on the bacterial reference database SILVA (SSU Ref database v.119), at a bootstrap cutoff set at 80 %. Remaining sequences were classified using the reference database SILVA (SSU_Ref database v.119), at a bootstrap cutoff set at 80 %. Sequences affiliated to Bacteria and Archaea domains were selected and other sequences discarded. OTUs were defined at 98 % sequence identity using the automated protocol within Mothur. This also yielded a representative consensus sequence, for each OTU, chosen as the most abundant sequence in a given OTU, and used for subsequent sequence alignments.

2.10. Analysis of functional genes

Raw reads were processed using mothur software package v.1.33.2 (Kozich *et al.*, 2013), including length filtering and quality trimming, and allowing sequence lengths within 20 nucleotides of the expected length of the amplicon. USEARCH software (Edgar, 2010) was then used for clustering the obtained filtered reads. Sequences occurring only once within the complete dataset of gene amplicon libraries were considered artefactual and removed, whereas singletons in individual amplicon libraries were retained. Reads were clustered iteratively at progressively lower cut-off values, and the maximum cut-off value at which the number of retrieved OTUs stabilized (90 % for *cmuA* and *mxaF/xoxF*, 80 % for *mch*) was selected (Stacheter *et al.*, 2013). Consensus sequences for each OTU were those provided by mothur. These sequences were compared against a gene-specific database generated from GenBank using BLAST (http://blast.ncbi.nlm.nih.gov) for taxonomic identification. A cluster table for each gene and each sample was obtained and analyzed as described below to identify labeled OTUs.

2.11. Identification of labeled OTUs

Labeled OTUs (OTU detected specifically and dependently to the microcosm tested) were defined by following a previously reported protocol developed to minimize false positives in 'heavy' DNA fractions (Lueders *et al.*, 2004; Liu 2007; Dallinger and Horn 2014). Briefly, the relative abundances in the 'heavy' DNA fraction of a given [¹³C]-labeled microcosm and in its corresponding unlabeled microcosm (amended with the corresponding unlabeled ([¹²C]-compound), as well as the relative abundances in 'heavy' and 'light' DNA fractions of a given [¹³C]-labeled microcosm, were compared. For this study, the following criteria had to be met in order to define OTUs in microcosms amended with a [¹³C]-labeled compound as "labeled": (1) the relative abundance of the OTU in the 'heavy' DNA fraction of the [¹³C]-labeled microcosm was higher than in the 'heavy' DNA fraction of the corresponding unlabeled microcosm; (2) the relative abundance of the OTU was higher in the 'heavy' DNA fraction than

in the 'light' fraction of the [¹³C]-labeled microcosm; (3) the relative abundance of the OTU in the 'heavy' DNA fraction of the [¹³C]-labeled microcosm was ≥ 0.5 %; (4) the relative abundance difference of the OTU between 'heavy' and 'light' DNA fractions was ≥ 0.1 % higher in the [¹³C]-labeled microcosm than in the corresponding unlabeled microcosm.

2.12. Statistical and phylogenetic analyses

Richness (as Sobs), Shannon index, and Simpson diversity indices were determined using the mothur routine as described (http://www.mothur.org/wiki/MiSeq_SOP), and with PAST software (http://folk.uio.no/ohammer/past). Shannon *H* index was $H = -\Sigma p_j \ln p_j$ and Simpson diversity $D_2 = 1/\Sigma p_j^2$, where p_j is the OTU relative abundance. Relationships between sequence datasets in different DNA fractions and microcosms were investigated by two-dimensional NMDS (non-metric multidimensional scaling) within mothur and visualized with Kaleidagraph (Synergy Software, Reading, PA, USA). Heatmaps were produced with R software (version 3.2.1, 2015-06-18) using packages ggplot2, RODBC, reshape, plyr and RColorBrewer. Consensus sequences of labeled OTUs were aligned with known reference sequences, and the alignment was used for Maximum likelihood analysis based on the Tamura-Nei model using MEGA 7 software (Tamura et Nei 1993; Kumar, Stecher, et Tamura 2016). Tree representations were generated and plotted within MEGA7.

3. <u>Results</u>

3.1. Microcosm setup for detection of CH₃Cl-utilizing methylotrophs in forest soil

CH₃Cl, a growth substrate for a minority of specialized methylotrophic microorganisms, is known to be naturally produced in top soils and especially in tree vegetated ecosystems such as forests. On the other hand, CH₃OH, another C₁ growth substrate of methylotrophs, is produced in much larger amounts in the same environments. A stable isotope labeling approach using the [¹³C] isotopologue of CH₃Cl was chosen to identify microbial populations that assimilate carbon of CH₃Cl in temperate forest soils in the presence and absence of CH₃OH. Forest soil microcosms amended with [¹³C]-CH₃Cl or [¹³C]-CH₃OH were set up in order to (i) identify CH₃Cl-utilizers in forest soil, and ii) assess the impact of CH₃OH on CH₃Cl utilization and the corresponding soil microbial methylotroph community. Starting from sieved and homogeneized forest soil pre-activated for about 2 weeks by addition of 18 mM

CH₃Cl, a total of four [¹³C]-labeled amendment conditions was tested on the same sieved soil: [¹³C]-CH₃Cl, [¹³C]-CH₃Cl with unlabeled CH₃OH, [¹³C]-CH₃OH with unlabeled CH₃Cl, and [¹³C]-CH₃OH (Table 4.1). Three control microcosms were treated with equivalent unlabeled carbon amendments, and another control microcosm was incubated without carbon amendment (Teble 4.1). Comparison with unlabeled microcosms was required for definition of labeled OTUs of used gene markers (see below, Section 4.3.3). CH₃Cl and/or CH₃OH were supplemented with 5 pulses of 18 mM unlabeled or [¹³C]-labeled C₁ compound over a period of 23 days (Figure 4.2).



Figure 4.2. Carbon mineralization in the SIP experiment

(A) Time-course of total ($[^{12}C]$ -CO₂ and $[^{13}C]$ -CO₂) production of CO₂ in the 8 microcosms. When only CH₃Cl or CH₃OH was added (top and bottom panels), squares indicate CO₂ production from $[^{13}C]$ -labeled C₁ compounds, and circles production from unlabeled compounds, respectively. When both CH₃OH and CH₃Cl were added (middle panel), squares represent the production from $[^{13}C]$ -CH₃Cl and unlabeled CH₃OH, triangles production from unlabeled CH₃Cl and $[^{13}C]$ -CH₃OH, and circles production from unlabeled compounds, respectively. Data for the control microcosm (no carbon addition, crosses) are given in all three graphs for comparison. The 5 carbon pulse time points are indicated (vertical arrow). Significant differences in microcosm duplicates were observed after the fourth pulse (data not shown). Thus, DNA extractions and PCR amplifications were performed after the third carbon pulse was consumed (curved arrow). (B) Carbon mineralization as $[^{13}C]$ -labeled CO₂ is shown

only for the 4 microcosms to which $[^{13}C]$ -labeled carbon was added. Data are shown at the time-point of DNA extraction, as the total $[^{13}C]$ -labeled CO₂ after 3 carbon pulses (54 mM $[^{13}C]$ -labeled carbon supplemented in total). Vertical bars show standard deviation of 2 biological replicates.

3.2. Mineralization and assimilation of chloromethane in forest soil microcosms

When methylotrophic microorganisms present in the microcosm actively utilize amended CH_3CI and CH_3OH as energy source, this leads to CO_2 production (Figure 4.1; also detailed for strains of the genus *Methylobacterium* in Chapter 3). CO_2 production in microcosms was determined by gas chromatography (Figure 4.2A). As expected, a net increase in CO_2 production was observed in all microcosms in response to carbon supplementation (Figure 4.2A). Overall, a trend towards larger CO_2 release upon supplemental CH_3OH than upon supplemental CH_3CI was observed. There was no significant difference in CO_2 production in microcosms to which labeled or unlabeled compounds were added, and thus the use of the [¹³C] isotope did not affect carbon metabolism overall. Observed rates of CO_2 production in microcosms were in the range of 0.2-0.3 mmol g_{soil}^{-1} day⁻¹ for microcosms to which CH₃CI and CH₃OH were added.

Amendment with [¹³C]-labeled compounds allowed to estimate the proportion of the carbon amendment that was used for energy production, based on produced [¹³C]-CO₂ (Figure 4.2). The errors in the determinations are large, but the calculations suggest that about 10 mM of the total added 54 mM [¹³C]-CH₃Cl were converted to [¹³C]-CO₂ in the [¹³C]-CH₃Cl amended microcosm. Thus, about 80 % of the supplemented carbon was assimilated into biomass. A significant increase in the extent of mineralization from labeled CH₃Cl (about 50 %) was observed when unlabeled CH₃OH was also provided. In the reverse case of microcosms amended with [¹³C]-CH₃OH, however, the situation was different. Only a minor increase in mineralization was observed in the microcosm to which unlabeled CH₃Cl was provided together with [¹³C]-CH₃OH. Moreover, methanol was mineralized to a significantly larger extent than CH₃Cl when provided alone.

3.3. Taxa associated with chloromethane utilization in forest soils

Taxonomical identification was based on sequence analysis of an approximately 460 nucleotide-long segment of the 16S rRNA gene hypervariable region that was PCR amplified.

The amplicons from heavy, middle and light gradient fractions retrieved of each of the 8 microcosms (Table 4.2) were sequenced by Illumina MiSeq, generating tens of thousands of sequences per amplicon (Table 4.3; Table S4.1).

Table 4.3. Number of filtered amplicon sequences obtained from heavy DNA fractions								
Microcosm	rrsA	стиА	mxaF_xoxF	mch				
[¹³ C]-CH ₃ Cl	46110	4282	786	55892				
[¹³ C]-CH ₃ Cl + CH ₃ OH	178042	3778	1534	38644				
CH₃Cl + [¹³ C]-CH₃OH	76690	614	918	52586				
[¹³ C]-CH ₃ OH	98332	1886	588	43088				
CH₃CI	72522	370	2616	27732				
CH₃Cl + CH₃OH	47236	922	862	44356				
CH₃OH	90822	246	2946	26518				
Control	100032	184	2710	26804				

See Table S4.1 for details of numbers of raw and filtered reads obtained in all DNA fractions.

The analysis of the recovered diversity in all DNA fractions at the 98 % sequence identity level suggested that, with one possible exception (i.e. the heavy fraction of the microcosm amended with [¹³C]-CH₃OH and CH₃Cl, see below, section 4.3.4), datasets clustered according in a self-consistent pattern (Figure 4.3).



Figure 4.3. Clustering analysis of 16S rRNA gene amplicon sequence datasets from all microcosms

Relationships between sequence datasets were investigated by two-dimensional NMDS (nonmetric multidimensional scaling) analysis within mothur. [¹³C]-labeled microcosms and unlabeled microcosms are shown with full and empty symbols, respectively: microcosms supplemented with CH₃Cl only (squares); with [¹³C]-labeled CH₃Cl and CH₃OH (upward pointing triangles); CH₃Cl with [¹³C]-labeled CH₃OH (downward pointing triangles); with unlabeled CH₃Cl and CH₃OH (diamonds); with CH₃OH only (circles); and the unamended control (asterisks). Heavy, middle, and light fractions are shown in red, orange and green, respectively. Stress: 13.6 %.

A large and similar effect on community structure (Figure 4.3) was noticed on heavy fractions of microcosms labeled with [13 C]-CH₃Cl alone, or together with unlabeled CH₃OH. The effect of addition of [13 C]-CH₃OH alone was also prominent. Computed diversity indices of all DNA fractions then confirmed the observation from cluster analysis (Figure 4.3) that there was no significant difference in the recovered diversity of light fractions of all microcosms, as well as the unamended control (Table 4.4). The light fractions of [13 C]-labeled microcosms, as well as the middle fractions that were not investigated further, were also little affected by the amendments (Figure 4.3). In contrast, a significant decrease in diversity was observed in the heavy fractions of microcosms labeled with [13 C]-CH₃Cl alone or together with unlabeled CH₃OH, and also in the microcosm to which CH₃OH alone was added (Table 4.4).

Microcosm	SIP fraction	Sobs ^a	Shannon indexª	Simpson diversity ^a
[¹³ C]-CH ₃ Cl	Н	1762	5.95	44
	L	2870	7.82	223
$[^{13}C]$ -CH ₃ Cl + CH ₃ OH	Н	1629	6.11	36
	L	3150	7.55	354
CH ₃ Cl + [¹³ C]-CH ₃ OH	Н	1567	5.13	12
	Η ^b	1566	6.52	120
	L	3062	7.86	283
[¹³ C]-CH ₃ OH	Н	1215	4.58	7
	L	3148	7.40	322
Unamended control	Н	1804	6.45	102
	L	2904	7.39	217

Table 4.4. Diversity indices for 16S rrnA amplicon sequences obtained from heavy and light	ıt
DNA fractions	

^a Calculated at the 98 % sequence identity level. See Materials and Methods for definitions. Simpson diversity is considered a conservative measure of the effective number of phylotypes (Haegeman *et al.*, 2014).

^b Computed after removal of the *Curtobacterium* genus-level OTU from Microbacteriaceae, detected at 29.8 % relative abundance in the heavy DNA fraction of the [¹³C]-CH₃OH labeled microcosm, but at 0.6 % relative abundance in all other 23 DNA fractions (see text).

After clustering at 98 %, a total of 117 family-level 16S RNA OTUs were detected in the eight microcosms (Table 4.2, Figure S4.1). However, the large set of sequence data obtained by Illumina MiSeq of PCR amplicons (Table S4.1) makes it difficult to assess directly which taxa are associated with CH₃Cl and/or CH₃OH utilization (Figures S4.1 to S4.4). The specific procedure for data analysis described in Material and Methods, basing on the approach described by Liu *et al.*, (2011), allowed to define and identify labeled taxa (OTUs) in a conservative manner. In total and across all [¹³C]-labeled microcosms, only 5 families were defined as labeled (Figure 4.4).





Only microcosms exposed to [¹³C]-labeling are shown. In total, 5 of the recovered 117 familylevel OTUs satisfied the criteria defined for a [¹³C]-labeled OTU (see Materials and Methods). Labeled OTUs in each microcosm are indicated by a star (see also Table 4.4). A 6th OTU (*Microbacteriaceae*), detected as highly abundant in the heavy fraction of the [¹²C]-CH₃Cl/[¹³C]-CH₃OH microcosm only, and thereby predicted to be labeled, is considered potentially doubtful (see text). Dotted lines in this microcosm indicate relative abundances of OTUs if this *Microbacteriaceae* OTU is included.

These were Beijerinckiaceae (Alphaproteobacteria), three families of Actinobacteria (i.e. Acidothermaceae, Pseudonocardiaceae and *Streptomycetaceae*) and а minor. uncharacterized family-level OTU belonging to the phylum TM7 (Table 4.5). A 6th OTU (Microbacteriaceae) was detected at lower than 0.6 % relative abundance in 22 gradient fractions, including the heavy fraction of the equivalent unlabeled microcosm, as well all other fractions of unlabeled and labeled microcosms amended with both CH₃Cl and CH₃OH. However, the Microbacteriaceae OTU was detected at very high (29.8 %) relative abundance in one heavy fraction of the [¹²C]-CH₃Cl/[¹³C]-CH₃OH microcosm, and thereby predicted to be labeled. Although this label may be considered as doubtful, it might indicate that these organisms are chloromethane degraders that dissimilate but not assimilate CH₃Cl (see Discussion below).

Four out of five family-level labeled taxa correspond to frequently labelled genera. *Methylovirgula, Acidothermus,* and *Streptomyces* represent over 95 % of the highly dominant

labeled family-level OTUs *Beijerinckiaceae*, *Actinomycetales* and *Streptomycetaceae*, respectively (Table 4.5). In the microcosm supplemented only with [¹³C]-CH₃Cl, 3 of the 5 family-level OTUs (*Acidothermaceae*, *Beijerinckiaceae* and *Streptomycetaceae*) made for about 80 % of the total relative OTU abundance (Table 4.5).

	Relative abundance (%) in heavy fractions of [¹³ C]-labeled microcosms ^a								
Conuc (family)									
Genus (turniny)	[¹³ C]-CH ₃ Cl	[¹³ C]-CH ₃ Cl +	CH₃Cl +	[¹³ C]-CH₃OH					
		CH₃OH	[¹³ C]-CH₃OH						
Methylovirgula	27.0	29.9		58.5					
(Beijerinckiaceae)	(27.9)	(31.1)	(13.5)	(59.4)					
Streptomyces	29.9								
(Streptomycetaceae)	(29.9)	(26.1)	(23.8)	(11.4)					
Acidothermus	12.3	7.2	5.9						
(Acidothermaceae)	(12.9)	(7.3)	(6.0)	(2.8)					
Pseudonocardia			4.8						
(Pseudonocardiaceae)	(3.5)	(3.1)	(5.3)	(1.4)					
Curtobacterium			29.8						
(Microbacteriaceae) ^b	(0.3)	(0.6)	(30.0)	(0.2)					
TM7 ^c	(0.3)	(1.3)	(0.8)	(0.3)					
(Total of family-level taxa)	(74.5)	(68.9)	(79.4)	(75.3)					

Table 4.5. Dominant	genera within	labeled family	v-level	¹³ C	OTUs

^a Percentage of total filtered reads affiliated by GreenGenes to the same dominant genus (featuring several OTUs at >98 % sequence identity) in the 5 family-level taxa defined as labeled in all microcosms (in brackets: % of total reads; labeled family-level taxa in each microcosm are shown in bold).

^b The family-level OTU *Microbacteriaceae* was detected at 30 % relative abundance only in the heavy DNA fraction of this microcosm labeled with [¹³C]-CH₃OH, and at ≤0.6 % relative abundance in the heavy fraction of the corresponding unlabeled microcosm as well as in all other 22 DNA fractions that were sequenced (see text and Figure 4.2).

^c Uncharacterized family-level taxon (>80 % sequence identity) within the TM7 phylum.

In the microcosm supplemented only with [¹³C]-CH₃OH, in contrast, only *Beijerinckiaceae* was labeled and amounted to an even larger percentage than in the other labeled microcosms. The minor abundant family-level labeled OTU of phylum TM7 was only labeled in the [¹³C]-CH₃Cl treated microcosm that was additionally supplemented with unlabeled CH₃OH. The labeled OTU affiliated with *Pseudonocardiacae* was a dominant taxon in all CH₃Cl-amended

microcosms, but was only detected as labeled in the microcosm supplemented with $[^{13}C]$ -CH₃OH together with unlabeled CH₃Cl.

Labeled OTUs differed significantly from the closest type strains as well as from previously described CH₃Cl-degrading strains (Figure 4.5).



0.02

Figure 4.5. Phylogenetic relationships between partial sequences of the 16S RNA gene of labeled OTUs and of reference chloromethane-utilizing strains

Genus-level labeled OTUs are in bold. Their selected consensus 16S rRNA gene sequences (464 nucleotides) were aligned with those of corresponding type strains including *Acidothermus cellulolyticus* 11B^T (Acc. No. JQ899226), *Curtobacterium ginsengisoli*^T (EF587758), *Nocardiodes caeni* MN8^T (FJ423551), *Methylovirgula ligni* BSW863^T (FM252034), *Pseudonocardia chloroethenivorans* SL-1^T (AF454510), *Streptomyces rishiriensis* NRRL B-3239^T (EF178682), and those of characterized CH₃Cl-degrading strains (underlined) such as *M. extorquens* CM4 (AF198624, Doronina *et al.*, 1996), *Hyphomicrobium* sp. MC1 (NR_074190, Hartmans *et al.*, 1986, Vuilleumier *et al.*, 2011), *Hyphomicrobium* sp. AT3 (FN667865, Nadalig *et al.*, 2011), *Leisingera methylohalidovorans* strain DSM 14336 (DQ915607, Schaefer *et al.*, 2002), *Aminobacter ciceronei* strain IMB-1 (AF034798, Hancock *et al.*, 1998), and *Nocardioides* sp. SAC-4 (F279792.1, McAnulla *et al.*, 2001). A Maximum Likelihood phylogenetic tree representation was constructed from the alignment basing on the Tamura-Nei model (Tamura *et al.*, 1993).

Chapter 4

3.4. Diversity of *cmu* pathway chloromethane utilizers in forest soil microcosms

Chloromethane: cobalamin methyltransferase *cmuA*, the key gene biomarker for CH₃Cl consumption, was amplified with new primers specifically designed for this study (see Chapter 2). Based on the amplicon sequence data in the heavy fractions (Table 4.3), OTUs in all fractions were defined and analyzed as for the 16S rRNA gene, but with a cut-off of 90 % sequence identity. In total, 8 OTUs were recovered, among which 5 satisfied the criteria defined in this work for labeled OTUs (see Materials and Methods, Table 4.2, Figure 4.6 and S4.2).



Figure 4.6. Affiliation and relative abundance of cmuA OTUs

OTUs shown in bold were defined as labeled according to the criteria defined in Materials and Methods. **(A)** Phylogenetic analysis of *cmuA* amplicon sequences. The tree was constructed from a 422 nucleotide sequence alignment of consensus sequences of the 8 labeled OTUs defined for the PCR amplicons obtained from microcosm heavy DNA fractions (OTU_cmuA_X; X ranging from 1 to 8); sequences of uncultured organisms detected in previous studies on CH₃Cl-utilization in three different soils (Gibbet Hill, Tocil Wood and Warwickshire from Miller *et al.*, 2004) and Plymouth coastal water enrichments (Schäfer *et al.*, 2005); and sequences from reference CH₃Cl degrading strains (*Aminobacter lissarensis, Aminobacter* sp. IMB-1, *Hyphomicrobium* sp. MC1, *Hyphomicrobium* sp. AT3 and *M. extorquens* CM4; underlined), as well as other strains of known genome sequence containing the *cmuA* gene. Branch lengths

are proportional to percentage sequence differences. Bootstrap was performed with 1,000 replicates. **(B)** Heatmap of the relative abundance of the 8 *cmuA* OTUs detected in the heavy DNA fractions in [13 C]-labeled microcosms. The total number of reads after PCR amplification of the *cmuA* gene (Table 4.2) is also shown.

Consensus sequences of these 8 OTUs were compared to *cmuA* sequences obtained in a previous SIP study of soil with [¹³C]-CH₃Cl (Borodina *et al.*, 2005), and to the sequences of isolated strains of known genome sequence harboring a *cmuA* gene, several of which were shown to be CH₃Cl-degraders (Figure 4.6A). The 8 OTUs belong to 3 distinct groups. All labeled OTUs are closely related to the sequences of known CH₃Cl-degraders including *M. extorquens* CM4 (>99 % identity) or *Hyphomicrobium* sp. MC1, itself closely related to sequences and CH₃Cl-degrading strains previously obtained in the SIP CH₃Cl study of soil (Borodina *et al.*, 2005). In the CH₃Cl condition, OTU_*cmuA*_2 and OTU_*cmuA*_3 were labeled and among the most dominant (relative abundance of 63.7 and 24 %, respectively). When CH₃OH was added to [¹³C]-CH₃Cl, only OTU_*cmuA*_2 was still defined as labeled, and was even more abundant than in the microcosm amended only with CH₃Cl (over 83 % of all sequences, Figure 4.6B). In the microcosm supplemented with [¹³C]-CH₃OH, in contrast, OTU_*cmuA*_3, OTU_*cmuA*_4, OTU_*cmuA*_6 were detected as labeled, whereas OTU_cmuA_2 was not, and OTU_cmuA_6 was found as abundant (35 % of reads). In the [¹³C]-CH₃OH, OTU_cmuA_2, 3, 4 and 6 were detected, with a large abundance of OTU_*cmuA*_2 (64 %), but only OTU_cmuA_3 was labeled.

3.5. Diversity of methanol utilizers in forest soil microcosms

Sequences of the MDH amplicon in the heavy DNA fraction of [¹³C]-labeled microcosms (Table 4.3) were grouped in 6 OTUs at the chosen cut-off of 90 % sequence identity (Table 4.2, Figure 4.7 and Figure S 4.3). Closest sequences to MDH_OTU_6 are from *xoxF* genes from *Hyphomicrobium* strains, whereas the 5 other OTUs cluster closely together, and are more closely related to *xoxF* sequences of *Methylobacterium* strains (Figure 4.7A). Thus, only *xoxF*-like sequences were recovered (Figure 4.7), despite the fact that primer pairs used to generate amplicons of genes encoding CH₃OH dehydrogenase (MDH) had the potential to amplify both *mxaF* and *xoxF* types of the target gene for a catalytic subunit of the enzyme (Table 4.2).



Figure 4.7. Affiliation and relative abundance of MDH OTUs

(A) Phylogenetic analysis of methanol dehydrogenase (MDH) amplicon sequences was performed with 46 unique 549 nucleotide long sequences including *xoxF* and *mxaF* sequences from reference strains (known CH₃Cl-degraders are underlined) and consensus sequences for OTUs (OTU_MDH) defined in heavy DNA fractions at the >90 % sequence identity level. All recovered MDH OTUs were defined as labeled according to the criteria defined in Materials and Methods. No *mxaF* or *xoxF* sequences are detected in CH₃Cl-degraders *Aminobacter lissarensis, Aminobacter* sp. IMB-1 and *Hyphomicrobium* sp. AT3. The tree was obtained from the alignment by the Maximum Likelihood method (Tamura Nei model, Tamura *et al.,* 1993). Branch lengths are proportional to sequence differences. Bootstrap was performed with 1,000 replicates. (B) Heatmap of the relative abundance of OTUs in heavy DNA fractions of [¹³C]-labeled microcosms.

The 6 MDH OTUs were defined as labeled in one or several of the [¹³C]-labeled microcosms. OTU_MDH_2, OTU_MDH_3 and OTU_MDH_6 were labeled in the [¹³C]-CH₃Cl treatment, with OTU_MDH_3 the most abundant OTU detected (69.7 %, see Table S4.2). In the condition in which unlabeled CH₃OH was added on top of [¹³C]-CH₃Cl to forest soil, in contrast, OTU_MDH_3 and OTU_MDH_6 were not labeled, but the other OTUs were (Figure 4.7B). Of these, OTU_MDH_2 and OTU_MDH_4 were also labeled in the reciprocal condition (i.e. addition of [¹³C]-labeled CH₃OH and unlabeled CH₃Cl). OTU_MDH_2 and OTU_MDH_6, were

detected at low abundance (3 %), while OTU_MDH_4 had higher abundance in the microcosm to which only $[^{13}C]$ -CH₃OH was added.

Analysis of *mch* homologs showed an overall higher diversity compared to other studied functional genes. At the cut-off value of 80 % (Table 4.2, Stacheter *et al.*, 2013), 16 OTUs were defined, more than for the other two functional genes (Tables 4.2 and S4.2). Only 4 of the 16 OTUs were defined as labeled (OTU_mch_3, OTU_mch_9, OTU_mch_10 and OTU_mch_11; Figure 4.8 and Figure S 4.4).



Figure 4.8. Affiliation and relative abundance of mch OTUs

(A) Phylogenetic analysis of methenyl cyclohydrolase (*mch*) amplicon sequences was performed with a set of 47 unique 430 nucleotide long sequences including sequences from reference strains (known CH₃Cl-degraders are underlined), sequences of isolates from a previous SIP study on CH₃OH utilization in top soils (Stacheter *et al.*, 2013), and consensus sequences for *mch* OTUs defined in heavy DNA fractions at the >80 % sequence identity level. OTUs defined as labeled according to the criteria defined in Materials and Methods are shown in bold. The tree was obtained from the alignment by the Maximum Likelihood method (Tamura Nei model, Tamura *et al.*, 1993). Branch lengths are proportional to sequence differences. Bootstrap was performed with 1,000 replicates. (B) Heatmap of the relative

abundance of recovered OTUs in heavy DNA fractions of [¹³C]-labeled microcosms. Labeled OTUs are shown in bold.

In the [¹³C]-CH₃Cl treatment, labeled OTU_mch_10 and OTU_mch_11 represent more than 90 % of *mch* sequences. These two OTUs had similar abundances (Table S4.2) and were also labeled in the corresponding treatment to which unlabeled CH₃OH was added together with [¹³C]-CH₃Cl. Conversely, when CH₃OH was labeled but not CH₃Cl, genotypes OTUs_mch_10 and OTUs_mch_11 remained labeled, but their relative abundance decreased compared to that of OTU_mch_9 (Figure 4.8B). Finally, in the microcosm to which only [¹³C]-labeled CH₃OH was added, the same 3 OTUs were labeled and also represented together over 80 % of the total *mch* sequences, although their relative abundances changed in the [¹³C]-CH₃OH treatment. The minor frequent genotype OTU_mch_3 was only labeled in this treatment.

4. Discussion

In this study, forest soil samples were pre-incubated with unlabeled CH₃Cl to allow subsequent efficient CH₃Cl assimilation, and identify potential chloromethane degraders. Notably, the chosen experimental strategy also prevents oxidation products (labeled CO₂) to be subsequently fixed by taxa unable to directly assimilate CH₃Cl. Indeed, out of the five labeled OTUs in [¹³C]-CH₃Cl and [¹³C]-CH₃OH amended microcosms, four are clearly not autotrophs, although some strains of *Pseudonocardiaceae* capable of utilizing CH₃OH have been described as facultative autotrophs (Grostern and Alvarez-Cohen, 2013). Our data suggest that the experimental setup with alternating cycles of labeling and aeration prevented detectable cross feeding via [¹³C]-CO₂ between taxa. During CH₃Cl labeling, a dilution of the labeling with unlabeled CH₃OH present in the forest soil might have occurred, although the high concentration of the C₁ compound used for amendment would in theory help minimize this process. Moreover, CO₂ production was very similar in all microcosms, as well as [¹³C]-CO₂ production from CH₃Cl and CH₃OH (Figure 4.2). Incidentally, this clearly indicates that soil microbiomes have much higher capacity for oxidation of carbon nutrients than that required for prevailing ambient concentrations of both methanol and chloromethane. In other words, obtaining evidence for differential production of CO₂ from CH₃Cl or CH₃OH would require even higher concentrations than the concentrations used in this study. Conversely, the used experimental setup, involving preincubation with high concentrations of chloromethane and relatively high substrate concentrations of methylotrophic substrates, may have led to overseeing active microorganisms growing with CH₃Cl and/or CH₃OH at ambient concentrations – a situation currently typical of the limitations of the SIP approach (Coyotzi *et al.*, 2016).

4.1. New insights into CH₃Cl utilizers in forest soil

One might have speculated that pre-incubation with CH₃Cl might have narrowed down the diversity compared to the enrichment and selection isolation protocols used previously (Miller et al., 2004; Borodina et al., 2005; Nadalig et al., 2011). However, the detected taxonomic diversity in soil microcosms was quite high and generally unaffected by further pulses of CH₃OH and CH₃Cl (Table 4.4, Figure 4.2). Moreover, among the 117 different taxonomic OTUs described at the family-level based on the 16S rRNA gene, only 5 could be conservatively defined as labeled (Table 4.5). They belong to the three phyla Proteobacteria, Actinobacteria, and TM7. The latter is a major bacterial lineage which has been recalcitrant to cultivation and isolation until recently (He et al., 2015). All five differentially detected OTUs have been previously isolated or detected in soils, but until now, were not associated with CH₃Cl degradation. It was also striking to find that these family-level OTUs, with the possible exception of the poorly defined TM7 phylum OTU, essentially consisted of sequences that could be ascribed to a highly dominant genus (Table 4.5). Hence, the main detected genus within Alphaproteobacteria could be tentatively affiliated to the genus Methylovirgula largely found in acidic forest soil and decaying deadwood (Hoppe et al., 2015). Streptomyces is found predominantly in soils and decaying plant material and taxonomically diverse in forest soils (Adams et al., 2011; Amore et al., 2012). Pseudonocardiaceae representatives have also been isolated from soils (Warwick et al., 1994). Acidothermus comprises obligate aerobes, some with cellulolytic activity, and found in soils (Talia et al., 2012). Strikingly, the genus Hyphomicrobium, previously shown to dominate CH₃Cl-utilizing communities (Borodina et al., 2005) and for which strains were also readily isolated from soils (Borodina et al., 2005) and the phyllosphere (Nadalig et al., 2011), was not prominent and not detected as labeled, although present in the present study (Figure S4.1). Pseudonocardiaceae feature the closest known cultured CH₃Cl-degrader, affiliated with the genus Norcardiodes: strain SAC-4, a forest top soil isolate unable to utilize CH₃OH for growth (McAnulla *et al.*, 2001). Thus, the diversity of currently known cultivated CH₃Cl-degraders is likely not representative of microbial CH₃Cl utilization in soils (Figure 4.5), as suggested previously (Schäfer *et al.*, 2007).

Of the five genus-level labeled phylotypes that were identified, only *Methylovirgula*, *Streptomyces*, and *Acidothermus* were labeled in microcosms that were solely amended with $[^{13}C]$ -CH₃Cl. This clearly demonstrates that strains from these taxa have the ability to assimilate CH₃Cl as a carbon source. Moreover, representatives of the TM7 phylum were found to be labeled in microcosms supplemented with $[^{13}C]$ -CH₃Cl in the presence of unlabeled CH₃OH, although they were detected at low relative abundance (0.3 to 1.3 %). This suggests that strains belonging to the TM7 phylum are also able to utilize CH₃Cl as a carbon source. Conversely, strains associated with the labeled family-level taxon *Pseudonocardiaceae* may rather be associated with strains incapable of carbon assimilation from CH₃Cl, since this phylotype was only labeled in the microcosm amended with $[^{13}C]$ -CH₃OH in presence of $[^{12}C]$ -CH₃Cl.

Hence, the relative abundance of the five-labeled phylotypes in the four microcosms amended with $[^{13}C]$ -CH₃Cl and/or CH₃OH (Figure 4.4; Table 4.5), revealed three major metabolic profiles associated with dominant phylotypes with regard to utilization of CH₃Cl in the investigated forest soil:

- *Streptomyces* and *Acidothermus*: CH₃Cl amendments alone or combined with CH₃OH always confer high relative abundance. Organisms that use only CH₃Cl and not CH₃OH for growth, as found for the cultivable *Aminobacter* species (Coulter *et al.*, 1999), label with [¹³C]-CH₃Cl. A particularity was observed in the case of *Acidothermus*. This OTU was labeled with [¹³C]-CH₃OH when unlabeled CH₃Cl was amended, which suggests that CH₃Cl may rather represent a source of energy for such organisms, and that CH₃OH is preferentially used as the carbon source for growth.

- *Methylovirgula*: CH₃Cl amendment lowers relative abundance. *Methylovirgula* was described to use CH₃OH produced during wood decomposition (Hope *et al.*, 2014, Kielak *et al.*, 2016). To date, CH₃Cl utilization in a *Methylovirgula* strain has not been described and, surprisingly, no genomes from this genus are yet publically available. Otherwise, *Methylovirgula* was defined as labeled in [¹³C]-CH₃Cl treatments

supplemented or not with unlabeled CH₃OH, as in the [¹³C]-CH₃OH condition. Its absence as a labeled taxon in the treatment with [¹³C]-CH₃Cl and [¹²C]-CH₃OH suggest that *Methylovirgula* representatives prefer CH₃OH over CH₃Cl as a carbon source. Cultivated *Alphaproteobacteria* which can utilize both CH₃OH and CH₃Cl simultaneously as carbon and energy source for growth have been described, and include well-characterized *Hyphomicrobium* and *Methylobacterium* strains (McDonald *et al.*, 2001). It can be speculated that the combination of CH₃Cl and CH₃OH could decrease growth rate of representatives of *Methylovirgula* strains, as indeed was observed *in vitro* with *Methylobacterium extorquens* CM4 (Sandro Roselli PhD thesis, 2009), and explain why *Methylovirgula* was not detected as a labeled OTU in the condition in which unlabeled CH₃Cl was added together with [¹³C]-CH₃OH.

- *Pseudonocardiaceae* and TM7 exhibit an intermediate metabolic profile compared to the other two profiles described above. No labeling was detected in treatments that were solely supplemented with one of the two C₁ compounds. Utilization of CH₃Cl as electron donor only but not as carbon source could potentially increase labeling with CH₃OH as found for *Pseudonocardia*. An extreme case of this scenario may be associated with the very high relative abundance of the *Microbacteriaceae* OTU, essentially consisting of the *Curtobacterium* genus, and only detected in amended microcosms with both [¹³C]-CH₃OH and unlabeled CH₃Cl (Figure 4.4, Table 4.5). Considering the very high (30 %) abundance this OTU reached under this condition, however, it is surprising that no increase whatsoever was observed for this OTU in any fraction of the corresponding unlabeled microcosm (<0.6 %, data not shown). The intriguing abundance profile of this OTU in microcosms of Steigerwald forest soil under different conditions will now need to be ascertained by independent approaches such as qPCR.

In the reverse scenario, use of CH₃Cl as carbon source, and CH₃OH for energy could increase labeling with CH₃Cl, as found for the TM7 phylum. The TM7 phylum was only found labeled in treatments in which [¹³C]-CH₃Cl was added together with unlabeled CH₃OH. Thus, methylotrophic organisms affiliated to the TM7 phylum might utilize CH₃Cl more for carbon assimilation, and CH₃OH more for energy production. The low abundance of the TM7 phylum family-level OTU also suggests that corresponding strains assimilate carbon from CH₃Cl

227

inefficiently or at a slow rate. A synergistic metabolism of CH₃OH and CH₃Cl utilization may also be envisaged in this context. This was reported for the marine methylotrophic bacterium HTCC2181 that used CH₃Cl as carbon source only in the presence of CH₃OH (Halsey *et al.*, 2012).

As a final note, metabolic inhibition by chloromethane cannot be ruled out. Use of unlabeled CH₃Cl could have a toxic effect, potentially resulting in reduction of labeling. For instance, chloromethane was demonstrated to inhibit ammonia monooxygenase, an essential enzyme of the obligate chemolithoautotrophic aerobe, *Nitrosomonas europaea*, which uses ammonia as its sole natural energy source (Rasche *et al.*, 1991; Keener and Arp, 1993). Conversely, as a fortuitous substrate of particular methane monooxygenase, chloromethane affects methylotrophic growth of *Methylomicrobium album* GB8 on CH₃OH (Han et Semrau, 2000).

4.2. *cmuA* OTUs and metabolic diversity in the environment

Eight *cmuA* OTUs were detected in this study, and five were labeled under different conditions (Figure 4.6B). The labeled *cmuA* OTUs were closely related to each other and clustered with cultured Alphaproteobacteria sequences of Methylobacterium and Hyphomicrobium as well as a few cmuA genotypes of uncultured bacteria that were previously detected in soil samples (Figure 4.6A; Miller et al., 2004: Borodina et al., 2005). OTU_cmuA_2 was detected as the most abundant OTU in all treatments. Taken together, in the studied forest soil, these data may be explained by the fact that few microorganism harbor cmuA-related sequences or alternatively, that the *cmuA* primers are inefficient to detect the *cmuA* diversity present in soils. Another explanation would be that *cmuA*-independent CH₃Cl utilization pathway predominate within CH₃Cl-degraders in soil (Nadalig et al., 2014). While OTU cmuA 3 was labeled in the microcosm amended with [13C]-CH₃Cl alone, only OTU_cmuA_2 was labeled when CH₃OH was also added. Conversely, in the microcosm where [¹³C]-CH₃OH and unlabeled CH₃Cl were added, the diversity of *cmuA* OTUs increased. Interestingly, OTUs cmuA 2, 3 and 4 were labeled in the microcosm treated with [¹³C]-CH₃OH alone. Taken together, these results tentatively suggest that *cmuA* OTUs may cover the four phylotypes of CH_3Cl -degraders: (i) strains with OTU_cmuA_2 or OTU_cmuA_3 may be able to use both CH₃Cl and CH₃OH as carbon and energy source; (ii) strains with OTU_cmuA_4 may be able to use CH₃OH as carbon and energy source, and CH₃Cl as energy source; (iii) strains with OTU_cmuA_6 may assimilate carbon from CH₃OH and use CH₃Cl as energy source; (iv) low abundance strains with OTU_cmuA_8 only found labeled in the microcosm labeled with $[^{13}C]$ -CH₃OH and to which CH₃Cl was also added may use CH₃OH as carbon source and CH₃Cl only for energy.

4.3. Chloromethane-associated methylotrophy

Soil methylotrophs were assessed using two types of biomarker genes, mxaF/xoxF for methanol oxidation, and mch for tetrahydrofolate-dependent C₁-interconversion reactions. In heavy fractions of the [¹³C]-CH₃Cl microcosms, OTUs MDH 2, 3 and 6, with a large dominance of the OTU_MDH_3 (relative abundance of 69.7 %) were labeled (Figure 4.7). MDH diversity shifted in response to CH₃OH supplementation. Thus, CH₃Cl utilization impacts the diversity of detected methanol dehydrogenase phylotypes, with labeled MDH OTUs all affiliated to xoxFsequences in this study (Figure 4.8). Similar effects of CH_3CI were also observed in the analysis of the 4 labeled mch OTUs. OTU mch 3 is most similar to the mch sequence of an uncultured organism from forest soil (Stacheter et al., 2013), whereas the other 3 labeled OTUs are clustered together but not affiliated to previously reported mch sequences of isolates or metagenomic datasets (Figure 4.8A). Methylotrophs harboring OTU mch 9 may use CH₃Cl only in presence of CH₃OH, whereas CH₃OH appears to affect the relative abundance of OTU_mch_10. On the contrary, the OTU_mch_3 was detected only in the [13C]-CH₃OH microcosm, suggesting that strains of this phenotype cannot use CH₃Cl. The obtained results are reflected into the phylogenetic tree, where OTU mch 9, 10 and 11 are closely related, whereas OTU_mch_3 is relatively divergent (Figure 4.8). Whatever the basis of these observations may be, the diversity of OTUs associated with genes involved in methylotrophy metabolism, as well as their dynamics as a function of amendment with unlabeled or labeled CH₃Cl, clearly suggest that the diversity of methylotrophic communities is affected by chloromethane.

5. <u>Conclusion</u>

This study shows for the first time that chloromethane consumption in forest soil may be affected by the key methylotrophic substrate methanol. This study also provides the first deep coverage exploration of the bacterial diversity associated with chloromethane metabolism in soil. A main outcome of this study is the uncovering of new taxa associated with CH₃Cl utilization, including specific genera from *Alphaproteobacteria* and *Actinomycetales* for which

no degrading strains have yet been isolated or associated with CH₃Cl consumption in previous studies. Conversely, taxa corresponding to previously isolated CH₃Cl-degraders, such as *Hyphomicrobium* and *Methylobacterium*, were not detected as effective CH₃Cl-utilizers in the studied forest soil, despite the fact that *cmuA* genes closely related to those of corresponding isolated were detected. This suggests the possibility of active organisms in soils with known chloromethane utilization genes acquired by horizontal transfer. Future efforts will require the use of complementary approaches to improve knowledge of CH₃Cl-utilizers in the environment, including experiments without CH₃Cl pre-incubation and the use of concentrations of CH₃Cl closer to those prevailing in the soil environment. Moreover, further work is required to address the relevance of the detected labeled OTUs in the environment and to characterize new isolates from corresponding taxa, which may also feature as yet unknown CH₃Cl degradation pathways other than the canonical *cmu* pathway.

6. Supplemental materials

Table S4.1. Summary of sequence data obtained in the SIP experiment

		16S rRNA rrsA gene reads		Functional gene reads							
Microcosm	DNA fraction ^a		Total	Filtered reads Raw reads ^b	_	Total filtered reads ^c			Filtered reads		
WICIOCOSIII		Raw reads ^b	filtered reads ^c		стиА	mch	mxaF_xoxF	стиА	mch	mxaF_xoxF	
	Н	47530		46110	96190				4282	55892	786
[¹³ C]-CH₃Cl	М	111914	306404	107946	58556	7438	130172	3356	2056	37524	1118
	L	157600		152348	54408				1100	36756	1452
	Н	183860		178042	74270				3778	38644	1534
	М	107282	331804	103360	83720	10320	124876	4794	4384	48252	1692
CH3OH	L	52098		50402	57828				2158	37980	1568
	Н	79510		76690	76850				614	52586	918
	М	173144	349990	167078	74756	11448	143134	3800	5584	44510	1394
[C]-CH3OH	L	110046		106222	72968				5250	46038	1488
	Н	100600		98332	72356				1886	43088	588
[¹³ C]-CH₃OH	М	94430	227354	91136	66856	6144	115594	2862	3384	41322	1086
	L	39066		37886	45940				874	31184	1188
	Н	74708		72522	57694				370	27732	2616
CH₃Cl	М	107742	248254	103850	77772	5438	125532	6692	1988	51800	2216
	L	74522		71882	70190				3080	46000	1860
	Н	49080		47236	75074				922	44356	862
CH ₃ OH + CH ₃ Cl	М	84564	219992	81572	46122	6524	118428	3844	2230	28196	1300
	L	93876		91184	70152				3372	45876	1682
	Н	93620		90822	50286				246	26518	2946
CH₃OH	М	51602	244402	49984	63712	4578	98696	5434	1872	43490	1422
	L	106682		103596	43526				2460	28688	1066
	Н	102994		100032	50886				184	26804	2710
Control	М	82628	240724	79894	64516	4690	113260	6154	876	42884	1772
	L	62760		60798	65778				3630	43572	1672

^a H, M and L: heavy, middle and light DNA fractions respectively

^b Sequences containing the intact barcode sequence and the forward primer sequence with less than 3 mismatches

^c Sum of filtered reads in fractions H, M and L for each microcosm

^d For 16S rRNA sequences, filtered reads were obtained from raw reads by following the SOP in mothur (www.mothur.org/wiki/MiSeq_SOP). Amplicons of functional genes were processed as described in Materials and Methods. Briefly, amplicons differing by more than 20 nucleotides from the expected size, sequences with a quality score lower than 20, and singletons were all removed. As a final step, OTUs for which the reference sequence did not match the target gene by BLAST were also discarded.

Table S4.2. Relative abundance (%) and labeled status of functional gene OTUs in heavy DNA fractions of [¹³C]-labeled microcosms

OTU/microcosm ^a	[¹³ C]-CH₃Cl	[¹³ C]-CH₃Cl + CH₃OH	CH₃Cl + [¹³ C]-CH₃OH	[¹³ C]-CH₃OH
OTU_cmuA_1	6.1	0.5	0.0	0.0
OTU_cmuA_2	63.8	83.8	42.7	64.1
OTU_cmuA_3	24.0	12.9	18.2	26.6
OTU_cmuA_4	4.0	1.4	3.3	7.4
OTU_cmuA_5	0.1	NR ^b	NR	NR
OTU_cmuA_6	2.1	1.4	35.2	1.9
OTU_cmuA_7	LR	LR	LR	LR
OTU_cmuA_8	LR	LR	0.7	LR
OTU_MDH_1	13.2	28.3	26.4	25.5
OTU_MDH_2	3.6	12.4	5.9	11.6
OTU_MDH_3	69.7	29.3	46.0	23.1
OTU_MDH_4	7.1	15.4	12.0	23.5
OTU_MDH_5	3.8	13.7	9.2	13.3
OTU_MDH_6	2.5	0.9	0.7	3.1
OTU_mch_1	0.1	0.1	LR	LR
OTU_mch_2	0.1	0.2	LR	LR
OTU_mch_3	0.2	0.3	0.5	3.7
OTU_mch_4	7.2	9.5	8.2	1.0
OTU_mch_5	0.1	0.1	0.1	LR
OTU_mch_6	LR	0.1	0.1	LR
OTU_mch_7	1.0	1.7	1.1	0.2
OTU_mch_8	LR	0.1	LR	LR
OTU_mch_9	LR	0.2	52.5	27.2
OTU_mch_10	32.3	31.9	15.4	23.5
OTU_mch_11	59.1	55.9	22.3	44.3
OTU_mch_12	LR	LR	LR	LR
OTU_mch_13	LR	LR	LR	LR
OTU_mch_14	LR	LR	LR	LR
OTU_mch_15	LR	LR	LR	LR
OTU_mch_16	LR	LR	LR	LR

^a OTUs defined as labeled in a given microcosm are shown in bold; ^b NR, no reads; ^c LR, low read number, relative abundance less than 0.04 %.



Figure S4.1. Relative abundance of family-level taxa in amplicon libraries of the 16S rRNA gene

Families (OTUs with >80 % sequence identity) which were defined as labeled by comparison of heavy DNA fractions of the labeled and corresponding unlabeled microcosms are framed.





H, M and L represent heavy, middle and light DNA fractions in the 8 microcosms. OTUs were defined at the 90 % sequence identity level, and OTUs defined as labeled (see Materials and Methods) are shown in bold.



Figure S4.3. Relative abundance of OTUs in amplicon libraries of the *mxaF* and *xoxF* genes H, M and L represent heavy, middle and light DNA fractions in the 8 microcosms. All OTUs at the 90 % sequence identity level were defined as labeled according to the defined criteria (see Materials and Methods) in microcosms incubated with [¹³C]-labeled compounds.





H, M and L represent heavy, middle and light DNA fractions from the 8 microcosms. OTUs at the 80 % sequence identity level shown in bold were defined as labeled as described in Materials and Methods.

Chapitre 5

Conclusion and perspectives

Chapter 5. Conclusions and perspectives

In this final Chapter, the major conclusions of my PhD studies are presented, and discussed in the context of the future experiments that I propose (Figure 5.1).



Figure 5.1. Overview of research avenues to assess the role of bacteria as chloromethane emission filters in forest soil

Four major tasks are proposed (ovals), and color-coded for experiments performed *in labo* (pink) or *in situ* (green). These tasks are based on culture-dependent and independent methods. As suggested by the arrows, these research axes are interconnected in an interactive observe/test/validate workflow.

In lab controlled experiments with cultivated model strains are needed for a better understanding of chloromethane metabolism by the different bacterial chloromethanedegrading isolates which are available. For instance, one can assess if these isolates are able to use chloromethane as the sole source of carbon and energy, as a source of carbon only, or as a source of energy only, and whether co-utilization of other carbon compounds found in forest soils occurs in the context of chloromethane degradation.

In-depth knowledge of catabolic pathways and adaptation to growth with chloromethane is needed to discover alternative biomarkers to obtain a full picture of the entire diversity of bacterial chloromethane degradation in forest soil.

Exploring without a *priori* the diversity of chloromethane-utilizers *in situ* using molecularbased techniques and discovering as yet unknown degradation pathways is also needed to uncover chloromethane-degrading ecotypes active *in situ* in their entirety.

Biomarkers are biological measurable indicators that are representative of some biological state, function or condition. For the questions at hand, biomarkers that report on microbial filtering of chloromethane in soils are needed.

1.News insights in chloromethane utilization deduced from the
study of the model strain *M. extorquens* CM4

1.1. In-depth study of the bacterial adaptive response to chloromethane utilization

Analysis of the genes that were found with higher transcript abundance in *M. extorquens* CM4 cultures grown with chloromethane compared to methanol allowed the identification of key genes in the adaptive response to growth with chloromethane. Hereafter, I will focus on three genes that were not correlated with growth with chloromethane previously.

- The *hppA* gene encoding a membrane-bound proton translocating pyrophosphatase. The energy of pyrophosphate hydrolysis may serve as the driving force for proton extrusion to build up a proton motive, force as demonstrated in other bacteria (Lopez-Marques *et al.*, 2004). The proton produced upon chloromethane dechlorination in the cytoplasm will be excreted to help maintain intracellular pH build up a proton motive force. Cells may thereby also be able to coexcrete chloride ions, as suggested by the study of *hppA* homologs demonstrated to contribute to salt resistance that were retrieved from the metagenome of brines and moderate-salinity rhizosphere (Mirete *et al.*, 2015). In the dichloromethane-degrading strain of the same species, *M. extorquens* DM4, *hppA* was also found with higher transcript abundance in cultures grown with dichloromethane. Thus, the *hppA* gene appears to be involved in adaptation to dechlorination reactions during growth with chlorinated methanes chloromethane and dichloromethane.

- **The** *pnt* **gene cluster encoding the transhydrogenase PntA**. The Pnt complex of PntAA, PntAB and PntB subunits allows the regeneration of NADP⁺ by transfer of reducing equivalents between NAD(H) and NADP(H) for translocation of protons across the membrane, as demonstrated in *M. extorquens* AM1 (Chou *et al.*, 2015). Basing on this study, it can be speculated that the NADP⁺ pool is a limiting factor for growth with chloromethane, since two molecules of NADP⁺ are needed in two successive reactions of the *cmu* pathway catalyzed by MetF2 and FoID (Fig. 3.1, Chapter 3). If this hypothesis is correct, NADP⁺ pools are predicted to be low in chloromethane-utilizing methylotrophs grown with chloromethane, as compared to other carbon substrates.

- The gene cluster encoding the acetone carboxylase subunits AcxA, AcxB, AcxC. There is growing evidence for a possible link between the *acxABC* gene cluster and chloromethane utilization. We observed higher expression of all three *acxABC* genes in the transcriptome of the chloromethane-grown strain. Previous comparative genomics revealed that the *acx* cluster is part of the shared variable genome of chloromethane-degrading strains of known genome sequence *M. extorquens* CM4, *Hyphomicrobium* sp. MC1, and in part for *Roseovarius* sp. 217 (Roselli *et al.*, 2013; Vuilleumier *et al.*, 2011). Strain CM4 was previously demonstrated to grow with acetone as the sole source of carbon and energy (Roselli *et al.*, 2013), but this has not been tested in other chloromethane-utilizers. Whether acetone, in comparison with other C₂ substrates, has a specific positive growth effect on growth with chloromethane and not with other C₁ compounds may be assessed in *M. extorquens* CM4 and other chloromethane-utilizing isolates. [¹³C]-methanol labeling and liquid chromatography mass spectrometry analyses have been used in the past to study the different fates of the carbon from two substrates that are co-utilized in *M. extorquens* (Peyraud *et al.*, 2012). A metabolomic approach using labeled [¹³C]-acetone could be envisaged to characterize metabolic intermediates for chloromethane utilization in presence of acetone (Tang *et al.*, 2012).

In the investigated conditions, the transcriptional responses of *M. extorquens* strains to growth with chloromethane or dichloromethane were mostly unrelated. However, high-throughput transcriptomic analysis only captures an averaged global snapshot of the response of bacteria in terms of gene expression at a given point in time. In my PhD project, all RNAs were extracted from cultures in mid-log growth phase in the presence of a unique source of carbon and energy. Assessing transcriptional patterns under growth conditions of chloromethane and methanol co-utilization, including high and low concentrations of growth substrates, may uncover genes not previously associated with growth with chloromethane, but relevant under environmental conditions.

1.2. Search of chloromethane-utilization-dependent regulatory elements

A bacterial cell needs to adapt to varying environments and nutrient availability, by quick, sensitive and effective responses (Francez-Charlot *et al.*, 2015; Balleza *et al.*, 2009). This may involve regulation processes at the transcriptional and translational levels, but also modulation of the metabolite pool (Figure 5.2). Relationships between gene and transcript abundance and corresponding protein-encoded processes are commonly assumed, yet rarely

observed (Rocca *et al.*, 2015). Basic research on regulatory elements and mechanisms that control chloromethane-utilization is still needed to establish predictive links between genomic, transcriptomic and proteomic data in ecosystem processes.



Figure 5.2. Central questions for future in lab experiments with model strains

Up to this day, the master regulator involved in transcriptional induction of the *cmu* gene remains unknown. Neither have the chloromethane-dependent genetic elements (promoter and associated regulating factors) been characterized. Different approaches are proposed to better understand the adaptive response to chloromethane utilization, at the DNA level (orange), transcriptional level (green), proteomic level (purple), and combined with a metabolite analysis (grey). For instance, dosage of NADP⁺ pools may address the question whether NADP⁺ pool is a limiting factor for growth with chloromethane. All the proposed approaches, including the comparison of data obtained in cultures grow with chloromethane alone, or provided with other C_1 or multi-carbon substrates, are needed to discover specific regulatory elements of chloromethane utilization.

The protein encoded by *fmdB* nearby *cmuA* was proposed as a putative regulator of chloromethane utilization in *Hyphomicrobium chloromethanicum* strain CM2, based on its very low amino acid identity (24 %) with the *Methylophilus methylotrophus* FmdB regulator protein (Wyborn *et al.*, 1996; McAnulla *et al.*, 2001). To our knowledge, no FmdB-like protein has been experimentally investigated. The corresponding gene in *M. extorquens* CM4 is also located adjacent to *cmuA*, and had increased transcript abundance in cultures grown with

chloromethane compared to methanol (Chapter 3). Among possible experiments that would help in finding the regulator of *cmu* genes, a targeted knockout of putative regulators would be a straightforward experiment to test their role in chloromethane-dependent gene regulation and growth. Unfortunately, as recently discovered, reverse genetics by allelic gene replacement does not work with M. extorquens CM4, contrary to other tested Methylobacterium strains (Marx et al., 2008), and unpublished data from both Strasbourg and Idaho laboratories. Also, knockout mutants have not been obtained so far in strain MC1, which belongs to the Hyphomicrobium genus, known to be recalcitrant to genetic analysis. Thus, another strategy to characterize the master regulator of the *cmu* pathway is needed. Developing a CRISPR/Cas9 system (Jakočiūnas et al., 2016) for Methylobacterium would be of great interest, but this still remains to be achieved. Alternatively, an experimental evolution approach of chloromethane utilization could also be envisaged. A previous experimental evolution was successfully developed for dichloromethane utilization. After transfer of a synthetic dcmA-based gene cassette for dichloromethane utilization, M. extorquens strains initially unable to degrade dichloromethane acquired the ability to grow on dichloromethane as the sole carbon and energy source (Michener et al., 2014). Accordingly, a synthetic gene cassette for chloromethane-degradation containing a minimal set of *cmu* pathway-encoding genes was constructed (Michener, Vuilleumier, Bringel, Marx, unpublished data). This cmu gene cassette was introduced by transformation into methylotrophic strains unable to grow with chloromethane, and the ability of the obtained transconjugants to grow with chloromethane was tested. This synthetic chloromethane-utilization gene cassette may be used to study the effect of genes of interest, including the *fmdB* gene, and by doing so, to assess their role in chloromethane utilization (Figure 5.2).

Post-transcriptional regulation has recently been increasingly demonstrated to be involved in modulation of gene expression in changing environments. It involves regulation elements which are for instance implied in translation of RNA into protein or RNA stability (Van Assche *et al.*, 2015). The majority of known mechanisms of regulation of translation target the translation initiation process (Duval *et al.*, 2015). Identification of 5' untranslated regions (UTRs) of mRNA would be a prerequisite to explore possible post-transcriptional regulation mechanisms. To find and annotate 5'UTRs, transcriptional start sites (TSS) and N-termini of proteins (to determine the translational start) need to be experimentally determined (as outlined in Figure 5.3).

243
The gene promoter region is composed of the TSS numbered +1 by convention, with two upstream sets of hexanucleotides-long consensus sequences elements approximately located at -10 and -35 nucleotides with an optimal spacing of 17 bp. The identification of TSS is now possible using high throughput sequencing in an approach called differential RNAseq (dRNAseq), with selective sequencing of primary transcripts (Sharma et al., 2010) (Figure 5.3A). In this PhD work, a dRNAseq approach has been initiated for *M. extorquens* CM4 cultures grown with chloromethane compared to methanol, as well as for *M. extorquens* DM4 grown with DCM compared to methanol. RNAs from *M. extorquens* CM4 and DM4 cultures grown with methanol or chlorinated methanes were extracted (a total of eight biological independent samples corresponding to 4 different conditions). Then, the DNase-treated RNA extracts were sent to Vertis Biotechnologie AG (Freising, Germany) for cDNA library construction and sequencing. The dRNAseq data of eight TSS banks will be mapped soon to the corresponding genomes in collaboration with S. Cruveiller (Génoscope, Evry, France). Comparative TSS mapping in closely-related strains is expected to provide a large overview of what is conserved, what is strain specific (Dugar et al., 2013), and what is specific to growth with chloromethane.

The identification of the N-terminal position of proteins may be performed by the "dN-TOP" approach (Figure 5.3B). The N-terminome protein analysis by the "dN-TOP" approach has already been validated in *M. extorquens* DM4 grown with dichloromethane and with methanol in collaboration with C. Carapito (Laboratoire de Spectrométrie de Masse BioOrganique or LSMBO, Strasbourg, France).

The combined RNAseq and proteomic approaches will improve genome annotation and our understanding of genome expression of chloromethane-degrading reference strains grown in controlled lab conditions of C_1 and chloromethane utilization.



Figure 5.3. Exploration of chloromethane-dependent regulation mechanisms in model strains

A) Transcription Start Site (TSS) identification using dRNAseq (Sharma and Vogel 2014). Sequencing yields enrichment patterns of the cDNA coverage at the 5' end of primary transcripts which can be used to annotate TSS. The method is based on the fact that the bacterial cellular RNA pools consists of processed or cleaved transcripts characterized by a 5'monophosphate end (5'P), which is digested by the 5'P-dependent terminator exonuclease (TEX), while primary transcripts which have 5' triphosphate (5'PPP) or, less frequently, a 5' hydroxyl (5'OH) group are resistant to TEX (Evguenieva-Hackenberg and Klug 2011). Two libraries are prepared in parallel. Library – (in blue) captures all RNAs. Library + (in red) has been enriched for primary transcripts (Sharma and Vogel, 2014);

B) N-terminome protein analysis by the "dN-TOP" approach. The trimethoxyphenyl phosphonium (TMPP) labeling approach is an efficient N-terminomic approach that allows the characterization of both N-terminal and internal peptides in a single experiment (Bertaccini *et al.*, 2013);

C) Integrative data analysis will, among other discoveries, facilitate in the annotation of 5'UTR. The difference of the +1 of the transcription compared to N-terminal position gives access to the 5' UTR of mRNAs, involved in post transcriptional regulation (Duval *et al.*, 2015; Van Assche *et al.*, 2015).

2. Identification of chloromethane-utilizing ecotypes in forest soil microcosms

The DNA-SIP approach gives access to active microorganisms independently of their abundance in soil and their cultivability (Coyotzi *et al.*, 2016). During my PhD thesis, I used [¹³C]-labeled chloromethane, alone or in combination with [¹³C]-labeled methanol to assess the role of co-utilization of different key substrates of methylotrophic metabolism (Chapter 4), and thereby identified at least three new phylotypes associated with utilization of chloromethane as a carbon and/or energy source in forest soil. In addition to DNA-SIP, RNAs have been extracted from the labeled microcosms and reverse transcribed into cDNA. RNA-SIP is more sensitive than DNA-SIP, does not require cell division to labeled RNAs, and has already been used for microbial soil studies (Leng *et al*, 2016). Unfortunately, with the cDNA used as template, no PCR amplification was observed with primer sets targeting *cmuA*, *mch* and *mxaF/xoxF*, and I did not have the chance to attempt these experiments again.

For efficient [¹³C]-labeling of chloromethane-utilizers in forest soil, a pre-incubation step prior to [¹³C]-labeling was performed. In addition, high chloromethane concentrations were used. Within this experimental setup, underestimation of the diversity of chloromethane-utilizers is a possibility. A new SIP labeling strategy has been proposed recently in the literature using H₂¹⁸O (Rettedal *et al.*, 2015). H₂¹⁸O is perhaps the only universal "substrate" targeting all active microorganisms in a given environment. A dual labeling with low [¹³C]-CH₃Cl and H₂¹⁸O would in principle be more effective to assess among the active microorganisms the diversity of chloromethane utilizers at the prevailing low ambient forest soil chloromethane concentrations. Use of high levels of labeled [¹³C]-CH₃Cl concentrations nevertheless allowed the identification of new chloromethane-associated phylotypes among families that have not been described so far to feature representatives capable of degrading chloromethane (Chapter 4). To validate and characterize if and how these phylotypes utilize chloromethane, additional experiments need to be performed. Cultural approaches may give access to these phylotypes for detailed characterization of their metabolism in the laboratory. For instance, physiological characteristics such as generation times upon growth with chloromethane and co-utilization abilities with other carbon sources could easily be assessed. Dehalogenation kinetics can be estimated by dosage of chloride production in resting-cells as previously performed in the laboratory for different chloromethane-degrading isolates of the phyllosphere (Nadalig et al., 2011; S. Roselli PhD thesis, 2009). Additional molecular tools such as PCR amplification or Southern hybridization with a *cmuA* probe may provide information about cmuA occurrence and cmu gene organization. Evidence for new pathways of chloromethane utilization can be suggested by chloromethane isotope fractionation signatures (H and C), as demonstrated for Leisingera methylohalidovorans MB2 (Nadalig et al., 2014). Genome sequencing may also help predict methylotrophic central metabolism and inform on the presence of the *cmu* pathway. If genetic tools for the new isolates are available or can be developed (gene manipulation, cloning, site-directed mutagenesis, transformation, conjugation), they will greatly facilitate the identification and characterization of new genes involved in growth with chloromethane, as indeed was the case for characterization of the cmu pathway in strain CM4 (Vannelli et al., 1998, 1999). However, many strains of interest may not be cultivable under laboratory conditions and are not amenable to genetic investigations. Thus, global culture-independent approaches based on direct content studies of proteins (metaproteomics), RNAs (metatranscriptomics), and DNA (metagenomics) are increasingly successful in many environmental studies (Blagodatskaya et al., 2013). In addition and in the case of a soil environment such as the forest soil investigated here, Fluorescence Hybridization In Situ (FISH) has also been used successfully to quantify taxa abundance (Dumont and Murrell 2005). To target active microorganisms in soil, a combination of FISH and microautoradiography (MAR) called FISH-MAR has been developed (Hirsch et al., 2010). This technique involves a short incubation time (to avoid cross feeding) of the environmental sample with radioactively labeled substrate. Then, if a bacterium is detected with both the fluorescent and radioactive probes under the microscope, this implies that it consumes the labeled substrate. Notably, the FISH-MAR technique is more sensitive than DNA-SIP, since the radiotracer is incorporated into all macromolecules (nucleic acids, proteins, lipids) and not only into nucleic acids (Wagner et al., 2006). However, soil is a complex and heterogeneous system, with autofluorescent properties and large amounts of particles of different sizes, so that its direct imaging is challenging (Vos et al., 2013). In addition, the use of radiotracers is expensive, requires specialized training, registered facilities, dedicated instruments, and generates potential hazardous wastes. An alternative way to avoid radioelement-based experiments is to combine FISH with flow cytometry for single-cell isolation and study of individual cells (Kalyuzhnaya et al., 2006). Nevertheless, for single-cell approaches to work with soil samples, soil particles need to be minimized to a minimal amount for efficient image analysis. Coupling phylogenetic identity (with FISH) and metabolic function (using stable isotope labeling incubation experiment) of single cells is possible for complex bacterial communities from the environment. In particular, those labeling technics coupled with a highresolution secondary ion mass spectrometry (NanoSims) has been recently developped to investigate soils. This allows to link a high resolution microscopy with isotopy, providing information on molecular and isotopic compositions of the investigated microorganisms in their native environment as already described in marine environments (Polerecky *et al.*, 2012). However, NanoSIMS has only recently been applied to studies of bacterial activity in soil ecosystems, when it has been possible to efficiently dissociate bacterial cells from soil particles (Eichorst *et al.*, 2015). Ideally, single-cell studies of incubations with [¹³C]-CH₃Cl followed by NanoSIMS mass spectrometry analysis could be used to investigate the role of bacteria in chloromethane consumption in forest soil with unprecedented detail.

3. Assessing the diversity of chloromethane-utilizing pathway in soil

Besides the well-characterized oxic degradation of chloromethane mediated by the cmu pathway (Vannelli et al., 1999), at least one other pathway for chloromethane assimilation under oxic conditions exists. No cmuA-like homologs was detected in the genome of Leisingera methylohalidovorans, a strain affiliated to Alphaproteobacteria and isolated from a marine environment, and able to grow with chloromethane (Schaefer 2002; Nadalig et al., 2014). How L. methylohalidovorans and possibly related labeled chloromethane-utilizing ecotypes in forest soil grow with chloromethane, the genes involved and their connection to methylotrophic metabolism, remains to be assessed. More generally, a quantitative PCR (qPCR) analysis targeting 16S rRNA-encoding gene and *cmuA* gene to quantify in native soil samples and $[^{13}C]$ -CH₃Cl-incubated soil, the ratio between *cmuA* copy number as a proxy for *cmuA*-dependent chloromethane utilization potential and 16S rRNA copy number as a proxy for bacterial abundance of labeled ecotypes, remains to be performed. The absence of correlation for a given ecotype active in chloromethane-degradation in forest soil could then be taken to suggest a *cmuA*-independent pathway for chloromethane utilization. Methods based on DNA amplification to assess the microbial diversity in forest soil are quite challenging, since humic acids bind DNA and interfere with Taq polymerase in PCR amplification reactions, and should be combined with "omics" approaches (Coyotzi et al., 2016). Thus, funds permitting, high throughout sequencing of metagenomes extracted from microbial communities of native forest soil or [¹³C]-heavy fractions of SIP samples, should represent an approach of choice to allow the direct capture and analysis of soil microbial communities (Lee *et al.*, 2004). This global approach would enable the inventory of chloromethane utilization genes diversity in forest soil, and give access to *cmuA* gene in its entirety (including canonical and more divergent sequences), as well as the search of alternative chloromethane-degrading pathway-encoding signature genes.

However, the abundance of a gene within a metagenome does not suffice to assess its expression (RNA-level information) and the associated enzymatic activity (protein-level information). A direct quantification of CmuA protein may constitute more convincing evidence for occurrence of *in situ* chloromethane dehalogenation in forest soil. Mass spectrometry is indeed a very sensitive technique (Arsène-Ploetze *et al.*, 2015), which has already been applied in investigations of forest soil (Mancuso *et al.*, 2015). Molecular culture-independent methods such as this one may provide novel information about relevant gene products and proteins (Bastida *et al.*, 2009). In addition, the bacterial fluorescent bioreporter developed by the team (Figure 1.7) (Farhan UI Haque *et al.*, 2013) may help to identify hotspots of chloromethane production in forest soil, and increase the chance to identify chloromethane degraders.

4. <u>Need for new functional biomarkers for evaluation of the</u> microbial chloromethane sinks

Several functional genes have been used as biomarkers of methylotrophic metabolism in environmental studies, including in forest soil. For example, genes *pmoA* and *mmoX*, encoding membrane-associated and soluble methane monooxygenases, respectively, were used as biomarkers to establish the diversity of methanotrophs involved in trichloroethylene degradation in soil (Shukla *et al.*, 2009).

With regard to chloromethane utilization, the *cmuA* gene is still the best functional gene marker available. Nevertheless, there is growing evidence of *cmuA*-independent chloromethane utilization (Nadalig *et al.*, 2014), confirming that the gene *cmuA* cannot adequately cover the entire diversity of chloromethane degraders. In some environments in which other methyl halides are present, *cmuA* may be inadequate as a specific biomarker for chloromethane. For instance, *cmuA* will also be associated to bacterial bromomethane utilization in marine environments (Cox *et al.*, 2012). Thus, a larger toolbox of biomarkers of microbial chloromethane degradation indication is required to investigate the diversity and distribution of chloromethane-utilizing microorganisms in the environment. In Chapter 4,

potential new chloromethane-utilizing taxa were identified. Culture-based, or cultureindependent characterization of genes involved in chloromethane utilization by such ecotypes may provide new biomarkers which might be more representative of the diversity of chloromethane utilizers in forest soil and complementary to the "historical" *cmuA* gene marker.

Defining novel biomarkers for the soil environment appears especially worthwhile considering the now strong evidence for bacteria-driven chloromethane degradation in soil (Borodina *et al.*, 2005; Miller *et al.*, 2004; and Chapter 4), and its implications for the role of bacteria in mitigating chloromethane emissions from soil to the atmosphere. The diversity and abundance of active chloromethane-utilizing ecotypes remains to be further investigated with in-depth genetic investigations. Finally, quantitative evaluation of the microbial contribution of terrestrial chloromethane sinks may be facilitated by chromatography/mass spectrometry/isotope ratio mass spectrometry (GC-MS-IRMS) and measurements of δ^{13} C and δ^{2} H values of chloromethane (Keppler *et al.*, 2005; Greule *et al.*, 2012, Nadalig *et al.*, 2014), as in the ongoing ANR-DFG Chlorofilter project associating the Keppler group in Heidelberg with both partner laboratories of my PhD project.

Bibliography

- Adams, A.S., Jordan, M.S., Adams, S.M., Suen, G., Goodwin, L.A., Davenport, K.W., Currie, C.R., and Raffa, K.F. (2011). Cellulose-degrading bacteria associated with the invasive woodwasp Sirex noctilio. ISME J. 5, 1323–1331.
- Amore, A., Pepe, O., Ventorino, V., Birolo, L., Giangrande, C., and Faraco, V. (2012). Cloning and recombinant expression of a cellulase from the cellulolytic strain *Streptomyces* sp. G12 isolated from compost. Microb. Cell Factories *11*, 1.
- Anke, H., and Weber, R.W.S. (2006). White-rots, chlorine and the environment a tale of many twists. Mycologist *20*, 83–89.
- Antony, C.P., Doronina, N.V., Boden, R., Trotsenko, Y.A., Shouche, Y.S., and Murrell, J.C. (2012). *Methylophaga lonarensis* sp. nov., a moderately haloalkaliphilic methylotroph isolated from the soda lake sediments of a meteorite impact crater. Int. J. Syst. Evol. Microbiol. *62*, 1613–1618.
- Arsène-Ploetze, F., Bertin, P.N., and Carapito, C. (2015). Proteomic tools to decipher microbial community structure and functioning. Environ. Sci. Pollut. Res. *22*, 13599–13612.
- Bai, Y., Müller, D.B., Srinivas, G., Garrido-Oter, R., Potthoff, E., Rott, M., Dombrowski, N., Münch, P.C., Spaepen, S., Remus-Emsermann, M., *et al.* (2015). Functional overlap of the *Arabidopsis* leaf and root microbiota. Nature *528*, 364–369.
- Balleza, E., López-Bojorquez, L.N., Martínez-Antonio, A., Resendis-Antonio, O., Lozada-Chávez,
 I., Balderas-Martínez, Y.I., Encarnación, S., and Collado-Vides, J. (2009). Regulation by
 transcription factors in bacteria: beyond description. FEMS Microbiol. Rev. 33, 133–151.
- Bastida, F., Moreno, J.L., Nicolás, C., Hernández, T., and García, C. (2009). Soil metaproteomics: a review of an emerging environmental science. Significance, methodology and perspectives. Eur. J. Soil Sci. *60*, 845–859.
- Bastviken, D., Thomsen, F., Svensson, T., Karlsson, S., Sandén, P., Shaw, G., Matucha, M., and Öberg, G. (2007). Chloride retention in forest soil by microbial uptake and by natural chlorination of organic matter. Geochim. Cosmochim. Acta *71*, 3182–3192.
- Beck, D.A.C., McTaggart, T.L., Setboonsarng, U., Vorobev, A., Goodwin, L., Shapiro, N., Woyke, T., Kalyuzhnaya, M.G., Lidstrom, M.E., and Chistoserdova, L. (2015). Multiphyletic origins of methylotrophy in *Alphaproteobacteria*, exemplified by comparative genomics of Lake Washington isolates: Genomics of alphaproteobacterial methylotrophs. Environ. Microbiol. *17*, 547–554.
- Bertaccini, D., Vaca, S., Carapito, C., Arsène-Ploetze, F., Van Dorsselaer, A., and Schaeffer-Reiss, C. (2013). An improved stable isotope N-terminal labeling approach with light/heavy TMPP to automate proteogenomics data validation: dN-TOP. J. Proteome Res. 12, 3063–3070.
- Blagodatskaya, E., and Kuzyakov, Y. (2013). Active microorganisms in soil: Critical review of estimation criteria and approaches. Soil Biol. Biochem. *67*, 192–211.

- Blom, D., Fabbri, C., Connor, E.C., Schiestl, F.P., Klauser, D.R., Boller, T., Eberl, L., and Weisskopf, L. (2011). Production of plant growth modulating volatiles is widespread among rhizosphere bacteria and strongly depends on culture conditions: Volatilemediated impact of bacteria on *Arabidopsis thaliana*. Environ. Microbiol. 13, 3047–3058.
- Boden, R., Thomas, E., Savani, P., Kelly, D.P., and Wood, A.P. (2008). Novel methylotrophic bacteria isolated from the River Thames (London, UK). Environ. Microbiol. *10*, 3225–3236.
- Bonfante, P., and Anca, I.A. (2009). Plants, mycorrhizal fungi, and bacteria: A network of interactions. Annu. Rev. Microbiol. *63*, 363–383.
- Borodina, E., Cox, M.J., McDonald, I.R., and Murrell, J.C. (2005). Use of DNA-stable isotope probing and functional gene probes to investigate the diversity of methyl chlorideutilizing bacteria in soil. Environ. Microbiol. *7*, 1318–1328.
- Bosch, G., Skovran, E., Xia, Q., Wang, T., Taub, F., Miller, J.A., Lidstrom, M.E., and Hackett, M. (2008). Comprehensive proteomics of *Methylobacterium extorquens* AM1 metabolism under single carbon and nonmethylotrophic conditions. Proteomics *8*, 3494–3505.
- Brantl, S. (2007). Regulatory mechanisms employed by cis-encoded antisense RNAs. Curr. Opin. Microbiol. 10, 102–109.
- Bringel, F., and Couée, I. (2015). Pivotal roles of phyllosphere microorganisms at the interface between plant functioning and atmospheric trace gas dynamics. Front. Microbiol. 6.
- Brophy, J.A., and Voigt, C.A. (2016). Antisense transcription as a tool to tune gene expression. Mol. Syst. Biol. *12*, 854–854.
- Brunner, W., Staub, D., and Leisinger, T. (1980). Bacterial degradation of dichloromethane. Appl. Environ. Microbiol. *40*, 950–958.
- Campbell, M.M., and Sederoff, R.R. (1996). Variation in lignin content and composition (mechanisms of control and implications for the genetic improvement of plants). Plant Physiol. *110*, 3.
- Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S.M., Betley, J., Fraser, L., Bauer, M., *et al.* (2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J. *6*, 1621–1624.
- Chang, H., Replogle, J.M., Vather, N., Tsao-Wu, M., Mistry, R., and Liu, J.M. (2015). A *cis* regulatory antisense RNA represses translation in *Vibrio cholerae* through extensive complementarity and proximity to the target locus. RNA Biol. *12*, 136–148.
- Chen, Y., and Murrell, J.C. (2010). When metagenomics meets stable-isotope probing: progress and perspectives. Trends Microbiol. *18*, 157–163.
- Chistoserdova, L., Vorholt, J.A., Thauer, R.K., and Lidstrom, M.E. (1998). C₁ transfer enzymes and coenzymes linking methylotrophic bacteria and methanogenic *Archaea*. Science *281*, 99–102.

- Chistoserdova, L., Chen, S.-W., Lapidus, A., and Lidstrom, M.E. (2003). Methylotrophy in *Methylobacterium extorquens* AM1 from a genomic point of view. J. Bacteriol. *185*, 2980–2987.
- Chistoserdova, L., Laukel, M., Portais, J.-C., Vorholt, J.A., and Lidstrom, M.E. (2004). Multiple formate dehydrogenase enzymes in the facultative methylotroph *Methylobacterium extorquens* AM1 are dispensable for growth on methanol. J. Bacteriol. *186*, 22–28.
- Chistoserdova, L., Crowther, G.J., Vorholt, J.A., Skovran, E., Portais, J.C., and Lidstrom, M.E. (2007). Identification of a fourth formate dehydrogenase in *Methylobacterium extorquens* AM1 and confirmation of the essential role of formate oxidation in methylotrophy. J. Bacteriol. 189, 9076–9081.
- Chistoserdova, L., Kalyuzhnaya, M.G., and Lidstrom, M.E. (2009). The expanding world of methylotrophic metabolism. Annu. Rev. Microbiol. *63*, 477–499.
- Chistoserdova, L. (2011). Modularity of methylotrophy, revisited. Environ. Microbiol. 13, 2603–2622.
- Chistoserdova, L., and Lidstrom, M. (2013). Aerobic methylotrophic prokaryotes. In The Prokaryotes, pp. 267–285.
- Chou, H.H., Marx, C.J., and Sauer, U. (2015). Transhydrogenase promotes the robustness and evolvability of *Escherichia coli* deficient in NADPH production. PLoS Genet. *11*, e1005007.
- Chu, F., and Lidstrom, M.E. (2016). XoxF acts as the predominant methanol dehydrogenase in the Type I methanotroph *Methylomicrobium buryatense*. J. Bacteriol. *198*, 1317–1325.
- Clerbaux, C., Cunnold, D.M., Anderson, J., Engel, A., Fraser, P.J., Mahieu, E., Manning, A., Miller, J., Montzka, S.A., Nassar, R., *et al.* (2007). Long-lived organic compounds (Chapter 1). Sci. Assess. Ozone Deplet. 2006 1–63.
- Coulter, C., Hamilton, J.T.G., McRoberts, W.C., Kulakov, L., Larkin, M.J., and Harper, D.B. (1999). Halomethane: bisulfide/halide ion methyltransferase, an unusual corrinoid enzyme of environmental significance isolated from an aerobic methylotroph using chloromethane as the sole carbon source. Appl. Environ. Microbiol. *65*, 4301–4312.
- Cox, M.L., Sturrock, G.A., Fraser, P.J., Siems, S.T., Krummel, P.B., and O'Doherty, S. (2003). Regional sources of methyl chloride, chloroform and dichloromethane identified from AGAGE observations at Cape Grim, Tasmania, 1998–2000. J. Atmospheric Chem. 45, 79– 99.
- Cox, M.J., Schäfer, H., Nightingale, P.D., McDonald, I.R., and Murrell, J.C. (2012). Diversity of methyl halide-degrading microorganisms in oceanic and coastal waters. FEMS Microbiol. Lett. *334*, 111–118.
- Coyotzi, S., Pratscher, J., Murrell, J.C., and Neufeld, J.D. (2016). Targeted metagenomics of active microbial populations with stable-isotope probing. Curr. Opin. Biotechnol. 41, 1–8.

- Crowther, G.J., Kosály, G., and Lidstrom, M.E. (2008). Formate as the main branch point for methylotrophic metabolism in *Methylobacterium extorquens* AM1. J. Bacteriol. *190*, 5057–5062.
- Cui, J., Good, N.M., Hu, B., Yang, J., Wang, Q., Sadilek, M., and Yang, S. (2016). Metabolomics revealed an association of metabolite changes and defective growth in *Methylobacterium extorquens* AM1 overexpressing *ecm* during growth on methanol. PloS one *11*, e0154043.
- Dallinger, A., and Horn, M.A. (2014). Agricultural soil and drilosphere as reservoirs of new and unusual assimilators of 2,4-dichlorophenol carbon: CAP-SIP of 2,4-DCP degraders in soil and drilosphere. Environ. Microbiol. *16*, 84–100.
- David, M., Bringel, F., Pagni, M., Gilmartin, N., BOUBAKRI, H., Nadalig, T., Simonet, P., Vogel, T., Vuilleumier, S. (2008). Diversité et évolution des déshalogénases bactériennes : détection bioinformatique et perspectives de recherche. In: *7ème colloque national "Ressources génétiques"* (p. 83-94). Presented at 7ème colloque national "Ressources génétiques", Strasbourg, FRA (2008-10-13).
- Debussche, L., Thibaut, D., Cameron, B., Crouzet, J., and Blanche, F. (1993). Biosynthesis of the corrin macrocycle of coenzyme B₁₂ in *Pseudomonas denitrificans*. J. Bacteriol. *175*, 7430–7440.
- Dekker, R.F., Vasconcelos, A.-F.D., Barbosa, A.M., Giese, E.C., and Paccola-Meirelles, L. (2001). A new role for veratryl alcohol: regulation of synthesis of lignocellulose-degrading enzymes in the ligninolytic ascomyceteous fungus, *Botryosphaeria* sp.; influence of carbon source. Biotechnol. Lett. 23, 1987–1993.
- DeLeon-Rodriguez, N., Lathem, T.L., Rodriguez-R, L.M., Barazesh, J.M., Anderson, B.E., Beyersdorf, A.J., Ziemba, L.D., Bergin, M., Nenes, A., and Konstantinidis, K.T. (2013). Microbiome of the upper troposphere: Species composition and prevalence, effects of tropical storms, and atmospheric implications. Proc. Natl. Acad. Sci. U. S. A. 110, 2575– 2580.
- Delmotte, N., Knief, C., Chaffron, S., Innerebner, G., Roschitzki, B., Schlapbach, R., von Mering,
 C., and Vorholt, J.A. (2009). Community proteogenomics reveals insights into the physiology of phyllosphere bacteria. Proc. Natl. Acad. Sci. *106*, 16428–16433.
- Denman, S.E., Tomkins, N.W., and McSweeney, C.S. (2007). Quantitation and diversity analysis of ruminal methanogenic populations in response to the antimethanogenic compound bromochloromethane: Monitoring of rumen methanogenic *Archaea*. FEMS Microbiol. Ecol. 62, 313–322.
- Donnelly, M.I., and Dagley, S. (1980). Production of methanol from aromatic acids by *Pseudomonas putida*. J. Bacteriol. 142, 916–924.
- Doronina, N.V., Sokolov, A.P., and Trotsenko, Y.A. (1996). Isolation and initial characterization of aerobic chloromethane-utilizing bacteria. FEMS Microbiol. Lett. *142*, 179–183.
- Dourado, M.N., Aparecida Camargo Neves, A., Santos, D.S., and Araújo, W.L. (2015). Biotechnological and agronomic potential of endophytic pink-pigmented methylotrophic *Methylobacterium* spp.. BioMed Res. Int. 2015, 1–19.

- Dugar, G., Herbig, A., Förstner, K.U., Heidrich, N., Reinhardt, R., Nieselt, K., and Sharma, C.M. (2013). High-resolution transcriptome maps reveal strain-specific regulatory features of multiple *Campylobacter jejuni* isolates. PLoS Genet. *9*, e1003495.
- Dumont, M.G., and Murrell, J.C. (2005). Stable isotope probing—linking microbial identity to function. Nat. Rev. Microbiol. *3*, 499–504.
- Dumont, M.G., Pommerenke, B., Casper, P., and Conrad, R. (2011). DNA-, rRNA- and mRNAbased stable isotope probing of aerobic methanotrophs in lake sediment: Stable isotope probing of methanotrophs. Environ. Microbiol. *13*, 1153–1167.
- Duval, M., Simonetti, A., Caldelari, I., and Marzi, S. (2015). Multiple ways to regulate translation initiation in bacteria: Mechanisms, regulatory circuits, dynamics. Biochimie *114*, 18–29.
- Edgar, R.C. (2010). Search and clustering orders of magnitude faster than BLAST. Bioinformatics *26*, 2460–2461.
- Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., and Knight, R. (2011). UCHIME improves sensitivity and speed of chimera detection. Bioinformatics *27*, 2194–2200.
- Eichorst, S.A., Strasser, F., Woyke, T., Schintlmeister, A., Wagner, M., and Woebken, D. (2015). Advancements in the application of NanoSIMS and Raman microspectroscopy to investigate the activity of microbial cells in soils. FEMS Microbiol. Ecol. *91*, fiv106.
- Erb, T.J., Rétey, J., Fuchs, G., and Alber, B.E. (2008). Ethylmalonyl-CoA mutase from *Rhodobactsr sphaeroides* defines a new subclade of conzyme B12-dependent Acyl-CoA mutases. J. Biol. Chem. 283, 32283–32293.
- Evershed, R.P., Crossman, Z.M., Bull, I.D., Mottram, H., Dungait, J.A., Maxfield, P.J., and Brennand, E.L. (2006). ¹³C-Labelling of lipids to investigate microbial communities in the environment. Curr. Opin. Biotechnol. *17*, 72–82.
- Evguenieva-Hackenberg, E. (2005). Bacterial ribosomal RNA in pieces: rRNA fragmentation in bacteria. Mol. Microbiol. *57*, 318–325.
- Evguenieva-Hackenberg, E., and Klug, G. (2011). New aspects of RNA processing in prokaryotes. Curr. Opin. Microbiol. 14, 587–592.

Fall, R., and Benson, A. (1996). Leaf methanol-the simplest natural product from plants. Trends in Plant Science, 1(9), 296-301.

- Farhan UI Haque, M., Nadalig, T., Bringel, F., Schaller, H., and Vuilleumier, S. (2013). Fluorescence-based bacterial bioreporter for specific detection of methyl halide emissions in the environment. Appl. Environ. Microbiol. 79, 6561–6567.
- Fiddaman, P.J., and Rossall, S. Effect of substrate on the production of antifungal volatiles from *Bacillus subtilis*. J. Appli. Bacteriol. *76*, 395-405.
- Fitriyanto, N. A., Mako F., Mika M., Ambar P., Tomonori I., et Keiichi Kawai. (2011). Molecular structure and gene analysis of Ce³⁺-Induced methanol dehydrogenase of *Bradyrhizobium* Sp. MAFF211645. J. Biosco. Bioeng. *111*, 613-617.

- Forczek, S.T., Laturnus, F., Doležalová, J., Holík, J., and Wimmer, Z. (2015). Emission of climate relevant volatile organochlorines by plants occurring in temperate forests. Plant Soil Environ. *61*, 103–108.
- Francez-Charlot, A., Frunzke, J., Reichen, C., Ebneter, J.Z., Gourion, B., and Vorholt, J.A. (2009). Sigma factor mimicry involved in regulation of general stress response. Proc. Natl. Acad. Sci. *106*, 3467–3472.
- Francez-Charlot, A., Kaczmarczyk, A., Fischer, H.-M., and Vorholt, J.A. (2015). The general stress response in *Alphaproteobacteria*. Trends Microbiol. *23*, 164–171.
- Frascari, D., Zanaroli, G., and Danko, A.S. (2015). In situ aerobic cometabolism of chlorinated solvents: a review. J. Hazard. Mater. *283*, 382–399.
- Freedman, D.L., Swamy, M., Bell, N.C., and Verce, M.F. (2004). Biodegradation of chloromethane by *Pseudomonas aeruginosa* strain NB1 under nitrate-reducing and aerobic conditions. Appl. Environ. Microbiol. 70, 4629–4634.
- Frey-Klett, P., Garbaye, J., and Tarkka, M. (2007). The mycorrhiza helper bacteria revisited. New Phytol. *176*, 22–36.
- Galbally, I.E., and Kirstine, W. (2002). The production of methanol by flowering plants and the global cycle of methanol. J. Atmospheric Chem. *43*, 195–229.
- Gälli, R., and Leisinger, T. (1985). Specialized bacterial strains for the removal of dichloromethane from industrial waste. Conserv. Recycl. *8*, 91–100.
- Georg, J., and Hess, W.R. (2011). Cis-antisense RNA, another level of gene regulation in bacteria. Microbiol. Mol. Biol. Rev. *75*, 286–300.
- Goodwin, K.D., Lidstrom, M.E., and Oremland, R.S. (1997). Marine bacterial degradation of brominated methanes. Environ. Sci. Technol. *31*, 3188–3192.
- Gourion, B., Rossignol, M., and Vorholt, J.A. (2006). A proteomic study of *Methylobacterium extorquens* reveals a response regulator essential for epiphytic growth. Proc. Natl. Acad. Sci. *103*, 13186–13191.
- Greule, M., Huber, S.G., and Keppler, F. (2012). Stable hydrogen-isotope analysis of methyl chloride emitted from heated halophytic plants. Atmos. Environ. *62*, 584–592.
- Gribble, G.W. (2003). The diversity of naturally produced organohalogens. Chemosphere 52, 289–297.
- Grostern, A., and Alvarez-Cohen, L. (2013). RubisCO-based CO₂ fixation and C₁ metabolism in the *Actinobacterium Pseudonocardia dioxanivorans* CB1190: Autotrophic growth by *Pseudonocardia dioxanivorans*. Environ. Microbiol. *15*, 3040-3053.
- Gruffaz, C., Muller, E.E.L., Louhichi-Jelail, Y., Nelli, Y.R., Guichard, G., and Bringel, F. (2014). Genes of the N-methylglutamate pathway are essential for growth of *Methylobacterium extorquens* DM4 with monomethylamine. Appl. Environ. Microbiol. *80*, 3541–3550.

- Guenther, A.B., Jiang, X., Heald, C.L., Sakulyanontvittaya, T., Duhl, T., Emmons, L.K., and Wang, X. (2012). The model of emissions of gases and aerosols from nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions. Geosci. Model Dev. 5, 1471–1492.
- Guo, X., and Lidstrom, M.E. (2008). Metabolite profiling analysis of *Methylobacterium extorquens* AM1 by comprehensive two-dimensional gas chromatography coupled with time-of-flight mass spectrometry. Biotechnol. Bioeng. *99*, 929–940.
- Halsey, K.H., Carter, A.E., and Giovannoni, S.J. (2012). Synergistic metabolism of a broad range of C₁ compounds in the marine methylotrophic bacterium HTCC2181: C₁ metabolism in the marine isolate HTCC2181. Environ. Microbiol. *14*, 630–640.
- Hamilton, J.T.G. (2003). Chloride methylation by plant pectin: an efficient environmentally significant process. Science *301*, 206–209.
- Han, J.-I., and Semrau, J.D. (2000). Chloromethane stimulates growth of *Methylomicrobium album* BG8 on methanol. FEMS Microbiol. Lett. *187*, 77–81.
- Hancock, T.L.C., Costello, A.M., Lidstrom, M.E., and Oremland, R.S. (1998). Strain IMB-1, a novel bacterium for the removal of methyl bromide in fumigated agricultural soils. Appl. Environ. Microbiol. *64*, 2899–2905.
- Harper, D., and Kennedy, J. (1986). Effect of growth conditions on halomethane production by *Phellinus* species: biological and environmental implications. Microbiology *132*, 1231–1246.
- Harper, D., Kennedy, J., and Hamilton, J.T. (1988). Chloromethane biosynthesis in poroid fungi. Phytochemistry 27, 3147–3153.
- Harper, D.B. (2000). The global chloromethane cycle: biosynthesis, biodegradation and metabolic role. Nat. Prod. Rep. *17*, 337–348.
- Harper, D.B., Hamilton, J.T.G., Ducrocq, V., Kennedy, J.T., Downey, A., and Kalin, R.M. (2003).
 The distinctive isotopic signature of plant-derived chloromethane: possible application in constraining the atmospheric chloromethane budget. Chemosphere 52, 433–436.
- Hartmans, S., Schmuckle, A., Cook, A.M., and Leisinger, T. (1986). Methyl chloride: naturally occurring toxicant and C-1 growth substrate. Microbiology *132*, 1139–1142.
- He, X., McLean, J.S., Edlund, A., Yooseph, S., Hall, A.P., Liu, S.Y., Dorrestein, P.C., Esquenazi, E., Hunter, R.C., Cheng, G., *et al.* (2015). Cultivation of a human-associated TM7 phylotype reveals a reduced genome and epibiotic parasitic lifestyle. Proc. Natl. Acad. Sci. *112*, 244–249.
- Herlemann, D.P., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J.J., and Andersson, A.F. (2011). Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. ISME J. 5, 1571–1579.
- Herrmann, A.M., Ritz, K., Nunan, N., Clode, P.L., Pett-Ridge, J., Kilburn, M.R., Murphy, D.V., O'Donnell, A.G., and Stockdale, E.A. (2007). Nano-scale secondary ion mass spectrometry

A new analytical tool in biogeochemistry and soil ecology: A review article. Soil Biol.
 Biochem. 39, 1835–1850.

- Hines, M.E., Crill, P.M., Varner, R.K., Talbot, R.W., Shorter, J.H., Kolb, C.E., and Harriss, R.C. (1998). Rapid consumption of low concentrations of methyl bromide by soil bacteria. Appl. Environ. Microbiol. *64*, 1864–1870.
- Hirsch, P.R., Mauchline, T.H., and Clark, I.M. (2010). Culture-independent molecular techniques for soil microbial ecology. Soil Biol. Biochem. 42, 878–887.
- Holmes, A.J., Costello, A., Lidstrom, M.E., and Murrell, J.C. (1995). Evidence that participate methane monooxygenase and ammonia monooxygenase may be evolutionarily related. FEMS Microbiol. Lett. *132*, 203–208.
- Hope, P.R., Bohmann, K., Gilbert, M.T.P., Zepeda-Mendoza, M.L., Razgour, O., and Jones, G. (2014). Second generation sequencing and morphological faecal analysis reveal unexpected foraging behaviour by *Myotis nattereri* (*Chiroptera, Vespertilionidae*) in winter. Front. Zool. 11, 1.
- Hoppe, B., Krger, K., Kahl, T., Arnstadt, T., Buscot, F., Bauhus, J., and Wubet, T. (2015). A pyrosequencing insight into sprawling bacterial diversity and community dynamics in decaying deadwood logs of *Fagus sylvatica* and *Picea abies*. Sci. Rep. *5*, 9456.
- Horiuchi, J., Badri, D.V., Kimball, B.A., Negre, F., Dudareva, N., Paschke, M.W., and Vivanco, J.M. (2007). The floral volatile, methyl benzoate, from snapdragon (*Antirrhinum majus*) triggers phytotoxic effects in *Arabidopsis thaliana*. Planta 226, 1–10.
- Hu, L., Yvon-Lewis, S.A., Liu, Y., Salisbury, J.E., and O'Hern, J.E. (2010). Coastal emissions of methyl bromide and methyl chloride along the eastern Gulf of Mexico and the east coast of the United States: Coastal emissions of CH₃Br and CH₃Cl. Glob. Biogeochem. Cycles *24*.
- Huang, Y.Y., Tanaka, M., Vecchio, D., Garcia-Diaz, M., Chang, J., Morimoto, Y., and Hamblin, M.R. (2012). Photodynamic therapy induces an immune response against a bacterial pathogen. Expert Rev. Clin. Immunol. *8*, 479–494.
- Huang, B., Lei, C., Wei, C., and Zeng, G. (2014). Chlorinated volatile organic compounds (Cl-VOCs) in environment — sources, potential human health impacts, and current remediation technologies. Environ. Int. *71*, 118–138.
- Hung, W.L., Wade, W.G., Boden, R., Kelly, D.P., and Wood, A.P. (2011). Facultative methylotrophs from the human oral cavity and methylotrophy in strains of *Gordonia*, *Leifsonia*, and *Microbacterium*. Arch. Microbiol. *193*, 407–417.
- Insam, H., and Seewald, M.S.A. (2010). Volatile organic compounds (VOCs) in soils. Biol. Fertil. Soils 46, 199–213.
- Jacob, D.J. (2005). Global budget of methanol: Constraints from atmospheric observations. J. Geophys. Res. 110.
- Jakočiūnas, T., Jensen, M.K., and Keasling, J.D. (2016). CRISPR/Cas9 advances engineering of microbial cell factories. Metab. Eng. *34*, 44–59.

- Janssen, D.B., Dinkla, I.J.T., Poelarends, G.J., and Terpstra, P. (2005). Bacterial degradation of xenobiotic compounds: evolution and distribution of novel enzyme activities. Environ. Microbiol. *7*, 1868–1882.
- Jehmlich, N., Schmidt, F., von Bergen, M., Richnow, H.-H., and Vogt, C. (2008). Protein-based stable isotope probing (Protein-SIP) reveals active species within anoxic mixed cultures. ISME J. 2, 1122–1133.
- Kai, M., Crespo, E., Cristescu, S.M., Harren, F.J.M., Francke, W., and Piechulla, B. (2010). Serratia odorifera: analysis of volatile emission and biological impact of volatile compounds on Arabidopsis thaliana. Appl. Microbiol. Biotechnol. 88, 965–976.
- Kalyuzhnaya, M.G., Lidstrom, M.E., and Chistoserdova, L. (2004). Utility of environmental primers targeting ancient enzymes: methylotroph detection in Lake Washington. Microb. Ecol. *48*, 463–472.
- Kalyuzhnaya, M.G., Zabinsky, R., Bowerman, S., Baker, D.R., Lidstrom, M.E., and Chistoserdova, L. (2006). Fluorescence in situ hybridization-flow cytometry-cell sortingbased method for separation and enrichment of type I and type II methanotroph populations. Appl. Environ. Microbiol. 72, 4293–4301.
- Kalyuzhnaya, M.G., Hristova, K.R., Lidstrom, M.E., and Chistoserdova, L. (2008). Characterization of a novel methanol dehydrogenase in representatives of *Burkholderiales*: implications for environmental detection of methylotrophy and evidence for convergent Evolution. J. Bacteriol. 190, 3817–3823.
- Käserer, H. (1906). Über die oxidation des wasserstoves und des methans durch mikroorganismen. Cent. Bakteriol Parasitenkd Infekt. Hyg Abt II *15*, 573–576.
- Kaviraj, A., Unlu, E., Gupta, A., and El Nemr, A. (2014). Biomarkers of environmental pollutants. BioMed Res. Int. 2014, 1–2.
- Kayser, M.F., and Vuilleumier, S. (2001). Dehalogenation of dichloromethane by dichloromethane dehalogenase/glutathione S-transferase leads to formation of DNA adducts. J. Bacteriol. *183*, 5209–5212.
- Kayser, M.F., Ucurum, Z., and Vuilleumier, S. (2002). Dichloromethane metabolism and C₁ utilization genes in *Methylobacterium* strains. Microbiology *148*, 1915–1922.
- Keener, W., and Arp, D. (1993). Kinetic studies of ammonia monooxygenase Inhibition in *Nitrosomonas europaea* by hydrocarbons and halogenated hydrocarbons in an optimized whole-cell assay. Appl Environ Microbiol 59, 2501–2510.
- Keiblinger, K.M., Wilhartitz, I.C., Schneider, T., Roschitzki, B., Schmid, E., Eberl, L., Riedel, K., and Zechmeister-Boltenstern, S. (2012). Soil metaproteomics – Comparative evaluation of protein extraction protocols. Soil Biol. Biochem. 54, 14–24.
- Keppler, F., Borchers, R., Elsner, P., Fahimi, I., Pracht, J., and Schöler, H.F. (2003). Formation of volatile iodinated alkanes in soil: results from laboratory studies. Chemosphere 52, 477–483.

- Keppler, F., Harper, D.B., Röckmann, T., Moore, R.M., and Hamilton, J.T.G. (2005). New insight into the atmospheric chloromethane budget gained using stable carbon isotope ratios. Atmospheric Chem. Phys. *5*, 2403–2411.
- Kiefer, P., Portais, J.C., and Vorholt, J.A. (2008). Quantitative metabolome analysis using liquid chromatography–high-resolution mass spectrometry. Anal. Biochem. *382*, 94–100.
- Kiefer, P., Delmotte, N., and Vorholt, J.A. (2011). Nanoscale ion-pair reversed-phase HPLC–MS for sensitive metabolome analysis. Anal. Chem. *83*, 850–855.
- Kielak, A.M., Scheublin, T.R., Mendes, L.W., van Veen, J.A., and Kuramae, E.E. (2016). Bacterial community succession in pine-wood decomposition. Front. Microbiol. 7.
- Knief, C., Frances, L., and Vorholt, J.A. (2010). Competitiveness of diverse *Methylobacterium* strains in the phyllosphere of *Arabidopsis thaliana* and identification of representative models, including *M. extorquens* PA1. Microb. Ecol. 60, 440–452.
- Knief, C., Delmotte, N., Chaffron, S., Stark, M., Innerebner, G., Wassmann, R., von Mering, C., and Vorholt, J.A. (2012). Metaproteogenomic analysis of microbial communities in the phyllosphere and rhizosphere of rice. ISME J. 6, 1378–1390.
- Kögel-Knabner, I. (2002) The macromolecular organic composition of plant and microbial residues as inputs to soil organic matter. Soil Biology and Biochemistry, *34*, 139-162.
- Kohler-Staub, D., and Leisinger, T. (1985). Dichloromethane dehalogenase of *Hyphomicrobium* sp. strain DM2. J. Bacteriol. *162*, 676–681.
- Kolb, S. (2009). Aerobic methanol-oxidizing Bacteria in soil. FEMS Microbiol. Lett. 300, 1–10.
- Kolb, S., and Stacheter, A. (2013). Prerequisites for amplicon pyrosequencing of microbial methanol utilizers in the environment. Front. Microbiol. *4*, 268.
- Korotkova, N., and Lidstrom, M.E. (2001). Connection between poly-hydroxybutyrate biosynthesis and hrowth on C₁ and C₂ compounds in the methylotroph *Methylobacterium extorquens* AM1. J. Bacteriol. *183*, 1038–1046.
- Koskimäki, J.J., Pirttilä, A.M., Ihantola, E.-L., Halonen, O., and Frank, A.C. (2015). The Intracellular scots pine shoot symbiont *Methylobacterium extorquens* DSM13060 aggregates around the host nucleus and encodes eukaryote-like proteins. mBio *6*, e00039-15.
- Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., and Schloss, P.D. (2013). Development of a dual-Index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. Appl. Environ. Microbiol. 79, 5112–5120.
- Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. Mol. Biol. Evol. msw054.
- Kurylo, M., and Rodriguez, J. (1998). Short-lived ozone-related compounds. (Scientific assessment of ozone depletion). Chapter 2

- La Roche, S.D., and Leisinger, T. (1990). Sequence analysis and expression of the bacterial dichloromethane dehalogenase structural gene, a member of the glutathione S-transferase supergene family. J. Bacteriol. *172*, 164–171.
- Laukel, M., Rossignol, M., Borderies, G., Völker, U., and Vorholt, J.A. (2004). Comparison of the proteome of *Methylobacterium extorquens* AM1 grown under methylotrophic and nonmethylotrophic conditions. Proteomics *4*, 1247–1264.
- Lee, S.W., Won, K., Lim, H.K., Kim, J.C., Choi, G.J., and Cho, K.Y. (2004). Screening for novel lipolytic enzymes from uncultured soil microorganisms. Appl. Microbiol. Biotechnol. *65*, 720–726.
- Leng, L., Chang, J., Geng, K., Lu, Y., and Ma, K. (2015). Uncultivated Methylocystis Species in Paddy Soil Include Facultative Methanotrophs that Utilize Acetate. Microb. Ecol. 70, 88–96.
- Lidstrom, M. (2006). Aerobic methylotrophic procaryotes. In The Prokaryotes, pp. 618–634.
- Lin, C., Owen, S., and Penuelas, J. (2007). Volatile organic compounds in the roots and rhizosphere of *Pinus* spp. Soil Biol. Biochem. *39*, 951–960.
- Liu, X.S. (2007). Getting started in tiling microarray analysis. PLoS Comput. Biol. 3.
- Llorens-Rico, V., Cano, J., Kamminga, T., Gil, R., Latorre, A., Chen, W.H., Bork, P., Glass, J.I., Serrano, L., and Lluch-Senar, M. (2016). Bacterial antisense RNAs are mainly the product of transcriptional noise. Sci. Adv. *2*, e1501363–e1501363.
- Llusia, J., Peñuelas, J., Guenther, A., and Rapparini, F. (2013). Seasonal variations in terpene emission factors of dominant species in four ecosystems in NE Spain. Atmos. Environ. 70, 149–158.
- Loew, O. (1892). Ueber einen *Bacillus*, welcher ameisensäure und formaldehyd assimiliren kann. Cent. Für Bakteriol. Parasitenkd. Infekt. Hyg. Abt. II *12*, 462–465.
- Lopez-Marques, R.L., Perez-Castineira, J.R., Losada, M., and Serrano, A. (2004). Differential regulation of soluble and membrane-bound inorganic pyrophosphatases in the photosynthetic bacterium *Rhodospirillum rubrum* provides insights into pyrophosphate-based stress bioenergetics. J. Bacteriol. *186*, 5418–5426.
- Lueders, T., Manefield, M., and Friedrich, M.W. (2003). Enhanced sensitivity of DNA- and rRNA-based stable isotope probing by fractionation and quantitative analysis of isopycnic centrifugation gradients: Quantitative analysis of SIP gradients. Environ. Microbiol. *6*, 73–78.
- Mägli, A., Messmer, M., and Leisinger, T. (1998). Metabolism of dichloromethane by the strict anaerobe *Dehalobacterium formicoaceticum*. Appl. Environ. Microbiol. *64*, 646–650.
- Mancuso, S., Taiti, C., Bazihizina, N., Costa, C., Menesatti, P., Giagnoni, L., Arenella, M., Nannipieri, P., and Renella, G. (2015). Soil volatile analysis by proton transfer reactiontime of flight mass spectrometry (PTR-TOF-MS). Appl. Soil Ecol. 86, 182–191.

- Marison, I.W., and Attwood, M.M. (1982). A possible alternative mechanism for the oxidation of formaldehyde to formate. Microbiology *128*, 1441–1446.
- Marx, C.J., O'Brien, B.N., Breezee, J., and Lidstrom, M.E. (2003). Novel methylotrophy genes of *Methylobacterium extorquens* AM1 Identified by using transposon mutagenesis including a putative dihydromethanopterin reductase. J. Bacteriol. *185*, 669–673.
- Marx, C.J., Dien, S.J.V., and Lidstrom, M.E. (2005). Flux analysis uncovers key role of functional redundancy in formaldehyde metabolism. PLoS Biol *3*, e16.
- Marx, C.J. (2008). Development of a broad-host-range *sacB*-based vector for unmarked allelic exchange. BMC Res. Notes 1, 1.
- Marx, C.J., Bringel, F., Chistoserdova, L., Moulin, L., Farhan Ul Haque, M., Fleischman, D.E., Gruffaz, C., Jourand, P., Knief, C., Lee, M.C., *et al.* (2012). Complete genome sequences of six strains of the genus *Methylobacterium*. J. Bacteriol. *194*, 4746–4748.
- McAnulla, C., Woodall, C.A., McDonald, I.R., Studer, A., Vuilleumier, S., Leisinger, T., and Murrell, J.C. (2001). Chloromethane utilization gene cluster from *Hyphomicrobium chloromethanicum* strain CM2T and development of functional gene Probes to detect halomethane-degrading bacteria. Appl. Environ. Microbiol. *67*, 307–316.
- McCulloch, A., and Aucott, M.L. (1999). Global emissions of hydrogen chloride and chloromethane from coal combustion, incineration and industrial activities: reactive chlorine emissions inventory. J. Geophys. Res-Atmos. *104*, 8391-8403.
- McCulloch, A., Midgley, P.M., and Ashford, P. (2003). Releases of refrigerant gases (CFC-12, HCFC-22 and HFC-134a) to the atmosphere. Atmos. Environ. *37*, 889–902.
- McDonald, I.R., and Murrell, J.C. (1997). The methanol dehydrogenase structural gene *mxaF* and its use as a functional gene probe for methanotrophs and methylotrophs. Appl. Environ. Microbiol. *63*, 3218–3224.
- McDonald, I.R., Doronina, N.V., Trotsenko, Y.A., McAnulla, C., and Murrell, J.C. (2001). *Hyphomicrobium chloromethanicum* sp. nov. and *Methylobacterium chloromethanicum* sp. nov., chloromethane-utilizing bacteria isolated from a polluted environment. Int. J. Syst. Evol. Microbiol. *51*, 119–122.
- McDonald, I.R., Radajewski, S., and Murrell, J.C. (2005). Stable isotope probing of nucleic acids in methanotrophs and methylotrophs: A review. Org. Geochem. *36*, 779–787.
- McDonald, I.R., Bodrossy, L., Chen, Y., and Murrell, J.C. (2008). Molecular Ecology Techniques for the Study of Aerobic Methanotrophs. Appl. Environ. Microbiol. 74, 1305–1315.
- Michener, J.K., Vuilleumier, S., Bringel, F., and Marx, C.J. (2014a). Phylogeny poorly predicts the utility of a challenging horizontally transferred gene in *Methylobacterium* strains. J. Bacteriol. *196*, 2101–2107.
- Michener, J.K., Neves, A.C., Vuilleumier, S., Bringel, F., and Marx, C.J. (2014b). Effective use of a horizontally-transferred pathway for dichloromethane catabolism requires post–transfer refinement. Elife *3*, e04279.

- Miller, L.G., Warner, K.L., Baesman, S.M., Oremland, R.S., McDonald, I.R., Radajewski, S., and Murrell, J.C. (2004). Degradation of methyl bromide and methyl chloride in soil microcosms: Use of stable C isotope fractionation and stable isotope probing to identify reactions and the responsible microorganisms. Geochim. Cosmochim. Acta 68, 3271– 3283.
- Mirete, S., Mora-Ruiz, M.R., Lamprecht-Grandío, M., de Figueras, C.G., Rosselló-Móra, R., and González-Pastor, J.E. (2015). Salt resistance genes revealed by functional metagenomics from brines and moderate-salinity rhizosphere within a hypersaline environment. Front. Microbiol. *6*.
- Montzka, S.A., and Fraser, P. (2003). Relative contributions of greenhouse gas emissions to global warming.
- Montzka, S.A., Dlugokencky, E.J., and Butler, J.H. (2011). Non-CO₂ greenhouse gases and climate change. Nature 476, 43–50.
- Moore, R.M., Gut, A., and Andreae, M.O. (2005). A pilot study of methyl chloride emissions from tropical woodrot fungi. Chemosphere *58*, 221–225.
- Muller, E.E.L., Hourcade, E., Louhichi-Jelail, Y., Hammann, P., Vuilleumier, S., and Bringel, F. (2011a). Functional genomics of dichloromethane utilization in *Methylobacterium extorquens* DM4. Environ. Microbiol. *13*, 2518–2535.
- Muller, E.E.L., Bringel, F., and Vuilleumier, S. (2011b). Dichloromethane-degrading bacteria in the genomic age. Res. Microbiol. *162*, 869–876.
- Nacke, H., Fischer, C., Thürmer, A., Meinicke, P., and Daniel, R. (2014). Land use type significantly affects microbial gene transcription in soil. Microb. Ecol. *67*, 919–930.
- Nadalig, T., Farhan Ul Haque, M., Roselli, S., Schaller, H., Bringel, F., and Vuilleumier, S. (2011). Detection and isolation of chloromethane-degrading bacteria from the *Arabidopsis thaliana* phyllosphere, and characterization of chloromethane utilization genes: chloromethane-degrading bacteria from *Arabidopsis thaliana*. FEMS Microbiol. Ecol. 77, 438–448.
- Nadalig, T., Greule, M., Bringel, F., Vuilleumier, S., and Keppler, F. (2013). Hydrogen and carbon isotope fractionation during degradation of chloromethane by methylotrophic bacteria. MicrobiologyOpen *2*, 893–900.
- Nadalig, T., Greule, M., Bringel, F., Keppler, F., and Vuilleumier, S. (2014). Probing the diversity of chloromethane-degrading bacteria by comparative genomics and isotopic fractionation. Front. Microbiol. *5*. 523
- Nagatoshi, Y., and Nakamura, T. (2007). Characterization of three halide methyltransferases in *Arabidopsis thaliana*. Plant Biotechnol. *24*, 503–506.
- Nayak, D.D., and Marx, C.J. (2014). Genetic and phenotypic comparison of facultative methylotrophy between *Methylobacterium extorquens* strains PA1 and AM1. PLoS ONE *9*, e107887.

- Nercessian, O., Noyes, E., Kalyuzhnaya, M.G., Lidstrom, M.E., and Chistoserdova, L. (2005). Bacterial populations active in metabolism of C₁ compounds in the sediment of Lake Washington, a freshwater lake. Appl. Environ. Microbiol. *71*, 6885–6899.
- Neufeld, J.D., Vohra, J., Dumont, M.G., Lueders, T., Manefield, M., Friedrich, M.W., and Murrell, J.C. (2007). DNA stable-isotope probing. Nat. Protoc. *2*, 860–866.
- Neufeld, J.D., Chen, Y., Dumont, M.G., and Murrell, J.C. (2008). Marine methylotrophs revealed by stable-isotope probing, multiple displacement amplification and metagenomics. Environ. Microbiol. *10*, 1526–1535.
- Öberg, G.M. (2003). The biogeochemistry of chlorine in soil. In natural production of organohalogen compounds, (Springer), pp. 43–62.
- Ochsner, A.M., Sonntag, F., Buchhaupt, M., Schrader, J., and Vorholt, J.A. (2015). *Methylobacterium extorquens*: methylotrophy and biotechnological applications. Appl. Microbiol. Biotechnol. *99*, 517–534.
- Oderbolz, D.C., Aksoyoglu, S., Keller, J., Barmpadimos, I., Steinbrecher, R., Skjøth, C.A., Plaß-Dülmer, C., and Prévôt, A.S.H. (2013). A comprehensive emission inventory of biogenic volatile organic compounds in Europe: improved seasonality and land-cover. Atmospheric Chem. Phys. *13*, 1689–1712.
- O'Doherty, S., Simmonds, P., Cunnold, D., and Wang, H. (2001). *In situ* chloroform measurements at advanced global atmospheric gases experiment atmospheric research stations from 1994 to 1998. J. Geophys. Res-Atmos. *106*, 20429–20444.
- Okubo, Y., Skovran, E., Guo, X., Sivam, D., and Lidstrom, M.E. (2007). Implementation of microarrays for *Methylobacterium extorquens* AM1. OMICS J. Integr. Biol. *11*, 325–340.
- Patt, T.E., Cole, G.C., and Hanson, R.S. (1976). *Methylobacterium*, a new genus of facultatively methylotrophic bacteria. Int. J. Syst. Evol. Microbiol. *26*, 226–229.
- Peano, C., Pietrelli, A., Consolandi, C., Rossi, E., Petiti, L., Tagliabue, L., De Bellis, G., and Landini, P. (2013). An efficient rRNA removal method for RNA sequencing in GC-rich bacteria. Microb. Inform. Exp. *3*, 1.
- Peñuelas, J., Asensio, D., Tholl, D., Wenke, K., Rosenkranz, M., Piechulla, B., and Schnitzler, J.P. (2014). Biogenic volatile emissions from the soil: Biogenic volatile emissions from the soil. Plant Cell Environ. 37, 1866–1891.
- Peyraud, R., Kiefer, P., Christen, P., Massou, S., Portais, J.C., and Vorholt, J.A. (2009). Demonstration of the ethylmalonyl-CoA pathway by using ¹³C metabolomics. Proc. Natl. Acad. Sci. 106, 4846–4851.
- Peyraud, R., Schneider, K., Kiefer, P., Massou, S., Vorholt, J.A., and Portais, J.C. (2011). Genome-scale reconstruction and system level investigation of the metabolic network of *Methylobacterium extorquens* AM1. BMC Syst. Biol. 5, 1.
- Peyraud, R., Kiefer, P., Christen, P., Portais, J.C., and Vorholt, J.A. (2012). Co-consumption of methanol and succinate by *Methylobacterium extorquens* AM1. PLoS ONE *7*, e48271.

- Pham, V.H.T., and Kim, J. (2012). Cultivation of unculturable soil bacteria. Trends Biotechnol. 30, 475–484.
- Polerecky, L., Adam, B., Milucka, J., Musat, N., Vagner, T., and Kuypers, M.M.M. (2012). Look@NanoSIMS - a tool for the analysis of nanoSIMS data in environmental microbiology. Environ. Microbiol. 14, 1009–1023.
- Prinn, R., Weiss, R., and Fraser, P. (2000). A history of chemically and radiatively important gases in air from ALE/GAGE/AGAGE. J. Geophys. Res. *105*, 17751–17792.
- Prosser, J.I. (2015). Dispersing misconceptions and identifying opportunities for the use of 'omics' in soil microbial ecology. Nat. Rev. Microbiol. 13, 439–446.
- Radajewski, S., Webster, G., Reay, D.S., Morris, S.A., Ineson, P., Nedwell, D.B., Prosser, J.I., and Murrell, J.C. (2002). Identification of active methylotroph populations in an acidic forest soil by stable-isotope probingc. Microbiology *148*, 2331–2342.
- Rasche, M.E., Hyman, M.R., and Arp, D.J. (1991). Factors limiting aliphatic chlorocarbon degradation by *Nitrosomonas europaea*: cometabolic inactivation of ammonia monooxygenase and substrate specificity. Appl. Environ. Microbiol. *57*, 2986–2994.
- Rappé, M.S., and Giovannoni, S.J. (2003). The uncultured microbial majority. Annu. Rev. Microbiol. *57*, 369–394.
- Reaser, B.C., Yang, S., Fitz, B.D., Parsons, B.A., Lidstrom, M.E., and Synovec, R.E. (2016). Nontargeted determination of ¹³C-labeling in the *Methylobacterium extorquens* AM1 metabolome using the two-dimensional mass cluster method and principal component analysis. J. Chromatogr. A 1432, 111–121.
- Redeker, K.R., Wang, N.Y., Low, J.C., McMillan, A., Tyler, S.C., and Cicerone, R.J. (2000). Emissions of methyl halides and methane from rice paddies. Science *290*, 966–969.
- Redeker, K.R., Treseder, K.K., and Allen, M.F. (2004). Ectomycorrhizal fungi: a new source of atmospheric methyl halides? Glob. Change Biol. *10*, 1009–1016.
- Redeker, K.R., and Kalin, R.M. (2012). Methyl chloride isotopic signatures from Irish forest soils and a comparison between abiotic and biogenic methyl halide soil fluxes. Glob. Change Biol. 18, 1453–1467.
- Rettedal, E.A., and Brözel, V.S. (2015). Characterizing the diversity of active bacteria in soil by comprehensive stable isotope probing of DNA and RNA with H_2 ¹⁸O. MicrobiologyOpen *4*, 208–219.
- Rhew, R.C., Østergaard, L., Saltzman, E.S., and Yanofsky, M.F. (2003). Genetic control of methyl halide production in *Arabidopsis*. Curr. Biol. *13*, 1809–1813.
- Rhew, R.C., and Abel, T. (2007). Measuring simultaneous production and consumption fluxes of methyl chloride and methyl bromide in annual temperate grasslands. Environ. Sci. Technol. *41*, 7837–7843.

- Rhew, R.C., Chen, C., Teh, Y.A., and Baldocchi, D. (2010). Gross fluxes of methyl chloride and methyl bromide in a California oak-savanna woodland. Atmos. Environ. 44, 2054–2061.
- Rigamonte, T.A., Pylro, V.S., and Duarte, G.F. (2010). The role of mycorrhization helper bacteria in the establishment and action of ectomycorrhizae associations. Braz. J. Microbiol. *41*, 832–840.
- Rocca, J.D., Hall, E.K., Lennon, J.T., Evans, S.E., Waldrop, M.P., Cotner, J.B., Nemergut, D.R., Graham, E.B., and Wallenstein, M.D. (2015). Relationships between protein-encoding gene abundance and corresponding process are commonly assumed yet rarely observed. ISME J. 9, 1693–1699.
- Roche, S.D.L., and Leisinger, T. (1991). Identification of *dcmR*, the regulatory gene governing expression of dichloromethane dehalogenase in *Methylobacterium* sp. strain DM4. J. Bacteriol. *173*, 6714–6721.
- Roselli, S. (2009). Génomique fonctionnelle de la dégradation microbienne du chlorométhane. Thèse de doctorat, Université de Strasbourg.
- Roselli, S., Nadalig, T., Vuilleumier, S., and Bringel, F. (2013). The 380 kb pCMU01 plasmid encodes chloromethane utilization genes and redundant genes for vitamin B12- and tetrahydrofolate-dependent chloromethane metabolism in *Methylobacterium extorquens* CM4: A proteomic and bioinformatics study. PLoS ONE *8*, e56598.
- Ruecker, A., Weigold, P., Behrens, S., Jochmann, M., Laaks, J., and Kappler, A. (2014). Predominance of biotic over abiotic formation of halogenated hydrocarbons in hypersaline sediments in Western Australia. Environ. Sci. Technol. *48*, 9170–9178.
- Rüffer, M. (2013). Spatial distribution of *cmuA*-genotypes and CH₃Cl oxidation in a mixed hardwood forest soil (Steigerwald). Research proposal.
- Sailaukhanuly, Y., Sárossy, Z., Carlsen, L., and Egsgaard, H. (2014). Mechanistic aspects of the nucleophilic substitution of pectin. On the formation of chloromethane. Chemosphere *111*, 575–579.
- Saito, T., and Yokouchi, Y. (2008). Stable carbon isotope ratio of methyl chloride emitted from glasshouse-grown tropical plants and its implication for the global methyl chloride budget. Geophys. Res. Lett. *35*.
- Saito, T., Yokouchi, Y., Kosugi, Y., Tani, M., Philip, E., and Okuda, T. (2008). Methyl chloride and isoprene emissions from tropical rain forest in Southeast Asia. Geophys. Res. Lett. *35*.
- Schade, G., and Goldstein, A. (2001). Fluxes of oxygenated volatile organic compounds from a ponderosa pine plantation. J. Geophys. Res. *106*, 3111–3123.
- Schaefer, J.K. (2002). *Leisingera methylohalidivorans* gen. nov., sp. nov., a marine methylotroph that grows on methyl bromide. Int. J. Syst. Evol. Microbiol. *52*, 851–859.
- Schäfer, H., McDonald, I.R., Nightingale, P.D., and Murrell, J.C. (2005). Evidence for the presence of a CmuA methyltransferase pathway in novel marine methyl halide-oxidizing bacteria: novel marine methyl halide-oxidizing bacteria. Environ. Microbiol. *7*, 839–852.

- Schäfer, H., Miller, L.G., Oremland, R.S., and Murrell, J.C. (2007). Bacterial cycling of methyl halides. Adv Appl Microbiol. *61*, 307-346.
- Schauffler, S.M. (2003). Chlorine budget and partitioning during the stratospheric aerosol and gas experiment (SAGE) III ozone loss and validation experiment (SOLVE). J. Geophys. Res. *108*.
- Schink, B., and Zeikus, J. (1980). Microbial methanol formation: a major end product of pectin metabolism. Curr. Microbiol. 6, 387–389.
- Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski, R.A., Oakley, B.B., Parks, D.H., Robinson, C.J., *et al.* (2009). Introducing mothur: Opensource, platform-independent, community-supported Software for describing and comparing microbial communities. Appl. Environ. Microbiol. 75, 7537–7541.
- Schmidt, S., Christen, P., Kiefer, P., and Vorholt, J.A. (2010). Functional investigation of methanol dehydrogenase-like protein XoxF in *Methylobacterium extorquens* AM1. Microbiology 156, 2575–2586.
- Schmidt, P.A., Bálint, M., Greshake, B., Bandow, C., Römbke, J., and Schmitt, I. (2013). Illumina metabarcoding of a soil fungal community. Soil Biol. Biochem. *65*, 128–132.
- Sharma, C.M., Hoffmann, S., Darfeuille, F., Reignier, J., Findeiß, S., Sittka, A., Chabas, S., Reiche, K., Hackermüller, J., Reinhardt, R., *et al.* (2010). The primary transcriptome of the major human pathogen *Helicobacter pylori*. Nature *464*, 250–255.
- Sharma, C.M., and Vogel, J. (2014). Differential RNA-seq: the approach behind and the biological insight gained. Curr. Opin. Microbiol. 19, 97–105.
- Shorter, J., Kolb, C., and Crill, P. (1995). Rapid degradation of atmospheric methyl bromide in soil. Nature *377*, 717–719.
- Shukla, A.K., Vishwakarma, P., Upadhyay, S.N., Tripathi, A.K., Prasana, H.C., and Dubey, S.K. (2009). Biodegradation of trichloroethylene (TCE) by methanotrophic community. Bioresour. Technol. 100, 2469–2474.
- Skovran, E., Palmer, A.D., Rountree, A.M., Good, N.M., and Lidstrom, M.E. (2011b). XoxF is required for expression of methanol dehydrogenase in *Methylobacterium extorquens* AM1. J. Bacteriol. *193*, 6032–6038.
- Šmejkalová, H., Erb, T.J., and Fuchs, G. (2010). Methanol assimilation in *Methylobacterium extorquens* AM1: demonstration of all enzymes and their regulation. PLoS ONE *5*, e13001.
- Söhngen, N.L. (1906). Uber bakterien welche methan ab kohlenstoffnahrung und energiequelle gerbrauchen. Zbl Bact Parasitenk *15*, 513–517.
- Stacheter, A., Noll, M., Lee, C.K., Selzer, M., Glowik, B., Ebertsch, L., Mertel, R., Schulz, D., Lampert, N., Drake, H.L and Kolb, S. (2013). Methanol oxidation by temperate soils and environmental determinants of associated methylotrophs. ISME J. 7, 1051–1064.

- Studer, A., Vuilleumier, S., and Leisinger, T. (1999). Properties of the methylcobalamin: H4folate methyltransferase involved in chloromethane utilization by *Methylobacterium* sp. strain CM4. Eur. J. Biochem. *264*, 242–249.
- Studer, A., Stupperich, E., Vuilleumier, S., and Leisinger, T. (2001). Chloromethane: tetrahydrofolate methyl transfer by two proteins from *Methylobacterium chloromethanicum* strain CM4. Eur. J. Biochem. FEBS *268*, 2931–2938.
- Studer, A., McAnulla, C., Buchele, R., Leisinger, T., and Vuilleumier, S. (2002). Chloromethaneinduced genes define a third C₁ utilization pathway in *Methylobacterium chloromethanicum* CM4. J. Bacteriol. *184*, 3476–3484.
- Talia, P., Sede, S.M., Campos, E., Rorig, M., Principi, D., Tosto, D., Hopp, H.E., Grasso, D., and Cataldi, A. (2012). Biodiversity characterization of cellulolytic bacteria present on native Chaco soil by comparison of ribosomal RNA genes. Res. Microbiol. *163*, 221–232.
- Tang, J.K.-H., You, L., Blankenship, R.E., and Tang, Y.J. (2012). Recent advances in mapping environmental microbial metabolisms through 13C isotopic fingerprints. J. R. Soc. Interface 9, 2767–2780.
- Tamura, K., and Nei, M. (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. Mol. Biol. Evol. 10, 512–526.
- Taubert, M., Grob, C., Howat, A.M., Burns, O.J., Dixon, J.L., Chen, Y., and Murrell, J.C. (2015).
 XoxF encoding an alternative methanol dehydrogenase is widespread in coastal marine environments: *xoxF* in costal marine environments. Environ. Microbiol. *17*, 3937–3948.
- Temkiv, T.Š., Finster, K., Hansen, B.M., Nielsen, N.W., and Karlson, U.G. (2012). The microbial diversity of a storm cloud as assessed by hailstones. FEMS Microbiol. Ecol. *81*, 684–695.
- Traunecker, J., Preuss, A., and Diekert, G. (1991a). Isolation and characterization of a methyl chloride utilizing, strictly anaerobic bacterium. Arch. Microbiol. *156*, 416–421.
- United Nations Environment Programm (UNEP) (2005). Production and consumption of ozone depleting substances under the Montreal protocol 1986-2004.
- Van Aken, B. (2004). *Methylobacterium populi* sp. nov., a novel aerobic, pink-pigmented, facultatively methylotrophic, methane-utilizing bacterium isolated from poplar trees (Populus deltoidesxnigra DN34). Int. J. Syst. Evol. Microbiol. *54*, 1191–1196.
- Van Assche, E., Van Puyvelde, S., Vanderleyden, J., and Steenackers, H.P. (2015). RNA-binding proteins involved in post-transcriptional regulation in bacteria. Front. Microbiol. *6*.
- Van der Klei, I.J., Yurimoto, H., Sakai, Y., and Veenhuis, M. (2006). The significance of peroxisomes in methanol metabolism in methylotrophic yeast. Biochim. Biophys. Acta BBA Mol. Cell Res. *1763*, 1453–1462.
- VanDien, S.J., Marx, C.J., O'Brien, B.N., and Lidstrom, M.E. (2003). Genetic characterization of the carotenoid biosynthetic pathway in *Methylobacterium extorquens* AM1 and isolation of a colorless mutant. Appl. Environ. Microbiol. *69*, 7563–7566.

- Van Dijk, E.L., Jaszczyszyn, Y., and Thermes, C. (2014). Library preparation methods for nextgeneration sequencing: Tone down the bias. Exp. Cell Res. *322*, 12–20.
- Vannelli, T., Studer, A., Kertesz, M., and Leisinger, T. (1998). Chloromethane metabolism by *Methylobacterium* sp. Strain CM4. Appl. Environ. Microbiol. *64*, 1933–1936.
- Vannelli, T., Messmer, M., Studer, A., Vuilleumier, S., and Leisinger, T. (1999). A corrinoiddependent catabolic pathway for growth of a *Methylobacterium* strain with chloromethane. Proc. Natl. Acad. Sci. *96*, 4615–4620.
- Vorholt, J.A., Chistoserdova, L., Stolyar, S.M., Thauer, R.K., and Lidstrom, M.E. (1999). Distribution of tetrahydromethanopterin-dependent enzymes in methylotrophic bacteria and phylogeny of methenyl tetrahydromethanopterin cyclohydrolases. J. Bacteriol. 181, 5750–5757.
- Vorholt, J.A. (2002). Cofactor-dependent pathways of formaldehyde oxidation in methylotrophic bacteria. Arch. Microbiol. *178*, 239–249.
- Vorholt, J.A. (2012). Microbial life in the phyllosphere. Nat. Rev. Microbiol. 10, 828–840.
- Vos, M., Wolf, A.B., Jennings, S.J., and Kowalchuk, G.A. (2013). Micro-scale determinants of bacterial diversity in soil. FEMS Microbiol. Rev. *37*, 936–954.
- Vu, H.N., Subuyuj, G.A., Vijayakumar, S., Good, N.M., Martinez-Gomez, N.C., and Skovran, E. (2016). lanthanide-dependent regulation of methanol oxidation systems in *Methylobacterium extorquens* AM1 and their contribution to methanol growth. J. Bacteriol. *198*, 1250–1259.
- Vuilleumier, S., Ivoš, N., Dean, M., and Leisinger, T. (2001). Sequence variation in dichloromethane dehalogenases/glutathione S-transferases. Microbiology 147, 611–619.
- Vuilleumier, S. (2002). Coping with a halogenated one-carbon diet: aerobic dichloromethanemineralising bacteria. In Biotechnology for the Environment, Focus on Biotechnology Serie, pp. 105–131.
- Vuilleumier, S., Chistoserdova, L., Lee, M.-C., Bringel, F., Lajus, A., Zhou, Y., Gourion, B., Barbe, V., Chang, J., Cruveiller, S., et al. (2009). Methylobacterium genome sequences: a reference blueprint to investigate microbial metabolism of C₁ compounds from natural and industrial sources. PLoS ONE 4, e5584.
- Vuilleumier, S., Nadalig, T., Farhan Ul Haque, M., Magdelenat, G., Lajus, A., Roselli, S., Muller, E.E.L., Gruffaz, C., Barbe, V., Medigue, C., *et al.* (2011). Complete genome sequence of the chloromethane-degrading *Hyphomicrobium* sp. strain MC1. J. Bacteriol. *193*, 5035– 5036.
- Wagner, M., Nielsen, P.H., Loy, A., Nielsen, J.L., and Daims, H. (2006). Linking microbial community structure with function: fluorescence in situ hybridization-microautoradiography and isotope arrays. Curr. Opin. Biotechnol. *17*, 83–91.
- Wang, Z., Gerstein, M., and Snyder, M. (2009). RNA-Seq: a revolutionary tool for transcriptomics. Nat. Rev. Genet. 10, 57–63.

- Warneke, C., Karl, T., and Judmaier, H. (1999). Acetone, methanol, and other partially oxidized volatile organic emissions from dead plant matter by abiologicalp rocesses: Significance for atmospheric HOx chemistry. Global Biogeochemical Cycles *13*, 9–17.
- Warwick, S., Bowen, T., McVeigh, H., and Embley, T.M. (1994). A phylogenetic analysis of the family *Pseudonocardiaceae* and the genera *Actinokineospora* and *Saccharothrix* with 16S rRNA sequences and a proposal to combine the genera *Amycolata* and *Pseudonocardia* in an emended genus *Pseudonocardia*. Int. J. Syst. Evol. Microbiol. 44, 293–299.
- Watling, R., and Harper, D.B. (1998). Chloromethane production by wood-rotting fungi and an estimate of the global flux to the atmosphere. Mycol. Res. *102*, 769–787.
- White, R.H. (1982). Biosynthesis of methyl chloride in the fungus *Phellinus pomaceus*. Arch. Microbiol. *132*, 100–102.
- Whittenbury, R., Phillips, K.C., and Wilkinson, J.F. (1970). Enrichment, isolation and some properties of methane-utilizing bacteria. Microbiology *61*, 205–218.
- Wohlfahrt, G., Amelynck, C., Ammann, C., Arneth, A., Bamberger, I., Goldstein, A.H., Gu, L., Guenther, A., Hansel, A., Heinesch, B., *et al.* (2015). An ecosystem-scale perspective of the net land methanol flux: synthesis of micrometeorological flux measurements. Atmospheric Chem. Phys. *15*, 7413–7427.
- Wishkerman, A., Gebhardt, S., McRoberts, C.W., Hamilton, J.T.G., Williams, J., and Keppler, F. (2008). Abiotic methyl bromide formation from vegetation, and its strong dependence on temperature. Environ. Sci. Technol. 42, 6837–6842.
- Wuosmaa, A., and Hager, L. (1990). Methyl chloride transferase: a carbocation route for biosynthesis of halometabolites. Science 249, 160–162.
- Wyborn, N.R., Mills, J., Williams, S.G., and Jones, C.W. (1996). Molecular characterisation of formamidase from Methylophilus methylotrophus. Eur. J. Biochem. 240, 314–322.
- Xiao, X. (2008). Optimal estimation of the surface fluxes of chloromethanes using a 3-D global atmospheric chemical transport model. Massachusetts Institute of Technology.
- Xie, S., Sun, W., Luo, C., and Cupples, A.M. (2011). Novel aerobic benzene degrading microorganisms identified in three soils by stable isotope probing. Biodegradation 22, 71–81.
- Yang, S., Hoggard, J.C., Lidstrom, M.E., and Synovec, R.E. (2013). Comprehensive discovery of 13C labeled metabolites in the bacterium *Methylobacterium extorquens* AM1 using gas chromatography–mass spectrometry. J. Chromatogr. A *1317*, 175–185.
- Yoshida, Y., Wang, Y., Zeng, T., and Yantosca, R. (2004). A three-dimensional global model study of atmospheric methyl chloride budget and distributions. J. Geophys. Res. *109*.
- Yoshida, Y., Wang, Y., Shim, C., Cunnold, D., Blake, D.R., and Dutton, G.S. (2006). Inverse modeling of the global methyl chloride sources. J. Geophys. Res. 111.

- Yokouchi, Y., Saito, T., Ishigaki, C., and Aramoto, M. (2007). Identification of methyl chlorideemitting plants and atmospheric measurements on a subtropical island. Chemosphere *69*, 549–553.
- Yurimoto, H. (2009). Molecular basis of methanol-inducible gene expression and its application in the methylotrophic yeast *Candida boidinii*. Biosci. Biotechnol. Biochem. *73*, 793–800.
- Zahn, K., Inui, M., and Yukawa, H. (2000). Divergent mechanisms of 5'23S rRNA IVS processing in the α-*proteobacteria*. Nucleic acids research *28*, 4623–4633.

APPENDICES





UNIVERSITE DE STRASBOURG

RESUME DE LA THESE DE DOCTORAT

Discipline : Sciences du vivant

Spécialité : Aspects moléculaires et cellulaires de la biologie

Présentée par : Pauline CHAIGNAUD

Cotutelle de thèse entre l'Université de Strasbourg et l'Université de Bayreuth (Allemagne)

Titre : Le rôle des bactéries dans le filtrage du chlorométhane, un gaz destructeur de la couche d'ozone – Des souches modèles aux communautés microbiennes de sols forestiers

Unité de Recherche : UMR7156 UNISTRA CNRS

Laboratoire de Génétique Moléculaire, Génomique, Microbiologie (GMGM) Département "Micro-organismes, génomes, environnement" Equipe "Adaptations et Interactions Microbiennes dans l'Environnement" 28 rue Goethe, 67083 Strasbourg

Directrice de thèse : Françoise BRINGEL, Directrice de Recherche au CNRS Localisation: Laboratoire GMGM, 28 rue Goethe, 67000 Strasbourg

Co-Directeur de thèse : Steffen KOLB, Directeur d'institut Landscape Biogeochemistry (LBG) Localisation: ZALF, Leibniz Centre for Agricultural Landscape Research, Müncheberg, Allemagne

ECOLES DOCTORALES :

ED - Sciences de l'Homme et des sociétés	ED 269 - Mathématiques, sciences de
ED 99 – Humanités	l'information et de l'ingénieur
ED 101 – Droit, sciences politique et histoire	ED 270 – Théologie et sciences religieuses
ED 182 – Physique et chimie physique	ED 413 – Sciences de la terre, de l'univers et de l'environnement
🗌 ED 221 – Augustin Cournot	🔀 ED 414 – Sciences de la vie et de la santé
ED 222 - Sciences chimiques	

1. Introduction

Le chlorométhane (CH₃Cl) est un composé organique volatile, responsable de 16 % de la dégradation de l'ozone stratosphérique, liée aux composés chlorés [1]. Ses sources et ses puits sont surtout naturels [2,3]. Les échanges de chlorométhane entre les écosystèmes terrestres et l'atmosphère peuvent être modulés par les microorganismes capables de dégrader le chlorométhane, notamment ceux du sol [4,5]. De fait, le sol forestier constitue un puits important estimé à 137 Gg. an⁻¹ [6,7]. Cependant, le rôle de filtre biologique des bactéries dans les flux d'émission de chlorométhane dans l'atmosphère reste mal évalué dans le budget global du chlorométhane. A ce jour, la seule voie métabolique connue de dégradation du chlorométhane, appelée voie cmu pour « ChloroMethane Utilization », a été caractérisée en condition oxique chez Methylobacterium extorquens CM4 [8,9]. Cette bactérie est méthylotrophe, c'est-à-dire capable d'utiliser des composés réduits sans liaison carbonecarbone comme unique source de carbone et d'énergie tels que le chlorométhane et le méthanol (CH₃OH), qui est le substrat de référence de la méthylotrophie. Les gènes de la voie *cmu* sont portés par le plasmide pCMU01, essentiel à l'utilisation du chlorométhane, qui contient également des gènes impliqués dans le métabolisme des cofacteurs associés à la voie *cmu*, le tétrahydrofolate (H₄F) et un cofacteur corrinoïde [10]. Parmi les gène *cmu*, le gène *cmuA* codant la chlorométhane-méthyltransférase est essentiel à la croissance sur chlorométhane et conservé chez d'autres souches chlorométhane-dégradantes [11]. C'est pourquoi, le gène cmuA est utilisé comme gène fonctionnel marqueur de la dégradation bactérienne du chlorométhane et sa présence a été mise en évidence au sein de communautés d'Alphaproteobacteria chlorométhane-dégradantes d'environnements marins et de sols [4,11–14]. L'utilisation microbienne du chlorométhane comme unique source de carbone et d'énergie dans le sol est restreinte par le fait que les concentrations ambiantes sont faibles, de l'ordre de 1 pM de chlorométhane par g de sol [15]. Pour activement contribuer au filtrage du chlorométhane émis même en faible quantité, les bactéries chlorométhane-dégradante du sol pourraient en plus du chlorométhane utiliser un autre composé en C1 présent en plus fortes concentrations dans le sol, tel que méthanol dont la concentration est estimée à 1 nM.g⁻¹ [16]. Un des objectifs de mon projet de thèse était de tester cette hypothèse en caractérisant des communautés bactériennes méthylotrophes de sols forestiers capables d'utiliser le chlorométhane seul ou en présence de méthanol.

2. Projet de thèse

Mon projet de thèse s'est articulé autour de deux volets : de l'étude d'une souche modèle cultivée au laboratoire vers l'étude de communautés bactériennes chlorométhane-dégradantes en microcosmes de sols forestiers. Le premier volet a consisté à approfondir nos connaissances fondamentales de la réponse adaptative au niveau transcriptionnel de la souche modèle M. extorquens CM4 à la croissance sur le chlorométhane et le méthanol ajoutés comme unique source de carbone et d'énergie. Pour mieux cibler la réponse spécifique à la croissance sur le chlorométhane, une approche de transcriptomique comparative par RNA sequencing a été réalisée chez Methylobacterium extorquens CM4, mais également chez une souche proche phylogénétiquement, Methylobacterium extorquens DM4, capable d'utiliser le dichlorométhane (CH₂Cl₂) et non le chlorométhane comme unique source de carbone et d'énergie. Les transcriptomes des deux souches en condition de croissance sur des méthanes chlorés (chlorométhane ou dichlorométhane) ont été comparés à ceux obtenus en croissance sur le méthanol, et ceci en collaboration avec le Génoscope. Dans le deuxième volet de mes travaux, une approche de type Stable Isotope Probing (SIP) [17] a été appliquée à des microcosmes préparés à partir de sol forestier, incubés avec du méthanol et/ou du chlorométhane marqués avec du ¹³C, afin d'être incorporés spécifiquement dans l'ADN de communautés actives dans l'assimilation de ces composés. L'ADN enrichi en ¹³C (ADN « lourd ») a été séparé de l'ADN non marqué (« léger » portant du ¹²C) par ultracentifugation. A partir de l'ADN « lourd », la diversité bactérienne est estimée par séquençage de produits PCR d'un marqueur taxonomique (gène codant l'ARNr 16S) et de marqueurs fonctionnels ciblant la voie *cmu* (gène *cmuA*), la dégradation du méthanol (gène codant une sous-unité d'une méthanol déshydrogénase) et le métabolisme méthylotrophe central (gène mch) [16].

3. Résultats

3.1. Réponse transcriptionnelle d'une bactérie modèle à la croissance sur le chlorométhane en condition contrôlée au laboratoire. Parmi les 6276 CDS du génome de *M. extorquens* CM4, 137 ont été détectés comme différentiellement abondants uniquement en chlorométhane, avec comme attendu, les gènes *cmu* et d'autres gènes impliqués dans le métabolisme et le transport de cofacteurs essentiels à l'utilisation du chlorométhane par la voie *cmu* portée par le plasmide pCMU01. De plus, des gènes communs aux génomes de *M. extorquens* (core génome) pourraient jouer un rôle potentiel dans l'expulsion de protons produits lors de la déshalogénation (gène *hppA* codant une pyrophosphatase translocatrice de protons), la

régénération du NADP (gène *pnt* codant une « membrane-bound transhydrogenase ») et le métabolisme C1 du H₄F (gènes *gcvPHT* du complexe de clivage de la glycine). L'ensemble de ces données de transcriptomique, en accord avec des travaux précédents [18], démontre le rôle central du plasmide pCMU01 porteur de gènes plus exprimés en condition de culture sur chlorométhane non seulement pour les gènes essentiels de la voie *cmu*, mais aussi d'une vingtaine de gènes ayant des homologues portés par le chromosome, qui eux ne sont pas différentiellement exprimés. Enfin, l'utilisation du chlorométhane semble peu corrélée à celle du dichlorométhane (12 gènes du core génome ont des transcrits plus abondants en culture sur méthanes chlorés comparé au méthanol). Un 1^{er} article à soumettre sous peu rassemble les résultats du travail effectué à Strasbourg.

3.2. Caractérisation de la diversité des communautés bactériennes dégradant le chlorométhane dans des microcosmes de sol forestier. Les résultats du séquençage des fractions enrichies en ¹³C ont permis de mettre en évidence la présence de nouvelles communautés bactériennes capables d'utiliser le méthanol et le chlorométhane. Des *Alphaproteobacteria* du genre *Methylovirgula* ont été détectées comme majoritaires dans la communauté bactérienne capable d'utiliser soit le chlorométhane, soit le méthanol comme source de carbone. L'ajout simultané de méthanol et de chlorométhane a permis la caractérisation d'une nouvelle famille de bactéries assimilatrices de chlorométhane apparentées aux *Actinomycetales* du genre *Actinothermus*. L'analyse des séquences du gène *cmuA* a révélé qu'une partie des OTUs (Operational Taxonomic Unit) identifiées sont proches phylogénétiquement du gène *cmuA* de *Methylobacterium extorquens* CM4, alors que d'autres sont affiliées à des souches non cultivables de l'environnement. Un 2^e article en préparation rassemblera les résultats des travaux réalisés à Bayreuth.

4. Conclusions et perspectives

L'étude du transcriptome par du RNAseq, de deux souches phylogénétiquement proches, *Methylobacterium extorquens* CM4 et DM4, a permis une meilleure compréhension de l'implication des génomes communs et spécifiques dans l'adaptation liée à la croissance en présence de méthanol ou de chlorométhane, démontrant l'importance des génomes variables dans cette adaptation. Cependant, les éléments et les mécanismes de régulation liés au métabolisme des composés en C₁ restent encore inconnus. Dans le but de les identifier, des banques directionnelles pour l'identification des sites d'initiation de la transcription ont été préparées et sont actuellement en cours de séquençage. L'approche SIP a permis une première évaluation de la diversité des communautés bactériennes chlorométhane-dégradantes, avec ou sans ajout de méthanol, de sols forestiers. Des bactéries du genre *Methylovirgula* ou *Acidothermus* ont été nouvellement identifiés comme actives dans l'assimilation du chlorométhane dans des microcosmes de sols forestiers. Des approches non culturales (métagénomique; marquage FISH pour « Fluorescence In Situ Hybridation »; PCR quantitative) couplées à des approches culturales permettraient à l'avenir de mieux estimer l'abondance, la diversité, les voies métaboliques impliquées dans le filtrage du chlorométhane par les communautés microbiennes du sol.

5. Bibliographie

- 1. Montzka S, Reimann S, O'Doherty S, Engel A, Krüger K, Sturges WT. Ozone-depleting substances (ODSs) and related chemicals. 2011; Available: http://oceanrep.geomar.de/10405/2/03-Chapter_1.pdf
- Cox MJ, Schäfer H, Nightingale PD, McDonald IR, Murrell JC. Diversity of methyl halide-degrading microorganisms in oceanic and coastal waters. FEMS Microbiol Lett. 2012;334: 111–118. doi:10.1111/j.1574-6968.2012.02624.x
- Keppler F, Harper DB, Röckmann T, Moore RM, Hamilton JTG. New insight into the atmospheric chloromethane budget gained using stable carbon isotope ratios. Atmospheric Chem Phys. 2005;5: 2403– 2411.
- Borodina E, Cox MJ, McDonald IR, Murrell JC. Use of DNA-stable isotope probing and functional gene probes to investigate the diversity of methyl chloride-utilizing bacteria in soil. Environ Microbiol. 2005;7: 1318– 1328.
- Clerbaux C, Cunnold DM, Anderson J, Engel A, Fraser PJ, Mahieu E, et al. Long-lived organic compounds (Chapter 1). Sci Assess Ozone Deplet 2006. 2007; Available: http://orbi.ulg.ac.be/handle/2268/34986
- Harper DB. The global chloromethane cycle: biosynthesis, biodegradation and metabolic role (1982 to January 2000). Nat Prod Rep. 2000;17: 337–348. doi:10.1039/a809400d
- Yoshida Y. A three-dimensional global model study of atmospheric methyl chloride budget and distributions. J Geophys Res. 2004;109. doi:10.1029/2004JD004951
- Vannelli T, Studer A, Kertesz M, Leisinger T. Chloromethane metabolism by *Methylobacterium* sp. strain CM4. Appl Environ Microbiol. 1998;64: 1933–1936. doi:05/1998; 64(5):1933-6.
- Vannelli T, Messmer M, Studer A, Vuilleumier S, Leisinger T. A corrinoid-dependent catabolic pathway for growth of a *Methylobacterium* strain with chloromethane. Proc Natl Acad Sci. 1999;96: 4615–4620. doi:10.1073/pnas.96.8.4615
- 10. Messmer M, Wohlfarth G, Diekert G. Methyl chloride metabolism of the strictly anaerobic, methyl chlorideutilizing homoacetogen strain MC. Arch Microbiol. 1993;160: 383–387.
- Nadalig T, Farhan Ul Haque M, Roselli S, Schaller H, Bringel F, Vuilleumier S. Detection and isolation of chloromethane-degrading bacteria from the *Arabidopsis thaliana* phyllosphere, and characterization of chloromethane utilization genes: chloromethane-degrading bacteria from *A. thaliana*. FEMS Microbiol Ecol. 2011;77: 438–448. doi:10.1111/j.1574-6941.2011.01125.x
- Miller LG, Warner KL, Baesman SM, Oremland RS, McDonald IR, Radajewski S, et al. Degradation of methyl bromide and methyl chloride in soil microcosms: Use of stable C isotope fractionation and stable isotope probing to identify reactions and the responsible microorganisms. Geochim Cosmochim Acta. 2004;68: 3271–3283. doi:10.1016/j.gca.2003.11.028
- Nadalig T, Greule M, Bringel F, Keppler F, Vuilleumier S. Probing the diversity of chloromethane-degrading bacteria by comparative genomics and isotopic fractionation. Front Microbiol. 2014;5. doi:10.3389/fmicb.2014.00523
- Schäfer H, McDonald IR, Nightingale PD, Murrell JC. Evidence for the presence of a CmuA methyltransferase pathway in novel marine methyl halide-oxidizing bacteria: Novel marine methyl halide-oxidizing bacteria. Environ Microbiol. 2005;7: 839–852. doi:10.1111/j.1462-2920.2005.00757.x
- Harper DB, Hamilton JTG, Ducrocq V, Kennedy JT, Downey A, Kalin RM. The distinctive isotopic signature of plant-derived chloromethane: possible application in constraining the atmospheric chloromethane budget. Chemosphere. 2003;52: 433–436. doi:10.1016/S0045-6535(03)00206-6
- Stacheter A, Noll M, Lee CK, Selzer M, Glowik B, Ebertsch L, et al. Methanol oxidation by temperate soils and environmental determinants of associated methylotrophs. ISME J. 2013;7: 1051–1064. doi:10.1038/ismej.2012.167
- 17. Neufeld JD, Dumont MG, Vohra J, Murrell JC. Methodological Considerations for the Use of Stable Isotope Probing in Microbial Ecology. Microb Ecol. 2007;53: 435–442. doi:10.1007/s00248-006-9125-x
- Roselli S, Nadalig T, Vuilleumier S, Bringel F. The 380 kb pCMU01 plasmid encodes chloromethane utilization genes and redundant genes for vitamin B₁₂- and tetrahydrofolate-dependent chloromethane metabolism in *Methylobacterium extorquens* CM4: a proteomic and bioinformatics study. PLoS ONE. 2013;8: e56598. doi:10.1371/journal.pone.0056598





PhD thesis abstract and key words

Pauline CHAIGNAUD

Bacteria as chloromethane sinks – from model strains to forest soil communities

Chloromethane (CH₃Cl) is a volatile organic compound responsible for over 15% of stratospheric ozone degradation due to chlorinated compounds. It is mainly produced by living and decaying plants. Bacteria utilizing CH₃Cl as sole carbon and energy source for growth were shown to be involved in the filtering of CH₃Cl emissions to the atmosphere. This biological process remains to be quantified in the environment, especially for forest soil, a major CH₃Cl sink. The *cmuA* gene is used as a biomarker of bacterial CH₃Cl degradation in environmental studies. It encodes a CH₃Cl methyltransferase essential for bacterial growth by the *cmu* (chloromethane utilization) pathway for growth with CH₃Cl and the only one characterized so far. My thesis project had a double aim: i) In depth studies of CH₃Cl adaptation of a model methylotrophic bacterium, *Methylobacterium extorquens* strain CM4; ii) Exploration of bacterial CH₃Cl-utilizers in forest

An RNAseq study of strain CM4 has shown that growth with CH₃Cl leads to a difference of transcription of 137 genes in its 6.2 Mb genome compared to growth with methanol (CH₃OH). Among those, genes of the *cmu* pathway and other genes involved in the metabolism of essential cofactors for CH₃Cl utilization by this pathway, are all plasmid pCMU01-encoded. Paralogous genes located on the chromosome were not differentially expressed. On the other hand, other chromosomal genes potentially involved in extruding protons generated during CH₃Cl deshalogenation (*hppA*), NADP⁺ regeneration (*pnt*), or in the cofactor tetrahydrofolate metabolism (*gcvPHT*) were differentially expressed.

The diversity of CH₃Cl-degrading bacteria in forest soil of the German natural park of Steigerwald was studied in microcosms using stable isotope probing. Microorganisms able to assimilate labeled [¹³C]-CH₃Cl incorporate this heavy carbon isotope in their DNA. Sequence analysis of the PCR-amplified 16S RNA encoding gene from [¹³C]-DNA fractions uncovered phylotypes of the genus *Methylovirgula* and of the order of the *Actinomycetales*, which were not associated with bacterial CH₃Cl degradation so far. In contrast, PCR-amplified sequences of *cmuA* and other genes of methylotrophic metabolism were closely related to known CH₃Cl-degrading isolates. These results suggest that bacteria containing genes of the *cmu* pathway contribute to biological filtering of CH₃Cl in forest soil. Future experiments coupling molecular and culture methods will aim to discover new CH₃Cl-degrading pathways and to characterize the abundance and diversity of CH₃Cl-degradation metabolism in soil and other environmental compartments.

<u>Key words</u>: chloromethane, bacterial dechlorination, methylotrophy, *Methylobacterium extorquens* CM4, bacterial communities, RNA sequencing, stable isotope probing, forest soil, microcosms



Pauline CHAIGNAUD



Le rôle des bactéries dans le filtrage du chlorométhane, un gaz destructeur de la couche d'ozone –

des souches modèles aux communautés microbiennes de sols forestiers

Die Rolle von Bakterien als Filter für Chlormethan, ein Ozonschicht-schädigendes Gas –

Von Modellorganismen zu mikrobiellen Gemeinschaften im Waldboden

Le chlorométhane (CH₃Cl) est un composé organique volatile responsable de plus de 15 % de la dégradation de l'ozone stratosphérique due aux composés chlorés. Il est produit majoritairement par les plantes vivantes ou en décomposition. Les bactéries capables d'utiliser le CH₃Cl comme source de carbone pour leur croissance peuvent jouer un rôle de filtre dans les émissions de CH₃Cl vers l'atmosphère. Ce processus biologique reste à quantifier dans l'environnement, notamment pour les sols forestiers considérés comme un puits majeur de ce composé. Dans les études environnementales, le gène *cmuA* est utilisé comme biomarqueur de la dégradation bactérienne du CH₃Cl. Il code une chlorométhane méthyltransférase essentielle à la croissance bactérienne avec le CH₃Cl par la voie *cmu* (pour chloromethane utilization), la seule caractérisée à ce jour. Mon projet de thèse avait un double objectif : i) l'approfondissement des connaissances de l'adaptation au CH₃Cl chez une bactérie méthylotrophe modèle, *Methylobacterium extorquens* CM4; ii) l'exploration de la diversité des bactéries CH₃Cl-dégradantes de sols forestiers.

L'étude RNAseq chez la souche CM4 a montré que la croissance avec le CH₃Cl s'accompagne de différences dans la transcription de 137 gènes de son génome (6.2 Mb) par rapport à sa croissance sur le méthanol (CH₃OH). Les gènes de la voie *cmu*, ainsi que d'autres gènes impliqués dans le métabolisme de cofacteurs essentiels à l'utilisation du CH₃Cl par cette voie et eux aussi portés par le plasmide pCMU01 de la souche, en font partie. Les paralogues de ces gènes localisés sur le chromosome ne sont quant à eux pas différentiellement exprimés. En revanche, d'autres gènes du chromosome, potentiellement impliqués dans l'excrétion de protons produits lors de la déshalogénation (*hppA*), la régénération du NADP⁺ (*pnt*), ou le métabolisme du cofacteur tétrahydrofolate (gènes *gcvPHT*), le sont.

L'étude de la diversité des bactéries CH₃Cl-dégradantes de sol forestier de la réserve naturelle de Steigerwald (Allemagne) a été réalisée sur des microcosmes par une approche de « Stable Isotope Probing ». Les micro-organismes capables d'assimiler le CH₃Cl marqué au [¹³C] incorporent cet isotope lourd du carbone dans leur ADN. L'analyse des séquences amplifiées par PCR des gènes codant l'ARN 16S des fractions d'ADN enrichies en [¹³C] a permis de mettre en évidence de nouveaux phylotypes, du genre *Methylovirgula* et de l'ordre des *Actinomycetales*, distincts de ceux auxquelles les souches dégradant le CH₃Cl isolées jusqu'ici sont affiliées. En revanche, les séquences du gène *cmuA* et d'autres gènes du métabolisme méthylotrophe obtenues par PCR à partir de l'ADN enrichi en [¹³C] sont très proches de celles des souches CH₃Cl-dégradantes connues. Les résultats obtenus suggèrent ainsi que des bactéries ayant acquis par transfert horizontal les gènes de dégradation de la voie *cmu* ou ne possédant pas de gène *cmuA* contribuent au filtrage biologique du CH₃Cl des sols forestiers. A l'avenir, le couplage de différentes méthodes moléculaires et des approches culturales visera à découvrir de nouvelles voies microbiennes de l'utilisation du CH₃Cl, et à caractériser l'abondance et la diversité des métabolismes impliqués dans la dégradation du CH₃Cl dans les sols et d'autres compartiments environnementaux.

<u>Mots-clés</u> : chlorométhane, déchloration bactérienne, méthylotrophie, *Methylobacterium extorquens* CM4, communautés bactériennes, RNA sequencing, stable isotope probing, sols forestiers, microcosmes

Chloromethane (CH₃Cl Chlormethan (CH₃Cl) ist eine volatile organische Verbindung (BVOC), die für mehr als 15% des stratosphärischen Ozonabbaus verantwortlich ist. Es wird vor allem von lebenden und toten Pflanzenteilen gebildet. Bakterien, die CH₃Cl als Kohlenstoff- und Energiequelle für ihr Wachstum nutzen, sind an der Reduktion der CH₃Cl -Emissionen in die Atmosphäre beteiligt. Dieser biologische Prozess wurde bislang in der Umwelt nicht quantifiziert, insbesondere in Waldböden, die eine wichtige biologische Senke darstellen. Das *cmuA*-Gen wird als Biomarker zur Detektion von bakterieller CH₃Cl -Abbau in der Umwelt eingesetzt. Es kodiert für eine CH₃Cl -Methyltransferase, die essentiell für Wachstums mittels des *cmu* (<u>c</u>hloro<u>m</u>ethane <u>u</u>tilization)-Stoffwechsel ist – der bislang einzig bekannte Abbau-Stoffwechselweg. Mein Promotionsprojekt verfolgte zwei Ziele: i) Hochauflösende Untersuchungen am Transkriptom zur Anpassung an die CH₃Cl -Nutzung im Model-Bakterium *Methylobacterium extorquens* Stamm CM4; ii) Erfassung von bakteriellen CH₃Cl -Nutzern in einem Waldboden.

In einer RNAseq-Studie am Stamm CM4 konnte gezeigt werden, dass Wachstum auf CH₃Cl zu unterschiedlichen Transkription von 137 Genen des 6.2 Mb-Genom im Vergleich mit Wachstum auf Methanol (CH₃OH) führte. Von diesen Genen sind alle Gene des *cmu*-Stoffwechselweges und die Gene zur Synthese essentieller Kofaktoren auf dem Plasmid pCMU01 kodiert. Paraloge, die sich auf dem Chromosom befinden, wurden nicht differentiell exprimiert. Andererseits wurden chromosomale Gene, die potentiell am Export von während der Dehalogenierung gebildeten Protonen (*hppA*), an der NADP⁺-Regeneration (*pnt*) oder an der Synthese des Kofaktors Tetrathydrofolat (*gcvPHT*) beteiligt sind, differentiell exprimiert.

Im Zweiten Teil des Promotionsprojektes wurde die Biodiversität der CH₃Cl-abbauenden Bakterien in einem Waldboden im deutschen Naturpark Steigerwald mittels Stabiler-Isotopen-Beprobung in Mikrokosmen untersucht. Mikroorganismen, die [¹³C]-CH₃Cl assimilierten, inkorporierten dieses stabile Isotop in ihrer DNA. Sequenz-Analysen markierter DNA anhand des Genmarkers 16S-rRNA-Gen ergaben Phylotypen der Gattung *Methylovirgula* und der Ordnung *Actinomycetales* - Bakterien, die bislang nicht mit CH₃Cl-Abbau in Zusammenhang gebracht werden konnten. Dahingegen waren die markierten Genotypen des Gens *cmuA* und anderer Gene aus dem C1-Stoffwechsel nah zu bekannten Genotypen bereits kultivierter Reinkulturen verwandt. Diese Ergebnisse legen nahe, dass Bakterien, die Gene des *cmu*-Stoffwechsels besitzen, diese durch horizontalen Gentransfer aquiriert haben bzw. dass es CH₃Cl-abbauende Bakterien gibt, die nicht den *cmu*-Stoffwechselweg besitzen. Zukünftige Untersuchungen, die molekulare und Kultivierungsmethoden kombinieren werden, sollen neue CH₃Cl-Abbau-Stoffwechselwege identifizieren und darüber hinaus die Abundanz und Diversität der CH₃Cl-Abbaustoffwechsel in Böden und anderen Umwelten aufklären.

<u>Stichworte</u>: Chlormethan, bakterielle Dechlorinierung, Methylotrophie, *Methylobacterium extorquens* CM4, bakterielle Gemeinschaften, RNA-Sequenzierung, Stabile-Isotopen-Beprobung, Waldboden, Mikrokosmen