



**HAL**  
open science

# Mécanismes de coexistence de deux espèces sympatriques d'otaries, *Arctocephalus australis* et *Otaria byronia*, à Punta San Juan, au Pérou

Susana Cardenas Alayza

► **To cite this version:**

Susana Cardenas Alayza. Mécanismes de coexistence de deux espèces sympatriques d'otaries, *Arctocephalus australis* et *Otaria byronia*, à Punta San Juan, au Pérou. Sciences agricoles. Université Montpellier; Universidad Peruana Cayetano Heredia, 2021. Français. NNT : 2021MONTG077 . tel-03604632

**HAL Id: tel-03604632**

**<https://theses.hal.science/tel-03604632v1>**

Submitted on 10 Mar 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**THÈSE POUR OBTENIR LE GRADE DE DOCTEUR  
DE L'UNIVERSITÉ DE MONTPELLIER**

**En Écologie fonctionnelle**

**École doctorale GAIA**

**Unité mixte de recherche MARBEC**

**En partenariat international avec l'Universidad Peruana Cayetano Heredia, PÉROU**

**Mécanismes de coexistence de deux espèces sympatriques  
d'otaries, *Arctocephalus australis* et *Otaria byronia*, à Punta  
San Juan, au Pérou**

**Présentée par Susana CÁRDENAS ALAYZA  
Le 21 décembre 2021**

**Sous la direction de Yann TREMBLAY  
et Dimitri GUTIÉRREZ**

**Devant le jury composé de**

Jean-Yves GEORGES, Directeur de recherche, CNRS  
Arnaud BERTRAND, Directeur de recherche, IRD, MARBEC  
Patricia MAJLUF, Professeur, Universidad Peruana Cayetano Heredia  
Carlos ZAVALAGA, Professeur, Universidad Científica del Sur  
Dimitri GUTIERREZ, Professeur, Universidad Peruana Cayetano Heredia  
Carey Elizabeth KUHN, Chargée de recherche, NOAA  
Yann TREMBLAY, Chargé de recherche, IRD, MARBEC

Président du jury  
Membre du jury  
Membre du jury  
Membre du jury  
Co-directeur de thèse  
Rapporteur  
Co-directeur de thèse



**UNIVERSITÉ  
DE MONTPELLIER**

## Avant-Propos

Cette thèse de doctorat a été financée par la Escuela Doctoral Franco Peruana en Ciencias de la Vida, Universidad Peruana Cayetano Heredia et l'Ecole Doctorale GAIA Biodiversité, Agriculture, Environnement, Terre, Eau, Université de Montpellier, France. Au Pérou, ce travail a été réalisé sous la direction du Dr Dimitri Gutierrez au Laboratorio de Ciencias del Mar de l'Universidad Peruana Cayetano Heredia à Lima. En France, sous la direction du Dr. Yann Tremblay à l'Institut de Recherche pour le Développement (IRD) - UMR MARBEC au sein de la station marine de Sète.

La collecte de données à Punta San Juan, Pérou, a été possible grâce à des accords de collaboration inter-institutionnelle entre l'Universidad Peruana Cayetano Heredia, Servicio Nacional de Areas Naturales Protegidas del Peru (SERNANP) du Ministerio del Ambiente (MINAM) et AgroRural du Ministerio Nacional de Agricultura y Riego (MINAGRI). Nous tenons à remercier le SERNANP pour l'accès à la réserve de Punta San Juan et AgroRural pour l'utilisation des installations de terrain.

Nous remercions la Chicago Zoological Society, le Saint Louis Zoo, le Kansas City Zoo et le Woodland Park Zoo pour le financement des opérations du programme Punta San Juan, qui sont essentielles à la collecte des données à long terme analysées dans cette enquête. Nous remercions également le Chicago Board of Trade Endangered Species Fund, la Society for Marine Mammalogy's Grant-in-Aid of Research, la Paul M. Angel Foundation et la famille Feay pour leur soutien financier. Je remercie également le fonds de mobilité MARBEC pour l'aide qu'il m'a apportée sous forme de bourse de voyage pendant mes études de doctorat.

Enfin, nous remercions tous les bénévoles, les stagiaires et le personnel du programme Punta San Juan du Centro para la Sostenibilidad Ambiental de l'Universidad Peruana Cayetano Heredia qui ont participé aux travaux sur le terrain, à la collecte des échantillons et à la préparation de cette étude.

## Remerciements

This work could not have been possible by so many people, it will be hard to make sure they can all fit in these pages. Although the doctoral thesis has been a special time dedicated to learning, answering research questions, study design, data analysis, statistical modelling, writing and publishing... This investigation also carries within it information collected on fur seals and sea lions in Punta San Juan since 2001, which has involved the help from many people. So, here we go...

An important place to start is to thank all the staff at the Punta San Juan Program. Over the past years, since I have become the Program Director, I thank my team for being open to my ideas and plans for collecting information, samples and tagging animals in the field. And, also for continuing the work while I have been busy with this PhD. The Punta San Juan Program is only possible by motivated people that are willing to learn and innovate for research and conservation. For this I thank past and present field staff Marco Cardeña, Paulo Colchao, Daniel Torres, Juan Carlos Jerí; Research Assistants María José Ganoza, Karla Calderón, Lyanne Ampuero and administrative/operations staff Ken Rojas and Fernando Nishio who have also been an important part of “getting things done”. And of course, a long list of incredible interns and volunteers that through the years have made sample collection and analyses possible as part of their training in the Punta San Juan Program.

For Chapter 1 (counts) and Chapter 2 (diet), I would like to thank the dedication of the Punta San Juan field staff who have counted pinnipeds and collected samples monthly, weekly (or even daily!) with care and meticulousness, in a way that we can trust the information. For this, first and foremost, I thank Patricia Majluf, who has set a high standard for long-term field research and monitoring programs in Peru through the establishment of the Punta San Juan Program. I have the luxury of being able to continue what she started and, I hope, that this academic contribution gives justice and can help answer some of the questions we have been discussing for years. Scat sample collection and analysis is also a big part of the job in pinniped trophic ecology research. For this I extend a special thanks to Marco Cardeña, Walter Díaz, Franco Sandoval, Paulo Guerrero, Marco Quispe, Paulo Colchao and Daniel Torres for their years of dedication to complete the job. Chapter 2 would not have been possible without the incredible dedication and perseverance of Daniel Torres, who has spent a countless number of hours cleaning samples and identifying fish otoliths in recent years, and teaching others how to get these tasks done as part of his position in the PSJ Program. I would also

like to thank Leticia Escobar and the team of scat volunteers we organized at the beginning of my PhD to work through the sea lion scat samples which we all know is -literally- a mountain of work. I started working on tracking pinnipeds (Chapter 3) when I returned to Peru after obtaining my master's on fur seal population dynamics, that left me with more questions than answers (which I consider positive!). I wanted to know where fur seals were obtaining their food and if they were competing with sea lions. When I started this journey, who would have thought that those old refurbished satellite tags kindly donated from Andrew Trites back in 2013, would turn out to be the first tagged South American sea lions in Peru ?! Thanks Andrew, for the leap of faith. Chapter 3 has involved a lot of people in the field. I would like to thank my *partner in tagging*, wildlife veterinarian Mike Adkesson. With Mike we have organized many field seasons with pinnipeds in PSJ with multiple objectives. Together we've obtained funding to make these field seasons in Peru possible, imported a countless amount of equipment (and people!) to Peru; as well as contacted experts to prepare for our field seasons to get things right. As we learned how to do this, we have trained many biologists, veterinarians, veterinary technicians and together with them, we continue to learn. My deepest thanks to all the teams that have worked with us in the field during the 2013-2019 PSJ pinniped seasons included in this thesis. I would like to give special remembrance to Gwen Jankowski, an incredible person and veterinarian who quickly connected with the magic of Punta San Juan and whom with I worked closely tagging female fur seals in PSJ. Gwen, you are deeply missed and I know you would be so interested in knowing about all these results. Here is to you, friend!

For Chapter 4 (isotopic signals), work with whiskers was also triggered by Andrew Trites, who connected me with Amy Hirons to explore possible El Niño signals. This gave birth to Mickie Edwards masters project (Frontiers article) and started what has become a larger collaboration with Nova Southeastern University. Sample preparation for Chapter 4 was possible thanks to the support from Amy Hirons, Mickie Edwards and all the volunteers and students at the Hiron's lab which I am grateful to. The analyses of the data has been a collaboration with Valentina Franco-Trecu, whom I am forever in debt to for having the time and patience for teaching me about analyses of stable isotope data with repeated measures, while nurturing what today I consider a great friendship.

This doctoral thesis started out as a project that was in the back of my mind since many years ago; but it took me a while to accept that while working on developing the Punta San Juan Program, I never had enough time to get all the analyses done in the depth that they required. Thanks to the

unexpected roads that life takes you down, this PhD suddenly became a reality. I would like to thank Yann Tremblay and Dimitri Gutiérrez, my supervisors who accepted to join me on this research (and administrative!) adventure to develop this thesis. Even if that meant a team of three, working most of the time in either two (or three) countries (bear in mind, this was in pre-pandemic era). Regardless, I always felt they respected my opinion and let me lead the project, moving it forward with their support. I now consider the latter crucial, to face the real world of academia and project management. I would also like to thank the professors who served on my examination committee at UPCH, Wilfredo Gonzáles, Armando Valdés and Patricia Majluf as well as the committee members at Université de Montpellier Hervé Demarcq, Jean-Yves Georges and Simon Benhamou, who through their different different optics have given me valuable advice to improve and advance this research project forward.

This doctorate program involved the incredible possibility of moving to France, during which I lived in Sète for 1.5 years. During this time, while I worked in Marbec, I met some pretty amazing people whom through different gestures and conversations have helped me. I would especially like to thank Babbette Gunther, Samantha Cox, Philippine Chambault, Magali Combes (and the amazing yellow Kangoo!), Alexandra Diallo, Miriam Brandt, Geraldine Pérez, Alexandre Mignucci, Laura Mannocci. I would also like to thank my dearest friends from UBC -Tiphainne Jeanniard du Dot, Ruth Joy and Rowenna Gryba- whom with I reconnected during this PhD and have been an incredible source of support and inspiration. May the WD sessions never stop! Thank you to Basak Uysal, whom with after casually meeting in a class in U.Montpellier, became eachothers PhD accountability partners and have been mutually supporting eachother since. Thank you to “mi tía Guri” who helped me quickly find an affordable way to live back in Lima in the middle of the pandemic when all I had to do was continue with this thesis. A special timeless thank you to my parents for always supporting me while chasing after my dreams, even if it involved working with fur seals and sea lions in guano (for many years). I would like to thank my sister for always lending a helping hand, letting me pick her brain and giving critical advice on the science -of life-! Thank you to her and James for the walks, long saturday lunches, the dancing and listening to all my updates and frustrations with research. And last but definetly not least, to Nico and Pepa, my amazing support team. They are the backbone of this work. Thank you for believing in me and understanding all the hours I had to dedicate to this. My stability, my fun, my strength, my laughter, my imagination, my freedom and my love is always growing thanks to them. And, I am sure, this is just the beginning... My deepest thanks to you all.

## Résumé de Thèse

Comprendre comment des espèces sympatriques ayant des besoins similaires coexistent est une question majeure en écologie. Une communauté où le partage des ressources ou le chevauchement des niches est plus important accueillera plus d'espèces qu'une autre où il l'est moins. À cet égard, le principe d'exclusion compétitive prédit que les espèces doivent avoir des mécanismes de ségrégation pour coexister. Dans le système du courant de Humboldt (HCS) au Pérou, les lions de mer sud-américains (SASL, *Otaria byronia*) coexistent avec les otaries à fourrure sud-américaines (SAFS, *Arctocephalus australis*). Cependant, le partitionnement trophique, temporel et spatial lors de la recherche de nourriture reste largement inconnu. Cette étude se déroule dans la réserve nationale de Punta San Juan (PSJ) (15° 22'S, 75° 12'W) au Pérou, où se trouvent d'importantes colonies de reproduction des deux espèces d'otariidés, ce qui en fait un cadre idéal pour étudier les mécanismes de coexistence. L'objectif principal de cette thèse est décrire et de comparer les niches écologiques des espèces d'otariidés sympatriques au Pérou, pour déterminer la différenciation/le chevauchement des niches intraspécifiques et interspécifiques. Ceci sera abordé à travers trois objectifs spécifiques qui incluent décrire et de comparer (i) l'écologie trophique des pinnipèdes sympatriques; (ii) les variables environnementales qui caractérisent les niches écologiques et (iii) les stratégies de recherche de nourriture (dimensions spatiales et temporelles) au sein des niches écologiques des pinnipèdes sympatriques, selon l'espèce et le sexe. Dans le Chapitre 1, l'analyse des trajectoires de la population montre comment sur une période de 20 ans, les deux populations de pinnipèdes sont maintenant en déclin. Une diminution des juvéniles dans la colonie, suggère que la limitation des ressources est le principal facteur contribuant au déclin actuel des populations. Dans le Chapitre 2, la ségrégation alimentaire interspécifique a été détectée, grâce à l'analyse des parties dures dans les fécès (2015-2018). Les SAFS se nourrissent principalement d'anchois du Pérou (*Engraulis ringens*) et de céphalopodes tandis que les SASL consomment surtout des galathées (*Pleuroncodes monodon*) et des anchois du Pérou, en plus petites proportions. Au cours de cette période d'étude, un chevauchement trophique a été détecté pendant un événement El Niño extraordinaire, ce qui suggère que les changements climatiques peuvent exacerber la compétition. Dans le Chapitre 3, nous avons analysé les localisations de 35 otariidés (18 SASL, 17 SAFS ; 12 femelles, 23 mâles) équipés de balises satellites (2013-2017) pour déterminer les stratégies de recherche de nourriture. Les mâles se séparent dans l'espace et dans le temps; tandis que les femelles présentent un chevauchement spatial

mais disposent de mécanismes de ségrégation temporelle. Les variables environnementales associées suggèrent des caractéristiques d'habitat pélagique et côtier chez les otaries à fourrure et les lions de mer femelles, respectivement. Par contre, les mâles présentent une plus grande variabilité interindividuelle qui devrait être exploré davantage. Quoi qu'il en soit, les résultats démontrent que les stratégies de recherche de nourriture multiples à différentes échelles contribuent à la répartition des ressources. Au Chapitre 4, la différenciation de niche à long terme a été étudiée en examinant les signatures isotopiques archivées en série dans les vibrisses SAFS et SASL (2005-2019). Les otariidés ont étendu la largeur de leur niche (à l'exception des femelles SASL) en réponse à l'augmentation du niveau moyen de la mer et au réchauffement des températures de surface de la mer. Le déclin général des signaux  $\delta^{15}\text{N}$  suggère un appauvrissement de la base du réseau alimentaire, avec des effets bottom-up qui ont impacté les consommateurs tertiaires. Ces résultats aident à expliquer la diminution actuelle de la population et peuvent servir de base pour étendre les recommandations pour la conservation des populations d'otariidés dans le HCS.

**Mots clés** : coexistence ; ségrégation ; pinnipèdes ; écologie trophique ; niches ; système du courant de Humboldt



## Thesis Abstract

Understanding how sympatric species with similar requirements coexist is a major issue in ecology. A community with more resource sharing or greater niche overlap, will support more species than one with less. In this regard, the principle of competitive exclusion predicts that species will exhibit segregation mechanisms to coexist. In the Humboldt Current System (HCS) in Peru, South American sea lions (SASL, *Otaria byronia*) coexist with South American fur seals (SAFS, *Arctocephalus australis*). However, trophic, temporal and spatial partitioning while foraging remains unclear. This study takes place in Punta San Juan (PSJ) national reserve (15 ° 22'S, 75 ° 12'W) in Peru, there are important breeding colonies of the two species of otariids, making it an ideal setting to study coexistence mechanisms. The main goal of this thesis is to describe and compare the ecological niches of sympatric otariids in Peru, to determine intraspecific and interspecific niche differentiation/overlap. This will be addressed through three specific goals which include to describe and compare (i) the trophic ecology of sympatric pinnipeds; (ii) environmental variables that characterize ecological niches; and (iii) the foraging strategies (spatial and temporal dimensions) within the ecological niches of the sympatric pinnipeds, according to species and sex. In Chapter 1, analysis of population trajectories show how over a 20 year period both pinniped populations are now in a state of decline. A decrease in juvenile age-classes in the colony, suggests that resource limitation is a main contributing factor for current population decline. In Chapter 2, interspecific dietary segregation was detected, through the analysis of hard parts in scats (2015-2018). SAFS feed primarily on Peruvian anchovy (*Engraulis ringens*) and cephalopods while SASL consume mostly red squat lobsters (*Pleuroncodes monodon*) and Peruvian anchovy, in smaller proportions. In this study period, trophic overlap was detected during an Extraordinary El Niño event, suggesting that climatic changes can exacerbate competition. In Chapter 3, I analyzed the locations of 35 otariids (18 SASL, 17 SAFS; 12 females, 23 males) equipped with satellite tags (2013-2017) to determine foraging strategies. Males segregate in space and time; whereas females exhibit spatial overlap but have temporal segregation mechanisms. Associated environmental variables suggest pelagic and coastal habitat features in female fur seals and sea lions, respectively. Whereas, males have higher interindividual variability that should be further explored. Regardless, results demonstrate how multiple foraging strategies at different scales aid resource partitioning. In Chapter 4, long-term niche differentiation was investigated by examining the isotopic signatures serially archived in whiskers

SAFS and SASL (2005-2019). Otariids expanded their niche width (except SASL females) as a response to the increase in average sea level and warmer sea surface temperatures. The overall decline in the  $\delta^{15}\text{N}$  signals suggests an impoverished baseline of the food web, with bottom-up effects that has impacted tertiary consumers. These findings help explain the current state of population decline and can serve as a basis to extend recommendations for conservation of otariid populations in HCS.

**Keywords:** coexistence; segregation; pinnipeds; trophic ecology; niches; Humboldt Current System

## Resumen de Tesis

Entender como especies simpátricas con requerimientos similares coexisten constituye un tema de gran envergadura en ecología. Una comunidad con más recursos compartidos o mayor solapamiento de nichos, soportará más especies que una con menos. En este sentido, el principio de exclusión competitiva predice que las especies mostrarán mecanismos de segregación para coexistir. En el Sistema de la Corriente de Humboldt (HCS) en Perú, los lobos marinos chuscos (SASL, *Otaria byronia*) coexisten con los lobos marinos finos (SAFS, *Arctocephalus australis*). Sin embargo, la distribución trófica, temporal y espacial durante la búsqueda de alimento no está claramente definida. Este estudio se desarrolla en la reserva nacional de Punta San Juan (PSJ) (15° 22'S, 75° 12'W) en Perú, donde se encuentran importantes colonias reproductivas de las dos especies de otáridos; haciéndolo un entorno ideal para estudiar la coexistencia. El presente estudio postula describir y comparar los nichos ecológicos de las dos especies de lobos marinos simpátricos en Perú, para determinar si existe diferenciación / sobreposición interespecífica e intraespecífica de nichos. Esto se abordará a través de tres objetivos específicos que incluyen describir y comparar: (i) los nichos tróficos; (ii) las variables ambientales; y (iii) evaluar si existen estrategias de forrajeo distintas (dimensiones espaciales y temporales) empleadas por los pinnípedos simpátricos en relación con las condiciones ambientales, según especie y sexo. En el Capítulo 1, el análisis de las trayectorias poblacionales muestra cómo después de un período de 20 años, actualmente, ambas poblaciones se encuentran en un estado de declive. La reducción de individuos juveniles en la colonia, sugiere que la limitación de recursos es uno de los principales factores que contribuyen al declive actual. En el Capítulo 2, se detectó la segregación alimentaria interespecífica, a través del análisis de partes duras en fecas (2015-2018). Los SAFS se alimentan principalmente de anchoveta (*Engraulis ringens*) y cefalópodos, mientras que los SASL consumen principalmente múnida (*Pleuroncodes monodon*) y anchoveta, en menor proporción. Durante este periodo de estudio, se detectó un solapamiento trófico durante un evento extraordinario de El Niño, lo cual sugiere que los cambios climáticos pueden exacerbar la competencia. En el Capítulo 3, analicé las localizaciones de 35 otáridos (18 SASL, 17 SAFS; 12 hembras, 23 machos) equipados con transmisores satelitales (2013-2017) para determinar sus estrategias de forrajeo. Los machos se segregan en el espacio y en el tiempo; mientras que las hembras muestran un solapamiento espacial pero tienen mecanismos de segregación temporal. Las variables ambientales asociadas sugieren afinidad al hábitat pelágico y costero en las hembras de lobo

fino y lobo chusco, respectivamente. Por su parte, los machos presentan una mayor variabilidad interindividual que debería ser explorada en mayor profundidad. Independientemente, los resultados demuestran cómo las múltiples estrategias de forrajeo a diferentes escalas facilitan la partición de los recursos. En el Capítulo 4, se investigó la diferenciación de nicho a largo plazo examinando las firmas isotópicas archivadas en las vibrisas de SAFS y SASL (2005-2019). Los otáridos incrementaron la amplitud de su nicho (excepto las hembras del SASL) como respuesta al aumento del nivel medio del mar y a las temperaturas más cálidas de la superficie del mar. La disminución general de las señales  $\delta^{15}\text{N}$  sugiere un empobrecimiento de la base de la red trófica, con efectos acumulativos que impactan a los consumidores terciarios. Estos resultados ayudan a explicar el estado actual de declive de las poblaciones, y pueden servir de base para ampliar las recomendaciones de conservación de las poblaciones de otáridos en HCS.

**Palabras clave:** coexistencia; segregación; pinnípedos; ecología trófica; nichos; Sistema de la Corriente de Humboldt

## Publications découlant de ce travail de thèse

### *Article « peer-reviewed » publié*

**Cárdenas-Alayza S**, Gutiérrez D, Tremblay Y (2021) Trends in sympatric otariid populations suggest resource limitations in the Peruvian Humboldt Current System. *Mar Environ Res*:105349. <https://doi.org/10.1016/j.marenvres.2021.105349> (**Chapitre 1**)

### *Accepté pour publication*

Edwards, M.R., **Cárdenas-Alayza S**, Adkesson, M.J., Daniels-Adulahad, M, Hirons, A.C. Peruvian fur seals as Archivists of El Niño Southern Oscillation Effects. Manuscript accepted for publication to *Frontiers in Marine Science* (**Chapitre 4**)

### *En révision*

**Cárdenas-Alayza S**, Torres, D.A., Gutiérrez D, Tremblay Y. Resource partitioning as a mechanism for trophic segregation in sympatric otariids from the productive upwelling Peruvian Humboldt Current System. Manuscript submitted for publication to *Austral Ecology*. (**Chapitre 2**)

**Cárdenas-Alayza S**, Adkesson, M.J., Gutiérrez D, Demarcq, H, Tremblay Y. Multiple strategies for segregation during foraging in sympatric otariids of the Peruvian upwelling Humboldt Current System. Manuscript submitted for publication to *Marine Ecology Progress Series* (**Chapitre 3**)

**Cárdenas-Alayza S**, Adkesson, M.J., Edwards, M.R., Hirons, A.C., Gutiérrez D, Tremblay Y, Franco-Trecu, V. Sympatric otariids increase trophic segregation in response to warming ocean conditions in Peruvian Humboldt Current System. Manuscript submitted for publication to *PLOS ONE* (**Chapitre 4**)

# Table des Matières

Avant-Propos.....	ii
Remerciements .....	iii
Résumé de Thèse.....	vi
Thesis Abstract.....	viii
Resumen de Tesis.....	x
Publications découlant de ce travail de thèse .....	xii
Table des Matières .....	xiii
Index des Illustrations .....	xviii
Index des Tableaux.....	xxii
Résumé exécutif en Français .....	xxiv
<b>1. Introduction</b> .....	xxiv
<b>2. Objectifs de la thèse</b> .....	xxiv
<b>3. Structure de la thèse</b> .....	xxvi
<b>4. Matériel et méthodes</b> .....	xxviii
4.1 Chapitre 1 : Tendances de la population .....	xxviii
4.2 Chapitre 2 : Composition du régime alimentaire.....	xxix
4.3 Chapitre 3 : Stratégies de recherche alimentaire .....	xxxii
4.4 Chapitre 4 : Niche isotopique .....	xxxiii
<b>5. Résultats</b> .....	xxxiv
<b>6. Conclusion et perspectives</b> .....	xxxviii
<b>INTRODUCTION GÉNÉRALE</b> .....	1
<b>1. Introduction</b> .....	2
<b>2. Theoretical Framework</b> .....	5
2.1 Ecological niche .....	5
2.2 Competition and coexistence.....	6
2.3 Sympatry in otariids.....	7
2.4 Interspecific segregation.....	8
2.5 Intraspecific segregation.....	9
2.6 Otariids' foraging .....	10

2.7 Body size and oxygen storage .....	11
2.8 Peruvian Humboldt Current System.....	12
2.9 A warming climate .....	14
2.10 Punta San Juan study site.....	16
<b>3. Thesis Objectives.....</b>	<b>17</b>
<b>4. Structure of the Thesis .....</b>	<b>17</b>
<b>CHAPITRE 1: Trends in sympatric otariid populations suggest resource limitations in the Peruvian Humboldt Current System.....</b>	<b>21</b>
Résume .....	22
Abstract .....	23
<b>1. Introduction .....</b>	<b>24</b>
<b>2. Materials and Methods .....</b>	<b>29</b>
2.1 Study Site and Data Collection.....	29
2.2 Abundance time series.....	30
2.3 Age class proportions .....	31
2.4 Biomass time series .....	32
<b>3. Results .....</b>	<b>33</b>
3.1 Abundance time series.....	33
3.2 Age-class proportions .....	34
3.3 Biomass time series .....	37
<b>4. Discussion.....</b>	<b>39</b>
4.1 Changes in abundance, density and population trajectories .....	39
4.2 Contributing factors of population fluctuations in Peruvian atariids .....	36
4.3 Natural history traits that influence interspecific competition in otariids .....	38
4.4 Current and future trends .....	40
APPENDIX A1 .....	41
APPENDIX A2 .....	42
<b>CHAPITRE 2: Resource partitioning as a mechanism for trophic segregation in sympatric otariids from the productive upwelling Peruvian Humboldt Current System.....</b>	<b>43</b>
Résumé.....	44
Abstract .....	45
<b>1. Introduction .....</b>	<b>46</b>

<b>2. Materials and Methods</b> .....	48
2.1 Study area and sampling .....	48
2.2 Diet composition.....	50
2.3 Trophic indices .....	51
2.4 Environmental conditions .....	52
2.5 Statistical analyses .....	52
<b>3. Results</b> .....	53
3.1 Study area and sampling .....	53
3.2 Diet composition.....	54
3.3 Trophic indices .....	56
3.4 Clustering.....	61
3.5 Environmental conditions and trophic indices .....	65
<b>4. Discussion</b> .....	65
4.1 Resource partitioning and trophic overlap.....	66
4.2 Empty scats.....	67
4.3 Previous research .....	68
4.4 General limitations in methodology .....	70
4.5 Environmental conditions and trophic indices .....	71
<b>APPENDIX B1</b> .....	73
<b>APPENDIX B2</b> .....	75
<b>APPENDIX B3</b> .....	76
<b>APPENDIX B4</b> .....	77
<b>CHAPITRE 3: Multiple strategies for segregation during foraging in sympatric otariids of the Peruvian upwelling Humboldt Current System</b> .....	78
<b>Résumé</b> .....	79
<b>Abstract</b> .....	80
<b>1. Introduction</b> .....	81
<b>2. Materials and Methods</b> .....	83
2.1 Data collection.....	83
2.2 Track construction .....	85
2.3 Foraging trips and events.....	85
<b>3. Utilization distributions</b> .....	86



4. Foraging by hour of day .....	86
5. Associated environmental conditions .....	86
6. Results .....	89
6.1 Body mass.....	89
6.2 Trip distance and duration .....	91
6.3 Utilization distributions .....	92
6.4 Foraging by hour of day .....	93
6.5 Associated environmental conditions .....	97
7. Discussion.....	100
7.1 Foraging tips and utilization distributions .....	100
7.2 Foraging by hour of day .....	102
7.3 Associated environmental conditions .....	103
APPENDIX C1 .....	105
APPENDIX C2 .....	106
APPENDIX C3 .....	107
APPENDIX C4 .....	108
APPENDIX C5 .....	109
APPENDIX C6 .....	110
CHAPITRE 4: Sympatric otariids increase trophic segregation in response to warming ocean conditions in Peruvian Humboldt Current System.....	111
Résumé.....	112
Abstract .....	113
1. Introduction .....	114
2. Materials and Methods .....	117
2.1 Study site and sample collection .....	117
2.2 Sample analysis .....	117
2.3 Data analysis.....	118
2.4 Environment: sea level and sea temperature .....	118
2.5 Study groups.....	119
2.6 Isotopic signatures .....	119
2.7 Influence of the environment, species and sex on isotopic composition.....	119

3. Isotopic niche.....	120
4. Results .....	120
4.1 Study groups .....	120
4.2 Influence of environment on isotopic composition .....	122
4.3 Isotopic niche size, overlap and community metrics according to period.....	126
5. Discussion.....	129
APPENDIX D1 .....	134
APPENDIX D2 .....	135
DISCUSSION GÉNÉRALE .....	136
1. Interspecific and intraspecific segregation of the trophic niche (diet composition and isotopic signals).....	137
2. Foraging strategies (spatial and temporal dimensions) .....	139
3. Environmental conditions associated to foraging.....	142
4. Final remarks .....	144
5. Future Recommendations .....	147
Références Bibliographiques .....	149
Références Iconographiques .....	168

# Index des Illustrations

**Figure I. 1** Diagram showing resource partitioning among 11 species of anole lizards found on the island of Puerto Rico. Each species occupies a different type or elevation of vegetation. The habitat is further partitioned by the amount of sunlight and moisture available. Image by Eva Horne modified from (Williams, 1983). Source: <https://www.khanacademy.org/science/ap-biology/ecology-ap/community-ecology/a/niches-competition> ..... 3

**Figure I. 2** Competitive exclusion example showing *Paramecium caudatum* grow well individually, but when they compete for the same resources, *P. aurelia* outcompetes *P. caudatum*. Image from: Fowler et al., 2013. Source: <https://openstax.org/books/concepts-biology/pages/19-4-community-ecology>..... 6

**Figure I. 3** Schematic representation of Lotka-Volterra competition. Image from D’Andrea et al., 2019 ..... 7

**Figure I. 4** Morphology of seafloor along the coast of Peru. Punta San Juan study area located in box C, region with a very narrow continental shelf. Image from IMARPE, 2014a..... 13

**Figure I. 5** A) Example of the distribution of the shallow oxygen minimum zone in the Southeast Pacific Ocean. Source: <http://omz.udec.cl> ; B) Example of the oxycline delimiting the vertical distribution of plankton and anchovy schools in relation to sea bottom in Peru. Image from IRD /Marbec. Source: <https://books.openedition.org/irdeditions/docannexe/image/34324/img-5.jpg>..... 14

**Figure I. 6** Location of Punta San Juan peninsula (left) on the south coast of Peru, South America (right). Image modified from S. Cárdenas-Alayza, 2012. .... 17

**Figure I. 7** Study species and groups included in this study: South American fur seal (*Arctocephalus australis*) A) adult female with pup; B) territorial male and South American sea lion (*Otaria byronia*) C) adult female with pup; D) sub-adult males. Images provided by Punta San Juan Program, CSA-UPCH..... 18

**Figure 1. 1** Time series of the total number South American sea lions (SASL, open circles) and South American fur seals (SAFS, filled diamonds) counted on the coast of Peru (top) and Punta San Juan study site (bottom) between 1965-2019. Dashed grey vertical lines delimit the population decline caused by the 1997-98 ENSO. Time series were constructed with official counts from surveys obtained from various sources (Arias-Schreiber & Rivas, 1998; IMARPE, 2012, 2013, 2014b, 2015, 2016, 2017, 2018; Majluf & Trillmich, 1981; Muck & Fuentes, 1987; SERNANP, 2016; Tovar & Fuentes, 1984). .. 28

**Figure 1. 2** Top panel shows 3-day interpolated counts (black line) with trend of abundance (red line) for A) SAFS and B) SASL collected in Punta San Juan, Peru. Bottom panel shows wavelet spectrum with horizontal line indicating the annual signal at 365 days for C) SAFS and D) SASL abundance time series. .... 35

**Figure 1. 3.** Boxplots of the weighted proportion (product of mean mass and number of individuals) of each age-class for SAFS and SASL during breeding season (BS, white) and non-breeding season (non BS, grey) during March 2012 - December 2019. .... 36

**Figure 1. 4** Time series of the proportion of age-classes for SAFS (green) and SASL (orange) constructed from weekly counts (circles) with smooth curve (line) with 95% CI (grey area). Age-class counts collected at Punta San Juan otariid breeding sites between November 2003 - December 2019..... 38

**Figure 1. 5** Stacked area chart of constructed biomass time series in tons (t) estimated from otariid abundance in Punta San Juan for SAFS (light gray) and SASL (dark grey) during years 2001-2019..... 39

**Figure A. 1** Multi-panel plot of South American fur seals counts by age-class according to day of year collected at the main breeding site S3 in Punta San Juan, Peru. Classes include: Adult Females, Juveniles, Pups, Subadult Males and Territorial Males. Total is the sum of all the age-classes. Blue line is a fitted loess model to depict the trend in abundance throughout the year. Data points include all study years available for site. .... 41

**Figure A. 2** Multi-panel plot of South American sea lions counts by age-class according to day of year collected at the main breeding site N7 in Punta San Juan, Peru. Classes include: Adult Females, Juveniles, Pups, Subadult Males and Territorial Males. Total is the sum of all the age-classes. Blue line is a fitted loess model to depict the trend in abundance throughout the year. Data points include all study years available for site. .... 42

**Figure 2. 1** Bar graphs of A) percent number (%) of taxa consumed by each species during the study period and B) stacked % IIMP values for each prey item over the entire study period for South American fur seal (SAFS, green) and South American sea lion (SASL, blue). Prey items (N = 22) detected are listed organized in alphabetical order. Values for IIMP >1% are annotated in each otariid species color. .... 57

**Figure 2. 2** Stacked bar graphs of A) Percent number (%) by taxa; B) Frequency of occurrence (%) by prey item. South American fur seals (SAFS, top panels) and South American sea lions (SASL, bottom panels) for each year (2015-2018) and season (sum = summer, fal = fall, win = winter, spr = spring). ... 59

**Figure 2. 3** Stacked bar graphs of C) IIMP >1% by prey items and D) IIMP (%) by ecological groups South American fur seals (SAFS, top panels) and South American sea lions (SASL, bottom panels) for each year (2015-2018) and season (sum = summer, fal = fall, win = winter, spr = spring)...... 60

**Figure 2. 4** Dendrogram using agglomerative hierarchical clustering with Ward's method and Bray-Curtis distances, showing two separate clusters for South American sea lion (SASL, blue) and South American fur seal (SAFS, green). Two SASL seasons (SASL-2015-fal and SASL-2015-sum) are inside the SAFS cluster (red asterisks)..... 62

**Figure 2. 5** NMDS spider plot (stress = 0.08) showing two separate clusters for each otariid species. Each node represents a sampling season for South American sea lion (SASL, blue) and South American fur seal (SAFS, green). Relevant prey items for principal dimensions (axis 1 and axis 2), overlaid in dark red. .... 63

**Figure 2. 6** Ranked abundance curves for prey items of South American sea lion (SASL, blue) and South American fur seal (SAFS, green) samples during A) seasons with trophic overlap (summer 2015 and fall 2015) and B) seasons without trophic overlap. Labels shown for top-five ranking prey items.... 64

**Figure B1. 1** Boxplots of abundance of South American sea lions (SASL) in site N7 (2015-2018) according to age class during breeding (BS, January -March) and non breeding months (non BS, April – December) according to age class. .... 73

**Figure B1. 2** Boxplots of abundance of South American fur seals (SAFS) in site S3 (2015-2018) according to age class according to age class during breeding (BS, October - December) and non breeding months (non BS, January – September). .... 74

<b>Figure B3. 1</b> Boxplots of the seasonal A) SST and B) SSTA for study period 2015-2018 collected in San Juan de Marcona, source: DHN - Peru. ....	76
<b>Figure B4. 1</b> Spearman correlation charts ( $p < 0.05$ ) for trophic and environmental indices for A) South American sea lions, SASL and B) South American fur seals, SAFS. Positive correlation scores in shades of blue and negative in shades of red. ....	77
<b>Figure 3. 1</b> Tracks for A) SAFS and B) SASL in the study area. SAFS females (orange), SASL females (aquamarine); SAFS males (brown) and SASL males (teal). Yellow circle indicates the Punta San Juan, Peru breeding colony, location of otariid instrumentation. ....	88
<b>Figure 3. 2</b> Boxplot of body mass (kg) in y-axis according to species and sex groups in x-axis (ANOVA, $p < 0.05$ ). SAFS females (orange), SASL females (aquamarine); SAFS males (brown) and SASL males (teal). ....	90
<b>Figure 3. 3</b> Boxplots of A) distance (km) and B) duration (days) of foraging trips according to sex (x-axis). Colors represent species: SAFS (orange) and SASL (teal). ....	91
<b>Figure 3. 4</b> Maps with filled contours of Utilization Distributions (UD) for 50% core foraging range (dotted line) and 95% home range (continuous line) for A) SAFS females (orange) and SASL females (aquamarine); B) SAFS males (brown) and SASL males (teal); C) SAFS females (orange) and SAFS males (brown); D) SASL females (aquamarine) and SASL males (teal). Black lines represent coastline and isobaths for depths of 2,000, 4,000 and 6,000 meters. ....	95
<b>Figure 3. 5</b> Density plots of the foraging events (y-axis) by hour of day (x-axis) for A) SAFS females (orange) and SASL females (aquamarine); B) SAFS males (brown) and SASL males (teal); C) SAFS females (orange) and SAFS males (brown); D) SASL females (aquamarine) and SASL males (teal). Shaded areas indicate dark hours (0 - 5, 18 - 23 hours) in study area. ....	96
<b>Figure 3. 6</b> Probability of foraging events (y-axis 1 and 2) in relation to ranges of environmental variables (x-axis 1 and 2) for A) SAFS females (orange) and SASL females (aquamarine) and B) SAFS males (brown) and SASL males (teal). Colors of labels on x- and y-axis are indicative of values for each species and sex group. ....	99
<b>Figure C2. 1</b> Smoother response for probability of foraging (y-axis) according to hour of day (x-axis) for A) SAFS females, B) SAFS males, C) SASL females and D) SASL males from General Additive Mixed-Models. Significant relationships ( $p < 0.01$ ) were found for A) SAFS females and B) SASL males. Icons represent daylight (sun) and dark hours (moon). ....	106
<b>Figure C4. 1</b> Environmental conditions (SST, Chl-a, Front Proximity, Front Gradient and DC) associated to female SAFS (orange) and SASL (aquamarine) foraging events. A) Boxplots depicting significant differences (***) indicates $p < 0.01$ . B) Density curves show distribution of frequencies. SST between SAFS and SASL is not significantly different (N.S.).....	108
<b>Figure C4. 2</b> Environmental conditions (SST, Chl-a, Front Proximity, Front Gradient and DC) associated to male SAFS (brown) and SASL (teal) foraging events. A) Boxplots depicting significant differences (***) indicates $p < 0.01$ . B) Density curves showing distribution of frequencies.....	108

**Figure C6. 1** Correlation chart (Spearman rho,  $p=0.05$ ) between the environmental covariates (SST, Chl-a, Front Proximity, Front Gradient, Distance to Coast) for A) SAFS females and B) SASL females. SST not included in for SAFS females due to 39% missing values..... 110

**Figure C6. 2** Correlation chart (Spearman rho,  $p=0.05$ ) between the environmental covariates (SST, Chl-a, Front Proximity, Front Gradient, Distance to Coast) for A) SAFS males and B) SASL males .... 110

**Figure 4. 1** Principal components analysis of six morphometric measures (skull girth, axillary girth, snout to tail base, brachial girth, axilla to fore flipper, tail base to hind flipper) for each of the groups. SAFS females (pink,  $N = 35$ ), SASL females (green,  $N= 22$ ), SAFS males (blue,  $N=12$ ) and SASL males (orange,  $N=29$ ) ..... 121

**Figure 4. 2** Time series for A) monthly sea level records from San Juan de Marcona bay, horizontal lines show Period 1 (P1, mean = 3.42 cm) and Period 2 (P2, mean = 8.06 cm), B)  $\delta^{13}C$  and C)  $\delta^{15}N$  isotopic signatures for all otariid groups. .... 124

**Figure 4. 3** Standard Ellipse Area (75%) for A) Period 1 and B) Period 2. Black lines delineate community convex hull between centroids of each group. .... 127

**Figure D1. 1** Standard Ellipse Area estimated for each group. Labels represent Community (Period 1 and 2) and Group (1 = SAFS females, 2 = SAFS males, 3 = SASL females, 4 = SASL males). The black points correspond to the mean standard ellipse area for each group, red cross is the standard ellipse area corrected for small sample size. Grey and white boxed areas reflect the 95, 75 and 50% confidence intervals..... 134

**Figure D2. 1** Total area of convex hull estimated for each community or period. Community 1 = Period 1, Community 2 = Period 2. The black points correspond to the mean standard ellipse area for each group while the grey and white boxed areas reflect the 95, 75 and 50% confidence intervals..... 135

**Figure D. 1** Diagram summarizing main findings to explain current population decline in Peruvian otariids in context of two contrasting environmental scenarios: cool (blue) and warm (red) conditions. 144

# Index des Tableaux

<b>Table 1. 1.</b> Median proportion of age-class per species with inter-quartile range (IQR). Mean mass and standard deviation ( $\pm$ SD) in kilograms. Sample size and source used for each age-class. ....	36
<b>Table 2. 1</b> Number of collected and proportion of scat samples with content (in brackets) for South American fur seal (SAFS) and South American sea lion (SASL) per year (2015 - 2018), season (sum = summer, fal = fall, win = winter and spr = spring) and total.....	54
<b>Table 2. 2</b> Species Richness (S), Levin’s Trophic Niche Breadth ( $B_i \pm 95\%$ CI) and Trophic Level (TL) for South American fur seal (SAFS) and South American sea lion (SASL) per year (2015 - 2018), season (sum = summer, fal = fall, win = winter and spr = spring) and Mean $\pm$ SD for the study period. .	58
<b>Table B2. 1</b> List of prey items identified in the samples of South American fur seals (SAFS) and South American sea lions (SASL) and the habitat group category assigned.....	75
<b>Table 3. 1</b> Number of individuals (N ind), number of trips (N trips), trip duration (days), and trip distance (km) for each species-sex group included in this study. Results reported as mean $\pm$ SD. ....	92
<b>Table 3. 2</b> Area in km <sup>2</sup> of 25%, 50%, 75% and 95% Utilization Distributions (UD) for each species and sex group.....	92
<b>Table 3. 3</b> Proportion overlap for each group for core area (50% UD) and home range (95 % UD). Values represent the proportion overlap for the group in the rows by the group in the column. Proportion overlap $> 0.70$ in bold.....	93
<b>Table 3. 4</b> Results from the General Additive Mixed-Models to test for effect of Hour in the probability of foraging events in SAFS females, SASL females, SAFS males and SASL males. N represents the number of observations per group. ....	93
<b>Table 3. 5</b> Results from the forward selection process of covariates that explain environment during foraging using a General Additive Mixed-Models. Bold indicates best fitting model and N represents the number of observations per group. Significant Chi square values have a (*) for p-values $< 0.01$ and (.) for p-values $< 0.05$ . SST was not included in the SAFS females models due to missing values in 39% of foraging locations.....	98
<b>Table C1. 1</b> Species, sex, individual identification (ID) deployment date, weight (kg), length (cm), number of days tracked and number of trips recorded for each individual. ....	105
<b>Table C3. 1</b> Descriptive statistics for the environmental covariates extracted for foraging event locations for each species and sex group. ....	107
<b>Table C5. 1</b> Deviance explained by univariate General Additive Models (GAM) of each environmental covariate explored in response to foraging events. N = number of observations included in each group. ....	109



**Table 4. 1** Isotopic signatures of Peruvian otariid whiskers according to periods investigated in present study. Number of whiskers (N), mean length (mm) by species and sex. Mean  $\pm$  SD of the stable isotope values ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) during two periods according to sea level. N fragments: sample size for species and sex. .... 123

**Table 4. 2** Linear Mixed-Effects Models (LMM) for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values. LMM for  $\delta^{13}\text{C}_{\text{cor}}$  and  $\delta^{15}\text{N}_{\text{cor}}$  whisker values including group (sex-species), period and their interactions as fixed effects and identity and whisker portion as random effect (continuous autocorrelation function). Estimates and p-values (in brackets) are shown for each variable. In bold we show the selected model by the Akaike Information Criterion (AIC). (\*\*\*) denotes  $p < 0.01$  ..... 125

**Table 4. 3** Corrected Standard Ellipse Area (SEAc) for each period and group estimated with 75% maximum likelihood. .... 126

**Table 4. 4** Percent overlap between Peruvian otariid groups according to period. Values indicate the percent overlap for the group in the row. .... 128

**Table 4. 5** Layman metrics estimated according to period.  $\delta^{13}\text{C}$  range = distance range of  $\delta^{13}\text{C}$  values,  $\delta^{15}\text{N}$  range = distance range of  $\delta^{15}\text{N}$  values, TA = Total area of convex hull, CD = the mean distance to centroid from means, MNND = mean nearest neighbor distance of the means, SDNND = standard deviation of the nearest neighbor distance. .... 128



# Résumé exécutif en Français

## 1. Introduction

La compréhension de la coexistence d'espèces sympatriques ayant des besoins similaires est un thème important en écologie. La compétition est un processus écologique influencé par la sélection naturelle, qui façonne les espèces dans le temps présent, en fonction des conditions de leur environnement écologique. La compétition peut compromettre l'adaptation des espèces concernées et peut finalement conduire à « l'exclusion compétitive » d'une espèce (Pacala & Roughgarden, 1985). La façon dont les espèces des communautés écologiques se partagent les ressources disponibles est un facteur déterminant de la diversité des espèces. Une communauté où le partage des ressources ou le chevauchement des niches est plus important abritera plus d'espèces qu'une communauté où le chevauchement des niches est moindre. Ainsi, les espèces coexistantes doivent différer dans leurs exigences écologiques d'au moins une quantité minimale pour éviter l'exclusion compétitive (Pianka, 1974).

Cette étude porte sur deux populations sympatriques de prédateurs supérieurs d'otariidés : l'otarie à fourrure d'Amérique du Sud (SAFS, *Arctocephalus australis*) et les lions de mer d'Amérique du Sud (SASL, *Otaria byronia*), qui semblent avoir des besoins trophiques similaires. Après des déclin de population causés par une pénurie alimentaire extrême due à l'oscillation El Niño (ENSO) de 1997-98, les otaries d'Amérique du Sud au Pérou ont augmenté régulièrement, avec une population estimée à plus de 130 000 individus (IMARPE, 2014b). En comparaison, les SAFS au Pérou n'ont pas été en mesure de se rétablir dans la même mesure, atteignant à peine une population d'environ 11 000 individus (IMARPE, 2014b). Cette étude utilise des données recueillies via l'utilisation de marques électroniques, via des comptages et via des échantillons de fécès et de vibrisses collectés à Punta San Juan (PSJ, Latitude : 15°22') au sud du Pérou, où ces deux espèces partagent leur site de reproduction. A PSJ, les populations reproductrices se concentrent jusqu'à 5.000 SAFS et 12.000 SASL pendant la période d'étude de cette thèse. Au Pérou, ces espèces présentent une ségrégation temporelle dans leurs saisons de reproduction, qui s'étendent sur 1-3 mois. L'allochronie, qui est reconnue comme un moteur majeur de la spéciation, peut être considérée comme facilitant la partition des ressources pendant les périodes énergétiquement critiques (Taylor & Friesen, 2017). Au Pérou, les SAFS se reproduisent d'octobre à décembre et les SASL de janvier à mars. Cependant, l'allaitement de la

progéniture peut être prolongée par rapport à d'autres espèces d'otariidés, et le moment du sevrage est flexible, allant de 6 à 36 mois, avec une grande variabilité interannuelle (Majluf, 1987; Soto, 2004). Ceci génère un chevauchement temporel dans la période de forte demande énergétique pendant laquelle les femelles doivent trouver de la nourriture pour assurer la survie de leurs jeunes.

Pour comprendre le niveau de compétition entre ces deux espèces, il est nécessaire de mesurer le chevauchement et/ou la ségrégation entre les niches trophiques ( la différenciation de niche). Ceci ne sera possible qu'en déterminant les mécanismes d'exploitation et de partition des ressources des espèces dans leur environnement. Lorsque les espèces vivent dans le même habitat, elles peuvent se ségréger de différentes façons : séparation spatiale (par exemple, en se nourrissant à différentes profondeurs ou dans différents lieux géographiques) ; séparation temporelle des ressources (par exemple, en utilisant des ressources disponibles à différents moments de la journée ou à différentes saisons) ou ségrégation alimentaire (par exemple, en consommant différents types de proies) (Begon et al., 2006). Dans ce contexte, il semble évident que les conditions environnementales sont cruciales à prendre en compte car elles influencent tous ces paramètres, pouvant ainsi contraindre les différentes trajectoires évolutives des espèces en co-habitation.

Une caractéristique importante influençant le mode de recherche de nourriture chez les otariidés peut être la productivité marine locale et son effet sur la disponibilité des proies. Il se peut que, sauf dans les régions très productives, une petite taille favorise la recherche de proies très disparates mais denses près de la surface (zooplancton, petits poissons ou calmars), alors que les mammifères à respiration aérienne, de plus grande taille, sont mieux adaptés à la recherche de proies plus uniformément réparties mais moins denses tel que dans le benthos (gros poissons ou poulpes). Cela coïncide avec la tendance à la consommation d'une plus grande masse de proies par les otaries benthiques en quête de nourriture et à la proportion plus importante de temps passé en mer à plonger par rapport aux otaries à fourrure épipelagiques plus petites (Arnould & Costa, 2006). Des études plus récentes ont montré que ce schéma n'est pas toujours la norme, et que certaines populations d'otaries vivant dans des zones d'upwelling présentent une stratégie alimentaire pélagique (Hückstädt et al., 2014), que des otaries à fourrure sont des plongeurs benthiques (Arnould & Hindell, 2001) et que certaines espèces présentent une combinaison de plongées peu profondes et profondes en fonction des besoins métaboliques et de la disponibilité locale des proies (Harcourt et al., 2002; Villegas-Amtmann et al., 2008). Ceci démontre la plasticité des individus aux conditions spatiales et temporelles locales, et

qui peuvent également varier selon le sexe et l'âge (Weise & Costa, 2007).

Ainsi, le principal objectif de cette étude est de décrire les niches écologiques de espèces d'otariidés sympatriques afin de déterminer le degré de chevauchement des niches intraspécifiques et interspécifiques dans l'environnement marin du système du Courant de Humboldt (HCS). Il est particulièrement important de comprendre les mécanismes de partitionnement qui permettent la coexistence des populations de prédateurs. Les prédateurs affectent généralement les écosystèmes par le biais d'un contrôle top-down et impactent ainsi les populations d'espèces proies. Ainsi, le maintien des populations de prédateurs est essentiel pour préserver la « santé des écosystèmes » (abondance, diversité). Des évaluations précises des stratégies de recherche alimentaire sont essentielles pour comprendre la fonction des prédateurs et la structure des écosystèmes. Les écosystèmes et les réseaux trophiques étant très dynamiques, le couplage des informations traditionnelles sur le régime alimentaire (par exemple, les parties dures identifiées dans les fécès) avec les données sur la localisation de la recherche de nourriture peut ajouter une dimension spatiale et temporelle aux estimations du régime alimentaire (Jeanniard du Dot et al., 2017). De plus, de longues séries temporelles de comptages de population et de signatures isotopiques des vibrisses fourniront une fenêtre temporelle différente pour explorer les habitudes trophiques des SAFS et des SASL et leur conséquences sur les populations. Pour ce faire, l'étude actuelle propose de combiner une série de jeux de données pour décrire et comprendre l'écologie trophique des deux espèces à différentes échelles temporelles et spatiales afin d'expliquer les tendances divergentes actuelles des populations d' otariidés sympatriques à Punta San Juan.

## 2. Objectifs de la thèse

L'objectif principal de cette thèse est de décrire et de comparer les niches écologiques des espèces d'otariidés sympatriques *Arctocephalus australis* et *Otaria byronia* qui coexistent à Punta San Juan ; pour déterminer la différenciation/le chevauchement des niches intraspécifiques et interspécifiques. Trois objectifs spécifiques permettront d'atteindre ces objectifs. Le premier est de décrire et de comparer l'écologie trophique des pinnipèdes sympatriques en fonction de l'espèce et du sexe. Le deuxième est de décrire et de comparer les variables environnementales qui caractérisent les niches écologiques selon l'espèce et le sexe. Enfin, le troisième objectif est de décrire et de comparer les stratégies de recherche de nourriture (dimensions spatiales et temporelles) selon l'espèce et le sexe.

### 3. Structure de la thèse

Pour mettre en contexte l'état actuel des population des espèces étudiées, l'abondance et la biomasse des espèces sont analysées pour comprendre la compétition interspécifique potentielle à PSJ. Pour aborder l'écologie trophique, le couplage des informations traditionnelles sur le régime alimentaire (par exemple, les parties dures récupérées dans les fèces), les données sur les lieux de recherche de nourriture et les informations sur les isotopes stables sont explorés dans le but de mieux comprendre l'utilisation de l'environnement marin par les mâles et les femelles de la SAFS et de la SASL, à différentes résolutions. Dans le cadre de chaque analyse, des descriptions de l'environnement sont incluses qui aident à caractériser les conditions pour chaque ensemble de données trophiques. En somme, l'étude actuelle propose de combiner une série de jeux de données, divisés en quatre chapitres, afin de faire la lumière sur les stratégies de recherche alimentaire des deux espèces et des deux sexes à différentes échelles temporelles et spatiales pour expliquer les tendances actuelles des populations.

Dans le **chapitre 1**, j'ai analysé les trajectoires des populations de SASL et de SAFS dans le site de reproduction sympatrique de PSJ, au Pérou, entre 2001-2019 et discuté des mécanismes de compétition et/ou de limitation des ressources.

Dans le **chapitre 2**, j'ai comparé la composition saisonnière du régime alimentaire de SAFS et SASL, sur 15 saisons entre 2015 et 2018 (qui comprenaient des conditions ENSO et non-ENSO) pour évaluer si le partage des ressources existe dans les populations d'otariidés sympatriques à PSJ .

Dans le **chapitre 3**, j'ai analysé les comportements en mer de 18 SASL et 17 SAFS (12 femelles, 23 mâles) équipés de balises satellites à PSJ, au Pérou, au cours de 2013-2017. Je détermine si la ségrégation temporelle et spatiale entre les espèces et les groupes de sexe est influencée par les conditions morphologiques (masse corporelle), comportementales ou environnementales pendant la recherche de nourriture.

Dans le **chapitre 4**, j'ai évalué l'effet des événements de réchauffement marin sur la ségrégation de niche inter- et intra-spécifique à long terme. Pour cela, j'ai analysé les signatures isotopiques sur les vibrisses collectées dans PSJ de SAFS (55 femelles et 21 mâles) et SASL (14 femelles et 22 mâles) pour déterminer la ségrégation de niche isotopique interspécifique sur la base des valeurs  $\delta^{13}\text{C}$  et  $\delta^{15}\text{N}$

organisées à une échelle mensuelle pour 2005 - 2019.

## **4. Matériel et méthodes**

### ***4.1 Chapitre 1 : Tendances de la population***

#### ***4.1.1 Échantillonnage***

Le littoral de PSJ comprend 20 sites, qui sont des plages et des criques toutes utilisées pour la reproduction et/ou comme sites de repos par SASL et SAFS. Un programme de recherche à long terme (Punta San Juan Program) qui surveille les populations d'otariidés in-situ a été mis en place. L'accès à la base de données du programme PSJ a été accordé pour les besoins de cette étude. A PSJ, des comptages terrestres des sites accessibles d'otariidés sont effectués au moins une fois par semaine depuis 2001 pour suivre la dynamique des populations. Pour cette étude, nous avons accédé aux données de comptage hebdomadaires pendant les années 2001 à 2019. Les comptages ont eu lieu entre 6h00 et 9h00 par des observateurs équipés de jumelles (10x50) et de compteurs à main au sommet de falaises de 8 à 30 mètres surplombant les plages. Les sites accessibles sont comptés pour l'abondance totale et les catégories de classe d'âge, pour chaque espèce. Des séries chronologiques de l'abondance des populations de SAFS et SASL à PSJ et aux sites d'étude seront préparées afin de décrire la dynamique des populations de chaque espèce tout au long de la période d'étude. L'analyse des trajectoires de population sera effectuée à l'aide de techniques de décomposition en ondelette afin de refléter les tendances actuelles de l'abondance des espèces au fil du temps.

#### ***4.1.2 Analyse des données***

Pour extraire la tendance à long terme dans les séries temporelles, nous avons utilisé une décomposition en ondelettes avec une ondelette mère "morlet", car elle permet de surmonter les problèmes de non-stationnarité dans une série temporelle en effectuant une décomposition locale du signal à l'échelle temporelle. Cette méthode donne la possibilité d'étudier l'évolution temporelle avec différentes composantes rythmiques. J'ai identifié l'échelle de longueur d'onde correspondant à une année en utilisant l'analyse des ondelettes dans le paquet R WaveletComp v. 1.1 pour extraire la tendance de l'abondance pour les deux espèces. La tendance des séries chronologiques d'abondance, la densité et les ratios d'abondance ont été comparés entre les deux espèces. Les phases

d'augmentation ou de diminution des tendances ont été déterminées à l'aide de dérivées de premier ordre. Les points de retournement permettent de détecter la date du changement de direction de la trajectoire de la population d'une espèce. La différence des dérivées de premier ordre a été utilisée pour comprendre quelle espèce était en augmentation par rapport à l'autre.

## 4.2 *Chapitre 2 : Composition du régime alimentaire*

### 4.2.1 *Échantillonnage*

Grâce à la surveillance à long terme et à la recherche sur place à PSJ, nous avons pu collecter des échantillons de 2015 à 2018, qui comprenaient des conditions ENSO et non-ENSO. Au total, 3 325 échantillons de fèces (1 484 SAFS et 1 841 SASL) ont été collectés. Les échantillons collectés ont été trempés dans de l'eau savonneuse pendant 3 à 24 heures afin de faciliter la dissolution des matières fécales. Les durées variaient en fonction de la concentration de matière organique dans les échantillons. Le contenu a été passé à travers des tamis emboîtés de 1,00 mm et 0,297 mm de maille. Les restes durs comprenaient des otolithes, des becs de céphalopodes et des restes de crustacés, qui ont tous été soigneusement retirés. Les otolithes et les becs de céphalopodes qui étaient érodés à plus de 50 % ont été jetés, en suivant Tollit et al., 1997. Les otolithes sagittaux ont été identifiés au niveau des espèces en suivant García-Godos (2001). Les becs de céphalopodes ont été identifiés comme un seul groupe taxonomique ("Céphalopodes"). Pour les crustacés, une seule espèce (*Pleuoncodes monodon*) a été identifiée et comptée (en utilisant le céphalothorax). L'importance des proies a été évaluée en utilisant le pourcentage du nombre (*PN*), le pourcentage de la fréquence d'occurrence (*FO*) et le pourcentage de l'indice d'importance (*IIMP*). La richesse en espèces (*S*) a été estimée pour chaque saison d'échantillonnage en regroupant toutes les années d'une saison et pour chaque saison et année indépendante. Je pars du principe que chaque échantillon fécal représente un individu de la colonie, et donc que l'échantillonnage des excréments représente un échantillon aléatoire de la population. Afin de déterminer si les individus d'une saison d'échantillonnage donnée présentent différents degrés d'expertise en matière d'alimentation, l'indice standardisé Levin  $B_i$  (Krebs, 1999) a été estimé pour distinguer si chaque espèce est un consommateur spécialiste ou généraliste. Le niveau trophique (*TL*) a été calculé pour chaque espèce et chaque saison. Les enregistrements saisonniers mensuels de la température de surface de la mer (SST) ont été collectés à moins de 5 km des colonies d'otariidés, et ont été utilisés pour estimer les séries chronologiques des anomalies (SSTA) afin de

caractériser les conditions environnementales pendant la période d'étude.

#### 4.2.2 *Analyses des données*

Pour déterminer les différences d'association de la composition des proies consommées par saison et par espèce, des matrices avec les espèces (colonnes) et les saisons (lignes) utilisant les valeurs IIMP ont été construites. Le regroupement hiérarchique agglomérant a été estimé à l'aide de la méthode de Ward (Ward & Hook, 1963) et les valeurs  $p$  pour chaque groupe ont été calculées avec un rééchantillonnage par bootstrap (1 000 itérations) à l'aide du paquet `vegan` du logiciel statistique R. Pour interpréter la position des saisons d'échantillonnage et des éléments de proie entre les deux espèces dans un nombre réduit de dimensions, nous avons utilisé l'échelle multidimensionnelle non métrique (NMDS,  $\text{stress} < 0,5$ ) sur les matrices standardisées. Pour tester les hypothèses, la normalité a été évaluée avec le test de Shapiro-Wilk ( $p < 0,05$ ) et l'homogénéité de la variance a été vérifiée avec le test de Levene ( $p < 0,05$ ). Les différences entre les groupes ont été évaluées par ANOVA pour les données paramétriques et par Kruskal-Wallis pour les données non paramétriques ( $p < 0,05$ ). Les tests post-hoc Tukey et Chi-carré ont été utilisés pour les comparaisons multiples par paire ( $p < 0,05$ ) pour les données paramétriques et non-paramétriques, respectivement.

Afin de déterminer s'il existait une corrélation avec les conditions environnementales, la proportion de fèces vides, la PN par taxa (poissons, céphalopodes et crustacés), la richesse spécifique ( $S$ ), la largeur trophique ( $Bi$  de Levin) et le niveau trophique ( $TL$ ) estimés pour SAFS et SASL ont été corrélés avec les métriques SST et SSTA à l'aide d'une matrice de corrélation (rang de Spearman,  $r_s$ ,  $p < 0,05$ ) avec les packages `rstatix` et `corrplot` du logiciel statistique R (R Core Team, 2018).

## 4.3 *Chapitre 3 : Stratégies de recherche alimentaire*

### 4.3.1 *Échantillonnage*

Des balises satellites ont été déployées sur 35 adultes ou sub-adultes entre 2013-2017 dans PSJ. Dix-sept SAFS (8 femelles et 9 mâles) et dix-huit SASL (4 femelles et 14 mâles) ont été capturés et instrumentés dans le cadre de cette étude. Les femelles adultes allaitaient des nouveau-nés avant le marquage et sont donc estimées âgées de 4 ans et plus. Alors que les mâles SASL ont été catégorisés sur le terrain comme des mâles sub-adultes, qui sont des individus matures sur le plan reproductif, mais qui ne contrôlent pas encore les harems de femelles et dont l'âge est estimé entre 5 et 8 ans. Les mâles adultes des SAFS ont été sélectionnés parmi les mâles en possession de territoires et ont donc un âge estimé de 9+ ans.

### 4.3.2 *Analyse des données*

Les localisations possibles ont été modélisées à l'aide d'une méthode de marche aléatoire corrélée, selon Tremblay et al. (2009). Une trajectoire moyenne et sa distribution d'erreur associée ont été calculées pour décrire la trajectoire de chaque individu. Les positions terrestres ont été filtrées en utilisant une combinaison de 'buffer' circulaires de 5 km sur les colonies de pinnipèdes connues le long de la côte du Pérou et un seuil temporel basé sur des résultats précédents (Ganoza, 2016; Hückstädt et al., 2014). La distance totale (km) et la durée en mer (jours) ont été estimées pour chaque voyage. Les résultats sont rapportés en moyenne  $\pm$  SD obtenus par chaque espèce et groupe de sexe. Les différences interspécifiques et intraspécifiques ont été testées pour chacune de ces variables (Kruskal Wallis,  $p < 0,05$ ).

J'ai estimé le temps de résidence (RT) en tant qu'indice de recherche de nourriture pour comprendre le temps passé (en heures) pour la recherche en zones restreintes par les individus dans des zones de rayon "r" (en km) et le temps maximum (en heures) pendant lequel ils ont activement exploré ou recherché de la nourriture, en suivant Barraquand et Benhamou (2008). Pour déterminer la meilleure échelle, plusieurs rayons de taille et seuils de temps ont été testés pour chaque individu. Les rayons testés étaient de l'ordre de 1 km, et de 5 à 45 km pour chaque animal et des durées maximales de 3, 6, 8 et 12 heures ont été testées. Les séries de RT présentant le contraste le plus élevé ont été



sélectionnées et les segments ont été catégorisés en tant que temps de résidence élevé (HRT) ou faible (LRT) selon la méthode de segmentation de Lavielle (Lavielle, 2005), en utilisant le paquet `adehabitatLT` dans R (Calenge, 2006).

J'ai utilisé des estimations de densité kernel pour calculer la distribution de l'utilisation (UD) ; avec 'kernels' de 25%, 50%, 75% et 95% UD avec le paquet R `adehabitatHR` (Calenge, 2006). La proportion de chevauchement du 'home range' (95% UD) et des 'core areas' (50% UD) a été estimée et rapportée dans une gamme allant de 0 (aucun chevauchement) à 1 (chevauchement complet). J'ai ensuite déterminé la densité des événements de recherche de nourriture par heure de la journée pour évaluer s'il existe une ségrégation interspécifique entre l'occurrence de la recherche de nourriture par heure de la journée pour chaque sexe. Pour vérifier s'il existe des modèles différents par espèce et par groupe de sexe, nous avons construit un modèle mixte additif général (GAMM, package R `mgcv`) pour déterminer si l'heure de la journée avait un effet significatif sur les événements de recherche de nourriture qui se sont produits dans des blocs d'une heure selon chaque espèce et chaque groupe de sexe (Wood, 2011).

Pour déterminer les variables environnementales qui expliquent le mieux les conditions environnementales pendant la recherche de nourriture, nous avons effectué un processus de sélection avant pour chaque espèce et chaque sexe, en construisant un modèle GAMM avec une distribution binomiale (variable de réponse 1,0) et des 'cubic splines' pour modéliser les relations non paramétriques. L'identité de l'animal a été définie comme effet aléatoire et une structure d'erreur `corAR1` pour tenir compte de la corrélation temporelle. J'ai évalué la colinéarité et les valeurs manquantes. Pour éviter un ajustement excessif, les nœuds ont été fixés à 5. La sélection des modèles a été effectuée par un processus de sélection avancée qui consistait à évaluer si l'ajout d'une nouvelle covariable produisait une amélioration significative du modèle, évaluée par des tests ANOVA (Chi carré,  $p < 0,05$ ).

## 4.4 *Chapitre 4 : Niche isotopique*

### 4.4.1 *Échantillonnage*

Les vibrisses ont été collectées sur 76 SAFS (55 femelles et 21 mâles) et 36 SASL (14 femelles et 22 mâles). Les vibrisses, y compris le follicule, ont été enlevées manuellement. Tous les animaux ont été jugés en bonne santé par un vétérinaire sur la base des résultats de l'examen physique et de l'évaluation vétérinaire des paramètres sanguins de routine. Les poids corporels ont été obtenus en utilisant un trépied et une balance de terrain à la centaine de grammes près (Mini Crane Scale 300 kg, OCS-L).

Les enregistrements mensuels du niveau de la mer (cm) pour la latitude 15° Sud, emplacement de notre zone d'étude entre janvier 2005 et décembre 2019, ont été extraits du dépôt de produits océaniques Copernicus (<https://marine.copernicus.eu/>). Les enregistrements mensuels de la température de surface de la mer (SST) recueillis par la base navale de la baie de San Juan de Marcona (Dirección de Hidrografía y Navegación, DHN), située à moins de 5 km du site d'échantillonnage pendant la période d'étude. Nous avons construit une série temporelle de SST pour chaque mois de la période de quinze ans.

### 4.4.2 *Analyses des données*

Premièrement, j'ai utilisé des modèles linéaires à effets mixtes (LMM) ((Pinheiro & Bates, 2000) pour examiner l'effet de la masse corporelle sur les valeurs  $\delta^{13}\text{C}$  et  $\delta^{15}\text{N}$  de chaque sexe, en utilisant l'individu (ID) comme effet aléatoire. Puisque la masse corporelle au moment de la collecte des vibrisses est affectée par les habitudes alimentaires des derniers mois, j'ai inclus dans le LMM les valeurs des trois derniers fragments analysés de chaque individu, intégrant ainsi les informations sur l'alimentation des six derniers mois. Deuxièmement, j'ai utilisé le LMM pour examiner les différences sur les valeurs moyennes de  $\delta^{13}\text{C}$  et  $\delta^{15}\text{N}$  entre les groupes (espèce et sexe), les périodes, et leurs interactions comme effets fixes. Les modèles incluaient l'identité comme effet aléatoire pour tenir compte des mesures répétées de chaque variable de réponse sur les différents fragments. Des ellipses bayésiennes d'isotopes stables en R (SIBER) (Jackson et al., 2011) ont été estimées pour mesurer la largeur de la niche isotopique à 75 % pour SAFS et SASL. Les valeurs isotopiques moyennes pour

chaque individu ont été utilisées pour estimer les ellipses puisque les fragments de vibrisses sont corrélés dans le temps et ne respectent pas l'indépendance de l'échantillon. Cette méthode est une version bayésienne de la métrique de Layman (Layman et al., 2007) qui peut intégrer des incertitudes telles que les biais d'échantillonnage et les petites tailles d'échantillon dans la métrique de niche (Jackson et al., 2011). Basée sur la simulation Monte Carlo à chaîne de Markov, l'approche SIBER obtient des mesures d'incertitude pour construire les paramètres des ellipses d'une manière similaire au bootstrap. Les zones d'ellipse standard corrigées pour la petite taille de l'échantillon (SEAc) ont été utilisées pour comparer les différents groupes sexe-espèce (femelles SAFS, femelles SASL, mâles SAFS, mâles SASL) dans l'espace isotopique au cours des deux périodes et ont estimé la zone de chevauchement de leur niche isotopique en utilisant les zones d'ellipse standard du maximum de vraisemblance (SEAc). Les valeurs estimées pour le chevauchement de chaque groupe sont exprimées en pourcentage.

Le niveau de la mer et la SST étaient corrélés pendant la période d'étude (Spearman,  $r=0.74$ ,  $p<0.001$ ), pour éviter la redondance, j'ai choisi le niveau de la mer comme variable environnementale car il reflète les processus à plus long terme causés par l'expansion de la chaleur dans la colonne d'eau. En revanche, en raison du forçage atmosphérique à grande échelle, la SST n'est pas un bon indicateur des changements d'intensité de l'upwelling dans un contexte de réchauffement (Demarcq, 2009).

## 5. Résultats

Dans le chapitre 1, j'ai analysé les tendances de l'abondance sur une période de 20 ans et j'ai constaté qu'en moyenne, 83% de toute la biomasse des otariidés est composée par la SASL et 17% par la SAFS; ce qui fait de la SASL l'espèce "dominante" actuelle du site d'étude (la SASL représente actuellement 89% de la biomasse totale des otariidés). De plus, les résultats de l'évaluation de la population révèlent un déclin actuel des deux populations, suggérant une limitation des ressources dans l'environnement marin, qui a commencé en novembre 2014 chez SAFS et en mars 2017 chez SASL.

Dans le chapitre 2, j'ai déterminé que la ségrégation alimentaire interspécifique existe en raison du partage de trois principales proies communes : L'anchois du Pérou, les céphalopodes et les galathées. La consommation de ces proies dans des proportions différentes par chaque espèce se sépare en deux

groupes contenant les combinaisons des ressources de proies consommées pendant les échantillonnages saisonniers consécutifs de la période d'étude 2015-2018. En général, les SAFS consomment systématiquement d'anchois du Pérou (42,8 %) et des céphalopodes (34,7 %), avec des composantes plus faibles de myctophidés (7,4 %), de galathées (5 %) et de onze autres espèces de poissons. Le régime alimentaire de la SASL est principalement composé de galathées (69,9%) et d'anchois du Pérou (19,1%), avec des quantités moindres de céphalopodes (4,7%), de capucettes (4,6%) et de cinq autres espèces de poissons. La classification des proies en groupes écologiques, basée sur la littérature publiée sur les habitudes des proies, a montré que les proies de la SAFS appartiennent aux groupes écologiques pélagique (52,35%) et démersal-pélagique (47,25%). Quant aux proies de la SASL, elles appartiennent aux groupes démersal-pélagique (69,44 %) et pélagique (23,88 %). Les espèces benthiques démersales ne sont pas importantes dans la composition du régime alimentaire des otariidés péruviens (1-1,5%). Cela suggère que la majorité des proies proviennent de la zone pélagique, mais des méthodes plus raffinées sont nécessaires pour comprendre l'utilisation verticale ou temporelle des zones de recherche de nourriture des otariidés (utilisation d'enregistreurs de plongées). De plus, une différence significative importante a été trouvée entre l'étendue de la niche basée sur l'identification des parties dures, avec SAFS ( $0,26 \pm 0,13$ ) atteignant le double du *Bi* de Levin (larguer de niche) que SASL ( $0,14 \pm 0,16$ ). En général, le régime alimentaire de la SAFS tendait également à présenter une plus grande richesse en espèces que celui de la SASL (chapitre 2). En comparant les espèces consommées, bien que la composition du régime alimentaire de SAFS se concentre sur la consommation d'anchois et de céphalopodes, pendant de nombreuses saisons, il a ajouté de nouvelles espèces au régime alimentaire, expliquant l'élargissement de la largeur de la niche ; contrairement à une richesse en espèces moins variable dans le régime alimentaire de SASL, basé de manière constante sur de grandes quantités de galathées et d'anchois.

Les résultats du chapitre 3 montrent que les otariidés sympatriques du HCS ont de multiples stratégies de ségrégation lors de la recherche de nourriture dans l'environnement pélagique et que la masse corporelle n'est pas le seul moteur de la ségrégation dans l'espace et le temps, mais plutôt une combinaison de contraintes énergétiques en fonction du sexe, des rôles reproducteurs liés à des ressources alimentaires spécifiques. J'ai constaté que, bien qu'ils soient de plus petite taille, les femelles et les mâles de la SAFS effectuent des voyages de plus longue distance et de plus longue durée que les deux sexes de SASL, rejetant notre hypothèse selon laquelle les animaux plus grands effectueraient des voyages plus longs. En termes de superficie, les SASL couvrent des 'home range'

et des 'core area' plus étendus, mais ils le font en effectuant davantage de déplacements consécutifs plus courts que les SAFS. Cela révèle une utilisation différente des zones de recherche de nourriture au niveau interspécifique. Les SASL ont des 'home range' plus étendus en tant que groupe, couvrant une plus grande zone parallèle au littoral qui enveloppe le 'home range' des SAFS. En revanche, les SAFS se déplacent plus loin au large pendant les voyages d'alimentation vers des zones spécifiques. En accord avec nos prédictions basées sur des études précédentes portant sur la ségrégation sexuelle des otariidés, les femelles des deux espèces montrent des voyages de recherche de nourriture plus limités en termes de distance parcourue, de 'home range' et de 'core areas' en comparaison avec les mâles des deux espèces (Page et al., 2006; Staniland & Robinson, 2008). Enfin, bien que la dispersion post-reproduction soit courante chez les pinnipèdes mâles, cette étude montre que les otariidés péruviens ne partent pas en migration, mais subviennent à leurs demandes énergétiques dans une zone locale de 55 000 à 106 000 km<sup>2</sup>. Ces données sont la preuve que ces espèces sympatriques ont évolué et se sont adaptées pour obtenir leurs ressources localement dans les eaux d'upwelling productives du HCS, occupant des « rookeries » comme PSJ toute l'année.

Deux échelles sont explorées et discutées pour la ségrégation dans la dimension temporelle au chapitre 3. Les otariidés péruviens ont une ségrégation temporelle de 1 à 3 mois dans leurs saisons de reproduction (SAFS se reproduit d'octobre à décembre et SASL de janvier à mars). C'est un mécanisme de ségrégation, connu sous le nom d'allochronie, qui permet de répartir les ressources pendant les périodes énergétiquement critiques (Taylor & Friesen, 2017). Étant donné que la lactation est prolongée chez les espèces de latitude tempérée et que le moment du sevrage est flexible (il est signalé qu'il a lieu entre 6 et 36 mois), nous considérons qu'il est utile d'explorer les mécanismes de ségrégation pendant que les femelles en lactation cherchent de la nourriture pour allaiter leurs petits. A une échelle plus petite, j'ai trouvé une recherche de nourriture nocturne chez les femelles SAFS (pic au crépuscule) et diurne chez les mâles SASL (pic à l'aube).

Les variables environnementales associées aux événements de recherche de nourriture évalués chez les pinnipèdes sympatriques dans le chapitre 3 ont révélé des différences interspécifiques et intraspécifiques. Lorsqu'on les compare, les caractéristiques de l'habitat des femelles SASL correspondent à un habitat côtier qui est caractérisé par une distance à la côte (DC) plus faible, une température de surface de la mer (SST) plus faible, une proximité des fronts plus faible et des valeurs de chlorophylle-a (Chl-a) plus élevées. Alors que les femelles SAFS sont associées à un domaine plus

pélagique caractérisé par une SST plus élevée, une plus grande proximité des fronts, des gradients de fronts plus élevés et bien sûr, des valeurs de DC plus élevées. Parmi les mâles, les modèles les mieux adaptés pour les mâles SAFS comprenaient la SST, la Chl-a et la proximité des fronts, tandis que les mâles SASL comprenaient toutes les covariables environnementales. Les mâles SASL utilisent des zones où la SST est plus faible, la Chl-a est plus faible, la proximité des fronts est plus faible et la DC est plus faible, ce qui pourrait être le résultat de stratégies individuelles. Les gammes des variables environnementales ciblées par les mâles SAFS et SASL se chevauchent fortement; les conclusions doivent donc être prises avec précaution et d'autres études explorant les stratégies individuelles doivent être menées. Je suggère également d'incorporer des variables environnementales supplémentaires pour mieux expliquer l'environnement de recherche de nourriture des mâles, telles que celles liées à la structure verticale de la colonne d'eau.

Dans le chapitre 4, j'ai constaté une réorganisation de nos quatre groupes d'étude d'otariidés dans l'espace isotopique en réponse à l'augmentation du niveau moyen de la mer et au réchauffement des températures de surface de la mer à partir de novembre 2013 (Période 2). Les groupes ont élargi la largeur de leur niche (à l'exception des femelles SASL), se sont séparés dans l'espace isotopique et ont réduit le signal  $\delta^{15}\text{N}$  pendant la période plus chaude. Un déclin du  $\delta^{15}\text{N}$  des otariidés suggère des valeurs de base appauvries de  $\delta^{15}\text{N}$  avec des effets bottom-up impactant les consommateurs tertiaires, un changement de régime alimentaire vers des proies de niveau trophique inférieur, ou les deux. Le chevauchement des niches était réduit entre tous les groupes d'otariidés au cours de la Période 2, ce qui confirme le passage temporel d'un assemblage plus redondant (niches isotopiques similaires) au cours de la période plus froide (Période 1) à un assemblage plus diversifié sur le plan trophique (niches isotopiques divergentes) au cours de la période plus chaude (Période 2). Cet ensemble de données à long terme montre un changement dans l'organisation trophique qui est aligné temporellement avec le déclin des populations d'otaries de Punta San Juan détecté au chapitre 1. De plus, la rupture en deux périodes de conditions environnementales variables explique les changements dans la réorganisation trophique de la communauté d'otariidés. Au cours de la Période 2, les SASL femelles présentent la plus grande chute de  $\delta^{15}\text{N}$  de 2.4, tandis que les valeurs de  $\delta^{15}\text{N}$  des groupes mâles diminuent de 1.5 et les SAFS femelles diminuent légèrement de  $\delta^{15}\text{N}$  de 1.0. Nous considérons que le changement de  $\delta^{15}\text{N}$  au cours de la Période 2 reflète un effet ascendant dans le réseau trophique, lié aux habitudes alimentaires de chaque espèce et sexe, en réponse aux changements de l'environnement.

## 6. Conclusion et perspectives

Les populations peuvent être séparées en spécialistes lorsqu'elles ont des niches plus étroites ou en généralistes lorsqu'elles ont des niches plus larges (Begon et al., 2006). Les populations généralistes avec des niches plus larges sont probablement favorisées dans les environnements hétérogènes et stables, alors que les populations spécialistes avec des niches plus étroites sont favorisées dans les écosystèmes hautement dynamiques et les environnements homogènes (Kassen, 2002). Ainsi, pendant la période plus chaude, les résultats suggèrent que les assemblages de proies sont plus diversifiés, contrairement aux agrégations denses de proies favorisées par les eaux froides et riches en nutriments dans les HCS ; ce qui favorise une réponse différente chez les otariidés, qui se reflète dans leurs stratégies de recherche de nourriture, la composition de leur régime alimentaire et la dynamique de leur population.

De toutes les espèces et les sexes, les femelles SASL montrent la plus grande variation en réponse au scénario de réchauffement, réduisant à la fois la largeur de la niche isotopique et les valeurs  $\delta^{15}\text{N}$ . Cela suggère que ce groupe a une plus grande plasticité pour s'adapter aux conditions environnementales changeantes, comme cela a été rapporté chez d'autres otariidés (Amador-Capitanachi et al., 2020). Sachant que le SASL est l'espèce dominante au Pérou, une plasticité plus élevée chez les femelles adultes permet d'expliquer la biomasse actuelle et plus importante de la population de SASL rapportée pour PSJ et le Pérou, contrairement à la population de SAFS, plus petite et en déclin plus prononcé (Cárdenas-Alayza et al., 2021). Ainsi, la plasticité trophique, améliorée par l'alimentation principalement sur une source de nourriture plus disponible (*P. monodon*) qui n'a pas de pression de pêche commerciale, peut être actuellement plus avantageuse pour la croissance de la population de SASL.

Les résultats montrent que les otariidés sympatriques dans le HCS montrent actuellement un certain niveau de ségrégation à travers les indices trophiques et pendant la recherche de nourriture en mer, en partie associé au rôle reproducteur, à la taille du corps et à l'espèce. Sur la base de nos séries chronologiques d'isotopes stables à long terme, les signaux isotopiques (Edwards et al., 2021) et les mécanismes de ségrégation trophique ont changé après novembre 2014. Les mécanismes de ségrégation n'ont de sens que si les proies sont partiellement limitées et partitionnées dans l'espace.


Dans le cas d'une très grande disponibilité des proies, la coexistence pourrait avoir lieu sans qu'il y ait besoin de ségrégation. Cependant, en cas de pénurie de nourriture, les mécanismes de ségrégation pourraient soutenir la coexistence, en tirant parti des capacités différentielles à faire face à une telle situation, ce qui pourrait être le cas de la période plus chaude (période 2 au chapitre 4). Dans ce dernier cas, les différences observées dans les écologies de recherche de nourriture peuvent dépeindre une situation où une espèce surpasse une autre, ce qui ne pourrait pas être considéré comme un mécanisme de coexistence. Dans ce contexte, la plasticité des femelles de SASL, par opposition au groupe de femelles de SAFS, moins flexible, peut aider à expliquer que SASL soit l'espèce dominante des deux espèces sympatriques dans HCS. Enfin, dans le cadre de cette étude, nous proposons une série de recommandations pour les recherches futures afin de continuer à démêler les sources de ségrégation pour expliquer la coexistence des otariidés péruviens et d'autres prédateurs dans le HCS.

À ce jour, il est bien documenté que l'anchois péruvien (*Engraulis ringens*) est fortement exploité et constitue une source de compétition pour plusieurs prédateurs dans le HCS. Cependant, nous n'avons pas connaissance d'une exploitation à grande échelle de galathées (*P. monodon*) au Pérou qui limite la disponibilité de cette proie pour les prédateurs. Si cela devait se produire, cela pourrait gravement affecter la disponibilité de cette ressource proie pour les SASL et pourrait entraîner d'autres changements dans le régime alimentaire de cette espèce, et augmenter la concurrence entre les deux espèces d'otariidés sympatriques comme cela a été mis en évidence dans d'autres régions ((Drago et al., 2009; Szteren et al., 2018).



# INTRODUCTION GÉNÉRALE

---



"WHAT WORLD LIES BEYOND THAT STORMY SEA  
I DO NOT KNOW,  
BUT EVERY OCEAN HAS A DISTANT SHORE,  
AND I SHALL REACH IT."

*- Cesare Pavese*



## 1. Introduction


Understanding how sympatric species with similar requirements coexist, is an important theme in ecology. Competition is an ecological processes permeated by natural selection, that shapes species in present time, in reference to the conditions of their ecological setting. Competition can compromise the fitness of the species involved and may ultimately lead to the competitive exclusion of one species (Pacala & Roughgarden, 1985). The ways in which species within ecological communities partition available resources among themselves is a major determinant of the species diversity. A community with more resource sharing or greater niche overlap, will support more species than one with less niche overlap. Thus, coexisting species must differ in their ecological requirements by at least some minimal amount to avoid competitive exclusion (Pianka, 1974).

This study focuses on two sympatric top predator populations of otariid species South American fur seals (SAFS, *Arctocephalus australis*) and South American sea lions (SASL, *Otaria byronia*), that appear to have similar trophic requirements. After population declines caused by extreme food shortage due to 1997-98 El Niño Southern Oscillation (ENSO), SASL in Peru have been growing steadily, with an estimated population of over ca. 130,000 individuals (IMARPE, 2014b). In comparison SAFS in Peru have not been able to recover in the same extent, barely reaching a population of ca. 11,000 individuals (IMARPE, 2014b). This study focuses on information collected via instrumentation of animals, counts and samples collected at Punta San Juan (PSJ, Latitude: 15°22') in south Peru, where these two species share a common coastal haul-out site for reproduction. In PSJ, breeding populations concentrate up to 5,000 SAFS and 12,000 SASL within the 2013-2019 study period of this thesis. In Peru, these species have temporal segregation in their breeding seasons, which are 1-3 months apart. Allochrony, which is recognized as a major driver behind speciation, can be considered to facilitate resource partition during energetically critical times (Taylor & Friesen, 2017). In Peru, SAFS breed during October-December and SASL breed during January-March. However, lactation of offspring can be extended compared to other otariid species, and time of weaning is flexible with reports of 6 - 36 months, with high interannual variability (Majluf, 1987; Soto, 2004). This generates a temporal overlap in the time of high energetic demand during which females have to ensure the survival of their young.

To understand the level of competition between these two species, it is necessary to understand the



overlap and/or segregation between trophic niches (e.g., niche differentiation)). This will only be possible by determining the species mechanisms of exploitation and resource partitioning in their environment. When species are living in the same habitat, the ways in which they can partition their resources are: spatial separation (e.g. feeding in different depths or geographic locations); temporal separation of resources (e.g. utilizing resources that are available at different times of day or different seasons) or dietary segregation (e.g. consumption of different prey types) (Begon et al., 2006). However, it is crucial to take into account the environmental context of coexisting species; in which varying conditions can produce new responses before patterns are completely established (Figure I.1).



**Figure I. 1** Diagram showing resource partitioning among 11 species of anole lizards found on the island of Puerto Rico. Each species occupies a different type or elevation of vegetation. The habitat is further partitioned by the amount of sunlight and moisture available. Image by Eva Horne modified from (Williams, 1983). Source: <https://www.khanacademy.org/science/ap-biology/ecology-ap/community-ecology/a/niches-competition>

Another important influential feature on the foraging mode in otariids may be local marine productivity and its effect on prey availability. It may be that, except in very productive regions, smaller body size favors foraging on a highly patchy but dense prey resource near the surface



(zooplankton, small fish or squid) compared to larger body sized air breathing mammals are better adapted for searching for more evenly distributed but less dense prey resource on the benthos (larger fishes or octopus). This coincides with the trend of greater mass of prey items being consumed by benthic foraging sea lions and the greater proportions of time spent at sea diving compared to smaller epipelagic fur seals (Arnould & Costa, 2006). More recent studies have elucidated that this pattern is not always the norm, and there are some sea lion populations that inhabit upwelling areas that display pelagic foraging (Hückstädt et al., 2014), fur seals that are benthic divers (Arnould & Hindell, 2001) and some otariids that display a combination of shallow and deep dives according to metabolic needs and local prey availability (Harcourt et al., 2002; Villegas-Amtmann et al., 2008). This demonstrates the plasticity in the adaptations of populations and individuals to local spatial and temporal conditions, and which may also vary according to sex and age (Weise & Costa, 2007).

Thus, the main goal of this study is to describe the ecological niches of sympatric otariid species to determine intraspecific and interspecific niche differentiation / overlap in the marine environment of the HCS. It is particularly important to understand the partitioning mechanisms that drive coexistence of predator populations. Predators usually affect ecosystems through top-down control and behaviorally mediate impacts on prey species. Thus, the maintenance of predator population is key to preserve healthy ecosystems. Accurate assessments of foraging ecology are essential to help understand the function of predators and structure of ecosystems. As ecosystems and trophic webs are highly dynamic, the coupling of traditional diet information (e.g. hard parts identified in scats) with foraging location data can add a spatial and temporal dimension to dietary estimates (Jeanniard du Dot et al., 2017). Furthermore, long time series of population counts and isotopic signatures from whiskers will provide a different temporal windows to explore trophic habits in SAFS and SASL during 15 to 20 year timeframes. To achieve this, the current study proposes to combine a suite of datasets and analytical tools that will offer insight into the trophic dimension of both species at different temporal and spatial scales to explain current divergent trends in population abundance of sympatric otariid predators.



## 2. Theoretical Framework

### 2.1 *Ecological niche*

A niche is an  $n$ -dimensional hypervolume which defines the limits within which a given species can survive and reproduce for a number of environmental factors, including both conditions and resources (Hutchinson & MacArthur, 1959). The resource dimensions of a species niche can sometimes be represented in similar manner to that adopted for conditions, with lower and upper limits within which a species can thrive. Thus, a predator may only be able to detect and handle prey between lower and upper limits of size. Together, resources and conditions, define a species niche.


The ecological niche of a species needs to be multidimensional. In an  $n$ -dimensional hypervolume,  $n$  is the number of dimensions that make up the niche. Usually a species has a larger ecological niche in the absence of competitors and predators. Thus, there are a certain number of combinations of conditions and resources that can allow a species to maintain a viable population, but only if it is not being adversely affected by competitors (Hutchinson & MacArthur, 1959).

All environments are heterogeneous. The extent to which an environment is heterogeneous depends on the scale of the organism that senses it. There may also be gradients in space (e.g. depth) or gradients in time (e.g. seasonal cycles). Gradients can be directional (e.g. accumulation of pollutants) or erratic (e.g. tsunami) as well. When a species is exposed to variations in the physical factors of the environment, the two species that interact can change and produce alterations in the life of the other, and each may generate selective forces that direct the evolution of the other (Begon et al., 2006). Behavior and metabolism constrain the animal within a narrow niche and deny it access to what otherwise may appear as a suitable alternative food source (Begon et al., 2006). Temperature, for example, limits the growth and reproduction of all organisms, but different organisms tolerate different ranges of temperature. Thus, in this example, the temperature range is one dimension of the organisms' ecological niche.



## 2.2 Competition and coexistence

From a historical evolutionary point of view, natural selection have shaped species. The ecological effects are broad and species may be eliminated from a habitat by competition from individuals of other species or reductions in fecundity and survival appear (Begon et al., 2006). Under current ecological conditions, competition and resources availability shape species *in vivo*. In this context, competition can compromise the fitness of the species involved and may ultimately lead to the competitive exclusion of one species (Figure I.2), following the Competitive Exclusion Principle (Hardin, 1960; Pacala & Roughgarden, 1985).




**Figure I. 2** Competitive exclusion example showing *Paramecium caudatum* grow well individually, but when they compete for the same resources, *P. aurelia* outcompetes *P. caudatum*. Image from: Fowler et al., 2013. Source: <https://openstax.org/books/concepts-biology/pages/19-4-community-ecology>

When two species compete, individuals of one or both species may suffer reductions in fecundity or survivorship. The fittest individuals of each species may be those that escape competition because they utilize the habitat in ways that differ from most of those adopted by individuals of the other species. Natural selection can favor such individuals and eventually the population can consist entirely of them. The two species become more different from one another, compete less and can evolve to coexist (Begon et al., 2006). The Lotka-Volterra model predicts the stable coexistence of competitors in situations where interspecific competition is, for both species, less significant than





intraspecific competition (Begon et al., 2006). Thus, niche differentiation will tend to concentrate competitive effects more within species than between them. The Lotka-Volterra model and the Competitive Exclusion Principle imply that complete competitors cannot coexist and species must have ecological differentiation as a necessary condition for coexistence (Figure I.3, Hardin, 1960).



**Figure I. 3** Schematic representation of Lotka-Volterra competition. Image from D'Andrea et al., 2019

### 2.3 *Sympatry in otariids*

Sympatric species are those species whose breeding ranges and distribution overlap and in which individuals can coexist without hybridizing. Thus, it is defined as the occurrence of two or more species breeding in the same location (Cain, 1953). If two non-interbreeding populations occupy the same geographical territory and the same ecological niche, they are sympatric. Many species of fur seals and sea lions occur in sympatry throughout the world (Bailleul et al., 2005; Dellinger & Trillmich, 1999; Franco-Trecu et al., 2012; Jeglinski et al., 2013; Page et al., 2005; Villegas-Amtmann et al., 2013; Waite, Trumble, et al., 2012). Areas of sympatry, however, represent only a small proportion of the allopatric or non-breeding ranges. The duration of coexistence in sympatric otariid seals is not known, and it is possible that sympatry has resulted from recent population recoveries and recolonization following the cessation of commercial sealing (Costa et al., 2007). If sympatry is a result of recent events, it is possible that divergence is not yet evident or that competitive exclusion is currently occurring.

Some authors suggest that the differences in the population dynamics of sea lions and fur seals are



associated with the observed divergent trends of epipelagic and benthic foraging (Arnould & Costa, 2006). All species of otariid seals throughout the world were subject to extensive and, in most cases, excessive hunting pressure during the eighteenth and nineteenth centuries. By the late 1800s, however, most species had acquired total legislative protection or were subject to only regulated harvests. Despite this protection, populations of various sea lion species have experienced very little recovery and in some cases are declining; whereas fur seal species have generally experienced rapid population recovery rates (Costa et al., 2007; Wickens & York, 1997).

## 2.4 *Interspecific segregation*

Interspecific competition, defined as reciprocal negative effects of one species on another, is an important process determining the structure of natural communities (Chase et al., 2002). This can occur either directly or indirectly mediated by changes in resource availability. Lotka–Volterra competition theory states that coexistence of two species is possible when the per capita effects of intraspecific competition on per capita rates of population growth are greater than those of interspecific competition (Chase et al., 2002; Chesson & Warner, 1981; Schoener, 1983).

In conditions of limited resources, competition between species implies a reduction in some population attributes, such as growth, survival or fecundity (Begon et al., 2006). Closely related species with similar life-history strategies often share similar niches. Non-migratory, central-place foraging species with overlapping ranges may compete for similar resources, such as prey or breeding sites. Among the sympatric marine mammal species with similar life-history traits and foraging habits are the otariids: fur seals and sea lions. In general, where fur seals and sea lions live in sympatry, the fur seal population is typically larger and they appear to outcompete sea lions. Many studies have examined potential competition between sympatric fur seal and sea lions, and their results are mixed. Some have found ecological segregation with no trophic overlap (Antonelis et al., 1990; Aurioles-Gamboa & Camacho-Ríos, 2007; Dellinger & Trillmich, 1999; Franco-Trecu et al., 2012), while others have found some dietary overlap (Páez-Rosas & Aurioles-Gamboa, 2010; Page et al., 2005; Waite, Burkanov, et al., 2012).

Recent studies that include stable isotope analysis have elucidated findings that provide new insights into understanding how sympatric species exposed to strong interspecific competition can develop





foraging strategies that may decrease competition and facilitate survival in stochastic environments with variable resource availability. Individual long-term dietary specialization from two sympatric otariid species using isotopic ratios in whiskers was analyzed in SAFS and SASL of the Uruguayan coast showed that fur seals had a larger trophic niche resulting from generalist individuals, whereas sea lions have a smaller trophic niche, from individuals with higher degree of trophic specialization (Franco-Trecu et al., 2014). Also, a study comparing stable isotopes in Galapagos sea lions (*Zalophus wollebaeki*) and Galapagos fur seals (*Arctocephalus galapagoensis*), found that differences in habitat use. *Z. wollebaeki* was persistent in repeating the use of the same benthic / inshore feeding grounds, whereas *A. galapagoensis* feeds on pelagic / offshore prey items; while both species isotopic signatures vary according to oceanographic conditions (El Niño, La Niña, normal years) (Páez-Rosas et al., 2012).

## 2.5 *Intraspecific segregation*

Sexual differentiation in the ecological niches in pinniped species is expected due to high dimorphism between sex. Males are originally thought to have higher diving capacity to exploit larger three dimensional foraging areas, and considering that they are not involved in postnatal care, they could feed farther away from the rookery than females and exploit wider ranges and more exclusive foraging habitats (González-Suárez & Cassini, 2014). Stable isotope studies are challenging this assumption and demonstrating that this is not always the case. Several recent studies demonstrate the role of the local environment as a major driver of resource partitioning in otariid populations (Drago et al., 2015).

Individual long-term dietary specialization of male and females sympatric otariid species using isotopic ratios in whiskers was analyzed in sympatric South American fur seals *A. australis* and South American sea lions *O. byronia* on the Uruguayan coast. Sexual dimorphism in body size, present in both species, suggests that males will exploit larger preys often at higher trophic levels. However, in the study,  $\delta^{15}\text{N}$  values showed that both sexes of sea lions and fur seals consumed preys at similar trophic levels. Females in both species, showed wider ranges of  $\delta^{15}\text{N}$  that are associated with a higher consumption of prey at lower trophic levels than males. This differentiation was interpreted by the additional constraint on female foraging trips, since this group is obligated to return to shore to nurse



their pup (Franco-Trecu et al., 2014).

In terms of differences in  $\delta^{13}\text{C}$ , different signals between sexes may reflect physiological (body mass) or behavioral (parental care) differences. An expected result of the same beforementioned study was that males, who are not involved in postnatal care, could use their higher diving capacity to exploit larger, deeper and more distant foraging areas than females. This would lead to wider ranges of a  $\delta^{13}\text{C}$  in more exclusive foraging habitats. Study results showed that sea lion males had a lower  $\delta^{13}\text{C}$  values than females; and the opposite was true for fur seals. Although the range of  $\delta^{13}\text{C}$  was wider for males of both species, there was a large overlap in the overall areas of isotopic niches between the sexes of both sea lions and fur seals. Thus, males despite not being linked to the colonies, have a similar foraging behavior to females off the coast of Uruguay (Franco-Trecu et al., 2014).

## 2.6 *Otariids' foraging*

Animals should forage in a way that optimizes their energy intake and, ultimately, fitness. Multiple factors influence foraging behavior, including predation risk and prey availability. Air-breathing mammals and birds foraging underwater are additionally constrained by the duration of their dives as they need to supply oxygen stores frequently (Sigler et al., 2009). The foraging behaviors of top marine predators are related to the energetic requirements of individuals and environmental features (Costa & Gales, 2003). Indeed, according to optimal foraging theory, organisms are supposed to adopt foraging behaviors that optimize fitness for given environmental conditions. In the marine environment, food resources are distributed heterogeneously in space and time and their distribution is related to oceanographic features. Distribution and behavior of top marine predators are related to physical and biological features (bathymetry, sea-surface temperature, primary productivity), as found in seabirds (Bost et al., 1997; Guinet et al., 1997) and pinnipeds (Georges, Bonadonna, et al., 2000; Jessopp et al., 2004).

Many pinniped studies in the 1980s and 1990s were conducted on lactating females, since this is the age-class with highest energetic constraint, and to increase in the recovery of biologging devices compared to males. Results from these studies have grouped sea lions and fur seals to have two broadly divergent foraging patterns. Lactating sea lions generally undertake short trips (1-2 days) foraging mostly on the benthos of continental shelf areas where they have a continuous dive pattern



with no diel variation, foraging mostly on benthic or demersal prey in continental shelf areas (Riet-Sapriza et al., 2013; Thompson et al., 1998; Werner & Campagna, 1995). In contrast, lactating fur seals generally undertake longer trips (4-23 days) foraging mostly on vertically migrating prey in oceanic frontal structures or continental shelf-edges with upwelling regions (Arnould & Costa, 2006), during which diving is mostly nocturnal. This mode of foraging is hereafter referred to as “epipelagic” foraging and mainly occurs in oceanic frontal structures or continental shelf-edges with upwelling regions (Gentry & Kooyman, 1987). The dives occur in the deep scattering layer, with a pronounced diel variation in depth that reflects the vertical migration of their prey (Boyd & Croxall, 1992; Francis et al., 1998; Georges, Tremblay, et al., 2000).

## 2.7 *Body size and oxygen storage*

A main feature that differentiates fur seals and sea lions is their fur layer. The insulating feature of fur seal integument traps a layer of air within the fur and is inefficient at great depths preventing them from foraging as deep as sea lions (Arnould & Costa, 2006; Gentry & Kooyman, 1987). Also, it has been suggested that the larger body size of sea lions compared to fur seals results in greater oxygen storage capabilities, enabling them to aerobically dive for longer periods and reach higher depths (Costa et al., 1998) and can have implications on the metabolic and reproductive output. The Aerobic Dive Limit (ADL) is a model to understand if pinnipeds and seabirds are performing dives that approach their maximum aerobic capacity. ADL is defined as the maximum dive duration before blood lactic acid levels rises as a result of an increase in anaerobic metabolism, and is therefore a function of both oxygen stores and oxygen consumption (metabolic rate).


Calculated ADL (cADL) is a commonly used proxy for ADL, and is calculated from estimates of volumes of usable oxygen stores divided by oxygen consumption in a particular species (Butler, 2006). Exceeding cADL has been reported for some benthic and deep diving otariids. There is evidence indicating that the diving capacity of fur seals measured in their aerobic dive limit is relatively smaller (1.6 to 1.7 min) compared to sea lions (2.3 to 3.8 min), which imposes a constraint for the depth fur seals are capable of reaching during foraging dives (Costa et al., 2001). It has been proposed that deep diving strategies can put the population at risk if individuals are unable to cope with environmental variability and fluctuations. Thus, otariids that display benthic foraging usually



exhibit smaller population sizes and lower population growth rates compared with those that feed pelagically. A revision on reproductive outputs showed that benthic feeding otariids have been documented to have significant lower birth rates compared to epipelagic feeders (Arnould & Costa, 2006). As mentioned, recent studies have elucidated that the ‘benthic sea lions and pelagic fur seals’ pattern is not always the norm, demonstrating the plasticity in the adaptations of populations and individuals to local spatial and temporal conditions (Villegas-Amtmann et al. 2008, Schwarz et al. 2021). Other studies (Hückstädt et al., 2014) have found a weak relationship between pinniped size and total oxygen stores amongst individuals with different foraging strategies (pelagic and benthic). Thus, local environment is an important driver in diving performance among pinnipeds shaping foraging strategies that needs to be explored in each ecosystem.

## 2.8 *Peruvian Humboldt Current System*

The Humboldt Current System (HCS) in Peru is recognized as the most productive of the Eastern Boundary Upwelling Systems in the world (Bakun & Weeks, 2008). As other upwelling ecosystems, cold nutrient rich water rises to the euphotic layer by a combination of the South east trade winds and the Coriolis force that fuel Ekman transport, giving rise to primary productivity in the upper pelagic and nearshore areas. Peru’s coast is characterized by a narrow continental shelf, reaching over 1000 m in less than 50 km offshore (Figure I.4). The shallow productive upwelling waters and shallow thermocline are related to an upper minimum oxygen layer at similar depths of approximately 50-80m on average, constraining expansion of vertical habitat (Figure I.5, Bertrand et al., 2010; Demarcq, 2009; Echevin et al., 2008). These features limit the potential habitat for otariid prey to be linked to the benthos. Thus, in this productive shallow environment foraging on highly patchy, but dense prey resources near the surface, may preclude the advantages of searching for more evenly distributed, but less dense, prey resources on the benthos, constrained by bottom water hypoxia.



**Figure I. 4** Morphology of seafloor along the coast of Peru. Punta San Juan study area located in box C, region with a very narrow continental shelf. Image from IMARPE, 2014a.


Furthermore, HCS is characterized by recurring El Niño Southern Oscillation (ENSO) events, which can vary in intensity and duration. ENSO is known as a cycle of alternating warm El Niño and cold La Niña events and is the most prominent climate signal on Earth (McPhaden et al., 2006). When HCS is affected by warming events, such as the El Niño Southern Oscillation (ENSO) or Kelvin waves, food web composition is altered (Tam et al., 2008). These changes impact the success of predator foraging events and, depending on their intensity and duration, can decrease reproductive output and survival rates in top predators (Majluf, 1991; Trillmich & Dellinger, 1991). Extraordinary ENSO events have triggered significant changes in the food web composition causing nutritional stress and mortality in many species (Bond & Lavers, 2014; Forcada et al., 2006; Oliveira, 2011; Sprogis et al., 2018).

The ENSO events that occurred in 1982-83 and 1997-98, were categorized as ‘Extraordinary’ and are





well documented for dramatically altering species composition and reducing marine biomass (Barber & Chavez, 1983; Tam et al., 2008). These climatic events were strong and felt worldwide, changing the availability of prey items and causing declines in reproduction and survival of top marine predator populations around the globe (Milena Arias-Schreiber & Rivas, 1998; Bond & Lavers, 2014; Cárdenas-Alayza, 2012; Forcada et al., 2006; Sprogis et al., 2018; Trathan et al., 2007). For example, the 1997-98 ENSO event caused a reduction of wildlife along Peru's coast with 75% and 80% reductions in the populations of SAFS and SASL, respectively (Oliveira et al., 2006, 2012). Since 1998 populations have recovered to an extent, but are currently in a state of decline (Cárdenas-Alayza et al., 2021).



**Figure I. 5** A) Example of the distribution of the shallow oxygen minimum zone in the Southeast Pacific Ocean. Source: <http://omz.udec.cl> ; B) Example of the oxycline delimiting the vertical distribution of plankton and anchovy schools in relation to sea bottom in Peru. Image from IRD /Marbec. Source: <https://books.openedition.org/irdeditions/docannexe/image/34324/img-5.jpg>

## 2.9 A warming climate

A warming climate is currently altering global ecosystem structure and driving species distributions to higher latitudes, thereby leading to altered interspecific interactions with unspecified consequences (du Pontavice et al., 2020; Walther et al., 2002). Climate driven shifts are more pronounced at the poles where deviations from long-term climatic variables are more striking (Kortsch et al., 2015; Parmesan & Yohe, 2003). However, in the past decade, a series of warming events have taken place



in the Pacific Ocean that may affect the organization of predator communities in the Peruvian HCS. Long-term warming climatic/oceanic events, or heatwaves known as “The Blob”, have impacted the Pacific Ocean in recent years (Bond et al., 2015). The Blob was first detected in the Gulf of Alaska in 2013-2014 and is a result of a persistent pattern of higher than normal rates of sea level pressure and weak coastal winds. In the past decade in Peru, El Niño conditions occurred between 2014 and 2015 (Graco et al., 2016), and was followed by the strong El Niño event of 2015-2016 that affected the entire Pacific Ocean (Robinson, 2016). This event was then followed by the effects of ‘El Niño costero’ that was a more localized event detected in the El Niño 1+2 region between January and February of 2017 (Paulino et al., 2019). During these years, studies suggest that physical anomalies of these warming ocean events generated a biological response similar to those of El Niño (Espinoza-Morriberón et al., 2017). All the beforementioned events involve the rise of sea level forced by the passage of remotely generated and coastal trapped waves in response to heat expansion (Ryan & Noble, 2002). The thermocline, nearshore depth of chlorophyll-a, and the nutricline deepened, decreasing nutrient enrichment in the euphotic zone. Weak winds caused the reduction in vertical nutrient fluxes to the depleted euphotic zone. A deeper concentration of chlorophyll-a decreased phytoplankton biomass, causing a shift in community structure toward smaller phytoplankton which can have bottom-up effects on food and energy availability at higher trophic levels (Zaba & Rudnick, 2016).


Pinniped researchers have found that body-size declines in aquatic ecosystems is a universal response to climate change. Body size directly affects energy and water requirements for thermoregulation, energy, mass acquisition and utilization rates and life-history characteristics (Daufresne et al., 2009). For example, Galapagos fur seals from Fernandina Island, in Galapagos Archipelago, that are significantly smaller than sympatric Galapagos sea lions exhibited lower foraging effort when compared. Although sympatric, fur seals are less successful; it is possible that their low plasticity in foraging behavior, expressed by their lower individual variability, has contributed to their overall smaller population size. Compared to the high plasticity in sea lions, fur seals are more impacted by variations in prey abundance, such as during ENSO events (Villegas-Amtmann et al., 2013). Oceanographic processes are changing as a result of climate change and direct anthropogenic influence, altering features of the current pelagic habitat. This, in synergy with competition for prey with fisheries can severely hamper future recoveries of otariids after future ENSO events.



## 2.10 *Punta San Juan study site*

Within the HCS there are few sites where otariids are found in important sympatric colonies. Punta San Juan (PSJ, 15°22'S, 75°12'W), a peninsula on the southern coast of Peru, is one such site. PSJ is a peninsula protected from land by a 1.2 km concrete wall that is part of a marine coastal national reserve system called 'Reserva Nacional Sistema de Islas, Islotes y Puntas Guaneras' (RNSIIPG, Spanish acronym). The PSJ peninsula consists of 20 sites, which are beaches and coves all used for breeding and/or as haul-out sites by SASL and SAFS (Figure I.6). PSJ supports important breeding rookeries of both species with an annual range of 6-12,000 SASL and 2-5,000 SAFS over the past 5 years (Cárdenas-Alayza et al., 2021). In PSJ, SAFS and SASL colonies are found year round, making it an ideal setting for interspecific comparison studies. In PSJ operates the Punta San Juan Program, a unique long-term research program that monitors otariid populations *in-situ* for over 20 years. This has permitted the routine collection of scats and counts as well as research projects that include instrumentation, morphometrics and whisker collection of pinnipeds, analyzed for this doctoral thesis project.





**Figure I. 6** Location of Punta San Juan peninsula (left) on the south coast of Peru, South America (right). Image modified from S. Cárdenas-Alayza, 2012.

### 3. Thesis Objectives


The main goal of this thesis is to describe and compare the ecological niches of sympatric otariid species *Arctocephalus australis* and *Otaria byronia* that coexist in Punta San Juan; and determine intraspecific and interspecific niche differentiation/overlap. This will be addressed through three specific goals. The first, is to describe and compare the trophic ecology of sympatric pinnipeds according to species and sex. The second is to describe and compare the environmental variables that characterize ecological niches between the sympatric pinnipeds according to species and sex. Finally, the third objective is to describe and compare the foraging strategies (spatial and temporal dimensions) within the ecological niches of the sympatric pinnipeds according to species and sex.

### 4. Structure of the Thesis

To contextualize the current state of the study species, population trends in abundance and biomass of sympatric study species are analyzed to understand the potential interspecific competition at PSJ. To address trophic ecology, the coupling of traditional diet information (e.g., hard parts recovered in scats), foraging location data and stable isotopes information are explored with the aim to better



understand the use of the marine environment by male and female SAFS and SASL, at different resolutions. As part of each analysis, descriptions of the environment are included that help characterize the conditions for each trophic dataset. In sum, the current study proposes to combine a suite of datasets, divided into four data chapters, in order to shed light into the foraging ecology of both species and sexes (Figure I.7) at different temporal and spatial scales to explain current population trends.



**Figure I. 7** Study species and groups included in this study: South American fur seal (*Arctocephalus australis*) A) adult female with pup; B) territorial male and South American sea lion (*Otaria byronia*) C) adult female with pup; D) sub-adult males. Images provided by Punta San Juan Program, CSA-UPCH.



## **Chapter 1 – Trends in sympatric pinniped populations suggest resource limitations in the Peruvian Humboldt Current System**

I evaluate population trajectories of Peruvian sympatric otariids and discuss mechanisms for competition and/or resource limitation. In this chapter, I analyze population trajectories of SASL and SAFS in a sympatric breeding site in Punta San Juan, Peru between 2001-2019. By using a combination of abundance time series, age-class proportions and biomass time series within and between species. This is considered as a first step towards exploring if larger body mass (SASL), a proxy for higher consumption of prey resources, can offer a greater competitive advantage in comparison to a smaller species (SAFS) to sustain population abundance in a common environmental setting.

## **Chapter 2 – Resource partitioning as a mechanism for trophic segregation in sympatric otariids from the productive upwelling Peruvian Humboldt Current System**

To understand if resource partitioning contributes towards sympatric breeding of otariids in Peru's dynamic HCS I compared seasonal diet composition SAFS and SASL, over 15 seasons between 2015 and 2018 which included ENSO and non-ENSO conditions. The objective of this research is to assess if resource partitioning exists in the sympatric otariid populations at PSJ over a four year study period. My hypothesis is that for sympatry to occur both otariids species should exploit food resources differently, being resource partitioning a potential mechanism to avoid competitive exclusion. Furthermore, this chapter updates knowledge on resource consumption by each species and contribute towards the understanding of the current foraging strategies of sympatric otariids in the highly productive and dynamic HCS.

## **Chapter 3 - Multiple strategies for segregation during foraging in sympatric otariids of the Peruvian upwelling Humboldt Current system**

I analyzed locations of 18 SASL and 17 SAFS (12 females, 23 males) equipped with satellite tags in Punta San Juan, Peru during 2013-2017. I determine if temporal and spatial segregation among species-sex groups is influenced by morphological (body mass), behavioral or environmental conditions during foraging. Due to larger body mass in SASL, sexual dimorphism and different




parental care, I expect SASL will travel longer distances, duration and cover larger areas than SAFS; and that males of both species will also travel longer distances, duration and cover larger areas in comparison to females. Tracks were constructed, analyzed and behaviors categorized to understand when individuals were actively exploiting a prey patch with the ultimate goal to understand spatial and temporal overlap and the associated environmental conditions (Sea surface temperature, Chlorophyll-a, Distance from Coast, Proximity to Fronts and Front Gradient) that characterize foraging events identified at sea.

#### **Chapter 4 - Sympatric otariids increase trophic segregation in response to warming ocean conditions in Peruvian Humboldt Current System**

Recent ocean warming events (2014 – 2017) that can decrease and impoverish prey biomass have occurred in HCS. In this context, my aim was to assess the effect of warming events on long-term inter- and intra-specific niche segregation. Whisker samples from SAFS (55 females and 21 males) and SASL (14 females and 22 males) collected in Punta San Juan, Peru were used to determine interspecific isotopic niche segregation based on  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values organized at a monthly scale for 2005 – 2019. I determined and compared niche width of species-sex groups and their overlap to assess response to environmental changes.

## CHAPITRE 1

*Trends in sympatric otariid populations  
suggest resource limitations in the  
Peruvian Humboldt Current System*



**Cárdenas-Alayza S, Gutiérrez D, Tremblay Y. (2021)** Trends in sympatric otariid populations suggest resource limitations in the Peruvian Humboldt Current System. *Mar Environ Res*:105349.

<https://doi.org/10.1016/j.marenvres.2021.105349>





## Résumé

Les espèces sympatriques développent des mécanismes pour éviter la compétition et coexister. Dans le système du courant de Humboldt (HCS), les populations d'otaries sud-américaines (SASL, *Otaria byronia*) et d'otaries à fourrure sud-américaines (SAFS, *Arctocephalus australis*) fluctuent principalement en raison des événements ENSO et de la disponibilité des proies. Nous évaluons les trajectoires des populations d'otariidés sympatriques péruviens et discutons des mécanismes de compétition et/ou de limitation des ressources. À cette fin, nous avons analysé les trajectoires de population de SASL et SAFS dans un site de reproduction sympatrique à Punta San Juan, au Pérou, entre 2001-2019. L'analyse en ondelettes a été utilisée pour extraire les tendances et les dérivés afin d'estimer les taux et les points de retournement. Les proportions des classes d'âge et les séries chronologiques de biomasse ont été construites à partir de comptages hebdomadaires et évaluées. Les deux populations montrent une phase de croissance et un déclin ultérieur. SAFS a commencé le déclin  $\sim 2.25$  ans avant et à un taux 1.5 fois plus rapide que SASL. La diminution de la classe d'âge des juvéniles suggère que la limitation des ressources est le principal facteur contribuant au déclin actuel de la population. Les espèces sympatriques développent des mécanismes pour éviter la compétition et coexister. Dans le système du courant de Humboldt (HCS), les populations de lions de mer sud-américains (SASL, *Otaria byronia*) et d'otaries à fourrure sud-américaines (SAFS, *Arctocephalus australis*) fluctuent principalement en raison des événements ENSO et de la disponibilité des proies. Nous évaluons les trajectoires des populations d'otariidés sympatriques péruviens et discutons des mécanismes de compétition et/ou de limitation des ressources. À cette fin, nous avons analysé les trajectoires de populations sympatriques de SASL et SAFS à Punta San Juan, au Pérou, entre 2001-2019. L'analyse en ondelettes a été utilisée pour extraire les tendances et les dérivés afin d'estimer les taux et les points de retournement. Les proportions des classes d'âge et les séries chronologiques de biomasse ont été construites et évaluées à partir de comptages hebdomadaires. Les deux populations montrent une phase de croissance et un déclin ultérieur. SAFS a commencé le déclin  $\sim 2.25$  ans avant et à un taux 1.5 fois plus rapide que SASL. La diminution de la classe d'âge des juvéniles suggère que la limitation des ressources est le principal facteur contribuant au déclin actuel de la population.

**Mots-clés :** Pérou ; Punta San Juan ; dynamique de population ; sympatrie ; otarie à fourrure ; lion de mer ; *Otaria byronia* ; *Arctocephalus australis* ; pinniped



## Abstract

Sympatric species evolve mechanisms to avoid competition and coexist. In the Humboldt Current System (HCS), populations of South American sea lions (SASL, *Otaria byronia*) and South American fur seals (SAFS, *Arctocephalus australis*) fluctuate mostly due to ENSO events and prey availability. We evaluate population trajectories of Peruvian sympatric otariids and discuss mechanisms for competition and/or resource limitation. For this purpose, we analyzed population trajectories of SASL and SAFS in a sympatric breeding site in Punta San Juan, Peru between 2001-2019. Wavelet analysis was used to extract trends and derivatives to estimate rates and turning points. Age-class proportions and biomass times series were constructed from weekly counts and evaluated. Both populations show a growth phase and subsequent decline. SAFS started decline ~2.25 years before and at a rate 1.5 times faster than SASL. Decrease in juvenile age-class suggests that resource limitation is the main contributing factor for current population decline.

**Keywords:** Peru; Punta San Juan; population decline; competition; fur seal; sea lion; *Otaria byronia*; *Arctocephalus australis*; pinniped



## 1. Introduction

Understanding how species coexist is an important theme in ecology. Coexistence theory is a framework to understand how competitor traits can maintain species diversity and prevent competitive exclusion, even among similar species in ecologically similar habitats (Chesson & Kuang 2008). Competition can compromise the fitness of one of the species involved, and may ultimately lead to its exclusion (Pacala & Roughgarden, 1985). Under the resource-utilization niche concept, two species cannot occupy the same ecological niche without exerting strong competition on each other (Schoener 1974). The Lotka–Volterra competition model states that coexistence of two species is possible when the competitive effect that a species has on another species (interspecific competition) is less than the competitive effect that it has on its own species (intraspecific competition) (Chesson & Warner 1981, Schoener 1983, Chase et al. 2002). Furthermore, competitive interactions are known to be stronger between morphologically and phylogenetically closely related sympatric species (Schoener 1983, Loveridge & Macdonald 2003, Di Bitetti et al. 2009).

Otariids, composed of fur seals and sea lions, occur in sympatry in a number of upwelling regions in the world and have many similar life-history traits and foraging habits (Dellinger & Trillmich 1999, Bailleul et al. 2005, Page et al. 2005, Franco-Trecu et al. 2012, Waite et al. 2012b, Jeglinski et al. 2013, Villegas-Amtmann et al. 2013a). Coexisting species are expected to develop strategies for niche differentiation when resources become scarce, to mitigate the reduction in population attributes, such as growth, survival or fecundity (Begon et al. 2006). The duration of coexistence in sympatric otariid seals is not known, and it is possible that sympatry has resulted from recent population recoveries posterior to the period of commercial sealing (Arnould & Costa 2006). If sympatry is a result of recent events, it is possible that divergence is not yet evident or that competitive exclusion is now occurring. In sympatric fur seals and sea lions, past research have found mixed results with regards to competition and segregation. Contrary to the situation in Peru, in many locations fur seal populations are typically larger and they appear to outcompete sea lions (Wickens & York 1997). Researchers studying sympatric otariids have found trophic segregation, differing foraging strategies and contrasting population trends as mechanisms that explain coexistence (Páez-Rosas et al. 2012, Franco-Trecu et al. 2014, Pablo-Rodríguez et al. 2016). More recent studies demonstrate that intraspecific segregation, driven by interindividual specialization, takes place in colonial species





(such as otariids) that are constrained to limited foraging ranges that overlap with conspecifics in similar niches (Jeglinski et al. 2013, Villegas-Amtmann et al. 2013b, Schwarz et al. 2021).

Some authors suggest that trophic segregation between sea lions and fur seals is associated with the observed divergent trends in epipelagic and benthic foraging characteristic of each group (Arnould & Costa 2006) and argue that body size is a central feature that characterizes foraging modes between larger sized benthic foraging sea lions and smaller epipelagic foraging fur seals. In those studies, sea lions showed greater oxygen storage capabilities; enabling them to aerobically dive longer and deeper resulting in higher metabolic and reproductive outputs (Costa et al. 1998). However, other studies have also demonstrated that sea lions of different age-classes (Jeglinski et al. 2013, Villegas-Amtmann et al. 2013b) and within the same age-class (Schwarz et al. 2021) display specialized types of benthic, pelagic and nocturnal epipelagic foraging. Other studies have shown that foraging modes can vary regionally in the same species, therefore demonstrating the importance of prey availability as a driver for foraging mode (Hückstädt et al. 2016).

The Humboldt Current System (HCS) in Peru is recognized as the most productive of the Eastern Boundary Upwelling Systems in the world (Bakun & Weeks 2008). As other upwelling ecosystems, cold nutrient rich water rises to the euphotic layer by a combination of the South east trade winds and the Coriolis force that fuel Ekman transport, giving rise to primary productivity in the upper pelagic and nearshore areas. Peru's coast is characterized by a narrow continental shelf, shallow productive upwelling waters and a relatively shallow thermocline related to an upper minimum oxygen layer at similar depths of approximately 50-80m, on average (Echevin et al. 2008, Demarcq 2009, Bertrand et al. 2010). These features limit the potential habitat for otariid prey to be linked to the benthos. Thus, in this productive shallow environment foraging on highly patchy, but dense prey resources near the surface, may preclude the advantages of searching for more evenly distributed, but less dense, prey resources on the benthos (Arnould & Costa 2006), which are also constrained by bottom water hypoxia.

Furthermore, HCS is characterized by recurring El Niño Southern Oscillation (ENSO) events, which can vary in intensity and duration. ENSO is known as a cycle of alternating warm El Niño and cold La Niña events and is the most prominent climate signal on Earth (McPhaden et al. 2006). When HCS is affected by an ENSO, increments of sea surface temperature and reduction of primary productivity




occur in the marine environment, altering distribution of habitats and associated prey biomass. The ENSO events that occurred in 1982-83 and 1997-98, were categorized as 'Extraordinary' and are well documented for dramatically altering species composition and reducing marine biomass (Barber & Chavez 1983, Tam et al. 2008). These climatic signals were strong and felt worldwide, changing the availability of prey items and causing declines in reproduction and survival of top marine predator populations around the globe (Arias-Schreiber & Rivas 1998, Forcada et al. 2006, Oliveira et al. 2006, Trathan et al. 2007, Oliveira 2011, Cárdenas-Alayza 2012, Bond & Lavers 2014, Sprogis et al. 2018). In Peru two species of otariids coexist that have breeding colonies along the coast, the South American sea lion (SASL) and South American fur seal (SAFS). Analysis of population trends of coexisting species helps understand population status and response to ecosystem conditions. Otariid surveys for the coast of Peru exist from mid 1960s, providing information on the abundance of these populations, but had heterogeneous sampling effort. Since the mid 1990s, systematic annual surveys are conducted at main colonies and during each species breeding season (February/March for SASL and November/December for SAFS) by scientists from Instituto del Mar del Peru (IMARPE) to monitor their status and fluctuations (Arias-Schreiber & Rivas 1998, Oliveira et al. 2006). Regardless, variability in Peru otariid continues to be observed (Figure 1.1). In the 1970s and early 1980s, historical records show a similar abundance for SASL and SAFS of ~ 20-25,000 individuals. However, in the late 1980s and 1990s SASL were more numerous than SAFS. Furthermore, a decline of 75% and 80% is documented for both SAFS and SASL, respectively due to the 1997-98 ENSO (Oliveira et al. 2006, 2012b). Surveys post ENSO reflect population recoveries for both species but with a larger SASL abundance. Similar fluctuations are observed until 2014, after which, there is a decline in SAFS.

Existence of sympatric colonies of SAFS and SASL implies that some mechanism currently operates (or has in the past), to permit interspecific coexistence in the HCS ecosystem. Sympatry of SAFS and SASL occur at only a few locations along the distribution ranges of our study species in the HCS. Punta San Juan (PSJ), a peninsula on the south coast of Peru is a sympatric site and is considered a historically important breeding center for otariids in Peru. The effects of the 1997-98 ENSO also impacted this breeding site, with local reductions of ~87% in both species. In PSJ, abundance declined from 2,751 to 345 in SAFS from November 1996 to November 1999 and from 12,415 to 1,495 in February 1997 to February 1999 in SASL (Arias-Schreiber & Rivas 1998). Since the decline, weekly



counts of otariids were conducted to monitor population recovery. Although high resolution time series of marine megafauna populations are scarce; at this location, weekly counts permit to study fluctuations in abundance, which can help understand interspecific and intraspecific interactions in a common ecological context. Thus, we consider that the long time series collected at PSJ (which accounts roughly for 10-50% of the total Peruvian populations in both species in this only site) provide an adequate setting to study the coexistence of SAFS and SASL and is representative of their dynamics in Peru.

The main goal of this study is to describe population trajectories over the last 20 years, in a sympatric otariid colony in Peru and secondly, to discuss mechanisms (e.g. competition and/or resource limitation), possibly explaining these trends. In terms of the population trajectories, we expect a first phase of fast population recovery post 1997-98 ENSO, followed by a subsequent stabilization in populations' size. Given the larger body size and population size of SASL on the coast of Peru we hypothesize that the abundance and biomass of SASL dominates over SAFS, in PSJ during our study period. By using a combination of abundance time series, age-class proportions and biomass time series within and between species, we consider this is a first step towards exploring if larger body mass (SASL), a proxy for higher consumption of prey resources, can offer a greater competitive advantage in comparison to a smaller species (SAFS) to sustain population abundance in a common environmental setting.



**Figure 1. 1** Time series of the total number South American sea lions (SASL, open circles) and South American fur seals (SAFS, filled diamonds) counted on the coast of Peru (top) and Punta San Juan study site (bottom) between 1965-2019. Dashed grey vertical lines delimit the population decline caused by the 1997-98 ENSO. Time series were constructed with official counts from surveys obtained from various sources (Arias-Schreiber & Rivas, 1998; IMARPE, 2012, 2013, 2014b, 2015, 2016, 2017, 2018; Majluf & Trillmich, 1981; Muck & Fuentes, 1987; SERNANP, 2016; Tovar & Fuentes, 1984).



## 2. Materials and Methods

### 2.1 *Study Site and Data Collection*

Punta San Juan (15°22'S, 75°12'W) is a peninsula protected from land by a 1.2 km concrete wall that is part of a marine coastal national reserve system called 'Reserva Nacional Sistema de Islas, Islotes y Puntas Guaneras' (RNSIIPG, Spanish acronym). The PSJ coastline consists of 20 sites, which are beaches and coves all used for breeding and/or as haul-out sites by SASL and SAFS. In PSJ operates the Punta San Juan Program, a long-term research program that monitors otariid and seabird populations *in-situ*. Access to the PSJ Program database was granted for purpose of this study. At PSJ, land-based counts of accessible otariid sites are conducted at least once per week since 2001 to monitor population dynamics. For this study we accessed the count data during years 2001 to 2019, which have a minimum frequency of once per week. Counts took place between 0600-0900 hours by observers with 10x50 binoculars and tally counters on top of 8 – 30 meter cliffs overlooking otariid sites. Accessible sites are counted for total abundance and age-class categories, for each species. Gaps in count data occurred when roosting seabirds made it impossible to access the cliff-edge to view beaches and coves. More frequent counts available during the breeding seasons (January thru March for SASL and October thru December for SAFS) at selected breeding sites, were also included to enhance time series. Due to the long-term nature of this dataset (19 years), counters changed over time and beaches were counted by different observers. Standard error of the mean count between three simultaneous observers ranges between 3-4% for SAFS and 3-13% for SASL. Satellite images with 2.8m resolution (image courtesy of CNES/Airbus 31 October 2019 via Google Earth) were used to estimate the area of the main breeding sites for each species. SAFS main breeding site was estimated through *in-situ* measurements collected during low tide in 1999 when animal abundance was zero and used to validate use of satellite images. Finally, a time series of mean monthly density was estimated during the breeding season (January – March for SASL and October – December for SAFS) by dividing mean monthly number of otariids by the area of the site where animals were counted on land and reported as number of individuals per square meter (ind /m<sup>2</sup>).



## 2.2 *Abundance time series*

In otariid breeding sites, the ratios of age-classes change throughout the year. Most evident shifts occur during the breeding season when territorial males, adult females and pups increase and subadult males and juveniles decrease. Breeding season for SAFS and SASL last for approximately 3 months (SAFS: October – December; SASL: January - March). Early in the breeding season males arrive to compete and hold territories; subsequently females arrive to give birth and there is rise in female and pup numbers. Females generally stay with the pup for 5-13 days, depending on the species. After females complete their post-partum fasting period, they will be at sea for varying periods and return to land to nurse the pup. Intra-annual dynamics in each species and age-class was examined (Appendix A1-A2). Due to the intra-annual dynamics due to breeding and non-breeding seasons as well as the long-term population fluctuations, we detrended the abundance time series with the goal to extract an annual trend of abundance for analysis and remove all the high-frequencies' signal in the data.

Total abundance for PSJ was estimated from the sum of all sites individually counted on a given date between 2001 and 2019, for each species. We verified that interannual dynamics of study sites was the same as dynamic of the sum of all sites. Because counts varied on a daily to weekly scale, these were linearly interpolated to generate a time series with an equal sampling interval of every 3 days. To detrend the abundance time series and be able to extract the values we used a wavelet trend decomposition with a “morlet” mother wavelet, since it overcomes the problems of non-stationarity in a time series by performing a local time-scale decomposition of the signal. This method gives the possibility to investigate the temporal evolution of a times series with different rhythmic components. We identified the wavelength scale corresponding to one year using wavelet analysis in WaveletComp R package v. 1.1 to extract the trend of abundance for both species. To test the study hypothesis, the trend of abundance time series, density and ratios of abundance will be compared between both species. Increasing or decreasing phase of trends, were determined with first order derivatives. Turning points will permit to detect when the date of change in direction of a species population trajectory occurred. Difference of first order derivatives was used to understand which species was increasing over the other.



## 2.3 *Age class proportions*

Time series for the proportion of each age-class were constructed using counts available for the main breeding sites of PSJ for SASL and SAFS, respectively. For this, we selected data sets from the main breeding sites in PSJ that are exclusive to one of the two species year-round. Age-class counts from beach site S3 were collected for SAFS between November 2003 to December 2019 ( $n = 977$  observations). For SASL, there was a change in the beach site used to estimate age-classes in years 2012-2013 to improve the number of times the sea lion breeding site could be accessed over the year by observers. Therefore, we constructed a single time series combining counts from beach site S7.S8 for years 2003-2013 and beach N7 for years 2012-2019 ( $n = 666$  observations), including a mean proportion between the two sites for overlapping dates. The pattern in the proportions was inspected carefully to verify that age-classes did not vary abruptly due to site change.

Age-class categories for this study includes: pups, juveniles, adult females, subadult males and territorial males. These are based on the categories used in the field at PSJ for counting animals, based on distinguishable anatomical features and behaviors (Vaz-Ferreira and Ponce de León, 1987). A ‘juvenile’ category was created in this study to group the ‘yearling’ (1 year old) and ‘immature’ (2-5 years old) categories used in the field. To estimate age-class proportions we only used the dates that considered all the beforementioned categories. Median proportions and interquartile range (IQR) were estimated .

Mean mass  $\pm$  SD of otariids according to age classes were estimated using PSJ Program database. Mass was collected at PSJ between during otariid tagging and health assessment fieldwork that involved weighing live animals during the study period. Mass of adult age classes were collected during the breeding season while pupping was taking place. Meanwhile, pup mass was collected during pup round-ups, approximately 2 months after the peak of pupping occurs in each species (April for SASL and January for SAFS). Mass for the SAFS subadult male age-class was obtained from the PSJ Program necropsy records of individuals that stranded dead in PSJ due to interactions with longline fisheries, and is reported here. These animals were confirmed to have died from interaction with fishing gear, carcasses were in good body condition and therefore we consider their mass accurately represents their age-class. The only records of mass for dead juveniles available in PSJ were in emaciated body condition, probably due to nutritional stress, thus we do not consider they





represent mass of healthy live individuals.

If values for the mean mass of an age-class were not available from PSJ, these were obtained from literature from conspecific or similar sized age-class otariids. We used reported mass of juveniles (Hückstädt et al. 2016); and territorial males (George-Nascimento et al., 1985) SASL collected in Chile. To our knowledge, no published values exist for mass of juvenile SAFS in the Pacific Ocean and in the Atlantic, studies either report captive animals or small sample sizes (<5) for an class that represents many ages and high variance. We therefore applied the reported mass of conspecific Northern fur seals (*Callorhinus ursinus*) from a study involving live captures and grouped mass of animals of ages 1-2, 2-3, 3-4 and 4-5 years old for the juvenile age class (Baker et al., 1994). To verify our hypothesis about body size, we tested for differences between mean mass of age-classes between species. To evaluate if there is a source of competition between age-classes during breeding and non-breeding season (e.g., overlap in the proportion of territorial males and adult females in non-breeding season) we tested for independence between the weighted age-class proportions (product of proportion and mean mass for each age-class) within each species and season. This was done by applying a t-test or Mann Whitney Wilcoxon Test ( $p < 0.05$ ) after verifying data distribution. Finally, we describe any shifts in the trends of age-class proportion time series to explain the mechanisms behind the population trajectory for each species.

## 2.4 *Biomass time series*

Biomass time series were constructed for each species in tons (t) by calculating the product of the abundance time series trend, the median age-class proportion and the mean mass per age-class. A time series for total otariid biomass was created from the sum of the SASL and SAFS biomass time series to determine the proportion of biomass that pertains to each species. To test our hypothesis, the percentages of each of the species biomass over time are used to describe interspecific competition and evaluate if SASL are dominating over SAFS. All analysis were done using R software version 3.6.2.





## 3. Results

### 3.1 *Abundance time series*

After removing high frequency signal from count data, annual trends of abundance were extracted for each species. This revealed population trajectories in PSJ that vary between a range of 829 - 4,550 individuals for SASL (Figure 1.2A) and 291 – 3,443 individuals for SAFS (Figure 1.2B) between 2001 to 2019. During this study period mean density of individuals during breeding was  $0.40 \pm 0.23$  ind /m<sup>2</sup> for SASL (January-March) and  $0.12 \pm 0.08$  ind /m<sup>2</sup> for SAFS (October-December).

Throughout the abundance time series, there is always a higher abundance of SASL in comparison to SAFS, with a median ratio of 1.65 SASL : 1 SAFS. The ratio of SASL to SAFS starts at a maximum ratio of 2.84 SASL : 1 SAFS, followed by decreasing phase that hits a minimum of 1.23 SASL : 1 SAFS in October 2012. Then, a final increase phase that rises until a ratio of 2.75 SASL : 1 SAFS in December 2019.


In general, both species started with a positive increasing phase and then both declined toward a negative phase, with differing dates at each turning point. Turning points were detected as the date when a change from positive to negative phases occurred. First, SAFS changed its phase from positive to negative in November 2014. Then, 2.25 years after (in March 2017), SASL changed its phase and started to decline. Linear regressions ( $p < 0.05$ ) were adjusted to standardized data of the (negative) declining phase for each species. Slopes were significantly different from zero in both species ( $R^2 = 0.99$ ,  $p < 0.05$ ); being slope incline of SAFS ( $B_0 = -179.35 \pm SE 0.21$ ) steeper than SASL slope ( $B_0 = -97.33 \pm SE 0.23$ ).

The difference between first orders derivatives describes which of the species had a higher growth rate, at a given time, in comparison to the other. We used these values to organize the time series into three different periods. Sea lions had a greater growth rate in comparison to SAFS, during the first period (January 2001 - June 2007) and in the third period (March 2012 - December 2019). SAFS had a stronger growth rate in comparison to SASL during the second period (June 2007 - March 2012).



### 3.2 *Age-class proportions*

Median age-class proportions (Median, IQR) and mean mass (Mean  $\pm$  SD) estimated for the entire time series are reported in Table 1. Due to the non-normal distribution of datasets, we applied the Mann Whitney Wilcoxon Test ( $p < 0.05$ ) with Bonferroni p-adjustment. Significant differences were found between mean mass of all age-class proportions within species and between species ( $p < 0.05$ ), with SASL being of a significant larger size (x1.5 larger) than SAFS (Table 1). Between the adult sexes, territorial male SAFS are 2.23 larger than adult female SAFS and territorial male SASL are 3.74 times larger than adult female SASL.




**Figure 1. 2** Top panel shows 3-day interpolated counts (black line) with trend of abundance (red line) for A) SAFS and B) SASL collected in Punta San Juan, Peru. Bottom panel shows wavelet spectrum with horizontal line indicating the annual signal at 365 days for C) SAFS and D) SASL abundance time series.

**Table 1. 1.** Median proportion of age-class per species with inter-quartile range (IQR). Mean mass and standard deviation ( $\pm$  SD) in kilograms. Sample size and source used for each age-class.

	Age-class	Proportion of age-class		Mass of age-class (kg)		
		Median	IQR	Mean $\pm$ SD	<i>n</i>	Source
<b>SASL</b>	Pups	0.16	0.18	13.98 $\pm$ 0.62	77	This study
	Juveniles	0.23	0.17	78.88 $\pm$ 17.68	9	Hückstädt et al 2016
	Adult females	0.38	0.19	82.74 $\pm$ 11.31	23	This study
	Subadult males	0.12	0.16	173.78 $\pm$ 40.96	44	This study
	Territorial males	0.05	0.08	310	-	George-Nascimento et al 1985
<b>SAFS</b>	Pups	0.36	0.13	8.75 $\pm$ 0.65	171	This study
	Juveniles	0.26	0.10	24.88 $\pm$ 14.4	5606	Baker et al 1994
	Adult females	0.33	0.09	48.56 $\pm$ 8.08	91	This study
	Subadult males	0.03	0.02	66.53 $\pm$ 8.56	6	This study
	Territorial males	0.04	0.03	108.64 $\pm$ 14.20	26	This study



Significant differences were found between weighted age-class proportions (product of mean mass and number of individuals) of each species between breeding and non-breeding seasons (Mann Whitney Wilcoxon Test,  $p < 0.05$ ), during the final period of the study (March 2012 - December 2019) (Figure 1.3), suggesting low intraspecific competition between age-classes at this scale.



**Figure 1. 3.** Boxplots of the weighted proportion (product of mean mass and number of individuals) of each age-class for SAFS and SASL during breeding season (BS, white) and non-breeding season (non BS, grey) during March 2012 - December 2019.




Time series of age-class proportions with smooth loess curve and 95% CI were compared during November 2003 – December 2019 (Figure 1.4). Evident shifts in proportion of age-classes include i) increase in adult females (+0.2 SASL and +0.1 SAFS) between 2016 - 2018; ii) decrease in juveniles (-0.40 SASL and -0.30 SAFS) from 2015 – 2019; iii) decrease in sub adult males (-0.10 SASL and -0.05 SAFS) between 2007 - 2015; iv) decrease in territorial males (-0.10 SASL) between 2008 - 2013. All the other ages class proportions are relatively stable. These trends suggest that the population decline in recent years in both species is associated to a decrease in the juvenile and, to a lesser extent, the sub-adult male age-classes at PSJ.

### ***3.3 Biomass time series***


Total otariid biomass time series was in the range of 83.16 – 506.72 t with a mean of  $335.26 \pm 139.95$  t and reached its maxima in March of 2016. Throughout the time series 82.61 %  $\pm$  3.75 of total biomass is composed on average by SASL (range: 77.72 - 88.97 %), and 17.39 %  $\pm$  3.75 by SAFS (range: 11.03 – 22.28 %). This trend continues until the end of the study period, with 88.63 % of otariid biomass comprised by SASL and 11.36% by SAFS in the end of 2019. SASL occupy a biomass in the range of 73.99 – 406.25 t with a mean  $272.97 \pm 108.61$  t, reaching its maxima in March 2017. Whereas SAFS occupy a biomass of 9.17 – 108.41 t with a mean of  $62.30 \pm 33.21$  t, reaching its maxima in November 2014 (Figure 1.5).





**Figure 1. 4** Time series of the proportion of age-classes for SAFS (green) and SASL (orange) constructed from weekly counts (circles) with smooth curve (line) with 95% CI (grey area). Age-class counts collected at Punta San Juan otariid breeding sites between November 2003 - December 2019.





**Figure 1. 5** Stacked area chart of constructed biomass time series in tons (t) estimated from otariid abundance in Punta San Juan for SAFS (light gray) and SASL (dark grey) during years 2001-2019.

## 4. Discussion

### 4.1 *Changes in abundance, density and population trajectories*

Since the effects of the Extraordinary ENSO in 1997-98 that caused otariid mortality and dispersal, it took SAFS 13 years and SASL 15 years to reach peak abundances similar to levels in 1980s and 1990s. Decomposition of time series of counts and biomass during years 2001-2019 shows that both species had a growth phase, a turning point and are currently in a declining phase. Our analyses reveal



that SAFS abundance began to decline  $\sim 2.25$  years before SASL and at a rate 1.5 times faster than SASL. The current phase of decline in SASL was unexpected and was only detectable with an in-depth analysis of the population trajectories and associated variables. In this study we also found a decrease in the juvenile and (to a lesser extent) in subadult male age-class proportions in both SASL and SAFS that is contributing to the recent decline detected in both species (Figure 1.4). Decrease in juveniles and subadult males can be attributed to an increase in mortality or dispersal of these age-classes, exacerbating failure of recruitment in PSJ. Unfortunately, we do not have information to adequately address mortality or dispersal. Furthermore, in our study we did not find any evidence for intraspecific competition at population level (Table 1.1, Figure 1.3). However it is possible that intraspecific competition can be operating at another scale, such as in the overlap of resource utilization or in use of foraging grounds, that should be further explored.

The trend in biomass in this study between species shows that, on average, 83% of all otariid biomass is composed by SASL and 17% by SAFS, making SASL the current ‘dominating’ species, as hypothesized. In the final phase of the time series 89% of total otariid biomass belongs to SASL, marking a large difference in the current composition between the two sympatric otariid species that inhabit the PSJ study site. Total SASL abundance is 1.23 - 2.84 times larger in comparison to SAFS, with an median ratio of 1.65 throughout the entire time series. We think this can be related to the different mechanisms of congregation employed by each species, evidenced at PSJ. The aggregation of individuals on land is different when we examine raw counts of otariids at PSJ over time (Figure 1.2A-B). Following recolonization, raw counts show that SAFS increments with what appears to be an exponential rate at PSJ (Cárdenas-Alayza, 2012) that grows accumulating individuals at PSJ. Meanwhile, SASL shows pulse-like increments with seasonal peaks of abundance that can vary widely between years. However, as SAFS abundance increments plateau and then decrease, populations pulses of SASL have been up to four times the abundance of SAFS (Figure 1.2A-B). This variability in the recruitment of individuals in a given breeding season has played an important role in the population growth of SASL at PSJ.

A model for male dispersal proposed by Giardino et al. (2016) developed in the Atlantic population of SASL shows that male sea lions disperse between winter haulouts and breeding sites. Haulouts can be spatially segregated from central breeding areas during the non-breeding season, and are followed by male dispersal towards breeding sites in consecutive years. Males tend to have strong site fidelity



to both breeding and haul-out sites maintaining functional connectivity and gene flow throughout their range (Giardino et al., 2016). This same phenomenon contributes towards the recruitment of new individuals at breeding sites and can also be occurring in Peru, with PSJ being a reproductive site where individuals that haul-out at different sites congregate to breed. Furthermore, as evidenced from the raw counts and intra-annual dynamics (Appendix A2), SASLs leave the PSJ breeding rookery decreasing their abundance during non-breeding months. In contrast, SAFS female, juveniles and pups are found year-round at PSJ to complete extended lactation periods (up to 36 months), during which females, juveniles and pups show site fidelity towards lactation sites (Majluf, 1989). Thus, high dispersal rates in SASL seem to be key for augmenting population growth in a given breeding season permitting the recruitment of more individuals for breeding. However, the decline in juvenile and subadult age-classes suggests that PSJ is not a successful site for the recruitment of these age-class in both otariid species in recent years.

Otariid breeding sites tend to show seasonal age-class proportions that reflect the roles of age-classes and sexual segregation in their polygynous mating system (Bartholomew, 1970). In this study we found that PSJ breeding sites are occupied by different proportions of age-classes during breeding and non-breeding season in both species, with females, pups and juveniles being the most important age-classes throughout the year. Intra-annual dynamics show that from July until December, the abundance of all SASL age-classes decrease to very low and can reach zero abundance in PSJ. In contrast to SAFS, which is present year round, and has a slight decrease in July-August (Appendix A1). The wide variability in SASL in comparison to SAFS over the study period is promoted by the arrival of larger number of animals in a given season and the higher density at the breeding sites. Interestingly SAFS show a higher proportion of pups : female ratio than SASL (Figure 1.4), which suggests that female condition may be better in SAFS. However, this can be explained by the high site fidelity of breeding female SAFS at this site, where experienced breeders return to pup (Majluf, 1989). Whereas in SASL, it is possible that a bulk of the SASL females are young females.

In this study we found that SASL density ( $0.40 \pm 0.23$  ind /  $m^2$ ) at PSJ during breeding was 3-4 times SAFS density ( $0.12 \pm 0.08$  ind /  $m^2$ ). This can be explained by the difference in the social tactics intrinsic to each species. Soto and Trites (2011) reported a lek-like mating system in SASL with the highest female : male ratio recorded for this species in Ballestas Islands in Peru, about 250km from our study site. Large abundances of females congregate to breed at what some authors identify as



population “hotspots”. These high density aggregations have been explained as a means to enhance mate selection (Soto & Trites, 2011). Territorial males line up at the tide line to display themselves to reproductive females. Females benefit from male competition and avoid subadult male harassment while remaining in close proximity to foraging sites (Bradbury & Gibson, 1983; Soto & Trites, 2011). Although there are no studies on the SASL mating system at PSJ, we suppose that the social tactics described above also occurs and PSJ is an important site for SASL breeding (S. Cárdenas-Alayza, personal observation).

We know from historical records that the SAFS population ranged between 9,644 - 14,303 and SASL between 3,249 – 5,836 at the PSJ study site in 1978-79 (Figure 1.1). This shows that PSJ is able to support larger abundances of these species on land and in the surrounding marine environment. Interestingly, during the post 1997-98 ENSO recolonization process, SAFS gathered in lower densities than SASL. Pre ENSO density of SAFS at PSJ was reported between 0.5 - 1 ind /m<sup>2</sup> (Majluf, 1989), considerably higher than average current density of  $0.12 \pm 0.08$  ind / m<sup>2</sup> found in this study. In the 1980s and 1990s, 48% pup mortality was recorded in SAFS in PSJ. This was caused by maladaptive aggression of females towards pups, attributed to the high density aggregations (Harcourt, 1992a, 1992b; Majluf, 1992). Furthermore, SAFS currently occupies many more beaches for breeding in PSJ in lower density, in comparison to the pre 1997-98 ENSO era (Cárdenas-Alayza, 2007). Other studies have demonstrated that habitat selection for breeding sites by SAFS and SASL varies in Peru, and is driven by access to water or shade for thermoregulation (Stevens & Boness, 2003). Therefore, availability of suitable habitat features will also play a role in the occupation of breeding sites in a given area so that each species can efficiently operate mating tactics characteristic of species social system.



## 4.2 *Contributing factors of population fluctuations in Peruvian atariids*

All species of otariid seals throughout the world were once subject to extensive and, in most cases, excessive hunting pressure during the eighteenth, nineteenth and twentieth centuries. By the late 1800s, however, most species had acquired total legislative protection or were subject to only regulated managed harvests (Wickens & York, 1997). Despite this protection, populations of various sea lion species have experienced very little recovery, and in some cases, are declining, and fur seal species have generally experienced rapid population recovery rates (Costa et al., 2007; Wickens & York, 1997). In Peru, we have the opposite case, a smaller fur seal population and a more abundant sea lion population. The current lack of recovery of SAFS in Peru may be explained in part by little time for SAFS population to recover from historical commercial exploitation, in synergy with recurring ENSO events and competition with fisheries and other predators (Majluf, 1991; Oliveira, 2011).

In Peru, SAFS were under more exploitation pressure due to the interest driven by the higher value of their pelts in the international fur trade market. By early 1900s, fur seal colonies in Peru were reported as small and isolated, describing this species as “close to the brink of extinction” (Murphy, 1925). The first commercial ban for otariids was legislated in Peru in 1946, prohibiting culling of adult female and pup SAFS (Majluf & Reyes, 1989). In 1950s fur seal abundance continued to be low which resulted in an extension of this ban to include all SAFS age-classes and also female SASL. Exploitation of otariids was reactivated in 1967 until early 1970s. Finally, in 1976 commercial exploitation of both species was banned in Peru (Tovar & Fuentes, 1984).

In mid 1950s began the development of the industrial fishery for Peruvian anchovy (*Engraulis ringens*) in Peru, and has turned into the largest monospecific fishery in the world capturing between 4-12 million tons of fish annually (Chavez et al., 2008). Direct links have been found between anchoveta biomass and population abundance of other top marine predators in HCS, such as seabirds, that rely almost exclusively on this marine resource for growth, reproduction and survival (Barbraud et al., 2018; S. Bertrand et al., 2012; Passuni et al., 2016). In synergy with ENSO conditions, overfishing can have acute effects on the survival of predator populations in HCS (Oliveira, 2011).



After the effects of the 1997-98 ENSO, a mere abundance of 6,257 SAFS was recorded on the coast of Peru from a previous count of 24,481 in 1996 (Arias-Schreiber & Rivas, 1998). After some time, nationwide 2006 survey showed that the population had recovered to an abundance of 15,137 SAFS (IMARPE, 2006). However, the SAFS population recovery was also limited geographically. Of the entire current abundance, 80% resides at five sites on the entire coast of Peru (IMARPE, 2013). In comparison, 44,725 SASL were counted in Peru in 1999 after the 1997-98 ENSO, from a previous abundance of 143,713 in early 1997 (Arias-Schreiber & Rivas, 1998; Oliveira, 2011). In 2006, recovery of SASL was documented with a count of 118,220 (IMARPE, 2006). In these years, SASL, breeding colonies and haulouts were already widespread along the coast compared to SAFS. For both species, authors suggest that recovery is due to the re-establishment of prey populations, the lack of strong ENSO events and migration from individuals in the colonies in northern Chile (Cárdenas-Alayza, 2012; Oliveira et al., 2012), but the fact that SASL was not reduced to such a small abundance seems fundamental to their recovery potential.

Our results show that SAFS recently declined at a rate 1.5 faster than SASL. The steeper decline in SAFS, is possibly linked to reduced recruitment in breeding adults at PSJ in recent years in combination with juvenile dispersal and mortality from this site. In our study, the sympatric otariids are exposed to the same environmental conditions, reason why the difference in this response can also be linked to intrinsic species traits that respond differently to the changes in the environment. A populations risk of extinction will depend both on the deterministic (harvest, commercial exploitation, natural selection, etc.) and stochastic (environmental, genetic, etc.) processes. The latter are more pronounced at small population sizes and can accelerate extinction process started by deterministic factors (Palstra & Ruzzante, 2008). Previous studies have shown that the effective population ( $N_e$ ) is used as a tool to evaluate viable populations. High  $N_e$  values retain high levels of genetic diversity and reduce probability of effects from inbreeding depression (Oliveira et al., 2006, 2012; Wright, 1990). Whereas a population with a low  $N_e$ , is susceptible to genetic drift and less apt to respond to selection (Oliveira, 2011).

After the 1997-98 ENSO, SAFS have a critical  $N_e$  value of 2,153 (Oliveira et al., 2006) while SASL  $N_e$  is 7,715 (Oliveira et al., 2012). According to Oliveira et al (2009) population reductions after the 1997-98 ENSO, may have caused a significant genetic bottleneck in the Pacific population of SAFS and SASL, with loss of genetic diversity and presumably some loss of evolutionary potential for the



species (Oliveira et al., 2009, 2012). Furthermore, the rate of loss of genetic diversity via genetic drift is greater in populations where  $N_e$  is small and migration is limited, as is the case for SAFS. This can be evidenced in the information gathered through Peru's nationwide surveys, where >90% of SAFS breeding population is mostly restricted to latitude 15 -17° South, with PSJ supporting at least 40% of entire Peru SAFS population. Meanwhile, SASL population is distributed continuously along the coast, with important breeding sites in several latitudes (6, 13-14, 15 and 17° South) (IMARPE, 2016, 2017, 2018). Wider distribution of breeding sites with high density aggregations of SASL, suggest that migration and dispersal rates in SASL are higher than SAFS, resulting in higher connectivity. Therefore, even if there was genetic loss due to a bottleneck from a strong ENSO event, SASL has mechanisms to counterbalance loss of genetic diversity with temporal gene flow through migration. The difference in gene flow between these two species in HCS can be a key factor explaining the limited response to achieve population growth under scenarios of environmental change in SAFS in comparison to SASL suggested in this study, and should be further studied.

### ***4.3 Natural history traits that influence interspecific competition in otariids***

Body size of SASL is overall 1.5 larger in all age-classes compared to SAFS. In this study, although we hypothesize that a larger body size (SASL) could serve as advantageous to consume larger/ more prey, the decline in population abundance of both SAFS and SASL suggests that a resource limitation is affecting trajectories of both species. Between the sexes, territorial male SAFS are 2.23 larger than adult females and SASL males are 3.74 times larger than adult SASL females. Although at population level we did not find evidence for segregation between age-classes and sex, mechanism to avoid competition may be operating at a different scale. To elucidate differences and how these species are exploiting the environment, further studies need to be conducted to test intra and interspecific trophic niche segregation to understand how these sympatric otariid species exploit the HCS.

In our study we compared otariid biomass as a possible outcome of interspecific competition in PSJ and found that 83 % of total average biomass is composed by SASL throughout our time series. This can be explained by a the difference of 1.6 - 3.3 in mass between SASL and SAFS; in combination with a high density mating system in SASL versus lower density aggregations in SAFS. Researchers





have examined potential competition between sympatric fur seal and sea lions at other locations and have found in some cases that trophic segregation is a mechanism to explain differing foraging strategies and contrasting population trends (Franco-Trecu et al., 2014; Pablo-Rodríguez et al., 2016; Páez-Rosas et al., 2012). Some authors suggest that difference in divergent trends is explained through epipelagic and benthic foraging, being the latter characteristic of sea lions that have larger body size and diving capacity (Arnould & Costa, 2006). Other studies have also demonstrated that different age-classes within the same sea lion population have specialized benthic and epipelagic foraging modes (Jeglinski et al., 2013; Villegas-Amtmann et al., 2013). Therefore, otariid foraging ecology in PSJ needs to be evaluated at interspecific and intraspecific level to understand how the different groups are exploiting the local environment.

In the coast off PSJ, the continental shelf is extremely narrow limiting the access to benthic habitat. Furthermore, previous studies in Chile have demonstrated that SASL are epipelagic foragers with mean dives of  $29.0 \pm 34.8$ m, when they are in an upwelling environment (Hückstädt et al., 2016). Therefore, we expect that competition for similar prey items may be taking place within the upper limit of the oxygen minimum zone off the Peru coast, which is usually located  $\sim 50$  m in depth (A. Bertrand et al., 2010) making the foraging habitat of pelagic prey readily available for both otariid species diving capacity, independent of differences in body size.

It is well documented that the HCS in Peru holds a world-leading monospecific forage fishery in terms of landings of Peruvian anchovy, sustaining  $>5$  million tons per year on average since 1960s (Bakun & Weeks, 2008; Chavez et al., 2008). In Peruvian otariids, the number of pups born, mean birth dates and anchovy biomass were found to have direct relationships in SASL and SAFS reproductive success, demonstrating that anchovy is a key resource in fulfilling the energetic demands of both species (Cárdenas-Alayza, 2012; Soto et al., 2004). Other studies have evaluated diet composition between these sympatric otariids in the decades of 1980s and late 1990s in south Peru, but conclusions for resource partitioning are not clear. Vásquez (1995) reported that otariid diets were both based on fish and cephalopods; with Peruvian anchovy (*Engraulis ringens*) and Jack Mackerel (*Trachurus murphyi*) as the main consumed species. Afterwards, Arias-Schreiber (2000) found that Peruvian anchovy, squids and a crustacean, the Red squat lobster (*Pleuroncodes monodon*), were ranked as important prey items shared by the sympatric species. Based on this, we understand that anchovy is an important prey shared by both otariids during these study periods, but mechanisms for



trophic segregation are not clearly explained. Furthermore, prey availability may have changed in the last 20 years. Therefore, resource partitioning research needs to be updated and further explored in these sympatric species to disentangle current resource utilization in relation to population trajectories.


#### 4.4 *Current and future trends*

In the final 5 years of our time series, both sympatric otariid populations were found to be in a state of decline. SAFS were the first to decline in 2014, which coincides with a decrease in the anchovy biomass estimated for the coast of Peru during this period. Anchovy biomass dropped from 11,201 tons in 2013 to 3,799 tons in 2014 (Zavala et al., 2019). In 2014 anchovy landings summed 1,302 tons, leaving an available biomass of 2,497 tons for all predators and other fisheries to consume, which may have posed limitations for growth and maintenance of predator populations. Warmer oceanographic conditions in HCS persisted and coincided with a reduced anchovy biomass from 2014 to 2017 (Zavala et al., 2019). Changes in oceanographic conditions can shrink and expand available habitat for prey items forcing new competition scenarios between coexisting species. Thus, under a scenario of reduced anchovy biomass we consider it is possible that prey availability is limiting population sizes at PSJ and can be responsible for the declines observed in the juvenile and subadult age-class for both SAFS and SASL; as younger individuals do not possess full diving capabilities.

Predictions for the future of HCS include scenarios with shoaling of the oxycline (Brochier et al., 2013), compressing the pelagic habitat that sustains the abundant biomass of small pelagic fish currently in the system (A. Bertrand et al., 2011). If this occurs, it will increase competition between all the predators that depend on pelagic resources, unless they adapt foraging strategies under new conditions. Under this hypothetical predicted scenario and our findings, we suspect SASL will be the more resilient of the two sympatric otariid species that will thrive in HCS, given its population recovery in the past 20 years. Therefore, to propose adequate management for the future of these species, it is important to better understand current underlying foraging strategies used by otariids of HCS and determine if trophic competition rules over space and time between species. These studies are key to tease apart the underlying mechanisms behind the current population trajectories revealed by the present study.




# APPENDIX A1



**Figure A. 1** Multi-panel plot of South American fur seals counts by age-class according to day of year collected at the main breeding site S3 in Punta San Juan, Peru. Classes include: Adult Females, Juveniles, Pups, Subadult Males and Territorial Males. Total is the sum of all the age-classes. Blue line is a fitted loess model to depict the trend in abundance throughout the year. Data points include all study years available for site.




## APPENDIX A2



**Figure A. 2** Multi-panel plot of South American sea lions counts by age-class according to day of year collected at the main breeding site N7 in Punta San Juan, Peru. Classes include: Adult Females, Juveniles, Pups, Subadult Males and Territorial Males. Total is the sum of all the age-classes. Blue line is a fitted loess model to depict the trend in abundance throughout the year. Data points include all study years available for site.

## CHAPITRE 2


*Resource partitioning as a mechanism  
for trophic segregation in sympatric  
otariids from the productive upwelling  
Peruvian Humboldt Current System*



**Cárdenas-Alayza S, Torres, D.A., Gutiérrez D, Tremblay Y.** Resource partitioning as a mechanism for trophic segregation in sympatric otariids from the productive upwelling Peruvian Humboldt Current System.

Manuscript submitted for publication to Austral Ecology.






## Résumé

Le partage des ressources peut contribuer à la coexistence des espèces en réduisant la compétition trophique. Pour comprendre si le partage des ressources contribue à la reproduction sympatrique des otariidés dans le système dynamique du courant de Humboldt (HCS) au Pérou, nous avons comparé la composition saisonnière du régime alimentaire des otaries à fourrure sud-américaines (SAFS, *Arctocephalus australis*) et des lions de mer sud-américains (SASL, *Otaria byronia*) sur 15 saisons entre 2015 et 2018 pour déterminer s'il existe un mécanisme détectable de partage des ressources au niveau de la population. La composition du régime alimentaire a été estimée à partir de l'identification de restes durs de proies obtenus à partir de 3,325 féces collectés dans des colonies à Punta San Juan, au Pérou. Le régime alimentaire des SAFS est principalement composé d'anchois du Pérou (42,8 %) et de céphalopodes (34,7 %), avec de plus petites composantes de myctophidés (7,4 %), de galathées (5 %) et de 11 autres espèces de poissons. Le régime alimentaire de la SASL est principalement composé de galathées (69,9 %) et d'anchois du Pérou (19,1 %), avec de plus petites quantités de céphalopodes (4,7 %), de capucettes (4,6 %) et de cinq autres espèces de poissons. Dans cette étude, les deux prédateurs ont été classés comme spécialistes et une ségrégation trophique interspécifique a été trouvée dans 13 des 15 saisons échantillonnées. Un chevauchement trophique a été détecté au cours de l'été austral et de l'automne 2015 pendant un événement El Niño extraordinaire, au cours duquel les otariidés ont consommé des proportions similaires d'anchois du Pérou (55% SAFS, 48,6% SASL), de céphalopodes (21,7% SAFS, 13,6% SASL) et de galathées (9% SAFS, 2% SASL). En conclusion, l'analyse démontre un partitionnement des ressources basé sur l'abondance des proies primaires consommées par deux espèces d'otariidés et aide à expliquer la ségrégation trophique chez les otariidés dans le courant du Humboldt. En outre, les résultats montrent que les changements climatiques peuvent déclencher une réduction brutale des proies et exacerber la compétition interspécifique.

**Mots clés :** écologie de la recherche de nourriture ; régime alimentaire ; otarie à fourrure ; lion de mer ; Pérou ; ségrégation trophique




## Abstract

Resource partitioning can contribute to species coexistence by reducing interspecific competition. To understand if resource partitioning contributes towards sympatric breeding of otariids in Peru's dynamic Humboldt Current System (HCS), we compared seasonal diet composition of South American fur seals (SAFS, *Arctocephalus australis*) and South American sea lions (SASL, *Otaria byronia*) over 15 seasons between 2015 and 2018 to determine if there is a detectable mechanism for resource partitioning at population level. Diet composition was estimated from identification of hard remains of prey items obtained from 3,325 scats collected from colonies at Punta San Juan, Peru. SAFS diet is primarily composed of Peruvian anchovy (42.8%) and cephalopods (34.7%); with smaller components of myctophids (7.4%), red squat lobsters (5%), and 11 other fish species. SASL diet is mainly composed of red squat lobster (69.9%) and Peruvian anchovy (19.1%); with smaller amounts of cephalopods (4.7%), silversides (4.6%), and five other fish species. In this study, both predators were classified as specialists and interspecific trophic segregation was found in 13 of the 15 seasons sampled. Trophic overlap was detected in the austral summer and fall of 2015 during an extraordinary El Niño event, in which otariids consumed similar proportions of Peruvian anchovy (55% SAFS, 48.6% SASL), cephalopods (21.7% SAFS, 13.6% SASL), and red squat lobsters (9% SAFS, 2% SASL). In conclusion, the analysis demonstrates resource partitioning based on the abundance of primary prey items consumed by two otariid species and helps explain trophic segregation in otariids of HCS. Furthermore, results show that climatic changes can trigger abrupt reduction of prey items and exacerbate interspecific competition.

**Keywords:** foraging ecology; diet; fur seal; sea lion; Peru






## 1. Introduction

The ways in which species partition available resources is a major determinant of ecological communities and coexistence. The competitive exclusion principle states that two species in the same habitat cannot have exactly the same niche and coexist in stable form (Begon et al., 2006). Thus, coexisting species must differ in their ecological requirements by at least some minimal amount to avoid competitive exclusion (Pianka, 1974). Otariids (fur seals and sea lions) occur in sympatry in many locations around the globe. Research on sympatric otariids have found differing mechanisms of trophic segregation that explain coexistence (Franco-Trecu et al., 2014; Pablo-Rodríguez et al., 2016; Páez-Rosas et al., 2012). For instance, trophic segregation among otariids has been associated with divergent trends in epipelagic and benthic foraging modes (Arnould & Costa, 2006). Larger body size in sea lions has been linked to the consumption of large demersal fish and cephalopods in the coastal realm, which are accessible by benthic foraging and increased diving time (Costa & Gales, 2003; S. L. Fowler et al., 2006). In contrast, fur seals are often epipelagic feeders that forage on smaller prey resources with highly patchy distributions near the surface (e.g., zooplankton, small fish, or squid) (Arnould & Costa, 2006).


In the Peruvian Humboldt Current System (HCS) two species of otariids coexist, the South American fur seal (SAFS, *Arctocephalus australis*) and the South American sea lions (SASL, *Otaria byronia*). Existence of sympatric colonies of SAFS and SASL implies that a mechanism exists that permits their interspecific coexistence in the HCS ecosystem. Within the HCS there are few sites where otariids are found in important sympatric colonies. Punta San Juan (PSJ, 15°22'S, 75°12'W), a peninsula on the southern coast of Peru, is one such site. PSJ supports important breeding rookeries of both species with an annual range of 6-12,000 SASL and 2-5,000 SAFS over the past 5 years (Cárdenas-Alayza et al., 2021). In PSJ, SAFS and SASL colonies are found year round. In SAFS, territorial males and adult females increment during breeding months (October to December) and pups are born. In SASL, territorial males, adult females and pups increase during breeding months (January to March). In both species, numbers decrease during the remainder of the year, during which females, pups and juveniles make up the bulk of the age-classes during non-breeding months (Cárdenas-Alayza et al., 2021). During breeding, females arrive pregnant and all viable births to newborn pups occur in a lapse of 2-3 months (Cárdenas-Alayza, 2012; Soto, 2004). After pupping, females remain on land with the pup for 5-13 days and then begin alternating cycles of 2-4 days



foraging at sea followed by 2-3 days of nursing on shore (Ganoza, 2016; Majluf, 1987; Soto, 2004). Long lactation that can last up to 36 months as a mechanism to enhance offspring survival during variable oceanographic conditions have been reported in both species in Peruvian HCS (Majluf, 1987; Soto et al., 2004). As in other central place breeding otariids, females are energetically constrained to proximate foraging grounds for nursing and offspring survival in contrast to males that can move freely between different rookeries and foraging grounds (González-Suárez & Cassini, 2014; Hückstädt et al., 2014; Oliveira et al., 2017).

The Peruvian HCS is recognized as the world's most productive upwelling system in terms of fish biomass (Bakun & Weeks, 2008). The coastline is characterized by a narrow continental shelf and a relatively shallow thermocline related to the upper boundary of the oxygen minimum layer at approximately 50-80m (A. Bertrand et al., 2010; Demarcq, 2009; Echevin et al., 2008). These features limit the habitat for prey biomass of small pelagic fish such as the Peruvian anchovy (*Engraulis ringens*), making them available in high densities for marine predators (A. Bertrand et al., 2011). However, productivity in the HCS decreases when warming events occur, such as the El Niño Southern Oscillation (ENSO) or Kelvin waves, which can alter food web composition (Tam et al., 2008). These changes impact the success of predator foraging events and, depending on their intensity and duration, can decrease reproductive output and survival rates in top predators (Majluf, 1991; Trillmich & Dellinger, 1991). Extraordinary ENSO events have triggered significant changes in the food web composition causing nutritional stress and mortality in many species (A. L. Bond & Lavers, 2014; Forcada et al., 2006; Oliveira, 2011; Sprogis et al., 2018). For example, the 1997-98 ENSO event caused a reduction of wildlife along Peru's coast with 75% and 80% reductions in the populations of SAFS and SASL, respectively (Oliveira et al., 2006, 2012). Since 1998 populations have recovered to an extent, but are currently in a state of decline (Cárdenas-Alayza et al., 2021).

Identification of hard remains from scat samples is the most widely used technique to identify resource utilization in marine vertebrates. Although the technique has many biases, its wide use permits making interesting comparisons over time. The majority of previous research on the diet composition of otariids within HCS has focused of SAFS and SALS independently (Majluf, 1987; Paredes & Arias-Schreiber, 1999; Sarmiento-Devia et al., 2020; Zavalaga et al., 1998). However, some few studies have evaluated diet composition between SAFS and SASL concurrently in Peru in the 1980s and late 1990s. The first study conducted by Vásquez (1995), in late 1980s reports otariid diets were based on fish and cephalopods; with the Peruvian anchovy as the main species consumed by both pinnipeds followed by Jack Mackerel (*Trachurus murphyi*). One decade later, Arias-




Schreiber (2003) confirms that Peruvian anchovy and squid were main prey items shared by both species, with also red squat lobsters (*Pleuroncodes monodon*) comprising an important part of the diet for SASL. Recent studies in northern Chile also confirms the red squat lobster as a dominating prey item in the diet of SASL, followed by Peruvian anchovy, and squid (Sarmiento-Devia et al., 2020; Sielfeld et al., 2018). However, these recent studies argue that consumption of crustaceans by SASL is influenced by changes in the food web due to ENSO events, and that this resource reflects a coping mechanism when Peruvian anchovy is lacking (Sielfeld et al., 2018). However, there are no studies with consecutive sampling to test if this pattern is true.

Through long term monitoring and onsite research at PSJ, we were able to collect samples from 2015 to 2018, which included ENSO and non-ENSO conditions. The objective of this research was to assess if resource partitioning exists in the resident sympatric otariid populations of SAFS and SASL at PSJ over a four year study period. Our hypothesis is that for sympatry to occur both otariids species exploit food resources differently, being resource partitioning the mechanism to avoid competitive exclusion. Furthermore, we expect to update current patterns of resource consumption by each species and contribute towards the understanding of the current foraging strategies of sympatric otariids in the highly productive and dynamic HCS.

## 2. Materials and Methods

### 2.1 *Study area and sampling*


A total of 3,325 scat samples (1,484 SAFS and 1,841 SASL) were collected at Punta San Juan (PSJ) between January 2015 and December 2018 study period. PSJ (15°22'S, 75°11'W) is a 54 Ha peninsula in Peru with 20 beaches and coves occupied by SASL and SAFS. Samples were collected by personnel of the Punta San Juan Program, CSA, U. Peruana Cayetano Heredia under authorization of the RNSIIPG, (Reserva Nacional Sistema de Islas, Islotes y Puntas Guaneras) administered by SERNANP (Servicio Nacional de Areas Naturales Protegidas). Information and samples for this study were collected in the main breeding rookeries for each species that contain 40-60% of total PSJ population, (beach N7 for SASL and beach S3 for SAFS). During the study period, counts by age-class were carried out in each of the breeding rookeries 2-4 times per month through direct observation from cliff edges using 10x50 binoculars between 0600-0900 hours by specialised personnel. Counts during breeding season (BS, January – March in SASL and October – December



in SAFS) and non-breeding season (non BS, April - December in SASL and January - September in SAFS) were examined for the entire study period (see Appendix B1).

This research did not require invasive sample collection from live animals. Samples were collected at beaches exclusively used by each study species, making it impossible to confuse samples between species. All samples were collected from the ground on a given day of the month, leaving 21-31 days between samplings. Sample collection occurred between 1100 - 1500 hours; a time of day when otariids move towards the shoreline to thermoregulate; thus disturbance towards the colony is minimized. Samples were chosen based on their appearance and tenderness in aims to collect complete, recently defecated faeces. Samples were collected for 40 months for SAFS and for 35 months for SASL. No samples were collected in Spring 2016 for SASL. Samples were stored in plastic bags and labelled. Collected samples were soaked in soap water for 3 - 24 hours to aid dissolving faecal matter. Times varied according to concentration of organic matter in samples. Contents were passed through nested sieves of 1.00 mm and 0.297 mm mesh sizes. Hard remains included otoliths, cephalopod beaks, and crustacean remains, which were all carefully removed. Otoliths and cephalopod beaks that were eroded in >50% were discarded, following Tollit et al. 1997. Sagittal otoliths were identified to species level following García-Godos (2001). Right/left otoliths or upper/lower mandibles were not considered since frequency of occurrence and relative abundance indices (IIMP) were used for diet composition analyses. Cephalopod beaks were identified as one taxonomic group ("Cephalopods"). For crustaceans, only one species (*Pleuroncodes monodon*) was identified and counted using the cephalothorax. Fish otoliths were stored under dry conditions, cephalopod beaks were stored in 70° ethanol, and all crustacean cephalothorax were discarded. Hard parts were not measured.

In order to ensure that sample size is not biased and is sufficient to correctly represent the diversity of prey consumed, we calculated cumulative diversity in a given sampling season and verified the minimum number of samples required to reach an asymptote. If so, this means that additional samples do not modify the diversity index significantly, indicating a sufficient sample size (Ferry et al., 1997). This was evaluated by calculating the accumulated average and standard deviation of diversity curves generated from matrices of prey abundance data per season. We estimated the Shannon-Wiener Index ( $H'$ ) (Krebs, 1999) with 1,000 random permutations and considered an error of 5% (MATLAB program, v.2019b). Each curve is considered asymptotic if at least two previous values to sample trophic diversity ( $H'$ ) are in the range of  $H' \pm 0.05H'$ . This permutation routine was adapted from Ferry, Clark and Cailliet (1997) and has been applied to test sample size adequacy in other species



(Adams, 2004; Koen Alonso et al., 2002; Páez-Rosas & Aurióles-Gamboa, 2010; Porras-Peters et al., 2008).


## 2.2 Diet composition

The importance of prey was evaluated using the percent number (*PN*), percent frequency of occurrence (*FO*) and the percent Importance Index (*IIMP*). *PN* was used to describe otariid diets according to taxa (fish, cephalopods, and crustaceans), measuring the number of hard remains of taxon *i* in the total number of hard remains from all prey found in scat samples; for the entire study period and by sampling season. *FO* was used to describe the occurrence of a prey item in each of the otariids diet per season, and *IIMP* was used to estimate the importance of each prey item per sampling unit (scat) following García-Rodríguez (1999) and García-Rodríguez & Aurióles-Gamboa (2004). This measure estimates the probability of finding a given prey species in a scat sample, through the following equation:

### **Equation 1.**

$$IIMP_i = \frac{1}{U} \sum_{j=1}^u \frac{X_{ij}}{X_j}$$

Where  $X_{ij}$  is the number of occurrences of species *i* in scat *j*,  $X_j$  is the total number of identifiable structures in scat *j*, *u* is the number of scats in which the taxon *i* was found and *U* is the number of scat samples where prey items were found. This index generates results from 0 to 1 that is then converted into percentages ( $IIMP_i \times 100$ ). In this study, we will consider the prey items that contribute an  $IIMP \geq 1\%$  (García-Rodríguez & Aurióles-Gamboa, 2004; Koen Alonso et al., 2000). *IIMP* was developed for analysis of hard structures in pinniped scats and is used in this study as the main assessor for prey composition in this study because it is less sensitive to changes in the number of prey in an individual scat (Amador-Capitanachi et al., 2020; García-Rodríguez & Aurióles-Gamboa, 2004; Páez-Rosas & Aurióles-Gamboa, 2010). The *IIMP* allows each scat to contribute an equal amount of information, whereas *PN* can be dominated by a few scats with a large number of a single prey-taxon hard remains and *FO* does not reflect the abundance of prey items in the samples (García-Rodríguez & de la Cruz-Agüero, 2011). With *IIMP* values, ranked abundances were constructed to evaluate the composition of all prey items ranked by their contribution of abundance per each sampling. Also, prey items were classified in ecological groups according to their main habitat /



ecology extracted from FishBase (<http://www.fishbase.org>; Froese and Pauly, 2019). The ecological groups considered were: Pelagic, Demersal-Pelagic, Benthic-Demersal and Demersal, following Koen Alonso *et al.* (2001).

## 2.3 Trophic indices

Species Richness ( $S$ ) was estimated for each sampling season by pooling all years in a season and for each independent season and year. We assume that each faecal sample represents one individual in the rookery, and therefore scat sampling represents a random sample of the population. To determine whether individuals in a given sampling season demonstrate different degrees of feeding expertise, the standardized Levin  $B_i$  Index (Krebs, 1999) was estimated to distinguish if each species is a specialist or generalist consumer.

### **Equation 2.**

$$B_i = \frac{1}{n-1} \left\{ \left( \frac{1}{\sum IIMP_{ij}^2} \right) - 1 \right\}$$

Where  $B_i$  is the trophic breadth,  $IIMP_{ij}$  is the proportion of the prey item  $j$  in the diet of predator  $i$ , and  $n$  is the number of prey species in the diet. The standardized  $B_i$  is an index that generates results from 0 to 1. Predators are characterized as a specialist when values are less than 0.6 and as a generalist when values are  $\geq 0.6$ . Confidence intervals ( $\pm 95$ ) were estimated by applying  $\pm 1.96\sqrt{\sigma_{B_i}^2}$  (Krebs, 1999).


Trophic Level ( $TL$ ) was calculated for each species and season (Eq. 3) to identify the trophic level based on its diet composition and the trophic level of their prey (Christensen & Pauly, 1992).

### **Equation 3.**

$$TL = 1 + \left( \sum_{j=1}^n DC_{ij} \right) (TL_j)$$

Where  $TL_i$  is the prey's trophic level;  $DC_{ij}$  is the proportion of the prey  $i$  in the diet of the predator  $j$  and  $n$  is the total number of prey species in the diet. Trophic level values for prey species were extracted from FishBase (<http://www.fishbase.org>; Froese and Pauly, 2019). Cephalopod and crustacean values were extracted from literature (Espinoza *et al.*, 2017; Tam *et al.*, 2008). For cephalopods we used the  $TL$  value for *Loligo gahi*, since it is the only species identified in some of





the randomly selected beak squids in our samples and has been previously identified for these study species (Vásquez 1995; Arias-Schreiber 2003; Sielfeld et al. 2018; Sarmiento-Devia et al. 2020).


## 2.4 *Environmental conditions*

Monthly seasonal sea surface temperature (SST) records collected by the naval base at San Juan de Marcona bay (Dirección de Hidrografía y Navegación, DHN), located less than 5km from the otariid colonies, were used to characterize the environmental conditions during the study period. We constructed a time series of sea surface temperature anomalies (SSTA) by subtracting the monthly record to the long term mean for the complete time series (1976 - 2018) for each month. To compare environmental conditions with scat sampling seasons, we grouped SST and SSTA for austral summer (January - March), fall (April - June), winter (July - September) and spring (October - December), and estimated mean SST and SSTA values.

## 2.5 *Statistical analyses*

To determine differences in association of the composition of the prey items that were consumed per season and species, matrices with species (columns) and seasons (rows) using *IIMP* values were constructed. It is important to note that *IIMP* is already a form of standardized data because it weighs the contribution of each prey item according to the sample size. Since there are many zeros in the prey items matrix, it was first transformed with  $\log(n+1)$  prior to analysis. Bray-Curtis distances between 0-1 were estimated between the sampling seasons (Q-mode). Agglomerative hierarchical clustering was estimated using Ward's method (Ward & Hook, 1963) and p-values for each cluster were calculated with bootstrap resampling (1,000 iterations) using the *vegan* package in R statistical software (R Core Team, 2018). To interpret the position of sampling seasons and prey items between the two species in a reduced number of dimensions, we used Non-Metric Multidimensional Scaling (NMDS, stress < 0.5) on the standardized matrices. To test assumptions, normality was evaluated using Shapiro-Wilk's Test ( $p < 0.05$ ) and homogeneity of variance was checked with Levene's Test ( $p < 0.05$ ). Differences between groups were evaluated with ANOVA for parametric data and Kruskal-Wallis for non-parametric data ( $p < 0.05$ ). Tukey and Chi-square were used as post-hoc tests for multiple pairwise comparisons ( $p < 0.05$ ) for parametric and non-parametric data, respectively. To determine if there were correlation with environmental conditions, proportion of empty scats, *PN*






by taxa (fish, cephalopod and crustaceans), species richness ( $S$ ), trophic breadth (Levin's  $B_i$ ), and trophic level ( $TL$ ) estimated for SAFS and SASL were correlated with SST and SSTA metrics using a correlation matrix (Spearman rank,  $r_s$ ,  $p < 0.05$ ) with packages `rstatix` and `corrplot` in R statistical software (R Core Team, 2018).

## 3. Results

### 3.1 *Study area and sampling*

Of all samples collected, 80.06% ( $n = 2,662$ ) had identifiable hard remains; the remainder were recorded as empty scats. The proportion of SASL with hard remains (93.10%,  $n=1,714$ ) was significantly larger (Chi-square test,  $X^2 = 17.3$ ,  $p < 0.05$ ) than the proportion of SAFS with hard remains (63.88%,  $n=948$ ). This pattern was the same in all years, except 2015 (Chi-square test,  $X^2 = 0.337$ ,  $p = 0.561$ ), when the proportion of empty scats was not significantly different between species (Table 1). Adequate resolution to capture the prey diversity was obtained by grouping samples for three consecutive months during the 2015 - 2018 study period for austral summer (January - March), fall (April - June), winter (July - September) and spring (October - December). A total of 31 matrices were constructed for 16 sampling seasons for SAFS and 15 for SASL. Required samples per season for SAFS were in range of 35 - 120 and 27 - 171 for SASL. Therefore, we grouped monthly sampling into seasons, so sample size was always above the number of required samples to determine diet diversity for both SAFS and SASL.




**Table 2. 1** Number of collected and proportion of scat samples with content (in brackets) for South American fur seal (SAFS) and South American sea lion (SASL) per year (2015 - 2018), season (sum = summer, fal = fall, win = winter and spr = spring) and total.

<b>Year</b>	<b>Season</b>	<b>SAFS</b>	<b>SASL</b>
	2015-sum	62 (0.66)	30 (0.43)
	2015-fal	97 (0.63)	97 (0.53)
	2015-win	70 (0.69)	50 (0.96)
	2015-spr	40 (0.70)	151 (0.99)
<b>2015</b>		269	328
	2016-sum	129 (0.74)	183 (0.94)
	2016-fal	111 (0.77)	129 (0.98)
	2016-win	101 (0.78)	183 (0.99)
	2016-spr	40 (0.68)	-
<b>2016</b>		381	495
	2017-sum	120 (0.38)	120 (1.00)
	2017-fal	60 (0.38)	131 (0.98)
	2017-win	110 (0.55)	86 (1.00)
	2017-spr	50 (1.00)	60 (0.92)
<b>2017</b>		340	397
	2018-sum	126 (0.52)	189 (0.97)
	2018-fal	121 (0.83)	185 (0.97)
	2018-win	125 (0.63)	188 (0.93)
	2018-spr	122 (0.67)	59 (1.00)
<b>2018</b>		494	621
<b>Total</b>		<b>1,484</b>	<b>1,841</b>

### 3.2 Diet composition


SAFS hard remains were composed of 56.47% fish (n = 13,403 otoliths), 30.70% cephalopods (n = 7,286 beaks) and 12.84% crustaceans (n = 3,047 cephalothorax). Meanwhile SASL samples were composed of 6.55% fishes (n = 12,189 otoliths), 1.60% cephalopods (n = 2,981 beaks) and 91.85%



crustaceans (n = 171,034 cephalothorax) (Figure 2.1A).

The proportion of fishes (Chi-square test,  $X^2 = 7.90$ ,  $p < 0.05$ ) and cephalopods (Chi-square test,  $X^2 = 9.15$ ,  $p < 0.05$ ) in the hard remains of SAFS samples were ten to thirty times higher than in SASL. Significant differences in fishes were found for all years except for year 2015 (Chi-square test,  $X^2 = 0.00$ ,  $p = 1.00$ ). The proportion of fishes in diet was significantly different between SAFS and SASL in spring (Chi-square test,  $X^2 = 4.60$ ,  $p < 0.05$ ) and winter ( $X^2 = 5.33$ ,  $p < 0.05$ ), but not in fall (Chi-square test,  $X^2 = 1.33$ ,  $p = 0.25$ ) and summer (Chi-square test,  $X^2 = 1.33$ ,  $p = 0.25$ ). The proportion of the cephalopods was always higher in SAFS in comparison to SASL samples in summer (Chi-square test,  $X^2 = 5.33$ ,  $p < 0.05$ ), fall (Chi-square test,  $X^2 = 4.10$ ,  $p < 0.05$ ), winter (Chi-square test,  $X^2 = 5.33$ ,  $p < 0.05$ ) and spring (Chi-square test,  $X^2 = 4.60$ ,  $p < 0.05$ ). Finally, the proportion of crustaceans found in SAFS samples was seven times lower than those found in SASL (Chi-square test,  $X^2 = 13.70$ ,  $p < 0.05$ ). This pattern was found in all years except for year 2015, when no significant differences were found for crustaceans ( $p = 0.77$ ). The proportion of crustaceans in diet of SAFS and SASL was significantly different in spring (Chi-square test,  $X^2 = 4.50$ ,  $p < 0.05$ ) and winter seasons (Chi-square test,  $X^2 = 5.33$ ,  $p < 0.05$ ) but not in summer (Chi-square test,  $X^2 = 1.71$ ,  $p = 0.19$ ) and fall (Chi-square test,  $X^2 = 2.08$ ,  $p = 0.15$ ).



A total of 22 prey items were identified in this study; 19 for SAFS and 15 for SASL. With 22 species accounting for >1% of *IIMP*. When all years and seasons were pooled, the main prey items in the diet of SAFS were 42.8% *Engraulis ringens*, 34.7% cephalopods, 7.4% *Leuroglossus. stilbius*, 5% *Pleuroncodes monodon*, 3.8% *Anchoa nasus*, 2.1% *Sphyraena idiaestes*, 1.6% *Odontesthes regia*, and 1.1% *Merluccius gayi*. The main prey items in the diet of SASL were 69.9% *P. monodon*, 19.1% *E. ringens*, 4.7% cephalopods, and 4.6% *O. regia* (Figure 2.1B). The complete list of prey item *IIMP* values for each species, year, and season is available as Appendix B2. Average percent FO for top five prey items during the entire study period for SAFS were 37.0% *Engraulis ringens*, 32.5% cephalopods, 8.0% *Leuroglossus. stilbius*, 5% *Pleuroncodes monodon* and 6.3% *Anchoa nasus*. Meanwhile average percent FO for top five prey items during the entire study period for SASL were 43.8% *P. monodon*, 30.1% *E. ringens*, 12.6% cephalopods, 4.9% *O. regia* and 3.5% *N. crockeri*. Seasonal variations in PN per taxa and *FO* and *IIMP* for prey items are shown in Figure 2.2-2.3. Ecological grouping reveals that SAFS consumed prey items that were 52.6% Pelagic, 47.3% Demersal-Pelagic and 1.5% Demersal-Benthic. Meanwhile SASL consumed prey items that were



23.9% Pelagic, 69.4% Demersal-Pelagic and 1.4% Demersal-Benthic (Figure 2.3D).

### 3.3 *Trophic indices*



Species richness ( $S$ ) for all SAFS samples ( $n = 1,484$ ) was estimated at 19 species and for all SASL samples ( $n = 1,841$ ) at 15 species, with a maximum of 6 species per sample for either SAFS or SASL. No seasonal difference in  $S$  was found for SAFS ( $12.06 \pm 0.82$  for summer,  $11.36 \pm 0.67$  for fall,  $12.13 \pm 0.84$  for winter and  $12.97 \pm 0.18$  for spring) or SASL ( $10.52 \pm 0.65$  in summer,  $9.29 \pm 0.66$  in fall,  $12.82 \pm 0.88$  in winter and  $11.72 \pm 0.51$  in spring) (Table 2). Levin's  $B_i$  for both species was  $< 0.6$ , categorizing both species as “specialists”. Regardless, SASL has a more narrow trophic niche breadth ( $B_i = 0.11 \pm 0.12$ ) than SAFS ( $B_i = 0.26 \pm 0.13$ ) (Table 2). Mean  $TL$  for SAFS was  $4.30 \pm 0.15$  and  $4.55 \pm 0.10$  for SASL, which categorizes them both as tertiary consumers ( $TL > 4$ ), with no significant differences between species (Table 2).



**Figure 2. 1** Bar graphs of A) percent number (%) of taxa consumed by each species during the study period and B) stacked % IIMP values for each prey item over the entire study period for South American fur seal (SAFS, green) and South American sea lion (SASL, blue). Prey items (N = 22) detected are listed organized in alphabetical order. Values for IIMP > 1% are annotated in each otariid species color.



**Table 2. 2** Species Richness (*S*), Levin's Trophic Niche Breadth (*Bi* ± 95 %CI) and Trophic Level (*TL*) for South American fur seal (SAFS) and South American sea lion (SASL) per year (2015 - 2018), season (sum = summer, fal = fall, win = winter and spr = spring) and Mean ± SD for the study period.

Season	<i>S</i>		<i>Bi</i> ± 95% CI		<i>TL</i>	
	SAFS	SASL	SAFS	SASL	SAFS	SASL
2015-sum	8	4	0.18 ± 0.033	0.34 ± 0.125	4.12	4.30
2015-fal	8	7	0.29 ± 0.028	0.41 ± 0.080	4.28	4.51
2015-win	8	8	0.19 ± 0.031	0.15 ± 0.009	4.19	4.39
2015-spr	8	7	0.39 ± 0.050	0.23 ± 0.013	4.49	4.44
2016-sum	11	8	0.35 ± 0.032	0.05 ± 0.002	4.53	4.61
2016-fal	11	9	0.20 ± 0.015	0.07 ± 0.003	4.26	4.61
2016-win	11	12	0.19 ± 0.019	0.01 ± 0.001	4.19	4.67
2016-spr	6	-	0.31 ± 0.063	-	4.46	-
2017-sum	7	8	0.26 ± 0.031	0.03 ± 0.001	4.44	4.64
2017-fal	3	7	0.62 ± 0.179	0.04 ± 0.002	4.28	4.64
2017-win	11	9	0.27 ± 0.029	0.02 ± 0.001	4.40	4.68
2017-spr	8	9	0.15 ± 0.017	0.06 ± 0.002	4.46	4.61
2018-sum	8	9	0.34 ± 0.039	0.07 ± 0.002	4.31	4.53
2018-fal	6	7	0.15 ± 0.014	0.07 ± 0.002	4.11	4.58
2018-win	7	12	0.09 ± 0.013	0.05 ± 0.001	4.05	4.54
2018-spr	8	6	0.22 ± 0.023	0.09 ± 0.004	4.31	4.55
<b>Mean ± SD</b>	<b>8 ± 2.17</b>	<b>8 ± 2.07</b>	<b>0.26 ± 0.13</b>	<b>0.14 ± 0.16</b>	<b>4.31 ± 0.15</b>	<b>4.55 ± 0.11</b>




**Figure 2. 2** Stacked bar graphs of A) Percent number (%) by taxa; B) Frequency of occurrence (%) by prey item. South American fur seals (SAFS, top panels) and South American sea lions (SASL, bottom panels) for each year (2015-2018) and season (sum = summer, fal = fall, win = winter, spr = spring).







**Figure 2. 3** Stacked bar graphs of C) IIMP >1% by prey items and D) IIMP (%) by ecological groups South American fur seals (SAFS, top panels) and South American sea lions (SASL, bottom panels) for each year (2015-2018) and season (sum = summer, fal = fall, win = winter, spr = spring).





### 3.4 Clustering



The hierarchical clustering shows two clear separate clusters for each species. The only exception are summer and fall 2015 with an overlap between SASL and SAFS (Figure 2.4). There are no clear patterns for other seasons and/or years. Non-metric Multidimensional Scaling (NMDS) plot (stress = 0.08) reinforces the separation between SAFS and SASL (Figure 2.4). *E. ringens* is an important prey for both species, due to its central position between the two clusters. Cephalopods are closely associated with the SAFS cluster, whereas *P. monodon* is associated to the SASL cluster. Finally, *O. regia* is positioned close to the sampling season nodes of SASL that correspond to summer 2015 and fall 2015. Fish species *Isacia conceptionis*, *Paralabrax humeralis*, *Sciacena deliciosa*, and *Normanichthys crockeri* are associated with SASL; whereas a larger diversity of other fish species *Merluccius gayi*, *Lampanyctus parvicauda*, *Anchoa nasus*, *Leuroglossus stilbius*, *Strongylura exilis*, *Sardinops sagax sagax*, *Fodiator rostratus*, *Sphyræna idiaestes*, *Scomber japonicus*, and *Scomberesox saurus scombroides* are more closely associated with SAFS. Abundance rank curves show that during seasons with trophic overlap (summer and fall 2015), SAFS and SASL diets were not significantly different ( $p < 0.05$ ) and were composed of 55% and 48.6% *E. ringens*, plus 21.7% and 13.6% cephalopods, respectively. Furthermore, during these seasons SASL consumed 33% of *O. regia* and SAFS consumed 5.6% *L. stilbius* and 4.6% *A. nasus*. In these seasons, *P. monodon* consumption was 9% and 2%, respectively for SAFS and SASL diets (Figure 2.5).




**Figure 2. 4** Dendrogram using agglomerative hierarchical clustering with Ward's method and Bray-Curtis distances, showing two separate clusters for South American sea lion (SASL, blue) and South American fur seal (SAFS, green). Two SASL seasons (SASL-2015-fal and SASL-2015-sum) are inside the SAFS cluster (red asterisks).



**Figure 2. 5** NMDS spider plot (stress = 0.08) showing two separate clusters for each otariid species. Each node represents a sampling season for South American sea lion (SASL, blue) and South American fur seal (SAFSL, green). Relevant prey items for principal dimensions (axis 1 and axis 2), overlaid in dark red.



**Figure 2. 6** Ranked abundance curves for prey items of South American sea lion (SASL, blue) and South American fur seal (SAFS, green) samples during A) seasons with trophic overlap (summer 2015 and fall 2015) and B) seasons without trophic overlap. Labels shown for top-five ranking prey items.




### 3.5 *Environmental conditions and trophic indices*

Throughout the entire study period, SST ranged between 13.20 and 17.70 °C. Mean SST throughout the study period was  $16.06 \pm 0.45$  °C in summer,  $15.52 \pm 0.50$  °C in fall,  $14.28 \pm 0.56$  °C in winter and  $14.56 \pm 0.43$  °C in spring. SSTA ranged between -1.00 to 1.44 °C. Mean SSTA during the study period was  $0.21 \pm 0.48$  °C in summer,  $0.54 \pm 0.49$  °C in fall,  $0.30 \pm 0.31$  °C in winter and  $0.32 \pm 0.31$  °C in spring. In general, SST was significantly different (ANOVA,  $F(1,15) = 9.54$ ,  $p < 0.01$ ), allowing to treat each season independently throughout the study period. However, Tukey multiple comparison of pairwise means showed that fall-summer ( $p = 0.38$ ) and spring-winter ( $p = 0.84$ ) were not significantly different to each other in some years (summer–fall 2015, 2017, 2018 and winter–spring 2017, 2018). Since the six months grouping could not be applied to the entire study period, we kept the 3-month seasonal sampling resolution to compare trophic indices and environmental conditions for all seasons and years (Appendix B3).

For SAFS (N = 16 seasons), significant correlations with scores  $> 0.50$  were found between PN of cephalopods and TL ( $r_s = 0.76$ ); TL and Bi ( $r_s = 0.64$ ); Bi and mean SST ( $r_s = 0.54$ ); and between PN of cephalopods and PN of fish ( $r_s = -0.74$ ), PN of fish and TL ( $r_s = -0.64$ ), and PN cephalopods and crustaceans ( $r_s = -0.55$ ). For SASL (N = 15 seasons), significant correlations with scores  $> 0.50$  were found between PN of fish and Bi ( $r_s = 0.93$ ), PN crustaceans and TL ( $r_s = 0.82$ ), PN of fish and empty scats ( $r_s = 0.72$ ), and proportion of empty scats and Bi ( $r_s = 0.65$ ); and between PN of crustaceans and PN of fish ( $r_s = -0.96$ ), TL and Bi ( $r_s = -0.90$ ), PN of crustaceans and Bi ( $r_s = -0.90$ ), PN of fish and TL ( $r_s = -0.89$ ), PN of crustaceans and proportion of empty scats ( $r_s = -0.7$ ), proportion of empty scats and TL ( $r_s = -0.72$ ), and Species Richness and Bi ( $r_s = -0.63$ ). For SASL, no significant correlations with scores  $> 0.50$  were found between trophic and environmental indices (Appendix B4).

## 4. Discussion

This study demonstrates that trophic segregation exists between populations of sympatric SAFS and SASL at PSJ, Peru primarily due to the resource partitioning of three main




shared prey items: Peruvian anchovy, cephalopods, and red squat lobster. Both otariids share Peruvian anchovy as an important prey item and complement their diets with two other important items: cephalopods in SAFS and red squat lobsters in SASL. The consumption of these prey items in different proportions by each otariid explains the trophic segregation detected through the hard part remains found in scats collected during the consecutive seasonal samplings in the 2015-2018 study period. As part of the study, we also detected that periods of higher trophic overlap can occur, when similar proportions of *E. ringens* and cephalopods were consumed by both species and interspecific competition could be exacerbated.

#### 4.1 *Resource partitioning and trophic overlap*

In previous studies conducted in sympatric otariid colonies in Peru, Peruvian anchovy (*E. ringens*) made up 30 - 40% of the diet of both SAFS and SASL and was reported as the cause for interspecific trophic overlap between these sympatric otariids (Vásquez 1995, Arias-Schreiber 2003). As expected, *E. ringens* continues to be an important prey item that is widely available in the HCS system. However, in our study, SAFS consistently consumed anchovy and cephalopods, whereas SASL consumed anchovies and red squat lobsters. Increased in availability and consumption of red squat lobsters within the HCS system following the 1997-98 El Niño (Gutiérrez et al. 2008, Kiko et al. 2015) has made this item an important part of SASL diet (Espinoza et al. 2017).

During austral summer and fall of 2015 in which trophic overlap was detected, SASL did not consume *P. monodon* in the same proportions as other seasons, but instead increased consumption of *E. ringens* and included *O. regia*. This may be due to a reduced availability of *P. monodon* during months of the 2015/16 El Niño in the preferred habitat where sea lions forage as compared to SAFS. Oceanographic acoustic studies have shown that *P. monodon* have preference for cold coastal water masses, since they ensure good conditions for feeding, reproductive success and recruitment. However, during Extraordinary El Niño events such as in 2015-16, cold water contracts into the coast, decreasing the habitat for *P. monodon* and decreasing availability of the resource (Yuffra et al. 2018). Therefore, it is possible that *P. monodon* was not available for SASL






foraging range in the 2015 summer and fall seasons.

Foraging habitat of SAFS and SASL has been mostly studied in the Atlantic coast of South America and has found that SASL consumes demersal and benthic prey types while SAFS consumes pelagic prey (Riet-Sapriza et al. 2013, Baylis et al. 2016). In this study, ecological groups revealed that Demersal-Benthic species are not important in neither of the Peruvian otariids' diet composition (1-1.5%). SAFS diet is mostly composed of prey items from Pelagic (52.35%) and Demersal-Pelagic (47.25%) ecological groups. Meanwhile SASL's prey items are mostly Demersal-Pelagic (69.44%) and also include smaller amounts of Pelagic (23.88%) groups. This can be explained by the limited access to the benthic habitat for otariids i.e., in PSJ depth drops to >1000 m in <50km from shore. However, differences may also be attributed to sexual trophic segregation, as evidenced in SASL in Patagonia, where females have a more pelagic diet than males (Drago et al. 2009). Further studies using methods to explore intraspecific differences (e.g., stable isotopes) and depth (e.g., time depth recorders) are recommended in HCS.

## 4.2 *Empty scats*

In this study we found significant differences between the proportion of empty scats in SAFS compared to SASL. We also found a direct relationship between empty scats and the proportion of fish consumed, which can be related with longer offshore foraging trips (Naya et al. 2002, Franco-Trecu et al. 2013, Baylis et al. 2014). We consider this an important indicator for differences foraging locations between fur seals (offshore) and sea lions (inshore), given that the three important prey items (*E. ringens*, *P. monodon* and Cephalopods) have pelagic ecological habits. This relation was also found in SAFS from the Atlantic coast of Uruguay, where the number of empty scats from females was related to longer foraging trips far away from the rookery (Franco-Trecu et al. 2013). An observational study at PSJ estimated that foraging trip duration of adult female SAFS attending new born pups was 3 - 4 days on average, within a range of 1 - 9 days (Ganoza 2016). Whereas studies in adult female SASL have reported foraging trips by adult females of 1-2 days (Soto et al. 2006). Thus, the high proportion of SAFS empty scats is possibly due the evacuation of otoliths in the ocean prior to returning to the colony.



### 4.3 Previous research


Due to the differences in the access to the benthic habitat for otariids in the Atlantic Ocean (extensive shelf with gradual decline) versus the Pacific Ocean (narrow shelf with steep decline), here we focus on comparing our study with previous research in sympatric sites in the Pacific Ocean conducted 2-3 decades earlier to analyse possible fluctuations in the availability of prey consumed over time. Vásquez (1995) evaluated samples from SAFS and SASL in the same study location (PSJ) between 1986 - 1989. Back then red squat lobster was not identified as an important prey item for either of the otariids. However, Pacific Jack Mackerel (*Trachurus murphyi*, previously identified as *T. symmetricus*) was ranked as the second most frequent prey item after Peruvian anchovy with 17% and 15% occurrence in diets of SAFS and SASL, respectively. In our study, zero occurrence of *T. murphyi* was found. Also, Vásquez reported an occurrence of <1% in cephalopods for both SAFS and SASL diets and no occurrence of *P. monodon*. In contrast, we found cephalopods as an important prey item for SAFS and *P. monodon* as a dominating prey item for SASL. Overall, our findings suggest an increase in the availability of red squat lobsters and cephalopods in the HCS and a decrease in *T. murphyi* as a possible shifts in the diet composition of otariids over the past 25 years, that should be further explored.

Another important study for comparison is Arias-Schreiber (2003), that analysed samples from Punta Coles (17°42'S, 71° 23'W). The location of study sites undoubtedly can influence availability of prey resources; however, tagged SAFS and SASL from PSJ have been sighted in Punta Coles demonstrating animal dispersal between these sites (Arias-Schreiber 2000). Arias-Schreiber (2003) found that both otariids consumed mainly teleost fish, with no significant differences in fish diversity consumed. In total 61 fish species were identified, 41 fish species in SAFS and 51 in SASL. In our study, fish consumption was significantly higher in SAFS than in SASL; with 19 species in SAFS and 15 in SASL making a total of only 22 unique fish species identified for both otariids. Arias-Schreiber (2003) found that 20% of SAFS and 32% of SASL samples contained cephalopod beaks, whereas in our study cephalopod consumption was significantly different between otariids, with 32% in SAFS in comparison to 1.6% in SASL. In regards to crustaceans, Arias-Schreiber (2003) found 8% of SAFS and 60% of SASL samples contained red squat



lobster; similar to the 5% and 70% for SAFS and SASL respectively, found in our study. Overall, Arias-Schreiber (2003) results are similar to the current diet composition found in sympatric otariids in our study, suggesting long-term changes in trophic assemblages after the Extraordinary 1997-98 El Niño.

Trophic level did not reveal any significant differences between consumers. Similarity in the overall TL score is because the most abundant prey items have similar TL values (*E. ringens* TL = 2.9, *P. monodon* TL = 3.2 and cephalopods TL = 3.5). In terms of trophic niche breadth, we found that both species are categorized as specialists (Levin's  $B_i < 0.6$ ), which contrasts with previous studies that classified species as generalists (Naya et al. 2002, Arias-Schreiber 2003, Sarmiento-Devia et al. 2020). This can be explained by the high proportions of 1-2 main prey items in each of the predator diets, in combination with the lower diversity of fish species found in our study. We also found higher trophic niche breadth for SAFS than SASL. This also differs from previous studies in the Peruvian HCS, that found that SASL consumed more fish species in previous years (Vásquez 1995, Arias-Schreiber 2003). Changes in the SASL diet coincides with higher variability and dietary shifts reported in SASL Atlantic populations (Drago et al. 2009) in comparison to more stable diets in Atlantic SAFS (Vales et al. 2020).




#### 4.4 *General limitations in methodology*

The analysis of hard remains in scats is one of the most widely used technique for estimating the diet of pinnipeds (Tollit et al., 2006, 1997). However, there are a number of recognized problems with this method, the main limitation are not being able to identify soft shell prey (e.g., crabs), variation in recovery rates and observation errors associated with structures used for identification (Tollit et al. 1997). In this study, we acknowledge that all these limitations are inherent to the study. However, we also consider this error is consistent between study species, location, and sampling season, making long-term temporospatial comparisons valid.

Previous studies in SAFS and SASL in HCS have identified cephalopod beaks belonging to the Teuthidae family (Majluf 1987, Vásquez 1995, Arias-Schreiber 2003, Sarmiento-Devia et al. 2020). However, other species, such as *Octopus mimus* have been detected in other locations (Sielfeld et al. 2018). Therefore, another limitation is the lack of species identification in the cephalopod prey item category. An additional mechanism for interspecific resource partitioning may also exist in prey sizes targeted. Thus, we recommend measuring hard parts to estimate differences in prey size consumed to determine if other measures of segregation in prey selection exist.


Low recovery of otoliths in otariid scats has been demonstrated through experimental feeding trials with captive otariids that measure the gut transit time of otoliths (Dellinger & Trillmich 1988, Tollit et al. 1997). In an experiment with captive SAFS and California sea lions (*Zalophus californianus*), Dellinger and Trillmich (1988) found mean recovery rates of 38.5% with high interindividual variation (1.25 - 87.3% in sea lions and 3.8 - 80% in fur seals), and a mean passage time in the gut of 22 hours (with initial defecation time of only 4 hours post feeding trial). How these data compare to free-ranging pinnipeds remains uncertain, but we acknowledge it as a limitation of our methods. Regardless, we consider the large sample size per season included in our study, along with the consecutive sampling over a four year period, strengthens our data as a reliable method to capture long term patterns and changes in the hard remains of prey items at a population scale.



#### 4.5 *Environmental conditions and trophic indices*


Correlations between environmental condition indices (SST and SSTA) and trophic indices show that increases in SST and SSTA are related to wider trophic niche breadth in SAFS, elucidating that during warmer conditions there is inclusion of more prey pelagic / oceanic items, in agreement with previous studies conducted during ENSO conditions (Arias-Schreiber 2003). This suggests that a mechanism of SAFS to cope with warmer conditions is to increase the species richness as a response to the increase of tropical water species in the HCS (Majluf 1987, Paredes & Arias-Schreiber 1999, Arias-Schreiber 2003). In SASL however, we did not find a clear relationship between changes in resource consumption in regard to SST or SSTA.

During February and March of 2015 El Niño conditions promoted the influx of superficial warm water masses towards coastal Peruvian waters, with the peak of positive SSTA extending from April to June 2015 in the El Niño +1.2 region (<http://met.igp.gob.pe/datos/icen.txt>). The 2015-16 El Niño was categorized as one of the most intense El Niño events in the historical record (L'Heureux et al. 2017). The metrics of the 2015-16 El Niño were based on the SSTA measured in the El Niño 3.4 region (5°N - 5°S, 170° - 120°W) that is ~1,800 km from the Punta San Juan (15°S, 72°W). Values for SST and SSTA used in this study were collected 5 km from PSJ. Seasonal SSTA values showed warming in the first quarter of 2016 and last 3 quarters of 2015. Thus, warming of the surface layer farther offshore could have already affected prey distribution earlier in 2015. Also in 2015 there were unusual reports of oceanic and tropical species found inshore, increased proportion of *E. ringens* juveniles in anchovy purse seine fisheries (IMARPE 2015) and decrease in available biomass of *P. monodon*, evidencing changes in coastal water masses (Yuffra et al. 2018). Although this does not completely explain the trophic overlap in summer and fall of 2015 between SAFS and SASL in the first semester of 2015, it reflects how oceanographic changes triggered by warm Kelvin waves can alter spatial and temporal food web composition, including prey base assemblages.




Increases in the population of *P. monodon* in Peru has occurred since mid-1990s, when the range of this species expanded from central Chile towards Peru taking advantage of the northern expansion of cold-water conditions (Gutiérrez et al. 2008). This, in combination with the shallow oxycline (< 100 m), has led authors to hypothesize a complete pelagic lifestyle for *P. monodon* in the Peruvian HCS takes place, making it available prey for otariids and seabirds (Espinoza et al. 2017). However, other studies state that vertical migration from the oxygenated surface waters towards the hypoxic to anoxic layer and back occur as well (Kiko et al. 2015). Therefore, further studies on day/night foraging in these otariids is recommended. From a predator perspective, the incidence of *P. monodon* in the diet of SASL proves that this prey item is accessible to sea lions in the oxygenated layer; supporting sea lion colonies in PSJ that range between 3,000 - 12,000 individuals and between 75 - 150,000 in Peru (Cárdenas-Alayza et al. 2021). This reveals the importance of the understudied red squat lobster in the fulfilling the energetic demands of SASL, the larger and most abundant of the two otariids in Peruvian HCS. In Falkland Islands (Malvinas), evidence of red lobster consumption are linked to nearshore feeding, due to nearby shoaling locations (Baylis et al. 2014). Furthermore, when predators feed on crustaceans, carapaces tend to dominate the scats evidencing consumption (Staniland 2002).

It has been well documented that the Peruvian HCS contains one of the world's largest monospecific fisheries based on Peruvian anchovy, along with other fisheries that target cephalopods and other small pelagic fish species. Many of these fisheries have been overcapitalized and are decreasing available biomass to dangerous levels for sustainability (Fréon et al. 2008, De la Puente et al. 2020). To present date, we are unaware of any large-scale exploitation of *P. monodon* in Peru that limits the availability of this prey item for predators. However, if this were to occur, it may severely affect the availability of this prey resource for SASL and could lead to further changes in the diet of this species, and increase competition between the two sympatric otariid species as evidenced in other regions (Drago et al. 2009, Szteren et al. 2018)





APPENDIX B1




**Figure B1. 1** Boxplots of abundance of South American sea lions (SASL) in site N7 (2015-2018) according to age class during breeding (BS, January -March) and non breeding months (non BS, April – December) according to age class.






**Figure B1. 2** Boxplots of abundance of South American fur seals (SAFS) in site S3 (2015-2018) according to age class according to age class during breeding (BS, October - December) and non breeding months (non BS, January – September).




## APPENDIX B2

**Table B2. 1** List of prey items identified in the samples of South American fur seals (SAFS) and South American sea lions (SASL) and the habitat group category assigned.


Scientific name	English common name	Habitat	SAFS	SASL
<b>1</b>	Cephalopods	Demersal-Pelagic	X	X
<b>2</b> <i>Anchoa nasus</i>	Longnose anchovy	Pelagic	X	X
<b>3</b> <i>Aphos prosus</i>	Banded toadfish	Demersal-Benthic		X
<b>4</b> <i>Engraulis ringens</i>	Peruvian anchovy	Pelagic	X	X
<b>5</b> <i>Fodiator rostratus</i>	Sharpchin flyingfish	Pelagic	X	
<b>6</b> <i>Haliocerhes dispilus</i>	Chameleon wrasse	Demersal-Benthic	X	
<b>7</b> <i>Isacia conceptionis</i>	Cabinza grunt	Demersal-Pelagic	X	X
<b>8</b> <i>Lampanyctus parvicauda</i>	Slimtail lampfish	Demersal-Pelagic	X	
<b>9</b> <i>Leuroglossus stilbius</i>	California smoothtongue	Demersal-Pelagic	X	X
<b>10</b> <i>Merluccius gayi</i>	South Pacific hake	Demersal-Benthic	X	X
<b>11</b> <i>Mugil cephalus</i>	Flathead grey mullet	Pelagic		X
<b>12</b> <i>Normanichthys crockeri</i>	Mote sculpin	Demersal-Benthic	X	X
<b>13</b> <i>Odontesthes regia</i>	Chilean silverside	Pelagic	X	X
<b>14</b> <i>Paralabrax humeralis</i>	Peruvian rock seabass	Demersal-Pelagic	X	X
<b>15</b> <i>Pleuroncodes monodon</i>	Red squat lobster	Demersal-Pelagic	X	X
<b>16</b> <i>Prionotus stephanophris</i>	Lumptail searobin	Demersal-Benthic		X
<b>17</b> <i>Sardinops sagax sagax</i>	South American pilchard	Pelagic	X	
<b>18</b> <i>Sciacena deliciosa</i>	Lorna drum	Demersal-Benthic	X	X
<b>19</b> <i>Scomberesox saurus</i> <i>scombroides</i>	King gar	Pelagic	X	
<b>20</b> <i>Scomber japonicus</i>	Chub mackerel	Pelagic	X	X
<b>21</b> <i>Sphyraena idiaestes</i>	Pelican barracuda	Pelagic	X	
<b>22</b> <i>Strongylura exilis</i>	California needlefish	Pelagic	X	




## APPENDIX B3



**Figure B3. 1** Boxplots of the seasonal A) SST and B) SSTA for study period 2015-2018 collected in San Juan de Marcona, source: DHN - Peru.




# APPENDIX B4



**Figure B4. 1** Spearman correlation charts ( $p < 0.05$ ) for trophic and environmental indices for A) South American sea lions, SASL and B) South American fur seals, SAFS. Positive correlation scores in shades of blue and negative in shades of red.

## CHAPITRE 3

*Multiple strategies for segregation  
during foraging in sympatric otariids of  
the Peruvian upwelling Humboldt  
Current System*



**Cárdenas-Alayza S, Adkesson, M.J., Gutiérrez D, Demarcq, H, Tremblay Y.** Multiple strategies for segregation during foraging in sympatric otariids of the Peruvian upwelling Humboldt Current System.

Manuscript submitted for publication to Marine Ecology Progress Series



## Résumé

Le principe d'exclusion compétitive prédit que les espèces doivent avoir des mécanismes de ségrégation pour coexister. Dans le système du courant de Humboldt, les lions de mer d'Amérique du Sud (*Otaria byronia*; SASL) coexistent avec les otaries à fourrure d'Amérique du Sud (*Arctocephalus australis*; SAFS) ; cependant, la ségrégation temporelle et/ou spatiale lors de la recherche de nourriture reste inconnue. Nous avons analysé les localisations de 35 adultes (18 SASL, 17 SAFS ; 12 femelles, 23 mâles) équipés de balises satellites à Punta San Juan, Pérou (2013-2017) afin de déterminer leurs stratégies respectives de recherche de nourriture. Nous avons évalué : 1) la distance et la durée des trajets de recherche de nourriture ; 2) les distributions d'utilisation (UD) à 25, 50, 75 et 95 % ; 3) la recherche de nourriture par heure de la journée et 4) l'association à des variables environnementales. Les trajectoires ont été interpolés à intervalles réguliers (e/30min) en utilisant une marche aléatoire corrélée et le temps de résidence a été estimé pour déterminer les phases potentielles de nourrissage. La proportion de chevauchement et l'analyse de similarité ont été utilisées pour distinguer les emplacements dans les zones centrales (50%) et les domaines vitaux (95%). Des GAMM ont été construits pour déterminer si l'heure du jour avait un effet sur les événements de recherche de nourriture et pour évaluer l'association avec des variables environnementales. Les résultats montrent que la durée et la distance entre les groupes d'espèces-sexes sont significativement différentes, sauf pour la durée des déplacements entre les sexes chez les SASL. Toutes les UD des femelles se chevauchent, tandis que celles des mâles ne se chevauchent que dans 25% des cas. L'effet de l'heure du jour était significatif pour les femelles SAFS et les mâles SASL. En général, les variables environnementales associées suggèrent des caractéristiques d'habitat pélagique et côtier chez les femelles otaries à fourrure et les lions de mer, respectivement. Les mâles se séparent dans l'espace et dans le temps, tandis que les femelles présentent un chevauchement spatial mais disposent de mécanismes de ségrégation temporelle. Les résultats démontrent que les stratégies de recherche de nourriture multiples à différentes échelles contribuent au partage des ressources.

**Mots clés :** ségrégation interspécifique; coexistence ; otarie à fourrure ; lion de mer ; système du courant de Humboldt ; stratégie de recherche alimentaire



## Abstract

The principle of competitive exclusion predicts that species exhibit segregation mechanisms to coexist. In the Humboldt Current System, South American sea lions (SASL, *Otaria byronia*) coexist with South American fur seals (SAFS, *Arctocephalus australis*); however, temporal and spatial partitioning while foraging remains unknown. We analyzed locations of 35 adults (18 SASL, 17 SAFS; 12 females, 23 males) equipped with satellite tags in Punta San Juan, Peru (2013-2017) to determine their respective foraging strategies. We evaluated: 1) distance and duration of foraging trips; 2) 25, 50, 75 and 95% utilization distributions (UD); 3) foraging by hour of day and 4) association of environmental variables. Regular interval tracks ( $e/30\text{min}$ ) were modeled using correlated random walk and residence time was estimated. Proportion overlap and analysis of similarity were used to discriminate locations in core areas (50%) and home ranges (95%). GAMMs were built to determine if hour of day had an effect on foraging events and to evaluate association with environmental conditions. Results show that duration and distance between species-sex groups are significantly different, except for trip duration between SASL sexes. All female UD overlap, while males UD overlap in 25% only. Effect of hour of day was significant for SAFS females and SASL males. In general, associated environmental variables suggest pelagic and coastal habitat features in female fur seals and sea lions, respectively. Males segregate in space and time; whereas females exhibit spatial overlap but have temporal segregation mechanisms. Results demonstrate how multiple foraging strategies at different scales aid resource partitioning.

**Keywords:** interspecific segregation, coexistence, fur seal, sea lion, Humboldt Current System





## 1. Introduction

Coexisting species that share similar morphological traits, may compete for resources (Hutchinson & MacArthur, 1959). Competition can compromise the fitness of the species involved and may ultimately lead to the competitive exclusion of one species, especially when resources are limited. The principal of competitive exclusion predicts that coexisting species will exhibit resource partitioning (spatial, temporal or dietary segregation) and occupy different ecological niches (Pianka, 1974; Schoener, 1974). However, the Lotka-Volterra model predicts that stable coexistence of competitors is possible when interspecific competition is, for both species, less significant than intraspecific competition (Begon et al., 2006). In these scenarios, niche differentiation will tend to concentrate competitive effects more within species than between them. The Lotka-Volterra model and the Competitive Exclusion Principle, therefore imply that any amount of niche differentiation will allow the stable coexistence of competitors. The ways in which species within ecological communities partition available resources among themselves is a major determinant of the diversity of coexisting species. A community with more resource sharing or greater niche overlap, will clearly support more species than one with lower niche overlap. Thus, coexisting species must differ in their ecological requirements by at least some minimal amount to avoid competitive exclusion (Pianka, 1974).

Among the sympatric marine mammal species with similar life-history traits and foraging habits are the otariids: fur seals and sea lions. In general, where fur seals and sea lions live in sympatry, the fur seal population is typically larger and they appear to outcompete sea lions. Many studies in both northern and southern latitudes have examined potential competition between sympatric fur seal and sea lions, and their results are mixed. Some have found segregation with no trophic overlap (Antonelis et al., 1990; Aurióles-Gamboa & Camacho-Ríos, 2007; Dellinger & Trillmich, 1999; Franco-Trecu et al., 2012), some have determined spatial segregation (Baylis et al., 2016, 2018; Schwarz et al., 2021; Villegas-Amtmann & Costa, 2017) while others have found dietary overlap (Páez-Rosas & Aurióles-Gamboa, 2010; Page et al., 2005; Waite, Burkanov, et al., 2012).

In the southern Pacific Ocean, South American fur seals (SAFS) and South American sea lions (SASL) can be found in sympatry along Peru's coastline. The present study is based in Punta San Juan, Peru, a protected area which supports important breeding colonies of both species in sympatry



year round, making it a suitable location to study coexistence (Cárdenas-Alayza et al., 2021). South American fur seals and sea lions are sexually dimorphic otariid species, with differential parental care between the sexes. Reproductive roles can influence habitat selection and differential intraspecific foraging behavior in various ways. Foraging behavior of adult females (3-4+ years old) are constrained to provisioning for offspring, limiting the duration while searching for food. Whereas adult males are free to extend their foraging trips in terms of duration and distance; except during the breeding season, when territorial males are competing for copulations, which may require a time of tenure on shore (Majluf, 1987; Soto & Trites, 2011). Furthermore, mass and size affect oxygen stores and metabolic rates which in turn have an impact on diving/foraging capabilities. Therefore, in our study species, sexual segregation plays a major role in obtaining nutritional demands.

Previous studies describe SAFS in Peru as epipelagic nocturnal foragers that dive in the range of 11-30m (Trillmich et al., 1987). However, at present, there are no published data on the spatial use patterns of SAFS in this ecosystem. Studies using depth sensors have reported that in many locations along their range, SASL feed on benthic resources on the continental shelf (Riet-Sapriza et al., 2013; Rodríguez et al., 2013; Thompson et al., 1998; Werner & Campagna, 1995). However, in the north of Chile pelagic foraging was reported by juvenile male SASL, with average dive depths of 30m (Hückstädt et al., 2014, 2016). Therefore, we consider it is possible that both species are partitioning resources in the pelagic domain. In a highly pelagic environment such as the south Peruvian coast of HCS, where the combination of a narrow continental shelf (80km from shore) and a shallow oxycline (50-80m) compress the habitat for potential prey (A. Bertrand et al., 2010, 2011) we hypothesize pelagic foraging strategies exist for both species combined with other forms of temporal and spatial segregation.

To determine how male and female SAFS and SASL segregate in the HCS, we compare feeding trips of 35 individuals (18 SASL, 17 SAFS) at Punta San Juan, Peru equipped with satellite transmission tags to determine interspecific and intraspecific differences in the temporal, spatial, and environmental features associated to foraging events. Specific objectives include determination of interspecific and intraspecific differences in: 1) distance and duration of foraging trips; 2) utilization distributions for home range and core areas; 3) time of foraging by hour of day; and 4) environmental conditions associated with foraging events for each of the four species and sex groups.



We aim to determine if temporal and spatial segregation among species and sex groups is influenced by morphological (body mass), behavioral or environmental conditions during foraging. SASL are considerably larger in body size than fur seals and both species have a strong sexual dimorphism, with males being much larger than females (Ralls & Mesnick, 2002). We consequently expect that SASL will travel longer distances, forage for longer durations, and cover larger areas than SAFS. Furthermore, we anticipate that males of both species will travel longer distances, forage for longer durations, and cover larger areas in comparison to females. All females in this study were lactating pups when tagged. Due to energetic constraints linked to maternal attendance in both species, we expect females will have reduced ranges in terms of distance, duration and consequently, area covered compared to males. In terms of temporal segregation, based on Trillmich et al. (1991), we expect that fur seals will forage mostly in the night hours to capture superficial available prey during diel vertical migration of the most abundant prey item in HCS, Peruvian anchovy (*Engraulis ringens*), whereas we expect SASL to forage indifferently throughout the day or night hours as shown in other sea lion species (Riet-Sapriza et al., 2013; Schwarz et al., 2021; Villegas-Amtmann et al., 2008). Due to the above mentioned energetic constraints linked to offspring survival, we also expect females of both species to be more coastal (inshore) and males more pelagic (offshore) as evidenced in other otariids (Page et al., 2006; Staniland & Robinson, 2008). In HCS, the coastal upwelling translates to lower sea surface temperatures (SST), higher productivity (assessed by higher concentrations of chlorophyll-a), less distance from coast, lower proximity to thermal fronts, and lower front gradients; and the opposite for pelagic habitats, which we expect to be more associated to male foraging behavior (Echevin et al., 2008; Gutiérrez et al., 2011).

## 2. Materials and Methods

### 2.1 *Data collection*

Satellite transmitting tags were deployed on 35 adult pinnipeds between 2013-2017 in Punta San Juan (PSJ), Peru (15°22'S, 75°12'W, Figure 3.1). Seventeen SAFS (8 females and 9 males) and eighteen SASL (4 females and 14 males) were captured and instrumented as part of this study. Adult females were nursing newborn pups prior to tagging and are therefore estimated to be 4+ years old. Adult



male SAFS were selected from bulls holding tenure at territories and therefore are an estimated age of 9+ years old and their feeding trips reflect post-tenure feeding trips. Whereas SASL males were categorized in the field as sub-adult males, which are reproductively mature individuals, but do not control female harems yet and are estimated to be within 5-8 years old. Adult females were instrumented during the peak of pupping during their respective breeding season: November for SAFS and February for SASL. Whereas males were all instrumented in November.

Anesthesia of individual SASL and male SAFS was induced using a combination of midazolam, butorphanol, and medetomidine administered via plastic dart as previously reported (Adkesson, Chinnadurai, Balko, Jankowski, Langan, et al., 2019; Adkesson, Chinnadurai, Balko, Jankowski, Meegan, et al., 2019). Female SAFS were captured using a hoop net and then anesthetized with isoflurane gas (1 - 5% to effect) mixed with oxygen as previously described (Jankowski *et al* 2015). Anesthesia was performed by a board-certified specialist in zoological medicine. Morphometrics and physiological parameters were recorded. Body weights were obtained using a tripod and field scale to the nearest 100 grams (Mini Crane Scale 300 kg, OCS-L). All animals were determined to be in good health by a veterinarian based on physical examination and veterinary assessment of routine blood parameters. Antagonist drugs (flumazenil, naltrexone, and atipamezole) were administered and all animals returned safely back into the colony or immediate surroundings.

All samples and methodology were approved by the Peruvian government under research permits Resolución Jefatural No. 09-2013, No. 024-2014, No. 008-2015- and 019-2016-SERNANP-RNSIIPG issued by the Peruvian National Service of Natural Protected Areas (Spanish acronym SERNANP) and the Peruvian Ministry of the Environment (Spanish acronym MINAM). Procedures were approved by the Ethics Committee of Universidad Peruana Cayetano Heredia (Constancia #005-02-18). During restraint, each individual was instrumented with satellite transmitting tags (Spot 5, Spot 6 and Mk 110 from Wildlife Computers) that communicate with the Argos satellite system, programmed to relay 4-6 transmissions per day. Animal locations and associated errors were estimated by calculating the distance between tags and the satellites present during the time of message transmission. Geographically referenced sightings of satellite tagged individuals at the Punta San Juan site were added to each animal's location record.



## ***2.2 Track construction***

A range of possible locations were modelled using a forward particle correlated random walk method following Tremblay et al. (2009). Fifty particles were projected every 30 minutes based on animal speed and Argos location class quality to inform model. One average track and its associated error distribution were computed to describe each individual's track. Land locations were filtered out by using a combination of 5km circle buffers on known pinniped colonies along the coast of Peru (including islands), a bathymetry gridded map (GEBCO v. 2020) and a limit for a minimum number of 15 hours or more to consider a foraging trip at sea, based on previous findings (Ganoza, 2016; Hückstädt et al., 2014) (Figure 3.1). Individuals were tracked for 23-78 days. A summary of individual identification, deployment date, weight, length, number of days, and number of trips for each individual is included in the Appendix C1.

## ***2.3 Foraging trips and events***

Total distance (km) and duration at sea (days) was estimated for each trip. Results are reported in mean  $\pm$  SD obtained per each species and sex group. Interspecific and intraspecific differences were tested for each of these variables (Kruskal Wallis,  $p < 0.05$ ). We estimated residence time (RT) as a foraging index to understand the time spent (in hours) for area-restricted search by individuals in areas of radius "r" (in km) and maximum time (in hours) during which they actively explored or foraged, following Barraquand and Benhamou (2008). Individuals are presumed to be in "foraging" mode based on movements that include high tortuosity, slower speeds, and repeated entries in/out of an area. Whereas low tortuosity and high speeds are presumed to be in transit or commuting between foraging areas and/or haul-outs. To determine the best scale, multiple size radii and time thresholds were tested for each individual. Radii tested were in the range of 1km, and 5 - 45 each 5 km for each animal and maximum time of 3, 6, 8 and 12 hours were tested. An RT series was constructed for each animal and radii, resulting in 40 possible RT series per individual. The RT series with the highest contrast was selected and segments were categorized as either high (HRT) or low (LRT) time following Lavielle segmentation method (Lavielle, 2005), using the adehabitatLT package in R (Calenge, 2006).



### 3. Utilization distributions

We used kernel density estimations to calculate the utilization distribution (UD); 25%, 50%, 75% and 95% UD kernels (grid size =200 ) with R package *adehabitatHR* (Calenge, 2006). Proportion of overlap of the home range (95% UD) and core areas (50% UD) were estimated using *adehabitatHR* package and reported in ranges from 0 (no overlap) to 1 (complete overlap). Bhattacharyya's affinity (BA) for all locations where presumed foraging activity took place was estimated. BA values were used to construct a dissimilarity matrix and analysis of similarity (ANOSIM) based on 999 permutations, to test the similarity between groups for each probable UD (25%, 50%, 75%, 95%) using package *vegan* in R. ANOSIM ranges from -1 to 1, with 0 indicating a random grouping. Finally, we tested for significant differences between populations according to UD based on ANOSIM statistic values ( $p < 0.05$ ).

### 4. Foraging by hour of day

We first explored the density of foraging events by hour of day to assess if there is interspecific segregation between the occurrence of foraging by hour of day for each sex. To test if there were different patterns per species and sex group, we constructed a general additive mixed-model (GAMM, *mgcv* R package) to determine if hour of day had a significant effect on the foraging events that occurred in 1 hour blocks according to each species and sex group (Wood, 2011). We used a binomial distribution, with a 1,0 response variable for foraging and hour of day as the explanatory variable. Since the data is analyzed in 24, 1 hour cycles, we chose a cyclic penalized cubic regression spline as a smoother. Animal ID served to assign random effects and a continuous time index for each foraging trip was constructed and assigned to indicate a *corAR1* temporal correlation structure. In the our study area and study period, changes in the time of sunrise and sunset are less than our sampling interval of 1 hour blocks, thus we did not include day length in our analysis.


### 5. Associated environmental conditions

We extracted five environmental variables for each location during feeding trips from remote sensor databases using R package *raster*. Environmental variables include: Distance to the Coast (DC,



Source: NOAA ETOPO1, 1 arc minute), Sea Surface Temperature (SST, Source: MODIS, 4 km), Chl-a concentration (Source: MODIS-Aqua, 4 km), Proximity to thermal fronts (km) and Gradient of thermal fronts ( $^{\circ}\text{C}/\text{km}$ ) both in a  $5 \times 5$  pixels range (4km/ pixel). Thermal fronts were constructed from daily composite SST maps (MODIS, 4 km) following Roa-Pascuali et al. (2015). Due to cloud coverage, some dates and locations do not have associated data. We extracted values during HRT locations (presumed foraging events) and compared significant differences between species (Wilcoxon, Test,  $p < 0.001$ ). Then, to explore the preferred environmental conditions during foraging, we fit univariate binomial General Additive Model (GAM) using cubic spline smoothers to estimate nonparametric functions for HRT. Finally, to determine which variables best explained environmental conditions during foraging, we conducted a forward selection process for each species and sex group, constructing a GAMM model with binomial distribution (1,0 response variable) and cubic-splines smoothers to model non-parametric relationships. Animal id was set as random effect and a corAR1 error structure to account for temporal correlation. If more than 25% of locations lacked information, the covariate was not included in model selection. We then assessed for collinearity between remaining candidate explanatory variables and did not include covariates that had a Spearman rank correlation  $> 0.8$  ( $p < 0.05$ ). Finally, we only kept locations that had information for all environmental covariates included in the global model. To avoid overfitting, knots were set to 5. Model selection was done through a forward selection process which consisted of evaluating if the addition of a new covariate produced a significant improvement in the model, evaluated with ANOVA tests (Chi square ,  $p < 0.05$ ).






**Figure 3.1** Tracks for A) SAFS and B) SASL in the study area. SAFS females (orange), SASL females (aquamarine); SAFS males (brown) and SASL males (teal). Yellow circle indicates the Punta San Juan, Peru breeding colony, location of otariid instrumentation.



## 6. Results

### 6.1 *Body mass*

Body mass and length between all individuals have a strong positive linear relationship ( $r^2 = 0.91$ ,  $p < 0.05$ ) with sizes scaling up in the following order: SAFS females < SASL females < SAFS males < SASL males. We therefore compared groups using only body mass and found that all species and sex groups are significantly different from each other, allowing us to make comparisons among the four groups (2-way ANOVA with Tukey post-hoc,  $p < 0.01$ , Figure 3.2). Body mass of SASL males has the widest range since this age-class gathers sub adult males who are still growing at different rates with high interindividual variability, in contrast to the other groups that have less variable values between individuals in the group.




**Figure 3. 2** Boxplot of body mass (kg) in y-axis according to species and sex groups in x-axis (ANOVA,  $p < 0.05$ ). SAFS females (orange), SASL females (aquamarine); SAFS males (brown) and SASL males (teal).



## 6.2 Trip distance and duration

SAFS travelled longer distances during trips at sea than SASL (Table 3.1, Figure 3.3). Interspecific differences were found in trip distance between adult females (Kruskal-Wallis,  $K-W = 14.8$ ,  $n = 149$ ,  $p < 0.01$ ) and adult males (Kruskal-Wallis,  $K-W = 8.8$ ,  $n = 276$ ,  $p < 0.01$ ). Intraspecific differences were found in trip distance between SAFS (Kruskal-Wallis,  $K-W = 4.1$ ,  $n = 134$ ,  $p < 0.05$ ) and SASL (Kruskal-Wallis,  $K-W = 19.5$ ,  $n = 291$ ,  $p < 0.01$ ). Mean trip duration was also longer in SAFS than SASL (Table 3.1, Figure 3.3). Interspecific differences exist in trip duration between adult females (Kruskal-Wallis,  $K-W = 8.5$ ,  $n = 149$ ,  $p < 0.01$ ) and adult males (Kruskal-Wallis,  $K-W = 14.6$ ,  $n = 276$ ,  $p < 0.01$ ). Intraspecific differences exist in trip duration between SAFS (Kruskal-Wallis,  $K-W = 4.6$ ,  $n = 134$ ,  $p < 0.05$ ) but not between SASL trip duration (Kruskal-Wallis,  $K-W = 3.6$ ,  $n = 291$ ,  $p = 0.0592$ ).



**Figure 3. 3** Boxplots of A) distance (km) and B) duration (days) of foraging trips according to sex (x-axis). Colors represent species: SAFS (orange) and SASL (teal).



**Table 3. 1** Number of individuals (*N ind*), number of trips (*N trips*), trip duration (days), and trip distance (km) for each species-sex group included in this study. Results reported as mean  $\pm$  SD.

Group	<i>N ind</i>	<i>N trips</i>	Trip Duration (days)	Trip Distance (km)
SAFS Females	8	77	3.65 $\pm$ 3.38	221.81 $\pm$ 286.06
SASL Females	4	72	2.16 $\pm$ 2.02	81.76 $\pm$ 64.08
SAFS Males	9	57	8.32 $\pm$ 11.44	298.03 $\pm$ 277.97
SASL Males	14	219	2.87 $\pm$ 2.50	178.52 $\pm$ 184.09

### 6.3 Utilization distributions

In this study we consider Utilization Distribution at 95% as the home range and at 50% as the core area for foraging. SAFS home range area is smaller than SASL in both sexes, and female home range and core area is smaller than males in both species (Table 3.2). SAFS females home range and core area was overlapped in 0.43 - 0.50 by SASL females, whereas SASL females were overlapped 0.70 - 0.83 by SAFS females in 95% and 50 % UD, respectively. SAFS males home range and core area was overlapped in 0.85 - 0.71 by SASL males, whereas SASL males were overlapped 0.41 - 0.43 by SAFS males in 95% and 50 % UD, respectively (Table 3.3). In regards to intraspecific overlap, SAFS females were overlapped 0.83 - 1 by SAFS males, whereas SASL females were always completely overlapped by SASL males in 95% and 50 % UD, respectively (Table 3.3).

**Table 3. 2** Area in km<sup>2</sup> of 25%, 50%, 75% and 95% Utilization Distributions (UD) for each species and sex group.

Group	UD 25%	UD 50%	UD 75%	UD 95%
SAFS Females	172.48	1,074.52	2,779.49	9,166.06
SASL Females	333.46	1,645.23	4,931.07	14,450.21
SAFS Males	2,430.72	7,806.89	20,616.80	55,537.99
SASL Males	1,786.86	10,894.50	33,211.20	106,615.05

No significant differences were detected between the distribution of SAFS and SASL females for all UD (25% to 95%), confirming that all female foraging grounds overlap in space and are considered



similar by ANOSIM test. We found that SAFS and SASL males are only similar in the smallest UD area (25%), that is most proximate to the PSJ colony (Figure 3.4A-B). Between SAFS males and females, significant differences in UD occur at 75% and 95%; but 25% and 50% UD are not significantly different; suggesting sexual segregation in foraging grounds utilized at 75% and 95%. In contrast, in SASL males and females, only the home range (95% UD) is significantly different; meanwhile all other utilization distributions (25%-75%) overlap between the sexes (Figure 3.4C-D).

**Table 3. 3** Proportion overlap for each group for core area (50% UD) and home range (95 % UD). Values represent the proportion overlap for the group in the rows by the group in the column. Proportion overlap > 0.70 in bold.

		SAFS F	SAFS M	SASL F	SASL M
<b>95% UD</b>	SAFS F	-	<b>0.83</b>	0.43	<b>1.00</b>
	SAFS M	0.20	-	0.13	<b>0.85</b>
	SASL F	<b>0.70</b>	<b>0.82</b>	-	<b>1.00</b>
	SASL M	0.12	0.41	0.07	-
<b>50% UD</b>	SAFS F	-	<b>1.00</b>	0.50	<b>1.00</b>
	SAFS M	0.18	-	0.11	<b>0.71</b>
	SASL F	<b>0.83</b>	<b>1.00</b>	-	<b>1.00</b>
	SASL M	0.11	0.43	0.07	-

## 6.4 Foraging by hour of day

Hour of day had a significant effect ( $p < 0.01$ ) on the probability of foraging events in female SAFS and male SASL. Foraging events for female SAFS concentrates during night hours (0-5, 18 - 23 hours) and in SASL males in daylight hours (6 - 15 hours); whereas in the other groups the effect of hour of day was not identified as significant (Table 3.4, Figure 3.5, Appendix C2).


**Table 3. 4** Results from the General Additive Mixed-Models to test for effect of Hour in the probability of foraging events in SAFS females, SASL females, SAFS males and SASL males. N represents the number of observations per group.




---

<b>Group</b>	<b><i>N</i></b>	<b>Hour</b>	<b>p-value</b>	<b>Log-likelihood</b>	<b>df</b>	<b>Adj. R Sq.</b>
SAFS Females	<i>24,818</i>	6.208	<0.01	-56037.49	3.12	0.000619
SASL Females	<i>22,650</i>	0.056	0.339	-53855.38	1.16	1.24e-06
SAFS Males	<i>60,927</i>	0	0.908	-148151.6	1	-5.31e-10
SASL Males	<i>91,209</i>	5.532	<0.01	-216553.8	3.08	0.000254





**Figure 3.** 4 Maps with filled contours of Utilization Distributions (UD) for 50% core foraging range (dotted line) and 95% home range (continuous line) for A) SAFS females (orange) and SASL females (aquamarine); B) SAFS males (brown) and SASL males (teal); C) SAFS females (orange) and SAFS males (brown); D) SASL females (aquamarine) and SASL males (teal). Black lines represent coastline and isobaths for depths of 2,000, 4,000 and 6,000 meters.



**Figure 3. 5** Density plots of the foraging events (y-axis) by hour of day (x-axis) for A) SAFS females (orange) and SASL females (aquamarine); B) SAFS males (brown) and SASL males (teal); C) SAFS females (orange) and SAFS males (brown); D) SASL females (aquamarine) and SASL males (teal). Shaded areas indicate dark hours (0 - 5, 18 - 23 hours) in study area.



## 6.5 *Associated environmental conditions*


Mean values of environmental variables encountered while foraging are significantly different between species (Wilcoxon, Test,  $p < 0.001$ ), except for SST between females (ESM 2). Descriptive parameters of the environmental variable values (minimum, maximum, median, mean and SD) for locations where animals were presumed to be foraging are reported in the Appendix C3.

In HCS, higher DC (Distance to Coast), higher SST, lower Chl-a, increased Front Proximity and higher Front Gradient characterizes more oceanic waters and the opposite describes coastal waters. Univariate modelling of environmental variables shows that SAFS females are associated to higher SST, lower Chl-a, higher Front Proximity and higher Front Gradient and higher DC, characterizing more oceanic variables. In contrast, SASL females feature lower SST, higher Chl-a, lower Front Proximity and lower Front Gradient and lower DC. In females, SST, Front Proximity and Front Gradients showed inverse responses (Figure 3.6, Appendix C4). Among males, differences are not so evident. SAFS and SASL are positively and similarly associated to SST and Chl-a. However, SAFS males were associated to higher Front Proximity and Gradient and lower DC in comparison to SASL males (Figure 3.6, Appendix C4).

In preparation for the forward selection process, the only variable discarded was SST for SAFS female model, due to 39% missing information. All remaining covariates were checked for collinearity and had Spearman rank  $\rho < 0.55$  (Appendix C6). Throughout the GAMM forward selection process, in general, adding terms improved model performance (ANOVA,  $p < 0.01$ , Table 3.5). Best fitting models for SAFS females included Chl-a, Front Proximity, Front Gradient and DC. Best fitting models for SASL females and SASL males included all covariates and best model for SAFS males included only SST, Chl-a and Proximity to Fronts (Table 3.5).

**Table 3. 5** Results from the forward selection process of covariates that explain environment during foraging using a General Additive Mixed-Models. *Bold* indicates best fitting model and *N* represents the number of observations per group. Significant Chi square values have a (\*) for *p*-values < 0.01 and (.) for *p*-values < 0.05. SST was not included in the SAFS females models due to missing values in 39% of foraging locations.

Group	SST	Chl-a	Proximity to Front	Gradient of Front	DC	Log-likelihood	df	Adj. R sq
SAFS Females <i>N = 14,286</i>	-	432.8 (*)				-32,203.29	3.99	0.0494
	-	427.4 (*)	141.8 (*)			-32,328.98	6.98	0.0629
	-	608.2 (*)	138.9 (*)	231.6 (*)		-33,964.49	9.7	0.0932
	-	<b>580.1 (*)</b>	<b>174.3 (*)</b>	<b>282.8 (*)</b>	<b>608.1 (*)</b>	<b>-35,384.95</b>	<b>12.7</b>	<b>0.165</b>
SASL Females <i>N = 18,423</i>	217.7 (*)					-43,395.54	3.98	0.0287
	153.4 (*)	265.4 (*)				-44,243.44	6.95	0.0815
	194.2 (*)	290.3 (*)	253.5 (*)			-45,400.05	9.93	0.113
	167.5 (*)	124.8 (*)	72.23 (*)	325.30 (*)		-46,215.19	12.77	0.154
SAFS Males <i>N = 51,849</i>	<b>83.4 (*)</b>	<b>111.3 (*)</b>	<b>22.6 (*)</b>	<b>497.9 (*)</b>	<b>7035.2 (*)</b>	<b>-56,823.42</b>	<b>15.43</b>	<b>0.279</b>
	221.0 (*)					-126,784.5	3.96	0.0292
	220.1 (*)	1.3 (*)				-126,812.9	4.85	0.0291
	<b>310.5 (*)</b>	<b>2.5 (*)</b>	<b>1269.6 (*)</b>			<b>-133,128.4</b>	<b>8.06</b>	<b>0.124</b>
SASL Males <i>N = 65,040</i>	224.3 (*)	0.0	1051.0 (*)	298.9 (*)		-131,095.6	9.96	0.104
	249.5 (*)	0.9 (.)	710.5 (*)	300.1 (*)	62.7 (*)	-133,079.6	13.59	0.103
	71.35 (*)					-159,739.4	3.58	-0.005
	31.5 (*)	261.9 (*)				-161,470.9	6.56	-0.001
SASL Males <i>N = 65,040</i>	17.2 (*)	237.3 (*)	63.7 (*)			-161,359.8	9.35	0.006
	26.8 (*)	219.7 (*)	63.2 (*)	150.7 (*)		-164,919.2	12.17	0.009
	<b>67.8 (*)</b>	<b>161.8 (*)</b>	<b>245.4 (*)</b>	<b>597.0 (*)</b>	<b>2,111.3 (*)</b>	<b>-168,627.1</b>	<b>15.8</b>	<b>0.032</b>



**Figure 3. 6** Probability of foraging events (y-axis 1 and 2) in relation to ranges of environmental variables (x-axis 1 and 2) for A) SAFS females (orange) and SASL females (aquamarine) and B) SAFS males (brown) and SASL males (teal). Colors of labels on x- and y-axis are indicative of values for each species and sex group.





## 7. Discussion

The results of our study show that sympatric otariids in the Humboldt Current System have multiple strategies to segregate while foraging in the pelagic environment and that body mass is not the sole driver for segregation in space and time, but rather a combination of energetic constraints according to sex, reproductive roles linked to specific dietary resources.

### 7.1 *Foraging tips and utilization distributions*

We found that although they are smaller in size, both SAFS females and males, make trips of longer distance and duration in comparison to both sexes of SASL, rejecting our hypothesis that larger animals make longer trips. In terms of area, SASL of both sexes cover larger areas, both in the home ranges and core areas, but do so with more consecutive shorter trips compared to SAFS. This reveals different use of the foraging grounds at interspecific level. SASL have larger home ranges as a group, covering a larger area parallel to the coastline that envelopes the SAFS home range. In contrast, SAFS travel farther offshore during feeding trips to specific areas (Figure 3.1A).

In agreement with our predictions based on previous studies focused on sexual segregation of otariids, females of both species show more constrained foraging trips in terms of distance traveled, home ranges and core areas in comparison to males of both species (Page et al., 2006; Staniland & Robinson, 2008). Interestingly, neither sex of SASL show a considerable significant difference in the duration of their foraging trips (marginal p-value), albeit they are the species with the highest dimorphism. This result is in agreement with previous findings for this species in the Falkland Islands (Baylis et al., 2016). However, SASL sex-based differences should be further explored with the addition of tracking with depth recorders to better understand the foraging effort invested in diving while feeding.

In terms of spatial segregation, female locations are not significantly different throughout their home ranges or core areas. Among females, 0.70 - 0.83 of SASL UD was overlapped by SAFS; while 0.43 - 0.50 of SAFS UD was overlapped by SASL in 50 and 95% UD. These high proportions of overlap can be influenced by the main prey items reported for these species, red squat lobster (*Pleuroncodes monodon*) and Peruvian anchovy (*Engraulis ringens*). Red squat lobsters have been reported as a main prey for SASL based on identification of hard structures in scats (Cárdenas-Alayza et al., 2021;



Sarmiento-Devia et al., 2020; Sielfeld et al., 2018). Distribution of red squat lobsters are associated to cold coastal waters and found in the first 30 km from the coast (Yuffra et al., 2018), explaining shorter trips within the coastal habitat covered by SASL. Red squat lobsters and anchovies have a high degree of spatial overlap in areas 50 km from the coastline. However, Peruvian anchovies, the preferred prey item by SAFS and second preferred by SASL, expand further offshore (Gutiérrez et al., 2008). Anchovies make dense aggregation sites, probably driving the preferred foraging grounds for SAFS as already reported in other regions (Baylis et al., 2014; Franco-Trecu et al., 2013; Naya et al., 2002).

It is important to note that female fur seals and sea lions were not tracked simultaneously, or in the same calendar months/seasons. This study targeted females that were nursing recently newborn pups, and tracked SAFS females from November to January and SASL females from February to March. This is due to a two month difference in breeding seasons (SAFS breed during October-December and SASL breed during January-March). In Peru, these species have temporal segregation in their breeding seasons, contributing yet another mechanism for segregation known as allochrony to partition resources during energetically critical times, which has been postulated as one of the founding drivers behind speciation (Taylor & Friesen, 2017).

Also, in both species weaning of offspring is flexible and can be extended between 6 - 36 months, with high interannual variability (Majluf, 1987; Soto, 2004). Thus we consider that the spatial use patterns of lactating females in this study are representative of the most energetically constrained feeding trips during which the timing of feeding trips is critical towards the survival of small pups that do not withstand extended periods without provisions, compared to larger offspring (Campagna & Harcourt, 2021). Since trip duration, distance, and areas can change as offspring grow older, we recommend future studies to explore if female foraging behavior changes according to breeding season offspring number and age (Drago et al., 2010, 2015, 2021; Franco-Trecu et al., 2012).

In the spatial realm between males, the proportion of SAFS areas covered by the 50 and 95% UD are overlapped in large proportions by SASL (0.85 - 0.71). However, because of the distributions of foraging locations in space, the similarity of distributions after 999 iterations was only significant for the 25% UD (e.g., close to the PSJ colony). In male SAFS, the focalized number of foraging locations and the bimodal distribution in Distance to Coast (Appendix C4) suggest a repetitive exploration of specific foraging areas within the wider and enveloping SASL male range. In contrast, SASL males





make shorter and faster trips. Our results show that the varying lengths in distance and duration of foraging trips can be indicative of specialized foraging grounds, especially in SAFS males, suggesting potential foraging site fidelity in HCS. This has been reported in other locations and should be further explored for this study area (Baylis et al., 2017; Knox et al., 2018).

Since males are not constrained in their foraging trip distance and duration in comparison to females that have dependent offspring ashore, post-breeding dispersal is common in male pinnipeds. Regardless, this study shows that the South American otariids do not depart on a migration, but rather obtain their energetic demands in a local range of 55,000 - 106,000 km<sup>2</sup>. These data are evidence that these sympatric species have evolved and adapted to obtain their resources locally in the nearby productive upwelling waters of HCS, occupying rookeries like PSJ year round.

When comparing sexes among sea lions, the analyses of similarity suggests that core areas and home ranges are significantly similar (25 - 95% UD). In SAFS similarity occurs in 25 - 50% UD, but remaining 75 - 95% UD are significantly different. This supports the theory that intraspecific competition within SASL can be enhancing stable coexistence with SAFS by decreasing interspecific competition pressures, as postulated by the Lotka-Volterra model (Schoener, 1974). On the contrary, reduced intraspecific competition in SAFS could negatively affect adaptation to current interspecific pressures. This may be one of the confounding reasons behind the current lack of population recovery and of decreasing SAFS in Peru, that also exists in this same study population (Cárdenas-Alayza et al., 2021).

## ***7.2 Foraging by hour of day***

In this study we found that the probability of foraging events incremented in the night hours for SAFS females and in the daytime for SASL males. Partly validating our hypothesis of nocturnal foraging patterns in SAFS and daylight foraging in SASL. Nocturnal foraging in SAFS females may be a synergistic effect between the energetic constraints to provision for their young and the vertical diel migration of prey items for SAFS such as Peruvian anchovies, squids and myctophids, making them more available in the night hours (Ayón et al., 2008). Since we did not find a significant effect for hour of day in male SAFS, we reject our hypothesis that this pattern holds for all SAFS. Since males do not have energetic constraints by offspring, daytime foraging segregation in SASL males may be



a reflection of a reduction in competition with another otariid age-class, predator species or preference for a specific prey (Chilvers & Wilkinson, 2009). Recent studies have found same age-class in Galapagos sea lions (*Zalophus wollebaeki*) have different foraging modes (pelagic, benthic and nocturnal divers, Schwarz et al. 2021). Thus, it is possible that there is individual specialization; or that individual can also switch between strategies according to prey availability. Thus, we recommend this is further explored in this ecosystem. Future studies involving depth recorders would again help refine estimation of foraging, since the present analyses may have associated errors in the identification of foraging events based on locations alone.

### ***7.3 Associated environmental conditions***

In general, environmental variables associated with foraging events show interspecific differences. When compared, environmental habitat features of SASL females are linked to the coastal habitat that is characterized by lower DC, less Proximity to Fronts and Gradients and higher Chl-a values. Whereas SAFS females are associated to a more pelagic environment characterized by lower Chl-a, higher Proximity and Front Gradients and higher DC values. It is important to note that the range in SST amongst female locations was wide and not significantly different. Thus, SST is not recommended as an explanatory environmental variable to compare these groups. Evidence from hard parts analysis in PSJ reports that SAFS consume Peruvian anchovy and squids (Arias-Schreiber, 2003) that are characteristic of the pelagic habitat (Argüelles et al., 2012), in contrast to the more coastal red squat lobster (Yuffra et al., 2018), which is the primary prey item of SASL (Gutiérrez et al., 2008); lending support to the environmental variables associated with female foraging.

Among males, best fitting models for SAFS males included SST, Chl-a and Proximity to Fronts, whereas SASL males included all environmental covariates. SASL males have lower SST, lower Chl-a, lower Proximity to Fronts and lower DC, which could be a product of individual strategies. Ranges in the environmental variables targeted by SAFS and SASL males are highly overlapping; conclusions should be taken with caution and further studies exploring individual strategies should be conducted. We also suggest incorporating additional environmental variables to better explain male foraging environment, such as those related to the vertical structure of the water column.

Sympatric otariids in the Humboldt Current System show some level of segregation while foraging



at sea, partly associated with reproductive role, body size and species. Whether these segregation mechanisms have changed or not in the recent years is unknown, but answering this question is key to understanding the recent trends in population sizes observed in both species after the 1997-1998 El Niño recovery (Cárdenas-Alayza et al., 2021). To tackle the beforementioned we recommend further studies using trophic signatures that archive information from 4-5 years to capture potential trophic shifts in individuals (e.g., stable isotopes and/or fatty acid analyses).

Segregation mechanisms only make sense if prey are partly limited and spatially partitioned. In the case of very high availability of prey, coexistence could occur without much need of segregation. However, in the case of food shortage, segregation mechanisms could sustain coexistence, by taking advantage of differential abilities to cope with such a situation. In the latter case, the differences observed in the foraging ecologies may depict a situation of one species outcompeting another, which couldn't be considered as a coexistence mechanism. If the segregation we observed is stable in time and if both populations keep coexisting, it might indeed be induced by their ability to segregate. This raises the question of the dynamics of segregation mechanisms with respect to food availability, and calls for maintenance of the long term monitoring and research programs to compare these indices over time.




## APPENDIX C1

**Table C1. 1** Species, sex, individual identification (ID) deployment date, weight (kg), length (cm), number of days tracked and number of trips recorded for each individual.

No.	Species	Sex	ID	Date Deployment	Weight	Length	# Days	# Trips
1	SAFS	F	A00_157689	November 19, 2015	38.00	114.0	57.69	16
2	SAFS	F	A02_157694	November 19, 2015	47.80	124.0	23.19	6
3	SAFS	F	A03_157695	November 19, 2015	52.20	125.0	53.83	9
4	SAFS	F	A04_157696	November 19, 2015	56.00	131.0	75.38	11
5	SAFS	F	A05_157697	November 19, 2015	40.10	115.0	45.38	9
6	SAFS	F	A06_157698	November 20, 2015	47.60	121.0	49.40	15
7	SAFS	F	A07right_8	November 20, 2015	33.70	111.0	34.50	8
8	SAFS	F	44Y_157700	November 20, 2015	48.80	127.0	26.44	4
9	SASL	F	F01_157689	February 24, 2017	74.50	146.0	72.77	23
10	SASL	F	F02_157697	February 25, 2017	63.30	135.0	78.02	29
11	SASL	F	F04_157699	February 25, 2017	85.00	143.0	76.31	23
12	SASL	F	F05_157700	February 25, 2017	86.00	148.5	48.85	19
13	SAFS	M	36Y_143666	November 15, 2014	118.80	158.0	56.19	5
14	SAFS	M	39Y_143667	November 15, 2014	117.30	151.0	43.68	12
15	SAFS	M	41Y_143663	November 15, 2014	107.20	150.0	66.19	10
16	SAFS	M	C55_143664	November 13, 2016	113.30	148.0	60.15	3
17	SAFS	M	C56_143665	November 13, 2016	99.20	151.0	70.38	9
18	SAFS	M	C57_157694	November 13, 2016	106.60	147.0	50.40	3
19	SAFS	M	C58_157695	November 14, 2016	103.40	159.5	46.88	9
20	SAFS	M	C59_157697	November 14, 2016	102.40	158.0	40.87	3
21	SAFS	M	C60_157699	November 14, 2016	94.10	153.5	70.87	4
22	SASL	M	03V_134432	November 15, 2013	151.40	174.5	73.75	22
23	SASL	M	04V_134430	November 15, 2013	237.20	205.0	70.79	14
24	SASL	M	05V_134433	November 15, 2013	219.40	198.0	72.29	18
25	SASL	M	06V_134429	November 16, 2013	238.60	203.0	73.75	16
26	SASL	M	12V_134431	November 19, 2013	184.10	183.0	67.40	14
27	SASL	M	13V_141850	November 13, 2014	196.60	196.5	36.75	11
28	SASL	M	15V_141852	November 13, 2014	201.60	200.0	70.58	7
29	SASL	M	17V_141853	November 14, 2014	162.40	190.0	73.71	14
30	SASL	M	19V_157686	November 17, 2015	160.60	196.0	61.90	12
31	SASL	M	20V_157685	November 17, 2015	139.30	189.0	60.88	14
32	SASL	M	22V_157688	November 17, 2015	191.90	191.0	67.42	25
33	SASL	M	23V_157693	November 17, 2015	131.60	170.5	71.96	21
34	SASL	M	25V_157690	November 18, 2015	137.80	185.0	72.92	13
35	SASL	M	27V_157692	November 18, 2015	172.00	191.1	69.50	20

## APPENDIX C2



**Figure C2.1** Smoother response for probability of foraging (y-axis) according to hour of day (x-axis) for A) SAFS females, B) SAFS males, C) SASL females and D) SASL males from General Additive Mixed-Models. Significant relationships ( $p < 0.01$ ) were found for A) SAFS females and B) SASL males. Icons represent daylight (sun) and dark hours (moon).




## APPENDIX C3

*Table C3. 1 Descriptive statistics for the environmental covariates extracted for foraging event locations for each species and sex group.*


<b>Group</b>	<b>Environmental covariate</b>	<b>SST</b>	<b>Chl-a</b>	<b>Proximity to Front</b>	<b>Gradient of Front</b>	<b>Distance from Coast</b>
<b>SAFS Females</b>	<i>N obs</i>	4,408	6,294	5,852	5,862	6,348
	Min	19.28	0.71	0.01	1.28	4.00
	Max	23.03	5.92	0.11	93.50	57.00
	Median	21.29	2.15	0.04	9.88	19.00
	Mean	21.16	2.27	0.04	14.29	22.47
	SD	0.92	0.84	0.01	16.30	11.09
<b>SASL Females</b>	<i>N obs</i>	4,269	4,392	4,356	4,356	4,170
	Min	19.07	0.63	0.01	0.57	9.00
	Max	23.33	20.20	0.11	50.00	26.00
	Median	21.26	4.58	0.07	1.94	15.00
	Mean	21.12	5.85	0.07	3.69	13.93
	SD	0.84	3.88	0.01	7.00	4.16
<b>SAFS Males</b>	<i>N obs</i>	12,264	12,474	13,197	13,197	12,996
	Min	16.11	0.22	0.01	0.48	7.00
	Max	25.22	23.47	0.13	39.13	148.00
	Median	21.08	3.26	0.05	3.55	42.00
	Mean	20.97	3.95	0.05	7.48	51.20
	SD	1.39	3.38	0.02	8.30	34.27
<b>SASL Males</b>	<i>N obs</i>	14,223	17,367	18,036	18,075	18,345
	Min	15.41	0.18	0.00	0.64	4.00
	Max	26.11	25.23	0.11	98.57	147.00
	Median	20.71	2.05	0.03	8.92	23.00
	Mean	20.48	3.05	0.04	13.88	40.46
	SD	1.89	3.10	0.02	14.69	36.11



# APPENDIX C4



**Figure C4. 1** Environmental conditions (SST, Chl-a, Front Proximity, Front Gradient and DC) associated to female SAFS (orange) and SASL (aquamarine) foraging events. A) Boxplots depicting significant differences (\*\*\*) indicates  $p < 0.01$ . B) Density curves show distribution of frequencies. SST between SAFS and SASL is not significantly different (N.S.).



**Figure C4. 2** Environmental conditions (SST, Chl-a, Front Proximity, Front Gradient and DC) associated to male SAFS (brown) and SASL (teal) foraging events. A) Boxplots depicting significant differences (\*\*\*) indicates  $p < 0.01$ . B) Density curves showing distribution of frequencies.





## APPENDIX C5

**Table C5. 1** Deviance explained by univariate General Additive Models (GAM) of each environmental covariate explored in response to foraging events. *N* = number of observations included in each group.

Group	Environmental covariate	% Deviance
SAFS Females <i>N</i> = 14,286	SST	5.37%
	Chl-a	7.31%
	Proximity to Front	1.90%
	Gradient of Front	3.72%
	DC	6.37%
SASL Females <i>N</i> = 18,423	SST	3.95 %
	Chl-a	6.60 %
	Proximity to Front	4.29 %
	Gradient of Front	12.50 %
	DC	11.20 %
SAFS Males <i>N</i> = 51,849	SST	5.21 %
	Chl-a	2.02 %
	Proximity to Front	7.49 %
	Gradient of Front	3.82 %
	DC	2.75 %
SASL Males <i>N</i> = 65,040	SST	3.17%
	Chl-a	0.76 %
	Proximity to Front	0.98 %
	Gradient of Front	0.39 %
	DC	4.05 %



APPENDIX C6




Figure C6. 1 Correlation chart (Spearman rho,  $p=0.05$ ) between the environmental covariates (SST, Chl-a, Front Proximity, Front Gradient, Distance to Coast) for A) SAFS females and B) SASL females. SST not included in for SAFS females due to 39% missing values.





Figure C6. 2 Correlation chart (Spearman rho,  $p=0.05$ ) between the environmental covariates (SST, Chl-a, Front Proximity, Front Gradient, Distance to Coast) for A) SAFS males and B) SASL males

## CHAPITRE 4

*Sympatric otariids increase trophic segregation in response to warming ocean conditions in Peruvian Humboldt Current System*



**Cárdenas-Alayza S**, Adkesson, Edwards, M.R., M.R., Hirons, A.C. Gutiérrez D, Tremblay Y, Franco-Trecu, V. Sympatric otariids increase trophic segregation in response to warming ocean conditions in Peruvian Humboldt Current System.

Manuscript submitted for publication to PLOS ONE



## Résumé

La détermination des habitudes trophiques des communautés de prédateurs est essentielle pour comprendre les interactions interspécifiques et leurs réponses aux fluctuations environnementales. Les otaries à fourrure sud-américaines, *Arctocephalus australis* (SAFS) et les lions de mer Sud-Américains *Otaria byronia* (SASL), coexistent le long des côtes du Pérou. Récemment, des événements de réchauffement de l'océan (2014 - 2017) pouvant diminuer et appauvrir la biomasse des proies se sont produits dans le système péruvien du courant de Humboldt (HCS). Dans ce contexte, notre objectif était d'évaluer l'effet des événements de réchauffement sur la ségrégation de niche inter- et intra-spécifique à long terme. Nous avons collecté des vibrisses de SAFS (55 femelles et 21 mâles) et SASL (14 femelles et 22 mâles) à Punta San Juan, Pérou. Nous avons utilisé les valeurs  $\delta^{13}\text{C}$  et  $\delta^{15}\text{N}$  archivées en série dans les vibrisses pour construire une série chronologique mensuelle entre 2005 et 2019. À partir de la même série chronologique, nous avons utilisé l'élévation du niveau de la mer pour déterminer différentes conditions environnementales en utilisant une méthode de détection de ruptures. Les zones d'ellipse (SIBER) ont estimé la largeur de niche des groupes d'espèces-sexes et leur chevauchement. Nous avons détecté un changement moyen des conditions environnementales montrant deux périodes distinctes (P1 : janvier 2005 - octobre 2013 ; P2 : novembre 2013 - décembre 2019). La réduction du  $\delta^{15}\text{N}$  dans tous les groupes au cours de la période 2 suggère des valeurs de base appauvries avec des effets bottom-up, un changement vers la consommation de proies de niveau trophique inférieur, ou les deux. La réduction du chevauchement entre tous les groupes au cours de la période 2 laisse supposer un assemblage plus redondant au cours de la période 1, plus froide, et un assemblage plus ségrégué sur le plan trophique au cours de la période 2, plus chaude. Les femelles SASL montrent la plus grande variation en réponse au scénario de réchauffement (P2), réduisant à la fois la surface de l'ellipse et les valeurs moyennes de  $\delta^{15}\text{N}$ . La plasticité pour s'adapter aux environnements changeants et se nourrir sur une source de nourriture plus disponible en absence de pression de pêche pourrait être plus avantageuse pour les femelles SASL, bien que les effets trophiques bottom-up soient temporaires. Cela pourrait expliquer la taille plus importante de la population de SASL au Pérou, en comparaison à la population de SAFS, plus petite et en déclin.

**Mots clés :** niche isotopique ; otaries à fourrure ; lions de mer ; changement trophique



## Abstract

Determining trophic habits of predator communities is essential to measure interspecific interactions and response to environmental fluctuations. South American fur seals, *Arctocephalus australis* (SAFS) and sea lions *Otaria byronia* (SASL), coexist along the coasts of Peru. Recently, ocean warming events (2014 – 2017) that can decrease and impoverish prey biomass have occurred in the Peruvian Humboldt Current System (HCS). In this context, our aim was to assess the effect of warming events on long-term inter- and intra-specific niche segregation. We collected whisker from SAFS (55 females and 21 males) and SASL (14 females and 22 males) in Punta San Juan, Peru. We used  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values serially archived in otariid whiskers to construct a monthly time series for 2005 – 2019. From the same time series we used sea level rise to determine different environmental conditions using a change point analysis. Ellipse areas (SIBER) estimated niche width of species-sex groups and their overlap. We detected a mean change in environmental conditions showing two distinct periods (P1: January 2005 - October 2013; P2: November 2013 - December 2019). Reduction in  $\delta^{15}\text{N}$  in all groups during Period 2 suggests impoverished baseline values with bottom-up effects, a shift towards consuming lower trophic level prey, or both. Reduced overlap between all groups in P2 lends support of a more redundant assemblage during the colder P1 to a more trophically segregated assemblage during warmer P2. SASL females show the largest variation in response to the warming scenario (P2), reducing both ellipse area and  $\delta^{15}\text{N}$  mean values. Plasticity to adapt to changing environments and feeding on a more available food source without fishing pressure can be more advantageous for female SASL, albeit temporary trophic bottom-up effects. This helps explain larger population size of SASL at Peru, in contrast to the smaller and declining SAFS population.

**Keywords:** isotopic niche, fur seals, sea lions, trophic shift



## 1. Introduction

Individual foraging behavior and trophic interactions determine the flow of energy along the food web, and ultimately these drive population dynamics, community structure, and most ecosystem processes (Arim et al., 2010). However, unpredictable and highly variable environments can also affect trophic behavior (Yurkowski et al., 2018). Top predators, such as pinniped species, are considered structural components of communities and ecosystem functioning (Pauly et al., 1998). Determining trophic habits of predators in a community is essential to measure the potential interspecific niche overlap and strength of interactions as well as the effects of environment changes (Layman et al., 2007).

Two otariid species breed in sympatry along the coast of South America: the South American fur seal, *Arctocephalus australis* (SAFS), and the South American sea lion, *Otaria byronia* (SASL). Information from areas where these otariids coexist serves as an interesting model to assess interspecific segregation (Franco-Trecu et al., 2014; Riverón et al., 2021; Szteren et al., 2018). However, the majority of research has focused on the Atlantic Ocean where SAFS are recognized as pelagic and SASL as benthic; little is known about their segregation in the upwelling ecosystem of the Humboldt Current System (HCS) in the Pacific Ocean. In contrast to the Atlantic coast that has a gradually declining offshore shelf, Peru's coast is characterized by a narrow continental shelf and shallow productive upwelling waters with a relatively shallow thermocline related to an upper minimum oxygen layer at depths of approximately 50-80 m on average (Bertrand et al., 2010, 2011; Demarcq, 2009; Echevin et al., 2008). These features limit the potential habitat for otariid prey to be linked to the benthos (Hückstädt et al., 2014; Trillmich et al., 1987), making pelagic resources common for tertiary consumers in HCS (Arias-Schreiber, 2003; Sarmiento-Devia et al., 2020). Thus, in this productive, shallow environment foraging on highly patchy, dense prey resources near the surface may preclude the advantages of searching for more evenly distributed, but less dense, prey resources on the benthos (Arnould & Costa, 2006).

HCS is very dynamic since it is characterized by recurring El Niño Southern Oscillation (ENSO) events, which can vary in intensity and duration. ENSO is a coupled oceanic-atmospheric cycle of alternating warm El Niño and cold La Niña events, and it is the most prominent climate signal on Earth (McPhaden et al., 2006). When HCS is affected by an ENSO event, wind pressure drops, sea surface temperature rises, and reductions of primary productivity occur in the marine environment, altering distribution of habitats and associated prey biomass. 'Extraordinary' ENSO events that





occurred 1982-83 and 1997-98 are well documented for dramatically altering species composition, reducing zooplankton and fish biomass, and causing mortality in higher trophic levels (Barber & Chavez, 1983; Tam et al., 2008; Tovar et al., 1985). However, the effects on trophic adaptation during weaker events that do not cause direct mortality, but impact reproduction, juvenile survival, and weaning have not been thoroughly investigated.

A warming climate is currently altering global ecosystem structure and driving species distributions to higher latitudes, thereby leading to altered interspecific interactions with unspecified consequences (du Pontavice et al., 2020; Walther et al., 2002). Climate driven shifts are more pronounced at the poles where deviations from long-term climatic variables are more striking (Kortsch et al., 2015; Parmesan & Yohe, 2003). However, in the past decade a series of warming events have taken place in the Pacific Ocean that may affect the organization of predator communities in the Peruvian HCS. Long-term warming climatic/oceanic events, or heatwaves known as “The Blob”, have impacted the Pacific Ocean in recent years (N. A. Bond et al., 2015). The Blob was first detected in the Gulf of Alaska in 2013-2014 and is a result of a persistent pattern of higher than normal rates of sea level pressure and weak coastal winds. In Peru, El Niño conditions occurred between 2014 and 2015 (Graco et al., 2016), and were followed by the strong El Niño event of 2015-2016 that affected the entire Pacific Ocean (Robinson, 2016). This event was then followed by the effects of ‘El Niño costero’ that was a more localized event detected in the El Niño 1+2 region between January and February of 2017 (Paulino et al., 2019). During these years, studies suggest that physical anomalies of these warming ocean events generated a biological response similar to those of El Niño (Espinoza-Morriberón et al., 2017). All the aforementioned events involve the rise of sea level forced by the passage of remotely generated and coastal trapped waves in response to heat expansion (Ryan & Noble, 2002). The thermocline, nearshore depth of chlorophyll-a, and the nutricline deepened, decreasing nutrient enrichment in the euphotic zone. Weak winds caused the reduction in vertical nutrient fluxes to the depleted euphotic zone. A deeper concentration of chlorophyll-a decreased phytoplankton biomass, causing a shift in community structure toward smaller phytoplankton which can have bottom-up effects on food and energy availability at higher trophic levels (Zaba & Rudnick, 2016).

Upper trophic levels predators, like pinnipeds, act as sentinels to trophodynamic and species assemblage changes lower in the food web through their diet (Hazen et al., 2019). Stable isotopes analysis (SIA) of different tissues has emerged as a key method to investigate the differences in the





foraging ecology of pinnipeds (Newsome et al., 2010). The  $\delta^{15}\text{N}$  value is used to estimate the trophic position (Post et al., 2002) and  $\delta^{13}\text{C}$  values predict the carbon sources used by the predators (Bearhop et al., 2004; DeNiro & Epstein, 1978) at different temporal scales, depending on the moment of production and the turnover rate of the tissue analyzed (Dalerum & Angerbjörn, 2005). The SIA of metabolically inert tissues, such as tooth dentine and whisker keratin with continuous growth, represents sequential archives that allow inference of foraging strategies at the individual level over long periods of time (Franco-Trecu et al., 2014; Kernaléguen et al., 2012). Based on previous studies that estimated whisker growth rates in this species (Lima et al., 2019), each whisker gives information for up to 4.8 years (Edwards, 2018). In our study, long-term niche differentiation in individuals was investigated by examining the isotopic signature in whiskers of male and female South American fur seals and South American sea lions. We aim to understand how niche segregation between species is affected by changing environmental, morphological, and/or behavioral traits.

Several authors have reported relationships showing that larger animals tend to consume higher trophic level prey (Barnes et al., 2010; Segura et al., 2015). South American sea lions are considerably larger in body size than fur seals and both species have a strong sexual dimorphism, with males being much larger than females (Ralls & Mesnick, 2002); thus we expect that SASL males have highest values of  $\delta^{15}\text{N}$  and that SAFS males have higher values than SAFS females. In terms of  $\delta^{13}\text{C}$ , we expect groups to follow the inshore-offshore  $\delta^{13}\text{C}$  gradient already reported in northern HCS (Espinoza et al., 2017). Since otariid females perform feeding trips during lactation, which usually last ~11 months (Schulz & Bowen, 2005), we suspect they are restricted to foraging grounds close to the breeding colony compared to males (Le Boeuf et al., 2000; Meynier et al., 2008). Thus, we expect both otariid females to show more enriched  $^{13}\text{C}$  values, indicative of higher productivity since they remain constrained closer to the land-based colony. In contrast, we expect males to have a wider range of  $\delta^{13}\text{C}$  from foraging more “freely” between coastal and pelagic zones. As reported, the Peruvian coast has been affected by a decrease in the biomass of local prey and an impoverishment in lower trophic levels during 2014 - 2017 warming ocean events (Espinoza-Morriberón et al., 2017; Graco et al., 2016). On a long-term timescale, we expect to find a reduction in the trophic position of otariids and a higher overlap in isotopic niche space during warmer events when pelagic (cold-water) resources are less available. Finally, due to the beforementioned energetic constraints linked to maternal attendance, we expect females will reflect a narrow range of isotopic signatures compared to males and therefore be less flexible to adapt to changes in the environment.



## 2. Materials and Methods

### 2.1 *Study site and sample collection*

Whiskers were collected from lactating females and adult males chosen at random during pinniped health assessment campaigns between 2010 and 2019 at Punta San Juan (PSJ), a marine protected area located along the southern coast of Peru (15° 22' S, 75° 11' W). Whiskers were collected from 76 SAFS (55 females and 21 males) and 36 SASL (14 females and 22 males).

Anesthesia of individual SASL and male SAFS was induced using a combination of midazolam, butorphanol, and medetomidine administered via plastic dart as previously reported (Adkesson, Chinnadurai, Balko, Jankowski, Langan, et al., 2019; Adkesson, Chinnadurai, Balko, Jankowski, Meegan, et al., 2019). Female SAFS were captured using a hoop net and then anesthetized with isoflurane gas (1 - 5% to effect) mixed with oxygen as previously reported (Jankowski et al. 2015). Anesthesia was performed by a board-certified specialist in zoological medicine. Morphometrics and physiological parameters were recorded. Body weights were obtained using a tripod and field scale to the nearest 100 grams (Mini Crane Scale 300 kg, OCS-L). Whiskers, including the follicle, were removed manually. All animals were determined to be in good health by a veterinarian based on physical examination findings and veterinary assessment of routine blood parameters.

### 2.2 *Sample analysis*

Each whisker was scrubbed with an abrasive plastic pad and rinsed with deionized water to remove surface contaminants. Whiskers were thoroughly dried at 60° C for a minimum of 24 hours and then cut into 2.5 mm fragments from base (proximal) to tip (distal), obtaining between 10 and 62 fragments according to the total length of the whisker. Every other fragment was prepared for stable isotope analysis while the remaining segments were reserved. We then subsampled each portion to obtain 0.6 - 0.8 mg (high precision microbalance Mettler Toledo MX5, precision=1µg). Fragments were placed in individual tin capsules, pelletized, and then sent for stable isotope analysis.

Samples were combusted and analyzed for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  at the Smithsonian Institution's Museum Conservation Institute (Suitland, MD) using a Thermo Delta V Advantage mass spectrometer in continuous flow mode coupled to a Costech 4010 Elemental Analyzer (EA) via a Thermo ConFlo IV (CF-IRMS) to determine natural carbon and nitrogen abundance and their isotopic ratios with an



analytical precision of  $\pm 0.2$  ‰ for both isotopes. A set of standards were run for every 10-12 samples. The standards included USGS40 and USGS41 (L-glutamic acid) as well as Costech acetanilide. All samples and standards were run with the same parameters; this included an expected reproducibility of the standards  $< 0.2$  ‰ ( $1\sigma$ ) for both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ . Stable isotope values were expressed in terms of  $\delta$  and were reported relative to the standard reference material, Vienna Pee Dee Belemnite (VPDB) standard for  $\delta^{13}\text{C}$  and atmospheric air ( $\text{N}_2$ ) for  $\delta^{15}\text{N}$ . The resulting isotope ratios for each whisker segment were converted and reported to the conventional values delta ( $\delta$ ) in parts per thousand (‰), according to:  $\delta X = (R_{\text{sam}} / R_{\text{st}}) - 1$ , where X stands for  $^{13}\text{C}$  or  $^{15}\text{N}$ , R for  $^{13}\text{C}/^{12}\text{C}$  or  $^{15}\text{N}/^{14}\text{N}$ ,  $R_{\text{sam}}$  for isotope ratio of the sample, and  $R_{\text{st}}$  for isotope ratio of the standard.

### ***2.3 Data analysis***

The most recent growth is located at the base of the whisker, and an individual whisker can represent several years' growth (Ginter et al., 2012; Hirons, 2001; Hirons et al., 2001). Based on an otariid whisker growth rate of 0.08 mm/day (Lima et al. 2019), each 2.5 mm segment represented approximately 30 days.

### ***2.4 Environment: sea level and sea temperature***

Monthly records for sea level (cm) for latitude  $15^\circ$  South, location of our study area between January 2005 to December 2019, were extracted from the Copernicus ocean products repository (<https://marine.copernicus.eu/>). Monthly sea surface temperature (SST) records collected by the naval base at San Juan de Marcona bay (Dirección de Hidrografía y Navegación, DHN), located less than 5 km from the sampling site during the study period. We constructed a time series of SST for each month during the fifteen-year period. Since sea level and SST were correlated during the study period (Spearman,  $r=0.74$ ,  $p<0.001$ ), to avoid redundancy, we selected sea level as our environmental variable because it reflects longer term processes caused by heat expansion in the water column. In contrast, due to the large scale atmospheric forcing SST is not good indicator of changes in upwelling intensity in a warming context (Demarcq, 2009). Change point analysis was applied to the sea level monthly time series to distinguish between different consecutive and consistent periods of environmental conditions in the 2005-2019 time series using R package changepoint (Killick & Eckley, 2014).



## 2.5 *Study groups*


Principal components analysis (PCA) of six morphometric measures (skull girth, axillary girth, snout to tail base, brachial girth, axilla to fore flipper, tail base to hind flipper) collected for SAFS and SASL (N=98) between 2010 -2019 were accessed and grouped by species and sex groups. Measures were standardized and PCA was built to determine significant clusters based on the morphometric differences between groups.

## 2.6 *Isotopic signatures*

By analyzing monthly means of individual seals' isotopic signatures, a mean for both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  was acquired per individual (i.e. month, year). Across the 2,941 fragments from 112 whisker samples, a total of 14 years of stable isotope data were recorded.

## 2.7 *Influence of the environment, species and sex on isotopic composition*

First, we used Linear Mixed-Effects Models (LMM) (Pinheiro & Bates, 2000) to examine the effect of body mass on the  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values from each sex group, using individual ID as random effect. Since body mass at the time of whisker collection is affected by feeding habits during recent months, in the LMM we included values from the last three fragments analyzed from each individual, integrating feeding information about the last six months. Second, we used LMM to examine the differences on the average  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values between groups (species and sex), periods, and their interactions as fixed effects. Models included individual identity as a random effect to account for repeated measures of each response variable on the different fragments of each whisker. We used a continuous autocorrelation function (Pinheiro & Bates, 2000) to model the serial correlation of the set of values of the response variables at the individual level. We compared the global model (all fixed effects and their interactions) with models without interaction with the Akaike Information Criterion (AIC) using the  $\Delta\text{AIC} > 2$  criterion (Bolker, 2008). Selected models were subject to the customary residual analyses (results not shown) and were found to have a satisfactory fit.



### 3. Isotopic niche


Stable Isotope Bayesian Ellipses in R (SIBER) (Jackson et al., 2011) were estimated to measure the isotopic niche width at 75% for SAFS and SASL. Mean isotopic values for each individual were used to estimate the ellipses since whisker fragments are temporally correlated and do not comply with sample independence. This method is a Bayesian version of Layman metrics (Layman et al., 2007) that can incorporate uncertainties such as sampling biases and small sample sizes into niche metrics (Jackson et al., 2011). Based on Markov-Chain Monte Carlo simulation, the SIBER approach obtains measures of uncertainty to construct parameters of ellipses in a way similar to the bootstrap. Standard ellipse areas corrected for small sample size ( $SEAC$ ) were used to compare the different sex-species groups (SAFS females, SASL females, SAFS males, SASL males) in the isotopic space in both periods and estimated the area of overlap of their isotopic niche using Maximum Likelihood standard ellipse areas ( $SEAC$ ). Values estimated for the overlap of each group is expressed in percentage.

Then, Layman metrics were estimated for the otariid community (the four species-sex groups) by period. Layman metrics are  $\delta^{13}C$  range,  $\delta^{15}N$  range, total area of convex hull (TA), mean distance to centroid from means (CD), mean nearest neighbor distance of the means, (MNND), and the standard deviation of the nearest neighbor distance (SDNND) following Layman et al. (2007).

## 4. Results

### 4.1 *Study groups*

PCA results show that horizontal axis 1 explains 87.9% of all variation, forming four different groups for SAFS females, SAFS males, SASL females and SASL males. Vertical axis reveals that 4.7% of variation is explained by similar morphometrics between SAFS males and SASL females (Figure 4.1). Since the majority of variation is explained for each species and sex, four species-sex groups (SAFS females, SASL females, SAFS males and SASL males) are compared throughout our study. LMM from females indicate that there is a positive correlation between body mass and  $\delta^{15}N$  values, but only at intraspecific level (p-value < 0.001 in both cases). While SASL females are twice the size of SAFS, the former had 2.22 ‰ lower in  $\delta^{15}N$  values (p-value < 0.001). In the male LMM, no body mass effect on  $\delta^{15}N$  was detected for either of the two species (p-values > 0.05 in both cases).



**Figure 4. 1** Principal components analysis of six morphometric measures (skull girth, axillary girth, snout to tail base, brachial girth, axilla to fore flipper, tail base to hind flipper) for each of the groups. SAFS females (pink, N = 35), SASL females (green, N= 22), SAFS males (blue, N=12) and SASL males (orange, N=29)




## 4.2 *Influence of environment on isotopic composition*

The average lengths of the whiskers analyzed were  $123.97 \pm 11.72$  mm and  $175.63 \pm 325.23$  mm for SAFS and SASL, respectively. Mean isotopic values by group, sample size, and fragments number analyzed are shown in Table 4.1. We found significant change points in the sea level time series, dividing it into two periods: Period 1, from January 2005 – October 2013 with a mean of 3.42 cm, and Period 2, from November 2013 – December 2019 with a mean of 8.06 cm (Figure 4.2A). A similar change point was found for SST, with only 1 month difference, giving support to period breakdown. There were significant differences in the mean whisker  $\delta^{13}\text{C}$  content among species-sex groups in both periods analyzed (see interaction terms in Table 4.2). In Period 1 the average  $\delta^{13}\text{C}$  was significantly lower for SAFS females compared with SAFS males and SASL males and females. However, this difference is even greater in Period 2 due to the change in opposite direction of mean  $\delta^{13}\text{C}$  isotopic value of SAFS females in comparison to the other three groups (Figure 4.2, Table 4.2). Mean  $\delta^{15}\text{N}$  values significantly differed between species-sex group and periods, but without interaction. SASL females had the lowest values and showed the greatest change in Period 2, being the only species-sex group that decreases the N mean value (Figure 4.2, Table 4.2).



**Table 4. 1** Isotopic signatures of Peruvian otariid whiskers according to periods investigated in present study. Number of whiskers (*N*), mean length (mm) by species and sex. Mean  $\pm$  SD of the stable isotope values ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) during two periods according to sea level. *N* fragments: sample size for species and sex.


Species	Sex	<i>N</i>	Length (mm)	Period 1			Period 2			
				$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	<i>N</i> fragments	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	<i>N</i> fragments	
SAFS	F	55	115.68 $\pm$ 26.78	-14.32 $\pm$ 0.39	19.03 $\pm$ 0.94	1,002	-14.87 $\pm$ 0.47	17.99 $\pm$ 1.01	182	
SASL	F	14	150.71 $\pm$ 32.61	-13.58 $\pm$ 0.50	19.05 $\pm$ 1.13	71	-13.72 $\pm$ 0.73	16.66 $\pm$ 1.60	322	
SAFS	M	21	132.26 $\pm$ 40.34	-13.91 $\pm$ 0.35	20.20 $\pm$ 0.83	406	-14.27 $\pm$ 0.36	18.71 $\pm$ 1.30	125	
SASL	M	22	200.54 $\pm$ 60.23	-13.45 $\pm$ 0.43	20.04 $\pm$ 1.54	719	-13.36 $\pm$ 0.58	18.58 $\pm$ 1.84	114	
		112					2,198			743



**Figure 4. 2** Time series for A) monthly sea level records from San Juan de Marcona bay, horizontal lines show Period 1 (P1, mean = 3.42 cm) and Period 2 (P2, mean = 8.06 cm), B)  $\delta^{13}\text{C}$  and C)  $\delta^{15}\text{N}$  isotopic signatures for all otariid groups.

**Table 4. 2** Linear Mixed-Effects Models (LMM) for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values. LMM for  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  whisker values including group (sex-species), period and their interactions as fixed effects and identity and whisker portion as random effect (continuous autocorrelation function). Estimates and p-values (in brackets) are shown for each variable. In bold we show the selected model by the Akaike Information Criterion (AIC). (\*\*\*) denotes  $p < 0.01$

Model	Intercept	SAFS M	SASL F	SASL M	Period 2	SAFS M *Period2	SASL F *Period2	SASL M *Period2	AIC
<b><math>\delta^{13}\text{C} \sim \text{SpecieSex} * \text{Period}</math></b>	<b>-14.40</b> (***)	<b>0.35</b> (***)	<b>0.47 (***)</b>	<b>0.88 (***)</b>	<b>-0.34</b> (***)	<b>0.17 (0.14)</b>	<b>0.40 (***)</b>	<b>0.39 (***)</b>	<b>1300</b>
$\delta^{13}\text{C} \sim \text{SpecieSex} + \text{Period}$	-14.44 (***)	0.37 (***)	0.69 (***)	0.96 (***)	-0.11 (***)	-	-	-	1311
$\delta^{15}\text{N} \sim \text{SpecieSex} * \text{Period}$	18.91 (***)	1.14 (***)	-1.02 (***)	0.86 (***)	-0.56 (0.0034)	-0.0008 (0.99)	-0.03 (0.92)	0.34 (0.21)	6291
<b><math>\delta^{15}\text{N} \sim \text{SpecieSex} + \text{Period}</math></b>	<b>18.89</b> (***)	<b>1.13</b> (***)	<b>-1.10</b> (***)	<b>0.94 (***)</b>	<b>-0.47</b> (***)	-	-	-	<b>6287</b>




### 4.3 *Isotopic niche size, overlap and community metrics according to period*


Standard ellipse areas (SEAc) increased in male groups of both species between Period 1 and Period 2 (11.03% in SAFS M and 16.27% in SASL M). In females, SEAc remained stable in SAFS and decreased for SASL by 72.75% (Figure 4.3, Table 4.3). In Period 1, SASL female SEAc overlaps with all groups (66.37 - 92.99%) followed by SASL male SEAc that overlap with SASL females (30.90%) and SAFS males (22.63%) and SEAc of SAFS males and SAFS females that overlap in range of 12.85 - 24.58% and 10.85 - 15.48%, respectively. In Period 2 only two cases of overlap of  $\leq 1\%$  were detected (Table 4.4). All ellipses segregate in the isotopic space in Period 2, evidenced by expansion of Layman metrics, including  $\delta^{15}\text{N}$  range,  $\delta^{13}\text{C}$  range, TA, CD, MNND and SDNND (Table 4.5).

**Table 4.3** Corrected Standard Ellipse Area (SEAc) for each period and group estimated with 75% maximum likelihood.

Group	SEAc Period 1	SEAc Period 2
SAFS F	0.94	0.95
SAFS M	1.29	1.45
SASL F	4.88	1.33
SASL M	2.27	2.64



**Figure 4. 3** Standard Ellipse Area (75%) for A) Period 1 and B) Period 2. Black lines delineate community convex hull between centroids of each group.




**Table 4. 4** Percent overlap between Peruvian otariid groups according to period. Values indicate the percent overlap for the group in the row.

		SAFS F	SAFS M	SASL F	SASL M
<b>Period 1</b>	<b>SAFS F</b>	-	14.94	80.67	0
	<b>SAFS M</b>	10.85	-	92.99	22.63
	<b>SASL F</b>	15.48	24.58	-	30.90
	<b>SASL M</b>	0	12.85	66.37	-
<b>Period 2</b>	<b>SAFS F</b>	-	1.03	0	0
	<b>SAFS M</b>	0.67	-	0	0
	<b>SASL F</b>	0	0	-	0
	<b>SASL M</b>	0	0	0	-

**Table 4. 5** Layman metrics estimated according to period.  $\delta^{13}C$  range = distance range of  $\delta^{13}C$  values,  $\delta^{15}N$  range = distance range of  $\delta^{15}N$  values, TA = Total area of convex hull, CD = the mean distance to centroid from means, MNND = mean nearest neighbor distance of the means, SDNND = standard deviation of the nearest neighbor distance.

Layman Metrics	Period 1	Period 2
$\delta^{13}C$ range	0.94	1.41
$\delta^{15}N$ range	1.18	2.36
TA	0.67	1.91
CD	0.60	1.05
MNND	0.71	1.16
SDNND	0.05	0.45




## 5. Discussion

In this study, long-term niche differentiation in individuals was investigated by examining the isotopic signatures serially archived in whiskers of male and female South American fur seals and South American sea lions during 2005-2019 in the HCS. We found a reorganization of our four otariid study groups in the isotopic space as a response to the increase in average sea level and warmer sea surface temperatures from November 2013 onwards (Period 2). Groups expanded their niche width (except SASL females), segregated in isotopic space, and reduced the  $\delta^{15}\text{N}$  signal during the warmer period. Overall decline in otariid  $\delta^{15}\text{N}$  suggests impoverished baseline values of  $\delta^{15}\text{N}$  with bottom-up effects impacting tertiary consumers, a shift in diet towards lower trophic prey, or both. Overlap reduced between all otariid groups in Period 2, providing support of a temporal shift from a more redundant assemblage (similar isotopic niches) during the colder period to a more trophically diverse assemblage (divergent isotopic niches) during the warmer period.

Although we did find significant differences to compare the four species-sex groups according to morphometrics, size was not a driver for higher  $\delta^{15}\text{N}$  as hypothesized. In our study, differences in the isotopic signatures in both species are driven by environmental conditions and behavioral traits according to sex. Different behavioral constraints between sexes in both otariid species, may allow larger male individuals to invest more time and energy foraging at sea, possibly being more selective hunters to capture higher level prey, as evidenced in reports of diverse hunting strategies in male and subadult male Galapagos sea lions (*Zalophus wollebaeki*) (De Roy et al., 2021; Páez-Rosas et al., 2020). Interestingly, sexual differences are generally maintained, but all  $\delta^{15}\text{N}$  signatures decrease in Period 2, with female SASL showing the largest drop in  $\delta^{15}\text{N}$  of 2.4, while male groups  $\delta^{15}\text{N}$  values drop in 1.5 and female SAFS decrease slightly in  $\delta^{15}\text{N}$  by 1.0. We consider that the change in  $\delta^{15}\text{N}$  in Period 2 reflects a bottom-up effect in the trophic food web, linked to the dietary habits of each species and sex group, in response to changes in the environment.


In November 2013 an elevation in the average sea level time series was detected, giving rise to a second period of higher sea level and warmer sea surface temperatures. This change is in agreement with the propagation of physical anomalies that started in the North Pacific Ocean that forced a biological response in the vertical distribution of nutrients and the abundance of chlorophyll-a (Zaba & Rudnick, 2016). Concurrently, in mid-2014 off the coast of Peru, a moderate, coastal El Niño was reported and then followed by an Extraordinary El Niño in 2015-2016 (L'Heureux et al., 2017) and






subsequently followed by a costal El Niño in 2017, continuing to promote warm water assemblage. Warm water Kelvin waves that travel north-south, during which cold water gets “trapped” under, depresses upwelling and decreasing productivity in the euphotic zone (Graco et al., 2007). Arrival of these warm water waves changed oceanographic conditions in April 2014, deepening the thermocline, increasing nearshore depth of chlorophyll-a, and decreasing the nutricline which negatively impacted nutrient enrichment in the system (Graco et al., 2016). In the California Current, another upwelling eastern boundary current marine ecosystem, deeper chlorophyll-a decreased phytoplankton biomass and caused a shift in community structure towards a smaller phytoplankton species, which caused bottom-up effects on food and energy availability at higher trophic levels, that resulted in high mortality rates of red tuna crabs (*Pleuroncodes planipes*) and California sea lions (*Zalophus californianus*) (Zaba & Rudnick, 2016). Based on the long-term changes of the  $\delta^{15}\text{N}$  signal captured in otariid whiskers, we suggest similar effects occurred in the HCS off the coast of Peru.

The red squat lobster (*Pleuroncodes monodon*) inhabits the northern HCS and has been abundant since the mid-1990s (Gutiérrez et al., 2008). It is an important prey item for seabirds, mammals, and coastal fish (Jahncke & Goya, 1997; Zavalaga et al., 1998). This crustacean has been identified as the dominant prey item in SASL diet composition (~70%) based on scat analysis. Whereas, SAFS have been reported to consume mostly Peruvian anchovy (49%) and squids (30%), with red squat lobsters contributing only a minor amount (6%) to the diet (Arias-Schreiber, 2003; Sarmiento-Devia et al., 2020; Sielfeld et al., 2018; Vásquez, 1995). Due to inherent limitations in the methodology, trophic studies based on scat samples rely on manually collected samples from colonies, resulting in sampling from individuals of unknown sex and age class. When samples are collected at breeding sites, researchers commonly assume that these belong to adult females given they alone represent 40-60% of the colony abundance and spend considerable time onshore while nursing their pups, while males are fasting on land (García-Rodríguez & Aurióles-Gamboa, 2004; Páez-Rosas & Aurióles-Gamboa, 2010; Porrás-Peters et al., 2008). Having mentioned this, the steeper drop in  $\delta^{15}\text{N}$  signal during Period 2 in SASL females compared to SAFS females agrees with a higher consumption of a lower trophic prey item (crustaceans) in the former. Furthermore, during El Niño events, diatoms, the dominant phytoplankton in the HCS, are greatly impacted by the less enriched conditions, with a reported decrease of 25-60% of their biomass during these events, while less nutritious species (e.g. dinoflagellates) can temporarily occupy their niche (Espinoza-Morriberón et al., 2017). *P. monodon*




has a similar trophic position as anchovies, with phytoplankton, eggs, and larvae (and not anchovies) being the main source of their diet (Espinoza et al., 2017). Meanwhile, zooplankton is the main prey item of anchovies (Espinoza & Bertrand, 2008) and enforces bottom-up control (Ayón et al., 2008). Squids feed on mesopelagic fish (Argüelles et al., 2012). Thus, since SASL females are feeding directly on red squat lobsters aggregated on the coastal habitat, their  $\delta^{15}\text{N}$  signal can be more directly reduced in comparison to SAFS females feeding on anchovies and squid in the pelagic realm. Thus, the less pronounced bottom-up effect in the latter can be partly explained by the trophic food web interactions. However, we cannot rule out partial prey switching or variations in the proportions of resources consumed, which can also impact the  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  signals.


The  $\delta^{13}\text{C}$  isotopic signature serves as a proxy for the inshore /offshore location of the foraging grounds. We found interspecific differences in  $\delta^{13}\text{C}$  were more accentuated during the warmer period, with lower  $\delta^{13}\text{C}$  values in SAFS compared to SASL, independent of period. We did not find differences in  $\delta^{13}\text{C}$  explained by sex but followed the inshore-offshore gradient reported by Espinoza et al. (2017), with decreasing values in the gradient of coastal >neritic >oceanic zones. Therefore, trophic resource acquisition by SAFS is linked to more offshore location items in comparison to SASL. Furthermore, in the warming period, SAFS exhibit decreases in the  $\delta^{13}\text{C}$  signal towards offshore waters, while SASL remains in the same range. This agrees with the association of the biomass of the main prey item of SASL the red squat lobster with cold coastal waters that contract further inshore during warming events (Yuffra et al., 2018). Studies using hydro acoustic surveys along the coast of Peru found that distribution of Peruvian anchovy and red squat lobsters spatially overlap, with both prey species distributed close to the coast, but red squat lobsters concentrate closer towards shore, in the first 40 km (Gutiérrez et al., 2008). Whereas, Peruvian anchovies, the main prey item for SAFS, show larger range in  $\delta^{13}\text{C}$  values, indicative of a wider use of the pelagic habitat in comparison to red squat lobsters (Espinoza et al., 2017). Between sexes, males have higher  $\delta^{13}\text{C}$  than females, which may reflect inclusion of coastal/benthic foraging in males. The latter can be related to the requirement in females for homogenous prey patches that are available in the offshore environment in HCS to more efficiently optimize milk production to nurse pups, whereas males have more possibilities of exploring different foraging grounds and strategies (Krüger et al., 2014). The reduction in overlap of the isotopic niches between the two periods suggests that the response to warmer anomalies promoted trophic segregation between otariid predator communities, contrary to




what we hypothesized. When using Layman metrics, we found that the ranges of both isotopic signatures ( $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  ranges) and the total area occupied (TA) expanded in response to warmer conditions in Period 2 (Figure 4.3, Appendix D1). Furthermore, the lower levels of the trophic position in the otariid community during Period 2 was predominantly driven by a niche shift in all groups, with a significant contribution from SASL females. This supports the theory that a new basal resource with a more depleted  $\delta^{15}\text{N}$  took place in Period 2, as reported in other studies showing temporal niche shifts in marine predators (Amador-Capitanachi et al., 2020; Yurkowski et al., 2018). In general, populations can be separated into specialists when they have narrower niches or into generalists when they have broader niches (Begon et al., 2006). Generalist populations with broader niches are likely favored in heterogeneous and stable environments, whereas specialist populations with narrower niches are favored in highly dynamic ecosystems and homogeneous environments (Kassen, 2002). Thus, during the warmer period, it is possible that prey assemblages are more diverse in contrast with dense prey aggregations favored by nutrient rich, cold-water in HCS. Since TA serves as a proxy for the extent of trophic diversity within the community, its expansion (along with CD, NND and SDNND) reflects a more generalist community with looser packing of individuals in the isotopic niche space during Period 2 (Layman et al. 2007). Of all the groups, female SASL show the largest variation in response to the warming scenario, reducing both ellipse area and  $\delta^{15}\text{N}$  values. This suggests that this group has higher plasticity to adapt to changing environmental conditions, as reported in other otariids (Amador-Capitanachi et al., 2020). This helps explain current, larger population biomass of SASL reported for PSJ and Peru, in contrast to the smaller and steeper declining SAFS population (Cárdenas-Alayza et al., 2021). Plasticity and feeding on a more available food source that lacks commercial fishing pressure can be currently more advantageous for SASL. In the Atlantic Ocean SAFS forage in pelagic environments with abundant, but similar, prey. They are identified as a specialist population composed of generalist individuals. Whereas, benthic SASL forage in habitats with diverse, but less abundant prey, with more generalist populations composed of highly specialized individuals (Franco-Trecu et al., 2014; Riverón et al., 2021). However, this marine ecosystem is relatively stable compared to the dynamic upwelling HCS. In the Galapagos Islands, an upwelling system with more similarities to the HCS environment, segregation with low overlap exists between sympatric fur seals and sea lions. Galapagos fur seals (*Arctocephalus galapagoensis*) forage in the pelagic zone with a smaller population isotopic niche compared to




benthic foraging Galapagos sea lions (*Zalophus wollebaeki*) with larger isotopic niche (Páez-Rosas & Aurióles-Gamboa, 2014). However, in our study site in the Peruvian HCS, the bathymetry deepens very rapidly, reaching over 1000 m in less than 50 km offshore, leaving a much-reduced area for benthic foraging; this reason is why the contribution of specialized individuals employing benthic strategies is still put in question. Thus, individual specialization and the capacity to adapt different foraging strategies can be playing an important role in the variability between our study groups. We recommend future research explore individual specialization to understand exploitation mechanisms by individual otariids in the HCS and how individuals respond to changing environmental conditions.




# APPENDIX D1



**Figure D1. 1** Standard Ellipse Area estimated for each group. Labels represent Community (Period 1 and 2) and Group (1 = SAFS females, 2 = SAFS males, 3 = SASL females, 4 = SASL males). The black points correspond to the mean standard ellipse area for each group, red cross is the standard ellipse area corrected for small sample size. Grey and white boxed areas reflect the 95, 75 and 50% confidence intervals




## APPENDIX D2



**Figure D2. 1** Total area of convex hull estimated for each community or period. Community 1 = Period 1, Community 2 = Period 2. The black points correspond to the mean standard ellipse area for each group while the grey and white boxed areas reflect the 95, 75 and 50% confidence intervals

## DISCUSSION GÉNÉRALE

---



“FOR WHATEVER WE LOSE  
(LIKE A YOU OR A ME).  
IT’S ALWAYS OUR SELF  
WE FIND IN THE SEA.”

- E.E. Cummings






The main goal of this thesis is to describe and compare the ecological niches of sympatric otariid species South American fur seal (SAFS, *Arctocephalus australis*) and South American sea lion (SASL, *Otaria byronia*) that coexist in Punta San Juan; and determine intraspecific and interspecific niche differentiation/overlap. In the data chapters of this thesis I explore different dimensions of trophic ecology and surrounding environmental conditions to explain the current divergent population trends of sympatric otariids in Punta San Juan, reported in Chapter 1. In Chapter 1 analyzed trends of abundance for a 20 year period and found that on average, 83% of all otariid biomass is composed by SASL and 17% by SAFS; making SASL the current ‘dominating’ species in the study site (SASL is currently 89% of total otariid biomass). Furthermore, results from the population assessment reveal a current decline in both populations, suggesting a resource limitation in the marine environment, which started in November of 2014 in SAFS and in March 2017 in SASL.

## **1. Interspecific and intraspecific segregation of the trophic niche (diet composition and isotopic signals)**

To describe and compare the trophic ecology of sympatric pinnipeds, I assess the degree of interspecific and intraspecific trophic niche segregation, through the analysis of diet composition and isotopic signals between the sympatric pinnipeds SAFS and SASL in relation to environmental conditions. My main hypotheses was that larger body size relates to foraging strategies that facilitates consumption of larger prey items, linked to the benthos. Being SASL overall 1.5 times larger than SAFS in all age-sex classes, I hypothesized that SASL would be foraging in more coastally restrained areas to access the benthos to find larger prey, and SAFS would forage in pelagic locations (Chapter 3). I also expected that consumption of higher trophic level prey would be related to body size. This was assessed through the trophic levels of prey items consumed (Chapter 2) and through the trophic position of each species-sex group determined through the  $\delta^{15}\text{N}$  isotopic signature in individual whisker samples (Chapter 4).

Due to the dimorphism in otariids, I expected to find sexual segregation, when sample collection from individuals permitted this evaluation (Chapter 3 and Chapter 4). In this case, the hypotheses was that



lactating female foraging grounds should be more proximate to the colony in comparison to males, reflecting a reduction in the distance, time and area invested in the search for prey, to ensure pup survival on land (Chapter 3). In the isotopic signals, this would reflect on  $\delta^{13}\text{C}$  values of females closer to zero (more coastal) with a narrower range, indicative of more restricted area search for prey in comparison to males. I expected males could have more negative  $\delta^{13}\text{C}$  values with a broader range of values, reflecting wider search for food in the pelagic domain (Chapter 4).

For this objective, results have shown that trophic interspecific segregation exists at different levels. In Chapter 2 I determined that dietary interspecific segregation exists due to resource partitioning of three main shared prey items: Peruvian anchovy, cephalopods, and red squat lobster. The consumption of these prey items in different proportions by each otariid segregates into two clusters containing the combinations the prey resources consumed during consecutive seasonal samplings in the 2015-2018 study period. In general, SAFS consistently consumed Peruvian anchovy (42.8%) and cephalopods (34.7%); with smaller components of myctophids (7.4%), red squat lobsters (5%), and eleven other fish species. SASL diet is mainly composed of red squat lobster (69.9%) and Peruvian anchovy (19.1%); with smaller amounts of cephalopods (4.7%), silversides (4.6%), and five other fish species. The classification of prey items into ecological groups, based on published literature on habits of prey items, showed that SAFS prey items are from Pelagic (52.35%) and Demersal-Pelagic (47.25%) ecological groups. Meanwhile SASL's prey items are from Demersal-Pelagic (69.44%) and of Pelagic (23.88%) groups. Meanwhile, Demersal-Benthic species are not important in neither of the Peruvian otariids' diet composition (1-1.5%). This suggests that majority of prey items come from the pelagic zone, but more refined methods are required to understand the vertical or temporal use of foraging grounds of otariids (i.e., time depth recorders).

Average trophic level values of the prey items alone did not reveal any significant differences between trophic positions for SAFS ( $4.31 \pm 0.15$ ) and SASL ( $4.55 \pm 0.11$ ). This can be due to the variability in the sources of information from the literature and the wide range of study locations. However, isotopic signatures for  $\delta^{15}\text{N}$  for recent years (December 2014 - December 2019), did reveal a higher trophic position in males compared to females for both species. In females, a higher  $\delta^{15}\text{N}$  signal was found in SAFS compared to SASL, albeit the interspecific difference in body size (SAFS females:17.9; SAFS males:18.7 and SASL females: 16.6; SASL males: 18.5, Chapter 4). In Chapter 2, an important significant difference was found between niche breadth based on identification of



hard parts, with SAFS ( $0.26 \pm 0.13$ ) reaching double Levin's  $B_i$  than SASL ( $0.14 \pm 0.16$ ). In general, SAFS diet also tended to show higher species richness compared to SASL (Chapter 2). When comparing the species consumed, although SAFS diet composition concentrated on anchovy and cephalopod consumption, during many seasons it added new species to the diet, explaining the broadening of niche breadth; in contrast to a less variable species richness in SASL diet, based consistently on large quantities of red squat lobsters and anchovy.

The increase in unique species to SAFS were of pelagic domain (e.g., King gar (*Scomberesox saurus scombroides*), Pelican barracuda (*Sphyræna idiaestes*), California needlefish (*Strongylura exilis*), Sharpchin flyingfish (*Fodiator rostratus*), Appendix B2) in diet confirms that SAFS forage commonly further offshore. Whereas in SASL's unique species pertain to the coastal domain (e.g., Chameleon wrasse (*Haliogheres dispilus*), Banded toadfish (*Aphos prosus*), Appendix B2). In the isotopic realm, dietary segregation of a more offshore diet on anchovy and cephalopods in SAFS and a more inshore diet based on red squat lobsters and anchovy is backed up by the interspecific  $\delta^{13}\text{C}$  signals found (Chapter 4). The  $\delta^{13}\text{C}$  signal SAFS has a more negative (e.g. offshore) signal in comparison to SASL (e.g., inshore) in both sexes (SAFS females:  $-14.87 \pm 0.47$ , males:  $-14.27 \pm 0.36$ ; SASL females:  $-13.72 \pm 0.73$ , males:  $-13.36 \pm 0.58$ ).

## 2. Foraging strategies (spatial and temporal dimensions)

To respond to my objective focused on determination of distinct foraging strategies (spatial and temporal dimensions) employed by sympatric pinnipeds SAFS and SASL in relation to environmental conditions. To address this objective, my hypothesis was also based on body size. Since SASL are considerably larger in body size than SAFS, and both species have a strong sexual dimorphism, with males being much larger than females (Ralls & Mesnick, 2002). In Chapter 3, I expected that SASL will travel longer distances, forage for longer durations, and cover larger areas than SAFS. Also, in terms of temporal segregation, based on Trillmich et al. (1991), I predicted that fur seals will forage mostly in the night hours to capture superficial available prey during diel vertical migration of the most abundant prey item in HCS, Peruvian anchovy (*Engraulis ringens*). In contrast, I expected SASL to forage indifferently throughout the day or night hours as shown in other sea lion species (Riet-Sapriza et al., 2013; Schwarz et al., 2021; Villegas-Amtmann et al., 2008).

Due to the energetic constraints linked to parental roles and offspring survival, in Chapter 3 I expected



females of both species to be more coastal (inshore) and males more pelagic (offshore) as evidenced in other otariids (Page et al., 2006; Staniland & Robinson, 2008). Since in HCS, cool coastal upwelling translates to lower sea surface temperatures (SST), higher productivity (assessed by higher concentrations of chlorophyll-a), lower proximity to fronts, and lower front gradients in reduced distance from coast (Echevin et al., 2008; Gutiérrez et al., 2011) I expect this pattern to be related to female foraging habitat. Whereas the opposite features, that characterize pelagic habitats, I envisioned would be associated to male foraging. Furthermore, in Chapter 4, in terms of isotopic space I hypothesized that both otariid females will show  $\delta^{13}\text{C}$  values closer to zero indicative of higher productivity since they remain constrained closer to the land-based colony, where chlorophyll-a is higher. In contrast, males would have a wider range of  $\delta^{13}\text{C}$  from foraging more “freely” between coastal and pelagic zones.

Two scales are explored and discussed for segregation in the temporal dimension in Chapter 3. Peruvian otariids have a 1-3 month temporal segregation in their breeding seasons (SAFS breeds from October to December and SASL from January to March). This contributes yet another mechanism for segregation known as allochrony to partition resources during energetically critical times, which has been postulated as one of the founding drivers behind speciation (Taylor & Friesen, 2017). Since lactation is extended in temperate latitude species and time of weaning is flexible (reported to occur between 6-36 months), we consider it valuable to explore segregation mechanisms while lactating females are foraging to nurse their pups. In a smaller foraging trip scale, I found nocturnal foraging in SAFS females (incremental peak in dusk hours) and daytime foraging in SASL males (incremental peak in dawn hours). Nocturnal foraging in SAFS females may be a synergistic effect between the energetic constraints to provision for their young and the vertical diel migration of prey items for SAFS such as Peruvian anchovies, squids and myctophids, that are more available in the dusk to night hours (Ayón et al., 2008). Since we did not find a significant effect for hour of day in male SAFS, we reject our hypothesis that this pattern holds for all SAFS. Since males do not have energetic constraints linked to parental care, daytime foraging segregation in SASL males may be a reflection of a reduction in competition with another otariid age-class, predator species or preference for a specific prey (Chilvers & Wilkinson, 2009). In Chapter 2 we report a high consumption of red squat lobsters in SASL, therefore we consider this could be a preferred targeted prey for this temporal segregation. Studies on diel migration of Peruvian anchovies and red squat lobsters show a high



degree of overlap. However at hours of dawn, squat lobsters congregate in more dense aggregations closer to the surface in comparison to anchovies (Gutierrez et al., 2005), suggesting it can be the best time to increase the capturability of these prey patches. Studies in other sea lion species have found individuals in the same age-class in have different foraging modes (pelagic, benthic and nocturnal divers, Schwarz et al. 2021). Thus, it is possible that also in SASL there are individual specialization mechanisms; or that individuals can also switch between strategies according to surrounding prey availability. Thus, we recommend these scenarios are further explored in this ecosystem. Future studies involving depth recorders coupled with availability of local biomass of prey items would help refine validation of foraging along the water column, since present analyses may have associated errors in the identification of foraging events based on locations alone.

The results of Chapter 3 show that sympatric otariids in the HCS have multiple strategies to segregate while foraging in the pelagic environment and that body mass is not the sole driver for segregation in space and time, but rather a combination of energetic constraints according to sex, reproductive roles linked to specific dietary resources. We found that although they are smaller in size, both SAFS females and males, make trips of longer distance and duration in comparison to both sexes of SASL, rejecting our hypothesis that larger animals would make longer trips. In terms of area, SASL cover larger home ranges and core areas, but do so with more consecutive shorter trips compared to SAFS. This reveals different use of the foraging grounds at interspecific level. SASL have larger home ranges as a group, covering a larger area parallel to the coastline that envelopes the SAFS home range. In contrast, SAFS travel farther offshore during feeding trips to specific areas. In agreement with our predictions based on previous studies focused on sexual segregation of otariids, females of both species show more constrained foraging trips in terms of distance traveled, home ranges and core areas in comparison to males in both species (Page et al., 2006; Staniland & Robinson, 2008). Finally, although post-breeding dispersal is common in male pinnipeds this study shows that the South American otariids do not depart on a migration, but rather obtain their energetic demands in a local range of 55,000 - 106,000 km<sup>2</sup>. These data are evidence that these sympatric species have evolved and adapted to obtain their resources locally in the nearby productive upwelling waters of HCS, occupying rookeries like PSJ year round.



### 3. Environmental conditions associated to foraging

The environmental variables associated with foraging events evaluated in sympatric pinnipeds in Chapter 3 revealed interspecific and intraspecific differences. When compared, environmental habitat features of SASL females are linked to the coastal habitat that is characterized by lower distance to coast (DC), lower sea surface temperature (SST), lower Proximity to Fronts and higher chlorophyll-a (Chl-a) values. Whereas SAFS females are associated to a more pelagic domain characterized by higher SST, more proximity to fronts, higher front gradients and of course, higher DC values. Among males, best fitting models for SAFS males included SST, Chl-a and Proximity to Fronts, whereas SASL males included all environmental covariates. SASL males have lower SST, lower Chl-a, lower Proximity to Fronts and lower DC, which could be a product of individual strategies. Ranges in the environmental variables targeted by SAFS and SASL males are highly overlapping; thus conclusions should be taken with caution and further studies exploring individual strategies should be conducted. We also suggest incorporating additional environmental variables to better explain male foraging environment, such as those related to the vertical structure of the water column.

In Chapter 4, I set up a framework to explore  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  isotopic niche differentiation along longitudinal samples in whiskers, permitting to compare a fifteen year period (2005-2019), during which a change in environmental conditions were found. In this study I found a reorganization of our four otariid study groups in the isotopic space as a response to the increase in average sea level and warmer sea surface temperatures from November 2013 onwards (Period 2). Groups expanded their niche width (except SASL females), segregated in isotopic space, and reduced the  $\delta^{15}\text{N}$  signal during the warmer period. A decline in otariid  $\delta^{15}\text{N}$  suggests impoverished baseline values of  $\delta^{15}\text{N}$  with bottom-up effects impacted tertiary consumers, a shift in diet towards lower trophic prey, or both. Niche overlap was reduced between all otariid groups in Period 2, providing support of a temporal shift from a more redundant assemblage (similar isotopic niches) during the colder period (Period 1) to a more trophically diverse assemblage (divergent isotopic niches) during the warmer period (Period 2). Results from Chapter 4 are in agreement with other findings in stable isotope signals in SAFS whiskers that suggest having captured ENSO signals (Edwards et al., 2021).

The analyses of this long term dataset depicts a shift in the trophic organization that is aligned temporally with the decline of the Punta San Juan otariid populations detected in Chapter 1.





Furthermore, the break into two periods of varying environmental conditions explains the changes in the trophic reorganization in the otariid community. During Period 2, female SASL show the largest drop in  $\delta^{15}\text{N}$  of 2.4, while male groups  $\delta^{15}\text{N}$  values drop in 1.5 and female SAFS decrease slightly in  $\delta^{15}\text{N}$  by 1.0. We consider that the change in  $\delta^{15}\text{N}$  in Period 2 reflects a bottom-up effect in the trophic food web, linked to the dietary habits of each species and sex group, in response to changes in the environment.

In November 2013 an elevation in the average sea level time series for the study area was detected, giving rise to a second period of higher sea level and warmer sea surface temperatures. This change is in agreement with the propagation of physical anomalies that started in the North Pacific Ocean that forced a biological response in the vertical distribution of nutrients and the abundance of chlorophyll-a (Zaba & Rudnick, 2016). Concurrently, in mid-2014 off the coast of Peru, a moderate, coastal El Niño was reported and then followed by an Extraordinary El Niño in 2015-2016 (L'Heureux et al., 2017) and subsequently followed by a coastal El Niño in 2017, continuing to promote a warm assemblage. During these years, studies suggest that physical anomalies of these warming ocean events generated a biological response similar to those of El Niño (Espinoza-Morriberón et al., 2017). All the beforementioned events involve the rise of sea level forced by the passage of remotely generated and coastal trapped waves in response to heat expansion (Ryan & Noble, 2002). The thermocline, nearshore depth of chlorophyll-a, and the nutricline deepened, decreasing nutrient enrichment in the euphotic zone. Weak winds caused the reduction in vertical nutrient fluxes to the depleted euphotic zone. A deeper concentration of chlorophyll-a decreased phytoplankton biomass, causing a shift in community structure toward smaller phytoplankton which can have bottom-up effects on food and energy availability at higher trophic levels (Zaba & Rudnick, 2016).

Given these results and those of Edwards et al. (2021), we propose that there is a pattern in the response of otariid communities towards different climatic scenarios in HCS. In cold scenarios with low SST and low sea level there will be higher prey available of higher quality. This will promote population growth in otariids, with higher recruitment of females and juveniles, and the opposite during warm regimes. Due to more prey available, dietary segregation, spatial overlap, temporal segregation and segregation in isotopic space will be reduced, in comparison to an opposite warm scenario (Figure D.1).






		Cool	Warm
Environment	SST	↓	↑
	Sea level	↓	↑
Environment	Prey availability	↑	↓
	Prey quality	↑	↓
Chp 1	Female recruitment	↑	↓
	Juvenile recruitment	↑	↓
Chp 2	Dietary segregation	↓	↑
Chp 3	Spatial overlap	↓	↑
	Temporal segregation	↓	↑
Chp 4	Isotopic segregation	↓	↑

**Figure D. 1** Diagram summarizing main findings to explain current population decline in Peruvian otariids in context of two contrasting environmental scenarios: cool (blue) and warm (red) conditions.

## 4. Final remarks

Populations can be separated into specialists when they have narrower niches or into generalists when they have broader niches (Begon et al., 2006). Generalist populations with broader niches are likely favored in heterogeneous and stable environments, whereas specialist populations with narrower niches are favored in highly dynamic ecosystems and homogeneous environments (Kassen, 2002).



Thus, during the warmer period, results suggest that prey assemblages are more diverse in contrast with dense prey aggregations favored by nutrient rich, cold-water in HCS. Since total ellipse area serves as a proxy for the extent of trophic diversity within the community, its expansion during Period 2 reflects a more generalist community with looser packing of individuals in the isotopic niche space (Layman et al., 2007). From all the species and sex groups, female SASL show the largest variation in response to the warming scenario, reducing both isotopic niche width and  $\delta^{15}\text{N}$  values. This suggests that this group has higher plasticity to adapt to changing environmental conditions, as reported in other otariids (Amador-Capitanachi et al., 2020). Knowing that SASL is the dominating species in Peru, higher plasticity in adult females helps explain current, larger population biomass of SASL reported for PSJ and Peru, in contrast to the smaller and steeper declining SAFS population (Cárdenas-Alayza et al., 2021). Thus, trophic plasticity, enhanced by feeding primarily on a more available food source (*P. monodon*) that lacks commercial fishing pressure can be currently more advantageous for SASL population growth.

Results show that sympatric otariids in the HCS currently show some level of segregation through trophic indices and while foraging at sea, partly associated with reproductive role, body size and species. Based on our long term stable isotopes time series, trophic segregation mechanisms have changed after November 2014. Segregation mechanisms only make sense if prey are partly limited and spatially partitioned. In the case of very high availability of prey, coexistence could occur without much need of segregation. This could be the case in the previous colder Period 1, January 2005 – October 2014. However, in the case of food shortages, segregation mechanisms could sustain coexistence, by taking advantage of differential abilities to cope with such a situation, which could be the case of Period 2. In the latter case, the differences observed in the foraging ecologies may depict a situation of one species outcompeting another, which couldn't be considered as a coexistence mechanism. In this context, the plasticity in SASL females in contrast to the less flexible SAFS female group, can help explain SASL as the dominating of the two sympatric species in HCS, albeit environmental changes. This continues to raise the question of the dynamics of segregation mechanisms with respect to food availability in dynamic marine environments like HCS, and calls for maintenance of the long term monitoring and research programs.

In general, empirical understanding of competitive networks is limited by the absence of detailed information on the extent of resource partitioning and competition at a community scale, and the



effects of interactions among species that do not come into direct contact (Eurich et al., 2018). Thus, other potential competitors in HCS can also be contributing to reduced SAFS population. For example, many other predatory fish species have been reported to be feeding red squat lobsters, after the increase in its availability in the cold water regime. Species like dolphinfish (*Coryphaena hippurus*), Jack mackerel (*Trachurus murphyi*), bonito (*Sarda chilensis*) and several shark species could be a source of competition for items targeted by SASL (Alegre, 2015; Gonzalez-Pestana et al., 2021). This will increase in cases of shortage of this currently abundant prey item that is sustaining many predators.

Research on Jumbo squid (*Dosidicus gigas*) show how this voracious short-lived cephalopod with diel migration has increased its population biomass in the cold regime during the 2000s off the coast of Peru incrementing the size at maturity (Argüelles et al., 2008). Investigations on trophic ecology of this species report that the main contribution to diet composition is based on other cephalopods along with mesopelagic fishes, euphysiids and some cannibalistic feeding on smaller age-classes of *D. gigas*. Larger sized Jumbo squid consume larger prey items such as cephalopods, while smaller *D. gigas* consume lower trophic level prey (Alegre et al., 2014). Thus, the successful population growth of Jumbo squid in the past 20 years can be a new source of competition for species like SAFS, based on the similarities in their diet composition and offshore nocturnal habits. Potential competition between SAFS and *D. gigas* should be further explored to determine if recent changes in predator guilds can help explain reduced population abundance of SAFS.

It has been well documented that the Peruvian HCS contains one of the world's largest monospecific fisheries based on Peruvian anchovy, along with other fisheries that target cephalopods and other small pelagic fish species. Many of these fisheries have been overcapitalized and are decreasing available biomass to dangerous levels for sustainability, both at industrial and artisanal level (De la Puente et al., 2020; Fréon et al., 2008). To present date, it is well documented that Peruvian anchovy (*E. ringens*) is heavily exploited and is a source of competition for several predators in HCS. However, we are unaware of any large-scale exploitation of red squat lobsters (*P. monodon*) in Peru that limits the availability of this prey item for predators. However, if this were to occur, it may severely affect the availability of this prey resource for SASL and could lead to further changes in the diet of this species, and increase competition between the two sympatric otariid species as evidenced in other regions (Drago et al., 2009; Szteren et al., 2018). Thus, we recommend the



incorporation of trophic ecology studies of top predators to further inform the generation of quotas to improve the level of biomass left available in the oceans for the predator populations to recover from decline.

## 5. Future Recommendations

As part of this study, we would like to offer a series of recommendations for future research to continue to disentangle sources of segregation to explain the coexistence of Peruvian otariids, as follows:

- First, there is a need to continue to update resource partitioning information by identifying cephalopods to species level to determine how cephalopod species diversity impacts niche breadth. Furthermore, estimation of the size of fish and cephalopods consumed by otariids is recommended to determine if there is interspecific resource partitioning in prey sizes targeted by SAFS or SASL.
- In the realm of foraging strategies, I recommend to determine if female foraging behavior changes according to breeding or non-breeding season as well as offspring number and age (Drago et al., 2010, 2015, 2021; Franco-Trecu et al., 2012). This could help finetuning the critical time of energetic constraint in females according to species.
- Findings show varying lengths in distance duration of foraging trips according to species-sex groups. This can be indicative of the use of specialized foraging grounds in some groups, suggesting potential site fidelity to foraging grounds. This has been reported in other locations and should be further explored in HCS (Baylis et al., 2017; Knox et al., 2018).
- Ranges in the environmental variables targeted by SAFS and SASL males are highly overlapping and ranges are hard to interpret as a specific habitat selection. Further analyses that explore individual and/or group strategies is recommended.
- Given the dynamic environment in HCS and the differences observed in the foraging ecologies of



one species outcompeting another, the question of the dynamics of segregation mechanisms with respect to food availability persists and calls for maintenance of the long term monitoring and research programs to compare these indices over time. Monitoring programs on top predators should continue to collect information on diet composition, foraging behavior and populations dynamics.

- Given the variation in some groups (e.g., males) contribution of specialized individuals employing benthic strategies close to shore is still put in question. Thus, individual specialization and the capacity to adapt different foraging strategies can be playing an important role in the variability between study groups. I recommend future research to explore individual specialization to understand exploitation mechanisms into how individuals respond to changing environmental conditions. Deployment of time-depth recorders and collection of tissue samples that archive long term signals such as stable isotopes and fatty acid signatures should be prioritized.
  
- Ecological theory has progressed from a focus of pairwise interactions to a multifactorial perspective of mechanisms that drive competition at community level. To get a bigger picture of the interactions of SAFS and SASL in HCS, further studies that compare trophic ecologies with other potential predators in HCS (e.g., seabirds, squids, large predatory fish, elasmobranchs and cetaceans) are recommended to further establish roles and interactions between potential competing predators in HCS.

## Références Bibliographiques

- Adams, J. (2004). *Foraging Ecology and reproductive biology of Cassin's Auklet (Ptychoramphus a. aleuticus) in the California Channel Islands*. San Francisco State University, Moss Landing, California.
- Adkesson, M. J., Chinnadurai, S. K., Balko, J. A., Jankowski, G., Langan, J., Meegan, J. M., Cárdenas-Alayza, S., & Allender, M. C. (2019). Field anesthesia of South American sea lions (*Otaria byronia*) in Peru using Medetomidine, Midazolam and Butorphanol. *51st AAZV Annual Conference Proceedings -American Association of Zoo Veterinarians*, 78.
- Adkesson, M. J., Chinnadurai, S. K., Balko, J. A., Jankowski, G., Meegan, J. M., Cárdenas-Alayza, S., & Allender, M. C. (2019). Anesthesia of free-ranging Peruvian fur seals (*Arctocephalus australis*) using Medetomidine, Midazolam and Butorphanol. *Proc Zoo Wildlife Health Conference*, 57–58.
- Alegre, A. (2015). *Trophic ecology of jumbo squid and predatory fishes in the Northern Humboldt Current System*. Université de Montpellier.
- Alegre, A., Ménard, F., Tafur, R., Espinoza, P., Argüelles, J., Maehara, V., Flores, O., Simier, M., & Bertrand, A. (2014). Comprehensive model of jumbo squid *Dosidicus gigas* trophic ecology in the northern Humboldt current system. *PLoS ONE*, 9(1). <https://doi.org/10.1371/journal.pone.0085919>
- Amador-Capitanachi, M. J., Moreno-Sánchez, X. G., Ventura-Domínguez, P. D., Juárez-Ruiz, A., González-Rodríguez, E., Gálvez, C., Norris, T., & Elorriaga-Verplancken, F. R. (2020). Ecological implications of unprecedented warm water anomalies on interannual prey preferences and foraging areas of Guadalupe fur seals. *Marine Mammal Science*, 36(4), 1254–1270. <https://doi.org/10.1111/mms.12718>
- Antonelis, G. A., Stewart, B. S., & Perryman, W. (1990). Foraging characteristics of female northern fur seals (*Callorhinus ursinus*) and California sea lions (*Zalophus californianus*). *Canadian Journal of Zoology*, 68, 150–158.
- Argüelles, J., Lorrain, A., Cherel, Y., Graco, M., Tafur, R., Alegre, A., Espinoza, P., Taipe, A., Ayón, P., & Bertrand, A. (2012). Tracking habitat and resource use for the jumbo squid *Dosidicus gigas*: A stable isotope analysis in the Northern Humboldt Current System. *Marine Biology*, 159(9), 2105–2116. <https://doi.org/10.1007/s00227-012-1998-2>
- Argüelles, J., Tafur, R., Taipe, A., Villegas, P., Keyl, F., Dominguez, N., & Salazar, M. (2008). Size increment of jumbo flying squid *Dosidicus gigas* mature females in Peruvian waters, 1989-2004. *Progress in Oceanography*, 79(2–4), 308–312. <https://doi.org/10.1016/j.pocean.2008.10.003>

- Arias-Schreiber, M. (2003). *Prey spectrum and feeding behaviour of two sympatric pinnipeds (Arctocephalus australis and Otaria flavescens) in relation to the 1997/98 ENSO in southern Peru*. University of Bremen. Bremen, Germany.
- Arias-Schreiber, M., & Rivas, C. (1998). Distribución, tamaño y estructura de las poblaciones de lobos marinos *Arctocephalus australis* y *Otaria byronia* en el litoral peruano, en noviembre 1996 y marzo 1997. *Informe Progresivo Del Instituto Del Mar Del Perú*, 73, 17–32.
- Arim, M., Abades, S. R., Laufer, G., Loureiro, M., & Marquet, P. A. (2010). Food web structure and body size: Trophic position and resource acquisition. *Oikos*, 119(1), 147–153. <https://doi.org/10.1111/j.1600-0706.2009.17768.x>
- Arnould, J. P. Y., & Costa, D. P. (2006). Sea Lions in Drag, Fur Seals Incognito: Insights from the Otariid Deviants. In A. W. Trites, S. K. Atkinson, D. P. DeMaster, L. W. Fritz, T. S. Gelatt, L. D. Rea, & K. M. Wynne (Eds.), *Sea Lions of the World* (1st ed., pp. 309–324). Alaska Sea Grant College Program. <https://doi.org/https://doi.org/10.4027/slw.2006.21>
- Arnould, J. P. Y., & Hindell, M. A. (2001). Dive behaviour, foraging locations, and maternal-attendance of Australian fur seals (*Arctocephalus pusillus doriferus*). *Canadian Journal of Zoology*, 79(1), 35–48.
- Auriolos-Gamboa, D., & Camacho-Ríos, F. J. (2007). Diet and Feeding Overlap of Two Otariids, *Zalophus californianus* and *Arctocephalus townsendii*: Implications to Survive Environmental Uncertainty. *Aquatic Mammals*, 33(3), 315–326. <https://doi.org/10.1578/AM.33.3.2007.315>
- Ayón, P., Swartzman, G., Bertrand, A., Gutiérrez, M., & Bertrand, S. (2008). Zooplankton and forage fish species off Peru: Large-scale bottom-up forcing and local-scale depletion. *Progress in Oceanography*, 79(2–4), 208–214. <https://doi.org/10.1016/j.pocean.2008.10.023>
- Bailleul, F., Luque, S., Dubroca, L., Arnould, J. P. Y., & Guinet, C. (2005). Differences in foraging strategy and maternal behaviour between two sympatric fur seal species at the Crozet Islands. *Marine Ecology Progress Series*, 293(June), 273–282. <https://doi.org/10.3354/meps293273>
- Bakun, A., & Weeks, S. J. (2008). The marine ecosystem off Peru: What are the secrets of its fishery productivity and what might its future hold? *Progress in Oceanography*, 79(2–4), 290–299. <https://doi.org/10.1016/j.pocean.2008.10.027>
- Barber, R. T., & Chavez, F. P. (1983). Biological consequences of El Niño. *Science*, 222(4629), 1203–1210. <https://doi.org/10.1126/science.222.4629.1203>
- Barbraud, C., Bertrand, A., Bouchón, M., Chaigneau, A., Delord, K., Demarcq, H., Gimenez, O., Torero, M. G., Gutiérrez, D., Oliveros-Ramos, R., Passuni, G., Tremblay, Y., & Bertrand, S. (2018). Density dependence, prey accessibility and prey depletion by fisheries drive Peruvian



- seabird population dynamics. *Ecography*, 41(7), 1092–1102. <https://doi.org/10.1111/ecog.02485>
- Barnes, C., Maxwell, D., Reuman, D. C., & Jennings, S. (2010). Global patterns in predator-prey size relationships reveal size dependency of trophic transfer efficiency. *Ecology*, 91(1), 222–232. <https://doi.org/10.1890/08-2061.1>
- Barraquand, F., & Benhamou, S. (2008). Animal movements in heterogeneous landscapes: Identifying profitable places and homogeneous movement bouts. *Ecology*, 89(12), 3336–3348. <https://doi.org/10.1890/08-0162.1>
- Bartholomew, G. A. (1970). A model for the evolution of pinniped polygyny. *Evolution*, 24(3), 546–559.
- Baylis, A. M. M., Arnould, J. P. Y., & Staniland, I. J. (2014). Diet of South American fur seals at the Falkland Islands. *Marine Mammal Science*, 30(3), 1210–1219. <https://doi.org/10.1111/mms.12090>
- Baylis, A. M. M., Orben, R. A., Costa, D. P., Arnould, J. P. ., & Staniland, I. J. (2016). Sexual segregation in habitat use is smaller than expected in a highly dimorphic marine predator, the southern sea lion. *Marine Ecology Progress Series*, 554, 201–211.
- Baylis, A. M. M., Orben, R. A., Costa, D. P., Tierney, M., Brickle, P., & Staniland, I. J. (2017). Habitat use and spatial fidelity of male South American sea lions during the nonbreeding period. *Ecology and Evolution*, February, 1–11. <https://doi.org/10.1002/ece3.2972>
- Baylis, A. M. M., Tierney, M., Staniland, I. J., & Brickle, P. (2018). Habitat use of adult male South American fur seals and a preliminary assessment of spatial overlap with trawl fisheries in the South Atlantic. *Mammalian Biology*, 93, 76–81. <https://doi.org/10.1016/j.mambio.2018.07.007>
- Bearhop, S., Adams, C. E., Waldron, S., Fuller, R. A., & Macleod, H. (2004). Determining trophic niche width: A novel approach using stable isotope analysis. *Journal of Animal Ecology*, 73(5), 1007–1012. <https://doi.org/10.1111/j.0021-8790.2004.00861.x>
- Begon, M., Townsend, C. R., & Harper, J. L. (2006). Ecology: From Individuals to Ecosystems. In *Freshwater Biology* (fourth, Vol. 51, Issue 9). Blackwell Publishing Ltd. <https://doi.org/10.1111/j.1365-2427.2006.01592.x>
- Bertrand, A., Ballón, M., & Chaigneau, A. (2010). Acoustic observation of living organisms reveals the upper limit of the oxygen minimum zone. *PLoS ONE*, 5(4). <https://doi.org/10.1371/journal.pone.0010330>
- Bertrand, A., Chaigneau, A., Peraltilla, S., Ledesma, J., Graco, M., Monetti, F., & Chavez, F. P. (2011). Oxygen: A fundamental property regulating pelagic ecosystem structure in the coastal southeastern tropical pacific. *PLoS ONE*, 6(12), 2–9.

<https://doi.org/10.1371/journal.pone.0029558>

- Bertrand, S., Joo, R., Arbulu Smet, C., Tremblay, Y., Barbraud, C., & Weimerskirch, H. (2012). Local depletion by a fishery can affect seabird foraging. *Journal of Applied Ecology*, *49*(5), 1168–1177. <https://doi.org/10.1111/j.1365-2664.2012.02190.x>
- Bolker, B. M. (2008). *Ecological models and data in R*. Princeton University Press.
- Bond, A. L., & Lavers, J. L. (2014). Climate change alters the trophic niche of a declining apex marine predator. *Global Change Biology*, *20*(7), 2100–2107. <https://doi.org/10.1111/gcb.12554>
- Bond, N. A., Cronin, M. F., Freeland, H., & Mantua, N. (2015). Causes and impacts of the 2014 warm anomaly in the NE Pacific. *Geophysical Research Letters*, *42*(9), 3414–3420. <https://doi.org/10.1002/2015GL063306>
- Bost, C. A., Georges, J. Y., Guinet, C. A., Cherel, Y., & Putz, K. (1997). Foraging habitat and food intake of satellite-tracked king penguins during the austral summer at Crozet Archipelago. *Marine Ecology Progress Series*, *150*, 21–33.
- Boyd, I. L., & Croxall, J. P. (1992). Diving behaviour of lactating Antarctic fur seals. *Canadian Journal of Zoology*, *70*, 919–928.
- Bradbury, J. W., & Gibson, R. M. (1983). Leks and mate choice. *Mate Choice*, 109–138.
- Brochier, T., Echevin, V., Tam, J., Chaigneau, A., Goubanova, K., & Bertrand, A. (2013). Climate change scenarios experiments predict a future reduction in small pelagic fish recruitment in the Humboldt Current system. *Global Change Biology*, *19*(6), 1841–1853. <https://doi.org/10.1111/gcb.12184>
- Butler, P. J. (2006). Aerobic dive limit. What is it and is it always used appropriately? *Comparative Biochemistry*, *145*(December 2004), 1–6. <https://doi.org/10.1016/j.cbpa.2006.06.006>
- Cain, A. J. (1953). Geography, ecology and coexistence in relation to the biological definition of the species. *Evolution*, *7*(March), 76–83.
- Calenge, C. (2006). The package “adehabitat” for the R software: A tool for the analysis of space and habitat use by animals. *Ecological Modelling*, *197*(3–4), 516–519. <https://doi.org/10.1016/j.ecolmodel.2006.03.017>
- Campagna, C., & Harcourt, R. (2021). Ethology and Behavioral Ecology of Otariids and the Odobenid. In *Ethology and Behavioral Ecology of Marine Mammals Ser.* [https://www.proquest.com/books/ethology-behavioral-ecology-otariids-odobenid/docview/2542044510/se-2?accountid=11526%0Ahttps://hud.alma.exlibrisgroup.com/discovery/openurl?institution=44HUD\\_INST&vid=44HUD\\_INST:Services&sid=ProQ%3Aebookcentral1&genre=book&](https://www.proquest.com/books/ethology-behavioral-ecology-otariids-odobenid/docview/2542044510/se-2?accountid=11526%0Ahttps://hud.alma.exlibrisgroup.com/discovery/openurl?institution=44HUD_INST&vid=44HUD_INST:Services&sid=ProQ%3Aebookcentral1&genre=book&)

- Cárdenas-Alayza, S. (2007). *Preferencia de uso de sitios para pariciones por hembras de Arctocephalus australis (ZIMMERMANN, 1783) en la Reserva Guanera Punta San Juan, Perú*. Universidad Peruana Cayetano Heredia.
- Cárdenas-Alayza, S. (2012). *Prey Abundance and Population Dynamics of South American Fur Seals (Arctocephalus australis) in Peru*. (Issue May). The University of British Columbia.
- Cárdenas-Alayza, S., Gutiérrez, D., & Tremblay, Y. (2021). Trends in sympatric otariid populations suggest resource limitations in the Peruvian Humboldt Current System. *Marine Environmental Research*, 105349. <https://doi.org/10.1016/j.marenvres.2021.105349>
- Chase, J. M., Abrams, A., Grover, J. P., Diehl, S., Chesson, P., & Holt, R. D. (2002). *The interaction between predation and competition : a review and synthesis*. 302–315.
- Chavez, F. P., Bertrand, A., Guevara-Carrasco, R., Soler, P., & Csirke, J. (2008). The northern Humboldt Current System: Brief history, present status and a view towards the future. *Progress in Oceanography*, 79(2–4), 95–105. <https://doi.org/10.1016/j.pocean.2008.10.012>
- Chesson, P. L., & Warner, R. R. (1981). Environmental Variability Promotes Coexistence in Lottery Competitive Systems. *American Naturalist*, 117(6), 923–943.
- Chilvers, B. L., & Wilkinson, I. S. (2009). Diverse foraging strategies in lactating New Zealand sea lions. *Marine Ecology Progress Series*, 378, 299–308. <https://doi.org/10.3354/meps07846>
- Christensen, V., & Pauly, D. (1992). ECOPATH II: A software for balancing steady-state ecosystem models and calculating network characteristics \* Programming language. *Ecological Modelling*, 61(681), 169–185. [https://doi.org/10.1016/0304-3800\(92\)90016-8](https://doi.org/10.1016/0304-3800(92)90016-8)
- Costa, D. P., & Gales, N. J. (2003). Energetics of a Benthic diver: Seasonal foraging ecology of the Australian sea lion, *Neophoca cinerea*. *Ecological Monographs*, 73(1), 27–43.
- Costa, D. P., Gales, N. J., & Crocker, D. E. (1998). Blood volume and diving ability of the New Zealand sea lion, *Phocarcos hookeri*. *Physiological Zoology*, 71(2), 208–213. <https://doi.org/10.1086/515911>
- Costa, D. P., Gales, N. J., & Goebel, M. E. (2001). Aerobic dive limit : how often does it occur in nature ? *Comparative Biochemistry and Physiology*, 129, 771–783.
- Costa, D. P., Weise, M. I. J., & Arnould, J. P. Y. (2007). Potential influences of whaling on the status and trends of pinniped populations. In *Whales, Whaling, and Ocean Ecosystems* (pp. 344–359). University of California Press. <https://doi.org/10.1525/california/9780520248847.003.0027>
- D’Andrea, R., Riolo, M., & Ostling, A. M. (2019). Generalizing clusters of similar species as a signature of coexistence under competition. *PLoS Computational Biology*, 15(1), 1–19. <https://doi.org/10.1371/journal.pcbi.1006688>

- Dalerum, F., & Angerbjörn, A. (2005). Resolving temporal variation in vertebrate diets using naturally occurring stable isotopes. *Oecologia*, *144*(4), 647–658. <https://doi.org/10.1007/s00442-005-0118-0>
- Daufresne, M., Lengfellner, K., & Sommer, U. (2009). Global warming benefits the small in aquatic ecosystems. *Proc. Nat. Acad. Sci.*, *106*(31), 12788–12793.
- De la Puente, S., López de la Lama, R., Benavente, S., Sueiro, J. C., & Pauly, D. (2020). Growing Into Poverty: Reconstructing Peruvian Small-Scale Fishing Effort Between 1950 and 2018. *Frontiers in Marine Science*, *7*(August), 1–14. <https://doi.org/10.3389/fmars.2020.00681>
- De Roy, T., Espinoza, E. R., & Trillmich, F. (2021). Cooperation and opportunism in Galapagos sea lion hunting for shoaling fish. *Ecology and Evolution*, *11*(14), 9206–9216. <https://doi.org/10.1002/ece3.7807>
- Dellinger, T., & Trillmich, F. (1999). Fish prey of the sympatric Galapagos fur seals and sea lions: Seasonal variation and niche separation. *Canadian Journal of Zoology*, *77*(8), 1204–1216. <https://doi.org/10.1139/z99-095>
- Demarcq, H. (2009). Trends in primary production, sea surface temperature and wind in upwelling systems (1998-2007). *Progress in Oceanography*, *83*(1–4), 376–385. <https://doi.org/10.1016/j.pocean.2009.07.022>
- DeNiro, M. J., & Epstein, S. (1978). Influence of diet on the distribution of carbon isotopes in animals. *Geochimica et Cosmochimica Acta*, *42*, 495–506.
- Drago, M., Cardona, L., Crespo, E. A., García, N., Ameghino, S., & Aguilar, A. (2010). Change in the foraging strategy of female South American sea lions (Carnivora: Pinnipedia) after parturition. *Scientia Marina*, *74*(3), 589–598. <https://doi.org/10.3989/scimar.2010.74n3589>
- Drago, M., Cardona, L., Franco-Trecu, V., Riet-Sapriza, F. G., Crespo, E. A., García, N., & Inchausti, P. (2021). Relationship between the female attendance pattern and pup growth rate in the South American sea lion (Carnivora). *Scientia Marina*, *85*(2), 81–90. <https://doi.org/10.3989/scimar.05128.008>
- Drago, M., Crespo, E. A., Aguilar, A., Cardona, L., García, N., Dans, S. L., & Goodall, N. (2009). Historic diet change of the South American sea lion in Patagonia as revealed by isotopic analysis. *Marine Ecology Progress Series*, *384*, 273–286. <https://doi.org/10.3354/meps08017>
- Drago, M., Franco-Trecu, V., Zenteno, L., Szteren, D., Crespo, E. A., Riet-Sapriza, F. G., Oliveira, L. R., Machado, R., Inchausti, P., & Cardona, L. (2015). Sexual foraging segregation in South American sea lions increases during the pre-breeding period in the Río de la Plata plume. *Marine Ecology Progress Series*, *525*, 261–272. <https://doi.org/10.3354/meps11236>

- du Pontavice, H., Gascuel, D., Reygondeau, G., Maureaud, A., & Cheung, W. W. L. (2020). Climate change undermines the global functioning of marine food webs. *Global Change Biology*, *26*(3), 1306–1318. <https://doi.org/10.1111/gcb.14944>
- Echevin, V., Aumont, O., Ledesma, J., & Flores, G. (2008). The seasonal cycle of surface chlorophyll in the Peruvian upwelling system: A modelling study. *Progress in Oceanography*, *79*(2–4), 167–176. <https://doi.org/10.1016/j.pocean.2008.10.026>
- Edwards, M. R. (2018). *Peruvian Pinnipeds as Archivists of ENSO Effects Thesis of*. Nova Southeastern University.
- Edwards, M. R., Cárdenas-Alayza, S., Adkesson, M. J., Daniels-Abdulahad, M., & Hirons, A. (2021). Peruvian Fur Seals as Archivists of El Niño Southern Oscillation Effects. *Frontiers in Marine Science*.
- Espinoza-Morriberón, D., Echevin, V., Colas, F., Tam, J., Ledesma, J., Vásquez, L., & Graco, M. (2017). Impacts of El Niño events on the Peruvian upwelling system productivity. *Journal of Geophysical Research: Oceans*, *122*(5423), 2647–2651. <https://doi.org/10.1002/2016JC012439>.Received
- Espinoza, P., & Bertrand, A. (2008). Revisiting Peruvian anchovy (*Engraulis ringens*) trophodynamics provides a new vision of the Humboldt Current system. *Progress in Oceanography*, *79*(2–4), 215–227. <https://doi.org/10.1016/j.pocean.2008.10.022>
- Espinoza, P., Lorrain, A., Ménard, F., Cherel, Y., Tremblay-Boyer, L., Argüelles, J., Tafur, R., Bertrand, S., Tremblay, Y., Ayón, P., Munaron, J. M., Richard, P., & Bertrand, A. (2017). Trophic structure in the northern Humboldt Current system: new perspectives from stable isotope analysis. *Marine Biology*, *164*(4), 0. <https://doi.org/10.1007/s00227-017-3119-8>
- Eurich, J. G., McCormick, M. I., & Jones, G. P. (2018). Direct and indirect effects of interspecific competition in a highly partitioned guild of reef fishes. *Ecosphere*, *9*(8). <https://doi.org/10.1002/ecs2.2389>
- Ferry, L. A., Clark, S. L., & Cailliet, G. M. (1997). Food Habits of Spotted Sand Bass (*Paralabrax maculatofasciatus*, Serranidae) from Bahia Los Angeles, Baja California. *Bull. Southern California Acad. Sci.*, *96*(1), 1–21.
- Forcada, J., Trathan, P. N., Reid, K., Murphy, E. J., & Croxall, J. P. (2006). Contrasting population changes in sympatric penguin species in association with climate warming. *Global Change Biology*, *12*, 411–423. <https://doi.org/10.1111/j.1365-2486.2006.01108.x>
- Fowler, S. L., Costa, D. P., Arnould, J. P. Y., Gales, N. J., & Kuhn, C. E. (2006). Ontogeny of diving behaviour in the Australian sea lion: Trials of adolescence in a late bloomer. *Journal of Animal*



*Ecology*, 75(2), 358–367. <https://doi.org/10.1111/j.1365-2656.2006.01055.x>

Fowler, S., Roush, R., & Wise, J. (2013). *Concepts in Biology*. OpenStax.

Francis, J., Boness, D., & Ochoa-Acuña, H. (1998). A protracted foraging and attendance cycle in female Juan Fernández fur seals. *Marine Mammal Science*, 14(3), 552–574.

Franco-Trecu, V., Aurióles-Gamboa, D., Arim, M., & Lima, M. (2012). Prepartum and postpartum trophic segregation between sympatrically breeding female *Arctocephalus australis* and *Otaria flavescens*. *Journal of Mammalogy*, 93(2), 514–521. <https://doi.org/10.1644/11-MAMM-A-174.1>

Franco-Trecu, V., Aurióles-Gamboa, D., & Inchausti, P. (2014). Individual trophic specialisation and niche segregation explain the contrasting population trends of two sympatric otariids. *Marine Biology*, 161(3), 609–618. <https://doi.org/10.1007/s00227-013-2363-9>

Franco-Trecu, V., Drago, M., Riet-Sapriza, F. G., Parnell, A., Frau, R., & Inchausti, P. (2013). Bias in diet determination: Incorporating traditional methods in Bayesian mixing models. *PLoS ONE*, 8(11), 1–8. <https://doi.org/10.1371/journal.pone.0080019>

Fréon, P., Bouchon, M., Mullan, C., García, C., & Ñiquen, M. (2008). Interdecadal variability of anchoveta abundance and overcapacity of the fishery in Peru. *Progress in Oceanography*, 79(2–4), 401–412. <https://doi.org/10.1016/j.pocean.2008.10.011>

Froese, R., & Pauly, D. (2000). FishBase 2000: Concepts, design and data sources. In *ICLARM*.

Ganoza, M. J. (2016). *Patrones de atención maternal de hembras de lobo marino fino (Arctocephalus australis ssp.) en Punta San Juan, Marcona, Ica, Perú*. Universidad Peruana Cayetano Heredia, Lima, Peru.

García-Godos, I. (2001). Patrones Morfológicos del Otolito Sagitta de algunos Peces Oseos del Mar Peruano. *Boletín Del Instituto Del Mar Del Perú*, 20(1–2), 46.

García-Rodríguez, F. J. (1999). *Cambios espaciales y estacionales en la estructura trófica y consumo del lobo marino de California, Zalophus californianus, en la región de las grandes islas, Golfo de California*. Instituto Politécnico Nacional, La Paz, Mexico.

García-Rodríguez, F. J., & Aurióles-Gamboa, D. (2004). Spatial and temporal variation in the diet of the California sea lion (*Zalophus californianus*) in the Gulf of California, Mexico Pacific Ocean. *Fishery Bulletin*, 102, 47–62. <http://fishbull.noaa.gov/1021/garcia.pdf>

García-Rodríguez, F. J., & de la Cruz-Agüero, J. (2011). A comparison of indexes for prey importance inferred from otoliths and cephalopod beaks recovered from pinniped scats. *Journal of Fisheries and Aquatic Science*, 6(2), 186–193. <https://doi.org/10.3923/jfas.2011.186.193>

Gentry, R. L., & Kooyman, G. L. (1987). *Fur seals: maternal strategies on land and at sea*. Princeton University Press.

- Georges, J. Y., Bonadonna, F., & Guinet, C. (2000). Foraging habitat and diving activity of lactating subantarctic fur seals in relation to sea-surface temperatures at Amsterdam Island. *Marine Ecology Progress Series*, 196, 291–304.
- Georges, J. Y., Tremblay, Y., & Guinet, C. (2000). Seasonal diving behaviour in lactating subantarctic fur seals on Amsterdam Island. *Polar Biology*, 23, 59–69.
- Giardino, G. V., Mandiola, M. A., Bastida, J., Denuncio, P. E., Bastida, R. O., & Rodríguez, D. H. (2016). Travel for sex: Long-range breeding dispersal and winter haulout fidelity in southern sea lion males. *Mammalian Biology*, 81(1), 89–95. <https://doi.org/10.1016/j.mambio.2014.12.003>
- Ginter, C. C., DeWitt, T. J., Fish, F. E., & Marshall, C. D. (2012). Fused traditional and geometric morphometrics demonstrate pinniped whisker diversity. *PLoS ONE*, 7(4), 1–10. <https://doi.org/10.1371/journal.pone.0034481>
- Gonzalez-Pestana, A., Mangel, J. C., Alfaro-Córdova, E., Acuña-Perales, N., Córdova-Zavaleta, F., Segura-Cobeña, E., Benites, D., Espinoza, M., Coasaca-Céspedes, J., Jiménez, A., Pingo, S., Moscoso, V., Alfaro-Shigueto, J., & Espinoza, P. (2021). Diet, trophic interactions and possible ecological role of commercial sharks and batoids in northern Peruvian waters. *Journal of Fish Biology*, 98(3), 768–783. <https://doi.org/https://doi.org/10.1111/jfb.14624>
- González-Suárez, M., & Cassini, M. H. (2014). Variance in male reproductive success and sexual size dimorphism in pinnipeds: testing an assumption of sexual selection theory. *Mammal Review*, 44(2), 88–93.
- Graco, M., Correa, D., García, W., & Sarmiento, M. (2016). Impactos del ENSO en la biogeoquímica del sistema de afloramiento frente a Perú central, febrero 2013 - diciembre 2015. *Boletín Trimestral Oceanográfico Imarpe*, 2(1), 2–6.
- Graco, M., Ledesma, J., Flores, G., & Girón, M. (2007). Nutrientes, oxígeno y procesos biogeoquímicos en el sistema de surgencias de la corriente de Humboldt frente a Perú. *Revista Peruana de Biología*, 14(1), 117–128. <https://doi.org/10.15381/rpb.v14i1.2165>
- Guinet, C., Bost, C. A., Pierre, J., Georges, J. Y., Mouchot, M. C., & Jouventin, P. (1997). Foraging behaviour of satellite-tracked king penguins in relation to sea-surface temperatures obtained by satellite telemetry at Crozet Archipelago, a study during three austral summers. *Marine Ecology Progress Series*, 150, 11–20.
- Gutiérrez, D., Bouloubassi, I., Sifeddine, A., Purca, S., Goubanova, K., Graco, M., Field, D., Méjanelle, L., Velasco, F., Lorre, A., Salvatelli, R., Quispe, D., Vargas, G., Dewitte, B., & Ortlieb, L. (2011). Coastal cooling and increased productivity in the main upwelling zone off Peru since the mid-twentieth century. *Geophysical Research Letters*, 38(L07603), 1–7.



<https://doi.org/10.1029/2010GL046324>

- Gutierrez, M., Gerlotto, F., Vasquez, L., & Chipollini, A. (2005). Anchovy and munida spatio-temporal dynamic and interactions of diel cycles of aggregation. *ICES CM*, 1. <http://www.ices.dk/products/CMdocs/2005/U/U2205.pdf>
- Gutiérrez, M., Ramirez, A., Bertrand, S., Móron, O., & Bertrand, A. (2008). Ecological niches and areas of overlap of the squat lobster “munida” (*Pleuroncodes monodon*) and anchoveta (*Engraulis ringens*) off Peru. *Progress in Oceanography*, 79(2–4), 256–263. <https://doi.org/10.1016/j.pocean.2008.10.019>
- Harcourt, R. (1992a). Factors affecting early mortality in the South American fur seal (*Arctocephalus australis*) in Peru: density-related effects and predation. *Journal of Zoology*, 226(2), 259–270.
- Harcourt, R. (1992b). Maternal aggression in the South American fur seal in Peru. *Canadian Journal of Zoology*, 70(2), 320–325. <https://doi.org/10.1139/z92-048>
- Harcourt, R., Bradshaw, C. J. A., Dickson, K., & Davis, L. S. (2002). Foraging ecology of a generalist predator, the female New Zealand fur seal. *Marine Ecology Progress Series*, 227, 11–24.
- Hardin, G. (1960). The Competitive Exclusion Principle. *Science*, 131(3409), 1292–1297.
- Hazen, E. L., Abrahms, B., Brodie, S., Carroll, G., Jacox, M. G., Savoca, M. S., Scales, K. L., Sydeman, W. J., & Bograd, S. J. (2019). Marine top predators as climate and ecosystem sentinels. *Frontiers in Ecology and the Environment*, 17(10), 565–574. <https://doi.org/10.1002/fee.2125>
- Hirons, A. (2001). *Trophic dynamics of pinniped populations in Alaska using stable carbon and nitrogen isotope ratios*. University of Alaska Fairbanks.
- Hirons, A., Schell, D., & Finney, B. (2001). Temporal records of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  in North Pacific pinnipeds: inferences regarding environmental change and diet. *Oecologia*, 129(4), 591–601. <https://doi.org/10.1007/s004420100756>
- Hückstädt, L. A., Quiñones, R. A., Sepúlveda, M., & Costa, D. P. (2014). Movement and diving patterns of juvenile male South American sea lions off the coast of central Chile. *Marine Mammal Science*, 30(3), 1175–1183. <https://doi.org/10.1111/mms.12085>
- Hückstädt, L. A., Tift, M. S., Riet-Sapriza, F., Franco-Trecu, V., Baylis, A. M. M., Orben, R. A., Arnould, J. P. Y., Sepúlveda, M., Santos-Carvallo, M., Burns, J. M., & Costa, D. P. (2016). Regional variability in diving physiology and behavior in a widely distributed air-breathing marine predator, the South American sea lion (*Otaria byronia*). *Journal of Experimental Biology*, 219(15), 2320–2330. <https://doi.org/10.1242/jeb.138677>
- Hutchinson, G. E., & MacArthur, R. H. (1959). A theoretical ecological model of size distributions among species of animals. *The American Naturalist*, 93(869), 117–125.

- IMARPE. (2006). *Anuario Científico-Tecnológico IMARPE*.
- IMARPE. (2012). *Anuario científico tecnológico imarpe*.
- IMARPE. (2013). *Anuario Científico-Tecnológico IMARPE*.
- IMARPE. (2014a). *50 Años de Mar y Ciencia* (IMARPE (ed.); 1st Ed.).
- IMARPE. (2014b). *Anuario Científico-Tecnológico IMARPE*.
- IMARPE. (2015). *Anuario Científico-Tecnológico IMARPE*.
- IMARPE. (2016). *Anuario Científico-Tecnológico IMARPE*.
- IMARPE. (2017). *Anuario Científico-Tecnológico IMARPE*.
- IMARPE. (2018). *Anuario Científico-Tecnológico IMARPE*.
- Jackson, A. L., Inger, R., Parnell, A. C., & Bearhop, S. (2011). Comparing isotopic niche widths among and within communities : SIBER – Stable Isotope Bayesian Ellipses in R. *The Journal of Animal Ecology*, *80*, 595–602. <https://doi.org/10.1111/j.1365-2656.2011.01806.x>
- Jahncke, J., & Goya, E. (1997). Variación latitudinal y estacional en la dieta del guanay (*Leucocarbo bougainvillii*) y el piquero peruano (*Sula variegata*) en la costa peruana. *Boletín Del Instituto Del Mar Del Peru*, *16*(1), 23–41.
- Jeanniard du Dot, T., Thomas, A. C., Cherel, Y., Trites, A. W., & Guinet, C. (2017). Combining hard-part and DNA analyses of scats with biologging and stable isotopes can reveal different diet compositions and feeding strategies within a fur seal population. *Marine Ecology Progress Series*, *584*(December), 1–16. <https://doi.org/10.3354/meps12381>
- Jeglinski, J. W. E., Goetz, K. T., Werner, C., Costa, D. P., & Trillmich, F. (2013). Same size – same niche? Foraging niche separation between sympatric juvenile Galapagos sea lions and adult Galapagos fur seals. *Journal of Animal Ecology*, *82*, 694–706. <https://doi.org/10.1111/1365-2656.12019>
- Jessopp, M. J., Forcada, J., Reid, K., Trathan, P. N., & Murphy, E. J. (2004). Winter dispersal of leopard seals (*Hydrurga leptonyx*): environmental factors influencing demographics and seasonal abundance. *Journal of Zoology, London*, *263*, 251–258. <https://doi.org/10.1017/S0952836904005102>
- Kassen, R. (2002). The experimental evolution of specialists, generalists, and the maintenance of diversity. *Journal of Evolutionary Biology*, *15*(2), 173–190. <https://doi.org/10.1046/j.1420-9101.2002.00377.x>
- Kernaléguen, L., Cazelles, B., Arnould, J. P. Y., Richard, P., Guinet, C., & Cherel, Y. (2012). Long-term species, sexual and individual variations in foraging strategies of fur seals revealed by stable isotopes in whiskers. *PLoS ONE*, *7*(3). <https://doi.org/10.1371/journal.pone.0032916>

- Killick, R., & Eckley, I. A. (2014). Changepoint: An R package for changepoint analysis. *Journal of Statistical Software*, 58(3), 1–19. <https://doi.org/10.18637/jss.v058.i03>
- Knox, T. C., Baylis, A. M. M., & Arnould, J. P. Y. (2018). Foraging site fidelity in male Australian fur seals. *Marine Biology*, 165(7). <https://doi.org/10.1007/s00227-018-3368-1>
- Koen Alonso, M., Crespo, E. A., García, N. A., Pedraza, S. N., Mariotti, P. A., Berón Vera, B., & Mora, N. J. (2001). Food habits of *Dipturus chilensis* (Pisces: Rajidae) off Patagonia, Argentina. *ICES Journal of Marine Science*, 58(1), 288–297. <https://doi.org/10.1006/jmsc.2000.1010>
- Koen Alonso, M., Crespo, E. A., García, N. A., Pedraza, S. N., Mariotti, P. A., & Mora, N. J. (2002). Fishery and ontogenetic driven changes in the diet of the spiny dogfish, *Squalus acanthias*, in Patagonian waters, Argentina. *Environmental Biology of Fishes*, 63, 193–202. <https://doi.org/10.1023/A:1014229432375>
- Koen Alonso, M., Crespo, E. A., Pedraza, S. N., Garcia, N. A., & Coscarella, M. A. (2000). Food habits of the South American sea lion, *Otaria flavescens*, off Patagonia, Argentina. *Fishery Bulletin*, 98, 250–263.
- Kortsch, S., Primicerio, R., Fossheim, M., Dolgov, A. V., & Aschan, M. (2015). Climate change alters the structure of arctic marine food webs due to poleward shifts of boreal generalists. *Proceedings of the Royal Society B: Biological Sciences*, 282(1814). <https://doi.org/10.1098/rspb.2015.1546>
- Krebs, C. J. (1999). *Ecological Methodology* (Addison-Wesley Educational Publishers (ed.); 2nd ed.). Benjamin Cummings.
- Krüger, O., Wolf, J. B. W., Jonker, R. M., Hoffman, J. I., & Trillmich, F. (2014). Disentangling the contribution of sexual selection and ecology to the evolution of size dimorphism in pinnipeds. *Evolution*, 68(5), 1485–1496.
- L’Heureux, M. L., Takahashi, K., Watkins, A. B., Barnston, A. G., Becker, E. J., Di Liberto, T. E., Gamble, F., Gottschalck, J., Halpert, M. S., Huang, B., Mosquera-Vásquez, K., & Wittenberg, A. T. (2017). Observing and predicting the 2015/16 El Niño. *Bulletin of the American Meteorological Society*, 98(7), 1363–1382. <https://doi.org/10.1175/BAMS-D-16-0009.1>
- Lavielle, M. (2005). Using penalized contrasts for the change-point problem. *Signal Processing*, 85(8), 1501–1510. <https://doi.org/10.1016/j.sigpro.2005.01.012>
- Layman, C. A., Quatrocchi, J. P., Peyer, C. M., & Allgeier, J. E. (2007). Niche width collapse in a resilient top predator following ecosystem fragmentation. *Ecology Letters*, 10, 937–944. <https://doi.org/10.1111/j.1461-0248.2007.01087.x>
- Le Boeuf, B. J., Crocker, D. E., Costa, D. P., Blackwell, S. B., Webb, P. M., & Houser, D. S. (2000). Foraging ecology of northern elephant seals. *Ecological Monographs*, 70(3), 353–382.

[https://doi.org/10.1890/0012-9615\(2000\)070\[0353:FEONES\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2000)070[0353:FEONES]2.0.CO;2)

- Lima, R. C. De, Franco, V., Damián, T., Pablo, G. V., & Eduardo, I. (2019). Individual foraging specialization and sexual niche segregation in South American fur seals. *Marine Biology*, 166(32), 1–12. <https://doi.org/10.1007/s00227-019-3480-x>
- Majluf, P. (1987). *Reproductive ecology of female South American fur seals at Punta San Juan, Peru*. University of Cambridge, Cambridge, UK.
- Majluf, P. (1991). El Niño Effects on Pinnipeds in Peru. In F. Trillmich & K. Ono (Eds.), *Pinnipeds and El Niño* (pp. 55–65). Springer-Verlag.
- Majluf, P. (1992). Timing of births and juvenile mortality in the South American fur seal in Peru. *Journal of Zoology*, 227(3), 367–383.
- Majluf, P. (1989). Reproductive ecology of South American fur seals in Peru. *The Peruvian Upwelling Ecosystem and Interactions. ICLARM Conference Proceedings 18,438p. Instituto Del Mar Del Perú (IMARPE)*, 332–343.
- Majluf, P., & Reyes, J. C. (1989). The marine mammals of Peru: A review. *The Peruvian Upwelling Ecosystem: Dynamics and Interactions. ICLARM Conference Proceedings*.
- Majluf, P., & Trillmich, F. (1981). Distribution and abundance of sea lions (*Otaria byronia*) and fur seals (*Arctocephalus australis*) in Peru. *Zeitschrift Für Säugetierkunde*, 46.
- McPhaden, M. J., Zebiak, S. E., & Glantz, M. H. (2006). ENSO as an integrating concept in earth science. *Science*, 314(5806), 1740–1745. <https://doi.org/10.1126/science.1132588>
- Meynier, L., Morel, P. C. H., Chilvers, B. L., Mackenzie, D. D. S., MacGibbon, A., & Duignan, P. J. (2008). Temporal and sex differences in the blubber fatty acid profiles of the New Zealand sea lion *Phocarctos hookeri*. *Marine Ecology Progress Series*, 366, 271–279. <https://doi.org/10.3354/meps07617>
- Muck, P., & Fuentes, H. (1987). Sea lion and fur seal predation on the Peruvian anchoveta, 1953 to 1982. In D. Pauly & I. Tsukayama (Eds.), *The Peruvian anchoveta ad its upwelling ecosystem: Three decades of change. ICLARM Studies and Reviews 15*. (first, pp. 234–247). IMARPE/GTZ.
- Murphy, R. C. (1925). *Bird islands of Peru: the record of a sojourn on the west coast*. GP Putnam's Sons.
- Naya, D. E., Arim, M., & Vargas, R. (2002). Diet of South American fur seals (*Arctocephalus Australis*) in Isla de Lobos, Uruguay. *Marine Mammal Science*, 18(3), 734–745. <https://doi.org/10.1111/j.1748-7692.2002.tb01070.x>
- Newsome, S. D., Clementz, M. T., & Koch, P. L. (2010). Using stable isotope biogeochemistry to study marine mammal ecology. *Marine Mammal Science*, 26(3), 509–572.

<https://doi.org/10.1111/j.1748-7692.2009.00354.x>

- Oliveira, L. R. (2011). Vulnerability of South American Pinnipeds Under El Niño Southern Oscillation Events. In D. S. Casalegno (Ed.), *Global Warming Impacts - Case Studies on the Economy, Human Health, and on Urban and Natural Environments* (pp. 237–252). InTech. <https://doi.org/10.5772/25204>
- Oliveira, L. R., Arias-Schreiber, M., Meyer, D., & Morgante, J. S. (2006). Effective population size in a bottlenecked fur seal population. *Biological Conservation*, *131*(4), 505–509. <https://doi.org/10.1016/j.biocon.2006.02.017>
- Oliveira, L. R., Fraga, L. D., & Majluf, P. (2012). Effective population size for South American sea lions along the Peruvian coast: the survivors of the strongest El Niño event in history. *Journal of the Marine Biological Association of the United Kingdom*, *92*(8), 1–7. <https://doi.org/10.1017/S0025315411001871>
- Oliveira, L. R., Gehara, M. C. M., Fraga, L. D., Lopes, F., Túnez, J. I., Cassini, M. H., Majluf, P., Cárdenas-Alayza, S., Pavés, H. J., Crespo, E. A., García, N., De Castro, R. L., Hoelzel, A. R., Sepúlveda, M., Olavarría, C., Valiati, V. H., Quiñones, R., Pérez-Alvarez, M. J., Ott, P. H., & Bonatto, S. L. (2017). Ancient female philopatry, asymmetric male gene flow, and synchronous population expansion support the influence of climatic oscillations on the evolution of South American sea lion (*Otaria flavescens*). *PLoS ONE*, *12*(6). <https://doi.org/10.1371/journal.pone.0179442>
- Oliveira, L. R., Meyer, D., Hoffman, J., Majluf, P., & Morgante, J. S. (2009). Evidence of a genetic bottleneck in an El Niño affected population of South American fur seals, *Arctocephalus australis*. *Journal of the Marine Biological Association of the United Kingdom*, *89*(8), 1717–1725. <https://doi.org/10.1017/S0025315409000162>
- Pablo-Rodríguez, N., Aurióles-Gamboa, D., & Montero-Muñoz, J. L. (2016). Niche overlap and habitat use at distinct temporal scales among the California sea lions (*Zalophus californianus*) and Guadalupe fur seals (*Arctocephalus philippii townsendi*). *Marine Mammal Science*, *32*(2), 466–489. <https://doi.org/10.1111/mms.12274>
- Pacala, S. W., & Roughgarden, J. (1985). Populations Experiments with the Anolis of St. Maarten and St. Eustatius. *Ecology*, *66*(1), 129–141.
- Páez-Rosas, D., & Aurióles-Gamboa, D. (2010). Alimentary niche partitioning in the Galapagos sea lion, *Zalophus wollebaeki*. *Marine Biology*, *157*(12), 2769–2781. <https://doi.org/10.1007/s00227-010-1535-0>
- Páez-Rosas, D., & Aurióles-Gamboa, D. (2014). Spatial variation in the foraging behaviour of the



- Galapagos sea lions (*Zalophus wollebaeki*) assessed using scat collections and stable isotope analysis. *Journal of the Marine Biological Association of the United Kingdom*, 94(06), 1099–1107. <https://doi.org/10.1017/S002531541300163X>
- Páez-Rosas, D., Aurióles-Gamboa, D., Alava, J. J., & Palacios, D. M. (2012). Stable isotopes indicate differing foraging strategies in two sympatric otariids of the Galapagos Islands. *Journal of Experimental Marine Biology and Ecology*, 424–425, 44–52. <https://doi.org/10.1016/j.jembe.2012.05.001>
- Páez-Rosas, D., Vaca, L., Pepolas, R., Wollocombe, R., De Roy, T., & Rivas-Torres, G. (2020). Hunting and cooperative foraging behavior of Galapagos sea lion: An attack to large pelagics. *Marine Mammal Science*, 36(1), 386–391. <https://doi.org/10.1111/mms.12646>
- Page, B., Mckenzie, J., & Goldsworthy, S. D. (2005). Dietary resource partitioning among sympatric New Zealand and Australian fur seals. *Marine Ecology Progress Series*, 293, 283–302.
- Page, B., Mckenzie, J., Sumner, M. D., Coyne, M., & Goldsworthy, S. D. (2006). *Spatial separation of foraging habitats among New Zealand fur seals*. 323, 263–279. <https://doi.org/doi:10.3354/meps323263>
- Palstra, F. P., & Ruzzante, D. E. (2008). Genetic estimates of contemporary effective population size: What can they tell us about the importance of genetic stochasticity for wild population persistence? *Molecular Ecology*, 17(15), 3428–3447. <https://doi.org/10.1111/j.1365-294X.2008.03842.x>
- Paredes, R., & Arias-Schreiber, M. (1999). Dieta del Lobo fino (*Arctocephalus australis*) y Lobo chusco (*Otaria byonia*) en la costa peruana durante Mayo y Junio de 1999. *Imarpe*.
- Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, 421(6918), 37–42. <https://doi.org/10.1038/nature01286>
- Passuni, G., Barbraud, C., Chaigneau, A., Demarcq, H., Ledesma, J., Bertrand, A., Castillo, R., Perea, A., Mori, J., Viblan, V. A., Torres-Maita, J., & Bertrand, S. (2016). Seasonality in marine ecosystems: Peruvian seabirds, anchovy, and oceanographic conditions. *Ecology*, 97(1), 182–193. <https://doi.org/10.1890/14-1134.1>
- Paulino, C., Escudero, L., Alburqueque, E., & Xu, H. (2019). Characteristics of the El Niño Costero 2017 through satellite observations. *Bol Inst Mar Perú*, 34(1), 26.
- Pauly, D., Trites, A. W., Capuli, E., & Christensen, V. (1998). Diet composition and trophic levels of marine mammals. *ICES Journal of Marine Science*, 55(3), 467–481. <https://doi.org/10.1006/jmsc.1997.0280>
- Pianka, E. R. (1974). Niche Overlap and Diffuse Competition. *Proc. Nat. Acad. Sci.*, 71(5), 2141–2145.

- Pinheiro, J. C., & Bates, D. M. (2000). Linear mixed-effects models: basic concepts and examples. In *Mixed-effects models in S and S-Plus*. Springer.
- Porras-Peters, H., Aurióles-Gamboa, D., Cruz-Escalona, V. H., & Koch, P. L. (2008). Trophic level and overlap of sea lions (*Zalophus californianus*) in the Gulf of California, Mexico. *Marine Mammal Science*, 24(July), 554–576. <https://doi.org/10.1111/j.1748-7692.2008.00197.x>
- Post, D. M., Ecology, S., & Mar, N. (2002). Using Stable Isotopes to Estimate Trophic Position. *Ecology*, 83(3), 703–718.
- R Core Team. (2018). R: A Language and Environment for Statistical Computing. In *R Foundation for Statistical Computing* (3.6.2). [www.r-project.org](http://www.r-project.org)
- Ralls, K., & Mesnick, S. L. (2002). Sexual dimorphism. In J. Perrin, WF, Würsig, B., Thewissen (Ed.), *Encyclopedia of Marine Mammals*. Academic Press, San Diego.
- Riet-Sapiriza, F. G., Costa, D. P., Franco-Trecu, V., Marín, Y., Chocca, J., González, B., Beathyate, G., Louise Chilvers, B., & Hückstadt, L. A. (2013). Foraging behavior of lactating South American sea lions (*Otaria flavescens*) and spatial-temporal resource overlap with the Uruguayan fisheries. *Deep-Sea Research Part II: Topical Studies in Oceanography*, 88–89, 106–119. <https://doi.org/10.1016/j.dsr2.2012.09.005>
- Riverón, S., Raoult, V., Baylis, A. M. M., Jones, K. A., Slip, D. J., & Harcourt, R. G. (2021). Pelagic and benthic ecosystems drive differences in population and individual specializations in marine predators. *Oecologia*, 196(3), 891–904. <https://doi.org/10.1007/s00442-021-04974-z>
- Roa-Pascuali, L., Demarcq, H., & Nieblas, A. E. (2015). Detection of mesoscale thermal fronts from 4km data using smoothing techniques: Gradient-based fronts classification and basin scale application. *Remote Sensing of Environment*, 164, 225–237. <https://doi.org/10.1016/j.rse.2015.03.030>
- Robinson, C. J. (2016). Evolution of the 2014–2015 sea surface temperature warming in the central west coast of Baja California, Mexico, recorded by remote sensing. *Geophysical Research Letters*, 43(January 2000), 7066–7071. <https://doi.org/10.1002/2016GL069356>.Received
- Rodríguez, D. H., Dassis, M., Ponce de León, A., Barreiro, C., Farenga, M., Bastida, R. O., & Davis, R. W. (2013). Foraging strategies of Southern sea lion females in the La Plata River Estuary (Argentina-Uruguay). *Deep-Sea Research Part II: Topical Studies in Oceanography*, 88–89, 120–130. <https://doi.org/10.1016/j.dsr2.2012.07.012>
- Ryan, H. F., & Noble, M. (2002). Sea level response to ENSO along the central California coast: How the 1997-1998 event compares with the historic record. *Progress in Oceanography*, 54(1–4), 149–169. [https://doi.org/10.1016/S0079-6611\(02\)00047-2](https://doi.org/10.1016/S0079-6611(02)00047-2)



- Sarmiento-Devia, R., Sepúlveda, M., Pavez, G., Valdés, J., Canto, A., Orellana, M., & Oliva, D. (2020). Diet composition of an opportunistic predator from an upwelling area in the Southeastern Pacific. *Austral Ecology*, 45(8), 1145–1155. <https://doi.org/10.1111/aec.12944>
- Schoener, T. W. (1974). Resource partitioning in ecological communities. *Science*, 185(4145), 27–39.
- Schoener, T. W. (1983). Field experiments on interspecific competition. *American Naturalist*, 122(2), 240–285. <https://doi.org/10.1086/284133>
- Schulz, T. M., & Bowen, W. D. (2005). The evolution of lactation strategies in pinnipeds: A phylogenetic analysis. *Ecological Monographs*, 75(2), 159–177. <https://doi.org/10.1890/04-0319>
- Schwarz, J. F. L., Mews, S., DeRango, E. J., Langrock, R., Piedrahita, P., Páez-Rosas, D., & Krüger, O. (2021). Individuality counts: A new comprehensive approach to foraging strategies of a tropical marine predator. *Oecologia*, 195(2), 313–325. <https://doi.org/10.1007/s00442-021-04850-w>
- Segura, A. M., Franco-Trecu, V., Franco-Fraguas, P., & Arim, M. (2015). Gape and energy limitation determine a humped relationship between trophic position and body size. *Canadian Journal of Fisheries and Aquatic Sciences*, 72(2), 198–205. <https://doi.org/10.1139/cjfas-2014-0093>
- SERNANP. (2016). *Línea Base Biológica Terrestre y Marina de la Reserva Nacional Sistema de Islas, Islotes y Puntas Guaneras- Punta Coles (Ilo , Moquegua)* (SERNANP (ed.); First).
- Sielfeld, W., Barraza, J., & Amado, N. (2018). Local feeding patterns of the South American sea lion *Otaria byronia*: Case of Punta Patache, Northern Chile. *Revista de Biología Marina y Oceanografía*, 53(3), 307–319. <https://doi.org/10.22370/rbmo.2018.53.3.1356>
- Sigler, M. F., Tollit, D. J., Vollenweider, J. J., Thedinga, J. F., Csepp, D. J., Womble, J. N., Wong, M. A., Rehberg, M. J., & Trites, A. W. (2009). Steller sea lion foraging response to seasonal changes in prey availability. *Marine Ecology Progress Series*, 388, 243–261. <https://doi.org/10.3354/meps08144>
- Soto, K. H. (2004). *The Effects of Prey Abundance on the Diet, Maternal Attendance and Pup Mortality of the South American Sea Lion (Otaria flavescens) in Peru*. The University of British Columbia.
- Soto, K. H., & Trites, A. W. (2011). South American sea lions in Peru have a lek-like mating system. *Marine Mammal Science*, 27(2), 306–333. <https://doi.org/10.1111/j.1748-7692.2010.00405.x>
- Soto, K. H., Trites, A. W., & Arias-Schreiber, M. (2004). The effects of prey availability on pup mortality and the timing of birth of South American sea lions (*Otaria flavescens*) in Peru. *Journal of Zoology*, 264(4), 419–428. <https://doi.org/10.1017/S0952836904005965>
- Sprogis, K. R., Christiansen, F., Wandres, M., & Bejder, L. (2018). El Niño Southern Oscillation influences the abundance and movements of a marine top predator in coastal waters. *Global Change Biology*, 24(3), 1085–1096. <https://doi.org/10.1111/gcb.13892>

- Staniland, I. J., & Robinson, S. L. (2008). Segregation between the sexes: Antarctic fur seals, *Arctocephalus gazella*, foraging at South Georgia. *Animal Behaviour*, *75*, 1581–1590. <https://doi.org/10.1016/j.anbehav.2007.10.012>
- Stevens, M. A., & Boness, D. J. (2003). Influences of habitat features and human disturbance on use of breeding sites by a declining population of southern fur seals (*Arctocephalus australis*). *Journal of Zoology*, *260*(2), 145–152. <https://doi.org/10.1017/S0952836903003583>
- Szteren, D., Auriolles-Gamboa, D., Labrada-Martagón, V., Hernández-Camacho, C. J., & María, M. (2018). Historical age - class diet changes in South American fur seals and sea lions in Uruguay. *Marine Biology*, *165*(February). <https://doi.org/10.1007/s00227-018-3315-1>
- Tam, J., Taylor, M. H., Blaskovic, V., Espinoza, P., Michael Ballón, R., Díaz, E., Wosnitza-Mendo, C., Argüelles, J., Purca, S., Ayón, P., Quipuzcoa, L., Gutiérrez, D., Goya, E., Ochoa, N., & Wolff, M. (2008). Trophic modeling of the Northern Humboldt Current Ecosystem, Part I: Comparing trophic linkages under La Niña and El Niño conditions. *Progress in Oceanography*, *79*(2–4), 352–365. <https://doi.org/10.1016/j.pocean.2008.10.007>
- Taylor, R. S., & Friesen, V. L. (2017). The role of allochrony in speciation. *Molecular Ecology*, *26*(13), 3330–3342. <https://doi.org/10.1111/mec.14126>
- Thompson, D., Duck, C. D., McConnell, B. J., & Garrett, J. (1998). Foraging behaviour and diet of lactating female southern sea lions (*Otaria flavescens*) in the Falkland Islands. *Journal Zoology of London*, *246*, 135–146.
- Tollit, D. J., Steward, M. J. ., Thompson, P. M., Pierce, G. J., Santos, M. B., & Hughes, S. (1997). Species and size differences in the digestion of otoliths and beaks: implications for estimates of pinniped diet composition. *Canadian Journal of Fisheries and Aquatic Science*, *54*, 105–119. <https://doi.org/https://doi.org/10.1139/f96-264>
- Tovar, H., Cabrera, D., & Farfán del Pino, M. (1985). Impacto del Fenómeno “El Niño” en la población de lobos marinos en Punta San Juan. In “*El Niño*” *Su impacto en la fauna marina*. *Boletín IMARPE. Volumen extraordinario* (pp. 195–200).
- Tovar, H., & Fuentes, H. (1984). Magnitud poblacional de lobos marinos en el litoral peruano en marzo de 1984. *Informes Del Instituto Del Mar (Peru)*, *88*, 32pp.
- Trathan, P. N., Forcada, J., & Murphy, E. J. (2007). Environmental forcing and Southern Ocean marine predator populations: Effects of climate change and variability. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *362*(1488), 2351–2365. <https://doi.org/10.1098/rstb.2006.1953>
- Tremblay, Y., Robinson, P. W., & Costa, D. P. (2009). A Parsimonious Approach to Modeling Animal

- Movement Data. *PLoS ONE*, 4(3). <https://doi.org/10.1371/journal.pone.0004711>
- Trillmich, F., & Dellinger, T. (1991). The Effects of El Niño on Galapagos Pinnipeds. In F. Trillmich & K. Ono (Eds.), *Pinnipeds and El Niño* (pp. 66–74). Springer-Verlag.
- Trillmich, F., Kooyman, G. L., Majluf, P., & Sánchez-Griñan, M. (1987). Attendance and diving behavior of South American fur seals during El Niño in 1983. In R. L. Gentry & G. Kooyman (Eds.), *Fur Seals: Maternal Strategies on Land and at Sea* (pp. 153–167). Springer-Verlag.
- Vásquez, P. (1995). *Determinación de los hábitos alimentarios de Arctocephalus australis y Otaria byronia en Punta San Juan de Marcona, Ica, Perú*. Universidad Nacional Agraria La Molina, Lima, Peru.
- Villegas-Amtmann, S., & Costa, D. (2017). *Intraspecific variation in feeding strategies of Galapagos sea lions: A case of trophic specialization*. 1–16. <https://doi.org/10.1371/journal.pone.0185165>
- Villegas-Amtmann, S., Costa, D. P., Tremblay, Y., Salazar, S., & Aurióles-Gamboa, D. (2008). Multiple foraging strategies in a marine apex predator , the Galapagos sea lion *Zalophus wollebaeki*. *Marine Ecology Progress Series*, 363, 299–309. <https://doi.org/10.3354/meps07457>
- Villegas-Amtmann, S., Jeglinski, J. W. E. E., Costa, D. P., Robinson, P. W., & Trillmich, F. (2013). Individual Foraging Strategies Reveal Niche Overlap between Endangered Galapagos Pinnipeds. *PloS One*, 8(8). <https://doi.org/10.1371/journal.pone.0070748>
- Waite, J. N., Burkanov, V. N., & Andrews, R. D. (2012). Prey competition between sympatric Steller sea lions (*Eumetopias jubatus*) and northern fur seals (*Callorhinus ursinus*) on Lovushki Island, Russia. *Canadian Journal of Zoology*, 90(1), 110–127. <https://doi.org/10.1139/Z11-117>
- Waite, J. N., Trumble, S. J., Burkanov, V. N., & Andrews, R. D. (2012). Resource partitioning by sympatric Steller sea lions and northern fur seals as revealed by biochemical dietary analyses and satellite telemetry. *Journal of Experimental Marine Biology and Ecology*, 416–417, 41–54. <https://doi.org/10.1016/j.jembe.2012.02.009>
- Walther, G., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., Fromentin, J., I, O. H., & Bairlein, F. (2002). Ecological response to recent climate change. *Nature*, 416, 389–395.
- Ward, J. H., & Hook, M. E. (1963). Application of an hierarchical grouping procedure to a problem of grouping profiles. *Educational and Psychological Measurement*, 23(1), 69–81. <https://doi.org/10.1177/001316446302300107>
- Weise, M. J., & Costa, D. P. (2007). Total body oxygen stores and physiological diving capacity of California sea lions as a function of sex and age. *Journal of Experimental Biology*, 210, 278–289. <https://doi.org/10.1242/jeb.02643>
- Werner, R., & Campagna, C. (1995). Diving behaviour of lactating southern sea lions ( *Otaria*

- flavescens ) in Patagonia. *Canadian Journal of Zoology*, 73, 1975–1982.
- Wickens, P., & York, A. (1997). Comparative Population Dynamics of Fur Seals. *Marine Mammal Science*, 13(April), 241–292.
- Williams, E. E. (1983). *Lizard Ecology: Studies of a Model Organism* (R. B. Huey, E. R. Pianka, & T. W. Schoener (eds.)). Harvard University Pres.
- Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society. Series B: Statistical Methodology*, 73(1), 3–36. <https://doi.org/10.1111/j.1467-9868.2010.00749.x>
- Wright, S. (1990). Evolution in mendelian populations. *Bulletin of Mathematical Biology*, 52(1–2), 241–295. <https://doi.org/10.1007/BF02459575>
- Yuffra, M. S., Jimenez, G. H., Torero, M. G., & Bertrand, A. (2018). The spatio-temporal relationship of red squat lobster (*Pleuroncodes monodon*) with the superficial water masses of the Peruvian Sea between 1998 and 2016. *2017 IEEE/OES Acoustics in Underwater Geosciences Symposium, RIO Acoustics 2017, 2018-Janua*, 1–7. <https://doi.org/10.1109/RIOAcoustics.2017.8349737>
- Yurkowski, D. J., Hussey, N. E., Ferguson, S. H., & Fisk, A. T. (2018). A temporal shift in trophic diversity among a predator assemblage in a warming Arctic. *Royal Society Open Science*, 5(10). <https://doi.org/10.1098/rsos.180259>
- Zaba, K. D., & Rudnick, D. L. (2016). The 2014-2015 warming anomaly in the Southern California Current System observed by underwater gliders. *Geophysical Research Letters*, 43(3), 1241–1248. <https://doi.org/10.1002/2015GL067550>
- Zavala, R., Gutiérrez, D., Morales, R., Grunwaldt, A., Gonzales, N., Tam, J., Rodríguez, C., & Bucaram, S. (Eds.). (2019). *Avances del Perú en la Adaptación al Cambio Climático del Sector Pesquero y del Ecosistema Marino-Costero* (first). Banco Interamericano de Desarrollo.
- Zavalaga, C., Paredes, R., & Arias-Schreiber, M. (1998). Dieta del lobo fino (*Arctocephalus australis*) y del lobo chusco (*Otaria byronia*) en la costa sur del Perú, en febrero de 1998. *Informe Progresivo Del Instituto Del Mar Del Perú*, 79, 3–16.

## Références Iconographiques

Illustrations d'otariidés réalisées par Paulo Colchao et données à l'auteur pour être utilisées dans cette thèse avec autorisation.

Dessin sur les pages de titre réalisé via [www.canva.com/design](http://www.canva.com/design)