



Actinobacteria communities in natural and anthropogenic environments

Andrea Faitova

► To cite this version:

Andrea Faitova. Actinobacteria communities in natural and anthropogenic environments. Bacteriology. Université de Lyon; Univerzita Karlova (Prague), 2021. English. NNT: 2021LYSE1028 . tel-03667945

HAL Id: tel-03667945

<https://theses.hal.science/tel-03667945>

Submitted on 13 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

N°d'ordre NNT :
2021LYSE1028



THESE de DOCTORAT DE L'UNIVERSITE DE LYON

opérée au sein de
l'Université Claude Bernard Lyon 1

Ecole Doctorale N° 341
Ecosystèmes Evolution Modélisation Microbiologie

Spécialité de doctorat : Ecologie Microbienne
Discipline : BIOLOGIE

Soutenue publiquement le 11/02/2021, par :

Andrea Faitova (Buresova)

Actinobacteria communities in natural and anthropogenic environments

Devant le jury composé de :

JURADO Valme (Chercheure – Institute des ressources naturelles et d'agrobiologie de Seville, Espagne) - *Rapporteure*

KYSELKOVA Martina (Chercheure – Académie tchèque des Sciences – Prague, République Tchèque) - *Rapporteure*

HEUILIN Thierry (Directeur de Recherche CNRS rovence UMR 7265 – BIAM)-*Examineur*
MOËNNE-LOCCOZ Yvan (Professeur – Université Claude Bernard Lyon 1) - *Examineur*

PETRUSEK Adam (Professeur – Université Charles de Prague) - *Examineur*
STORCH David (Professeur – Université Charles de Prague) - *Examineur*

MARECKOVA Marketa (Chercheure – Institute de recherche sur la production végétale - Prague, République Tchèque) - *Directeurice de thèse*

RODRIGUEZ-NAVA Veronica (Professeure des Universités – Université Claude Bernard Lyon 1) - *Directeurice de thèse*

KOPECKY Jan (Chercheure – Institute de recherche sur la production végétale -Prague, République Tchèque) - *Invité*

Joint PhD. thesis between

Charles University in Prague

Faculty of Science

and

Université Claude Bernard Lyon 1

École Doctorale Evolution, Ecosystèmes, Microbiologie et Modélisation



To obtain the degree of Doctor in Natural Sciences from both universities (Cotutelle)

Mgr. Ing. Andrea Faitová (Burešová)

Actinobacteria communities in natural and anthropogenic environments

Thesis supervisors:

RNDr. Markéta Ságová-Marečková PhD. and Prof. Veronica Rodriguez-Nava

Prague, December 2020

Declaration

I hereby declare that I have written this thesis independently, using the listed references; or in cooperation with other paper co-authors. I have submitted neither this thesis, nor any of its parts, to acquire any other academic degree.

Prague, 3rd December 2020

Andrea Faitová (Burešová)

Contents

Abstract	10
Abstrakt (In Czech)	12
Résumé (In French)	14
Aim and outline of the thesis	16
Introduction	
○ Microbes in a changing world	21
○ <i>Actinobacteria</i> in terrestrial ecosystems	23
○ Environmental factors driving <i>Actinobacteria</i> communities	24
○ Techniques for <i>in situ Actinobacteria</i> determination	26
Part 1: Cave environments and microbial inhabitants	28
○ Human derived changes in show caves	30
○ Problematic of Lascaux Cave	32
○ <i>Actinobacteria</i> roles in caves	33
Part 2: Decomposition of plant litter and the climate change	35
○ Microbial colonizers of litter	36
○ <i>Actinobacteria</i> as one of the main litter decomposers	38
Summary and conclusions	42
References	47

Original research papers 60

-Part 1-

Paper I: Usefulness of *hsp65* marker as a complement to 16S rRNA sequences to demonstrate differences in *Actinobacteria* community according to anthropization of limestone caves and occurrence of visual marks in Lascaux Cave

-Part 2-

Paper II: Succession of microbial decomposers is determined by litter type but sites conditions drive decomposition rate

Paper III: Litter chemical quality and bacterial community structure influenced decomposition in acidic forest soil

Paper IV: Litter traits and rainfall reduction alter microbial litter decomposers: the evidence from three Mediterranean forests

Appendix

The thesis is based on the following papers:

- I. **Buresova A.**, Alonso L., Moënne-Loccoz Y., Kopecky J., Sagova-Mareckova M., Vautrin F., Rodriguez-Nava V., (*in prep.*), Usefulness of *hsp65* marker as a complement to 16S rRNA sequences to demonstrate differences in Actinobacteria community according to anthropization of limestone caves and occurrence of visual marks in Lascaux Cave, Molecular Ecology
- II. **Buresova A.**, Kopecky J., Hrdinkova V., Kamenik Z., Omelka M. and Sagova-Mareckova M., (2019), Succession of Microbial Decomposers Is Determined by Litter Type, but Site Conditions Drive Decomposition Rates, Applied and Environmental Microbiology, 85 (24): e01760-19
- III. **Buresova A.**, Tejnecky V., Kopecky J., Drabek O., Madrova P., Rerichova N., Omelka M., Krizova P., Nemecek K., Parr T. B., Ohno T. and Sagova-Mareckova M., (*under revision*), Litter chemical quality and bacterial community structure determined decomposition in acidic forest soil, European Journal of Soil Biology
- IV. Pereira S., **Buresova A.**, Kopecky J., Madrova P., Aupic-Samain A., Fernandez C., Baldy V. and Sagova-Mareckova M., (2019), Litter traits and rainfall reduction alter microbial litter decomposers: the evidence from three Mediterranean forests, FEMS Microbiology Ecology, 95 (12): fiz168

Authors' contribution

- I. Veronica Rodriguez-Nava and Yvan Moëgne-Loccoz prepared the original paper concept. Lise Alonso and Yvan-Moëgne Loccoz did sampling in caves. Lisa Alonso made extraction of DNA from samples and 16S rRNA amplicon sequencing. Andrea Buresova and Florian Vautrin created the *hsp65* reference database. Andrea Buresova amplified *hsp65* gene (PCR) for amplicon sequencing processed and analyzed the 16S rRNA and *hsp65* sequence data and a statistical analysis with help of Jan Kopecky. Andrea Buresova did drafting and writing the manuscript. Final editing of the manuscript was done by Veronica Rodriguez-Nava, Marketa Sagova-Mareckova, Yvan Moëgne-Loccoz and Andrea Buresova.
- II. Marketa Sagova-Mareckova, Jan Kopecky prepared the original paper concept and together with Veronica Hrdinková participated at field sampling and conducted mass loss measurement. Andrea Buresova conducted DNA extraction, amplicon sequencing, quantitative real-time PCR analysis, extracellular enzyme activities, prepared samples for nutrients content analysis and made extracts for HPLC analysis. Zdeněk Kameník was responsible for HPLC analysis and HPLC data were processed by Jan Kopecky. Sequence data were processed by Andrea Buresova and Jan Kopecky. Marek Omelka performed statistical analysis. Andrea Buresova was responsible for manuscript writing and final editing was performed by Marketa Sagova-Mareckova, Jan Kopecky and Andrea Buresova.
- III. Vaclav Tejnecky and Marketa Sagova-Mareckova and Jan Kopecky prepared the original paper concept. Together with Petra Krizova, Karel Nemecek and Nada Rerichova they participated at field sampling. Chemical analysis did Vaclav Tejnecky, Nada Rerichova, Ondrej Drabek and Karel Nemecek. Thomas B. Parr and Tsutomu Ohno conducted the measurements of humification index and aromaticity of studied litter. Pavla Madrova and Andrea Buresova extracted DNA from litter. Andrea Buresova did a microbiological analysis (Illumina Sequencing, ddPCR of bacteria and fungi) and with Jan Kopecky processed sequence data. The statistical analysis was conducted by Marek Omelka. Manuscript writing performed Andrea

Buresova and together with Marketa Sagova-Mareckova and Vaclav Tejnecky did final editing.

- IV. Virginia Baldy, Catherine Fernandez and Marketa Sagova-Mareckova prepared the original paper concept and together with Susana Pereira participated at a field sampling. Susana Pereira, Andrea Buresova and Pavla Madrova extracted DNA and conducted microbiological analysis (qPCR, Illumina MiSeq). Jan Kopecky performed analysis of sequence data. Adrianne Aupic-Samain did the chemical analysis of the leaves samples. Susana Pereira was responsible for manuscript writing and final editing was performed by Marketa Sagova-Mareckova and Virginia Baldy

On behalf of all the co-authors, I declare the participation of Andrea Faitová (Burešová) at research and papers writing as described above.

Marketa Sagova-Mareckova

Abbreviations

AMOVA: Analysis of molecular variance

ddPCR: Droplet digital polymerase chain reaction

DNA: Deoxyribonucleic Acid

HOMOVA: Homogeneity of molecular variance

hsp65: 65-kDa Heat shock protein

PCR: Polymerase chain reaction

qPCR: Quantitative real-time polymerase chain reaction

rRNA: Ribosomal ribonucleic acid

Acknowledgements

I would like to express my biggest thanks to my Czech supervisor **Markéta Marečková**, for her valuable leading, support and advising during my studies. Without her support I would not have opportunities to gain such valuable research experiences from our Czech laboratory at Crop Research Institute and Czech University of Life Science but also from great research institutions from abroad. I would like to thanks my French supervisor **Veronica Rodriguez-Nava** for her enthusiasm and great advices during my stay in France as well as during the article writing and thesis preparation. I am also grateful for financial support from her institution to participate at FEMS 2019 conference with a poster presentation. Special thanks are also for **Jan Kopecký**, since many of the analysis and experiments I performed would not be done without his help and advising.

I am pleased to have an opportunity to cooperate on the Lascaux Cave project with **Yvan Moënné-Loccoz**, and I thank him for his valuable comments on the article. Similarly, I would like to thanks to **Václav Tejnecký** for his cooperation on decomposition experiments and helping me with improvement of our article.

It was a pleasure to share the office with a fantastic researcher **Daria Rapoport**, whose support motivated me many times in improving my research work and never give up ☺. I also thank to **Tereza Patrmánová**, **Adam Šťovíček** and **Pavla Mádrová**, for their help and for creating friendly environment at our workplace. I also thank to all members from the French laboratory Bacterial Opportunistic Pathogens and Environment for their patience and help during my work.

My special thanks belong to the **Josef Hlavka Foundation**, **Campus France** (French Government) and **Charles University** (Mobility Fond, ERASMUS) for financial supports enabled me to study the doctorate within the Cotutelle program in France and in Czech Republic, perform research stay at foreign institutions or participate at research conferences abroad.

My deep thanks belong to my family, which supported me whole the 23 years of my studies.

Abstract

Actinobacteria are important bacterial group participating in various ecosystem processes particularly in the decomposition of complex organic compounds. Their abilities enable them to surviving in harsh conditions of oligotrophic habitats like lakes, deserts, cave walls or recalcitrant and resistant litter in soil, where *Actinobacteria* often dominate. Although certain biotic and abiotic factors were recognized to modulate *Actinobacteria* incidence in such habitats, the influence of anthropogenic pressure on their communities is scarcely known. The main objective of this thesis is therefore to determine differences of *Actinobacteria* communities under the direct (the human visitors changing microenvironment of caves, **part 1**) and indirect (climate change factors like altered precipitation or plant litter quality, **part 2**) anthropogenic influence in two habitats, plant litter in soil and cave walls, where *Actinobacteria* play important roles and dominate.

In a first part of the thesis we monitored *Actinobacteria* communities in French limestone caves walls differently affected by humans (pristine versus anthropized caves). For identification of important species like potential pathogens or pigments producing *Actinobacteria* using amplicon sequencing of environmental DNA (Illumina MiSeq), we firstly used a molecular marker gene *hsp65* coding for heat shock protein specific for *Actinobacteria*. Special attention was payed to anthropogenically most affected Lascaux Cave with Paleolithic paintings. There, a comparison of different rooms differently affected by a human-derived intervention as well as between visual dark marks and unmarked areas on the wall paintings were compared (**paper I**). In the second part, we monitored litter *Actinobacteria* communities during a decomposition process under manipulated precipitation (**paper IV**), on different litter type, quality and origin (**papers II, III, IV**) in different forests including Mediterranean oak and pine forests (**paper IV**), mountainous spruce and beech forests (**paper III**), a beech temperate forest (**paper II**) and also one grassland (**paper II**).

Our results show that *Actinobacteria* communities were strongly dependent to anthropized/pristine status of caves (**Part 1**) as well as climatic and litter quality changes during the decomposition (**Part 2**). In caves (**Part 1**), *Actinobacteria* community structure indicated the anthropogenic disturbance, because typical pristine and anthropogenic taxa identified according to the *hsp65* marker were recovered in relation to an anthropization status (**paper I**). Moreover, the dominance of *Streptomyces* was found in the area with visual dark marks

suggesting the marked areas were factor influencing this group (**paper I**). During decomposition (**part 2**), we found that *Actinobacteria* were **i**) affected by litter type regardless its origin but their dominance on recalcitrant litter type did not result in faster decomposition (**papers II, III, IV**), **ii**) not directly affected by climatic conditions (**paper III, IV**) but were site-specific (**papers II, III, IV**) with a potential to dominate introduced coniferous forests (**papers III, IV**), and **iii**) in decomposition process had opposite strategies to fungi, since were influenced by different conditions than fungi (**papers II, III, IV**). Overall, *Actinobacteria* respond to anthropogenic pressure on a community and species level and are also able to adapt to harsh conditions and thus, these changes leading to *Actinobacteria* persistence in ecosystems. Consequently, *Actinobacteria* might be considered as stress-tolerant microbes, which may also benefit from man-made disturbances.

Key words: *Actinobacteria*, caves, decomposition of plant litter, climate change, anthropization, *hsp65* marker, fungi

Abstrakt (In Czech)

Actinobacteria jsou významnou bakteriální skupinou účastnící se různých ekosystémových procesů, zejména při rozkladu komplexních organických sloučenin. Jejich schopnosti jim umožňují přežít v drsných podmínkách oligotrofních habitatů, jako jsou jezera, pouště, stěny jeskyní nebo odolný a nerozložitelný rostlinný opad v půdě, kde aktinobakterie často dominují. Ačkoliv jsou známy určité biotické a abiotické faktory ovlivňující výskyt aktinobakterií v těchto habitatech, vliv antropogenního tlaku na jejich společenstvo je zřídka znám. Hlavním cílem této práce je tedy určit rozdíly v aktinobakteriálních společenstvech pod přímým (lidští návštěvníci měnící mikroenvironmentální prostředí jeskyň, **část 1**) a nepřímým (faktory klimatické změny jako jsou změny srážek nebo kvality rostlinného opadu, **část 2**) vlivem člověka ve dvou habitatech, rostlinném opadu v půdě a stěnách jeskyň, kde aktinobakterie hrají významnou roli a kde dominují.

V první části disertace jsme sledovali aktinobakteriální společenstva na zdech francouzských vápencových jeskyní odlišně ovlivněných člověkem (přírodní versus antropizované jeskyně). K identifikaci významných druhů (např. potenciálních patogenů nebo pigmenty produkujících aktinobakterií) pomocí amplikonového sekvenování environmentální DNA (Illumina MiSeq), jsme poprvé použili gen molekulárního markeru *hsp65* kódující protein teplotního šoku (heat shock protein) specifický pro aktinobakterie. Zvláštní pozornost byla věnována antropogenně nejvíce zasažené jeskyni Lascaux s Paleolitickými malbami. V této jeskyni byly porovnány různé místnosti odlišně ovlivněné zásahem člověka a také byly porovnány tmavá místa mikrobiálního původu nacházející se na nástěnných malbách (**článek I**). Ve druhé části jsme se zaměřili na společenstva aktinobakterií během rozkladného procesu pod vlivem změny srážek (**článek IV**), na různém typu opadu s odlišným původem a s různou kvalitou (**články II, III, IV**) v různých lesích včetně středozevního bukového a borového lesa (**článek IV**), horského smrkového a bukového lesa (**článek III**), bukového lesa mírného pásma (**článek II**) a také stepi (**článek II**).

Naše výsledky ukazují, že aktinobakteriální společenstvo bylo silně závislé na antropogenním/přírodním stavu jeskyň (**část 1**) stejně jako na klimatických změnách a změnách v kvalitě opadu během rozkladu (**část 2**). Struktura společenstva aktinobakterií v jeskyních (**část 1**) indikovala antropogenní narušení, protože typické přírodní a antropogenní skupiny byly identifikované (pomocí *hsp65* markeru) právě ve vztahu k antropogennímu/přírodnímu

stavu jeskyň (**článek I**). Navíc bylo zjištěno, že *Streptomyces* dominují v oblastech s tmavými místy což naznačuje, že tato tmavá místa jsou faktorem ovlivňujícím tuto skupinu aktinobakterií (**článek I**). Během rozkladného procesu (**část 2**) jsme zjistili, že aktinobakterie **i**) byly ovlivněny typem opadu nehledě na jeho původu, ale jejich dominance na nerozložitelném opadu nevedla k rychlejšímu rozkladu (**články II, III, IV**), **ii**) nebyly přímo ovlivněny klimatickými podmínkami (**články II, IV**), ale byly habitat-specifické (**články II, III, IV**) s potenciálem dominovat zavlečeným jehličnatým lesům (**články III, IV**), a **iii**) během rozkladného procesu měly opačné strategie než houby, jelikož byly ovlivněny jinými podmínkami prostředí (**články II, III, IV**). Aktinobakterie reagují na antropogenní tlak na úrovni společenstva i druhu a jsou také schopné adaptovat se na nehostinné podmínky, a proto tyto změny vedou k perzistenci aktinobakterií v ekosystémech. V důsledku toho mohou být aktinobakterie považovány za mikroorganismy tolerantní vůči stresu, které mohou mít výhodu ze změn způsobených člověkem.

Klíčová slova: *Actinobacteria*, jeskyně, rozklad rostlinného opadu, klimatická změna, antropogenní vliv, *hsp65* marker, houby

Résumé (In French)

Les actinobactéries sont un groupe bactérien important participant dans plusieurs processus écosystémiques comme la décomposition de composés organiques. Les actinobactéries possèdent des nombreuses capacités leur permettant de survivre dans des conditions difficiles comme celles retrouvées dans des habitats oligotrophes tels que les lacs, déserts, parois des grottes ou la litière végétale récalcitrante du sol où les actinobactéries dominent. Bien que certains facteurs biotiques et abiotiques ont été reconnus comme modulateurs de l'incidence des actinobactéries dans ces écosystèmes, l'influence des conditions changeantes provoquées par la pression anthropique sur leurs communautés reste inconnue. L'objectif principal de cette thèse est donc de déterminer des différences dans les communautés d'actinobactéries sous l'influence anthropique directe (visiteurs humains modifiant le microenvironnement des grottes, partie 1) et indirecte (facteurs liées à des changements climatiques : précipitations ou qualité de la litière végétale, partie 2) dans deux habitats : litière végétale du sol et parois des grottes, où les actinobactéries jouent un rôle important et prédominant.

Dans une première partie de la thèse, nous avons suivi les communautés d'actinobactéries dans des parois de grottes calcaires françaises affectées différemment par l'homme (grottes vierges ou anthropisées). Pour la détection d'espèces potentiellement pathogènes ou productrices de pigments à partir d'ADN environnemental (Illumina MiSeq), nous avons utilisé le marqueur moléculaire *hsp65* spécifique pour le genre *Actinobacteria*. Une attention particulière a été portée à la grotte de Lascaux, abritant des peintures paléolithiques et étant une des plus affectées d'un point de vue anthropique. Une comparaison au niveau de plusieurs salles affectées de façon différente par l'intervention humaine ainsi qu'entre zones des mêmes salles avec des marques visuelles d'origine microbienne sur les peintures murales a été faite dans cette grotte (**article I**). Dans la deuxième partie, nous avons surveillé les communautés d'actinobactéries de la litière végétale au cours d'un processus de décomposition sous conditions modifiées prévues (précipitation modifiée (**article IV**), différents types, qualités et origines de litière, (**articles II, III, IV**) dans différentes forêts (chêne méditerranéen et pin (**article IV**)), épicéa de montagne et hêtre (**article III**), forêt tempérée de hêtre (**article II**) et prairie (**article II**)).

Nos résultats montrent que les communautés d'actinobactéries étaient fortement dépendantes du statut anthropisé/vierge des grottes (**partie 1**) ainsi que des changements climatiques et de qualité des litières pendant la décomposition (**partie 2**). Dans les grottes (**partie 1**), la structure des communautés d'actinobactéries a indiqué des perturbations anthropiques, où des taxons typiques d'environnements vierges ou anthropiques comprenant des espèces potentiellement pathogènes (identifiées selon le marqueur *hsp65*) ont été trouvés ayant un rapport avec l'anthropisation (**article I**). De plus, la domination du genre *Streptomyces* a été trouvée dans les zones avec des marques visuelles suggérant que les zones marquées sont un facteur affectant ce groupe (**article I**). Lors de la décomposition (**partie 2**), nous avons constaté que les actinobactéries **i**) étaient affectées par le type de litière quel que soit son origine, mais leur dominance sur de la litière récalcitrante n'a pas entraîné une décomposition plus rapide (**articles II, III, IV**), **ii**) n'étaient pas directement affectés par les conditions climatiques (**article III, IV**) mais plutôt spécifiques au site (**articles II, III, IV**) avec du potentiel pour dominer dans des forêts de conifères introduites (**articles III, IV**), et **iii**) étaient influencées par des conditions différentes à celles des champignons (**articles II, III, IV**). Dans l'ensemble, les actinobactéries répondent principalement aux changements anthropiques au niveau de communautés ou d'espèces et sont capables de s'adapter à ces changements conduisant à leur persistance dans les écosystèmes. Les Actinobactéries peuvent être mentionnées comme des microorganismes tolérants au stress et, dans une certaine mesure, comme bénéficiant des perturbations d'origine humaine.

Mots clés: Actinobactéries, grottes, décomposition des litières végétales, changement climatique, anthropisation, marqueur *hsp65*, champignons

Aim and outline of the thesis

The thesis focuses on communities of terrestrial *Actinobacteria* under the anthropogenic influence in French caves and during plant litter decomposition in forests and grassland habitats in the Czech Republic, France and Austria. The first part describes the direct influence of humans on *Actinobacteria* inhabiting wall surfaces of caves in France under different human impact. The aim was to show how *Actinobacteria* react to human disturbance and coincide with the stability of these habitats. Such intervention as a biocide treatment but also the high frequency of visitors, which represent the alteration of microclimate due to their metabolism and introduction of allochthonous microbes and nutrients, represent direct anthropogenic factors affecting autochthonous microbial inhabitants. The second part focuses on the indirect consequences of global climate changes on communities of soil and litter *Actinobacteria*. The aim was to show how forest and grassland *Actinobacteria*, as an important part of soil decomposer community, will respond to changes in litter quality and habitat properties but also to the reduction of rainfall as expected from the climate change. For more complex insight, the other important microbial decomposers within the community were also analyzed to see possible relationships typical for the respective environmental and substrate conditions.

The overall aim of the thesis was to uncover the ecology of *Actinobacteria* in habitats modified by human activities directly or indirectly. Using a combination of environments, where *Actinobacteria* play the dominant role we were able to determine the sensitivity of this group to anthropogenic changes in open and less stable (plant litter) versus isolated and relatively more stable (caves) habitats. Using the novel marker gene *hsp65* for analysis of *Actinobacteria* communities from eDNA, we were able to overcome the problem of unculturable strains in identifying the cave *Actinobacteria* to the species level in French caves, Lascaux, Rouffignac, Mouflon and Reille. That helped us to identify in detail the part of the community that was affected by human interventions or by the presence of visual dark marks (black stains and dark areas on cave walls) of microbial origin in the Lascaux Cave (**part 1**). In the part two, where we quantitatively and qualitatively measured the reaction of *Actinobacteria* to other decomposers and environmental factors, we firstly showed the abiotic but also biotic ones driving *Actinobacteria* activities during decomposition (**part 2**). The work further demonstrates the necessity to include *Actinobacteria* into environmental monitoring and prediction models as a one of the indicative microbial groups.

We focus on the following subjects:

- 1) Detailed analysis of *Actinobacteria* communities of four French caves and also individual rooms within the Lascaux Cave. The questions were: 1) Are *Actinobacteria* in caves shaped according to the anthropization status (presence of visitors, application of chemicals)? 2) Can we identify with a novel marker (*hsp65*) *Actinobacteria* species related specifically to anthropized/pristine habitats, for example potentially pathogenic and novel species?
- 2) Comparison of *Actinobacteria* between dark visual marks and unmarked areas on wall paintings within the Lascaux Cave. The questions were: Are visual marks in the highly anthropized Lascaux Cave a factor affecting *Actinobacteria* communities?
- 3) Quantitative and qualitative analysis of changes in *Actinobacteria* communities in relation to altered litter type in different forests and grassland. The questions were: Are *Actinobacteria* affected by changes in precipitation, litter quality and origin at the studied sites? Are *Actinobacteria* able to adapt to altered litter quality regardless of the environment or time of decomposition?
- 4) Determination of *Actinobacteria* successional patterns over the decomposition process and comparison with the other important microbial decomposers, especially fungi. The questions were: Are *Actinobacteria* initial, middle or late-stage decomposers? Do they have similar or opposite colonization patterns to other decomposers? Could be these patterns affected with changes in litter and environmental conditions?

After a short introduction which is divided into two sections (**part 1, part 2**), the thesis is concluded by general summary and conclusions of the obtained results. The main part of the thesis is also divided into two sections (**part 1, part 2**) and consists of four original research papers. In the appendix, my curriculum vitae containing all my publications and trainings can be found.

Outline of the original papers

This thesis consists of four papers (I-IV). The **paper I** focused on communities of *Actinobacteria* in French caves differently affected by anthropization (visited versus non-visited caves) with closer focus on Lascaux cave, which was disturbed to a high extent and thus already closed to visitors. Three papers (II-IV) focused on microbial communities during litter decomposition with closer focus on *Actinobacteria*. Within that, **paper II** describes one-year decomposition experiment with litters of different origin and quality at beech forest and grassland sites, while the **paper III** is focused on later stage decomposition (15 to 29 months of decomposition) of two highly recalcitrant litters in beech (native) and spruce (introduced) mountainous forests. The **paper IV** describes recalcitrant oak (native) and pine (introduced) litter decomposition under the altered precipitation conditions in Mediterranean forests.

Paper I (Buresova et al., in prep, Molecular Ecology) focused on *Actinobacteria* communities colonizing walls in French caves differently affected by humans (Lascaux, Rouffignac – anthropized, Mouflon, Reille – pristine). In addition, in Lascaux cave, samples of different rooms (entrance Sas-1, Passage banks, Passage inclined planes, Apse, Diaclase) were picked wall from visual marks (black stains and dark area) and unmarked area. DNA was extracted and amplicons of heat shock protein gene *hsp65* and *16S rRNA* gene fragments were sequenced (Illumina MiSeq). The taxonomical database for assignment of *hsp65* amplicons was designed. Comparison of 16S rRNA and *hsp65* markers showed higher variability of *hsp65* than 16S rRNA gene enabling higher resolution of *Actinobacteria* at species level especially for *Nocardia*, *Mycobacteria*, *Streptomyces* and others. *Actinobacteria* communities differed between Lascaux, Rouffignac and both pristine caves. The results showed that anthropization level affected *Actinobacteria* community structure where high diversity of *Nocardia* was typical for visited Rouffignac and *Mycobacteria* for already closed Lascaux cave. Both caves contained potentially pathogenic species which were identified thanks to the *hsp65* reference database. Pristine caves harbored high proportions of typical pristine caves dwellers *Gaiellales* similarly to Diaclase, which is considered as pristine part within the anthropized Lascaux cave. Concerning the visual marks in Lascaux cave, *Actinobacteria* were shaped according to these micro-habitats and domination of *Streptomyces* were the predominating group in the visual marks.

Paper II (Buresova et al. 2019, Applied and Environmental Microbiology) described *Actinobacteria* communities over the decomposition process of autochthonous and allochthonous litter at beech forest and grassland sites. Over the transplant litterbag decomposition experiment the mass loss of litter (*Fagus sylvatica* - lowest quality defined by C, N, P, autochthonous in beech forest and allochthonous in grassland; *Carex humilis* and *Astragalus exscapus* - middle and highest litter quality, respectively, autochthonous in grassland and allochthonous in beech forest), C, N and P changes, and hydrolytic and oxidative extracellular enzymes activities were measured. Bacterial community structure was determined by 16S rRNA gene amplicon sequencing (Illumina MiSeq), and quantities of fungi (as 18S rRNA gene copy numbers), *Actinobacteria* (16S rRNA gene), *Alphaproteobacteria* (16S rRNA gene) and *Firmicutes* (16S rRNA gene) were measured using quantitative real-time PCR. Extracellular enzyme activities were higher at forest site, where also decomposition was faster, but where *Actinobacteria* 16S rRNA genes were less abundant. *Actinobacteria*, contrary to fungi, preferred the litter regardless its quality and origin. The fastest decomposed litter at both sites was the most nutritious one, which however was not the preferred one for neither fungi nor *Actinobacteria*. Dynamics (i.e. variation over time) of *Actinobacteria* and fungi was highly litter specific but not site specific. *Actinobacteria* 16S rRNA gene abundance increased in later stages, similarly as for *Firmicutes*, and differing from fungi and *Alphaproteobacteria*. *Actinobacteria* and fungi were differently influenced by litter or site conditions, but had similar strategies in colonizing the litter over time. *Actinobacteria* should be more and fungi less affected by changes in complex site conditions, because *Actinobacteria* were site specialists and fungi site generalists. But *Actinobacteria* adapted to decompose the litter regardless its quality, and may be potentially less affected by changes in vegetation cover and resulting litter quality.

Paper III (Buresova et al., under revision, European Journal of Soil Biology) built on our previous results where *Actinobacteria* increased over the decomposition. This paper focused on late stage decomposition of recalcitrant beech (native) and spruce (introduced) litter under the mountainous climate. All abiotic conditions like a parent rock and climate were identical at for both sites since the forests were located next to each other on the hillside. A litterbag experiment was set and a later stage was determined to months 15, 19 and 29. Bacterial community structure (16S rRNA gene, Illumina Miseq), fungal and bacterial abundance (ITS region and 16S rRNA

gene, respectively, ddPCR), micronutrients, macronutrients and characteristics of organic matter were measured over that period in the litter. Nutrients were also measured in soil profiles. Our results showed that fungal and bacterial abundance did not change between the sites with different litter and was therefore affected by other than litter characteristics, for example site conditions. Contrary to *Proteobacteria*, *Actinobacteria* dominated in late stages on the more recalcitrant introduced spruce litter which however did not lead to a higher decomposition rate.

Paper IV (Pereira et al 2019, FEMS Microbiology Ecology) focused on *Actinobacteria* community changes under the reduction of rainfall during decomposition of allochthonous and autochthonous litter in Mediterranean two deciduous and one coniferous forests. The results show that, contrary to fungi, *Actinobacteria* abundances (16S rRNA gene, qPCR) were not affected by reduced precipitation. *Actinobacteria* dominated at each site in deciduous litter, while coniferous litter was colonized by fungi with higher preference, especially in deciduous forests (18S rRNA gene, qPCR). In coniferous forest, both microbial groups dominated on deciduous litter with lower lignin content and the highest availability of nutrients. It suggests that in coniferous forest both groups had similar strategies and colonized the more nutritious deciduous litter. *Actinobacteria* were not affected by rainfall reduction but were affected by litter quality.

Introduction

Microbes in a changing world

Human derived changes on earth resulted in a new geological epoch called the Anthropocene, which consists of three stages called paleoanthropocene, industrial revolution and great acceleration (Gillings and Paulsen, 2014). During those stages the widespread of agriculture and industry together with the high population growth, consumption of energy and resources represented a new selection pressure for the natural ecosystems. Consequently, the loss of biodiversity, elevated production of greenhouse gases and chemical pollution are some of the main outcomes challenging the stability of the nature nowadays (Gillings and Paulsen, 2014; Zhu and Penuelas, 2020).

The human impact influenced microbial communities across many biomes and might be reflected in various measures, such as stability, diversity and functioning of microbial communities (Zhu and Penuelas, 2020). Such human impacts might have direct and indirect influences. For example, the pollution of environment by toxic substances (Abdu et al., 2017) and eutrophication (Schobben et al., 2016) can directly negatively affect microbes. On the contrary, the changes in vegetation cover (Prieto et al., 2019), precipitation events and temperature (Suseela et al., 2014) due to a climate changes like increase CO₂ production are examples how human affect microbes indirectly.

The alteration of temperatures and precipitation regimes represented one of the predicted effects of climate change with direct consequences to microbes. It was suggested that the warming and higher frequencies of freeze/thaw cycles might promote animal and plant pathogens and suppress decomposers of organic carbon compounds as well as those involved in N-cycling (Garcia et al., 2020). Another finding revealed that thawing of permafrost might unleash danger pathogens and promote carbon release from soil, which has in turn a negative feedback for the climate change (Gross, 2019). The promotion of long drought with more intense rain events were expected also in arid habitats and might promote further soil degradation together with decrease of microbial diversity (Maestre et al., 2012; Meisner et al., 2018).

On the other hand, plant diversity and biomass alteration changes are another indirect consequences of climate change which can modulate the activity, biomass and structure of microbial communities (Lange et al., 2015). The plant diversity decreased and proportions of structural components of plant tissues might slow down the decomposition. The climate change scenarios through warming and drought might affect the nutrient content in plant tissues and thus palatability for microbes (Rosenblatt, 2018). These and other climate-related changes might in turn result in settlement of stress tolerance communities (Garcia et al., 2020).

Human activity might directly affect microbial communities by organic and inorganic pollution by biocides, heavy metals but also excess of nutrients. The high concentration of heavy metals (Lee et al., 2002; Száková et al., 2016) or pesticides (Pose-Juan et al., 2017) in soil might decrease of microbial biomass together with enzymatic activities as a reaction to toxic stress. However, although some microbes might be sensitive to heavy metals, others are resistant or adapt over time (Chen et al., 2020). Simultaneously, pesticide amendments might shift microbial community towards high proportions of microbes capable to degrade the respective organic pollutant (Bragança et al., 2019). Also, the excess of nutrients reached by fertilization was associated with shifts in microbial diversity as a reaction to eutrophication. In soil, nitrogen fertilization might shift the microbial community towards dominance of copiotrophic microbial taxa like *Firmicutes*, *Proteobacteria* and *Zygomycota* contrary to the oligotrophic like *Acidobacteria* (Francioli et al., 2016). Moreover, high nutrient amendments might promote mineralization of stable and recalcitrant organic matter, which is known as a “priming effect”, (Meier and Bowman, 2008) and thus promote emission of CO₂ and CH₄ to the atmosphere.

Some studies focused only on diversity indices of bacterial or fungal community or overall microbial biomass (Száková et al., 2016; Wagg et al., 2019), which however could not be a sufficient indicator of which changes we can expect in relation to anthropogenic pressure. We apply as a fact, that more diverse community maintain more functions (Delgado-Baquerizo et al., 2016) but since microbes are not functionally equivalent (Bray et al., 2012; Wertz et al., 2006) the crucial question should be, if such diverse community contain specific taxa responsible for respective modulations in ecosystems. Moreover in this context arise a question what are the main microbial groups to which these changes are significant (McGuire and Treseder, 2010). At the same time, does these groups maintain key roles in ecosystems? If so, which feedbacks can be expected through modulation of their abundances, proportions and

interactions in community and activity? Then, we have to consider such key groups jointly with other community members, since with the interactions in the community we can completely gain the biotic and abiotic factors affecting their functioning (Bani et al., 2018).

Actinobacteria in terrestrial ecosystems

Actinobacteria are phylogenetically diverse and ubiquitous gram positive class of bacteria further characterized by high content of cytosine and guanine in their genomic DNA (Barka et al., 2016). *Actinobacteria* sustain important ecosystem functions since among other activities they participate in carbon cycling and nutrient transformation. They possess useful enzymes and also produce metabolites of various functions, for example antibiotics and thus have broad applications in biotechnology and biomedicine (Sharma et al., 2014). Since we face the problems of antibiotic resistance and new diseases, which call for exploration of novel searching for unknown *Actinobacteria* especially from underexplored habitats with capacity to produce new metabolites are one of the main goals nowadays (Rangseekaew and Pathom-Aree, 2019). Due to *Actinobacteria* high application potential many studies focus on *Actinobacteria* production of therapeutics, antifungal substances, enzymes or degradation of pollutants but much fewer are concerned about their ecological demands affecting their distribution and functioning in natural habitats.

High frequencies of *Actinobacteria* were monitored in various types of terrestrial, aquatic ecosystems and in the atmosphere. In terrestrial ecosystems, free living *Actinobacteria* are highly abundant especially in soil, where they count 10^6 to 10^9 cells per gram (Barka et al., 2016). Since *Actinobacteria* have broad adaptations to live in oligotrophic conditions *Actinobacteria* live epilithically and endolithically on building surfaces, rocks, stone monuments and statues (McNamara et al., 2006; Mihajlovski et al., 2017). Their advantages include the capability to produce resistant spores as a reaction to harsh conditions (B.-Z. Fang et al., 2017; Makhalanyane et al., 2015). Moreover, *Actinobacteria* produce pigments like melanin to be protected against high radiation, temperature extremes and free radicals (Manivasagan et al., 2013). That enables their survival in extreme habitats such as deserts and semideserts (Makhalanyane et al., 2015), alkaline saline soils (Shi et al., 2019), polluted soils (Alvarez et al., 2017), permafrost soils (Aszalós et al., 2020; Hansen et al., 2007), subterranean habitats (De Mandal et al., 2017; Hershey et al., 2018).

Besides that, *Actinobacteria* are important symbionts but also pathogens of plants and humans. In plants they can for example caused potatoes common scabs for which the *Streptomyces* genera are responsible (Marketa Sagova-Mareckova et al., 2015). Within the *Clavibacter* genera are plant pathogenic species like *C. michiganensis subsp. sepedonicus* causing bacterial root rot of potatoe or *C. michiganensis subsp. michiganensis* causing bacterial wilt and canker of tomato (Bentley et al., 2008). For humans, many diseases including tuberculosis, cystic fibrosis, pulmonary disease, nocardiosis and others caused *Actinobacteria* genera like for example *Nocardia* (*N. pneumoniae*, *N. farcinica*, *N. asteroides*) (Boiron et al., 1992; Taj-Aldeen et al., 2013) and *Mycobacterium* (*M. tuberculosis*, *M. abscessus*, *M. avium*) (Gagneux, 2018; To et al., 2020). *Rhodococcus equi* is an example of animal pathogen, namely of horses, ruminants and pigs, that might causes opportunistic infections for humans (Vázquez-Boland and Meijer, 2019).

Environmental factors driving *Actinobacteria* communities

Actinobacteria, their diversity, activity and abundance are highly dependent on pH, moisture and temperature. The pH was considered to be the main factor affecting *Actinobacteria* community. In the study performed in different soils of temperate forest and meadow *Actinobacteria* abundance negatively correlated to pH (Sagova-Mareckova et al., 2011). Some *Actinobacteria* were considered acidophilic including the new species *Trebonia kvetii* isolated from highly acidic soil (Rapoport et al., 2020). Similarly, acidic soil pH favorited some *Actinobacteria* genera for example *Acidimicrobium*, but alkaline soil pH favorited genera such as *Nocardia*, *Micrococcus* and *Mycobacterium* (Jenkins et al., 2009). In addition, another study based on *Actinobacteria* from rock surfaces suggested they most grew on media with neutral pH (B.-Z. Fang et al., 2017).

Aquatic *Actinobacteria* were found in free water and sediments of saline and fresh water habitats where pH (Newton et al., 2007), UV-radiation (Rasuk et al., 2017), nutrients (Holmfeldt et al., 2009) and various other factors affected their community structure (Kurtbóke, 2017; Sharma et al., 2014). Therefore, *Actinobacteria* have also adaptation to live in aquatic ecosystems, but these are not the subject of the thesis. Terrestrial *Actinobacteria* prefer humid habitats but well oxidized since they are generally considered as aerobes or microaerophiles since their counts in waterlogged habitats declined to about 10^2 or 10^3 per dry gram of soil

(Kurtbóke, 2017). Yet, hydration of arid soil after the dry period was linked to *Actinobacteria* abundance decline with subsequent increase after the habitat recovery as the oxygen availability increased (Št'oviček et al., 2017). Similarly in the Mediterranean forest, *Actinobacteria* abundance were not influenced by the rainfall exclusion (Pereira et al., 2019). That suggested terrestrial *Actinobacteria* were well adapted to dry conditions because their proportions significantly increased with dryness but not only in warm but also in cold climates including Arctic soil (Rego et al., 2019). Yet, some *Actinobacteria* dominated waterlogged habitats (Kopecky et al., 2011).

Optimal growth temperature for *Actinobacteria* is between 25 – 30°C however, thermophilic and psychrophilic *Actinobacteria* also exist (Barka et al., 2016; Li et al., 2010). The temperatures might have determining effect on the size and composition of *Actinobacteria* communities not only directly but also indirectly through the litter quality and substrate availability (Jenkins et al., 2009; Kurtbóke, 2017). *Actinobacteria* seasonal changes between winter and summer showed, that higher diversity with higher proportions of unknown taxa were found in winter (M. Sagova-Mareckova et al., 2015). Similar results were found in deciduous forest, where the known *Actinobacteria* taxa were higher in winter in upper horizons where the highest amount of organic matter was located (Kopecky et al., 2011). For example, in the grassland and forest sites *Actinobacteria* proportions increased especially in July and then in January more visibly in grassland, however it again seemed to be dependent on nutrients availability (Buresova et al., 2019). *Actinobacteria* might dominate high nutritious habitat (Francioli et al., 2016), but as was mentioned earlier, their competitive advantage represented also the capability to overcome oligotrophic conditions and to utilize recalcitrant substrates (Barton and Jurado, 2007).

Although it is certain that *Actinobacteria* communities are driven by environmental forces, different factors affect *Actinobacteria* and a single resolute pattern cannot be recognized. Above that *Actinobacteria* participated in many biotic interactions so might be suppressed by fungal pathogen or facilitated by the assembly of microbes or plants, for which *Actinobacteria* mediated available sources of energy by degradation of resistant compounds (Radha et al., 2017). Cooperation or competition of *Actinobacteria* with other microbial community members might be mediated by secondary metabolites production, however in nature has been scarcely investigated (Miao and Davies, 2010).

Techniques for *in situ* *Actinobacteria* determination

Culture dependent analyses of microbial diversity represented a big problem in the past since only the cultivable bacteria were recovered (Penny R. Hirsch et al., 2010). The cultivation approach is limited not only due to missing knowledge on requirements of the microbes for growth conditions and nutrients but also for the absence of cooperating microbes or their metabolites, which allow growth of the other ones (Lewis, 2009). Moreover, these techniques are time-consuming contrary to culture independent methods (Penny R. Hirsch et al., 2010).

With the expansion of *in situ* molecular techniques, the ecological role and community structure of microorganisms including *Actinobacteria* can be done *in situ* (Barka et al., 2016). The gene coding the small ribosomal subunit (16S rRNA gene) is the most frequently used molecule for description of prokaryotic microbial communities and molecular taxonomy. Currently, the next generation sequencing is the preferred sequencing method but even its high throughput capacities generally cannot resolve individual sequences at the species level (Fox et al., 1992; Penny R Hirsch et al., 2010). Moreover, this gene cannot be used for issuing abundance because for example in *Actinobacteria* there are on average 3.1 ± 1.7 16S rRNA copies per genome depending on the particular genus/species, while in *Firmicutes* it is on average 5.8 ± 2.8 copies and in *Proteobacteria* from 2.2 ± 1.3 copies in *Alphaproteobacteria* to 5.8 ± 2.8 copies in *Gammaproteobacteria* (Větrovský and Baldrian, 2013). Above that a single bacterium can harbor diverse copies of the gene, while identical gene copies can be found in different bacteria, which complicates the molecular taxonomical identification especially to the species level (Pei et al., 2010).

Alternatively, highly conserved protein coding genes from primary metabolism were proposed for identification of closely related bacteria including *Actinobacteria*. Generally, genes coding *rpoB* (beta subunit of RNA polymerase), *gyrA*, *gyrB* (A, B subunit of DNA gyrase), *secA* (ATPase secretory preprotein translocase), *atpD* (beta subunit of ATP synthase F1), *gap* (glyceraldehyde-3-phosphate dehydrogenase), *pnp* (polyribonucleotide nucleotidyltransferase), *ftsZ* (tubulin-like GTP-binding protein) or *dnaK* (chaperone Hsp70 in DNA biosynthesis), as well as the 23S rRNA gene are used (Gtari et al., 2015; Martens et al., 2008). *rpoB* coding beta subunit of DNA polymerase exists in a single copy, but it is not conserved enough to be amplified by universal primers (Ogier et al., 2019; Vos et al., 2012). For *Actinobacteria*, *sodA*, *hsp65*, the *recA* gene encoding DNA recombination and repair

systems were used (Adékambi and Drancourt, 2004). DNA gyrase subunit B encoding gene *gyrB* was used for differentiation of *Nocardia* species, however only some species were more distinct according to *gyrB* than 16S rRNA (Takeda et al., 2010). Simultaneously, *hsp65* gene coding heat shock protein is present in genome in a single copy and has been used as a molecular marker for identifying a large number of *Actinobacteria* isolates (Rodríguez-Nava et al., 2007, 2006), but has never been sequenced from *in situ* samples (from environmental DNA). Although such markers might employed drawbacks like lacking databases for assigning the taxonomy, they might be used in combination with other housekeeping genes in Multi Locus Sequence Analysis (MLSA) studies (Diancourt et al., 2005; Ogier et al., 2019). Usually five or more genes were sequenced for definition species within the genus as for example for *Frankia alni* gene markers *atpI*, *ftsZ*, *dnaK*, *gyrA* and *secA* were used (Nouioui et al., 2019). For genetic relatedness between *Ensifer* species (*Alphaproteobacteria*) ten genes encoding *atpD*, *dnaK*, *gap*, *glnA*, *gltA*, *gyrB*, *pnp*, *recA*, *rpoB* and *thrC* were used (Martens et al., 2008). However, recently is more reliable method for species identification the whole genome sequencing (Verma et al., 2013).

Part 1: Cave environments and microbial inhabitants

Caves are underexplored habitats unique in its stable microclimatic conditions and isolation from the surrounding environment. The isolation from the sunlight results in the absence of phototrophic primary production (Tomczyk-Żak and Zielenkiewicz, 2015). The chemoheterotrophic or chemolithotrophic microbes living in caves are under the pressure due to nutrient deficit, to which they are adapted by alternative nutrient acquisition strategies (Barton and Jurado, 2007; Ortiz et al., 2014). This selective pressure consequently resulted in unique microbial communities (De Mandal et al., 2017). Microbes might face harsh conditions, which might cause high DNA damage due to high calcium content, which resulted in overabundance of DNA-repair enzymes (Ortiz et al., 2014). Cooperative microbiological interactions predominate over competition in such habitats, since the limited inorganic and complex organic nutrients are unlikely to be acquired by a single organism due to high energy demand versus little energy gain in those processes. Those living conditions consequently explain the high microbial diversity (Barton and Jurado, 2007), and low microbial biomass ($<10^6$ cells/g) observed in caves (Hershey and Barton, 2018).

In caves, the microbial communities are driven by the type of habitat like the rock wall, sediment, water, air (Wu et al., 2015). It was even suggested that the sample type is a stronger determinant of the variability of cave microbial communities than the differences between caves (Zhu et al., 2019). The most ubiquitous bacterial phyla in caves were found *Proteobacteria* and *Actinobacteria* (Tomczyk-Żak and Zielenkiewicz, 2015) while other highly abundant groups were *Firmicutes*, *Verrucomicrobia* or *Acidobacteria* (De Mandal et al., 2017; Zhu et al., 2019). The presence of Archaea were frequently found in caves, such as *Thaumarchaeota*, most likely thanks to their various CO₂ fixation pathways (Ortiz et al., 2014). Fungi in caves were found to be parasites of animals or decomposers especially of organics, which entered the cave externally. In that case, the organics were covered by conidia of such fungi like *Aspergillus*, *Penicillium* or *Mucor*, which were at the same time generally the most frequently reported genera from caves together with *Histoplasma* or *Geomyces* (Vanderwolf et al., 2013).

Some of the most ubiquitous cave taxa could be also found in soil, which is explained by the seeding hypothesis suggesting that these microbes were carried by the percolating water into a newly created cave. However, the long-time impact and selective pressure sorted for

microbes with unique adaptations (Hershey and Barton, 2018) and precisely those adaptations lead to the high proportions of unknown and endemic microbes observed in caves (Lavoie et al., 2017). Those included for example *Actinobacteria* species *Hoyosella altamirensis* and *Nocardia altamirensis*, which were firstly isolated from Altamira cave wall (Jurado et al., 2009, 2008) and *Nocardia jejuensis* from a Korean cave (Lee, 2006). More recently, *Nonomurea cavernae* and *Bacillus antri* were examples of bacteria isolated from a karst cave of China (B. Z. Fang et al., 2017; Rao et al., 2019). Also, novel fungal species were isolated from caves like three highly oligotrophic fungal species *Cephalotrichum guizhouense*, *Cephalotrichum laeve* and *Cephalotrichum oligotrophicum* (Jiang et al., 2017) or filamentous fungi with high cellulolytic activity (Paula et al., 2019).

Some cave microbes represent a source of novel secondary metabolites used as therapeutics. *Cervimyxin*, *Xiakemycin*, *Hypogeamicin* and *Huanglongmycin* are examples of new antibiotic and anticancer metabolites produced by *Actinobacteria* (Rangseekaew and Pathom-Aree, 2019). Interestingly, some caves are also reservoirs of antibiotic resistance or pathogenic microbes. Their presence was in the past attributed to the human activity, which later became controversial since they were found also in natural and isolated caves without any human intervention (Bhullar et al., 2012; Jurado et al., 2010).

The microbial inhabitants of caves are driven by biotic and abiotic conditions and in turn modify the inner environment since some act as constructive or destructive agents. More specifically, microbial mineral precipitation in caves resulted in the formation of speleothems, calcium carbonate, iron precipitation and others are driven by number of processes (Banerjee and Joshi, 2013; Tomczyk-Żak and Zielenkiewicz, 2015). In the Altamira cave, the mineral deposits were found to be associated with microbial colonies, while the biomineralization was driven by microenvironmental conditions like high humidity, temperature and CO₂ production (Cuezva et al., 2009). Also, the pH of biofilm strongly influenced calcium carbonate dissolution and precipitation. The pH increase was associated with precipitation and thus carbonate formation, while pH decrease caused carbonate dissolution (Banerjee and Joshi, 2013).

Some studies suggest *Actinobacteria* and fungi to be one of the crucial microbes inducing mineral deposition (Tomczyk-Żak and Zielenkiewicz, 2015; Vanderwolf et al., 2013). However, although *Actinobacteria* and *Proteobacteria* dominated speleothems of Karstic caves, metabolically dominating groups associated with calcium carbonate precipitation were

Proteobacteria and *Firmicutes* (Dhami et al., 2018). Another study pointed on the most important role in mineral deposition of those cyanobacteria, algae and fungi able to create biofilms and mats (Banerjee and Joshi, 2013).

Microbes might colonize cave surfaces and create a biofilm, which is made of microbial consortia held together by polymers leading to their protection and effective use of energy sources (Mulec et al., 2015). The biofilms may be colorless but also highly pigmented, which might indicate its functional composition and reflect the microenvironmental factors driving its creation (Banerjee and Joshi, 2013). The green biofilm consists of green algae and cyanobacteria in the areas where light, artificial or natural light in the entrance of cave are present (Urzi et al., 2010). The yellow biofilm in lava tubes was linked to percolation of organic matter originated from lignin degradation containing yellow colored phenolic compounds. In this type of biofilm, *Proteobacteria* followed by *Actinobacteria* dominated, however, the latter group was metabolically more active (Gonzalez-Pimentel et al., 2018). Black and white pigmented biofilms were found to have alkaline pH and thus, are contrary to yellow biofilms responsible for speleothems induced by precipitation (Portillo and Gonzalez, 2010). Dark color of biofilms might be caused by algae producing black pigments like *Nostoc* and melanized fungi like *Aspergillus niger*, *Acremonium* or *Ochroconis* (Alonso et al., 2018; Popović et al., 2015). Although these dark pigments like melanin might have protective functions for example against free radicals their roles in cave biofilms are not completely clear (Agostinho and Nosanchuk, 2017; Mulec et al., 2015).

Human derived changes in show caves

Pristine caves are fragile habitats, where human derived intervention might cause disruption of structure and functioning of microbial communities. However, microbes are to some extent resilient to the influence from human activity and are able to mitigate the impact (Tomczyk-Żak and Zielenkiewicz, 2015). That was demonstrated for example by the effect of human urine, which represented enrichment of nitrogen as well as introduction of commensal organisms (Johnston et al., 2012). As a result, the indigenous cave microbiome including the endemic species took an advantage of nitrogen and organic compounds enrichment without alteration of their community composition (Johnston et al., 2012). Another study suggested,

that microbial community composition remained unchanged, but the enrichment by exogenous nutrients in show caves increased microbial biomass (Adetutu et al., 2012).

Unfortunately, the situation was changing rapidly once the cave became open to public suggesting that the level of anthropization might be the crucial determinant of the cave microbiome over other environmental factors (Moldovan et al., 2020). Extensive physical human impact can also began with installation of walkways, lights and other utilities and more drastically continued with high frequency of visitors (Bercea et al., 2019). The visitors directly changed the inner microclimate by their metabolism, including production of CO₂, H₂O, and heat and by consumption of O₂ (Hoyos et al., 1998). Moreover, each visitor brought to a cave not only external organic and inorganic matter on their shoes or clothes but also exogenous microbes like human or animal-associated commensals or pathogens, representing new potential colonisators and thus, aliens for true microbial cave dwellers (Leuko et al., 2017). At the end, the remediation of human impact by physico-chemical and biological treatments might even more drastically affect the cave microbiome in a negative way (Martin-Sanchez et al., 2015).

Some cave microbes might react to the human induced changes by coloration, creation of microbial mats and by deterioration of rock walls and historical paintings induced by pH decrease, production of acids and enzymes (Banerjee and Joshi, 2013; Carlo et al., 2016). Indeed, the problem with wall paintings endanger by bacteria and fungi try to deal with such world renowned caves like Altamira, Roman tombs, Lascaux and others (Martin-Sanchez et al., 2015; Schabereiter-Gurtner et al., 2002). The secondary calcium deposits of white color (moonmilk) endangered the Etruscan wall paintings in Italian tomb, where *Proteobacteria*, *Acidobacteria* and *Actinobacteria* dominated (Cirigliano et al., 2018). It was a similar case for moonmilk deposits on wall paintings of Altamira cave, where *Proteobacteria* and *Actinobacteria* dominated (Portillo and Gonzalez, 2011). The paintings of ancient Egyptian tombs suffered from color changes performed by *Actinobacteria* species, namely *Streptomyces*, by oxalic and citric acids, pigments like melanin, karotenoids and hydrogen sulphide production (Abdel-Haliem et al., 2013). On the contrary to that study, the melanin in black stains on wall Paleolithic paintings of Lascaux Cave were found to originated from fungi, namely from species from *Ochroconis* genera (De La Rosa et al., 2017).

The human contamination can also result in the presence of bacterial, fungal and viral opportunistic pathogens, which can affect the health especially of immunosuppressed visitors or those with lower defense (Carlo et al., 2016; Jurado et al., 2010). The anthropogenically contaminated species such as *Escherichia* and *Lysinibacillus* might enter the cave with heavy rains (Davis et al., 2020), others can enter the cave with visitors (Bercea et al., 2019) or with animals (Jurado et al., 2010). Bacterial opportunistic pathogens caused various skin, brain and/or lung infections including for example those from groups *Actinobacteria* (*Nocardia*, *Mycobacterium*, *Rhodococcus*, *Kribella*), *Alphaproteobacteria* (*Aurantimonas*) and *Spirochaete* (Jurado et al., 2010). Known human pathogens *Staphylococcus aureus* and *Bacillus spp.* were detected in one of the largest show cave in Japan (Ando and Murakami, 2020). In Chinese cave, fungi related species associated with human and animal mycotoxicosis was isolated from dead bats (Karunarathna et al., 2020). Also Vanderwolf et al., (2013) mentioned in his review some cave pathogenic fungi like *Histoplasma capsulatum* and *Geomyces destructans*. *Histoplasma capsulatum* produce pulmonary histoplasmosis which was relatively frequent in caves explorers while the species from a genus *Geomyces* cause skin and nails human diseases (Jurado et al., 2010; Nováková, 2009).

Problematic of Lascaux Cave

The example of drastic human induced alteration of a cave microbiome is a case of Lascaux Cave in France (Martin-Sanchez et al., 2015). The Lascaux Cave is located in Dordogne department in south-western France and contains wall Paleolithic paintings. The cave was discovered in 1940 and in 1948 was open to public. In 1955 increased amount of CO₂, humidity and temperature in the cave due to high number of visits per day so the air conditioning machine was installed. In 1960 the green colonies with green algae and a calcite film coating from water condensation appeared along the walls. Since the microbial colonization progressed, the cave was closed in 1963 to a public. Consequently, the biocide treatment containing formaldehyde solution was applied, followed by antibiotic spraying which suppressed green colonies for some time. Interestingly, in 1966 lichens appeared in cave but bacteria and fungi remain stable (Bastian et al., 2010; Martin-Sanchez et al., 2015). However, in 2001 white fungal mycelia appeared at the walls while the treatment of 50% benzalkonium chloride dissolved in ethanol or water was unsuccessful for suppression of their proliferation (Martin-Sanchez et al.,

2015). More aggressive treatments were applied, which however resulted only in proliferation of white mucus of *Pseudomonas fluorescens* and five month later first black microbial colonies appeared on the walls of the cave (Bastian et al., 2007). The mechanical cleaning and new biocide treatments were used against the black stains, however without a long lasting success (Bastian et al., 2010).

Black stains represent recently the major problem for preservation of cave paintings. The microbial composition was therefore studied to use successful treatment for their elimination. Black stains were considered as most likely fungal origin, since the *Dematiaceous hyphomycetes* producing olivaceous to black colonies, together with species from genera *Verticillium* and *Scolecobasidium* were firstly isolated from stains (Fabiola Bastian et al., 2009). The melanin producing fungi *Scolecobasidium tshawytschae* appeared in stains most likely as a reaction to previous organic treatment since is typical for oil-contaminated soils (Bastian et al., 2010). More recently, the novel fungal species of the genus *Ochroconis*, *O. lascauxensis* and *O. anomala*, which produced melanin were isolated from the stains (De La Rosa et al., 2017). Based on sequencing approach, strong interactions between fungal and bacterial species (for example between *Aspergillus*, *Janibacter* and *Streptomyces*) were found in the black stains (Alonso et al., 2018), pointing on a possibility that more complex microbial relationships play a role in stains creation. The microbial concentration in islands of nutrients in the oligotrophic cave walls might lead to increase competition linked to antibiotic production, which stress fungi to melanin production. However, if these fungi are during the creation of the black stains influenced by other microbes or how they in turn affect microbiology of the caves remain unclear but might be crucial for a definitive black stain suppression.

Actinobacteria roles in caves

Actinobacteria are highly abundant in many caves worldwide (Tomczyk-Żak and Zielenkiewicz, 2015). The suggestion that harsh conditions of caves are for this group highly favorable over the surrounding habitat was supported by a study where *Actinobacteria* dominated lava caves bacterial community contrary to the aboveground soil (Lavoie et al., 2017). Moreover, *Actinobacteria* advantage lie in the abilities to adapt to starving conditions, which for them represent a strong selection factor leading to an evolution of novel species. It

was supported by the study from Hamed et al., (2019) who found that the highest number of novel *Actinobacteria* species were associated with the most oligotrophic part of the cave. *Actinobacteria* could fight against oligotrophy by production of extracellular hydrolytic and oxidative enzymes enable them to degrade even very recalcitrant organic compounds coming to the cave (Hamed et al., 2019; Le Roes-Hill and Prins, 2016). However, another effect of some oxidative enzymes like laccase might be oxidation of phenolic molecules with subsequent polymerization into dark melanin-like pigments (Al Khatib et al., 2018). Indeed, some *Actinobacteria* like *Streptomyces* (Abdel-Halim et al., 2013; Cuezva et al., 2012; Mitova et al., 2017), *Pseudonocardia* (Porca et al., 2012) or *Nocardioidea* (Groth et al., 1999) were found to be responsible for pigmented stain formation on walls of caves.

The caves harbor unique *Actinobacteria* communities, some *Actinobacteria* were mentioned to be typical for pristine caves like *Gaiellales* (Zhu et al., 2019) while other for anthropized caves, like *Pseudonocardiaceae* (Lavoie et al., 2017). However, some genera like *Streptomyces* were also found typical for both types of caves (Maciejewska et al., 2017). *Nocardiaceae* might be also considered typical pristine cave members (De Mandal et al., 2017), however some species from that genera are known human pathogens and might be introduced into cave by visitors (Jurado et al., 2010). Also other pathogenic *Actinobacteria* species like *Dietzia maris* or *Kocuria rosea* were recovered from the air of visited Slovenian cave (Mulec et al., 2017). Therefore, the presence and proportions of potential pathogens in the community associated with visited caves is important to be monitored in relation to visitor's health.

Part 2: Decomposition of plant litter and the climate change

Soil is the largest reservoir of organic carbon in terrestrial ecosystems and plant litter decomposition is a dynamic process where nutrients from litter are sequentially returning back to the soil (Berg, 2018; Moorhead and Sinsabaugh, 2006). This process is to a large extent affected by climate (temperature, precipitation) (Glassman et al., 2018; Jin et al., 2013), soil quality (pH, bedrock material, fertility) (Delgado-Baquerizo et al., 2015), litter characteristics (quality, quantity, accessibility) (Prieto et al., 2019) and decomposer organisms like soil fauna (Frouz et al., 2015) and microbes (Baker et al., 2018). Climate was the most important factor especially when considering its effect on a regional scale, while the litter played more prominent role on a lower spatial scale (Berg, 2000). However, some studies denied the climate factor as a primary control on the decomposition process (Bradford et al., 2014) or assumed that its role was dependent only with combination with other factors as for example microbial communities or litter quality (Glassman et al., 2018; Thiessen et al., 2013).

More specifically for the climate, microbial communities were differently affected by temperature, where bacteria shifted more strongly in relation to changes in climate characteristics than fungi (Glassman et al., 2018). Temperature sensitivity also depended on a litter recalcitrancy according to Arrhenius theory: the more recalcitrant the litter was, the more dependent it was on temperature (Thiessen et al., 2013). However, the more recent study in a field experiment showed that the more the organic matter was stabilized, the less it was sensitive to temperature, which directly contrasted with the Arrhenius theory (Moinet et al., 2020). Also a precipitation was found to have a predominant effect on the decomposition, since it eliminated the moisture limitation of microbes and stimulated their activity (Jarvis et al., 2007; Smith et al., 2010).

Concerning the litter quality, the litter of higher quality (i.e. high diversity of more nutritious litter, low C:N:P ratio) was colonized by more diverse microbial decomposers contrary to lower quality litter (high proportions of structural polymers like hemicellulose and lignin, higher C:N:P ratio) and resulted in faster decomposition (Bray et al., 2012; McDaniel et al., 2014). The litter quality changed over time and microbial decomposers reflected these changes (Buresova et al., 2019). According to the general point of view, first stages of decomposition were accompanied with a loss of nutrients and r-selected microbial taxa, while

the recalcitrant debris remained in late stages and were associated with K-selected microbial taxa (F Bastian et al., 2009; Philippot et al., 2010). Other study suggested that mostly earlier stages of decomposition (especially C and N loss) were regulated by biotic factors - microbial but also nematodes community, while late stages were driven by abiotic factors - initial litter quality and soil moisture (García-Palacios et al., 2016). More specifically for litter quality, the initial stage of decomposition was driven by P-related characteristics, while later stages were mostly controlled by C:N, phenolics:N, phenolics:P and lignin:N ratios (Ge et al., 2013). Other studies suggested that N and P might stimulated decomposition in earlier stages and inhibit decomposition in later stages due to creation of unavailable complexes (Ge et al., 2013; Vitousek et al., 2010). However, as mentioned in the study by Bani et al., (2018), the decomposition was a very complex and complicated process where all factors needed to be considered jointly for more reliable application of decomposition models.

The climatic models predicted changes in temperature as a reaction to greenhouse gas emissions and included temperature increase with rainfall reduction, however with more extreme precipitation events (Field et al., 2012). As a result, plant litter production and litter quality would be modified, which would have consequences on decomposition dynamics. As decomposition dynamics affected the nutrients availability in soil, the litter quality of new litter would be in turn modified, which shifted the balance of nutrient and C cycling (Prieto et al., 2019). However, it was indicated that direct climate changes in rainfall controlled the C and N mineralization to a greater extent than the indirect ones, namely CO₂-induced changes in litter quality (Jin et al., 2013). The impacts of climate changes might also directly affect decomposition, for example by inhibition of enzyme activities due to soil drought as a reaction to warming and reduced rainfall (Suseela et al., 2014). Although the rain event in dry soil increased decomposition activity more than intense rain event in wet soils (Lee et al., 2014), too high intensity of rain might reduce oxygen availability for decomposers (Schwartz et al., 2007) and slow down the decomposition. The direct changes in climate conditions influenced more bacterial than fungal litter community, which specifically impacted decomposition rates but also a type of degraded carbon (Glassman et al., 2018).

Microbial colonizers of litter

Fungi

Although soil fauna might contribute to a large extent to physical fragmentation of litter and its mixing with organic and mineral horizons (Frouz et al., 2013), microbes were the main mediators of plant litter decomposition (Bani et al., 2018). Fungi were always considered crucial in decomposition since they produced variety of extracellular enzymes including those which were able to decompose even very recalcitrant polymers like lignin from plant litter (Purahong et al., 2014). Fungi were divided into subcommunities according to different stages and function in decomposition. The early-stage subcommunity produced hydrolytic and oxidative enzymes while the later stage subcommunity contributed to laccase and peroxidase production (Purahong et al., 2016a). At the same time, earlier stages fungi were mostly identified as *Ascomycota* and later stages fungi as *Basidiomycota* (Purahong et al., 2016b). Concerning the abiotic factors affecting fungi during decomposition, fungal communities were less responsible to climate changes than bacteria (Glassman et al., 2018). It was contradictory to a study where a fungal community structure was alternated with changes in a high and a low soil moisture, however it did not affect the litter decomposition (Kaisermann et al., 2015).

In our study (Buresova et al., 2019), fungi were concluded as site generalists, but substrate specialists since they preferentially colonized one litter type. Indeed, fungi were mostly known to be driven by a litter quality since they could select for a certain litter type thanks to their colonizing abilities and selective translocation of potentially limiting nutrients by fungal hyphae (Song et al., 2010). Generally, fungi were known as dwellers of typical recalcitrant litter, like deadwood (Johnston et al., 2016). It was in accordance with a study, where fungi were explored to be earlier colonizers and their decomposition of leaf litter might facilitate a bacterial decomposition (Purahong et al., 2014). In spite of this, it was found that fungi were associated to a high quality C source, which contrasted the generally known hypothesis of their K-strategist orientation (Rousk and Frey, 2015). Simultaneously, high quality litter stimulated the production of fungal but not bacterial hydrolytic enzymes (Schneider et al., 2012).

Bacteria

As opposed to fungi, bacteria were believed to be r-strategic decomposers but latest studies showed that some bacteria were able to be K-strategists (Bani et al., 2018). Many bacteria including *Proteobacteria*, *Acidobacteria*, *Actinobacteria*, *Firmicutes* and *Bacteroidetes* were recovered from deadwood, where they most likely cooperated or competed during decomposition with fungi (Johnston et al., 2016). Although some bacterial groups might utilize easily accessible low-molecular weight compounds or scavenge the products of decomposition from other microbes (Šnajdr et al., 2011), others were able to directly participate in the process by a production of hydrolytic and oxidative enzymes (Větrovský et al., 2014). For example, glycoside hydrolases β -1,4-endoglucanase, cellobiohydrolase and β -d-glucosidase were produced by isolates from groups *Actinobacteria*, *Proteobacteria* and *Firmicutes* suggesting they were able to completely hydrolytically decompose cellulose (Woo et al., 2014). Within that, oxidases and peroxidases were highly active in *Gordonia* (*Actinobacteria*) and less active in *Aquitalea* (*Proteobacteria*), suggested these strains were able to degrade lignin although with different efficiency (Woo et al., 2014). Production of chitinase might be linked to N-rich substrates including fungal biomass. With this enzyme, bacteria might mediate decomposition of death fungal biomass (Tlaskal et al., 2016) or actively inhibited fungal growth (Lacombe-Harvey et al., 2018).

Actinobacteria as one of the main litter decomposers

Actinobacteria were one of the most dominating bacterial group during the plant litter decomposition in different ecosystems (Glassman et al., 2018; Johnston et al., 2016). Soil and litter dwelling *Actinobacteria* were advantageous during decomposition due to their abilities to better colonize the litter thanks to their mycelial growth (De Boer et al., 2005), extracellular enzymes (Le Roes-Hill and Prins, 2016) and secondary metabolites production. Secondary metabolites might serve as inhibiting or communicating molecules, which might modulate the relationships of community during the decomposition (Polkade et al., 2016).

Actinobacteria were known to be litter and soil quality dependent, which was supported by a study where *Actinobacteria* but not a fungal biomass increased after complex organic fertilizers addition to a soil (Jenkins et al., 2009). *Actinobacteria* were also promoted by

inorganic inputs including external S supply (Xu et al., 2016). However, N amendment in the late stages down-regulated *Actinobacteria* but not fungal abundance. It might be caused by limitation of C because *Actinobacteria* were found to regulate CO₂ emission derived from litter and were thus more than fungi dependent on C availability (Xu et al., 2016). Generally, it was presumed that *Actinobacteria* colonize low nutritious and recalcitrant substrates (Kurtbóke, 2017). Since the decomposition process was sorted into different stages according to actual quality of decomposing litter, *Actinobacteria* should therefore be associated with the later stages when recalcitrant residues remained (F Bastian et al., 2009). Some study indeed found *Actinobacteria* as late-stage decomposers, however they explained it as a result of *Actinobacteria* slow growth, which might indicate low contribution to C flow through microbial food web (López-Mondéjar et al., 2018).

Concerning the direct effect of climate change during decomposition, although *Actinobacteria* were generally considered highly seasonally responsible (Kopecky et al., 2011; M. Sagova-Mareckova et al., 2015), they could be stable during the altered climatic conditions. The example is a study by Tikhonova et al., (2019) where *Actinobacteria* dominated in a part of the microbial decomposers community, which remained stable during temperature increase. However, more studies are necessary to understand how *Actinobacteria* are during decomposition responsible to direct or indirect climate changes. Since they are subjected to variety of factors, including pH, the complexity of factors is necessary to take into account for understanding how climate change affects *Actinobacteria* community structure and biomass.

Extracellular hydrolytic and oxidative enzymes

Actinobacteria were found to produce variety of extracellular hydrolytic (Lacombe-Harvey et al., 2018) and oxidative enzymes (Le Roes-Hill and Prins, 2016; Woo et al., 2014). Production of such enzymes like for example chitinases, cellulases, β -glucanases and xylanases aided them in existing under different conditions thus enable them to inhabited multiple niches (Bennur et al., 2015). In recent years, *Actinobacteria* were interestingly found producing oxidative enzymes like peroxidases and laccase with potential application in biotechnological processes or bioremediation (Le Roes-Hill and Prins, 2016). Especially laccases have attracted much attention recently, because, contrary to fungal laccase, *Actinobacteria* laccases were active in a wider range of abiotic factors like pH and temperature (Fernandes et al., 2014). With laccase and other oxidases and peroxidases, *Actinobacteria* could decompose lignin, although

it was assumed that with lower efficiency than fungi (Bugg et al., 2011). With hydrolytic enzymes like cellobiohydrolase, β -glucosidase and β -xylosidase they could degrade the low-molecular weight compounds containing lignocellulosic polymers and potentially cooperated on a lignin degradation with other microbes like fungi (Johnston et al., 2016; Větrovský et al., 2014). Concerning other types of enzymes, *Actinobacteria* might acquire phosphate from organic esters by production of phosphatases (Lladó et al., 2017) or nitrogen from N-rich polymers like fungal hyphae by chitinases (Lacombe-Harvey et al., 2018). Moreover, some studies suggested that with chitinases *Actinobacteria* might suppress fungal growth (Gasmi et al., 2019).

The enzymology of bacteria in general is less explored and one study contemplated that bacteria break the lignin by oxidative enzymes and laccase due to protection against toxicity of phenolic compounds rather than to decompose lignin (Lladó et al., 2017). These questions might be important in relation to other stress like situations including environmental factors changes and low nutrient contents. Therefore, understanding enzyme production under stress conditions might be crucial in determining how carbon storage in soil would be affected under amplified climate conditions.

Relationship of main decomposers

Since the results predicting how abiotic factors explain *Actinobacteria* community structure and abundance are not always consistent, the biotic factors such as interaction with other community members might explain the rest of the unknown variability driving *Actinobacteria* (Bani et al., 2018; De Boer et al., 2005; Purahong et al., 2016b). If different microbial groups have a similar role in decomposition, their niches overlap. The mutual cooperation or competition, but also different sensitivity to environmental conditions and other variables, which regulated microbes, resulted in niche separation (Kielak et al., 2016; Leibold and McPeck, 2006). For example, *Actinobacteria* abundance decreased over decomposition from winter to spring, which was contrary to *Proteobacteria*. It might point on *Actinobacteria* seasonality over decomposition process together with niche separation over time with another dominant bacterial decomposer (Schneider et al., 2012). Similarly, *Actinobacteria* increased the most among other bacteria after litter addition to the soil, but more in a case, where fungi did not dominate (Malik et al., 2016). Simultaneously, it was found that when *Actinobacteria*

increased, especially after organic input to soil, a fungal abundance falls (Jenkins et al., 2009). It suggested a negative relationship between these two groups, a niche separation and a potential inhibition of fungi by *Actinobacteria*. On the contrary, *Actinobacteria* colonizing organic substrate might also facilitate a growth of microbes already present in the substrate suggesting a synergistic interaction (Rousk et al., 2008; Zhao et al., 2016). The mutual relationship between *Actinobacteria* and fungi has rarely been studied in a single experiment although *Actinobacteria* advantages during decomposition are generally similar for fungi, making *Actinobacteria* the main fungal competitors (De Boer et al., 2005).

Summary and conclusions

Actinobacteria community in anthropized and pristine caves

The thesis shows that *Actinobacteria* may serve as indicators of human derived contamination in caves since *Actinobacteria* community structure differed between anthropized and pristine caves (**paper I**). More precisely, *Actinobacteria* indicated the least affected caves (Mouflon, Reille) and one room within the Lascaux Cave (Diaclase) by high proportions of typical pristine taxa from the order *Gaiellales* (Zhu et al., 2019), the environmental clone *MB-A2-108_fa* (Kumar De, 2019; Zhang et al., 2019) and unknown *Actinobacteria* taxa (Rangseekaew and Pathom-Aree, 2019). Rouffignac Cave was identified as being under intermediate human-derived pressure due to the coexistence of *Actinobacteria* groups, which were earlier recognized as typical for pristine caves, namely *Jiangellaceae* (Rangseekaew and Pathom-Aree, 2019)) as well as typically anthropized, namely by observation of high richness but low evenness of *Nocardia* including potentially pathogenic species (Jurado et al., 2010; Modra et al., 2017; Roxburgh et al., 2004)). The Lascaux Cave was characterized by high proportions of *Mycobacteriaceae*, a group containing potentially pathogenic members (Jurado et al., 2010; Modra et al., 2017). These findings point to the shift of *Actinobacteria* community towards high diversity and proportions of potential human pathogens in the visited caves.

Use of the novel marker gene hsp65 for identification of Actinobacteria species

In the thesis, it is demonstrated that the *hsp65* gene marker had higher nucleotide polymorphism and higher molecular distances than the 16S rRNA gene especially on the lowest taxonomic level of *Actinobacteria* (**paper I**). We designed a database, where sequences of paralogous genes were recognized and marked but only groups from the class *Actinobacteria* were targeted by selective primers and were thus, successfully identified by the database (Telenti et al., 1993). Although the 16S rRNA marker identified more groups on higher taxonomic ranks than *hsp65* marker, the *hsp65* marker identified more species. Therefore, the *hsp65* gene could be used as a supportive marker complementary to the 16S rRNA based studies where identification of *Actinobacteria* to the species level is needed. However, the identification of *Actinobacteria* with the *hsp65* gene marker has a limitation because the

database has only a limited number of deposited sequences compared to the 16S rRNA reference databases and primers were not complementary to all *Actinobacteria* taxa.

Visual dark marks on walls of the highly anthropized Lascaux Cave in relation to Actinobacteria communities

The differences of *Actinobacteria* community between visual marks (black stains and dark areas) and unmarked areas on cave walls were demonstrated, while higher diversity was measured in visual marks (**paper I**). Furthermore, higher proportions of *Streptomyces* with the highest proportion of *S. mirabilis* were identified in the visual marks especially at the cave entrance. *Streptomyces* might contribute to pigment production as was monitored in other studies (Abdel-Haliem et al., 2013; Cuezva et al., 2012). Moreover, *Streptomyces* were dominating in pigmented areas of other caves like Altamira, Spain (Groth et al., 1999). These findings suggest that *Streptomyces* were influenced by visual marks and potentially, might be involved in the visual marks formation. However, more detailed studies on the pigment production or interaction with the pigment forming fungi with *Streptomyces* is needed for understanding their role.

Actinobacteria communities in relation to litter changes during decomposition

Concerning the decomposition process in temperate climate, *Actinobacteria* abundance was not favorized by any studied litter type or its origin. but their dynamics over one year were dependent on litter type and their abundance increased in later stages of decomposition (**paper II**). It implies that *Actinobacteria* were functionally redundant since their decomposition strategies function according to the litter quality rather than the litter type and origin (Glassman et al., 2018) (**paper II**). In the Mediterranean climate, *Actinobacteria* preferred deciduous litter with the highest nutrient content but also with structural compounds such as lignin, cellulose, hemicellulose regardless of its origin (**paper IV**). Similarly, in a mountainous forest, *Actinobacteria* dominated later-stage bacterial community on the more resistant spruce litter compared to the beech litter (**paper III**).

These results show that *Actinobacteria* were mostly favorized by a recalcitrant litter or stages, when the litter became recalcitrant (F Bastian et al., 2009; Bray et al., 2012; Bugg et al.,

2011; Fierer et al., 2007). Moreover, *Actinobacteria* were able to adapt to changes in a litter, however, their dominance did not lead to an increased rate of decomposition (**paper III, IV**). It shows that *Actinobacteria* were adaptable to litter alteration with a potential to dominate the recalcitrant litter but the changes did not promote decomposition and thus also carbon and nutrients release.

Actinobacteria adaptability to changes in habitat properties

In the thesis, *Actinobacteria* occurring in the litter were not affected by rainfall reduction in Mediterranean forests (**paper IV**) and also in the temperate mountainous forest, other than climate characteristics affected their abundance (**paper III**). The comparison of temperate grassland and beech forest showed that the grassland harbored higher abundance of *Actinobacteria* but the decomposition was overall faster at the forest site (**paper II**).

Therefore, not only climatic characteristics, but also another factor or complex site conditions affected *Actinobacteria* in these habitats (**paper II, III, IV**). *Actinobacteria* were considered slow-growing thus, their response to actual weather changes might be retarded (López-Mondéjar et al., 2018) or they might be resistant to weather changes due to the spore formation (Acosta-Martínez et al., 2014; B.-Z. Fang et al., 2017). When comparing the forest habitats, *Actinobacteria* were able to adapt to introduced coniferous forests in both Mediterranean and temperate mountainous climate since they finally dominated the spruce or pine bacterial community in comparison to the native beech or oak deciduous forests, respectively (**papers III, IV**). Generally, acidic and nutrient-limited coniferous forest might enable the increase of *Actinobacteria* proportions because they are more than other community members adaptable to such harsh conditions, again suggesting *Actinobacteria* as oligotrophic and adaptable taxa.

However, the dominance of *Actinobacteria* at the respective sites (grassland (**paper II**), coniferous forests (**papers III, IV**)) did not lead to more intense decomposition of litter contrary to deciduous forests (**paper II, III, IV**). These results suggest that climate change would affect *Actinobacteria* indirectly through habitat alteration and the altered habitats dominated by *Actinobacteria* might be linked to slower carbon release from the litter.

Actinobacteria as late-stage decomposers with opposite patterns to other decomposers

The succession of microbial decomposers during decomposition with closer focus on *Actinobacteria* and fungi has been rarely investigated. In the temperate beech forest and grassland, *Actinobacteria* were similarly to fungi driven by a litter quality but an overall abundance of fungi was comparable between the sites while *Actinobacteria* abundance was site specific. Moreover, *Actinobacteria* were similarly to *Firmicutes* late-stage colonizers but fungi and *Alphaproteobacteria* were initial stage litter colonizers (**paper II**). In the Mediterranean climate, *Actinobacteria* were not affected by rainfall reduction contrary to fungi (**paper IV**). There, *Actinobacteria* were favored by deciduous litter, while fungi mostly by coniferous litter except for the coniferous forest, where both groups coexisted in a high abundance at the same litter (**papers IV**). These findings point to the niche separation between *Actinobacteria* and fungi but also *Actinobacteria* and other decomposers during litter decomposition (*Firmicutes*, *Alphaproteobacteria*) (**paper II**).

Concerning the *Actinobacteria* and fungi, they had opposite strategies driven by different factors (**paper II, III, IV**) and their succession over time was opposite (**paper II**). However, in certain situations such as on the most palatable litter type in the coniferous forest (**paper IV**), intermediately recalcitrant litter type in the grassland (**paper II**)) their niches were overlapping (**paper II, IV**), which might point to a synergistic interaction (Meidute et al., 2008). Overall, *Actinobacteria* and fungi were litter quality dependent however, fungi seemed to be affected more by direct climatic factors (**papers III, IV**) and *Actinobacteria* more by indirect climatic factors such as habitat alteration (**papers II, III, IV**). Therefore, the climate derived changes leading to altered litter or precipitation might shift the fungal to actinobacterial ratio and thus the relationship between the two groups potentially leading to an altered decomposition process and carbon cycling.

Concluding remarks

From the results described above, I conclude that *Actinobacteria* communities might play important roles in anthropogenic and natural habitats and are directly and indirectly influenced by human-derived changes. The thesis might help in improving the study of *Actinobacteria* with a novel marker gene *hsp65*. We also suggest that *Actinobacteria* are adaptable members of microbial communities, particularly to man-induced changes in caves

and during decomposition processes on a plant litter. Consequently, this taxonomic group might be proposed as a bioindicator in monitoring of environmental quality or in prediction models. Thus, *Actinobacteria* deserve a closer attention and more studies showing how their activity is modified by human derived changes. Moreover, our results show that interactions between *Actinobacteria* and other microbial members, especially fungi, might play an important role in driving the ecology of *Actinobacteria* in the studied environments. Thus, new experiments that would uncover the positive or negative interactions between *Actinobacteria* and fungi are proposed for understanding their combined contribution to decomposition and carbon cycling.

References

- Abdel-Halim, M.E.F., Sakr, A.A., Ali, M.F., Ghaly, M.F., Sohlenkamp, C., 2013. Characterization of *Streptomyces* isolates causing colour changes of mural paintings in ancient Egyptian tombs. *Microbiol. Res.* 168, 428–437. <https://doi.org/10.1016/j.micres.2013.02.004>
- Abdu, N., Abdullahi, A.A., Abdulkadir, A., 2017. Heavy metals and soil microbes. *Environ. Chem. Lett.* 15, 65–84. <https://doi.org/10.1007/s10311-016-0587-x>
- Acosta-Martínez, V., Cotton, J., Gardner, T., Moore-Kucera, J., Zak, J., Wester, D., Cox, S., 2014. Predominant bacterial and fungal assemblages in agricultural soils during a record drought/heat wave and linkages to enzyme activities of biogeochemical cycling. *Appl. Soil Ecol.* 84, 69–82. <https://doi.org/10.1016/j.apsoil.2014.06.005>
- Adékambi, T., Drancourt, M., 2004. Dissection of phylogenetic relationships among 19 rapidly growing *Mycobacterium* species by 16S rRNA, hsp65, sodA, recA and rpoB gene sequencing. *Int. J. Syst. Evol. Microbiol.* 54, 2095–2105. <https://doi.org/10.1099/ijs.0.63094-0>
- Adetutu, E.M., Thorpe, K., Shahsavari, E., Bourne, S., Cao, X., Fard, R.M.N., Kirby, G., Ball, A.S., 2012. Bacterial community survey of sediments at Naracoorte Caves, Australia. *Int. J. Speleol.* 41, 137–147. <https://doi.org/10.5038/1827-806X.41.2.2>
- Agustinho, D.P., Nosanchuk, J.D., 2017. Functions of Fungal Melanins, in: Reference Module in Life Sciences. Elsevier. <https://doi.org/10.1016/b978-0-12-809633-8.12091-6>
- Al Khatib, M., Harir, M., Costa, J., Baratto, M., Schiavo, I., Trabalzini, L., Pollini, S., Rossolini, G., Basosi, R., Pogni, R., 2018. Spectroscopic Characterization of Natural Melanin from a *Streptomyces cyaneofuscatus* Strain and Comparison with Melanin Enzymatically Synthesized by Tyrosinase and Laccase. *Molecules* 23, 1916. <https://doi.org/10.3390/molecules23081916>
- Alonso, L., Creuzé-des-Châtelliers, C., Trabac, T., Dubost, A., Moëgne-Loccoz, Y., Pommier, T., 2018. Rock substrate rather than black stain alterations drives microbial community structure in the passage of Lascaux Cave. *Microbiome* 6, 216. <https://doi.org/10.1186/s40168-018-0599-9>
- Alvarez, A., Saez, J.M., Davila Costa, J.S., Colin, V.L., Fuentes, M.S., Cuozzo, S.A., Benimeli, C.S., Polti, M.A., Amoroso, M.J., 2017. Actinobacteria: Current research and perspectives for bioremediation of pesticides and heavy metals. *Chemosphere*. <https://doi.org/10.1016/j.chemosphere.2016.09.070>
- Ando, K., Murakami, T., 2020. Detection of human-associated bacteria in water from Akiyoshi-do Cave, Japan. *Water Environ. Res.* wer.1355. <https://doi.org/10.1002/wer.1355>
- Aszalós, J.M., Szabó, A., Megyes, M., Anda, D., Nagy, B., Borsodi, A.K., 2020. Bacterial Diversity of a High-Altitude Permafrost Thaw Pond Located on Ojos del Salado (Dry Andes, Altiplano-Atacama Region). *Astrobiology* 20, 754–765. <https://doi.org/10.1089/ast.2018.2012>
- Baker, N.R., Khalili, B., Martiny, J.B.H., Allison, S.D., 2018. Microbial decomposers not constrained by climate history along a Mediterranean climate gradient in southern California. *Ecology* 99, 1441–1452. <https://doi.org/10.1002/ecy.2345>
- Banerjee, S., Joshi, S.R., 2013. Insights into cave architecture and the role of bacterial biofilm. *Proc. Natl. Acad. Sci. India Sect. B - Biol. Sci.* 83, 277–290. <https://doi.org/10.1007/s40011-012-0149-3>
- Bani, A., Pioli, S., Ventura, M., Panzacchi, P., Borruso, L., Tognetti, R., Tonon, G., Brusetti, L., 2018. The role of microbial community in the decomposition of leaf litter and deadwood. *Appl. Soil Ecol.* 126, 75–84.
- Barka, E.A., Vatsa, P., Sanchez, L., Gaveau-Vaillant, N., Jacquard, C., Klenk, H.-P., Clément, C., Ouhdouch, Y., van Wezel, G.P., 2016. Taxonomy, Physiology, and Natural Products of Actinobacteria. *Microbiol. Mol. Biol. Rev.* 80, 1–43. <https://doi.org/10.1128/mmbr.00019-15>
- Barton, H.A., Jurado, V., 2007. What's Up Down There? Microbial Diversity in Caves Microorganisms in caves survive under nutrient-poor conditions and are metabolically versatile and unexpectedly diverse. *Microbe* 2, 132–138.

- Bastian, Fabiola, Alabouvette, C., Journal, O., 2009. Lights and shadows on the conservation of a rock art cave: the case of Lascaux cave International Journal of Speleology International Journal of Speleology Lights and shadows on the conservation of a rock art cave: The case of Lascaux Cave, International Journal of Speleology.
- Bastian, F, Bouziri, L., Nicolardot, B., Ranjard, L., 2009. Impact of wheat straw decomposition on successional patterns of soil microbial community structure. *Soil Biol. Biochem.* 41, 262–275.
- Bastian, F., Jurado, V., Novakova, A., Alabouvette, C., Saiz-Jimenez, C., 2010. The microbiology of Lascaux Cave. *Microbiology* 156, 644–652. <https://doi.org/10.1099/mic.0.036160-0>
- Bastian, F., Orial, G., François, A., Alabouvette, C., 2007. La grotte de Lascaux : un écosystème complexe où bactéries et champignons interagissent. *Biofutur* 26, 28–31.
- Bennur, T., Kumar, A.R., Zinjarde, S., Javdekar, V., 2015. *Nocardia* species: Incidence, ecological roles and adaptations. *Microbiol. Res.* 174, 33–47. <https://doi.org/10.1016/J.MICRES.2015.03.010>
- Bentley, S.D., Corton, C., Brown, S.E., Barron, A., Clark, L., Doggett, J., Harris, B., Ormond, D., Quail, M.A., May, G., Francis, D., Knudson, D., Parkhill, J., Ishimaru, C.A., 2008. Genome of the actinomycete plant pathogen *Clavibacter michiganensis* subsp. *sepedonicus* suggests recent niche adaptation. *J. Bacteriol.* 190, 2150–2160. <https://doi.org/10.1128/JB.01598-07>
- Bercea, S., Năstase-Bucur, R., Moldovan, O.T., Kenesz, M., Constantin, S., 2019. Yearly microbial cycle of human exposed surfaces in show caves. *Subterr. Biol.* 31, 1–14. <https://doi.org/10.3897/subtbiol.31.34490>
- Berg, B., 2018. Decomposing litter; limit values; humus accumulation, locally and regionally. *Appl. Soil Ecol.* 123, 494–508. <https://doi.org/10.1016/J.APSSOIL.2017.06.026>
- Berg, B., 2000. Litter decomposition and organic matter turnover in northern forest soils. *For. Ecol. Manage.* 133, 13–22. [https://doi.org/10.1016/S0378-1127\(99\)00294-7](https://doi.org/10.1016/S0378-1127(99)00294-7)
- Bhullar, K., Waglechner, N., Pawlowski, A., Koteva, K., Banks, E.D., Johnston, M.D., Barton, H.A., Wright, G.D., 2012. Antibiotic Resistance Is Prevalent in an Isolated Cave Microbiome. *PLoS One* 7, e34953. <https://doi.org/10.1371/journal.pone.0034953>
- Boiron, P., Provost, F., Chevrier, G., Dupont, B., 1992. Review of nocardial infections in France 1987 to 1990. *Eur. J. Clin. Microbiol. Infect. Dis.* 11, 709–714. <https://doi.org/10.1007/BF01989975>
- Bradford, M.A., Ii, R.J.W., Baldrian, P., Crowther, T.W., Maynard, D.S., Oldfield, E.E., Wieder, W.R., Wood, S.A., King, J.R., 2014. Climate fails to predict wood decomposition at regional scales. <https://doi.org/10.1038/NCLIMATE2251>
- Bragança, I., Mucha, A.P., Tomasino, M.P., Santos, F., Lemos, P.C., Delerue-Matos, C., Domingues, V.F., 2019. Deltamethrin impact in a cabbage planted soil: Degradation and effect on microbial community structure. *Chemosphere* 220, 1179–1186. <https://doi.org/10.1016/j.chemosphere.2019.01.004>
- Bray, S.R., Kitajima, K., Mack, M.C., 2012. Temporal dynamics of microbial communities on decomposing leaf litter of 10 plant species in relation to decomposition rate. *Soil Biol. Biochem.* 49, 30–37. <https://doi.org/10.1016/j.soilbio.2012.02.009>
- Bugg, T.D.H., Ahmad, M., Hardiman, E.M., Singh, R., 2011. The emerging role for bacteria in lignin degradation and bio-product formation. *Curr. Opin. Biotechnol.* 22, 394–400. <https://doi.org/10.1016/j.copbio.2010.10.009>
- Buresova, A., Kopecky, J., Hrdinkova, V., Kamenik, Z., Omelka, M., Sagova-Mareckova, M., 2019. Succession of microbial decomposers is determined by litter type, but site conditions drive decomposition rates. *Appl. Environ. Microbiol.* 85. <https://doi.org/10.1128/AEM.01760-19>
- Carlo, E. Di, Chisesi, R., Barresi, G., Barbaro, S., Lombardo, G., Rotolo, V., Sebastianelli, M., Travagliato, G., Palla, F., 2016. Fungi and Bacteria in Indoor Cultural Heritage Environments: Microbial-related Risks for Artworks and Human Health. *Environ. Ecol. Res.* 4, 257–264. <https://doi.org/10.13189/eer.2016.040504>
- Chen, X., Zhao, Y., Zhao, X., Wu, J., Zhu, L., Zhang, X., Wei, Z., Liu, Y., He, P., 2020. Selective pressures of

- heavy metals on microbial community determine microbial functional roles during composting: Sensitive, resistant and actor. *J. Hazard. Mater.* 398, 122858. <https://doi.org/10.1016/j.jhazmat.2020.122858>
- Cirigliano, A., Tomassetti, M.C., Di Pietro, M., Mura, F., Maneschi, M.L., Gentili, M.D., Cardazzo, B., Arrighi, C., Mazzoni, C., Negri, R., Rinaldi, T., 2018. Calcite moonmilk of microbial origin in the Etruscan Tomba degli Scudi in Tarquinia, Italy. *Sci. Rep.* 8, 1–10. <https://doi.org/10.1038/s41598-018-34134-y>
- Cuezva, S., Fernandez-Cortes, A., Porca, E., Pašić, L., Jurado, V., Hernandez-Marine, M., Serrano-Ortiz, P., Hermosin, B., Cañaveras, J.C., Sanchez-Moral, S., Saiz-Jimenez, C., 2012. The biogeochemical role of Actinobacteria in Altamira Cave, Spain. *FEMS Microbiol. Ecol.* 81, 281–290. <https://doi.org/10.1111/j.1574-6941.2012.01391.x>
- Cuezva, S., Sanchez-Moral, S., Saiz-Jimenez, C., Carlos Canaveras, J., 2009. Microbial Communities and Associated Mineral Fabrics in Altamira Cave, Spain. *Int. J. Speleol.* 38, 83–92.
- Davis, M.C., Messina, M.A., Nicolosi, G., Petralia, S., Baker, M.D., Mayne, C.K.S., Dinon, C.M., Moss, C.J., Onac, B.P., Garey, J.R., 2020. Surface runoff alters cave microbial community structure and function. *PLoS One* 15, e0232742. <https://doi.org/10.1371/journal.pone.0232742>
- De Boer, W., Folman, L.B., Summerbell, R.C., Boddy, L., 2005. Living in a fungal world: Impact of fungi on soil bacterial niche development. *FEMS Microbiol. Rev.* 29, 795–811. <https://doi.org/10.1016/j.femsre.2004.11.005>
- De La Rosa, J.M., Martin-Sanchez, P.M., Sanchez-Cortes, S., Hermosin, B., Knicker, H., Saiz-Jimenez, C., 2017. Structure of melanins from the fungi *Ochroconis lascauxensis* and *Ochroconis anomala* contaminating rock art in the Lascaux Cave. *Sci. Rep.* 7, 1–11. <https://doi.org/10.1038/s41598-017-13862-7>
- De Mandal, S., Chatterjee, R., Kumar, N.S., 2017. Dominant bacterial phyla in caves and their predicted functional roles in C and N cycle. *BMC Microbiol.* 17, 1–9. <https://doi.org/10.1186/s12866-017-1002-x>
- Delgado-Baquerizo, M., García-Palacios, P., Milla, R., Gallardo, A., Maestre, F.T., 2015. Soil characteristics determine soil carbon and nitrogen availability during leaf litter decomposition regardless of litter quality. *Soil Biol. Biochem.* 81, 134–142. <https://doi.org/10.1016/j.soilbio.2014.11.009>
- Delgado-Baquerizo, M., Maestre, F.T., Reich, P.B., Jeffries, T.C., Gaitan, J.J., Encinar, D., Berdugo, M., Campbell, C.D., Singh, B.K., 2016. Microbial diversity drives multifunctionality in terrestrial ecosystems. *Nat. Commun.* 7, 1–8. <https://doi.org/10.1038/ncomms10541>
- Dhami, N.K., Mukherjee, A., Watkin, E.L.J., 2018. Microbial diversity and mineralogical-mechanical properties of calcitic cave speleothems in natural and in vitro biomineralization conditions. *Front. Microbiol.* 9. <https://doi.org/10.3389/fmicb.2018.00040>
- Diancourt, L., Passet, V., Verhoef, J., Grimont, P.A.D., Brisse, S., 2005. Multilocus Sequence Typing of *Klebsiella pneumoniae* Nosocomial Isolates. *J. Clin. Microbiol.* 43, 4178–4182. <https://doi.org/10.1128/JCM.43.8.4178-4182.2005>
- Fang, B.-Z., Salam, N., Han, M.-X., Jiao, J.-Y., Cheng, J., Wei, D.-Q., Xiao, M., Li, W.-J., 2017. Insights on the Effects of Heat Pretreatment, pH, and Calcium Salts on Isolation of Rare Actinobacteria from Karstic Caves. *Front. Microbiol.* 8. <https://doi.org/10.3389/fmicb.2017.01535>
- Fang, B.Z., Hua, Z.S., Han, M.X., Zhang, Z.T., Wang, Y.H., Yang, Z.W., Zhang, W.Q., Xiao, M., Li, W.J., 2017. *Nonomurea cavernae* sp. Nov., a novel actinobacterium isolated from a karst cave sample. *Int. J. Syst. Evol. Microbiol.* 67, 4692–4697. <https://doi.org/10.1099/ijsem.0.002364>
- Fernandes, A.R.T., da Silveira, W.B., Passos, M.L.F., Zucchi, T.D., 2014. Laccases from Actinobacteria-What We Have and What to Expect. *SciRes. Adv. Microbiol.* 4, 285–296. <https://doi.org/10.4236/aim.2014.46035>
- Field, C.B., Barros, V., Stocker, T.F., Dahe, Q., Jon Dokken, D., Ebi, K.L., Mastrandrea, M.D., Mach, K.J., Plattner, G.K., Allen, S.K., Tignor, M., Midgley, P.M., 2012. Managing the risks of extreme events and disasters to advance climate change adaptation: Special report of the intergovernmental panel on climate change, Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation:

Special Report of the Intergovernmental Panel on Climate Change.
<https://doi.org/10.1017/CBO9781139177245>

- Fierer, N., Bradford, M.A., Jackson, R.B., 2007. Toward an ecological classification of soil bacteria. *Ecology* 88, 1354–1364. <https://doi.org/10.1890/05-1839>
- Fox, G.E., Wisotzkey, J.D., Jurtshuk, P., 1992. How Close Is Close: 16S rRNA Sequence Identity May Not Be Sufficient To Guarantee Species Identity. *Int. J. Syst. Bacteriol.* 42, 166–170. <https://doi.org/10.1099/00207713-42-1-166>
- Francioli, D., Schulz, E., Lentendu, G., Wubet, T., Buscot, F., Reitz, T., 2016. Mineral vs. organic amendments: Microbial community structure, activity and abundance of agriculturally relevant microbes are driven by long-term fertilization strategies. *Front. Microbiol.* 7, 1446. <https://doi.org/10.3389/fmicb.2016.01446>
- Frouz, J., Jílková, V., Cajthaml, T., Pižl, V., Tajovský, K., Háněl, L., Burešová, A., Šimáčková, H., Kolaříková, K., Franklin, J., Nawrot, J., Groninger, J.W., Stahl, P.D., 2013. Soil biota in post-mining sites along a climatic gradient in the USA: Simple communities in shortgrass prairie recover faster than complex communities in tallgrass prairie and forest. *Soil Biol. Biochem.* 67. <https://doi.org/10.1016/j.soilbio.2013.08.025>
- Frouz, J., Roubíčková, A., Heděnc, P., Tajovský, K., 2015. Do soil fauna really hasten litter decomposition? A meta-analysis of enclosure studies. *Eur. J. Soil Biol.* 68, 18–24. <https://doi.org/10.1016/j.ejsobi.2015.03.002>
- Gagneux, S., 2018. Ecology and evolution of *Mycobacterium tuberculosis*. *Nat. Rev. Microbiol.* <https://doi.org/10.1038/nrmicro.2018.8>
- García-Palacios, P., Shaw, E.A., Wall, D.H., Hättenschwiler, S., 2016. Temporal dynamics of biotic and abiotic drivers of litter decomposition. *Ecol. Lett.* 19, 554–563. <https://doi.org/10.1111/ele.12590>
- Garcia, M.O., Templer, P.H., Sorensen, P.O., Sanders-DeMott, R., Groffman, P.M., Bhatnagar, J.M., 2020. Soil Microbes Trade-Off Biogeochemical Cycling for Stress Tolerance Traits in Response to Year-Round Climate Change. *Front. Microbiol.* 11, 616. <https://doi.org/10.3389/fmicb.2020.00616>
- Gasmi, M., Kitouni, M., Carro, L., Pujic, P., Normand, P., Boubakri, H., 2019. Chitinolytic actinobacteria isolated from an Algerian semi-arid soil: development of an antifungal chitinase-dependent assay and GH18 chitinase gene identification. *Ann. Microbiol.* 69, 395–405. <https://doi.org/10.1007/s13213-018-1426-z>
- Ge, X., Zeng, L., Xiao, W., Huang, Z., Geng, X., Tan, B., 2013. Effect of litter substrate quality and soil nutrients on forest litter decomposition: A review. *Acta Ecol. Sin.* 33, 102–108. <https://doi.org/10.1016/j.chnaes.2013.01.006>
- Gillings, M.R., Paulsen, I.T., 2014. Microbiology of the anthropocene. *Anthropocene* 5, 1–8. <https://doi.org/10.1016/j.ancene.2014.06.004>
- Glassman, S.I., Weihe, C., Li, J., Albright, M.B.N., Looby, C.I., Martiny, A.C., Treseder, K.K., Allison, S.D., Martiny, J.B.H., 2018. Decomposition responses to climate depend on microbial community composition. *Proc. Natl. Acad. Sci. U. S. A.* 115, 11994–11999. <https://doi.org/10.1073/pnas.1811269115>
- Gonzalez-Pimentel, J.L., Miller, A.Z., Jurado, V., Laiz, L., Pereira, M.F.C., Saiz-Jimenez, C., 2018. Yellow coloured mats from lava tubes of La Palma (Canary Islands, Spain) are dominated by metabolically active Actinobacteria. *Sci. Rep.* 8, 1–11. <https://doi.org/10.1038/s41598-018-20393-2>
- Gross, M., 2019. Permafrost thaw releases problems. *Curr. Biol.* 29, R39–R41. <https://doi.org/10.1016/j.cub.2018.12.045>
- Groth, I., Vettermann, R., Schuetze, B., Schumann, P., Saiz-Jimenez, C., 1999. Actinomycetes in Karstic caves of northern Spain (Altamira and Tito Bustillo). *J. Microbiol. Methods* 36, 115–122. [https://doi.org/10.1016/S0167-7012\(99\)00016-0](https://doi.org/10.1016/S0167-7012(99)00016-0)
- Gtari, M., Ghodhbane-Gtari, F., Nouioui, I., Ktari, A., Hezbri, K., Mimouni, W., Sbissi, I., Ayari, A., Yamanaka, T., Normand, P., Tisa, L.S., Boudabous, A., 2015. Cultivating the uncultured: Growing the recalcitrant

- cluster-2 *Frankia* strains. *Sci. Rep.* 5. <https://doi.org/10.1038/srep13112>
- Hamed, J., Kafshnouchi, M., Ranjbaran, M., 2019. A Study on actinobacterial diversity of Hampoeil cave and screening of their biological activities. *Saudi J. Biol. Sci.* 26, 1587–1595. <https://doi.org/10.1016/j.sjbs.2018.10.010>
- Hansen, A.A., Herbert, R.A., Mikkelsen, K., Jensen, L.L., Kristoffersen, T., Tiedje, J.M., Lomstein, B.A., Finster, K.W., 2007. Viability, diversity and composition of the bacterial community in a high Arctic permafrost soil from Spitsbergen, Northern Norway. *Environ. Microbiol.* 9, 2870–2884. <https://doi.org/10.1111/j.1462-2920.2007.01403.x>
- Hershey, O.S., Barton, H.A., 2018. *The Microbial Diversity of Caves*. Springer, Cham, pp. 69–90. https://doi.org/10.1007/978-3-319-98852-8_5
- Hershey, O.S., Kallmeyer, J., Wallace, A., Barton, M.D., Barton, H.A., 2018. High microbial diversity despite extremely low biomass in a deep karst aquifer. *Front. Microbiol.* 9. <https://doi.org/10.3389/fmicb.2018.02823>
- Hirsch, Penny R., Mauchline, T.H., Clark, I.M., 2010. Culture-independent molecular techniques for soil microbial ecology. *Soil Biol. Biochem.* 42, 878–887. <https://doi.org/10.1016/J.SOILBIO.2010.02.019>
- Hirsch, Penny R., Mauchline, T.H., Clark, I.M., 2010. Culture-independent molecular techniques for soil microbial ecology. *Soil Biol. Biochem.* 42, 878–887. <https://doi.org/10.1016/j.soilbio.2010.02.019>
- Holmfeldt, K., Dziallas, C., Titelman, J., Pohlmann, K., Grossart, H.-P., Riemann, L., 2009. Diversity and abundance of freshwater *Actinobacteria* along environmental gradients in the brackish northern Baltic Sea. *Environ. Microbiol.* 11, 2042–2054. <https://doi.org/10.1111/j.1462-2920.2009.01925.x>
- Hoyos, M., Cañaveras, J.C., Sánchez-Moral, S., Sanz-Rubio, E., Soler, V., 1998. Microclimatic characterization of a karstic cave: Human impact on microenvironmental parameters of a prehistoric rock art cave (Candamo Cave, northern Spain). *Environ. Geol.* 33, 231–242. <https://doi.org/10.1007/s002540050242>
- Jarvis, P., Rey, A., Petsikos, C., Wingate, L., Rayment, M., Pereira, J., Banza, J., David, J., Miglietta, F., Borghetti, M., Manca, G., Valentini, R., 2007. Drying and wetting of Mediterranean soils stimulates decomposition and carbon dioxide emission: The “Birch effect,” in: *Tree Physiology*. Heron Publishing, pp. 929–940. <https://doi.org/10.1093/treephys/27.7.929>
- Jenkins, S.N., Waite, I.S., Blackburn, A., Husband, R., Rushton, S.P., Manning, D.C., O'Donnell, A.G., 2009. Actinobacterial community dynamics in long term managed grasslands. *Antonie van Leeuwenhoek, Int. J. Gen. Mol. Microbiol.* 95, 319–334. <https://doi.org/10.1007/s10482-009-9317-8>
- Jiang, J.-R., Cai, L., Liu, F., 2017. Oligotrophic fungi from a carbonate cave, with three new species of *Cephalotrichum*. *Mycology* 8, 164–177. <https://doi.org/10.1080/21501203.2017.1366370>
- Jin, V.L., Haney, R.L., Fay, P.A., Polley, H.W., 2013. Soil type and moisture regime control microbial C and N mineralization in grassland soils more than atmospheric CO₂-induced changes in litter quality. *Soil Biol. Biochem.* 58, 172–180. <https://doi.org/10.1016/j.soilbio.2012.11.024>
- Johnston, M.D., Muench, B.A., Banks, E.D., Barton, H.A., 2012. Human urine in Lechuguilla cave: the microbiological impact and potential for bioremediation. *J. Cave Karst Stud.* 74, 278–291. <https://doi.org/10.4311/2011MB0227>
- Johnston, S.R., Boddy, L., Weightman, A.J., 2016. Bacteria in decomposing wood and their interactions with wood-decay fungi. *FEMS Microbiol. Ecol.* 92, fiw179. <https://doi.org/10.1093/femsec/fiw179>
- Jurado, V., Boiron, P., Kroppenstedt, R.M., Laurent, F., Couble, A., Laiz, L., Klenk, H.-P., Gonzalez, J.M., Saiz-Jimenez, C., Mouniee, D., Bergeron, E., Rodriguez-Nava, V., 2008. *Nocardia altamirensis* sp. nov., isolated from Altamira cave, Cantabria, Spain. *Int. J. Syst. Evol. Microbiol.* 58, 2210–2214. <https://doi.org/10.1099/ijs.0.65482-0>
- Jurado, V., Kroppenstedt, R.M., Saiz-Jimenez, C., Klenk, H.P., Mouniée, D., Laiz, L., Couble, A., Pötter, G., Boiron, P., Rodríguez-Nava, V., 2009. *Hoyosella altamirensis* gen. nov., sp. nov., a new member of the order Actinomycetales isolated from a cave biofilm. *Int. J. Syst. Evol. Microbiol.* 59, 3105–3110.

<https://doi.org/10.1099/ij.s.0.008664-0>

- Jurado, V., Laiz, L., Rodriguez-Nava, V., Boiron, P., Hermosin, B., Sanchez-Moral, S., Saiz-Jimenez, C., 2010. Pathogenic and opportunistic microorganisms in caves. *Int. J. Speleol.* 39 (1), 15–24.
- Kaisermann, A., Maron, P.A., Beaumelle, L., Lata, J.C., 2015. Fungal communities are more sensitive indicators to non-extreme soil moisture variations than bacterial communities. *Appl. Soil Ecol.* 86, 158–164. <https://doi.org/10.1016/j.apsoil.2014.10.009>
- Karunaratna, S.C., Dong, Y., Karasaki, S., Tibpromma, S., Hyde, K.D., Lumyong, S., Xu, J., Sheng, J., Mortimer, P.E., 2020. Discovery of novel fungal species and pathogens on bat carcasses in a cave in Yunnan Province, China. *Emerg. Microbes Infect.* 9, 1554–1566. <https://doi.org/10.1080/22221751.2020.1785333>
- Kielak, A.M., Scheublin, T.R., Mendes, L.W., van Veen, J.A., Kuramae, E.E., 2016. Bacterial community succession in pine-wood decomposition. *Front. Microbiol.* 7, 231. <https://doi.org/10.3389/fmicb.2016.00231>
- Kopecky, J., Kyselkova, M., Omelka, M., Cermak, L., Novotna, J., Grundmann, G.L., Moëgne-Loccoz, Y., Sagova-Mareckova, M., 2011. Actinobacterial community dominated by a distinct clade in acidic soil of a waterlogged deciduous forest. *FEMS Microbiol. Ecol.* 78, 386–394. <https://doi.org/10.1111/j.1574-6941.2011.01173.x>
- Kumar De, A., 2019. Profiling bacterial diversity of B2 Cave, a limestone cave of Baratang, Andaman and Nicobar islands, India. *Proc. Indian Natl. Sci. Acad.* 85, 853–862. <https://doi.org/10.16943/ptinsa/2019/49589>
- Kurtböke, D., 2017. Ecology and habitat distribution of actinobacteria, in: *Biology and Biotechnology of Actinobacteria*. Springer International Publishing, pp. 123–149. https://doi.org/10.1007/978-3-319-60339-1_6
- Lacombe-Harvey, M.È., Brzezinski, R., Beaulieu, C., 2018. Chitinolytic functions in actinobacteria: ecology, enzymes, and evolution. *Appl. Microbiol. Biotechnol.* 102, 7219–7230. <https://doi.org/10.1007/s00253-018-9149-4>
- Lange, M., Eisenhauer, N., Sierra, C.A., Bessler, H., Engels, C., Griffiths, R.I., Mellado-Vázquez, P.G., Malik, A.A., Roy, J., Scheu, S., Steinbeiss, S., Thomson, B.C., Trumbore, S.E., Gleixner, G., 2015. Plant diversity increases soil microbial activity and soil carbon storage. *Nat. Commun.* 6, 6707. <https://doi.org/10.1038/ncomms7707>
- Lavoie, K.H., Winter, A.S., Read, K.J.H., Hughes, E.M., Spilde, M.N., Northup, D.E., 2017. Comparison of bacterial communities from lava cave microbial mats to overlying surface soils from Lava Beds National Monument, USA. *PLoS One* 12, e0169339. <https://doi.org/10.1371/journal.pone.0169339>
- Le Roes-Hill, M., Prins, A., 2016. Biotechnological Potential of Oxidative Enzymes from Actinobacteria, in: *Actinobacteria - Basics and Biotechnological Applications*. pp. 200–226. <https://doi.org/10.5772/61321>
- Lee, H., Fitzgerald, J., Hewins, D.B., McCulley, R.L., Archer, S.R., Rahn, T., Throop, H.L., 2014. Soil moisture and soil-litter mixing effects on surface litter decomposition: A controlled environment assessment. *Soil Biol. Biochem.* 72, 123–132. <https://doi.org/10.1016/j.soilbio.2014.01.027>
- Lee, I.S., Kim, O.K., Chang, Y.Y., Bae, B., Kim, H.H., Baek, K.H., 2002. Heavy metal concentrations and enzyme activities in soil from a contaminated Korean shooting range. *J. Biosci. Bioeng.* 94, 406–411. [https://doi.org/10.1016/S1389-1723\(02\)80217-1](https://doi.org/10.1016/S1389-1723(02)80217-1)
- Lee, S.D., 2006. *Nocardia jejuensis* sp. nov., a novel actinomycete isolated from a natural cave on Jeju Island, Republic of Korea. *Int. J. Syst. Evol. Microbiol.* 56, 559–562. <https://doi.org/10.1099/ij.s.0.63866-0>
- Leibold, M., McPeck, M., 2006. Coexistence of the niche and neutral perspectives in community ecology. *Ecology* 87, 1399–1410.
- Leuko, S., Koskinen, K., Sanna, L., D'Angeli, I.M., De Waele, J., Marcia, P., Moissl-Eichinger, C., Rettberg, P., 2017. The influence of human exploration on the microbial community structure and ammonia oxidizing

- potential of the Su Bentu limestone cave in Sardinia, Italy. *PLoS One* 12, e0180700. <https://doi.org/10.1371/journal.pone.0180700>
- Lewis, K., 2009. *Persisters, Biofilms, and the Problem of Cultivability*. Springer, Berlin, Heidelberg, pp. 203–216. https://doi.org/10.1007/978-3-540-85465-4_7
- Li, H.R., Yu, Y., Luo, W., Zeng, Y.X., 2010. *Marisediminicola antarctica* gen. nov., sp. nov., an actinobacterium isolated from the Antarctic. *Int. J. Syst. Evol. Microbiol.* 60, 2535–2539. <https://doi.org/10.1099/ijs.0.018754-0>
- Lladó, S., López-Mondéjar, R., Baldrian, P., 2017. Forest Soil Bacteria: Diversity, Involvement in Ecosystem Processes, and Response to Global Change. *Microbiol. Mol. Biol. Rev.* 81, e00063-16. <https://doi.org/10.1128/MMBR.00063-16>
- López-Mondéjar, R., Brabcová, V., Štursová, M., Davidová, A., Jansa, J., Cajthaml, T., Baldrian, P., 2018. Decomposer food web in a deciduous forest shows high share of generalist microorganisms and importance of microbial biomass recycling. *ISME J.* 12, 1768–1778. <https://doi.org/10.1038/s41396-018-0084-2>
- Maciejewska, M., Adam, D., Naômé, A., Martinet, L., Tenconi, E., Całusińska, M., Delfosse, P., Hanikenne, M., Baurain, D., Compère, P., Carnol, M., Barton, H.A., Rigali, S., 2017. Assessment of the Potential Role of *Streptomyces* in Cave Moonmilk Formation. *Front. Microbiol.* 8, 1181. <https://doi.org/10.3389/fmicb.2017.01181>
- Makhalanyane, T.P., Valverde, A., Gunnigle, E., Frossard, A., Ramond, J.-B., Cowan, D.A., 2015. Microbial ecology of hot desert edaphic systems. *FEMS Microbiol. Rev.* 39, 203–221. <https://doi.org/10.1093/femsre/fuu011>
- Malik, A.A., Chowdhury, S., Schlager, V., Oliver, A., Puissant, J., Vazquez, P.G.M., Jehmlich, N., von Bergen, M., Griffiths, R.I., Gleixner, G., 2016. Soil Fungal:Bacterial Ratios Are Linked to Altered Carbon Cycling. *Front. Microbiol.* 7, 1247. <https://doi.org/10.3389/fmicb.2016.01247>
- Manivasagan, P., Venkatesan, J., Sivakumar, K., Kim, S.K., 2013. Actinobacterial melanins: Current status and perspective for the future. *World J. Microbiol. Biotechnol.* 29, 1737–1750. <https://doi.org/10.1007/s11274-013-1352-y>
- Martens, M., Dawyndt, P., Coopman, R., Gillis, M., De Vos, P., Willems, A., 2008. Advantages of multilocus sequence analysis for taxonomic studies: A case study using 10 housekeeping genes in the genus *Ensifer* (including former *Sinorhizobium*). *Int. J. Syst. Evol. Microbiol.* 58, 200–214. <https://doi.org/10.1099/ijs.0.65392-0>
- Martin-Sanchez, P.M., Miller, A.Z., Saiz-Jimenez, C., 2015. Lascaux Cave: An Example of Fragile Ecological Balance in Subterranean Environments. *Microb. Life Cave Syst.* <https://doi.org/10.1515/9783110339888-015>
- McDaniel, M.D., Grandy, A.S., Tiemann, L.K., Weintraub, M.N., 2014. Crop rotation complexity regulates the decomposition of high and low quality residues. *Soil Biol. Biochem.* 78, 243–254. <https://doi.org/10.1016/J.SOILBIO.2014.07.027>
- McGuire, K.L., Treseder, K.K., 2010. Microbial communities and their relevance for ecosystem models: Decomposition as a case study. *Soil Biol. Biochem.* 42, 529–535. <https://doi.org/10.1016/j.soilbio.2009.11.016>
- McNamara, C.J., Perry IV, T.D., Bearce, K.A., Hernandez-Duque, G., Mitchell, R., 2006. Epilithic and endolithic bacterial communities in limestone from a Maya archaeological site. *Microb. Ecol.* 51, 51–64. <https://doi.org/10.1007/s00248-005-0200-5>
- Meidute, S., Demoling, F., Bååth, E., 2008. Antagonistic and synergistic effects of fungal and bacterial growth in soil after adding different carbon and nitrogen sources. *Soil Biol. Biochem.* 40, 2334–2343. <https://doi.org/10.1016/J.SOILBIO.2008.05.011>
- Meier, C.L., Bowman, W.D., 2008. Links between plant litter chemistry, species diversity, and below-ground ecosystem function. *Proc. Natl. Acad. Sci. U. S. A.* 105, 19780–19785.

<https://doi.org/10.1073/pnas.0805600105>

- Meisner, A., Jacquioud, S., Snoek, B.L., Hooven, F.C., Freedman, Z.B., 2018. Drought Legacy Effects on the Composition of Soil Fungal and Prokaryote Communities 9, 1–12. <https://doi.org/10.3389/fmicb.2018.00294>
- Miao, V., Davies, J., 2010. Actinobacteria: The good, the bad, and the ugly, in: Antonie van Leeuwenhoek, International Journal of General and Molecular Microbiology. Springer, pp. 143–150. <https://doi.org/10.1007/s10482-010-9440-6>
- Mihajlovski, A., Gabarre, A., Seyer, D., Bousta, F., Di Martino, P., 2017. Bacterial diversity on rock surface of the ruined part of a French historic monument: The Chaalis abbey. Int. Biodeterior. Biodegrad. 120, 161–169. <https://doi.org/10.1016/j.ibiod.2017.02.019>
- Mitova, M., Ivanova, M., Iliev, M., 2017. ACTINOBACTERIA AS A COMPONENT OF MICROBIAL COMMUNITY FORMED ON ROCK PAINTINGS OF MAGURA CAVE, BULGARIA 102, 127–138.
- Modra, H., Bartos, M., Hribova, P., Ulmann, V., Hubelova, D., Konecny, O., Gersl, M., Kudelka, J., Voros, D., Pavlik, I., 2017. Detection of mycobacteria in the environment of the Moravian Karst (Bull Rock Cave and the relevant water catchment area): the impact of water sediment, earthworm castings and bat guano. Vet. Med. (Praha). 62 (2017), 153–168. <https://doi.org/10.17221/126/2016-VETMED>
- Moinet, G.Y.K., Moinet, M., Hunt, J.E., Rumpel, C., Chabbi, A., Millard, P., 2020. Temperature sensitivity of decomposition decreases with increasing soil organic matter stability. Sci. Total Environ. 704, 135460. <https://doi.org/10.1016/j.scitotenv.2019.135460>
- Moldovan, O.T., Bercea, S., Năstase-Bucur, R., Constantin, S., Kenesz, M., Mirea, I.C., Petculescu, A., Robu, M., Arghir, R.A., 2020. Management of water bodies in show caves – A microbial approach. Tour. Manag. 78, 104037. <https://doi.org/10.1016/j.tourman.2019.104037>
- Moorhead, D.L., Sinsabaugh, R.L., 2006. A theoretical model of litter decay and microbial interaction. Ecol. Monogr. 76, 151–174. [https://doi.org/10.1890/0012-9615\(2006\)076\[0151:ATMOLD\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2006)076[0151:ATMOLD]2.0.CO;2)
- Mulec, J., Oarga-Mulec, A., Šturm, S., Tomazin, R., Matos, T., 2017. Spacio-Temporal Distribution and Tourist Impact on Airborne Bacteria in a Cave (Škocjan Caves, Slovenia). Diversity 9, 28. <https://doi.org/10.3390/d9030028>
- Mulec, J., Oarga-Mulec, A., Tomazin, R., Matos, T., 2015. Characterization and fluorescence of yellow biofilms in karst caves, southwest Slovenia. Int. J. Speleol. 44, 107–114. <https://doi.org/10.5038/1827-806X.44.2.1>
- Newton, R.J., Jones, S.E., Helmus, M.R., McMahon, K.D., 2007. Phylogenetic ecology of the freshwater Actinobacteria acI lineage. Appl. Environ. Microbiol. 73, 7169–7176. <https://doi.org/10.1128/AEM.00794-07>
- Nouioui, I., Ghodhbane-Gtari, F., Jando, M., Tisa, L.S., Klenk, H.P., Gtari, M., 2019. Frankia torreyi sp. nov., the first actinobacterium of the genus Frankia Brunchorst 1886, 174 AL isolated in axenic culture. Antonie van Leeuwenhoek, Int. J. Gen. Mol. Microbiol. 112, 57–65. <https://doi.org/10.1007/s10482-018-1131-8>
- Nováková, A., 2009. Microscopic fungi isolated from the Domica Cave system (Slovak Karst National Park, Slovakia). A review. Int. J. Speleol. 38, 71–82. <https://doi.org/10.5038/1827-806X.38.1.8>
- Ogier, J.C., Pagès, S., Galan, M., Barret, M., Gaudriault, S., 2019. RpoB, a promising marker for analyzing the diversity of bacterial communities by amplicon sequencing. BMC Microbiol. 19, 2–16. <https://doi.org/10.1186/s12866-019-1546-z>
- Ortiz, M., Legatzki, A., Neilson, J.W., Fryslie, B., Nelson, W.M., Wing, R.A., Soderlund, C.A., Pryor, B.M., Maier, R.M., 2014. Making a living while starving in the dark: metagenomic insights into the energy dynamics of a carbonate cave. ISME J. 8, 478–491. <https://doi.org/10.1038/ismej.2013.159>
- Paula, C.C.P.D., Montoya, Q. V., Meirelles, L.A., Farinas, C.S., Rodrigues, A., Seleghim, M.H.R., 2019. High cellulolytic activities in filamentous fungi isolated from an extreme oligotrophic subterranean environment (Catão cave) in Brazil. An. Acad. Bras. Cienc. 91, 9–11. <https://doi.org/10.1590/0001-3765201920180583>

- Pei, A.Y., Oberdorf, W.E., Nossa, C.W., Agarwal, A., Chokshi, P., Gerz, E.A., Jin, Z., Lee, P., Yang, L., Poles, M., Brown, S.M., Sotero, S., DeSantis, T., Brodie, E., Nelson, K., Pei, Z., 2010. Diversity of 16S rRNA genes within individual prokaryotic genomes. *Appl. Environ. Microbiol.* 76, 3886–3897. <https://doi.org/10.1128/AEM.02953-09>
- Pereira, S., Burešová, A., Kopecky, J., Mádrová, P., Aupic-Samain, A., Fernandez, C., Baldy, V., Sagova-Mareckova, M., 2019. Litter traits and rainfall reduction alter microbial litter decomposers: The evidence from three Mediterranean forests. *FEMS Microbiol. Ecol.* 95, fiz168. <https://doi.org/10.1093/femsec/fiz168>
- Philippot, L., Andersson, S.G.E., Battin, T.J., Prosser, J.I., Schimel, J.P., Whitman, W.B., Hallin, S., 2010. The ecological coherence of high bacterial taxonomic ranks. *Nat. Rev. Microbiol.* 8, 523–529. <https://doi.org/10.1038/nrmicro2367>
- Polkade, A. V., Mantri, S.S., Patwekar, U.J., Jangid, K., 2016. Quorum Sensing: An Under-Explored Phenomenon in the Phylum Actinobacteria. *Front. Microbiol.* 7, 131. <https://doi.org/10.3389/fmicb.2016.00131>
- Popović, S., Simić, G.S., Stupar, M., Unković, N., Predojević, D., Jovanović, J., Grbić, M.L., 2015. Cyanobacteria, algae and microfungi present in biofilm from Božana Cave (Serbia). *Int. J. Speleol.* 44, 141–149. <https://doi.org/10.5038/1827-806X.44.2.4>
- Porca, E., Jurado, V., Žgur-Bertok, D., Saiz-Jimenez, C., Pašić, L., 2012. Comparative analysis of yellow microbial communities growing on the walls of geographically distinct caves indicates a common core of microorganisms involved in their formation. *FEMS Microbiol. Ecol.* 81, 255–266. <https://doi.org/10.1111/j.1574-6941.2012.01383.x>
- Portillo, M.C., Gonzalez, J.M., 2011. Moonmilk Deposits Originate from Specific Bacterial Communities in Altamira Cave (Spain). *Microb. Ecol.* 61, 182–189. <https://doi.org/10.1007/s00248-010-9731-5>
- Portillo, M.C., Gonzalez, J.M., 2010. Differential effects of distinct bacterial biofilms in a cave environment. *Curr. Microbiol.* 60, 435–438. <https://doi.org/10.1007/s00284-009-9561-1>
- Pose-Juan, E., Igual, J.M., Sánchez-Martín, M.J., Rodríguez-Cruz, M.S., 2017. Influence of Herbicide Triasulfuron on Soil Microbial Community in an Unamended Soil and a Soil Amended with Organic Residues. *Front. Microbiol.* 8, 378. <https://doi.org/10.3389/fmicb.2017.00378>
- Prieto, I., Almagro, M., Bastida, F., Querejeta, J.I., 2019. Altered leaf litter quality exacerbates the negative impact of climate change on decomposition. *J. Ecol.* 1365-2745.13168. <https://doi.org/10.1111/1365-2745.13168>
- Purahong, W., Krüger, D., Buscot, F., Wubet, T., 2016a. Correlations between the composition of modular fungal communities and litter decomposition-associated ecosystem functions. *Fungal Ecol.* 22, 106–114. <https://doi.org/10.1016/J.FUNECO.2016.04.009>
- Purahong, W., Schlöter, M., Pecyna, M.J., Kapturska, D., Däumlich, V., Mital, S., Buscot, F., Hofrichter, M., Gutknecht, J.L.M., Krüger, D., 2014. Uncoupling of microbial community structure and function in decomposing litter across beech forest ecosystems in Central Europe. *Sci. Rep.* 4, 7014. <https://doi.org/10.1038/srep07014>
- Purahong, W., Wubet, T., Lentendu, G., Schlöter, M., Pecyna, M.J., Kapturska, D., Hofrichter, M., Krüger, D., Buscot, F., 2016b. Life in leaf litter: novel insights into community dynamics of bacteria and fungi during litter decomposition. *Mol. Ecol.* 25, 4059–4074. <https://doi.org/10.1111/mec.13739>
- Radha, T.K., Rao, D.L.N., Sreeramulu, K.R., 2017. Actinobacteria of arid and semi-arid soils: Antagonism to fungal pathogens and plant growth promoting potential. *J. Pure Appl. Microbiol.* 11, 1045–1052. <https://doi.org/10.22207/JPAM.11.2.47>
- Rangseekaew, P., Pathom-Aree, W., 2019. Cave Actinobacteria as Producers of Bioactive Metabolites. *Front. Microbiol.* 10, 387. <https://doi.org/10.3389/fmicb.2019.00387>
- Rao, M.P.N., Dong, Z.Y., Zhang, H., Niu, X.K., Zhang, K., Fang, B.Z., Xiao, M., Kang, Y.Q., Li, W.J., 2019. *Bacillus antri* sp. Nov., isolated from cave soil. *Int. J. Syst. Evol. Microbiol.* 69, 2335–2339.

<https://doi.org/10.1099/ijsem.0.003473>

- Rapoport, D., Sagova-Mareckova, M., Sedláček, I., Provazník, J., Králová, S., Pavlinic, D., Benes, V., Kopecky, J., 2020. *Trebonia kvetii* gen. nov., sp. nov., an acidophilic actinobacterium, and proposal of the new actinobacterial family treboniaceae fam. nov. *Int. J. Syst. Evol. Microbiol.* 70, 5106–5114. <https://doi.org/10.1099/ijsem.0.004388>
- Rasuk, M.C., Ferrer, G.M., Kurth, D., Portero, L.R., Fariás, M.E., Albarracín, V.H., 2017. UV-Resistant Actinobacteria from High-Altitude Andean Lakes: Isolation, Characterization and Antagonistic Activities. *Photochem. Photobiol.* 93, 865–880. <https://doi.org/10.1111/php.12759>
- Rego, A., Raio, F., Martins, T.P., Ribeiro, H., Sousa, A.G.G., Séneca, J., Baptista, M.S., Lee, C.K., Craig Cary, S., Ramos, V., Carvalho, M.F., Leão, P.N., Magalhães, C., 2019. Actinobacteria and cyanobacteria diversity in terrestrial Antarctic microenvironments evaluated by culture-dependent and independent methods. *Front. Microbiol.* 10, 1018. <https://doi.org/10.3389/fmicb.2019.01018>
- Rodríguez-Nava, V., Couble, A., Devulder, G., Flandrois, J.-P., Boiron, P., Laurent, F., 2006. Use of PCR-restriction enzyme pattern analysis and sequencing database for hsp65 gene-based identification of *Nocardia* species. *J. Clin. Microbiol.* 44, 536–46. <https://doi.org/10.1128/JCM.44.2.536-546.2006>
- Rodríguez-Nava, V., Khan, Z.U., Pötter, G., Kroppenstedt, R.M., Boiron, P., Laurent, F., 2007. *Nocardia coubleae* sp. nov., isolated from oil-contaminated Kuwaiti soil. <https://doi.org/10.1099/ijms.0.64815-0>
- Rosenblatt, A.E., 2018. Shifts in plant nutrient content in combined warming and drought scenarios may alter reproductive fitness across trophic levels. *Oikos* 127, 1853–1862. <https://doi.org/10.1111/oik.05272>
- Rousk, J., Demoling, L.A., Bahr, A., Bååth, E., 2008. Examining the fungal and bacterial niche overlap using selective inhibitors in soil. *FEMS Microbiol. Ecol.* 63, 350–358. <https://doi.org/10.1111/j.1574-6941.2008.00440.x>
- Rousk, J., Frey, S.D., 2015. Revisiting the hypothesis that fungal-to-bacterial dominance characterizes turnover of soil organic matter and nutrients. *Ecol. Monogr.* 85, 457–472. <https://doi.org/10.1890/14-1796.1>
- Roxburgh, S.H., Shea, K., Wilson, J.B., 2004. The intermediate disturbance hypothesis: Patch dynamics and mechanisms of species coexistence. *Ecology* 85, 359–371. <https://doi.org/10.1890/03-0266>
- Sagova-Mareckova, M., Cermak, L., Omelka, M., Kyselkova, M., Kopecky, J., 2015. Bacterial diversity and abundance of a creek valley sites reflected soil pH and season. *Open Life Sci.* 10, 61–70.
- Sagova-Mareckova, Marketa, Daniel, O., Omelka, M., Kristufek, V., Divis, J., Kopecky, J., 2015. Determination of Factors Associated with Natural Soil Suppressivity to Potato Common Scab. <https://doi.org/10.1371/journal.pone.0116291>
- Sagova-Mareckova, M., Omelka, M., Cermak, L., Kamenik, Z., Olsovska, J., Hackl, E., Kopecky, J., Hadacek, F., 2011. Microbial communities show parallels at sites with distinct litter and soil characteristics. *Appl. Environ. Microbiol.* 77, 7560–7567. <https://doi.org/10.1128/AEM.00527-11>
- Schabereiter-Gurtner, C., Saiz-Jimenez, C., Pinar, G., Lubitz, W., Rölleke, S., 2002. Altamira cave Paleolithic paintings harbor partly unknown bacterial communities. *FEMS Microbiol. Lett.* 211, 7–11. <https://doi.org/10.1111/j.1574-6968.2002.tb11195.x>
- Schneider, T., Keiblinger, K.M., Schmid, E., Sterflinger-Gleixner, K., Ellersdorfer, G., Roschitzki, B., Richter, A., Eberl, L., Zechmeister-Boltenstern, S., Riedel, K., 2012. Who is who in litter decomposition? Metaproteomics reveals major microbial players and their biogeochemical functions. *ISME J.* 6, 1749–1762. <https://doi.org/10.1038/ismej.2012.11>
- Schobben, M., Stebbins, A., Ghaderi, A., Strauss, H., Korn, D., Korte, C., 2016. Eutrophication, microbial-sulfate reduction and mass extinctions. *Commun. Integr. Biol.* <https://doi.org/10.1080/19420889.2015.1115162>
- Schwartz, E., Adair, K.L., Schuur, E. a., 2007. Bacterial community structure correlates with decomposition parameters along a Hawaiian precipitation gradient. *Soil Biol. Biochem.* 39, 2164–2167. <https://doi.org/10.1016/j.soilbio.2007.02.013>

- Sharma, M., Dangi, P., Choudhary, M., 2014. Actinomycetes: source, identification, and their applications. *Int. J. Curr. Microbiol. Appl. Sci.* 3, 801–832.
- Shi, S., Tian, L., Nasir, F., Bahadur, A., Batool, A., Luo, S., Yang, F., Wang, Z., Tian, C., 2019. Response of microbial communities and enzyme activities to amendments in saline-alkaline soils. *Appl. Soil Ecol.* 135, 16–24. <https://doi.org/10.1016/j.apsoil.2018.11.003>
- Smith, W.K., Gao, W., Steltzer, H., Wallenstein, M.D., Tree, R., 2010. Moisture availability influences the effect of ultraviolet-B radiation on leaf litter decomposition. *Glob. Chang. Biol.* 16, 484–495. <https://doi.org/10.1111/j.1365-2486.2009.01973.x>
- Šnajdr, J., Cajthaml, T., Valášková, V., Merhautová, V., Petránková, M., Spetz, P., Leppänen, K., Baldrian, P., 2011. Transformation of *Quercus petraea* litter: Successive changes in litter chemistry are reflected in differential enzyme activity and changes in the microbial community composition. *FEMS Microbiol. Ecol.* 75, 291–303. <https://doi.org/10.1111/j.1574-6941.2010.00999.x>
- Song, F., Fan, X., Song, R., 2010. Review of mixed forest litter decomposition researches. *Acta Ecol. Sin.* 30, 221–225. <https://doi.org/10.1016/j.chnaes.2010.06.006>
- Šťovíček, A., Azatyan, A., Soares, M.I.M., Gillor, O., 2017. The Impact of Hydration and Temperature on Bacterial Diversity in Arid Soil Mesocosms. *Front. Microbiol.* 8, 1078. <https://doi.org/10.3389/fmicb.2017.01078>
- Suseela, V., Tharayil, N., Xing, B., Dukes, J.S., 2014. Warming alters potential enzyme activity but precipitation regulates chemical transformations in grass litter exposed to simulated climatic changes. *Soil Biol. Biochem.* 75, 102–112. <https://doi.org/10.1016/j.soilbio.2014.03.022>
- Szaková, J., Burešová, A., Praus, L., García-Sánchez, M., Holečková, Z., Gabriel, J., Sysalová, J., Červenka, R., Komárek, J., Grohová, S., Tlustoš, P., 2016. The response of mercury (Hg) transformation in soil to sulfur compounds and sulfur-rich biowaste application. *Environ. Earth Sci.* 75, 584. <https://doi.org/10.1007/s12665-016-5387-x>
- Taj-Aldeen, S.J., Deshmukh, A., Doiphode, S., Wahab, A.A., Allangawi, M., Muzrkehi, A. Al, Klaassen, C.H., Meis, J.J., 2013. Molecular Identification and Susceptibility Pattern of Clinical *Nocardia* Species: Emergence of *Nocardia crassostreae* as an Agent of Invasive Nocardiosis. *Can. J. Infect. Dis. Med. Microbiol.* 24, e33–e38.
- Takeda, K., Kang, Y., Yazawa, K., Gono, T., Mikami, Y., 2010. Phylogenetic studies of *Nocardia* species based on *gyrB* gene analyses. *J. Med. Microbiol.* 59, 165–171. <https://doi.org/10.1099/jmm.0.011346-0>
- Telenti, A., Marchesi, F., Balz, M., Bally, F., Bottger, E.C., Bodmer, T., 1993. Rapid identification of mycobacteria to the species level by polymerase chain reaction and restriction enzyme analysis. *J. Clin. Microbiol.* 31, 175–178.
- Thiessen, S., Gleixner, G., Wutzler, T., Reichstein, M., 2013. Both priming and temperature sensitivity of soil organic matter decomposition depend on microbial biomass - An incubation study. *Soil Biol. Biochem.* 57, 739–748. <https://doi.org/10.1016/j.soilbio.2012.10.029>
- Tikhonova, E.N., Men'ko, E. V., Ulanova, R. V., Li, H., Kravchenko, I.K., 2019. Effect of Temperature on the Taxonomic Structure of Soil Bacterial Communities during Litter Decomposition. *Microbiol. (Russian Fed.)* 88, 781–785. <https://doi.org/10.1134/S0026261719060195>
- Tlaskal, V., Voříšková, J., Baldrian, P., 2016. Bacterial succession on decomposing leaf litter exhibits a specific occurrence pattern of cellulolytic taxa and potential decomposers of fungal mycelia. *FEMS Microbiol. Ecol.* 92, fiw177. <https://doi.org/10.1093/femsec/fiw177>
- To, K., Cao, R., Yegiazaryan, A., Owens, J., Venketaraman, V., 2020. General Overview of Nontuberculous Mycobacteria Opportunistic Pathogens: *Mycobacterium avium* and *Mycobacterium abscessus*. *J. Clin. Med.* 9, 2541. <https://doi.org/10.3390/jcm9082541>
- Tomczyk-Żak, K., Zielenkiewicz, U., 2015. Microbial Diversity in Caves. *Geomicrobiol. J.* 1–19. <https://doi.org/10.1080/01490451.2014.1003341>

- Urzi, C., De Leo, F., Bruno, L., Albertano, P., 2010. Microbial Diversity in Paleolithic Caves: A Study Case on the Phototrophic Biofilms of the Cave of Bats (Zuheros, Spain). *Microb. Ecol.* 60, 116–129. <https://doi.org/10.1007/s00248-010-9710-x>
- Vanderwolf, K.J., Malloch, D., McAlpine, D.F., Forbes, G.J., 2013. A world review of fungi, yeasts, and slime molds in caves Karen. *Int. J. Speleol.* 42, 77–96. <https://doi.org/10.5038/1827-806X.42.1.9>
- Vázquez-Boland, J.A., Meijer, W.G., 2019. The pathogenic actinobacterium *Rhodococcus equi*: what's in a name? *Mol. Microbiol.* 112, 1–15. <https://doi.org/10.1111/mmi.14267>
- Verma, M., Lal, D., Kaur, Jaspreet, Saxena, A., Kaur, Jasvinder, Anand, S., Lal, R., 2013. Phylogenetic analyses of phylum Actinobacteria based on whole genome sequences. *Res. Microbiol.* 164, 718–728. <https://doi.org/10.1016/j.resmic.2013.04.002>
- Větrovský, T., Baldrian, P., 2013. The Variability of the 16S rRNA Gene in Bacterial Genomes and Its Consequences for Bacterial Community Analyses. *PLoS One* 8, e57923. <https://doi.org/10.1371/journal.pone.0057923>
- Větrovský, T., Steffen, K.T., Baldrian, P., 2014. Potential of Cometary Transformation of Polysaccharides and Lignin in Lignocellulose by Soil Actinobacteria. *PLoS One* 9, e89108. <https://doi.org/10.1371/JOURNAL.PONE.0089108>
- Vitousek, P.M., Porder, S., Houlton, B.Z., Chadwick, O.A., 2010. Terrestrial phosphorus limitation: mechanisms, implications, and nitrogen–phosphorus interactions. *Ecol. Appl.* 20, 5–15. <https://doi.org/10.1890/08-0127.1>
- Vos, M., Quince, C., Pijl, A.S., de Hollander, M., Kowalchuk, G.A., 2012. A comparison of rpoB and 16S rRNA as markers in pyrosequencing studies of bacterial diversity. *PLoS One* 7, e30600. <https://doi.org/10.1371/journal.pone.0030600>
- Wagg, C., Schlaeppli, K., Banerjee, S., Kuramae, E.E., van der Heijden, M.G.A., 2019. Fungal-bacterial diversity and microbiome complexity predict ecosystem functioning. *Nat. Commun.* 10, 1–10. <https://doi.org/10.1038/s41467-019-12798-y>
- Wertz, S., Degrange, V., Prosser, J.I., Poly, F., Commeaux, C., Freitag, T., Guillaumaud, N., Roux, X. Le, 2006. Maintenance of soil functioning following erosion of microbial diversity. *Environ. Microbiol.* 8, 2162–2169. <https://doi.org/10.1111/j.1462-2920.2006.01098.x>
- Woo, H.L., Hazen, T.C., Simmons, B. a., DeAngelis, K.M., 2014. Enzyme activities of aerobic lignocellulolytic bacteria isolated from wet tropical forest soils. *Syst. Appl. Microbiol.* 37, 60–67. <https://doi.org/10.1016/j.syapm.2013.10.001>
- Wu, Y., Tan, L., Liu, W., Wang, B., Wang, J., Cai, Y., Lin, X., 2015. Profiling bacterial diversity in a limestone cave of the western Loess Plateau of China. *Front. Microbiol.* 6, 244. <https://doi.org/10.3389/fmicb.2015.00244>
- Xu, Y., Fan, J., Ding, W., Bol, R., Chen, Z., Luo, J., Bolan, N., 2016. Stage-specific response of litter decomposition to N and S amendments in a subtropical forest soil. *Biol. Fertil. Soils* 52, 711–724. <https://doi.org/10.1007/s00374-016-1115-7>
- Zhang, B., Wu, X., Tai, X., Sun, L., Wu, M., Zhang, W., Chen, X., Zhang, G., Chen, T., Liu, G., Dyson, P., 2019. Variation in Actinobacterial Community Composition and Potential Function in Different Soil Ecosystems Belonging to the Arid Heihe River Basin of Northwest China. *Front. Microbiol.* 10, 2209. <https://doi.org/10.3389/fmicb.2019.02209>
- Zhao, Y., Lu, Q., Wei, Y., Cui, H., Zhang, X., Wang, X., Shan, S., Wei, Z., 2016. Effect of actinobacteria agent inoculation methods on cellulose degradation during composting based on redundancy analysis. *Bioresour. Technol.* 219, 196–203. <https://doi.org/10.1016/j.biortech.2016.07.117>
- Zhu, H.Z., Zhang, Z.F., Zhou, N., Jiang, C.Y., Wang, B.J., Cai, L., Liu, S.J., 2019. Diversity, distribution and co-occurrence patterns of bacterial communities in a karst cave system. *Front. Microbiol.* 10, 1726. <https://doi.org/10.3389/fmicb.2019.01726>

Zhu, Y., Penuelas, J., 2020. Changes in the environmental microbiome in the Anthropocene. *Glob. Chang. Biol.* 26, 3175–3177. <https://doi.org/10.1111/gcb.15086>

Original research papers

-Part 1-

Paper I: Usefulness of *hsp65* marker as a complement to 16S rRNA sequences to demonstrate differences in Actinobacteria community according to anthropization of limestone caves and occurrence of visual marks in Lascaux Cave

Usefulness of *hsp65* marker as a complement to 16S rRNA sequences to demonstrate differences in *Actinobacteria* community according to anthropization of limestone caves and occurrence of visual marks in Lascaux Cave

Buresova Andrea^{1,2,3}, Kopecky Jan³, Sagova-Mareckova Marketa³, Alonso Lise¹, Vautrin Florian¹, Moëgne-Loccoz Yvan¹, Rodriguez-Nava Veronica¹

¹Univ Lyon, Université Claude Bernard Lyon 1, CNRS, INRAE, VetAgro Sup, UMR5557 Ecologie Microbienne, F-69622 Villeurbanne, France,

²Department of Ecology, Faculty of Science, Charles University in Prague, Vinicna 7, Prague, Czech Republic,

³Laboratory for Diagnostics, Epidemiology and Ecology of Microorganisms, Crop Research Institute, Drnovska 507, CZ-16106 Prague 6, Ruzyne, Czech Republic

Abstract

Actinobacteria are important cave inhabitants, but knowledge of how anthropization and anthropization-related visual marks affect this community on cave walls is missing. We compared *Actinobacteria* communities among four French limestone caves (Mouflon, Reille, Rouffignac, and Lascaux) ranging from pristine to anthropized and within Lascaux Cave between marked and unmarked areas on walls in different rooms (Sas-1, Passage, Apse, and Diaclase). In addition to the 16S rRNA gene marker, 441-bp fragments of a heat shock protein gene (*hsp65*) were used for the identification of *Actinobacteria* to the species level by Illumina MiSeq analysis. The *hsp65* marker revealed higher resolution for species and richness (at 99% OTUs cutoff) assessment than did 16S rRNA, which on the other hand identified more taxa on higher taxonomic ranks. *Actinobacteria* communities varied between Mouflon and Reille (both pristine), Rouffignac (anthropized/visited), and Lascaux (strongly anthropized and now closed to the public). Rouffignac displayed high diversity of *Nocardia*, including species with potential pathogeny (i.e. *N. carnea*, *N. paucivorans*, *N. abscessus*), and Lascaux high *Mycobacterium* abundance, whereas *Gaiellales* were typical in pristine caves and the Diaclase i.e. the least affected room of Lascaux Cave. Within Lascaux, the *Pseudonocardiaceae* dominated on unmarked walls and the *Streptomycetaceae* (especially *Streptomyces mirabilis*) on marked surfaces, raising questions on the possible role of *S. mirabilis* in the formation of visual marks. Our results show how the use of the *hsp65* marker, well beyond the resolution provided by 16S rRNA sequences, enabled for the first time to document species-level variations of the *Actinobacteria* community according to the extent of anthropogenic pressure. This approach proved effective when comparing different limestone caves or specific conditions within a cave.

Introduction

Limestone caves are considered to be rather isolated habitats that are extremely limited in organic carbon (Barton et al., 2007), which leads to the development of microbial strategies enabling adaptation to oligotrophic conditions and a high calcium content (Barton & Jurado, 2007; De Mandal, Chatterjee, & Kumar, 2017; Ortiz et al., 2014). Some of these caves contain Paleolithic artwork and thus are frequently visited by tourists (Alonso et al., 2019; Schabereiter-Gurtner, Saiz-Jimenez, Pinar, Lubitz, & Rölleke, 2002). As a result, human-derived dispersion of endogenous as well as exogenous nutrients and microorganisms inside the caves together with the human metabolism may change their internal microclimatic and microbiological conditions (Hoyos, Cañaveras, Sánchez-Moral, Sanz-Rubio, & Soler, 1998; Mulec, 2014). Microorganisms can be resilient to changes to some extent (Johnston, Muench, Banks, & Barton, 2012; Tomczyk-Żak & Zielenkiewicz, 2016). Community shifts may also take place, which deserves attention if it could favor microorganisms with biodeterioration ability (F. Bastian, Jurado, Novakova, Alabouvette, & Saiz-Jimenez, 2010; Fernandez-Cortes et al., 2011; Sánchez-Moral et al., 1999) or increase the proportion of species harboring strains with pathogenic potential (Bercea, Năstase-Bucur, Moldovan, Kenesz, & Constantin, 2019; Neral, Rajput, & Rai, 2015; Rajput, Rai, & Biswas, 2012). If it were the case, such shifts could represent a threat for cave conservation and the health of visitors, respectively.

Actinobacteria represent an important cave taxonomic group, from which many potentially pathogenic (Valme Jurado et al., 2010), biotechnologically relevant (Bhullar et al., 2012; Hamed, Kafshnouchi, & Ranjbaran, 2019; Syiemiong & Jha, 2019) and novel (Fang et al., 2017; V. Jurado et al., 2008; Valme Jurado, Kroppenstedt, et al., 2009; Rangseekaew & Pathom-aree, 2019) species have been found in these environments. *Actinobacteria* can dominate both in anthropized

(Gonzalez-Pimentel et al., 2018) and pristine caves (De Mandal et al., 2017), but some actinobacterial taxa are specific only one type of cave. For example, *Pseudonocardiaceae* are known as indicators of urban contamination (Miller et al., 2018) and dominate in some anthropized caves, but this dominance is most likely due to their ability to degrade exogenous organic compounds (Lavoie et al., 2017; Porca, Jurado, Žgur-Bertok, Saiz-Jimenez, & Pašić, 2012). This is partially true as well for *Streptomyetaceae*, which were also found among pristine cave inhabitants (Maciejewska et al., 2017). Furthermore, these taxonomic groups seem to be prominent in pigment-forming communities on the walls of certain show caves (Cuezva et al., 2012; Porca et al., 2012). In contrast, *Euzebyales* (Gonzalez-Pimentel et al., 2018), *Nocardiaceae* (De Mandal et al., 2017) and *Gaiellales* (Zhu et al., 2019) represent oligotrophic taxa that dominate in pristine caves, most likely because of their ability to degrade complex compounds which are indigenously present in the cave or fix CO₂.

Actinobacteria taxonomic identification in these environments has largely been based on culture-dependent approaches and 16S rRNA sequencing (De Mandal et al., 2017; Yasir, 2018). The culture-dependent approaches limits the coverage of the community and thus should be used only as a supportive determinant with molecular approaches (Hamedi et al., 2019; Long et al., 2019). The 16S rRNA gene is the marker most frequently used for bacterial systematics but generally cannot resolve individual sequences at the species level (Fox, Wisotzkey, & Jurtshuk, 1992; Hirsch, Mauchline, & Clark, 2010). In addition, identical 16S rRNA gene sequences can be found in different taxa or different 16S rRNA gene sequences in a single species (Větrovský & Baldrian, 2013). Other protein-coding genes with high sequence polymorphisms (i.e., *rpoB*, *gyrB*, *hsp65*) permit the distinction between closely-related strains. However, some of these protein-coding markers may also have their limits as it is the case of *rpoB* gene who is present twice is

some genus belonging to Actinobacteria, they only target limited number of *Actinobacteria* or no appropriate reference database is available for taxonomic identification (Gao & Gupta, 2012; Takeda, Kang, Yazawa, Gono, & Mikami, 2010; Vos, Quince, Pijl, de Hollander, & Kowalchuk, 2012). Nevertheless, the gene of a molecular chaperone, the 65-kDa heat-shock protein (*hsp65*), when amplified by specific primers (Telenti et al., 1993), has been used as a molecular marker for identifying a large number of *Actinobacteria* isolates (Rodríguez-Nava et al., 2006, 2007). This gene is present in *Actinobacteria* genomes as a single copy, unlike the 16S rRNA gene marker (Colaco & MacDougall, 2014), and *hsp65* is considered to be a good candidate for making detailed *Actinobacteria* community assessments from environmental DNA. Additionally, the amplified fragment length of *hsp65* (441 bp) is sufficient for amplicon sequencing.

Lascaux Cave underwent several events of microbial outgrowth since the 1960s, as well as (sometimes unsuccessful) chemical treatments, which resulted in the formation of various visual marks (including stains) and threatened Paleolithic artwork (F. Bastian et al., 2010; Martin-Sanchez, Miller, & Saiz-Jimenez, 2015). For the sake of cave conservation, Lascaux was closed to the public in 1963. Although the few microbiological studies conducted in this cave suggest that *Actinobacteria* is one of prevalent groups (Alonso et al., 2018, 2019; Martin-Sanchez, Nováková, Bastian, Alabouvette, & Saiz-Jimenez, 2012), a more complete investigation of the *Actinobacteria* community is lacking, especially at the species level. If we consider that key functional traits can be related to precise species within the *Actinobacteria*, we can hypothesize that such an assessment would allow a better discrimination of *Actinobacteria* communities according to ecological conditions.

The objective of this work was to assess the usefulness of *hsp65* marker (as a complement to 16S rRNA sequence data) for a deeper, species-level appraisal of the *Actinobacteria* community

in caves, and to show variations according to cave conditions and especially human-induced disturbances (Mammola, 2019). These aims were achieved by comparing *Actinobacteria* communities in pristine limestone caves (Reille and Mouflon Caves) and anthropized caves (Lascaux and Rouffignac Caves), and within Lascaux Cave of marked and unmarked areas in different rooms (i.e., Sas-1, Passage, Apse, and Diaclase). To this end, *Actinobacteria* were identified for the first time to the species level in cave environments, using metabarcoding analysis of the *hsp65* gene marker.

Materials and methods

Study site description and sample collection

Samples were collected in May - June 2016 from the walls of four limestone caves located in the Dordogne department of southwestern France: Lascaux (45°03'13" N 1°10'12"E, 235 m length); Rouffignac (45°00'31"N 0°59'16"E, 8,000 m length); Reille (45°07'09.5"N 1°07'11.2"E, 2,000 m length) and Mouflon (44°55'N 1°5'E, 280 m length) (Fig. 1A).




Fig. 1. Map of A. Dordogne area with the cave locations (Lascaux, Rouffignac, Reille and Mouflon; white squares – pristine and gray squares – anthropized). Map of B. Lascaux Cave (entrance Sas-1, Passage, Apse, and Diacase).

Lascaux and Rouffignac caves contain Paleolithic wall paintings and engravings and are listed as UNESCO World Heritage Sites. Both caves are anthropized, Rouffignac being open for touristic visits since 1959 (with up to 500 visitors per day) (5) and Lascaux being closed after 15 years of tourism (with up to 2000 visitors per day). Various visual marks have developed at different times in Lascaux Cave, and extensive chemical treatments have been applied (13,42, 44,

45). Mouflon and Reille Caves are in a pristine state (5). The entrance of Mouflon Cave was locked shortly after discovery, whereas Reille is occasionally explored by seasoned speleologists.

In Lascaux Cave, samples were taken from the walls of four distinct rooms representing different cave environments that were located progressively farther from the entrance in the following order: Sas-1, Passage, Apse and Diaclase (Fig. 1B). Sas-1 is a man-made, calcareous airlock entrance and is isolated by doors from the cave interior and exterior. The Passage is a central alley connecting different rooms of Lascaux Cave. In the Passage, samples were taken from two mineral substrates: clay deposits (banks, later referred to as Passage B) and near-vertical limestone walls (inclined planes, later referred to as Passage IP). In the Apse, samples were collected from limestone inclined planes. The Diaclase is a distinct compartment located below the main part of the cave and was least-affected by humans. It is separated from the Apse by a trap door, received limited biocide treatments, and was never open to tourists. Samples originated from limestone walls located immediately below the Apse, in an area of the Diaclase termed the Shaft of the Dead Man.

Within each room, samples were collected from visual marks and unstained surfaces, as described (Alonso et al., 2018). Approximately 50 mg of wall material samples (3-6 replicates) were collected using sterile scalpels and placed in liquid nitrogen for transportation to the laboratory, where they were stored at -80°C until DNA extraction. Overall, six samples from Rouffignac, six from Mouflon, five from Reille and 41 from Lascaux (five from Sas-1, six from Passage B, six from Passage IP, 18 from Apse and six from Diaclase) were used for subsequent microbial analysis (Table S1). Sampling was performed in accordance with the caves' rules and regulations.

DNA extraction and amplicon sequencing

DNA extraction was performed using the FastDNA SPIN Kit for Soil (MP Biomedicals, Illkirch, France) following the manufacturer's instructions and was adapted to the low sample amounts, as described (Alonso et al., 2018). The elution step was achieved using two volumes of 50 µl elution buffer for each sample.

For bacterial 16S rRNA gene amplification, we used the primers 341F (CCTACGGGNGGCWGCAG) and 805R (GACTACHVGGGTATCTAATCC), which target the V3-V4 region (428-bp fragments) (Herlemann et al., 2011). Amplification and sequencing were carried out by the Fasteris Company (Geneva, Switzerland) using the Illumina MiSeq Reagent Kit v3 (600 cycles) with the paired-end mode, resulting in 2×300 bp sequence reads, as described by Alonso et al. (Alonso et al., 2018).

The 441-bp fragments of the 65-kDa heat shock protein gene (*hsp65*) were amplified by PCR using the specific primers TB11 (ACCAACGATGGTGTGTCCAT) and TB12 (CTTGTCGAACCGCATACCCT) from Telenti et al. (Telenti et al., 1993). Amplification was carried out in a reaction mixture with a total volume of 25 µl in packaged PCR tubes (Ready-to-Go PCR Beads; Amersham Biosciences, Piscataway, NJ) (2.5 U of Taq polymerase puRe Taq, 10 mM Tris-HCl [pH 9], 50 mM KCl, 1.5 mM MgCl₂, and 200 µM of each deoxynucleoside triphosphate) with primers (0.2 – 0.4 µM, depending on the template concentration) and DNA (1 – 40 ng). The PCR cycling protocol consisted of an initial denaturation at 94°C for 5 min, then 35 cycles of denaturation (94°C for 60 s), annealing (55°C for 60 s), and elongation (72°C for 60 s) (Rodríguez-Nava et al., 2006). Sequencing was carried out (together with purification and quality control) by the Biofidal Company (Biofidal, Vaulx-en-Velin, France – <http://www.biofidal-lab.com>) using the Illumina MiSeq Flow Cell V3 with the paired-end mode, resulting in 2×300

bp sequence reads. Overall, 50 samples for 16S rRNA gene and 45 samples for *hsp65* were successfully sequenced, but sequences for both markers were obtained for only 37 samples (Table S1).

Processing and analysis of sequence data

The paired sequence reads from sequencing were merged using FLASH (Fast Length Adjustment of Short reads (Magoč & Salzberg, 2011)) with a maximum of 25% mismatches in the overlapping region. The sequences were then filtered and aligned using reference alignments of the *hsp65* gene sequences (this work) or using the 16S rRNA gene sequences from the Silva database (Quast et al., 2012). Chimeric sequences were removed using the integrated Vsearch tool (Rognes, Flouri, Nichols, Quince, & Mahé, 2016) according to the MiSeq standard operating procedure (MiSeq SOP, February 2018) (Kozich, Westcott, Baxter, Highlander, & Schloss, 2013) in Mothur v. 1.39.5 software (Schloss et al., 2009). Taxonomical assignments of sequence libraries were performed in Mothur using the *hsp65* reference database for the *hsp65* sequences and the recreated SEED database subset of the Silva Small Subunit rRNA Database, release 132 (Yilmaz et al., 2014), adapted for use in Mothur (https://mothur.org/w/images/a/a4/Sylva.seed_v132.tgz), as the reference database for the 16S rRNA gene sequences.

The sequences of plastids and mitochondria and those not classified in the domain Bacteria (from the 16S rRNA sequence library), as well as sequences identified as homologs for *hsp65* (*groEL*, in the database renamed as GROESL; see the *hsp65* reference database design section) (from the *hsp65* sequence library), were discarded. The sequence library was clustered into operational taxonomic units (OTUs) using the Uparse pipeline in Usearch v10.0.240 software (OTUs at a 97% cutoff) (35) and Mothur (OTUs at a 99% cutoff). The OTU table was further

processed using tools implemented in Mothur. For 16S rRNA genes, only sequences corresponding to *Actinobacteria* were used.

For comparison of *Actinobacteria* among the caves, 25 samples for *hsp65* and 24 samples for 16S rRNA genes were used. Apse and Diacase (both unmarked areas) were chosen as representative samples for Lascaux Cave except for core microbiome analysis, for which all the Lascaux Cave samples were included. To test the differences among locations in Lascaux Cave, 33 samples for *hsp65* and 38 samples for 16S rRNA were used. Among those, from marked areas 16 samples for *hsp65* and 20 samples for 16S rRNA were used. From unmarked areas 17 samples for *hsp65* and 18 samples for 16S rRNA were used (Table S1). For 16S rRNA marker, Sas-1 unmarked area samples are missing for the analysis, because low number of replicates were successfully sequenced. To obtain the highest number of replicates per sample, all samples, even when they were not common to both markers, were included in the analysis. 37 common samples for both markers were used for analyses comparing the usefulness of both markers (rarefaction analysis, comparison of diversity indices and number of taxa recovered at different taxonomy ranks).

Rarefaction analysis was performed to analyze the richness of the *Actinobacteria* OTUs based on two cutoffs (i.e., 97% and 99%) as a function of sample number. The alpha diversity indices (richness: Chao 1, evenness: Simpson Evenness, and diversity: Inverse Simpson) were calculated in Mothur (97%, 99% OTU cutoff). Significant differences among samples were calculated by ANOVA with Tukey's post hoc test ($P < 0.05$; Caves and locations in Lascaux Cave), t tests and F tests ($P < 0.05$; marked/unmarked areas) in Past 4.02 software (Hammer, Harper, & Ryan, 2001).

Bray-Curtis distance matrices describing the differences in bacterial community composition among individual samples were calculated. The *Actinobacterial* communities in the caves, in the different locations of Lascaux Cave, and between marked and unmarked areas within each location were compared by nonmetric multidimensional scaling (NMDS) according to both gene markers (97% OTU cutoff). Analysis of molecular variance (AMOVA) and homogeneity of molecular variance (HOMOVA) were calculated in Mothur with 100,000 iterations for the same factors (97% OTU cutoff).

Pairwise comparisons with Metastats (Segata et al., 2011) and biomarker searching analysis with Lefse (Segata et al., 2011) were calculated to identify the *Actinobacteria* OTUs that were significantly different among the respective caves, locations and marked/unmarked areas in Lascaux Cave (99% OTU cutoffs). Metastats was also used to compare the relative abundances of *Streptomyces* sequences among the marked/unmarked areas (*hsp65* marker). Venn diagrams were created in Mothur to test the number of OTUs that encompassed the core microbiome for the different caves and four rooms (e.g., Sas-1, Passage, Apse, and Diaclase) in Lascaux Cave (99% OTU cutoff).

Taxonomical classification of OTUs was performed with Mothur by using the *hsp65* and 16S rRNA reference databases, and the *hsp65* marker was verified by BLASTN in the online databases of the National Center for Biotechnology Information (NCBI; U.S. National Library of Medicine, MD, USA). The taxonomical composition of the core microbiome was constructed using the Krona tool (Ondov, Bergman, & Phillippy, 2011). The number of taxa recovered with the common samples by *hsp65* and 16S rRNA at different taxonomic ranks were calculated and displayed by Venn diagrams.

Co-occurrence networks were used to predict how marked/unmarked areas and different locations influenced the relative abundances of *Actinobacteria* in Lascaux Cave. Spearman correlation coefficients were computed in Mothur for Lascaux Cave OTUs and were categorized according to their significantly different proportions in the marked/unmarked areas regardless of the location in Lascaux Cave (Metastats) or for the Lascaux Cave locations (Lefse) (99% OTU cutoff). Only significant ($P < 0.03$) correlations higher than 0.8/0.8 and lower than -0.35/-0.5 for the *hsp65*/16S rRNA markers, respectively, were visualized in Gephi 0.9.2. (M. Bastian, Heymann, & Jacomy, 2009) by Fruchterman-Reingold spatialization (Fruchterman & Reingold, 1991). The average degree, eigenvector centrality and modularity were computed, and the minimum number of links was filtered to 5 (using a degree range filter).

Figures were plotted using the vegan package (Oksanen et al., 2018) in the R computing environment (R Core Team, 2018) and in Inkscape (v0.92; <http://www.inkscape.org>).

The Illumina MiSeq 16S rRNA and *hsp65* gene amplicon sequences were deposited in the NCBI Sequence Read Archive (www.ncbi.nlm.nih.gov/sra).

***In silico* analysis**

Determination of the differences in the variability of the *hsp65* and 16S rRNA partial gene sequences between pairs of *Actinobacteria* strains at different taxonomical levels (e.g., intraclass, intraorder, intrafamily, and intragenus) was performed on genomes retrieved from the Integrated Microbial Genomes and Microbiomes database (IMG; University of California, CA, USA, <https://img.jgi.doe.gov/>). For the analysis, sequences from the available genomes of 47 species from the order *Streptomycetales* (*Streptomyces*, *Kitakatospora*, Table S2) and 58 species from the order *Corynebacteriales* (family *Corynebacteriaceae*: *Corynebacterium*; family *Gordoniaceae*:

Gordonia; family *Mycobacteriaceae*: *Mycobacterium*; family *Nocardiaceae*: *Nocardia*,
Rhodococcus; Table S2) were used. Sequences were aligned and trimmed to the amplified length
of both gene markers in BioEdit 7.2.5 software (Hall, Biosciences, & Carlsbad, 2011). The
pairwise distances between sequences were calculated in Mothur v. 1.39.5 software (Schloss et al.,
2009).

***hsp65* reference database design**

For taxonomic assignment using the *hsp65* marker, the reference database of the
Actinobacteria hsp65 gene was constructed similarly to that of the *rpoB* marker in the study of
Ogier et al. (2019) (Ogier, Pagès, Galan, Barret, & Gaudriault, 2019). Sequences homologous to
the 65-kDA heat shock protein (*hsp65*) of the reference strain *Nocardia asteroides* ATCC 14759
(Rodríguez-Nava et al., 2006) amplified by the TB11 and TB12 primers (Telenti et al., 1993) were
collected in the online databases of IMG and NCBI. Paralogs of heat shock proteins, such as *groEL*
genes from *Escherichia coli* (Colaco & MacDougall, 2014; Duchêne, Kieser, Hopwood,
Thompson, & Mazodier, 1994; C. M. S. Kumar, Mande, & Mahajan, 2015), were identified using
maximum-likelihood, FastTree 2.1 (Price, Dehal, & Arkin, 2010) and were kept in the reference
database (renamed as GROESL sequences) as an outgroup that enabled amplified sequences
belonging to this variant to be discarded.

The *hsp65* database contained 5,165 sequences, of which 1,782 sequences were denoted as
GROESL at the time of the analysis. Evaluation of the TB11 and TB12 primer complementarity
with the sequences from the reference database was performed using the OligoAnalyzer 3.1 tool
(<http://www.idtdna.com/calc/analyzer>). The percent coverage of the primers of *Actinobacteria*
genomes from different classes and orders was determined with primer BLAST using the RefSeq

293 reference genomes (NCBI). Only targets with no mismatches for the last three nucleotides at the
294 3' end and no more than 4 overall mismatches were included.

Results

Sequence polymorphism of *hsp65* versus 16S rRNA genes for cave *Actinobacteria*

A total of 1,799,680 *hsp65* and 1,919,236 16S rRNA gene sequence reads were obtained for cave *Actinobacteria*. They were mapped to 968 (97% cutoff) and 4,265 OTUs (99% cutoff) for *hsp65*, vs only 299 (97% cutoff) and 718 OTUs (99% cutoff) for 16S rRNA genes.

The curves of 16S rRNA genes for both cutoffs were more skewed and thus indicated an accumulation of identical OTUs due to their repeated sampling, which did not increase with a stricter cutoff, in contrast to *hsp65* (Fig. 2A). These results are in accordance with the *in silico* analysis of *hsp65* and 16S rRNA partial gene sequences downloaded from genomes available on IMG, where the pairwise distances between *hsp65* sequences, even for closely related species, were higher than those between 16S rRNA genes (Fig. 2B).




Fig. 2. A. Rarefaction curves for the 37 common samples of the *hsp65* and 16S rRNA genes at the 97% and 99% OTU cutoffs. The X-axis indicates the number of samples, and the Y-axis indicates the number of OTUs. B. Pairwise molecular distances between sequences of the *hsp65* (Y axes) and 16S rRNA (X axes) genes among *Actinobacteria* species from different taxonomical levels with the R^2 level for each equation.

The *hsp65* primers targeted *hsp65* sequences from the reference database with lower numbers of mismatches compared to the outgroup (i.e., distant homologs renamed in the database as GROESL, *GroEl* of *Escherichia coli* and Chloroflexi members) (Table S3). However, the primers targeted only selected groups from the class *Actinobacteria* as can be seen from analysis with representative genomes from online database (Table S4). Moreover, the number of taxa identified at class or lower categories is shown in a Fig. 3. At the class level, 20% of the detected class with *hsp65* were also recovered with 16S rRNA, 32.3% it was for order 36.1% for family, 24.6% for genus-level taxa. Although the number of taxa detected uniquely by *hsp65* marker had increasing trend with the lower taxonomic category, contrary to 16S rRNA, *hsp65* detected overall lower number of taxa. However, on a species level overall 168 species were recovered only by *hsp65* marker, where the highest species number were found for *Streptomyces*, *Mycobacterium* and *Nocardia* genera (Table S5).




Fig 3. Venn diagrams showing the number of different *Actinobacteria* taxa recovered with *hsp65* (blue) or 16S rRNA (red) at different taxonomic ranks. (A) Class, (B) Order, (C) Family, (E) Genus.

This means that the 16S rRNA marker was more suited for broad taxonomic profiling of the *Actinobacteria* community. In comparison, the *hsp65* marker could distinguish between selected *Actinobacteria* species, which the 16S marker failed to achieve, and thus it completed the results from the 16S rRNA marker (Table S5).

***Actinobacteria* diversity in caves based on *hsp65* versus 16S rRNA genes**

According to the diversity analysis based on 37 common samples of both markers, the *hsp65* marker recorded higher richness (Chao-1 index) and evenness (Simpson evenness index) only when considering the 99% OTU cutoff than 16S rRNA marker (Table S6). The diversity analysis

based on all samples indicated that Rouffignac Cave exhibited the highest *Actinobacteria* richness (Chao 1 index, for the *hsp65* marker; Fig. S1) but lower evenness (Simpson evenness index; for both markers; Fig. S1) as well as diversity (Inverse Simpson index; for the 16S rRNA marker; Fig. S1) among other caves. In contrast, the Lascaux and Rouffignac Caves had lower evenness than the pristine Reille (for the 16S rRNA marker) and the Mouflon Caves (for the *hsp65* marker; Fig. S1). When comparing locations within Lascaux Cave, only the difference in *Actinobacteria* richness between Sas-1 and Passage B was significant (for the 16S rRNA marker; Fig. S1). The average diversity of the unmarked areas was higher than that of the marked areas for both markers, (Fig. S1). Therefore, when documenting diversity *hsp65* might uncovered the high richness in some of the samples which was not recorded by the 16S rRNA marker.

***Actinobacteria* community structure in caves based on *hsp65* versus 16S rRNA genes**

NMDS analysis was conducted to compare the *Actinobacteria* communities in caves (97% OTU cutoff). These communities did not differ significantly between pristine caves Reille and Mouflon (AMOVA), based on *hsp65* and 16S rRNA genes. NMDS distinguished three groups of communities corresponding to Lascaux, Rouffignac, and the two pristine caves Reille and Mouflon, which was further supported by AMOVA (Table S7 A). The AMOVA comparison of two groups: anthropized (Lascaux and Rouffignac) and pristine (Reille and Mouflon) caves showed, that they significantly differed according to both markers (*hsp65*: $F= 3.986$, $p < 0.001$, 16S rRNA: $F=8.610$, $p < 0.001$). Within Lascaux Cave, the *Actinobacteria* communities differed largely according to all locations (i.e. Sas-1, Passage B, Passage IP, Apse, Diacalse) with both markers, except that differences between Passage IP, Sas-1 (both markers) and Diacalse (*hsp65*) were not significant due to the low number of samples (Table S7 A). Overall, the effect of surface

alterations (i.e. marked vs unmarked areas) in Lascaux Cave was significant for both markers (Table S7 A), with higher *Actinobacteria* stability in unmarked than vs marked areas (Table S7 B). Passage B had the highest community stability compared to Passage IP, Apse and Diacalse (Table S7 B), which was reflected in the NMDS plot, where the marked and unmarked areas in Passage IP were the least separated from one another (Fig. 4).




Fig. 4. Sammon projection of nonmetric multidimensional scaling (NMDS) based on the Bray-Curtis distance matrices of the *hsp65* and 16S rRNA gene markers of A. different caves and B. Lascaux Cave locations and marked/unmarked areas. The F-values and P-values of overall AMOVA are indicated (a - locations, b - un/marked areas; 97% OTU cutoff).

Core and cave-specific *Actinobacteria* microbiomes based on *hsp65* versus 16S rRNA genes

Venn diagrams indicated that the number of OTUs shared between different caves (Rouffignac, Lascaux, Mouflon, Reille) was only 2 (0.04%) for the *hsp65* marker vs 72 (11%) for the 16S rRNA marker (Fig. S2). Similarly, the number of OTUs shared between different Lascaux rooms (Sas-1, Passage, Apse, Diaclase) reached 23 OTUs (0.89%) for the *hsp65* marker vs as many as 88 (15.9%) for the 16S rRNA marker. The core microbiome was constituted especially by *Pseudonocardiales* and *Streptomycetales*, as determined by *hsp65*, and *Gaiellales*, *Pseudonocardiales* and strain IMCC26256_ge, as determined by 16S rRNA genes (Fig. S3).

The pairwise comparison of cave conditions (Rouffignac, Lascaux's Apse and Diaclase, Mouflon, Reille) using significantly different OTUs (Metastats) showed that Rouffignac (based on *hsp65*) and Lascaux's Apse (based on 16S rRNA genes) were most enriched in OTUs, separating them from the other caves (Fig. 5). Among them, *Pseudonocardiaceae*, *Crosiella* and *Nocardia* were the most abundant taxa that distinguished Rouffignac (for both markers), while *Nocardioides*, *Rhodococcus*, *Pseudonocardia* distinguished Lascaux's Apse from the other caves and Diaclase. *Mycobacterium* was uniquely found in Lascaux's Apse according to both markers. Pristine caves were typified by *Streptomyces* (*hsp65* marker), *Gaiellales*, *Pseudonocardiales* and the environmental clones MB-A2-108_ge and IMCC26256_ge (16S rRNA genes)(Fig. 5).




Fig. 5. Significantly different OTUs between pairs of caves (Metastats, $P < 0.005$). For each cave, the number of OTUs that proportionally differed from other caves and taxonomically assigned OTUs proportionally the most abundant in the respective cave are indicated (99% OTU cutoff).

Proportions of cave *Actinobacteria* in sequences libraries based on *hsp65* versus 16S rRNA genes

Taxonomical analysis showed that *Jiangellaceae*, *Actinobacteria incertae sedis* *Brevibacteriaceae*, *Nocardiopsaceae* and *Gordoniaceae* were uniquely identified by *hsp65* marker but not by 16S rRNA marker. *Nocardioidaceae*, *Streptomycetaceae* and *Pseudonocardiaceae* dominated the sequence libraries of both gene markers, when considering all cave conditions together. *Nocardioidaceae* and *Streptomycetaceae* (according to *hsp65*) while *Nocardiaceae* and *Streptomycetaceae* (according to 16S rRNA marker) were redundant especially in Lascaux Cave. According to both gene markers, *Streptomycetaceae* were in a higher proportion in marked areas than in unmarked areas of the Lascaux cave (Fig. 6). With a closer focus on *Streptomyces* using

only the *hsp65* marker, which contrary to 16S rRNA can identify the species, *S. mirabilis*, *S. niveus*, and *S. fulvissimus* were the three species, which dominated in marked areas within Lascaux Cave (Metastats $P < 0.005$; Fig. S4). The opposite result was found for *Pseudonocardiaceae*, which were more proportionally abundant in Lascaux's unmarked areas than in marked areas except for Passage IP (by *hsp65*) and Passage B (by 16S rRNA marker) (Fig. 6). *Mycobacteriaceae* were found to be highly abundant in Lascaux Cave, and the *hsp65* marker also identified them in Mouflon. *Jiangellaceae* was a group that was typical in Rouffignac and Passage IP (*hsp65* marker). Closer analysis of *Nocardiaceae* with the *hsp65* marker uncovered the greatest richness of *Nocardia* species in Rouffignac (including *N. carnea*, *N. paucivorans*, *N. abscessus*). Moreover, high proportions of *N. jejuensis* in Lascaux and Mouflon, and *N. cummidelens* in Rouffignac, Apse and Reille were found (Fig. S5). Both pristine caves had similar compositions of *Actinobacteria* communities, especially *Gaiellaceae*, taxa related to the environmental clones MB-A2-108_fa and IMCC26256_fa (16S rRNA marker), and high proportions of unclassified *Actinobacteria* (*hsp65*). High proportions of unclassified *Actinobacteria* were similarly to pristine caves observed in Diaclase (Fig. 6).




Fig. 6. Average proportions of families in the *Actinobacteria* A. *hsp65* and B. 16S rRNA amplicon sequence libraries from different caves (Rouffignac, Lascaux, Mouflon and Reille), Lascaux Cave locations (Sas-1, Passage-B, Passage - IP, Apse and Diacalse) and marked (S)/unmarked areas (U) within Lascaux.

Co-occurrence of *Actinobacteria* in Lascaux Cave based on *hsp65* versus 16S rRNA genes

Co-occurrence networks showed the Spearman correlations between the relative abundances of the *Actinobacteria* OTUs in Lascaux Cave (Fig. 7). The co-occurrence networks identify OTUs significantly different for any of the Lascaux locations (calculated by Lefse) or for marked/unmarked areas (calculated by Metastats). More co-occurrence connections were found for *hsp65* than for 16S rRNA marker. For *hsp65* marker, more correlated OTUs were not indicated by Lefse or Metastats as typical for any of the Lascaux location or marked/unmarked area, respectively. Negative correlations between the OTUs revealed only 16S rRNA marker.

We found clusters of correlated OTUs dominating the unmarked areas that were also significantly different in the Apse and Diaclase based on both gene markers (Fig. 7). Additionally, OTUs clusters that dominated the marked areas were typical for the Diaclase and Sas-1 areas based on the 16S rRNA marker. The OTUs from such distinct areas (Diaclase and Sas-1) correlated with each other, suggesting that the marked areas factors affected the *Actinobacteria* communities in both locations (Fig. 7). However, the *hsp65* gene marker showed that the OTU clusters typical of marked areas were not typical for any of the Lascaux Cave locations. Finally, based on 16S rRNA marker, the highest number of correlations between OTUs that did not differ between the marked and unmarked areas was found for those that dominated in Passage B, Sas-1 and Diaclase (Fig. 7). For those OTUs, the location factor was more influential than the marked/unmarked areas. Networks based on both markers include variable actors (except the OTUs in unmarked areas of Apse and in Diaclase regardless marked/unmarked areas) which show that each marker cover different part of community and their interactions.




Fig. 7. Co-occurrence networks of *Actinobacteria* OTUs (A. *hsp65* marker, B. 16S rRNA

marker) between OTUs that were significantly different between marked (black) and unmarked (white) areas in Lascaux Cave and those that did not differ between these areas (gray) using Metastats ($P < 0.05$). The letters indicate OTUs that were specific for the respective Lascaux Cave locations (S, Sas-1; PB, Passage banks; PI, Passage inclined planes; A, Apse; and D, Diaclose) using Lefse ($P < 0.03$). Strong significant connections (Spearman's correlation > 0.8 and < 0.5 for 16S rRNA gene and < 0.35 for *hsp65* these thresholds are determined how?) are displayed (99% OTU cutoff).

Discussion

This study represents the first attempt to exploit the potential of *hsp65* as a taxonomic marker more discriminant than the 16S rRNA gene to document *Actinobacteria* diversity at the species level in cave environments. Indeed, Venn results showed a higher variability for *hsp65* sequences than for *Actinobacteria* 16S rRNA sequences, and a higher OTU richness (at the 99% cutoff) of *Actinobacteria* was obtained. Moreover, thanks to *hsp65* marker we evidenced 168 species which could not be identified by the same samples number of 16S rRNA marker. Therefore, lower interspecific variation of 16S rRNA might underestimates the true diversity which might be obtained by *hsp65* marker as was similarly found for other protein-coding gene, (namely *rpoB*) (Vos et al., 2012). However, overall more taxa on higher taxonomic ranks were identified by 16S rRNA than *hsp65* marker which might point on gaps in reference database which prevent complete taxonomic assignment of *hsp65* sequences. Another reason might be a problem with mismatch primers. As was shown by *in silico* analysis, the primers amplify the gene from diverse *Actinobacteria* genomes, but targets without perfect homology might be amplified with lower efficiency leading to underestimation of some taxa (Deagle, Jarman, Coissac, Pompanon, & Taberlet, 2014). It confirms that *hsp65* marker is suitable complement for 16S rRNA in high throughput sequencing methods, especially for specific taxonomic groups analysis into species level, but possible primer biases might be take into account especially for community analysis at higher taxonomic ranks.

In this study, the combination of *hsp65* and 16S rRNA markers showed that *Actinobacteria* diversity could vary according to anthropization in caves, at both higher (i.e., between caves) and lower (i.e., between locations within one cave) spatial scales. Indeed, the actinobacterial communities of anthropized (Lascaux and Rouffignac) and pristine caves (Reille and Mouflon)

differed. The occurrence of these caves in the same region and limestone vein probably facilitated this observation, as geographic distance (Barraclough, Balbi, & Ellis, 2012) and geological type (Zhu et al., 2019) are significant factors influencing cave communities. The core *Actinobacteria* microbiome was relatively small (especially for *hsp65*) but it included typical pristine cave taxa, indicating that at least part of the community was able to resist anthropogenic disturbances (Shade et al., 2012).

Mouflon and Reille are pristine caves with evenly distributed taxa in the *Actinobacteria* community, possibly as a result of their stable environments without man-made disturbances (Mammola, 2019). Typical pristine cave taxa were evidenced in Mouflon and Reille, such as *Gaiellales* (Zhu et al., 2019) and *Actinobacteria* related to the environmental clone MB-A2-108_ge (A. DE Kumar et al., 2019; Zhang et al., 2019) according to the 16S rRNA marker. High proportions of *Streptomyces* and unclassified *Actinobacteria* based on *hsp65* data suggest that each marker selectively identify partly a different *Actinobacteria* community. Environmental selection in oligotrophic habitats may promote high molecular evolution (Kuo & Ochman, 2009a; Sagova-Mareckova et al., 2015a) and result in a high number of novel species isolated from caves (A. DE Kumar et al., 2019). A similar result was also found for *Streptomyces* genera (Hamm et al., 2017), whose genomes may undergo a high evolution rate (Cheng et al., 2015). Interestingly, similar results (i.e. high proportions of *Streptomyces*) to those for pristine caves were obtained from Diaclose, which is an isolated site within the anthropized Lascaux Cave (Kuo & Ochman, 2009b; Sagova-Mareckova et al., 2015b), suggesting that it has remained at least partially protected from human impacts. Therefore, not only individual caves but also less-exposed locations within anthropized caves maintained a typical *Actinobacteria* cave community, which probably include novel taxa not described so far (Rangseekaew & Pathom-Aree, 2019).

Contrary to a previous study based on whole bacterial community (Alonso et al., 2019), we found a high richness but low evenness of *Actinobacteria* in the visited Rouffignac Cave compared to the pristine caves based on *hsp65* marker. The high richness of *Actinobacteria* in this cave might be a result of community response to a disturbance, since more diverse community is more resilient to stressful conditions (Shade et al., 2012). Our results are especially supported by the dominance of *Pseudonocardiaceae* (based on 16S rRNA sequences), an indicator of human contamination (Miller et al., 2018), as well as by the highest richness of *Nocardia* species (based on the *hsp65* marker). More specifically, in Lascaux and Mouflon dominated *N. jejuensis*, the species firstly isolated from a natural cave, while in Reille, Rouffignac and Apse dominated *N. cummidelens*, a species previously isolated from rocks of Altamira cave (Valme Jurado, Fernandez-Cortes, et al., 2009). However, *Nocardia* in Rouffignac uniquely included species with pathogenic potential, such as *N. carnea* (Boiron, Provost, Chevrier, & Dupont, 1992), *N. paucivorans* (Watanabe et al., 2006), *N. testacea* (Taj-Aldeen et al., 2013), and *N. abscessus* (Kageyama et al., 2004). Although we cannot know if these environmental bacteria present any pathogenic potential, as those found in human hosts, the enrichment of these species in caves points to a possible allochthonous input by tourists. In turn, it raises the issue of potential health concerns, as proposed previously (Valme Jurado et al., 2010), even though *Actinobacteria* disease problems related to cave frequentation have never been reported. In contrast, *Jiangellaceae* are a typical pristine cave family (Rangseekaew & Pathom-aree, 2019) and are another important group in Rouffignac Cave. According to the intermediate disturbance hypothesis (Roxburgh, Shea, & Wilson, 2004), the coexistence of typical pristine and anthropized taxa in one habitat may suggest that the cave is under intermediate environmental pressure.

The actinobacterial communities in caves were highly specific even within very close distances (81), since each Lascaux room hosted its own *Actinobacteria* populations. Lascaux Cave was typified by high abundances of *Mycobacterium*. Similar to *Nocardia* in Rouffignac, this group can signify external contamination (Valme Jurado et al., 2010; Modra et al., 2017). Confirming the previous results from Alonso et al. (2018) that were based on the whole bacterial community (Alonso et al., 2018), the different geological substrates of Lascaux's Passage had a stronger role than the occurrence of marks, but marks there (black stains) had formed years before. However, the effect of geological substrate contrasted with Diaclase and Apse because these two locations contained significantly different *Actinobacteria*, especially in unmarked areas, as shown by network analysis. It is possible that the isolation and distance from the cave entrance represent the best filter for alien microorganisms and enable the maintenance of the cave oligotrophic community, which is typical for unmarked areas (Cuezva, Sanchez-Moral, Saiz-Jimenez, & Cañaveras, 2009; Mammola, 2019).

Our results show that the *Actinobacteria* community, mainly *Streptomyces*, varied according to the presence of visual marks. Marked areas, where *Streptomycetaceae* dominated, exhibited much lower diversity than unmarked zones, where *Pseudonocardiaceae* were prevalent. Similarly to our study, the group *Pseudonocardiaceae* was confirmed as true cave rock dwellers (Zhu et al., 2019). However, contrary to our results, this group was typically found in pigmented zones of pristine caves (Lavoie et al., 2017). In addition, the members from the group *Pseudonocardiaceae* are degraders of complex molecules and indicators of anthropogenic contamination (Lavoie et al., 2017; Porca et al., 2012), which suggests that the unmarked areas were not completely oligotrophic (Tomczyk-Żak & Zielenkiewicz, 2016). However, the higher community diversity in unmarked areas than in marked areas indicates the prevalence of

cooperative relationships, which are typical for oligotrophic cave environments (Tomczyk-Żak & Zielenkiewicz, 2016).

In contrast, the diversity of marked areas might be reduced due to the competitive advantage of invading species (Hamm et al., 2017; Van Elsas et al., 2012). In isolated cave systems, invasions are expected from the cave entrance, similar to Altamira Cave, where *Streptomyces* is the most dominant group, especially in stains (Groth, Vettermann, Schuetze, Schumann, & Saiz-Jimenez, 1999). Moreover, they are generally known to be competitively effective because of their secondary metabolite production, which is supported by the prevalence of the antimicrobial producers *S. mirabilis*, with gray mycelium (El-Sayed, 2012), *S. niveus* (Kominek, 1972), and *S. fulvissimus* (Myronovskyi, Tokovenko, Manderscheid, Petzke, & Luzhetskyy, 2013) in stains. Hypothetically, *Streptomyces* might also contribute to pigment production, as indicated in different studies (Abdel-Haliem, Sakr, Ali, Ghaly, & Sohlenkamp, 2013; Cuezva et al., 2012), or might affect the pigment-forming fungi that are involved in black stain formation in Lascaux Cave (De La Rosa et al., 2017; Frey-Klett et al., 2011). Another study showed that *Streptomyces* cooperation with fungi in stains promoted mycelial extension and secondary metabolite production of fungi (Frey-Klett et al., 2011). A previous study showed the correlation between *Streptomyces* and pigment-forming fungi *Acremonium* and *Exophiala* in Lascaux Cave, although only in unmarked areas (Alonso et al., 2018). Therefore, *Streptomyces* role in black stains should be studied in more detail to evaluate if this group may represent a missing chain for understanding stain development in Lascaux Cave.

To conclude, the *hsp65* marker can be reliably used for differentiating *Actinobacteria* species present in environmental samples. However, in environmental studies, *hsp65* should functionate as a complement for 16S rRNA gene, since 16S rRNA gene encompass larger part of

573 Actinobacteria community but cannot resolve species. the *Actinobacteria* community at different
574 spatial levels reflected the natural quality of the caves and different locations within Lascaux Cave.
575 Anthropization was shown to shape *Actinobacteria* communities, which altered typical cave
576 communities and was exhibited by the indicator *Actinobacteria*. We first found that the marked
577 areas of Lascaux Cave are an important factor shaping *Actinobacteria* communities, with particular
578 reference to the *Streptomyces*. Finally, the study revealed the linkage between *Actinobacteria* in
579 anthropized caves and marked areas and may help promote the conservation of

References

- Abdel-Halim, M. E. F., Sakr, A. A., Ali, M. F., Ghaly, M. F., & Sohlenkamp, C. (2013). Characterization of *Streptomyces* isolates causing colour changes of mural paintings in ancient Egyptian tombs. *Microbiological Research*, 168(7), 428–437. doi: 10.1016/j.micres.2013.02.004
- Alonso, L., Creuzé-des-Châtelliers, C., Trabac, T., Dubost, A., Moëgne-Loccoz, Y., & Pommier, T. (2018). Rock substrate rather than black stain alterations drives microbial community structure in the passage of Lascaux Cave. *Microbiome*, 6(1), 216. doi: 10.1186/s40168-018-0599-9
- Alonso, L., Pommier, T., Kaufmann, B., Dubost, A., Chapulliot, D., Doré, J., ... Moëgne-Loccoz, Y. (2019). Anthropization level of Lascaux Cave microbiome shown by regional-scale comparisons of pristine and anthropized caves. *Molecular Ecology*, 28(14), 3383–3394. doi: 10.1111/mec.15144
- Barracough, T. G., Balbi, K. J., & Ellis, R. J. (2012). Evolving Concepts of Bacterial Species. *Evolution Biology*, 39, 148–157. doi: 10.1007/s11692-012-9181-8
- Barton, H. A., & Jurado, V. (2007). *What's Up Down There? Microbial Diversity in Caves Microorganisms in caves survive under nutrient-poor conditions and are metabolically versatile and unexpectedly diverse*. Retrieved from https://pdfs.semanticscholar.org/cc59/ea424434255719237886aa7de76b3c99b24e.pdf?_ga=2.138095863.618314355.1561451191-1159738399.1560451939
- Barton, H. A., Taylor, N. M., Kreate, M. P., Springer, A. C., Oehrle, S. A., & Bertog, J. L. (2007). The Impact of Host Rock Geochemistry on Bacterial Community Structure in Oligotrophic Cave Environments. *International Journal of Speleology*, 36 (2), 93–104. Retrieved from www.ij.speleo.it
- Bastian, F., Jurado, V., Novakova, A., Alabouvette, C., & Saiz-Jimenez, C. (2010). The microbiology of Lascaux Cave. *Microbiology*, 156(3), 644–652. doi: 10.1099/mic.0.036160-0
- Bastian, M., Heymann, S., & Jacomy, M. (2009). Gephi: An Open Source Software for Exploring and Manipulating Networks. *Proceedings of the Third International ICWSM Conference (2009)*, 361–362. doi: 10.1136/qshc.2004.010033
- Bercea, S., Năstase-Bucur, R., Moldovan, O. T., Kenesz, M., & Constantin, S. (2019). Yearly microbial cycle of human exposed surfaces in show caves. *Subterranean Biology*, 31, 1–14. doi: 10.3897/subtbiol.31.34490
- Bhullar, K., Waglechner, N., Pawlowski, A., Koteva, K., Banks, E. D., Johnston, M. D., ... Wright, G. D. (2012). Antibiotic Resistance Is Prevalent in an Isolated Cave Microbiome. *PLoS ONE*, 7(4), e34953. doi: 10.1371/journal.pone.0034953
- Boiron, P., Provost, F., Chevrier, G., & Dupont, B. (1992). Review of nocardial infections in France 1987 to 1990. *European Journal of Clinical Microbiology & Infectious Diseases*, 11(8), 709–714. doi: 10.1007/BF01989975
- Cheng, K., Rong, X., Pinto-Tomás, A. A., Fernández-Villalobos, M., Murillo-Cruz, C., & Huang, Y. (2015). Population genetic analysis of *Streptomyces albidoflavus* reveals habitat barriers to homologous recombination in the diversification of streptomycetes. *Applied and Environmental Microbiology*, 81(3), 966–975. doi: 10.1128/AEM.02925-14
- Colaco, C. A., & MacDougall, A. (2014). Mycobacterial chaperonins: the tail wags the dog. *FEMS Microbiology Letters*, 350(1), 20–24. doi: 10.1111/1574-6968.12276
- Cuezva, S., Fernandez-Cortes, A., Porca, E., Pašić, L., Jurado, V., Hernandez-Marine, M., ... Saiz-Jimenez, C. (2012). The biogeochemical role of Actinobacteria in Altamira Cave, Spain. *FEMS Microbiology Ecology*, 81(1), 281–290. doi: 10.1111/j.1574-6941.2012.01391.x
- Cuezva, S., Sanchez-Moral, S., Saiz-Jimenez, C., & Cañaveras, J. C. (2009). Microbial communities and associated mineral fabrics in Altamira Cave, Spain. *International Journal of Speleology*, 38(1), 83–92. doi: 10.5038/1827-806x.38.1.9
- De La Rosa, J. M., Martin-Sanchez, P. M., Sanchez-Cortes, S., Hermosin, B., Knicker, H., & Saiz-Jimenez, C. (2017).

625 Structure of melanins from the fungi *Ochroconis lascauxensis* and *Ochroconis anomala* contaminating rock art
626 in the Lascaux Cave. *Scientific Reports*, 7(1), 1–11. doi: 10.1038/s41598-017-13862-7

627 De Mandal, S., Chatterjee, R., & Kumar, N. S. (2017). Dominant bacterial phyla in caves and their predicted functional
628 roles in C and N cycle. *BMC Microbiology*, 17(1), 1–9. doi: 10.1186/s12866-017-1002-x

629 Deagle, B. E., Jarman, S. N., Coissac, E., Pompanon, F., & Taberlet, P. (2014). DNA metabarcoding and the
630 cytochrome *c* oxidase subunit I marker: not a perfect match. *Biology Letters*, 10(9), 20140562. doi:
631 10.1098/rsbl.2014.0562

632 Duchêne, A.-M., Kieser, H. M., Hopwood, D. A., Thompson, C. J., & Mazodier, P. (1994). Characterization of two
633 groEL genes in *Streptomyces coelicolor* A3 (2). *Gene*, 144(1), 97–101. doi: 10.1016/0378-1119(94)90210-0

634 El-Sayed, M. H. (2012). Di-(2-ethylhexyl) Phthalate, a Major Bioactive Metabolite with Antimicrobial and Cytotoxic
635 Activity Isolated from the Culture Filtrate of Newly Isolated Soil *Streptomyces* (*Streptomyces mirabilis* Strain
636 NSQu-25). *World Applied Sciences Journal*, 20(9), 1202–1212. doi: 10.5829/idosi.wasj.2012.20.09.2868

637 Fang, B.-Z., Salam, N., Han, M.-X., Jiao, J.-Y., Cheng, J., Wei, D.-Q., ... Li, W.-J. (2017). Insights on the Effects of
638 Heat Pretreatment, pH, and Calcium Salts on Isolation of Rare Actinobacteria from Karstic Caves. *Frontiers in*
639 *Microbiology*, 8. doi: 10.3389/fmicb.2017.01535

640 Fernandez-Cortes, A., Cuezva, S., Sanchez-Moral, S., Cañaveras, J. C., Porca, E., Jurado, V., ... Saiz-Jimenez, C.
641 (2011). Detection of human-induced environmental disturbances in a show cave. *Environmental Science and*
642 *Pollution Research*, 18(6), 1037–1045. doi: 10.1007/s11356-011-0513-5

643 Fox, G. E., Wisotzkey, J. D., & Jurtshuk, P. (1992). How Close Is Close: 16S rRNA Sequence Identity May Not Be
644 Sufficient To Guarantee Species Identity. *Int. J. Syst. Bacteriol.*, 42(1), 166–170. doi: 10.1099/00207713-42-1-
645 166

646 Frey-Klett, P., Burlinson, P., Deveau, A., Barret, M., Tarkka, M., & Sarniguet, A. (2011). Bacterial-fungal
647 interactions: hyphens between agricultural, clinical, environmental, and food microbiologists. *Microbiology and*
648 *Molecular Biology Reviews* : *MMBR*, 75(4), 583–609. doi: 10.1128/MMBR.00020-11

649 Fruchterman, T. M. J., & Reingold, E. M. (1991). Graph drawing by force-directed placement. *Software: Practice and*
650 *Experience*, 21(11), 1129–1164. doi: 10.1002/spe.4380211102

651 Gao, B., & Gupta, R. (2012). Phylogenetic framework and molecular signatures for the main clades of the phylum
652 Actinobacteria. *Microbiology and Molecular Biology Reviews*, 76(1), 66–112. Retrieved from
653 <http://mmb.asm.org/content/76/1/66.short>

654 Gonzalez-Pimentel, J. L., Miller, A. Z., Jurado, V., Laiz, L., Pereira, M. F. C., & Saiz-Jimenez, C. (2018). Yellow
655 coloured mats from lava tubes of La Palma (Canary Islands, Spain) are dominated by metabolically active
656 Actinobacteria. *Sci. Rep.*, 8(1), 1–11. doi: 10.1038/s41598-018-20393-2

657 Groth, I., Vettermann, R., Schuetze, B., Schumann, P., & Saiz-Jimenez, C. (1999). Actinomycetes in Karstic caves of
658 northern Spain (Altamira and Tito Bustillo). *Journal of Microbiological Methods*, 36(1–2), 115–122. doi:
659 10.1016/S0167-7012(99)00016-0

660 Hall, T., Biosciences, I., & Carlsbad, C. (2011). BioEdit: An important software for molecular biology. *GERF Bulletin*
661 *of Biosciences*, 2(June), 60–61. doi: 10.1002/prot.24632

662 Hamed, J., Kafshnouchi, M., & Ranjbaran, M. (2019). A Study on actinobacterial diversity of Hampoeil cave and
663 screening of their biological activities. *Saudi Journal of Biological Sciences*, 26(7), 1587–1595. doi:
664 10.1016/j.sjbs.2018.10.010

665 Hamm, P. S., Caimi, N. A., Northup, D. E., Valdez, E. W., Buecher, D. C., Dunlap, C. A., ... Porras-Alfaro, A. (2017).
666 *Western Bats as a Reservoir of Novel Streptomyces Species with Antifungal Activity*. doi: 10.1128/AEM.03057-
667 16

668 Hammer, Ø., Harper, D. A. T., & Ryan, P. D. (2001). *PAST: PALEONTOLOGICAL STATISTICS SOFTWARE*
669 *PACKAGE FOR EDUCATION AND DATA ANALYSIS*. 4(1), 1–9.

Herlemann, D. P. R., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J. J., & Andersson, A. F. (2011). Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME Journal*, 5(10), 1571–1579. doi: 10.1038/ismej.2011.41

Hirsch, P. R., Mauchline, T. H., & Clark, I. M. (2010). Culture-independent molecular techniques for soil microbial ecology. *Soil Biol. Biochem.*, 42(6), 878–887. doi: 10.1016/j.soilbio.2010.02.019

Hoyos, M., Cañaveras, J. C., Sánchez-Moral, S., Sanz-Rubio, E., & Soler, V. (1998). Microclimatic characterization of a karstic cave: human impact on microenvironmental parameters of a prehistoric rock art cave (Candamo Cave, northern Spain). In *Environmental Geology* (Vol. 33). Springer-Verlag. Retrieved from Springer-Verlag website: <https://link.springer.com/content/pdf/10.1007%2Fs002540050242.pdf>

Johnston, M. D., Muench, B. A., Banks, E. D., & Barton, H. A. (2012). Human urine in Lechuguilla cave: the microbiological impact and potential for bioremediation. *Journal of Cave and Karst Studies*, 74(3), 278–291. doi: 10.4311/2011MB0227

Jurado, V., Boiron, P., Kroppenstedt, R. M., Laurent, F., Couble, A., Laiz, L., ... Rodríguez-Nava, V. (2008). *Nocardia altamirensis* sp. nov., isolated from Altamira cave, Cantabria, Spain. *International Journal of Systematic and Evolutionary Microbiology*, 58(9), 2210–2214. doi: 10.1099/ijs.0.65482-0

Jurado, Valme, Fernandez-Cortes, A., Cuezva, S., Laiz, L., Cañaveras, J. C., Sanchez-Moral, S., & Saiz-Jimenez, C. (2009). The fungal colonisation of rock-art caves: Experimental evidence. *Naturwissenschaften*, 96(9), 1027–1034. doi: 10.1007/s00114-009-0561-6

Jurado, Valme, Kroppenstedt, R. M., Saiz-Jimenez, C., Klenk, H. P., Mounié, D., Laiz, L., ... Rodríguez-Nava, V. (2009). *Hoyosella altamirensis* gen. nov., sp. nov., a new member of the order Actinomycetales isolated from a cave biofilm. *International Journal of Systematic and Evolutionary Microbiology*, 59(12), 3105–3110. doi: 10.1099/ijs.0.008664-0

Jurado, Valme, Laiz, L., Rodríguez-Nava, V., Boiron, P., Hermosin, B., Sanchez-Moral, S., & Saiz-Jimenez, C. (2010). Pathogenic and opportunistic microorganisms in caves. *International Journal of Speleology*, 39 (1), 15–24. Retrieved from www.ij.speleo.it

Kageyama, A., Yazawa, K., Kudo, T., Taniguchi, H., Nishimura, K., & Mikami, Y. (2004). First Isolates of *Nocardia abscessus* from Humans and Soil in Japan. *Nippon Ishinkin Gakkai Zasshi*, 45(1), 17–21. doi: 10.3314/jjmm.45.17

Kominek, L. A. (1972). Biosynthesis of novobiocin by *Streptomyces niveus*. *Antimicrobial Agents and Chemotherapy*, 1(2), 123–134. doi: 10.1128/AAC.1.2.123

Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013). Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the miseq illumina sequencing platform. *Appl. Environ. Microbiol.*, 79(17), 5112–5120. doi: 10.1128/AEM.01043-13

Kumar, C. M. S., Mande, S. C., & Mahajan, G. (2015). Multiple chaperonins in bacteria--novel functions and non-canonical behaviors. *Cell Stress & Chaperones*, 20(4), 555–574. doi: 10.1007/s12192-015-0598-8

Kumar, A. DE, Muthiyan, R., Sunder, J., Bhattacharya, D., Kundu, A., & Dam Roy, S. (2019). *Profiling Bacterial Diversity of B2 Cave, A Limestone Cave of Baratang, Andaman and Nicobar Islands, India*. doi: 10.16943/ptinsa/2019/49589

Kuo, C. H., & Ochman, H. (2009a). Inferring clocks when lacking rocks: The variable rates of molecular evolution in bacteria. *Biology Direct*, 4(1), 35. doi: 10.1186/1745-6150-4-35

Kuo, C. H., & Ochman, H. (2009b). Inferring clocks when lacking rocks: The variable rates of molecular evolution in bacteria. *Biology Direct*, 4(1), 35. doi: 10.1186/1745-6150-4-35

Lavoie, K. H., Winter, A. S., Read, K. J. H., Hughes, E. M., Spilde, M. N., & Northup, D. E. (2017). Comparison of bacterial communities from lava cave microbial mats to overlying surface soils from Lava Beds National Monument, USA. *PLoS One*, 12(2). doi: 10.1371/journal.pone.0169339

Long, Y., Jiang, J., Hu, X., Zhou, J., Hu, J., & Zhou, S. (2019). Actinobacterial community in Shuanghe Cave using culture-dependent and -independent approaches. *World Journal of Microbiology and Biotechnology*, 35(10). doi: 10.1007/s11274-019-2713-y

Maciejewska, M., Adam, D., Naômé, A., Martinet, L., Tenconi, E., Calusinska, M., ... Rigali, S. (2017). Assessment of the potential role of *Streptomyces* in cave moonmilk formation. *Front. Microbiol.*, 8(JUN), 1–18. doi: 10.3389/fmicb.2017.01181

Maciejewska, M., Całusińska, M., Cornet, L., Adam, D., Pessi, I., Malchair, S., ... Rigali, S. (2018). High-Throughput Sequencing Analysis of the Actinobacterial Spatial Diversity in Moonmilk Deposits. *Antibiotics*, 7(2), 27. doi: 10.3390/antibiotics7020027

Magoč, T., & Salzberg, S. L. (2011). FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics*, 27(21), 2957–2963. doi: 10.1093/bioinformatics/btr507

Mammola, S. (2019). Finding answers in the dark: caves as models in ecology fifty years after Poulson and White. *Ecography*, 42(7), 1331–1351. doi: 10.1111/ecog.03905

Martin-Sanchez, P. M., Miller, A. Z., & Saiz-Jimenez, C. (2015). Lascaux Cave: An Example of Fragile Ecological Balance in Subterranean Environments. *Microbial Life of Cave Systems*. doi: 10.1515/9783110339888-015

Martin-Sanchez, P. M., Nováková, A., Bastian, F., Alabouvette, C., & Saiz-Jimenez, C. (2012). Use of Biocides for the Control of Fungal Outbreaks in Subterranean Environments: The Case of the Lascaux Cave in France. *Environmental Science & Technology*, 46(7), 3762–3770. doi: 10.1021/es2040625

Miller, A. Z., Garcia-Sanchez, A. M., Martin-Sanchez, P. M., Costa Pereira, M. F., Spangenberg, J. E., Jurado, V., ... Saiz-Jimenez, C. (2018). Origin of abundant moonmilk deposits in a subsurface granitic environment. *Sedimentology*, 65(5), 1482–1503. doi: 10.1111/sed.12431

Modra, H., Bartos, M., Hribova, P., Ulmann, V., Hubelova, D., Konecny, O., ... Pavlik, I. (2017). Detection of mycobacteria in the environment of the Moravian Karst (Bull Rock Cave and the relevant water catchment area): the impact of water sediment, earthworm castings and bat guano. *Veterinárni Medicína*, 62 (2017)(No. 3), 153–168. doi: 10.17221/126/2016-VETMED

Mulec, J. (2014). Human impact on underground cultural and natural heritage sites, biological parameters of monitoring and remediation actions for insensitive surfaces: Case of Slovenian show caves. *Journal for Nature Conservation*, 22(2), 132–141. doi: 10.1016/J.JNC.2013.10.001

Myronovskiy, M., Tokovenko, B., Manderscheid, N., Petzke, L., & Luzhetskyy, A. (2013). Complete genome sequence of *Streptomyces fulvissimus*. *Journal of Biotechnology*, 168(1), 117–118. doi: 10.1016/j.jbiotec.2013.08.013

Neral, A., Rajput, Y., & Rai, V. (2015). Genotypic and Serotypic confirmations of Bacterial community to Kotumsar cave for occupational safety of cave workers and visitors from pathogenic threats. *International Journal of Occupational Safety and Health*, 5(1), 22–27. doi: 10.3126/ijosh.v5i1.16631

Ogier, J. C., Pagès, S., Galan, M., Barret, M., & Gaudriault, S. (2019). RpoB, a promising marker for analyzing the diversity of bacterial communities by amplicon sequencing. *BMC Microbiology*, 19(1). doi: 10.1186/s12866-019-1546-z

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., Mcglinn, D., ... Maintainer, H. W. (2018). *Package “vegan” Title Community Ecology Package*. Retrieved from <https://cran.ism.ac.jp/web/packages/vegan/vegan.pdf>

Ondov, B. D., Bergman, N. H., & Phillippy, A. M. (2011). Interactive metagenomic visualization in a Web browser. *BMC Bioinformatics*, 12(1), 385. doi: 10.1186/1471-2105-12-385

Ortiz, M., Legatzki, A., Neilson, J. W., Fryslie, B., Nelson, W. M., Wing, R. A., ... Maier, R. M. (2014). Making a living while starving in the dark: metagenomic insights into the energy dynamics of a carbonate cave. *The ISME Journal*, 8(2), 478–491. doi: 10.1038/ismej.2013.159

- Porca, E., Jurado, V., Žgur-Bertok, D., Saiz-Jimenez, C., & Pašić, L. (2012). Comparative analysis of yellow microbial communities growing on the walls of geographically distinct caves indicates a common core of microorganisms involved in their formation. *FEMS Microbiology Ecology*, 81(1), 255–266. doi: 10.1111/j.1574-6941.2012.01383.x
- Price, M. N., Dehal, P. S., & Arkin, A. P. (2010). FastTree 2 – Approximately Maximum-Likelihood Trees for Large Alignments. *PLoS ONE*, 5(3), e9490. doi: 10.1371/journal.pone.0009490
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., ... Glöckner, F. O. (2012). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research*, 41(D1), D590–D596. doi: 10.1093/nar/gks1219
- R Core Team. (2018). R Core Team, R: A language and environment for statistical computing. *R Foundation for Statistical Computing, Vienna, Austria*, R Foundation for Statistical Computing.
- Rajput, Y., Rai, V., & Biswas, J. (2012). Screening of Bacterial Isolates from Various Microhabitat Sediments of Kotumsar Cave: A Cogitation on Their Respective Benefits and Expected Threats for Complete Biosphere and Tourists. *Research Journal of Environmental Toxicology*, 6(1), 13–24.
- Rangseekaew, P., & Pathom-aree, W. (2019). Cave Actinobacteria as Producers of Bioactive Metabolites. *Frontiers in Microbiology*, 10, 387. doi: 10.3389/fmicb.2019.00387
- Rangseekaew, P., & Pathom-Aree, W. (2019). Cave Actinobacteria as Producers of Bioactive Metabolites. *Frontiers in Microbiology*, 10, 387. doi: 10.3389/fmicb.2019.00387
- Rodríguez-Nava, V., Couble, A., Devulder, G., Flandrois, J.-P., Boiron, P., & Laurent, F. (2006). Use of PCR-restriction enzyme pattern analysis and sequencing database for hsp65 gene-based identification of *Nocardia* species. *Journal of Clinical Microbiology*, 44(2), 536–546. doi: 10.1128/JCM.44.2.536-546.2006
- Rodríguez-Nava, V., Khan, Z. U., Pötter, G., Kroppenstedt, R. M., Boiron, P., & Laurent, F. (2007). *Nocardia coubleae* sp. nov., isolated from oil-contaminated Kuwaiti soil. doi: 10.1099/ijs.0.64815-0
- Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: a versatile open source tool for metagenomics. *PeerJ*, 4, e2584. doi: 10.7717/peerj.2584
- Roxburgh, S. H., Shea, K., & Wilson, J. B. (2004). The intermediate disturbance hypothesis: Patch dynamics and mechanisms of species coexistence. *Ecology*, 85(2), 359–371. doi: 10.1890/03-0266
- Sagova-Mareckova, M., Ulanova, D., Sanderova, P., Omelka, M., Kamenik, Z., Olsovska, J., & Kopecky, J. (2015a). Phylogenetic relatedness determined between antibiotic resistance and 16S rRNA genes in actinobacteria Ecological and evolutionary microbiology. *BMC Microbiology*, 15(1), 81. doi: 10.1186/s12866-015-0416-6
- Sagova-Mareckova, M., Ulanova, D., Sanderova, P., Omelka, M., Kamenik, Z., Olsovska, J., & Kopecky, J. (2015b). Phylogenetic relatedness determined between antibiotic resistance and 16S rRNA genes in actinobacteria Ecological and evolutionary microbiology. *BMC Microbiology*, 15(1), 81. doi: 10.1186/s12866-015-0416-6
- Sánchez-Moral, S., Soler, V., Cañaveras, J. ., Sanz-Rubio, E., Van Grieken, R., & Gysels, K. (1999). Inorganic deterioration affecting the Altamira Cave, N Spain: quantitative approach to wall-corrosion (solutional etching) processes induced by visitors. *Science of The Total Environment*, 243–244, 67–84. doi: 10.1016/S0048-9697(99)00348-4
- Schabereiter-Gurtner, C., Saiz-Jimenez, C., Pinar, G., Lubitz, W., & Rölleke, S. (2002). Altamira cave Paleolithic paintings harbor partly unknown bacterial communities. *FEMS Microbiology Letters*, 211(1), 7–11. doi: 10.1111/j.1574-6968.2002.tb11195.x
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., ... Weber, C. F. (2009). Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.*, 75(23), 7537–7541. doi: 10.1128/AEM.01541-09
- Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W. S., & Huttenhower, C. (2011). Metagenomic

- biomarker discovery and explanation. *Genome Biology*, 12(6), R60. doi: 10.1186/gb-2011-12-6-r60
- Shade, A., Peter, H., Allison, S. D., Baho, D. L., Berga, M., Bürgmann, H., ... Gilbert, J. (2012). *Fundamentals of microbial community resistance and resilience*. doi: 10.3389/fmicb.2012.00417
- Syiemiong, D., & Jha, D. K. (2019). Antibacterial potential of Actinobacteria from a Limestone Mining Site in Meghalaya, India. *Journal of Pure and Applied Microbiology*, 13(2), 789–802. doi: 10.22207/JPAM.13.2.14
- Taj-Aldeen, S. J., Deshmukh, A., Doiphode, S., Wahab, A. A., Allangawi, M., Muzrkchi, A. Al, ... Meis, J. J. (2013). Molecular Identification and Susceptibility Pattern of Clinical *Nocardia* Species: Emergence of *Nocardia crassostreae* as an Agent of Invasive Nocardiosis. *Canadian Journal of Infectious Diseases and Medical Microbiology*, 24(2), e33–e38. Retrieved from <https://www.hindawi.com/journals/cjidmm/2013/256025/>
- Takeda, K., Kang, Y., Yazawa, K., Gonoi, T., & Mikami, Y. (2010). Phylogenetic studies of *Nocardia* species based on *gyrB* gene analyses. *Journal of Medical Microbiology*, 59(2), 165–171. doi: 10.1099/jmm.0.011346-0
- Telenti, A., Marchesi, F., Balz, M., Bally, F., Bottger, E. C., & Bodmer, T. (1993). Rapid identification of mycobacteria to the species level by polymerase chain reaction and restriction enzyme analysis. *Journal of Clinical Microbiology*, 31(2), 175–178.
- Tomczyk-Żak, K., & Zielenkiewicz, U. (2016). Microbial Diversity in Caves. *Geomicrobiol. J.*, 33(1), 20–38. doi: 10.1080/01490451.2014.1003341
- Van Elsas, J. D., Chiurazzi, M., Mallon, C. A., Elhottova, D., Krištůfek, V., & Salles, J. F. (2012). Microbial diversity determines the invasion of soil by a bacterial pathogen. *Proceedings of the National Academy of Sciences of the United States of America*, 109(4), 1159–1164. doi: 10.1073/pnas.1109326109
- Větrovský, T., & Baldrian, P. (2013). The Variability of the 16S rRNA Gene in Bacterial Genomes and Its Consequences for Bacterial Community Analyses. *PLoS ONE*, 8(2), e57923. doi: 10.1371/journal.pone.0057923
- Vos, M., Quince, C., Pijl, A. S., de Hollander, M., & Kowalchuk, G. A. (2012). A comparison of *rpoB* and 16S rRNA as markers in pyrosequencing studies of bacterial diversity. *PLoS ONE*, 7(2). doi: 10.1371/journal.pone.0030600
- Watanabe, K., Shinagawa, M., Amishima, M., Iida, S., Yazawa, K., Kageyama, A., ... Mikami, Y. (2006). First Clinical Isolates of *Nocardia carnea*, *Nocardia elegans*, *Nocardia paucivorans*, *Nocardia puris* and *Nocardia takedensis* in Japan. *Nippon Ishinkin Gakkai Zasshi*, 47(2), 85–89. doi: 10.3314/jjmm.47.85
- Yasir, M. (2018). Analysis of bacterial communities and characterization of antimicrobial strains from cave microbiota. *Brazilian Journal of Microbiology*, 49(2), 248–257. doi: 10.1016/j.bjm.2017.08.005
- Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., ... Glöckner, F. O. (2014). The SILVA and “all-species Living Tree Project (LTP)” taxonomic frameworks. *Nucleic Acids Research*, 42(D1), 643–648. doi: 10.1093/nar/gkt1209
- Zhang, B., Wu, X., Tai, X., Sun, L., Wu, M., Zhang, W., ... Dyson, P. (2019). Variation in Actinobacterial Community Composition and Potential Function in Different Soil Ecosystems Belonging to the Arid Heihe River Basin of Northwest China. *Frontiers in Microbiology*, 10, 2209. doi: 10.3389/fmicb.2019.02209
- Zhu, H. Z., Zhang, Z. F., Zhou, N., Jiang, C. Y., Wang, B. J., Cai, L., & Liu, S. J. (2019). Diversity, distribution and co-occurrence patterns of bacterial communities in a karst cave system. *Frontiers in Microbiology*, 10(JULY). doi: 10.3389/fmicb.2019.01726

Supplementary Information for:

Usefulness of *hsp65* marker as a complement to 16S rRNA sequences to demonstrate differences in *Actinobacteria* community according to anthropization of limestone caves and occurrence of visual marks in Lascaux Cave

Buresova Andrea^{1,2,3}, Kopecky Jan³, Sagova-Mareckova Marketa³, Alonso Lise¹, Vautrin Florian¹, Moëgne-Loccoz Yvan¹, Rodriguez-Nava Veronica¹

¹Univ Lyon, Université Claude Bernard Lyon 1, CNRS, INRAE, VetAgro Sup, UMR5557 Ecologie Microbienne, F-69622 Villeurbanne, France

²Department of Ecology, Faculty of Science, Charles University in Prague, Vinicna 7, Prague, Czech Republic,

³Laboratory for Diagnostics and Epidemiology of Microorganisms, Crop Research Institute, Drnovska 507, CZ-16106 Prague 6, Ruzyně, Czech Republic

This PDF file includes:

Figs. S1 to S5

Tables S1 to S7




Fig. S1. Indices of alpha diversity (richness: Chao 1, evenness: Simpson evenness, diversity: inverse Simpson) for *Actinobacteria* communities among caves, different locations within Lascaux Cave, and marked/unmarked areas. Significant differences between groups are shown with letters (ANOVA and Tukey post hoc test; $P < 0.05$; 97% OTU cutoff).




Fig.

Fig S2. Venn diagrams showing unique and shared OTUs between A. different caves and B. different locations within Lascaux Cave based on *hsp65* and 16S rRNA gene markers (99% OTU cutoff).




Fig. S3. Taxonomical compositions of the core microbiome between A. caves and B. different locations within Lascaux Cave based on the *hsp65* and 16S rRNA gene markers (99% OTU cutoff).




Fig. S4. Average proportions of species within the genus *Streptomyces* in the *Actinobacteria* *hsp65* amplicon sequence libraries significantly different in marked (S) and unmarked areas (U) of Lascaux Cave (metastats, $P < 0.05$).




Fig. S5. Average proportions of genus *Nocardia* in the *Actinobacteria hsp65* gene amplicon sequence libraries in different caves (Rouffignac, Lascaux, Mouflon and Reille) and different locations within Lascaux Cave (Sas-1, Passage-B, Passage-IP, Apse and Diacase).

Table S1. Location of cave samples and list of gene markers sequenced using Illumina MiSeq

Sample ID	Cave name	Room	Area	Obtained sequences
Apse-S1	Lascaux	Apse	Visual mark	16S rRNA gene
Apse-S2	Lascaux	Apse		<i>hsp65</i> , 16S rRNA gene
Apse-S3	Lascaux	Apse		<i>hsp65</i> , 16S rRNA gene
Apse-U1	Lascaux	Apse	Unmarked area	<i>hsp65</i> , 16S rRNA gene
Apse-U2	Lascaux	Apse		<i>hsp65</i> , 16S rRNA gene
Apse-U3	Lascaux	Apse		<i>hsp65</i> , 16S rRNA gene
Apse-DZ1	Lascaux	Apse	Visual mark	16S rRNA gene
Apse-DZ2	Lascaux	Apse		<i>hsp65</i>
Apse-DZ3	Lascaux	Apse		<i>hsp65</i> , 16S rRNA gene
Apse-DZ4	Lascaux	Apse		16S rRNA gene
Apse-DZ5	Lascaux	Apse		<i>hsp65</i> , 16S rRNA gene
Apse-DZ6	Lascaux	Apse		<i>hsp65</i> , 16S rRNA gene
Apse-UDZ1	Lascaux	Apse	Unmarked area	16S rRNA gene
Apse-UDZ2	Lascaux	Apse		16S rRNA gene
Apse-UDZ3	Lascaux	Apse		<i>hsp65</i> , 16S rRNA gene
Apse-UDZ4	Lascaux	Apse		<i>hsp65</i> , 16S rRNA gene
Apse-UDZ5	Lascaux	Apse		16S rRNA gene
Apse-UDZ6	Lascaux	Apse		<i>hsp65</i> , 16S rRNA gene
Dia-S1	Lascaux	Diaclase	Visual mark	<i>hsp65</i> , 16S rRNA gene
Dia-S2	Lascaux	Diaclase		<i>hsp65</i> , 16S rRNA gene
Dia-S3	Lascaux	Diaclase		16S rRNA gene
Dia-U1	Lascaux	Diaclase	Unmarked area	<i>hsp65</i> , 16S rRNA gene
Dia-U2	Lascaux	Diaclase		<i>hsp65</i> , 16S rRNA gene
Dia-U3	Lascaux	Diaclase		<i>hsp65</i> , 16S rRNA gene
Passage B-S1	Lascaux	Passage banks	Visual mark	<i>hsp65</i> , 16S rRNA gene
Passage B-S2	Lascaux	Passage banks		<i>hsp65</i> , 16S rRNA gene
Passage B-S3	Lascaux	Passage banks		<i>hsp65</i> , 16S rRNA gene
Passage B-U1	Lascaux	Passage banks	Unmarked area	<i>hsp65</i> , 16S rRNA gene
Passage B-U2	Lascaux	Passage banks		<i>hsp65</i> , 16S rRNA gene
Passage B-U3	Lascaux	Passage banks		<i>hsp65</i> , 16S rRNA gene
Passage IP-S1	Lascaux	Passage inclined planes	Visual mark	<i>hsp65</i> , 16S rRNA gene
Passage IP-S2	Lascaux	Passage inclined planes		<i>hsp65</i> , 16S rRNA gene
Passage IP-S3	Lascaux	Passage inclined planes		<i>hsp65</i> , 16S rRNA gene
Passage IP-U1	Lascaux	Passage inclined planes	Unmarked area	<i>hsp65</i> , 16S rRNA gene
Passage IP-U2	Lascaux	Passage inclined planes		<i>hsp65</i> , 16S rRNA gene
Passage IP-U3	Lascaux	Passage inclined planes		16S rRNA gene
SAS-1-S1	Lascaux	Sas-1 (airlock-1 entrance zone, compartment 2)	Visual mark	<i>hsp65</i> , 16S rRNA gene
SAS-1-S2	Lascaux	Sas-1 (airlock-1 entrance zone, compartment 2)		<i>hsp65</i> , 16S rRNA gene
SAS-1-S3	Lascaux	Sas-1 (airlock-1 entrance zone, compartment 2)		<i>hsp65</i> , 16S rRNA gene
SAS-U1	Lascaux	Sas-1 (airlock-1 entrance zone, compartment 2)	Unmarked area	<i>hsp65</i>
SAS-U3	Lascaux	Sas-1 (airlock-1 entrance zone, compartment 2)		<i>hsp65</i>
Rouf1	Rouffignac	x	x	<i>hsp65</i> , 16S rRNA gene
Rouf2	Rouffignac			<i>hsp65</i>
Rouf3	Rouffignac			<i>hsp65</i>
Rouf4	Rouffignac			<i>hsp65</i> , 16S rRNA gene
Rouf5	Rouffignac			<i>hsp65</i> , 16S rRNA gene
Rouf6	Rouffignac			16S rRNA gene
Mouf1	Mouflon	x	x	16S rRNA gene
Mouf2	Mouflon			<i>hsp65</i> , 16S rRNA gene
Mouf3	Mouflon			<i>hsp65</i>
Mouf4	Mouflon			<i>hsp65</i> , 16S rRNA gene
Mouf5	Mouflon			<i>hsp65</i>
Mouf6	Mouflon			16S rRNA gene
Reil1	Reille	x	x	<i>hsp65</i>
Reil2	Reille			16S rRNA gene
Reil3	Reille			<i>hsp65</i> , 16S rRNA gene
Reil4	Reille			<i>hsp65</i> , 16S rRNA gene
Reil5	Reille			16S rRNA gene

Table S2. Genomes of *Actinobacteria* reference strains from the orders *Streptomycetales* and *Corynebacteriales* that were retrieved from the Integrated Microbial Genome and Microbiome database and used for *in silico* analysis – comparison of the *hsp65* and 16S rRNA partial gene sequences.

Order	Family	Genus	Species	Order	Family	Genus	Species
Corynebacteriales	Corynebacteriaceae	Corynebacterium	Corynebacterium ammoniagenes KCCM 40472	Streptomycetales	Streptomycetaceae	Streptomyces	Streptomyces agglomeratus 37742
			Corynebacterium atypicum R2070				Streptomyces albus NK660
			Corynebacterium aurimucosum ATCC 700975				Streptomyces albus J1074
			Corynebacterium callunae DSM 20147				Streptomyces autolyticus CGMCC0516
			Corynebacterium camporealensis DSM 44610				Streptomyces avermitilis MA-4680
			Corynebacterium crudilactis JZ16				Streptomyces bingchengensis BCW-1
			Corynebacterium diphtheriae PW8				Streptomyces cattleya DSM 46488
			Corynebacterium glutamicum CP				Streptomyces cattleya NRRL 8057
			Corynebacterium pseudotuberculosis 1002B				Streptomyces clavuligerus F613-1
			Corynebacterium pseudotuberculosis MB14				Streptomyces coelicolor A3(2)
			Corynebacterium ulcerans 131001				Streptomyces collinus Tu 365
			Corynebacterium xerosis DSM 20743				Streptomyces flavogriseus IAF 45
			Gordonia polyisoprenivorans VH2 DSM 44266				Streptomyces formicae KY5
							Streptomyces fulvissimus DSM 40593
							Streptomyces globisporus C-1027
							Streptomyces griseus subsp. griseus NBRC13350
	Gordoniaceae	Gordonia	Gordonia sp. 1D				Streptomyces hygroscopicus jinggangensis 5008
			Gordonia sp. KTR9				Streptomyces hygroscopicus jinggangensis TL01
			Gordonia sp. QH-11				Streptomyces katrae S3/4
			Gordonia sp. YC-JH1				Streptomyces leeuwenhoekii DSM 42122
			Gordonia terrae 3612				Streptomyces lincolnensis NRRL 2936
							Streptomyces lividans TK24
							Streptomyces lunae lactis MM109
							Streptomyces lydicus A02
	Mycobacteriaceae	Mycobacterium	Mycobacterium avium 2285 (R)				Streptomyces lydicus GS93/23
			Mycobacterium avium paratuberculosis E1				Streptomyces malaysiensis DSM 4137
			Mycobacterium bovis 1595				Streptomyces niveus SCSIO 3406
			Mycobacterium canettii CIPT 140070008				Streptomyces pactum KLBMP 5084
			Mycobacterium indicus pranii MTCC 9506				Streptomyces parvulus 2297
			Mycobacterium intracellulare 1956				Streptomyces pluripotens MUSC 137
			Mycobacterium kansasii 662				Streptomyces pristinaespiralis HCCB 10218
			Mycobacterium lepraemurium Hawaii				Streptomyces punicisabiei TW1S1
			Mycobacterium leprae TN NC 002677				Streptomyces purpureus KA281
			Mycobacterium liflandii 128FXT				Streptomyces reticuli TUE45
			Mycobacterium marinum M				Streptomyces sampsonii KJ40
			Mycobacterium marseillense FLAC0026				Streptomyces scabiei 87.22
			Mycobacterium shigaense UN-152				Streptomyces sp. PAMC26508
			Mycobacterium simiae MO323				Streptomyces sp. S10(2016)
			Mycobacterium tuberculosis 0A092DS				Streptomyces sp. SirexAA-E
			Mycobacterium tuberculosis 26105				Streptomyces sp. Tu6071
	Nocardiaceae	Nocardia	Nocardia brasiliensis ATCC 700358				Streptomyces sp. XZHG99
			Nocardia brasiliensis FDAARGOS 352				Streptomyces venezuelae Shinobu 719
			Nocardia cyriacigeorgica MDA3349				Streptomyces violaceusniger Tu 4113
			Nocardia farcinica IFM 10152				Streptomyces xiamenensis 318
			Nocardia nova SH22a				Streptomyces xiamenensis MCCC 1A01550
			Nocardia seriolae EM150506				Streptomyces xinghaiensis S187
			Nocardia seriolae UTF1				
			Nocardia terpenica NC YFY NT001				
	Nocardiaceae	Rhodococcus	Rhodococcus aetherivorans IcdP1			Kitakatospora	Kitakatospora setae KM-6054
			Rhodococcus coprophilus NCTC 10994				
			Rhodococcus equi 103S				
			Rhodococcus erythropolis CCM2595				
			Rhodococcus erythropolis PR4				
			Rhodococcus fascians D188				
			Rhodococcus hoagii DSSKP-R-001				
			Rhodococcus opacus B4				
			Rhodococcus opacus PD630				
			Rhodococcus pyridinivorans SB3094				
			Rhodococcus qingshengii dj1-6-2				
			Rhodococcus rhodochrous NCTC 10210				
			Rhodococcus ruber P14				
			Rhodococcus sp. BCP1				
			Rhodococcus sp. RHA1				

Table S3. Percentage of complementarity between the hsp65-specific forward (TB11) and reverse (TB12) primers and the hsp65 sequence database with respect to the number of mismatches.

	Number of mismatches	Match to <i>hsp65</i> [%]	Match to GROESL [%]
Forward primer	0	30.8	0
	1	74.8	0
	2	86	4.2
	3	98.1	56.2
	4	99.1	85.4
	5	100	100
Reverse primer	0	6.5	0
	1	44.9	0
	2	89.7	52.1
	3	96.3	64.6
	4	100	81.2
	5	100	89.6

Table S4. Percent coverage of the primers to the *Actinobacteria* RefSeq representative genomes from NCBI of different classes and orders. Only targets with no mismatches on the last three nucleotides of the 3' end and no more than 4 overall mismatches were included.

Class	Order	Numb. of RefSeq representative genomes	Percentual primer matches
Acidimicrobiia		8	38
Coriobacteriia		63	10
Nitrospirillum		5	0
Rubrobacteria		7	43
Thermoleophilum		7	14
Actinobacteria			
	Acidothermales	1	0
	Actinomycetales	75	27
	Actinopolysporales	8	100
	Bifidobacteriales	87	68
	Catenulisporales	3	100
	Corynebacteriales	473	83
	Frankiales	18	94
	Geodermatophilales	34	100
	Glycomycetales	13	77
	Kineosporiales	5	80
	Micrococcales	482	53
	Micromonosporales	106	100
	Nakamurellales	5	20
	Propionibacteriales	120	63
	Pseudonocardiales	151	74
	Streptomycetales	348	95
	Streptosporangiales	112	88

Table S5. Number of identified Actinobacteria species by hsp65 marker for respective genus

Family	Genus (number of recovered species)			
<i>Nocardiaceae</i>	<i>Nocardia</i> (24)	<i>Rhodococcus</i> (10)		
<i>Streptomycetaceae</i>	<i>Streptomyces</i> (30)	<i>Kitasatospora</i> (1)		
<i>Mycobacteriaceae</i>	<i>Mycobacterium</i> (26)			
<i>Pseudonocardiaceae</i>	<i>Amycolatopsis</i> (7)	<i>Pseudonocardia</i> (4)	<i>Lentzea</i> (2)	<i>Actinomycetospora</i> (1)
	<i>Kibdelosporangium</i> (1)	<i>Thermocrispum</i> (1)	<i>Saccharothrix</i> (1)	<i>Alloactinosynnema</i> (1)
	<i>Sciscionella</i> (1)	<i>Actinoalloteichus</i> (1)	<i>Actinokineospora</i> (1)	
<i>Norcardioidaceae</i>	<i>Nocardioidea</i> (6)	<i>Kribbella</i> (2)	<i>Pimelobacter</i> (1)	
	<i>Marmoricola</i> (1)	<i>Aeromicrobium</i> (1)		
<i>Streptosporangiaceae</i>	<i>Microbispora</i> (1)	<i>Microtetrastroma</i> (1)	<i>Nonomuraea</i> (1)	<i>Planomonospora</i> (1)
	<i>Sinosporangium</i> (1)	<i>Streptosporangium</i> (1)		
<i>Micromonosporaceae</i>	<i>Micromonospora</i> (3)	<i>Longispora</i> (1)	<i>Catelliglobospora</i> (1)	<i>Plantactinospora</i> (1)
<i>Intrasporangiaceae</i>	<i>Janibacter</i> (3)	<i>Knoellia</i> (1)		
<i>Actinomycetaceae</i>	<i>Actinomyces</i> (3)			
<i>Jiangellaceae</i>	<i>Jaingella</i> (3)			
<i>Microbacteriaceae</i>	<i>Agromyces</i> (1)	<i>Microcella</i> (1)		
<i>Propionibacteriaceae</i>	<i>Cutibacterium</i> (1)	<i>Propionibacterium</i> (1)		
<i>Geodermatophilaceae</i>	<i>Geodermatophilus</i> (1)	<i>Modestobacter</i> (1)		
<i>Micrococcaceae</i>	<i>Micrococcus</i> (1)	<i>Rothia</i> (1)		
<i>Dermabacteraceae</i>	<i>Brachybacterium</i> (2)			
<i>Brevibacteriaceae</i>	<i>Brevibacterium</i> (2)			
<i>Corynebacteriaceae</i>	<i>Corynebacterium</i> (2)			
<i>Catenulisporaceae</i>	<i>Catenulispora</i> (1)			
<i>Dermacoccaceae</i>	<i>Kytococcus</i> (1)			
<i>Actinobacteria_incertain_sedis</i>	<i>Thermobispora</i> (1)			
<i>Nakamurellaceae</i>	<i>Nakamurella</i> (1)			
<i>Sporichthyaceae</i>	<i>Sporichthya</i> (1)			
<i>Nocardiopsaceae</i>	<i>Thermobifida</i> (1)			

Table S6. T test and F test for comparison of means and variation of diversity indices for 37 common samples of *hsp* 65 and 16S rRNA markers at 97% and 99% OTUs cutoffs. For each marker averages and standard deviations are indicated.

		<i>hsp</i> 65	16S rRNA	t test		F test	
				t	P-value	F	P-value
99% OTU							
	Chao-1	260.78 ± 137.051	182.17 ± 82.018	2.984	0.003	2.745	0.017
	Simpson evenness	0.286 ± 0.246	0.089 ± 0.048	4.72	<0.001	26.81	<0.001
	Inverse Simpson	6.57 ± 4.974	14.07 ± 6.289	5.664	<0.001	1.597	0.153
97% OTU							
	Chao-1	75.74 ± 42.262	89.115 ± 36.349	1.45	0.148	1.315	0.588
	Simpson evenness	0.38 ± 0.264	0.148 ± 0.105	4.908	<0.001	6.281	0.002
	Inverse Simpson	4.089 ± 2.742	9.106 ± 4.443	5.824	<0.001	2.637	0.013

Table S7. Differences between Actinobacteria communities between A. caves, B. different locations within Lascaux Cave, and marked and unmarked areas calculated by analysis of molecular variance (AMOVA) and homogeneity of molecular variance (HOMOVA) for the hsp65 and 16S rRNA gene markers (97% OTUs cutoff).

A. AMOVA														
homogeneity of molecular variance (ANOVA test) for the haplo2 and 100 mtDNA gene markers (2770 of 0.05 cutoff):														
hsp 65	Cave				Lascaux cave location						Surface area			
	Lascaux		Rouffignac		Reille		Sas-1		Passage B		Passage IP		Apse	
	Df	F, p	Df	F, p	Df	F, p	Df	F, p	Df	F, p	Df	F, p	Df	F, p
	Rouffignac	13	5.72***					Passage B	10	7.858**				
	Reille	11	3.614**	7	4.617			Passage IP	9	3.978	10	4.419**		
	Mouflon	12	6.265**	8	8.033*	6	4.736	Apse	16	4.505***	17	5.903***	16	2.572***
								Diaclase	9	4.518	10	5.635**	9	2.786
	Overall	20	5.317***					Overall	32	4.279***				3.193***
16S rRNA														
	Rouffignac	15	5.79***					Passage B						
	Reille	15	5.43***	7	6.263			Passage IP	8	4.863	11	9.013**		
	Mouflon	15	5.36**	7	6.329	7	2.260	Apse	19	4.09***	22	8.115***	22	2.788**
								Diaclase	8	2.723	11	5.79**	11	2.902**
	Overall	23	5.274***					Overall	37	4.656***				3.005***
Bonferromi pair-wise err. rate: 0.008														
0.008 > * p > 0.005 > ** p > 0.001 > ***														
Exp-wise err. rate: 0.05														
0.05 > * p > 0.01 > ** p > 0.001 > P														
B. HOMOVA														
hsp 65	Cave				Lascaux cave location						Surface area			
	B, p		Reille		Sas-1		Passage B		Passage IP		Apse		B, p	
	Df	F, p	Df	F, p	Df	F, p	Df	F, p	Df	F, p	Df	F, p	Df	F, p
	Rouffignac	(0.193)						Passage B						
	Lascaux	(0.303)	0.235*					Passage IP	0.006					
	Reille	(0.342)	0.193	0.010				Apse	0.246	0.367				
	Mouflon	(0.160)	0.026	0.359	0.295			Diaclase	0.270*	0.433*	0.005			
			Overall	0.573*				Overall	0.126	0.206	0.020	0.010		
16S rRNA														
	Rouffignac	(0.145)						Passage B						
	Lascaux	(0.276)	0.380					Passage IP	0.012					
	Reille	(0.23)	0.133	0.034				Apse	0.166	0.531*				
	Mouflon	(0.199)	0.062	0.105	0.014			Diaclase	0.364	1.227*	0.079			
			Overall	0.432*				Overall	0.415	1.166*	0.134*	0.031		
Bonferromi pair-wise err. rate: 0.008 > * p														
Exp-wise err. rate: 0.05 > p*														

-Part 2-

Paper II: Succession of microbial decomposers is determined by litter type but sites conditions drive decomposition rate



Succession of Microbial Decomposers Is Determined by Litter Type, but Site Conditions Drive Decomposition Rates

A. Buresova,^{a,b,c} J. Kopecky,^a V. Hrdinkova,^a Z. Kamenik,^d M. Omelka,^e M. Sagova-Mareckova^{a,f}

^aEpidemiology and Ecology of Microorganisms, Crop Research Institute, Prague, Czech Republic

^bDepartment of Ecology, Faculty of Science, Charles University, Prague, Czech Republic

^cEcologie Microbienne, Université Claude Bernard Lyon 1, Villeurbanne, France

^dLaboratory for Biology of Secondary Metabolism, Institute of Microbiology, Academy of Sciences of the Czech Republic, Prague, Czech Republic

^eDepartment of Probability and Mathematical Statistics, Faculty of Mathematics and Physics, Charles University in Prague, Prague, Czech Republic

^fDepartment of Microbiology, Nutrition and Dietetics, Faculty of Agrobiology, Food and Natural Resources, Czech University of Life Science in Prague, Prague, Czech Republic

ABSTRACT Soil microorganisms are diverse, although they share functions during the decomposition of organic matter. Thus, preferences for soil conditions and litter quality were explored to understand their niche partitioning. A 1-year-long litterbag transplant experiment evaluated how soil physicochemical traits of contrasting sites combined with chemically distinct litters of sedge (S), milkvetch (M) from a grassland, and beech (B) from forest site decomposition. Litter was assessed by mass loss; C, N, and P contents; and low-molecular-weight compounds. Decomposition was described by the succession of fungi, *Actinobacteria*, *Alphaproteobacteria*, and *Firmicutes*; bacterial diversity; and extracellular enzyme activities. The M litter decomposed faster at the nutrient-poor forest site, where the extracellular enzymes were more active, but microbial decomposers were not more abundant. *Actinobacteria* abundance was affected by site, while *Firmicutes* and fungi by litter type and *Alphaproteobacteria* by both factors. *Actinobacteria* were characterized as late-stage substrate generalists, while fungi were recognized as substrate specialists and site generalists, particularly in the grassland. Overall, soil conditions determined the decomposition rates in the grassland and forest, but successional patterns of the main decomposers (fungi and *Actinobacteria*) were determined by litter type. These results suggest that shifts in vegetation mostly affect microbial decomposer community composition.

IMPORTANCE Anthropogenic disturbance may cause shifts in vegetation and alter the litter input. We studied the decomposition of different litter types under soil conditions of a nutrient-rich grassland and nutrient-poor forest to identify factors responsible for changes in the community structure and succession of microbial decomposers. This will help to predict the consequences of induced changes on the abundance and activity of microbial decomposers and recognize if the decomposition process and resulting quality and quantity of soil organic matter will be affected at various sites.

KEYWORDS succession, enzyme activities, forest, grassland, organic matter

Microbial community structure and activity reflect changes of biotic (soil biota) and abiotic (climate, soil chemistry, and litter quality) factors as well as ecosystem functioning (1, 2). In particular, the complex environmental characteristics of a given site influence the pool of microbial species at the regional scale, while litter quality affects the dynamics of microorganisms coming from that pool at the local scale (3, 4).

It is still debatable whether environmental conditions or litter quality plays a larger role in decomposition (5). The transfer of litter between sites can differentiate between

Citation Buresova A, Kopecky J, Hrdinkova V, Kamenik Z, Omelka M, Sagova-Mareckova M. 2019. Succession of microbial decomposers is determined by litter type, but site conditions drive decomposition rates. *Appl Environ Microbiol* 85:e01760-19. <https://doi.org/10.1128/AEM.01760-19>.

Editor Claire Vieille, Michigan State University

Copyright © 2019 American Society for Microbiology. All Rights Reserved.

Address correspondence to M. Sagova-Mareckova, marketa.sagova@gmail.com.

Received 31 July 2019

Accepted 3 October 2019

Accepted manuscript posted online 11 October 2019

Published 27 November 2019

these two factors because site will have a predominant effect on decomposition if the local communities decompose the litter to which they are preadapted faster (6). Yet, if the litter type affects decomposition regardless of the site, the microbes are functionally redundant with respect to litter type (7). Therefore, to understand the factors driving microbial decomposition, connections between local and regional scales need to be considered over time, together with various litter types and origin (5, 8).

The roles of biotic and abiotic factors change over time as the litter is sequentially decomposed. As the chemical composition of litter changes, the associated microbial community and extracellular enzyme activities also change (9–11). The differential response of microbes to the changing litter chemistry is the basis for sorting microbes into specific guilds (10, 12). More specifically, a nutritious organic substrate is connected to a wide range of nonspecialized microbes, while recalcitrant organic substrates are decomposed only by a limited number of taxa, which can break the resistant organic structures (3, 13). It has been proposed that the identification of microbes from the late stage and/or specialized guilds should be the first step in predicting the consequences of environmental change connected to decomposition processes in temperate areas (5, 14).

Fungi have long been considered the main microbial decomposers catalyzing the decomposition of complex substrates (9, 15). Most studies focused on their decomposing abilities, while factors affecting them were found to be variable (10, 16, 17). However, the role of bacteria in decomposition has also been recently highlighted (7, 11). Particularly, *Actinobacteria*, *Proteobacteria*, *Firmicutes*, and *Bacteroidetes* produce extracellular enzymes and are active in substrate degradation (5, 18). Moreover, *Actinobacteria* are able to produce lignolytic enzymes and secondary metabolites and form filaments, which are strategies similar to those of fungi (19–23). Therefore, *Actinobacteria* may compete with fungi (15), but those two groups have rarely been studied together during the decomposition process. Their coexistence patterns, such as niche overlap and/or separation (3, 15) under the influence of various abiotic factors, could be a missing biotic predictor of decomposition (5, 15, 24).

The aim of this study was to explore the interplay of factors affecting the abundances and succession of main decomposers. We hypothesized that main decomposers will be site specific but will selectively inhabit different litter and shift their enzyme activities during the decomposition stages. In particular, we wanted to determine the importance of cross-kingdom coexistence and successional patterns of microbial communities in influencing the effects of litter type and soil condition on decomposition processes.

RESULTS

Low-molecular-weight compounds (measured by high-performance liquid chromatography [HPLC]) shifted over time (Fig. 1A). In January and March 2011, compounds were separated according to litter type, while in later months they were separated according to sites. The HPLC profiles of the milkvetch (M) litter type in January and March 2011 significantly correlated with C content ($R = 6.377$, $P < 0.001$) and most extracellular enzymes, such as β -glucosidase ($R = 7.491$, $P < 0.001$), acid phosphatase ($R = 7.776$, $P < 0.001$), and chitinase ($R = 6.152$, $P < 0.001$), according to the permutation test. Of all bacterial groups, the abundance of *Actinobacteria* most strongly correlated with the late-stage HPLC profile ($R = 10.569$, $P < 0.001$). *Actinobacteria* significantly correlated with soil water potential ($R = 2.347$, $P = 0.004$) and temperature ($R = 4.871$, $P < 0.001$).

Bacterial communities on the beech (B) and sedge (S) litter types separated from that on the M litter type, especially in January and March 2011 (Fig. 1B). Bacterial communities also shifted from the initial to late stages of decomposition (confirmed by Sammon's multidimensional scaling; see Fig. S1 in the supplemental material). Bacterial communities from a later month of decomposition significantly correlated with the abundances of *Actinobacteria* ($R = 2.5940$, $P < 0.001$), *Firmicutes* ($R = 1.2363$, $P < 0.001$), and bacterial diversity ($R = 0.9875$, $P < 0.001$) according to the permutation test. From January




FIG 1 RDA of low-molecular-weight compounds (A, Manhattan distance) and bacterial communities (B, Bray-Curtis distance) from beech (circle), milkvetch (square), and sedge (triangle) litters at the forest (full symbol) and grassland (empty symbol) sites during the decomposition experiment (5 replicates). Numbers 1 to 7 (1 to 4, respectively, for bacterial communities) correspond to the statistically significant linear variables. Blue arrows represent extracellular enzyme activities.

2011, bacterial communities on the M litter type were significantly correlated with C ($R = 1.3441$, $P < 0.001$) and most extracellular enzyme activities, such as β -glucosidase ($R = 1.3134$, $P < 0.001$), cellobiohydrolase ($R = 0.5805$, $P < 0.001$), chitinase ($R = 0.5606$, $P < 0.001$), and acid phosphatase ($R = 0.4410$, $P = 0.007$). Bacterial communities on the B and S litter types in January 2011 significantly correlated with laccase ($R = 0.8082$, $P < 0.001$). The bacterial communities on the M litter from March significantly correlated with N ($R = 0.7699$, $P < 0.001$) and P ($R = 0.5463$, $P = 0.002$). Late-stage bacterial communities significantly correlated with soil water tension ($R = 0.7697$, $P < 0.001$) and temperature ($R = 0.3895$, $P = 0.007$).

Litter mass loss. Loss of dry mass significantly differed between sites in all decomposition stages. At the forest site (F), the average residual dry mass was about 0.5 g lower than the grassland site (G) (ANOVA, $P < 0.001$) (see Table S1 in the supplemental material). Milkvetch (M) had the highest average mass loss (ANOVA, $P < 0.001$) (Table S1), while that of sedge (S) was significantly higher than beech (B) (ANOVA, $P < 0.001$) (Table S1) at both sites. Time courses of dry mass loss were similar for the same litter type regardless of the site (Fig. 2, P_{perm} [i.e., the P value from permutational multivariate analysis of variance using distance matrices {PERMANOVA}] < 0.001 , Table S1), while dispersion did not differ between the litter types (Fig. 2; Table S1) (P_{disp} [i.e., the P value from homogeneity of multivariate dispersions using dissimilarity measures] = 0.382). No significant differences were observed between the indigenous and exogenous origin of the litter (i.e., M and S litters did not decompose faster than B litter in the grassland and B litter did not decompose faster than M and S litters in the forest).

Carbon, nitrogen, and phosphorus contents in the litter. Carbon (C), nitrogen (N), and phosphorus (P) contents in litter significantly differed between the forest and grassland sites during decomposition (see Fig. S2 in the supplemental material). The N and C contents in all litter types were significantly higher at the forest site than the grassland site (ANOVA, $P < 0.001$) (Table S1), while P content showed the opposite relationship (ANOVA, $P = 0.001$) (Table S1). C content did not differ between litter types




FIG 2 Average dry mass of beech (brown), milkvetch (yellow), and sedge (green) litters at the grassland and forest sites during the decomposition experiment (\pm SD, 5 replicates). The x axis indicates sampling months (i.e., initial, before litterbag burial in November 2010 and then in 2-month periods from January 2011; overall, 12 months). The P values indicate differences between profiles' distances (PERMANOVA; $P_{Perm.}$) and dispersion (dispersion test, $P_{Disp.}$).

at both sites, while N and P contents significantly differed, with the highest content measured for the M litter ($P < 0.001$ for N, $P < 0.001$ for P) (Table S1).

HPLC analysis of extractable low-molecular-weight compounds. HPLC profiles significantly differed between sites and litter types within each site ($P < 0.001$ for all sampling times and litter types) (Table S2).

Extracellular enzyme activities. Overall, the activities of extracellular enzymes were significantly higher in the forest than in the grassland (ANOVA, β -glucosidase, $P < 0.001$; cellobiohydrolase, $P = 0.001$; β -xylosidase, $P < 0.001$; chitinase, $P < 0.001$; glucuronidase, $P = 0.003$; acid phosphatase, $P < 0.001$; arylsulfatase, $P < 0.001$; α -glucosidase, $P = 0.001$; laccase, $P = 0.007$). Manganese peroxidase activity did not differ between the sites. Overall, higher enzyme activities were measured for the M than the S or B litters for β -glucosidase ($M > B$ and S , $P < 0.001$), cellobiohydrolase ($M > S$, $P < 0.005$; $M > B$, $P < 0.001$; $S > B$, $P < 0.001$), β -xylosidase ($M > S$, $P = 0.015$; $M > B$, $P < 0.001$; $S > B$, $P < 0.001$), and acid phosphatase ($M > B$ and S , $P < 0.001$). The activities on the M and S litters were not statistically different but were higher than those of the B litter for chitinase ($P < 0.001$), glucuronidase ($P < 0.001$), and α -glucosidase ($P < 0.001$). The S litter had the highest laccase activity ($P = 0.007$) (Table S1; Fig. 3).

Time courses of enzyme activities for chitinase, arylsulfatase, laccase, and manganese peroxidase were similar for the same litter type regardless of the site, while dispersion did not differ between the litter types (chitinase, $P_{Perm.} = 0.005$, $P_{Disp.} = 0.914$; arylsulfatase, $P_{Perm.} = 0.004$, $P_{Disp.} = 0.052$; laccase, $P_{Perm.} = 0.009$, $P_{Disp.} = 0.796$; and manganese peroxidase, $P_{Perm.} < 0.001$, $P_{Disp.} = 0.385$) (Table S1).

Bacterial communities. A total of 9,366,330 16S rRNA gene sequences were obtained, of which 6,898,021 (i.e., 73.6%) were mapped to 13,987 operational taxo-




FIG 3 Extracellular enzyme activities on beech (brown), milkvetch (yellow), and sedge (green) litters at the forest and grassland sites during the decomposition experiment (\pm SD, 5 replicates). The x axes indicate sampling months (January 2011 to January 2012, i.e., 2 to 14 months after litterbag burial). The P values indicate differences between profiles' distances (PERMANOVA; $P_{\text{Perm.}}$) and dispersion (dispersion test, $P_{\text{Disp.}}$).




FIG 4 Average proportions of phyla in the bacterial 16S rRNA gene amplicon sequence libraries (5 replicates). Labels on the horizontal axis indicate sampling months (January to November 2011, i.e., 2 to 12 months after litterbag burial) and experimental site (G, grassland; F, forest).

nomic units (OTUs). The numbers per sample ranged between 12,630 and 78,406 sequences, with a median at 41,137 sequences.

Bacterial communities differed significantly between the forest and grassland sites in the first two sampling months for the M litter ($P = 0.009$ in January 2011, $P = 0.008$ in March) (Table S3) and in the last two sampling months for the B and S litters ($P = 0.008$ and $P = 0.006$, respectively, in July, and both species $P = 0.009$ in November 2011) (see Table S3 in the supplemental material). The bacterial community on the M litter significantly differed from that on the B and S litters (analysis of molecular variance [AMOVA], $P < 0.001$ and $P = 0.003$ at the grassland and forest sites, respectively, for both litter types) (see Table S4 in the supplemental material), with the bacterial communities not differing between the latter two litter types.

The proportions of phyla in the bacterial 16S rRNA gene amplicon sequence libraries were the highest for *Proteobacteria*, followed by *Actinobacteria*, *Bacteroidetes*, *Firmicutes*, and *Verrucomicrobia* (Fig. 4). These proportions increased over time for *Actinobacteria* (except on the B litter in November 2011 in the forest) and *Firmicutes* (especially from January to March 2011) and decreased for *Proteobacteria* and *Bacteroidetes* from January to November 2011 (Fig. 4). *Acidobacteria* proportions were the highest in the late sampling months on the M and S litters (Fig. 4) and in soil (see Fig. S3 in the supplemental material) in January and November 2011. *Verrucomicrobia* proportions were the highest in the middle sampling months (Fig. 4). Within the phylum *Actinobacteria* (see Fig. S4 in the supplemental material), the families *Streptomyetaceae* (mostly on S plants) and *Mycobacteriaceae* increased at both sites, while the proportions of *Pseudonocardiaceae*, *Gaiellaceae*, and *Promicromonosporaceae* increased mostly in the grassland, and the order *Frankiales* increased in the forest. The Trebon clade proportion increased in November 2011 on the S litter in the forest (Fig. 4).

The number of OTUs (cutoff, 97%) significantly contributing to differences between the sites was the lowest for the B litter and the highest for the M litter at both sites according to metastats (see Fig. S5 in the supplemental material). According to the rarefaction curves, the highest number of OTUs in January and March 2011 was found at the grassland site, especially for the B and S litter types. The number of OTUs was similar for all litter types for both sites in July, while the highest number of OTUs in November was measured for the M followed by S and B litter types at the grassland site (see Fig. S6 in the supplemental material). Rarefaction analysis showed that bacterial diversity differed between the sites (ANOVA; $F = 12.86$, $P < 0.001$). Bacterial diversity




FIG 5 Abundances of fungi (18S rRNA genes), *Actinobacteria*, *Alphaproteobacteria*, and *Firmicutes* (16S rRNA genes) assayed by qPCR on beech, milkvetch, and sedge at the forest and grassland sites during the decomposition experiment (\pm SD, 5 replicates). The x axes indicate sampling months (i.e., initial in November 2010 and then in 2-month periods from January 2011; overall, 14 months).

increased on all litter types over decomposition except for the S and B litters in the forest, where the diversity declined from July to November 2011 (see Fig. S7 in the supplemental material).

Abundance of soil microorganisms. Overall, *Alphaproteobacteria* gene copies (16S rRNA) were more abundant in the grassland than at the forest but only for January and March 2011 (ANOVA, $P = 0.009$) (Table S1) and on the M than the B and S litters (both $P < 0.001$). *Actinobacteria* gene copies (16S rRNA) were more abundant in the grassland than at the forest (ANOVA, $P < 0.001$) (Table S1) but did not differ between litter types. Fungi (18S rRNA) and *Firmicutes* (16S rRNA) gene copies did not differ between the sites. *Firmicutes* were more abundant on the M than the B and S litters (ANOVA, both $P < 0.001$) (Table S1). Fungi were the least abundant on the B litter compared with both the S (ANOVA, $P < 0.001$) (Table S1) and M litters ($P = 0.015$) (Fig. 5).

The time courses of *Actinobacteria* ($P_{Perm} = 0.021$; $P_{Disp} = 0.387$) and fungi ($P_{Perm} < 0.001$; $P_{Disp} = 0.505$) were similar for the same litter type regardless of site, while dispersion did not differ between the litter types.

DISCUSSION

Decomposition rate, extracellular enzyme activities, and litter chemistry were specific according to sites and within both sites according to litter type but not litter origin. Fungi, *Actinobacteria*, *Firmicutes*, *Proteobacteria*, *Bacteroidetes*, *Acidobacteria*, and *Verrucomicrobia* each had its own role in the decomposition process. Their successional patterns differed based on their requirements for site conditions or litter type. *Actinobacteria* were identified as late-stage site-specific substrate generalists, which was the opposite pattern to fungi. The dynamics of these two groups together with bacterial community composition were driven by litter type and were synchronized with the mass loss of the most nutritious litter of milkvetch (M).

Overall, the forest site had higher decomposition rates and enzyme activities than the grassland site. The beech forest soils are considered to be an environment limited by low soil N and P (25). Thus, nutrient limitations might stimulate microbes to produce enzymes to gain the deficient nutrients from complex substrates (26). This is also supported by our finding that P acquisition for all substrates, as well as enzyme activities, was significantly higher at the forest site (27, 28). In contrast, the nutrient-rich but warmer and drier soil with diverse sources of litter at the grassland site (29) probably sufficiently covered the demands of soil microbes (30). Consequently, a higher diversity of bacterial communities was found on the grassland litter. (31).

The profiles of the low-molecular-weight compounds derived from the litter substrates represented the molecular environment during decomposition and were characteristic of each site (29) and as well as each decomposition stage within a site (32). Over time, the litter types separated mostly according to the site as opposed to the litter types (redundancy analysis [RDA]), which demonstrates how decomposition transforms litter to a humus type specific for a given site (33).

The bacterial communities for the sedge (S) and beech (B) litters considerably diverged between sites but only in the later stages of decomposition when they were composed mostly of bacteria representing the indigenous soil communities of the sites. In contrast, the associated bacterial communities on the M litter at the two sites converged over time as the nutrient-rich M litter was sequentially exhausted (4, 10). A uniform community, therefore, reflected the final stage of decomposition for the M litter, while it possibly represented only the initial depolymerization of lignin for the S and B litters (34). Furthermore, the higher-quality M litter showed the largest shifts in bacterial OTUs over time, as well as the higher decomposition rates and highest activity of hydrolytic enzymes compared with the B and S litters across both sites (35). Thus, local-scale decomposition was driven by the local community according to the litter type (33) without any effect of endogenous or exogenous litter origin (i.e., M and S litters originating at the grassland site and beech at the forest site), contrary to the home-field advantage theory (31).

Quantitative analysis based on 16S and 18S rRNA gene abundances suggested discriminating driving factors for various microbial groups during litter decomposition. Specifically, *Firmicutes* and *Alphaproteobacteria* preferentially colonized the M litter, but colonized *Firmicutes* at both sites and *Alphaproteobacteria* (only for January and March 2011) at the grassland site. Fungi were affected more and *Actinobacteria* less by litter quality, similarly as in boreal peat soil (36). Therefore, *Actinobacteria* are substrate generalists influenced by environmental conditions, and consequently, they can specifically reflect site soil characteristics (37). In contrast, the site generalism of fungi supports this group's resiliency to environmental factors (17). However, that contrasts with the generally acknowledged assumption of fungi sensitivity to soil pH (38). Thus, in this study, the relationship between fungi and bacteria was probably neutral because fungi were pH dependent only if they grew in competition with bacteria (39).

The dynamics of *Actinobacteria* and fungi, certain enzymatic activities (arylsulfatase, chitinase, laccase, and manganese peroxidase), and mass loss were more similar for one

litter type between sites than the three litter types were similar within each site. Most of those enzymes target complex organic substrates and were found to be (except manganese peroxidase) produced by both *Actinobacteria* and fungi (20, 40–42). The ability of different soil communities to carry out decomposition of the same substrate over time regardless of the site implies functional redundancy (43). Such dynamics were, however, not the case for *Alphaproteobacteria* and *Firmicutes*, which are, therefore, likely functionally distinct (7). The results indirectly suggest that the dynamics of *Actinobacteria* and fungi were driven by the same factor, i.e., litter type (19).

The decomposition process is often considered to be separated into three stages characterized by typical microbial guilds and their functions (11). In the present study, the initial stage (from January to March 2011) was characterized by high extracellular enzyme activities, especially at the forest site. The bacterial communities on the M litter type were positively affected by available N and P, suggesting that bacteria require a lower C:N but higher N:P ratio (44), especially after intense enzyme production which likely leads to substantial losses of C and N (9). In this stage, a specific form of low-molecular-weight compounds was recorded on the M litter (45). In addition, the nutrient-rich M litter had the most dynamic bacterial communities, with high incidences of fungi and *Alphaproteobacteria* (according to quantitative PCR [qPCR]). Since fungi and *Alphaproteobacteria* are known for their ability to depolymerize recalcitrant compounds (46), their initial dominance may correspond with carbon depolymerization of the M litter. Meanwhile, *Bacteroidetes* (according to amplicon sequencing) occurred on all litter types at both sites. *Bacteroidetes* are also known for their decomposing activity, and their presence in all litter types suggested promotion by litter addition (47) regardless of its quality and origin.

The middle stage of decomposition was prevalently characterized by cellulose decomposition (48), even though in the current study it was documented only at grassland site from the increased activity of cellulolytic enzymes in May for the M litter and in July for the S litter. In this stage, *Verrucomicrobia* and *Proteobacteria* make up the highest proportion of taxa on all litter types, which suggests that they are substrate generalists and polysaccharide decomposers (49). Fungi decreased and were replaced by bacteria at both sites possibly because bacteria are scavengers benefitting from the initial cellulose degradation by fungi (48).

Finally, bacterial diversity increased especially on the M litter type over time, which suggested an increased number of ecological niches when the soil and litter communities combine (9, 50). The gradual incorporation of litter to the soil is supported by the relative increase of *Acidobacteria* and *Actinobacteria*, also increased later in the decomposition process, which corresponds to their relative abundance in soil, especially *Actinobacteria* (51). The increase of *Actinobacteria*, as well as *Firmicutes*, on the M and S litter types (according to both qPCR and relative proportion in sequence libraries) implies that late-stage decomposition is associated with Gram-positive K-strategist microbial groups (10, 52).

Actinobacteria were substrate generalists that adapt to litter of different decomposability (19, 53). *Actinobacteria* quantity, together with overall bacterial diversity, was positively correlated with the structure of the late-stage bacterial communities, suggesting they have a specific function in that decomposition stage. This conclusion was also supported by the late proportional increase of the *Actinobacteria* families *Mycobacteriaceae* and *Streptomyetaceae*, of which some genera are soil saprotrophs (54) or known for lignin breakdown (46). These families are also recognized for antimicrobial activity, which may be utilized for competitor suppression or as a synergic communication to overcome the small energetic return from recalcitrant compounds (15, 23). The relative abundance of *Frankiales* also increased in the late stages but only at the forest site, suggesting the advantage of the N fixation metabolism typical for this group in decomposing recalcitrant substrates in infertile soil (53). Interestingly, we also found an increase in the proportion of the Trebon clade later in decomposition at the forest site but only on the S litter, supporting previous findings of that clade's adaptation to low pH forest soil with recalcitrant humus (55). Finally, *Actinobacteria* might have a strong impact on litter-derived humus formation through modulation of the molecular environment and interactions within the decomposer community.

Niche partitioning is related to changes in environmental conditions (35) and might result from previous competition between species regulating their coexistence (56). This may be demonstrated by the different abundance maxima of the studied microbial groups, especially fungi and *Actinobacteria*, on the rich M litter (57). On the poorer B and S litters, the abundance curves of *Actinobacteria*, fungi, and *Alphaproteobacteria* were congruent, which may represent a synergistic interaction (58, 59). In spite of fungi being the initial decomposers, they dominated over *Actinobacteria* in the later stages at the forest site on the S litter, habitats favorable for fungi and unfavorable for *Actinobacteria*. This is in accordance with the higher fitness of specialists (fungi) at their optimal environmental conditions relative to generalists (*Actinobacteria*) at the same site (60). The S litter at the grassland site represents a niche overlap for *Actinobacteria* and fungi since it is the most favorable habitat of both. Interestingly, in this situation, *Actinobacteria* dominated over fungi, which might suggest a greater contribution of *Actinobacteria* to ecosystem functioning (61).

Site conditions had the biggest impact on the progress of decomposition, measured as litter mass loss, litter chemistry, and enzyme activities. The site conditions also determined bacterial diversity, although the litter seemed to create specific niches according to the type and stage of decomposition. Finally, the site conditions were most important for the total abundance of *Actinobacteria*. In contrast, litter type determined the abundance of fungi, while the abundance of *Alphaproteobacteria* was related to both factors. Also, litter type influenced the succession of fungi and *Actinobacteria* regardless of the site. Consequently, the observed patterns revealed that the decomposition process is independent of litter type and, therefore, may be relatively stable with respect to shifts of vegetation due to disturbance. However, the niches created by various litter types and decomposition stages may have an impact on the dynamics of the main microbial decomposers and that may lead to changes in organic matter quality and sequestration.

MATERIALS AND METHODS

Litterbag experiment. A grassland site situated on the western slope of Oblik Hill in the Bohemian Massif Central, Czech Republic, and a forest site in Dorotheer Wald, Neustift near Vienna, Austria, were chosen for the litterbag experiment. Soil humidity and temperature differed between the sites (see Fig. S8 in the supplemental material). The forest site was dominated by beech (*Fagus sylvatica*), with acidic soil (pH 5.3) and soil organic matter content of 11.7%. The grassland site was dominated by diverse forbs (29), with alkaline soil (pH 7.9) and soil organic matter content of 11.0%.

For the litter transplant experiment, we used beech leaves (B), collected in November 2010 at the forest site from beech trees before litter fall, and aboveground parts of a sedge (S; *Carex humilis*) and milkvetch (M; *Astragalus exscapus*) both cut from living plants in November 2010 at the grassland site. We chose these litter types according to their quality and decomposability as described by their C:N:P ratio. B had the highest, M the lowest, and S an intermediate C:N:P ratio (Fig. S2). Polyester litterbags of size 10 × 10 cm with a mesh size of 1 mm were filled with 7 g of dried litter, with 70 litterbags prepared for each litter type. The remaining litter was stored at −70°C for future analysis and was considered the initial sampling time (initial). Overall, 105 litterbags (35 litterbags from each litter type) were placed under the top 5 cm of the soil organic layer at each site in November 2010 (Fig. 6).

A total of 50 ml of surrounding soil and litterbag samples were collected in 2-month intervals; overall, 7 samplings were performed (January, March, May, July, September, and November 2011; and January 2012). Each litterbag was weighed to assess the weight loss during the decomposition process. The M litter was completely decomposed by January 2012 and no material remained in the litterbags. Therefore, analysis of the M litter type for this sampling time is missing (Fig. 6).

The soil and litter samples were cooled during transportation and homogenized and divided into subsamples upon arrival at the laboratory. Litter water content was assessed by the gravimetric method with oven drying (62). For microbial analysis, the soil and litter samples were placed in 2-ml sterile plastic tubes and stored at −70°C until further analysis. For high-performance liquid chromatography (HPLC) analysis, 100-ml glass bottles were filled with the soil and litter samples, overlaid with a methanol-water-acetic acid mixture (80:19:1 [vol/vol/vol]), and stored at −20°C. The remaining soil and litter samples were placed in plastic bags for an analysis of pH; organic carbon content (soil samples); and C, N, and P contents (litter samples) (Fig. 6).

Soil analysis. Soil moisture tension and temperature were recorded at both sites by a Microlog SP1 datalogger (EMS, Brno, Czech Republic) equipped with a gypsum block sensor. From the recorded values, mean temperature and humidity were calculated for every month from November 2010 to March 2012. Soil pH was measured by a Multi 350 WTW glass electrode in a soil water extract (5 g of soil was extracted with 20 ml of distilled water and set for 12 hours at room temperature). Soil organic matter content was estimated by combustion in an oven at 550°C to a constant weight.




FIG 6 Diagram showing the design of the litterbag experiment on decomposition of beech (B), milkvetch (M), and sedge (S) at the grassland (G) and forest (F) sites over 1 year (January to November 2011, i.e., 2 to 12 months after litterbag burial).

Carbon, nitrogen, and phosphorus analysis in the litter. Total phosphorus (P) content was determined colorimetrically in 20 mg of dried plant litter sequentially decomposed with HNO_3 and HClO_4 using the ammonium molybdate-ascorbic acid method on a QuikChem 8500 flow injection analysis system (Lachat Instruments, Hach Company, Loveland, CO, USA) at the Botanical Institute in Třeboň, Czech Republic (63). Total carbon (C) and nitrogen (N) contents were determined in 2 g of dried litter by high-temperature combustion, followed by gas separation in a Vario Max CNS element analyzer (Elementar Analysensysteme, Hanau, Germany) by Aquatest a.s. (Prague, Czech Republic).

HPLC analysis of soluble low-molecular-weight soil metabolites. Low-molecular-weight compounds of the soil and litter samples were analyzed by high performance liquid chromatography (HPLC) (64). The compounds were extracted in methanol-water-acetic acid (80:19:1 [vol/vol/vol]) and then filtered and vacuum concentrated. The concentrated solution was fractionated using Amberlite XAD

TABLE 1 Primers used for qPCR and for MiSeq Illumina sequencing

Procedure	Primer	Sequence, 5'–3'	Concn (μM)	Target	Annealing temp (°C)	Reference
qPCR	EUB518 ^a	ATTACCGCGGCTGCTGG	0.2	Bacteria, 16S rRNA	56	76
	Act235	CGCGGCTATCAGCTTGTTG	0.2	Actinobacteria, 16S rRNA	56	77
	S-C-aProt-0528-a-S-19	CGGTAATACGRAGGGRGYT	0.4	Alphaproteobacteria, 16S rRNA	61	67
	S-C-aProt-0689-a-A-21	CBAATATCTACGAATTTCACCT	0.4	Alphaproteobacteria, 16S rRNA	61	67
	S-P-Firm-0352-a-S-18	CAGCAGTAGGGAATCTTC	0.2	Firmicutes, 16S rRNA	57	67
	S-P-Firm-0525-a-A-18	ACCTACGTATTACCGCGG	0.2	Firmicutes, 16S rRNA	57	67
	FF390	CGATAACGAACGAGACCT	0.6	Fungi, 18S rRNA	50	78
	FR1	AICCATTCATCGGTAIT	0.6	Fungi, 18S rRNA	50	78
Sequencing	CS1-515F	ACACTGACGACATGGTTCTA- CAGTGCCAGCMGCGCGGTAA		Bacteria, 16S rRNA		68
	CS2-806R	TACGGTAGCAGAGACTTGGT CTGGACTACHVGGGTWCTAAT		Bacteria, 16S rRNA		68

^aEub518 was used as a reverse primer for Act235.

1180 (5 g aqueous suspension; 2-cm-diameter glass column, equilibrated with water). A total of 50 ml of water (MilliQ quality) was used as the effluent for the first (hydrophilic) fraction and 50 ml of absolute ethanol for the second (lipophilic) fraction. The ethanolic fraction was evaporated and diluted in methanol at a concentration of 5 mg ml⁻¹. The analysis was performed using the Acquity ultraperformance liquid chromatography (UPLC) system with a 2996 photodiode array (PDA) detector (Waters, Milford, MA, USA) with a Phenomenex Luna column (5 mm, C18[2], 100 Å, 100 × 4.6 mm) equipped with a SecurityGuard cartridge (C18, 4 × 3.0-mm internal diameter [ID]; Phenomenex, Torrance, CA, USA). The separation was run in an air-conditioned room at 20°C, with a flow rate of 1.0 ml min⁻¹. Solvent A was ultrapure water, while solvent B was acetonitrile. The gradient started with 10% of solvent B for 1 min and then increased linearly to 100% of B within 8 min. The final concentration was held for a further 3.5 min. A total of 20 μl of each sample was injected. The spectra were recorded from 194 to 800 nm in 1-nm intervals. Data points were recorded twice per second and downloaded using MassLynx software (Waters) in text format for further processing.

Enzyme extraction and assays. The activities of the extracellular hydrolytic enzymes α-glucosidase (EC 3.2.1.3), 1,4-β-xylosidase (EC 3.2.1.37), 1,4-β-glucosidase (EC 3.2.1.21), acid phosphatase (EC 3.1.3.1), chitinase (EC 3.2.1.52), cellobiohydrolase (EC 3.2.1.91), glucuronidase (EC 3.2.1.31), and arylsulfatase (EC 3.1.6.1) were directly measured using methylumbelliferone-derived fluorogenic substrates in aliquots of litter samples homogenized in 50 mM sodium acetate buffer at pH 5.0 (65).

For the extracellular oxidative enzymes laccase (EC 1.10.3.2) and manganese peroxidase (EC 1.11.1.13), aliquots were desalted to separate inhibitory molecules. The activity of laccase was measured by monitoring the oxidation of 2,20-azinobis-3-ethylbenzothiazoline-6-sulfonic acid in a citrate-phosphate buffer (pH 5.0) (66). Manganese peroxidase activity was assayed in a succinate-lactate buffer (100 mM, pH 4.5) using 3-methyl-2-benzothiazolinone hydrazone and 3,3-dimethylaminobenzoic acid, which were oxidatively coupled by the enzyme. The results were compared to the activities of samples without manganese (for manganese peroxidase, the addition of manganese sulfate was substituted by an equimolar amount of EDTA). One unit of enzyme activity is defined as the amount of enzyme forming 1 μM reaction product per minute. Enzyme activities were measured in three analytical replicates and expressed per gram of plant litter dry mass (48).

DNA extraction and microbial community analysis. DNA was extracted from litter and soil samples according to Sagova-Mareckova et al. (67). The method is based on bead beating and phenol-chloroform extraction. The samples were purified by incubation with acetyltrimethylammonium bromide, followed by chloroform extraction and incubation with CaCl₂, and finally cleaned with the GeneClean turbo kit (MP Biomedicals, Santa Ana, CA, USA). DNA extracts were stored at -70°C until further analysis.

Illumina MiSeq and data processing. Bacterial 16S rRNA gene fragments, including the variable region V4, were amplified by PCR using universal primers with overhang adapters (Table 1) (68). The construction of amplicon libraries and sequencing using the MiSeq sequencer (Illumina, San Diego, CA, USA) were done at the DNA Services Facility, Research Resources Center, University of Illinois (Chicago, IL, USA). The resulting paired sequence reads were merged, filtered, and aligned using a reference alignment from the Silva database (69) and chimera checked using the integrated Vsearch tool (70) according to the MiSeq standard operation procedure (MiSeq standard operating procedure [SOP], February 2018 [68] in Mothur v. 1.39.5 [70]). A taxonomical assignment of sequence libraries was performed in Mothur using the nonredundant subset of the SILVA small subunit rRNA database, release 132 (71), adapted for use in Mothur (https://mothur.org/w/images/3/32/Silva.nr_v132.tgz), as the reference database. Sequences of plastids and mitochondria and those not classified in the domain *Bacteria* were discarded.

The sequence library was clustered into OTUs (97%) using Usearch v. 10.0.240 (72), and the OTU table was further processed using tools implemented in Mothur. Distance matrices describing the differences in community composition between individual samples were calculated using Bray-Curtis dissimilarity (73). Analysis of molecular variance (AMOVA) (74) was based on a matrix of the Bray-Curtis distances. Metastats (75) analyses were used to detect differentially represented OTUs. Rarefaction analysis was performed with the rarefaction.single function in Mothur, and areas under the rarefaction curves were

calculated using the trapezoidal rule in R version 3.4.4 (76). Figures were created using Inkscape (v. 0.92; <http://www.inkscape.org>).

Quantitative real-time PCR. Quantification of the 16S rRNA gene in *Actinobacteria*, *Alphaproteobacteria*, and *Firmicutes* and the 18S rRNA gene of fungi in the DNA extracted from litter samples was performed with the primers listed in Table 1 (77–79).

Quantitative PCR was performed using the StepOnePlus real-time PCR system (Applied Biosystems, Foster City, CA, USA) using 96-well plates with GoTaq qPCR master mix (Promega, Madison, WI, USA) containing SYBR green as a double-stranded DNA (dsDNA) binding dye. The reaction mixture contained in total a volume of 15 μ l with 1 \times GoTaq qPCR master mix, primers (for concentrations see Table 1), and 2-ng DNA sample. For all microbial groups, the PCR cycling protocol consisted of initial denaturation at 95°C for 10 min, followed by 45 cycles of denaturation (95°C for 15 s), annealing (for respective temperature see Table 1), and elongation (72°C for 30 s). Standards containing a known number of copies of the target genes 16S rRNA from *Streptomyces europaeiscabiei* DSM 41802 (for *Actinobacteria*), *Bartonella*-like clone (for *Alphaproteobacteria*), and *Paenibacillus* clone (for *Firmicutes*) and 18S rRNA from a *Penicillium* sp. clone (for fungi) were serially diluted to calibrate each qPCR melting curve and were recorded to ensure qPCR specificity. Baseline and threshold calculations were performed with StepOne v. 2.2.2. Inhibition was tested by serial DNA dilutions from each site. All qPCR measurements were done in duplicate.

Statistical analysis. All statistical calculations were done in the R computing environment (76). The effect of sites and litter types on bacterial community composition and the litter molecular parameters during the decomposition process were measured as the relationship between the distance matrices of the HPLC profiles of low-molecular-weight compounds (HPLC) and bacterial communities using a generalized distance covariance (80). The HPLC profiles were baseline corrected using the PTW package (81), and distances between profiles were calculated using Manhattan metrics. The distance matrices of the HPLC profiles and bacterial communities (exported from Mothur) were plotted by RDA projection and Sammon's multidimensional scaling using the MASS package (82). Correlations between linear variables, bacterial community, and HPLC profiles were analyzed by permutation test for capscale under a reduced model using the Bray-Curtis (bacterial communities) and Manhattan (HPLC) dissimilarities (with 999 permutations) in the VEGAN package in R (83).

To analyze the overall quantitative differences of linear variabilities (dry mass loss; abundances of microbial groups; enzyme activities; and C, N, and P contents in litter) between sites and litter types (within sites), a linear model with the additive effect of time, site, and litter type was calculated (84).

We used two tests based on the distance between the profiles as measured by correlation coefficients (see p. 319 of reference 85) to analyze the qualitative similarities between the linear variables. Permutational multivariate analysis of variance (PERMANOVA; significance values denoted here as P_{perm}) (86) tests whether the within-group distances (same litter type) are on average shorter than the between-group distances (between different litter types, aimed at detection of different mean profiles). The dispersion test (significance values denoted here as P_{Disp}), introduced in Gijbels and Omelka (87), checks whether the average within-group distances differ among groups (aiming at detecting different dispersions of the samples). For manganese peroxidase, a Manhattan distance instead of the distance based on the correlation coefficient was used due to zero values in some profiles.

All linear variables (dry mass loss; microbial gene quantities; C, N, and P content; and enzyme activities, except manganese peroxidase) were logarithmically transformed (with negative values of enzymes replaced by 1 before transformation) to make their distribution more similar to a normal distribution.

Data availability. The Illumina MiSeq 16S rRNA gene amplicon sequences have been deposited in the NCBI Sequence Read Archive under accession no. [SRP158384](https://www.ncbi.nlm.nih.gov/sra/SRP158384) and are available under BioProject accession number [PRJNA485178](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA485178).

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/AEM.01760-19>.

SUPPLEMENTAL FILE 1, PDF file, 0.8 MB.

ACKNOWLEDGMENTS

The work was funded by the Czech Science Foundation (grant number 15-01312S); Ministry of Education, Youth and Sports of the Czech Republic, European Regional Development Fund-Project number CZ.02.1.01/0.0/0.0/16_019/0000845 and grant number LTC17075; and Ministry of Agriculture of the Czech Republic, institutional support number MZE-RO0418.

We declare no conflict of interest.

REFERENCES

1. Purahong W, Krüger D, Buscot F, Wubet T. 2016. Correlations between the composition of modular fungal communities and litter decomposition-associated ecosystem functions. *Fungal Ecol* 22:106–114. <https://doi.org/10.1016/j.funeco.2016.04.009>.
2. Lladó S, López-Mondéjar R, Baldrian P. 2017. Forest soil bacteria: diversity, involvement in ecosystem processes, and response to global change. *Microbiol Mol Biol Rev* 81:e00063-16. <https://doi.org/10.1128/MMBR.00063-16>.

3. Fukami T. 2015. Historical contingency in community assembly: integrating niches, species pools, and priority effects. *Annu Rev Ecol Syst* 46:1–23. <https://doi.org/10.1146/annurev-ecolsys-110411-160340>.
4. McGuire KL, Treseder KK. 2010. Microbial communities and their relevance for ecosystem models: decomposition as a case study. *Soil Biol Biochem* 42:529–535. <https://doi.org/10.1016/j.soilbio.2009.11.016>.
5. Bani A, Pioli S, Ventura M, Panzacchi P, Borroso L, Tognetti R, Tonon G, Brusetti L. 2018. The role of microbial community in the decomposition of leaf litter and deadwood. *Appl Soil Ecol* 126:75–84. <https://doi.org/10.1016/j.apsoil.2018.02.017>.
6. Fraser LH, Hockin AD. 2013. Litter decomposition rates of two grass species along a semi-arid grassland-forest ecocline. *J Arid Environ* 88: 125–129. <https://doi.org/10.1016/j.jaridenv.2012.07.009>.
7. Glassman SI, Weihe C, Li J, Albright MBN, Looby CI, Martiny AC, Treseder KK, Allison SD, Martiny JBH. 2018. Decomposition responses to climate depend on microbial community composition. *Proc Natl Acad Sci U S A* 115:11994–11999. <https://doi.org/10.1073/pnas.1811269115>.
8. García-Palacios P, Shaw EA, Wall DH, Hättenschwiler S. 2016. Temporal dynamics of biotic and abiotic drivers of litter decomposition. *Ecol Lett* 19:554–563. <https://doi.org/10.1111/ele.12590>.
9. Purahong W, Wubet T, Lentendu G, Schlöter M, Pecyna MJ, Kapturska D, Hofrichter M, Krüger D, Buscot F. 2016. Life in leaf litter: novel insights into community dynamics of bacteria and fungi during litter decomposition. *Mol Ecol* 25:4059–4074. <https://doi.org/10.1111/mec.13739>.
10. Bray SR, Kitajima K, Mack MC. 2012. Temporal dynamics of microbial communities on decomposing leaf litter of 10 plant species in relation to decomposition rate. *Soil Biol Biochem* 49:30–37. <https://doi.org/10.1016/j.soilbio.2012.02.009>.
11. Tlaskal V, Voříšková J, Baldrian P. 2016. Bacterial succession on decomposing leaf litter exhibits a specific occurrence pattern of cellulolytic taxa and potential decomposers of fungal mycelia. *FEMS Microbiol Ecol* 92:fw177. <https://doi.org/10.1093/femsec/fw177>.
12. Wertz S, Degrange V, Prosser JI, Poly F, Commaux C, Freitag T, Guillaumaud N, Roux XL. 2006. Maintenance of soil functioning following erosion of microbial diversity. *Environ Microbiol* 8:2162–2169. <https://doi.org/10.1111/j.1462-2920.2006.01098.x>.
13. Purahong W, Schlöter M, Pecyna MJ, Kapturska D, Däumlich V, Mital S, Buscot F, Hofrichter M, Gutknecht JLM, Krüger D. 2014. Uncoupling of microbial community structure and function in decomposing litter across beech forest ecosystems in Central Europe. *Sci Rep* 4:7014. <https://doi.org/10.1038/srep07014>.
14. Moorhead DL, Sinsabaugh RL. 2006. A theoretical model of litter decay and microbial interaction. *Ecol Monogr* 76:151–174. [https://doi.org/10.1890/0012-9615\(2006\)076\[0151:ATMOLD\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2006)076[0151:ATMOLD]2.0.CO;2).
15. De Boer W, Folman LB, Summerbell RC, Boddy L. 2005. Living in a fungal world: impact of fungi on soil bacterial niche development. *FEMS Microbiol Rev* 29:795–811. <https://doi.org/10.1016/j.femsre.2004.11.005>.
16. Baldrian P, Zrůstová P, Tlaskal V, Davidová A, Merhautová V, Vrška T. 2016. Fungi associated with decomposing deadwood in a natural beech-dominated forest. *Fungal Ecol* 23:109–122. <https://doi.org/10.1016/j.funeco.2016.07.001>.
17. A'Bear AD, Jones TH, Kandler E, Boddy L. 2014. Interactive effects of temperature and soil moisture on fungal-mediated wood decomposition and extracellular enzyme activity. *Soil Biol Biochem* 70:151–158. <https://doi.org/10.1016/j.soilbio.2013.12.017>.
18. Fierer N, Bradford MA, Jackson RB. 2007. Toward an ecological classification of soil bacteria. *Ecology* 88:1354–1364. <https://doi.org/10.1890/05-1839>.
19. Větrovský T, Steffen KT, Baldrian P. 2014. Potential of cometabolic transformation of polysaccharides and lignin in lignocellulose by soil Actinobacteria. *PLoS One* 9:e89108. <https://doi.org/10.1371/journal.pone.0089108>.
20. Fernandes ART, da Silveira WB, Passos MLF, Zucchi TD. 2014. Laccases from Actinobacteria—what we have and what to expect. *Adv Microbiol* 04:285–296. <https://doi.org/10.4236/aim.2014.46035>.
21. Sagova-Mareckova M, Zadorova T, Penizek V, Omelka M, Tejnecky V, Pruchova P, Chuman T, Drabek O, Buresova A, Vanek A, Kopecky J. 2016. The structure of bacterial communities along two vertical profiles of a deep colluvial soil. *Soil Biol Biochem* 101:65–73. <https://doi.org/10.1016/j.soilbio.2016.06.026>.
22. Abdelmohsen UR, Krkovic T, Balasubramanian S, Kamel MS, Quinn RJ, Hentschel U. 2015. Elicitation of secondary metabolism in Actinomycetes. *Biotechnol Adv* 33:798–811. <https://doi.org/10.1016/j.biotechadv.2015.06.003>.
23. Sharma M, Dangi P, Choudhary M. 2014. Actinomycetes: source, identification, and their applications. *Int J Curr Microbiol Appl Sci* 3:801–832.
24. Fukami T, Dickie IA, Paula Wilkie J, Paulus BC, Park D, Roberts A, Buchanan PK, Allen RB. 2010. Assembly history dictates ecosystem functioning: evidence from wood decomposer communities. *Ecol Lett* 13:675–684. <https://doi.org/10.1111/j.1461-0248.2010.01465.x>.
25. Schmidt M, Veldkamp E, Corre MD. 2015. Tree species diversity effects on productivity, soil nutrient availability and nutrient response efficiency in a temperate deciduous forest. *For Ecol Manage* 338:114–123. <https://doi.org/10.1016/j.foreco.2014.11.021>.
26. Fontaine S, Mariotti A, Abbadie L. 2003. The priming effect of organic matter: a question of microbial competition? *Soil Biol Biochem* 35: 837–843. [https://doi.org/10.1016/S0038-0717\(03\)00123-8](https://doi.org/10.1016/S0038-0717(03)00123-8).
27. Mokany K, Raison RJ, Prokushkin AS. 2006. Critical analysis of root:shoot ratios in terrestrial biomes. *Glob Chang Biol* 12:84–96. <https://doi.org/10.1111/j.1365-2486.2005.001043.x>.
28. Lange M, Eisenhauer N, Sierra CA, Bessler H, Engels C, Griffiths RI, Mellado-Vázquez PG, Malik AA, Roy J, Scheu S, Steinbeiss S, Thomson BC, Trumbore SE, Gleixner G. 2015. Plant diversity increases soil microbial activity and soil carbon storage. *Nat Commun* 6:6707. <https://doi.org/10.1038/ncomms7707>.
29. Sagova-Mareckova M, Omelka M, Cermak L, Kamenik Z, Olsovska J, Hackl E, Kopecky J, Hadacek F. 2011. Microbial communities show parallels at sites with distinct litter and soil characteristics. *Appl Environ Microbiol* 77:7560–7567. <https://doi.org/10.1128/AEM.00527-11>.
30. McDaniel MD, Grandy AS, Tiemann LK, Weintraub MN. 2014. Crop rotation complexity regulates the decomposition of high and low quality residues. *Soil Biol Biochem* 78:243–254. <https://doi.org/10.1016/j.soilbio.2014.07.027>.
31. John MGS, Orwin KH, Dickie IA. 2011. No “home” versus “away” effects of decomposition found in a grassland-forest reciprocal litter transplant study. *Soil Biol Biochem* 43:1482–1489. <https://doi.org/10.1016/j.soilbio.2011.03.022>.
32. Ristok C, Luettel KN, Franke K, Scherer-Lorenzen M, Niklaus PA, Wessjohann LA, Bruehlheide H. 2017. Leaf litter diversity positively affects the decomposition of plant polyphenols. *Plant Soil* 419:305–317. <https://doi.org/10.1007/s11104-017-3340-8>.
33. Solly EF, Schöning I, Boch S, Kandler E, Marhan S, Michalzik B, Müller J, Zscheischler J, Trumbore SE, Schirmpf M. 2014. Factors controlling decomposition rates of fine root litter in temperate forests and grasslands. *Plant Soil* 382:203–218. <https://doi.org/10.1007/s11104-014-2151-4>.
34. Dungait JAJ, Hopkins DW, Gregory AS, Whitmore AP. 2012. Soil organic matter turnover is governed by accessibility not recalcitrance. *Glob Chang Biol* 18:1781–1796. <https://doi.org/10.1111/j.1365-2486.2012.02665.x>.
35. Kielak AM, Scheublin TR, Mendes LW, Van Veen JA, Kuramae EE. 2016. Bacterial community succession in pine-wood decomposition. *Front Microbiol* 7:231. <https://doi.org/10.3389/fmicb.2016.00231>.
36. Peltoniemi K, Straková P, Fritze H, Iräizoz PA, Pennanen T, Laiho R. 2012. How water-level drawdown modifies litter-decomposing fungal and actinobacterial communities in boreal peatlands. *Soil Biol Biochem* 51: 20–34. <https://doi.org/10.1016/j.soilbio.2012.04.013>.
37. Sagova-Mareckova M, Cermak L, Omelka M, Kyselkova M, Kopecky J. 2015. Bacterial diversity and abundance of a creek valley sites reflected soil pH and season. *Open Life Sci* 10:61–70. <https://doi.org/10.1515/biol-2015-0007>.
38. Höglberg MN, Höglberg P, Myrold DD. 2007. Is microbial community composition in boreal forest soils determined by pH, C-to-N ratio, the trees, or all three? *Oecologia* 150:590–601. <https://doi.org/10.1007/s00442-006-0562-5>.
39. Rousk J, Brookes PC, Bååth E. 2009. Contrasting soil pH effects on fungal and bacterial growth suggest functional redundancy in carbon mineralization. *Appl Environ Microbiol* 75:1589–1596. <https://doi.org/10.1128/AEM.02775-08>.
40. Malý S, Fiala P, Reininger D, Obdržálková E. 2014. The relationships among microbial parameters and the rate of organic matter mineralization in forest soils, as influenced by forest type. *Pedobiologia (Jena)* 57:235–244. <https://doi.org/10.1016/j.pedobi.2014.09.003>.
41. Chen H, Liu J, Li D, Xiao K, Wang K. 2019. Controls on soil arylsulfatase activity at a regional scale. *Eur J Soil Biol* 90:9–14. <https://doi.org/10.1016/j.ejsobi.2018.11.001>.
42. Vrsanska M, Voberkova S, Langer V, Palovcickova D, Moulick A, Adam V, Kopel P, Vrsanska M, Voberkova S, Langer V, Palovcickova D, Moulick A, Adam V, Kopel P. 2016. Induction of laccase, lignin peroxidase and

- manganese peroxidase activities in white-rot fungi using copper complexes. *Molecules* 21:1553. <https://doi.org/10.3390/molecules21111553>.
43. Banerjee S, Kirkby CA, Schmutter D, Bissett A, Kirkegaard JA, Richardson AE. 2016. Network analysis reveals functional redundancy and keystone taxa amongst bacterial and fungal communities during organic matter decomposition in an arable soil. *Soil Biol Biochem* 97:188–198. <https://doi.org/10.1016/j.soilbio.2016.03.017>.
 44. Güsewell S, Gessner MO. 2009. N:P ratios influence litter decomposition and colonization by fungi and bacteria in microcosms. *Funct Ecol* 23: 211–219. <https://doi.org/10.1111/j.1365-2435.2008.01478.x>.
 45. van Hees PAW, Jones DL, Finlay R, Godbold DL, Lundström US. 2005. The carbon we do not see—the impact of low molecular weight compounds on carbon dynamics and respiration in forest soils: a review. *Soil Biol Biochem* 37:1–13. <https://doi.org/10.1016/j.soilbio.2004.06.010>.
 46. Bugg TDH, Ahmad M, Hardiman EM, Singh R. 2011. The emerging role for bacteria in lignin degradation and bio-product formation. *Curr Opin Biotechnol* 22:394–400. <https://doi.org/10.1016/j.copbio.2010.10.009>.
 47. Sauvadet M, Chauvat M, Cluzeau D, Maron PA, Villenave C, Bertrand I. 2016. The dynamics of soil micro-food web structure and functions vary according to litter quality. *Soil Biol Biochem* 95:262–274. <https://doi.org/10.1016/j.soilbio.2016.01.003>.
 48. Šnajdr J, Cajthaml T, Valášková V, Merhautová V, Petránková M, Spetz P, Leppänen K, Baldrian P. 2011. Transformation of *Quercus petraea* litter: successive changes in litter chemistry are reflected in differential enzyme activity and changes in the microbial community composition. *FEMS Microbiol Ecol* 75:291–303. <https://doi.org/10.1111/j.1574-6941.2010.00999.x>.
 49. Cardman Z, Arnosti C, Durbin A, Ziervogel K, Cox C, Steen AD, Teske A. 2014. Verrucomicrobia are candidates for polysaccharide-degrading bacterioplankton in an arctic fjord of Svalbard. *Appl Environ Microbiol* 80:3749–3756. <https://doi.org/10.1128/AEM.00899-14>.
 50. Jackson CR. 2003. Changes in community properties during microbial succession. *Oikos* 101:444–448. <https://doi.org/10.1034/j.1600-0706.2003.12254.x>.
 51. Štursová M, Žifčáková L, Leigh MB, Burgess R, Baldrian P. 2012. Cellulose utilization in forest litter and soil: identification of bacterial and fungal decomposers. *FEMS Microbiol Ecol* 80:735–746. <https://doi.org/10.1111/j.1574-6941.2012.01343.x>.
 52. Bastian F, Bouziri L, Nicolardot B, Ranjard L. 2009. Impact of wheat straw decomposition on successional patterns of soil microbial community structure. *Soil Biol Biochem* 41:262–275. <https://doi.org/10.1016/j.soilbio.2008.10.024>.
 53. Gao B, Gupta R. 2012. Phylogenetic framework and molecular signatures for the main clades of the phylum Actinobacteria. *Microbiol Mol Biol Rev* 76:66–112. <https://doi.org/10.1128/MMBR.05011-11>.
 54. Kopecký J, Kyselkova M, Omelka M, Cermak L, Novotna J, Grundmann G, Moëne-Loccoz Y, Sagova-Mareckova M. 2011. Environmental mycobacteria closely related to the pathogenic species evidenced in an acidic forest wetland. *Soil Biol Biochem* 43:697–700. <https://doi.org/10.1016/j.soilbio.2010.11.033>.
 55. Kopecký J, Kyselkova M, Omelka M, Cermak L, Novotna J, Grundmann GL, Moëne-Loccoz Y, Sagova-Mareckova M. 2011. Actinobacterial community dominated by a distinct clade in acidic soil of a waterlogged deciduous forest. *FEMS Microbiol Ecol* 78:386–394. <https://doi.org/10.1111/j.1574-6941.2011.01173.x>.
 56. Leibold M, McPeck M. 2006. Coexistence of the niche and neutral perspectives in community ecology. *Ecology* 87:1399–1410. [https://doi.org/10.1890/0012-9658\(2006\)87\[1399:COTNAN\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[1399:COTNAN]2.0.CO;2).
 57. Rousk J, Demoling LA, Bahr A, Bååth E. 2008. Examining the fungal and bacterial niche overlap using selective inhibitors in soil. *FEMS Microbiol Ecol* 63:350–358. <https://doi.org/10.1111/j.1574-6941.2008.00440.x>.
 58. Johnston SR, Boddy L, Weightman AJ. 2016. Bacteria in decomposing wood and their interactions with wood-decay fungi. *FEMS Microbiol Ecol* 92:fiw179. <https://doi.org/10.1093/femsec/fiw179>.
 59. Meidute S, Demoling F, Bååth E. 2008. Antagonistic and synergistic effects of fungal and bacterial growth in soil after adding different carbon and nitrogen sources. *Soil Biol Biochem* 40:2334–2343. <https://doi.org/10.1016/j.soilbio.2008.05.011>.
 60. Monard C, Gantner S, Bertilsson S, Hallin S, Stenlid J. 2016. Habitat generalists and specialists in microbial communities across a terrestrial-freshwater gradient. *Sci Rep* 6:37719. <https://doi.org/10.1038/srep37719>.
 61. Strickland MS, Rousk J. 2010. Considering fungal:bacterial dominance in soils—methods, controls, and ecosystem implications. *Soil Biol Biochem* 42:1385–1395. <https://doi.org/10.1016/j.soilbio.2010.05.007>.
 62. Carter MR, Gregorich EG. 2008. Soil sampling and methods of analysis, second ed. Taylor & Francis Group, LLC, USA.
 63. Olsen SR. 1954. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. US Department of Agriculture, Washington, DC.
 64. Kameník Z, Hadacek F, Marečková M, Ulanova D, Kopecký J, Chobot V, Plháčková K, Olšovská J. 2010. Ultra-high-performance liquid chromatography fingerprinting method for chemical screening of metabolites in cultivation broth. *J Chromatogr A* 1217:8016–8025. <https://doi.org/10.1016/j.chroma.2010.08.031>.
 65. Baldrian P. 2009. Microbial enzyme-catalyzed processes in soils and their analysis. *Plant Soil Environ* 55:370–378. <https://doi.org/10.17221/134/2009-PSE>.
 66. Bourbonnais R, Paice MG. 1990. Oxidation of non-phenolic substrates. *FEBS Lett* 267:99–102. [https://doi.org/10.1016/0014-5793\(90\)80298-w](https://doi.org/10.1016/0014-5793(90)80298-w).
 67. Sagova-Mareckova M, Cermak L, Novotna J, Plhacova K, Forstova J, Kopecky J. 2008. Innovative methods for soil DNA purification tested in soils with widely differing characteristics. *Appl Environ Microbiol* 74: 2902–2907. <https://doi.org/10.1128/AEM.02161-07>.
 68. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl Environ Microbiol* 79:5112–5120. <https://doi.org/10.1128/AEM.01043-13>.
 69. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013. The SILVA ribosomal RNA gene database project: improved data processing and Web-based tools. *Nucleic Acids Res* 41:D590–D596. <https://doi.org/10.1093/nar/gks1219>.
 70. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. 2016. VSEARCH: a versatile open source tool for metagenomics. *PeerJ* 4:e2584. <https://doi.org/10.7717/peerj.2584>.
 71. Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J, Ludwig W, Glöckner FO. 2014. The SILVA and “all-species Living Tree Project (LTP)” taxonomic frameworks. *Nucleic Acids Res* 42: D643–D648. <https://doi.org/10.1093/nar/gkt1209>.
 72. Edgar RC. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat Methods* 10:996–998. <https://doi.org/10.1038/nmeth.2604>.
 73. Bray JR, Curtis JT. 1957. An ordination of the upland forest communities of southern Wisconsin. *Ecol Monogr* 27:325–349. <https://doi.org/10.2307/1942268>.
 74. Martin AP. 2002. Phylogenetic approaches for describing and comparing the diversity of microbial phylogenetic approaches for describing and comparing the diversity of microbial communities. *Appl Environ Microbiol* 68:3673–3682. <https://doi.org/10.1128/aem.68.8.3673-3682.2002>.
 75. White JR, Nagarajan N, Pop M. 2009. Statistical methods for detecting differentially abundant features in clinical metagenomic samples. *PLoS Comput Biol* 5:e1000352. <https://doi.org/10.1371/journal.pcbi.1000352>.
 76. R Core Team. 2018. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
 77. Muyzer G, de Waal EC, Uitterlinden AG. 1993. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl Environ Microbiol* 59:695–700.
 78. Stach JEM, Maldonado LA, Ward AC, Goodfellow M, Bull AT. 2003. New primers for the class Actinobacteria: application to marine and terrestrial environments. *Environ Microbiol* 5:828–841. <https://doi.org/10.1046/j.1462-2920.2003.00483.x>.
 79. Prévost-Bouré CN, Christen R, Dequiedt S, Mougel C, Lelièvre M, Jolivet C, Shahbazkia HR, Guillou L, Arrouays D, Ranjard L. 2011. Validation and application of a PCR primer set to quantify fungal communities in the soil environment by real-time quantitative PCR. *PLoS One* 6:e24166. <https://doi.org/10.1371/journal.pone.0024166>.
 80. Omelka M, Hudecová Š. 2013. A comparison of the Mantel test with a generalised distance covariance test. *Environmetrics* 24:449–460. <https://doi.org/10.1002/env.2238>.
 81. Bloemberg TG, Gerretzen J, Lunshof A, Wehrens R, Buydens L. 2013. Warping methods for spectroscopic and chromatographic signal alignment: a tutorial. *Anal Chim Acta* 781:14–32. <https://doi.org/10.1016/j.aca.2013.03.048>.
 82. Venables WN, Ripley BD. 2002. Random and mixed effects, p 271–300. *In* Modern applied statistics with S. Springer, New York, NY.
 83. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E,

- Wagner H. 2019. vegan: community ecology package. R package version 2.5-6. <https://CRAN.R-project.org/package=vegan>.
84. Hothorn T, Everitt BS. 2014. A handbook of statistical analyses using R. Chapman and Hall/CRC, New York, NY. <https://doi.org/10.1201/b17081>.
85. Legendre P, Legendre L. 2012. Numerical ecology, 3rd English ed. Elsevier Science BV, Amsterdam, the Netherlands.
86. McArdle BH, Anderson MJ. 2001. Fitting multivariate models to community data: a comment on distance-based redundancy analysis. *Ecology* 82:290–297. <https://doi.org/10.2307/2680104>.
87. Gijbels I, Omelka M. 2013. Testing for homogeneity of multivariate dispersions using dissimilarity measures. *Biometrics* 69:137–145. <https://doi.org/10.1111/j.1541-0420.2012.01797.x>.

Supplementary Information for

Decomposition was driven by site conditions, while dynamics of *Actinobacteria* and fungi reflected a litter type

Buresova A., Kopecky J., Hrdinkova V., Kamenik Z., Omelka M., Sagova-Mareckova M.

Corresponding author: Marketa Sagova-Mareckova
Email: marketa.sagova@gmail.com

This PDF file includes:

- Supplementary text
- Figs. S1 to S10
- Tables S1 to S5
- References for supplementary text citations

Supplementary Text

Materials and Methods

Litterbag experiment

A grassland site (G) situated at the western slope of Oblik hill in Bohemian Massive Central, Czech Republic and a forest site (F) in Dorotheer Wald, Neustift near Wien, Austria were chosen for the litterbag experiment. Soil pH, vegetation, soil organic carbon content, soil humidity and temperature contrasted the sites (Fig. S1.). The F site was dominated by beech (*Fagus sylvatica*), with acidic soil (pH 5.3) and soil organic matter content 11.7%. The G site was dominated by diverse forbs (1), with alkaline soil (pH 7.9) and soil organic matter content 11.0%.

For the experiment, we used leaves of beech (B, *Fagus sylvatica*) from *Fagaceae* family (collected in November 2010 at the forest site), and aboveground parts of sedge (S, *Carex humilis*) from *Cyperaceae* family, and milkvetch (M, *Astragalus exscapus*) from *Fabaceae* family (both collected in November 2010 at the grassland site). We chose these litter types according to their quality and decomposability described by C:N:P ratio. B disposed of the highest, M the lowest and S the intermediate C:N:P ratio in its tissues (Fig. S2).

The soil and litterbag samples were collected in two-month intervals; overall 7 samplings were performed (January 2011, March 2011, May 2011, July 2011, September 2011, November 2011 and January 2012). Each litterbag was weighted to assess the weight loss during the decomposition process. M litter was completely decomposed in January 2012 and no material remained in the litterbags. Therefore, analysis for this sampling time for M litter type are missing.

The soil and litter samples were cooled during transportation, homogenized and divided into subsamples upon arrival to the laboratory. For microbial analysis, the soil and litter samples were placed in 2 ml sterile plastic tubes and stored at -70°C until a further analysis. For high performance liquid chromatography (HPLC) analysis 100 ml glass bottles were filled with the soil and litter samples, overlaid with a mixture of methanol-water-acetic acid (80:19:1, vol/vol/vol) and stored in -20°C. The remaining soil and litter samples were placed in plastic bags for analysis of pH, organic carbon content (soil samples) and C, N and P contents (litter samples).

Soil analysis

Soil moisture tension and temperature were recorded at both sites by a datalogger Microlog SP1 (EMS, Brno, Czech Republic) equipped with a gypsum block sensor. From recorded values, mean temperature and humidity for every month from November 2010 to March 2012 were calculated. The soil pH was measured by a Multi 350 WTW glass electrode in a soil water extract (5 g of soil was extracted with 20 ml of distilled water and set for 12 hours at a room temperature). The soil organic matter content was estimated by combustion in an oven at 550°C to the constant weight.

HPLC analysis of soluble low-molecular-weight soil metabolites

The extracts of soil and litter samples in a methanol-water-acetic acid were filtered and vacuum concentrated. The concentrated solution was fractionated using Amberlite XAD 1180 (5 g of aqueous suspension; 2 cm diameter of a glass column, equilibrated with water). 50 ml of water (MilliQ quality) was used as the effluent for the first (hydrophilic) fraction; 50 ml of absolute ethanol was used for the second (lipophilic) fraction. The ethanolic fraction was evaporated and diluted in a methanol at a concentration of 5 mg ml⁻¹. The analysis was performed using Acquity UPLC system with the 2996 PDA detector (Waters, Milford, MA, USA). The column was a Phenomenex Luna 5 mm C18(2) 100 Å, 100 × 4.6 mm, equipped with a SecurityGuard cartridge C18, 4 × 3.0 mm ID (Phenomenex, Torrance, CA, USA). The separation was run in the air-conditioned room at 20°C, and the flow rate was 1.0 ml min⁻¹. Solvent A was ultrapure water; solvent B was acetonitrile. The gradient started with 10% of solvent B for 1 min and then increased linearly to 100% of B within 8 min. The final concentration was held for further 3.5 min. 20 µl of each sample were injected. The spectra were recorded from 194 to 800 nm in 1 nm intervals. Data points recorded twice per second were exported from a MassLynx software (Waters) in a text format for further processing.

Enzyme extraction and assays

The activities of extracellular hydrolytic enzymes α – glucosidase (3.2.1.3), 1,4- β - xylosidase (EC 3.2.1.37), 1,4- β -glucosidase (EC 3.2.1.21), acid phosphatase (EC 3.1.3.1), chitinase (EC 3.2.1.52), cellobiohydrolase (EC 3.2.1.91), glucuronidase

(3.2.1.31) and arylsulphatase (3.1.6.1) were measured using methylumbellyferone-derived fluorogenic substrates directly in aliquots of litter samples homogenized in 50 mM sodium acetate buffer at pH 5.0 (2).

For extracellular oxidative enzymes laccase (EC 1.10.3.2) and manganese peroxidase (EC 1.11.1.13), aliquots were desalted to separate inhibitory molecules. The activity of laccase was measured by monitoring the oxidation of 2,20-azinobis-3-ethylbenzothiazoline-6-sulfonic acid in a citrate-phosphate buffer (pH 5.0) (3). The manganese peroxidase activity was assayed in a succinate-lactate buffer (100 mM, pH 4.5) using 3-Methyl-2-benzothiazolinone hydrazone and 3,3-dimethylaminobenzoic acid, which were oxidatively coupled by the enzyme. The results were corrected by activities of samples without manganese (for MnP) – the addition of manganese sulphate was substituted by an equimolar amount of EDTA. One unit of enzyme activity is defined as the amount of enzyme forming 1 μ M of reaction product per min. Enzyme activities were measured in three analytical replicates and expressed per g plant litter dry mass (4).

DNA extraction and purification

DNA was extracted from 0.1 g of homogenized litter and 0.5 g of soil samples (only from those collected in January 2011 and November 2011) by method SK described by Sagova-Mareckova *et al.* (2008). The method is based on bead-beating and phenol/chloroform extraction. The samples were purified by incubation with acetyltrimethylammonium bromide followed by chloroform extraction and incubation with CaCl_2 , and finally cleaned with GeneClean Turbo kit (MP Biomedicals, Santa Ana, CA, USA). DNA extracts were stored at -70 °C until the further analysis.

Quantitative real-time PCR

Quantification of 16S rRNA gene of *Actinobacteria*, *Alphaproteobacteria*, *Firmicutes* and 18S rRNA gene of fungi in the extracted DNA from litter samples were performed with primers listed in Table S1(6–8).

Quantitative PCR were performed using a StepOnePlus Real-Time PCR System (Applied Biosystems, Foster City, CA, USA) using 96-well plates with GoTaq qPCR Master Mix (Promega, Madison, WI, USA) containing SYBR Green as a double-stranded DNA (dsDNA) binding dye. The reaction mixture contained in a total

volume of 15 µl: 1× GoTaq qPCR Master Mix, primers (for concentrations see Table S1.) and 2 ng DNA sample. For all microbial groups the PCR cycling protocol consisted of initial denaturation at 95°C for 10 min, followed by 45 cycles of denaturation (95°C for 15s), annealing (for respective temperature see Table S1.) and elongation (72°C for 30s). Standards containing known number of copies of the target genes 16S rRNA from *Streptomyces europaeiscabiei* DSM 41802 (for *Actinobacteria*), *Bartonella-like* clone (for *Alphaproteobacteria*), *Paenibacillus* clone (for *Firmicutes*) and 18S rRNA from *Penicillium sp.* clone (for fungi), were serially diluted to calibrate each qPCR melting curve and were recorded to ensure qPCR specificity. Baseline and threshold calculations were performed with the StepOne v. 2.2.2 software. The inhibition was tested by serial DNA dilutions from each site. All qPCR measurements were done in duplicate.

Illumina MiSeq and data processing

Bacterial 16S rRNA gene fragments including the variable region V4 were amplified by PCR using universal primers with overhang adapters (Table S1.) (9). Construction of amplicon libraries and sequencing using the MiSeq sequencer (Illumina, San Diego, USA) were done at the DNA Services Facility, Research Resources Center, University of Illinois (Chicago, USA). The resulting paired sequence reads were merged, filtered, aligned using a reference alignment from the Silva database (10), and chimera checked using integrated Vsearch tool [12] according to the MiSeq standard operation procedure (MiSeq SOP, February 2018; (9)) in Mothur v. 1.39.5 software (11). A taxonomical assignment of sequence libraries was performed in Mothur using the non-redundant subset of Silva Small Subunit rRNA Database, release 132 (12) adapted for use in Mothur (https://mothur.org/w/images/3/32/Silva.nr_v132.tgz) as the reference database. Sequences of plastids, mitochondria, and those not classified in the domain *Bacteria* were discarded.

The sequence library was clustered into OTUs using the Usearch v10.0.240 software (13), and the OTU table was further processed using tools implemented in the Mothur software. Distance matrices describing the differences in community composition between individual samples were calculated using the Yue-Clayton theta calculator (14). Analysis of molecular variance (AMOVA; (15)) was based on a

matrix of Yue-Clayton theta distances. Metastats (16) analyses were used to detect differentially represented OTUs. Rarefaction analysis was performed with rarefaction.single function in Mothur, and areas under the rarefaction curves were calculated using trapezoidal rule in the R version 3.4.4 software (17). Figures were created using Inkscape (v0.92; <http://www.inkscape.org>).

The Illumina MiSeq 16S rRNA gene amplicon sequences have been deposited in the NCBI Sequence Read Archive under accession no. SRP158384 (<https://www.ncbi.nlm.nih.gov/sra/SRP158384>) and are available under BioProject no. PRJNA485178.




Fig. S1. Monthly averages of soil moisture tension and soil temperature measured by datalogger during litterbag experiment at the forest (black) and the grassland (white) sites.




Fig. S2. Carbon, nitrogen and phosphorus content in dry weight of beech (black circles), milkvetch (grey squares) and sedge (white triangles) at the forest and the grassland sites during decomposition experiment \pm SD (5 replicates). The x axes indicate sampling time (Initial - Jan, i.e. initial C,N and P content and litterbag sampling after 2 - 14 months).




Fig. S3. Average proportions of families within the phylum *Actinobacteria* in the bacterial 16S rRNA gene amplicon sequence libraries (5 replicates).

Labels at the horizontal axis indicate type of sampled plant litter (beech, milkvetch, and sedge), sampling time (Jan - Nov, i.e. after 2 - 12 months), and experimental site (G - grassland, F - forest).




Fig. S4. Average proportions of families within the phylum *Firmicutes* in the bacterial 16S rRNA gene amplicon sequence libraries (5 replicates).

Labels at the horizontal axis indicate type of sampled plant litter (beech, milkvetch, and sedge), sampling time (Jan - Nov, i.e. after 2 - 12 months), and experimental site (G - grassland, F - forest).




Fig. S5. Average proportions of families within the phylum *Proteobacteria* in the bacterial 16S rRNA gene amplicon sequence libraries (5 replicates) .

Labels at the horizontal axis indicate type of sampled plant litter (beech, milkvetch, and sedge), sampling time (Jan - Nov, i.e. after 2 - 12 months), and experimental site (G - grassland, F - forest).




Fig. S6. Average proportions of families within the phylum *Bacteroidetes* in the bacterial 16S rRNA gene amplicon sequence libraries (5 replicates).

Labels at the horizontal axis indicate type of sampled plant litter (beech, milkvetch, and sedge), sampling time (Jan - Nov, i.e. after 2 - 12 months), and experimental site (G - grassland, F - forest).




Fig. S7. Average proportions (5 replicates) of phyla in the bacterial 16S rRNA gene amplicon sequence libraries from the soil sampled at the experimental sites (G - grassland, F - forest) during the first (Jan) and the sixth (Nov) litter sampling.




Fig. S8. Average proportions of families within the phylum *Verrucomicrobia* in the bacterial 16S rRNA gene amplicon sequence libraries (5 replicates).

Labels at the horizontal axis indicate type of sampled plant litter (beech, milkvetch, and sedge), sampling time (Jan - Nov, i.e. after 2 - 12 months), and experimental site (G - grassland, F - forest).




Fig. S9. Changes of composition of bacterial communities during plant litter decomposition assayed by Illumina amplicon sequencing at the grassland (A, open symbols) and the forest (B, grey symbols) sites. Numbers indicate the OTUs significantly contributing to the difference between samples in pairwise comparison (Metastats, $p < 0.05$).




Fig. S10. Rarefaction analysis of 16S rRNA gene amplicon sequence libraries. The columns represent areas below the rarefaction curves \pm SD (5 replicates) for the entire bacterial libraries and for the subsets of *Actinobacteria*, *Alphaproteobacteria* and *Firmicutes*. The x axis indicates sampling time (Jan - Nov, i.e. litterbag samplings after 2 - 12 months).

Table S1. Primers used for qPCR and for MiSeq Illumina sequencing

Experiment	Primer	Sequence [5'-3']	Target	Conc.	Annealing temp.	Ref.
qPCR	EUB518 ^a	ATTACCGGGGCTGCTGG	Bacteria, 16S rRNA	0.2 µM	56 °C	[6]
	Act235	CGCGGCCTATCAGCTTGTG	Actinobacteria, 16S rRNA	0.2 µM	56 °C	[7]
	S-C-aProt-0528-a-S-19	CGGTAATACGRAGGGRGYT	Alphaproteobacteria, 16S rRNA	0.4 µM	61 °C	[9]
	S-C-aProt-0689-a-A-21	CBAATATCTACGAATTYCACCT	Alphaproteobacteria, 16S rRNA	0.4 µM	61 °C	[9]
	S-P-Firm-0352-a-S-18	CAGCAGTAGGGAATCTTC	Firmicutes, 16S rRNA	0.2 µM	57 °C	[9]
	S-P-Firm-0525-a-A-18	ACCTACGTATTACCGCGG	Firmicutes, 16S rRNA	0.2 µM	57 °C	[9]
	FF390	CGATAACGAAACGAGACCT	Fungi, 18S rRNA	0.6 µM	50 °C	[8]
	FR1	AICCATTCAAATCGGTAIT	Fungi, 18S rRNA	0.6 µM	50 °C	[8]
Sequencing	CS1-515F	ACACTGACGACATGGTTCTACAGTGCCAGCMGCCGCGGTAA	Bacteria, 16S rRNA			[10]
	CS2-806R	TACGGTAGCAGAGACTTGGTCTGGACTACHVGGGTWTCTAAT	Bacteria, 16S rRNA			[10]

^aEub518 was used as a reverse primer for Act235.

Table S2. Pairwise quantitative comparison of the experimental sites (F – forest, G – grassland) and plant substrates (B – beech, M – milkvetch, S – sedge) assayed by ANOVA using linear model with the additive effects, correlation coefficients of profiles determined by PERMANOVA using distance matrices (P-permanova) and variability of the profiles assayed by testing for homogeneity of multivariate dispersions using dissimilarity measures (P-disp).

	ANOVA		CORRELATION COEFFICIENT	
	Sites	Plant substrates	P-permanova	P-disp
Dry weight loss	G < F***	M > S***, M > B***, S > B***	0.00	0.382
N	G < F***	M > B***, S***	0.002	0.000
C	G < F***	-	0.140	0.383
P	G > F**	M > S***, M > B***, S > B*	0.000	0.005
Actinobacteria	G > F***	-	0.021	0.387
Fungi	-	M**, S*** > B	0.000	0.505
Alphaproteobacteria ^a	G > F**	M > B***, S***	0.029	0.002
Firmicutes	-	M > B***, S***	0.852	0.006
β - Glucosidase	G < F***	M > B***, S***	0.651	0.932
Cellobiohydrolase	G < F**	M > S**, M > B***, S > B***	0.278	0.867
β - Xylosidase	G < F***	M > S*, M > B***, S > B***	0.213	0.001
N-acetylglucosaminidase (Chitinase)	G < F***	M***, S*** > B	0.005	0.914
Glucuronidase	G < F**	M***, S*** > B	0.125	0.000
Acid Phosphatase	G < F***	M > B***, S***	0.067	0.208
Arylsulphatase	G < F***	M***, S*** > B	0.004	0.052
α - Glucosidase	G < F**	M***, S*** > B	0.191	0.018
Laccase	G < F**	S > B**	0.009	0.796
Mn - peroxidase	-	-	0	0.385

For ANOVA: * P < 0.05; ** P < 0.01; *** P < 0.001

For correlation coefficient: Bonferroni-corrected critical P value < 0.0035

^aonly for the first two sampling points (from November to January)

Table S3. Differences (indicated by P – values) of low-molecular-weighted compounds at beech (B), milkvetch (M) and sedge (S) between experimental sites (Forest and grassland) at sampling times (January 2011 – January 2012) assayed by non-parametric multivariate analysis using distance matrix.

	B	M	S	OVERALL
January 2011	0.008	0.000	0.000	<0.001
March	0.002	0.001	0.001	<0.001
May	0.004	0.001	0.001	<0.001
July	0.001	0.001	0.001	<0.001
September	0.003	0.001	0.001	<0.001
November	0.030	0.009	0.018	<0.001
January 2012	0.005	NA	0.008	<0.001

Bonferroni-corrected threshold P value < 0.016

Table S4. Differences (indicated by P – values) of bacterial community at beech (B), milkvetch (M) and sedge (S) between experimental sites (forest and grassland) at sampling times (January 2011 – January 2012) assayed by non-parametric multivariate analysis using Yue-Clayton distance matrix.

	B	M	S	OVERALL
January 2011	0.098	0.009	0.418	0.002
March	0.092	0.008	0.161	<0.001
July	0.008	0.064	0.006	<0.001
November	0.009	0.039	0.009	<0.001

Bonferroni-corrected threshold P value < 0.016

Table S5. AMOVA analysis of differences in composition of bacterial communities at beech (B), milkvetch (M) and sedge (S) at the forest and the grassland sites (F-values).

Grassland				Forest			
	df	M	S		df	M	S
B	1	4.39381*	1.09189	B	1	2.67483*	1.29985
M	1		4.56859*	M	1		3.05833*

* $P < 0.016$ (Bonferroni-corrected threshold)

References for supplementary text citations

1. Sagova-Mareckova M, Omelka M, Cermak L, Kamenik Z, Olsovska J, Hackl E, Kopecky J, Hadacek F. 2011. Microbial communities show parallels at sites with distinct litter and soil characteristics. *Appl Environ Microbiol* 77:7560–7567.
2. Baldrian P. 2009. Microbial enzyme-catalyzed processes in soils and their analysis. *Plant, Soil Environ* 55:370–378.
3. Bourbonnais R, Paice MG. 1990. Oxidation of non-phenolic substrates. *FEBS Lett* 267:99–102.
4. Šnajdr J, Cajthaml T, Valášková V, Merhautová V, Petránková M, Spetz P, Leppänen K, Baldrian P. 2011. Transformation of *Quercus petraea* litter: Successive changes in litter chemistry are reflected in differential enzyme activity and changes in the microbial community composition. *FEMS Microbiol Ecol* 75:291–303.
5. Sagova-Mareckova M, Cermak L, Novotna J, Plhackova K, Forstova J, Kopecky J. 2008. Innovative methods for soil DNA purification tested in soils with widely differing characteristics. *Appl Environ Microbiol* 74:2902–2907.
6. Muyzer G, de Waal EC, Uitterlinden AG. 1993. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl Environ Microbiol* 59:695–700.
7. Stach JEM, Maldonado LA, Ward AC, Goodfellow M, Bull AT. 2003. New primers for the class Actinobacteria: Application to marine and terrestrial environments. *Environ Microbiol* 5:828–841.
8. Prévost-Bouré CN, Christen R, Dequiedt S, Mougél C, Lelièvre M, Jolivet C, Shahbazkia HR, Guillou L, Arrouays D, Ranjard L. 2011. Validation and application of a PCR primer set to quantify fungal communities in the soil environment by real-time quantitative PCR. *PLoS One* 6.
9. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the miseq illumina sequencing platform. *Appl Environ Microbiol* 79:5112–5120.
10. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2012. The SILVA ribosomal RNA gene database project:

- improved data processing and web-based tools. *Nucleic Acids Res* 41:D590–D596.
11. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. 2016. VSEARCH: a versatile open source tool for metagenomics. *PeerJ* 4:e2584.
 12. Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J, Ludwig W, Glöckner FO. 2014. The SILVA and “all-species Living Tree Project (LTP)” taxonomic frameworks. *Nucleic Acids Res* 42:643–648.
 13. Edgar RC. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat Methods* 10:996–998.
 14. Yue JC, Clayton MK. 2005. A similarity measure based on species proportions. *Commun Stat - Theory Methods* 34:2123–2131.
 15. Martin AP. 2002. Phylogenetic Approaches for Describing and Comparing the Diversity of Microbial Phylogenetic Approaches for Describing and Comparing the Diversity of Microbial Communities. *DNA Seq* 68:3673–3682.
 16. White JR, Nagarajan N, Pop M. 2009. Statistical Methods for Detecting Differentially Abundant Features in Clinical Metagenomic Samples. *PLoS Comput Biol* 5:e1000352.
 17. R Core Team. 2018. R Core Team, R : A language and environment for statistical computing. R Found Stat Comput Vienna, Austria R Foundation for Statistical Computing.

Paper III: Litter chemical quality and bacterial community structure influenced decomposition in acidic forest soil

Litter chemical quality and bacterial community structure influenced decomposition in acidic forest soil

Authors

Andrea Buresova ^{1,3,4}, Vaclav Tejnecky ², Jan Kopecky ¹, Ondrej Drabek ², Pavla Madrova ¹, Nada Rerichova ², Marek Omelka ⁵, Petra Krizova ², Karel Nemecek ², Thomas B. Parr ⁶, Tsutomu Ohno ⁷, Marketa Sagova-Mareckova ⁸

Corresponding author: Vaclav Tejnecky, email: tejnecky@af.czu.cz

¹ Epidemiology and Ecology of Microorganisms, Crop Research Institute, Drnovska 507, Prague, Czech Republic

² Department of Soil Science and Soil Protection, Faculty of Agrobiological Sciences, Czech University of Life Sciences Prague, Kamýcka 129, Prague, Czech Republic

³ Department of Ecology, Faculty of Science, Charles University in Prague, Vinicna 7, Prague, Czech Republic

⁴ Microbial Ecology, University Claude Bernard Lyon 1, UMR CNRS 5557, INRA 1418, Villeurbanne, France

⁵ Department of Probability and Mathematical Statistics, Faculty of Mathematics and Physics, Charles University in Prague, Sokolovska 68, Prague 8, Czech Republic

⁶ Department of Biology, University of Oklahoma, 730 Van Vleet Oval, Norman, OK, USA

⁷ School of Food and Agriculture, University of Maine, 5763 Rogers Hall, Orono, ME, USA

⁸ Department of Microbiology, Dietetics and Nutrition, Faculty of Agrobiological Sciences, Czech University of Life Sciences Prague, Kamýcka 129, Prague, Czech Republic

29 **Abstract**

30 The study was designed to see if naturally occurring beech forests uptake different nutrients than
31 artificially planted spruce forests and if the structure of microbial communities participating in
32 decomposition corresponds to the soil types developed under the respective tree species. This corresponds
33 to previous observations that the soil under spruce forest is typically dominated by podzol, while the soil
34 under beech forest undergoes higher mineralization, which leads to moder humus and the development of
35 cambisol. A litterbag experiment was performed with beech and spruce litter placed respectively to beech
36 and spruce forests. In the beech forest, the observed litter decomposition rate reached the exponential phase
37 after 15 months, while in the spruce forest a steeper decrease was noted only 29 months after the litter
38 burial. Thus, the study focused on the period between 15 and 29 months to observe the exponential stages
39 of recalcitrant organic matter transformation. In this period, the chemical composition of two litters was
40 distinguished by higher contents of Mn and Ca in beech and higher content of Fe, S, N and P in spruce.
41 On top of that, the beech litter released higher amount of dissolved organic carbon and its associated
42 bacterial community was enriched with r-selected taxa, that correlated positively with organic acids and
43 cations. In contrast, the more acidic spruce litter was dominated by K-selected acidophilic community and
44 its turnover rate was slower, resulting in increased carbon sequestration. The higher pH and humus quality
45 of beech cambisol also correlates with its observed higher resilience to disturbances by acidification, pests
46 or climate change in mountainous environments.

47

48 **Keywords**

49 Soil type, Humus, Elemental composition, DOC, Bacteria, Fungi

50

51

52 1. Introduction

53 Human-introduced spruce forests are more vulnerable to environmental changes than natural beech
54 forests in the Czech Republic. The spruce forests vulnerability is even increased in acidic soils, which are
55 limited by availability of nutrients [1] and potential mobilization of toxic forms of Al [2,3]. In such
56 environments, the litter represents a major source of Ca, K, Mg, P and organic carbon (OC) in the upper
57 layer of soil [4].

58 The precise chemical composition of litter depends mostly on the requirements of particular plant
59 species and its nutrient uptake and allocation to tissues [5,6]. The nutrient content and composition of
60 organic compounds in litter further establish the microbial community corresponding to the litter type [7].
61 Microbial decomposers return the litter nutrients back to soil by activities mostly regulated by soil
62 chemistry, but also by local climate, i.e. precipitation, which have further impact on soil development and
63 environmental stability [8].

64 In particular, the elevated concentration of some litter components may speed up [9] or slow down
65 the decomposition rate [10]. For example, N and P stimulate decomposition rate in earlier stages but inhibit
66 its rate in later stages due to formation of complexes, which are recalcitrant for microbes [10,11]. Litter
67 decomposition is further dependent on the amount of dissolved organic carbon (DOC) released from litter
68 and its character, as the low level of DOC with its elevated aromaticity (A_{254} ; absorbance at 254 nm) and
69 complexity of organic compounds increase the litter recalcitrance [12]. Finally, the decomposition is also
70 predicted by humification index (HIX) of the decomposed litter, which determines how the particular litter
71 contributes to the humus formation [13].

72 Decomposition process over time may be illustrated as an asymptotic function. The upper
73 threshold of the function - limit value of decomposition – represents the phase, when the substrate is no
74 longer decomposed and remains stabilized in the soil [14]. The lag-time shows how long it takes a litter

75 to enter the exponential stage of decomposition. The long lag-time eventually implicates that old organic
76 matter is not necessarily stabilized and recalcitrant but that other mechanisms keep it from the
77 decomposition [15]. Later stages of litter decomposition are typically less explored due to longer time
78 required for experiments. Yet, they are needed mostly to evaluate chemical traits of recalcitrant litter and
79 its contribution to humus and soil formation.

80 Consequently, the contribution of different biotic and abiotic factors to low limit value or long lag-
81 time of recalcitrant litter is not well studied, although it is important for understanding which mechanism
82 and microbial decomposers participate on the release of sequestered nutrients. Furthermore, it is not
83 known, what is the role of different tree species in creating specific soil chemical conditions, and their
84 further influence on the composition and response of the decomposing microbial community [16,17].

85 Bacteria are generally less studied decomposers in forest soils compared to fungi, particularly in
86 later stages of decomposition [18]. That might be due to the previous association of bacterial
87 decomposition with readily degradable nutrient rich litter. However, some recent studies found that after
88 the initial stage of decomposition, bacterial community showed increased diversity and different structure
89 than in the early stages of decomposition [19]. Moreover, bacterial phyla *Actinobacteria* and *Firmicutes*,
90 which represent K-selected decomposers, dominated in stages characterized by high proportion of a
91 recalcitrant organic matter [20].

92 Thus, our study aimed in evaluating differences between chemical conditions and microbial
93 communities of native beech and introduced spruce litter during the decomposition stage following the
94 initial stage in two forests soils. The forests are located next to each other in the Jizera Mountains, Czech
95 Republic. They share an acid bedrock (porphyric granite) and mountainous climate with intense
96 precipitation of 800–1700 mm per year [16,21]. We hypothesized that the beech litter will promote faster
97 decomposition with a more abundant and diverse microbial community. In contrast, the spruce litter will

98 be preserved against decomposition by relatively high acidity and sequestered C, which will support less
99 abundant decomposers possibly dominated by a few taxa. The comparison consisted of detailed litter
100 chemical analyses (DOC, HIX, A₂₅₄), low molecular mass organic acids (LMMOA), N, P, Ca, Mn, Fe
101 together with the observation of the microbial succession on beech leaves and spruce needles using
102 quantification of bacteria (16S rRNA, ddPCR) and fungi (ITS region, ddPCR) and bacterial community
103 composition (16S rRNA, Illumina MiSeq) after months 15, 19 and 29 of decomposition. That knowledge
104 was further applied to resolve factors which support higher decomposition rate and more diverse microbial
105 community in the beech forest soil and to find the basis of the humus quality and soil type formation at
106 the two sites.

2. Materials and Methods

2.1. Litter bag experiment

The litterbag experiment was carried out in a beech forest (European beech, *Fagus sylvatica* L.) and a spruce forest (Norway spruce, *Picea abies* [L.] Karst.), which grow next to each other on the Palicnik hill located in the Jizera Mountains (Czech Republic; 50.8673125N, 15.2544103E), affected by acidification in the past [16]. The altitude of studied plots ranged from 635 (bottom limit) to 680 m a.s.l. (upper limit). The bedrock of both forest types is a medium-grained porphyric granite and granodiorite of the Upper Carboniferous age. The soil under spruce was classified as the Entic and Haplic Podzols and the soil under beech as the Aluminic Cambisol [22]. Mor or moder are respectively the prevailing types of humus [23]. Mean annual temperatures 7.33 °C, 9 °C and 8.7 °C and mean annual precipitations 922 mm, 683 mm and 696 mm were measured in years 2013, 2014 and 2015, respectively, in Liberec region, where the experiment was conducted [24]. Diameter at breast height of living trees was 534.6 (\pm 153.4) mm in beech forest and 405.8 (\pm 112.4) mm in spruce forest, canopy closure was 87 % in beech and 40 % in spruce forest and total area of rock was 811.9 m² in beech forest and 10.9 m² in spruce forest [23]. The experimental layout is showed in Supplementary Fig. S1.

Soil samples were collected from the surface litter (L), fermentation (F), humified (H) organic and organo-mineral (humic, A) horizons. Samples from subsurface B horizons (cambic or spodic) were also collected. Each sample was thoroughly mixed. The soil samples were analyzed in a “fresh” state representing actual soil moisture and in dry and sieved (< 2mm) state. In total, 8 beech and 7 spruce soil pits were described and sampled in September 2015.

Newly fallen spruce needles and beech leaves were collected in November 2011. Part of the collected litter was directly used for chemical analysis (denoted as the initial litter). Overall, 120 bags were made of polyamide with 99 μ m mesh size and 50 μ m fibre diameter were filled with 5 g of dry beech

(BL) or spruce (SL) litter. Sixty bags with BL were placed in the beech forest and sixty with SL in the spruce forest 1 m apart between litter (L) and fermentation (F) organic horizons in April 2013. Three litterbags from each litter type were collected during each sampling period in May 2013, June 2013, July 2013, August 2013, September 2013, October 2013, November 2013, July 2014, November 2014 and September 2015 (i.e. months 1, 2, 3, 4, 5, 6, 7, 15, 19 and 29 from the start of the experiment). After the collecting, samples were weighed and frozen for further analysis. Mass loss of litter and chemical analyses were assessed for all samples but only samples collected in months 15, 19 and 29 were chosen for further microbiological analysis because at those dates, an increased rate of decomposition was observed. The stage was thus termed exponential.

2.2. Chemical analysis

2.2.1. Soil analysis

Fresh soil samples were extracted by deionised water and pH and DOC were measured according to Tejnecky et al. [16,17]. Cation exchange capacity (CEC) and exchangeable elements (Ca_{ex} , Al_{ex} , H^+_{ex}) were determined according to Cools and De Vos [25] and base saturation (BS in %) was calculated as a ratio between CEC and base cations (Na^+ , K^+ , Ca^{2+} and Mg^{2+}).

2.2.2. Litter analyses

Water extracts were made from each defrosted sample (1 g of thawed sample with 30 ml deionized H_2O). Water extraction represents easily available elements or substances and preserves sample matrices [26]. The suspension was then centrifuged at 4000 g for 15 min; finally, extracts were filtered through a 0.45 μm nylon membrane filter (Sigma-Aldrich, MO, USA). In aqueous extracts, the following chemical

parameters were determined: pH, main inorganic anions (NO_3^- , NO_2^- , SO_4^{2-} and PO_4^{3-} , Cl^- , F^-) and low-molecular mass organic acids (LMMOA, lactate, acetate, formate and oxalate) by means of ion chromatography (IC, ICS 1600; Dionex; CA, USA) with suppressed conductivity. Cations (NH_4^+ , K^+ , Mg^{2+} , Na^+ , Ca^{2+}) were determined by means of ion chromatography (ICS 90, Dionex; CA, USA) and the contents of selected elements (P, Al, Fe, K, Ca, Mn) by inductively coupled plasma-optical emission spectrometer (ICP-OES, DUO iCap 7000; Thermo Scientific, MA, USA). The contents of N_{tot} , C_{tot} , S_{tot} were determined by Elementary Analyzer Flash 2000 NCS (Thermo Scientific, UK). For C_{tot} , S_{tot} , N_{tot} , 9 replicates were used for SL but 7 replicates were used for BL. The respective part of each sample was oven dried (40°C), homogenized and digested in Teflon vessels with the addition of HNO_3 (0.5 g of samples and 5-10 ml 65 % HNO_3). In the obtained solution, the contents of selected elements (P_{tot} , Ca_{tot} , Mg_{tot} , K_{tot} , Al_{tot} , Fe_{tot} and Mn_{tot}) were determined by ICP-OES (DUO iCap 7000, Thermo Scientific, MA, USA). Quality of digestion and analysis were controlled using standard reference materials (NIST SRM 1575a Pine Needles and NCS DC 73351 Tea). The contents of elements and substances were calculated to dry mass.

Dissolved organic carbon (DOC), absorbance at 254 nm (A_{254}) and humification index (HIX) were determined in water extracts of litter- using 0.2500 g of homogenized litter mixed with 12.5 ml of deionized water incubated for 24 h at 4 °C on an orbital shaker. The 4 °C condition was selected to minimize microbial DOM alteration during extraction [27]. Suspensions were centrifuged at 3000 g for 25 min before vacuum filtering through 0.4 µm polycarbonate membrane filter (Whatman, England). The DOC concentration in the extracts was determined by high temperature catalytic oxidation using a Shimadzu TOC-V analyzer (Kyoto, Japan). UV A_{254} was obtained using a Shimadzu UV-1800 spectrophotometer and a 1-cm quartz cuvette.

174 All litter extracts were diluted with deionized water to set the absorbance at 254 nm to 0.10 to
175 minimize inner-filtration effects in obtaining the fluorescence spectra [28]. Fluorescence spectra were
176 obtained using a Hitachi F-4500 spectrofluorometer (Tokyo, Japan). The instrumental parameters were:
177 excitation (EX) and emission (EM) slits, 5 nm; response time, 0.5 s; and scan speed, 2400 nm min⁻¹. The
178 EM spectra were obtained using 240 nm for EX and EM recorded from 300 to 600 nm. Correction factors
179 were applied according to the manufacturer's instructions, emission scans were blank subtracted, then
180 Raman Normalized, and then corrected for dilution in R 3.5.0 [29]. The HIX was calculated as ratio of the
181 sum of fluorescence intensities from 435 to 480 nm divided by sum of intensities from 300 to 345 nm and
182 435 to 480 nm [28].

183 From initial litter samples, total contents of Ca_{tot}, Mn_{tot}, Fe_{tot}, K_{tot}, Mg_{tot}, Na_{tot}, Al_{tot}, DOC, A₂₅₄ and
184 HIX were assessed.

185

186 **2.3. DNA extraction and amplicon sequencing**

187 DNA was extracted from 0.1 g of homogenized plant litter by method described in Sagova-
188 Mareckova et al. [30]. The method is based on bead-beating and phenol/chloroform extraction. The
189 samples were purified by incubation with acetyltrimethylammonium bromide followed by chloroform
190 extraction and incubation with CaCl₂. DNA samples were finally cleaned with GeneClean Turbo kit (MP
191 Biomedicals, CA, USA) and afterwards stored at -70 °C until further analysis. Fragments of the bacterial
192 16S rRNA gene including the variable region V3-V4 were amplified from the DNA samples by PCR,
193 using bacteria specific forward primer 341F (CCTACGGGAGGCAGCAG) and universal reverse primer
194 806R (GGACTACHVGGGTWTCTAAT). Construction of amplicon libraries and sequencing using
195 MiSeq sequencer (Illumina, San Diego, USA) were done at the DNA Services Facility, Research
196 Resources Center, University of Illinois (Chicago, USA).

2.4. Processing of sequence data

The resulting paired sequence reads were merged, filtered, aligned using reference alignment from the Sylva database [31], and chimera checked using integrated Vsearch tool[32] following the MiSeq standard operation procedure (Miseq SOP, February 2018; [33]) in Mothur v. 1.39.5 software [34]. A taxonomic assignment of sequence libraries was performed in Mothur using the recreated SEED database subset of Sylva Small Subunit rRNA Database, release 132 [35] adapted for use in Mothur (https://mothur.org/w/images/a/a4/Sylva.seed_v132.tgz) as the reference database. Sequences of plastids, mitochondria, and those not classified in the domain *Bacteria* were discarded. The sequence library was clustered into operational taxonomic units (OTUs) using the Uparse pipeline in Usearch v10.0.240 software [36], and the OTU table was further processed using tools implemented in the Mothur software. Rarefaction curves were used to detect observed OTUs for respective sequence numbers. Bray-Curtis distance matrices describing the differences in bacterial community composition between individual samples were calculated. The Illumina MiSeq 16S rRNA gene amplicon sequences were deposited in the NCBI Sequence Read Archive (www.ncbi.nlm.nih.gov/sra) as BioProject No.: PRJNA494609.

2.5. Quantification of bacteria and fungi

Quantification of bacteria (16S rRNA genes) and fungi (ITS region) from extracted DNA were done by droplet digital PCR (ddPCR). For gene amplification primers 16Seu27f (AGAGTTTGATCMTGGCKCAG) [37] and mixture of 783r-a, (CTACCAGGGTATCTAATCCTG), 783r-b (CTACCGGGGTATCTAATCCCG) and 783r-c (CTACCGGGGTATCTAATCCGG) [38] were

219 used for quantification of a bacterial 16S rRNA gene and primers ITS1F
220 (CTTGGTCATTTAGAGGAAGTAA) [39] and ITS4 (TCCTCCGCTTATTGATATGC) [40] were used
221 for quantification of a fungal ITS region. Reactions were prepared with QX200 EvaGreen Digital PCR
222 Supermix (Bio-Rad Laboratories, CA, USA) with 400 μ M final concentration of forward and reverse
223 primers. After generating droplets in QX200™ Droplet Generator (Bio-Rad Laboratories, CA, USA) the
224 PCR amplifications were performed (annealing temperatures: 56°C for bacteria and 54°C for fungi) and
225 droplets were detected fluorescently in Droplet Reader (Bio-Rad Laboratories, CA, USA). Results were
226 analyzed in QuantaSoft 1.7.4.0917 (Bio-Rad Laboratories, CA, USA).

227

228 **2.6. Statistical analysis**

229 Based on the observation of increased rate of decomposition, samples from months 15, 19 and 29
230 of each litter (n = 9) were used for statistical analysis (except for several analyses of mass loss and a few
231 selected chemical traits in the Supplementary Material). The effect of litter type and time during the whole
232 course of the experiment (from months 1 to 29) on mass loss was calculated by two-way ANOVA. Pair-
233 wise differences in mass loss between each month of decomposition for BL and SL were separately
234 calculated by ANOVA and Tukey's post-hoc test. Mass loss, chemical and microbiological variables
235 (abundance of bacteria 16S rRNA and fungi ITS region) from the three months of exponential stage
236 decomposition period on the BL (n = 9) and SL (n = 9) litter and chemical variables from the initial litter
237 (n = 3) were compared by Welch version of the two-sample t-test [41], which does not require equal
238 variances for both groups. The effect of time was not statistically analyzed.

239 Rarefaction analysis was used to demonstrate bacterial diversity expressed as the number of observed
240 OTUs with sequence numbers. Analysis of molecular variance (AMOVA) was calculated based on a
241 matrix of Bray-Curtis distances [42]. Metastats [43] analysis was used to show differences between

bacterial communities of BL and SL and to detect the differentially represented OTUs between the two litter types. The outcome was also visualized as heat maps. Maximum-likelihood phylogram of the OTU representative sequences was constructed using FastTree 2 [44]. Figures were created using Interactive Tree of Life online tool (<http://itol.embl.de>; [45], and Inkscape (v0.92; <http://www.inkscape.org>). The Bray-Curtis distance matrix was plotted by non-metric multidimensional scaling [46] using the Mass package v. 7.3-51.5 [47]. Vectors of environmental variables which significantly correlated to bacterial community were fitted to the nmDS plot in R using function envfit from the package vegan v. 2.5-6 [48]. All statistical calculations were done in the R computing environment [29] or in Mothur v. 1.39.5 software [34]. Figures were plotted with the help of vegan package [49].

3. Results

3.1. Soil chemical characteristics

Spruce forest soils were more acidic ($\text{pH} = 3.82 \pm 0.29$) than beech forest soils ($\text{pH} = 4.08 \pm 0.27$) except for F horizons, where pH were not statistically different (Table 1). H^+_{ex} represents exchangeable pool of H^+ and significantly differed between beech and spruce soil profile except the B horizon (Table 1). In the uppermost L horizon, the content of $\text{Ca}^{2+}_{\text{ex}}$, base saturation (BS), cation exchange capacity (CEC) and dissolved organic carbon (DOC) were significantly higher in the beech forest (Table 1). $\text{Al}^{3+}_{\text{ex}}$ was the lowest in L horizon and did not differ between the two forest soils (Table 1). Litter (L) organic horizons thickness under beech was on average 2.7 cm and 1.8 cm under spruce and bulk density under beech was 0.015 g.cm^{-3} and 0.048 g.cm^{-3} under spruce.

Table 1. Spruce and beech soil horizons (L - litter, F - fermentation, H - humic, A - organic and organo-mineral and B - subsurface cambic or spodic horizon) differences in pH, dissolved organic carbon (DOC), base saturation (BS), cation exchange capacity (CEC) and exchangeable cations (H⁺ex, Al³⁺ex, Ca²⁺ex). Average values and standard deviations (SD), degrees of freedom (Df), t-values and p-values are indicated (t-test).

	Horizon	Beech		Spruce		T- test		
		Average	SD	Average	SD	Df	t - value	p - value
pH	L	4.99	0.21	4.59	0.12	13	4.39	< 0.001
	F	4.20	0.29	3.90	0.43	13	1.62	0.129
	H	4.15	0.22	3.79	0.11	13	3.96	0.002
	A	4.32	0.19	3.93	0.08	12	5.10	< 0.001
	B	4.51	0.10	4.32	0.09	12	3.77	0.003
DOC [mg kg ⁻¹]	L	69.43	20.20	34.36	38.42	13	2.26	0.042
	F	678.0	358.1	473.8	250.8	13	1.26	0.230
	H	305.8	115.5	461.4	132.4	13	-2.43	0.030
	A	143.6	58.99	218.1	73.42	12	-2.10	0.058
	B	57.80	17.41	27.30	15.36	12	3.48	0.005
CEC [meq kg ⁻¹]	L	185.3	27.6	115.3	6.962	13	6.50	< 0.001
	F	135.0	18.2	148.5	6.515	13	-1.85	0.087
	H	100	11.44	122.5	12.64	13	-3.64	0.003
	A	80.58	14.36	84.81	7.703	12	-0.69	0.505
	B	46.72	10.65	50.49	9.712	12	-0.69	0.502
BS [%]	L	84.73	4.13	74.62	6.44	13	3.67	0.003
	F	47.29	14.71	40.60	5.99	13	1.12	0.283
	H	14.71	2.17	15.40	2.15	13	-0.62	0.548
	A	9.753	2.007	9.943	2.106	12	-0.17	0.866
	B	9.589	3.210	8.644	3.079	12	0.56	0.585
H ⁺ _{ex} [meq kg ⁻¹]	L	3.924	0.906	15.76	2.57	13	-12.23	< 0.001
	F	13.80	6.073	31.37	2.96	13	-6.95	< 0.001
	H	9.080	2.566	21.03	2.59	13	-8.95	< 0.001
	A	5.913	1.589	14.63	1.56	12	-10.37	< 0.001
	B	2.833	0.463	3.270	0.891	12	-1.15	0.270
Al ³⁺ _{ex} [meq kg ⁻¹]	L	2.899	6.388	7.916	5.171	13	-1.65	0.122
	F	46.30	17.44	47.37	11.30	13	-0.14	0.892
	H	70.19	9.06	73.10	10.20	13	-0.59	0.568
	A	62.70	11.71	54.15	7.97	12	1.59	0.137
	B	37.84	9.21	40.57	7.97	12	-0.59	0.565
Ca ²⁺ _{ex} [meq kg ⁻¹]	L	125.4	21.51	60.36	12.24	13	7.04	< 0.001
	F	47.10	22.68	39.09	10.36	13	0.86	0.407
	H	6.063	1.531	6.055	1.784	13	0.01	0.992
	A	2.668	0.582	2.238	0.732	12	1.218	0.247
	B	1.651	0.506	0.949	0.187	12	3.48	0.005

3.2. Litter mass loss

Litter mass loss was affected by factors of time as well as litter type (Table S1.) The average litter mass decreased from month 1 to month 29 by about 22.4 % in beech and 12.4 % in spruce forests ($p < 0.001$, Fig 1). In month 15, beech litter (BL) entered the exponential decomposition rate with the highest differences in mass loss between beech and spruce (SL) litter recorded in the last month 29 (Fig. 1). The mass loss most significantly differed from previous months in months 19 and 29 for BL and in month 29 for SL (Fig. 1). The mass loss was significantly different between BL and SL in months 15 to 29 (Table S2 A).




Fig. 1. The average residual mass loss (\pm SD) of beech (yellow) and spruce (green) during litter bag decomposition experiment ($n = 3$). The x axe indicates sampling time.

3.3. Litter chemical characteristics

In the initial litter, significantly higher contents of Ca_{tot} , Mn_{tot} , K_{tot} , Mg_{tot} and Na_{tot} were measured for BL, while higher, Al_{tot} , S_{tot} , N_{tot} , C_{tot} were measured for SL (Table S2 B). Differences for selected elements (N_{tot} , S_{tot} , P_{tot} , Fe_{tot} , Ca_{tot} , Mn_{tot}) from month 1 to month 29 were specific for each litter type (Fig.

280 S2; Table S3). A particularly fast decrease of Ca and Mn was observed in beech, and P in spruce in the
281 beginning of the decomposition process (Fig. S2).

282 In the exponential stage litter, (month 15, 19 and 29) contents of Ca_{tot} , Mn_{tot} and pH were
283 significantly higher for BL, while Fe_{tot} , Fe_{w} , P_{tot} , S, N were significantly higher for SL (Table S2 A). Yet,
284 most of the studied chemical parameters ($\text{NO}_3^-_{\text{w}}$, $\text{NO}_2^-_{\text{w}}$, $\text{SO}_4^{2-}_{\text{w}}$, $\text{PO}_4^{3-}_{\text{w}}$, Cl^-_{w} , F^-_{w} , $\text{NH}_4^+_{\text{w}}$, K^+_{w} , $\text{Mg}^{2+}_{\text{w}}$,
285 Na^+_{w} , $\text{Ca}^{2+}_{\text{w}}$, lactate_w, acetate_w, formate_w, oxalate_w, P_{w} , Al_{w} , Al_{tot} , Mn_{w} , Ca_{w} , K_{w} , K_{tot} , DOC, HIX) did
286 not significantly differ between the two litter types throughout the experiment (Table S2 A).

287

288 3.4. Microbial characteristics

289 Abundances of bacteria (16S rRNA gene copies) and fungi (ITS region gene copies) did not
290 significantly differ between BL and SL in months 15, 19 and 29 of decomposition (Table S2 A) but
291 composition of bacterial communities (Illumina MiSeq) differed significantly between BL and SL (Table
292 S4).

293 More specifically, the bacterial community composition was analyzed using a total of 665,606 16S
294 rRNA gene sequence reads were mapped to 2052 OTUs. Bacterial communities were represented by
295 highest proportions of *Proteobacteria*, followed by *Actinobacteria*, *Firmicutes*, *Acidobacteria* and
296 *Bacteroidetes* on BL and by *Actinobacteria*, followed by *Proteobacteria*, *Acidobacteria*, *Firmicutes* and
297 *Bacteroidetes* on SL (Fig. 2). Proportions of *Acidobacteria*, *Actinobacteria*, *Bacteroidetes*, *Firmicutes*,
298 *Patescibacteria* and *Proteobacteria* significantly differed between BL and SL (Table S4.)





Fig. 2. Average proportions of phyla in the bacterial 16S rRNA gene amplicon sequence libraries on beech (B) and spruce (S) litter (n = 3) in different sampled time points.

Proportions of *Proteobacteria* families *Acetobacteraceae* and *Burkholderiaceae* were higher on SL than on BL, while *Bradyrhizobiaceae* and unclassified *Gammaproteobacteria* were higher on BL than on SL (Fig. S3). Proportions of all *Actinobacteria* families were higher on SL than on BL (Fig. S4). Highest proportions of *Solirubrobacteraceae*, followed by Trebon clade, *Mycobacteriaceae*, were found on both litter types, but proportionally lower were on BL (Fig. S4). Proportions of *Firmicutes* families decreased in the following period with both litter types, while they were overall higher (especially in month 15 and in month 29) on BL than on SL. The family *Bacillaceae 1* followed by *Bacillaceae 2*, *Paenibacillaceae 1* and unclassified *Bacillales* dominated on BL, while *Alicyclobacillaceae* dominated on SL (Fig. S5).

The phylogenetic tree was built from 839 OTUs differing significantly between communities associated with SL and BL (Metastats, $p < 0.05$). Out of those, proportions of 294 OTUs were higher on

313 SL and 545 OTUs on BL. OTUs corresponding to *Actinobacteria*, *Acidobacteria*, *Firmicutes* and *Cand.*
314 *Saccharibacteria* were proportionally higher on SL, while *Proteobacteria*, *Bacteroidetes*,
315 *Verrucomicrobia*, *Planctomycetes* and *Parcubacteria* on BL (Fig. 3).



316 **Fig. 3.** Phylogenetic tree of OTUs that significantly differed between the communities associated with
317 beech (yellow) and spruce (green) litter after 15, 19 and 29 months of decomposition (Metastats, $p < 0.05$,
318 $n = 9$). The phylogeny was inferred by maximum likelihood analysis of representative sequences of each
319 OTU.


320

321 Changes in the composition of bacterial communities showed that higher number of OTUs
322 significantly contributing to differences between months 15 and 29 were found on BL (Metastats, $p <$
323 0.05 , Fig. 4). The number of OTUs significantly contributing to differences between litter types was the
324 highest in month 29 and the lowest in month 19 (Metastats, $p < 0.05$). Rarefaction curves demonstrated

325 that bacterial diversity was higher for the BL than the SL in months 15 and 19 (Fig. S6).

326

327 Heat maps (supported by Metastats) showed differences between BL and SL between months 15
 328 and 29 (Table S5 A, B). On BL proportions of 155 OTUs differed (Metastats, $p < 0.05$), out of which 123
 329 OTUs proportions increased and 32 OTUs decreased. On SL, proportions of only 67 OTUs differed, out
 330 of which 30 increased and 37 decreased during that period. The most abundant bacterial groups separating
 331 samples from month 15 and month 29 were on BL especially *Actinobacteria*, *Proteobacteria*,
 332 *Acidobacteria*, *Verrucomicrobia*, *Bacteroidetes*, *Firmicutes* and *WPS-2* (Table S5 A), while on SL it was
 333 *Acidobacteria*, *Actinobacteria*, *Proteobacteria*, *Patescibacteria* and *Firmicutes* (Table S5 B).




334 **Fig. 4.** Changes of composition of bacterial communities during plant litter decomposition on beech (B,
 335 yellow circle) and spruce (S, green triangle) after months 15, 19 and 29 assayed by Illumina amplicon
 336 sequencing. Numbers indicate the OTUs significantly contributing to the difference between samples in
 337 pairwise comparison (Metastats, $p < 0.05$, $n = 3$).

338 **3.5. NMDS analysis**

339 NMDS distinguished bacterial communities into two groups corresponding to BL and SL (Fig. 5).
 340 The bacterial community on BL most strongly positively correlated to DOC, oxalate, K_w , followed by
 341 A_{254} , Mn_{tot} , pH, Ca_{tot} , Ca_w , Mg^{2+}_w and bacterial diversity (Fig. 5). The bacterial community on SL most

342 strongly positively correlated to P_{tot} , followed by Fe_w , N_{tot} , S_{tot} , Fe_{tot} (Fig. 5).
 343 Composition of bacterial communities at both sites significantly correlated to pH, Ca_{tot} , Ca^{2+}_w ,
 344 Mg^{2+}_w , Fe_{tot} , Fe_w , Mn_w , Mn_{tot} , P_{tot} , $PO_4^{3-}_w$, K_{tot} , K^+_w , N_{tot} , S_{tot} , Al_{tot} , DOC, A_{254} , Acetate_w, Oxalate_w, HIX,
 345 mass loss and bacterial diversity (Table S6).



346
 347 **Fig. 5.** Sammon's projection showing distance and relationship between bacterial communities on beech
 348 (yellow circle) and spruce (green triangle) after months 15 (full symbol), 19 (empty symbol outlined by a
 349 thick line) and 29 (empty symbol outlined by a thin line) of decomposition. Linear vectors represent
 350 chemical and microbiological variables significantly correlating with the structure of bacterial community
 351 ($p < 0.001$ red vectors; $0.001 < p < 0.01$ blue vectors; $0.01 < p < 0.05$ grey vectors). A_{254} indicates
 352 aromaticity, HIX humification index and DOC dissolved organic carbon.

353
 354

4. Discussion

Abundances of bacteria and fungi did not differ between beech (BL) and spruce litter (SL) but different bacterial communities were established on the two litter types after 15, 19 and 29 months of decomposition, when it reached the exponential stage. The litter mainly differed by the length of time to reach the exponential stage but also by chemistry of litter and its changes over the following period. That complied with previous findings that the succession of bacterial communities rather than abundance reflected chemical differences between litter types [50,51] and that this was transferred to situations of the later decomposition stage [52,53].

More specifically, beech was typical by accumulation of Ca and Mg in leaves, which led to concentration of those elements in shallow soil layers and consequently to improvement of soil quality. That activity is called the base-pump strategy, which describes how trees selectively uptake certain nutrients from lower horizons by their roots and via litter fall increase their concentration in top soil horizons [5]. In contrast, spruce needles accumulated N and S, which probably originated in anthropogenic acidification (in form of HNO_3^- and SO_4^{2-} , respectively) occurring in the past [16]. Also, the fresh needles were enriched in Fe and Al, possibly due to preferential uptake of those elements from the granite bedrock by spruce [21,54].

On beech litter, bacterial communities were most strongly correlated to DOC, oxalate and K_w . Total DOC increased after the initial stage of decomposition and exposed higher degree of aromatic compounds (A_{254}) [12]. Therefore, the correlation of BL community to DOC demonstrates the litter degradability, which enables microbes to utilize released C specific to a given stage of decomposition. The positive correlation between LMMOA (low molecular mass organic acids), especially oxalate and Ca^{2+} to BL community can be explained by microbe solubilization of Ca from poorly dissolved biominerals e.g. calcium oxalate [55]. Additionally, BL bacterial community correlation to Mn_{tot} may

378 reflect the activity of manganese peroxidase, since Mn^{2+} is an essential component of this ligninolytic
379 enzyme [4,9]. The positive relationship of organic compounds such as DOC, A_{254} and LMMOA with the
380 BL community together with higher weight loss supports the observed more complete decomposition of
381 this substrate and confirms that BL was subjected to a more advanced stage of decay probably entering
382 the exponential stage of decomposition [15].

383 On spruce litter, a positive relationship of SL bacterial community to P, Fe, N and S showed that
384 these elements were not only higher there but also available to bacterial inhabitants. More precisely, P
385 was probably not precipitated by basic cations as it happened in the BL [17] and was intensively depleted
386 from SL over time. Similarly, Fe often occurs in organo-mineral complexes [56] or complexes with P-
387 PO_4^{3-} [57], which are resistant to degradation. However, the correlation of SL community to both Fe and
388 P suggests that the community may be enriched with bacteria, which can solubilize those complexes
389 possibly using organic acids or siderophores [58]. It is typical for activities of *Actinobacteria*, which
390 indeed dominated SL. A positive correlation to N may reflect either increased N litter concentration or
391 possibly also N fixation by specialized bacteria [59]. Nitrogen and P typically promote decomposition in
392 earlier stages, and thus their relatively high content may enable a gradual shift of SL into the exponential
393 decomposition stage, which was already partially noticeable [10]. Above that, N together with S belong
394 among the main agents causing soil acidification and thus, might explain the community shift towards
395 acidophilic taxa, such as a proportional increase of *Acidobacteria* [16,18].

396 Interestingly, no effect of Al on bacterial communities was determined on either litter type,
397 suggesting that bacterial communities may be protected from its toxicity by three different processes; i.e.
398 while acidic SL might release Al to lower organic horizons, Al on BL might be bound by cations, inorganic
399 anions (SO_4^{2-} , F^-) and organic compounds (e.g. fulvic acid, DOC and organic acids) [60,61], which
400 function as natural complexation agents eliminating Al toxicity [3] or that the bacteria are adapted by

401 resistance developed over the time of exposure [62].

402 Differing content and availability of nutrients of the two litter types corresponded to bacterial
403 functional groups represented by typical taxa. *Proteobacteria*, *Bacteroidetes*, *Verrucomicrobia* dominated
404 on BL. Especially, *Betaproteobacteria* and *Bacteroidetes* are considered r-selected, preferentially utilizing
405 easily accessible substrates in balanced environment [63]. SL was typical by higher proportion of
406 *Actinobacteria*, *Acidobacteria* and *Firmicutes*. *Actinobacteria* and *Firmicutes* are typical K-selected litter
407 decomposers [64] and *Acidobacteria* are a group, which preferentially inhabits low pH forest soils [18,65].

408 The most pronounced differences between the communities on the two studied litter types were
409 found in *Firmicutes*, of which *Bacillaceae* and *Paenibacilaceae* had higher proportions on BL, while
410 *Alicyclobacilales* dominated on SL. Out of those, *Bacilli* and *Paenibacilli* are highly abundant in organic
411 soil horizons, where they participate at decomposition of calcareous substrates. They are also involved in
412 the N cycle, including N fixation and P solubilization in soil [66]. In comparison, the family
413 *Alicyclobacilaceae* suggests shifting of the SL community towards acidophilic members since its
414 members were described as thermo-acidophilic [67]. However, *Bacillaceae* as well as *Alicyclobacilaceae*
415 significantly decrease over the exponential stage on BL and SL suggesting a correlation with nutrients
416 typical for each litter type.

417 Other important exponential stage decomposers were *Actinobacteria*. *Actinobacteria* may promote
418 availability of some limiting nutrients because they produce an array of enzymes, organic acids and
419 siderophores, which participate in solubilization of organic matter and soil complexes [68]. In particular,
420 a higher proportion of *Solirubrobacteraceae* was observed on SL and they belong among plant growth
421 promoting bacteria [69]. Additionally, Trebon clade, a new clade of *Actinobacteria* occurring in acidic
422 soils [70] had higher relative abundance on SL.

423 The different modes of pedogenesis observed under spruce or beech [16] resulted from both litter

chemistry and bacterial community composition, which together provide a particular potential for decomposing activities and explain the diverging decomposition rates [71]. Furthermore, on the BL, higher litter Ca_{tot} positively correlated to soluble C, C turnover and diversity of soil microbes, which may further improve the development of soil horizons [72]. In contrast, SL with lower amounts of Mn and Ca, higher amounts of Fe_{tot} , S_{tot} , N_{tot} and P_{tot} and low pH, reduced the rate of microbial decomposition, resulting in higher accumulation of lower quality litter in the soil. More specifically, lower pH in SL can lead to increased mobility of Al, Mn and Fe, which leach from the L horizon to deeper parts of soil profile causing the differentiation of soil profiles – creating podzol (e.g. Sauer et al. [73]). It has been observed previously that even small differences between the quality of humus and soil chemistry (pH) led to increasing podsolization process in cambisols [74]. Similar observation was previously reported from the Giant Mountains (Czechia) [75]. Thus, soils with different characteristics are developed under the two tree species and their different microbial communities and decomposition rate may have consequences in the stability and resilience of beech and spruce forests.

5. Conclusions

Our study compared the litter decomposition rate of beech litter (BL) and spruce litter (SL). The BL displayed lower C accumulation and higher release of aromatic DOC after 15, 19 and 29 months of decomposition, while SL showed lower pH and higher amount of Fe, S, N and P, resulting in accumulation of organic matter and thus, C sequestration in the soil.

The BL had the lag-time of decomposition delayed until month 15 but after that period the decomposition was relatively fast and supported by readily available compounds, while promoting the growth of r-selected taxa. On the contrary, SL was described as an oligotrophic environment typical with

446 K-selected taxa, which together with acidic conditions caused that the SL substrate to remain in the lag-
447 phase during the entire experiment.

448 The litter chemical quality and specific decomposition pathways determined by the presence of
449 soil microorganisms with different life strategies lead to a consequent transformation of litter to humus
450 with different quality in beech and spruce forests. Finally, that process results to the development of soils,
451 which undergo podsolization under spruce but not beech.

452 **6. Funding**

453 Research was supported by the Czech University of Life Sciences Prague (internal project n.
454 [21130/1312/3131](#)), Ministry of Education, Youth and Sports of the CR (projects No.
455 CZ.02.1.01/0.0/0.0/16_019/0000845 and LTC17075), Czech Science Foundation (grant no. 15-01312S),
456 and Ministry of Agriculture of the CR (Institutional Project RO 0418).

References

- [1] K.C. Rice, J.S. Herman, Acidification of Earth: An assessment across mechanisms and scales, *Appl. Geochemistry*. 27 (2012) 1–14. <https://doi.org/10.1016/J.APGEOCHEM.2011.09.001>.
- [2] T. Fukami, I.A. Dickie, J. Paula Wilkie, B.C. Paulus, D. Park, A. Roberts, P.K. Buchanan, R.B. Allen, Assembly history dictates ecosystem functioning: evidence from wood decomposer communities, *Ecol. Lett.* 13 (2010) 675–684. <https://doi.org/10.1111/j.1461-0248.2010.01465.x>.
- [3] L. Borůvka, A. Nikodem, O. Drábek, P. Vokurková, V. Tejnecký, L. Pavlů, Assessment of soil aluminium pools along three mountainous elevation gradients, *J. Inorg. Biochem.* 103 (2009) 1449–1458. <https://doi.org/10.1016/j.jinorgbio.2009.07.022>.
- [4] B. Berg, Litter decomposition and organic matter turnover in northern forest soils, *For. Ecol. Manage.* 133 (2000) 13–22. [https://doi.org/10.1016/S0378-1127\(99\)00294-7](https://doi.org/10.1016/S0378-1127(99)00294-7).
- [5] G. van der Heijden, E. Dambrine, B. Pollier, B. Zeller, J. Ranger, A. Legout, Mg and Ca uptake by roots in relation to depth and allocation to aboveground tissues: results from an isotopic labeling study in a beech forest on base-poor soil, *Biogeochemistry*. 122 (2015) 375–393. <https://doi.org/10.1007/s10533-014-0047-2>.
- [6] Z. Wang, A. Gottlein, G. Bartonek, Effects of growing roots of Norway spruce and European beech on rhizosphere soil solution chemistry, *J. Plant Nutr. Soil Sci.* 164 (2001) 35–41.
- [7] J. Rousk, S.D. Frey, Revisiting the hypothesis that fungal-to-bacterial dominance characterizes turnover of soil organic matter and nutrients, *Ecol. Monogr.* 85 (2015) 457–472. <https://doi.org/10.1890/14-1796.1>.
- [8] A. Bani, S. Pioli, M. Ventura, P. Panzacchi, L. Borruso, R. Tognetti, G. Tonon, L. Brusetti, The role of microbial community in the decomposition of leaf litter and deadwood, *Appl. Soil Ecol.* 126 (2018) 75–84. www.elsevier.com/locate/apsoil (accessed April 10, 2018).

- 480 [9] B. Berg, B. Erhagen, M.-B. Johansson, M. Nilsson, J. Stendahl, F. Trum, L. Vesterdal, Manganese
481 in the litter fall-forest floor continuum of boreal and temperate pine and spruce forest ecosystems -
482 A review, *For. Ecol. Manage.* 358 (2015) 248–260. <https://doi.org/10.1016/j.foreco.2015.09.021>.
- 483 [10] X. Ge, L. Zeng, W. Xiao, Z. Huang, X. Geng, B. Tan, Effect of litter substrate quality and soil
484 nutrients on forest litter decomposition: A review, *Acta Ecol. Sin.* 33 (2013) 102–108.
485 <https://doi.org/10.1016/j.chnaes.2013.01.006>.
- 486 [11] P.M. Vitousek, S. Porder, B.Z. Houlton, O.A. Chadwick, Terrestrial phosphorus limitation:
487 mechanisms, implications, and nitrogen–phosphorus interactions, *Ecol. Appl.* 20 (2010) 5–15.
488 <https://doi.org/10.1890/08-0127.1>.
- 489 [12] S.L. Mosier, E.S. Kane, D.L. Richter, E.A. Lilleskov, M.F. Jurgensen, A.J. Burton, S.C. Resh,
490 Interactive effects of climate change and fungal communities on wood-derived carbon in forest
491 soils, *Soil Biol. Biochem.* 115 (2017) 297–309. <https://doi.org/10.1016/J.SOILBIO.2017.08.028>.
- 492 [13] A. Don, K. Kalbitz, Amounts and degradability of dissolved organic carbon from foliar litter at
493 different decomposition stages, *Soil Biol. Biochem.* 37 (2005) 2171–2179.
494 <https://doi.org/10.1016/J.SOILBIO.2005.03.019>.
- 495 [14] B. Berg, A.V. De Santo, F.A. Rutigliano, A. Fierro, G. Ekbohm, Limit values for plant litter
496 decomposing in two contrasting soils - Influence of litter elemental composition, *Acta Oecologica.*
497 24 (2003) 295–302. <https://doi.org/10.1016/j.actao.2003.08.002>.
- 498 [15] M. Kleber, P.S. Nico, A. Plante, T. Filley, M. Kramer, C. Swanston, P. Sollins, Old and stable soil
499 organic matter is not necessarily chemically recalcitrant: Implications for modeling concepts and
500 temperature sensitivity, *Glob. Chang. Biol.* 17 (2011) 1097–1107.
501 <https://escholarship.org/content/qt8q25s86p/qt8q25s86p.pdf> (accessed October 2, 2018).

- [16] V. Tejnecký, M. Bradová, L. Borůvka, K. Němeček, O. Šebek, A. Nikodem, J. Zenáhlíková, J. Rejzek, O. Drábek, Profile distribution and temporal changes of sulphate and nitrate contents and related soil properties under beech and spruce forests, *Sci. Total Environ.* 442 (2013) 165–171. <https://doi.org/10.1016/j.scitotenv.2012.10.053>.
- [17] V. Tejnecký, N. Řerichová, M. Brandová, K. Němeček, H. Santrucková, C. Ash, O. Drábek, Litter decomposition as a source of active phosphates in spruce and beech mountainous forests affected by acidification, *Procedia Earth Planet. Sci.* 10 (2014) 130–132. <https://doi.org/10.1016/j.proeps.2014.08.043>.
- [18] S. Lladó, R. López-Mondéjar, P. Baldrian, Forest Soil Bacteria: Diversity, Involvement in Ecosystem Processes, and Response to Global Change, *Microbiol. Mol. Biol. Rev.* 81 (2017) e00063-16. <https://doi.org/10.1128/MMBR.00063-16>.
- [19] V. Tlaskal, J. Voříšková, P. Baldrian, Bacterial succession on decomposing leaf litter exhibits a specific occurrence pattern of cellulolytic taxa and potential decomposers of fungal mycelia, *FEMS Microbiol. Ecol.* 92 (2016) fiw177. <https://doi.org/10.1093/femsec/fiw177>.
- [20] A. Buresová, J. Kopecký, V. Hrdinková, Z. Kameník, M. Omelka, M. Šagová-Marecková, Succession of microbial decomposers is designated by litter type, but site conditions drive decomposition rates., *Appl. Environ. Microbiol.* (2019). <https://doi.org/10.1128/AEM.01760-19>.
- [21] J. Chaloupský, J. Červenka, J. Jetel, F. Kralík, J. Libalová, E. Pichová, J. Pokorný, K. Posmourný, J. Sekyra, O. Šrbený, K. Salanský, J. Šrámek, J. Václ, *Geologie Krkonos a Jizerských hor*, 1., Moravské tiskarské závody, 1989.
- [22] I.W.G. WRB, World reference base for soil resources 2006, *World Soil Resour. Reports.* 103 (2006). <http://www.fao.org> (accessed February 11, 2019).

- 524 [23] M. Bradová, V. Tejnecký, L. Borůvka, K. Němeček, C. Ash, O. Šebek, M. Svoboda, J. Zenáhlíková,
525 O. Drábek, The variations of aluminium species in mountainous forest soils and its implications to
526 soil acidification, *Environ. Sci. Pollut. Res.* 22 (2015) 16676–16687.
527 <https://doi.org/10.1007/s11356-015-4855-2>.
- 528 [24] Czech Hydrometeorological Institute, Historical data - Territorial air temperature and precipitation,
529 (n.d.). <http://portal.chmi.cz/historicka-data/pocasi/zakladni-informace?l=en>.
- 530 [25] N. Cools, B. De Vos, Sampling and Analysis of Soil. In: UNECE ICP Forests Programme
531 Coordinating Centre (ed.): Manual on methods and criteria for harmonized sampling, assessment,
532 monitoring and analysis of the effects of air pollution on forests, Thünen Institute of Forest
533 Ecosystems, Eberswalde, Germany, 2016. <https://doi.org/10.1063/1.2761021>.
- 534 [26] P. Hubova, V. Tejnecky, C. Ash, L. Boruvka, O. Drabek, Low-Molecular-Mass Organic Acids in
535 the Forest Soil Environment, *Mini. Rev. Org. Chem.* 14 (2017) 75–84.
536 <https://www.ingentaconnect.com/contentone/ben/mroc/2017/00000014/00000001/art00010>
537 (accessed October 3, 2018).
- 538 [27] L.X. Zhou, J.W.C. Wong, Microbial Decomposition of Dissolved Organic Matter and Its Control
539 during a Sorption Experiment, *J. Environ. Qual.* 29 (2000) 1852.
540 <https://doi.org/10.2134/jeq2000.00472425002900060017x>.
- 541 [28] T. Ohno, Fluorescence inner-filtering correction for determining the humification index of
542 dissolved organic matter, *Environ. Sci. Technol.* 36 (2002) 742–746.
543 <https://doi.org/10.1021/es0155276>.
- 544 [29] R Core Team, R Core Team, R : A language and environment for statistical computing., R Found.
545 Stat. Comput. Vienna, Austria. (2018) R Foundation for Statistical Computing.

- 546 [30] M. Sagova-Mareckova, L. Cermak, J. Novotna, K. Plhackova, J. Forstova, J. Kopecky, Innovative
547 methods for soil DNA purification tested in soils with widely differing characteristics, *Appl.*
548 *Environ. Microbiol.* 74 (2008) 2902–2907. <https://doi.org/10.1128/AEM.02161-07>.
- 549 [31] C. Quast, E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, F.O. Glöckner, The
550 SILVA ribosomal RNA gene database project: improved data processing and web-based tools,
551 *Nucleic Acids Res.* 41 (2012) D590–D596. <https://doi.org/10.1093/nar/gks1219>.
- 552 [32] T. Rognes, T. Flouri, B. Nichols, C. Quince, F. Mahé, VSEARCH: a versatile open source tool for
553 metagenomics, *PeerJ.* 4 (2016) e2584. <https://doi.org/10.7717/peerj.2584>.
- 554 [33] J.J. Kozich, S.L. Westcott, N.T. Baxter, S.K. Highlander, P.D. Schloss, Development of a dual-
555 index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the miseq
556 illumina sequencing platform, *Appl. Environ. Microbiol.* 79 (2013) 5112–5120.
557 <https://doi.org/10.1128/AEM.01043-13>.
- 558 [34] P.D. Schloss, S.L. Westcott, T. Ryabin, J.R. Hall, M. Hartmann, E.B. Hollister, R.A. Lesniewski,
559 B.B. Oakley, D.H. Parks, C.J. Robinson, J.W. Sahl, B. Stres, G.G. Thallinger, D.J. Van Horn, C.F.
560 Weber, Introducing mothur: Open-source, platform-independent, community-supported software
561 for describing and comparing microbial communities, *Appl. Environ. Microbiol.* 75 (2009) 7537–
562 7541. <https://doi.org/10.1128/AEM.01541-09>.
- 563 [35] P. Yilmaz, L.W. Parfrey, P. Yarza, J. Gerken, E. Pruesse, C. Quast, T. Schweer, J. Peplies, W.
564 Ludwig, F.O. Glöckner, The SILVA and “all-species Living Tree Project (LTP)” taxonomic
565 frameworks, *Nucleic Acids Res.* 42 (2014) 643–648. <https://doi.org/10.1093/nar/gkt1209>.
- 566 [36] R.C. Edgar, UPARSE: highly accurate OTU sequences from microbial amplicon reads, *Nat.*
567 *Methods.* 10 (2013) 996–998. <https://doi.org/10.1038/nmeth.2604>.

- [37] L. Čermák, J. Kopecký, J. Novotná, M. Omelka, N. Parkhomenko, K. Plháčková, M. Ságová-Marečková, Bacterial communities of two contrasting soils reacted differently to lincomycin treatment, *Appl. Soil Ecol.* 40 (2008) 348–358. <https://doi.org/10.1016/j.apsoil.2008.06.001>.
- [38] M. Sakai, A. Matsuka, T. Komura, S. Kanazawa, Application of a new PCR primer for terminal restriction fragment length polymorphism analysis of the bacterial communities in plant roots, *J. Microbiol. Methods.* 59 (2004) 81–89. <https://doi.org/10.1016/J.MIMET.2004.06.005>.
- [39] M. Gardes, T.D. Bruns, ITS primers with enhanced specificity for basidiomycetes - application to the identification of mycorrhizae and rusts, *Mol. Ecol.* 2 (1993) 113–118. <https://doi.org/10.1111/j.1365-294X.1993.tb00005.x>.
- [40] T.J. White, T. Bruns, S. Lee, J.W. Taylor, Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics, *Acad. Press. Inc., New York.* (1990) 315–322. https://scholar.google.cz/scholar?hl=cs&as_sdt=0%2C5&q=White+TJ%2C+Bruns+T%2C+Lee+S+%26+Taylor+J+%281990%29+PCR+protocols.+Amplification+and+Direct+Sequencing+of+Fungal+Ribosomal+RNA+Genes+for+Phylogenetics+%28Inniss+MA%2C+Gelfand+DH%2C+Sinsky+JJ+%26+Whi (accessed February 5, 2019).
- [41] B.L. Welch, On the comparison of several mean values: an alternative approach, *Biometrika.* 38 (1951) 330–336. <https://doi.org/10.1093/biomet/38.3-4.330>.
- [42] A.P. Martin, Phylogenetic Approaches for Describing and Comparing the Diversity of Microbial Communities, *DNA Seq.* 68 (2002) 3673–3682. <https://doi.org/10.1128/AEM.68.8.3673>.
- [43] J.R. White, N. Nagarajan, M. Pop, Statistical methods for detecting differentially abundant features in clinical metagenomic samples., *PLoS Comput. Biol.* 5 (2009) e1000352. <https://doi.org/10.1371/journal.pcbi.1000352>.

- 590 [44] M.N. Price, P.S. Dehal, A.P. Arkin, FastTree 2 - Approximately maximum-likelihood trees for
591 large alignments, PLoS One. 5 (2010) e9490. <https://doi.org/10.1371/journal.pone.0009490>.
- 592 [45] I. Letunic, P. Bork, Interactive tree of life (iTOL) v3: an online tool for the display and annotation
593 of phylogenetic and other trees, Nucleic Acids Res. 44 (2016) W242–W245.
594 <https://doi.org/10.1093/nar/gkw290>.
- 595 [46] W.N. Venables, B.D. Ripley, Random and Mixed Effects, in: Mod. Appl. Stat. with S. Springer,
596 New York, NY, Springer, New York, NY, 2002: pp. 271–300. [https://doi.org/10.1007/978-0-387-](https://doi.org/10.1007/978-0-387-21706-2_10)
597 [21706-2_10](https://doi.org/10.1007/978-0-387-21706-2_10).
- 598 [47] B. Ripley, B. Venables, D.M. Bates, K. Hornik, A. Gebhardt, D. Firth, Package “MASS,” 2019.
599 <http://www.stats.ox.ac.uk/pub/MASS4/> (accessed April 2, 2020).
- 600 [48] J. Oksanen, F.G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. Mcglinn, P.R. Minchin, R.B.
601 O’hara, G.L. Simpson, P. Solymos, M. Henry, H. Stevens, E. Szoecs, H.W. Maintainer, Package
602 “vegan” Title Community Ecology Package Version 2.5-6, 2019.
- 603 [49] J. Oksanen, F.G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. Mcglinn, P.R. Minchin, R.B.
604 O’hara, G.L. Simpson, P. Solymos, M. Henry, H. Stevens, E. Szoecs, H.W. Maintainer, Package
605 “vegan” Title Community Ecology Package, 2018.
606 <https://cran.ism.ac.jp/web/packages/vegan/vegan.pdf> (accessed October 5, 2018).
- 607 [50] J.C. Frankland, Fungal succession — unravelling the unpredictable, Mycol. Res. 102 (1998) 1–15.
608 <https://doi.org/10.1017/S0953756297005364>.
- 609 [51] M. Sauvadet, M. Chauvat, D. Cluzeau, P.A. Maron, C. Villenave, I. Bertrand, The dynamics of soil
610 micro-food web structure and functions vary according to litter quality, Soil Biol. Biochem. 95
611 (2016) 262–274. <https://doi.org/10.1016/j.soilbio.2016.01.003>.

- 612 [52] T. Fukami, Historical Contingency in Community Assembly: Integrating Niches, Species Pools,
613 and Priority Effects, *Annu. Rev. Ecol. Evol. Syst.* 46 (2015) 1–23. [https://doi.org/10.1146/annurev-](https://doi.org/10.1146/annurev-ecolsys-110411-160340)
614 [ecolsys-110411-160340](https://doi.org/10.1146/annurev-ecolsys-110411-160340).
- 615 [53] L.C. Cline, D.R. Zak, Initial colonization, community assembly and ecosystem function: fungal
616 colonist traits and litter biochemistry mediate decay rate, *Mol. Ecol.* 24 (2015) 5045–5058.
617 <https://doi.org/10.1111/mec.13361>.
- 618 [54] J. Kratina, Rozdíly Ve Vlastnostech Horských Lesních Půd Na Kyselých a Bazických Matečných
619 Horninách, *Geol. Výzkumy Na Moravě a ve Slezsku*. 17 (2010).
- 620 [55] J.M. Dauer, S.S. Perakis, Calcium oxalate contribution to calcium cycling in forests of contrasting
621 nutrient status, *For. Ecol. Manage.* 334 (2014) 64–73.
622 <https://doi.org/10.1016/J.FORECO.2014.08.029>.
- 623 [56] J.P. Boudot, A.B. Hadj, R. Steiman, F. Seigle-Murandi, Biodegradation of synthetic organo-
624 metallic complexes of iron and aluminium with selected metal to carbon ratios, *Soil Biol. Biochem.*
625 21 (1989) 961–966. [https://doi.org/10.1016/0038-0717\(89\)90088-6](https://doi.org/10.1016/0038-0717(89)90088-6).
- 626 [57] F. Lang, J. Bauhus, E. Frossard, E. George, K. Kaiser, M. Kaupenjohann, J. Krüger, E. Matzner,
627 A. Polle, J. Prietzel, H. Rennenberg, N. Wellbrock, Phosphorus in forest ecosystems: New insights
628 from an ecosystem nutrition perspective, *J. Plant Nutr. Soil Sci.* 179 (2016) 129–135.
629 <https://doi.org/10.1002/jpln.201500541>.
- 630 [58] R.C. Hider, X. Kong, Chemistry and biology of siderophores, *Nat. Prod. Rep.* 27 (2010) 637–657.
631 <https://doi.org/10.1039/b906679a>.

- [59] D. Finn, K. Page, K. Catton, E. Strounina, M. Kienzle, F. Robertson, R. Armstrong, R. Dalal, Effect of added nitrogen on plant litter decomposition depends on initial soil carbon and nitrogen stoichiometry, *Soil Biol. Biochem.* 91 (2015) 160–168. <https://doi.org/10.1016/J.SOILBIO.2015.09.001>.
- [60] O. Drábek, I. Kipkoech Kiplagat, M. Komárek, V. Tejnecký, L. Borůvka, Study of Interactions between Relevant Organic Acids and Aluminium in Model Solutions Using HPLC and IC, *Orig. Pap. Soil Water Res.* 10 (2015) 172–180. <https://doi.org/10.17221/256/2014-SWR>.
- [61] O. Drabek, L. Mladkova, L. Boruvka, J. Szakova, A. Nikodem, K. Nemecek, Comparison of water-soluble and exchangeable forms of Al in acid forest soils, in: *J. Inorg. Biochem.*, Elsevier Inc., 2005: pp. 1788–1795. <https://doi.org/10.1016/j.jinorgbio.2005.06.024>.
- [62] R. Garcidueñas Pifia, C. Cervantes, Microbial interactions with aluminium, *Bio Met.* 9 (1996) 311–316.
- [63] L. Philippot, S.G.E. Andersson, T.J. Battin, J.I. Prosser, J.P. Schimel, W.B. Whitman, S. Hallin, The ecological coherence of high bacterial taxonomic ranks, *Nat. Rev. Microbiol.* 8 (2010) 523–529. <https://doi.org/10.1038/nrmicro2367>.
- [64] F. Bastian, L. Bouziri, B. Nicolardot, L. Ranjard, Impact of wheat straw decomposition on successional patterns of soil microbial community structure, *Soil Biol. Biochem.* 41 (2009) 262–275. <https://www.sciencedirect.com/science/article/pii/S0038071708003696> (accessed April 20, 2018).
- [65] M. Štursová, L. Žifčáková, M.B. Leigh, R. Burgess, P. Baldrian, Cellulose utilization in forest litter and soil: identification of bacterial and fungal decomposers, *FEMS Microbiol. Ecol.* 80 (2012) 735–746. <https://doi.org/10.1111/j.1574-6941.2012.01343.x>.

- 654 [66] I. Mandic-Mulec, P. Stefanic, J. Van Elsas, Ecology of Bacillaceae, *Bact. Spore from Mol. to Syst.*
655 (2015) 59–85. <https://doi.org/10.1128/microbiolspec.TBS-0017-2013>.
- 656 [67] E. Ciuffreda, A. Bevilacqua, M. Sinigaglia, M. Corbo, E. Ciuffreda, A. Bevilacqua, M. Sinigaglia,
657 M.R. Corbo, Alicyclobacillus spp.: New Insights on Ecology and Preserving Food Quality through
658 New Approaches, *Microorganisms*. 3 (2015) 625–640.
659 <https://doi.org/10.3390/microorganisms3040625>.
- 660 [68] W. Wang, Z. Qiu, H. Tan, L. Cao, Siderophore production by actinobacteria, *BioMetals*. 27 (2014)
661 623–631. <https://doi.org/10.1007/s10534-014-9739-2>.
- 662 [69] I.H. Franke-Whittle, L.M. Manici, H. Insam, B. Stres, Rhizosphere bacteria and fungi associated
663 with plant growth in soils of three replanted apple orchards, *Plant Soil*. 395 (2015) 317–333.
664 <https://doi.org/10.1007/s11104-015-2562-x>.
- 665 [70] J. Kopecky, M. Kyselkova, M. Omelka, L. Cermak, J. Novotna, G.L. Grundmann, Y. Moënne-
666 Loccoz, M. Sagova-Mareckova, Actinobacterial community dominated by a distinct clade in acidic
667 soil of a waterlogged deciduous forest, *FEMS Microbiol. Ecol.* 78 (2011) 386–394.
668 <https://doi.org/10.1111/j.1574-6941.2011.01173.x>.
- 669 [71] J.A.J. Dungait, D.W. Hopkins, A.S. Gregory, A.P. Whitmore, Soil organic matter turnover is
670 governed by accessibility not recalcitrance, *Glob. Chang. Biol.* 18 (2012) 1781–1796.
671 <https://doi.org/10.1111/j.1365-2486.2012.02665.x>.
- 672 [72] J. Frouz, V. Jílková, T. Cajthaml, V. Pižl, K. Tajovský, L. Háněl, A. Burešová, H. Šimáčková, K.
673 Kolaříková, J. Franklin, J. Nawrot, J.W. Groninger, P.D. Stahl, Soil biota in post-mining sites along
674 a climatic gradient in the USA: Simple communities in shortgrass prairie recover faster than
675 complex communities in tallgrass prairie and forest, *Soil Biol. Biochem.* 67 (2013).
676 <https://doi.org/10.1016/j.soilbio.2013.08.025>.

- 677 [73] D. Sauer, H. Sponagel, M. Sommer, L. Giani, R. Jahn, K. Stahr, Podzol: Soil of the year 2007. A
678 review on its genesis, occurrence, and functions, *J. Plant Nutr. Soil Sci.* 170 (2007) 581–597.
679 <https://doi.org/10.1002/jpln.200700135>.
- 680 [74] V. Brahy, H. Titeux, B. Delvaux, Incipient podzolization and weathering caused by complexation
681 in a forest Cambisol on loess as revealed by a soil solution study, *Eur. J. Soil Sci.* 51 (2000) 475–
682 484. <https://doi.org/10.1046/j.1365-2389.2000.00324.x>.
- 683 [75] S. Dlouhá, L. Borůvka, L. Pavlů, V. Tejnecký, O. Drábek, Comparison of Al speciation and other
684 soil characteristics between meadow, young forest and old forest stands., *J. Inorg. Biochem.* 103
685 (2009) 1459–64. <https://doi.org/10.1016/j.jinorgbio.2009.07.024>.
- 686

Supplementary Information for


**Litter chemical quality and bacterial community structure influenced decomposition
in acidic forest soil**

Andrea Buresova, Vaclav Tejnecky, Jan Kopecky, Ondrej Drabek, Pavla Madrova, Nada
Rerichova, Marek Omelka, Petra Krizova, Karel Nemecek, Thomas B. Parr, Tsutomu
Ohno, Marketa Sagova-Mareckova


Corresponding author: Vaclav Tejnecky,
Email: tejnecky@af.czu.cz

This PDF file includes:
Figs. S1 to S6
Tables S1 to S6


Supplementary Figure S1. Spruce and beech experimental forest sites located at Palicnik hill (635 - 680 a.s.l.) in Jizera mountains in the Czech Republic. The size of the investigated area was 2925 m² in beech and 3300 m² in spruce forest. 60 bags filled with beech litter were placed to beech forest and 60 bags filled with spruce litter were placed to spruce forest. DBH of living trees was 534.6 (± 153.4) in beech forest and 405.8 (± 112.4) in spruce forest, total area of rock was 811.9 m² in beech forest and 10.9 m² in spruce forest and canopy closure was 87% in beech and 40% in spruce forest.



Beech forest




Spruce forest




- ◆ Tree
- * Litter bags position

- Rock and surface runoff
- Crown projection


Supplementary Figure S2. Changes in the contents of selected elements in course of beech (yellow) and spruce (dark green) litter decomposition.




Supplementary Figure S3. Average proportions of groups at the family level within the phylum *Proteobacteria* in the bacterial 16S rRNA gene amplicon sequence libraries (n=3).




Supplementary Figure S4. Average proportions of groups at the family level within the phylum *Actinobacteria* in the bacterial 16S rRNA gene amplicon sequence libraries (n=3).



Supplementary Figure S5. Average proportions of groups at the family level within the phylum *Firmicutes* in the bacterial 16S rRNA gene amplicon sequence libraries (n=3).



Supplementary Figure S6. Rarefaction analysis of the sequence libraries (n=3).



Supplementary Table S1. Results of two-way ANOVA calculating the influence of factor of time and litter type (beech and spruce litter), and their interactions on mass loss (degrees of freedom (df), F and p value are indicated).

	Sum of sqrs	df	Mean square	F-value	p-value
Time:	9429	10	942.9	15.13	<0.001
Litter:	1070	1	1070	17.17	<0.001
Interaction:	3309	10	330.9	5.31	<0.001
Within:	2741	44	62.3		
Total:	16548	65			

Supplementary Table S2A. Results of t-test (degrees of freedom (Df), t-values and p-values) comparing total and water extractable content of selected elements and compounds, pH, dissolved organic carbon (DOC), aromaticity (A_{254}), humification index (HIX) and mass loss (average values and standard deviations are indicated) between beech and spruce litter in months 15, 19 and 29 of decomposition.

			Beech	Spruce	T-test		
					Df	t-value	p-value
Mass loss	[% of initial mass]	Average SD	49.69 20.18	75.92 8.278	14	-3.39	0.004
C _{tot}	[%]	Average SD	45.64 1.701	45.78 0.978	13	-0.23	0.823
N _{tot}	[%]	Average SD	1.289 0.336	1.812 0.12	13	-4.36	0.001
S _{tot}	[%]	Average SD	0.0653 0.0204	0.115 0.0179	13	-5.17	<0.001
P _{tot}	[mg.kg ⁻¹]	Average SD	523.9 160.1	829.7 51.24	10.04	-5.47	<0.001
Fe _{tot}	[mg.kg ⁻¹]	Average SD	466.7 231.7	1592 754.4	9.54	-4.27	0.002
Mn _{tot}	[mg.kg ⁻¹]	Average SD	699.1 381.7	203.2 49.87	8.27	4.26	0.003
Ca _{tot}	[mg.kg ⁻¹]	Average SD	6348 4000	2522 384.2	8.12	2.43	0.041
K _{tot}	[mg.kg ⁻¹]	Average SD	667.7 309.0	479.6 164.4	11.69	1.49	0.163
Al _{tot}	[mg.kg ⁻¹]	Average SD	1542 1447	1103 328.4	8.49	0.7	0.500
Mg _{tot}	[mg.kg ⁻¹]	Average SD	250.3 137.7	238.3 72.45	12.27	0.24	0.811
PO ₄ ³⁻ _w	[μmol.kg ⁻¹]	Average SD	5758 3968	9602 2608	15.55	-2.02	0.061
Fe _w	[μmol.kg ⁻¹]	Average SD	207.1 95.48	386.0 106.9	15.96	-3.44	0.003
Mn _w	[μmol.kg ⁻¹]	Average SD	2175 2045	638.2 375.1	8.38	1.87	0.097
Ca ²⁺ _w	[μmol.kg ⁻¹]	Average SD	11744 8737	5381 2007	8.60	1.68	0.129
K ⁺ _w	[μmol.kg ⁻¹]	Average SD	7124 3390	6387 1509	11.13	0.48	0.644

SO ₄ ²⁻ _w	[μmol.kg ⁻¹]	Average SD	1012 630.0	956.1 382.4	14.14	0.24	0.814
Al _w	[μmol.kg ⁻¹]	Average SD	1133 803.8	1157 476.4	10.79	-0.06	0.950
Mg ²⁺ _w	[μmol.kg ⁻¹]	Average SD	1733 1524	1229 580.5	9.51	0.76	0.464
Cl _w	[μmol.kg ⁻¹]	Average SD	1340 971.0	848.7 210.7	9.26	1.46	0.177
NH ₄ ⁺ _w	[μmol.kg ⁻¹]	Average SD	6586 3528	10311 5019	12.74	-1.32	0.212
NO ₃ ⁻ _w	[μmol.kg ⁻¹]	Average SD	343.0 398.8	343.3 330.6	15.76	0	0.999
F _w	[μmol.kg ⁻¹]	Average SD	1199 765.1	953.1 451.0	12.05	0.46	0.657
Oxalate _w	[μmol.kg ⁻¹]	Average SD	283.7 172.2	196.9 60.90	9.51	1.24	0.244
Acetate _w	[μmol.kg ⁻¹]	Average SD	3231 2875	1549 1501	12.58	1.12	0.284
Lactate _w	[μmol.kg ⁻¹]	Average SD	482.7 291.7	335.2 203.9	12.40	0.8	0.439
Formate _w	[μmol.kg ⁻¹]	Average SD	3138 2197	2914 2949	15.86	0.19	0.852
LMMOA _w	[μmol.kg ⁻¹]	Average SD	7657 4673	5260 3582	11.96	0.98	0.346
pH		Average SD	4.446 0.261	4.041 0.133	12.48	4.08	0.001
DOC	[mmol.kg ⁻¹]	Average SD	852.0 298.7	619.8 46.59	8.31	2.05	0.073
A ₂₅₄		Average SD	0.201 0.092	0.116 0.0125	8.29	2.34	0.047
HIX		Average SD	0.648 0.165	0.719 0.032	8.5	-1.07	0.316
Bacteria	[gene copies.ng DNA ⁻¹]	Average SD	37221 17719	37003 7704	13.87	0.03	0.977
Fungi	[gene copies.ng DNA ⁻¹]	Average SD	5700 3655	7112 3364	15.97	-0.77	0.451

Supplementary Table S2B. Results of t-test (degrees of freedom (Df), t-values and p-values) comparing the total content of selected elements, dissolved organic carbon (DOC), aromaticity (A_{254}) and humification index (HIX) in initial beech and spruce litter.

		Beech		Spruce		T-test		
		Average	SD	Average	SD	Df	t-value	p-value
N_{tot}	[%]	0.793	0.0714	1.527	0.040	4	-15.489	0.0001
C_{tot}	[%]	46.39	0.328	47.49	0.258	4	-4.583	0.010
S_{tot}	[%]	0.061	0.003	0.112	0.002	4	-24.99	<0.001
Mn_{tot}	[mg.kg ⁻¹]	1079	36.55	206.3	10.23	4	39.83	<0.001
Fe_{tot}	[mg.kg ⁻¹]	131.2	3.139	1438	310.9	4	-7.280	0.001
Ca_{tot}	[mg.kg ⁻¹]	8709	507.3	2042	2.840	4	22.77	<0.001
Mg_{tot}	[mg.kg ⁻¹]	649.4	11.74	232.2	12.91	4	41.41	<0.001
K_{tot}	[mg.kg ⁻¹]	2312	2.896	852.8	60.86	4	41.50	<0.001
Na_{tot}	[mg.kg ⁻¹]	152.4	0.240	64.96	7.547	4	20.06	<0.001
Al_{tot}	[mg.kg ⁻¹]	132.7	1.119	961.4	115.0	4	-12.48	<0.001
DOC	[mmol.kg ⁻¹]	9333	174.8	11458	575.5	4	-6.118	0.004
A_{254}		0.096	0.001	0.112	0.002	4	-9.260	<0.001
HIX		0.350	0.001	0.513	0.012	4	-21.95	<0.001

Supplementary Table S3. Chemical variables changes over the time (from the 1st to 29th month of decomposition). B_0 indicates average value for time 0 and b_1 indicates average percentual increase/decrease if the time from the beginning increase twice. P value indicates if the value of the precedent month is not significantly different.

	SL			BL		
	b_0	b_1	p-value	b_0	b_1	p-value
N_{tot}	1.37	3.5	0.001	0.73	7.6	< 0.001
C_{tot}	46.48	-0.2	0.553	46.22	-0.3	0.202
S_{tot}	0.1	1.8	0.176	0.08	-1.3	0.478
Ca_{tot}	1804	9.7	0.027	15424	-20.1	< 0.001
Fe_{tot}	643.9	6.4	0.281	81.89	36.5	< 0.001
Mn_{tot}	200.6	5	0.244	1488	-17.1	0.001
P_{tot}	2224	-13.7	< 0.001	435.7	5.7	0.024

Supplementary Table S4. Analysis of molecular variance (AMOVA) between composition of bacterial communities and subcommunities of individual phyla on beech and spruce litter in months 15, 19 and 29 of the decomposition. F - statistic and p - values are indicated (Df = 17).

	F-stat.	P-value
Bacterial communities	11.8	<0.001
<i>Acidobacteria</i>	5.93	<0.001
<i>Actinobacteria</i>	9.41	<0.001
<i>Bacteroidetes</i>	6.36	<0.001
<i>Firmicutes</i>	52.4	<0.001
<i>Patescibacteria</i>	3.98	<0.001
<i>Proteobacteria</i>	7.85	<0.001

Supplementary Table S5. Average frequencies (n = 3) of selected OTUs significantly contributing to separation of bacterial communities in the samples collected after 15 and 29 months.

A. The most abundant OTUs significantly separating the beech litter samples B15 and B29 (Metastats, $p < 0.05$).

OTU	B15	B19	B29	Phylum	Class	Order	Family	Genus
OTU1	4.030	5.903	12.883	Proteobacteria	Alphaproteobacteria	Rhizobiales	Xanthobacteraceae	uncl.
OTU12	1.337	1.177	3.363	Proteobacteria	Alphaproteobacteria	Rhizobiales	Beijerinckiaceae	Roseiarcus
OTU17	0.430	1.220	2.227	Actinobacteria	Acidimicrobiia	uncl.		
OTU26	0.113	0.260	1.443	Proteobacteria	Alphaproteobacteria	Rhizobiales	Xanthobacteraceae	uncl.
OTU35	0.203	0.747	1.290	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	uncl.
OTU62	0.507	0.593	1.420	Proteobacteria	Gammaaproteobacteria	JG36-TzT-191	uncl.	
OTU140	0.163	0.400	1.063	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	uncl.
OTU76	0.087	0.220	0.823	Acidobacteria	Acidobacteriia	Solibacterales	Solibacteraceae Subgroup 3	Bryobacter
OTU172	0.027	0.400	0.667	Actinobacteria	Actinobacteria	Trebon clade	uncl.	uncl.
OTU82	0.100	0.170	0.657	Acidobacteria	Acidobacteriia	Solibacterales	Solibacteraceae Subgroup 3	Cand. Solibacter
OTU24	0.450	0.537	0.937	Actinobacteria	Acidimicrobiia	uncl.		
OTU93	0.007	0.077	0.447	WPS-2	uncl.			
OTU100	0.030	0.153	0.453	Proteobacteria	Deltaproteobacteria	Myxococcales	Polyangiaceae	Pajaroellobacter
OTU65	0.040	0.110	0.427	Actinobacteria	Thermoleophila	Solirubrobacterales	Solirubrobacteraceae	uncl.
OTU51	0.367	0.220	0.747	Proteobacteria	Alphaproteobacteria	Caulobacterales	Caulobacteraceae	uncl.
OTU124	0.073	0.107	0.400	Acidobacteria	Acidobacteriia	Acidobacteriales	Koribacteraceae	Cand. Koribacter
OTU121	0.007	0.087	0.273	Acidobacteria	Acidobacteriia	Solibacterales	Solibacteraceae Subgroup 3	Bryobacter
OTU228	0.193	0.103	0.450	Proteobacteria	Alphaproteobacteria	Caulobacterales	Caulobacteraceae	uncl.
OTU63	0.063	0.237	0.307	Acidobacteria	Acidobacteriia	Solibacterales	Solibacteraceae Subgroup 3	Cand. Solibacter
OTU129	0.243	0.193	0.477	Proteobacteria	Alphaproteobacteria	Rhizobiales	Beijerinckiaceae	Roseiarcus
OTU41	0.027	0.233	0.260	Actinobacteria	Acidimicrobiia	uncl.		
OTU220	0.023	0.127	0.243	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	uncl.
OTU149	0.000	0.117	0.207	Acidobacteria	Acidobacteriia	Acidobacteriales	Acidobacteriaceae Subgroup 1	Acidicapsa
OTU171	0.007	0.000	0.213	WPS-2	uncl.			
OTU233	0.023	0.037	0.223	Proteobacteria	Alphaproteobacteria	Rhizobiales	Xanthobacteraceae	uncl.
OTU135	0.000	0.010	0.180	Planctomycetes	Planctomycetacia	Isosphaerales	Isosphaeraceae	uncl.
OTU130	0.013	0.050	0.170	Acidobacteria	Acidobacteriia	Solibacterales	Solibacteraceae Subgroup 3	uncl.
OTU174	0.000	0.013	0.147	Proteobacteria	Gammaaproteobacteria	uncl.		
OTU1513	0.007	0.000	0.153	Actinobacteria	Acidimicrobiia	uncl.		
OTU637	0.037	0.087	0.177	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	uncl.
OTU178	0.003	0.123	0.143	Planctomycetes	Planctomycetacia	Isosphaerales	Isosphaeraceae	Aquisphaera
OTU103	0.000	0.190	0.133	Acidobacteria	Acidobacteriia	Subgroup 2	uncl.	
OTU97	0.037	0.203	0.167	Actinobacteria	Actinobacteria	Trebon clade	uncl.	uncl.
OTU289	0.003	0.023	0.127	Acidobacteria	Acidobacteriia	Solibacterales	Solibacteraceae Subgroup 3	Bryobacter
OTU269	0.007	0.120	0.130	Proteobacteria	Alphaproteobacteria	Elsterales	uncl.	
OTU247	0.020	0.150	0.143	Acidobacteria	Acidobacteriia	Acidobacteriales	Acidobacteriaceae Subgroup 1	Occallatibacter
OTU184	0.000	0.023	0.123	Acidobacteria	Acidobacteriia	Subgroup 2		
OTU518	0.020	0.133	0.140	Actinobacteria	Acidimicrobiia	uncl.		
OTU344	0.043	0.060	0.153	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	uncl.
OTU290	0.000	0.040	0.100	Bacteroidetes	Bacteroidia	Spingobacteriales	Spingobacteriaceae	Mucilaginibacter

OTU71	0.117	0.017	0.017	Proteobacteria	Alphaproteobacteria	Caulobacteriales	Caulobacteraceae	uncult.
OTU306	0.120	0.157	0.007	Patescibacteria	Saccharimonadia	Saccharimonadales	uncl.	
OTU529	0.153	0.080	0.033	Proteobacteria	Alphaproteobacteria	Rickettsiales	SM2D12	
OTU482	0.130	0.230	0.000	Proteobacteria	Gammaaproteobacteria	Enterobacteriales	Enterobacteriaceae	Escherichia-Shigella
OTU54	0.173	0.330	0.027	Bacteroidetes	Bacteroidia	Sphingobacteriales	Sphingobacteriaceae	Mucilaginibacter
OTU243	0.217	0.163	0.010	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	uncult.
OTU163	0.243	0.073	0.033	Proteobacteria	Alphaproteobacteria	Rhodospirillales	uncult.	
OTU267	0.223	0.040	0.003	Acidobacteria	Acidobacteriia	Acidobacteriales	Acidobacteriaceae Subgroup 1	Terriglobus
OTU177	0.253	0.097	0.030	Verrucomicrobia	Verrucomicrobiae	Methylocidiphilales	Methylocidiphilaceae	uncult.
OTU203	0.320	0.210	0.087	Verrucomicrobia	Verrucomicrobiae	Chthoniobacteriales	Chthoniobacteraceae	Chthoniobacter
OTU78	0.477	0.903	0.207	Bacteroidetes	Bacteroidia	Sphingobacteriales	Sphingobacteriaceae	Mucilaginibacter
OTU37	0.593	0.263	0.303	WPS-2	uncl.			
OTU49	0.387	0.080	0.090	Acidobacteria	Acidobacteriia	Acidobacteriales	Acidobacteriaceae Subgroup 1	Granulicella
OTU57	0.367	0.090	0.060	Proteobacteria	Alphaproteobacteria	Sphingomonadales	Sphingomonadaceae	Sphingomonas
OTU6	0.580	0.490	0.157	Actinobacteria	Actinobacteria	Corynebacteriales	Mycobacteriaceae	Mycobacterium
OTU86	0.710	0.050	0.220	Proteobacteria	Alphaproteobacteria	Sphingomonadales	Sphingomonadaceae	Novosphingobium
OTU2012	0.673	0.540	0.173	Actinobacteria	Actinobacteria	Micrococcales	Microbacteriaceae	uncl.
OTU55	0.557	0.570	0.053	Verrucomicrobia	Verrucomicrobiae	Pedospaerales	Pedospaeraceae	uncl.
OTU2	22.287	18.457	3.007	Firmicutes	Bacilli	Bacillales	Bacillaceae	Bacillus

B. The most abundant OTUs significantly separating the spruce litter samples S15 and S29 (Metastats, $p < 0.05$).

OTU	S15	S19	S29	Phylum	Class	Order	Family	Genus
OTU3	2.227	2.773	6.557	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	uncl.
OTU1	1.927	3.357	3.540	Proteobacteria	Alphaproteobacteria	Rhizobiales	Xanthobacteraceae	uncl.
OTU10	1.007	1.987	2.430	Acidobacteria	Acidobacteria	Acidobacterales	Acidobacteriaceae Subgroup 1	Granulicella
OTU83	0.570	0.993	1.587	Proteobacteria	Gammaproteobacteria	Betaproteobacteriales	Burkholderiaceae	Burkholderia-Caballeronia-Paraburkholderia
OTU22	0.430	0.287	1.413	Acidobacteria	Acidobacteria	Solibacterales	Solibacteraceae Subgroup 3	Bryobacter
OTU36	0.173	0.413	0.807	Proteobacteria	Gammaproteobacteria	Betaproteobacteriales	Burkholderiaceae	Burkholderia-Caballeronia-Paraburkholderia
OTU522	0.300	0.210	0.827	Acidobacteria	Acidobacteria	Solibacterales	Solibacteraceae Subgroup 3	Bryobacter
OTU64	0.217	0.373	0.660	Acidobacteria	Acidobacteria	Acidobacterales	Acidobacteriaceae Subgroup 1	Acidipila
OTU61	0.147	0.217	0.510	Proteobacteria	Alphaproteobacteria	Rhizobiales	Beijerinckiaceae	uncl.
OTU148	0.007	0.053	0.323	Proteobacteria	Gammaproteobacteria	Betaproteobacteriales	Burkholderiaceae	Burkholderia-Caballeronia-Paraburkholderia
OTU1768	0.333	0.127	0.637	Actinobacteria	Thermoleophila	Solirubrobacterales	Solirubrobacteraceae	Conexibacter
OTU88	0.200	0.077	0.500	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	Acidiphilium
OTU75	0.207	0.093	0.490	Actinobacteria	Acidimicrobiia	IMCC26256	uncl.	uncl.
OTU46	0.167	0.230	0.447	Proteobacteria	Alphaproteobacteria	Rhizobiales	Beijerinckiaceae	Methylocapsa
OTU156	0.043	0.160	0.213	Acidobacteria	Acidobacteria	Acidobacterales	Acidobacteriaceae Subgroup 1	Acidipila
OTU172	0.230	0.110	0.353	Actinobacteria	Actinobacteria	Trebon clade	uncl.	uncl.
OTU219	0.017	0.030	0.127	Acidobacteria	Acidobacteria	Acidobacterales	Acidobacteriaceae Subgroup 1	uncl.
OTU543	0.037	0.020	0.143	Acidobacteria	Acidobacteria	Acidobacterales	Acidobacteriaceae Subgroup 1	uncl.
OTU769	0.123	0.147	0.020	Patescibacteria	Saccharimonadia	Saccharimonadales	uncl.	uncl.
OTU1674	0.187	0.577	0.077	Proteobacteria	Alphaproteobacteria	Micropepsales	Micropepsaceae	uncl.
OTU432	0.130	0.030	0.017	Actinobacteria	Thermoleophila	Solirubrobacterales	67-14	uncl.
OTU128	0.187	0.470	0.073	Patescibacteria	Saccharimonadia	Saccharimonadales	uncl.	uncl.
OTU703	0.133	0.060	0.007	Patescibacteria	Saccharimonadia	Saccharimonadales	uncl.	uncl.
OTU482	0.127	0.160	0.000	Proteobacteria	Gammaproteobacteria	Enterobacterales	Enterobacteriaceae	Escherichia-Shigella
OTU147	0.263	0.180	0.117	Actinobacteria	Acidimicrobiia	IMCC26256	uncl.	uncl.
OTU351	0.210	0.073	0.047	Firmicutes	Bacilli	Bacillales	Alicyclobacillaceae	Alicyclobacillus
OTU278	0.410	0.137	0.150	Actinobacteria	Thermoleophila	Solirubrobacterales	Solirubrobacteraceae	Conexibacter
OTU296	0.380	0.110	0.093	Actinobacteria	Actinobacteria	Corynebacteriales	Mycobacteriaceae	Mycobacterium
OTU32	1.227	0.700	0.830	Actinobacteria	Actinobacteria	Trebon clade	uncl.	uncl.
OTU68	0.617	0.637	0.177	Proteobacteria	Alphaproteobacteria	Micropepsales	Micropepsaceae	uncl.
OTU19	1.403	1.130	0.887	Acidobacteria	Acidobacteria	Acidobacterales	Acidobacteriaceae Subgroup 1	Granulicella
OTU72	0.863	0.717	0.047	Patescibacteria	Saccharimonadia	Saccharimonadales	uncl.	uncl.
OTU167	1.633	0.590	0.643	Actinobacteria	Thermoleophila	Solirubrobacterales	Solirubrobacteraceae	Conexibacter
OTU8	5.317	1.053	1.057	Firmicutes	Bacilli	Bacillales	Alicyclobacillaceae	Alicyclobacillus

Supplementary Table S6. Statistical correlations (coefficient of determination (R^2) and p - values) of total and water extractable content of selected elements and compounds, pH, dissolved organic carbon (DOC), aromaticity (A_{254}), humification index (HIX), abundance of bacteria, fungi and bacterial diversity (Bact. div) with bacterial communities on beech and spruce litter after months 15, 19 and 29 of decomposition (number of permutations = 99999).

	C _{tot}	N _{tot}	S _{tot}	P _{tot}	Fe _{tot}	Mn _{tot}	Ca _{tot}	K _{tot}	Al _{tot}	Mg _{tot}
R ²	0.310	0.473	0.591	0.629	0.535	0.601	0.476	0.422	0.476	0.338
p-value	0.062	0.008	0.002	0.001	0.004	0.002	0.009	0.017	0.006	0.046

	PO ₄ ³⁻ _w	Fe _w	Mn _w	Ca ²⁺ _w	K ⁺ _w	SO ₄ ²⁻ _w	Mg ²⁺ _w	Cl _w	NH ₄ ⁺ _w	NO ₃ ⁻ _w
R ²	0.408	0.589	0.370	0.493	0.602	0.189	0.569	0.084	0.172	0.226
p-value	0.016	0.001	0.032	0.007	0.001	0.207	0.002	0.526	0.242	0.142

	Oxalate _w	Acetate _w	Lactate _w	Formate _w	LMMOA _w
R ²	0.613	0.460	0.356	0.017	0.195
p-value	0.001	0.011	0.037	0.883	0.198

	pH	DOC	A ₂₅₄	HIX	Bacteria	Fungi	Bact. div.	Mass loss
R ²	0.542	0.702	0.638	0.421	0.301	0.091	0.578	0.376
p-value	0.003	< 0.001	0.001	0.015	0.069	0.487	0.002	0.030

Paper IV: Litter traits and rainfall reduction alter microbial
litter decomposers: the evidence from three Mediterranean
forests

RESEARCH ARTICLE

Litter traits and rainfall reduction alter microbial litter decomposers: the evidence from three Mediterranean forests

S. Pereira¹, A. Burešová^{2,3,4}, J. Kopecký⁵, P. Mádrová⁵, A. Aupic-Samain¹, C. Fernandez¹, V. Baldy^{1,*} and M. Sagova-Mareckova⁵

¹Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE UMR 7263, Marseille, France, ²Department of Ecology, Faculty of Science, Charles University in Prague, Vinicna 7, Prague, Czech Republic, ³Ecologie Microbienne, Université Claude Bernard Lyon 1, UMR CNRS 5557, INRA 1418, Villeurbanne, France, ⁴Laboratory for Diagnostics and Epidemiology of Microorganisms, Crop Research Institute, Drnovska 507, CZ-16106 Prague 6, Ruzyně, Czech Republic and ⁵Czech University of Life Sciences, Faculty of Agrobiological Sciences, Department of Microbiology, Nutrition and Dietetics, Kamýcká 129, 165 00 Praha 6 - Suchbátka, Czech Republic

*Corresponding author: Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE UMR 7263, Marseille, France. E-mail: virginie.baldy@imbe.fr

One sentence summary: This study addresses the impact of climate change, forest sites and tree species on microbial communities and litter decomposition process in three Mediterranean forests.

Editor: Taina Pennanen

ABSTRACT

The objective of the study was to evaluate changes in microbial communities with the predicted arrival of new species to Mediterranean forests under projected intensification of water stress conditions. For that, litter from three Mediterranean forests dominated respectively by *Quercus pubescens* Willd., *Quercus ilex* L. and *Pinus halepensis* Mill. were collected, and placed to their 'home' forest but also to the two other forests under natural and amplified drought conditions (i.e. rainfall reduction of 30%). Quantitative PCR showed that overall, actinobacteria and total bacteria were more abundant in *Q. pubescens* and *Q. ilex* than in *P. halepensis* litter. However, the abundance of both groups was dependent on the forest sites: placement of allochthonous litter to *Q. pubescens* and *P. halepensis* forests (i.e. *P. halepensis* and *Q. pubescens*, respectively) increased bacterial and fungal abundances, while no effect was observed in *Q. ilex* forest. *P. halepensis* litter in *Q. pubescens* and *Q. ilex* forests significantly reduced actinobacteria (A/F) and total bacteria (B/F) to fungi ratios. The reduction of rainfall did not influence actinobacteria and bacteria but caused an increase of fungi. As a result, a reduction of A/F ratio is expected with the plant community change towards the dominance of spreading *P. halepensis* under amplified drought conditions.

Keywords: litter decomposition; fungi; actinobacteria; Mediterranean forest; climate change

INTRODUCTION

Microorganisms are key actors in nutrient cycling and carbon turnover during litter decomposition (Gessner et al. 2010;

Garcia-Pausas and Paterson 2011). At a global scale, it is estimated that the soil C coming from soil microbial biomass is about 16.7 Pg C, which represents approximately 2.3–2.4% of total organic C found in soil in the top 30 cm (Xu, Thornton

Received: 15 December 2018; Accepted: 23 October 2019

© FEMS 2019. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com

and Post 2013). Heterotrophic respiration resulting from microbial litter decomposing activity can account for 10–90% of the CO₂ efflux from soils and substantially affect the atmospheric CO₂ concentration (Manzoni et al. 2012). However, annual reduction of rainfall by up to 30% predicted for the Mediterranean basin (Giorgi and Lionello 2008; IPCC 2014; Hertig and Trambly 2017) may significantly impact microbial abundance and structure, and thus, also the overall C dynamics (Wardle and Putten 2002).

Studies have already demonstrated shifts in plant community composition in response to the changed climatic conditions (Morales et al. 2007; Bertrand et al. 2011; Guiot and Cramer 2016). In southern France, dominating downy oak (*Quercus pubescens* Willd.) is expected to be replaced by more drought tolerant species: the sclerophylls' evergreen holm oak (*Quercus ilex* L.) and Aleppo pine (*Pinus halepensis* Mill.) (Barbero et al. 1990; Fortunel et al. 2009). Additionally, the increase of water stress may induce the synthesis of plant secondary metabolites (e.g. phenolics) known for their phytotoxic activities and anti-appetizing effects, which limit leaf consumption by fauna (Chomel et al. 2014). This investment in defence mechanisms over primary metabolic processes may result on the one hand to the decrease of primary productivity (Ogaya and Peñuelas 2007) and thus litter inputs, and on the other hand, in the reduction of litter quality through the increase of recalcitrant organic compounds (i.e. lignin, wax, polyphenolics) (Ormeño et al. 2006; Hättenschwiler and Jørgensen 2010; Sagova-Mareckova et al. 2011). Moreover, the reduction of moisture level may induce physiological stress and limit dispersal of microorganisms, enzymatic activities and substrate diffusion (Bouskill et al. 2013). Thus, the reduction of moisture levels and aboveground litter quality and quantity may impose new community dynamics belowground (Lunghini et al. 2013; Cleveland et al. 2014) and lead to a shift of soil microbial communities to more drought tolerant species, capable of using recalcitrant C sources and resistant to desiccation (Yuste et al. 2011; Peñuelas et al. 2012; Allison et al. 2013). Actinobacteria are known to thrive in low resource environments (Fierer, Bradford and Jackson 2007) and low soil moisture conditions (Yuste et al. 2011; De Vries and Shade 2013; Acosta-Martinez et al. 2014) reaching high abundance in arid soils (Okoro et al. 2009). They are well protected by a strong, thick interlinked peptidoglycan cell wall, which explains their high resistance to low osmotic potential conditions (Davet 2004). They produce enzymes able to degrade a wide range of biopolymers, whose building blocks including aromatic compounds (McCarthy and Williams 1992) have a vital role in organic matter turnover and humus formations (Anandan, Dharumadurai and Manogaran 2016). They produce a variety of secondary metabolites (Anandan et al. 2016), through which they interact with other microbial groups (Lewin et al. 2016). However, little is known about how microbial communities and actinobacteria in particular, would be affected by future reduction of rainfall, both directly by decrease of moisture level and indirectly through the plant species shift. In general, litter types sharing a common history with local microbial communities (autochthonous) have faster decomposition rates than litter types coming from distant (allochthonous) habitats (Gholz et al. 2000; Strickland et al. 2009). Consequently, changes in biochemical composition of local litter with the arrival of new species may constrain microbial activity and impact negatively litter decomposition (Schimel, Balser and Wallenstein 2007).

In here, we examine how reduction of rainfall and change in plant species may affect belowground microbial community's (actinobacteria, fungi and total bacteria) abundance and

structure and consequently also the litter decomposition process. For that, we performed a litter transplanting experiment between three main Mediterranean forests (*Q. pubescens* Willd., *Q. ilex* L., and *P. halepensis* Mill.) occurring in Southeast of France, under natural and amplified drought conditions for simulating the 30% rainfall reduction. We hypothesised that the arrival of new litter under amplified drought conditions will have a negative impact on microbial abundance and litter decomposition.

MATERIALS AND METHODS

Sites description

This study was conducted at the three forest sites in the Mediterranean region in southern France: the *Q. pubescens* forest at Oak Observatory at the 'Observatoire de Haute Provence' (O₃HP; Santonja et al. 2015a), the *Q. ilex* forest at Puéchabon (Limousin et al. 2008) and the mixed *P. halepensis* forest at Font-Blanche (Table 1). During the study period, the mean annual temperature was 12.1°C, 13.7°C and 14.0°C, and the cumulative annual precipitation was 609 mm, 794 mm and 632 mm, respectively (Table 1; Supplementary Figure 1, Supporting Information). There were no significant differences between sites in the mean annual temperature and cumulative annual precipitation during the study period (One-way ANOVA, $P < 0.05$). Each site included a control plot (natural drought—ND) and a rain exclusion plot (amplified drought—AD) to mimic the future reduction of rainfall predicted by the climatic model for the Mediterranean Region (about -30% yearly; Table 1 and Supplementary Figure 2, Supporting Information).

Experimental design and sample processing

In 2014, senescent leaves were collected in the control plot of the three forests. For that, litter traps were used during the abscission period that occurred from October to November for the broadleaves (*Q. ilex* and *Q. pubescens*) and from June to September for the needles (*P. halepensis*). Leaves and needles were air dried and stored at room temperature in paper boxes until the beginning of the experiment.

Litter decomposition was assessed by using the litter bag method (Swift et al. 1979). Four-mm mesh litter bags (20 × 20 cm) containing 10 g (air-dried) of the senescent material were used to perform the experiment. Litter transplants were made between each site for the three species considered, i.e. a litter bag containing the litter of each species placed on each forest site, under two drought conditions: natural (ND) and amplified (AD) (see Supplementary Figure 3, Supporting Information).

Thus, the experiment consisted in 18 modalities corresponding to the 3 forests *Q. ilex*, *Q. pubescens* and *P. halepensis* × 3 plant species (*Q. ilex*, *Q. pubescens* and *P. halepensis*) × 2 drought conditions (ND and AD). In total, 126 litter bags (18 modalities × 1 sampling date × 7 replicates) were analysed. Litter bags were placed perpendicularly to the gutters system in Aleppo pine and holm oak forests and under the rain exclusion device in the downy oak forest, by using a transect in blocks (7 columns × 6 lines), equidistantly one from each other (1 m distance between the 7 columns and 0.6 m between the 6 lines). Transects were E-W oriented. They were disposed in the ground floor after the removal of litter layer and fixed to the soil with galvanised nails to prevent movement by animals or wind. Litter layer was then replaced over the litterbags.

Litter bags were retrieved after 1 year (December 2015). Litterbags were placed in zipped plastic bags to prevent the loss of

Table 1. Main characteristics of the three forest sites selected for this study (for the study period: December 2014 to December 2015).

Forests	<i>Q. pubescens</i> Willd. Oak Observatory at the Observatoire de Haute Provence	<i>Q. ilex</i> L. Puéchabon	Mixed <i>P. halepensis</i> Mill. Font-Blanche
Location	43° 56' 115" N, 050 42' 642" E	43° 44' 29"N, 3° 35' 45"E	43° 14' 27" N, 5° 40' 45" E
Altitude a.s.l. (m)	650	270	425
MAT (Mean Annual Temperature) (°C)	12.25	13.71	14.01
MAP (Mean Annual Precipitation) (mm)	609	794	632
Soil type	pierric calcosol	rhodo-chromic luvisol	leptosol
Soil texture	clay	clay loam	clay
Soil pH	6.76	6.6	6.8
Surface rocks cover (%)	23	75	50
Dominant tree species	<i>Q. pubescens</i> Willd.	<i>Q. ilex</i> L.	mixed <i>P. halepensis</i> Mill./ <i>Q. ilex</i> L.
Other dominant plant species	<i>Acer monspessulanum</i> L. <i>Cotinus coggygia</i> Scop.	<i>Buxus sempervirens</i> L. <i>Phyllirea latifolia</i> L. <i>Pistacia terebinthus</i> L. <i>Juniperus oxycedrus</i> L.	<i>Quercus coccifera</i> L. <i>Phyllirea latifolia</i> L.
Tree density (stems/ ha)	3503	4500	3368
Forest structure	even-age (70 years)	even-age (74 years)	uneven-age (61 years)
Rain exclusion system dimensions (m ²)	300	140	625
Rain exclusion (%)	33	30	30

biological material. In the laboratory, the soil and other debris were removed from the litter bags. Then, they were freeze-dried (Lyovac GT2) for 72 hours and the remaining dry mass (%) was calculated. A 1 g of dry mass aliquote was collected, grounded to powder (Retsch® MM400, 30 Hz during 30s) and kept at -20°C for posteriori microbial DNA analysis.

Initial litter chemical characteristics

At the beginning of the experiment, 5 replicates from each leaf species collected from trees in control plots were grounded to powder (Retsch® MM400, 30 Hz during 30s). Carbon (C) and nitrogen (N) were determined by thermal combustion on a Flash EA 1112 series C/N elemental analyser (Thermo-Scientific, U.S.A). Phosphorus (P) and cations (Ca, K, Mg and Na) were extracted from 80 mg of ground litter sample with 8 ml of nitric acid and 2 ml of H₂O₂ at 175°C for 40 min using a microwave digestion system (Ethos One, Milestone SRL, Sorisole, Italy). After the mineralisation, each sample was adjusted to 50 ml with demineralised water. Phosphorus concentration was measured at 720 nm using a microplate reader (Victor, Perkin Elmer, Waltham, MA, USA). Cations were determined using an atom absorption spectrometer (AAS, iCE 3000 series, Thermo-Scientific, China). Leaf structural compounds (lignin, cellulose and hemicellulose) and water-soluble compounds (WSC) concentrations were determined according to the van Soest extraction protocol (Van Soest and Wine 1967), using a fiber analyser (Fibersac 24; Ankom®, Macedon, NJ, USA). Total phenolic concentrations were measured by colorimetry according to the method of Peñuelas et al. (1996) using gallic acid as standard. Aqueous extracts were made by dissolving 0.25 g of litter powder in 20 ml of 70% aqueous methanol solution with shaking for 1 hour and filtered (0.45 µm). A total of 25 ml of extracts obtained were then mixed with 0.25 ml of Folin-Ciocalteu reagent (Folin and Denis 1915), 0.5 ml of saturated aqueous Na₂CO₃ (to stabilize the colorimetric reaction) and 4 ml of distilled water. After 60 min, the reaction was completed and the concentration of phenolics was measured at 765 nm using a UV / Vis spectrophotometer (Thermoscientific, U.S.A.).

To determine the water holding capacity (WHC), twenty-one freshly abscised leaves (3 species x 5 replicates) were soaked in distilled water for 24 hours, drained and weighted (wet weight). The dry weight was measured after drying the samples for 48 hours at 60°C. Water holding capacity (in %) was calculated according to the formula: (wet weight/dry weight) × 100 (Santonja et al. 2015b). Specific leaf area (SLA) was determined by using the Image J software (<https://imagej.nih.gov/ij/>, MA, USA). Specific leaf area was calculated as the ratio between leaf area and leaf dry weight.

DNA extraction

Environmental DNA from leaf litter samples was extracted with the NucleoSpin® soil kit (Macherey-Nagel, Düren, Germany), following the standard protocol with a slight adjustment. In brief, approximately 250 mg of dry litter was added to the NucleoSpin Bead tubes. Lysis conditions were performed by using buffer SL1 (700 µl, vortex for 5 min, room temperature, 11 000 x g for 2 min), and buffer SL3 (150µl, vortex 5 min, 0–4°C, 14 000 g for 1 min). This procedure was repeated twice using the same sample to optimise the DNA extraction. The filtration of the supernatant was performed in the NucleoSpin Inhibitor Removal Column followed by 11 000 x g for 1 min. DNA binding was done in the Nucleo Spin Soil column after adding 250 µl of Buffer SB to the supernatant followed by 11 000 x g for 1 min. A series of successive DNA washing was performed: 500 µl buffer SB (11 000 x g, 3 sec), 550 µl buffer SW1 (11 000 x g, 3 sec), 700 µl SW2 (2 sec vortex, 11 000 x g, 3 sec). At each step, flow-through were discarded. Finally, DNA was eluted in 30 µl of Buffer SE followed by incubation at room temperature for 1 min and 11 000 x g for 30 seconds and kept at -20°C until further analysis. DNA quantification was performed in the ND-1000 spectrophotometer (Nanodrop Technology, Wilmington, DE).

Quantitative real-time PCR analysis

Abundances of total bacteria, actinobacteria and fungi were assessed using a quantitative real-time PCR analysis (qPCR)

Table 2. Initial leaf litter quality of each litter species: *Q. pubescens*, *Q. ilex* and *P. halepensis*, the values are mean \pm SE ($n = 5$) and the results of the one-way ANOVA to test differences between initial litter species are indicated (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ and different letters denote significant differences among species with $a > b > c$.)

	df	<i>Q. pubescens</i>			<i>Q. ilex</i>		<i>P. halepensis</i>		F-ratio	
Carbon	2	462.6 \pm 2.65	c		478.1 \pm 1.4	b	516.1 \pm 1.6	a	198.80	***
Nitrogen	2	6.4 \pm 0.2	b		9.6 \pm 0.2	a	5.4 \pm 0.1	c	146.30	***
Phosphorus	2	1.9 \pm 0.1	b		3.5 \pm 0.2	a	1.6 \pm 0.1	b	64.6	***
Calcium	2	32.9 \pm 0.8	a		25.1 \pm 0.6	b	18.4 \pm 0.2	c	168.1	***
Potassium	2	0.8 \pm 0.01	c		1.8 \pm 0.1	a	0.95 \pm 0.01	b	370.7	***
Magnesium	2	2.8 \pm 0.2	a		1.4 \pm 0.01	b	1.5 \pm 0.02	c	104.8	***
Sodium	2	0.04 \pm 0.00	b		0.11 \pm 0.00	a	0.11 \pm 0.00	a	149.6	***
Total phenolics	2	40.8 \pm 2.6	a		32.9 \pm 1.58	b	38.6 \pm 1.2	ab	4.7	*
Lignin	2	273.3 \pm 9.0	b		337.4 \pm 5.5	a	302.6 \pm 10.6	b	13.8	***
Cellulose	2	159.1 \pm 8.0	b		205 \pm 4.2	a	150.8 \pm 12.9	b	10.3	**
Hemicellulose	2	274.2 \pm 22.5	a		270.0 \pm 16.2	ab	206.6 \pm 10.4	b	4.7	*
Water-soluble compounds	2	293.4 \pm 12.1	b		187.6 \pm 10.2	c	340.0 \pm 7.1	a	60.9	***
Specific leaf area	2	133.1 \pm 0.4	a		51.5 \pm 0.9	c	83.5 \pm 2.2	b	690.7	***
Water holding capacity	2	146.9 \pm 1.0	a		137.2 \pm 1.0	b	113.9 \pm 1.03	c	285.5	***

method. Microbial abundance was quantified as copy number of a target gene. Partial 16S rRNA gene sequences were amplified from the total bacteria using primers COM1 (5'-CAG CAGCCGCGTAATAC-3') and 769R (5'-ATCCTGTTGMTMCCCV CRC-3', annealing temperature 59°C) (Dorn-In et al. 2015), and from actinobacteria using primers S-P-Acti-1154-a-S-19 (5'-GRD ACYGCCGGGTAACT-3' annealing temperature 59°C) and S-P-Acti-1339-a-A-18 (5'-TCWGCATTACTAGCGAC-3') (Pfeiffer et al. 2014). Partial 18S rRNA genes from fungi were amplified with primers FF390 (5'-CGATAACGAACGAGACCT-3') and FR1 (5'-A ICCATTCAATCGGTAIT-3', annealing temperature 50°C) (Vainio and Hantula 2000). All PCR reactions were done 15 μ l total volume containing 1 \times SYBR® Green Gotaq PCR Master Mix (Promega), primers at 600 nM for total bacteria and fungi, and 200 nM for actinobacteria, and 0.2–2 ng environmental DNA. The cycling parameters were: 10 min at 95°C followed by 45 cycles at 95°C (15 sec), 59°C (30 sec), 72°C (30 sec). Melting curves were recorded to ensure qPCR specificity, and gel electrophoresis analyses were performed to confirm the expected size of amplified products. All qPCR reactions were run in triplicate on StepOne Plus Real-Time PCR System (Applied Biosystems, Foster City, CA). To quantify the target gene copy numbers, standard dilution curves of known concentrations of previously cloned standards of the target genes were determined. Baseline and threshold calculations and data analysis were performed with the StepOne v. 2.2.2 software.

The primer pair recommended by Dorn-In et al. (2015) for amplification of bacterial 16S rRNA gene fragment from plant samples, Com1/769R (Rastogi et al. 2010; Fredriksson, Hermansson and Wilén 2013) was used for quantitative real-time PCR analysis of the total bacteria. The primer pair does not amplify 16 rRNA genes from chloroplasts, however, consequently also its coverage of the domain Bacteria is incomplete. In silico analysis using TestPrime 1.0 (Klindworth et al. 2013) against the Silva database, release 132 (www.arb-silva.de), showed 75% overall coverage of Bacteria with varying representation of the phyla, e.g. 88% Proteobacteria, 81.9% Firmicutes, 93.6% Bacteroidetes, but only 56.2% Actinobacteria, 20.5% Acidobacteria and 17.5% Verrucomicrobia. Therefore, the results can be used as a proxy of bacterial abundance when comparing between samples, but not to compare with qPCR quantification of individual bacterial phyla or lower taxonomic units using group-specific primers.

Statistical analysis

All statistical analyses were performed in R v.3.1.3 software (R Development Core Team 2017). One-way analysis of variance (ANOVA) was used to determine differences between the initial parameters of litter plant species and between forest sites for each litter species and microbial group. Three-way ANOVA was also used to test the effects of forest sites (S), litter species (L) and drought treatment (D) on litter mass loss and microbial abundance (total bacteria (B), actinobacteria (A), fungi (F) and the respective B/F and A/F ratios, followed by post-hoc Tukey tests. Microbial abundance was log-transformed to fit the assumption of normality and homoscedasticity.

RESULTS

Initial litter composition

Initial litter characteristics varied significantly among species (Table 2, One-way ANOVA, $P < 0.05$). *Quercus ilex* showed a significantly higher N (9.62 mg g⁻¹) and P (3.48 mg g⁻¹) contents than the two other species, particularly *P. halepensis* (5.36 mg g⁻¹ and 1.56 mg g⁻¹), which resulted in a lowest C/N and C/P ratio for *Q. ilex* and highest ratios for *P. halepensis* (Table 2). Regarding the cations, *Q. pubescens* showed significantly higher content of Ca (32.90 mg g⁻¹) and Mg (2.83 mg g⁻¹), and lower content of K (0.84 mg g⁻¹) and Na (0.04 mg g⁻¹) than the two other species. Concerning the structural compounds of the leaves, *Q. ilex* had higher lignin (337.38 mg g⁻¹) and cellulose content (204.99 mg g⁻¹) than the two other species. Finally, *Q. pubescens* showed higher total phenolics content compared to *Q. ilex*, and the highest specific leaf area (133.13 mg g⁻¹) and water holding capacity (146.90 mg g⁻¹). *Pinus halepensis* showed higher water-soluble compounds (340.02 mg g⁻¹) and lower water holding capacity (113.89 mg g⁻¹).

Litter mass loss

After 357 days of decomposition, litter mass loss was statistically different among sites and plant species with interaction effects (Table 3). On average, significantly higher mass loss occurred at *Q. pubescens* forest (36.94%) in comparison with *Q. ilex* forest (33.88%) and *P. halepensis* forest (33.80%). Regarding the litter, *P. halepensis* (36.81%) and *Q. ilex* (35.07%) lost significantly

Table 3. ANOVA table of F-Ratio and P-values for the effects of forest sites (S), litter species (L) and drought conditions (D) on the mass loss, microbial biomass (actinobacteria, bacteria and fungi) and actinobacteria/fungi (A/F) and bacteria/fungi (B/F) ratios.

	df	Mass loss		Actino (A)		Bacteria (B)		Fungi (F)		A/F		B/F	
		F	P	F	P	F	P	F	P	F	P	F	P
Sites (S)	2	8.81	0.0008	33.92	<0.0001	8.86	0.0003	30.38	<0.0001	6.11	0.0030	11.78	<0.0001
Litter species (L)	2	8.11	0.0022	16.31	<0.0001	4.58	0.0122	2.41	0.0945	53.55	<0.0001	11.34	<0.0001
Drought (D)	1	3.70	0.0571	0.09	0.7602	1.91	0.1688	6.20	0.0143	12.08	0.0007	11.70	0.2828
SxL	4	7.03	<0.0001	5.69	0.0003	20.56	<0.0001	21.01	<0.0001	7.67	<0.0001	11.49	<0.0001
SxD	2	0.43	0.6488	3.00	0.0539	7.25	0.0011	3.82	0.0251	0.18	0.8368	5.43	0.0056
LxD	2	0.14	0.8724	3.83	0.0246	1.36	0.2593	1.12	0.3313	3.34	0.0392	0.11	0.8973
SxLxD	4	0.10	0.9839	1.11	0.3540	2.72	0.0335	1.45	0.2229	2.12	0.0836	1.17	0.3293

higher mass than *Q. pubescens* (32.72%). Concerning interactions, *P. halepensis* lost most of its mass at *Q. pubescens* and *P. halepensis* forests compared to the *Q. ilex* forest (Fig. 1A). No significant effect of rain exclusion was found on the litter mass loss (Table 3).

Microbial abundance and structure

Abundances of actinobacteria, total bacteria and fungi differed significantly according to forest sites, when control and amplified drought conditions were combined (Table 3, Fig. 1). The highest abundances of all microbial groups were observed at *P. halepensis* forest and lowest at *Q. ilex* forest. However, it differed according to plant litter species producing a site x species interaction effect (Table 3; Fig. 1). Actinobacteria and total bacteria had the highest abundance in *Q. ilex* litter and lowest in *P. halepensis* at *Q. ilex* and *P. halepensis* forests, respectively (Fig. 1B and C). Roughly, higher abundance of fungi was observed in *P. halepensis* litter at *Q. pubescens* forest, *Q. ilex* litter at *Q. ilex* forest and *Q. pubescens* litter at *P. halepensis* forest (Fig. 1D). Regarding the actinobacteria to fungi (A/F) ratio, it was significantly higher in *Q. pubescens* and *Q. ilex* litter and lower in *P. halepensis* litter at *Q. pubescens* and *Q. ilex* forests (Fig. 1E). Total bacteria to fungi (B/F) ratio was significantly higher in *Q. ilex* litter and lowest in *P. halepensis* litter, at *Q. ilex* and *P. halepensis* forests (Fig. 1F).

Overall, there was a significant increase of fungal abundance and a decrease of A/F ratio under amplified drought (AD) in comparison with natural conditions (Table 3). No main effect of drought conditions was observed for actinobacteria, total bacteria and bacteria to fungi (B/F) ratio (Table 3). However, an interaction effect was observed between site and drought conditions for total bacteria, fungi and total bacteria to fungi ratio (Table 3; see Supplementary Table 1, Supporting Information). Thus, we observed a significant increase of total bacterial (Fig. 2A) and fungal abundance (Fig. 2B) under AD conditions at the *P. halepensis* forest, and a decrease of B/F ratio in *Q. pubescens* forest (Fig. 2C).

DISCUSSION

In this study, we evaluated how reduction of rainfall and change in plant species may affect belowground microbial community's abundance and structure and consequently the litter decomposition process in three dominant Mediterranean forests of southern France. Our findings are that (i) actinobacteria and total bacteria were more affected by changes of plant species than by the decrease of water availability; (ii) fungi were positively impacted by the reduction of rainfall and (iii) the future arrival of *P. halepensis* litter to *Q. pubescens* and *Q. ilex* forest and rainfall

reduction could alter the relative proportion of fungi and bacteria in the community hence having implication in microbial decomposers activity.

Effects of litter quality on microbial communities and mass loss

Our results showed the importance of leaf litter initial litter with respect to microbial composition and litter mass loss dynamics. We observed higher abundance of actinobacteria and total bacteria in oak litter than in pine. This is in line with previous studies, which showed a preference of bacteria for high quality litter (i.e. low C/N, C/P, lignin: P and lignin/N ratio) (Hodge, Robinson and Fitter 2000; De Boer et al. 2005; Romani et al. 2006). Both oaks *Q. pubescens* and *Q. ilex* are characterised by high N and P content, which are efficiently utilised by bacterial enzymes (Romani et al. 2006) and then are consistently considered key factors in litter decomposition and good predictors of litter mass loss (Moro and Domingo 2000; Garcia-Pausas, Casals and Romanya 2004; Güsewell and Gessner 2009). However, our results only partially corroborate with this by showing that *Q. ilex* and *P. halepensis* litter had higher mass loss than *Q. pubescens*. That is possibly due to higher initial content of total phenolics determined in *Q. pubescens* compared to *Q. ilex*, which are known to inhibit litter microbial communities and N mineralisation, during the early stage of decomposition (Lambers 1993; Souto, Chiapusio and Pellissier 2000; Ormeño et al. 2006; Chomel et al. 2014). This may explain the lower mass loss of its litter in comparison with *Q. ilex* and *P. halepensis*. The high mass loss of *P. halepensis* litter may be also explained by the leaching of high content of water-soluble compounds (Tripathi and Singh 1992; Kaushal et al. 2012) and high Na content (Gressel et al. 1995), which both are known to positive correlated litter decay. Gressel et al. (1995) observed that salt accumulation in *P. halepensis* needles enhanced their breakdown compared to *Q. coccifera* leaves in a typical Mediterranean mixed woodland.

Effects of litter-site interaction on litter decomposition processes

The influence of plant species on litter decomposition varied according to the forest sites as shown by strong litter-site interaction. At *Q. pubescens* forest, we observed the highest abundance of fungi associated with *P. halepensis* litter, associated to a reduction of the A/F ratio. This is consistent with the findings by Hodge et al. (2000) and Wardle et al. (2004) who showed that fungi are favoured by litter with high C/N ratio. In addition, fungi




Figure 1. Leaf litter mass loss (A) and associated leaf litter microbial decomposers abundances (actinobacteria (B), bacteria (C), fungi (D)), actinobacteria to fungi (E) and bacteria to fungi (F) ratios, for the three forests and litter species (control and amplified drought conditions were combined). Values are mean \pm SE (n = 14). Different letters denote significant differences between litter species within forest sites (e.g. both drought conditions combined), with a > b > c. DM = litter dry mass. Qp = *Q. pubescens*, Qi = *Q. ilex*, Ph = *P. halepensis*.

are rather considered specialists and are more equipped enzymatically to assimilate recalcitrant material than more generalist bacteria (Møller, Miller and Kjølner 1999; Romaní et al. 2006). The high abundance of total bacteria observed in *P. halepensis* litter may be explained by the synergistic relationship between bacteria and fungi because bacteria may benefit by the presence of fungi, which may provide them resources that bacteria were not able to acquire on their own (Romaní et al. 2006). At *Q. ilex* forest, microorganisms were more numerous with *Q. ilex* litter in comparison with the two other species, which suggests the high specificity of decomposers for decomposing the autochthonous litter in this forest (Ayres et al. 2009). However, this difference in microbial abundance did not lead to a significant difference in mass loss of *Q. ilex* litter in its own forest

compared to the two other forests. This may be due to the presence of decomposers able to use different sources of carbon. Decomposers can also develop preferences for some resources they are not used to (St John et al. 2011). Taking into account the three litter species, our results showed that *P. halepensis* forest had generally higher microbial abundance (actinobacteria, total bacteria and fungi) than other sites, possibly because this is the only mixed woodland out of the three studied forests and it was demonstrated that an increase of plant diversity promotes litter microbial growth (Chapman and Newman 2010; Santonja et al. 2017). Indeed, litter dissimilarity may decrease resource competition between functional groups and thus increase the abundance, activity, and hence efficiency of decomposers (Barbe et al. 2017).




Figure 2. Effects of the two drought conditions (ND = natural drought; AD = amplified drought) on the total bacterial (A), fungal (B) abundances and bacteria to fungi ratio (C). Barplots are mean \pm SE. Significant effects are indicated with respective symbols * for $P < 0.05$ and ** for $P < 0.01$. *Qp* = *Q. pubescens*, *Qi* = *Quercus ilex*, *Ph* = *P. halepensis*.

Effects of rainfall reduction on litter mass loss and microbial community

Contrary to our expectations, amplified drought conditions (AD) had no effect on litter mass loss after 1 year of decomposition. This may be explained by the extremely dry year during which the study was performed (Saunier et al. 2018). Actinobacteria were not directly affected by AD conditions, contrary to previous studies (Bouskill et al. 2013; Felsmann et al. 2015). Nevertheless, a reduction in the A/F ratio was observed, probably due to the positive effect of AD conditions on fungal abundance. The increase of fungal abundance under drier conditions was also observed in other Mediterranean forests (Hawkes et al. 2011; Sherman, Sternberg and Steinberger 2012; Allison et al. 2013; Alster et al. 2013). It is well recognised that fungi are more resistant to drought than bacteria (De Vries and Shade 2013; Guhr et al. 2015) and might get competitive advantage over bacteria during dry periods (Six 2012; Yuste et al. 2011). Moreover, the

expansion of fungal hyphae reported by some studies under drought periods (Allison et al. 2013; Zeglin et al. 2013) may result from the reduction of predatory pressure (i.e. grazers activity or number) under water stressed conditions (Davet 2004; Bapiri, Băăth and Rousk 2010; Barnard, Osborne and Firestone 2013). The observed decrease of the A/F ratio under AD conditions in *Q. pubescens* litter was possibly related to the persistence of phenolics due to the decrease of leaching processes under AD conditions, which delayed microbial colonisation by decomposers in that litter (Chomel et al. 2014). Contrary to fungi, total bacteria were not affected by changes in drought conditions. Bacterial biomass stability under drought was also observed by Hartmann et al. (2017), and it was explained by independence of bacteria on water film for dispersal (Mohammadipanah and Wink 2016). Nevertheless, those results varied according to forest site and litter type. For example, in the *P. halepensis* forest, an increase of total bacteria biomass was observed under AD conditions, which may be explained by (i) natural selection of the most drought-tolerant bacterial groups (Barnard, Osborne and Firestone 2013) and/or (ii) bacteria resilience to drought resulting from their predominating r-selected life strategy, which enables rapid recovery after a period of disturbance (i.e. drought) (Barnard, Osborne and Firestone 2013; Meisner, Rousk and Băăth 2015).

CONCLUSIONS

Our results showed that microbial litter decomposers respond differently to intensification of water stress conditions in the Mediterranean forests. Actinobacteria and total bacteria were more negatively affected by plant litter changes (*Q. ilex* and *Q. pubescens* higher than *P. halepensis*) than by the direct rainfall reduction. In contrary, fungal communities seemed to be more tolerant and even favored to water limitation. Also, the potential resilience of bacterial communities was proposed, while fungal response seemed to reflect a higher degree of adaptation to decreased water availability. Finally, the arrival of *P. halepensis* litter to a *Q. pubescens* and *Q. ilex* forest may change litter microbial communities, leading to a decrease of A/F and B/F ratios and promoting fungi and certain groups of bacteria, which are more capable of using complex carbon sources. Therefore, the intensification of drought conditions in Mediterranean forest may introduce a fungi dominated community over bacteria, by either the capability of degrading more recalcitrant litter or/and tolerance to future climate conditions.

SUPPLEMENTARY DATA

Supplementary data are available at FEMSEC online.

ACKNOWLEDGMENTS

We thank all the people that contributed in the set-up of the experiment and in the field campaigns especially to Sylvie Dupouyet, Jean-Philippe Orts and Jean-Marc Ourcival. We thank to Anne Haguénauer from Endoume Marine Station (Marseille, France) for her efficient orientation in the lab and whole team of CRI—Crop Research Institute (Prague, Czech Republic)—for their unconditional help and support. The three experimental sites were annually supported by the research infrastructure AnaEE-France (ANR-11-INBS-0001) and by Allenvi through the SOERE F-ORE-T. We acknowledge the se of the COOPERATE database (Reiter IM, Castagnoli G, Rotureau A 2015 "Cooperate database", <https://cooperate.obs-hp.fr/db>) run by the CNRS

FR3098 ECCOREV and OSU-Pytheas with support from the Labex OT-Med. We also thank the French Region PACA and Europe for the PhD grant attributed to Susana Pereira.

AUTHOR CONTRIBUTIONS

SP, MM and VB wrote the manuscript. VB and CF conceived and designed the experiments. SP, AB, AA, and PM performed the experiments. SP, JK, MM, and VB analysed the data.

FUNDING

This research was financially supported by the Agence Nationale pour la Recherche (ANR) with the project SecPriMe2 (ANR-12-BSV7-0016-01), the BioDivMeX Mistrals program and partially by the Ministry of Education, Youth and Sports of the Czech Republic, European Regional Development Fund-Project No. CZ.02.1.01/0.0/0.0/16.019/0000845 and grant No. LTC17075 and Ministry of Agriculture of the Czech Republic, institutional support MZE-RO0418.

This work is a contribution to Labex OT-Med (ANR-11-LABX-0061) and has received funding from Excellence Initiative of Aix-Marseille University—A*MIDEX, a French ‘Investissements d’Avenir’ programme.

Conflict of interest. None declared.

REFERENCES

- Acosta-Martinez V, Cotton J, Gardner T et al. Predominant bacterial and fungal assemblages in agricultural soils during a record drought/heat wave and linkages to enzyme activities of biogeochemical cycling. *Appl Soil Ecol* 2014;**84**:69–82.
- Allison SD, Lu Y, Weihe C et al. Microbial abundance and composition influence litter decomposition response to environmental change. *Ecology* 2013;**94**:714–25.
- Alster CJ, German DP, Lu Y et al. Microbial enzymatic responses to drought and to nitrogen addition in a southern California grassland. *Soil Biol Biochem* 2013;**64**:68–79.
- Anandan R, Dharumadurai D, Manogaran GP. An introduction to actinobacteria. In: *Actinobacteria - Basics and Biotechnological Applications*. Intech, 2016.
- Ayres E, Steltzer H, Berg S et al. Soil biota accelerate decomposition in high-elevation forests by specializing in the breakdown of litter produced by the plant species above them. *J Ecol* 2009;**97**:901–12.
- Bapiri A, Bååth E, Rousk J. Drying–rewetting cycles affect fungal and bacterial growth differently in an arable soil. *Microb ecol* 2010;**60**:419–28.
- Barbe L, Jung V, Prinzing A et al. Functionally dissimilar neighbors accelerate litter decomposition in two grass species. *New Phytologist* 2017;**214**:1092–102.
- Barbero M, Bonin G, Loisel R et al. Changes and disturbances of forest ecosystems caused by human activities in the western part of the mediterranean basin. *Vegetatio* 1990;**87**:151–73.
- Barnard RL, Osborne CA, Firestone MK. Responses of soil bacterial and fungal communities to extreme desiccation and rewetting. *The ISME journal* 2013;**7**:2229.
- Bertrand R, Lenoir J, Piedallu C et al. Changes in plant community composition lag behind climate warming in lowland forests. *Nature* 2011;**479**:517.
- Bouskill NJ, Lim HC, Borglin S et al. Pre-exposure to drought increases the resistance of tropical forest soil bacterial communities to extended drought. *The ISME journal* 2013;**7**:384.
- Chapman SK, Newman GS. Biodiversity at the plant–soil interface: microbial abundance and community structure respond to litter mixing. *Oecologia* 2010;**162**:763–9.
- Chomel M, Fernandez C, Bousquet-Mélou A et al. Secondary metabolites of *Pinus halepensis* alter decomposer organisms and litter decomposition during afforestation of abandoned agricultural zones. *J Ecol* 2014;**102**:411–24.
- Cleveland CC, Reed SC, Keller AB et al. Litter quality versus soil microbial community controls over decomposition: a quantitative analysis. *Oecologia* 2014;**174**:283–94.
- Davet P. *Microbial Ecology of Soil and Plant Growth*. Boca Raton: CRC Press, 2004.
- De Boer W, Folman LB, Summerbell RC et al. Living in a fungal world: impact of fungi on soil bacterial niche development. *FEMS Microbiol Rev* 2005;**29**:795–811.
- De Vries FT, Shade A. Controls on soil microbial community stability under climate change. *Front microbiol* 2013;**4**:265.
- Dorn-In S, Bassitta R, Schwaiger K et al. Specific amplification of bacterial DNA by optimized so-called universal bacterial primers in samples rich of plant DNA. *J microbiol meth* 2015;**113**:50–6.
- Felsmann K, Baudis M, Gimbel K et al. Soil bacterial community structure responses to precipitation reduction and forest management in forest ecosystems across Germany. *PLOS ONE* 2015;**10**:e0122539.
- Fierer N, Bradford MA, Jackson RB. Toward an ecological classification of soil bacteria. *Ecology* 2007;**88**:1354–64.
- Folin O, Denis WA. Colorimetric method for the determination of phenols (and phenol derivatives) in urine. *J Biol Chem* 1915;**22.2**:305–8.
- Fortunel C, Garnier E, Joffre R et al. Leaf traits capture the effects of land use changes and climate on litter decomposability of grasslands across Europe. *Ecology* 2009;**90**:598–611.
- Fredriksson NJ, Hermansson M, Wilén B-M. The choice of PCR primers has great impact on assessments of bacterial community diversity and dynamics in a wastewater treatment plant. *PLOS ONE* 2013;**8**:e76431.
- Garcia-Pausas J, Casals P, Romanya J. Litter decomposition and faunal activity in Mediterranean forest soils: effects of N content and the moss layer. *Soil Biol Biochem* 2004;**36**:989–97.
- Garcia-Pausas J, Paterson E. Microbial community abundance and structure are determinants of soil organic matter mineralisation in the presence of labile carbon. *Soil Biol Biochem* 2011;**43**:1705–13.
- Gessner MO, Swan CM, Dang CK et al. Diversity meets decomposition. *Trends ecol evol* 2010;**25**:372–80.
- Gholz HL, Wedin DA, Smitherman SM et al. Long-term dynamics of pine and hardwood litter in contrasting environments: toward a global model of decomposition. *Glob Change Biol* 2000;**6**:751–65.
- Giorgi F, Lionello P. Climate change projections for the Mediterranean region. *Global Planet Change* 2008;**63**:90–104.
- Gressel N, Inbar Y, Singer A et al. Chemical and spectroscopic properties of leaf litter and decomposed organic matter in the Carmel Range, Israel. *Soil Biol Biochem* 1995;**27**:23–31.
- Guhr A, Borken W, Spohn M et al. Redistribution of soil water by a saprotrophic fungus enhances carbon mineralization. *P Natl A Sci* 2015;**112**:14647–51.
- Guiot J, Cramer W. Climate change: the 2015 paris agreement thresholds and Mediterranean basin ecosystems. *Science* 2016;**354**:465–8.
- Güsewell S, Gessner MO. N:P ratios influence litter decomposition and colonization by fungi and bacteria in microcosms. *Funct Ecol* 2009;**23**:211–9.

- Hartmann M, Brunner I, Hagedorn F et al. A decade of irrigation transforms the soil microbiome of a semi-arid pine forest. *Mol ecol* 2017;**26**:1190–206.
- Hawkes CV, Kivlin SN, Rocca JD et al. Fungal community responses to precipitation. *Glob Change Biol* 2011;**17**: 1637–45.
- Hertig E, Trambly Y. Regional downscaling of Mediterranean droughts under past and future climatic conditions. *Global Planet Change* 2017;**151**:36–48.
- Hodge A, Robinson D, Fitter A. Are microorganisms more effective than plants at competing for nitrogen? *Trends plant sci* 2000;**5**:304–8.
- Hättenschwiler S, Jørgensen HB. Carbon quality rather than stoichiometry controls litter decomposition in a tropical rain forest. *J Ecol* 2010;**98**:754–63.
- IPCC. *Climate Change 2013 - The Physical Science Basis*. Cambridge: Cambridge University Press, 2014.
- Kaushal R, Verma KS, Chaturvedi OP et al. Leaf litter decomposition and nutrient dynamics in four multipurpose tree species. *Range Manag Agrofor* 2012;**33**:20–7.
- Klindworth A, Pruesse E, Schweer T et al. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic acids res* 2013;**41**:e1–e1.
- Lambers H. Rising CO₂, secondary plant metabolism, plant-herbivore interactions and litter decomposition. In: *CO₂ and Biosphere*. Dordrecht: Springer, 1993, 263–71.
- Lewin GR, Carlos C, Chevrette MG et al. Evolution and ecology of Actinobacteria and their bioenergy applications. *Annu rev microbiol* 2016;**70**:235–54.
- Limousin JM, Rambal S, Ourcival JM et al. Modelling rainfall interception in a Mediterranean *Quercus ilex* ecosystem: lesson from a throughfall exclusion experiment. *J Hydrol* 2008;**357**:57–66.
- Lunghini D, Granito VM, Di Lonardo DP et al. Fungal diversity of saprotrophic litter fungi in a Mediterranean maquis environment. *Mycologia* 2013;**105**:1499–515.
- Manzoni S, Schimel JP, Porporato A. Responses of soil microbial communities to water stress: results from a meta-analysis. *Ecology* 2012;**93**:930–8.
- McCarthy AJ, Williams ST. Actinomycetes as agents of biodegradation in the environment: a review. *Gene* 1992;**115**:189–92.
- Meisner A, Rousk J, Bååth E. Prolonged drought changes the bacterial growth response to rewetting. *Soil Biol Biochem* 2015;**88**:314–22.
- Mohammadipanah F, Wink J. Actinobacteria from arid and desert habitats: diversity and biological activity. *Front microbiol* 2016;**6**:1541.
- Morales P, Hickler T, Rowell DP et al. Changes in European ecosystem productivity and carbon balance driven by regional climate model output. *Global Change Biol* 2007;**13**:108–22.
- Moro MJ, Domingo F. Litter decomposition in four woody species in a Mediterranean climate: weight loss, N and P dynamics. *Ann Bot* 2000;**86**:1065–71.
- Møller J, Miller M, Kjølner A. Fungal-bacterial interaction on beech leaves: influence on decomposition and dissolved organic carbon quality. *Soil Biol Biochem* 1999;**31**:367–74.
- Ogaya R, Peñuelas J. Tree growth, mortality, and above-ground biomass accumulation in a holm oak forest under a five-year experimental field drought. *Plant Ecol* 2007;**189**:291–9.
- Okoro CK, Brown R, Jones AL et al. Diversity of culturable actinomycetes in hyper-arid soils of the Atacama Desert, Chile. *Anton van Lee* 2009;**95**:121–33.
- Ormeño E, Baldy V, Ballini C et al. Effects of environmental factors and leaf chemistry on leaf litter colonization by fungi in a Mediterranean shrubland. *Pedobiologia* 2006;**50**: 1–10.
- Peñuelas J, Estiarte M, Kimball BA et al. Variety of responses of plant phenolic concentration to CO₂ enrichment. *J Exp Bot* 1996;**47**:1463–7.
- Peñuelas J, Rico L, Ogaya R et al. Summer season and long-term drought increase the richness of bacteria and fungi in the foliar phyllosphere of *Quercus ilex* in a mixed Mediterranean forest. *Plant Biol* 2012;**14**:565–75.
- Pfeiffer S, Pastar M, Mitter B et al. Improved group-specific primers based on the full SILVA 16S rRNA gene reference database. *Environ Microbiol* 2014;**16**:2389–407.
- Rastogi G, Tech JJ, Coaker GL et al. A PCR-based toolbox for the culture-independent quantification of total bacterial abundances in plant environments. *J Microbiol Meth* 2010;**83**:127–32.
- R Core Team. R: A language and environment for statistical computing. 2017. <https://www.R-project.org/>
- Romaní AM, Fischer H, Mille-Lindblom C et al. Interactions of bacteria and fungi on decomposing litter: differential extracellular enzyme activities. *Ecology* 2006;**87**:2559–69.
- Sagova-Mareckova M, Omelka M, Cermak L et al. Microbial communities show parallels at sites with distinct litter and soil characteristics. *Appl Environ Microbiol* 2011;**77**:7560–7.
- Santonja M, Baldy V, Fernandez C et al. Potential shift in plant communities with climate change: outcome on litter decomposition and nutrient release in a Mediterranean oak forest. *Ecosystems* 2015b;**18**:1253–68.
- Santonja M, Fernandez C, Gauquelin T et al. Climate change effects on litter decomposition: intensive drought leads to a strong decrease of litter mixture interactions. *Plant and Soil* 2015a;**393**:69–8.
- Santonja M, Rancon A, Fromin N et al. Plant litter diversity increases microbial abundance, fungal diversity, and carbon and nitrogen cycling in a Mediterranean shrubland. *Soil Biol Biochem* 2017;**111**:124–34.
- Saunier A, Ormeño E, Havaux M et al. Resistance of native oak to recurrent drought conditions simulating predicted climatic changes in the Mediterranean region. *Plant Cell Environ* 2018;**41**:2299–312.
- Schimel J, Balser TC, Wallenstein M. Microbial stress-response physiology and its implications for ecosystem function. *Ecology* 2007;**88**:1386–94.
- Sherman C, Sternberg M, Steinberger Y. Effects of climate change on soil respiration and carbon processing in Mediterranean and semi-arid regions: an experimental approach. *Eur J Soil Biol* 2012;**52**:48–58.
- Six J. Fungal friends against drought. *Nat Clim Change* 2012;**2**:234–5.
- Souto XC, Chiapusio G, Pellissier F. Relationships between phenolics and soil microorganisms in spruce forests: significance for natural regeneration. *J Chem Ecol*. 2000;**26**: 2025–34.
- St. John MG, Orwin KH, Dickie IA. No “home” versus “away” effects of de-composition found in a grassland-forest reciprocal litter transplant study. *Soil Biol Biochem* 2011;**43**: 1482–9.
- Strickland MS, Lauber C, Fierer N et al. Testing the functional significance of microbial community composition. *Ecology* 2009;**90**:441–51.
- Swift MJ, Heal OW, Anderson JM et al. *Decomposition in Terrestrial Ecosystems*. Berkeley: University of California Press, 1979.

- Tripathi SK, Singh KP. Nutrient immobilization and release patterns during plant decomposition in a dry tropical bamboo savanna, India. *Biol Fert Soils* 1992;**14**:191–9.
- Vainio EJ, Hantula J. Direct analysis of wood-inhabiting fungi using denaturing gradient gel electrophoresis of amplified ribosomal DNA. *Mycol Res* 2000;**104**:927–36.
- Van Soest PJ, Wine RH. Use of detergents in the analysis of fibrous feeds: IV. Determination of plant cell-wall constituents. *J of official analytical chemists* 1967;**50**: 50–5.
- Wardle DA, Bardgett RD, Klironomos JN et al. Ecological linkages between aboveground and belowground biota. *Science* 2004;**304**:1629–33.
- Wardle DA, Van der Putten WH. Biodiversity, ecosystem functioning and above-ground-below-ground linkages. *Biodiversity and Ecosystem Functioning: Synthesis and Perspectives* 2002.
- Xu X, Thornton PE, Post WM. A global analysis of soil microbial biomass carbon, nitrogen and phosphorus in terrestrial ecosystems. *Global Ecol Biogeogr* 2013;**22**:737–49.
- Yuste JC, Peñuelas J, Estiarte M et al. Drought-resistant fungi control soil organic matter decomposition and its response to temperature. *Global Change Biol* 2011;**17**:1475–86.
- Zeglin LH, Bottomley PJ, Jumpponen A et al. Altered precipitation regime affects the function and composition of soil microbial communities on multiple time scales. *Ecology* 2013;**94**:2334–45.

Supplementary information for

Litter traits and rainfall reduction alter microbial litter decomposers: the evidence from three Mediterranean forests

S. Pereira¹, A. Burešová^{2,3,4}, J. Kopecký⁵, P. Mádrová⁵, A. Aupic-Samain¹,
C. Fernandez¹, V. Baldy¹, and M. Sagova-Mareckova⁵

¹Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE UMR 7263, Marseille, France,

²Department of Ecology, Faculty of Science, Charles University in Prague, Vinicna 7, Prague, Czech Republic,

³Ecologie Microbienne, Université Claude Bernard Lyon 1, UMR CNRS 5557, INRA 1418, Villeurbanne, France,

⁴Laboratory for Diagnostics and Epidemiology of Microorganisms, Crop Research Institute, Drnovska 507, CZ-16106 Prague 6, Ruzyne, Czech Republic and


⁵Czech University of Life Sciences, Faculty of Agrobiology, Food and Natural Resources, Department of Microbiology, Nutrition and Dietetics, Kamýcká 129, 165 00 Praha 6 - Suchbátka, Czech Republic

Supplementary Table 1 - Summary table of microbial abundance (expressed as 10^{10} gene copy number g^{-1} litter dry mass) for each litter species at each forest site and drought condition (ND = Natural drought; AD = amplified drought). One-way ANOVAs were performed for differences among litter species per drought condition and forest site. F-Ratio are indicated and P-values with the respective symbols * for $P < 0.05$, ** for $P < 0.01$, and *** for $P < 0.001$. Different letters denote significant differences among litter species with $a > b$. T-tests was performed to test the effect of drought conditions for each species at each forest site. T-test are indicated and significant P-values ($P < 0.05$) are denoted in bold.

			Forest sites			F	P
			<i>Q. pubescens</i>	<i>Q. ilex</i>	<i>P. halepensis</i>		
<i>Quercus pubescens</i>	Actinobacteria	ND	49.55 ± 17.89 ab	15.52 ± 4.23 b	74.75 ± 17.67 a	6.05	**
		AD	17.27 ± 5.13 b	7.15 ± 2.94 b	85.62 ± 14.22 a	19.95	***
		<i>t</i> -test <i>p</i> -value	-2.1757 ns	-1.9232 ns	0.93659 ns		
	Bacteria	ND	11.86 ± 3.10 a	10.41 ± 2.76 a	24.58 ± 4.25 a	33.51	ns
		AD	8.12 ± 1.92 b	6.73 ± 3.94 b	53.70 ± 8.17 a	19.54	***
		<i>t</i> -test <i>p</i> -value	-0.65657 ns	-1.3221 ns	2.7689 0.0176		
	Fungi	ND	0.30 ± 0.09 b	0.26 ± 0.05 b	1.08 ± 0.17 a	13.91	***
		AD	0.33 ± 0.13 b	0.16 ± 0.05 b	2.26 ± 0.23 a	33.35	***
		<i>t</i> -test <i>p</i> -value	0.089708 ns	-1.559 ns	4.1807 0.0018		
	Ratio A/F	ND	171.78 ± 40.21 a	58.32 ± 10.71 b	71.30 ± 18.25 b	5.60	*
		AD	62.77 ± 14.26 a	38.00 ± 5.39 a	37.91 ± 5.47 a	2.34	ns
		<i>t</i> -test <i>p</i> -value	-3.391 0.0054	-1.6586 ns	-1.5571 ns		
	Ratio B/F	ND	24.48 ± 4.57 a	48.18 ± 10.46 a	35.92 ± 6.42 a	1.91	ns
		AD	32.56 ± 5.13 a	29.48 ± 7.56 a	23.59 ± 2.65 a	0.67	ns
		<i>t</i> -test <i>p</i> -value	-0.91979 ns	-0.83125 ns	0.16864 ns		
<i>Quercus ilex</i>	Actinobacteria	ND	63.26 ± 18.05 a	30.64 ± 8.66 a	28.44 ± 6.39 a	0.94	ns
		AD	39.9 ± 10.89 ab	19.45 ± 5.14 b	50.03 ± 8.88 a	5.33	*
		<i>t</i> -test <i>p</i> -value	-0.47229 ns	-0.63645 ns	1.8596 ns		
	Bacteria	ND	18.66 ± 12.96 a	35.92 ± 9.87 a	17.00 ± 2.31 a	0.01	ns
		AD	7.27 ± 1.92 b	24.97 ± 5.84 b	55.01 ± 11.58 a	15.89	***
		<i>t</i> -test <i>p</i> -value	-0.56905 ns	-0.44153 ns	3.4846 0.0061		
	Fungi	ND	0.47 ± 0.14 a	0.47 ± 0.12 a	0.39 ± 0.06 a	29.61	ns
		AD	0.54 ± 0.09 a	0.41 ± 0.14 a	0.90 ± 0.17 a	4.00	*
		<i>t</i> -test <i>p</i> -value	0.9151 ns	-0.34763 ns	3.2994 0.0073		
	Ratio A/F	ND	129.45 ± 22.32 a	63.52 ± 9.21 b	72.76 ± 11.76 b	38.21	*
		AD	70.58 ± 12.05 a	53.67 ± 5.53 a	59.00 ± 6.49 a	0.83	ns
		<i>t</i> -test <i>p</i> -value	-2.2672 0.0445	-0.78997 ns	-0.45703 ns		
	Ratio B/F	ND	29.55 ± 10.32 b	77.12 ± 15.33 a	46.96 ± 7.00 ab	7.28	**
		AD	12.96 ± 2.31 b	73.53 ± 14.96 a	64.72 ± 12.05 a	27.71	***
		<i>t</i> -test <i>p</i> -value	-1.9048 ns	-0.19647 ns	1.2028 ns		
<i>Pinus halepensis</i>	Actinobacteria	ND	17.53 ± 5.56 a	5.57 ± 1.79 b	17.98 ± 4.8 a	5.73	*
		AD	27.18 ± 2.79 a	6.58 ± 1.44 b	23.74 ± 7.53 a	14.48	***
		<i>t</i> -test <i>p</i> -value	1.5554 ns	0.43757 ns	0.64576 ns		
	Bacteria	ND	32.19 ± 4.07 a	6.59 ± 1.86 b	5.77 ± 0.89 b	29.41	***
		AD	19.02 ± 1.91 a	17.52 ± 3.94 a	10.85 ± 0.82 a	29.38	ns
		<i>t</i> -test <i>p</i> -value	-2.88 0.0185	2.7371 0.0228	2.5945 0.0291		
	Fungi	ND	1.01 ± 0.1 a	0.23 ± 0.05 b	0.41 ± 0.07 b	15.48	***
		AD	1.47 ± 0.41 a	0.41 ± 0.07 b	0.59 ± 0.08 b	9.75	**
		<i>t</i> -test <i>p</i> -value	0.97788 ns	2.1384 ns	1.7302 ns		
	Ratio A/F	ND	17.56 ± 5.16 b	22.63 ± 1.88 ab	45.56 ± 9.66 a	7.50	**
		AD	24.90 ± 5.37 a	16.79 ± 3.51 a	43.48 ± 13.61 a	30.13	ns
		<i>t</i> -test <i>p</i> -value	0.98631 ns	-1.4713 ns	-0.1249 ns		
	Ratio B/F	ND	32.42 ± 3.27 a	28.16 ± 3.91 ab	19.43 ± 6.12 b	4.24	*
		AD	18.67 ± 5.31 b	44.85 ± 7.37 a	19.67 ± 3.29 b	6.13	**
		<i>t</i> -test <i>p</i> -value	-2.2045 ns	2.0002 ns	0.034746 ns		

40

41



42

43 **Supplementary Figure 1** – Ombrothermic diagrams at a) O₃HP, b) Puéchabon and c) Font-

44 Blanche study sites between December 2014 to December 2015. Bar represents the mean

45 monthly precipitation (mm) in black (ND= natural drought) and in white (AD = amplified

46 drought). The curve represents the mean monthly temperature (°C). At each forest site, drought

period is indicated by the horizontal arrow and the percentage represents the proportion of excluded rain compared to the natural drought plot.

a)




b)



c)



Supplementary Figure 2 – Rain exclusion devices in the three forest studies (a) *Q. pubescens* forest at O₃HP, (b) *Quercus ilex* at Puéchabon and (c) *P. halepensis* at Font-Blanche.



Supplementary Figure 3 - Litterbags experimental design: 1) leaf litter of the three tree species (*Q. pubescens*, *Q. ilex* and *P. halepensis*) coming from natural drought (ND) were redistributed and placed 2) in the three different forests (O₃HP, Puéchabon and Font-Blanche, respectively), under natural drought (ND) and amplified drought (AD) conditions .

Appendix

Curriculum vitae Mgr. Ing. Andrea Faitová

Date of birth: 28th May, 1991

Contact: Něvská 583, Praha 9, 196 00, Czech Republic

Phone: +420 775 931 831

E-mail: Andrea.Buresova@seznam.cz

Current positions

- **Young researcher** at research team Epidemiology and Ecology of Microorganisms
Crop Research Institute in Prague
Czech Republic
- **Ph.D. student** (Cotutelle)
Charles University in Prague and University Claude Bernard Lyon 1

Education

September 2015 - February 2021: PhD. in Microbial Ecology (Cotutelle)

Thesis title: Actinobacteria communities in natural and anthropogenic environments

Department of Ecology, Faculty of Science, Charles University in Prague (CZ)

Supervisor: RNDr. Marketa Sagova-Mareckova, PhD.

Microbial Ecology, doctoral school Evolution, Ecosystèmes, Microbiologie et Modélisation, University Claude Bernard Lyon 1 (FR)

Supervisor: Prof. Veronica Rodriguez-Nava

September 2013 – May 2016: M.Sc. in Waste Use and Management

Czech University of Life Sciences in Prague

Thesis Title: The effect of sulfur compounds on mercury mobility and activity of microorganisms in soil

Supervisor: prof. Ing. Jiřina Száková, CSc.

September 2013 – May 2015: M.Sc. in Ecology, Department of Ecology

Faculty of Science, Charles University in Prague

Thesis Title: Composition and activity of microbial community during decomposition of plant litter on two contrasting localities

Supervisor: RNDr. Marketa Sagova-Mareckova, PhD.

September 2010 – September 2013: B.Sc. in Soil science in Ecology and Environmental protection

Faculty of Science, Charles University in Prague

Thesis Title: Bioturbation and its role in soil formation

Supervisor: Prof. Ing. Mgr. Jan Frouz, CSc

Scholarships

- **May 2017** - French Government scholarship for Cotutelle studies
- **April 2016** – French Government scholarship for scientific internship
- **December 2014** – Merit scholarship at Charles University in Prague

Publications

2019

Buresova A., Kopecky J., Hrdinkova V., Kamenik Z., Omelka M. and Sagova-Mareckova M., (2019), Succession of Microbial Decomposers Is Determined by Litter Type, but Site Conditions Drive Decomposition Rates, *Applied and Environmental Microbiology*, 85 (24): e01760-19

Pereira S., **Buresova A.**, Kopecky J., Madrova P., Aupic-Samain A., Fernandez C., Baldy V. and Sagova-Mareckova M., (2019), Litter traits and rainfall reduction alter microbial litter decomposers: the evidence from three Mediterranean forests, *FEMS Microbiology Ecology*, 95 (12): fiz168

2016

Sagova-Mareckova M, Zádorová T, Penížek V, Omelka M, Tejnecký V, Pruchova P, Chuman T, Drábek O, **Burešová A**, Vaněk A, Kopecky J, (2016), The structure of microbial communities along two vertical profiles of a deep colluvial soil, *Soil Biology and Biochemistry*, 101, 65-73

Szákóvá J, **Burešová A**, Praus L, García-Sánchez M, Holečková Z, Gabriel J, Sysalová J, Komárek J, Grohová J, Tlustoš P, (2016), The response of mercury (Hg) transformation in soil on sulfur and sulfur-rich biowaste application, *Environmental Earth Sciences*, 75:584

2013

Frouz J, Jílková V, Cajthaml T, Pižl V, Tajovský K, Háněl K, **Burešová A**, Šimáčková H, Kolaříková K, Franklin J, Nawrot J, Groninger JW, Stahl PD. 2013. Soil biota in post-mining sites along a climatic gradient in the USA: Simple communities in shortgrass prairie recover faster than complex communities in tallgrass prairie and forest, *Soil Biology and Biochemistry*, 57, 212 – 225.

Contributions at scientific meetings and conferences

December 2020: Poster presentation at international conference ACTINO 2020 (online conference)

June 2019: Poster presentation at international conference FEMS 2019 in Glasgow, Scotland

May 2017: Oral presentation at ISME Meeting of Young Soil Microbial Ecologists, Prague

December 2015: Poster presentation at international conference Ecology of Soil Microorganisms, Prague, Czech Republic

Scientific internship

October 2017 - December 2017: Ben-Gurion University of the Negev, Israel

Research topic: Bacterial response to altered precipitation regimes in desert soil

June 2016 – July 2016: Microbial Ecology, Université Claude Bernard Lyon 1, France

Research topic: Phylogenetic diversity of *Actinobacteria* colonizing Paleolithic wall paintings of Lascaux cave, France

Main research interests

- Ecology of terrestrial *Actinobacteria* (soil, caves)
- Microbial decomposition of organic matter and extracellular enzyme production in soil

Methods experience

DNA, RNA extraction, PCR, qPCR, ddPCR, cloning, sequencing, primer design

Extracellular hydrolytic and oxidative enzyme activity assessment in soil and plant substrate

Working with microbial cultures, *Actinobacteria* isolation

Fumigation and microbial biomass carbon extraction

Extraction of low molecular weight substances for HPLC analysis

Bioinformatic analysis of microbial community: next-generation sequencing, taxonomic database design, molecular markers evaluation, phylogenetic analysis (Mothur, UPARSE, MEGA, Galaxy)

Statistical data analysis (STATISTICA, Past, Canoco, R)

Other qualifications

Languages: English (B2)

French (B1)

Driving license B category