



**HAL**  
open science

# Etude d'un problème de contrôle optimal convexe mal posé

Frédéric Abergel

► **To cite this version:**

Frédéric Abergel. Etude d'un problème de contrôle optimal convexe mal posé. Mathématiques [math]. Université Paris-Sud (1970-2019), 1986. Français. NNT: . tel-04140685

**HAL Id: tel-04140685**

**<https://theses.hal.science/tel-04140685v1>**

Submitted on 26 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

63448

ORSAY  
n° d'ordre :

UNIVERSITE DE PARIS-SUD  
CENTRE D'ORSAY

# THESE

présentée

Pour obtenir

Le ..... TITRE ..... de DOCTEUR ..... EN MATHEMATIQUES

PAR

Frédéric ABERGEL

---



SUJET : Etude d'un problème de contrôle optimal convexe mal posé.

soutenue le ..... 9 Juin 1986 ..... devant la Commission d'examen

MM. ..... R. TEMAM ..... Président

..... C. BARDOS .....

..... F. MIGNOT .....

.....



## Remerciements

J'aimerais avant tout exprimer à Roger Temam ma profonde gratitude pour la possibilité qu'il m'a offerte de réaliser cette thèse sous sa direction, ainsi que pour l'aide précieuse qu'il m'a plus d'une fois apportée au cours de ce travail.

Il m'est de même très agréable de remercier Claude Bardos et Fulbert Mignot, qui m'ont fait l'honneur de s'intéresser à mon travail, et ont bien voulu participer au jury de soutenance.

Je tiens aussi à remercier Jean-Paul Bourguignon et Yves Meyer pour les fructueuses indications qu'ils m'ont données.

Une des dernières étapes de la réalisation de cette thèse en a été la transformation en un article lisible par tous, grâce au travail de Mme Thaler. Qu'elle en soit remerciée ici.



Frédéric ABERGEL

RESUME DE THESE

Soit  $\Omega$  un ouvert borné de l'espace euclidien  $\mathbb{R}^N$ ,  $N \geq 3$ , dont la frontière  $\Gamma$  est la réunion  $\Gamma_0 \cup \Gamma_1$  de deux variétés de classe  $C^\infty$ , compactes connexes et disjointes.  $\Omega$  est supposé être localement d'un seul côté de  $\Gamma$ . Soit  $\alpha, \alpha', \beta, \beta'$  quatre réels strictement positifs. Notre propos est d'étudier le problème de contrôle optimal correspondant aux données suivantes :

\* L'ensemble des contrôles admissibles  $\mathcal{U}_{ad}$  est

$$\mathcal{U}_{ad} = \{(v_0, v_1) \in (L^2(\Gamma_0))^2, v_0(x) \text{ (resp. } v_1(x)) \in [-\alpha, \beta] \text{ (resp. } [-\alpha', \beta']) \text{ p.p. } x \in \Gamma_0\}.$$

\* L'état  $z$  du système vérifie les conditions :

$$(c_0) \quad z \in L^2(\Omega) ; \Delta z = 0 ; z = v_0 \text{ sur } \Gamma_0 ; \frac{\partial z}{\partial \nu} = v_1 \text{ sur } \Gamma_0.$$

\* La fonction coût, que l'on cherche à minimiser, est donnée par

$$j(z, v_0, v_1) = \frac{1}{2} \|z - z_d\|_{L^2(\Omega)}^2 + \frac{1}{2} \|v_0\|_{L^2(\Gamma_0)}^2 + \frac{1}{2} \|v_1\|_{L^2(\Gamma_0)}^2$$

( $z_d$  étant un élément fixe de  $L^2(\Omega)$ ).

On voit par des méthodes classiques que ce problème de contrôle admet une solution unique. Toutefois ce problème est mal posé, dans la mesure où l'état adjoint ne peut être déterminé par les méthodes usuelles ce qui rend difficile l'écriture des conditions d'optimalité. Nous sommes amenés ici à étudier directement ce problème de contrôle optimal comme un problème d'optimisation convexe dans l'espace

$$W = \{z \in L^2(\Omega), \Delta z \in L^2(\Omega), (z, \frac{\partial z}{\partial \nu}) \in (L^2(\Gamma_0))^2\}$$

dont la formulation est évidemment la suivante :

$$(\mathcal{P}) \quad \text{Inf} \left\{ \frac{1}{2} \|z - z_d\|_{L^2(\Omega)}^2 + \frac{1}{2} \|z\|_{L^2(\Gamma_0)}^2 + \frac{1}{2} \left\| \frac{\partial z}{\partial \nu} \right\|_{L^2(\Gamma_0)}^2 \right\}$$

pour  $z \in W$ ,  $\Delta z = 0$ ,  $z(x)$  (resp.  $\frac{\partial z}{\partial \nu}(x)$ )  $\in [-\alpha, \beta]$  (resp.  $[-\alpha', \beta']$ ) pour presque tout  $x \in \Gamma_0$ .

La détermination de l'état adjoint de la solution de ce problème est équivalente à la recherche de solutions du problème dual de  $(\mathcal{P})$ , soit  $(\mathcal{P}^*)$  :

$$(\mathcal{P}^*) \quad \text{Sup} \left\{ -\frac{1}{2} \|\Delta q\|_{L^2(\Omega)}^2 + \int_{\Omega} \Delta q \cdot z_d \, dx - \int_{\Gamma_0} h_{\alpha, \beta} \left( -\frac{\partial q}{\partial \nu} \right) d\Gamma - \int_{\Gamma_0} h_{\alpha', \beta'}(q) d\Gamma \right\}$$

pour  $q \in W$ ,  $q = \frac{\partial q}{\partial \nu} = 0$  sur  $\Gamma_1$ , et  $h_{\alpha, \beta}$  définie par :

$$h_{\alpha, \beta}(s) \begin{cases} = \frac{s^2}{2}, & s \in [-\alpha, \beta] \\ = \beta \left( s - \frac{\beta}{2} \right), & s \geq \beta \\ = -\alpha \left( s + \frac{\alpha}{2} \right), & s \leq -\alpha. \end{cases}$$

L'étude des problèmes  $(\mathcal{P})$  et  $(\mathcal{P}^*)$  : existence de solutions, système des conditions d'optimalité, va donc constituer la majeure partie de notre travail.

Dans la Section I, nous utilisons les résultats classiques d'optimisation convexe, dans le cas coercif sci, pour établir l'existence et l'unicité de la solution  $z_0$  de  $(\mathcal{P})$ , ainsi que l'égalité des extremas de  $(\mathcal{P})$  et  $(\mathcal{P}^*)$ .

Etant donné la définition de  $h_{\alpha, \beta}$ , nous ne savons pas si  $(\mathcal{P}^*)$  admet des solutions dans  $W$ , par manque de coercivité. L'étude des suites maximisantes de  $(\mathcal{P}^*)$  nous montre que ces dernières vérifient les conditions suivantes :

$$(c_1) \quad \Delta q_n \text{ bornée dans } L^2(\Omega),$$

$$(c_2) \quad (q_n, \frac{\partial q_n}{\partial \nu}) \text{ bornée dans } M_1(\Gamma_0) \times M_1(\Gamma_0),$$

où  $M_1(\Gamma_0)$  désigne l'ensemble des mesures bornées sur  $\Gamma_0$ . Nous souhaitons donc étendre l'ensemble des éléments admissibles pour  $(\mathcal{P}^*)$  à un espace dans lequel les suites maximisantes de  $(\mathcal{P}^*)$  seront bornées.

Avant de définir précisément l'extension souhaitée du problème  $(\mathcal{P}^*)$ , nous donnons dans la Section II quelques résultats d'approximation

et de régularisation de fonctions et de mesures sur une variété riemannienne compacte, puis nous définissons les fonctionnelles  $\mu \rightarrow \int h(\mu)$ , où  $\mu$  est une mesure bornée, et  $h$  une fonction convexe vérifiant certaines conditions. La Section III est consacrée à l'étude des représentations intégrales des solutions d'un problème de Neumann avec données au bord qui sont des mesures. En particulier, nous précisons l'appartenance de ces fonctions ainsi que de leurs traces sur  $\Gamma$ , à certains espaces  $L^p$ . Nous sommes en mesure, dans la Section IV, de définir l'espace  $BT(\Omega)$  ("bounded trace") qui constituera le domaine généralisé d'étude pour le problème  $(\mathcal{P}^*)$ ; de manière informelle,  $BT(\Omega)$  sera défini comme l'ensemble des fonctions définies sur  $\Omega$ , à laplacien dans  $L^2(\Omega)$ , et dont la dérivée normale appartient à  $M_1(\Gamma_0) \times H^{1/2}(\Gamma_1)$  (\*). Les résultats de la section III nous permettent de caractériser les inclusions de  $BT(\Omega)$  dans certains espaces de Lebesgue sur  $\Omega$  et  $\Gamma$ , ainsi que de donner des résultats d'approximation de fonctions de  $BT(\Omega)$  par des éléments de  $C^{\infty}(\bar{\Omega})$ .

Grâce à la notion de fonctionnelle convexe d'une mesure bornée, étudiée à la Section II, nous pouvons introduire, à la Section V, le problème généralisé  $(\mathcal{Q}^*)$  :

$$(\mathcal{Q}^*) \quad \text{Sup} \left\{ -\frac{1}{2} \|\Delta u\|_{L^2(\Omega)}^2 + \int_{\Omega} \Delta u \cdot z_d - \int_{\Gamma_0} h_{\alpha, \beta} \left( -\frac{\partial u}{\partial \nu} \right) - \int_{\Gamma_0} h_{\alpha', \beta}(u) d\Gamma \right\}$$

pour  $u \in BT(\Omega)$ ,  $u = \frac{\partial u}{\partial \nu} = 0$  sur  $\Gamma_1$ . Ici, " $\int_{\Gamma_0} h_{\alpha, \beta} \left( -\frac{\partial u}{\partial \nu} \right)$ " représente

l'intégrale sur  $\Gamma_0$  de la mesure bornée  $h_{\alpha, \beta} \left( -\frac{\partial u}{\partial \nu} \right)$ . Le principal résultat de cette section, obtenu par une méthode directe d'approximation est l'égalité suivante :  $\text{Inf}(\mathcal{P}) = \text{Sup}(\mathcal{P}^*) = \text{Sup}(\mathcal{Q}^*)$ .

La Section VI achève l'étude des problèmes  $(\mathcal{P})$  et  $(\mathcal{Q}^*)$ , et rassemble donc les résultats les plus importants de notre travail. Nous y démontrons l'existence d'un état adjoint, c'est-à-dire une solution dans  $BT(\Omega)$  du problème  $(\mathcal{Q}^*)$ ; ensuite, nous décrivons le système d'optimalité, qui donne les conditions nécessaires et suffisantes pour qu'un élément  $z$  de  $W$  soit l'état optimal. Ces résultats sont obtenus par la méthode directe du calcul des variations, dont l'utilisation est rendue possible grâce à une "formule de Green" généralisée liant un élément  $z$  admissible pour  $(\mathcal{P})$  et un élément  $U$  de  $BT(\Omega)$ .

Dans la Section VII, nous proposons une famille de problèmes perturbés  $(\mathcal{P}_{\epsilon})$ ,  $\epsilon > 0$ , pour laquelle nous prouvons les propriétés suivantes:

- (1) Il y a existence et unicité de l'état optimal  $z_{\epsilon}$ , ainsi que de l'état adjoint  $p_{\epsilon}$ ,
- (2)  $z_{\epsilon}$  converge vers  $z_0$  pour la topologie forte de  $W$ ,



- (3) la famille  $(p_\epsilon)_{\epsilon > 0}$  admet des points d'accumulation pour la topologie faible de  $BT(\Omega)$  qui sont des états adjoints de  $z_0$ .

Finalement, la Section VIII est consacrée à l'étude d'un problème similaire à  $(\mathcal{P})$ , correspondant au cas d'une fonction coût sur la frontière.

$$(\mathcal{R}) \quad \text{Inf} \left\{ \frac{1}{2} \|z - y_d\|_{L^2(\Gamma_1)}^2 + \frac{1}{2} \|z\|_{L^2(\Gamma_0)}^2 + \frac{1}{2} \left\| \frac{\partial z}{\partial v} \right\|_{L^2(\Gamma_0)}^2 \right\}$$

pour  $z \in W$ ,  $\Delta z = 0$ ,  $z \in L^2(\Gamma_1)$ ,  $z(x)$  (resp.  $\frac{\partial z}{\partial v}(x)$ )  $\in [-\alpha, \beta]$  (resp.  $[-\alpha', \beta']$ ) pour presque tout  $x \in \Gamma_0$ ;  $y_d$  est une fonction donnée dans  $L^2(\Gamma_1)$ .

Nous reconduisons quasiment pas à pas la méthode utilisée pour l'étude de  $(\mathcal{P})$ , ce qui nous permet d'obtenir, dans l'espace adapté  $BT_1(\Omega)$ , l'existence de solutions généralisées du problème dual  $(\mathcal{R}^*)$ , ainsi que le système d'optimalité de ce nouveau problème de contrôle optimal.

---

Note de bas de page

(\*) La condition " $\frac{\partial u}{\partial v}|_{\Gamma_1} \in H^{1/2}(\Gamma_1)$ ", techniquement nécessaire, n'impose

pas de restrictions, étant donné le critère d'admissibilité " $\mu = \frac{\partial u}{\partial v} = 0$  sur  $\Gamma_1$ ".

## SOMMAIRE

1. Exposé et formulation variationnelle du problème (P)  
les problèmes  $(P)$  et  $(P^*)$ : égalité des extremas, existence et unicité d'un état optimal  $z_0$ .
2. Quelques résultats techniques indispensables à l'étude de (P):
  - a) Résultats d'approximation et de régularisation sur une variété riemannienne.
  - b) Solutions d'un problème de Neumann avec des données mesures .
3. L'espace  $BT(\Omega)$ : premières propriétés.
4. Généralisation à  $BT(\Omega)$  du problème  $(P^*)$ : le problème  $(Q^*)$   
Egalité des extremas de  $(P^*)$  et  $(Q^*)$ .
5. Existence dans  $BT(\Omega)$  d'états adjoints pour l'état optimal  $z_0$ . Conditions nécessaires et suffisantes d'optimalité .
6. Etude d'un problème de contrôle optimal similaire à (P): existence d'un état optimal unique, existence d'états adjoints généralisés, conditions d'optimalité.
7. Pénalisation du problème (P) par une famille de problèmes sans contraintes  $(P_\epsilon)$ ; étude de la convergence vers (P).



## INTRODUCTION

When one is interested in problems of optimal control, the existence, and even uniqueness of an optimal state are not quite satisfactory ; one also requires some informations about the existence of an adjoint state, so as to establish the optimality system of the problem. When such informations are missing, the problem is said non-well posed [L], and one has to find a suitable way of overcoming this difficulty. In this work, our purpose is to study a non-well-posed problem of convex optimal control, by developing a method of generalized duality for a pair of conjugate problems in convex optimization. Our processes are based on some previous results for the study of plasticity (R. Temam [T], R. Kohn-R. Temam [KT]). Let us now present the mathematical setting of the problem we want to study. Let  $\Omega$  be an open bounded set of the euclidean space  $\mathbb{R}^N$ ,  $N \geq 3$ , whose boundary is the union  $\Gamma_0 \cup \Gamma_1$  of two disjoint connected  $(N-1)$  dimensional  $C^\infty$  submanifolds of  $\mathbb{R}^N$  and let  $(\alpha, \alpha', \beta, \beta')$  be four strictly positive real numbers.

We set :  $\mathcal{U}_{\alpha, \beta} = \{q \in L^2(\Gamma_0), q(x) \in [-\alpha, \beta] \text{ for a.e } x \in \Gamma_0\}$

The problem we want to study is :

to find  $(z_0, u_0, u_1) \in L^2(\Omega) \times \mathcal{U}_{\alpha, \beta} \times \mathcal{U}_{\alpha', \beta'}$  minimizing the cost function

$$J(z, u, v) = 1/2 \|z - z_0\|_{L^2(\Omega)}^2 + 1/2 \|u\|_{L^2(\Gamma_0)}^2 + 1/2 \|v\|_{L^2(\Gamma_0)}^2$$

(P)

the infimum being taken for  $z \in L^2(\Omega)$ ,  $\Delta z = 0$  in  $\Omega$ ,  $z = u$  on  $\Gamma_0$ ,

$$\partial z / \partial r = v \text{ on } \Gamma_0, (u, v) \in U_{\alpha, \beta} \times U_{\alpha', \beta'}$$

( $z_0$  a given function in  $L^2(\Omega)$ ).

If we set  $W = \{z \in L^2(\Omega), \Delta z \in L^2(\Omega), (z, (\partial z / \partial r)) \in L^2(\Gamma_0) \times L^2(\Gamma_0)\}$ , then (P) is obviously equivalent to the problem of convex minimization:

$$(P) \quad \text{Inf} \{ 1/2 |z - z_0|^2_{L^2(\Gamma_0)} + 1/2 |z|^2_{L^2(\Gamma_0)} + 1/2 |(\partial z / \partial r)|^2_{L^2(\Gamma_0)} \}$$

for  $z \in W, \Delta z = 0, (z, (\partial z / \partial r)) \in U_{\alpha, \beta} \times U_{\alpha', \beta'}$ .

The determination of an adjoint state and of the optimality system for (P) is equivalent to solving the conjugate problem of (P), [E-T], that is to say:

$$(P^*) \quad \text{Sup} \{ -1/2 |\Delta q|^2_{L^2(\Omega)} - \int_{\Gamma_0} h_{\alpha, \beta}(-\partial q / \partial r) d\Gamma - \int_{\Gamma_0} h_{\alpha', \beta'}(q) d\Gamma \}$$

for  $q \in W, q = \partial q / \partial r = 0$  on  $\Gamma_1$ .

$$h_{\alpha, \beta} \text{ is the function defined on } \mathbb{R}^1 \text{ by: } h_{\alpha, \beta}(s) \begin{cases} = (s^2/2), & \text{if } s \in [-\alpha, \beta] \\ = \beta(s - \beta/2), & \text{if } s \geq \beta \\ = -\alpha(s + \alpha/2), & \text{if } s \leq -\alpha \end{cases}$$

Problem (P) is easily seen to have a unique solution  $z_0$  in  $W$  [E.T].

Unfortunately, by lack of coercivity (see the definition of  $h_{\alpha, \beta}$ ), we do not know whether Problem (P\*) has a solution in  $W$ .

Nevertheless, one can remark that a minimizing sequence  $q_n$  of  $(P^*)$  is bounded in the following sense :

$$\begin{aligned} \Delta q_n & \text{ is bounded in } L^2(\Omega) \\ (q_n, (\partial q_n / \partial r)) & \text{ is bounded in } M_1(\Gamma_0) \times M_1(\Gamma_0) \end{aligned}$$

where  $M_1(\Gamma_0)$  is the set of bounded measures on  $\Gamma_0$ . As was the case for the problem considered by R. Temam in [T], that remark is the starting point of our method for studying problem  $(P^*)$ : as a matter of fact,  $M_1(\Gamma_0)$  has the good property of vague compactness for bounded sets, which makes it a nice substitute for the study of optimization problems that are coercive in the  $L^1$ -norm.

That is why we are led to introducing the space  $BT(\Omega)$ , which seems to be well adapted to the study of  $(P^*)$ . The main results for  $BT(\Omega)$ , extending some classical results for solutions of elliptic PDE [M1] are the following.

Theorem A :

a). Let  $u \in BT(\Omega)$ ;  $u$  satisfies the properties (i) to (iv)

- (i)  $u \in L^s(\Omega)$ , for  $1 \leq s < N/N-2$  ;
- (ii)  $\Delta u \in L^2(\Omega)$
- (iii)  $u|_{\Gamma} \in L^t(\Gamma_0) \times H^{3/2}(\Gamma_1)$ , for  $1 \leq t < N-1/N-2$
- (iv)  $\partial u / \partial r \in M_1(\Gamma_0) \times H^{1/2}(\Gamma_1)$ .

b) Let  $\tau_1$  be the (weak) topology defined on  $BT(\Omega)$  by :

$$\begin{aligned} u_n \rightarrow u \text{ for } \tau_1 \text{ if } f : \Delta u_n \rightarrow \Delta u \text{ in } L^2(\Omega) \text{ weakly} \\ \partial u_n / \partial r \rightarrow \partial u / \partial r \text{ in } M_1(\Gamma_0) \text{ vaguely and in } H^{1/2}(\Gamma_1) \text{ weakly.} \end{aligned}$$

Then every bounded set of  $BT(\Omega)$  is relatively compact for  $\tau_1$ .

By means of the notion of convex function of a measure, introduced by F. Demengel and R. Temam [DT1], we can define the generalized problem  $(Q^*)$ , which is an extension of  $(P^*)$  to  $BT(\Omega)$ . The study of  $(Q^*)$ , thanks to the generalized duality method, enables us to prove the following results, which solve Problem (P):

Theorem B :

a) There exist adjoint states, in  $BT(\Omega)$ , for the optimal state  $z_0$ ; they are the solutions of  $(Q^*)$ .

b) Let  $z$  be an admissible state;  $z$  is optimal iff there exists an adjoint state  $u \in BT(\Omega)$  satisfying the optimality conditions

- (i)  $\Delta u = -(z - z_0)$  in  $\Omega$
- (ii)  $h_{\alpha, \beta}(u) = u(\partial z / \partial r) - 1/2 |\partial z / \partial r|^2$  a.e. on  $\Gamma$ .
- (iii)  $h_{\alpha, \beta}(-\partial u / \partial r) = -z(\partial u / \partial r) - 1/2 |z|^2$  in  $M_1(\Gamma_0)$ .

We then turn to a study of a penalized form  $(P_\varepsilon)$  of  $(P)$ , having the property that its conjugate  $(P_\varepsilon^*)$  is coercive on  $W$ . The results we prove for  $(P_\varepsilon)$  and  $(P_\varepsilon^*)$  are summed up in the

Theorem C :

- a) There exists a unique optimal state (resp. adjoint state)  $z_\varepsilon$  (resp.  $p_\varepsilon$ ), solution in  $W$  of the optimization problem  $(P_\varepsilon)$  (resp.  $(P_\varepsilon^*)$ ).
- b)  $z_\varepsilon \rightarrow z_0$  in  $W$  as  $\varepsilon \rightarrow 0$ .
- c)  $\{p_\varepsilon\}_{\varepsilon > 0}$  has cluster points for the  $\tau_1$  - topology of  $BT(\Omega)$ , that are adjoint states of  $z_0$ .

Finally, we use our method to study another problem of optimal control (R), similar to (P), in which the cost function involves the boundary value  $z|_{\Gamma_1}$  of  $z$ . As for (P), we prove the existence of a generalized adjoint state for the optimal state, and establish the set of optimality conditions.

The article is organized as follows. Section I presents the first elementary results on  $(P)$  and  $(P^*)$ . Section II and III contains the technical results needed for the introduction of  $BT(\Omega)$  and the generalized problem  $(Q^*)$ . Section IV is devoted to the study of  $BT(\Omega)$ , while, in Section V, we define and study the generalized problem  $(Q^*)$ . In section VI, we develop the generalized duality method, and prove our main results for problem (P). Section VII presents the penalized problems  $(P_\varepsilon)$  and  $(P_\varepsilon^*)$ ; and, in Section VIII, we study problem (R).



## A NON WELL-POSED PROBLEM IN CONVEX OPTIMAL CONTROL.

### 0. Hypothesis, notations.

#### 0.1. Geometrical assumptions.

Let  $\Omega$  be a bounded open set of the euclidean linear space  $\mathbb{R}^N$ ,  $N \geq 3$ . Its boundary  $\Gamma$  is supposed to be the union  $\Gamma_0 \cup \Gamma$  of two  $(N-1)$  dimensional disjoint differentiable manifolds, each of them being compact, connected and indefinitely differentiable.

As usual, we shall write  $dx$  (resp.  $d\Gamma$ ) for the Lebesgue measure on  $\Omega$  (resp.  $\Gamma$ ), induced by the canonical riemannian structure of  $\mathbb{R}^N$ .

#### 0.2. Notations ; the function spaces.

We shall always consider real-valued functions, and generally use the classical notations of the theory of partial differential equations ( $\Delta, \partial/\partial x_i, \partial/\partial r, \dots$ ).

Let  $B$  be a Banach space ; the norm of an element  $x$  of  $B$  will be denoted by  $\|x\|_B$  ; if  $C, C^*$  are two conjugate topological vector spaces,  $\langle x, x^* \rangle_{C \times C^*}$  will stand for the value at  $x$  of the linear form  $x^*$ .

Let  $X$  be an indefinitely differentiable submanifold of  $\mathbb{R}^N$ , and  $\mu$  a measure on  $X$ . We shall write  $L^p(X; d\mu)$  ( $1 \leq p \leq +\infty$ ),  $C^k(X)$  ( $0 \leq k \leq +\infty$ ),  $L(X)$ ,  $H^s(X)$  ( $-\infty < s < +\infty$ ),  $M_1(X)$  for the spaces respectively defined as :

- \* (classes of) measurable functions whose  $p^{\text{th}}$  power is summable on  $X$  with respect to  $\mu$  (for  $p = \infty$ , essentially bounded measurable functions).

- \*  $k$  - times continuously differentiable functions on  $X$
- \* indefinitely differentiable compactly supported functions on  $X$ .
- \* distributions on  $X$  (dual of the former).
- \* Sobolev space of order  $s$  built on  $L^2(X, dx)$  ( $dx$  : Lebesgue measure on  $X$ ).
- \* Bounded measures on  $X$ ,  
each of these being endowed with its usual topology.  
When  $\mu$  is the Lebesgue measure on  $X$ , we shall write  $L^p(X)$  instead of  $L^p(X, d\mu)$ .

The Letter  $C$  will be allowed to represent as many different positive constants as necessary.

## **I. Exposition of the problem.**

### **1.1. Problem (P) and its variational formulation.**

Let us define the space

$$W = \{z \in L^2(\Omega), \Delta z \in L^2(\Omega) \mid (z, \partial z / \partial r) \in L^2(\Gamma_0) \times L^2(\Gamma_0)\},$$

which is a Hilbert space when endowed with the norm

$$\|z\|_W = \{\|z\|_{L^2(\Omega)}^2 + \|\Delta z\|_{L^2(\Omega)}^2 + \|z\|_{L^2(\Gamma_0)}^2 + \|\partial z / \partial r\|_{L^2(\Gamma_0)}^2\}^{1/2}.$$

**Remark 1.1.1.** According to [L2], p. 148,  $z|_{\Gamma_1}$  and  $\partial z / \partial r|_{\Gamma_1}$  are well-defined in  $H^{-1/2}(\Gamma_1)$  and  $H^{-3/2}(\Gamma_1)$ , the trace applications being linear continuous.

Let  $\alpha, \beta$  be two strictly positive real numbers ; we set :

$$U_{\alpha, \beta} = \{p \in L^2(\Gamma_0), p^+ = \text{Sup}(p, 0) \leq \beta, p^- = -\text{Inf}(p, 0) \leq \alpha \text{ a.e. on } \Gamma_0\}$$

$U_{\alpha, \beta}$  is a closed convex set of  $L^2(\Gamma_0)$  .

Given four such numbers  $\alpha, \alpha', \beta, \beta'$  , we want to study the following problem of optimal control.

to find  $(z_0, u_0, v_0) \in L^2(\Omega) \times U_{\alpha, \beta} \times U_{\alpha', \beta'}$  minimizing the cost function :

$$(P) \quad J(z, u, v) = 1/2 \|z - z_d\|_{L^2(\Omega)}^2 + 1/2 \|u\|_{L^2(\Gamma_0)}^2 + 1/2 \|v\|_{L^2(\Gamma_0)}^2$$

the infimum being taken for  $z \in L^2(\Omega)$ ,  $\Delta z = 0$  in  $\Omega$ ,  $z = u$  on  $\Gamma_0$ ,  $\partial z / \partial r = v$  on

$$\Gamma_0, (u, v) \in U_{\alpha, \beta} \times U_{\alpha', \beta'}$$

( $z_d$  is a given function in  $L^2(\Omega)$ ).

In order to establish the variational formulation of (P), we define the functional :

$$j(z) \begin{cases} = 1/2 (\|z - z_d\|_{L^2(\Omega)}^2 + \|z\|_{L^2(\Gamma_0)}^2 + \|\partial z / \partial r\|_{L^2(\Gamma_0)}^2) & \text{if } z \in W \\ \Delta z = 0 & \text{in } \Omega \\ (z, \partial z / \partial r) \in U_{\alpha, \beta} \times U_{\alpha', \beta'} & \\ = +\infty & \text{otherwise} \end{cases} \quad (1.1.1)$$

The functional  $j$  is strictly convex and coercive, that is :

$$\exists C > 0, \forall z \in W, j(z) \geq C \|z\|_W^2 \quad (1.1.2)$$

Moreover,  $j$  is lower semicontinuous ( $\&sc$ ) on  $W$ , for  $U_{\alpha, p}$  is a closed convex set. Problem (P) is obviously equivalent to the convex minimization problem  $(P)$ :

$$(P) \quad \inf_{z \in W} \{j(z)\} \quad (1.1.3)$$

We can now state our first result for (P):

Proposition 1.1.1. There exist a unique optimal state  $z_0$  of problem (P) in  $W$ .

Proof : This is a classical result ([E.T.], ch. II, Prop. 1.2) for the minimization of a strictly convex  $\&sc$  coercive functional on a Hilbert space.

The determination of an adjoint state for the optimal state  $z_0$ , and the establishment of the optimality system for (P) come down to solving the dual problem of (P), and writing the optimality conditions that the minimizer of (P) and a solution of its dual problem must satisfy.

## 1.2. The dual problem.

We shall now give the expression of the dual problem of (1.2.3), using the general duality frame of [E.T].

Let  $Y$  be the space  $(L^2(\Omega))^2 \times (L^2(\Gamma_0))^2$ ;  $Y$  will be identified with its conjugate  $Y^*$ .

Let  $\Lambda$  be the linear operator from  $W$  to  $Y$ , defined by :

$$\Lambda z = (z, \Delta z, z|_{\Gamma_0}, \partial z / \partial r|_{\Gamma_0}) \quad (1.2.1)$$

For  $z$  in  $W$ , and  $p = (p_i)_{1 \leq i \leq 4}$  in  $Y$ , we define the following functions :

$$F(z) = 0, \quad G(p) = \sum_{i=1}^4 G_i(p_i) \text{ with:}$$

$$G_1(p_1) = 1/2 \|p_1 - z_d\|_{L^2(\Omega)}^2 \quad (1.2.2)$$

$$G_2(p_2) \begin{cases} = 0, & \text{if } p_2 = 0 \\ = +\infty & \text{otherwise} \end{cases} \quad (1.2.3)$$

$$G_3(p_3) \begin{cases} = 1/2 \|p_3\|_{L^2(\Gamma_0)}^2, & \text{if } p_3 \in U_{\alpha, \beta} \\ = +\infty & \text{otherwise} \end{cases} \quad (1.2.4)$$

$$G_4(p_4) \begin{cases} = 1/2 \|p_4\|_{L^2(\Gamma_0)}^2, & \text{if } p_4 \in U_{\alpha, \beta} \\ = +\infty & \text{otherwise} \end{cases} \quad (1.2.5)$$

(P) may be written under the following form :

$$(P) \quad \inf_{z \in W} \{F(z) + G(\Lambda z)\} \quad (1.2.6)$$

and its dual problem is ([E.T], p. 61) :

$$(P^*) \quad \sup_{q \in Y^*} \{-G^*(-q) - F^*(\Lambda^*q)\} \quad (1.2.7)$$

$G^*$  (resp.  $F^*$ ) being the conjugate function of  $G$  (resp.  $F$ ), and  $\Lambda^*$ , the transposed operator of  $\Lambda$ .

Remark 1.2.1. The identification of  $Y$  and its conjugate, thanks to its hilbertian structure, permits to write  $(P^*)$  under the following form :

$$(P^*) \quad \sup_{p \in Y} \{-G^*(-p) - F^*(\Lambda^*p)\} \quad (1.2.8)$$

We now compute  $F^*$  and  $G^*$ .

Let  $h_{\alpha, \beta}$  be the function defined by :

$$h_{\alpha, \beta}(s) \begin{cases} = s^2/2, & \text{if } s \in [-\alpha, \beta] \\ = \beta(s - \beta/2), & \text{if } s \geq \beta \\ = -\alpha(s + \alpha/2), & \text{if } s \leq -\alpha \end{cases} \quad (1.2.9)$$

We can easily compute  $G^*$  ([E.T], p. 62) :

$G^*(q) = \sum_{i=1}^4 G_i^*(q_i)$ , with the following expressions for the  $G_i^*$ 's ;

$$G_1^*(q_1) = 1/2 \|q_1\|_{L^2(\Omega)}^2 + \int_{\Omega} q_1 \cdot z_d \, dx \quad (1.2.10)$$

$$G_2^*(q_2) = 0 \quad (1.2.11)$$

$$G_3^*(q_3) = \int_{\Gamma_0} h_{\alpha,\beta}(q_3) \, d\Gamma \quad (1.2.12)$$

$$G_4^*(q_4) = \int_{\Gamma_0} h_{\alpha,\beta}(q_4) \, d\Gamma \quad (1.2.13)$$

For  $F^*$ , we have ( $p \in Y$ ):

$$F^*(\wedge^* p) = \sup_{z \in W} \left\{ \int_{\Omega} p_1 z \, dx + \int_{\Omega} p_2 \Delta z \, dx + \int_{\Gamma_0} p_3 z \, d\Gamma + \int_{\Gamma_1} p_4 (\partial z / \partial r) \, d\Gamma \right\} \quad (1.2.14)$$

hence

$$F^*(\wedge^* p) \geq \sup_{z \in D(\Omega)} \left[ \int_{\Omega} p_1 z \, dx + \int_{\Omega} p_2 \Delta z \, dx + \int_{\Gamma_0} p_3 z \, d\Gamma + \int_{\Gamma_0} p_4 (\partial z / \partial r) \, d\Gamma \right],$$

$\geq$  (given that  $z = \partial z / \partial r = 0$  on  $\Gamma_0$ ),

$$\geq \begin{cases} 0 & \text{if } \Delta p_2 + p_1 = 0 \text{ in } \Omega \\ +\infty & \text{otherwise} \end{cases} \quad (1.2.15)$$

We now assume that  $\Delta p_2 + p_1 = 0$  in  $\Omega$ .

$$p_2 \text{ is such that : } p_2 \in L^2(\Omega), \Delta p_2 \in L^2(\Omega) \quad (1.2.16)$$

Thanks to [L1], p. 148, the following Green's formula is valid, for  $v$  in  $H^2(\Omega)$ :

$$\begin{aligned} \int_{\Omega} (\Delta p_2 \cdot v - \Delta v \cdot p_2) dx &= \langle \partial p_2 / \partial r, v \rangle_{H^{-3/2}(\Gamma) \times H^{3/2}(\Gamma)} \\ &\quad - \langle p_2, \partial v / \partial r \rangle_{H^{-1/2}(\Gamma) \times H^{1/2}(\Gamma)} \end{aligned} \quad (1.2.17)$$

Thus, we can derive the following inequalities :

$$F^*(\Lambda^*p) \geq \sup_{v \in H^2(\Omega)} \left[ \int_{\Omega} p_1 \cdot v dx + \int_{\Omega} p_2 \cdot \Delta v dx + \int_{\Gamma_0} p_3 \cdot v d\Gamma + \int_{\Gamma_0} p_4 \cdot \partial v / \partial r d\Gamma \right]$$

$$\geq \text{(Given (1.2.17))}$$

$$\geq \sup_{v \in H^2(\Omega)} \left[ \langle \partial v / \partial r, p_4 + p_2 \rangle_{H^{1/2}(\Gamma_0) \times H^{-1/2}(\Gamma_0)} \right]$$

$$+ \langle v, p_3 - \partial p_2 / \partial r \rangle_{H^{3/2}(\Gamma_0) \times H^{-3/2}(\Gamma_0)}$$

$$+ \langle \partial v / \partial r, p_2 \rangle_{H^{1/2}(\Gamma_1) \times H^{-1/2}(\Gamma_1)} + \langle v, \partial p_2 / \partial r \rangle_{H^{3/2}(\Gamma_1) \times H^{-3/2}(\Gamma_1)}]$$

$v, \partial v / \partial r$  being arbitrary in  $H^2(\Omega)$ , we finally get :



$$F^*(\wedge^*p) \geq \begin{cases} 0 & \text{if } \begin{aligned} \Delta p_2 + p_1 &= 0 \text{ in } \Omega \\ p_2 = \partial p_2 / \partial r &= 0 \text{ in } \Gamma_1 \\ p_4 + p_2 = p_3 - \partial p_2 / \partial r &= 0 \text{ on } \Gamma_0. \end{aligned} \\ +\infty & \text{otherwise} \end{cases} \quad (1.2.18)$$

$p_2$  satisfies the condition (1.2.16), and also :

$$\begin{aligned} p_2 = \partial p_2 / \partial r &= 0 \text{ on } \Gamma_1 \\ (p_2, \partial p_2 / \partial r) &\in [L^2(\Gamma_0)]^2 \end{aligned} \quad (1.2.19)$$

Let us admit (for the moment) the

Lemma 1.2.1. Let  $(z, p)$  belong to  $W \times W$ , such that  $(p, \partial p / \partial r)$  is in  $H^{3/2}(\Gamma_1) \times H^{1/2}(\Gamma_1)$ .

The following Green's formula is valid :

$$\begin{aligned} \int_{\Omega} (\Delta p \cdot z - \Delta z \cdot p) dx &= \int_{\Gamma_0} (z \cdot \partial p / \partial r - \partial z / \partial r \cdot p) d\Gamma \\ + \langle \partial p / \partial r, z \rangle_{H^{1/2}(\Gamma_1) \times H^{-1/2}(\Gamma_1)} &- \langle p, \partial z / \partial r \rangle_{H^{3/2}(\Gamma_1) \times H^{-3/2}(\Gamma_1)}. \end{aligned} \quad (1.2.20)$$

The expression of  $F^*$  obviously becomes :

$$F^*(\wedge^*p) \begin{cases} = 0 & \text{if } \begin{cases} \Delta p_2 + p_1 = 0 & \text{in } \Omega \\ p_2 = \partial p_2 / \partial r = 0 & \text{on } \Gamma_1 \\ p_4 + p_2 = p_3 - \partial p_2 / \partial r = 0 & \text{on } \Gamma_0. \end{cases} \\ = +\infty & \text{otherwise} \end{cases} \quad (1.2.21)$$

and the dual problem of (1.1.3) is then :

$$\text{Sup}_{p \in Y} \left\{ -1/2 \|p_2\|_{L^2(\Omega)}^2 + \int_{\Omega} p_1 \cdot z_d dx - \int_{\Gamma_0} h_{\alpha,\beta}(-p_3) d\Gamma - \int_{\Gamma_0} h_{\alpha,\beta}(-p_4) d\Gamma \right\} \quad (1.2.22)$$

$$\Delta p_2 + p_1 = 0 \text{ in } \Omega$$

$$p_2 = \partial p_2 / \partial r = 0 \text{ on } \Gamma_1$$

$$p_4 + p_2 = p_3 - \partial p_2 / \partial r = 0 \text{ on } \Gamma_0.$$

We can easily eliminate  $p_1, p_3, p_4$ , and reformulate (1.2.22) as follows :

$$(P^*) \text{ Sup}_{q \in W} \left\{ -1/2 \|\Delta q\|_{L^2(\Omega)}^2 - \int_{\Omega} \Delta q \cdot z_d dx - \int_{\Gamma_0} h_{\alpha,\beta}(-\partial q / \partial r) d\Gamma - \int_{\Gamma_0} h_{\alpha,\beta}(q) d\Gamma \right\}$$

$$q = \partial q / \partial r = 0 \text{ on } \Gamma_1 \quad (1.2.23)$$

We do not know whether there exists an adjoint state in  $W$  for the optimal state  $z_0$  : as a matter of fact,  $(P^*)$  is not coercive on  $W$  (see the definition of  $h_{\alpha,\beta}$ ), and we cannot state the existence of a solution of  $(P^*)$  in  $W$ . One of our major aims is to establish the existence of generalized solutions of

problem (1.2.23), and to relate them to the optimal state  $z_0$ . Right now, we prove Lemma 1.2.1, then we give the first results about the relationships between  $(P)$  and  $(P^*)$ .

Proof of Lemma 1.2.1 : Let  $(z,p)$  be as in the assumptions of Lemma 1.2.1. We now prove the existence of two sequences  $(z_n)_{n \in \mathbb{N}}$  and  $(p^n)_{n \in \mathbb{N}}$  of  $C^\infty(\overline{\Omega})^N$  approximating  $z$  and  $p$  respectively as follows :

$$\begin{aligned} z_n &\rightarrow z \text{ in } W & (1.2.24) \\ p^n &\rightarrow p \text{ in } W \end{aligned}$$

$$(p^n, (\partial p^n / \partial r)) \rightarrow (p, (\partial p / \partial r)) \text{ in } H^{3/2}(\Gamma_1) \times H^{1/2}(\Gamma_1). \quad (1.2.25)$$

Let  $\Phi$  be an element of  $D(\overline{\Omega})$ , identically equal to 0 (resp. 1) in a neighbourhood of  $\Gamma_0$  (resp.  $\Gamma_1$ ), such that  $0 \leq \Phi \leq 1$

We set

$$\begin{aligned} p &= \Phi p + (1 - \Phi) p = p_a + p_b \\ z &= \Phi z + (1 - \Phi) z = z_a + z_b \end{aligned}$$

We now show that  $z_a, z_b, p_a, p_b$  are in  $W$ . For instance, let us prove this result for  $z_a$ .

First of all,  $z_a$  obviously belongs to  $L^2(\Omega)$ .

We also have, in  $D'(\Omega)$ , the following equality, [Sch] ch. II :

$$\Delta z_a = \Delta z \cdot \Phi + \Delta \Phi \cdot z + 2 \sum_{i=1}^N \frac{\partial \Phi}{\partial x_i} \cdot \frac{\partial z}{\partial x_i} \quad (1.2.26)$$

It is now sufficient to prove that  $\frac{\partial \Phi}{\partial x_i} \cdot \frac{\partial z}{\partial x_i}$  belongs to  $L^2(\Omega)$ , for  $1 \leq i \leq N$ .

We know that  $\frac{\partial \Phi}{\partial x_i}$  belongs to  $D(\Omega)$ .

Moreover, we have ([L.M], ch. 2, § 2.3) that  $z$  is locally in  $H^2(\Omega)$ , that is to say :

$$\forall \Theta \in D(\Omega), (\Theta \cdot z) \in H^2(\Omega).$$

Writing  $\frac{\partial \Phi}{\partial x_i} \cdot \frac{\partial z}{\partial x_i} = \frac{\partial}{\partial x_i} (\frac{\partial \Phi}{\partial x_i} \cdot z) - \frac{\partial^2 \Phi}{\partial x_i^2} \cdot z$ , we can see immediately that  $\frac{\partial \Phi}{\partial x_i} \cdot \frac{\partial z}{\partial x_i}$  belongs to  $L^2(\Omega)$ . Therefore,  $z_a$  belongs to  $W$ , and so do  $z_b, p_a, p_b$ . In order to approximate  $p_a$ , we use the conditions [ $p_a \in L^2(\Omega)$ ,  $\Delta p_a \in L^2(\Omega)$ ,  $p_a = \partial p_a / \partial r = 0$  on  $\Gamma_1$ ,  $(p_a, \partial p_a / \partial r) \in H^{3/2}(\Gamma_0) \times H^{1/2}(\Gamma_0)$ ] to find a sequence  $(p_a^{(n)})_{n \in \mathbb{N}}$  of  $(D(\overline{\Omega}))^{\mathbb{N}}$  converging to  $p_a$  in  $H^2(\Omega)$  [L.M], ch. II, § 5).

For  $z_a$ , we use [L1], p. 148, and the condition  $(\text{Supp. } z_a) \cap \Gamma_0 = \emptyset$  to find a sequence  $z_a^{(n)}$  in  $D(\overline{\Omega})$  such that :

$$\lim_{n \rightarrow \infty} \|z_a^{(n)} - z_a\|_{L^2(\Omega)} = \lim_{n \rightarrow \infty} \|\Delta z_a^{(n)} - \Delta z_a\|_{L^2(\Omega)} = 0 \quad (1.2.27)$$

$$z_a^{(n)} = \partial z_a^{(n)} / \partial r = 0 \text{ on } \Gamma_0.$$

Thanks to Remark 1.1.1, we have :

$$(z_a^{(n)}, \partial z_a^{(n)} / \partial r) \rightarrow (z_a, \partial z_a / \partial r) \text{ in } H^{-1/2}(\Gamma_1) \times H^{-3/2}(\Gamma_1) \quad (1.2.28)$$

As for  $p_b$ , it satisfies the conditions  $(p_b \in L^2(\Omega), \Delta p_b \in L^2(\Omega), p_b = \partial p_b / \partial r = 0 \text{ on } \Gamma_1, (p_b, \partial p_b / \partial r) \in (L^2(\Gamma_0))^2)$ . To approximate such a function, we solve the Neumann problem :

$$\begin{aligned} \Delta u &= g_n \text{ in } L^2(\Omega) \\ u + \partial u / \partial r &= h_n \text{ on } \Gamma_0 \\ u + \partial u / \partial r &= 0 \text{ on } \Gamma_1 \end{aligned} \quad (1.2.29)$$

$(g_n)_{n \in \mathbb{N}}$  and  $(h_n)_{n \in \mathbb{N}}$  being such that :

$$g_n \in D(\Omega), \text{ and } \lim_{n \rightarrow \infty} \|g_n - \Delta p_b\|_{L^2(\Omega)} = 0 \quad (1.2.30)$$

$$h_n \in D(\Gamma_0), \text{ and } \lim_{n \rightarrow \infty} \|h_n - (p_b + \partial p_b / \partial r)\|_{L^2(\Gamma_0)} = 0 \quad (1.2.31)$$

According to [LM], ch.II, §7, Problem (1.2.29) admits a unique solution  $p_b^{(n)}$ , converging to  $p_b$  in  $H^{3/2}(\Omega)$ , such that  $\lim_{n \rightarrow \infty} \|\Delta p_b^{(n)} - \Delta p_b\|_{L^2(\Omega)} = 0$ .

We approximate  $z_b$  exactly as we did for  $p_b$ , and find the corresponding sequence  $z_b^{(n)}$ .

If we now set  $z_n = z_a^{(n)} + z_b^{(n)}$ ,  $p^n = p_a^{(n)} + p_b^{(n)}$ , we have obtained two sequences fulfilling the conditions (I.2.24) and (I.2.25); the Green's formula (I.2.20) for  $z$  and  $p$  now results from a straightforward passage to the limit in the Green's formula for  $z_n$  and  $p^n$ , and Lemma I.2.1 is proved.

### 1.3. Comparison of the extremas of $(P)$ and $(P^*)$ .

We now prove the following result :

Theorem I.3.1.  $(P)$  and  $(P^*)$  have the same extrema

$$-\infty < \text{Sup (I.2.23)} = \text{Inf (I.1.3)}. \quad (1.3.1)$$

Proof : For  $p$  in  $Y$ ,  $p = (p_i)_{1 \leq i \leq 4}$ , we consider the penalized problem :

$$\begin{aligned} (P_p) \quad \text{Inf}_{z \in W} \quad & \{ 1/2 \| p_1 - z + z_d \|^2_{L^2(\Omega)} + 1/2 \| z - p_3 \|^2_{L^2(\Gamma_0)} \\ & + 1/2 \| \partial z / \partial r - p_4 \|^2_{L^2(\Gamma_0)} \} \quad (1.3.2) \\ \Delta z = p_2 \quad & \text{in } \Omega \\ (z - p_3, \partial z / \partial r - p_4) \in & U_{\alpha, \beta} \times U_{\alpha', \beta'} \end{aligned}$$

We set  $\Phi(p) = \text{Inf } (P_p)$ .

We already know that  $\Phi(0) < +\infty$  ; using the method of [E.T], we now show

that  $\Phi$  is lsc at zero, which will prove Theorem 1.3.1.

Thanks to [E.T], p.34, we have :

$$\forall p \in Y, \exists ! z_p \in W, \Phi(p) = \inf_{z \in W} (F(z) + G(\Lambda z - p)) \quad (1.3.3)$$

$$\text{Let us suppose : } \quad \lim_{p \rightarrow 0} \Phi(p) < \Phi(0) \quad (1.3.4)$$

We then have :

$$\exists (p_n)_{n \in \mathbb{N}} \in Y^{\mathbb{N}}, \lim_{n \rightarrow \infty} (p_n) = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \Phi(p_n) = \lambda < \Phi(0) \quad (1.3.5)$$

We consider the sequence  $(z_n)_{n \in \mathbb{N}}$  such that  $\Phi(p_n) = F(z_n) + G(\Lambda z_n - p_n)$ . Given that  $p_n$  tends to zero,  $z_n$  is bounded in  $W$ . We can therefore choose a subsequence, still denoted by  $z_n$ , converging to an element  $y$  of  $W$  for the weak topology of  $W$ . As  $p_n$  tends to zero in  $Y$ ,

$$y \text{ satisfies the conditions } \left\{ \begin{array}{l} y \in W \\ \Delta y = 0 \text{ in } \Omega \\ (y, (\partial y / \partial r)) \in \mathcal{U}_{\alpha, \beta} \times \mathcal{U}'_{\alpha, \beta} \\ \text{(for } \mathcal{U}_{\alpha, \beta} \text{, being convex, is weakly closed).} \end{array} \right. \quad (1.3.6)$$

Therefore we have :

$$j(y) < +\infty \quad (1.3.7)$$

$$\begin{aligned} & 1/2(\|y - z\|_{L^2(\Omega)}^2 + \|y\|_{L^2(\Gamma_0)}^2 + \|\partial y / \partial r\|_{L^2(\Gamma_0)}^2) \\ & \leq \lim_{n \rightarrow \infty} (F(z_n) + G(\Lambda z_n - p_n)) < \Phi(0) \end{aligned} \quad (1.3.8)$$

but (1.3.7) and (1.3.8) are contradictory, because of the definition of  $\Phi$ . We have thus proved that  $\Phi$  is  $\ell$ sc at zero, and Theorem 1.3.1 is valid.



## II. Approximation and regularization on a riemannian manifold.

### 1. Some classical results.

We recall the following results, for which we refer to [B.G.M], [M.K.S], [D], [Min]. Let  $M$  be a compact connected  $\ell$ -dimensional riemannian manifold, indefinitely differentiable. The canonical measure on  $M$  is denoted by  $d\lambda$ . If  $M$  is a  $(N-1)$ -dimensional submanifold of  $R^N$ , then  $d\lambda$  is the Lebesgue measure on  $M$ .

One can define ([B.G.M], p.126 to 144) the Laplacean  $\Delta_M$ ;  $\Delta_M$  is a second order differential operator, self-adjoint with respect to the hilbertian structure of  $L^2(M;d\lambda)$ . Its spectrum is a sequence

$$0 = \omega_0 < \omega_1 < \dots < \omega_n \xrightarrow[n \rightarrow \infty]{} +\infty$$

of positive real numbers, associated with a family  $\{X_n\}_{n \in \mathbb{N}}$  of eigenfunctions, orthonormal in  $L^2(M;d\lambda)$  and dense in that space.

Following [B.G.M], [Min], one can construct a fundamental solution  $e(x,y,t)$  of the heat equation  $(\partial/\partial t + \Delta_M = 0)$  on  $M$ .

$e(x,y,t)$  satisfies the following conditions [M.K.S], [Min]:

$$e(x,y,t) \in C^\infty(M \times M \times ]0, +\infty[) \quad (II.1.1)$$

$$e(x,y,t) = \sum_{n=0}^{\infty} e^{-\omega_n t} X_n(x) X_n(y) \quad (II.1.2)$$

$$e(x,y,t) > 0 ; e(x,y,t) = e(y,x,t), (x,y,t) \in M^2 \times ]0, +\infty[ \quad (II.1.3)$$

$$\int_M e(x,y,t)d\lambda(x) = \int_M e(x,y,t)d\lambda(y) = 1 \quad (II.1.4)$$

$$\text{for } f \in \mathcal{C}(M) : \lim_{t \rightarrow 0} (\text{Sup}_{x \in M} |\int_M e(x,y,t) f(y)d\lambda(y) - f(x)|) = 0 \quad (II.1.5)$$

## 2. Approximation and regularization of a function defined on M.

Let  $f$  be an element of  $L^p(M; d\lambda)$ ,  $1 \leq p < +\infty$  ; we set

$$\forall x \in M, f_t(x) = \int_M e(x,y,t) f(y) dy \quad (II.2.1)$$

Thanks to (II.1.1),  $f_t$  belong to  $\mathcal{C}^\infty(M)$ .

We have the following results, whose proofs are given for the commodity of the reader :

Proposition II.2.1. if  $1 \leq p < +\infty$ , then  $\lim_{t \rightarrow 0} \|f - f_t\|_{L^p(M; d\lambda)} = 0$

Proposition II.2.2. if  $p = +\infty$ , then  $(f_t)_{t > 0}$  is such that :

$$\|f_t\|_{L^\infty(M; d\lambda)} \leq \|f\|_{L^\infty(M; d\lambda)} \quad (II.2.i)$$

$$f_t \rightarrow f \text{ for the weak } * \text{ topology of } L^\infty(M; d\lambda) \quad (II.2.ii)$$

Proof of II.2.1 : Let  $p$  be a real number,  $1 \leq p < +\infty$  .

We first recall that  $\mathcal{C}(M)$  is dense in  $L^p(M; d\lambda)$  for the norm topology.

If we suppose  $1 < p < +\infty$ , we have the following inequalities :

$$\begin{aligned}
 |f_t(x)| &= \left| \int_M e(x,y,t) f(y) d\lambda(y) \right| \\
 &\leq \text{(by Hölder's inequality)} \\
 &\leq \left( \int_M |e(x,y,t)| d\lambda(y) \right)^{1-1/p} \left( \int_M |e(x,y,t) f(y)|^p d\lambda(y) \right)^{1/p} \\
 &\leq \text{( Thanks to (II.1.3) and (II.1.4) )} \\
 &\leq \left\{ \int_M e(x,y,t) |f(y)|^p d\lambda(y) \right\}^{1/p}
 \end{aligned}$$

hence :

$$\int_M |f_t(x)|^p d\lambda(x) \leq \int_M \left\{ \int_M e(x,y,t) |f(y)|^p d\lambda(y) \right\} d\lambda(x) .$$

Using Fubini's Theorem, and (II.1.4), we get :

$$\|f_t\|_{L^p(M; d\lambda)} \leq \|f\|_{L^p(M; d\lambda)} \tag{II.2.2}$$

Let us now take  $\varepsilon > 0$  ; we first choose  $f'$  in  $\mathcal{C}(M)$ , such that  $\|f - f'\|_{L^p(M; d\lambda)} < \varepsilon/3$ . The function  $f'_t$  is defined as in (II.2.1).

We may write :

$$\begin{aligned}
 \|f - f_t\|_{L^p(M; d\lambda)} &\leq \|f_t - f'_t\|_{L^p(M; d\lambda)} + \|f'_t - f'\|_{L^p(M; d\lambda)} + \|f' - f\|_{L^p(M; d\lambda)} \\
 &\leq J_1 + J_2 + J_3
 \end{aligned}$$

Using (II.2.2), we have  $0 \leq J_1 \leq J_3 \leq \varepsilon/3$ . Thanks to (II.1.5), we can choose  $t_0$  such that  $J_2 < \varepsilon/3$  for  $t < t_0$  (remember that  $M$  is compact) and this ends the proof when  $1 < p < +\infty$ .

The inequality (II.2.2) being obvious if  $p=1$ , Proposition II.2.1 is proved.

Proof of II.2.2 : We first notice that (II.2.i) is a trivial consequence of (II.1.3) and (II.1.4).

To prove (II.2.ii), we have to show the following result :

for  $f \in L^\infty(M; d\lambda)$ , and  $g \in L^1(M; d\lambda)$ ,

$$\lim_{t \rightarrow 0} \int_M g(f_t - f) d\lambda = 0 \quad (\text{II.2.4})$$

We have :

$$\begin{aligned} \int_M g(x)[f_t(x) - f(x)]d\lambda(x) &= \int_M \left[ \int_M e(x,y,t)f(y)d\lambda(y) - f(x) \right] g(x)d\lambda(x) \\ &= \int_M g(x) \left\{ \int_M e(x,y,t)f(y)d\lambda(y) \right\} d\lambda(x) - \int_M g(x)f(x)d\lambda(x) \\ &= (\text{using Fubini's Theorem}) \\ &= \int_M f(y) \left[ \int_M e(x,y,t)g(x)d\lambda(x) - g(y) \right] d\lambda(y) \\ &= (\text{given (II.1.3)}) \\ &= \int_M f(x)[g_t(x) - g(x)]d\lambda(x) . \end{aligned}$$

Thanks to Proposition II.2.1, we have

$$\lim_{t \rightarrow 0} \|g_t - g\|_{L^1(M; d\lambda)} = 0 \quad (11.2.5)$$

and Proposition 11.2.2 is proved.

Remark 11.2.1 : for  $1 \leq p \leq +\infty$  we have :

$$\lim_{t \rightarrow 0} f_t(x) = f(x) \quad d\lambda - \text{almost everywhere on } M.$$

### 3. Approximation and regularization of a bounded measure on $M$ .

Let  $M_1(M)$  be the space of bounded measures on  $M[B]$ , and  $\mu$  an element of  $M_1(M)$ . We define a regularization  $\mu_t$  of  $\mu$  :

$$\forall x \in M, \quad \mu_t(x) = \int_M e(x, y, t) d\mu(y) \quad (11.3.1)$$

We know, as before, that  $\mu_t$  is indefinitely differentiable on  $M$ .

We now prove the

Proposition 11.3. Let  $\mu$  be in  $M_1(M)$ ; the family  $(\mu_t)_{t>0}$  defined by (11.3.1) converges to  $\mu$  for the vague topology of  $M_1(M)$ . Moreover, we have the following results :

$$\forall t \in ]0, +\infty[, \quad \int_M \mu_t(x) d\lambda(x) = \int_M \mu \quad (11.3.i)$$

if  $\mu$  is a positive measure, then so is  $\mu_t$  for every  $t$  in  $]0, +\infty[$   
(II.3.ii)

Proof of II.3.1 :

(II.3.i) and (II.3.ii) are obvious, thanks to (II.1.3) and (II.1.4).

Let  $f$  be in  $C(M)$ ; we have

$$\begin{aligned} \left| \int_M f(x) \mu_t(x) d\lambda(x) - \int_M f(x) d\mu(x) \right| &= \left| \int_M f(x) \left\{ \int_M e(x,y,t) d\mu(y) \right\} d\lambda(x) \right. \\ &\quad \left. - \int_M f(x) d\mu(x) \right| \\ &= \text{(using Fubini's Theorem)} \\ &= \left| \int_M \left\{ \int_M e(x,y,t) f(x) d\lambda(x) - f(y) \right\} d\mu(y) \right| \\ &\leq \| \mu \|_{M_1(M)} \sup_{y \in M} \left| \int_M e(x,y,t) f(x) d\lambda(x) - f(y) \right| . \end{aligned}$$

and Proposition II.3.1 is now a straightforward consequence of (II.1.5).

#### 4. Convex function of a bounded measure on $M$ .

Let  $h$  be a convex function of one real variable ; its conjugate  $h^*$  is defined by

$$\forall s \in \mathbb{R}, \quad h^*(s) = \sup_{t \in \mathbb{R}} [st - h(t)] \quad (II.4.1)$$

Let  $K = \{s \in \mathbb{R}, h^*(s) < +\infty\}$  be the domain of  $h^*$ .

We shall make, from now on, the following assumptions on  $h$  :

$$\exists C > 0, |h(s)| \leq C(1+|s|) \quad (11.4.2)$$

$$\exists C > 0, \forall s \in K, |h^*(s)| \leq C \quad (11.4.3)$$

Last, we define the "principal part"  $h_\infty$  of  $h$  as follows :

$$\forall s \in \mathbb{R}, h_\infty(s) = \sup_{t \in K} (s.t) = \lim_{t \rightarrow +\infty} h(st)/t \quad (11.4.4)$$

Let  $\mu$  be an element of  $M_1(M)$ .  $\mu$  admits a decomposition with respect to the measure  $d\lambda$  of the following form :

$$\mu = g d\lambda + \mu^s \quad (11.4.5)$$

where  $g$  is locally summable on  $M$ , and  $\mu^s$  is singular.

We denote by  $\Theta_s$  the function defined by  $\mu^s = \Theta_s |\mu^s|$  ([B]).

We can now define the measure  $h(\mu)$ , [D.T (1), (2)], [T] :

$$h(\mu) = (hog) d\lambda + (h_\infty \circ \Theta_s) |\mu^s| \quad (11.46)$$

We recall the following results [D.T (1), (2)], [T] :

(11.4.i)  $h(\mu)$  is a bounded measure, absolutely continuous with respect to  $\mu$  ; moreover, the application  $\mu \rightarrow h(\mu)$  is continuous for the strong topology of  $M_1(M)$ .

(II.4.ii)  $h(\mu)$  has the fundamental property :

$$\forall f \in C(M), \langle h(\mu), f \rangle = \sup_{\substack{g \in C^\infty(M) \\ h^*(g) \in L^1(M; d\lambda)}} \left\{ \int g.f.d\mu - \int_M h^*(g).f.d\lambda \right\}$$

(II.4.iii) If  $\mu_n \rightarrow \mu$  vaguely, and  $h(\mu_n) \rightarrow \nu$  vaguely, then  $h(\mu) \leq \nu$ .

$$\text{If furthermore } h \text{ is positive and } \lim_{n \rightarrow \infty} \int_M h(\mu_n).d\lambda \leq \int_M h(\mu),$$

then  $h(\mu) = \nu$ .

Remark II.4.1. In (II.4.7), the supremum may be taken indifferently in  $C^\infty(M)$  or  $C(M)$ .

We now assume that  $h$  is positive, and prove the following result :

Theorem II.4.1. Let  $h$  be a convex function satisfying the conditions (II.4.2), (II.4.3), such that  $h$  is positive. For any measure  $\mu$  in  $M_1(M)$ , there exists a sequence  $(\mu_n)_{n \in \mathbb{N}}$  in  $(C^\infty(M))^{\mathbb{N}}$ , such that  $(\mu_n, h(\mu_n))$  tends to  $(\mu, h(\mu))$  for the vague topology of  $M_1(M) \times M_1(M)$ .

Proof : We show that the family  $(\mu_t)_{t > 0}$  defined by (II.3.1) satisfies the required conditions.



Let  $g$  be in  $C^\infty(M)$ , such that  $h^*(g)$  belongs to  $L^1(M; d\lambda)$ .

We have :

$$\begin{aligned} \int_M g(x) \mu_t(x) d\lambda(x) - \int_M h^*(g) d\lambda &= \int_M \left[ \int_M e(x,y,t) d\mu(y) \right] g(x) d\lambda(x) \\ &\quad - \int_M h^*(g) d\lambda \tag{II.4.8} \\ &= (\text{using Fubini's Theorem}) \\ &= \int_M \left[ \int_M e(x,y,t) g(y) d\lambda(y) \right] d\mu(x) - \int_M h^*(g) d\lambda \end{aligned}$$

with (II.1.4) :

$$\begin{aligned} h^*\left(\int_M e(x,y,t) g(y) d\lambda(y)\right) &= \sup_{\gamma \in \mathbb{R}} \int_M [\gamma g(y) - h(y)] e(x,y,t) d\lambda(y) \\ &\leq (\text{Thanks to (II.1.3)}) \\ &\leq \int_M h^*(g)(y) e(x,y,t) d\lambda(y) \end{aligned}$$

and therefore

$$\begin{aligned} \int_M h^*\left(\int_M e(x,y,t) g(y) d\lambda(y)\right) d\lambda(x) &\leq \int_M d\lambda(x) \left(\int_M h^*(g)(y) e(x,y,t) d\lambda(y)\right) \\ &\leq (\text{with (II.1.4)}) \\ &\leq \int_M h^*(g) d\lambda \end{aligned}$$

hence (II.4.8) implies :

$$\begin{aligned} \int_M h^*(x) \mu_t(x) d\lambda(x) - \int_M h^*(g) d\lambda &\leq \int_M \left(\int_M e(x,y,t) g(y) d\lambda(y)\right) d\mu(x) - \\ &\quad - \int_M h^*\left(\int_M e(x,y,t) g(y) d\lambda(y)\right) d\lambda \end{aligned}$$

and we have :

$$\begin{aligned} \int_M g \mu_t d\lambda - \int_M h^*(g) d\lambda &\leq \text{Sup} \left\{ \int_M g' d\mu - \int_M h^*(g') d\lambda \right\} \\ &g' \in C^\infty(M) \\ &h^*(g') \in L^1(M, d\lambda) \\ &\leq \int_M h(\mu) . \end{aligned}$$

As we have  $\lim_{t \rightarrow \infty} \int_M h(\mu_t) \leq \int_M h(\mu)$ , we can extract a sequence  $(\mu_{(n)})_{n \in \mathbb{N}}$

satisfying the required conditions, thanks to (II.4.iii) .

In fact, a contradiction argument shows that the whole family  $(\mu_t)_{t > 0}$  tends to  $\mu$  in the sense of Theorem II.4.1.

Corollary II.4.2. The families  $(\mu_t)_{t > 0}$  and  $(h(\mu_t))_{t > 0}$  tends respectively towards  $\mu$  and  $h(\mu)$  in the sense of the tight convergence of measures.

**Proof :** The tight and vague convergences are equivalent, on a compact space, for positive measures.

### III. Integral representations for the solutions of a Neuman problem with irregular boundary data.

#### 1. The classical case.

We are interested in the study of the following Neumann problem :

Let  $f, g$  be two functions respectively defined on  $\Omega$  and  $\Gamma$  ;

$$\begin{aligned} \Delta u &= f \text{ in } \Omega \\ \text{to find } u \text{ solution of:} & \\ u + \partial u / \partial r &= g \text{ on } \Gamma \end{aligned} \tag{III.1.1}$$

We refer to [M<sub>1</sub>], ch. 1 to 3, for the following definitions and results.

Let  $F(x, y)$  be the Green function of Problem (II.1.1) ;  $F$  is defined in  $\overline{\Omega} \times \overline{\Omega} / \{(x, x), x \in \overline{\Omega}\}$ , indefinitely differentiable for  $x \neq y$  and symmetrical in  $(x, y)$ . We set  $H(x, y) = (1/(N-2)\sigma_N)(x-y)^{2-N}$ ,  $\sigma_N$  being the volume of the unit sphere in  $R^N$ .

$$\begin{aligned} F \text{ is such that :} & \quad F(x, y) - H(x, y) = O(|x-y|^{3-N}) \\ \forall (x, y) \in \overline{\Omega} \times \overline{\Omega}, \quad x \neq y & \quad \partial / \partial x_i [F - H](x, y) = O(|x-y|^{2-N}) \\ & \quad \partial^2 / \partial x_i \partial x_j [F - H](x, y) = O(|x-y|^{1-N}) \end{aligned} \tag{III.1.2}$$

(with the notations of Landau).

When  $f$  and  $g$  are summable respectively on  $\Omega$  and  $\Gamma$ , the following result is valid :

Proposition III.1.1. Problem (III.1.1) admits a unique solution  $u$  given by :

$$\forall x \in \Omega, \quad u(x) = - \int_{\Omega} F(x,y)f(y)dy + \int_{\Gamma} F(x,y)g(y)dy \quad (\text{III.1.3})$$

If  $(f,g)$  is in  $C^{\infty}(\overline{\Omega}) \times C^{\infty}(\Gamma)$ , we have  $[M_2]$  :

Proposition III.1.2. The function  $u$  defined by (III.1.3) belongs to  $C^{\infty}(\overline{\Omega})$ .

Last, we recall two theorems of Sobolev's ( $[M_2]$ , Th. 12.VII, [S]p.25-30)

Proposition III.1.3. The operator  $\lambda_1 : L^r(\Omega) \rightarrow C(\overline{\Omega})$

$$f \rightarrow \int_{\Omega} F(.,y)f(y)dy$$

is compact for  $r > N/2$ .

Proposition III.1.4. The operator  $\lambda_2 : L^q(\Gamma) \rightarrow C(\overline{\Omega})$

$$g \rightarrow \int_{\Gamma} F(.,y)g(y)dy$$

is compact, for  $q > (N-1)$ .

## 2. Neumann problem with data in $M_1(\Gamma)$ .

We now wish to study the following problem :

Let  $(f, \mu)$  belong to  $L^2(\Omega) \times M_1(\Gamma)$

$$\Delta u = f \text{ in } \Omega$$

to find  $u$  solution of: (III.2.1)

$$u + \partial u / \partial r = \mu \text{ in } M_1(\Gamma)$$

Let us define the function  $v$  by :

$$\forall x \in \Omega, v(x) = - \int_{\Omega} F(x, y) f(y) dy + \int_{\Gamma} F(x, y) d\mu(y) \quad (\text{III.2.2})$$

$v$  is obviously well defined in  $\Omega$ ; we shall prove that  $v$  is actually the unique solution of the problem (III.2.1).

We shall study separately the domain potential :

$$v_1 = - \int_{\Omega} F(., y) f(y) dy \quad (\text{III.2.3})$$

and the single layer potential

$$v_2 = \int_{\Gamma} F(., y) d\mu(y) \quad (\text{III.2.4})$$

### 3. The domain potential.

Using  $[M_1]$ , Th.12.VIII and Th.13.III, we know that  $v_1$  belongs to  $H^2(\Omega)$ . Furthermore,  $v_1$  is the solution of the homogeneous Neumann problem,

$$\begin{aligned} \Delta u &= f \text{ in } \Omega \\ u + \partial u / \partial r &= 0 \text{ on } \Gamma. \end{aligned} \tag{III.3.1}$$

The condition on  $\Gamma$  is true almost everywhere on  $\Gamma$  (actually, cf [LM], ch II in  $H^{1/2}(\Gamma)$ ).

#### 4. The single layer potential.

We first point out the fact that  $v_2$  is indefinitely differentiable on  $\Omega$ , and satisfies the condition :

$$\Delta v_2 = 0 \text{ in } \Omega \tag{III.4.1}$$

We now prove the following propositions :

Proposition III.4.1. The operator  $\lambda_1' : M_1(\Gamma) \rightarrow L^s(\Omega)$

$$\mu \rightarrow \int_{\Omega} F(.,y) d\mu(y)$$

is compact for  $1 \leq s < N/N-2$ .

Proposition III.4.2. The operator  $\lambda_2' : M_1(\Gamma) \rightarrow L^t(\Gamma)$

$$\mu \rightarrow \int_{\Gamma} F(.,y) d\mu(y)$$

[that is, the value on  $\Gamma$  of the function, defined on  $\Omega$ ,  $\int_{\Gamma} F(.,y) d\mu(y)$ ]

is compact for  $1 \leq t < N-1 / N-2$ .

Proposition III.4.3. The operator  $\lambda_3' : M_1(\Gamma) \rightarrow M_1(\Gamma)$

$$\mu \rightarrow \partial / \partial r \left[ \int_{\Gamma} F(.,y) d\mu(y) \right]$$

[that is, the normal derivative, in the distributional sense, of  $\int_{\Gamma} F(\cdot, y) d\mu(y)$ ]

is continuous, and the following equality is valid :

$$(\lambda_2' + \lambda_3')\mu = \nu \quad \text{in } M_1(\Gamma) \quad (\text{III.4.2})$$

The equalities (III.4.1) and (III.4.2) show that  $v_2$  is the solution of the Neumann problem :

$$\begin{aligned} \Delta u &= 0 \quad \text{in } \Omega \\ u + \partial u / \partial r &= \nu \quad \text{in } M_1(\Gamma) \end{aligned} \quad (\text{III.4.3})$$

Proof of Proposition III.4.1 :

Let  $(s, r)$  be a pair of conjugate exponents ( $1/r + 1/s = 1$ ), such that  $1 \leq s < N/N-2$ .

For  $g \in L^r(\Omega)$  ( $r > N/2$  in this case), we have :

$$\int_{\Omega} v_2(x) g(x) dx = \int_{\Omega} g(x) \left[ \int_{\Gamma} F(x, y) d\mu(y) \right] dx \quad (\text{III.4.4})$$

Proposition III.1.3 allows us to apply Fubini's theorem, and we get :

$$\begin{aligned} \left| \int_{\Omega} v(x) g(x) dx \right| &= \left| \int_{\Gamma} \left[ \int_{\Omega} g(x) F(x, y) dx \right] d\mu(y) \right| \\ &\leq C \|\mu\|_{M_1(\Gamma)} \|g\|_{L^r(\cdot)} \end{aligned}$$

A standard argument of functional analysis gives the desired conclusion, taking into account the fact that the adjoint of a compact operator is itself compact, and Proposition III.4.1 is proved.

Remark III.4.1. We can prove similarly that the partial derivatives  $\partial v_2 / \partial x_i$  ( $1 \leq i \leq N$ ) belong to  $L^\delta(\Omega)$ , for  $1 \leq \delta < N/N-1$ .

Proof of Proposition III.4.2 and III.4.3 :

We first remark that the boundary values  $v_2|_\Gamma$  and  $\partial v_2 / \partial r|_\Gamma$  are defined in the distributional sense on  $\Gamma$ . As a matter of fact,  $v_2$  belongs to  $L^s(\Omega)$ , for  $1 \leq s < N/N-2$ , and satisfies :  $\Delta v_2 = 0$  in  $\Omega$ . If we choose  $K$  such that  $L^s(\Omega)$  is imbedded in  $H^{-K}(\Omega)$  we can see that  $(v_2, \partial v_2 / \partial r)$  belongs to  $H^{-K-1/2}(\Gamma) \times H^{-K-3/2}(\Gamma)$  ([L.M], ch. II, Th. 6.5).

Let us now choose a sequence  $\mu_n$  of smooth functions tightly converging to  $\mu$  in  $M_1(\Gamma)$ , and set :

$$f_n = \int_\Gamma F(.,y) \mu_n(y) dy \quad (\text{III.4.5})$$

We know that  $f_n$  converges pointwise to  $v_2$  in  $\Omega$ , and that the family  $(f_n)_{n \in \mathbf{N}}$  is relatively compact in  $L^s(\Omega)$ . Hence there exists a subsequence, still denoted by  $f_n$ , converging to  $v_2$  in  $L^s(\Omega)$  for the norm topology ; according to [L.M], ch. II, the boundary values  $f_n, \partial f_n / \partial r$  converge respectively to  $v_2, \partial v_2 / \partial r$  in  $\mathcal{D}'(\Gamma)$ . The sequence  $\mu_n$  being bounded in  $L^1(\Gamma)$ , that proves (thanks to Sobolev's theorems, [M1] Th. 12.VII) that  $v_2, \partial v_2 / \partial r$  belong respectively to  $L^t(\Omega)$ , for  $1 \leq t < N-1/N-2$ , and  $M_1(\Gamma)$ , and that the operators  $\lambda'_2, \lambda'_3$  are continuous.

Moreover, the sequence  $f_n$  solves the Neumann problem :



$$\begin{aligned} \Delta f_n &= 0 \text{ in} \\ f_n + \partial f_n / \partial r &= \mu_n \text{ on } \Gamma \end{aligned} \tag{III.4.6}$$

which shows that  $v_2$  satisfies :

$$\begin{aligned} \Delta v_2 &= 0 \text{ in } \Omega \\ (\lambda'_2 + \lambda'_3)v &= \mu \text{ on } \Gamma \end{aligned} \tag{III.4.7}$$

The compactness of  $\lambda'_2$  is a straightforward consequence of the fact that  $\lambda'_2$  is the adjoint operator of  $\lambda_2$  (defined in Prop. III.1.4), which is itself a compact operator.

#### IV. A function space adapted to (P\*) .

##### 1. Definition of BT( $\Omega$ ).

Let  $(q_n)_{n \in \mathbb{N}}$  be a maximizing sequence of Problem (I.2.23) ; its satisfies the following conditions :

$$\Delta q_n \text{ is bounded in } L^2(\Omega) \quad (\text{IV.1.1})$$

$$q_n = \partial q_n / \partial r = 0 \text{ on } \Gamma_1 \quad (\text{IV.1.2})$$

$$(q_n, \partial q_n / \partial r) \text{ is bounded in } L^1(\Gamma_0) \quad (\text{IV.1.3})$$

Hence, it is natural to widen the class of admissible elements for Problem (I.2.23) so as to include functions having their Laplacean in  $L^2(\Omega)$ , and whose boundary value together with their normal derivative are bounded measures on  $\Gamma_0$ . As for the boundary value and normal derivative on  $\Gamma_1$ , we shall require that they should be sufficiently regular to allow a Green's formula to hold with any admissible state for Problem (P).

We consider the operator  $A$ , whose domain is  $L^2(\Omega) \times H^{1/2}(\Gamma_1) \times M_1(\Gamma_0)$ , defined by :

$$\text{for } (f, g, \mu) \in L^2(\Omega) \times H^{1/2}(\Gamma_1) \times M_1(\Gamma_0),$$

$$v = A(f, g, \mu) = - \int_{\Omega} F(., y) f(y) dy + \int_{\Gamma_1} F(., y) g(y) dy + \int_{\Gamma_0} F(., y) d(\mu) \quad (\text{IV.1.4})$$

According to Section III,  $v$  is the solution of the boundary value problem :

$$\begin{aligned}
\Delta u &= f \text{ in } \Omega \\
u + \partial u / \partial r &= g \text{ on } \Gamma_1 \\
u + \partial u / \partial r &= \psi \text{ on } \Gamma_0
\end{aligned}
\tag{IV.1.5}$$

the equality " $u + \partial u / \partial r = \psi$ " being understood in the sense of the distributions on  $\Gamma_0$ .

$\mathcal{A}$  is a one-to-one linear operator, and we define the space  $BT(\Omega)$  (BT stands for "bounded trace") as the range of  $\mathcal{A}$ .  $BT(\Omega)$  is a Banach space, when endowed with the graph norm :

$$\|u\|_{BT(\Omega)} = \|f\|_{L^2(\Omega)} + \|g\|_{H^{1/2}(\Gamma_1)} + \|\psi\|_{M_1(\Gamma_1)}, \text{ for } u = \mathcal{A}(f, g, \psi).$$

We shall now prove the following propositions :

Proposition IV.1.1. The space  $C^\infty(\bar{\Omega})$  is imbedded in  $BT(\Omega)$ .

Proposition IV.1.2 Let  $s$  be a real number,  $1 \leq s < N/N-2$ .

The embedding of  $BT(\Omega)$  into  $L^s(\Omega)$  is compact

Proposition IV.1.3. The operator  $\gamma_0^0$  (resp.  $\gamma_0^1$ ), which assigns to a function in  $C^\infty(\bar{\Omega})$  its boundary value on  $\Gamma_0$  (resp.  $\Gamma_1$ ) can be extended to a compact (resp. continuous) operator from  $BT(\Omega)$  to  $L^\tau(\Gamma_0)$ , for  $1 \leq \tau < N-1/N-2$  (resp.  $H^{3/2}(\Gamma_1)$ ).

Proposition IV.1.4. The operator  $\gamma_1^0$  (resp.  $\gamma_1^1$ ), which assigns to a function in  $C^\infty(\overline{\Omega})$  its normal derivative on  $\Gamma_0$  (resp.  $\Gamma_1$ ) can be extended to a continuous operator from  $BT(\Omega)$  to  $M_1(\Gamma_0)$  (resp.  $H^{1/2}(\Gamma_1)$ ).

Proof of Propositions IV.1.1 to IV.1.4:

For  $(f, g, \mu)$  in  $L^2(\Omega) \times H^{1/2}(\Gamma_1) \times M_1(\Gamma_0)$ , let  $u$  be equal to  $A(f, g, \mu)$ .

We set :  $u = u_1 + u_2$ .

Where  $u_1 = - \int_{\Omega} F(., y) f(y) dy + \int_{\Gamma_1} F(., y) g(y) dy$  (IV.1.6)

$u_2 = \int_{\Gamma_0} F(., y) d(y)$  (IV.1.7)

We first remark that  $u_1$  solves the Neumann problem

$$\begin{aligned} \Delta v &= f \text{ in } \Omega \\ v + \partial v / \partial r &= 0 \text{ on } \Gamma_0 \\ v + \partial v / \partial r &= g \text{ on } \Gamma_1, \end{aligned} \quad (\text{IV.1.8})$$

and therefore ([L.M], ch.II) belongs to  $H^2(\Omega)$ , which is obviously compactly imbedded in  $L^s(\Omega)$  for  $1 \leq s < N/N-2$ . Similarly, the trace (resp. normal derivative) of  $u_1$  on  $\Gamma$  belongs to  $H^{3/2}(\Gamma)$  (resp.  $H^{1/2}(\Gamma)$ ), and obviously  $H^{3/2}(\Gamma)$  is compactly imbedded in  $L^\tau(\Gamma)$ , for  $1 \leq \tau < N-1/N-2$ , while

$H^{1/2}(\Gamma)$  is embedded in  $M_1(\Gamma)$ .

Furthermore, one can see that, given an open set  $A$  such that  $\bar{A}$  does not intersect  $\Gamma_0$ ,  $u_2$  belongs to  $C^\infty(\bar{A})$ . The proof of Propositions IV.1.1 to IV.1.4 is now over, thanks to these remarks, and using Propositions III.1.2, III.4.1, III.4.2, III.4.3.

## 2. Approximation of a function in $BT(\Omega)$ by smooth functions.

We know that  $C^\infty(\bar{\Omega})$  is not dense in  $BT(\Omega)$  for the norm topology, for  $C^\infty(\Gamma)$  is not dense in  $M_1(\Gamma)$ .

Nevertheless, the following result is valid :

Theorem IV.2.1. Given  $u$  in  $BT(\Omega)$  and two numbers  $(s, \tau), 1 \leq s < N/N-2, 1 \leq \tau < (N-1)/N-2$ , there exists a sequence  $(\phi_n)_{n \in \mathbb{N}}$  in  $(C^\infty(\bar{\Omega}))^{\mathbb{N}}$  such that :

$$\phi_n \rightarrow u \text{ in } L^s(\Omega) \quad (\text{IV.2.i})$$

$$\Delta \phi_n \rightarrow \Delta u \text{ in } L^2(\Omega) \quad (\text{IV.2.ii})$$

$$(\gamma_0^0 \phi_n, \gamma_0^1 \phi_n) \rightarrow (\gamma_0^0 u, \gamma_0^1 u) \text{ in } L^\tau(\Gamma_0) \times H^{3/2}(\Gamma_1) \quad (\text{IV.2.iii})$$

$$\gamma_1^0 \phi_n \rightarrow \gamma_1^0 u \text{ vaguely in } M_1(\Gamma_0). \quad (\text{IV.2.iv})$$

$$\gamma_1^1 \phi_n \rightarrow \gamma_1^1 u \text{ in } H^{1/2}(\Gamma_1). \quad (\text{IV.2.v})$$

Proof : We set  $u = A(f, g, \mu)$ ,  $(f, g, \mu) \in L^2(\Omega) \times H^{1/2}(\Gamma_1) \times M_1(\Gamma_0)$ .

We consider a sequence  $f_n$  (resp.  $g_n$ ) of functions in  $C^\infty(\bar{\Omega})$  (resp.  $C^\infty(\Gamma_1)$ ), which tends to  $f$  (resp.  $g$ ) in  $L^2(\Omega)$  (resp.  $H^{1/2}(\Gamma_1)$ ).

We now choose  $(t_n)_{n \in \mathbb{N}}$ , a sequence of positive real number tending to zero, and we set  $\mu^{(n)} = \mu_{t_n}$ , where  $\mu_{t_n}$  is defined by (II.3.1).

The approximation sequence  $\Phi_n$  is then :

$$\Phi_n = - \int_{\Omega} F(., y) f_n(y) dy + \int_{\Gamma_1} F(., y) g_n(y) dy + \int_{\Gamma_0} F(., y) \mu^{(n)}(y) dy \quad (IV.2.1)$$

Thanks to Proposition III.1.2, we know that  $\Phi_n$  belongs to  $C^\infty(\bar{\Omega})$ .

Let us now split  $\Phi_n$  into

$$\Phi_n^{(1)} = - \int_{\Omega} F(., y) f_n(y) dy + \int_{\Gamma_1} F(., y) g_n(y) dy \quad (IV.2.2)$$

and

$$\Phi_n^{(2)} = \int_{\Gamma_0} F(., y) \mu^{(n)}(y) dy \quad (IV.2.3)$$

We already know that  $\Phi_n^{(1)}$  converges towards  $u_1$  (defined by (IV.1.6)) in  $H^2(\Omega)$ . Thanks to Propositions IV.2.2 and IV.2.3, we can extract from  $\Phi_n^{(2)}$  a subsequence, still denoted by  $\Phi_n^{(2)}$ , satisfying the following conditions :

$$\Phi_n^{(2)} \rightarrow u_2 \text{ in } L^5(\Omega) \quad (IV.2.4)$$

$$\gamma_0^0 \Phi_n^{(2)} \rightarrow \gamma_0^0 u_2 \text{ in } L^1(\Gamma_0) \quad (IV.2.5)$$

$$\gamma_1^0 \Phi_n^{(1)} \rightarrow \gamma_1^0 u_2 \text{ in } M_1(\Gamma_0) \text{ vaguely} \quad (\text{IV.2.6})$$

( $u_2$  is defined by (IV.1.7)).

Only two statements are now left to prove, namely :

$$\gamma_0^1 \Phi_n^{(2)} \rightarrow \gamma_0^1 u_2 \text{ in } H^{3/2}(\Gamma_1) \quad (\text{IV.2.7})$$

$$\gamma_1^1 \Phi_n^{(2)} \rightarrow \gamma_1^1 u_2 \text{ in } H^{1/2}(\Gamma_1)$$

Let us admit (for the moment) the following result :

**Lemma IV.2.2 :** Let  $A$  be an open set in  $\Omega$ , such that  $\bar{A} \cap \Gamma_0 = \emptyset$ . The family  $(\Phi_n^{(2)})_{n \in \mathbb{N}}$  is equicontinuous in  $\bar{A}$ , and so is any of the families of its successive derivatives.

We can now extract another subsequence, still denoted by  $\Phi_n^{(2)}$ , from  $\Phi_n^{(2)}$  which tends to  $u_2$  for the topology of  $C^2(\bar{A})$ , and a fortiori in  $H^2(A)$ , for  $\Omega$  is bounded.

The sequence  $\Phi_n = \Phi_n^{(1)} + \Phi_n^{(2)}$  satisfies the required conditions, and Theorem IV.2.1 is proved.

Proof of Lemma IV.2.2

Let  $(x, x')$  be in  $\bar{A} \times \bar{A}$ ;

we have :

$$|\Phi_n^{(2)}(x) - \Phi_n^{(2)}(x')| = \left| \int_{\Gamma_0} (F(x, y) - F(x', y)) \mu^{(n)}(y) dy \right|$$

$$\leq \int_{\Gamma_0} |F(x, y) - F(x', y)| |\mu^{(n)}(y)| dy$$

$\leq$  (with Theorem II.4.1)

$$\leq \|\mu\|_{M_1(\Gamma_0)} \sup_{y \in \Gamma_0} |F(x, y) - F(x', y)|$$

Hence, the equicontinuity of  $\Phi_n^{(2)}$  is obvious, for  $F(x, y)$  is indefinitely differentiable in  $\bar{A} \times \Gamma_0$ , and so Lemma IV.2.2 is proved.

Finally, we prove the following result, which will be of great use in the further study of Problem (I.2.23) :

Corollary IV.2.3. Let  $h$  be a convex function of one real variable satisfying the assumptions of Theorem II.4.1. The sequence  $(\Phi_n)_{n \in \mathbb{N}}$  defined by (IV.2.1) is such that :

$$h(y_1^0 \Phi_n) \rightarrow h(y_1^0 u) \text{ vaguely in } M_1(\Gamma_0) \quad (\text{IV.2.vi})$$



Proof : Thanks to Proposition III.4.3, we know that  $\gamma_1^0 \Phi_n = \mu^{(n)} - \gamma_0^0 \Phi_n$ .  
Using Theorems IV.2.1 and II.4.1, we know that  $h(\mu^{(n)})$  tends to  $h(\mu)$  vaguely in  $M_1(\Gamma_0)$ , while  $\gamma_0^0 \Phi_n$  tends to  $\gamma_0^0 u$  in  $L^1(\Gamma_0)$ , and this proves Corollary IV.2.3.

**V. Extension of  $(P^*)$  to the space  $BT(\Omega)$  ;  
the generalized problem  $(Q^*)$ .**

**1. Definition of  $(Q^*)$**

We recall the formulation of Problem (I.2.23)

$$(P^*) \quad \sup_{q \in W} \{-j^*(q)\} \quad (V.1.1)$$

where  $j^*$  is the convex functional defined on  $W$  by

$$j^*(q) \begin{cases} = 1/2 \|\Delta q\|_{L^2(\Omega)}^2 + \int_{\Omega} \Delta q \cdot z_d dx + \int_{\Gamma_0} h_{\alpha,\beta}(-\partial q / \partial r) d\Gamma \\ \quad + \int_{\Gamma_0} h_{\alpha,\beta}(q) d\Gamma \\ \text{if } q = \partial q / \partial r = 0 \text{ on } \Gamma_1 \\ = +\infty \text{ otherwise.} \end{cases} \quad (V.1.2)$$

Thanks to results of II and IV, we can see that  $j^*$  can be naturally extended to a functional  $J^*$  whose domain is  $BT(\Omega)$ . As a matter of fact, every element  $u$  of  $BT(\Omega)$  has a Laplacean  $\Delta u$  in  $L^2(\Omega)$ , a trace  $\gamma_0^0 u$  in  $L^1(\Gamma_0)$  and a normal derivative  $\gamma_1^0 u$  in  $M_1(\Gamma_0)$ . We may then define  $\int_{\Gamma_0} h_{\alpha,\beta}(-\gamma_1^0 u)$  as the integral on  $\Gamma_0$  of the bounded measure  $h_{\alpha,\beta}(-\gamma_1^0 u)$ .

Moreover, owing to the compactness of  $\Gamma_0$ , this integral is equal to the duality product  $\langle 1, h_{\alpha,\beta}(-\gamma_1^0 u) \rangle_{C(\Gamma_0) \times M_1(\Gamma_0)}$ .

We now define the maximization problem  $(Q^*)$ :

$$(Q^*) \quad \sup_{u \in BT(\Omega)} \{-J^*(u)\} \quad (V.1.3)$$

in which the functional  $J^*$  is defined as follows :

$$J^*(u) \begin{cases} = 1/2 \|\Delta u\|_{L^2(\Omega)}^2 + \int_{\Omega} \Delta u \cdot z_d \cdot dx + \int_{\Gamma_0} h_{\alpha, \beta}(-\gamma_1^0 u) \\ \quad + \int_{\Gamma_0} h_{\alpha, \beta}(\gamma_0^0 u) d\Gamma \\ \text{if } u = \partial u / \partial r = 0 \text{ on } \Gamma_1 \\ = +\infty \text{ otherwise.} \end{cases} \quad (V.1.4)$$

The space  $W$  is obviously imbedded in  $BT(\Omega)$ , and we have, for  $q$  in  $W$  :

$$j^*(q) = J^*(q) \quad (V.1.5)$$

As a consequence, the following inequality is satisfied :

$$\sup_{u \in BT(\Omega)} (-J^*(u)) \geq \sup_{q \in W} (-j^*(q)). \quad (V.1.6)$$

Actually, we shall now prove a stronger result :

Theorem V.1.1. The extremas of problems (V.1.3) and (I.2.23) are equal.

Proof : We first give an approximation result :

Lemma V.1.2. Let  $u$  be in  $BT(\Omega)$  : there exists a sequence  $(u_n)_{n \in \mathbb{N}}$  of  $(C^\infty(\bar{\Omega}))^N$  which tends to  $u$  in the sense of Theorem IV.2.1 and Corollary

IV.2.3, such that  $\gamma_0^{-1} u_n = \gamma_1^{-1} u_n = 0$  for every integer  $n$ .

Let us suppose that Lemma V.1.2 is proved.

For any  $u$  in  $BT(\Omega)$ , such that  $J^*(u) < +\infty$ , one can then find a sequence  $u_n$  of smooth functions, admissible for Problem (I.2.23), which satisfies:

$$\lim_{n \rightarrow \infty} j^*(u_n) = J^*(u) \quad (V.1.7)$$

That is obviously sufficient to prove Theorem V.1.1.

Proof of Lemma V.1.22:

We consider ([L.M], ch.II, §8) a family  $(\Gamma_1^\xi)_{0 \leq \xi \leq 1}$  of indefinitely differentiable surfaces parallel to  $\Gamma_1$  ( $\Gamma_1^0 = \Gamma_1$ ).

Let  $A_\xi$  be the open set limited by  $\Gamma_1$  and  $\Gamma_1^\xi$ , and  $A$  be the union  $\cup A_\xi$ ,  $0 \leq \xi \leq 1$ .  $A$  is supposed to be such that  $\bar{A} \cap \Gamma_0 = \emptyset$ .

The following result is then valid ([A], p.216):

Proposition V.1.3. There exists a family of truncating functions  $\{\psi_\xi\}_{0 \leq \psi_\xi \leq 1}$ , elements of  $C^\infty(\bar{\Omega})$ , satisfying the conditions:

$$0 \leq \psi_\xi \leq 1 ; \psi_\xi = 0 \text{ in } \bar{A}_{\xi/2} ; \psi_\xi = 1 \text{ in } \Omega / A_\xi \quad (V.1.8)$$

and such that

$$\forall v \in H^2(\Omega), v = \partial v / \partial r = 0 \text{ on } \Gamma_1, \lim_{\xi \rightarrow 0} \|(\psi_\xi - 1)v\|_{H^2(\Omega)} = 0 \quad (V.1.9)$$

In order to prove Lemma V.1.2, we shall truncate in the neighbourhood of  $\Gamma_1$  the functions  $\Phi_n$  constructed in Theorem IV.2.1.

We set :  $\Phi_{n,\xi} = \psi_\xi \Phi_n$  (V.1.10)

It is now sufficient to prove

Proposition V.1.4 : Let  $s$  be a real number,  $1 \leq s < N/N-2$ , the following assertion is true :  $\forall \varepsilon > 0, \exists \xi_0 \in ]0,1[, \exists n_0 \in \mathbb{N}$ ,

$$\forall n > n_0, \|\Phi_{n,\xi} - u\|_{L^s(\Omega)} < \varepsilon \quad (\text{V.1.i})$$

$$\|\Delta \Phi_{n,\xi} - \Delta u\|_{L^2(\Omega)} < \varepsilon \quad (\text{V.1.ii})$$

Proof of (V.1.i) :

We have

$$\begin{aligned} \|\Phi_{n,\xi} - u\|_{L^s(\Omega)} &= \left\{ \int_{\Omega} |\Phi_n(x) \psi_\xi(x) - u(x)|^s dx \right\}^{1/s} \\ &\leq \left\{ \int_{\Omega} |\Phi_n(x) \psi_\xi(x) - \Phi_n(x)|^s dx \right\}^{1/s} + \left\{ \int_{\Omega} |\Phi_n(x) - u(x)|^s dx \right\}^{1/s} \\ &\leq I_1 + I_2 \end{aligned}$$

Thanks to the definition of  $\Phi_n$ , we first choose  $n_1$  such that  $I_2 < \varepsilon/2$  for  $n > n_1$ .

For  $I_1$ , we remark that  $\Phi_n$  is bounded in  $L^{s_1}(\Omega)$ ,  $1 \leq s < s_1 < N/N-2$ , and therefore :

$$\begin{aligned} (I_1)^s &= \int_{\Omega} |1 - \Psi_{\xi}(x)|^s |\Phi_n(x)|^s dx \\ &\leq (\text{by Holder's inequality}) \\ &\leq \left( \int_{\Omega} |\Phi_n(x)|^{s_1} dx \right)^{s/s_1} \left( \int_{\Omega} |1 - \Psi_{\xi}(x)|^{s_1/s_1 - s} dx \right)^{s_1 - s / s_1} \\ &\leq C. (\text{mes } A_{\xi})^{s_1 - s / s_1}. \end{aligned}$$

$C$  being independent of  $n$ . We now choose  $\xi_1$  such that  $C(\text{mes } A_{\xi}) < \varepsilon/2$  for  $\xi \leq \xi_1$ , and (V.1.i) is proved.

Proof of (V.1.ii) :

$$\begin{aligned} \|\Delta(\Phi_n \Psi_{\xi}) - \Delta u\|_{L^2(\Omega)} &= \|\Delta(\Psi_{\xi} \Phi_n) - \Delta(\Psi_{\xi} u)\|_{L^2(\Omega)} + \|\Delta(\Psi_{\xi} u) - \Delta u\|_{L^2(\Omega)} \\ &\leq (\text{Thanks to Proposition V.1.3}) \\ &\leq \|\Psi_{\xi} \Phi_n - \Psi_{\xi} u\|_{H^2(A)} + \|\Delta \Phi_n - \Delta u\|_{L^2(\Omega)} + \|\Psi_{\xi} u - u\|_{H^2(A)} \\ &\leq J'_1 + J'_2 + J'_3 \end{aligned}$$

Proposition V.1.3 enables us to choose  $\xi_2$  such that  $J'_3 < \varepsilon/3$  for  $\xi \leq \xi_2$ . Owing to the definition of  $\Phi_n$ , we can choose  $n_2$  such that  $J'_2 < \varepsilon/3$  for  $n > n_2$ . Finally,  $\xi_2$  being fixed, we can find  $n_3$  such that  $J'_1 < \varepsilon/3$  for

$n > n_3$ , thanks to Lemma IV.2.2.

We have finally obtained :

$$\begin{aligned} \forall \varepsilon > 0, \quad \exists \xi_0 = \text{Inf}(\xi_1, \xi_2), \exists n_0 = \text{Sup}(n_1, n_2, n_3), \\ n > n_0 \Rightarrow \text{(V.1.i) and (V.1.ii)} \end{aligned} \quad \text{(V.1.11)}$$

and Lemma V.1.2 is proved.

Remark V.1.1. We have used, without mentioning it, the fact that  $BT(\Omega)$  is a local space (see Lemma I.2.1), and that  $u$  belongs to  $H^2(A)$ .

**VI. Existence of and adjoint state for Problem (P) ;**  
**System of optimality conditions.**

In this section, we give the main results of our work : the existence in  $BT(\Omega)$ , of an adjoint state for the optimal state of (P), and the optimality system for (P). We introduce a generalized duality between (P) and  $(Q^*)$ , which enables to use the direct method of calculus of variations and give very simple proofs for our results. Most of the latter are based on the generalized Green's formula hereafter.

**1. Green's formula.**

In this section, we prove the following result :

Proposition VI.1.1 Let  $(u,z)$  be an element of  $BT(\Omega) \times W$ ,  $z$  being an admissible state for problem (P). Then  $z$  satisfies the conditions :

$$\begin{aligned} z \in L^2(\Omega) ; \Delta z \in L^\infty(\Omega) ; (\gamma_0^0 z, \gamma_1^0 z) \in \mathcal{C}(\Gamma_0) \times L^\infty(\Gamma_0) ; (\gamma_1^0 z, \gamma_1^1 z) \\ \in H^{-1/2}(\Gamma_1) \times H^{-3/2}(\Gamma_1) \end{aligned}$$

and the Green's formula (V.1.i) is valid :

$$\int_{\Omega} (\Delta z \cdot u - \Delta u \cdot z) dx = \int_{\Gamma} ((\partial z / \partial r) \cdot u - (\partial u / \partial r) \cdot z) d\Gamma \quad (V.1.i)$$

Remark V.1.1 : The precise formulation of (V.1.i) is :



$$\int_{\Omega} (\Delta z \cdot u - \Delta u \cdot z) dx = \int_{\Gamma_0} \gamma_1^0 z \gamma_0^0 u \, d\Gamma + \langle \gamma_1^1 z, \gamma_0^1 u \rangle_{H^{-3/2}(\Gamma_1) \times H^{3/2}(\Gamma_1)} - \langle \gamma_0^0 z, \gamma_1^0 u \rangle_{C(\Gamma_0) \times M_1(\Gamma_0)} - \langle \gamma_0^1 z, \gamma_1^1 u \rangle_{H^{-1/2}(\Gamma_1) \times H^{-3/2}(\Gamma_1)}$$

Proof of Proposition III.1.1 :

We first point out the fact that all the expressions in (VI.1.1) are meaningful, and we proceed to prove an approximation result for  $z$ , namely :

Proposition VI.1.2. Let  $z$  belong to  $W$ ,  $z$  being admissible for Problem (I.1.3), and let  $\rho$  be a real number,  $2 \leq \rho < +\infty$ . There exists a sequence  $Z_n$  of functions of  $C^\infty(\overline{\Omega})$ , such that :

$$\Delta Z_n \rightarrow \Delta z \quad \text{in } L^\rho(\Omega); \quad (\text{VI.1.a})$$

$$Z_n \rightarrow z \quad \text{in } L^2(\Omega); \quad (\text{VI.1.b})$$

$$\gamma_1^1 Z_n \rightarrow \gamma_1^1 z \quad \text{in } H^{-1/2}(\Gamma_1) \quad (\text{VI.1.c})$$

$$\gamma_1^0 Z_n \rightarrow \gamma_1^0 z \quad \text{in } L^r(\Gamma_0), \text{ for } 1 \leq r < +\infty, \text{ and in } L^\infty(\Gamma_0) \text{ * -weakly :} \\ (\text{VI.1.d})$$

$$(\gamma_0^0 Z_n, \gamma_0^1 Z_n) \rightarrow (\gamma_0^0 z, \gamma_0^1 z) \quad \text{in } C(\Gamma_0) \times H^{-1/2}(\Gamma_1) \quad (\text{VI.1.e})$$

From this result, the Green's formula (VI.1.1) will be proved by a straight forward passage to the limit, using Theorem IV.2.1.

We start with a Lemma :

Lemma VI.1.3 We set  $\chi_\rho(\Omega) = \{v \in L^2(\Omega), \Delta v \in L^\rho(\Omega)\}$ , for  $2 \leq \rho < +\infty$  ;  $C^\infty(\overline{\Omega})$  is dense in  $\chi_\rho(\Omega)$  for the norm topology defined by :

$$\|v\|_{\chi_\rho(\Omega)} = \|v\|_{L^2(\Omega)} + \|\Delta v\|_{L^\rho(\Omega)} \quad (\text{VI.1.1})$$

Proof : Let  $b$  be a continuous linear form on  $\chi_\rho(\Omega)$ , whose restriction to  $C^\infty(\overline{\Omega})$  is the null form.

Thanks to the Hahn-Banach Theorem, there exists an element  $(f, g)$ , of  $L^2(\Omega) \times L^{\rho'}(\Omega)$ ,  $1/\rho + 1/\rho' = 1$ , such that

$$\langle v, b \rangle_{\chi_\rho(\Omega) \times [\chi_\rho(\Omega)]^*} = \int_{\Omega} f \cdot v \, dx + \int_{\Omega} g \cdot \Delta v \, dx \quad (\text{VI.1.2})$$

$b$  being null on  $C^\infty(\overline{\Omega})$ , we have, for  $v$  in  $D(\Omega)$  :

$$\int_{\Omega} [f \cdot v + g \cdot \Delta v] \, dx = 0 \quad (\text{VI.1.3})$$

and hence :

$$f + \Delta g = 0 \quad \text{in } D'(\Omega) \quad (\text{VI.1.4})$$

$g$  is then a function in  $L^\rho(\Omega)$ , whose Laplacean is in  $L^2(\Omega)$ ; we can define ([L.M], ch.II), in the distributional sense on  $\Gamma$ , its boundary values  $\gamma_0 g$  and  $\gamma_1 g$ .

For  $v$  in  $C^\infty(\overline{\Omega})$ , we shall have the Green's formula :

$$\int_{\Omega} (\Delta v \cdot g, \Delta g \cdot v) dx = \langle \gamma_1 v, \gamma_0 g \rangle_{D(\Gamma) \times D'(\Gamma)} - \langle \gamma_0 v, \gamma_1 g \rangle_{D(\Gamma) \times D'(\Gamma)} \quad (\text{VI.1.5})$$

but the assumptions on  $b$  now imply that  $\gamma_0 g = \gamma_1 g = 0$ .

From this, we obtain that  $g$  is in  $H^2(\Omega)$ , and we may write the Green's formula for  $v \in X_0(\Omega)$  ([L], p.148) :

$$\begin{aligned} \int_{\Omega} (g \cdot \Delta v - v \cdot \Delta g) &= \langle \gamma_0 g, \gamma_1 v \rangle_{H^{3/2}(\Gamma) \times H^{-3/2}(\Gamma)} \\ &- \langle \gamma_1 g, \gamma_0 v \rangle_{H^{1/2}(\Gamma) \times H^{-1/2}(\Gamma)} = 0 \end{aligned} \quad (\text{VI.1.6})$$

We may therefore conclude that  $b$  is the null form on  $X_0(\Omega)$ ; and  $C^\infty(\overline{\Omega})$  is dense in  $X_0(\Omega)$ .

Let us now prove Proposition VI.1.2.

Let  $z$  belong to  $W$ , such that  $j(z) < +\infty$ . We have necessarily :

$$\Delta z = 0 \text{ in } \Omega ; (\gamma_0^0 z, \gamma_0^1 z) \in (L^\infty(\Gamma_0))^2 \quad (\text{V.1.7})$$

Owing to the hypoellipticity of the Laplacean, we know that  $z$  is indefinitely differentiable in  $\Omega$ .

We now consider the same function  $\Phi$  as in the proof of Lemma I.1.1 and we set as before  $z = \Phi z_1 + (1 - \Phi) z_2$ .

$$= z_1 + z_2.$$

By using the fact that  $z$  and all its partial derivatives are bounded in any relatively compact open set of  $\Omega$ , we can show exactly as we did for Lemma I.2.1 that  $z_1$  and  $z_2$  are such that :

$$(z_1, z_2) \in (L^2(\Omega))^2 ; (\Delta z_1, \Delta z_2) \in (L^r(\Omega))^2 \quad (1 \leq r \leq +\infty) \quad (\text{VI.1.8})$$

In order to approximate  $z_2$ , we use Lemma VI.1.3 in an open set  $\omega$  whose closure does not intersect  $\Gamma_0$ , and obtain a sequence  $z_n^{(2)}$ .

For  $z_1$ , we have the additional conditions :

$$(\gamma_0^0 z_1, \gamma_1^0 z_1) = (\gamma_0^0 z, \gamma_1^0 z) \in L^\infty(\Gamma_0)^2 ; \gamma_1^0 z_1 = \gamma_1^1 z_1 = 0 \text{ on } \Gamma_1 \quad (\text{VI.1.9})$$

Therefore,  $z_1$  is the only solution of the Neumann problem :

$$\begin{aligned} \Delta z_1 &= \Delta(\Phi z) \text{ in } \Omega \\ z_1 + \partial z_1 / \partial r &= z + \partial z / \partial r \text{ on } \Gamma_0 \\ z_1 + \partial z_1 / \partial r &= 0 \text{ on } \Gamma_1 \end{aligned}$$

and this implies, using III, that  $z_1$  is defined by :

$$z_1 = - \int_{\Omega} F(y) [\Delta(\Phi z)](y) dy + \int_{\Gamma_0} F(y) [z + \partial z / \partial r](y) dy \quad (\text{VI.1.11})$$

Thanks to Proposition III.1.3, and [M<sub>1</sub>], Th.12.VIII,  $z_1$  is continuous on  $\overline{\Omega}$ ; if we use the approximation of  $(\gamma_0^0 z + \gamma_1^0 z)$  defined by (II.2.1), and any sequence of smooth functions converging to  $\Delta(\Phi z)$  in  $L^p(\Omega)$ , we can construct an approximating sequence  $Z_n^{(2)}$  which is easily shown to satisfy the conditions :

$$\begin{aligned} Z_n^{(1)} &\rightarrow z_1 \text{ in } C(\overline{\Omega}) ; \Delta Z_n^{(1)} \rightarrow \Delta z_1 \text{ in } L^p(\Omega) \\ (\gamma_1^0 Z_n^{(1)} ; \gamma_1^1 Z_n^{(1)}) &\rightarrow (\gamma_1^0 z_1, \gamma_1^1 z_1) \text{ in } L^r(\Gamma_0), \text{ for } 1 \leq r < +\infty, \\ &\text{and in } L^\infty(\Gamma_0) \text{ * - weakly} \end{aligned}$$

If we now set  $Z_n = Z_n^{(1)} + Z_n^{(2)}$ , it is clear that  $Z_n$  approximates  $z$  in the sense of Proposition V.1.2 ; we just have to take into account the fact that  $\gamma_0^0 Z_n^{(2)} = \gamma_1^0 Z_n^{(2)} = 0$  for every integer  $n$ .

Now, we can conclude the proof of Proposition VI.1.1 ; for this, we consider the sequence  $u_n$  of Theorem IV.2.1, with  $\tau=1$ ,  $1 \leq s < N/N-2$ , and the sequence  $Z_n$  for  $\rho = s/s-1$ .

We then have the classical Green's formula :

$$\int_{\Omega} (\Delta u_n Z_n - \Delta Z_n u_n) dx = \int_{\Gamma} ((\partial u_n / \partial r) Z_n - u_n (\partial Z_n / \partial r)) d\Gamma \quad (\text{VI.1.12})$$

and one can easily check that the conditions satisfied by  $u_n$  and  $Z_n$  ensures that equality (VI.1.12) is still valid after passing to the limit.

Remark VI.1.1 If we only make the assumption that  $\Delta z$  is bounded in  $\Omega$ , we can nevertheless prove analogous results. This is done by using the results of  $L^p$  regularity for the solutions of elliptic boundary value problems ([G.G]).

## 2. Existence in $BT(\Omega)$ of an adjoint state for $z_0$ .

First, we prove two auxiliary results

Lemma VI.2.1. Let  $(u, z)$  belong to  $BT(\Omega) \times W$ ,  $u$  (resp.  $z$ ) being admissible for Problem (V.1.3) (resp. I.1.3)). The following inequality is satisfied

$$-J^*(u) \leq j(z) - 1/2 \|\Delta u - (z - z_0)\|_{L^2(\Omega)}^2 \quad (\text{VI.2.1})$$

Proof : we recall that  $y_0^0 z$  is continuous on  $\Gamma_0$ , and that  $y_0^0(x)$  (resp.  $y_1^0(x)$ ) belongs to  $[-\alpha, \beta]$  (resp.  $[-\alpha', \beta']$ ) for almost every  $x$  in  $\Gamma_0$ .

From the definition of the conjugate of a convex function, and thanks to (II.4.7) we have :

$$h_{\alpha, \beta}(\gamma_0^0 u)(x) \geq (\gamma_0^0 u, \gamma_1^0 z)(x) - 1/2(\gamma_1^0 z(x))^2 \text{ a.e. on } \Gamma_0 \quad (\text{VI.2.2})$$

and

$$\begin{aligned} \langle h_{\alpha, \beta}(-\gamma_1^0 u), 1 \rangle_{M_1(\Gamma_0) \times C(\Gamma_0)} &\geq \langle -\gamma_1^0 u, \gamma_0^0 z \rangle_{M_1(\Gamma_0) \times C(\Gamma_0)} \\ &\quad - 1/2 \|\gamma_0^0 z\|_{L^2(\Gamma_0)}^2 \end{aligned} \quad (\text{VI.2.3})$$

Therefore, we obtain :

$$\begin{aligned} -J^*(u) &\leq -1/2 \|\Delta u\|_{L^2(\Omega)}^2 - \int_{\Omega} \Delta u \cdot z_d + 1/2 \|z\|_{L^2(\Gamma_0)}^2 + 1/2 \|\partial z / \partial r\|_{L^2(\Gamma_0)}^2 \\ &\quad + \int_{\Gamma_0} ((\partial u / \partial r) \cdot z - (\partial z / \partial r) \cdot u) d\Gamma \end{aligned} \quad (\text{VI.2.4})$$

$\leq$  (thanks to Proposition VI.1.1)

$$\begin{aligned} &\leq -1/2 \|\Delta u\|_{L^2(\Omega)}^2 - \int_{\Omega} \Delta u \cdot z_d dx + \int_{\Omega} (\Delta u \cdot z - \Delta z \cdot u) dx + \\ &\quad 1/2 \|\partial z / \partial r\|_{L^2(\Gamma_0)}^2 + 1/2 \|z\|_{L^2(\Gamma_0)}^2 \end{aligned}$$

$\leq$  (given that  $\Delta z = 0$  in  $\Omega$ )

$$\begin{aligned} &\leq 1/2 \|z - z_d\|_{L^2(\Omega)}^2 + 1/2 \|z\|_{L^2(\Gamma_0)}^2 + 1/2 \|\partial z / \partial r\|_{L^2(\Gamma_0)}^2 - \\ &\quad 1/2 \|\Delta u - (z - z_d)\|_{L^2(\Omega)}^2 \end{aligned}$$

and the last inequality is precisely the result we wished to prove.

We now define on  $BT(\Omega)$  the weak topology  $\tau_1$  as follows :

$$\begin{aligned}
& \Delta u_n \rightarrow \Delta u \text{ weakly in } L^2(\Omega) \\
u_n \rightarrow u \text{ for } \tau_1 \text{ if } & \begin{aligned} & \gamma_0^0 u_n \rightarrow \gamma_0^0 u \text{ in } L_1(\Gamma_0) \\ & \gamma_1^0 u_n \rightarrow \gamma_1^0 u \text{ vaguely in } M_1(\Gamma_0) \\ & (\gamma_0^1 u_n, \gamma_1^1 u_n) \rightarrow (\gamma_0^1 u, \gamma_1^1 u) \text{ in } L^2(\Gamma_1). \end{aligned}
\end{aligned} \tag{VI.2.5}$$

Thanks to the results of III and IV, we know that every bounded set of  $BT(\Omega)$  is relatively compact for the  $\tau_1$  topology.

For the  $\tau_1$  topology on  $BT(\Omega)$ , we have :

Lemma VI.2.2. The functional  $J^*$  is lower semi continuous on  $BT(\Omega)$  for  $\tau_1$ .

Proof : this result is obvious, thanks to (II.4.i), (II.4.iii) and to the lower semi continuity of the norm for the weak topology of  $L^2(\Omega)$ .

We can now prove the existence result for an adjoint state of  $z_0$  :

Theorem VI.2.3. There exists in  $BT(\Omega)$  an adjoint state for the optimal state  $z_0$  of (P) ; it is a solution of Problem (V.1.3). Moreover, any maximizing sequence of (1.2.23) has a cluster point for the  $\tau_1$  topology of  $BT(\Omega)$  which is an adjoint state of  $z_0$ . Last, any solution of (V.1.3) can be obtained as a cluster point for such a sequence.

Proof : Let  $(w_n)_{n \in \mathbb{N}}$  be a maximizing sequence of Problem (V.1.3). Thanks to Lemma VI.2.1,  $w_n$  is bounded in  $BT(\Omega)$ . Thus, it has a cluster point  $w$  for



the  $\tau_1$  topology, given Proposition III.4.1 and the definition of  $BT(\Omega)$ ; moreover  $w$  is admissible for Problem (V.1.3), for the convex set  $(\gamma_0^0 u = \gamma_1^1 u = 0 \text{ on } \Gamma_1)$  is closed for the  $\tau_1$  topology.

Owing to Lemma VI.2.2, we then have :

$$(-J^*(w)) \geq \lim_{n \rightarrow \infty} (-J^*(w_n)) = \text{Sup}(V.1.3) \quad (\text{VI.2.6})$$

and (VI.2.6) obviously implies that  $w$  is a solution of Problem (V.1.3). The second part of Theorem VI.2.3 is a consequence of Theorem V.1.1, while the last assertion is obvious, using the approximation result of Lemma V.1.2.

### 3. The optimality system.

We can now establish the set of optimality conditions for Problem (P).

Theorem VI.3.1. Let  $z$  be an admissible state for (P); then  $z$  is the optimal state if and only if there exists  $w$  in  $BT(\Omega)$  admissible for problem (V.1.3), and such that :

$$\Delta w = z - z_d \text{ in } \Omega \quad (\text{VI.3.i})$$

$$h_{\alpha, \beta}(\gamma_0^0 w) = \gamma_0^0 w \cdot \gamma_1^0 z - 1/2(\gamma_1^0 z)^2 \text{ almost everywhere on } \Gamma_0 \quad (\text{VI.3.ii})$$

$$h_{\alpha, \beta}(-\gamma_1^0 w) = -\gamma_1^0 w \cdot \gamma_0^0 z - 1/2(\gamma_0^0 z)^2 \text{ in } M_1(\Gamma_0) \quad (\text{VI.3.iii})$$

Moreover, if equalities (VI.3.i) to (VI.3.iii) are satisfied,  $z$  and  $w$  are a pair of solutions of the dual optimization problems  $(P)$  and  $(Q^*)$ .

Proof : If we look at the proof of Lemma VI.2.1, we remark that (VI.3.i), (VI.3.ii), (VI.3.iii) are nothing but the necessary and sufficient conditions for  $j(z)$  and  $(-J^*(w))$  to be equal.

## VII. Another problem of optimal control.

In this section, we apply the method of generalized duality to a problem of convex optimal control, in which the cost function involves the boundary value of the considered state.

### 1. Variational formulation.

We set

$W_1 = \{y \in W, y \in L^2(\Gamma_1)\}$  ;  $W_1$  is a Hilbert space when endowed with the norm :

$$\|y\|_{W_1} = \{\|y\|_{L^2(\Gamma_1)}^2 + \|y\|_W^2\}^{1/2}.$$

Moreover, the trace operator  $y \rightarrow \partial y / \partial r|_{\Gamma_1}$  is linear continuous from  $W_1$  into  $H^{-1}(\Gamma_1)$ , and  $W_1$  is imbedded in  $H^{1/2}(\Omega)$  ([L.M], ch. II). The problem we wish to study is :

to find  $(y_0, u_0, v_0) \in W_1 \times \mathcal{U}_{\alpha, \beta} \times \mathcal{U}_{\alpha, \beta}$ , minimizing the cost function

$$J_1(y, u, v) = 1/2 \{ \|y - y_d\|_{L^2(\Gamma_1)}^2 + \|u\|_{L^2(\Gamma_0)}^2 + \|v\|_{L^2(\Gamma_0)}^2 \}$$

(R)

the infimum being taken for  $y \in W_1$ ,  $\Delta y = 0$  in  $\Omega$ ,  $y = u$  on  $\Gamma_0$ ,

$$\partial y / \partial r = v \text{ on } \Gamma_0,$$

$$(u, v) \in \mathcal{U}_{\alpha, \beta} \times \mathcal{U}_{\alpha, \beta}.$$

( $y_d$  is a given function in  $L^2(\Gamma_1)$ ).

We define the functional

$$j_1(y) \begin{cases} = 1/2\{\|y-y_d\|_{L^2(\Gamma_1)}^2 + \|y\|_{L^2(\Gamma_0)}^2 + \|\partial y/\partial r\|_{L^2(\Gamma_0)}^2\} & \text{if } y \in W_1, \\ \Delta y = 0 \text{ in } \Omega, (y, \partial y/\partial r) \in \mathcal{U}_{\alpha,\beta} \times \mathcal{U}'_{\alpha,\beta} & \\ = +\infty & \text{otherwise.} \end{cases} \quad (\text{VII.1.1})$$

The variational formulation of Problem (R) is :

$$(R) \quad \inf_{y \in W_1} \{j_1(y)\} \quad (\text{VII.1.2})$$

Thanks to the results of [LM], ch.II, we know that  $j_1$  is coercive on  $W_1$  :

$$\exists C, \forall y \in W_1, j_1(y) \geq C\|y\|_{W_1}^2 \quad (\text{VII.1.3})$$

and therefore the following result is valid (see Section I):

Proposition VII.1.1. There exists a unique optimal state  $y_0$  of (R).

We now turn to the problem of existence of an adjoint state for  $y_0$ ; this leads us to studying the dual problem  $(R^*)$  of (R), which we now formulate.

We set  $Y_1 = L^2(\Omega) \times (L^2(\Gamma_0))^2 \times L^2(\Gamma_1)$ ;  $Y_1$  will be identified with its

conjugate  $Y_1^*$ .  $\Lambda_1$  is the operator from  $W_1$  to  $Y_1$  defined by

$$\Lambda_1(g) = (\Delta y, y|_{\Gamma_0}, \partial y / \partial \nu|_{\Gamma_0}, y|_{\Gamma_1}).$$

Let  $F, G$  be the functionals respectively defined on  $W_1$  and  $Y_1$  by :

$$F(y) = 0, \quad G(p) = \sum_{i=1}^4 G_i(p_i),$$

$$G_1(p_1) \quad \left\{ \begin{array}{l} = 0, \text{ if } p_1 = 0 \\ = +\infty \text{ otherwise} \end{array} \right. \quad (\text{VII.1.4})$$

$$G_2(p_2) \quad \left\{ \begin{array}{l} = 1/2 \|p_2\|_{L^2(\Gamma_0)}^2, \text{ if } p_2 \in \mathcal{U}_{\alpha, \beta} \\ = +\infty \text{ otherwise} \end{array} \right. \quad (\text{VII.1.5})$$

$$G_3(p_3) \quad \left\{ \begin{array}{l} = 1/2 \|p_3\|_{L^2(\Gamma_0)}^2, \text{ if } p_3 \in \mathcal{U}_{\alpha, \beta} \\ = +\infty \text{ otherwise} \end{array} \right. \quad (\text{VII.1.6})$$

$$G_4(p_4) = 1/2 \|p_4 - y_d\|_{L^2(\Gamma_1)}^2 \quad (\text{VII.1.7})$$

the dual problem  $(R^*)$  of Problem (VII.1.2) is then :

$$(R^*) \quad \sup_{q \in Y_1} \{-G^*(-q) - F^*(\Lambda^*, q)\} \quad (\text{VII.1.8})$$

We compute the conjugate functions  $F^*$  and  $G^*$  as in Section I ; for  $F^*$ ,

we shall need the following approximation result :

Lemma VII.1.2. Let  $(y,p)$  belong to  $W_1 \times W_1$ , such that  $\partial p/\partial r$  belongs to  $L^2(\Gamma_1)$ .

(i) there exist two sequences  $y_n$  and  $p'_n$  of smooth functions, such that :

$$\begin{aligned} p'_n &\rightarrow p \text{ in } H^{3/2}(\Omega) \\ \Delta p'_n &\rightarrow \Delta p \text{ in } L^2(\Omega) \end{aligned} \quad (\text{VII.1.9})$$

$$y_n \rightarrow y \text{ in } W_1 \quad (\text{VII.1.10})$$

(ii) The following Green's formula is valid :

$$\begin{aligned} \int_{\Omega} (\Delta p \cdot y - \Delta y \cdot p) dx = \\ \int_{\Gamma_0} ((\partial p/\partial r) y - (\partial y/\partial r) \cdot p) d\Gamma + \int_{\Gamma_1} \partial p/\partial r \cdot y d\Gamma \quad (\text{VII.1.11}) \\ - \langle p, (\partial y/\partial r) \rangle_{H^1(\Gamma_1) \times H^{-1}(\Gamma_1)} \end{aligned}$$

Proof : We first remark that the existence of  $p'_n$  has been shown in the proof of Lemma I.2.1. So as to approximate  $y$ , we write  $y = y_1 + y_2$ ,  $y_1$  and  $y_2$  being such that :

$$\begin{aligned} (y_1, \Delta y_1) &\in (L^2(\Omega))^2 \\ (y_1, (\partial y_1/\partial r)) &\in (L^2(\Gamma_0))^2 \quad (\text{VII.1.12}) \\ (\text{Supp } y_1) \cap \Gamma_1 &= \emptyset \end{aligned}$$

$$\begin{aligned}
(y_2, \Delta y_2) &\in (L^2(\Omega))^2 \\
y_2 &\in L^2(\Gamma_1) \\
\text{Supp}(y_2) \cap \Gamma_0 &= \emptyset
\end{aligned}
\tag{VII.1.13}$$

(cf. the proof of Lemma I.2.1).

The function  $y_1$  can be approximated in the same way as  $p$ ; for  $y_2$ , we first choose an indefinitely differentiable open set  $\omega_1$  of  $\Omega$ , whose closure does not intersect  $\Gamma_0$ , such that  $\text{supp}(y_1)$  is contained in  $\omega_1$ . We then solve a Dirichlet problem in  $\omega_1$ , namely:

$$\begin{aligned}
\Delta y_{2n} &= r_n \text{ in } \omega_1 \\
y_{2n} &= t_n \text{ on } \Gamma_1 \\
y_{2n} &= 0 \text{ on } \Gamma_0
\end{aligned}
\tag{VII.1.14}$$

$r_n$  (resp.  $t_n$ ) being a sequence of smooth functions approximating  $\Delta y_2$  (resp.  $y_2$ ) in  $L^2(\Omega)$  (resp.  $L^2(\Gamma_1)$ ). We know that  $y_{2n}$  tends to  $y_2$  in  $H^{1/2}(\omega_1)$ ,

while  $\partial y_{2n}/\partial r$  tends to  $\partial y_2/\partial r$  in  $H^{-1}(\Gamma_1)$ ; moreover,  $\Delta y_{2n}$  obviously tends to  $\Delta y_2$ . If we now set  $y_n = y_{1n} + y_{2n}$ , we obtain the sequence we were looking for, and Lemma VII.1.2 is proved.

The expression of  $F^*(\Lambda^*_1, q)$  is then easily shown to be :

$$F^*(\Lambda^*_1, q) \begin{cases} = 0 & \text{if } \Delta q_1 = 0 \text{ in } \Omega \\ & q_1 + q_3 = q_2 - \partial q_1 / \partial r = 0 \text{ on } \Gamma_0 \\ & q_4 - \partial q_1 / \partial r = q_1 = 0 \text{ on } \Gamma_1 \\ = +\infty & \text{otherwise.} \end{cases} \quad (\text{VII.1.15})$$

Computing the  $G_i^*$  is easy, and we find for  $(R^*)$  :

$$(R^*) \quad \begin{aligned} \text{Sup}_{q \in W_1} & \left\{ - \int_{\Gamma_0} h_{\alpha, \beta} (-\partial q / \partial r) d\Gamma - \int_{\Gamma_0} h_{\alpha, \beta}(q) d\Gamma \right. \\ & \left. - 1/2 \|\partial q / \partial r\|_{L^2(\Gamma_1)}^2 + \int_{\Gamma_1} (\partial q / \partial r) y_d d\Gamma \right\} \quad (\text{VII.1.16}) \\ & \Delta q = 0 \text{ in } \Omega \\ & q = 0 \text{ on } \Gamma_1 \\ & \partial q / \partial r \in L^2(\Gamma_1) \end{aligned}$$

The difficulty we meet here is obviously of the same kind as in Section I : the functional  $j_1^*$  defining  $(R^*)$  is not coercive on  $W_1$ , and hence, we do not know whether there exists an adjoint state in  $W_1$  for the optimal state  $y_0$ .

Nevertheless, we have the following comparison result :



Theorem VII.1.3. The extremas of Problems (VII.1.2) and (VII.1.16) are equal.

Proof : We consider the perturbations  $(R_p)$  of  $(R)$ , for  $p$  given in  $Y_1$  :

$$(R_p) \quad \text{Inf}_{z \in W_1} \{F'(z) + G'(\Lambda_1 z - p)\}$$

Given  $p^{(n)}$ , a sequence in  $Y_1$  converging to zero, let  $y^{(n)}$  be the solution of  $(R_{p^{(n)}})$ . As in the proof of Theorem I.3.1, we show that we can extract from  $y^{(n)}$  a subsequence weakly converging in  $W_1$  to the solution  $y_0$  of Problem (VII.1.2), the latter being therefore normal ([E.T]). Hence, the equality  $\text{Inf}(R) = \text{Sup}(R^*)$  is true.

## 2. A function space adapted to Problem (VII.1.16).

We now introduce a new function space, modelled on  $BT(\Omega)$ , on which we shall define a natural extension of  $(R^*)$ .

Let us first remark that a maximizing sequence  $(q_n)_{n \in \mathbb{N}}$  of  $(R^*)$  satisfies the conditions :

$$\Delta q_n = 0 \text{ in } \Omega ; (q_n, (\partial q_n / \partial r)) \text{ is bounded in } L^1(\Gamma_0)$$

$$\partial q_n / \partial r \text{ is bounded in } L^2(\Gamma_1) ; q_n = 0 \text{ on } \Gamma_1 \dots \quad (\text{VII.2.1})$$

Let  $A_1$  be the operator defined on  $L^2(\Omega) \times L^2(\Gamma_1) \times M_1(\Gamma_0)$  by :

$$A_1(f, g, \mu) = - \int_{\Omega} F(., y) f(y) dy + \int_{\Gamma_1} F(., y) g(y) dy + \int_{\Gamma_0} F(., y) d\mu(y) \quad (\text{VII.2.2})$$

where  $F$  is the Green function of the Neumann problem (III.1.1). We define the space  $BT_1(\Omega)$  as the range of  $A_1$ , endowed with the graph norm.  $BT_1(\Omega)$  is a Banach space, and its elements are all the functions defined on  $\Omega$  whose Laplacean is in  $L^2(\Omega)$ , and whose normal derivative on  $\Gamma_0$  (resp.  $\Gamma_1$ ) is a bounded measure (resp. in  $L^2(\Gamma_1)$ ). The properties of  $BT_1(\Omega)$  are very similar to those of  $BT(\Omega)$ ; actually, all the results of Section IV are valid for  $BT_1(\Omega)$ , after replacing  $H^{1/2}(\Gamma_1)$  and  $H^{3/2}(\Gamma_1)$  respectively by  $L^2(\Gamma_1)$  and  $H^1(\Gamma_1)$ .

The extended functional  $J_1^*$  is then defined on  $BT_1(\Omega)$  as follows :

$$J_1^*(q) \begin{cases} = \int_{\Gamma_0} h_{\alpha, \beta}(-\partial q / \partial r) + \int_{\Gamma_0} h_{\alpha, \beta}(q) d\Gamma + 1/2 \|\partial q / \partial r\|_{L^2(\Gamma_1)}^2 \\ \quad - \int_{\Gamma_1} \partial q / \partial r \cdot y_d d\Gamma \\ \\ \text{if } \Delta q = 0 \text{ in } \Omega, q = 0 \text{ on } \Gamma_1 \\ \\ = +\infty \text{ otherwise.} \end{cases} \quad (\text{VII.1.3})$$

As before,  $\int_{\Gamma_0} h_{\alpha, \beta}(-\partial q / \partial r)$  stands for the integral on  $\Gamma_0$  of the bounded measure  $h_{\alpha, \beta}(-\partial q / \partial r)$ .

The generalised problem ( $S^*$ ) is :

$$(S^*) \quad \text{Sup}_{q \in BT_1(\Omega)} \{-J_1^*(q)\} \quad (\text{III.2.4})$$

The following inequality is obvious :

$$\text{Sup (III.1.16)} \leq \text{Sup (VII.2.4)} \quad (\text{II.2.5})$$

In order to carry out the generalized duality method, as we did in Section VI, we shall need the following results :

Proposition III.2.1. Let  $(y, q)$  belong to  $W_1 \times BT_1(\Omega)$ ,  $y$  (resp.  $q$ ) being admissible for Problem (VII.1.2) (resp. (VII.2.4)). We have the Green's formula :

$$\begin{aligned} 0 &= \int_{\Omega} (\Delta q \cdot y - \Delta y \cdot q) dx = \langle \partial q / \partial r, y \rangle_{M_1(\Gamma_0) \times C(\Gamma_0)} \\ &\quad - \int_{\Gamma_0} \partial y / \partial r \cdot q \, d\Gamma + \int_{\Gamma_1} (\partial q / \partial r) \cdot y \, d\Gamma \\ &\quad - \langle q, (\partial y / \partial r) \rangle_{H^1(\Gamma_1) \times H^{-1}(\Gamma_1)} \end{aligned} \quad (\text{VII.2.6})$$

Proof : As for Lemma VI.1.1, all we need to prove is an approximation result for  $y$  and  $q$ . For  $q$ , we use the modified version of Theorem IV.2.1, with  $H^{3/2}(\Gamma_1)$  (resp.  $H^{1/2}(\Gamma_1)$ ) replaced by  $H^1(\Gamma_1)$  (resp.  $L^2(\Gamma_1)$ ); for  $y$ , we prove the existence of a sequence  $y'$  satisfying the conditions :

$$\begin{aligned}
y_n' &\rightarrow y_n \text{ in } L^2(\Omega); \Delta y_n' \rightarrow \Delta y_n \text{ in } L^\beta(\Omega) \text{ } (\beta \text{ a given number,} \\
&2 \leq \beta < +\infty); \\
y_n' &\rightarrow y \text{ in } C(\Gamma_0) \text{ and } L^2(\Gamma_1); \partial y_n' / \partial r \rightarrow \partial y / \partial r \text{ in } L^r(\Gamma_0), \\
&\text{for } 1 \leq r < +\infty, \text{ and } L^\infty(\Gamma_0) \text{ weakly;} \\
\partial y_n' / \partial r &\rightarrow \partial y / \partial r \text{ in } H^{-1}(\Gamma_1). \tag{VII.2.7}
\end{aligned}$$

To prove this, we set  $y = y_1 + y_2$  as in Lemma (VII.1.2) with  $y_1, y_2$  such that :

$$\begin{aligned}
\text{Supp}(y_1) \cap \Gamma_1 &= \emptyset \\
(y_1, \Delta y_1) &\in L^2(\Omega) \times L^\beta(\Omega) \tag{VII.2.8} \\
(y_1, \partial y_1 / \partial r) &\in L^\infty(\Gamma_0) \times L^\infty(\Gamma_0)
\end{aligned}$$

$$\begin{aligned}
\text{Supp}(y_2) \cap \Gamma_0 &= \emptyset \\
(y_2, \Delta y_2) &\in L^2(\Omega) \times L^\beta(\Omega) \tag{VII.2.9} \\
y_2 &\in L^2(\Gamma_1)
\end{aligned}$$

We approximate  $y_1$  as in the proof of Proposition VI.1.1 ; For  $y_2$  we solve a Dirichlet problem similar to (VII.1.14), with a sequence  $r_n$  converging to  $\Delta y_2$  in  $L^\beta(\Omega)$ .

The Green's formula (VII.2.6) is then proved, and both expressions appearing in it are null, thanks to :  $\Delta y = \Delta q = 0$  in  $\Omega$ .

### 3. Existence in $BT_1(\Omega)$ of an adjoint state for $y_0$ .

We first prove two auxiliary results :

Lemma VII.3.1. Let  $(y, q)$  be a couple of admissible elements for Problems (VII.1.2) and (VII.2.4) ; we have the following inequality :

$$-J_1^*(q) \leq j_1(y) \quad (\text{VII.3.1})$$

Proof : by the definition of  $h_{\alpha, \beta}, h_{\alpha, \beta}^*$ , and thanks to (II.4.ii), we have

$$\int_{\Gamma_0} h_{\alpha, \beta}(-\partial q / \partial r) \geq - \int_{\Gamma_0} (\partial q / \partial r) y \, d\Gamma - 1/2 \int_{\Gamma_0} (y)^2 \, d\Gamma \quad (\text{VII.3.2})$$

$$\int_{\Gamma_0} h_{\alpha, \beta}^*(q) \, d\Gamma \geq \int_{\Gamma_0} q (\partial y / \partial r) \, d\Gamma - 1/2 \int_{\Gamma_0} (\partial y / \partial r)^2 \, d\Gamma \quad (\text{VII.3.3})$$

and therefore we derive the following inequalities :

$$\begin{aligned} -J_1^*(q) &\leq -1/2 \|\partial q / \partial r\|_{L^2(\Gamma_0)}^2 + \int_{\Gamma_1} \partial q / \partial r \cdot y_d \, d\Gamma \\ &\quad + \int_{\Gamma_0} ((\partial q / \partial r) \cdot y - (\partial y / \partial r) \cdot q) \, d\Gamma + 1/2 \|y\|_{L^2(\Gamma_0)}^2 \\ &\quad + 1/2 \|\partial y / \partial r\|_{L^2(\Gamma_0)}^2 \\ &\leq 1/2 \|y - y_d\|_{L^2(\Gamma_0)}^2 + \int_{\Gamma_0} ((\partial q / \partial r) \cdot y - (\partial y / \partial r) \cdot q) \, d\Gamma + \int_{\Gamma_1} \partial q / \partial r \cdot y \, d\Gamma \\ &\quad + 1/2 \|y\|_{L^2(\Gamma_0)}^2 + 1/2 \|\partial y / \partial r\|_{L^2(\Gamma_0)}^2 \\ &\leq (\text{for } q=0 \text{ on } \Gamma_1) \\ &\leq j_1(y) + \int_{\Gamma} ((\partial q / \partial r) \cdot y - (\partial y / \partial r) \cdot q) \, d\Gamma \end{aligned}$$

≤ (by Proposition VII.2.1)

≤  $j_1(y)$ .

Lemma VII.3.2. Let  $\tau_1'$  be the weak topology defined on  $BT_1(\Omega)$  by :

$$\begin{aligned}
 & \Delta q_n \rightarrow \Delta q \text{ in } L^2(\Omega) \text{ weakly} \\
 & q_n \rightarrow q \text{ in } L^1(\Gamma_0) \text{ and } L^2(\Gamma_1). \\
 q_n \rightarrow q \text{ for } \tau_1' \text{ if } & \begin{aligned}
 & \partial q_n / \partial r \rightarrow \partial q / \partial r \text{ weakly in } L^2(\Gamma_1) \\
 & \partial q_n / \partial r \rightarrow \partial q / \partial r \text{ vaguely in } M_1(\Gamma_0).
 \end{aligned}
 \end{aligned} \tag{VII.3.4}$$

Then any bounded set of  $BT_1(\Omega)$  is compact for  $\tau_1'$  ; moreover, the functional  $J_1^*$  is lower semi continuous on  $BT_1(\Omega)$  endowed with the  $\tau_1'$  topology.

The proof of Lemma VII.3.2 is obvious.

We now state the main result for Problem (VII.2.4), namely :

Theorem VII.3.3. (i)  $\text{Inf (VII.1.2)} = \text{Sup (VII.1.16)} = \text{Sup(VII.2.4)}$

(ii) Problem (VII.2.4) has a solution in  $BT_1(\Omega)$ , which is an adjoint state of the optimal state  $y_0$ .

Proof : Theorem VII.3.3 is proved exactly as was Theorem VI.2.3, using the direct method of Calculus of Variations. The use of that method is made possible by Lemmas VII.3.1 and VII.3.2.

#### 4. System of optimality conditions.

We now present the system of optimality conditions for Problem (R).

Theorem VII.4.1. Let  $y$  be an admissible state for (R) ; then  $y$  is the optimal state  $y_0$  if and only if there exists  $q$  in  $BT_1(\Omega)$ , admissible for Problem  $(S^*)$ , such that  $(y,q)$  satisfy the following equalities :

$$h_{\alpha,\beta}(q) = q \cdot (\partial y / \partial r) - 1/2 (\partial y / \partial r)^2 \text{ on } \Gamma_0 \quad (\text{VII.4.i})$$

$$h_{\alpha,\beta}(-\partial q / \partial r) = -(\partial q / \partial r) \cdot y - 1/2 (y)^2 \text{ in } M_1(\Gamma_0) \quad (\text{VII.4.ii})$$

$$\partial q / \partial r = y - y_d \text{ on } \Gamma_1 \quad (\text{VII.4.iii})$$

Moreover, Equalities (VII.4.i) to (VII.4.iii) are equivalent to the condition that  $y$  (resp.  $q$ ) should be a solution of  $(R)$  (resp.  $(S^*)$ ).

Proof : obvious, thanks to lemma VII.3.1.

### VIII. Penalization of Problem (P).

We now return to the study of Problem (P), for which we propose a family of penalized problems  $(P_\varepsilon)_{\varepsilon>0}$ . We prove the existence and unicity in  $W$  of the optimal state  $z_\varepsilon$  of  $(P_\varepsilon)$  and of its adjoint state  $p_\varepsilon$ , and give the system of optimality conditions. We then study the asymptotic behaviour of the families  $(z_\varepsilon)_{\varepsilon>0}$  and  $(p_\varepsilon)_{\varepsilon>0}$  as  $\varepsilon$  tends to zero.

#### 1. Definition and variational formulation of $(P_\varepsilon)$ .

Let  $\alpha, \beta, \varepsilon$  be three strictly positive real numbers, we set

$${}_{\alpha}^{\beta}h_{\varepsilon}(s) \begin{cases} = s^2/2, & \text{if } s \in [-\alpha, \beta] \\ = \beta(s-\beta/2) + \varepsilon/2 (s-\beta)^2, & \text{if } s \geq \beta \\ = -\alpha(s+\alpha/2) + \varepsilon/2 (s+\alpha)^2, & \text{if } s \leq -\alpha \end{cases} \quad (\text{VIII.1.1})$$

${}_{\alpha}^{\beta}h_{\varepsilon}$  is a  $C^1$  convex function that converges to  $h_{\alpha, \beta}$  (def. (I.2.9)) as  $\varepsilon$  tends to zero.

Its conjugate function is given by :

$${}_{\alpha}^{\beta}h_{\varepsilon}^*(s) \begin{cases} = s^2/2, & \text{if } s \in [-\alpha, \beta] \\ = 1/2\varepsilon (s-\beta)^2 + \beta(s-\beta/2), & \text{if } s \geq \beta \\ = 1/2\varepsilon (s+\alpha)^2 - \alpha(s+\alpha/2), & \text{if } s \leq -\alpha \end{cases} \quad (\text{VII.1.2})$$



$h_{\alpha,\beta}^*$  is a  $C^1$  convex function converging pointwise to the conjugate  $h_{\alpha,\beta}^*$  of  $h_{\alpha,\beta}$ , defined by :

$$h_{\alpha,\beta}^*(s) \begin{cases} = s^2/2, & \text{if } s \in [-\alpha, \beta] \\ = +\infty & \text{otherwise} \end{cases} \quad (\text{VIII.1.3})$$

The penalized form of Problem (P) is :

to find  $(z_\varepsilon, u_{0\varepsilon}, v_{0\varepsilon}) \in W \times L^2(\Gamma_0) \times L^2(\Gamma_0)$  minimizing the cost function

$$(P_\varepsilon) \quad J_\varepsilon(z, u, v) = 1/2 \|z - z_d\|_{L^2(\Omega)}^2 + \int_{\Gamma_0} h_{\alpha,\beta}^*(u) d\Gamma + \int_{\Gamma_0} h_{\alpha,\beta}^*(v) d\Gamma \\ + 1/2\varepsilon \|\Delta z\|_{L^2(\Omega)}^2.$$

the infimum being taken for  $z \in W$ ,  $z = u$  on  $\Gamma_0$ ,  $\partial z / \partial r = v$  on  $\Gamma_0$  ( $z_d$  as in Section I).

Remark VIII.1.1  $(P_\varepsilon)$  is a problem without constraints.

Let us now give the variational formulation of  $(P_\varepsilon)$ .

In order to do so, we define on  $W$  the functional :

$$j_\varepsilon(z) = 1/2 \|z - z_d\|_{L^2(\Omega)}^2 + 1/2\varepsilon \|\Delta z\|_{L^2(\Omega)}^2 + \int_{\Gamma_0} h_{\alpha,\beta}^*(\partial z / \partial r) d\Gamma \\ + \int_{\Gamma_0} h_{\alpha,\beta}^*(z) d\Gamma \quad (\text{VIII.1.5})$$

and  $(P_\varepsilon)$  is equivalent to the minimization problem :

$$(P_\varepsilon) \quad \inf_{z \in W} \{j_\varepsilon(z)\} \quad (\text{VIII.1.6})$$

We have the following result [E.T]:

Proposition VIII.1. There exists a unique optimal state  $z_\varepsilon$  for  $(P_\varepsilon)$ .

The dual problem  $(P_\varepsilon^*)$  of  $(P_\varepsilon)$  is determined as in Section I, and its expression is

$$(P_\varepsilon^*) \quad \sup_{p \in W} \left\{ -\varepsilon/2 \|p\|_{L^2(\Omega)}^2 - 1/2 \|\Delta p\|_{L^2(\Omega)}^2 - \int_{\Omega} \Delta p \cdot z_\varepsilon dx \right. \\ \left. - \int_{\Gamma_0} \alpha^0 h_\varepsilon(-\partial p / \partial r) d\Gamma - \int_{\Gamma_0} \alpha^0 h_\varepsilon(p) d\Gamma \right\} \quad (\text{VIII.1.7})$$

$$p = \partial p / \partial r = 0 \quad \text{on } \Gamma_1$$

Thanks to the definition of  $\alpha^0 h_\varepsilon$ ,  $(P_\varepsilon^*)$  is a coercive problem on  $W$ ; hence, we can state:

Proposition VIII.1.1. There exists a unique adjoint state  $p_\varepsilon$  for the optimal state  $z_\varepsilon$ .

Moreover, the set of optimality conditions for  $z_\varepsilon$  and  $p_\varepsilon$  is easily obtained ([E.T], Th. 4.2 and Rem. 4.2), and is summed up in the following proposition:

Proposition VIII.1.2. An element  $z$  of  $w$  is an optimal state for  $(P_\varepsilon)$  if and only if there exists  $p$  in  $w$ , such that  $(z,p)$  satisfy the equalities :

$$\Delta p = z - z_d \text{ in } \Omega ; \quad (\text{VIII.1.i})$$

$$p = -1/\varepsilon \Delta z \text{ in } \Omega ; \quad (\text{VIII.1.ii})$$

$$\alpha' h_\varepsilon (-\partial p / \partial r) = -\partial p / \partial r \cdot z - \alpha' h_\varepsilon^*(z) \text{ on } \Gamma_0 \quad (\text{VIII.1.iii})$$

$$\alpha' h_\varepsilon(p) = p \cdot \partial z / \partial r - \alpha' h_\varepsilon^*(\partial z / \partial r) \text{ on } \Gamma_0 \quad (\text{VIII.1.iv})$$

Moreover, Equalities (VII.1.i) to (VIII.1.iv) are equivalent to the condition that  $z$  (resp.  $p$ ) should be equal to the solution  $z_\varepsilon$  (resp.  $p_\varepsilon$ ) of  $(P_\varepsilon)$  (resp.  $(P_\varepsilon^*)$ ).

## 2. Asymptotic behaviour of the optimal state $z_\varepsilon$ .

We now prove the :

Theorem VIII.2.1. The family  $\{z_\varepsilon\}_{\varepsilon > 0}$  tends to the optimal state  $z_0$  of  $(P)$  for the strong topology of  $W$  as  $\varepsilon$  tends to zero.

Proof :

We first notice that the family  $\{j_\varepsilon(z_\varepsilon)\}_{\varepsilon > 0}$  is decreasing and satisfies :

$$j(z_0) = j_\varepsilon(z_0) \geq j_\varepsilon(z_\varepsilon) \quad (\text{VIII.2.1})$$

hence, there exists a number  $\zeta$  with  $\zeta = \lim_{\varepsilon \rightarrow 0} (j_\varepsilon(z_\varepsilon))$ .

Thanks to (VIII.2.1),  $z_\varepsilon$  is bounded in  $W$ , and also satisfies :

$$\lim_{\varepsilon \rightarrow 0} \|\Delta z_\varepsilon\|_{L^2(\Omega)}^2 = 0 \quad (\text{VI.2.2})$$

We now extract a subsequence still denoted by  $z_\varepsilon$  which converges weakly in  $W$  towards an element  $z'_0$  of  $W$ . Let us now show that  $z'_0$  is admissible for problem (I.1.3).

First of all, we obviously have :

$$\Delta z'_0 = 0 \text{ in } \Omega \quad (\text{VIII.2.3})$$

Let us choose a fixed  $\eta$ ; we have :

$$\int_{\Gamma_0} \beta_\alpha h_\eta^*(z'_0) \leq \liminf_{\varepsilon \rightarrow 0} \int_{\Gamma_0} \beta_\alpha h_\eta^*(z_\varepsilon) d\Gamma \quad (\text{VIII.2.4})$$

thanks to the weak convergence of  $z_\varepsilon$  towards  $z'_0$ . But  $\beta_\alpha h_\varepsilon^*$  is a decreasing function of  $\varepsilon$ , and therefore we obtain from (VIII.2.4) :

$$\int_{\Gamma_0} \beta_\alpha h_\eta^*(z'_0) \leq \liminf_{\varepsilon \rightarrow 0} \int_{\Gamma_0} \beta_\alpha h_\varepsilon^*(z_\varepsilon) d\Gamma \quad (\text{VIII.2.5})$$

The sequence  $(\int_{\Gamma_0} \beta_\alpha h_\eta^*(z'_0) d\Gamma)$  is then bounded, and this necessarily implies that  $z'_0$  belongs to  $\mathcal{U}_{\alpha,\beta}$ ; we prove in the same way that  $\partial z'_0 / \partial r$  belongs to  $\mathcal{U}'_{\alpha,\beta}$ , and  $z'_0$  is therefore admissible for problem (I.1.3).

By lower semi continuity we have :

$$\begin{aligned} j(z'_0) &\leq \underline{\lim}_{\varepsilon \rightarrow 0} (1/2 \|z_\varepsilon - z_d\|_{L^2(\Omega)}^2 + 1/2 \|\partial z_\varepsilon / \partial r\|_{L^2(\Gamma_0)}^2 \\ &\quad + 1/2 \|z_\varepsilon\|_{L^2(\Gamma_0)}^2) \\ &\leq \underline{\lim}_{\varepsilon \rightarrow 0} (j_\varepsilon(z_\varepsilon)) \end{aligned}$$

and, using (VIII.2.1) :

$$j(z'_0) \leq \underline{\lim}_{\varepsilon \rightarrow 0} j_\varepsilon(z_\varepsilon) \leq \lim_{\varepsilon \rightarrow 0} j_\varepsilon(z_\varepsilon) \leq j(z_0) \quad (\text{VIII.2.6})$$

so that we finally obtain :

$$j(z'_0) = \lim_{\varepsilon \rightarrow 0} j_\varepsilon(z_\varepsilon) = j(z_0) \quad (\text{VIII.2.7})$$

Thus,  $z'_0$  and  $z_0$  are equal ; let us now prove the strong convergence of  $z_\varepsilon$  to  $z_0$ . We set :

$$\begin{aligned} H(z_0, z_\varepsilon) &= j(z_0) - 1/2 \|z_\varepsilon - z_d\|_{L^2(\Omega)}^2 - 1/2 \|z_\varepsilon\|_{L^2(\Gamma_0)}^2 \\ &\quad - 1/2 \|\partial z_\varepsilon / \partial r\|_{L^2(\Gamma_0)}^2 \end{aligned} \quad (\text{VIII.2.8})$$

$H(z_0, z_\varepsilon)$  is positive, and tends to zero with  $\varepsilon$ . Owing to the weak convergence of  $z_\varepsilon$  to  $z_0$ , we have :

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} (\|z_0\|_{L^2(\Omega)}^2 - \|z_\varepsilon\|_{L^2(\Omega)}^2) &= \lim_{\varepsilon \rightarrow 0} (\|z_0\|_{L^2(\Gamma_0)}^2 - \|z_\varepsilon\|_{L^2(\Gamma_0)}^2) \\ &= \lim_{\varepsilon \rightarrow 0} (\|\partial z_0 / \partial r\|_{L^2(\Gamma_0)}^2 - \|\partial z_\varepsilon / \partial r\|_{L^2(\Gamma_0)}^2) = 0 \end{aligned} \quad (\text{VIII.2.9})$$

and (VIII.2.9), together with the weak convergence of  $z_\varepsilon$ , is obviously enough to prove the strong convergence of  $z_\varepsilon$  towards  $z_0$ , for  $W$  is a Hilbert space.

Finally, we conclude by an easy contradiction argument that the whole family  $(z_\varepsilon)_{\varepsilon > 0}$  converges to  $z_0$ , and Theorem VIII.2.1. is proved.

### 3. Asymptotic behaviour of the adjoint state $p_\varepsilon$ .

Theorem VIII.3.1. The family  $\{p_\varepsilon\}_{\varepsilon > 0}$  has a cluster point in  $BT(\Omega)$  which is an adjoint state of the optimal state  $z_0$  of (P).

Proof : Thanks to the equality  $\text{Inf}(\text{VIII.1.6}) = \text{Sup}(\text{VIII.1.7})$ , we know that  $\{p_\varepsilon\}_{\varepsilon > 0}$  is bounded in  $BT(\Omega)$ , and therefore compact for the  $\tau_1$  topology.

Let  $w$  be a cluster point of this family for  $\tau_1$ ; the lower semi continuity of  $J^*$  ensures that  $w$  is a solution of Problem (V.1.3).

**BIBLIOGRAPHY**

- [A] R. ADAMS : Sobolev spaces, Academic press, 1975.
- [B] N. BOURBAKI : Integration, ch. 1.4, 2nd édition, Hermann, 1965.
- [BGM] M. BERGER, P. GAUDUCHON, E. MAZET : Spectre d'une variété riemannienne, Springer, 1971.
- [D] J. DIEUDONNE : Eléments d'Analyse, Vol. 4, Gauthier Villars, 1971.
- [DT 1] F. DEMENGEL, R. TEMAM : (1) Convex functions of a measure and applications, Indiana University Math. Journal, # 33, 1984.
- [DT 2] F. DEMENGEL, R. TEMAM : (2) Convex function of a measure and applications : the unbounded case (to appear).
- [ET] I. EKELAND, R. TEMAM : Analyse convexe et problèmes variationnels, Dunod, 1972.
- [GG] G. GEYMONAT, P. GRISVARD : Problèmes aux limites elliptiques dans  $L^\alpha$ , Université d'Orsay, 1964.
- [K] H. KARCHER : Riemannian center of mass and mollifier smoothing, CPAM, # 30, 1977.
- [KT] R. KOHN, R. TEMAM : Dual spaces of stresses and strains, Applied Maths and Optimization, Springer, 1983.
- [L] J.L. LIONS : Some methods in the mathematical analysis of systems and their control, Gordon Breach, 1981.
- [L 1] J.L. LIONS : Problèmes aux limites dans les équations aux dérivées partielles, Presses de l'Université de Montréal, 1982.
- [L 2] J.L. LIONS : Perturbations singulières dans les problèmes aux limites et en contrôle optimal, Springer, 1973.

- [LM] J.L. LIONS, E. MAGENES : Problèmes aux limites non homogènes et applications, vol. 1, Dunod, 1968.
- [M1] C. MIRANDA : Partial differential equations of elliptic type, 2nd edition, Springer, 1970.
- [M2] C. MIRANDA : Sulle proprietà di regolarità di certe trasformazioni integrali, Mem. Acad. Lincei, #7, 1963-1966.
- [Micr] S. MINAKSHISUNDARAN : Eigenfunctions on riemannian manifolds, J. Indian. Math. Soc., # 17, 1963.
- [MKS] H.P. Mc KEAN, I.M. SINGER : Curvature and the eigenvalues of the Laplacian, J. of diff. geometry, # 1, 1967.
- [S] S. SOBOLEV : Sur les équations aux dérivées partielles hyperboliques non linéaires, Edizioni Cremonese, 1961.
- [Sch] L. SCHWARTZ : Théorie des distributions, Hermann, 1966.
- [T] R. TEMAM : Problèmes mathématiques en plasticité, Gauthier Villars, 1983.