



HAL
open science

Lifting Algebraic Reasoning to Generalized Metric Spaces

Ralph Sarkis

► **To cite this version:**

Ralph Sarkis. Lifting Algebraic Reasoning to Generalized Metric Spaces. Logic in Computer Science [cs.LO]. Ecole normale supérieure de lyon - ENS LYON, 2024. English. NNT : 2024ENSL0025 . tel-04717268


HAL Id: tel-04717268

<https://theses.hal.science/tel-04717268v1>

Submitted on 1 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



THÈSE
en vue de l'obtention du grade de Docteur, délivré par
l'ÉCOLE NORMALE SUPÉRIEURE DE LYON

École Doctorale N°512
École Doctorale en Informatique et Mathématique de Lyon

Discipline : Informatique

Soutenue publiquement le 03/09/2024, par :

Ralph Sarkis

Lifting Algebraic Reasoning to Generalized Metric Spaces

Relèvement du raisonnement algébrique aux espaces métriques généralisés

Devant le jury composé de :

ADÁMEK, Jiří	Professeur Czech Technical University in Prague	Rapporteur
PLOTKIN, Gordon	Professeur University of Edinburgh	Rapporteur
B. B. LUCYSHYN-WRIGHT, Rory	Professeur associé Brandon University	Examineur
TASSON, Christine	Professeure ISAE Supareo	Examinatrice
MIO, Matteo	Chargé de recherche CNRS & ENS de Lyon	Directeur de thèse
VIGNUDELLI, Valeria	Chargée de recherche CNRS & ENS de Lyon	Examinatrice

Abstract

Algebraic reasoning is ubiquitous in mathematics and computer science, and it has been generalized to many different settings. In 2016, Mardare, Panangaden, and Plotkin introduced quantitative algebras, that is, metric spaces equipped with operations that are nonexpansive relative to the metric. They proved counterparts to important results in universal algebra, and in particular they provided a sound and complete deduction system generalizing Birkhoff's equational logic by replacing equality with equality up to ε . This allowed them to give algebraic axiomatizations for several important metrics like the Hausdorff and Kantorovich distances.

In this thesis, we make two modifications to Mardare et al.'s framework. First, we replace metrics with a more general notion that captures pseudometrics, partial orders, probabilistic metrics, and more. Second, we do not require the operations in a quantitative algebra to be nonexpansive. We provide a sound and complete deduction system, we construct free quantitative algebras, and we demonstrate the value of our generalization by proving that any monad on generalized metric spaces that lifts a monad on sets can be presented with a quantitative algebraic theory. We apply this last result to obtain an axiomatization for the Łukaszyk–Karmowski distance.

Résumé

On retrouve le raisonnement algébrique partout en mathématique et en informatique, et il a déjà été généralisé à pleins de contextes différents. En 2016, Mardare, Panangaden et Plotkin ont introduit les algèbres quantitatives, c'est-à-dire, des espaces métriques équipés d'opérations 1-lipschitzienne relativement à la métrique. Ils ont prouvées des homologues à des résultats importants en algèbre universelle, et en particulier ils ont donné un système de deduction correct et complet qui généralise la logique équationnelle de Birkhoff en remplaçant l'égalité par l'égalité à ε près. Ça leur a permis de donner une axiomatisation algébrique pour quelques métriques importantes comme la distance de Hausdorff et celle de Kantorovich.

Dans cette thèse, on modifie deux aspects du cadre de Mardare et al. Premièrement, on remplace les métriques par une notion plus générale qui englobe les pseudométriques, les ordres partiels, les métriques probabilistes, entre autres. Deuxièmement, on n'exige pas que les opérations de nos algèbres quantitatives soient lipshitzienne. On donne un système de deduction correct et complet, on construit les algèbres quantitatives libres, et on démontre la valeur de notre généralisation en prouvant que toute monade sur les espaces métriques généralisés qui est le relèvement d'une monade finitaire sur les ensembles peut être présentée par une théorie algébrique quantitative. On applique ce dernier résultat pour obtenir une axiomatisation de la distance de Łukaszyk–Karmowski.

Contents

Preface	5
0 Introduction	9
0.1 Universal Algebra and Monads	13
0.2 Generalized Metric Spaces	14
0.3 Universal Quantitative Algebra	17
1 Universal Algebra	21
1.1 Algebras	21
1.2 Terms and Equations	24
1.3 Free Algebras	34
1.4 Equational Logic	45
1.5 Monads	49
2 Generalized Metric Spaces	60
2.1 L-Spaces	60
2.2 Equational Constraints	70
2.3 The Categories GMet	77
3 Universal Quantitative Algebra	88
3.1 Quantitative Algebras	88
3.2 Free Quantitative Algebras	99
3.3 Quantitative Equational Logic	118
3.4 Quantitative Alg. Presentations	125
3.5 Lifting Presentations	129
4 Conclusion	140
4.1 Future Work	140
Bibliography	144

Preface

Tamacun

Rodrigo y Gabriela

In place of the traditional citations as epigraphs at the start of every chapter, I put (links to) music I enjoyed listening to while writing this manuscript.

This document was not optimized for printing. The two main reasons are:

1. I use a slightly customized version of the [tufte-book](#) document class [KWG15]. This puts the main body of text closer to the left margin and all footnotes^o in the right margin. This allows me to use a lot of footnotes throughout the text. I use them as if they were big parentheses, to add details, to digress, to add references, or to display diagrams. Printing with these margins can be complicated, and the text in the margins is a bit smaller.
2. I use the **knowldege** package [Col24]. This allows me to easily add hyperlinks towards the definition of a symbol or a term every time I use that symbol or term. In particular, if you want to start reading at say Chapter 3, you do not have to go over the notation introduced earlier, you can simply click on a symbol or word you don't recognize to see how it was defined. What is more, in the appendix¹, I put a draft of a book on category theory that I am writing, so there is no background section on categories, but every time I use a notion from that book (e.g. $\text{Hom}_{\mathbb{C}}(A, B)$, functor, natural transformation), the **knowldege** link will go there.² Combined with the links to results, equations, and references (like Theorem 3.80, (3.13), and [MPP16]) there are more than twenty thousand links in this document!

^o Like this one.

¹ Available at <https://ralphs16.github.io/src/Manuscript.pdf>.

² Test the links right now! Some PDF viewers are better than others to navigate a document with lots of links. Most have a navigation history so you can follow a sequence of links and get back to your original position by e.g. pressing ALT+P or the back button on a mouse. Some viewers also display a preview of the target of a hyperlink when you hover it, so there is no need to click.

With that said, if you would rather read on paper, I do not think there will be major difficulties since there is adequate numbering throughout the main text. However, I suggest you do not print the appendix (which is longer than the main text) because the links to the appendix are rarely numbered, and I did not make an index.

Notations and Conventions

Here are several standard and non-standard notations and conventions that I use throughout the text.

- Starting now, we will use the pronoun “we” when referring to us, author and readers. Occasionally, “I” will be used to refer to me (Ralph), and “we” will be used to refer to me and my supervisors Matteo and Valeria.

- We use the following abbreviations:³
 - **I.H.** indicates a step of a proof that relies on the induction hypothesis (which is often left implicit).
 - **resp.** stands for “respectively”.
 - **L.H.S.** stands for “left-hand side” (of an equation usually).
 - **R.H.S.** stands for “right-hand side”.
- When defining a function $f : A/\sim \rightarrow B$, whose domain is a quotient, by giving a value for $f(a)$ for each $a \in A$, we say it is **well-defined** if $f(a) = f(a')$ whenever $a \sim a'$.
- We sometimes have to deal with **proper classes** (see, e.g., [AHS06, §2.2]), i.e. collections of things that cannot be sets. We use classes to mean a collection that is either a set or a proper class.⁴
- We use the term **classical** to refer to universal algebra (the subject of Chapter 1), usually in opposition to universal quantitative algebra (the subject of Chapter 3).
- The two principal references [MPP16, FMS21] are given special colors to help to recognize them when reading.

³ Paired with a knowledge link going to this list.

⁴ Really the only reason we need classes is for the collection of all sets, so nothing very fancy.

Acknowledgements — Remerciements

J’ai la chance d’avoir des gens formidables dans ma vie sans qui cette thèse n’aurait pas vu le jour. Je les remercie pour leur soutien.

I am lucky to have wonderful people in my life without whom this thesis would not have seen the light of day. I thank them for their support.

To Matteo Mio, I could not have asked for a better initiation into research work than being your student. You shared your enthusiasm for asking seemingly innocuous questions and finding invaluable insights in their answers. You gave me down-to-earth advice on many subjects to hone my scientific skills. You made me feel good about sharing my ideas and my work. I believe that, thanks to your work philosophy, many hardships of a PhD student’s life have been mitigated.

To Valeria Vignudelli, your considerable involvement in the supervision of my thesis is greatly appreciated. You shared your experience in very helpful ways, at very helpful times. Your rigorous and thorough evaluation of my and other people’s work is an example I aspire to follow.

To Matteo and Valeria, I had so much fun working with you. I will fondly remember all the Mondays spent hacking away for hours on a white board that actively fights back with electric shocks until we reached near starvation.

To Jiří Adámek, Rory Lucyshyn-Wright, Gordon Plotkin, and Christine Tasson, for kindly agreeing to participate in my jury to review and evaluate my work. I admire your scientific achievements, and my ambition to make a good impression on you has surely improved the final version of this manuscript, with help from the reviewers’ insightful comments and questions.

Ma thèse a été financée par un contrat doctoral spécifique normalien.

To Prakash Panangaden, you have had a grand positive compounded impact on my life as a researcher. You showed me it is possible to do category theory, logic, and computer science at the same time; you recommended my wonderful supervisors; you hosted me for a semester of work closer to my family in Montréal; and you invited me to a fantastic workshop in Bellairs on the subject of my thesis (which also originates from a paper you coauthored). I am delighted to have been your student.

Aux membres de Plume, toutes nos discussions aux GDTs, aux Plume Beers, et à CHoCoLa ont fait grandir en moi un sentiment d'appartenance à notre communauté scientifique. Je remercie plus particulièrement Daniel Hirschhoff pour sa sympathie contagieuse; Marie Narducci pour sa bonne humeur infaillible; Russ Harmer pour son implication en tant que superviseur officiel temporaire; Alexandre Goy pour notre amitié express en début de doctorat; et Colin Riba et Michele Pagani pour leur confiance en moi pour leurs TDs. J'espère pouvoir entretenir une ambiance de travail comparable à celle de Plume dans mes prochaines missions.

To the members of my wider scientific community, all the people I have met, all the events I have attended, and all the papers and messages I have read have made me confident in my dream to pursue life as an academic. I particularly thank Fabio Zanasi for a research visit that kickstarted work towards a common objective; the participants and organizers of the 2022 ACT adjoint school for the opportunity to work closely with other young researchers; Tom Hirschowitz and Marie Kerjean for inviting me to present my work; and the CT Zulipchat for producing a constant stream of interesting discussions.

Aux étudiant-es des séminaires de théorie des catégories, votre enthousiasme pour nos cours m'a rendu fier d'enseigner cette matière qui me tient à cœur. L'écoute et les retours de cell-eux qui ont enseigné à leur tour avec mes conseils m'ont permis d'améliorer chaque année mes méthodes et mon bouquin.

Aux gens qui m'ont permis de faire de la médiation, grâce à vous, j'ai eu accès à un public stimulant et j'ai pu faire preuve de créativité dans ma pédagogie. Je remercie plus particulièrement les belles âmes de la MMI, Charlotte Avellaneda, Camille Beaudou, Olivier Druet, et Nina Gasking, pour les repas animés avant ou après mes interventions; et l'association Un Peu de Bon Science pour mes trois stages de recherches super enrichissants.

To the creators and contributors to open source software that I used to automate and accelerate the typesetting of this document: [L^AT_EX](#), the `knowldege` package, the `tufte-book` document class, the GUI commutative diagram editors [quiver](#) and [tikzcd-editor](#), and [VSCodium](#).

À mes ami-es de Lyon et de Montréal, que ce soit autour d'un filet de Spikeball, d'une table de dîner, d'un plateau de jeu, ou d'un verre au bar, j'ai toujours été en merveilleuse compagnie. Je remercie plus particulièrement Valentin avec qui mon amitié n'a pas faibli même après des années que j'ai passées outre-Atlantique; Nina qui partage mon amour pour la bouffe épicée et végane; et Alaa avec qui on discute du bon et du mauvais dans la culture libanaise. Tant de beaux moments avec tant de belles personnes.

À mon papa, je t'aime. Tu m'as transmis une curiosité insatiable, un pouvoir

précieux de relativisation, et une appréciation pour la routine et la discipline. Ces qualités font que j'ai passé ces trois dernières années à véritablement m'amuser en passant l'épreuve du doctorat. Les taboulés et chammehs illimités me manquent déjà.

À ma maman, je t'aime. Tu m'as transmis une curiosité insatiable, un amour pour la bonne bouffe et la cuisine, une ouverture d'esprit, et un mélange de culture fusionnel. Grâce à tous tes efforts pour rendre ma vie heureuse et facile, je ne serais jamais aussi résilient que toi. J'ai quand même hérité d'assez de ton caractère pour finir cette thèse indemne. C'est difficile parfois d'aller me coucher à 22h22 sans te faire de câlins.

À mon frère, je t'aime. Tu es la personne qui me comprend le mieux et tu le resteras toute notre vie. J'ai eu la chance de vivre avec toi pendant quelques mois durant ce doctorat (à Montréal et à Lyon), et ce furent de loin les meilleurs. Pour les ramens, les séries et vidéos YouTube bingées, les parties de squash, les questions absurdes, les discussions philosophiques, les inside jokes, et les souvenirs de notre enfance, je ne peux pas ne pas t'aimer.

À toute ma famille, mon frère, mes parents, mes grand-parents, mes tantes, mes oncles, et mes cousin-es, je vous aime. Vous m'avez tant aimé et maintenant que ce manuscrit est fini, ça m'attriste de voir que tout l'amour que je porte pour ma discipline n'est pas dirigé vers vous. Vous le méritez. Je pense que ce refrain de Beau Dommage est gravé dans mes pensées.

Ça ne vaut pas la peine
De laisser ceux qu'on aime
Pour aller faire tourner
Des ballons sur son nez.

0 Introduction

[Across the Stars](#)

John Williams and the London Symphony Orchestra

Most programmers write code **compositionally**.⁵ They write small lines of code that combine to make small functions that combine to make small files that combine to make a complete software. When studying the semantics of programs, we sometimes like to model these *combination* steps with algebraic operations.

This idea seems to originate in [SS71] and [GTWW77], and it continues to reverberate in current research, e.g. [TP97, HHL22, GMS⁺23]. It is referred to as **algebraic semantics**. We give only an informal account here to motivate the mathematics behind it.

If P , Q and Q' are programs, we can use $P;Q$ to represent the program that runs P then Q , and $\text{ifte}(P, Q, Q')$ to represent the program that runs P , then runs Q if the Boolean value of the output of P was True or Q' if it was False. We view the set of programs as an algebra where instead of the well-known operations like addition and multiplication, many new operations are allowed to combine programs. The set of available operations varies with the kind of programs that are studied, it is called the signature, and we say that operations in the signature are interpreted in the algebra of programs.

Furthermore, the set of behaviors of programs⁶ is also seen as an algebra for the same signature. Then, semantics is represented by a function from programs to behaviors which preserves the operations, namely, the combination of behaviors is the behavior of the combination. It is a homomorphism of algebras.

Oftentimes, one realizes that two different programs have the same behavior, for example $P;(Q;R)$ and $(P;Q);R$ or $P;Q$ and $\text{ifte}(P, Q, Q)$, so they should be considered **equal** (or **equivalent**). The bread and butter of algebraic semanticists is to find a (sound and complete) collection of simple equations (axioms) that make it possible to reason compositionally about program equivalence.⁷ Sometimes these axiomatizations help in designing (semi)-automatic procedures to answer the question “is P equal to Q ?”.

A famous example is combinatory logic, originating in [Cur29], which gives a computational model as powerful as the pure λ -calculus using four operations to combine programs and three equations between small programs.⁸ In this thesis, we detail two other well-known examples that model nondeterministic and probabilistic choices in Examples 1.78 and 1.79 respectively.

0.1 Universal Algebra and Monads	13
0.2 Generalized Metric Spaces	14
0.3 Universal Quantitative Algebra	17

⁵ Some don't (e.g. [code golfers](#)).

⁶ The word behavior can be understood in many ways that depend on what properties of the programs one is interested in.

⁷ For instance, with the equations above, we can infer that $\text{ifte}(P, Q; (R; S), (Q; R); S)$ and $P; (Q; (R; S))$ are equivalent.

⁸ see, e.g. [Mim20, §3.6.3].

Much of the work on algebraic semantics relies on the theoretical foundations of **universal algebra**, an old subject popularized by Birkhoff in [Bir33, Bir35]. Three of his major results are:

1. a logical system, called equational logic (Figure 1.3), that allows one to syntactically derive which equations are entailed by a set of axioms,
2. the construction of free algebras (Definition 1.34 and Proposition 1.49), and
3. the HSP (or variety) theorem [Wec92, §3.2, Theorem 21] which characterizes classes of algebras that can be defined with equations.

There is also tight connection between universal algebra and **monads on Set** (Definition 1.62) that can be exploited to study semantics with algebraic and categorical reasoning. For instance, nondeterminism can be modelled with the theory of semilattices and the powerset monad (Example 1.78), and probability can be modelled with the theory of convex algebras and the distribution monad (Example 1.79). These two examples and similar ones show up very often in the study of program semantics.⁹

⁹ See, e.g. [PPo1a, PPo1b, PPo2, BP15, BSS21, BSV22].

Since computers interact with humans (or the other way around), it makes sense to take into account the quirks of a human mind when studying the behavior of programs. For example, many standard data compression algorithms (in particular for image, audio, and video) are efficient at the cost of losing some small amount of information.¹⁰ In that situation (and others like it), program equivalence is too coarse of a relation, so researchers have to build more sensitive models to handle and compare **approximations** of programs.

¹⁰ Usually, users will not notice nor mind because of the inherent information degradation in the human perception process [SB06].

This makes the case for developing **quantitative algebraic semantics**. We view the set of programs as an algebra (we can still combine them) with a notion of distance (we can now compare them more finely than with equality). Intuitively, the distance between P and Q shall reflect the disparity in their behaviors, hence, the behaviors must come with a notion of distance too. For example, if P is a lossless compression algorithm, and Q is a lossy one, the distance between P and Q may be the fraction of the inputs (picked in a real-world dataset) wherein the outputs of P and Q noticeably differ.¹¹

¹¹ For metrics actually used in practice, see [LJ11].

We most commonly think of a distance as a number, but our formalization of distances (Definitions 2.11 and 2.32) will accommodate a large array of things to call distances, see Examples 2.13, 2.14, and 2.16.

If the field of algebraic semantics finds itself on universal algebra, there needs to be a quantitative version of this theoretical basis to support research in quantitative algebraic semantics.

The concept of extending algebraic reasoning to diverse settings is by no means novel, as evidenced by the following (inevitably) non-exhaustive list of references: [Ben68, Dub70, Gra75, BD80, Die80, Bur81, Bur82, BB92, KP93, Wea93, GP98, Pow99, Rob02, BV05, HP06, NP09, LP09, VK11, FH11, LR11, BBvT12, AMMU15, LW16, GP18, MU19, BG19, FMS21, Ros21, LP23, RT23, Ros24]. While these approaches excel in their generality and abstraction, it is at the cost of usability, even for

someone who is already familiar with universal algebra. More concrete solutions exist. We mention two that seem to be of particular interest to computer scientists.

If we equip the algebra of programs with a partial order, the question “is P equal to Q ?” becomes “is P less than Q ?”.¹² There is already a lot of work in universal algebra on partial orders [Blo76, ANR85, KV17, AFMS21, FMS21, ADV22, Sch22a, Sch22b].

If we equip the algebra of programs with a metric space, the question “is P equal to Q ?” becomes “are P and Q closer than ε from each other?”, where ε is a real number. There is already a lot of work in universal algebra on metric spaces [Wea95, MPP16, Hin16, MPP17, BMPP18, MPP18, MV20, BMPP21, MPP21, Ros21, MSV21, Adá22, MSV22, MSV23, ADV23b, Ros24, ?].

In this thesis, we make another attempt to generalize algebraic reasoning without straying too far from the classical setting. Our main inspirations are [MPP16], the seminal paper on quantitative algebras, and [FMS21], a vast generalization.¹³

In [MPP16], the authors study algebras equipped with a metric such that the interpretation of operations in the signature are nonexpansive. More precisely, they are metric spaces (A, d) with, for each n -ary operation op in the signature, an interpretation $\llbracket \text{op} \rrbracket : A^n \rightarrow A$ satisfying

$$\forall a, b \in A^n, d(\llbracket \text{op} \rrbracket(a_1, \dots, a_n), \llbracket \text{op} \rrbracket(b_1, \dots, b_n)) \leq \max_{1 \leq i \leq n} d(a_i, b_i). \quad (0.1)$$

This is a very natural condition because it is equivalent to saying that $\llbracket \text{op} \rrbracket$ is a morphism from $(A, d)^n$ to (A, d) in the category **Met** of metric spaces and nonexpansive maps, where $(A, d)^n$ denotes the n -wise categorical product.¹⁴

In [FMS21], the authors view **Met** as an instance of a category $\mathbf{Str}(\mathcal{S})$ of relational structures, see [FMS21, Example 3.5.(3)]. Without going into details, we can mention that the category **Poset** of partially ordered sets and monotone maps is another instance. Therefore, their work is general enough to cover both algebras equipped with a metric and algebras equipped with a partial order. Accordingly, a generalization of (0.1) is imposed on the interpretation of operations, namely, $\llbracket \text{op} \rrbracket$ is a morphism from A^n to A , where $A \in \mathbf{Str}(\mathcal{S})$.¹⁵

In both papers, there is a sound and complete logical system that generalizes Birkhoff’s equational logic, [MPP16] replaces equations with *quantitative inferences* and [FMS21] replaces equations with Σ -relations (where Σ is the signature). An explicit construction of free algebras equipped with a metric (resp. a relational structure) is given in [MPP16, Theorem 5.3] (resp. [FMS21, Theorem 4.18]). Later papers provided generalizations of the HSP theorem [MPP17, MU19, JMU24], and the connection with monads has been investigated in [FMS21, Adá22, ADV23b].

In [MPP16, §8–10], the authors use their logic to axiomatize well-known constructions on metrics. They show that the total variation distance (Example 3.92), the Kantorovich distance (Example 3.5), and the Hausdorff distance (Example 2.17) can all be defined as free algebras for some carefully chosen set of axioms. Ford et al. do the same for the metric completion in [FMS21, Example 4.8]. Many other so-called presentation results are found in, e.g. [MV20, BMPP21, MSV21, MSV22, ?], sometimes with applications to semantics.

¹² The meaning of $P \leq Q$ depends on what kind of programs and properties are studied.

¹³ I gave their references special colors to help recognize them when reading.

¹⁴ I would say this is the expected definition of “algebra over a metric space”, especially to those familiar with functorial semantics [Law63], or subsequent work in categorical algebra and categorical logic.

¹⁵ It is actually more complicated than that, because in [FMS21], operations come with an arity $\text{ar}(\text{op})$ that is not just a natural number but a whole relational structure itself (with some size conditions). This allows them to handle some *partial* operations, e.g. $x + y$ can be undefined.

While trying to axiomatize other interesting distances, we had to question some assumptions of [MPP16]. We learned about the ŁK distance (3.3) on probability distributions in [CKPR21], where they use it as an easier-to-compute alternative to the Kantorovich distance. We intended to simply adapt that axiomatization, but we quickly faced two obstacles.

First, the ŁK distance is not a metric, it is a diffuse metric [CKPR21, §4.2].¹⁶ In particular, the distance between a distribution and itself can be non-zero. Second, combining probability distributions like it is done for the Kantorovich distance (with convex combinations) is not nonexpansive in the sense of (0.1) with d being the ŁK distance.

The generality of [FMS21] is enough to overcome the first problem since the category of diffuse metrics is an instance of $\mathbf{Str}(\mathcal{K})$. However, we already said that they also work with (an analog to) the requirement of (0.1), so the second problem remains. In the present work, we introduce a framework that deals with both 1) distances that are not metrics and 2) operations that do not satisfy (0.1).¹⁷ Rejecting that assumption was previously done in [Wea93, Wea95, Hin16, Hin17, BBLM18a, AFMS21] in various different contexts.

We define generalized metrics to be distance functions valued inside an arbitrary complete lattice L ($d : A \times A \rightarrow L$) satisfying an arbitrary set of axioms expressed with quantitative equations (a variant of the quantitative inferences in [MPP16]).¹⁸ Then, our quantitative algebras (Definition 3.1) are simply algebras equipped with a generalized metric. Importantly, no further restriction is imposed on the operations in the algebra, and this allows us to axiomatize the ŁK distance in Example 3.102.

With this setting, we recover some of the classical results in universal algebra, and more. The major contributions, Items i., ii., and iv., already appear in [MSV23] with a different presentation and a fixed $L = [0, 1]$.

- i. We define quantitative equational logic (Figure 3.1), a logical system that is sound (Theorem 3.69) and complete (Theorem 3.76) relative to our quantitative algebras. It mirrors equational logic more closely than Mardare et al.'s logic¹⁹ without renouncing their fundamental idea to *merely* change equality with equality up to ε .
- ii. We construct the free quantitative algebras (Theorem 3.57) relative to any class of quantitative equations.²⁰ This induces a monad on the category \mathbf{GMet} of generalized metric space, and the quantitative algebras modelling the chosen class of quantitative equations coincide with the algebras for that monad (Theorem 3.80).
- iii. We provide a simple axiomatization of the set of probability distributions with the ŁK distance as a free quantitative algebra in Example 3.102.
- iv. In achieving Item iii., we prove a more general result (Theorem 3.98) which states that any monad lifting to \mathbf{GMet} (Definition 3.87) of a monad on \mathbf{Set} with an algebraic presentation also has a quantitative algebraic presentation (Definition 3.82), i.e. it can be axiomatized with quantitative equations.²¹ In

¹⁶ That is a relaxation of the usual axioms for metrics (see Definition 0.1). Diffuse metrics are also called dislocated metrics in [HS00].

¹⁷ The first time we did this was in [MSV22], and with [MSV23] and this thesis, we aim to simplify and broaden our initial proposal.

¹⁸ In particular, taking $L = [0, \infty]$ with the axioms of Definition 0.1 translated into quantitative equations yields metrics (see Example 2.34).

¹⁹ See the discussion in §0.3 and Example 3.70.

²⁰ We give a semantical and a syntactical construction (Definitions 3.45 and 3.73 respectively), and they are equivalent thanks to soundness and completeness of our logic.

²¹ This is related to [ADV22, §5], where *strongly finitary* monads on the category of posets are shown to be monad liftings of *finitary* monads on \mathbf{Set} .

particular, it yields a presentation for a monad on **Met** that is not captured by the framework of [MPP16] nor that of [FMS21] (Remark 3.101).

Apart from those technical contributions, our approach describes quantitative algebraic reasoning as a cleaner generalization of algebraic reasoning.²² This guided the outline of this manuscript which is divided in three chapters, one on classical algebraic reasoning, one on our tailored generalization of metric spaces, and one on combining these two chapters, **lifting algebraic reasoning to generalized metric spaces**. Let us now give more detailed introductions for each of these chapters.²³

0.1 Universal Algebra and Monads

With a bit of experience adding natural numbers together, you quickly notice that addition respects some *rules*. If you add n and m , you get the same thing as if you add m and n , no matter what numbers n and m are. If you add n and 0 , you obtain n . If you add n and m , then add k , you get the same thing as if you add n to the sum of m and k . We represent these rules with **equations**:

$$n + m = m + n \quad n + 0 = n \quad (n + m) + k = n + (m + k). \quad (0.2)$$

These equations also hold when n , m , and k belong to the integers or the real numbers. We can also replace addition with multiplication and 0 with 1 .

Since these rules apply in different contexts, mathematicians came up with an abstract definition of a commutative monoid: a set M with a function $+ : M \times M \rightarrow M$ (written infix) and an element $0 \in M$, such that for all $n, m, k \in M$, the equations above are true. The study of these abstract structures (and other variants like groups and rings) is extremely fruitful,²⁴ so much so that you probably learned about them in a first-year undergraduate mathematics course with “algebra” in its title.

With a bit of experience studying monoids, groups, and rings, you quickly notice the similarities in their definitions, and in the reasoning in proofs about them. The purpose of universal algebra is to formalize what they have in common, in order to investigate them all at once. We study an arbitrary **algebraic theory** instead of doing group theory, ring theory, etc.

An algebraic theory is a syntactic gadget that specifies one kind of algebraic structure with a signature Σ containing operation symbols, and a collection of equations E asserting that some sequences of symbols can be replaced by others. For instance, the theory of commutative monoids contains the symbol $+$, the symbol 0 , and the equations in (0.2).

The models of a theory derived from (Σ, E) are called (Σ, E) -algebras. They are sets in which you can combine elements as dictated by the operations in Σ in a way that respects the rules expressed by the equations in E . For instance, the models of the theory of commutative monoids are commutative monoids.²⁵

The flexibility of universal algebra was recognized as a powerful tool early on in the history of formal semantics of programming languages (at the least in [SS71]). We already saw that sequential composition $;$ and conditional branching ifte could

²² This is supported by Item iv. and Examples 3.70 and 3.71.

²³ You could skip these now and come back to each of the following sections when starting to read the corresponding chapter.

²⁴ I cannot do better than a euphemism here. Even narrowing to theoretical computer science, algebraic reasoning has many applications — there are two noteworthy international conferences with “algebra” and “computer science” in their names, CALCO [GS21] and RAMiCS [FGSW21]. Our story focuses on algebraic semantics only.

²⁵ Groups and rings are other similar examples of algebras, but fields are not because the property of all non-zero elements being units cannot be asserted with equations.

be modelled as algebraic operations. Let us mention two additional well-known examples which were the main source of examples for quantitative algebras.²⁶

To represent programs that use nondeterminism, we use a binary operation \oplus . If P and Q are programs, then $P \oplus Q$ nondeterministically chooses to run P or Q . The equations that govern the behavior of \oplus are

$$P \oplus P = P, \quad P \oplus Q = Q \oplus P, \quad \text{and} \quad P \oplus (Q \oplus R) = (P \oplus Q) \oplus R.$$

Briefly, they state that a nondeterministic choice is not affected by the order or multiplicity of the possible outcomes.²⁷

To represent programs that make decisions according to some probability distributions, we use a family of binary operations $+_p$ indexed by real numbers $0 < p < 1$. If P and Q are programs, then $P +_p Q$ is the program that runs P with probability p and Q with probability $1 - p$. For example, if P and Q return HEADS and TAILS respectively, then $P +_{0.5} Q$ is a **fair coin**. The equations look a lot like those for \oplus , for example $P +_p P = P$ for any p .²⁸

To fully grasp the last sentence of the paragraph on nondeterminism, it is crucial to note that the three equations we gave entail many more equations (for example $(Q \oplus P) \oplus (P \oplus Q) = P \oplus Q$). We can appreciate this from two equivalent angles. Semantically, an equation ϕ is entailed by a set of equations E if all the models of (Σ, E) satisfy ϕ . Syntactically, ϕ is entailed by E if it can be derived in equational logic (see Figure 1.3).²⁹

Yet another take on algebraic theories comes from category theory. Birkhoff [Bir35] had already realized that one can always freely generate (Σ, E) -algebras, and Lawvere [Law63] and Linton [Lin66] recognized this induces a monad $\mathcal{T}_{\Sigma, E}$ on the category of sets. They also showed there is a (partial) converse: any *finitary* monad on **Set** is presented by an algebraic theory.³⁰

Moggi first conveyed the applicability of monads (an abstract notion from category theory) in computer science in [Mog89, Mog91]. They became a valuable tool in semantics, and a monad paired with an algebraic presentation allows to combine categorical and equational reasoning. It can be very effective (even outside semantics) as shown in, e.g. [PP01a, PP01b, PP02, BP15, BHKR15, DPS18, PRSW20, BSS21, BSV22, ZM22, RZHE24].

In Chapter 1, we tell the story many times retold of universal algebra. We adopt a somewhat peculiar presentation of the material in order to replicate it more accurately in Chapter 3. We also give some examples of algebras, algebraic theories and algebraic presentations.

0.2 Generalized Metric Spaces

In many applications, deciding whether programs are equivalent or not is overly simplistic. We gave the example of compression algorithms, but let us give three more.

Artificial Intelligence. A lot of models in AI, especially in machine learning, rely on probabilistic reasoning to make decisions.³¹ For example, when a **classifier** is

²⁶ See [BSV22] for a more detailed account in the classical setting, and [MSV21] for the quantitative setting.

²⁷ See Example 1.78.

²⁸ See Example 1.79.

²⁹ The first account of this logic, and the equivalence between these two points of view are due to Birkhoff [Bir35], and we prove it in Theorems 1.55 and 1.60.

³⁰ i.e. any finitary monad is isomorphic to $\mathcal{T}_{\Sigma, E}$ for some Σ and E .

³¹ See, e.g. [CKPR21] which motivated Example 3.102.

fed an image, before deciding what the image depicts, it produces a probability distribution over things that could possibly be in that image. It goes like this:

$$\text{dogpic.jpg} \xrightarrow{\text{classify}} 89\% \text{ dog} + 6\% \text{ lion} + 2\% \text{ cat} + \dots \xrightarrow{\text{max}} \text{dog}.$$

Consider two different classifiers that consistently give the same (possibly correct) answer on the testing dataset. One might consider them to be equal, but a closer examination could reveal that one classifier is more confident than the other. In other words, the distributions produced by one classifier may be more concentrated than those produced by the other.³² Therefore, it makes sense to compare classifiers (more generally, AI models) by devising a notion of distance on the probability distributions they produce. We will give two examples of distances between distributions within our framework in Examples 3.85 and 3.102.

Quantitative Information Flow. When designing software that handles data containing private information, one often wants a balance between the privacy of the users and the utility provided. It makes sense to share the average grade for a class of 100 students, but not for a class of 5 students. With the methods developed in quantitative information flow [QIF20] (especially differential privacy [Dwo06]), we can compare the levels of confidentiality of different programs, before deciding what is the safest (most private) one to roll out.³³

Code Optimization. Consider the two pieces of pseudocode in Figure 1.

```

do
    x = Bernoulli(0.3)
    y = Bernoulli(0.3)
while (x == y)
return x

return Bernoulli(0.5)

```

For all intents and purposes, they are equivalent.³⁴ However, there is only a weak guarantee that the second program terminates (it does with probability 1). Still, if you are unable to run `Bernoulli(0.5)` for some reason, you would be perfectly happy to use the second program. If you want to have a strong guarantee of termination, you could interrupt the loop after, say, 1000 iterations and then return an arbitrary value (see Figure 2). Unfortunately, this breaks the equivalence with `Bernoulli(0.5)`, but

```

i = 0
do
    i = i + 1
    x = Bernoulli(0.3)
    y = Bernoulli(0.3)
while (x == y) AND i < 1000
return x

```

it is still appropriate to say that the two programs are close to each other (even if they

³² In particular, the distributions produced by the perfect classifier (one that knows the correct labels) are always fully concentrated at a single point.

³³ For now, this is only a potential application, we do not have concrete results in this direction.

Figure 1: Simulating a fair coin flip with a biased coin (with a weak guarantee of termination) using an idea of [vN51].

³⁴ If you throw a (possibly biased) coin twice and you get two different outcomes, the probability that the first outcome was HEADS is equal to the probability that it was TAILS, hence it is 0.5 (assuming throws are independent).

Figure 2: Simulating a fair coin flip with a biased coin (with a strong guarantee of termination).

are not equivalent), and that they would be even closer if we increase the maximum number of iterations. When some features are not available, or more realistically when their implementation is not efficient, it can be convenient to write code that approximates the specification but runs (faster).

A widespread alternative to equality that is inherently more fine-grained is metrics. The first definition of metric space (under the name “(E) classes”) is credited to Fréchet’s thesis [Fré06]. We give the definition that is now standard.³⁵

Definition 0.1 (Metric space). A **metric space** is a pair (A, d) comprising a set A and a function $d : A \times A \rightarrow [0, \infty)$ called the metric satisfying for all $a, b, c \in A$:

1. separation: $d(a, b) = 0 \Leftrightarrow a = b$,
2. symmetry: $d(a, b) = d(b, a)$, and
3. triangle inequality: $d(a, c) \leq d(a, b) + d(b, c)$.

For more than a 100 years now, metrics have been a good abstract formalization of what we intuitively understand to be **distances**. In particular, $d(a, b)$ is often called the distance between a and b . Therefore, instead of reasoning about program equivalence, we reason about **program distances**.³⁶

The study of distances between programs (especially those with probabilistic aspects) began in the previous century (see [vBo1] for a (relatively old) survey). While there is no international conference on the subject,³⁷ it is still a very active area of research (see, e.g. [CDL15, CDL17, BBLM18a, BMPP18, BBLM18b, BBKK18, MSV21, Pis21]).

In this literature, there is a recurring idea that positive real numbers are not always the best space to value distances in. Oftentimes, the value ∞ is allowed, where $d(a, b) = \infty$ means a and b are as far apart as they can be. Sometimes, distances are bounded above by 1, so $[0, 1]$ replaces $[0, \infty)$. In more exotic cases, it makes sense for $d(a, b)$ to not even be a number, it can be a set [ABH⁺12], a probability distribution [HR13], an element of a continuous semiring [LMMP13], or just a boolean value.

It is also common to remove or modify some axioms of Definition 0.1 to work with, e.g. pseudometric spaces [BBKK18] or ultrametric spaces [Esc99, Pis21].

It would be ideal if we could devise a definition that encapsulates all existing formal notions of distance. That is obviously not possible. Moreover, even the term “generalized metric space” is employed across various research communities with different meanings (see, e.g. [BvBR98, Bra00, LY16, Pis21]).

In [MPP16], the authors propose theoretical foundations for quantitative algebraic semantics. Their work allows to reason equationally about metrics. One of our contributions in [MSV22] was to show that you can handpick any subset of the axioms of metric spaces and carry out all the proofs of the original paper [MPP16] without much trouble.³⁸ I believe this was known to the authors of [MPP16], especially in light of the results of [FMS21] which morally do the same thing except for an even more general class of structures.

³⁵ Up to small variations. It is essentially equivalent to Fréchet’s definition, but uses different notation and terminology.

³⁶ In semantics, people also use the term behavioral distance/metric.

³⁷ Work on this definitely fits in QAPL, but the last meeting was in 2019 [AW20].

³⁸ Although there is a subtlety about the equality predicate that we explain in §0.3.

In this thesis, we propose yet another definition of generalized metric spaces that is as general as possible without requiring any additional technical machinery. In fact, if you read the present work being comfortable with the frameworks presented in [MPP16] or [MSV22], I believe you will not feel far from home.

We first define L-spaces (Definition 2.11) which are sets equipped with a distance function into a complete lattice L ($A, d : A \times A \rightarrow L$). The structure of a complete lattice allows comparing distances (say one is smaller or bigger than another), and to define a distance as an infimum of a set of bounds in L . That is enough to do quantitative algebraic reasoning in the sense of [MPP16].³⁹

Then, we describe a language to specify axioms one can put on L-spaces. We call such axioms quantitative equations (Definition 2.23). They are a restriction of quantitative equations that we define in Chapter 3, so we will motivate them in §0.3. Examples include separation, symmetry and triangle inequality from Definition 0.1, but also reflexivity, transitivity, and antisymmetry of a binary relation, the strong triangle inequality of ultrametric spaces, and many more. A generalized metric space is then an L-space that satisfies a fixed set of quantitative equations.

In Chapter 2, we give lots of examples including posets, preorders, metrics, pseudometrics, ultrametrics, etc. We also study some properties of the categories of generalized metric spaces⁴⁰ in preparation for Chapter 3 which essentially just combines the first and second chapter to do universal algebra on generalized metric spaces.

0.3 Universal Quantitative Algebra

The term *quantitative* is used in this thesis to refer to a notion of distance that *quantifies* how far apart two things are.⁴¹ Universal quantitative algebra is then a framework where one can reason about both equality and distances between algebraic terms (built out of variables and operations in a signature). The first paper on the subject is [MPP16]. Its theoretical contributions are three-fold.⁴²

The authors work in the category **Met** of extended metric spaces (distances valued in $[0, \infty)$) and nonexpansive maps — a function is nonexpansive if it never increases the distance of its inputs (2.3). First, they define a **quantitative algebra** to be a metric space (A, d) equipped with operations that are interpreted as nonexpansive functions $(A, d)^n \rightarrow (A, d)$, where $(A, d)^n$ denotes the n -wise categorical product of (A, d) with itself. Second, they develop an analog to Birkhoff’s equational logic to reason about properties of quantitative algebras, and they show it is sound and complete. Third, they show that free quantitative algebras always exist.

Let us briefly explain the logic presented in [MPP16]. At its core, there is the neat observation that the data of a metric $d : A \times A \rightarrow [0, \infty]$ can be equivalently given as a family of binary relations $\{R_\varepsilon^d \subseteq X \times X\}$ indexed by $\varepsilon \in [0, \infty]$ with some additional properties.⁴³ This point of view is not completely new, it can be glimpsed in [Wea95], [DLPS07, §1.2], [Ngu10, After Proposition 1], and [Con17]. However, in their quantitative equational logic, the authors of [MPP16] propose to take more seriously the point of view that the relation R_ε^d means “equality up to ε ”, and thus

³⁹ We say more on this in Remark 2.22.

⁴⁰ We get one category **GMet** for each complete lattice L and each collection of quantitative equations we decide to impose.

⁴¹ In contrast with the work on Girard’s *quantitative semantics* [Gir88, BE99] or Kesner and Ventura’s *quantitative types* [KV14] which aim to quantify the resource usage of a program.

⁴² The authors also admirably sell their results with several examples combining algebraic and metric reasoning to axiomatize well-known metrics, the Hausdorff distance which we treat more generally in Example 3.83, the Kantorovich distance (Example 3.5), and the total variation distance (Example 3.92).

⁴³ We prove a more general version in Proposition 2.21.

that we can reason about it kind of how we do for equality. In particular, they use the symbol $=_\varepsilon$.

Their logic closely resembles implicational logic (see, e.g. IL1–8 in [Wec92, p. 223–224]) where the equality predicate is replaced by a family of predicates $=_\varepsilon$ where ε is a positive real number.⁴⁴ The meaning of $s =_\varepsilon t$ is that for all possible assignments of variables, the interpretations of s and t are at distance at most ε . It is clearly reminiscent of the meaning of $s = t$ in universal algebra (that for all possible assignments of variables, the interpretations of s and t are equal). The shape of a generic judgment, called quantitative inference, is $\{s_i =_{\varepsilon_i} t_i\} \vdash s =_\varepsilon t$. It asserts that whenever the distance between the interpretations of s_i and t_i are below ε_i for each $i \in I$, the distance between the interpretations of s and t is below ε .

Here are a few inference rules in that logic.

$$\frac{}{\vdash t =_0 t} \text{REFL} \quad \frac{}{s =_\varepsilon t \vdash s =_{\varepsilon+\varepsilon'} t} \text{MAX} \quad \frac{\forall \phi \in \Gamma', \Gamma \vdash \phi \quad \Gamma' \vdash \psi}{\Gamma \vdash \psi} \text{CUT}$$

$$\frac{\{s_i =_\varepsilon t_i \mid 1 \leq i \leq n\} \vdash \text{op}(s_1, \dots, s_n) =_\varepsilon \text{op}(t_1, \dots, t_n)}{\text{NEXP}}$$

The first states that the distance between the interpretation of t and itself is always below 0 (hence equal to 0), this mirrors one side of the separation axiom of metric spaces. The second rule, quantified over all positive reals $\varepsilon, \varepsilon'$, states that if ε is an upper bound for the distance between (the interpretations of) s and t , then you can add any positive quantity, and it will remain an upper bound. The third is a cut rule that you always find in similar deductive systems,⁴⁵ and it simply reflects the semantics of \vdash being an implication.

The last one states that whenever the distance between s_i and t_i is bounded above by ε for each $i \in I$, so is the distance between $\text{op}(s_1, \dots, s_n)$ and $\text{op}(t_1, \dots, t_n)$. After unrolling some definitions, one verifies this is equivalent to the interpretation of op being nonexpansive with respect to the product metric (0.1).⁴⁶

The quantitative equational logic that we present in Figure 3.1 is adapted from the one in [MPP16] in three key ways.

1. In order to deal with quantitative algebras on generalized metric spaces, the predicates $=_\varepsilon$ are now indexed with quantities $\varepsilon \in L$, and the rules like **REFL** above are removed.⁴⁷ Without **REFL**, there is no predicate $=_\varepsilon$ that corresponds to equality. Thus, we have to reintroduce the predicate $=$, and add rules ensuring that it behaves like equality (it is a congruence).
2. We remove **NEXP**. As we foreshadowed, this rule and the requirement of (0.1) are not necessary to develop the theory of quantitative algebras. We first showed this in [MSV22], where we replaced these with a technical notion we called lifted signatures [MSV22, Definition 3.6] and a corresponding inference rule. In [MSV23] and here, we do not replace them with anything as it makes the base logic simpler. It is always possible to recover the nonexpansive property (or its variants from [MSV22]) by adding more axioms (see (3.9)).
3. In an effort to make a better parallel with equational logic, we slightly reduce the

⁴⁴ It is harmless to restrict to rational numbers if one cares about the size of the formal system.

⁴⁵ c.f. IL7 in [Wec92, p. 224] and *cut* in [CM22b, Definition 4.1.1].

⁴⁶ We mentioned this property of interpretations is very natural, but so is the **NEXP** rule: it says that the relation $=_\varepsilon$ is preserved by the operations (like a congruence, except is not necessarily an equivalence relation).

⁴⁷ We also remove rules that ensure the other side of separation, symmetry and triangle inequality.

expressiveness of the logic. The authors of [MPP16] already identified a special class of judgments whose terms in the premises are all variables, that is, their generic shape is $\{x_i =_{\varepsilon_i} y_i\} \vdash s =_{\varepsilon} t$. They call these *basic quantitative inferences*,⁴⁸ and they crucially rely on them to define free algebras⁴⁹ [MPP16, Theorem 5.1], and to prove variants of the HSP theorem [MPP17, Theorem 3.11].

The premises of a basic quantitative inference (predicates to the left of the turnstile \vdash) can equivalently be described with an L-space on the variables used.⁵⁰ Thus, our generic judgments are now written like $(X, d) \vdash s =_{\varepsilon} t$ or $(X, d) \vdash s = t$, where (X, d) is an L-space, where d is the largest distance that models the premises of the corresponding basic quantitative inference. We call these judgments quantitative equations as we believe they are the proper counterpart to equations in universal algebra.

Recall that quantitative equations also generalize the axioms of generalized metric spaces from §0.2. More accurately, the quantitative equations of Chapter 2 are instances of the quantitative equations of Chapter 3 when the signature is empty. That is morally the reason why we define generalized metric spaces with them.⁵¹

The first and third item can both be found, under guise of further abstraction, in [FMS21]. They deal with relational structures which are more general, but harder to link back to the equational reasoning we are used to in universal algebra. Our main advantage is that, while we can handle various notions of distances that are not metrics (e.g. ultrametrics and partial orders), our logic is not more complicated than [MPP16]’s. In fact, in a sense it is simpler because it only deals with basic quantitative inferences, yet it is still sound and complete.⁵²

To be impactful, one could say our logic is to [MPP16]’s logic as equational logic is to implicational logic. Indeed, what is a *basic* implication in implicational logic? It is a judgment of shape $\{x_i = y_i\} \vdash s = t$, where the terms in the premises are variables only. But this means the premises are trivial because if two variables are equal, you can use a single variable instead. Thus a basic implication is just an equation, and similarly, a basic quantitative inference is just a quantitative equation.

The second item seems to be novel. Although people had removed the non-expansive requirement in [Wea93, Wea95, Hin16, Hin17, AFMS21],⁵³ nobody had done it in the logical apparatus. We were inspired by the ad-hoc approach of [BBLM18a, BBLM18b].

Dismissing NEXP is necessary to prove Theorem 3.98, the main theorem in §3.5. The motivating applications of [MPP16] are presentation results for monads on **Met**. Briefly, they show how the distances induced by their logic (with different sets of axioms) coincide with popular distances used in semantics. Similar results were obtained in, e.g. [MSV21, BMPP18, BMPP21, MSV22], and they all have in common that they reuse a known algebraic presentation for a monad on **Set**. We show in Theorem 3.98 that this is always possible when the monad on **Met** is a monad lifting of the monad on **Set** (Definition 3.87).

When working with nonexpansive operations, or equivalently with the NEXP rule, the induced monads are automatically enriched.⁵⁴ We exhibit a monad lifting

⁴⁸ They require the set of premises to be finite, but that is not important for us.

⁴⁹ Consequently, their examples of axiomatizations only use basic quantitative inferences.

⁵⁰ See the discussion on syntactic sugar before Remark 2.28. This idea also appears in, e.g. [AFMS21, FMS21, Adá22, ADV23b].

⁵¹ We took a less elegant but more pragmatic approach in [MSV23, §8].

⁵² We say more on this in Remark 3.68.

⁵³ Unfortunately, we were not aware of these papers when we published [MSV22], and we did not cite them.

⁵⁴ This is proved in the metric context in [ADV23a, after Corollary 4.19], in the ordered context in [AFMS21, Proposition 4.6], and in the context of relational structures in [FMS21, Corollary 4.14].

that is not enriched in Example 3.88, and it is presented by a quantitative algebraic theory thanks to Theorem 3.98. This shows that our approach is more general (in one aspect) than [MPP16] and [FMS21].

A final benefit we can highlight is the way our simplifications make the story of universal quantitative algebra so similar to the story of universal algebra. In Chapter 3, the outline and many proofs from Chapter 1 are reprised to work with quantitative algebras. We also give some examples of quantitative algebras, quantitative algebraic theories, and quantitative algebraic presentations.

1 Universal Algebra

Concerto Al Andalus

Marcel Khalifé

1.1 Algebras	21
1.2 Terms and Equations	24
1.3 Free Algebras	34
1.4 Equational Logic	45
1.5 Monads	49

For a comprehensive introduction to the concepts and themes explored in this chapter, please refer to §0.1. Here, we only give a brief overview.

In this chapter, we cover the content on universal algebra and monads that we will need in the rest of the thesis. This material has appeared many times in the literature,⁵⁵ but for completeness (and, to be honest, for my own satisfaction) we take our time with it, although we assume the reader is comfortable with basic category theory (the material in the appendix). In Chapter 3, we will follow the outline of the current chapter to generalize the definitions and results to sets equipped with a notion of distance. Thus, many choices in our notations and presentation are motivated by the needs of Chapter 3.⁵⁶

Outline: In §1.1 and §1.2, we define algebras, terms, and equations over a signature of finitary operation symbols. In §1.3, we explain how to construct the free algebras for a given signature and class of equations. In §1.4, we give the rules for equational logic to derive equations from other equations, and we show it is sound and complete. In §1.5, we define monads and algebraic presentations for monads. We give examples all throughout, some small ones to build intuition and some bigger ones that will be important later.

1.1 Algebras

We said in §0.1 that groups and rings are both examples of algebras we want to understand. Groups and rings allow different kinds of combinations of elements, you can do $x \cdot (y + z)$ in a ring but not in a group. To specify which combinations are allowed, we use a signature, and essentially all of this chapter will be parametric over a signature denoted Σ .

Definition 1.1 (Signature). A **signature** is a set Σ whose elements, called **operation symbols**, each come with an **arity** $n \in \mathbb{N}$. We write $\text{op} : n \in \Sigma$ for a symbol op with arity n in Σ . With some abuse of notation, we also denote by Σ the functor $\Sigma : \mathbf{Set} \rightarrow \mathbf{Set}$ with the following action:⁵⁷

$$\Sigma(A) := \coprod_{\text{op}:n \in \Sigma} A^n \text{ on sets} \quad \text{and} \quad \Sigma(f) := \coprod_{\text{op}:n \in \Sigma} f^n \text{ on functions.}$$

⁵⁵ [Wec92] and [Bau19] are two of my favorite references on universal algebra, and both [Rie17, Chapter 5] and [BW05, Chapter 3] are great references for monads (the latter calls them *triples*).

⁵⁶ I hope this will not make this chapter too terse, but the payoff of simply copy-pasting proofs to obtain the generalized results is worth it.

⁵⁷ The set $\Sigma(A)$ can be identified with the set containing $\text{op}(a_1, \dots, a_n)$ for all $\text{op} : n \in \Sigma$ and $a_1, \dots, a_n \in A$. Then, the function $\Sigma(f)$ sends $\text{op}(a_1, \dots, a_n)$ to $\text{op}(f(a_1), \dots, f(a_n))$.

An algebra for a signature Σ is a structure where each operation symbol in Σ is associated to a concrete way to combine elements.

Definition 1.2 (Σ -algebra). A Σ -**algebra** (or just algebra) is a set A equipped with functions $\llbracket \text{op} \rrbracket_A : A^n \rightarrow A$ for every $\text{op} : n \in \Sigma$ called the **interpretation** of the symbol. We call A the **carrier** or **underlying** set, and when referring to an algebra, we will switch between using a single symbol \mathbb{A} ⁵⁸ or the pair $(A, \llbracket - \rrbracket_A)$, where $\llbracket - \rrbracket_A : \Sigma(A) \rightarrow A$ is the function sending $\text{op}(a_1, \dots, a_n)$ to $\llbracket \text{op} \rrbracket_A(a_1, \dots, a_n)$ (it compactly describes the interpretations of all symbols).

A **homomorphism** from \mathbb{A} to \mathbb{B} is a function $h : A \rightarrow B$ between the underlying sets of \mathbb{A} and \mathbb{B} that preserves the interpretation of all operation symbols in Σ , namely, for all $\text{op} : n \in \Sigma$ and $a_1, \dots, a_n \in A$,⁵⁹

$$h(\llbracket \text{op} \rrbracket_A(a_1, \dots, a_n)) = \llbracket \text{op} \rrbracket_B(h(a_1), \dots, h(a_n)). \quad (1.2)$$

The identity maps $\text{id}_A : A \rightarrow A$ and the composition of two homomorphisms are always homomorphisms, therefore we have a category whose objects are Σ -algebras and morphisms are Σ -algebra homomorphisms. We denote it by $\mathbf{Alg}(\Sigma)$.

This category is concrete over \mathbf{Set} with the forgetful functor $U : \mathbf{Alg}(\Sigma) \rightarrow \mathbf{Set}$ which sends an algebra \mathbb{A} to its carrier and a homomorphism to the underlying function between carriers.

Remark 1.3. In the sequel, we will rarely distinguish between the homomorphism $h : \mathbb{A} \rightarrow \mathbb{B}$ and the underlying function $h : A \rightarrow B$. Although, we may write Uh for the latter, when disambiguation is necessary.

Example 1.4. 1. Let $\Sigma = \{\text{p}:0\}$ be the signature containing a single operation symbol p with arity 0. A Σ -algebra is a set A equipped with an interpretation of p as a function $\llbracket \text{p} \rrbracket_A : A^0 \rightarrow A$. Since A^0 is the singleton $\mathbf{1}$, $\llbracket \text{p} \rrbracket_A$ is just a choice of element in A ,⁶⁰ so the objects of $\mathbf{Alg}(\Sigma)$ are pointed sets (sets with a distinguished element). Moreover, instantiating (1.2) for the symbol p , we find that a homomorphism from \mathbb{A} to \mathbb{B} is a function $h : A \rightarrow B$ sending the distinguished point of A to the distinguished point of B . We conclude that $\mathbf{Alg}(\Sigma)$ is the category \mathbf{Set}_* of pointed sets and functions preserving the points.

2. Let $\Sigma = \{\text{f}:1\}$ be the signature containing a single unary operation symbol f . A Σ -algebra is a set A equipped with an interpretation of f as a function $\llbracket \text{f} \rrbracket_A : A \rightarrow A$.

For example, we have the Σ -algebra whose carrier is the set of integers \mathbb{Z} and where f is interpreted as “adding 1”, i.e. $\llbracket \text{f} \rrbracket_{\mathbb{Z}}(k) = k + 1$. We also have the integers modulo 2, denoted by \mathbb{Z}_2 , where $\llbracket \text{f} \rrbracket_{\mathbb{Z}_2}(k) = k + 1 \pmod{2}$.

The fact that a function $h : A \rightarrow B$ satisfies (1.2) for the symbol f is equivalent to the following commutative square.

$$\begin{array}{ccc} A & \xrightarrow{h} & B \\ \llbracket \text{f} \rrbracket_A \downarrow & & \downarrow \llbracket \text{f} \rrbracket_B \\ A & \xrightarrow{h} & B \end{array}$$

⁵⁸ We will try to match the symbol for the algebra and the one for the underlying set only modifying the former with `\mathbb{A}`.

⁵⁹ Equivalently, h makes the following square commute:

$$\begin{array}{ccc} \Sigma(A) & \xrightarrow{\Sigma(f)} & \Sigma(B) \\ \llbracket - \rrbracket_A \downarrow & & \downarrow \llbracket - \rrbracket_B \\ A & \xrightarrow{f} & B \end{array} \quad (1.1)$$

This amounts to an equivalent and more concise definition of $\mathbf{Alg}(\Sigma)$: it is the category of algebras for the signature functor $\Sigma : \mathbf{Set} \rightarrow \mathbf{Set}$ [Awo10, Definition 10.8].

⁶⁰ Hence, we often call 0-ary symbols **constants**.

We conclude that $\mathbf{Alg}(\Sigma)$ is the category whose objects are endofunctions and whose morphisms are commutative squares as above.⁶¹ There is a homomorphism is_odd from \mathbb{Z} to \mathbb{Z}_2 that sends k to $k \pmod{2}$, that is, to 0 when it is even and to 1 when it is odd.

3. Let $\Sigma = \{\cdot : 2\}$ be the signature containing a single binary operation symbol. A Σ -algebra is a set A equipped with an interpretation $\llbracket \cdot \rrbracket_A : A \times A \rightarrow A$. Such a structure is often called a **magma**, and it is part of many more well-known algebraic structures like groups, rings, monoids, etc. While every group has an underlying Σ -algebra⁶² and every group homomorphism is a homomorphism of Σ -algebras, not every Σ -algebra underlies a group since $\llbracket \cdot \rrbracket_A$ is not required to be associative for example. In other words, the category of groups \mathbf{Grp} is a subcategory of $\mathbf{Alg}(\Sigma)$.

We now turn to subalgebras and products of algebras.

Definition 1.5 (Subalgebra). Given $\mathbb{A} \in \mathbf{Alg}(\Sigma)$, a **subalgebra** of \mathbb{A} is a subset $B \subseteq A$ that is closed under the operations in Σ , namely, for any $\text{op} : n \in \Sigma$ and $b_1, \dots, b_n \in B$, $\llbracket \text{op} \rrbracket_A(b_1, \dots, b_n) \in B$. It quickly follows that $\llbracket - \rrbracket_B : \Sigma(B) \rightarrow B$ can be defined as a (co)restriction of $\llbracket - \rrbracket_A$, making $\mathbb{B} = (B, \llbracket - \rrbracket_B)$ into a Σ -algebra and the inclusion $B \hookrightarrow A$ into a homomorphism.

Remark 1.6. The inclusion of a subalgebra is always an injective homomorphism, and reasoning up to isomorphisms we can always view an injective homomorphism as an inclusion of a subalgebra. This relies on the fact that isomorphisms in $\mathbf{Alg}(\Sigma)$ are precisely the bijective homomorphisms.⁶³

Example 1.7. 1. All the standard notions of submonoids, subgroups, subrings, etc. are examples of subalgebras.

2. With the signature $\Sigma = \{p : 0\}$, the subalgebras of a Σ -algebra/pointed set \mathbb{A} are all its subsets that contain the distinguished point, and the latter is the distinguished point inside the subalgebra.
3. For any homomorphism $h : \mathbb{A} \rightarrow \mathbb{B}$, the image $h(A) = \{h(a) \mid a \in A\}$ is closed under the operations by definition, thus it is a subalgebra of \mathbb{B} .

Products of algebras are defined using the usual categorical definition. Namely, they are the products in the category $\mathbf{Alg}(\Sigma)$, we show those always exist by giving their construction.

Lemma 1.8. Let $\{\mathbb{A}_i = (A_i, \llbracket - \rrbracket_i) \mid i \in I\}$ be a family of Σ -algebras indexed by I . We define the algebra $\mathbb{A} = (A, \llbracket - \rrbracket_A)$ with $A = \prod_{i \in I} A_i$ being the cartesian product, and $\llbracket - \rrbracket_A$ being defined coordinatewise: for all $a_1, \dots, a_n \in A$ and $\text{op} : n \in \Sigma$,

$$\llbracket \text{op} \rrbracket_A(a_1, \dots, a_n) = \langle \llbracket \text{op} \rrbracket_i(\pi_i(a_1), \dots, \pi_i(a_n)) \rangle_{i \in I}. \quad (1.3)$$

Then \mathbb{A} is the product $\prod_{i \in I} \mathbb{A}_i$, with $\pi_i : \mathbb{A} \rightarrow \mathbb{A}_i$ being the projection of the cartesian product.

⁶¹ For more categorical thinkers, we can also identify $\mathbf{Alg}(\Sigma)$ with the functor category $[\mathbf{BN}, \mathbf{Set}]$ from the delooping of the (additive) monoid \mathbb{N} to the category of sets. Briefly, it is because a functor $\mathbf{BN} \rightarrow \mathbf{Set}$ is completely determined by where it sends $1 \in \mathbb{N}$.

⁶² In fact, every group has an underlying algebra for the signature $\{\cdot : 2, e : 0, (-)^{-1} : 1\}$.

⁶³ *Quick Proof.* If $h : \mathbb{A} \rightarrow \mathbb{B}$ is a homomorphism and $h^{-1} : \mathbb{B} \rightarrow \mathbb{A}$ is its inverse, we show that it is a homomorphism as well using $h \circ h^{-1} = \text{id}_B$, then (1.2), and finally $h^{-1} \circ h = \text{id}_A$:

$$\begin{aligned} & h^{-1}(\llbracket \text{op} \rrbracket_B(b_1, \dots, b_n)) \\ &= h^{-1}(\llbracket \text{op} \rrbracket_B(h \circ h^{-1}(b_1), \dots, h \circ h^{-1}(b_n))) \\ &= h^{-1} \circ h(\llbracket \text{op} \rrbracket_B(h^{-1}(b_1), \dots, h^{-1}(b_n))) \\ &= \llbracket \text{op} \rrbracket_A(h^{-1}(b_1), \dots, h^{-1}(b_n)). \end{aligned}$$

Proof. First, we need to show that every π_i is a homomorphism, but this is exactly the meaning of (1.3). Next, let $f_i : X \rightarrow \mathbb{A}_i$ be a family of homomorphisms. The pairing $\langle f_i \rangle_{i \in I} : X \rightarrow A$ is defined by $\langle f_i \rangle_{i \in I}(x) = \langle f_i(x) \rangle_{i \in I} \in A$. It is the only function satisfying $\pi_i \circ \langle f_i \rangle_{i \in I} = f_i$ inside **Set**, so it is the only candidate for a homomorphism $X \rightarrow \mathbb{A}$ in **Alg**(Σ) that satisfies $\pi_i \circ \langle f_i \rangle_{i \in I} = f_i$.⁶⁴

Let us show this pairing is a homomorphism with the following derivation:

$$\begin{aligned} \langle f_i \rangle_{i \in I}(\llbracket \text{op} \rrbracket_X(x_1, \dots, x_n)) &= \langle f_i(\llbracket \text{op} \rrbracket_X(x_1, \dots, x_n)) \rangle_{i \in I} \\ &= \langle \llbracket \text{op} \rrbracket_i(f_i(x_1), \dots, f_i(x_n)) \rangle_{i \in I} \\ &= \langle \llbracket \text{op} \rrbracket_i(\pi_i \circ \langle f_i \rangle_{i \in I}(x_1), \dots, \pi_i \circ \langle f_i \rangle_{i \in I}(x_n)) \rangle_{i \in I} \\ &= \llbracket \text{op} \rrbracket_A(\langle f_i(x_1) \rangle_{i \in I}(a_1), \dots, \langle f_i(x_n) \rangle_{i \in I}(a_1)). \quad \square \end{aligned}$$

⁶⁴ Uniqueness holds because $U : \mathbf{Alg}(\Sigma) \rightarrow \mathbf{Set}$ is faithful.

Example 1.9. Arguably the most famous example of a product of algebras is the cartesian plane \mathbb{R}^2 . Equipped with $+$ and 0 , the real numbers form an abelian group, and the product of $(\mathbb{R}, +, 0)$ with itself is \mathbb{R}^2 where $(0, 0)$ is the zero element and $(x, y) + (x', y') = (x + x', y + y')$.

1.2 Terms and Equations

While a group is a Σ -algebra for a naturally chosen signature $\{\cdot : 2, e : 0, (-)^{-1} : 1\}$, we saw that **Grp** is merely a subcategory of **Alg**(Σ) because groups are special kinds of Σ -algebras. They satisfy some properties like associativity of the binary operation which are not true in all Σ -algebras. In this section, we study the kind of properties that can be stated as equations.

If we want to say that a binary operation \cdot is interpreted as a commutative operation, we could write

$$\forall a, b \in A, \quad \llbracket \cdot \rrbracket_A(a, b) = \llbracket \cdot \rrbracket_A(b, a).$$

To say that \cdot is associative, we write

$$\forall a, b, c \in A, \quad \llbracket \cdot \rrbracket_A(\llbracket \cdot \rrbracket_A(a, b), c) = \llbracket \cdot \rrbracket_A(a, \llbracket \cdot \rrbracket_A(b, c)),$$

and as you can see, it gets hard to read very quickly. We make our life easier by defining the interpretation of Σ -terms which are syntactic gadgets built by iterating the symbols in Σ .

Definition 1.10 (Term). Let Σ be a signature and A be a set.⁶⁵ We denote with $\mathcal{T}_\Sigma A$ the set of Σ -terms built syntactically from A and the operation symbols in Σ , i.e. the set inductively defined by

$$\frac{a \in A}{a \in \mathcal{T}_\Sigma A} \quad \text{and} \quad \frac{\text{op} : n \in \Sigma \quad t_1, \dots, t_n \in \mathcal{T}_\Sigma A}{\text{op}(t_1, \dots, t_n) \in \mathcal{T}_\Sigma A}. \quad (1.4)$$

We identify elements $a \in A$ with the corresponding terms $a \in \mathcal{T}_\Sigma A$, and we also identify (as outlined in Footnote 57) elements of $\Sigma(A)$ with terms in $\mathcal{T}_\Sigma A$ containing exactly one occurrence of an operation symbol.⁶⁶

⁶⁵ In the sequel, unless otherwise stated, Σ will be an arbitrary signature.

⁶⁶ Note that any constant $p : 0 \in \Sigma$ belongs to all $\mathcal{T}_\Sigma A$ by the second rule defining $\mathcal{T}_\Sigma A$.

The assignment $A \mapsto \mathcal{T}_\Sigma A$ can be turned into a functor $\mathcal{T}_\Sigma : \mathbf{Set} \rightarrow \mathbf{Set}$ by inductively defining, for any function $f : A \rightarrow B$, the function $\mathcal{T}_\Sigma f : \mathcal{T}_\Sigma A \rightarrow \mathcal{T}_\Sigma B$ as follows:⁶⁷

$$\frac{a \in A}{\mathcal{T}_\Sigma f(a) = f(a)} \quad \text{and} \quad \frac{\text{op} : n \in \Sigma \quad t_1, \dots, t_n \in \mathcal{T}_\Sigma A}{\mathcal{T}_\Sigma f(\text{op}(t_1, \dots, t_n)) = \text{op}(\mathcal{T}_\Sigma f(t_1), \dots, \mathcal{T}_\Sigma f(t_n))}. \quad (1.5)$$

Lemma 1.11. *The action of \mathcal{T}_Σ is functorial, namely, for any $A \xrightarrow{f} B \xrightarrow{g} C$, $\mathcal{T}_\Sigma \text{id}_A = \text{id}_{\mathcal{T}_\Sigma A}$ and $\mathcal{T}_\Sigma(g \circ f) = \mathcal{T}_\Sigma g \circ \mathcal{T}_\Sigma f$.*

Proof. We proceed by induction for both equations.⁶⁸ For any $a \in A$, we have $\mathcal{T}_\Sigma \text{id}_A(a) = \text{id}_A(a) = a$ and

$$\mathcal{T}_\Sigma(g \circ f)(a) = (g \circ f)(a) = \mathcal{T}_\Sigma g(\mathcal{T}_\Sigma f(a)).$$

For any $t = \text{op}(t_1, \dots, t_n)$, we have

$$\mathcal{T}_\Sigma \text{id}_A(\text{op}(t_1, \dots, t_n)) \stackrel{(1.5)}{=} \text{op}(\mathcal{T}_\Sigma \text{id}_A(t_1), \dots, \mathcal{T}_\Sigma \text{id}_A(t_n)) \stackrel{\text{I.H.}}{=} \text{op}(t_1, \dots, t_n),$$

and

$$\begin{aligned} \mathcal{T}_\Sigma(g \circ f)(t) &= \mathcal{T}_\Sigma(g \circ f)(\text{op}(t_1, \dots, t_n)) \\ &= \text{op}(\mathcal{T}_\Sigma(g \circ f)(t_1), \dots, \mathcal{T}_\Sigma(g \circ f)(t_n)) && \text{by (1.5)} \\ &= \text{op}(\mathcal{T}_\Sigma g(\mathcal{T}_\Sigma f(t_1)), \dots, \mathcal{T}_\Sigma g(\mathcal{T}_\Sigma f(t_n))) && \text{I.H.} \\ &= \mathcal{T}_\Sigma g(\text{op}(\mathcal{T}_\Sigma f(t_1), \dots, \mathcal{T}_\Sigma f(t_n))) && \text{by (1.5)} \\ &= \mathcal{T}_\Sigma g \mathcal{T}_\Sigma f(\text{op}(t_1, \dots, t_n)). && \text{by (1.5)} \quad \square \end{aligned}$$

Example 1.12. 1. With $\Sigma = \{p:0\}$, a Σ -term over A is either an element of A or the constant p . For a function $f : A \rightarrow B$, the function $\mathcal{T}_\Sigma f$ sends a to $f(a)$ and p to itself. The functor \mathcal{T}_Σ is then naturally isomorphic to the maybe functor sending A to $A + \mathbf{1}$.

2. With $\Sigma = \{f:1\}$, a Σ -term over A is either an element of A or a term $f(f(\dots f(a)))$ for some a and a finite number of iterations of f . For a function $f : A \rightarrow B$, the function $\mathcal{T}_\Sigma f$ replaces a with $f(a)$ and does not change the number of iterations of f . The functor \mathcal{T}_Σ is then naturally isomorphic to the functor sending A to $\mathbb{N} \times A$.

3. With $\Sigma = \{\cdot:2\}$, a Σ -term is either an element of A or any expression formed by *multiplying* elements of A together like $a \cdot b$, $a \cdot (b \cdot c)$, $((a \cdot a) \cdot c) \cdot (b \cdot c)$ and so on when $a, b, c \in A$.⁶⁹

As we said above, any element in A is a term in $\mathcal{T}_\Sigma A$. We will denote this embedding with $\eta_A^\Sigma : A \rightarrow \mathcal{T}_\Sigma A$, in particular, we will write $\eta_A^\Sigma(a)$ to emphasize that we are dealing with the term a and not the element of A . For instance, the base case of the definition of $\mathcal{T}_\Sigma f$ in (1.5) becomes

$$\frac{a \in A}{\mathcal{T}_\Sigma f(\eta_A^\Sigma(a)) = \eta_B^\Sigma(f(a))}.$$


⁶⁷ In words, $\mathcal{T}_\Sigma f$ replaces a with $f(a)$ and does nothing to operation symbols nor the structure of the term. In particular, $\mathcal{T}_\Sigma f$ acts as identity on constants.

⁶⁸ Many proofs in this chapter are by induction until some point where we will have enough results to efficiently use commutative diagrams.

⁶⁹ We write \cdot infix as is very common. The parentheses are formal symbols to help delimit which \cdot is taken first. They are necessary because the interpretation of \cdot is not necessarily associative so $a \cdot (b \cdot c)$ and $(a \cdot b) \cdot c$ can be interpreted differently in some Σ -algebras.

This is exactly what it means for the family of maps $\eta_A^\Sigma : A \rightarrow \mathcal{T}_\Sigma A$ to be natural in A ,⁷⁰ in other words that $\eta^\Sigma : \text{id}_{\text{Set}} \Rightarrow \mathcal{T}_\Sigma$ is a natural transformation. We can mention now that it will be part of some additional structure on the functor \mathcal{T}_Σ (a monad). The other part of that structure is a natural transformation $\mu^\Sigma : \mathcal{T}_\Sigma \mathcal{T}_\Sigma \Rightarrow \mathcal{T}_\Sigma$ that is more easily described using trees.

For an arbitrary signature Σ , we can think of $\mathcal{T}_\Sigma A$ as the set of rooted trees whose leaves are labelled with elements of A and whose nodes with n children are labelled with n -ary operation symbols in Σ . This makes the action of a function $\mathcal{T}_\Sigma f$ fairly straightforward: it applies f to the labels of all the leaves as depicted in Figure 1.1.



⁷⁰ As a commutative square:

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \eta_A^\Sigma \downarrow & & \downarrow \eta_B^\Sigma \\ \mathcal{T}_\Sigma A & \xrightarrow{\mathcal{T}_\Sigma f} & \mathcal{T}_\Sigma B \end{array} \quad (1.6)$$

Figure 1.1: Applying $\mathcal{T}_\Sigma f$ to $b \cdot (a \cdot c)$ yields $f(b) \cdot (f(a) \cdot f(c))$.

This point of view is particularly helpful when describing the **flattening** of terms: there is a natural way to see a Σ -term over Σ -terms over A as a Σ -term over A . This is carried out by the map $\mu_A^\Sigma : \mathcal{T}_\Sigma \mathcal{T}_\Sigma A \rightarrow \mathcal{T}_\Sigma A$ which takes a tree T whose leaves are labelled with trees T_1, \dots, T_n to the tree T where instead of the leaf labelled T_i , there is the root of T_i with all its children and their children and so on (we “glue” the tree T_i at the leaf labelled T_i). Figure 1.2 shows an example for $\Sigma = \{\cdot : 2\}$. More formally, μ_A^Σ is defined inductively by:

$$\mu_A^\Sigma(\eta_{\mathcal{T}_\Sigma A}^\Sigma(t)) = t \text{ and } \mu_A^\Sigma(\text{op}(t_1, \dots, t_n)) = \text{op}(\mu_A^\Sigma(t_1), \dots, \mu_A^\Sigma(t_n)). \quad (1.7)$$




Figure 1.2: Flattening of a term.

The use of the word “natural” above is not benign, μ^Σ is actually a natural transformation.

Lemma 1.13. *The family of maps $\mu_A^\Sigma : \mathcal{T}_\Sigma \mathcal{T}_\Sigma A \rightarrow \mathcal{T}_\Sigma A$ is natural in A .*

Proof. We need to prove that for any function $f : A \rightarrow B$, $\mathcal{T}_\Sigma f \circ \mu_A^\Sigma = \mu_B^\Sigma \circ \mathcal{T}_\Sigma \mathcal{T}_\Sigma f$.⁷¹ It makes sense intuitively: we should get the same result when we apply f to all the leaves before or after flattening. Formally, we use induction.

⁷¹ As a commutative square:

$$\begin{array}{ccc} \mathcal{T}_\Sigma \mathcal{T}_\Sigma A & \xrightarrow{\mathcal{T}_\Sigma \mathcal{T}_\Sigma f} & \mathcal{T}_\Sigma \mathcal{T}_\Sigma B \\ \mu_A^\Sigma \downarrow & & \downarrow \mu_B^\Sigma \\ \mathcal{T}_\Sigma A & \xrightarrow{\mathcal{T}_\Sigma f} & \mathcal{T}_\Sigma B \end{array} \quad (1.8)$$

For the base case (i.e. terms in the image of $\eta_{\mathcal{T}_E A}^\Sigma$), we have

$$\begin{aligned} \mu_B^\Sigma(\mathcal{T}_E \mathcal{T}_E f(\eta_{\mathcal{T}_E A}^\Sigma(t))) &= \mu_B^\Sigma(\eta_{\mathcal{T}_E B}^\Sigma(\mathcal{T}_E f(t))) && \text{by (1.6)} \\ &= \mathcal{T}_E f(t) && \text{by (1.7)} \\ &= \mathcal{T}_E f(\mu_A^\Sigma(\eta_{\mathcal{T}_E A}^\Sigma(t))). && \text{by (1.7)} \end{aligned}$$

For the inductive step, we have

$$\begin{aligned} \mu_B^\Sigma(\mathcal{T}_E \mathcal{T}_E f(\text{op}(t_1, \dots, t_n))) &= \mu_B^\Sigma(\text{op}(\mathcal{T}_E \mathcal{T}_E f(t_1), \dots, \mathcal{T}_E \mathcal{T}_E f(t_n))) && \text{by (1.5)} \\ &= \text{op}(\mu_B^\Sigma(\mathcal{T}_E \mathcal{T}_E f(t_1)), \dots, \mu_B^\Sigma(\mathcal{T}_E \mathcal{T}_E f(t_n))) && \text{by (1.7)} \\ &= \text{op}(\mathcal{T}_E f(\mu_A^\Sigma(t_1)), \dots, \mathcal{T}_E f(\mu_A^\Sigma(t_n))) && \text{I.H.} \\ &= \mathcal{T}_E f(\text{op}(\mu_A^\Sigma(t_1), \dots, \mu_A^\Sigma(t_n))) && \text{by (1.5)} \\ &= \mathcal{T}_E f(\mu_A^\Sigma(\text{op}(t_1, \dots, t_n))). && \text{by (1.7)} \quad \square \end{aligned}$$

By definition, we have that $\mu^\Sigma \cdot \eta^\Sigma \mathcal{T}_E$ is the identity transformation $\mathbb{1}_{\mathcal{T}_E} : \mathcal{T}_E \Rightarrow \mathcal{T}_E$.⁷² In words, we say that seeing a term trivially as a term over terms then flattening it yields back the original term. Another similar property is that if we see all the variables in a term trivially as terms and flatten the resulting term over terms, the result is the original term. Formally:

Lemma 1.14. *For any set A , $\mu_A^\Sigma \circ \mathcal{T}_E \eta_A^\Sigma = \text{id}_{\mathcal{T}_E A}$, hence $\mu^\Sigma \cdot \mathcal{T}_E \eta^\Sigma = \mathbb{1}_{\mathcal{T}_E}$.*

Proof. We proceed by induction. For the base case, we have

$$\mu_A^\Sigma(\mathcal{T}_E \eta_A^\Sigma(\eta_A^\Sigma(a))) \stackrel{(1.6)}{=} \mu_A^\Sigma(\eta_{\mathcal{T}_E A}^\Sigma(\eta_A^\Sigma(a))) \stackrel{(1.7)}{=} \eta_A^\Sigma(a).$$

For the inductive step, if $t = \text{op}(t_1, \dots, t_n)$, we have

$$\begin{aligned} \mu_A^\Sigma(\mathcal{T}_E \eta_A^\Sigma(t)) &= \mu_A^\Sigma(\mathcal{T}_E \eta_A^\Sigma(\text{op}(t_1, \dots, t_n))) \\ &= \mu_A^\Sigma(\text{op}(\mathcal{T}_E \eta_A^\Sigma(t_1), \dots, \mathcal{T}_E \eta_A^\Sigma(t_n))) && \text{by (1.5)} \\ &= \text{op}(\mu_A^\Sigma(\mathcal{T}_E \eta_A^\Sigma(t_1)), \dots, \mu_A^\Sigma(\mathcal{T}_E \eta_A^\Sigma(t_n))) && \text{by (1.7)} \\ &= \text{op}(t_1, \dots, t_n) = t && \text{I.H.} \quad \square \end{aligned}$$

Trees also make the depth of a term a visual concept. A term $t \in \mathcal{T}_E A$ is said to be of **depth** $d \in \mathbb{N}$ if the tree representing it has depth d .⁷³ We give an inductive definition:

$$\text{depth}(a) = 0 \text{ and } \text{depth}(\text{op}(t_1, \dots, t_n)) = 1 + \max\{\text{depth}(t_1), \dots, \text{depth}(t_n)\}.$$

A term of depth 0 is a term in the image of η_A^Σ . A term of depth 1 is an element of $\Sigma(A)$ seen as a term (recall Footnote 57).

In any Σ -algebra \mathbb{A} , the interpretations of operation symbols give us an element of A for each element of $\Sigma(A)$. Therefore, we get a value in A for all terms in $\mathcal{T}_E A$ of depth 0 or 1 (the value associated to $\eta_A^\Sigma(a)$ is a). Using the inductive definition of $\mathcal{T}_E A$, we can extend these interpretations to all terms: abusing notation, we define the function $\llbracket - \rrbracket_A : \mathcal{T}_E A \rightarrow A$ by⁷³

$$\frac{a \in A}{\llbracket a \rrbracket_A = a} \quad \text{and} \quad \frac{\text{op} : n \in \Sigma \quad t_1, \dots, t_n \in \mathcal{T}_E A}{\llbracket \text{op}(t_1, \dots, t_n) \rrbracket_A = \llbracket \text{op} \rrbracket_A(\llbracket t_1 \rrbracket_A, \dots, \llbracket t_n \rrbracket_A)}. \quad (1.9)$$

⁷² We write \cdot to denote the vertical composition of natural transformations and juxtaposition (e.g. $F\phi$ or ϕF to denote the action of functors on natural transformations), namely, the component of $\mu^\Sigma \cdot \eta^\Sigma \mathcal{T}_E$ at A is $\mu_A^\Sigma \circ \eta_{\mathcal{T}_E A}^\Sigma$ which is $\text{id}_{\mathcal{T}_E A}$ by (1.7).

⁷³ i.e. the longest path from the root to a leaf has d edges. In Figure 1.2, the depth of T and T_1 is 1, the depth of T_2 is 0 and the depth of $\mu_A^\Sigma T$ is 2.

⁷³ For categorical thinkers, $\mathcal{T}_E A$ is essentially defined to be the initial algebra for the endofunctor $\Sigma + A : \mathbf{Set} \rightarrow \mathbf{Set}$ sending X to $\Sigma(X) + A$. Any Σ -algebra $(A, \llbracket - \rrbracket_A)$ defines another algebra for that functor $\llbracket [-]_A, \text{id}_A \rrbracket : \Sigma(A) + A \rightarrow A$. Then, the extension of $\llbracket [-]_A$ to terms is the unique algebra morphism drawn below.

$$\begin{array}{ccc} \Sigma(\mathcal{T}_E A) + A & \dashrightarrow & \Sigma(A) + A \\ \downarrow & & \downarrow \llbracket [-]_A, \text{id}_A \rrbracket \\ \mathcal{T}_E A & \dashrightarrow & A \end{array}$$

The vertical arrow on the left is basically (1.4).

This allows to further extend the interpretation $\llbracket - \rrbracket_A$ to all terms $\mathcal{T}_\Sigma X$ over some set of variables X , provided we have an assignment of variables $\iota : X \rightarrow A$, by precomposing with $\mathcal{T}_\Sigma \iota$. We denote this interpretation with $\llbracket - \rrbracket_A^\iota$:

$$\llbracket - \rrbracket_A^\iota = \mathcal{T}_\Sigma X \xrightarrow{\mathcal{T}_\Sigma \iota} \mathcal{T}_\Sigma A \xrightarrow{\llbracket - \rrbracket_A} A. \quad (1.10)$$

Example 1.15. In the signature $\Sigma = \{f : 1\}$ and over the variables $X = \{x\}$, we have (amongst others) the terms $t = \text{ff}x$ and $s = \text{fff}x$. If we compute the interpretation of t and s in \mathbb{Z} and \mathbb{Z}_2 ,⁷⁴ we obtain for any assignment $\iota : X \rightarrow \mathbb{Z}$ (resp. $\iota : X \rightarrow \mathbb{Z}_2$):

$$\llbracket t \rrbracket_{\mathbb{Z}}^\iota = \iota(x) + 2 \quad \llbracket s \rrbracket_{\mathbb{Z}}^\iota = \iota(x) + 3 \quad \llbracket t \rrbracket_{\mathbb{Z}_2}^\iota = \iota(x) \quad \llbracket s \rrbracket_{\mathbb{Z}_2}^\iota = \iota(x) + 1 \pmod{2}.$$

By definition, a homomorphism preserves the interpretation of operation symbols. We can prove by induction that it also preserves the interpretation of arbitrary terms. Namely, if $h : \mathbb{A} \rightarrow \mathbb{B}$ is a homomorphism, then the following square commutes.⁷⁵

$$\begin{array}{ccc} \mathcal{T}_\Sigma A & \xrightarrow{\mathcal{T}_\Sigma h} & \mathcal{T}_\Sigma B \\ \llbracket - \rrbracket_A \downarrow & & \downarrow \llbracket - \rrbracket_B \\ A & \xrightarrow{h} & B \end{array} \quad (1.11)$$

The converse is (almost trivially) true, if (1.11) commutes, then we can quickly see (1.1) commutes by embedding $\Sigma(A)$ into $\mathcal{T}_\Sigma A$ and $\Sigma(B)$ into $\mathcal{T}_\Sigma B$. It follows readily that for all homomorphisms $h : \mathbb{A} \rightarrow \mathbb{B}$ and all assignments $\iota : X \rightarrow A$,

$$h \circ \llbracket - \rrbracket_A^\iota = \llbracket - \rrbracket_B^{h \circ \iota}. \quad (1.12)$$

Coming back to associativity, instead of writing $\llbracket \cdot \rrbracket_A(a, \llbracket \cdot \rrbracket_A(b, c))$, we can now write $\llbracket a \cdot (b \cdot c) \rrbracket_A$, and it looks cleaner. Moreover, instead of considering a different term for each choice of $a, b, c \in A$, we can consider the term $x \cdot (y \cdot z)$ over a set of variables $\{x, y, z\}$ and quantify over all the possible assignments $\{x, y, z\} \rightarrow A$. We obtain the following definition.

Definition 1.16 (Equation). An **equation** over a signature Σ is a triple comprising a set X of variables called the **context**, and a pair of terms $s, t \in \mathcal{T}_\Sigma X$. We write these as $X \vdash s = t$.

A Σ -algebra \mathbb{A} **satisfies** an equation $X \vdash s = t$ if for any assignment of variables $\iota : X \rightarrow A$, $\llbracket s \rrbracket_A^\iota = \llbracket t \rrbracket_A^\iota$. We use ϕ and ψ to refer to equations, and we write $\mathbb{A} \models \phi$ when \mathbb{A} satisfies ϕ . We also write $\mathbb{A} \models^\iota \phi$ when the equality $\llbracket s \rrbracket_A^\iota = \llbracket t \rrbracket_A^\iota$ holds for a particular assignment $\iota : X \rightarrow A$ and not necessarily for all assignments.

Remark 1.17. Our notation for equations is not standard because many authors do not bother writing the context of an equation and suppose it contains exactly the variables used in s and t . That is theoretically sound for universal algebra, but it will not remain so when we generalize to universal quantitative algebras. Thus, we make the context explicit in our equations as is done in [Wec92] or [Bau19] with the notations $\forall X. s = t$ and $X \mid s = t$ respectively.⁷⁶ We use the turnstile \vdash to match the convention in the literature on quantitative algebras (e.g. [MPP16] and [FMS21]).

⁷⁴ Recall their Σ -algebra structures from Example 1.4.

⁷⁵ *Quick proof.* If $t = a \in A$, then both paths send it to $h(a)$. If $t = \text{op}(t_1, \dots, t_n)$, then

$$\begin{aligned} h(\llbracket t \rrbracket_A) &= h(\llbracket \text{op} \rrbracket_A(\llbracket t_1 \rrbracket_A, \dots, \llbracket t_n \rrbracket_A)) \\ &= \llbracket \text{op} \rrbracket_B(h(\llbracket t_1 \rrbracket_A), \dots, h(\llbracket t_n \rrbracket_A)) \\ &= \llbracket \text{op} \rrbracket_B(\llbracket \mathcal{T}_\Sigma h(t_1) \rrbracket_B, \dots, \llbracket \mathcal{T}_\Sigma h(t_n) \rrbracket_B) \\ &= \llbracket \text{op}(\mathcal{T}_\Sigma h(t_1), \dots, \mathcal{T}_\Sigma h(t_n)) \rrbracket_B \\ &= \llbracket \mathcal{T}_\Sigma h(t) \rrbracket_B. \end{aligned}$$

⁷⁶ Only finite contexts are used in [Wec92] and [Bau19]. We say a bit more on this in Remark 1.61

Example 1.18 (Associativity). With the signature $\Sigma = \{\cdot : 2\}$ and the context $X = \{x, y, z\}$, the equation $\phi = X \vdash x \cdot (y \cdot z) = (x \cdot y) \cdot z$ ⁷⁷ asserts that the interpretation of \cdot is associative. To prove that, suppose $\mathbb{A} \models \phi$, we need to show that for any $a, b, c \in A$,

$$\llbracket \cdot \rrbracket_A(a, \llbracket \cdot \rrbracket_A(b, c)) = \llbracket \cdot \rrbracket_A(\llbracket \cdot \rrbracket_A(a, b), c). \quad (1.13)$$

Let $s = x \cdot (y \cdot z)$ and $t = (x \cdot y) \cdot z$. Observe that the L.H.S. is the interpretation of s under the assignment $\iota : X \rightarrow A$ sending x to a , y to b and z to c , that is, we have $\llbracket \cdot \rrbracket_A(a, \llbracket \cdot \rrbracket_A(b, c)) = \llbracket s \rrbracket_A^\iota$. Under the same assignment, the interpretation of t is the R.H.S. Since $\mathbb{A} \models X \vdash s = t$, $\llbracket s \rrbracket_A^\iota = \llbracket t \rrbracket_A^\iota$, and we conclude (1.13) holds.⁷⁸

Example 1.19. Here are some other simple examples of equations.

- $x, y \vdash x \cdot y = y \cdot x$ states that the interpretation binary operation \cdot is commutative.
- $x, y, z, w \vdash x \cdot y = y \cdot x$ also states that (the interpretation of) \cdot is commutative, but it has some extra unused variables in the context.⁷⁹
- $x \vdash x \cdot x = x$ states that the binary operation \cdot is idempotent.
- $x \vdash fx = fx$ states that the unary operation f is idempotent.
- $x \vdash p = x$ states that the constant p is equal to all elements in the algebra (this means the algebra is a singleton).
- $x, y \vdash x = y$ states that all elements in the algebra are equal (this means the algebra is either empty or a singleton).

Using the fact that interpretations are preserved by homomorphisms (1.12), we can describe how satisfaction is also preserved. Very naively, one would want to say that if $h : \mathbb{A} \rightarrow \mathbb{B}$ is a homomorphism and $\mathbb{A} \models \phi$, then $\mathbb{B} \models \phi$. That is not true.⁸⁰ It is morally because there can be many more assignments into \mathbb{B} than there are into \mathbb{A} . Nevertheless, the naive statement is true on a per-assignment basis.

Lemma 1.20. *Let ϕ be an equation with context X . If $h : \mathbb{A} \rightarrow \mathbb{B}$ is a homomorphism and $\mathbb{A} \models^\iota \phi$ for an assignment $\iota : X \rightarrow A$, then $\mathbb{B} \models^{h \circ \iota} \phi$.*

Proof. Let ϕ be the equation $X \vdash s = t$, we have

$$\begin{aligned} \mathbb{A} \models^\iota \phi &\iff \llbracket s \rrbracket_A^\iota = \llbracket t \rrbracket_A^\iota && \text{definition of } \models \\ &\implies h(\llbracket s \rrbracket_A^\iota) = h(\llbracket t \rrbracket_A^\iota) \\ &\iff \llbracket s \rrbracket_B^{h \circ \iota} = \llbracket t \rrbracket_B^{h \circ \iota} && \text{by (1.12)} \\ &\iff \mathbb{B} \models^{h \circ \iota} \phi. && \text{definition of } \models \quad \square \end{aligned}$$

However, if h is surjective, then any assignment $\iota : X \rightarrow B$, can be factored through an assignment of $h^{-1} \circ \iota : X \rightarrow A$, where h^{-1} is the right inverse of h , i.e. $h \circ h^{-1} = \text{id}_B$. It follows that the naive result holds when h is surjective.

Lemma 1.21. *Let ϕ be an equation with context X . If $h : \mathbb{A} \rightarrow \mathbb{B}$ is a surjective homomorphism and $\mathbb{A} \models \phi$, then $\mathbb{B} \models \phi$.*

⁷⁷ Alternatively, we may write ϕ omitting brackets:

$$x, y, z \vdash x \cdot (y \cdot z) = (x \cdot y) \cdot z.$$

⁷⁸ It is also clear from this argument that any Σ -algebra \mathbb{A} where $\llbracket \cdot \rrbracket_A$ is associative satisfies ϕ .

⁷⁹ This is allowed, but it is always possible to remove unused variables in the context (see Remark 1.61).

⁸⁰ For any Σ which does not contain constants, there is an initial Σ -algebra \mathbb{I} whose carrier is the empty set \emptyset (the interpretation of operations is completely determined because $\Sigma(\emptyset) = \emptyset$ and there is only one function $\emptyset^n \rightarrow \emptyset$). The unique function $\emptyset \rightarrow B$ is always a homomorphism $\mathbb{I} \rightarrow \mathbb{B}$ because (1.1) trivially commutes. While \mathbb{I} satisfies all equations (vacuously), it is clearly possible that \mathbb{B} does not.

Proof. For any assignment $\iota : X \rightarrow B$, we know that $\mathbb{A} \models^{h^{-1} \circ \iota} \phi$ by hypothesis. Then, by Lemma 1.20, we get $\mathbb{B} \models^{h \circ h^{-1} \circ \iota} \phi$, which in turn means $\mathbb{B} \models^{\iota} \phi$ because $h \circ h^{-1} = \text{id}_B$. \square

Moreover, inspecting the proof of Lemma 1.20, we note that the only implication that is not an equivalence in the derivation becomes an equivalence when h is injective. We conclude that for any $\iota : X \rightarrow A$, if \mathbb{B} satisfies ϕ under $h \circ \iota$, then \mathbb{A} satisfies it under ι . Thus,

Lemma 1.22. *Let ϕ be an equation with context X . If $h : \mathbb{A} \rightarrow \mathbb{B}$ is an injective homomorphism and $\mathbb{B} \models \phi$, then $\mathbb{A} \models \phi$.⁸¹*

The last three results stem from the nice interaction between interpretations and homomorphism (1.12). The flattening also interacts well with interpretations in the following sense.

Lemma 1.23. *For any Σ -algebra \mathbb{A} , the following square commutes.⁸²*

$$\begin{array}{ccc} \mathcal{T}_\Sigma \mathcal{T}_\Sigma A & \xrightarrow{\mathcal{T}_\Sigma \llbracket - \rrbracket_A} & \mathcal{T}_\Sigma A \\ \mu_A^\Sigma \downarrow & & \downarrow \llbracket - \rrbracket_A \\ \mathcal{T}_\Sigma A & \xrightarrow{\llbracket - \rrbracket_A} & A \end{array} \quad (1.14)$$

Proof. We proceed by induction. For the base case, we have

$$\llbracket \mu_A^\Sigma(\eta_A^\Sigma(t)) \rrbracket_A \stackrel{(1.7)}{=} \llbracket t \rrbracket_A \stackrel{(1.9)}{=} \llbracket \eta_A^\Sigma(\llbracket t \rrbracket_A) \rrbracket_A \stackrel{(1.6)}{=} \llbracket \mathcal{T}_\Sigma \llbracket - \rrbracket_A(\eta_A^\Sigma(t)) \rrbracket_A.$$

For the inductive step, if $t = \text{op}(t_1, \dots, t_n)$, then

$$\begin{aligned} \llbracket \mu_A^\Sigma(t) \rrbracket_A &= \llbracket \text{op}(\mu_A^\Sigma(t_1), \dots, \mu_A^\Sigma(t_n)) \rrbracket_A && \text{by (1.7)} \\ &= \llbracket \text{op} \rrbracket_A (\llbracket \mu_A^\Sigma(t_1) \rrbracket_A, \dots, \llbracket \mu_A^\Sigma(t_n) \rrbracket_A) && \text{by (1.9)} \\ &= \llbracket \text{op} \rrbracket_A (\llbracket \mathcal{T}_\Sigma \llbracket - \rrbracket_A(t_1) \rrbracket_A, \dots, \llbracket \mathcal{T}_\Sigma \llbracket - \rrbracket_A(t_n) \rrbracket_A) && \text{I.H.} \\ &= \llbracket \text{op}(\mathcal{T}_\Sigma \llbracket - \rrbracket_A(t_1), \dots, \mathcal{T}_\Sigma \llbracket - \rrbracket_A(t_n)) \rrbracket_A && \text{by (1.9)} \\ &= \llbracket \mathcal{T}_\Sigma \llbracket - \rrbracket_A(\text{op}(t_1, \dots, t_n)) \rrbracket_A && \text{by (1.5)} \\ &= \llbracket \mathcal{T}_\Sigma \llbracket - \rrbracket_A(t) \rrbracket_A. && \square \end{aligned}$$

Remark 1.24. To see Lemma 1.23 in another way, notice that (1.14) looks a lot like (1.11), but the map on the left is not the interpretation on an algebra. Except it is! Indeed, we can give a trivial (or syntactic) interpretation of $\text{op} : n \in \Sigma$ on the set $\mathcal{T}_\Sigma A$ by letting $\llbracket \text{op} \rrbracket_{\mathcal{T}_\Sigma A}(t_1, \dots, t_n) = \text{op}(t_1, \dots, t_n)$. Then, we can verify by induction⁸³ that $\llbracket - \rrbracket_{\mathcal{T}_\Sigma A} : \mathcal{T}_\Sigma \mathcal{T}_\Sigma A \rightarrow \mathcal{T}_\Sigma A$ is equal to μ_A^Σ . We conclude that Lemma 1.23 says that for any algebra, $\llbracket - \rrbracket_A$ is a homomorphism from $(\mathcal{T}_\Sigma A, \llbracket - \rrbracket_{\mathcal{T}_\Sigma A})$ to \mathbb{A} .

In light of this remark, we mention two very similar results: given a set A , μ_A^Σ is a homomorphism between $\mathcal{T}_\Sigma \mathcal{T}_\Sigma A$ and $\mathcal{T}_\Sigma A$, and given a function $f : A \rightarrow B$, $\mathcal{T}_\Sigma f$ is a homomorphism between $\mathcal{T}_\Sigma A$ and $\mathcal{T}_\Sigma B$.

⁸¹ In particular, all the equations satisfied by an algebra are satisfied by all its subalgebras.

⁸² In words, given a term in $\mathcal{T}_\Sigma \mathcal{T}_\Sigma A$, you obtain the same result if you interpret its flattening in \mathbb{A} , or if you interpret the term obtained by first interpreting all the ‘‘inner’’ terms.

This also generalizes to terms in $\mathcal{T}_\Sigma \mathcal{T}_\Sigma X$. Indeed, given an assignment, $\iota : X \rightarrow A$, we can either flatten a term and interpret it under ι , or we can interpret all the inner terms under ι , then interpret the result, as shown in (1.15).

$$\begin{array}{ccccc} & & \mathcal{T}_\Sigma \llbracket - \rrbracket'_A & & \\ & & \text{(1.10)} & & \\ \mathcal{T}_\Sigma \mathcal{T}_\Sigma X & \xrightarrow{\mathcal{T}_\Sigma \mathcal{T}_\Sigma \iota} & \mathcal{T}_\Sigma \mathcal{T}_\Sigma A & \xrightarrow{\mathcal{T}_\Sigma \llbracket - \rrbracket'_A} & \mathcal{T}_\Sigma A \\ \mu_X^\Sigma \downarrow & \text{(1.8)} & \mu_A^\Sigma \downarrow & \text{(1.14)} & \downarrow \llbracket - \rrbracket_A \\ \mathcal{T}_\Sigma X & \xrightarrow{\mathcal{T}_\Sigma \iota} & \mathcal{T}_\Sigma A & \xrightarrow{\llbracket - \rrbracket_A} & A \\ & & \text{(1.10)} & & \\ & & \llbracket - \rrbracket'_A & & \end{array} \quad (1.15)$$

⁸³ Or we can compare (1.7) and (1.9) to see they become the same inductive definition in this instance.

Lemma 1.25. For any function $f : A \rightarrow B$, the following squares commute.⁸⁴

$$\begin{array}{ccc} \mathcal{T}_\Sigma \mathcal{T}_\Sigma \mathcal{T}_\Sigma A & \xrightarrow{\mathcal{T}_\Sigma \mu_A^\Sigma} & \mathcal{T}_\Sigma \mathcal{T}_\Sigma A \\ \mu_{\mathcal{T}_\Sigma A}^\Sigma \downarrow & & \downarrow \mu_A^\Sigma \\ \mathcal{T}_\Sigma \mathcal{T}_\Sigma A & \xrightarrow{\mu_A^\Sigma} & \mathcal{T}_\Sigma \end{array} \quad (1.16)$$

$$\begin{array}{ccc} \mathcal{T}_\Sigma \mathcal{T}_\Sigma A & \xrightarrow{\mathcal{T}_\Sigma \mathcal{T}_\Sigma B} & \mathcal{T}_\Sigma \mathcal{T}_\Sigma B \\ \mu_A^\Sigma \downarrow & & \downarrow \mu_B^\Sigma \\ \mathcal{T}_\Sigma & \xrightarrow{\mathcal{T}_\Sigma f} & \mathcal{T}_\Sigma B \end{array} \quad (1.17)$$

Another consequence of (1.16) is that if you have a term in $\mathcal{T}_\Sigma^n A$ for any $n \in \mathbb{N}$, there are $(n-1)!$ ways to flatten it⁸⁵ by successively applying an instance of $\mathcal{T}_\Sigma^i \mu_{\mathcal{T}_\Sigma^j A}^\Sigma$ with different i and j (i.e. flattening at different levels inside the term), but all these ways lead to the same end result in $\mathcal{T}_\Sigma A$. It is like when you have an expression built out of additions with possibly lots of nested bracketing, you can compute the sums in any order you want, and it will give the same result. That property of addition is a consequence of associativity, hence one also says μ^Σ is associative.

Going back to **Grp** as a subcategory of **Alg**(Σ) with $\Sigma = \{\cdot, 2, e, 0, (-)^{-1} : 1\}$, we can precisely identify it as the full subcategory containing only algebras that satisfy the following equations:

$$\begin{array}{l} x, y, z \vdash x \cdot (y \cdot z) = (x \cdot y) \cdot z \\ x \vdash x \cdot e = x \quad x \vdash x \cdot x^{-1} = e \\ x \vdash e \cdot x = x \quad x \vdash x^{-1} \cdot x = e \end{array} \quad (1.18)$$

We can tell a similar story with **Ring**, **Mon**, **Ab**, and so on. These special subcategories are called varieties.

Definition 1.26 (Variety). Given a class E of equations, we say \mathbb{A} satisfies E and write $\mathbb{A} \models E$ if $\mathbb{A} \models \phi$ for all $\phi \in E$.⁸⁶ A (Σ, E) -**algebra** is a Σ -algebra that satisfies E . We define **Alg**(Σ, E), the category of (Σ, E) -algebras, to be the full subcategory of **Alg**(Σ) containing only those algebras that satisfy E . A **variety** is a category equal to **Alg**(Σ, E) for some class of equations E .

There is an evident forgetful functor $U : \mathbf{Alg}(\Sigma, E) \rightarrow \mathbf{Set}$ which is the composition of the inclusion functor $\mathbf{Alg}(\Sigma, E) \rightarrow \mathbf{Alg}(\Sigma)$ and $U : \mathbf{Alg}(\Sigma) \rightarrow \mathbf{Set}$.⁸⁷

It is never the case in practice that E is a proper class, it is usually a finite or countable set, even recursively enumerable. Still, nothing breaks when E is a class, and we will need this generality in one our main contributions (Theorem 3.98).

Example 1.27. 1. With $\Sigma = \{p : 0\}$, there are morally only four different equations:⁸⁸

$$\vdash p = p, \quad x \vdash x = x, \quad x \vdash p = x, \quad \text{and} \quad x, y \vdash x = y,$$

where we write nothing before the turnstile (\vdash) instead of the empty set \emptyset .

Any algebra \mathbb{A} satisfies the first two equations because $\llbracket p \rrbracket_A^\iota = \llbracket p \rrbracket_A^\iota$, where $\iota : \emptyset \rightarrow A$ is the only possible assignment, and $\llbracket x \rrbracket_A^\iota = \iota(x) = \llbracket x \rrbracket_A^\iota$ for all $\iota : \{x\} \rightarrow A$. If \mathbb{A} satisfies the third, it means that A is empty or a singleton because for any $a, b \in A$, the assignments $\iota_a = x \mapsto a$ and $\iota_b = x \mapsto b$ give us⁸⁹

$$a = \iota_a(x) = \llbracket x \rrbracket_A^{\iota_a} = \llbracket p \rrbracket_A^{\iota_a} = \llbracket p \rrbracket_A^{\iota_b} = \llbracket x \rrbracket_A^{\iota_b} = \iota_b(x) = b.$$

⁸⁴ *Proof.* We have already shown both these squares commute. Indeed, (1.16) is an instance of (1.14) where we identify μ_A^Σ with the interpretation $\llbracket - \rrbracket_{\mathcal{T}_\Sigma A}$ as explained in Remark 1.24, and (1.17) is the naturality square (1.8).

⁸⁵ There is 1 way to flatten a term in $\mathcal{T}_\Sigma^2 A$ to one in $\mathcal{T}_\Sigma A$, and there are $n-1$ ways to flatten from $\mathcal{T}_\Sigma^n A$ to $\mathcal{T}_\Sigma^{(n-1)} A$. By induction, we find $(n-1)!$ possible combinations of flattening $\mathcal{T}_\Sigma^n A \rightarrow \mathcal{T}_\Sigma A$.

⁸⁶ Similarly for satisfaction under a particular assignment ι :

$$\mathbb{A} \models^\iota E \iff \forall \phi \in E, \mathbb{A} \models^\iota \phi.$$

⁸⁷ We will denote all the forgetful functors with the symbol U unless we need to emphasize the distinction. However, thanks to the knowledge package, you can click on (or hover) that symbol to check exactly which forgetful functor it is referring to.

⁸⁸ Let us not formally argue about that here, but your intuition on equality and the fact that terms in $\mathcal{T}_\Sigma X$ are either $x \in X$ or p should be enough to convince you.

⁸⁹ We find $a = b$ for any $a, b \in A$ and A contains at least one element, the interpretation of the constant p , so A is a singleton.

If \mathbb{A} satisfies the fourth equation, it is also empty or a singleton because for any $a, b \in A$, the assignment ι sending x to a and y to b gives us

$$a = \iota(x) = \llbracket x \rrbracket'_A = \llbracket y \rrbracket'_A = \iota(y) = b.$$

Therefore,⁹⁰ there are only two varieties in that signature, either $\mathbf{Alg}(\Sigma, E)$ is all of $\mathbf{Alg}(\Sigma)$, or it contains only the empty set and the singletons.

⁹⁰ Modulo the argument about these being all the possible equations over Σ .

2. With $\Sigma = \{+ : 2, e : 0\}$, there are many more possible equations, but the following three are well-known:

$$x, y, z \vdash x + (y + z) = (x + y) + z, \quad x, y \vdash x + y = y + x, \quad \text{and} \quad x \vdash x + e = x. \quad (1.19)$$

We already saw in Example 1.18 that the first asserts associativity of the interpretation of $+$. With a similar argument, one shows that the second asserts $\llbracket + \rrbracket$ is commutative, and the third asserts $\llbracket e \rrbracket$ is a neutral element (on the right) for $\llbracket + \rrbracket$.⁹¹ Moreover, note that a homomorphism of Σ -algebras from \mathbb{A} to \mathbb{B} is any function $h : A \rightarrow B$ that satisfies

$$\forall a, a' \in A, \quad h(\llbracket + \rrbracket_A(a, a')) = \llbracket + \rrbracket_B(h(a), h(a')) \quad \text{and} \quad h(\llbracket e \rrbracket_A) = \llbracket e \rrbracket_B.$$

Namely, a homomorphism preserves the addition and its neutral element. Thus, letting E be the set containing the equations in (1.19), we find that $\mathbf{Alg}(\Sigma, E)$ is the category \mathbf{CMon} of commutative monoids and monoid homomorphisms.

⁹¹ i.e. if \mathbb{A} satisfies $x \vdash x + e = x$, then for all $a \in A$,

$$\llbracket a + e \rrbracket_A = a.$$

By commutativity, we also get $\llbracket e + a \rrbracket_A = a$.

3. We can add a unary operation symbol $-$ to get $\Sigma = \{+ : 2, e : 0, - : 1\}$, and add the equation $x \vdash x + (-x) = e$ to those in (1.19),⁹² and we can show that $\mathbf{Alg}(\Sigma, E)$ is the category \mathbf{Ab} of abelian groups and group homomorphisms. Note that E is not the set of equations we used to define \mathbf{Grp} with an additional commutativity equation because when the binary operation is commutative, some equations are redundant. We infer that it is possible for (Σ, E) and (Σ', E') to define the same variety (or isomorphic varieties), even if $\Sigma = \Sigma'$ and $E \neq E'$.

⁹² While the signature has changed between the two examples, the equations of (1.19) can be understood over both signatures because they concern terms constructed using the symbols common to both signatures.

One fundamental result in universal algebra characterizes varieties as the subcategories of $\mathbf{Alg}(\Sigma)$ that are closed under some simple constructions. It follows from Lemma 1.21 that all varieties is closed under homomorphic images, meaning that if $h : \mathbb{A} \rightarrow \mathbb{B}$ is a homomorphism in $\mathbf{Alg}(\Sigma)$ and $\mathbb{A} \in \mathbf{Alg}(\Sigma, E)$, then the image $h(A)$ is also in $\mathbf{Alg}(\Sigma, E)$.⁹³ It follows from Lemma 1.22 that all varieties are closed under subalgebras, meaning that all subalgebras of $\mathbb{A} \in \mathbf{Alg}(\Sigma, E)$ also belongs to $\mathbf{Alg}(\Sigma, E)$.⁹⁴ Finally, varieties are also closed under products.

⁹³ Because h can be seen as a surjective homomorphism $\mathbb{A} \rightarrow h(A)$, so $\mathbb{A} \models E$ implies $h(A) \models E$.

⁹⁴ Because the inclusion of a subalgebra \mathbb{B} in \mathbb{A} is an injective homomorphism, so $\mathbb{A} \models E$ implies $\mathbb{B} \models E$.

Lemma 1.28. *Let ϕ be an equation with context X , $\{\mathbb{A}_i = (A_i, \llbracket - \rrbracket_i) \mid i \in I\}$ be a family of algebras indexed by I , and $\mathbb{A} = \prod_{i \in I} \mathbb{A}_i$ be their product as described in Lemma 1.8. For any assignment $\iota : X \rightarrow A$,*

$$\mathbb{A} \models' \phi \Leftrightarrow \forall i \in I, \mathbb{A}_i \models^{\pi_i \circ \iota} \phi. \quad (1.20)$$

Consequently, if every \mathbb{A}_i satisfies ϕ , then so does \mathbb{A} .⁹⁵

⁹⁵ This readily follows from (1.20) because for any assignment $\iota : X \rightarrow A$, every $\pi_i \circ \iota$ is an assignment $X \rightarrow A_i$. Then, by hypothesis $\mathbb{A}_i \models^{\pi_i \circ \iota} \phi$ holds for every $i \in I$, and we conclude that \mathbb{A} satisfies ϕ by (3.11).

Proof. Because each π_i is a homomorphism, we can use Lemma 1.20 for the forward direction (\Rightarrow). For the converse (\Leftarrow), if ϕ is $X \vdash s = t$, then $\pi_i(\llbracket s \rrbracket'_A) = \pi_i(\llbracket t \rrbracket'_A)$ holds by satisfaction under each $\pi_i \circ \iota$ and by (1.12). This means that the interpretations under ι of s and t agree on all coordinates, hence they must coincide, i.e. $\mathbb{A} \models^t \phi$. \square

Birkhoff's variety or HSP theorem states that these closure properties characterize the varieties in $\mathbf{Alg}(\Sigma)$.⁹⁶

Theorem 1.29 (Birkhoff). *A subcategory of $\mathbf{Alg}(\Sigma)$ is a variety if and only if it is closed under homomorphic images, subalgebras, and products.*

We proved the forward direction, and we refer the reader to [Wec92, §3.2.3, Theorem 21] for the converse. We can also mention that Birkhoff's variety theorem has been generalized many times in very abstract settings, see e.g. [Bar92, Bar94, Bar02, MU19, JMU24], [Man76, §3.3], and [AHS06, Corollary 16.17].

A single variety can be defined with different classes of equations, but among different classes of equations over the same signature that define the same variety, there is a largest one.

Definition 1.30 (Algebraic theory). Given a class E of equations over Σ , the **algebraic theory** generated by E , denoted by $\mathfrak{Th}(E)$, is the class of equations (over Σ) that are satisfied in all (Σ, E) -algebras:⁹⁷

$$\mathfrak{Th}(E) = \{X \vdash s = t \mid \forall \mathbb{A} \in \mathbf{Alg}(\Sigma, E), \mathbb{A} \models X \vdash s = t\}.$$

Formulated differently, $\mathfrak{Th}(E)$ contains the equations that are semantically entailed by E , namely $\phi \in \mathfrak{Th}(E)$ if and only if

$$\forall \mathbb{A} \in \mathbf{Alg}(\Sigma), \quad \mathbb{A} \models E \implies \mathbb{A} \models \phi. \quad (1.21)$$

Of course, $\mathfrak{Th}(E)$ contains all of E ,⁹⁸ but also many more equations like $x \vdash x = x$ which is satisfied by any algebra. We will see in §1.4 how to find which equations are entailed by others.

It is easy to see that 1) $\mathbf{Alg}(\Sigma, E) = \mathbf{Alg}(\Sigma, E')$ implies $\mathfrak{Th}(E) = \mathfrak{Th}(E')$, 2) $E \subseteq \mathfrak{Th}(E)$, and 3) $\mathbf{Alg}(\Sigma, \mathfrak{Th}(E)) = \mathbf{Alg}(\Sigma, E)$. It follows that $\mathfrak{Th}(E)$ is the maximal class of equations defining the variety $\mathbf{Alg}(\Sigma, E)$.

Example 1.31. If E contains the equations in (1.19), then $\mathfrak{Th}(E)$ will contain all the equations that every commutative monoid satisfies. Here is a non-exhaustive list:

- $x \vdash e + x = x$ says that $\llbracket e \rrbracket$ is a neutral element on the left for $\llbracket + \rrbracket$ which is true because, by equations in (1.19), $\llbracket e \rrbracket$ is neutral on the right and $\llbracket + \rrbracket$ is commutative.
- $z, w \vdash z + w = w + z$ also states commutativity of $\llbracket + \rrbracket$ but with different variable names.
- $x, y, z, w \vdash (x + w) + (x + z) + (x + y) = ((x + x) + x) + (y + (z + (e + w)))$ is just a random equation that can be shown using the properties of commutative monoids.⁹⁹

⁹⁶ It is stated with some different notations and terminology in [Bir35, Theorem 10].

⁹⁷ Note that, even if E is a set, there is no guarantee that $\mathfrak{Th}(E)$ is a set (in fact it never is) because the collection of all equations is a proper class (because the contexts can be any set).

⁹⁸ Because a (Σ, E) -algebra satisfies E by definition.

⁹⁹ We will see in §1.4 how to systematically generate all the equations in $\mathfrak{Th}(E)$.

1.3 Free Algebras

Very briefly, the free (Σ, E) -algebra on X is the least constrained Σ -algebra which “contains” X and satisfies E . It necessarily satisfies all the equations in $\mathfrak{Th}(E)$ as well, but it does not satisfy any other equation $X \vdash s = t$ that is not also satisfied by all (Σ, E) -algebras. We will prove it always exist and we start with an example.

Example 1.32 (Words). Let $\Sigma_{\text{Mon}} = \{\cdot, \cdot, e, 0\}$, $X = \{a, b, \dots, z\}$ be the set of (lowercase) letters in the Latin alphabet, and X^* be the set of finite words using only these letters.¹⁰⁰ There is a natural Σ_{Mon} -algebra structure on X^* where \cdot is interpreted as concatenation, i.e. $\llbracket \cdot \rrbracket_{X^*}(u, v) = uv$, and e as the empty word ε . This algebra satisfies the equations defining a monoid given in (1.22).¹⁰¹

$$E_{\text{Mon}} = \{x, y, z \vdash x \cdot (y \cdot z) = (x \cdot y) \cdot z, \quad x \vdash x \cdot e = x, \quad x \vdash e \cdot x = x\}. \quad (1.22)$$

In fact, X^* is the *free* monoid over X . This means that for any other $(\Sigma_{\text{Mon}}, E_{\text{Mon}})$ -algebra \mathbb{A} and any function $f : X \rightarrow \mathbb{A}$, there exists a unique homomorphism $f^* : X^* \rightarrow \mathbb{A}$ such that $f^*(x) = f(x)$ for all $x \in X \subseteq X^*$.¹⁰² This can be summarized in the following diagram, where X^* denotes both the set of words and the monoid.

$$\begin{array}{ccc}
 & \text{in Set} & \text{in Alg}(\Sigma_{\text{Mon}}, E_{\text{Mon}}) \\
 X & \xleftrightarrow{\quad} & X^* \\
 & \searrow f & \downarrow f^* \\
 & & \mathbb{A} \\
 & & \leftarrow U \\
 & & X^* \\
 & & \downarrow f^* \\
 & & \mathbb{A}
 \end{array} \quad (1.23)$$

A consequence of (1.23) which makes the idea of freeness more concrete is that X^* satisfies an equation $X \vdash s = t$ if and only if all $(\Sigma_{\text{Mon}}, E_{\text{Mon}})$ -algebras satisfy it.¹⁰³ In other words, X^* only satisfies the equations it *needs* to satisfy.

The free $(\Sigma_{\text{Mon}}, E_{\text{Mon}})$ -algebra over any set is always¹⁰⁴ the set of finite words over that set with \cdot and e interpreted as concatenation and the empty word respectively.

At a first look, X^* does not seem correlated to the operation symbols in Σ_{Mon} and the equations in E_{Mon} , so it may seem hopeless to generalize this construction of free algebra for an arbitrary Σ and E . It is possible however to describe the algebra X^* starting from Σ_{Mon} and E_{Mon} .

Recall that $\mathcal{T}_{\Sigma_{\text{Mon}}} X$ is the set of all terms constructed with the symbols in Σ_{Mon} and the elements of X .¹⁰⁵ Since we want the interpretation of e to be a neutral element for the interpretation of \cdot , we could identify many terms together like e and $e \cdot e$, in fact whenever a term has an occurrence of e , we can remove it with no effect on its interpretation in a $(\Sigma_{\text{Mon}}, E_{\text{Mon}})$ -algebra. Similarly, since we want \cdot to be interpreted as an associative operation, we could identify $r \cdot (s \cdot m)$ and $(r \cdot s) \cdot m$, and more generally, we can rearrange the parentheses in a term with no effect on its interpretation in a $(\Sigma_{\text{Mon}}, E_{\text{Mon}})$ -algebra.

Squinting a bit, you can convince yourself that a Σ_{Mon} -term over X considered modulo occurrences of e and parentheses is the same thing as a finite word in X^* .¹⁰⁶ Under this correspondence, we find that the interpretation of \cdot on X^* (which was concatenation) can be realized syntactically by the symbol \cdot . For example, the

¹⁰⁰ We are talking about words in a mathematical sense, so X^* contains weird stuff like $aczlp$ and the empty word ε .

¹⁰¹ It does not satisfy $x, y \vdash x \cdot y = y \cdot x$ asserting commutativity because ab and ba are two different words.

¹⁰² f^* sends $x_1 \dots x_n$ to $\llbracket f(x_1) \cdot (f(x_2) \cdot \dots \cdot f(x_n)) \rrbracket_{\mathbb{A}}$.

¹⁰³ The forward direction uses Lemma 1.20 with ι being the inclusion $X \hookrightarrow X^*$ and h being f^* . The converse direction is trivial since we know X^* belongs to $\text{Alg}(\Sigma_{\text{Mon}}, E_{\text{Mon}})$.

¹⁰⁴ We have to say “up to isomorphism” here if we want to be fully rigorous. Let us avoid this bulkiness here and later in most places where it can be inferred.

¹⁰⁵ For instance, it contains $e, e \cdot e, a \cdot a, a \cdot (e \cdot u)$, and so on.

¹⁰⁶ For instance, both $r \cdot (s \cdot m)$ and $(r \cdot s) \cdot m$ become the word rsm and $e, e \cdot e$ and $e \cdot (e \cdot e)$ all become the empty word.

concatenation of the words corresponding to $r \cdot r$ and $u \cdot p$ is the word corresponding to $(r \cdot r) \cdot (u \cdot p)$. The interpretation of e in X^* is the empty word which corresponds to e . We conclude that the algebra X^* could have been described entirely using the syntax of Σ_{Mon} and equations in E_{Mon} .

We promptly generalize this to other signatures and sets of equations. Fix a signature Σ and a class E of equations over Σ . For any set X , we can define a binary relation \equiv_E on Σ -terms¹⁰⁷ that contains the pair (s, t) whenever the interpretation of s and t coincide in any (Σ, E) -algebra. Formally, we have for any $s, t \in \mathcal{T}_\Sigma X$,

$$s \equiv_E t \iff X \vdash s = t \in \mathfrak{Th}(E). \quad (1.24)$$

We now show \equiv_E is a congruence relation on $\mathcal{T}_\Sigma X$.¹⁰⁸

Lemma 1.33. *For any set X , the relation \equiv_E is reflexive, symmetric, transitive, and satisfies for any $\text{op} : n \in \Sigma$ and $s_1, \dots, s_n, t_1, \dots, t_n \in \mathcal{T}_\Sigma X$,*

$$(\forall 1 \leq i \leq n, s_i \equiv_E t_i) \implies \text{op}(s_1, \dots, s_n) \equiv_E \text{op}(t_1, \dots, t_n). \quad (1.25)$$

Proof. Briefly, reflexivity, symmetry, and transitivity all follow from the fact that equality satisfies these properties, and (1.25) follows from the fact that operation symbols are interpreted as *deterministic* functions (a unique output for each input), so they preserve equality. We detail this below.

(Reflexivity) For any $t \in \mathcal{T}_\Sigma X$, and any Σ -algebra \mathbb{A} , $\mathbb{A} \models X \vdash t = t$ because it holds that $\llbracket t \rrbracket'_A = \llbracket t \rrbracket'_A$ for all $\iota : X \rightarrow A$.

(Symmetry) For any $s, t \in \mathcal{T}_\Sigma X$ and $\mathbb{A} \in \mathbf{Alg}(\Sigma)$, if $\mathbb{A} \models X \vdash s = t$, then $\mathbb{A} \models X \vdash t = s$. Indeed, if $\llbracket s \rrbracket'_A = \llbracket t \rrbracket'_A$ holds for all ι , then $\llbracket t \rrbracket'_A = \llbracket s \rrbracket'_A$ holds too. Symmetry follows because if all (Σ, E) -algebras satisfy $X \vdash s = t$, then they also satisfy $X \vdash t = s$.

(Transitivity) For any $s, t, u \in \mathcal{T}_\Sigma X$, if all (Σ, E) -algebras satisfy $X \vdash s = t$ and $X \vdash t = u$, then they also satisfy $X \vdash s = u$.¹⁰⁹ Transitivity follows.

(1.25) For any $\text{op} : n \in \Sigma$, $s_1, \dots, s_n, t_1, \dots, t_n \in \mathcal{T}_\Sigma X$, and $\mathbb{A} \in \mathbf{Alg}(\Sigma)$, if \mathbb{A} satisfies $X \vdash s_i = t_i$ for all i , then for any assignment $\iota : X \rightarrow A$, we have $\llbracket s_i \rrbracket'_A = \llbracket t_i \rrbracket'_A$ for all i . Hence,

$$\begin{aligned} \llbracket \text{op}(s_1, \dots, s_n) \rrbracket'_A &= \llbracket \text{op} \rrbracket_A(\llbracket s_1 \rrbracket'_A, \dots, \llbracket s_n \rrbracket'_A) && \text{by (1.9)} \\ &= \llbracket \text{op} \rrbracket_A(\llbracket t_1 \rrbracket'_A, \dots, \llbracket t_n \rrbracket'_A) && \forall i, \llbracket s_i \rrbracket'_A = \llbracket t_i \rrbracket'_A \\ &= \llbracket \text{op}(t_1, \dots, t_n) \rrbracket'_A && \text{by (1.9),} \end{aligned}$$

which means $\mathbb{A} \models X \vdash \text{op}(s_1, \dots, s_n) = \text{op}(t_1, \dots, t_n)$. This was true for all Σ -algebras, so we can use the same arguments as above to conclude (1.25). \square

This lemma shows \equiv_E is in particular an equivalence relation, so we can define terms modulo E . Given Σ , E , and X , let $\mathcal{T}_{\Sigma, E} X = \mathcal{T}_\Sigma X / \equiv_E$ denote the set of Σ -terms modulo E . We will write $[-]_E : \mathcal{T}_\Sigma X \rightarrow \mathcal{T}_{\Sigma, E} X$ for the canonical quotient map, so $[t]_E$ is the equivalence class of t in $\mathcal{T}_{\Sigma, E} X$.

This yields a functor $\mathcal{T}_{\Sigma, E} : \mathbf{Set} \rightarrow \mathbf{Set}$ which sends a function $f : X \rightarrow Y$ to the unique function $\mathcal{T}_{\Sigma, E} f$ making (1.26) commute, i.e. satisfying $\mathcal{T}_{\Sigma, E} f([t]_E) = [\mathcal{T}_\Sigma f(t)]_E$. By definition, $[-]_E$ is also a natural transformation from \mathcal{T}_Σ to $\mathcal{T}_{\Sigma, E}$.

$$\begin{array}{ccc} \mathcal{T}_\Sigma X & \xrightarrow{[-]_E} & \mathcal{T}_{\Sigma, E} X \\ \mathcal{T}_\Sigma f \downarrow & & \downarrow \mathcal{T}_{\Sigma, E} f \\ \mathcal{T}_\Sigma Y & \xrightarrow{[-]_E} & \mathcal{T}_{\Sigma, E} Y \end{array} \quad (1.26)$$

¹⁰⁷ We omit the set X from the notation as it would be more bulky than illuminating.

¹⁰⁸ A **congruence** on a Σ -algebra \mathbb{A} is an equivalence relation $\sim \subseteq A \times A$ on the carrier satisfying for all $\text{op} : n \in \Sigma$ and $a_1, \dots, a_n, b_1, \dots, b_n \in A$:

$$(\forall i, a_i \sim b_i) \implies \llbracket \text{op} \rrbracket_A(a_1, \dots, a_n) \sim \llbracket \text{op} \rrbracket_A(b_1, \dots, b_n).$$

¹⁰⁹ Just like for symmetry, it is because for any $\mathbb{A} \in \mathbf{Alg}(\Sigma)$ and $\iota : X \rightarrow A$, $\llbracket s \rrbracket'_A = \llbracket t \rrbracket'_A$ with $\llbracket t \rrbracket'_A = \llbracket u \rrbracket'_A$ imply $\llbracket s \rrbracket'_A = \llbracket u \rrbracket'_A$.

Definition 1.34 (Term algebra, semantically). The **term algebra** for (Σ, E) on X is the Σ -algebra whose carrier is $\mathcal{T}_{\Sigma, E}X$ and whose interpretation of $\text{op} : n \in \Sigma$ is¹¹⁰

$$\llbracket \text{op} \rrbracket_{\text{TX}}(\llbracket t_1 \rrbracket_E, \dots, \llbracket t_n \rrbracket_E) = \llbracket \text{op}(t_1, \dots, t_n) \rrbracket_E. \quad (1.27)$$

We denote this algebra by $\mathbb{T}_{\Sigma, E}X$ or simply TX .

A main motivation behind this definition is that it makes $[-]_E : \mathcal{T}_{\Sigma}X \rightarrow \mathcal{T}_{\Sigma, E}X$ a homomorphism,¹¹¹ namely, (1.28) commutes.

$$\begin{array}{ccc} \mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma} X & \xrightarrow{\mathcal{T}_{\Sigma}[-]_E} & \mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma, E} X \\ \mu_X^{\Sigma} \downarrow & & \downarrow \llbracket - \rrbracket_{\text{TX}} \\ \mathcal{T}_{\Sigma} X & \xrightarrow{[-]_E} & \mathcal{T}_{\Sigma, E} X \end{array} \quad (1.28)$$

Remark 1.35. We can understand Definition 1.34 a bit more abstractly. If \mathbb{A} is a Σ -algebra and $\sim \subseteq A \times A$ is a congruence, then the quotient A/\sim inherits a Σ -algebra structure defined as in (1.27) ($[a]$ denotes the equivalence class of a in A/\sim):

$$\llbracket \text{op} \rrbracket_{A/\sim}(\llbracket a_1 \rrbracket, \dots, \llbracket a_n \rrbracket) = \llbracket \llbracket \text{op} \rrbracket_A(a_1, \dots, a_n) \rrbracket.$$

Then, $\mathbb{T}_{\Sigma, E}X$ is the quotient of the algebra $\mathcal{T}_{\Sigma}X$ defined in Remark 1.24 by the congruence \equiv_E . From this point of view, one can give an equivalent definition of \equiv_E as the smallest congruence on $\mathcal{T}_{\Sigma}X$ such that the quotient satisfies E .¹¹²

It is very easy to *compute* in the term algebra because all operations are realized syntactically, that is, only by manipulating symbols. Let us first look at the interpretation of Σ -terms in TX , i.e. the function $\llbracket - \rrbracket_{\text{TX}} : \mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma, E} X \rightarrow \mathcal{T}_{\Sigma, E} X$. It was defined inductively to yield¹¹³

$$\llbracket \eta_{\mathcal{T}_{\Sigma, E} X}^{\Sigma}(\llbracket t \rrbracket_E) \rrbracket_{\text{TX}} = \llbracket t \rrbracket_E \text{ and } \llbracket \text{op}(t_1, \dots, t_n) \rrbracket_{\text{TX}} = \llbracket \text{op} \rrbracket_{\text{TX}}(\llbracket t_1 \rrbracket_{\text{TX}}, \dots, \llbracket t_n \rrbracket_{\text{TX}}). \quad (1.29)$$

Remark 1.36. In particular, when E is empty, the set $\mathcal{T}_{\Sigma, \emptyset} X$ is $\mathcal{T}_{\Sigma} X$ quotiented by \equiv_{\emptyset} , and one can show that \equiv_{\emptyset} is equal to equality ($=$), i.e. $\mathfrak{Th}(\emptyset)$ only contains equation of the form $X \vdash t = t$.¹¹⁴ Therefore, $\mathcal{T}_{\Sigma, \emptyset} X = \mathcal{T}_{\Sigma} X$. Moreover, since $[-]_{\emptyset}$ is the identity map, we find that (1.27) becomes the definition of the interpretations given in Remark 1.24, so $\mathbb{T}_{\Sigma, \emptyset} X$ is the algebra on $\mathcal{T}_{\Sigma} X$ we had defined. Also, we find the interpretation of terms $\llbracket - \rrbracket_{\mathbb{T}_{\Sigma, \emptyset} X}$ is the flattening.¹¹⁵

Example 1.37. Let $\Sigma = \Sigma_{\text{Mon}}$ and $E = E_{\text{Mon}}$ be the signature and equations defining monoids as explained in Example 1.32. We saw informally that $\mathcal{T}_{\Sigma, E} X$ is in correspondence with the set X^* of finite words over X , and we already have a monoid structure on X^* .¹¹⁶ Thus, we may wonder whether the term algebra TX describes the same monoid. Let us compute the interpretation of $u \cdot (v \cdot w)$ where $u = uu$, $v = vv$ and $w = www$ are words in $X^* \cong \mathcal{T}_{\Sigma, E} X$. First we use the inductive definition:

$$\llbracket u \cdot (v \cdot w) \rrbracket_{\text{TX}} = \llbracket \cdot \rrbracket_{\text{TX}}(\llbracket u \rrbracket_{\text{TX}}, \llbracket v \cdot w \rrbracket_{\text{TX}}) = \llbracket \cdot \rrbracket_{\text{TX}}(\llbracket u \rrbracket_{\text{TX}}, \llbracket \cdot \rrbracket_{\text{TX}}(\llbracket v \rrbracket_{\text{TX}}, \llbracket w \rrbracket_{\text{TX}})).$$

¹¹⁰ This is well-defined (i.e. invariant under change of representative) by (1.25).

¹¹¹ Indeed, (1.27) looks exactly like (1.2) with $h = [-]_E$, $\mathbb{A} = \mathcal{T}_{\Sigma} X$ and $\mathbb{B} = \text{TX}$.

¹¹² Namely, if $\mathcal{T}_{\Sigma} X/\sim$ satisfies E , then $\equiv_E \subseteq \sim$.

¹¹³ where $t \in \mathcal{T}_{\Sigma} X$, $\text{op} : n \in \Sigma$, and $t_1, \dots, t_n \in \mathcal{T}_{\Sigma, E} X$.

¹¹⁴ Any other equation $X \vdash s = t$, where s and t are not the same term, is not satisfied by the Σ -algebra $\mathcal{T}_{\Sigma} X$ because the assignment $\eta_X^{\Sigma} : X \rightarrow \mathcal{T}_{\Sigma} X$ yields

$$\llbracket s \rrbracket_{\mathcal{T}_{\Sigma} X}^{\eta_X^{\Sigma}} = s \neq t = \llbracket t \rrbracket_{\mathcal{T}_{\Sigma} X}^{\eta_X^{\Sigma}}.$$

¹¹⁵ By Remark 1.24, or by comparing (1.29) when $E = \emptyset$ and the definition of μ_X^{Σ} (1.7).

¹¹⁶ The interpretation of \cdot and e is concatenation and the empty word.

Next, we choose a representative for $u, v, w \in \mathcal{T}_{\Sigma, E}X$ and apply the base step of the inductive definition:

$$\llbracket u \cdot (v \cdot w) \rrbracket_{\text{TX}} = \llbracket \cdot \rrbracket_{\text{TX}}(\llbracket u \cdot u \rrbracket_E, \llbracket \cdot \rrbracket_{\text{TX}}(\llbracket v \cdot v \rrbracket_E, \llbracket w \cdot (w \cdot w) \rrbracket_E)).$$

Finally, we can apply (1.27) a couple of times to find

$$\llbracket u \cdot (v \cdot w) \rrbracket_{\text{TX}} = \llbracket \cdot \rrbracket_{\text{TX}}(\llbracket u \cdot u \rrbracket_E, \llbracket (v \cdot v) \cdot (w \cdot (w \cdot w)) \rrbracket_E) = \llbracket (u \cdot u) \cdot ((v \cdot v) \cdot (w \cdot (w \cdot w))) \rrbracket_E,$$

which means that the word corresponding to $\llbracket u \cdot (v \cdot w) \rrbracket_{\text{TX}}$ is $uuvvwww$, i.e. the concatenation of u , v and w .

In general (for other signatures), what happens when applying $\llbracket - \rrbracket_{\text{TX}}$ to some big term in $\mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma, E}X$ can be decomposed in three steps.

1. Apply the inductive definition until you have an expression built out of many $\llbracket \text{op} \rrbracket_{\text{TX}}$ and $\llbracket c \rrbracket_{\text{TX}}$ where $\text{op} \in \Sigma$ and c is an equivalence class of Σ -terms.
2. Choose a representative for each such classes (i.e. $c = [t]_E$).
3. Use (1.27) repeatedly until the result is just an equivalence class in $\mathcal{T}_{\Sigma, E}X$.

Working with terms in $\mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma, E}X$ as trees whose leaves are labelled in $\mathcal{T}_{\Sigma, E}X$, $\llbracket - \rrbracket_{\text{TX}}$ replaces each leaf by the tree corresponding to a representative for the equivalence class of the leaf's label, and then returns the equivalence class of the resulting tree. In this sense, $\llbracket - \rrbracket_{\text{TX}}$ looks a lot like the flattening μ_X^{Σ} except it deals with equivalence classes of terms. This motivates the definition of $\mu_X^{\Sigma, E}$ to be the unique function making (1.30) commute.¹¹⁷

$$\begin{array}{ccc} \mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma, E}X & \xrightarrow{\llbracket - \rrbracket_{\text{TX}}} & \mathcal{T}_{\Sigma, E}X \\ & \searrow \llbracket - \rrbracket_E & \nearrow \mu_X^{\Sigma, E} \\ & \mathcal{T}_{\Sigma, E} \mathcal{T}_{\Sigma, E}X & \end{array} \quad (1.30)$$

The first thing we showed when defining μ_X^{Σ} was that it yielded a natural transformation $\mu^{\Sigma} : \mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma} \Rightarrow \mathcal{T}_{\Sigma}$. We can also do this for $\mu^{\Sigma, E}$.

Proposition 1.38. *The family of maps $\mu_X^{\Sigma, E} : \mathcal{T}_{\Sigma, E} \mathcal{T}_{\Sigma, E}X \rightarrow \mathcal{T}_{\Sigma, E}X$ is natural in X .*

Proof. We need to prove that for any function $f : X \rightarrow Y$, the square below commutes.

$$\begin{array}{ccc} \mathcal{T}_{\Sigma, E} \mathcal{T}_{\Sigma, E}X & \xrightarrow{\mathcal{T}_{\Sigma, E} \mathcal{T}_{\Sigma, E} f} & \mathcal{T}_{\Sigma, E} \mathcal{T}_{\Sigma, E}Y \\ \mu_X^{\Sigma, E} \downarrow & & \downarrow \mu_Y^{\Sigma, E} \\ \mathcal{T}_{\Sigma, E}X & \xrightarrow{\mathcal{T}_{\Sigma, E} f} & \mathcal{T}_{\Sigma, E}Y \end{array} \quad (1.31)$$

We can pave the following diagram.¹¹⁸

¹¹⁷ This guarantees $\mu_X^{\Sigma, E}$ satisfies the following equations that look like the inductive definition of μ_X^{Σ} in (1.7): for any $t \in \mathcal{T}_{\Sigma}X$, $\mu_X^{\Sigma, E}(\llbracket [t]_E \rrbracket_E) = [t]_E$, and for any $\text{op} : n \in \Sigma$ and $t_1, \dots, t_n \in \mathcal{T}_{\Sigma}X$,

$$\mu_X^{\Sigma, E}(\llbracket \text{op}([t_1]_E, \dots, [t_n]_E) \rrbracket_E) = \llbracket \text{op}(t_1, \dots, t_n) \rrbracket_E.$$

Thanks to Remark 1.36, we can immediately see that $\mu_X^{\Sigma, \emptyset} = \mu_X^{\Sigma}$ because $\llbracket - \rrbracket_{\emptyset}$ is the identity and $\llbracket - \rrbracket_{\mathcal{T}_{\Sigma, \emptyset}X} = \mu_X^{\Sigma}$.

¹¹⁸ By paving a diagram, we mean to build a large diagram out of smaller ones, showing all the smaller ones commute, and then concluding the bigger must commute. We often refer parts of the diagram with letters written inside them, and explain how each of them commutes one at a time.

$$\begin{array}{ccccc}
 \mathcal{T}_\Sigma \mathcal{T}_{\Sigma,E} X & \xrightarrow{[-]_E} & \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} X & \xrightarrow{\mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} f} & \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} Y \\
 \downarrow [-]_E & \searrow \mathcal{T}_\Sigma \mathcal{T}_{\Sigma,E} f & \nearrow & \nearrow & \downarrow \mu_Y^{\Sigma,E} \\
 & & \mathcal{T}_\Sigma \mathcal{T}_{\Sigma,E} Y & & \\
 & & \nearrow [-]_E & & \\
 & & & & \\
 \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} X & \xrightarrow{\mu_X^{\Sigma,E}} & \mathcal{T}_{\Sigma,E} X & \xrightarrow{\mathcal{T}_{\Sigma,E} f} & \mathcal{T}_{\Sigma,E} Y \\
 & & \searrow [-]_{\text{TX}} & \searrow [-]_{\text{TY}} & \\
 & & & &
 \end{array}$$

All of (a), (b) and (d) commute by definition. In more details, (a) is an instance of (1.26) with X replaced by $\mathcal{T}_{\Sigma,E} X$, Y by $\mathcal{T}_{\Sigma,E} Y$ and f by $\mathcal{T}_{\Sigma,E} f$, and both (b) and (d) are instances of (1.30). To show (c) commutes, we draw another diagram that looks like a cube with (c) as the front face.

$$\begin{array}{ccccc}
 \mathcal{T}_\Sigma \mathcal{T}_\Sigma X & \xrightarrow{\mathcal{T}_\Sigma \mathcal{T}_\Sigma f} & \mathcal{T}_\Sigma \mathcal{T}_\Sigma Y & & \\
 \downarrow \mu_X^\Sigma & \searrow \mathcal{T}_\Sigma [-]_E & \downarrow \mu_Y^\Sigma & \searrow \mathcal{T}_\Sigma [-]_E & \\
 \mathcal{T}_\Sigma \mathcal{T}_{\Sigma,E} X & \xrightarrow{\mathcal{T}_\Sigma \mathcal{T}_{\Sigma,E} f} & \mathcal{T}_\Sigma \mathcal{T}_{\Sigma,E} Y & & \\
 \downarrow [-]_{\text{TX}} & \searrow \mathcal{T}_\Sigma f & \downarrow [-]_{\text{TY}} & \searrow [-]_E & \\
 \mathcal{T}_\Sigma X & \xrightarrow{\mathcal{T}_\Sigma f} & \mathcal{T}_\Sigma Y & & \\
 \downarrow [-]_E & \searrow & \downarrow [-]_E & \searrow & \\
 \mathcal{T}_{\Sigma,E} X & \xrightarrow{\mathcal{T}_{\Sigma,E} f} & \mathcal{T}_{\Sigma,E} Y & &
 \end{array}$$

We can show all the other faces commute, and then use the fact that $\mathcal{T}_\Sigma [-]_E$ is surjective (i.e. epic) to conclude that the front face must also commute.¹¹⁹ The first diagram we paved implies (1.31) commutes because $[-]_E$ is epic. \square

The front face of the cube is interesting on its own, it says that for any function $f : X \rightarrow Y$, $\mathcal{T}_{\Sigma,E} f$ is a homomorphism from $\mathcal{T}_{\Sigma,E} X$ to $\mathcal{T}_{\Sigma,E} Y$. We redraw it below for future reference.

$$\begin{array}{ccc}
 \mathcal{T}_\Sigma \mathcal{T}_{\Sigma,E} X & \xrightarrow{\mathcal{T}_\Sigma \mathcal{T}_{\Sigma,E} f} & \mathcal{T}_\Sigma \mathcal{T}_{\Sigma,E} Y \\
 \llbracket - \rrbracket_{\text{TX}} \downarrow & & \downarrow \llbracket - \rrbracket_{\text{TY}} \\
 \mathcal{T}_{\Sigma,E} X & \xrightarrow{\mathcal{T}_{\Sigma,E} f} & \mathcal{T}_{\Sigma,E} Y
 \end{array} \quad (1.32)$$

Stating it like this may remind you of Lemma 1.23 and Remark 1.24. We will need a variant of Lemma 1.23 for $\mathcal{T}_{\Sigma,E}$, but there is a slight obstacle due to types. Indeed, given a Σ -algebra \mathbb{A} we would like to prove a square like in (1.33) commutes.

However, the arrows on top and bottom do not really exist, the interpretation $\llbracket - \rrbracket_A$ takes terms over A as input, not equivalence classes of terms. The quick fix is to assume that \mathbb{A} satisfies the equations in E . This means that $\llbracket - \rrbracket_A$ is well-defined on equivalence class of terms because if $[s]_E = [t]_E$, then $A \vdash s = t \in \mathfrak{Th}(E)$, so \mathbb{A}

¹¹⁹ In more details, the left and right faces commute by (1.28), the bottom and top faces commute by (1.26), and the back face commutes by (1.8).

The function $\mathcal{T}_\Sigma [-]_E$ is surjective (i.e. epic) because $[-]_E$ is (it is a canonical quotient map) and functors on **Set** preserve epimorphisms (if we assume the axiom of choice). Thus, it suffices to show that $\mathcal{T}_\Sigma [-]_E$ pre-composed with the bottom path or the top path of the front face gives the same result.

Now it is just a matter of going around the cube using the commutativity of the other faces. Here is the complete derivation (we write which face was used as justifications for each step).

$$\begin{aligned}
 & \mathcal{T}_{\Sigma,E} f \circ \llbracket - \rrbracket_{\text{TX}} \circ \mathcal{T}_\Sigma [-]_E && \text{left} \\
 & = \mathcal{T}_{\Sigma,E} f \circ [-]_E \circ \mu_X^\Sigma && \text{bottom} \\
 & = [-]_E \circ \mathcal{T}_\Sigma f \circ \mu_X^\Sigma && \text{back} \\
 & = [-]_E \circ \mu_Y^\Sigma \circ \mathcal{T}_\Sigma \mathcal{T}_\Sigma f && \text{right} \\
 & = \llbracket - \rrbracket_{\text{TY}} \circ \mathcal{T}_\Sigma [-]_E \circ \mathcal{T}_\Sigma \mathcal{T}_\Sigma f && \text{right} \\
 & = \llbracket - \rrbracket_{\text{TY}} \circ \mathcal{T}_\Sigma \mathcal{T}_{\Sigma,E} f \circ \mathcal{T}_\Sigma [-]_E && \text{top}
 \end{aligned}$$

$$\begin{array}{ccc}
 \mathcal{T}_\Sigma \mathcal{T}_{\Sigma,E} A & \xrightarrow{\mathcal{T}_\Sigma \llbracket - \rrbracket_A} & \mathcal{T}_\Sigma A \\
 \llbracket - \rrbracket_{\text{TA}} \downarrow & & \downarrow \llbracket - \rrbracket_A \\
 \mathcal{T}_{\Sigma,E} A & \xrightarrow{\llbracket - \rrbracket_A} & A
 \end{array} \quad (1.33)$$

satisfies that equation, and taking the assignment $\text{id}_A : A \rightarrow A$, we obtain

$$\llbracket s \rrbracket_A = \llbracket s \rrbracket_A^{\text{id}_A} = \llbracket t \rrbracket_A^{\text{id}_A} = \llbracket t \rrbracket_A.$$

When \mathbb{A} is a (Σ, E) -algebra, we abusively write $\llbracket - \rrbracket_A$ for the interpretation of terms and equivalence classes of terms as in (1.34).

$$\begin{array}{ccc} \mathcal{T}_\Sigma A & \xrightarrow{\llbracket - \rrbracket_A} & A \\ & \searrow \llbracket - \rrbracket_E & \nearrow \llbracket - \rrbracket_A \\ & \mathcal{T}_{\Sigma, E} A & \end{array} \quad (1.34)$$

Lemma 1.39. *For any (Σ, E) -algebra \mathbb{A} , the square (1.33) commutes.*

Proof. Consider the following diagram that we can view as a triangular prism whose front face is (1.33). Both triangles commute by (1.34), the square face at the back and on the left commutes by (1.28), and the square face at the back and on the right commutes by (1.14). With the same trick as in the proof of Proposition 1.38 using the surjectivity of $\mathcal{T}_\Sigma[-]_E$, we conclude that the front face commutes.¹²⁰

$$\begin{array}{ccccc} & & \mathcal{T}_\Sigma \mathcal{T}_\Sigma A & & \\ & \mathcal{T}_\Sigma[-]_E \swarrow & \downarrow \mathcal{T}_\Sigma \llbracket - \rrbracket_A & \searrow \mathcal{T}_\Sigma[-]_A & \\ \mathcal{T}_\Sigma \mathcal{T}_{\Sigma, E} A & \xrightarrow{\quad} & \mathcal{T}_\Sigma A & \xrightarrow{\quad} & \mathcal{T}_\Sigma A \\ & \downarrow \llbracket - \rrbracket_{\text{TA}} & \downarrow \mu_A^\Sigma & \downarrow \llbracket - \rrbracket_A & \\ & \mathcal{T}_{\Sigma, E} A & \mathcal{T}_\Sigma A & \mathcal{T}_\Sigma A & \\ & \swarrow \llbracket - \rrbracket_E & \downarrow \llbracket - \rrbracket_A & \searrow \llbracket - \rrbracket_A & \\ & \mathcal{T}_{\Sigma, E} A & \xrightarrow{\quad} & A & \\ & & \llbracket - \rrbracket_A & & \end{array}$$

□

An important consequence of Lemma 1.23 was (1.16) saying that flattening is a homomorphism from $\mathbb{T}_{\Sigma, \emptyset} \mathbb{T}_{\Sigma, \emptyset} A$ to $\mathbb{T}_{\Sigma, \emptyset} A$. This is also true when E is not empty, i.e. $\mu_A^{\Sigma, E}$ is a homomorphism from $\mathbb{T}A$ to $\mathbb{T}A$.

Lemma 1.40. *For any set A , the following square commutes.*

$$\begin{array}{ccc} \mathcal{T}_\Sigma \mathcal{T}_{\Sigma, E} \mathcal{T}_{\Sigma, E} A & \xrightarrow{\mathcal{T}_\Sigma \mu_A^{\Sigma, E}} & \mathcal{T}_\Sigma \mathcal{T}_{\Sigma, E} A \\ \llbracket - \rrbracket_{\mathbb{T}A} \downarrow & & \downarrow \llbracket - \rrbracket_{\mathbb{T}A} \\ \mathcal{T}_{\Sigma, E} \mathcal{T}_{\Sigma, E} A & \xrightarrow{\mu_A^{\Sigma, E}} & \mathcal{T}_{\Sigma, E} A \end{array} \quad (1.35)$$

Proof. We prove it exactly like Lemma 1.39 with the following diagram.¹²¹

$$\begin{array}{ccccc} & & \mathcal{T}_\Sigma \mathcal{T}_\Sigma \mathcal{T}_{\Sigma, E} A & & \\ & \mathcal{T}_\Sigma[-]_E \swarrow & \downarrow \mathcal{T}_\Sigma \mu_A^{\Sigma, E} & \searrow \mathcal{T}_\Sigma[-]_{\mathbb{T}A} & \\ \mathcal{T}_\Sigma \mathcal{T}_{\Sigma, E} \mathcal{T}_{\Sigma, E} A & \xrightarrow{\quad} & \mathcal{T}_\Sigma \mathcal{T}_{\Sigma, E} A & \xrightarrow{\quad} & \mathcal{T}_\Sigma \mathcal{T}_{\Sigma, E} A \\ & \downarrow \llbracket - \rrbracket_{\mathbb{T}A} & \downarrow \mu_A^{\Sigma, E} & \downarrow \llbracket - \rrbracket_{\mathbb{T}A} & \\ & \mathcal{T}_{\Sigma, E} \mathcal{T}_{\Sigma, E} A & \mathcal{T}_{\Sigma, E} A & \mathcal{T}_{\Sigma, E} A & \\ & \swarrow \llbracket - \rrbracket_E & \downarrow \llbracket - \rrbracket_{\mathbb{T}A} & \searrow \llbracket - \rrbracket_{\mathbb{T}A} & \\ & \mathcal{T}_{\Sigma, E} \mathcal{T}_{\Sigma, E} A & \xrightarrow{\mu_A^{\Sigma, E}} & \mathcal{T}_{\Sigma, E} A & \end{array}$$

¹²⁰ Here is the complete derivation.

$$\begin{aligned} & \llbracket - \rrbracket_A \circ \llbracket - \rrbracket_{\mathbb{T}A} \circ \mathcal{T}_\Sigma[-]_E && \\ & = \llbracket - \rrbracket_A \circ \llbracket - \rrbracket_E \circ \mu_A^\Sigma && \text{left} \\ & = \llbracket - \rrbracket_A \circ \mu_A^\Sigma && \text{bottom} \\ & = \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma \llbracket - \rrbracket_A && \text{right} \\ & = \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma[-]_E && \text{top} \end{aligned}$$

Then, since $\mathcal{T}_\Sigma[-]_E$ is epic, we conclude that $\llbracket - \rrbracket_A \circ \llbracket - \rrbracket_{\mathbb{T}A} = \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma \llbracket - \rrbracket_A$.

¹²¹ The top and bottom faces commute by definition of $\mu_A^{\Sigma, E}$ (1.30), the back-left face by (1.28), and the back-right face by (1.14).

Then, $\mathcal{T}_\Sigma[-]_E$ is epic, so the following derivation suffices.

$$\begin{aligned} & \mu_A^{\Sigma, E} \circ \llbracket - \rrbracket_{\mathbb{T}A} \circ \mathcal{T}_\Sigma[-]_E && \\ & = \mu_A^{\Sigma, E} \circ \llbracket - \rrbracket_E \circ \mu_{\Sigma, E}^\Sigma && \text{left} \\ & = \llbracket - \rrbracket_{\mathbb{T}A} \circ \mu_{\Sigma, E}^\Sigma && \text{bottom} \\ & = \llbracket - \rrbracket_{\mathbb{T}A} \circ \mathcal{T}_\Sigma \llbracket - \rrbracket_{\mathbb{T}A} && \text{right} \\ & = \llbracket - \rrbracket_{\mathbb{T}A} \circ \mathcal{T}_\Sigma \mu_A^{\Sigma, E} \circ \mathcal{T}_\Sigma[-]_E && \text{top} \end{aligned}$$

□

In a moment, we will show that $\mathbb{T}_{\Sigma,E}X$ is not only a Σ -algebra, but also a (Σ, E) -algebra. This requires us to talk about satisfaction of equations, hence about the interpretation of terms in some $\mathcal{T}_E Y$ under an assignment $\sigma : Y \rightarrow \mathcal{T}_{\Sigma,E}X$.¹²² By the definition $\llbracket - \rrbracket_{\mathbb{T}X}^\sigma = \llbracket - \rrbracket_{\mathbb{T}X} \circ \mathcal{T}_E \sigma$, and our informal description of $\llbracket - \rrbracket_{\mathbb{T}X}$, we can infer that $\llbracket t \rrbracket_{\mathbb{T}X}^\sigma$ is the equivalence class of the term t where all occurrences of the variable y have been substituted by a representative of $\sigma(y)$.

In particular, this means that under the assignment $\sigma : X \rightarrow \mathcal{T}_{\Sigma,E}X$ that sends a variable x to its equivalence class $[x]_E$, the interpretation of a term $t \in \mathcal{T}_E X$ is $[t]_E$.¹²³ We prove this formally below.

Lemma 1.41. *Let $\sigma = X \xrightarrow{\eta_X^\Sigma} \mathcal{T}_E X \xrightarrow{[-]_E} \mathcal{T}_{\Sigma,E}X$ be an assignment. Then, $\llbracket - \rrbracket_{\mathbb{T}X}^\sigma = [-]_E$.*

Proof. We proceed by induction. For the base case, we have

$$\begin{aligned} \llbracket \eta_X^\Sigma(x) \rrbracket_{\mathbb{T}X}^\sigma &= \llbracket \mathcal{T}_E \sigma(\eta_X^\Sigma(x)) \rrbracket_{\mathbb{T}X} && \text{by (1.10)} \\ &= \llbracket \mathcal{T}_E [-]_E(\mathcal{T}_E \eta_X^\Sigma(\eta_X^\Sigma(x))) \rrbracket_{\mathbb{T}X} && \text{Lemma 1.11} \\ &= \llbracket \mathcal{T}_E [-]_E(\eta_{\mathcal{T}_E X}^\Sigma(\eta_X^\Sigma(x))) \rrbracket_{\mathbb{T}X} && \text{by (1.6)} \\ &= \llbracket \eta_{\mathcal{T}_{\Sigma,E}X}^\Sigma([\eta_X^\Sigma(x)]_E) \rrbracket_{\mathbb{T}X} && \text{by (1.6)} \\ &= [\eta_X^\Sigma(x)]_E && \text{by (1.29)} \end{aligned}$$

For the inductive step, if $t = \text{op}(t_1, \dots, t_n)$, we have

$$\begin{aligned} \llbracket t \rrbracket_{\mathbb{T}X}^\sigma &= \llbracket \mathcal{T}_E \sigma(t) \rrbracket_{\mathbb{T}X} && \text{by (1.10)} \\ &= \llbracket \mathcal{T}_E \sigma(\text{op}(t_1, \dots, t_n)) \rrbracket_{\mathbb{T}X} \\ &= \llbracket \text{op}(\mathcal{T}_E \sigma(t_1), \dots, \mathcal{T}_E \sigma(t_n)) \rrbracket_{\mathbb{T}X} && \text{by (1.5)} \\ &= \llbracket \text{op} \rrbracket_{\mathbb{T}X} (\llbracket \mathcal{T}_E \sigma(t_1) \rrbracket_{\mathbb{T}X}, \dots, \llbracket \mathcal{T}_E \sigma(t_n) \rrbracket_{\mathbb{T}X}) && \text{by (1.29)} \\ &= \llbracket \text{op} \rrbracket_{\mathbb{T}X} ([t_1]_E, \dots, [t_n]_E) && \text{I.H.} \\ &= [\text{op}(t_1, \dots, t_n)]_E. && \text{by (1.27)} \quad \square \end{aligned}$$

We will denote that special assignment $\eta_X^{\Sigma,E} = [-]_E \circ \eta_X^\Sigma : X \rightarrow \mathcal{T}_{\Sigma,E}X$.¹²⁴ A quick corollary of the previous lemma is that for any equation ϕ with context X , ϕ belongs to $\mathfrak{Th}(E)$ if and only if the algebra $\mathbb{T}_{\Sigma,E}X$ satisfies it under the assignment $\eta_X^{\Sigma,E}$. This comes back to Example 1.32 where we said that freeness of X^* means it satisfies all and only the equations in $\mathfrak{Th}(E_{\text{Mon}})$. Instead here, we do not know yet that $\mathbb{T}X$ is free (we have not even proved it satisfies E yet), but we can already show it satisfies only the necessary equations, and freeness will follow.

Lemma 1.42. *Let $s, t \in \mathcal{T}_E X$, $X \vdash s = t \in \mathfrak{Th}(E)$ if and only if $\mathbb{T}_{\Sigma,E}X \models_{\eta_X^{\Sigma,E}} X \vdash s = t$.¹²⁵*

The interaction between μ^Σ and η^Σ is mimicked by $\mu^{\Sigma,E}$ and $\eta^{\Sigma,E}$.

¹²² We used ι before for assignments, but when considering assignments into (equivalence classes of) terms, we prefer using σ because we will adopt a different attitude with them (see Definition 1.44).

¹²³ The representative chosen for $\sigma(x)$ is x so the term t is not modified.

¹²⁴ Note that $\eta_X^{\Sigma,E}$ becomes a natural transformation $\text{id}_{\text{Set}} \rightarrow \mathcal{T}_{\Sigma,E}$ because it is the vertical composition $[-]_E \cdot \eta^\Sigma$.

¹²⁵ *Proof.* By Lemma 1.41, we have

$$\llbracket s \rrbracket_{\mathbb{T}X}^{\eta_X^{\Sigma,E}} = [s]_E \text{ and } \llbracket t \rrbracket_{\mathbb{T}X}^{\eta_X^{\Sigma,E}} = [t]_E,$$

then by definition of \equiv_E , $X \vdash s = t \in \mathfrak{Th}(E)$ if and only if $[s]_E = [t]_E$.

Lemma 1.43. *The following diagram commutes.*

$$\begin{array}{ccccc}
 \mathcal{T}_{\Sigma,E}X & \xrightarrow{\eta_{\mathcal{T}_{\Sigma,E}X}^{\Sigma,E}} & \mathcal{T}_{\Sigma,E}\mathcal{T}_{\Sigma,E}X & \xleftarrow{\mathcal{T}_{\Sigma,E}\eta_X^{\Sigma,E}} & \mathcal{T}_{\Sigma,E}X \\
 & \searrow \text{id}_{\mathcal{T}_{\Sigma,E}X} & \downarrow \mu_X^{\Sigma,E} & \swarrow \text{id}_{\mathcal{T}_{\Sigma,E}X} & \\
 & & \mathcal{T}_{\Sigma,E}X & &
 \end{array}$$

Proof. For the triangle on the left, we pave the following diagram.

$$\begin{array}{ccccc}
 & & \eta_{\mathcal{T}_{\Sigma,E}X}^{\Sigma,E} & & \\
 & & \curvearrowright & & \\
 & & \text{(a)} & & \\
 \mathcal{T}_{\Sigma,E}X & \xrightarrow{\eta_{\mathcal{T}_{\Sigma,E}X}^{\Sigma,E}} & \mathcal{T}_{\Sigma}\mathcal{T}_{\Sigma,E}X & \xrightarrow{[-]_E} & \mathcal{T}_{\Sigma,E}\mathcal{T}_{\Sigma,E}X \\
 & \searrow \text{id}_{\mathcal{T}_{\Sigma,E}X} & \downarrow \mu_X^{\Sigma,E} & \swarrow \text{(c)} & \\
 & & \mathcal{T}_{\Sigma,E}X & & \\
 & & \text{(b)} \quad \llbracket - \rrbracket_{\text{TX}} & &
 \end{array} \quad (1.36)$$

Showing (1.36) commutes:

(a) Definition of $\eta_X^{\Sigma,E}$.

(b) Definition of $\llbracket - \rrbracket_{\text{TX}}$ (1.29).

(c) Definition of $\mu_X^{\Sigma,E}$ (1.30).

For the triangle on the right, we show that $[-]_E = \mu_X^{\Sigma,E} \circ \mathcal{T}_{\Sigma,E}\eta_X^{\Sigma,E} \circ [-]_E$ by paving (1.37), and we can conclude since $[-]_E$ is epic that $\text{id}_{\mathcal{T}_{\Sigma,E}X} = \mu_X^{\Sigma,E} \circ \mathcal{T}_{\Sigma,E}\eta_X^{\Sigma,E}$.

$$\begin{array}{ccccc}
 & & \mathcal{T}_{\Sigma,E}\eta_X^{\Sigma,E} & & \\
 & & \curvearrowright & & \\
 & & \text{(a)} & & \\
 \mathcal{T}_{\Sigma}X & \xrightarrow{[-]_E} & \mathcal{T}_{\Sigma,E}X & \xrightarrow{\mathcal{T}_{\Sigma,E}\eta_X^{\Sigma,E}} & \mathcal{T}_{\Sigma,E}\mathcal{T}_{\Sigma}X & \xrightarrow{\mathcal{T}_{\Sigma,E}[-]_E} & \mathcal{T}_{\Sigma,E}\mathcal{T}_{\Sigma,E}X \\
 & \searrow \text{id}_{\mathcal{T}_{\Sigma}X} & \downarrow \mu_X^{\Sigma} & \swarrow \text{(c)} & \downarrow \mu_X^{\Sigma,E} & \swarrow \text{(f)} & \\
 & & \mathcal{T}_{\Sigma}X & \xrightarrow{[-]_E} & \mathcal{T}_{\Sigma,E}X & & \\
 & & \text{(d)} & \text{(e)} & \text{(f)} & & \\
 & & \mu_X^{\Sigma} & \llbracket - \rrbracket_{\text{TX}} & & &
 \end{array} \quad (1.37)$$

Showing (1.37) commutes:

(a) Definition of $\eta_X^{\Sigma,E}$ and functoriality of $\mathcal{T}_{\Sigma,E}$.

(b) Naturality of $[-]_E$ (1.26).

(c) Naturality of $[-]_E$ again.

(d) Definition of μ_X^{Σ} (1.7).

(e) By (1.28).

(f) By (1.30).

□

We single out another special case of interpretation in a term algebra when E is empty (recall from Remark 1.36 that $\mathbb{T}_{\Sigma,\emptyset}X$ is the algebra on $\mathcal{T}_{\Sigma}X$ whose interpretation of op applies op syntactically).

Definition 1.44 (Substitution). Given a signature Σ , an empty set of equations, and an assignment $\sigma : Y \rightarrow \mathcal{T}_{\Sigma}X$,¹²⁶ we call $\llbracket - \rrbracket_{\text{TX}}^{\sigma}$ the **substitution map**, and we denote it by $\sigma^* : \mathcal{T}_{\Sigma}Y \rightarrow \mathcal{T}_{\Sigma}X$. We saw in Remark 1.36 that $\llbracket - \rrbracket_{\text{TX}} = \mu_X^{\Sigma}$, thus substitution is

$$\sigma^* = \mathcal{T}_{\Sigma}Y \xrightarrow{\mathcal{T}_{\Sigma}\sigma} \mathcal{T}_{\Sigma}\mathcal{T}_{\Sigma}X \xrightarrow{\mu_X^{\Sigma}} \mathcal{T}_{\Sigma}X. \quad (1.38)$$

In words, σ^* replaces the occurrences of a variable y by $\sigma(y)$.¹²⁷

¹²⁶ We can identify $\mathcal{T}_{\Sigma}X$ with $\mathcal{T}_{\Sigma,\emptyset}X$ because \equiv_{\emptyset} is the equality relation.

¹²⁷ You may be more familiar with the notation $t[\sigma(y)/y]$ (e.g. from substitution in the λ -calculus). An inductive definition can also be given: for any $y \in Y$, $\sigma^*(\eta_Y^{\Sigma}(y)) = \sigma(y)$, and

$$\sigma^*(\text{op}(t_1, \dots, t_n)) = \text{op}(\sigma^*(t_1), \dots, \sigma^*(t_n)).$$

That simple description makes substitution a little special, and the following result has even deeper implications. It morally says that substitution preserves the satisfaction of equations.¹²⁸

Lemma 1.45. *Let $Y \vdash s = t$ be an equation, $\sigma : Y \rightarrow \mathcal{T}_\Sigma X$ an assignment, and \mathbb{A} a Σ -algebra. If \mathbb{A} satisfies $Y \vdash s = t$, then it also satisfies $X \vdash \sigma^*(s) = \sigma^*(t)$.*

Proof. Let $\iota : X \rightarrow \mathbb{A}$ be an assignment, we need to show $\llbracket \sigma^*(s) \rrbracket'_A = \llbracket \sigma^*(t) \rrbracket'_A$. Define the assignment $\iota_\sigma : Y \rightarrow \mathbb{A}$ that sends $y \in Y$ to $\llbracket \sigma(y) \rrbracket'_A$, we claim that $\llbracket - \rrbracket'_A = \llbracket \sigma^*(-) \rrbracket'_A$. The lemma then follows because by hypothesis, $\llbracket s \rrbracket'_A = \llbracket t \rrbracket'_A$. The following derivation proves our claim.

$$\begin{aligned}
\llbracket - \rrbracket'_A &= \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma(\iota_\sigma) && \text{by (1.10)} \\
&= \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma(\llbracket \sigma(-) \rrbracket'_A) && \text{definition of } \iota_\sigma \\
&= \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma(\llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma \iota \circ \sigma) && \text{by (1.10)} \\
&= \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma \mathcal{T}_\Sigma \iota \circ \mathcal{T}_\Sigma \sigma && \text{Lemma 1.11} \\
&= \llbracket - \rrbracket_A \circ \mu_A^\Sigma \circ \mathcal{T}_\Sigma \mathcal{T}_\Sigma \iota \circ \mathcal{T}_\Sigma \sigma && \text{by (1.14)} \\
&= \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma \iota \circ \mu_Y^\Sigma \circ \mathcal{T}_\Sigma \sigma && \text{by (1.8)} \\
&= \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma \iota \circ \sigma^* && \text{by (1.38)} \\
&= \llbracket \sigma^*(-) \rrbracket'_A. && \text{by (1.10)} \quad \square
\end{aligned}$$

We are finally ready to show that $\mathbb{T}_{\Sigma,E}A$ is a (Σ, E) -algebra.¹²⁹

Proposition 1.46. *For any set A , the term algebra $\mathbb{T}_{\Sigma,E}A$ satisfies all the equations in E .*

Proof. Let $X \vdash s = t$ belong to E and $\iota : X \rightarrow \mathbb{T}_{\Sigma,E}A$ be an assignment. We need to show that $\llbracket s \rrbracket'_{\mathbb{T}A} = \llbracket t \rrbracket'_{\mathbb{T}A}$. We factor ι into¹³⁰

$$\iota = X \xrightarrow{\eta_X^{\Sigma,E}} \mathcal{T}_{\Sigma,E}X \xrightarrow{\mathcal{T}_{\Sigma,E}\iota} \mathcal{T}_{\Sigma,E}\mathcal{T}_{\Sigma,E}A \xrightarrow{\mu_A^{\Sigma,E}} \mathbb{T}_{\Sigma,E}A.$$

Now, Lemma 1.42 says that the equation is satisfied in $\mathbb{T}X$ under the assignment $\eta_X^{\Sigma,E}$, i.e. that $\llbracket s \rrbracket'_{\mathbb{T}X} = \llbracket t \rrbracket'_{\mathbb{T}X}$. We also know by Lemma 1.20 that homomorphisms preserve satisfaction, so we can apply it twice using the facts that $\mathcal{T}_{\Sigma,E}\iota$ and $\mu_A^{\Sigma,E}$ are homomorphisms (by (1.32) and (1.35) respectively) to conclude that

$$\llbracket s \rrbracket'_{\mathbb{T}A} = \llbracket s \rrbracket'_{\mathbb{T}A} \circ \mu_A^{\Sigma,E} \circ \mathcal{T}_{\Sigma,E}\iota \circ \eta_X^{\Sigma,E} = \llbracket t \rrbracket'_{\mathbb{T}A} \circ \mu_A^{\Sigma,E} \circ \mathcal{T}_{\Sigma,E}\iota \circ \eta_X^{\Sigma,E} = \llbracket t \rrbracket'_{\mathbb{T}A}. \quad \square$$

We now know that $\mathbb{T}_{\Sigma,E}X$ belongs to $\mathbf{Alg}(\Sigma, E)$. In order to tie up the parallel with Example 1.32, we will show that $\mathbb{T}_{\Sigma,E}X$ is the free (Σ, E) -algebra over X .

Definition 1.47 (Free object). Let \mathbf{C} and \mathbf{D} be categories, $U : \mathbf{D} \rightarrow \mathbf{C}$ be a functor between them, and $X \in \mathbf{C}_0$. A **free object** on X (relative to U) is an object $Y \in \mathbf{D}_0$ along with a morphism $i \in \text{Hom}_{\mathbf{C}}(X, UY)$ such that for any object $A \in \mathbf{D}_0$ and morphism $f \in \text{Hom}_{\mathbf{C}}(X, UA)$, there exists a unique morphism $f^* \in \text{Hom}_{\mathbf{D}}(Y, A)$ such that $Uf^* \circ i = f$. This is summarized in the following diagram.¹³¹

¹²⁸ We will give more intuition on Lemma 1.45 when we define equational logic.

¹²⁹ All the work we have been doing finally pays off.

¹³⁰ This factoring is correct because

$$\begin{aligned}
\iota &= \text{id}_{\mathbb{T}_{\Sigma,E}A} \circ \iota \\
&= \mu_A^{\Sigma,E} \circ \eta_{\mathbb{T}_{\Sigma,E}A}^{\Sigma,E} \circ \iota && \text{Lemma 1.43} \\
&= \mu_A^{\Sigma,E} \circ \mathcal{T}_{\Sigma,E}\iota \circ \eta_X^{\Sigma,E}. && \text{naturality of } \eta^{\Sigma,E}
\end{aligned}$$

¹³¹ This is almost a copy of (1.23).

$$\begin{array}{ccc}
 \text{in } \mathbf{C} & & \text{in } \mathbf{D} \\
 X \xrightarrow{i} UY & & Y \\
 \searrow f & \dashrightarrow Uf^* \xleftarrow{U} & \downarrow f^* \\
 & UA & A
 \end{array} \quad (1.39)$$

Proposition 1.48. *Free objects are unique up to isomorphism, namely, if Y and Y' are free objects on X , then $Y \cong Y'$.¹³²*

Proposition 1.49. *For any set X , the term algebra $\mathbb{T}_{\Sigma,E}X$ is the free (Σ, E) -algebra on X .*

Proof. Let \mathbb{A} be another (Σ, E) -algebra and $f : X \rightarrow \mathbb{A}$ a function. We claim that $f^* = \llbracket - \rrbracket_{\mathbb{A}} \circ \mathcal{T}_{\Sigma,E}f$ is the unique homomorphism making the following commute.

$$\begin{array}{ccc}
 \text{in } \mathbf{Set} & & \text{in } \mathbf{Alg}(\Sigma, E) \\
 X \xrightarrow{\eta_X^{\Sigma,E}} \mathcal{T}_{\Sigma,E}X & & \mathbb{T}X \\
 \searrow f & \dashrightarrow f^* \xleftarrow{U} & \downarrow f^* \\
 & A & \mathbb{A}
 \end{array}$$

First, f^* is a homomorphism because it is the composite of two homomorphisms $\mathcal{T}_{\Sigma,E}f$ (by (1.32)) and $\llbracket - \rrbracket_{\mathbb{A}}$ (by Lemma 1.39 since \mathbb{A} satisfies E). Next, the triangle commutes by the following derivation.

$$\begin{aligned}
 \llbracket - \rrbracket_{\mathbb{A}} \circ \mathcal{T}_{\Sigma,E}f \circ \eta_X^{\Sigma,E} &= \llbracket - \rrbracket_{\mathbb{A}} \circ \eta_A^{\Sigma,E} \circ f && \text{naturality of } \eta^{\Sigma,E} \\
 &= \llbracket - \rrbracket_{\mathbb{A}} \circ [-]_E \circ \eta_A^{\Sigma} \circ f && \text{definition of } \eta^{\Sigma,E} \\
 &= \llbracket - \rrbracket_{\mathbb{A}} \circ \eta_A^{\Sigma} \circ f && \text{by (1.34)} \\
 &= f && \text{definition of } \llbracket - \rrbracket_{\mathbb{A}} \text{ (1.9)}
 \end{aligned}$$

Finally, uniqueness follows from the inductive definition of $\mathbb{T}X$ and the homomorphism property. Briefly, if we know the action of a homomorphism on equivalence classes of terms of depth 0, we can infer all of its action because all other classes of terms can be obtained by applying operation symbols.¹³³ \square

Once we have free objects, we have an adjunction, and once we have an adjunction, we have a monad, the most wonderful mathematical object in the world (objectively). Unfortunately, our universal algebra spiel is not finished yet, we will get back to monads shortly.

Abstract Equations

Before moving to equational logic, we quickly go over an equivalent and more categorical definition of equations that will help us argue for our generalization in Chapter 3.¹³⁴

Definition 1.50. An **abstract equation**¹³⁵ in $\mathbf{Alg}(\Sigma)$ is a surjective homomorphism $e : \mathcal{T}_{\Sigma}X \rightarrow \mathbb{Y}$, with the algebra structure on $\mathcal{T}_{\Sigma}X$ given in Remark 1.24. We say that

¹³² Very abstractly: a free object on X is the same thing as an initial object in the comma category $\Delta(X) \downarrow U$, and initial objects are unique up to isomorphism.

¹³³ Formally, let $f, g : \mathbb{T}X \rightarrow \mathbb{A}$ be two homomorphisms such that for any $x \in X$, $f[x]_E = g[x]_E$, then, we can show that $f = g$. For any $t \in \mathcal{T}_{\Sigma}X$, we showed in Lemma 1.41 that $[t]_E = \llbracket t \rrbracket_{\mathbb{T}X}^{\Sigma,E}$. Then using (1.12), we have

$$f[t]_E = \llbracket t \rrbracket_{\mathbb{A}}^{f \circ \eta_X^{\Sigma,E}} = \llbracket t \rrbracket_{\mathbb{A}}^{g \circ \eta_X^{\Sigma,E}} = g[t]_E,$$

where the second inequality follows by hypothesis that f and g agree on equivalence classes of terms of depth 0.

¹³⁴ Similar definitions (when instantiated to Σ -algebras) appear in [MU19, Definition 3.3], [AHS06, 16.16(1)], and [JMU24, Definition 4.2].

¹³⁵ We use the terminology from [JMU24].

an algebra \mathbb{A} **satisfies** e if for any assignment $\iota : X \rightarrow A$, the function $\llbracket - \rrbracket_A^{\iota}$ factors through e in $\mathbf{Alg}(\Sigma)$:

$$\llbracket - \rrbracket_A^{\iota} = \mathcal{T}_{\Sigma} X \xrightarrow{e} \mathbb{Y} \xrightarrow{h} \mathbb{A}.$$

We say that \mathbb{A} satisfies a class of abstract equations if it satisfies all of its elements.

An abstract equation cannot be directly translated to an equation, but it can be translated to a set of equations. Briefly, given $e : \mathcal{T}_{\Sigma} X \rightarrow \mathbb{Y}$, X can be seen as the context, and the factorization $\llbracket - \rrbracket_A^{\iota} = h \circ e$ means that the interpretation of two terms s and t must coincide whenever $e(s) = e(t)$. Therefore, an algebra satisfying e will satisfy all the equations $X \vdash s = t$ with $s, t \in \mathcal{T}_{\Sigma} X$ and $e(s) = e(t)$.

Conversely, any equation ϕ can be translated to an abstract equation by noting that the canonical quotient $[-]_{\{\phi\}} : \mathcal{T}_{\Sigma} X \rightarrow \mathcal{T}_{\Sigma, \{\phi\}} X$ is a surjective homomorphism, and $[s]_{\{\phi\}} = [t]_{\{\phi\}}$ is true precisely when all $(\Sigma, \{\phi\})$ -algebras satisfy $X \vdash s = t$.

We give more details in the two following proofs.

Proposition 1.51. *If E is a class of abstract equations, then there is a class E° of equations such that \mathbb{A} satisfies E if and only if it satisfies E° .*

Proof. Given a class E of abstract equations, we define E° to contain the equation $X \vdash s = t$ for every $e : \mathcal{T}_{\Sigma} X \rightarrow \mathbb{Y}$ in E such that $e(s) = e(t)$. An algebra satisfies E if and only if it satisfies E° .

(\Rightarrow) If \mathbb{A} satisfies E and $X \vdash s = t \in E^{\circ}$ comes from $e : \mathcal{T}_{\Sigma} X \rightarrow \mathbb{Y}$ in E , then for any assignment $\iota : X \rightarrow A$, the factorization $\llbracket - \rrbracket_A^{\iota} = h \circ e$ implies $\llbracket s \rrbracket_A^{\iota} = \llbracket t \rrbracket_A^{\iota}$ because $e(s) = e(t)$. Thus, \mathbb{A} satisfies $X \vdash s = t$. This works for every equation in E° , hence \mathbb{A} satisfies E° .

(\Leftarrow) If \mathbb{A} satisfies E° and $e : \mathcal{T}_{\Sigma} X \rightarrow \mathbb{Y}$ is in E . Then, for any assignment $\iota : X \rightarrow A$, since e is surjective, we can define a function $h : Y \rightarrow A$ by $h(y) = \llbracket t_y \rrbracket_A^{\iota}$ with $t_y \in e^{-1}(y)$. Surjectivity of e means h is defined on all of Y , and the choice of t_y does not matter because for any other $t'_y \in e^{-1}(y)$, we have $e(t_y) = y = e(t'_y)$, so \mathbb{A} satisfies $X \vdash t_y = t'_y$ which in turn means $\llbracket t_y \rrbracket_A^{\iota} = \llbracket t'_y \rrbracket_A^{\iota}$. By definition, we have $\llbracket - \rrbracket_A^{\iota} = h \circ e$, but it remains to check that h is a homomorphism. For any $\text{op} : n \in \Sigma$ and $y_1, \dots, y_n \in Y$, pick $t_i \in e^{-1}(y_i)$ for each i , then we have

$$\begin{aligned} h(\llbracket \text{op} \rrbracket_Y(y_1, \dots, y_n)) &= h(\llbracket \text{op} \rrbracket_Y(e(t_1), \dots, e(t_n))) && \text{definition of } t_i \\ &= h \circ e(\text{op}(t_1, \dots, t_n)) && e \text{ is a homomorphism} \\ &= \llbracket \text{op}(t_1, \dots, t_n) \rrbracket_A^{\iota} && \text{definition of } h \\ &= \llbracket \text{op} \rrbracket_A(\llbracket t_1 \rrbracket_A^{\iota}, \dots, \llbracket t_n \rrbracket_A^{\iota}) && \text{by (1.15)} \\ &= \llbracket \text{op} \rrbracket_A(h \circ e(t_1), \dots, h \circ e(t_n)) && \text{definition of } h \\ &= \llbracket \text{op} \rrbracket_A(h(y_1), \dots, h(y_n)). && \text{definition of } t_i \quad \square \end{aligned}$$

Proposition 1.52. *If E is a class of equations, then there is a class E^{\bullet} of abstract equations such that \mathbb{A} satisfies E if and only if it satisfies E^{\bullet} .*

Proof. Given an equation $\phi = X \vdash s = t$, we let ϕ^{\bullet} be the surjective homomorphism $[-]_{\{\phi\}} : \mathcal{T}_{\Sigma} X \rightarrow \mathcal{T}_{\Sigma, \{\phi\}} X$. An algebra satisfies ϕ if and only if it satisfies ϕ^{\bullet} .

(\Rightarrow) If $\mathbb{A} \models \phi$, then for any assignment $\iota : X \rightarrow A$, we have the following unique factorization because $\mathcal{T}_{\Sigma, \{\phi\}}$ is the free $(\Sigma, \{\phi\})$ -algebra, and $\mathbb{A} \in \mathbf{Alg}(\Sigma, \{\phi\})$:

$$\begin{array}{ccc} X & \xrightarrow{\eta_X^{\Sigma, \{\phi\}}} & \mathcal{T}_{\Sigma, \{\phi\}} X \\ & \searrow \iota & \downarrow h \\ & & A \end{array}$$

We can further decompose with another factorization because $\mathcal{T}_{\Sigma} X$ is the free Σ -algebra and $\mathbb{A} \in \mathbf{Alg}(\Sigma)$.¹³⁶

$$\begin{array}{ccccc} & & \eta_X^{\Sigma, \{\phi\}} & & \\ & \nearrow & & \searrow & \\ X & \xrightarrow{\eta_X^{\Sigma}} & \mathcal{T}_{\Sigma} X & \xrightarrow{\phi^{\bullet}} & \mathcal{T}_{\Sigma, \{\phi\}} X \\ & \searrow & \downarrow h & \nearrow & \downarrow h \\ & & A & & A \end{array}$$

(a) ϕ^{\bullet} (b) $\llbracket - \rrbracket'_A$ (c)

Hence, $\llbracket - \rrbracket'_A$ factors through ϕ^{\bullet} .

(\Leftarrow) If \mathbb{A} satisfies ϕ^{\bullet} , then for any $\iota : X \rightarrow A$, the factorization $\llbracket - \rrbracket'_A = h \circ [-]_{\{\phi\}}$ means that $\llbracket s \rrbracket'_A$ and $\llbracket t \rrbracket'_A$ coincide because $[s]_{\{\phi\}} = [t]_{\{\phi\}}$.¹³⁷ Thus, $\mathbb{A} \models \phi$.

Now, given a class E of equations, it is clear that \mathbb{A} satisfies E if and only if it satisfies $E^{\bullet} = \{\phi^{\bullet} \mid \phi \in E\}$. \square

We conclude that equations and abstract equations are equivalent in terms of expressiveness. We will see in Propositions 3.62 and 3.63 how this generalizes to quantitative equations.

1.4 Equational Logic

We were happy that interpretations in the term algebra are computed syntactically, but there is a big caveat. Everything is done modulo \equiv_E which was defined in (1.24) to basically contain all the equations in $\mathfrak{Th}(E)$, that is, all the equations semantically entailed by E . Thanks to Lemma 1.42, if we want to know whether $X \vdash s = t$ is in $\mathfrak{Th}(E)$, it is enough to check if the free (Σ, E) -algebra $\mathbb{T}X$ satisfies it, but that is a circular argument since the carrier $\mathcal{T}_{\Sigma, E} X$ is defined via \equiv_E .

Equational logic is a deductive system which produces an alternative definition of the free algebra, relying only on syntax. In short, the rules of equational logic allow to syntactically derive all of $\mathfrak{Th}(E)$ starting from E .

In Lemma 1.33, we proved that \equiv_E is a congruence (i.e. reflexive, symmetric, transitive, and invariant under operations), and in Lemma 1.45 we showed \equiv_E is also preserved by substitutions. This can help us syntactically derive $\mathfrak{Th}(E)$ because, for instance, if we know $X \vdash s = t \in E$, we can conclude $X \vdash t = s \in \mathfrak{Th}(E)$ by symmetry. If we know $x, y \vdash f(x) = f(y) \in E$, then we can conclude $X \vdash f(s) = f(t) \in \mathfrak{Th}(E)$ for

¹³⁶ (a) commutes by definition of $\eta_X^{\Sigma, \{\phi\}}$ and the definition $\phi^{\bullet} = [-]_{\{\phi\}}$. (b) commutes by definition of $\llbracket - \rrbracket'_A$. (c) commutes because $\llbracket - \rrbracket'_A$ is the unique homomorphism such that $\llbracket - \rrbracket'_A \circ \eta_X^{\Sigma} = \iota$, but $h \circ \phi^{\bullet}$ is also such a homomorphism by the previous factorization.

¹³⁷ This is because $\phi = X \vdash s = t$ belongs to $\mathfrak{Th}(\{\phi\})$.

$$\begin{array}{c}
\frac{}{X \vdash t = t} \text{REFL} \qquad \frac{X \vdash s = t}{X \vdash t = s} \text{SYMM} \qquad \frac{X \vdash s = t \quad X \vdash t = u}{X \vdash s = u} \text{TRANS} \\
\\
\frac{\text{op} : n \in \Sigma \quad \forall 1 \leq i \leq n, X \vdash s_i = t_i}{X \vdash \text{op}(s_1, \dots, s_n) = \text{op}(t_1, \dots, t_n)} \text{CONG} \\
\\
\frac{\sigma : Y \rightarrow \mathcal{T}_E X \quad Y \vdash s = t}{X \vdash \sigma^*(s) = \sigma^*(t)} \text{SUB}
\end{array}$$

Figure 1.3: Rules of equational logic over the signature Σ , where X and Y can be any set, and s, t, u, s_i and t_i can be any term in $\mathcal{T}_E X$ (or $\mathcal{T}_E Y$ for SUB). As indicated in the premises of the rules CONG and SUB, they can be instantiated for any n -ary operation symbol, and for any function σ respectively.

any terms $s, t \in \mathcal{T}_E X$ by substituting x with s and y with t . This can be summarized with the inference rules of **equational logic** in Figure 1.3.

The first four rules are fairly simple, and they essentially say that equality modulo E is a congruence relation. The SUB rule looks a bit more complicated, it is named after the function σ^* used in the conclusion which we called substitution. Intuitively, it reflects the fact that variables in the context Y are universally quantified. If you know $Y \vdash s = t$ holds, then you can replace each variable $y \in Y$ by $\sigma(y)$ (which may even be a complex term using new variables in X), and you can prove that $X \vdash \sigma^*(s) = \sigma^*(t)$ holds. We did this in Lemma 1.45, and the argument to extract from there is that the interpretation of $\sigma^*(t)$ under some assignment $\iota : X \rightarrow A$ is equal to the interpretation of t under the assignment ι_σ sending $y \in Y$ to the interpretation of $\sigma(y)$ under ι . Since satisfaction of $Y \vdash s = t$ means satisfaction under any assignment (this is where universal quantification comes in), we conclude that $X \vdash \sigma^*(s) = \sigma^*(t)$ must be satisfied.

If you have written sequences of computations to solve a mathematical problem, you are already familiar with the essence of doing proofs in equational logic. The rigorous details of such proofs can be formalized with the following definition.

Definition 1.53 (Derivation). A **derivation**¹³⁸ of $X \vdash s = t$ in equational logic with axioms E (a class of equations) is a finite rooted tree such that:

- all nodes are labelled by equations,
- the root is labelled by $X \vdash s = t$,
- if an internal node (not a leaf) is labelled by ϕ and its children are labelled by ϕ_1, \dots, ϕ_n , then there is a rule in Figure 1.3 which concludes ϕ from ϕ_1, \dots, ϕ_n , and
- all the leaves are either in E or instances of REF, i.e. an equation $Y \vdash u = u$ for some set Y and $u \in \mathcal{T}_E Y$.

Example 1.54. We write a derivation with the same notation used to specify the inference rules in Figure 1.3. Consider the signature $\Sigma = \{+ : 2, e : 0\}$ with E containing the equations defining commutative monoids in (1.19). Here is a derivation of

¹³⁸ Many other definitions of derivations exist, and our treatment of them will not be 100% rigorous.

$x, y, z \vdash x + (y + z) = z + (x + y)$ in equational logic with axioms E .

$$\frac{\frac{x, y, z \vdash x + (y + z) = (x + y) + z \in E}{x, y, z \vdash x + (y + z) = z + (x + y)} \text{TRANS}}{\frac{\sigma = \begin{array}{l} x \mapsto x + y \\ y \mapsto z \end{array} \quad \frac{}{x, y \vdash x + y = y + x} \in E}{x, y, z \vdash (x + y) + z = z + (x + y)} \text{SUB}} \text{TRANS}$$

Given any class of equations E , we denote by $\mathfrak{Th}'(E)$ the class of equations that can be proven from E in equational logic, i.e. $\phi \in \mathfrak{Th}'(E)$ if and only if there is a derivation of ϕ in equational logic with axioms E .

Our goal now is to prove that $\mathfrak{Th}'(E) = \mathfrak{Th}(E)$. We say that equational logic is sound and complete for (Σ, E) -algebras. Less concisely, soundness means that whenever equational logic proves an equation ϕ with axioms E , ϕ is satisfied by all (Σ, E) -algebras, and completeness says that whenever an equation ϕ is satisfied by all (Σ, E) -algebras, there is a derivation of ϕ in equational logic with axioms E .

Soundness is a straightforward consequence of earlier results.¹³⁹

Theorem 1.55 (Soundness). *If $\phi \in \mathfrak{Th}'(E)$, then $\phi \in \mathfrak{Th}(E)$.*

Proof. In the proof of Lemma 1.33, we proved that each of REFL, SYMM, TRANS, and CONG are sound rules for a fixed arbitrary algebra. Namely, if $\mathbb{A} \in \mathbf{Alg}(\Sigma)$ satisfies the equations on top, then it satisfies the one on the bottom. Lemma 1.45 states the same soundness property for SUB. This implies a weaker property: if all (Σ, E) -algebras satisfy the equations on top, then they satisfy the one on the bottom.¹⁴⁰

Now, if $\phi \in \mathfrak{Th}'(E)$ was proven using equational logic and the axioms in E , then since all $\mathbb{A} \in \mathbf{Alg}(\Sigma, E)$ satisfy all the axioms, by repeatedly applying the weaker property above for each rule in the derivation, we find that all $\mathbb{A} \in \mathbf{Alg}(\Sigma, E)$ satisfy ϕ , i.e. $\phi \in \mathfrak{Th}(E)$. \square

Completeness is the harder direction, and there are many ways to prove it.¹⁴¹ We will define an algebra exactly like TX but using the equality relation induced by $\mathfrak{Th}'(E)$ instead of \equiv_E which was induced by $\mathfrak{Th}(E)$. We then show that algebra is a (Σ, E) -algebra, and by construction, it will imply $\mathfrak{Th}(E) \subseteq \mathfrak{Th}'(E)$.

Fix a signature Σ and a class E of equations over Σ . For any set X , we can define a binary relation \equiv'_E on Σ -terms¹⁴² that contains the pair (s, t) whenever $X \vdash s = t$ can be proven in equational logic. Formally, we have for any $s, t \in \mathcal{T}_\Sigma X$ (c.f. (1.24)),

$$s \equiv'_E t \iff X \vdash s = t \in \mathfrak{Th}'(E). \quad (1.40)$$

We can show \equiv'_E is a congruence relation.

Lemma 1.56. *For any set X , the relation \equiv'_E is reflexive, symmetric, transitive, and for any $\text{op} : n \in \Sigma$ and $s_1, \dots, s_n, t_1, \dots, t_n \in \mathcal{T}_\Sigma X$,¹⁴³*

$$(\forall 1 \leq i \leq n, s_i \equiv'_E t_i) \implies \text{op}(s_1, \dots, s_n) \equiv'_E \text{op}(t_1, \dots, t_n). \quad (1.41)$$

Proof. This is immediate from the presence of REFL, SYMM, TRANS, and CONG in the rules of equational logic. \square

¹³⁹ In the story we are telling, the rules of equational logic were designed to be sound because we knew some properties of \equiv_E already. In general, we may use intuitions when defining rules of a logic, and later prove soundness to confirm them, or realize that soundness does not hold and infirm them.

¹⁴⁰ This is a standard theorem of first order logic:

$$(\forall A.(PA \Rightarrow QA)) \Rightarrow (\forall A.PA \Rightarrow \forall A.QA)$$

¹⁴¹ The original proof of Birkhoff [Bir35, Theorem 10] relies on constructing free algebras. Several later proofs (e.g. [Wec92, Theorem 29]) rely on a theory of congruences.

¹⁴² Again, we omit the set X from the notation.

¹⁴³ i.e. \equiv'_E is a congruence on the Σ -algebra $\mathcal{T}_\Sigma X$ defined in Remark 1.24.

We write $\lambda_{-}\int_E : \mathcal{T}_E X \rightarrow \mathcal{T}_E X / \equiv'_E$ for the canonical quotient map, so $\lambda t \int_E$ is the equivalence class of t modulo the congruence \equiv'_E induced by equational logic.

Definition 1.57 (Term algebra, syntactically). The *new* term algebra for (Σ, E) on X is the Σ -algebra whose carrier is $\mathcal{T}_E X / \equiv'_E$ and whose interpretation of $\text{op} : n \in \Sigma$ is defined by¹⁴⁴

$$\llbracket \text{op} \rrbracket_{\mathbb{T}'X}(\lambda t_1 \int_E, \dots, \lambda t_n \int_E) = \lambda \text{op}(t_1, \dots, t_n) \int_E. \quad (1.42)$$

We denote this algebra by $\mathbb{T}'_{\Sigma, E} X$ or simply $\mathbb{T}'X$.

With soundness (Theorem 1.55) of equational logic, completeness would mean this alternative definition of the term algebra coincides with $\mathbb{T}X$. First, we have to show that $\mathbb{T}'X$ belongs to $\mathbf{Alg}(\Sigma, E)$ like we did for $\mathbb{T}X$ in Proposition 1.46, and we prove a technical lemma before that.

Lemma 1.58. *Let $\iota : Y \rightarrow \mathcal{T}_E X / \equiv'_E$ be an assignment. For any function $\sigma : Y \rightarrow \mathcal{T}_E X$ satisfying $\lambda \sigma(y) \int_E = \iota(y)$ for all $y \in Y$, we have $\llbracket - \rrbracket_{\mathbb{T}'X}^{\iota} = \lambda \sigma^*(-) \int_E$.¹⁴⁵*

Proof. We proceed by induction. For the base case, we have by definition of the interpretation of terms (1.9), definition of σ , and definition of σ^* (1.38),

$$\llbracket \eta_Y^{\Sigma}(y) \rrbracket_{\mathbb{T}'X}^{\iota} \stackrel{(1.9)}{=} \iota(y) = \lambda \sigma(y) \int_E \stackrel{(1.38)}{=} \lambda \sigma^*(\eta_Y^{\Sigma}(y)) \int_E.$$

For the inductive step, we have

$$\begin{aligned} \llbracket \text{op}(t_1, \dots, t_n) \rrbracket_{\mathbb{T}'X}^{\iota} &= \llbracket \text{op} \rrbracket_{\mathbb{T}'X}(\llbracket t_1 \rrbracket_{\mathbb{T}'X}^{\iota}, \dots, \llbracket t_n \rrbracket_{\mathbb{T}'X}^{\iota}) && \text{by (1.9)} \\ &= \llbracket \text{op} \rrbracket_{\mathbb{T}'X}(\lambda \sigma^*(t_1) \int_E, \dots, \lambda \sigma^*(t_n) \int_E) && \text{I.H.} \\ &= \lambda \text{op}(\sigma^*(t_1), \dots, \sigma^*(t_n)) \int_E && \text{by (1.42)} \\ &= \lambda \sigma^*(\text{op}(t_1, \dots, t_n)) \int_E. && \text{definition of } \sigma^* \quad \square \end{aligned}$$

Proposition 1.59. *For any set X , $\mathbb{T}'X$ satisfies all the equations in E .*

Proof. Let $Y \vdash s = t$ belong to E and $\iota : Y \rightarrow \mathcal{T}_E X / \equiv'_E$ be an assignment. By the axiom of choice,¹⁴⁶ there is a function $\sigma : Y \rightarrow \mathcal{T}_E X$ satisfying $\lambda \sigma(y) \int_E = \iota(y)$ for all $y \in Y$. Thanks to Lemma 1.58, it is enough to show $\lambda \sigma^*(s) \int_E = \lambda \sigma^*(t) \int_E$.¹⁴⁷ Equivalently, by definition of $\lambda_{-}\int_E$ and $\mathfrak{Th}'(E)$, we can just exhibit a derivation of $X \vdash \sigma^*(s) = \sigma^*(t)$ in equational logic with axioms E . This is rather simple because that equation can be proven with the SUB rule instantiated with $\sigma : Y \rightarrow \mathcal{T}_E X$ and the equation $Y \vdash s = t$ which is an axiom. \square

Completeness of equational logic readily follows.

Theorem 1.60 (Completeness). *If $\phi \in \mathfrak{Th}(E)$, then $\phi \in \mathfrak{Th}'(E)$.*

Proof. Write $\phi = X \vdash s = t \in \mathfrak{Th}(E)$. By Proposition 1.59 and definition of $\mathfrak{Th}(E)$, we know that $\mathbb{T}'X \models \phi$. In particular, $\mathbb{T}'X$ satisfies ϕ under the assignment

$$\iota = X \xrightarrow{\eta_X^{\Sigma}} \mathcal{T}_E X \xrightarrow{\lambda_{-}\int_E} \mathcal{T}_E X / \equiv'_E,$$

¹⁴⁴ This is well-defined (i.e. invariant under change of representative) by (1.41).

¹⁴⁵ This result looks like a stronger version of Lemma 1.41 for $\mathbb{T}'X$. Morally, they are both saying that interpretation of terms in $\mathbb{T}X$ or $\mathbb{T}'X$ is just a syntactical matter.

¹⁴⁶ Choice implies the quotient map $\lambda_{-}\int_E$ has a right inverse $r : \mathcal{T}_E X / \equiv'_E \rightarrow \mathcal{T}_E X$, and we can then set $\sigma = r \circ \iota$.

¹⁴⁷ By Lemma 1.58, it implies

$$\llbracket s \rrbracket_{\mathbb{T}'X}^{\iota} = \lambda \sigma^*(s) \int_E = \lambda \sigma^*(t) \int_E = \llbracket t \rrbracket_{\mathbb{T}'X}^{\iota},$$

and since ι was an arbitrary assignment, we conclude that $\mathbb{T}'X \models Y \vdash s = t$.

namely, $\llbracket s \rrbracket_{\mathbb{T}'X} = \llbracket t \rrbracket_{\mathbb{T}'X}$. Moreover with $\sigma = \eta_X^\Sigma$, we can show σ satisfies the hypothesis of Lemma 1.58 and $\sigma^* = \text{id}_{\mathcal{T}_E X}$,¹⁴⁸ thus we conclude

$$\wr s \wr_E = \llbracket s \rrbracket_{\mathbb{T}'X} = \llbracket t \rrbracket_{\mathbb{T}'X} = \wr t \wr_E.$$

This implies $s \equiv'_E t$ which in turn means $X \vdash s = t$ belongs to $\mathfrak{Th}'(E)$. \square

Note that because $\mathbb{T}X$ and $\mathbb{T}'X$ were defined in the same way in terms of $\mathfrak{Th}(E)$ and $\mathfrak{Th}'(E)$ respectively, and since we have proven the latter to be equal, we obtain that $\mathbb{T}X$ and $\mathbb{T}'X$ are the same algebra.¹⁴⁹

Remark 1.61. We have used the axiom of choice in proving completeness of equational logic, but that is only an artifact of our presentation that deals with arbitrary contexts. Since terms are finite and operation symbols have finite arities, we can make do with only finite contexts (which removes the need for choice). Formally, one can prove by induction on the derivation that a proof of $X \vdash s = t$ can be transformed into a proof of $\text{FV}\{s, t\} \vdash s = t$ which uses only equations with finite contexts.¹⁵⁰ You can also verify semantically that \mathbb{A} satisfies $X \vdash s = t$ if and only if it satisfies $\text{FV}\{s, t\} \vdash s = t$ essentially because the extra variables have no effect on the quantification of the free variables in s and t nor on the interpretation.

We mention now two related results for the sake of comparison when we introduce quantitative equational logic. First, for any set X and variable y , the following inference rules are derivable in equational logic.

$$\frac{X \vdash s = t}{X \cup \{y\} \vdash s = t} \text{ ADD} \qquad \frac{X \vdash s = t \quad y \notin \text{FV}\{s, t\}}{X \setminus \{y\} \vdash s = t} \text{ DEL}$$

In words, ADD says that you can always add a variable to the context, and DEL says you can remove a variable from the context when it is not used in the terms of the equations. Both these rules are instances of SUB. For the first, take σ to be the inclusion of X in $X \cup \{y\}$ (it may be the identity if $y \in X$). For the second, let σ send y to whatever element of $X \setminus \{y\}$ and all the other elements of X to themselves¹⁵¹, then since y is not in the free variables of s and t , $\sigma^*(s) = s$ and $\sigma^*(t) = t$.

Second, we allowed the collection of equations E generating an algebraic theory $\mathfrak{Th}(E)$ to be a proper class, and that is really not common. Oftentimes, a countable set of variables $\{x_1, x_2, \dots\}$ is assumed, and equations are defined only with a context contained in that set. With this assumption, the collection of all equations is a set, and so are E and $\mathfrak{Th}(E)$. This has no effect on expressiveness since for any equation $X \vdash s = t$, there is an equivalent equation $X' \vdash s' = t'$ with $X' \subseteq \{x_1, x_2, \dots\}$.¹⁵²

1.5 Monads

Our presentation of universal algebra used the language of category theory, e.g. functors, naturality, commutative diagrams, etc. Both these fields of mathematics were born within a decade of each other¹⁵³ with a similar goal: abstracting the way mathematicians use mathematical objects in order to apply one general argument

¹⁴⁸ We defined \wr precisely to have $\wr \sigma(x) \wr_E = \wr x \wr$. To show $\sigma^* = \eta_X^\Sigma$ is the identity, use (1.38) and the fact that $\mu^\Sigma \cdot \mathcal{T}_E \eta^\Sigma = \mathbb{1}_{\mathcal{T}_E}$ (Lemma 1.14).

¹⁴⁹ It is good to keep in mind these two equivalent definitions of the free (Σ, E) -algebra on X . It means you can prove s equals t in $\mathbb{T}X$ by exhibiting a derivation of $X \vdash s = t$ in equational logic, or you can prove $s \neq t$ by exhibiting an algebra that satisfies E but not $X \vdash s = t$.

¹⁵⁰ We denoted by $\text{FV}\{s, t\}$ the set of **free variables** used in s and t . This can be defined inductively as follows:

$$\begin{aligned} \text{FV}\{\eta_X^\Sigma(x)\} &= \{x\} \\ \text{FV}\{\text{op}(t_1, \dots, t_n)\} &= \text{FV}\{t_1\} \cup \dots \cup \text{FV}\{t_n\} \\ \text{FV}\{t_1, \dots, t_n\} &= \text{FV}\{t_1\} \cup \dots \cup \text{FV}\{t_n\}. \end{aligned}$$

Note that $\text{FV}\{-}$ applied to a finite set of terms is always finite.

¹⁵¹ When X is empty, the equations on the top and bottom of DEL coincide, so the rule is derivable.

¹⁵² We already know $X \vdash s = t$ is equivalent to $\text{FV}\{s, t\} \vdash s = t$, and since the context of the latter is finite, we have a bijection $\sigma : \text{FV}\{s, t\} \cong \{x_1, \dots, x_n\}$. Then the SUB rule instantiated with σ and σ^{-1} proves the desired equivalence.

¹⁵³ [Bir33, Bir35] and [EM45] were the seminal papers for universal algebra and category theory respectively. Birkhoff and MacLane even wrote an undergraduate textbook together [MB99].

to many specific cases.¹⁵⁴ One could argue (looking at today’s practicing mathematicians) that category theory was more successful. This is why a portion of this manuscript is spent on monads, a more categorical formulation of the content in universal algebra which became popular in computer science after Moggi’s work [Mog89, Mog91] using monads to abstract computational effects.

There is another categorical approach to universal algebra introduced by Lawvere [Law63] and first popularized in the computer science community by Hyland, Plotkin, and Power [PPo1a, PPo1b, HPPo6, HPo7]. We will stick to monads because most of the literature on quantitative algebras does, and because I am not sure yet how the generalizations we contributed port to Lawvere’s approach.¹⁵⁵

Definition 1.62 (Monad). A **monad** on a category \mathbf{C} is a triple (M, η, μ) made up of an endofunctor $M : \mathbf{C} \rightarrow \mathbf{C}$ and two natural transformations $\eta : \text{id}_{\mathbf{C}} \Rightarrow M$ and $\mu : M^2 \Rightarrow M$, called the **unit** and **multiplication** respectively, that make (1.43) and (1.44) commute in $[\mathbf{C}, \mathbf{C}]$ (the category of endofunctors on \mathbf{C}).¹⁵⁶

$$\begin{array}{ccc} M & \xrightarrow{M\eta} & M^2 & \xleftarrow{\eta M} & M \\ & \searrow & \downarrow \mu & \swarrow & \\ & \mathbb{1}_M & M & & \mathbb{1}_M \end{array} \quad (1.43) \qquad \begin{array}{ccc} M^3 & \xrightarrow{\mu M} & M^2 \\ M\mu \downarrow & & \downarrow \mu \\ M^2 & \xrightarrow{\mu} & M \end{array} \quad (1.44)$$

We often refer to the monad (M, η, μ) simply with M .

In this chapter we will mostly talk about monads on **Set**, but it is good to keep some arguments general for later. Here are some very important examples (for computer scientists and especially for this manuscript).

Example 1.63 (Maybe). Suppose \mathbf{C} has (binary) coproducts and a terminal object $\mathbf{1}$, then $(- + \mathbf{1}) : \mathbf{C} \rightarrow \mathbf{C}$ is a monad. It is called the **maybe monad** (the name “option monad” is also common).¹⁵⁷ We write inl^{X+Y} (resp. inr^{X+Y}) for the coprojection of X (resp. Y) into $X + Y$.¹⁵⁸ First, note that for a morphism $f : X \rightarrow Y$,

$$f + \mathbf{1} = [\text{inl}^{Y+\mathbf{1}} \circ f, \text{inr}^{Y+\mathbf{1}}] : X + \mathbf{1} \rightarrow Y + \mathbf{1}.$$

The components of the unit are given by the coprojections, i.e. $\eta_X = \text{inl}^{X+\mathbf{1}} : X \rightarrow X + \mathbf{1}$, and the components of the multiplication are

$$\mu_X = [\text{inl}^{X+\mathbf{1}}, \text{inr}^{X+\mathbf{1}}, \text{inr}^{X+\mathbf{1}}] : X + \mathbf{1} + \mathbf{1} \rightarrow X + \mathbf{1}.$$

Checking that (1.43) and (1.44) commute is an exercise in reasoning with coproducts. It is more interesting to give the intuition in **Set** where $+$ is the disjoint union and $\mathbf{1}$ is the singleton $\{*\}$.¹⁵⁹

- $X + \mathbf{1}$ is the set X with an additional (fresh) element $*$,
- the function $f + \mathbf{1}$ acts like f on X and sends the new element $* \in X$ to the new element $* \in Y$,
- the unit $\eta_X : X \rightarrow X + \mathbf{1}$ is the injection (sending $x \in X$ to itself),

¹⁵⁴ This is very close to a goal of mathematics as a whole: abstracting the way nature works in order to apply one general argument to many specific cases, c.f. Cheng calling category theory the “mathematics of mathematics” [Che16].

¹⁵⁵ In the paper introducing quantitative algebra [MPP16], the authors already mentioned enriched Lawvere theories [Pow99]. The works of Hyland and Power [HPo6], Nishizawa and Power [NPo9], Lucyshyn-Wright and Parker [LW16, LP23], and Rosický [Ros24] are also relevant.

¹⁵⁶ I also recommend Marsden’s series of blog posts on monads for a relatively light and comprehensive survey: <https://stringdiagram.com/2022/05/17/hello-monads/>.

¹⁵⁷ It is also called the lift monad in [Jac16, Example 5.1.3.2].

¹⁵⁸ These notations are common in the community of programming language research, they stand for *injection left* (resp. *right*). We may omit the superscript.

¹⁵⁹ This intuition should carry over well to many categories where the coproduct and terminal objects have similar behaviors.

- the multiplication μ_X acts like the identity on X and sends the two new elements of $X + \mathbf{1} + \mathbf{1}$ to the single new element $X + \mathbf{1}$,
- one can check (1.43) and (1.44) commute by hand because (briefly) $x \in X$ is always sent to $x \in X$ and $*$ is always sent to $*$.

The fresh element $*$ is often seen as a terminating state, so the maybe monad models the most basic computational effect: termination. Even when no other observation can be made on states of a program, one can distinguish between states by looking at their execution traces which may or may not contain $*$.¹⁶⁰

Example 1.64 (Powerset). The covariant **non-empty finite powerset** functor $\mathcal{P}_{\text{ne}} : \mathbf{Set} \rightarrow \mathbf{Set}$ sends a set X to the set of non-empty finite subsets of X which we denote by $\mathcal{P}_{\text{ne}}X$. It acts on functions just like the usual powerset functor, i.e. given a function $f : X \rightarrow Y$, $\mathcal{P}_{\text{ne}}f$ is the direct image function, it sends $S \subseteq X$ to $f(S) = \{f(x) \mid x \in S\}$. It is clear that $f(S)$ is non-empty and finite when S is non-empty and finite.

One can show \mathcal{P}_{ne} is a monad with the following unit and multiplication:¹⁶¹

$$\eta_X : X \rightarrow \mathcal{P}_{\text{ne}}(X) = x \mapsto \{x\} \text{ and } \mu_X : \mathcal{P}_{\text{ne}}(\mathcal{P}_{\text{ne}}(X)) \rightarrow \mathcal{P}_{\text{ne}}(X) = F \mapsto \bigcup_{s \in F} s.$$

Again as early as in Moggi's papers, the powerset monad was used to model nondeterministic computations (see also [VW06, KS18, BSV19, GPA21]). A set $S \in \mathcal{P}_{\text{ne}}X$ is seen as all the possible states at a point in the execution. We assume that S is finite for convenience, and that it is non-empty because an empty set of possible states would mean termination which can already be modelled with the maybe monad.¹⁶²

Example 1.65 (Distributions). The functor $\mathcal{D} : \mathbf{Set} \rightarrow \mathbf{Set}$ sends a set X to the set of **finitely supported distributions** on X :¹⁶³

$$\mathcal{D}(X) := \{\varphi : X \rightarrow [0, 1] \mid \sum_{x \in X} \varphi(x) = 1 \text{ and } \varphi(x) \neq 0 \text{ for finitely many } x\text{'s}\}.$$

We call $\varphi(x)$ the **weight** of φ at x and let $\text{supp}(\varphi)$ denote the **support** of φ , that is, $\text{supp}(\varphi)$ contains all the elements $x \in X$ such that $\varphi(x) \neq 0$.¹⁶⁴ On morphisms, \mathcal{D} sends a function $f : X \rightarrow Y$ to the function between sets of distributions defined by

$$\mathcal{D}f : \mathcal{D}X \rightarrow \mathcal{D}Y = \varphi \mapsto \left(y \mapsto \sum_{x \in X, f(x)=y} \varphi(x) \right). \quad (1.45)$$

In words, the weight of $\mathcal{D}f(\varphi)$ at y is equal to the total weight of φ on the preimage of y under f .¹⁶⁵

One can show that \mathcal{D} is a monad with unit $\eta_X = x \mapsto \delta_x$, where δ_x is the **Dirac** distribution at x (the weight of δ_x is 1 at x and 0 everywhere else), and multiplication

$$\mu_X = \Phi \mapsto \left(x \mapsto \sum_{\varphi \in \text{supp}(\Phi)} \Phi(\varphi)\varphi(x) \right). \quad (1.46)$$

¹⁶⁰ This was already known to Moggi who used different terminology in [Mog91, Example 1.1].

¹⁶¹ Note that $\{x\}$ is non-empty and finite, and so is $\bigcup_{s \in F} s$ whenever F and all $s \in F$ are non-empty and finite. Thus, we can define \mathcal{P}_{ne} as a submonad of the full powerset monad in, e.g. [Jac16, Example 5.1.3.1].

¹⁶² Also, the maybe monad can be *combined* with any other monad, see e.g. [MSV21, Corollary 5].

¹⁶³ We will simply call them distributions.

¹⁶⁴ We often write $\varphi(S)$ for the total weight of φ on all of $S \subseteq X$.

¹⁶⁵ The distribution $\mathcal{D}f(\varphi)$ is sometimes called the **pushforward** of φ .

In words, the weight $\mu_X(\Phi)$ at x is the average of $\varphi(x)$ weighted by $\Phi(\varphi)$ for all distributions in the support of Φ .¹⁶⁶

Moggi only hinted at the distribution monad being a good model for computations that rely on random/probabilistic choices. For fleshed out research based on \mathcal{D} and variants, see, e.g. [Gra88, JP89, RP02, VW06, Sta17, SW18, BSV19].

Monads have been a popular categorical approach to universal algebra¹⁶⁷ thanks to a result of Linton [Lin66, Proposition 1] stating that any algebraic theory gives rise to a monad. Given a signature Σ and a class E of equations, we already implicitly described the monad Linton constructed, it is the triple $(\mathcal{T}_{\Sigma,E}, \eta^{\Sigma,E}, \mu^{\Sigma,E})$.

Proposition 1.66. *The functor $\mathcal{T}_{\Sigma,E} : \mathbf{Set} \rightarrow \mathbf{Set}$ defines a monad on \mathbf{Set} with unit $\eta^{\Sigma,E}$ and multiplication $\mu^{\Sigma,E}$. We call it the **term monad** for (Σ, E) .*

Proof. We have done most of the work already.¹⁶⁸ We showed that $\eta^{\Sigma,E}$ and $\mu^{\Sigma,E}$ are natural transformations of the right type in Footnote 124 and Proposition 1.38 respectively, and we showed the appropriate instance of (1.43) commutes in Lemma 1.43. It remains to prove (1.44) commutes which, instantiated here, means proving the following diagram commutes for every set A .

$$\begin{array}{ccc} \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} A & \xrightarrow{\mathcal{T}_{\Sigma,E} \mu_A^{\Sigma,E}} & \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} A \\ \mu_{\mathcal{T}_{\Sigma,E} A}^{\Sigma,E} \downarrow & & \downarrow \mu_A^{\Sigma,E} \\ \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} A & \xrightarrow{\mu_A^{\Sigma,E}} & \mathcal{T}_{\Sigma,E} A \end{array}$$

It follows from the following paved diagram.¹⁶⁹

$$\begin{array}{ccccc} \mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} A & \xrightarrow{\mathcal{T}_{\Sigma} \mu_A^{\Sigma,E}} & \mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma,E} A & & \\ \downarrow \llbracket - \rrbracket_{\text{TTA}} & \searrow & \swarrow & \downarrow \llbracket - \rrbracket_{\text{TA}} & \\ & \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} A & \xrightarrow{\mu_{\mathcal{T}_{\Sigma,E} A}^{\Sigma,E}} & \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} A & \\ & \swarrow & \searrow & \swarrow & \\ \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} A & \xrightarrow{\mu_A^{\Sigma,E}} & \mathcal{T}_{\Sigma,E} A & & \end{array}$$

(a) $\llbracket - \rrbracket_E$ (b) $\llbracket - \rrbracket_E$ (c) $\llbracket - \rrbracket_{\text{TA}}$ (d)

Note that when E is empty, we get a monad $(\mathcal{T}_{\Sigma}, \eta^{\Sigma}, \mu^{\Sigma})$.¹⁷⁰ □

Linton also showed that from a monad M , you can build a theory whose corresponding term monad is isomorphic to M [Lin69, Lemma 10.1]. This however relied on a more general notion of theory. We will not go over the details here, rather we will introduce the necessary concepts to talk about our main examples on \mathbf{Set} : $(- + 1)$, \mathcal{R}_{re} , and \mathcal{D} . First, we introduce algebras for a monad.

Definition 1.67 (M -algebra). Let (M, η, μ) be a monad on \mathbf{C} , an M -**algebra** is a pair (A, α) comprising an object $A \in \mathbf{C}_0$ and a morphism $\alpha : MA \rightarrow A$ such that (1.47) and (1.48) commute.

¹⁶⁶ It was Giry [Gir82] who first studied probabilities through the categorical lens with a monad with inspiration from Lawvere [Law62], \mathcal{D} is a discrete version of Giry's original construction. (See [Jac16, Example 5.1.3.4].)

¹⁶⁷ See [HP07] for a thorough survey on categorical approaches to universal algebra.

¹⁶⁸ In fact, we have done it twice because we showed that $\mathbb{T}_{\Sigma,E} A$ is the free (Σ, E) -algebra on A for every set A , and that automatically yields (through abstract categorical arguments) a monad sending A to the carrier of $\mathbb{T}_{\Sigma,E} A$, i.e. $\mathcal{T}_{\Sigma,E} A$.

¹⁶⁹ We know that (a), (b) and (c) commute by (1.30), (1.26), and (1.30) respectively. This means that (d) pre-composed by the epimorphism $\llbracket - \rrbracket_E$ yields the outer square. Moreover, we know the outer square commutes by (1.35), therefore, (d) must also commute.

¹⁷⁰ Here is an alternative proof that \mathcal{T}_{Σ} is a monad. We showed η^{Σ} and μ^{Σ} are natural in (1.6) and (1.8) respectively. The right triangle of (1.43) commutes by definition of μ^{Σ} (1.7), the left triangle commutes by Lemma 1.14, and the square (1.44) commutes by (1.16).

$$\begin{array}{ccc}
 A & \xrightarrow{\eta_A} & MA \\
 & \searrow \text{id}_A & \downarrow \alpha \\
 & & A
 \end{array} \quad (1.47) \qquad
 \begin{array}{ccc}
 MMA & \xrightarrow{\mu_A} & MA \\
 M\alpha \downarrow & & \downarrow \alpha \\
 MA & \xrightarrow{\alpha} & A
 \end{array} \quad (1.48)$$

We call A the carrier and we may write only α to refer to an M -algebra.

Definition 1.68 (Homomorphism). Let (M, η, μ) be a monad and (A, α) and (B, β) be two M -algebras. An M -algebra **homomorphism** or simply M -homomorphism from α to β is a morphism $h : A \rightarrow B$ in \mathbf{C} making (1.49) commute.

$$\begin{array}{ccc}
 MA & \xrightarrow{Mh} & MB \\
 \alpha \downarrow & & \downarrow \beta \\
 A & \xrightarrow{h} & B
 \end{array} \quad (1.49)$$

The composition of two M -homomorphisms is an M -homomorphism and id_A is an M -homomorphism from (A, α) to itself, thus we get a category of M -algebras and M -homomorphisms called the **Eilenberg–Moore category** of M , and denoted by $\mathbf{EM}(M)$.¹⁷¹ Since $\mathbf{EM}(M)$ was built from objects and morphisms in \mathbf{C} , there is an obvious forgetful functor $U^M : \mathbf{EM}(M) \rightarrow \mathbf{C}$ sending an M -algebra (A, α) to its carrier A , and an M -homomorphism to its underlying morphism.

Example 1.69. We will see some more concrete examples in a bit, but we can mention now that the similarities between the squares in the definitions of a monad (1.44), of an algebra (1.48), and of a homomorphism (1.49) have profound consequences. First, for any A , the pair (MA, μ_A) is an M -algebra because (1.50) and (1.51) commute by the properties of a monad.¹⁷²

$$\begin{array}{ccc}
 MA & \xrightarrow{\eta_{MA}} & MMA \\
 \text{id}_{MA} \searrow & & \downarrow \mu_A \\
 & & MA
 \end{array} \quad (1.50) \qquad
 \begin{array}{ccc}
 MMA & \xrightarrow{\mu_{MA}} & MMA \\
 M\mu_A \downarrow & & \downarrow \mu_A \\
 MMA & \xrightarrow{\mu_A} & MA
 \end{array} \quad (1.51)$$

Furthermore, for any M -algebra $\alpha : MA \rightarrow A$, (1.48) (reflected through the diagonal) precisely says that α is a M -homomorphism from (MA, μ_A) to (A, α) . After a bit more work,¹⁷³ we can conclude that (MA, μ_A) is the free M -algebra (relative to $U^M : \mathbf{EM}(M) \rightarrow \mathbf{Set}$).

The terminology suggests that (Σ, E) -algebras and $\mathcal{T}_{\Sigma, E}$ -algebras are the same thing.¹⁷⁴ Let us check this, obtaining a large family of examples at the same time.

Proposition 1.70. *There is an isomorphism $\mathbf{Alg}(\Sigma, E) \cong \mathbf{EM}(\mathcal{T}_{\Sigma, E})$.*

Proof. Given a (Σ, E) -algebra \mathbb{A} , we already explained in (1.34) how to obtain a function $\llbracket - \rrbracket_{\mathbb{A}} : \mathcal{T}_{\Sigma, E} \mathbb{A} \rightarrow \mathbb{A}$ which sends $[t]_E$ to the interpretation of the term t under the trivial assignment $\text{id}_{\mathbb{A}} : \mathbb{A} \rightarrow \mathbb{A}$.¹⁷⁵ Let us verify that $\llbracket - \rrbracket_{\mathbb{A}}$ is a $\mathcal{T}_{\Sigma, E}$ -algebra. We need to show the following instances of (1.47) and (1.48) commutes.

$$\begin{array}{ccc}
 A & \xrightarrow{\eta_A^{\Sigma, E}} & \mathcal{T}_{\Sigma, E} A \\
 & \searrow \text{id}_A & \downarrow \llbracket - \rrbracket_A \\
 & & A
 \end{array} \qquad
 \begin{array}{ccc}
 \mathcal{T}_{\Sigma, E} \mathcal{T}_{\Sigma, E} A & \xrightarrow{\mu_A^{\Sigma, E}} & \mathcal{T}_{\Sigma, E} A \\
 \mathcal{T}_{\Sigma, E} \llbracket - \rrbracket_A \downarrow & & \downarrow \llbracket - \rrbracket_A \\
 \mathcal{T}_{\Sigma, E} A & \xrightarrow{\llbracket - \rrbracket_A} & A
 \end{array}$$

¹⁷¹ Named after the authors of the article introducing that category [EM65].

¹⁷² (1.50) is the component at A of the right triangle in (1.43), and (1.51) is the component at A of (1.44).

¹⁷³ Given an M -algebra (A', α') and a function $f : A \rightarrow A'$, we can show $\alpha' \circ Mf$ is the unique M -homomorphism such that $\alpha' \circ Mf \circ \eta_A = f$.

¹⁷⁴ Also, Example 1.69 starts to confirm this, if we compare it with Remark 1.24, and Lemma 1.25.

¹⁷⁵ That is well-defined because \mathbb{A} satisfies all the equations in $\mathfrak{Th}(E)$.

The triangle commutes by definitions,¹⁷⁶ and the square commutes by the following diagram.

$$\begin{array}{ccccc}
\mathcal{T}_\Sigma \mathcal{T}_{\Sigma,E} A & \xrightarrow{\mathcal{T}_\Sigma [-]_A} & \mathcal{T}_\Sigma A & & \\
\downarrow [-]_{\mathcal{T}A} & \searrow [-]_E & \swarrow [-]_E & & \downarrow [-]_A \\
& \mathcal{T}_{\Sigma,E} \mathcal{T}_{\Sigma,E} A & \xrightarrow{\mathcal{T}_{\Sigma,E} [-]_A} & \mathcal{T}_{\Sigma,E} A & \\
& \swarrow \mu_A^{\Sigma,E} & \searrow [-]_A & & \\
\mathcal{T}_{\Sigma,E} A & \xrightarrow{[-]_A} & A & &
\end{array}
\begin{array}{l}
\text{(a)} \\
\text{(b)} \\
\text{(c)} \\
\text{(d)}
\end{array}$$

Since the outer rectangle commutes by Lemma 1.39, (a) commutes by naturality of $[-]_E$ (1.26), (b) commutes by definition of $\mu_A^{\Sigma,E}$ (1.30), and (d) commutes by (1.34), we can conclude that (c) commutes because $[-]_E$ is epic.

We also already explained in Footnote 75 that any homomorphism $h : \mathbb{A} \rightarrow \mathbb{B}$ makes the outer rectangle below commute.

$$\begin{array}{ccccc}
\mathcal{T}_\Sigma A & \xrightarrow{\mathcal{T}_\Sigma h} & \mathcal{T}_\Sigma B & & \\
\downarrow [-]_A & \searrow [-]_E & \swarrow [-]_E & & \downarrow [-]_B \\
& \mathcal{T}_{\Sigma,E} A & \xrightarrow{\mathcal{T}_{\Sigma,E} h} & \mathcal{T}_{\Sigma,E} A & \\
& \swarrow [-]_A & \searrow [-]_A & & \\
A & \xrightarrow{h} & B & &
\end{array}
\begin{array}{l}
\text{(a)} \\
\text{(b)} \\
\text{(c)} \\
\text{(d)}
\end{array}$$

Since (a), (b), and (d) commute by naturality of $[-]_E$, (1.34), and (1.34) respectively, we conclude that (c) commutes again because $[-]_E$ is epic. This means h is a $\mathcal{T}_{\Sigma,E}$ -homomorphism.

We obtain a functor¹⁷⁷ $P : \mathbf{Alg}(\Sigma, E) \rightarrow \mathbf{EM}(\mathcal{T}_{\Sigma,E})$ sending $\mathbb{A} = (A, [-]_A)$ to $(A, \alpha_{\mathbb{A}})$ where $\alpha_{\mathbb{A}} = [-]_A : \mathcal{T}_{\Sigma,E} A \rightarrow A$ (we give it a different name to make the sequel easier to follow).

In the other direction, given an algebra $\alpha : \mathcal{T}_{\Sigma,E} A \rightarrow A$, we define an algebra \mathbb{A}_α with the interpretation of $\text{op} : n \in \Sigma$ given by¹⁷⁸

$$[[\text{op}]]_\alpha(a_1, \dots, a_n) = \alpha[\text{op}(a_1, \dots, a_n)]_E, \quad (1.52)$$

and we can prove by induction that $[[t]]_\alpha = \alpha[t]_E$ for any Σ -term t over A (note that we use the $\mathcal{T}_{\Sigma,E}$ -algebra properties of α).¹⁷⁹ Now, if $h : (A, \alpha) \rightarrow (B, \beta)$ is a $\mathcal{T}_{\Sigma,E}$ -homomorphism, then h is a homomorphism from \mathbb{A}_α to \mathbb{B}_β because for any $\text{op} : n \in \Sigma$ and $a_1, \dots, a_n \in A$, we have

$$\begin{aligned}
h([[\text{op}]]_\alpha(a_1, \dots, a_n)) &= h(\alpha[\text{op}(a_1, \dots, a_n)]_E) && \text{by (1.52)} \\
&= \beta(\mathcal{T}_{\Sigma,E} h[\text{op}(a_1, \dots, a_n)]_E) && \text{by (1.49)} \\
&= \beta[\mathcal{T}_\Sigma h(\text{op}(a_1, \dots, a_n))]_E && \text{by (1.26)} \\
&= \beta[\text{op}(h(a_1), \dots, h(a_n))]_E && \text{by (1.5)} \\
&= [[\text{op}]]_\beta(h(a_1), \dots, h(a_n)). && \text{by (1.52)}
\end{aligned}$$

¹⁷⁶ We have $[[\eta_A^{\Sigma,E}(a)]]_A = [[a]_E]_A = [[a]]_A = a$.

¹⁷⁷ Checking functoriality is trivial because P acts like the identity on morphisms.

¹⁷⁸ For readability, we write $\alpha[-]$ instead of $\alpha([-])$.

¹⁷⁹ For the base case, we have

$$[[a]]_\alpha \stackrel{(1.9)}{=} a \stackrel{(1.47)}{=} \alpha[\eta_A^\Sigma(a)]_E = \alpha[a]_E.$$

For the inductive step, let $t = \text{op}(t_1, \dots, t_n) \in \mathcal{T}_\Sigma A$:

$$\begin{aligned}
[[t]]_\alpha &= [[\text{op}(t_1, \dots, t_n)]]_\alpha \\
&= [[\text{op}]]_\alpha([[t_1]]_\alpha, \dots, [[t_n]]_\alpha) && (1.9) \\
&= [[\text{op}]]_\alpha(\alpha[t_1]_E, \dots, \alpha[t_n]_E) && \text{I.H.} \\
&= \alpha[\text{op}(\alpha[t_1]_E, \dots, \alpha[t_n]_E)]_E && (1.52) \\
&= \alpha[\mathcal{T}_\Sigma \alpha(\text{op}([t_1]_E, \dots, [t_n]_E))]_E && (1.5) \\
&= \alpha(\mathcal{T}_{\Sigma,E} \alpha[\text{op}([t_1]_E, \dots, [t_n]_E)])_E && (1.26) \\
&= \alpha(\mu_A^{\Sigma,E}[\text{op}([t_1]_E, \dots, [t_n]_E)])_E && (1.47) \\
&= \alpha[\text{op}(t_1, \dots, t_n)]_E && (1.30) \\
&= \alpha[t]_E.
\end{aligned}$$

We obtain a functor $P^{-1} : \mathbf{EM}(\mathcal{T}_{\Sigma,E}) \rightarrow \mathbf{Alg}(\Sigma, E)$ sending (A, α) to \mathbb{A}_α .

Finally, we need to check that P and P^{-1} are inverses to each other, i.e. that $\alpha_{\mathbb{A}_\alpha} = \alpha$ and $\mathbb{A}_{\alpha_{\mathbb{A}}} = \mathbb{A}$. For the former, $\alpha_{\mathbb{A}_\alpha}$ is defined to be the interpretation $\llbracket - \rrbracket_\alpha$ extended to terms modulo E , which we showed in Footnote 179 acts just like α . For the latter, we need to show that $\llbracket - \rrbracket_{\alpha_{\mathbb{A}}}$ and $\llbracket - \rrbracket_A$ coincide. Using Footnote 179 for the first equation and the definition of $\alpha_{\mathbb{A}}$ for the second, we have

$$\llbracket t \rrbracket_{\alpha_{\mathbb{A}}} = \alpha_{\mathbb{A}}[t]_E = \llbracket t \rrbracket_A.$$

Therefore, P and P^{-1} are inverses, thus $\mathbf{Alg}(\Sigma, E)$ and $\mathbf{EM}(\mathcal{T}_{\Sigma,E})$ are isomorphic.¹⁸⁰ \square

Remark 1.71. This result (along with the construction of free (Σ, E) -algebras in Proposition 1.49) means that $U : \mathbf{Alg}(\Sigma, E) \rightarrow \mathbf{Set}$ is a (strictly) **monadic** functor. I decided not to define or discuss monadic functors in this document in order to have fewer prerequisites,¹⁸¹ and because I like to exhibit the explicit isomorphism between categories of algebras. MacLane proves Proposition 1.70 using a monadicity theorem in [Mac71, §VI.8, Theorem 1].

What about algebras for other monads? Are they algebras for some signature Σ and equations E ?

Example 1.72 (Maybe). In \mathbf{Set} , a $(- + \mathbf{1})$ -algebra is a function $\alpha : A + \mathbf{1} \rightarrow A$ making the following diagrams commute.

$$\begin{array}{ccc} A & \xrightarrow{\eta_A} & A + \mathbf{1} \\ & \searrow \text{id}_A & \downarrow \alpha \\ & & A \end{array} \qquad \begin{array}{ccc} A + \mathbf{1} + \mathbf{1} & \xrightarrow{\mu_A} & A + \mathbf{1} \\ \alpha + \mathbf{1} \downarrow & & \downarrow \alpha \\ A + \mathbf{1} & \xrightarrow{\alpha} & A \end{array}$$

Reminding ourselves that η_A is the inclusion in the left component, the triangle commuting enforces α to act like the identity function on all of A . We can also write $\alpha = [\text{id}_A, \alpha(*)]$.¹⁸² The square commuting adds no constraint. Thus, an algebra for the maybe monad on \mathbf{Set} is just a set with a distinguished point. Let $h : A \rightarrow B$ be a function, commutativity of (1.53) is equivalent to $h(\alpha(*)) = \beta(*)$. Hence, a $(- + \mathbf{1})$ -homomorphism is a function that preserves the distinguished point.

Seeing the distinguished point of a $(- + \mathbf{1})$ -algebra as the interpretation of a constant, we recognize that the category $\mathbf{EM}(- + \mathbf{1})$ is isomorphic to the category $\mathbf{Alg}(\Sigma)$ where $\Sigma = \{\rho : 0\}$ contains a single constant.¹⁸³

Another option to recognize $\mathbf{EM}(- + \mathbf{1})$ as a category of algebras is via monad isomorphisms.

Definition 1.73 (Monad morphism). Let (M, η^M, μ^M) and (N, η^N, μ^N) be two monads on \mathbf{C} . A **monad morphism** from M to N is a natural transformation $\rho : M \Rightarrow N$ making (1.54) and (1.55) commute.¹⁸⁴

$$\begin{array}{ccc} \text{id}_{\mathbf{C}} & & \\ \eta^M \downarrow & \searrow \eta^N & \\ M & \xrightarrow{\rho} & N \end{array} \quad (1.54)$$

$$\begin{array}{ccc} MM & \xrightarrow{\rho \circ \rho} & NN \\ \mu^M \downarrow & & \downarrow \mu^N \\ M & \xrightarrow{\rho} & N \end{array} \quad (1.55)$$

¹⁸⁰ Observe that the functors P and P^{-1} commute with the forgetful functors because they do not change the carriers of the algebras.

¹⁸¹ I became comfortable with monadicity theorems relatively late into my PhD, so I think avoiding them keeps things more accessible.

¹⁸² We identify the element $\alpha(*) \in A$ with the function $\alpha(*) : \mathbf{1} \rightarrow A$ picking out that element.

$$\begin{array}{ccc} A + \mathbf{1} & \xrightarrow{h + \mathbf{1}} & B + \mathbf{1} \\ [\text{id}_A, \alpha(*)] \downarrow & & \downarrow [\text{id}_B, \beta(*)] \\ A & \xrightarrow{h} & B \end{array} \quad (1.53)$$

¹⁸³ Notice, again, that this isomorphism would commute with the forgetful functors to \mathbf{Set} because the carriers are unchanged.

¹⁸⁴ Recall that $\rho \circ \rho$ denotes the horizontal composition of ρ with itself, i.e.

$$\rho \circ \rho = \rho N \cdot M \rho = N \rho \cdot \rho M.$$

As expected ρ is called a monad isomorphism when there is a monad morphism $\rho^{-1} : N \Rightarrow M$ satisfying $\rho \cdot \rho^{-1} = \mathbb{1}_N$ and $\rho^{-1} \cdot \rho = \mathbb{1}_M$. In fact, it is enough that all the components of ρ are isomorphisms in \mathbf{C} to guarantee ρ is a monad isomorphism.¹⁸⁵

Example 1.74. For the signature $\Sigma = \{p:0\}$, the term monad \mathcal{T}_Σ is isomorphic to $- + \mathbf{1}$. Indeed, recall that a Σ -term over A is either an element of A or p , this yields a bijection $\rho_A : \mathcal{T}_\Sigma A \rightarrow A + \mathbf{1}$ that sends any element of A to itself and p to $* \in \mathbf{1}$. To verify that ρ is a monad morphism, we check these diagrams commute.¹⁸⁶

$$\begin{array}{ccc} \mathcal{T}_\Sigma A & \xrightarrow{\rho_A} & A + \mathbf{1} \\ \mathcal{T}_\Sigma f \downarrow & & \downarrow f + \mathbf{1} \\ \mathcal{T}_\Sigma B & \xrightarrow{\rho_B} & B + \mathbf{1} \end{array} \quad (1.56)$$

$$\begin{array}{ccc} A & & \\ \eta_A^\Sigma \downarrow & \searrow \eta_A & \\ \mathcal{T}_\Sigma A & \xrightarrow{\rho_A} & A + \mathbf{1} \end{array} \quad (1.57)$$

$$\begin{array}{ccc} \mathcal{T}_\Sigma \mathcal{T}_\Sigma A & \xrightarrow{\rho_{\mathcal{T}_\Sigma A} \circ (\rho_A + \mathbf{1})} & A + \mathbf{1} + \mathbf{1} \\ \mu_A^\Sigma \downarrow & & \downarrow \mu_A \\ \mathcal{T}_\Sigma A & \xrightarrow{\rho_A} & A + \mathbf{1} \end{array} \quad (1.58)$$

We obtain a monad isomorphism between the maybe monad and the term monad for the signature $\Sigma = \{p:0\}$. We can recover the isomorphism between the categories of algebras from Example 1.72 with the following result.

Proposition 1.75. *If $\rho : M \Rightarrow N$ is a monad morphism, then there is a functor $-\rho : \mathbf{EM}(N) \rightarrow \mathbf{EM}(M)$. If ρ is a monad isomorphism, then $-\rho$ is also an isomorphism.*

Proof. Given an N -algebra $\alpha : NA \rightarrow A$, we show that $\alpha \circ \rho_A : MA \rightarrow A$ is an M -algebra by paving the following diagrams.

$$\begin{array}{ccc} A & \xrightarrow{\eta_A^M} & MA \\ & \searrow \eta_A^N \text{ (a)} & \downarrow \rho_A \\ & & NA \\ & \searrow \text{id}_A \text{ (b)} & \downarrow \alpha \\ & & A \end{array} \quad \begin{array}{ccc} MMA & \xrightarrow{\mu_A^M} & MA \\ M\rho_A \downarrow & & \downarrow \rho_A \\ MNA & \xrightarrow{\rho_{NA}} & NNA \xrightarrow{\mu_A^N} NA \\ M\alpha \downarrow \text{ (d)} & & N\alpha \downarrow \text{ (e)} \\ MA & \xrightarrow{\rho_A} & NA \xrightarrow{\alpha} A \end{array} \quad (1.59)$$

Moreover, if $h : A \rightarrow B$ is an N -homomorphism from α to β , then it is also a M -homomorphism from $\alpha \circ \rho_A$ to $\beta \circ \rho_B$ by the paving below.¹⁸⁷

$$\begin{array}{ccc} MA & \xrightarrow{Mh} & MB \\ \rho_A \downarrow & & \downarrow \rho_B \\ NA & \xrightarrow{Nh} & NB \\ \alpha \downarrow & & \downarrow \beta \\ A & \xrightarrow{h} & B \end{array}$$

We obtain a functor $-\rho : \mathbf{EM}(N) \rightarrow \mathbf{EM}(M)$ taking an algebra (A, α) to $(A, \alpha \circ \rho_A)$ and a homomorphism $h : (A, \alpha) \rightarrow (B, \beta)$ to $h : (A, \alpha \circ \rho_A) \rightarrow (B, \beta \circ \rho_B)$.

Furthermore, it is easy to see that $-\rho = \text{id}_{\mathbf{EM}(M)}$ when $\rho = \mathbb{1}_M$ is the identity monad morphism, and that for any other monad morphism $\rho' : N \Rightarrow L$, $-(\rho' \cdot \rho) =$

¹⁸⁵ One checks that natural isomorphisms are precisely the natural transformations whose components are all isomorphisms, and that the inverse of a monad morphism is a monad morphism.

¹⁸⁶ All of them commute essentially because ρ_A and both multiplications act like the identity on A .

Showing (1.59) commutes:

(a) By (1.54).

(b) By (1.47) for $\alpha : NA \rightarrow A$.

(c) By (1.55), noting that $(\rho \circ \rho)_A = \rho_{NA} \circ M\rho_A$.

(d) Naturality of ρ .

(e) By (1.48) for $\alpha : NA \rightarrow A$.

¹⁸⁷ The top square commutes by naturality of ρ and the bottom square commutes because h is an N -homomorphism (1.49).

$(-\rho) \circ (-\rho')$.¹⁸⁸ Thus, when ρ is a monad isomorphism with inverse ρ^{-1} , $-\rho^{-1}$ is the inverse of $-\rho$, so $-\rho$ is an isomorphism. \square

With the monad isomorphism $\mathcal{T}_\Sigma \cong - + \mathbf{1}$ of Example 1.74, we obtain an isomorphism $\mathbf{EM}(- + \mathbf{1}) \cong \mathbf{EM}(\mathcal{T}_\Sigma)$, and composing it with the isomorphism of Proposition 1.70 $\mathbf{EM}(\mathcal{T}_\Sigma) \cong \mathbf{Alg}(\Sigma)$ (instantiating $E = \emptyset$), we get back the result from Example 1.72 that algebras for the maybe monad are the same thing as algebras for the signature with a single constant.

In general, we now know that $\mathcal{T}_{\Sigma,E} \cong M$ implies $\mathbf{EM}(M) \cong \mathbf{Alg}(\Sigma, E)$, but constructing a monad isomorphism (and showing it is one) is not always the easiest thing to do.¹⁸⁹ There is a converse implication, but it requires a restriction to isomorphisms of categories that commute with the forgetful functors to **Set**. Anyways, that is a mild condition we foreshadowed.

Proposition 1.76. *If $P : \mathbf{EM}(N) \rightarrow \mathbf{EM}(M)$ is a functor such that $U^M \circ P = U^N$, then there is a monad morphism $\rho : M \rightarrow N$. If P is an isomorphism, then so is ρ .*

Proof. Quick corollary of [BW05, Chapter 3, Theorem 6.3]. \square

This motivates the following definition which states that a monad M is presented by (Σ, E) when it is isomorphic to the term monad $\mathcal{T}_{\Sigma,E}$ or, thanks to Proposition 1.76 and Proposition 1.70, when M -algebras on A and (Σ, E) -algebras on A are identified.

Definition 1.77 (Set presentation). Let M be a monad on **Set**, an **algebraic presentation** of M is signature Σ and a class of equations E along with a monad isomorphism $\rho : \mathcal{T}_{\Sigma,E} \cong M$. We also say M is presented by (Σ, E) .

We chose to state the definition with the monad isomorphism it makes some arguments in §3.5 quicker. Showing that a monad is presented by (Σ, E) can be done in many ways that are equivalent to building a monad isomorphism.¹⁹⁰

We have proven in Example 1.74 that $\Sigma = \{p : 0\}$ and $E = \emptyset$ is an algebraic presentation for the maybe monad on **Set**. Here is a couple of additional examples.

Example 1.78 (Powerset). The powerset monad \mathcal{P}_{ne} is presented by the theory of **semilattices** (Σ_S, E_S) ,¹⁹¹ where $\Sigma_S = \{\oplus : 2\}$ and E_S contains the following equations stating that \oplus is idempotent, commutative and associative respectively.

$$x \vdash x = x \oplus x \quad x, y \vdash x \oplus y = y \oplus x \quad x, y, z \vdash x \oplus (y \oplus z) = (x \oplus y) \oplus z$$

This means there is a monad isomorphism $\mathcal{T}_{\Sigma_S, E_S} \cong \mathcal{P}_{ne}$.

Another thing we obtain from this isomorphism is that for any set X , interpreting \oplus as union on $\mathcal{P}_{ne}X$ (i.e. $(S, T) \mapsto S \cup T$) yields the free semilattice on X .¹⁹²

Example 1.79 (Distributions). The distribution monad \mathcal{D} is presented by the theory of **convex algebras** (Σ_{CA}, E_{CA}) where $\Sigma_{CA} = \{+_p : 2 \mid p \in (0, 1)\}$ and E_{CA} contains the following equations for all $p, q \in (0, 1)$.

$$x \vdash x = x +_p x \quad x, y \vdash x +_p y = y +_{1-p} x$$

¹⁸⁸ In other words, the assignments $M \mapsto \mathbf{EM}(M)$ and $\rho \mapsto -\rho$ becomes a functor from the category of monads on **C** and monad morphisms to the category of categories (ignoring size issues).

¹⁸⁹ For instance, the isomorphism of categories of algebras in Example 1.72 is definitely clearer than the isomorphism of monads in Example 1.74.

¹⁹⁰ We already gave one with Proposition 1.76, and you can also read some great discussions in Remark 3.6 and §4.2 in [BSV22].

¹⁹¹ Usually, when we say “theory of X ”, we mean that X s are the algebras for that theory. For instance, semilattices are the (Σ_S, E_S) -algebras. After some unrolling, we get the more common definition of a semilattice, that is, a set with a binary operation that is idempotent, commutative, and associative.

¹⁹² It is relatively easy to show that union is idempotent, commutative, and associative, freeness is more difficult but follows from the algebraic presentation, and the fact that $(\mathcal{P}_{ne}X, \mu_X)$ is the free \mathcal{P}_{ne} -algebra (recall Example 1.69).

$$x, y, z \vdash (x +_p y) +_q z = x +_{pq} + (y +_{\frac{p(1-q)}{1-pq}} z)$$

The free convex algebra on X can now be seen as $\mathcal{D}X$ with $+_p$ interpreted as the usual convex combination, that is,¹⁹³

$$\llbracket \varphi +_p \psi \rrbracket_{\mathcal{D}X} = p\varphi + (1-p)\psi = (x \mapsto p\varphi(x) + (1-p)\psi(x)). \quad (1.60)$$

Remark 1.80. Not all monads on **Set** have an algebraic presentation.¹⁹⁴ The monads that can be presented by a signature with finitary operation symbols are aptly called **finitary monads**. They can be characterized as the monads whose underlying functor preserve limits of a certain shape and size, see e.g. [Bor94, Proposition 4.6.2].

In Chapter 3, we will need to relate monads on different categories, we give some background on that here.

Definition 1.81 (Lax monad morphism). Let (T, η^T, μ^T) be a monad on \mathbf{D} , and (M, η^M, μ^M) be a monad on \mathbf{C} . A **lax monad morphism** from T to M is a pair (F, λ) comprising a functor $F : \mathbf{C} \rightarrow \mathbf{D}$, and a natural transformation $\lambda : TF \Rightarrow FM$ making (1.61) and (1.62) commute.¹⁹⁵

$$(1.61) \quad \begin{array}{ccc} F & & \\ \eta^T F \downarrow & \searrow F\eta^M & \\ TF & \xrightarrow{\lambda} & FM \end{array} \quad (1.62) \quad \begin{array}{ccccc} TTF & \xrightarrow{T\lambda} & TFM & \xrightarrow{\lambda M} & FMM \\ \mu^T F \downarrow & & \downarrow F\mu^M & & \\ TF & \xrightarrow{\lambda} & FM & & \end{array}$$

Proposition 1.82. If $(F, \lambda) : T \rightarrow M$ is a lax monad morphism, then there is a functor $F - \circ \lambda : \mathbf{EM}(M) \rightarrow \mathbf{EM}(T)$ sending an M -algebra $\alpha : MA \rightarrow A$ to $F\alpha \circ \lambda_A : TFA \rightarrow A$, and an M -homomorphism $h : A \rightarrow B$ to $Fh : FA \rightarrow FB$.¹⁹⁶

Proof. We need to show that $F\alpha \circ \lambda$ is a T -algebra whenever α is an M -algebra. We pave the following diagrams showing (1.47) and (1.48) commute respectively.

$$(1.64) \quad \begin{array}{ccc} FA & \xrightarrow{\eta_{FA}^T} & TFA \\ \downarrow F\eta_A^M & \searrow (a) & \downarrow \lambda_A \\ FMA & & FMA \\ \downarrow F\alpha & \searrow (b) & \downarrow F\alpha \\ FA & & FA \end{array} \quad \begin{array}{ccc} TFA & \xrightarrow{\mu_{FA}^T} & TFA \\ T\lambda_A \downarrow & (c) & \downarrow \lambda_A \\ TFMA & \xrightarrow{\lambda_{MA}} & FMMA \xrightarrow{F\mu_A^M} & FMA \\ TF\alpha \downarrow & (d) & FM\alpha \downarrow & (e) & \downarrow F\alpha \\ TFA & \xrightarrow{\lambda_A} & FMA & \xrightarrow{F\alpha} & FA \end{array}$$

Next, we need to show that when $h : A \rightarrow B$ is an M -homomorphism from α to β , then Fh is a T -homomorphism from $F\alpha \circ \lambda_A$ to $F\alpha \circ \lambda_B$. We pave the following diagram where (a) commutes by naturality of λ and (b) by applying F to (1.49).

$$\begin{array}{ccc} TFA & \xrightarrow{TFh} & TFB \\ \lambda_A \downarrow & (a) & \downarrow \lambda_B \\ FMA & \xrightarrow{FMh} & FMB \\ F\alpha \downarrow & (b) & \downarrow F\beta \\ FA & \xrightarrow{Fh} & FB \end{array}$$

¹⁹³ For later, we will write \bar{p} for $1-p$.

¹⁹⁴ For example, the *full* powerset monad does not, although it still has an algebraic flavor as its algebras are in correspondence with complete sup-lattices, see e.g. [Bor94, Proposition 4.6.5].

¹⁹⁵ Note the similarities with Definition 1.73, lax monad morphisms generalize monad morphisms to monads on different base categories. The terminology comes from e.g. [LS12, Rie13], but the name monad functor was originally used in [Str72] and the direction of morphisms is sometimes reversed.

¹⁹⁶ By definition, the functor $F - \circ \lambda$ lifts F along the forgetful functors, namely, it makes (1.63) commute.

$$(1.63) \quad \begin{array}{ccc} \mathbf{EM}(M) & \xrightarrow{F - \circ \lambda} & \mathbf{EM}(T) \\ U^M \downarrow & & \downarrow U^T \\ \mathbf{C} & \xrightarrow{F} & \mathbf{D} \end{array}$$

Showing (1.64) commutes:

- (a) By (1.61).
- (b) Apply F to (1.47).
- (c) By (1.62).
- (d) Naturality of λ .
- (e) Apply F to (1.48).

□

There are two special cases of lax monad morphisms. When T and M are on the same category \mathbf{C} and $F = \text{id}_{\mathbf{C}}$, a lax monad morphism is just a monad morphism from T to M ,¹⁹⁷ and then the proof above reduces to the proof of Proposition 1.75. When λ_A is an identity morphism for every A , i.e. $TF = FM$, we say that M is a monad lifting of T along F . That notion is central to §3.5, where we redefine it in a more specific setting.

Our goal for the next two chapters is to make all the results here more general by considering carriers to be generalized metric spaces, i.e. sets with a notion of distance. In Chapter 2 we define what we mean by distance, and in Chapter 3, we define quantitative algebras, quantitative equational logic, and quantitative algebraic presentations analogously to the definitions above.

¹⁹⁷ Sometimes, authors introduce lax monad morphisms with the name monad morphism, and take our notion of monad morphism as a particular instance. Some authors also use the name monad map for either notion.

2 Generalized Metric Spaces

The Homeless Wanderer

Emahoy Tsegué-Maryam Guèbrou

2.1 L-Spaces	60
2.2 Equational Constraints	70
2.3 The Categories GMet	77

For a comprehensive introduction to the concepts and themes explored in this chapter, please refer to §0.2. Here, we only give a brief overview.

In this chapter, we give our definition of generalized metric spaces which is different from the many definitions already in the literature.¹⁹⁸ Once again, we take our time with this material in preparation for the next chapter, introducing many examples and disseminating some insights along the way. While the content of Chapter 1 can safely be skipped before reading the current chapter, our main point here is the definition of quantitative equation (Definition 2.23) as an answer to the question “How do we impose constraints on distances with the familiar syntax of equations?”, thus it makes sense to be comfortable with equational reasoning before reading what follows.

Outline: In §2.1, we define complete lattices and relations valued in a complete lattice, we also give an equivalent definition that justifies the syntax of quantitative equations. In §2.2, we define quantitative equations and the categories of generalized metric spaces which are parametrized by collections of quantitative equations. In §2.3, we study the properties that all categories of generalized metric spaces have.

2.1 L-Spaces

Chapter 1 is titled *Universal Algebra* and Chapter 3 is titled *Universal Quantitative Algebra*. In order to go from the former to the latter, we will explain what we mean by *quantitative*. In the original paper on quantitative algebras [MPP16], and in many other works on quantitative program semantics,¹⁹⁹ the **quantities** considered are, more often than not, positive real numbers. In [MSV22, MSV23], we worked with quantities inside $[0, 1]$. In this document, we will abstract away from real numbers, thinking of quantities as things you can compare and say whether one is bigger or smaller than another. You can do that with positive real numbers thanks to the usual ordering \leq , but it has a crucial property that we exploit, it is *complete* in the (informal) sense that you can always find the smallest quantity of a set of real numbers. Formally it is a complete lattice.²⁰⁰

Definition 2.1 (Complete lattice). A **complete lattice** is a partially ordered set²⁰¹ (or

¹⁹⁸ See e.g. [BvBR98, Bra00, Pis21].

¹⁹⁹ e.g. [Kwio7, vBW01, KyKK⁺21, ZK22].

²⁰⁰ Small caveat: we need to add ∞ or work with an upper bound (see Examples 2.2 and 2.3).

²⁰¹ i.e. L is a set and $\leq \subseteq L \times L$ is a binary relation on L that is reflexive, transitive and antisymmetric.

poset) (L, \leq) where all subsets $S \subseteq L$ have an infimum and supremum denoted by $\inf S$ and $\sup S$ respectively. In particular, L has a **bottom element** $\perp = \inf L = \sup \emptyset$ and a **top element** $\top = \sup L = \inf \emptyset$ that satisfy $\perp \leq \varepsilon \leq \top$ for all $\varepsilon \in L$. We use L to refer to the lattice and its underlying set, and we call its elements **quantities**.²⁰²

Let us describe two central (for this thesis) examples of complete lattices.

Example 2.2 (Unit interval). The **unit interval** $[0, 1]$ is the set of real numbers between 0 and 1. It is a poset with the usual order \leq (“less than or equal”) on numbers. It is usually an axiom in the definition of \mathbb{R} that all non-empty bounded subsets of real numbers have an infimum and a supremum. Since all subsets of $[0, 1]$ are bounded (by 0 and 1), we conclude that $([0, 1], \leq)$ is a complete lattice with $\perp = 0$ and $\top = 1$.

Later in this section, we will see elements of $[0, 1]$ as distances between points of some space. It would make sense, then, to extend the interval to contain values bigger than 1. Still because a complete lattice must have a top element there must be a number above all others. We could either stop at some arbitrary $0 \leq B \in \mathbb{R}$ and consider $[0, B]$, or we can consider ∞ to be a number as done below.²⁰³

Example 2.3 (Extended interval). Similarly to the unit interval, the **extended interval** is the set $[0, \infty]$ of positive real numbers extended with ∞ , and it is a poset after asserting $\varepsilon \leq \infty$ for all $\varepsilon \in [0, \infty]$. It is also a complete lattice because non-empty bounded subsets of $[0, \infty)$ still have an infimum and supremum, and if a subset is not bounded above or contains ∞ , then its supremum is ∞ . We find that 0 is bottom and ∞ is top.

It is the prevailing custom to consider distances valued in the extended interval.²⁰⁴ In our papers [MSV21, MSV22, MSV23], we worked with the unit interval, but in theory, there is no difference since $[0, 1]$ and $[0, \infty]$ are isomorphic as complete lattices.²⁰⁵ In practice, one can use additional structure and properties that are not preserved by this isomorphism (like adding quantities).

Remark 2.4. The first two examples are both **quantales** [HST14, §II.1.10], informally, complete lattices where quantities can be added together in a way that preserves the order and the “smallest” quantities. It is also quite common in the literature on quantitative programming semantics to generalize from real numbers to elements of a quantale.²⁰⁶ Since none of the results we establish require dealing with addition, we will work at the level of generality of complete lattices (no difficulty arises from this abstraction), even though many of the following examples are quantales.

There are many other interesting complete lattices, although (unfortunately) they are less often viewed as possible places to value distances.

Example 2.5 (Booleans). The **Boolean lattice** B is the complete lattice containing only two elements, bottom and top. Its name comes from the interpretation of \perp as a false value and \top as a true value which makes the infimum act like an AND and the supremum like an OR.

Example 2.6 (Extended natural numbers). The set \mathbb{N}_∞ of natural numbers extended with ∞ is a sublattice of $[0, \infty]$.²⁰⁷ Indeed, it is a poset with the usual order and

²⁰² The name quantity is not standard, we use this terminology only in the context of our work.

²⁰³ If one needs negative distances, it is also possible to work with any interval $[A, B]$ with $A \leq B \in \mathbb{R}$, or even $[-\infty, \infty]$. We will stick to $[0, 1]$ and $[0, \infty]$.

²⁰⁴ In fact, $[0, \infty]$ with the reverse order and additional structure is also famous under the name *Lawvere quantale* because of Lawvere’s seminal paper [Law02]. In that work, he used the **quantale** structure on $[0, \infty]$ to give a categorical definition very close to that of a metric.

²⁰⁵ Take the mapping $x \mapsto \frac{1}{1-x} - 1$ from $[0, 1]$ to $[0, \infty]$ with $\frac{1}{0} - 1 = \infty$. It is bijective, monotone and preserves infimums.

²⁰⁶ e.g. [DGY19, GP21, GD23, FSW⁺23].

²⁰⁷ As expected, a (complete) **sublattice** of (L, \leq) is a set $S \subseteq L$ closed under taking infimums and supremums. Note that the top and bottom of S need not coincide with those of L . For instance $[0, 1]$ is a sublattice of $[0, \infty]$, but $\top = 1$ in the former and $\top = \infty$ in the latter.

the infimum and supremum of a subset of natural numbers is either itself a natural number or ∞ (when the subset is empty or unbounded respectively).

Example 2.7 (Powerset lattice). For any set X , we denote the powerset of X by $\mathcal{P}(X)$. The inclusion relation \subseteq between subsets of X makes $\mathcal{P}(X)$ a poset. The infimum of a family of subsets $S_i \subseteq X$ is the intersection $\bigcap_{i \in I} S_i$, and its supremum is the union $\bigcup_{i \in I} S_i$. Hence, $\mathcal{P}(X)$ is a complete lattice. The bottom element is \emptyset and the top element is X .

It is well-known that subsets of X correspond to functions $X \rightarrow \{\perp, \top\}$.²⁰⁸ Endowing the two-element set with the complete lattice structure of \mathbf{B} is what yields the complete lattice structure on $\mathcal{P}(X)$. The following example generalizes this construction.

Example 2.8 (Function space). Given a complete lattice (L, \leq) , for any set X , we denote the set of functions from X to L by L^X . The pointwise order on functions defined by

$$f \leq_* g \iff \forall x \in X, f(x) \leq g(x)$$

is a partial order on L^X . The infimums and supremums of families of functions are also computed pointwise. Namely, given $\{f_i : X \rightarrow L\}_{i \in I}$, for all $x \in X$:

$$(\inf_{i \in I} f_i)(x) = \inf_{i \in I} f_i(x) \quad \text{and} \quad (\sup_{i \in I} f_i)(x) = \sup_{i \in I} f_i(x).$$

This makes L^X a complete lattice. The bottom element is the function that is constant at \perp , and the top element is the function that is constant at \top .

As a special case of function spaces, it is easy to show that when X is a set with two elements, L^X is isomorphic (as complete lattices) to the product $L \times L$.

Example 2.9 (Product). Let (L, \leq_L) and (K, \leq_K) be two complete lattices. Their **product** is the poset $(L \times K, \leq_{L \times K})$ on the cartesian product of L and K with the order defined by

$$(\varepsilon, \delta) \leq_{L \times K} (\varepsilon', \delta') \iff \varepsilon \leq_L \varepsilon' \text{ and } \delta \leq_K \delta'. \quad (2.1)$$

It is a complete lattice where the infimums and supremums are computed coordinatewise, namely, for any $S \subseteq L \times K$,²⁰⁹

$$\begin{aligned} \inf S &= (\inf\{\pi_L(c) \mid c \in S\}, \inf\{\pi_K(c) \mid c \in S\}) \text{ and} \\ \sup S &= (\sup\{\pi_L(c) \mid c \in S\}, \sup\{\pi_K(c) \mid c \in S\}). \end{aligned}$$

The bottom (resp. top) element of $L \times K$ is the pairing of the bottom (resp. top) elements of L and K . i.e. $\perp_{L \times K} = (\perp_L, \perp_K)$ and $\top_{L \times K} = (\top_L, \top_K)$.

The following example is also based on functions, and it appears in several works on generalized notions of distances, e.g. [Fla97, HR13].

²⁰⁸ A subset $S \subseteq X$ is sent to the characteristic function χ_S , and a function $f : X \rightarrow \mathbf{B}$ is sent to $f^{-1}(\top)$. We say that $\{\perp, \top\}$ is the subobject classifier of \mathbf{Set} .

Taking $L = \mathbf{B}$, we find that $\mathcal{P}(X)$ and \mathbf{B}^X are isomorphic as complete lattices under the usual correspondence. Namely, pointwise infimums and supremums become intersections and unions respectively. For example, if $\chi_S, \chi_T : X \rightarrow \mathbf{B}$ are the characteristic functions of $S, T \subseteq X$, then

$$\begin{aligned} \inf\{\chi_S, \chi_T\}(x) = \top &\iff \chi_S(x) = \chi_T(x) = \top \\ &\iff x \in S \text{ and } x \in T \\ &\iff x \in S \cap T. \end{aligned}$$

²⁰⁹ Where π_L and π_K are the projections from $L \times K$ to L and K respectively.

Example 2.10 (CDF). A **cumulative distribution function**²¹⁰ (or CDF for short) is a function $f : [0, \infty] \rightarrow [0, 1]$ that is monotone (i.e. $\varepsilon \leq \delta \implies f(\varepsilon) \leq f(\delta)$) and satisfies

$$f(\delta) = \sup\{f(\varepsilon) \mid \varepsilon < \delta\}. \quad (2.2)$$

Intuitively, (2.2) says that f cannot abruptly change value at some $x \in [0, \infty]$, but it can do that “after” some x .²¹¹ For instance, out of the two functions below, only $f_{>1}$ is a CDF.

$$f_{\geq 1} = x \mapsto \begin{cases} 0 & x < 1 \\ 1 & x \geq 1 \end{cases} \quad f_{>1} = x \mapsto \begin{cases} 0 & x \leq 1 \\ 1 & x > 1 \end{cases}$$

We denote by $\text{CDF}([0, \infty])$ the subset of $[0, 1]^{[0, \infty]}$ containing all CDFs, it inherits a poset structure (pointwise ordering), and we can show it is a complete lattice.²¹²

Let $\{f_i : [0, \infty] \rightarrow [0, 1]\}_{i \in I}$ be a family of CDFs. We will show the pointwise supremum $\sup_{i \in I} f_i$ is a CDF, and that is enough since having all supremums implies having all infimums [DP02, Theorem 2.31].

- If $\varepsilon \leq \delta$, since all f_i s are monotone, we have $f_i(\varepsilon) \leq f_i(\delta)$ for all $i \in I$ which implies

$$\left(\sup_{i \in I} f_i\right)(\varepsilon) = \sup_{i \in I} f_i(\varepsilon) \leq \sup_{i \in I} f_i(\delta) = \left(\sup_{i \in I} f_i\right)(\delta).$$

- For any $\delta \in [0, \infty]$, we have

$$\left(\sup_{i \in I} f_i\right)(\delta) = \sup_{i \in I} f_i(\delta) = \sup_{i \in I} \sup_{\varepsilon < \delta} f_i(\varepsilon) = \sup_{\varepsilon < \delta} \sup_{i \in I} f_i(\varepsilon) = \sup_{\varepsilon < \delta} \left(\sup_{i \in I} f_i\right)(\varepsilon).$$

Nothing prevents us from defining CDFs on other domains, and we will write $\text{CDF}(L)$ for the complete lattice of functions $L \rightarrow [0, 1]$ that are monotone and satisfy (2.2). This is a concrete instance of a more general fact that the set of Scott-continuous functions with the pointwise order has all supremums computed pointwise (see, e.g. [GHK⁺03, Lemma II-2.5]).

Definition 2.11 (L-space). Given a complete lattice L and a set A , an **L-relation** on A is a function $d : A \times A \rightarrow L$. We call the pair (A, d) an **L-space**, and A its **carrier** or **underlying set**. We will also use a single bold-face symbol \mathbf{A} to refer to an L-space with underlying set A and L-relation $d_{\mathbf{A}}$.²¹³

A **nonexpansive** map from \mathbf{A} to \mathbf{B} is a function $f : A \rightarrow B$ between the underlying sets of \mathbf{A} and \mathbf{B} that satisfies

$$\forall x, x' \in A, \quad d_{\mathbf{B}}(f(x), f(x')) \leq d_{\mathbf{A}}(x, x'). \quad (2.3)$$

The identity maps $\text{id}_A : A \rightarrow A$ and the composition of two nonexpansive maps are always nonexpansive²¹⁴, therefore we have a category whose objects are L-spaces and morphisms are nonexpansive maps. We denote it by **LSpa**.

This category is concrete over **Set** with the forgetful functor $U : \text{LSpa} \rightarrow \text{Set}$ which sends an L-space \mathbf{A} to its carrier and a morphism to the underlying function between carriers.

²¹⁰ Although cumulative *sub*distribution function might be preferred.

²¹¹ This property is often called *right-continuity*.

²¹² Note however that $\text{CDF}([0, \infty])$ is not a sublattice of $[0, 1]^{[0, \infty]}$ because the infimums are not always taken pointwise. For instance, given $0 < n \in \mathbb{N}$, define f_n by (see them on [Desmos](#))

$$f_n(x) = \begin{cases} 0 & x \leq 1 - \frac{1}{n} \\ nx & 1 - \frac{1}{n} < x < 1 \\ 1 & 1 \leq x \end{cases}$$

The pointwise infimum of $\{f_n\}_{n \in \mathbb{N}}$ clearly sends everything below 1 to 0 and everything above and including 1 to 1, so it does not satisfy $f(1) = \sup_{\varepsilon < 1} f(\varepsilon)$. We can find the infimum with the general formula that defines infimums in terms of supremums:

$$\inf_{n > 0} f_n = \sup\{f \in \text{CDF}([0, \infty]) \mid \forall n > 0, f \leq_* f_n\}.$$

We find that $\inf_{n > 0} f_n = f_{>1}$.

²¹³ We will often switch between referring to spaces with \mathbf{A} or $(A, d_{\mathbf{A}})$, and we will try to match the symbol for the space and the one for its underlying set only modifying the former with `mathbf{b}`.

²¹⁴ Fix three L-spaces \mathbf{A} , \mathbf{B} and \mathbf{C} with two nonexpansive maps $f : A \rightarrow B$ and $g : B \rightarrow C$, we have by nonexpansiveness of g then f :

$$\begin{aligned} d_{\mathbf{C}}(gf(a), gf(a')) &\leq d_{\mathbf{B}}(f(a), f(a')) \\ &\leq d_{\mathbf{A}}(a, a'). \end{aligned}$$

Remark 2.12. In the sequel, we will not distinguish between the morphism $f : \mathbf{A} \rightarrow \mathbf{B}$ and the underlying function $f : A \rightarrow B$. Although we may write Uf for the latter when disambiguation is necessary.

Instantiating \mathbf{L} for different complete lattices, we can get a feel for what the categories \mathbf{LSpa} look like. We also give concrete examples of \mathbf{L} -spaces.

Example 2.13 (Binary relations). When $\mathbf{L} = \mathbf{B}$, a function $d : A \times A \rightarrow \mathbf{B}$ is the same thing as a subset of $A \times A$, which is the same thing as a binary relation on A .²¹⁵ Then, a \mathbf{B} -space is a set equipped with a binary relation and we choose to have, as a convention, $d(a, a') = \perp$ when a and a' are related and $d(a, a') = \top$ when they are not.²¹⁶ A nonexpansive map from \mathbf{A} to \mathbf{B} is a function $f : A \rightarrow B$ such that for any $a, a' \in A$, $f(a)$ and $f(a')$ are related when a and a' are. When a and a' are not related, $f(a)$ and $f(a')$ might still be related.²¹⁷ The category \mathbf{BSpa} is well-known under different names, **EndoRel** in [Vig23], **Rel** in [AHS06] (although that name is more commonly used for the category where relations are morphisms) and **2Rel** in my book. Here are a couple of fun examples of \mathbf{B} -spaces:

1. **Chess.** Let P be the set of positions on a [chessboard](#) (a2, d6, f3, etc.) and $d_B : P \times P \rightarrow \mathbf{B}$ send a pair (p, q) to \perp if and only if q is accessible from p in one bishop's move. The pair (P, d_B) is an object of \mathbf{BSpa} . Let d_Q be the \mathbf{B} -relation sending (p, q) to \perp if and only if q is accessible from p in one queen's move. The pair (P, d_Q) is another object of \mathbf{BSpa} . The identity function $\text{id}_P : P \rightarrow P$ is nonexpansive from (P, d_B) to (P, d_Q) because whenever a bishop can go from p to q , a queen can too. However, it is not nonexpansive from (P, d_Q) to (P, d_B) because e.g. a queen can go from a1 to a2 but a bishop cannot.²¹⁸ One can check that any rotation of the chessboard is nonexpansive from (P, d_B) to itself and from (P, d_Q) to itself. And since nonexpansive maps compose, any rotation is also nonexpansive from (P, d_B) to (P, d_Q) .

2. **Siblings.** Let H be the set of all humans (me, Paul Erdős, my brother Paul, etc.) and $d_S : H \times H \rightarrow \mathbf{B}$ send (h, k) to \perp if and only if h and k are full siblings.²¹⁹ The pair (H, d_S) is an object of \mathbf{BSpa} . Let $d_=$ be the \mathbf{B} -relation sending (h, k) to \perp if and only if h and k are the same person. The pair $(H, d_=)$ is another object of \mathbf{BSpa} . The function $f : H \rightarrow H$ sending h to their biological mother is nonexpansive from (H, d_S) to $(H, d_=)$ because whenever h and k are full siblings, they have the same biological mother.

Example 2.14 (Distances). The main examples of \mathbf{L} -spaces in this thesis are $[0, 1]$ -spaces or $[0, \infty]$ -spaces. These are sets A equipped with a function $d : A \times A \rightarrow [0, 1]$ or $d : A \times A \rightarrow [0, \infty]$, and we can usually understand $d(a, a')$ as the distance between two points $a, a' \in A$. With this interpretation, a function is nonexpansive when applying it never increases the distances between points.²²⁰ Let us give several examples of $[0, 1]$ - and $[0, \infty]$ -spaces:

1. **Euclidean.** Probably the most famous distance in mathematics is the **Euclidean distance** on real numbers $d : \mathbb{R} \times \mathbb{R} \rightarrow [0, \infty] = (x, y) \mapsto |x - y|$. The distance

²¹⁵ Hence, the terminology \mathbf{L} -relation also appearing in e.g. [HST14, GD23].

²¹⁶ This convention might look backwards, but it makes sense with the morphisms.

²¹⁷ Note that this interpretation of nonexpansiveness depends on our just chosen convention. Swapping the meaning of $d(a, a') = \top$ and $d(a, a') = \perp$ is the same thing as taking the opposite order on \mathbf{B} (i.e. $\top \leq \perp$), namely, morphisms become functions $f : A \rightarrow B$ such that for any $a, a' \in A$, $f(a)$ and $f(a')$ are *not* related when neither are a and a' .

²¹⁸ In other words, the set of valid moves for a bishop is included in the set of valid moves for a queen, but not vice versa.

²¹⁹ Full siblings share the same biological parents.

²²⁰ This is a justification for the term nonexpansive. In the setting of distances being real-valued, another popular term is 1-Lipschitz.

between any two points is unbounded, but it is never ∞ . The pair (\mathbb{R}, d) is an object of $[0, \infty]\mathbf{Spa}$. It is also very common to study subsets of \mathbb{R} , like \mathbb{Q} or $[0, 1]$, with the Euclidean distance appropriately restricted. We say that (\mathbb{Q}, d) and $([0, 1], d)$ are subspaces of (\mathbb{R}, d) . In general, a **subspace** of an L-space \mathbf{A} is a subset $B \subseteq A$ equipped with the L-relation $d_{\mathbf{A}}$ restricted to B , i.e. $d_{\mathbf{B}} = B \times B \hookrightarrow A \times A \xrightarrow{d_{\mathbf{A}}} L$.

Multiplication by $r \in \mathbb{R}$ is a nonexpansive function $r \cdot - : (\mathbb{R}, d) \rightarrow (\mathbb{R}, d)$ if and only if r is between -1 and 1 . Intuitively, a function $f : (\mathbb{R}, d) \rightarrow (\mathbb{R}, d)$ is nonexpansive when its derivative at any point is between -1 and 1 .²²¹

²²¹ The derivatives might not exist, so this is just an informal explanation.

- Collaboration.** Let H be the set of humans again. A **collaboration chain** between two humans h and k is a sequence of scientific papers P_1, \dots, P_n such that h is a coauthor of P_1 , k is a coauthor of P_n , and P_i and P_{i+1} always have at least one common coauthor. The collaboration distance d between two humans h and k is the length of a shortest collaboration chain.²²² For instance $d(\text{me}, \text{Paul Erdős}) = 4$ as computed by csauthors.net on February 20th 2024:

me [PS21] D. Petrişan [GPR16] M. Gehrke [EGP07] M. Ern e [EE86] P. Erdős

²²² As conventions, the length of a chain is the number of papers, not humans. Also, $d(h, k) = \infty$ when no such chain exists between h and k , except when $h = k$, then $d(h, h) = 0$ (or we could say it is the length of the empty chain from h to h).

The pair (H, d) is a $[0, \infty]$ -space, but it could also be seen as a \mathbb{N}_{∞} -space (because the length of a chain is always an integer).

- Hamming.** Let W be the set of words of the English language. If two words u and v have the same number of letters, the Hamming distance $d(u, v)$ between u and v is the number of positions in u and v where the letters do not match.²²³ When u and v are of different lengths, we let $d(u, v) = \infty$, and we obtain a $[0, \infty]$ -space (W, d) . (It is also a \mathbb{N}_{∞} -space.)

²²³ For instance $d(\text{carrot}, \text{carpet}) = 2$ because these words differ only in two positions, the second and third to last ($r \neq p$ and $o \neq e$).

As Example 2.14 comes with many important intuitions, we will often call an L-relation $d : X \times X \rightarrow L$ a **distance function** and $d(x, y)$ the **distance** from x to y , even when L is neither $[0, 1]$ nor $[0, \infty]$.

Remark 2.15. The asymmetry in the terminology “distance from x to y ” is justified because, in general, nothing guarantees $d(x, y) = d(y, x)$. Since language is processed in a sequential order, we cannot even get rid of this asymmetry, but I feel like “distance *between* x and y ” would be more appropriate if we required $d(x, y) = d(y, x)$.

Example 2.16. We give more examples of L-spaces to showcase the potential of our abstract framework.

- Diversions.**²²⁴ Let J be the set of products available to consumers inside a vending machine (including a “no purchase” option), the second-choice diversion $d(p, q)$ from product p to product q is the fraction of consumers that switch from buying p to buying q when p is removed (or out of stock) from the machine. That fraction is always contained between 0 and 1, so we have a function $d : J \times J \rightarrow [0, 1]$ which makes (J, d) an object of $[0, 1]\mathbf{Spa}$.²²⁵

²²⁴ This example takes inspiration from the diversion matrices in [CMS23], where the authors consider the automobile market in the U.S.A. instead of a vending machine.

²²⁵ Even though d is valued in $[0, 1]$, calling it a distance function does not fit our intuition because when $d(p, q)$ is big, it means the products p and q are probably very similar.

2. **Rank.** Let P be the set of web pages available on the internet. In [BP98], the authors introduce an algorithm to measure the importance of a page $p \in P$ giving it a rank $R(p) \in [0, 1]$. This data can be compiled into a function $d_R : P \times P \rightarrow \mathbf{B}$ which sends (p, q) to \perp if and only if $R(p) \leq R(q)$, so d_R compares the ranks of web pages. This yields a B-space (P, d_R) .²²⁶

The rank of a page varies over time (it is computed from the links between all web pages which change quite frequently), so if we let T be the set of instants of time, we can define $d'_R(p, q)$ to be the function of type $T \rightarrow \mathbf{B}$ which sends t to the Boolean value of $R(p) \leq R(q)$ computed at time t . This makes (P, d'_R) into a \mathbf{B}^T -space.

In order to create a search engine, we also need to consider the input of the user looking for some web page.²²⁷ If U is the set of possible user inputs, we can define $d''_R(p, q)$ to depend on U and T , so that (P, d''_R) is a $\mathbf{B}^{U \times T}$ -space.

3. **Collaboration (bis).** In Example 2.14, we defined the collaboration distance $d : H \times H \rightarrow \mathbb{N}_\infty$ that measures how far two people are from collaborating on a scientific paper. We can define a finer measure by taking into account the total number of people involved in the collaboration. It allows us to say you are closer to Erdős if you wrote a paper with him and no one else than if you wrote a paper with him and two additional coauthors. The distance d' is now valued in $\mathbb{N}_\infty \times \mathbb{N}_\infty$,²²⁸ the first coordinate of $d'(h, k)$ is $d(h, k)$ the length of the shortest collaboration chain between h and k , and the second coordinate of $d'(h, k)$ is the smallest total number of authors in a collaboration chain of length $d(h, k)$. For instance, according to csauthors.net on February 20th 2024, there are only two chains of length four between me and Erdős, both involving (the same) seven people, hence $d'(\text{me}, \text{Paul Erdős}) = (4, 7)$.

4. **Bisimulation for CTS.** A conditional transition system (CTS) [ABH⁺12, Example 2.5] is a labelled transition system with a semantics different from the usual one. Instead of following transitions when the label matches an input, some label is chosen before the execution, and only those transitions which have the chosen label remain possible. Formulated differently, it is a family of transition systems on the same set of states indexed by a set of labels. If X is the set of states, and L is the set of labels, we can define a $\mathcal{P}(L)$ -relation $d : X \times X \rightarrow \mathcal{P}(L)$ by²²⁹

$$d(x, y) = \{\ell \in L \mid x \text{ and } y \text{ are not bisimilar when } \ell \text{ is chosen}\}.$$

Here is one last example further making the case for working over an abstract complete lattice. We also revisit it in Examples 3.4 and 3.83.

Example 2.17 (Hausdorff distance). Given an L-relation d on a set X , we define the L-relation d^\uparrow on non-empty finite subsets of X :

$$\forall S, T \in \mathcal{P}_{\text{ne}} X, \quad d^\uparrow(S, T) = \sup \left\{ \sup_{x \in S} \inf_{y \in T} d(x, y), \sup_{y \in T} \inf_{x \in S} d(x, y) \right\}.$$

²²⁶ The set P equipped with the function $R : P \rightarrow [0, 1]$ is not a $[0, 1]$ -space, but it is a *fuzzy set* in the sense of Castelnovo and Miculan [CM22a]. Their work shows how to reason with algebraic structures on fuzzy sets instead of L-spaces like we do here.

²²⁷ The rank of a Wikipedia page about [ramen](#) will be lower when the user inputs “[Genre Humaine](#)” than when they input “[Ramen_Lord](#)”.

²²⁸ There may be cases where $d'(h, k) = (4, 7)$ (a long chain with few authors) and $d'(h, k') = (2, 16)$ (a short chain with many authors). Then, with the product of complete lattices defined in Example 2.9, we could not compare the two distances. This is unfortunate in this application, so we may want to consider a different kind of product of complete lattices. The **lexicographical order** on $\mathbb{N}_\infty \times \mathbb{N}_\infty$ is

$$(\varepsilon, \delta) \leq_{\text{lex}} (\varepsilon', \delta') \Leftrightarrow \varepsilon \leq \varepsilon' \text{ or } (\varepsilon = \varepsilon' \text{ and } \delta \leq \delta').$$

In words, you use the order on the first coordinates, and only when they are equal, you use the order on the second coordinates.

If L and K are complete lattices, $(L \times K, \leq_{\text{lex}})$ is a complete lattice where the infimum is not computed pointwise, but rather

$$\inf S = (\inf \pi_L S, \sup \{\varepsilon \mid \forall s \in S, (\inf \pi_L S, \varepsilon) \leq s\}).$$

²²⁹ More details in [ABH⁺12, §Definitions C.1 and C.2].

This distance is a variation of a metric defined by Hausdorff in [Hau14].²³⁰ It measures how far apart two subsets are in three steps. First, we postulate that a point $x \in S$ and T are as far apart as x and the closest point $y \in T$. Then, the distance from S to T is as big as the distance between the point $x \in S$ furthest from T . Finally, to obtain a symmetric distance, we take the maximum of the distance from S to T and from T to S . As we expect from any interesting optimization problem, there is a dual formulation given by the L-relation d^\downarrow .²³¹

$$\forall S, T \in \mathcal{R}_{\text{re}}X, d^\downarrow(S, T) = \inf \left\{ \sup_{(x,y) \in C} d(x, y) \mid C \subseteq X \times X, \pi_1(C) = S, \pi_2(C) = T \right\}$$

To compare two sets with the second method, you first need a binary relation C on X that covers all and only the points of S and T in the first and second coordinates respectively. Borrowing the terminology from probability theory, we call C a **coupling** of S and T , it is a subset of $X \times X$ whose *marginals* are S and T . According to a coupling C , the distance between S and T is the biggest distance between a pair in C . Amongst all couplings of S and T , we take the one achieving the smallest distance to define $d^\downarrow(S, T)$.

The first punchline of this example is that the two L-relations d^\uparrow and d^\downarrow coincide.

Lemma 2.18. *For any $S, T \in \mathcal{R}_{\text{re}}X$, $d^\uparrow(S, T) = d^\downarrow(S, T)$.²³²*

Proof. (\leq) For any coupling $C \subseteq X \times X$, for each $x \in S$, there is at least one $y_x \in T$ such that $(x, y_x) \in C$ (because $\pi_1(C) = S$) so

$$\sup_{x \in S} \inf_{y \in T} d(x, y) \leq \sup_{x \in S} d(x, y_x) \leq \sup_{(x,y) \in C} d(x, y).$$

After a symmetric argument, we find that $d^\uparrow(S, T) \leq \sup_{(x,y) \in C} d(x, y)$ for all couplings, the first inequality follows.

(\geq) For any $x \in S$, let $y_x \in T$ be a point in T that attains the infimum of $d(x, y)$,²³³ and note that our definition ensures $d(x, y_x) \leq d^\uparrow(S, T)$. Symmetrically define x_y for any $y \in T$ and let $C = \{(x, y_x) \mid x \in S\} \cup \{(x_y, y) \mid y \in T\}$. It is clear that C is a coupling of S and T , and by our choices of y_x and x_y , we ensured that

$$\sup_{(x,y) \in C} d(x, y) \leq d^\uparrow(S, T),$$

therefore we found a coupling witnessing that $d^\downarrow(S, T) \leq d^\uparrow(S, T)$ as desired. \square

The second punchline of this example comes from instantiating it with the complete lattice \mathbf{B} . Recall that a B-relation d on X corresponds to a binary relation $R_d \subseteq X \times X$ where x and y are related if and only if $d(x, y) = \perp$. This seemingly backwards convention makes it so that nonexpansive functions are those that preserve the relation. Let us be careful about it while describing R_{d^\uparrow} and R_{d^\downarrow} .

Given $S, T \in \mathcal{R}_{\text{re}}X$ and $x \in S$, notice that $\inf_{y \in T} d(x, y) = \perp$ if and only if $d(x, y) = \perp$ for at least one y , or equivalently, if x is related by R_d to at least one $y \in T$. This means the infimum behaves like an existential quantifier. Dually, the supremum acts like a universal quantifier yielding²³⁴

²³⁰ Hausdorff considered positive real valued distances and compact subsets.

²³¹ The notation was inspired by [BBKK18]. We write $\pi_1(C)$ for $\{x \in X \mid \exists (x, y) \in C\}$ and similarly for π_2 . (We should really write $\mathcal{R}_{\text{re}}\pi_1(C)$ and $\mathcal{R}_{\text{re}}\pi_2(C)$.)

²³² Hardly adapted from [M 11, Proposition 2.1].

²³³ It exists because T is non-empty and finite.

²³⁴ Symmetrically,

$$\sup_{y \in T} \inf_{x \in S} d(x, y) = \perp \Leftrightarrow \forall y \in T, \exists x \in S, (x, y) \in R_d.$$

$$\sup_{x \in S} \inf_{y \in T} d(x, y) = \perp \iff \forall x \in S, \exists y \in T, (x, y) \in R_d.$$

Combining with its symmetric counterpart, and noting that a binary universal quantification is just an AND, we find that (S, T) belongs to R_{d^\uparrow} if and only if

$$\forall x \in S, \exists y \in T, (x, y) \in R_d \text{ and } \forall y \in T, \exists x \in S, (x, y) \in R_d. \quad (2.4)$$

We call R_{d^\uparrow} the Egli–Milner extension of R_d as in, e.g. [WS20, GPA21].

Given a coupling C of S and T , $\sup_{(x,y) \in C} d(x, y)$ can only equal \perp when all pairs $(x, y) \in C$ are related by R_d . Then, if a coupling $C \subseteq R_d$ exists, the infimum of d^\downarrow will be \perp . Therefore, S and T are related by R_{d^\downarrow} if and only if

$$\exists C \subseteq R_d, \pi_1(C) = S \text{ and } \pi_2(C) = T. \quad (2.5)$$

The relation R_{d^\downarrow} is sometimes called the Barr lifting of R_d [Baro6].

Our proof above yields the equivalence between (2.4) and (2.5).²³⁵

While the categories \mathbf{BSpa} , $[0, 1]\mathbf{Spa}$ and $[0, \infty]\mathbf{Spa}$ are interesting on their own, they contain subcategories which are more widely studied. For instance, the category \mathbf{Poset} of posets and monotone maps is a full subcategory of \mathbf{BSpa} where we only keep B-spaces (X, d) where the binary relation corresponding to d is reflexive, transitive and antisymmetric. Similarly, a $[0, \infty]$ -space (X, d) where the distance function satisfies the triangle inequality $d(x, z) \leq d(x, y) + d(y, z)$ and reflexivity $d(x, x) \leq 0$ is known as a Lawvere metric space [Lawo2].

The next section lays out the language we will use to state conditions as those above on L-spaces. The syntax is heavily inspired by the syntax of equations in universal algebra, the binary predicate $=$ for equality is joined by a family of binary predicates $=_\varepsilon$ indexed by the quantities in L. That idea comes from the original work of Mardare, Panangaden, and Plotkin on quantitative algebras [MPP16], and it implicitly relies on the following equivalent definition of L-spaces (the equivalent definition is not due to Mardare et al., see the discussion in §0.3).

Definition 2.19 (L-structure). Given a complete lattice L, an L-structure²³⁶ is a set X equipped with a family of binary relations $R_\varepsilon \subseteq X \times X$ indexed by $\varepsilon \in L$ satisfying

- **monotonicity** in the sense that if $\varepsilon \leq \varepsilon'$, then $R_\varepsilon \subseteq R_{\varepsilon'}$, and
- **continuity** in the sense that for any I -indexed family of elements $\varepsilon_i \in L$,²³⁷

$$\bigcap_{i \in I} R_{\varepsilon_i} = R_\delta, \text{ where } \delta = \inf_{i \in I} \varepsilon_i.$$

Intuitively $(x, y) \in R_\varepsilon$ should be interpreted as bounding the distance from x to y above by ε . Then, monotonicity means the points that are at a distance below ε are also at a distance below ε' when $\varepsilon \leq \varepsilon'$. Continuity means the points that are at a distance below a bunch of bounds ε_i are also at a distance below the infimum of those bounds $\inf_{i \in I} \varepsilon_i$.

The names for these conditions come from yet another equivalent definition.²³⁸

²³⁵ That equivalence is folklore and has probably been given as exercise to many students in a class on bisimulation or coalgebras.

²³⁶ We borrow the name “structure” from model theorists. The more general notion of relational structure is used in [FMS21, Par22, Par23]. Also, our L-structures are both more and less general than the \mathcal{L}_S -structures of [Con17].

²³⁷ By monotonicity, $R_\delta \subseteq R_{\varepsilon_i}$ so the inclusion $R_\delta \subseteq \bigcap_{i \in I} R_{\varepsilon_i}$ always holds. Also, continuity implies monotonicity because $\varepsilon \leq \varepsilon'$ implies

$$R_\varepsilon \cap R_{\varepsilon'} = R_{\inf\{\varepsilon, \varepsilon'\}} = R_\varepsilon,$$

which means $R_\varepsilon \subseteq R_{\varepsilon'}$. Still, we keep monotonicity explicit for better exposition.

²³⁸ This time more directly equivalent.

Organizing the data of an L-structure into a function $R : L \rightarrow \mathcal{P}(X \times X)$ sending ε to R_ε , we can recover monotonicity and continuity by seeing $\mathcal{P}(X \times X)$ as a complete lattice like in Example 2.7. Indeed, monotonicity is equivalent to R being a monotone function between the posets (L, \leq) and $(\mathcal{P}(X \times X), \subseteq)$, and continuity is equivalent to R preserving infimums. Seeing L and $\mathcal{P}(X \times X)$ as posetal categories, we can simply say that R is a continuous functor.²³⁹

A morphism between two L-structures $(X, \{R_\varepsilon\})$ and $(Y, \{S_\varepsilon\})$ is a function $f : X \rightarrow Y$ satisfying

$$\forall \varepsilon \in L, \forall x, x' \in X, (x, x') \in R_\varepsilon \implies (f(x), f(x')) \in S_\varepsilon. \quad (2.6)$$

This should feel similar to nonexpansive maps.²⁴⁰ Let us call **LStr** the category of L-structures.

We give one trivial example, before proving that L-structures are just L-spaces.

Example 2.20. A consequence of continuity (take $I = \emptyset$) is that R_\top is the full binary relation $X \times X$. Therefore, taking $L = 1$ to be a singleton where $\perp = \top$, a 1-structure is only a set (there is no choice for R), and a morphism is only a function (the implication in (2.6) is always true because $S_\varepsilon = Y \times Y$). In other words, **1Str** is isomorphic to **Set**. Instantiating the next result (Proposition 2.21) means that **1Spa** is also isomorphic to **Set**, this is clear because there is only one function $d : X \times X \rightarrow 1$ for any set X . This example is relatively important because it means the theory we develop later over an arbitrary category of L-spaces specializes to the case of **Set**.²⁴¹

Proposition 2.21. *For any complete lattice L , the categories **LSpa** and **LStr** are isomorphic.*²⁴²

Proof. Given an L-relation (X, d) , we define the binary relations $R_\varepsilon^d \subseteq X \times X$ by

$$(x, x') \in R_\varepsilon^d \iff d(x, x') \leq \varepsilon. \quad (2.7)$$

This family satisfies monotonicity because for any $\varepsilon \leq \varepsilon'$ we have

$$(x, x') \in R_\varepsilon^d \xrightarrow{(2.7)} d(x, x') \leq \varepsilon \implies d(x, x') \leq \varepsilon' \xrightarrow{(2.7)} (x, x') \in R_{\varepsilon'}^d.$$

It also satisfies continuity because if $(x, x') \in R_{\varepsilon_i}$ for all $i \in I$, then $d(x, x') \leq \varepsilon_i$ for all $i \in I$. By definition of infimum, we must have $d(x, x') \leq \inf_{i \in I} \varepsilon_i$, hence $(x, x') \in R_{\inf_{i \in I} \varepsilon_i}$. We conclude the forward inclusion (\subseteq) of continuity holds, the converse (\supseteq) follows from monotonicity.

Any nonexpansive map $f : (X, d) \rightarrow (Y, \Delta)$ in **LSpa** is also a morphism between the L-structures $(X, \{R_\varepsilon^d\})$ and $(Y, \{R_\varepsilon^\Delta\})$ because for all $\varepsilon \in L$ and $x, x' \in X$,

$$(x, x') \in R_\varepsilon^d \xrightarrow{(2.7)} d(x, x') \leq \varepsilon \xrightarrow{(2.3)} \Delta(f(x), f(x')) \leq \varepsilon \xrightarrow{(2.7)} (f(x), f(x')) \in R_\varepsilon^\Delta.$$

It follows that the assignment $(X, d) \mapsto (X, \{R_\varepsilon^d\})$ is a functor $F : \mathbf{LSpa} \rightarrow \mathbf{LStr}$ acting trivially on morphisms.

Given an L-structure $(X, \{R_\varepsilon\})$, we define the function $d_R : X \times X \rightarrow L$ by

$$d_R(x, x') = \inf \{ \varepsilon \in L \mid (x, x') \in R_\varepsilon \}.$$

²³⁹ Limits in a posetal category are always computed by taking the infimum of all the points in the diagram, so preserving limits and preserving infimums is the same thing.

²⁴⁰ In words, (2.6) reads as: if x and x' are at a distance below ε' then so are $f(x)$ and $f(x')$.

²⁴¹ See Example 3.70.

²⁴² This result is a stripped down version of [MPP17, Theorem 4.3]. A more general version also appears in [FMS21, Example 3.5.(4)]. Another similar result is shown in [Par22, Appendix]. The core idea, (2.7) and (2.8), also appears in [Con17, Theorem A] and [LR17, Example 4.5.(3)].

Taking $L = \mathbf{B}$, Proposition 2.21 gives back our interpretation of **BSpa** as the category **2Rel** from Example 2.13. Indeed, a B-structure is just a set X equipped with a binary relation $R_\perp \subseteq X \times X$ (because R_\top is required to equal $X \times X$), and morphisms of B-structures are functions that preserve that binary relation. This also justifies our weird choice of $d(x, y) = \perp$ meaning x and y are related.

Note that monotonicity and continuity of the family $\{R_\varepsilon\}$ imply²⁴³

$$d_R(x, x') \leq \varepsilon \iff (x, x') \in R_\varepsilon. \quad (2.8)$$

This allows us to prove that a morphism $f : (X, \{R_\varepsilon\}) \rightarrow (Y, \{S_\varepsilon\})$ is nonexpansive from (X, d_R) to (Y, d_S) because for all $\varepsilon \in \mathbf{L}$ and $x, x' \in X$, we have

$$d_R(x, x') \leq \varepsilon \stackrel{(2.8)}{\iff} (x, x') \in R_\varepsilon \stackrel{(2.6)}{\implies} (f(x), f(x')) \in S_\varepsilon \stackrel{(2.8)}{\iff} d_S(f(x), f(x')) \leq \varepsilon,$$

hence putting $\varepsilon = d_R(x, x')$, we obtain $d_S(f(x), f(x')) \leq d_R(x, x')$. It follows that the assignment $(X, \{R_\varepsilon\}) \mapsto (X, d_R)$ is a functor $G : \mathbf{LStr} \rightarrow \mathbf{LSpa}$ acting trivially on morphisms.

Observe that (2.7) and (2.8) together say that $R_\varepsilon^{d_R} = R_\varepsilon$ and $d_{R^d} = d$, so F and G are inverses to each other on objects. Since both functors do nothing to morphisms, we conclude that F and G are inverses to each other, and that $\mathbf{LSpa} \cong \mathbf{LStr}$. \square

This result is central in our treatment of L-spaces because it allows us to specify an L-relation through the (binary) truth value of a family of predicates $=_\varepsilon$. In other words, we can reason equationally about L-spaces.

Remark 2.22. The upshot of Proposition 2.21 is that the structure of a complete lattice is enough to do quantitative algebraic reasoning.²⁴⁴ Still, in practice, \mathbf{L} often has more structure. If you need to state the triangle inequality (2.12), then you need a way of adding distances/quantities. A frequent choice made by researchers is to let \mathbf{L} be a quantale see e.g. [CH06, Pis21]. Often, this is for the theoretical convenience of seeing a metric space as an enriched category as suggested in [Law02].²⁴⁵ In closely related work [CM22a], Castelnovo and Miculan require \mathbf{L} to be a frame (a special kind of quantale).

2.2 Equational Constraints

It is often the case that one wants to impose conditions on the L-spaces they consider. For instance, recall that when \mathbf{L} is $[0, 1]$ or $[0, \infty]$, L-spaces are sets with a notion of distance between points. Starting from our intuition on the distance between points of the space we live in, people have come up with several abstract conditions to enforce on distance functions. For example, we can restate (with a slight modification²⁴⁶) the axioms defining metric spaces (Definition 0.1).

First, symmetry says that the distance from x to y is the same as the distance from y to x :

$$\forall x, y \in X, \quad d(x, y) = d(y, x). \quad (2.9)$$

Reflexivity, also called indiscernibility of identicals, says that the distance between x and itself is 0 (i.e. the smallest distance possible):

$$\forall x \in X, \quad d(x, x) = 0. \quad (2.10)$$

Identity of indiscernibles, also called Leibniz's law, says that if two points x and y are at distance 0, then x and y must be the same:

$$\forall x, y \in X, \quad d(x, y) = 0 \implies x = y. \quad (2.11)$$

²⁴³ The converse implication (\Leftarrow) is by definition of infimum. For (\Rightarrow), continuity says that

$$R_{d_R(x, x')} = \bigcap_{\varepsilon \in \mathbf{L}, (x, x') \in R_\varepsilon} R_\varepsilon,$$

so $R_{d_R(x, x')}$ contains (x, x') , then by monotonicity, $d_R(x, x') \leq \varepsilon$ implies R_ε also contains (x, x') .

²⁴⁴ This point will be strengthened when we develop the theory of quantitative algebras over an arbitrary complete lattice in Chapter 3.

²⁴⁵ The book [HST14] explores the theoretical foundations of this approach.

²⁴⁶ The separation axiom is now divided in two, (2.10) and (2.11).

Finally, the triangle inequality says that the distance from x to z is always smaller than the sum of the distances from x to y and from y to z :

$$\forall x, y, z \in X, \quad d(x, z) \leq d(x, y) + d(y, z). \quad (2.12)$$

There are also very famous axioms on B-spaces (X, d) that arise from viewing the binary relation corresponding to d as some kind of order on elements of X . They are abstraction of properties of the “smaller or equal” order \leq on, say, real numbers.

First, reflexivity says that any element x is related to itself, i.e. $x \leq x$. Translating back to the B-relation, this is equivalent to:

$$\forall x \in X, \quad d(x, x) = \perp. \quad (2.13)$$

Antisymmetry says that if both (x, y) and (y, x) are in the order relation, then they must be equal:²⁴⁷

$$\forall x, y \in X, \quad d(x, y) = \perp = d(y, x) \implies x = y. \quad (2.14)$$

²⁴⁷ i.e. if $x \leq y$ and $y \leq x$, then $x = y$.

Finally, transitivity says that if (x, y) and (y, z) belong to the order relation, then so does (x, z) :²⁴⁸

$$\forall x, y, z \in X, \quad d(x, y) = \perp = d(y, z) \implies d(x, z) = \perp. \quad (2.15)$$

²⁴⁸ i.e. if $x \leq y$ and $y \leq z$, then $x \leq z$.

We can immediately notice that all the axioms (2.9)–(2.15) start with a universal quantification of variables. Another thing to note is that we never actually needed to talk about equality between distances. For instance, the equation $d(x, y) = d(y, x)$ in the axiom of symmetry (2.9) can be replaced by two inequalities $d(x, y) \leq d(y, x)$ and $d(y, x) \leq d(x, y)$, and moreover since x and y are universally quantified, only one of these inequalities is necessary:

$$\forall x, y \in X, \quad d(x, y) \leq d(y, x). \quad (2.16)$$

If we rely on the equivalence between L-spaces and L-structures (Proposition 2.21), we can transform (2.16) into a family of implications indexed by all $\varepsilon \in L$:²⁴⁹

$$\forall x, y \in X, \quad (y, x) \in R_\varepsilon^d \implies (x, y) \in R_\varepsilon^d. \quad (2.17)$$

²⁴⁹ Recall that $(x, y) \in R_\varepsilon^d$ is the same thing as $d(x, y) \leq \varepsilon$. Hence, (2.16) and (2.17) are equivalent because requiring $d(x, y)$ to be smaller than $d(y, x)$ is equivalent to requiring all upper bounds of $d(y, x)$ (in particular $d(y, x)$ itself) to also be upper bounds of $d(x, y)$.

Starting from the triangle inequality (2.12) and applying the same transformations that got us from (2.9) to (2.17), we obtain a family of implications indexed by two quantities $\varepsilon, \delta \in L$:²⁵⁰

$$\forall x, y, z \in X, \quad (x, y) \in R_\varepsilon^d \text{ and } (y, z) \in R_\delta^d \implies (x, z) \in R_{\varepsilon+\delta}^d. \quad (2.18)$$

²⁵⁰ You can try proving how (2.12) and (2.18) are equivalent if the process of going from the former to the latter was not clear to you.

The last conceptual step is to make the L.H.S. of the implication part of the universal quantification. That is, instead of saying “for all x and y , if P then Q ”, we say “for all x and y such that P , Q ”. We do this by introducing a syntax very similar to the equations of universal algebra. We fix a complete lattice (L, \leq) , but you can keep in mind the examples $L = [0, 1]$ and $L = [0, \infty]$.

Definition 2.23 (Quantitative equation).²⁵¹ A **quantitative equation** (over L) is a tuple comprising an L -space \mathbf{X} called the **context**, two elements $x, y \in X$, and optionally a quantity $\varepsilon \in L$. We write these as $\mathbf{X} \vdash x = y$ when no ε is given or $\mathbf{X} \vdash x =_\varepsilon y$ when it is given.

An L -space \mathbf{A} **satisfies** a quantitative equation

- $\mathbf{X} \vdash x = y$ if for any nonexpansive assignment $\hat{I} : \mathbf{X} \rightarrow \mathbf{A}$, $\hat{I}(x) = \hat{I}(y)$.
- $\mathbf{X} \vdash x =_\varepsilon y$ if for any nonexpansive assignment $\hat{I} : \mathbf{X} \rightarrow \mathbf{A}$, $d_{\mathbf{A}}(\hat{I}(x), \hat{I}(y)) \leq \varepsilon$.

We use ϕ and ψ to refer to a quantitative equation, and we sometimes call them simply equations. We write $\mathbf{A} \models \phi$ when \mathbf{A} satisfies ϕ ,²⁵² and we also write $\mathbf{A} \models^{\hat{I}} \phi$ when the equality $\hat{I}(x) = \hat{I}(y)$ or the bound $d_{\mathbf{A}}(\hat{I}(x), \hat{I}(y)) \leq \varepsilon$ holds for a particular assignment $\hat{I} : \mathbf{X} \rightarrow \mathbf{A}$ (and not necessarily for all assignments).

Remark 2.24. The authors of [MPP16] introduced the symbol $=_\varepsilon$ to represent “equality up to ε ”, and it is at the basis of their theory of algebras over metric spaces. Just like $=$ is understood as equality in any set, we understand $=_\varepsilon$ as the relation R_ε in any L -structure. Therefore, under the assignment \hat{I} inside \mathbf{A} , $x =_\varepsilon y$ becomes $\hat{I}(x) R_\varepsilon^{\hat{I}} \hat{I}(y)$, which in turn means $d_{\mathbf{A}}(\hat{I}(x), \hat{I}(y)) \leq \varepsilon$. Unlike $=$, $=_\varepsilon$ is not a symmetric notation, but that is convenient for us as arbitrary L -spaces are not required to satisfy $d_{\mathbf{A}}(x, y) = d_{\mathbf{A}}(y, x)$.

Let us illustrate the definition of quantitative equations with an example.

Example 2.25 (Symmetry). We want to translate (2.17) into a quantitative equation. A first approximation would be replacing the relation R_ε^d with our new syntax $=_\varepsilon$ to obtain something like

$$x, y \vdash y =_\varepsilon x \implies x =_\varepsilon y.$$

We are not allowed to use implications like this, so we have to implement the last step mentioned above by putting the premise $y =_\varepsilon x$ into the context. This means we need to quantify over variables x and y with a bound ε on the distance from y to x .

Note that when defining satisfaction of a quantitative equation, the quantification happens at the level of assignments $\hat{I} : \mathbf{X} \rightarrow \mathbf{A}$. Hence, we have to find a context \mathbf{X} such that nonexpansive assignments $\mathbf{X} \rightarrow \mathbf{A}$ correspond to choices of two elements in \mathbf{A} with the same bound ε on their distance.

Let the context \mathbf{X} be the L -space with two elements x and y such that $d_{\mathbf{X}}(y, x) = \varepsilon$ and all other distances are \top . A nonexpansive assignment $\hat{I} : \mathbf{X} \rightarrow \mathbf{A}$ is just a choice of two elements $\hat{I}(x), \hat{I}(y) \in \mathbf{A}$ satisfying $d_{\mathbf{A}}(\hat{I}(y), \hat{I}(x)) \leq \varepsilon$.²⁵³ For all of these, we have to impose the condition $d_{\mathbf{A}}(\hat{I}(x), \hat{I}(y)) \leq \varepsilon$. Therefore, our quantitative equation is

$$\mathbf{X} \vdash x =_\varepsilon y. \tag{2.19}$$

For a fixed quantity $\varepsilon \in L$, an L -space \mathbf{A} satisfies (2.19) if and only if it satisfies (2.17). Hence,²⁵⁴ if \mathbf{A} satisfies that quantitative equation for all $\varepsilon \in L$, then it satisfies (2.9), i.e. the distance $d_{\mathbf{A}}$ is symmetric.

²⁵¹ The name quantitative equation will be reclaimed in Definition 3.8 for a more general notion. See also Remark 3.9.

²⁵² Of course, satisfaction generalizes straightforwardly to sets of quantitative equations, i.e. if \hat{E} is a class of quantitative equations, $\mathbf{A} \models \hat{E}$ means $\mathbf{A} \models \phi$ for all $\phi \in \hat{E}$.

²⁵³ Indeed, since \top is the top element of L , the other values of $d_{\mathbf{X}}$ being \top means that they impose no further condition on $d_{\mathbf{A}}$.

²⁵⁴ Recall our argument in Footnote 249.

In practice, defining the context like this is more cumbersome than need be, so we will define some syntactic sugar to remedy this. Before that, we take the time to do another example.

Example 2.26 (Triangle inequality). With $L = [0, 1]$ or $L = [0, \infty]$, let the context \mathbf{X} be the L -space with three elements x, y and z such that $d_{\mathbf{X}}(x, y) = \varepsilon$ and $d_{\mathbf{X}}(y, z) = \delta$, and all other distances are \top .²⁵⁵ A nonexpansive assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$ is just a choice of three elements $a = \hat{\iota}(x), b = \hat{\iota}(y), c = \hat{\iota}(z) \in A$ such that $d_{\mathbf{A}}(a, b) \leq \varepsilon$ and $d_{\mathbf{A}}(b, c) \leq \delta$. Hence, if \mathbf{A} satisfies

$$\mathbf{X} \vdash x =_{\varepsilon + \delta} z, \quad (2.20)$$

it means that for any such assignment, $d_{\mathbf{A}}(a, c) \leq \varepsilon + \delta$ also holds. We conclude that \mathbf{A} satisfies (2.18). If \mathbf{A} satisfies $\mathbf{X} \vdash x =_{\varepsilon + \delta} z$ for all $\varepsilon, \delta \in L$, then \mathbf{A} satisfies the triangle inequality (2.12).

Remark 2.27. There is a small caveat above. If we are in $L = [0, 1]$ and $\varepsilon = 1$ and $\delta = 1$, then $\varepsilon + \delta = 2 \notin [0, 1]$, so the predicate $x =_{\varepsilon + \delta} z$ is not allowed. There are two easy fixes that we never explicit. You can either define a truncated addition so that $\varepsilon + \delta = 1$ whenever their sum is really above 1,²⁵⁶ or you can quantify over ε and δ such that $\varepsilon + \delta \leq 1$. Indeed, every $[0, 1]$ -space satisfies $\mathbf{X} \vdash x =_1 z$ because 1 is a global upper bound for the distance between points, thus when $\varepsilon + \delta > 1$, there is no difference between having that equation or not as an axiom.

Notice that in the contexts of Examples 2.25 and 2.26, we only needed to set one or two distances and all the others where the maximum they could be \top . In our **syntactic sugar** for quantitative equations, we will only write the distances that are important (using the syntax $=_{\varepsilon}$), and we understand the underspecified distances to be as high as they can be. For instance, (2.19) will be written²⁵⁷

$$y =_{\varepsilon} x \vdash x =_{\varepsilon} y, \quad (2.21)$$

and (2.20) will be written

$$x =_{\varepsilon} y, y =_{\delta} z \vdash x =_{\varepsilon + \delta} z. \quad (2.22)$$

In this syntax, we call **premises** everything on the left of the turnstile \vdash and **conclusion** what is on the right.


More generally, when we write $\{x_i =_{\varepsilon_i} y_i\}_{i \in I} \vdash x =_{\varepsilon} y$ (resp. $\{x_i =_{\varepsilon_i} y_i\}_{i \in I} \vdash x = y$), it corresponds to the quantitative equation $\mathbf{X} \vdash x =_{\varepsilon} y$ (resp. $\mathbf{X} \vdash x = y$), where the context \mathbf{X} contains the variables in²⁵⁸

$$X = \{x, y\} \cup \{x_i \mid i \in I\} \cup \{y_i \mid i \in I\},$$

and the L -relation is defined for $u, v \in X$ by²⁵⁹

$$d_{\mathbf{X}}(u, v) = \inf\{\varepsilon \mid u =_{\varepsilon} v \in \{x_i =_{\varepsilon_i} y_i\}_{i \in I}\}.$$

²⁵⁵ Here is a depiction of \mathbf{X} , where the label on an arrow is the distance from the source to the target of that arrow:



²⁵⁶ This operation is well-known in fuzzy logic, under different names like *bounded sum*, *strong disjunction*, or *t-conorm*, see, e.g. [CHN11, Chapter 1, §2.2.2].

²⁵⁷ We can understand this syntax as putting back the information in the context into an implication. For instance, you can read (2.21) as “if the distance from y to x is bounded above by ε , then so is the distance from x to y ”. You can read (2.22) as “if the distance from y to x is bounded above by ε and the distance from y to z is bounded above by δ , then the distance from x to z is bounded above by $\varepsilon + \delta$ ”.

²⁵⁸ Note that the x_i s, y_i s, x and y need not be distinct. In fact, x and y almost always appear in the x_i s and y_i s.

²⁵⁹ In words, the distance from u to v is the smallest value ε such that $u =_{\varepsilon} v$ was a premise. If no such premise occurs, the distance from u to v is \top . It is rare that u and v appear several times together (because $u =_{\varepsilon} v$ and $u =_{\delta} v$ can be replaced with $u =_{\inf\{\varepsilon, \delta\}} v$), but our definition allows it.

Remark 2.28. The judgments (or quantitative inferences) in the logic of [MPP16] with an empty signature coincide with our syntactic sugar. We showed those are a formally equivalent to quantitative equations in [MSV23, Lemma 8.4], but there is a special case we want to discuss.

In [MPP16, Definition 2.1], their axiom (Arch) is equivalent, in the presence of their axiom (Max), to²⁶⁰

$$\{x =_{\varepsilon_i} y \mid i \in I\} \vdash x =_{\inf_{i \in I} \varepsilon_i} y.$$

Now, if we apply our translation to obtain a quantitative equation as in Definition 2.23, we get $\mathbf{X} \vdash x =_{\varepsilon} y$, where $d_{\mathbf{X}}(x, y) = \varepsilon = \inf_{i \in I} \varepsilon_i$ and all other distances are \top . This quantitative equation is obviously always satisfied,²⁶¹ so it makes sense to have it as an axiom, but it seems we are loosing a bit of information. That is, the original axiom looks like it ensures the continuity property of Definition 2.19. In fact, that axiom has several names in different papers, one of which is `CONT`. In the version of quantitative equational logic we propose in this thesis (Figure 3.1), there is an inference rule `CONT` (rather than an axiom) that ensures continuity.

Here are some more translations of famous properties into quantitative equations written with the syntactic sugar:

- reflexivity (of a metric) (2.10) becomes $x \vdash x =_0 x$,²⁶²
- Leibniz's law (2.11) becomes $x =_0 y \vdash x = y$,
- reflexivity (of an order) (2.13) becomes $x \vdash x =_{\perp} x$,
- antisymmetry (2.14) becomes $x =_{\perp} y, y =_{\perp} x \vdash x = y$, and
- transitivity (2.15) becomes $x =_{\perp} y, y =_{\perp} z \vdash x =_{\perp} z$.

Remark 2.29. The translations of (2.10) and (2.13) look very close. In fact, noting that 0 is the bottom element of $[0, 1]$ and $[0, \infty]$, the quantitative equation $x \vdash x =_{\perp} x$ can state the reflexivity of a distance in $[0, 1]$ or $[0, \infty]$ or the reflexivity of a binary relation.

Similarly, in the translation of the triangle inequality (2.22), if we let ε and δ range over \mathbf{B} and interpret $+$ as an OR, we get three vacuous quantitative equations,²⁶³ and the translation of (2.15) above. So transitivity and triangle inequality are the same under this abstract point of view.²⁶⁴

Let us emphasize one thing about contexts of quantitative equations: they only give constraints that are upper bounds for distances.²⁶⁵ In particular, it can be very hard to operate on the quantities in \mathbf{L} non-monotonically. For instance, we will see (after Definition 2.40) that we cannot read $x =_{\varepsilon_1} y, y =_{\varepsilon_2} z, y =_{\varepsilon_3} y \vdash x =_{\varepsilon_1 + \varepsilon_2 - \varepsilon_3} z$ as saying that $d(x, z) \leq d(x, y) + d(y, z) - d(y, y)$,²⁶⁶ and one quick explanation is that subtraction is not a monotone operation on $[0, \infty] \times [0, \infty]$. Another consequence is that an equation ϕ will always entail ψ when the latter has a *stricter* context (i.e. when the upper-bounds in the premises are smaller).²⁶⁷ We prove a more general version of this below.

²⁶⁰ See [MSV23, Remark 4.3].

²⁶¹ For any nonexpansive assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$, $d_{\mathbf{A}}(\hat{\iota}(x), \hat{\iota}(y)) \leq d_{\mathbf{X}}(x, y) = \varepsilon$.

²⁶² As further sugar, we also write x instead of $x =_{\top} x$ to the left of the turnstile \vdash to say that the variable x is in the context without imposing any constraint. For instance, the context of $x, y \vdash x = y$ has two variables x and y and all distances are \top . Thus, if \mathbf{A} satisfies $x, y \vdash x = y$, then \mathbf{A} is either empty or a singleton.

²⁶³ When either ε or δ equals \top , $\varepsilon + \delta = \top$, but when the conclusion of a quantitative equation is $x =_{\top} z$, it is always satisfied.

²⁶⁴ These observations were probably folkloric since at least the original publication of [Lawo2] in 1973.

²⁶⁵ Well, if you consider the opposite order on \mathbf{L} , they now give lower bounds. What is important is that they only speak about one of them.

²⁶⁶ Assume $\mathbf{L} = [0, \infty]$ and $d(y, y)$ may be non-zero.

²⁶⁷ For example, if \mathbf{A} satisfies $x =_{1/2} y \vdash x = y$, then it satisfies $x =_{1/3} y \vdash x = y$. This says that if all distances between distinct points are above $1/2$, then they are also above $1/3$.

Lemma 2.30. *Let $f : \mathbf{X} \rightarrow \mathbf{Y}$ be a nonexpansive map. If \mathbf{A} satisfies $\mathbf{X} \vdash x = y$ (resp. $\mathbf{X} \vdash x =_\varepsilon y$), then \mathbf{A} satisfies $\mathbf{Y} \vdash f(x) = f(y)$ (resp. $\mathbf{Y} \vdash f(x) =_\varepsilon f(y)$).²⁶⁸*

Proof. Any nonexpansive assignment $\hat{l} : \mathbf{Y} \rightarrow \mathbf{A}$ yields a nonexpansive assignment $\hat{l} \circ f : \mathbf{X} \rightarrow \mathbf{A}$. By hypothesis, we have

$$\mathbf{A} \models^{\hat{l} \circ f} \mathbf{X} \vdash x = y \quad (\text{resp. } \mathbf{A} \models^{\hat{l} \circ f} \mathbf{X} \vdash x =_\varepsilon y),$$

which means $\hat{l}(f(x)) = \hat{l}(f(y))$ (resp. $d_{\mathbf{A}}(\hat{l}(f(x)), \hat{l}(f(y))) \leq \varepsilon$). Thus, we conclude

$$\mathbf{A} \models^{\hat{l}} \mathbf{Y} \vdash f(x) = f(y) \quad (\text{resp. } \mathbf{A} \models^{\hat{l}} \mathbf{Y} \vdash f(x) =_\varepsilon f(y)). \quad \square$$

Let us continue this list of examples for a while, just in case it helps a reader that is looking to translate an axiom into a quantitative equation. We will also give some results later which could imply that a reader's axiom cannot be translated in this language.

Example 2.31. For any complete lattice \mathbf{L} .

1. The **strong triangle inequality** states that $d(x, z) \leq \max\{d(x, y), d(y, z)\}$,²⁶⁹ it is equivalent to the satisfaction of the following family of quantitative equations

$$\forall \varepsilon, \delta \in \mathbf{L}, \quad x =_\varepsilon y, y =_\delta z \vdash x =_{\sup\{\varepsilon, \delta\}} z. \quad (2.23)$$

2. We can impose that all distances are below a **global upper bound** $\varepsilon \in \mathbf{L}$ (i.e. $d(x, y) \leq \varepsilon$) with the quantitative equation²⁷⁰

$$x, y \vdash x =_\varepsilon y. \quad (2.24)$$

3. We can *almost* impose a **global lower bound** $\varepsilon \in \mathbf{L}$ on distances. What we can do instead is impose a strict lower bound on distances that are not self-distances (i.e. $\forall x \neq y, d(x, y) > \varepsilon$). To achieve this with an equation, we ensure the equivalent property that whenever $d(x, y)$ is smaller than ε , then $x = y$:²⁷¹

$$x =_\varepsilon y \vdash x = y. \quad (2.25)$$

Let $\mathbf{L} = [0, 1]$ or $\mathbf{L} = [0, \infty]$.

1. Given a positive number $b > 0$, the **b -triangle inequality** states that $d(x, z) \leq b(d(x, y) + d(y, z))$,²⁷² it is equivalent to the satisfaction of

$$\forall \varepsilon, \delta \in \mathbf{L}, \quad x =_\varepsilon y, y =_\delta z \vdash x =_{b(\varepsilon + \delta)} z. \quad (2.26)$$

2. The **rectangle inequality** states that $d(x, w) \leq d(x, y) + d(y, z) + d(z, w)$,²⁷³ it is equivalent to the satisfaction of

$$\forall \varepsilon_1, \varepsilon_2 \in \mathbf{L}, \quad x =_{\varepsilon_1} y, y =_{\varepsilon_2} z, z =_{\varepsilon_3} w \vdash x =_{\varepsilon_1 + \varepsilon_2 + \varepsilon_3} w. \quad (2.27)$$

Let $\mathbf{L} = \mathbf{B}$.

²⁶⁸ In particular, when written with syntactic sugar, you can reduce the quantities in the premises of

$$\{x =_{\varepsilon_i} y \mid i \in I\} \vdash x =_{\inf_{i \in I} \varepsilon_i} y$$

to obtain

$$\{x =_{\varepsilon'_i} y \mid i \in I\} \vdash x =_{\inf_{i \in I} \varepsilon'_i} y,$$

with $\varepsilon'_i \leq \varepsilon$. If \mathbf{X} and \mathbf{X}' respectively denote the context described by these sets of premises, then they have the same carrier, and the identity function will be nonexpansive from \mathbf{X} to \mathbf{X}' . Therefore, if \mathbf{A} satisfies the first quantitative equation, it satisfies the second.

²⁶⁹ This property is used in defining ultrametrics [Rut96].

²⁷⁰ For instance $[0, 1]$ -spaces are $[0, \infty]$ -spaces that satisfy $x, y \vdash x =_1 y$.

²⁷¹ We can also do a non-strict lower bound (i.e. $\forall x \neq y, d(x, y) \geq \varepsilon$) by considering the family of equations $x =_\delta y \vdash x = y$ for all $\delta < \varepsilon$.

²⁷² This property is used in defining b -metrics [KP22, Definition 1.1].

²⁷³ This property is used in defining g.m.s. in [Braoo, Definition 1.1].

1. A binary relation R on $X \times X$ is said to be **functional** if there are no two distinct $y, y' \in X$ such that $(x, y) \in R$ and $(x, y') \in R$ for a single $x \in X$. This is equivalent to satisfying

$$x =_{\perp} y, x =_{\perp} y' \vdash y = y'. \quad (2.28)$$

2. We say $R \subseteq X \times X$ is **injective** if there are no two distinct $x, x' \in X$ such that $(x, y) \in R$ and $(x', y) \in R$ for a single $y \in X$.²⁷⁴ This is equivalent to satisfying

$$x =_{\perp} y, x' =_{\perp} y \vdash x = x'. \quad (2.29)$$

3. We say $R \subseteq X \times X$ is **circular** if whenever (x, y) and (y, z) belong to R , then so does (z, x) (compare with transitivity (2.15)). This is equivalent to satisfying

$$x =_{\perp} y, y =_{\perp} z \vdash z =_{\perp} x. \quad (2.30)$$

We now turn to the study of subcategories of **LSpa** that are defined via quantitative equations. Given a class \hat{E} of quantitative equations, we can define a full subcategory of **LSpa** that contains only those L-spaces that satisfy \hat{E} , this is the category $\mathbf{GMet}(L, \hat{E})$ whose objects we call generalized metric spaces or spaces for short. We also write $\mathbf{GMet}(\hat{E})$ or \mathbf{GMet} when the complete lattices L or the class \hat{E} are fixed or irrelevant. There is an evident forgetful functor $U : \mathbf{GMet} \rightarrow \mathbf{Set}$ which is the composition of the inclusion functor $\mathbf{GMet} \rightarrow \mathbf{LSpa}$ and $U : \mathbf{LSpa} \rightarrow \mathbf{Set}$.²⁷⁵

The terminology generalized metric space appears quite a lot in the literature with different meanings (e.g. [BvBR98, Bra00, Pis21]), so I expect many will navigate to this definition before reading what is above. Catering to these readers, let us redefine what we mean by generalized metric space.

Definition 2.32 (Generalized metric space). A **generalized metric space** or **space** is a set X along with a function $d : X \times X \rightarrow L$ into a complete lattice L such that (X, d) satisfies some constraints expressed by a fixed collection of quantitative equations.

When $L = [0, \infty]$, examples include metrics [Fré06], ultrametrics [Rut96], pseudometrics, quasimetrics [Wil31a], semimetrics [Wil31b], b -metrics [KP22], the generalized metric spaces of [Bra00], dislocated metrics [HS00] also called diffuse metrics in [CKPR21], the generalized metric spaces of [BvBR98] which are the metric spaces of [Law02], etc.

When $L = B$ (the Boolean lattice), examples include posets, preorders, equivalence relations, partial (or restricted) equivalence relations [Sco76], graphs, etc.

The most notable examples of generalized metric spaces are posets and metric spaces, they form the categories **Poset** and **Met**.

Example 2.33 (Poset). The category of partially ordered sets and monotone maps is the full subcategory of **BSpa** with all B-spaces satisfying reflexivity, antisymmetry, and transitivity stated as quantitative equations:²⁷⁶

$$\hat{E}_{\mathbf{Poset}} = \{x \vdash x =_{\perp} x, x =_{\perp} y, y =_{\perp} x \vdash x = y, x =_{\perp} y, y =_{\perp} z \vdash x =_{\perp} z\}.$$

²⁷⁴ Equivalently, the opposite (or converse) of R is functional. You may want to formulate totality or surjectivity of a binary relation with quantitative equations, but you will find that difficult. We show in Example 2.47 that it is not possible.

²⁷⁵ Recall that while we use the same symbol for both forgetful functors, you can disambiguate them with the `knowldege` hyperlinks.

²⁷⁶ Examples of posets include any set of numbers (e.g. $\mathbb{N}, \mathbb{Q}, \mathbb{R}$) equipped with the usual (non-strict) order \leq , and $\mathcal{P}_{\text{inc}} X$ with the inclusion order.

In practice, it would be useful to replace the symbol for $=_{\perp}$ with \leq so the axioms become the more familiar

$$\hat{E}_{\text{Poset}} = \{x \vdash x \leq x, x \leq y, y \leq x \vdash x = y, x \leq y, y \leq z \vdash x \leq z\}.$$

Example 2.34 (Met). The category of **metric spaces** and nonexpansive maps is the full subcategory of $[0, 1]\mathbf{Spa}$ (taking $[0, \infty]$ works just as well) with all $[0, 1]$ -spaces satisfying symmetry, reflexivity, identity of indiscernibles and triangle inequality stated as quantitative equations:²⁷⁷ \hat{E}_{Met} contains all the following

$$\begin{aligned} \forall \varepsilon \in [0, 1], \quad & y =_{\varepsilon} x \vdash x =_{\varepsilon} y \\ & \vdash x =_0 x \\ & x =_0 y \vdash x = y \\ \forall \varepsilon, \delta \in [0, 1], \quad & x =_{\varepsilon} y, y =_{\delta} z \vdash x =_{\varepsilon+\delta} z. \end{aligned}$$

²⁷⁷ Examples of metric spaces include $[0, 1]$ with the Euclidean distance from Example 2.14, the Kantorovich distance from Example 3.5, and the total variation distance from Example 3.92.

2.3 The Categories \mathbf{GMet}

In this section, we prove some basic results about the categories of generalized metric spaces. We fix a complete lattice L and a class of quantitative equations \hat{E} throughout, and denote by \mathbf{GMet} the category of L -spaces that satisfy \hat{E} . The goal here is mainly to become familiar with L -spaces and quantitative equations, so not everything will be useful later. This also means we will avoid using abstract results (that we prove later) which can (sometimes drastically) simplify some proofs.²⁷⁸

We also take some time to identify a few (well-known) conditions on L -spaces that *cannot* be expressed via quantitative equations. These proofs are always in the same vein, we know \mathbf{GMet} has some property, we show the class of L -spaces with a condition does not have that property, hence that condition is not expressible as a class of quantitative equations.

Products

The category \mathbf{GMet} has all products. We prove this in three steps. First, we find the terminal object, second we show \mathbf{LSpa} has all products, and third we show the products of L -spaces which all satisfy some quantitative equation also satisfies that quantitative equation.

Proposition 2.35. *The category \mathbf{GMet} has a terminal object.*

Proof. The terminal object $\mathbf{1}$ in \mathbf{LSpa} is relatively easy to find,²⁷⁹ it is a singleton $\{*\}$ with the L -relation $d_{\mathbf{1}}$ sending $(*, *)$ to \perp . Indeed, for any L -space \mathbf{X} , we have a function $! : X \rightarrow *$ that sends any x to $*$, and because $d_{\mathbf{1}}(*, *) = \perp \leq d_{\mathbf{X}}(x, x')$ for any $x, x' \in X$, $!$ is nonexpansive. We obtain a morphism $! : \mathbf{X} \rightarrow \mathbf{1}$, and since any other morphism $\mathbf{X} \rightarrow \mathbf{1}$ must have the same underlying function,²⁸⁰ $!$ is the unique morphism of this type.

Since \mathbf{GMet} is a full subcategory of \mathbf{LSpa} , it is enough to show $\mathbf{1}$ is in \mathbf{GMet} to conclude it is the terminal object in this subcategory. We can do this by showing $\mathbf{1}$

²⁷⁸ For instance, we will see that $U : \mathbf{GMet} \rightarrow \mathbf{Set}$ is a right adjoint, so it has many nice properties which we could use in this section.

²⁷⁹ Again, many abstract results could help guide our search, but it is enough to have a bit of intuition about L -spaces.

²⁸⁰ Because $\{*\}$ is terminal in \mathbf{Set} .

satisfies absolutely all quantitative equations, and in particular those of \hat{E} .²⁸¹ Let \mathbf{X} be any L-space, $x, y \in X$, and $\varepsilon \in L$. As we have seen above, there is only one assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{1}$, and it sends x and y to $*$. This means

$$\hat{\iota}(x) = * = \hat{\iota}(y) \quad \text{and} \quad d_{\mathbf{1}}(\hat{\iota}(x), \hat{\iota}(y)) = d_{\mathbf{1}}(*, *) = \perp \leq \varepsilon.$$

Therefore, $\mathbf{1}$ satisfies both $\mathbf{X} \vdash x = y$ and $\mathbf{X} \vdash x =_{\varepsilon} y$. We conclude $\mathbf{1} \in \mathbf{GMet}$. \square

Proposition 2.36. *The category \mathbf{LSpa} has all products.*

Proof. Let $\{\mathbf{A}_i = (A_i, d_i) \mid i \in I\}$ be a family of L-spaces indexed by I . We define the L-space $\mathbf{A} = (A, d)$ with carrier $A = \prod_{i \in I} A_i$ (the cartesian product of the carriers) and L-relation $d : A \times A \rightarrow L$ defined by the following supremum:²⁸²

$$\forall a, b \in A, \quad d(a, b) = \sup_{i \in I} d_i(a_i, b_i). \quad (2.31)$$

For each $i \in I$, we have the evident projection $\pi_i : \mathbf{A} \rightarrow \mathbf{A}_i$ sending $a \in A$ to $a_i \in A_i$, and it is nonexpansive because, by definition, for any $a, b \in A$,

$$d_i(a_i, b_i) \leq \sup_{i \in I} d_i(a_i, b_i) = d(a, b).$$

We will show that \mathbf{A} with these projections is the product $\prod_{i \in I} \mathbf{A}_i$.

Let \mathbf{X} be some L-space and $f_i : \mathbf{X} \rightarrow \mathbf{A}_i$ be a family of nonexpansive maps. By the universal property of the product in \mathbf{Set} , there is a unique function $\langle f_i \rangle : X \rightarrow A$ satisfying $\pi_i \circ \langle f_i \rangle = f_i$ for all $i \in I$. It remains to show $\langle f_i \rangle$ is nonexpansive from \mathbf{X} to \mathbf{A} . For any $x, x' \in X$, we have²⁸³

$$d(\langle f_i \rangle(x), \langle f_i \rangle(x')) = \sup_{i \in I} d_i(f_i(x), f_i(x')) \leq d_{\mathbf{X}}(x, x').$$

Note that a particular case of this construction for I being empty is the terminal object $\mathbf{1}$ from Proposition 2.35. Indeed, the empty cartesian product is the singleton, and the empty supremum is the bottom element \perp . \square

In order to show that satisfaction of a quantitative equation is preserved by the product of L-spaces, we first prove a simple lemma.²⁸⁴

Lemma 2.37. *Let ϕ be a quantitative equation with context \mathbf{X} . If $f : \mathbf{A} \rightarrow \mathbf{B}$ is a nonexpansive map and $\mathbf{A} \models^{\hat{\iota}} \phi$ for a nonexpansive assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$, then $\mathbf{B} \models^{f \circ \hat{\iota}} \phi$.*

Proof. There are two very similar cases. If ϕ is of the form $\mathbf{X} \vdash x = y$, we have²⁸⁵

$$\mathbf{A} \models^{\hat{\iota}} \phi \iff \hat{\iota}(x) = \hat{\iota}(y) \implies f\hat{\iota}(x) = f\hat{\iota}(y) \iff \mathbf{B} \models^{f \circ \hat{\iota}} \phi.$$

If ϕ is of the form $\mathbf{X} \vdash x =_{\varepsilon} y$, we have²⁸⁶

$$\mathbf{A} \models^{\hat{\iota}} \phi \iff d_{\mathbf{A}}(\hat{\iota}(x), \hat{\iota}(y)) \leq \varepsilon \implies d_{\mathbf{B}}(f\hat{\iota}(x), f\hat{\iota}(y)) \leq \varepsilon \iff \mathbf{B} \models^{f \circ \hat{\iota}} \phi. \quad \square$$

Proposition 2.38. *If all L-spaces \mathbf{A}_i satisfy a quantitative equation ϕ , then $\prod_{i \in I} \mathbf{A}_i \models \phi$.*

²⁸¹ Which defined \mathbf{GMet} at the start of this section.

²⁸² For $a \in A$, let a_i be the i th coordinate of a .

²⁸³ The equation holds because the i th coordinate of $\langle f_i \rangle(x)$ is $f_i(x)$ by definition of $\langle f_i \rangle$, and the inequality holds because for all $i \in I$, $d_i(f_i(x), f_i(x')) \leq d_{\mathbf{X}}(x, x')$ by nonexpansiveness of f_i .

²⁸⁴ It may remind you of Lemma 1.20 which states the same result for homomorphism and non-quantitative equations.

²⁸⁵ The equivalences hold by definition of \models .

²⁸⁶ The equivalences hold by definition of \models , and the implication holds by nonexpansiveness of f .

Proof. Let $\mathbf{A} = \prod_{i \in I} \mathbf{A}_i$ and \mathbf{X} be the context of ϕ . It is enough to show that for any assignment $\hat{l} : \mathbf{X} \rightarrow \mathbf{A}$, the following equivalence holds:²⁸⁷

$$\left(\forall i \in I, \mathbf{A}_i \models^{\pi_i \circ \hat{l}} \phi \right) \iff \mathbf{A} \models^{\hat{l}} \phi. \quad (2.32)$$

The proposition follows because if $\mathbf{A}_i \models \phi$ for all $i \in I$, then the L.H.S. holds for any \hat{l} , hence the R.H.S. does too, and we conclude $\mathbf{A} \models \phi$. Let us prove (2.32).

(\Rightarrow) Consider the case $\phi = \mathbf{X} \vdash x = y$. The satisfaction $\mathbf{A}_i \models^{\pi_i \circ \hat{l}} \phi$ means $\pi_i \hat{l}(x) = \pi_i \hat{l}(y)$. If it is true for all $i \in I$, then we must have $\hat{l}(x) = \hat{l}(y)$ by universality of the product, thus we get $\mathbf{A} \models^{\hat{l}} \phi$. In case $\phi = \mathbf{X} \vdash x =_\varepsilon y$, the satisfaction $\mathbf{A}_i \models^{\pi_i \circ \hat{l}} \phi$ means $d_{\mathbf{A}_i}(\pi_i \hat{l}(x), \pi_i \hat{l}(y)) \leq \varepsilon$. If it is true for all $i \in I$, we get $\mathbf{A} \models \phi$ because

$$d_{\mathbf{A}}(\hat{l}(x), \hat{l}(y)) = \sup_{i \in I} d_{\mathbf{A}_i}(\pi_i \hat{l}(x), \pi_i \hat{l}(y)) \leq \varepsilon.$$

(\Leftarrow) Apply Lemma 2.37 for all π_i . □

Corollary 2.39. *The category \mathbf{GMet} has all products, and they are computed like in \mathbf{LSpa} .*²⁸⁸

Unfortunately, this means that the notion of metric space originally defined in [Fré06], and incidentally what the majority of mathematicians calls a metric space, is not an instance of generalized metric space as we defined them. Since they only allow finite distances, some infinite products do not exist.²⁸⁹ In general, if one wants to bound the distance above by some quantity $B \in \mathbf{L}$, this can be done with the equation $x, y \vdash x =_B y$, but the value B is still allowed as a distance. For instance $[0, 1]\mathbf{Spa}$ is the full subcategory of $[0, \infty]\mathbf{Spa}$ defined by the equation $x, y \vdash x =_1 y$.

Arguably, this is only a superficially negative result since it is already common in parts of the literature (e.g. [BvBR98, Law02, HST14]) to allow infinite distances because the resulting category of metric spaces has better properties (like having infinite products and coproducts).

Let us give two other conditions on $[0, \infty]$ -spaces, arising in the definition of partial metrics [Mat94, Definition 3.1], which are not preserved under (finite) products.

Definition 2.40. A $[0, \infty]$ -space (A, d) is called a **partial metric space** if it satisfies the following conditions:²⁹⁰

$$\forall a, b \in A, \quad a = b \iff d(a, a) = d(a, b) = d(b, b) \quad (2.33)$$

$$\forall a, b \in A, \quad d(a, a) \leq d(a, b) \quad (2.34)$$

$$\forall a, b \in A, \quad d(a, b) = d(b, a) \quad (2.35)$$

$$\forall a, b, c \in A, \quad d(a, c) \leq d(a, b) + d(b, c) - d(b, b) \quad (2.36)$$

These conditions look similar to what we were able to translate into equations before, but the first and last are problematic.²⁹¹

For (2.33), note that the forward implication is trivial, but for the converse, we would need to compare three distances at once inside the context, which seems impossible because the context only individually bounds distances by above. For

²⁸⁷ When I is empty, the L.H.S. of (2.32) is vacuously true, and the R.H.S. is true since \mathbf{A} is the terminal L-space which we showed satisfies all quantitative equations in Proposition 2.35.

²⁸⁸ We showed that products in \mathbf{LSpa} of objects in \mathbf{GMet} also belong to \mathbf{GMet} , it follows that this is also their products in \mathbf{GMet} because the latter is a full subcategory of \mathbf{LSpa} .

²⁸⁹ For instance let \mathbf{A}_n be the metric space with two points $\{a, b\}$ at distance $n > 0 \in \mathbb{N}$ from each other. Then $\mathbf{A} = \prod_{n > 0 \in \mathbb{N}} \mathbf{A}_n$ exists in $[0, \infty]\mathbf{Spa}$ as we have just proven, but

$$d_{\mathbf{A}}(a^*, b^*) = \sup_{n > 0 \in \mathbb{N}} d_{\mathbf{A}_n}(a, b) = \sup_{n > 0 \in \mathbb{N}} n = \infty,$$

which means \mathbf{A} is not a metric space in the sense of Definition 0.1.

²⁹⁰ There is some ambiguity in what $+$ and $-$ means when dealing with ∞ (the original paper [Mat94] supposes distances are finite), but it is irrelevant for us.


²⁹¹ We can translate (2.34) into $x =_\varepsilon y \vdash x =_\varepsilon x$, and (2.35) is just symmetry which we can translate into $y =_\varepsilon x \vdash x =_\varepsilon y$.

(2.36), the problem comes from the minus operation on distances which will not interact well with upper bounds. Indeed, if we naively tried something like

$$x =_{\varepsilon_1} y, y =_{\varepsilon_2} z, y =_{\varepsilon_3} z \vdash x =_{\varepsilon_1 + \varepsilon_2 - \varepsilon_3} z,$$


we could always take ε_3 huge (even ∞) and make the distance between x and z as close to 0 as we would like (provided we can take ε_1 and ε_2 finite).

These are just informal arguments, but thanks to Corollary 2.39, we can prove formally that these conditions are not expressible as (classes of) quantitative equations. Let **A** and **B** be the $[0, \infty]$ -spaces pictured below (the distances are symmetric).²⁹²



²⁹² The numbers on the lines indicate the distance between the ends of the line, e.g. $d_A(a_1, a_1) = 0$, $d_A(a_1, a_3) = 1$, and $d_B(b_2, b_3) = 10$.

We can verify (by exhaustive checks) that **A** and **B** are partial metric spaces. If we take their product inside $[0, \infty]\mathbf{Spa}$, we find the following $[0, \infty]$ -space (some distances are omitted) which does not satisfy (2.33) nor (2.36).²⁹³



²⁹³ For (2.33), the three points in the middle row $\{a_2b_1, a_2b_2, a_2b_3\}$ are all at distance 10 from each other and from themselves while not being equal. For (2.36), we have (on the diagonal)

$$d_A(a_1b_1, a_3b_3) = 15, \text{ and} \\ d_A(a_1b_1, a_2b_2) + d_A(a_2b_2, a_3b_3) - d_A(a_2b_2, a_2b_2) = 10,$$

but $15 > 10$.

We infer that there is no class \hat{E} of quantitative equations such that $\mathbf{GMet}([0, \infty], \hat{E})$ is the full subcategory of $[0, \infty]\mathbf{Spa}$ containing all the partial metric spaces.²⁹⁴

Coproducts

The case of coproducts in \mathbf{GMet} is more delicate. While \mathbf{LSpa} has coproducts, they do not always satisfy the equations satisfied by each of their components.

Proposition 2.41. *The category \mathbf{GMet} has an initial object.*

Proof. The initial object \emptyset in \mathbf{LSpa} is the empty set with the only possible L-relation $\emptyset \times \emptyset \rightarrow L$ (the empty function). The empty function $f : \emptyset \rightarrow X$ is always nonexpansive from \emptyset to X because (2.3) is vacuously satisfied.

²⁹⁴ It is still possible that the category of partial metrics and nonexpansive maps is identified with some $\mathbf{GMet}(L, \hat{E})$ for some cleverly picked L and \hat{E} . That would mean (infinite) products of partial metrics exist but they are not computed with supremums.

Just as for the terminal object, since **GMet** is a full subcategory of **LSpa**, it suffices to show \emptyset is in **GMet** to conclude it is initial in this subcategory. We do this by showing \emptyset satisfies absolutely all quantitative equations, and in particular those of \hat{E} . This is easily done because when \mathbf{X} is not empty,²⁹⁵ there are no assignments $\mathbf{X} \rightarrow \emptyset$, so \emptyset vacuously satisfies $\mathbf{X} \vdash x = y$ and $\mathbf{X} \vdash x =_\varepsilon y$. \square

Proposition 2.42. *The category **LSpa** has all coproducts.*

Proof. We just showed the empty coproduct (i.e. the initial object) exists. Let $\{\mathbf{A}_i = (A_i, d_i) \mid i \in I\}$ be a family of L-spaces indexed by a non-empty set I . We define the L-space $\mathbf{A} = (A, d)$ with carrier $A = \coprod_{i \in I} A_i$ (the disjoint union of the carriers) and L-relation $d : A \times A \rightarrow \mathbb{L}$ defined by:²⁹⁶

$$\forall a, b \in A, \quad d(a, b) = \begin{cases} d_i(a, b) & \exists i \in I, a, b \in A_i \\ \top & \text{otherwise} \end{cases}.$$

For each $i \in I$, we have the evident coprojection $\kappa_i : \mathbf{A}_i \rightarrow \mathbf{A}$ sending $a \in A_i$ to its copy in A , and it is nonexpansive because, by definition, for any $a, b \in A_i$, $d(a, b) = d_i(a, b)$.²⁹⁷ We show \mathbf{A} with these coprojections is the coproduct $\coprod_{i \in I} \mathbf{A}_i$.

Let \mathbf{X} be some L-space and $f_i : \mathbf{A}_i \rightarrow \mathbf{X}$ be a family of nonexpansive maps. By the universal property of the coproduct in **Set**, there is a unique function $[f_i] : A \rightarrow X$ satisfying $[f_i] \circ \kappa_i = f_i$ for all $i \in I$. It remains to show $[f_i]$ is nonexpansive from \mathbf{A} to \mathbf{X} . For any $a, b \in A$, suppose a belongs to A_i and b to A_j for some $i, j \in I$, then we have²⁹⁸

$$d_{\mathbf{X}}([f_i](a), [f_i](b)) = d_{\mathbf{X}}(f_i(a), f_j(b)) \leq \begin{cases} d_i(a, b) & i = j \\ \top & \text{otherwise} \end{cases} = d(a, b).$$

\square

Because the distance between elements in different copies does not depend on the original spaces, it is easy to construct a quantitative equation that is not preserved by coproducts of L-spaces. For instance, even if all \mathbf{A}_i satisfy $x, y \vdash x =_\varepsilon y$ for some fixed $\varepsilon \neq \top \in \mathbb{L}$,²⁹⁹ the coproduct $\coprod_{i \in I} \mathbf{A}_i$ in **LSpa** does not satisfy it because some distances are $\top > \varepsilon$.

Still, **GMet** always has coproducts as we will show in Corollary 3.60, but they are not computed like in **LSpa**, and they are not that easy to define.³⁰⁰

Isometries

Since the forgetful functor $U : \mathbf{LSpa} \rightarrow \mathbf{Set}$ preserves isomorphisms, we know that the underlying function of an isomorphism in **LSpa** is a bijection between the carriers. What is more, we show in Proposition 2.44 it must preserve distances on the nose, it is called an isometry.

Definition 2.43 (Isometry). A nonexpansive map $f : \mathbf{X} \rightarrow \mathbf{Y}$ is an **isometry** if³⁰¹

²⁹⁵ The context of a quantitative equation cannot be empty because the variables, say x and y , must belong to the context.

²⁹⁶ In words, \mathbf{A} is the L-space with a copy of each \mathbf{A}_i where the L-relation sends two points in different copies to \top (intuitively, the copies are completely unrelated inside \mathbf{A}).

²⁹⁷ Hence κ_i is even an isometric embedding.

²⁹⁸ The first equation holds by definition of $[f_i]$ (it applies f_i to elements in the copy of A_i). The inequality holds by nonexpansiveness of f_i which is equal to f_j when $i = j$. The second equation is the definition of d .

²⁹⁹ i.e. there is an upper bound smaller than \top on all distances in all \mathbf{A}_i .

³⁰⁰ In many cases like **Met** and **Poset**, they are computed like in **LSpa**.

³⁰¹ The inequality in (2.3) is replaced by an equation.

$$\forall x, x' \in X, \quad d_{\mathbf{Y}}(f(x), f(x')) = d_{\mathbf{X}}(x, x'). \quad (2.37)$$

If furthermore f is injective, we call it an **isometric embedding**.³⁰² If $f : \mathbf{X} \rightarrow \mathbf{Y}$ is an isometric embedding, we can identify \mathbf{X} with the subspace of \mathbf{Y} containing all the elements in the image of f . Conversely, the inclusion of a subspace of \mathbf{Y} in \mathbf{Y} is always an isometric embedding.

Proposition 2.44. *In \mathbf{GMet} , isomorphisms are precisely the bijective isometries.*

Proof. We show a morphism $f : \mathbf{X} \rightarrow \mathbf{Y}$ has an inverse $f^{-1} : \mathbf{Y} \rightarrow \mathbf{X}$ if and only if it is a bijective isometry.

(\Rightarrow) Since the underlying functions of f and f^{-1} are inverses, they must be bijections. Moreover, using (2.3) twice, we find that for any $x, x' \in X$,³⁰³

$$d_{\mathbf{X}}(x, x') = d_{\mathbf{X}}(f^{-1}f(x), f^{-1}f(x')) \leq d_{\mathbf{Y}}(f(x), f(x')) \leq d_{\mathbf{X}}(x, x'),$$

thus $d_{\mathbf{X}}(x, x') = d_{\mathbf{Y}}(f(x), f(x'))$, so f is an isometry.

(\Leftarrow) Since f is bijective, it has an inverse $f^{-1} : Y \rightarrow X$ in \mathbf{Set} , but we have to show f^{-1} is nonexpansive from \mathbf{Y} to \mathbf{X} . For any $y, y' \in Y$, by surjectivity of f , there are $x, x' \in X$ such that $y = f(x)$ and $y' = f(x')$, then we have

$$\begin{aligned} d_{\mathbf{X}}(f^{-1}(y), f^{-1}(y')) &= d_{\mathbf{X}}(f^{-1}f(x), f^{-1}f(x')) \\ &= d_{\mathbf{X}}(x, x') \\ &\stackrel{(2.37)}{=} d_{\mathbf{Y}}(f(x), f(x')) = d_{\mathbf{Y}}(y, y'). \end{aligned}$$

Hence f^{-1} is nonexpansive, it is even an isometry. \square

In particular, this means, as is expected, that isomorphisms preserve the satisfaction of quantitative equations. We can show a stronger statement: any isometric embedding reflects the satisfaction of quantitative equations.³⁰⁴

Lemma 2.45. *Let $f : \mathbf{Y} \rightarrow \mathbf{Z}$ be an isometric embedding between L-spaces and ϕ a quantitative equation, then*

$$\mathbf{Z} \models \phi \implies \mathbf{Y} \models \phi. \quad (2.38)$$

Proof. Let \mathbf{X} be the context of ϕ . Any nonexpansive assignment $\hat{l} : \mathbf{X} \rightarrow \mathbf{Y}$ yields an assignment $f \circ \hat{l} : \mathbf{X} \rightarrow \mathbf{Z}$. By hypothesis, we know that \mathbf{Z} satisfies ϕ for this particular assignment, namely,

$$\mathbf{Z} \models^{f \circ \hat{l}} \phi. \quad (2.39)$$

We can use this and the fact that f is an isometric embedding to show $\mathbf{Y} \models^{\hat{l}} \phi$. There are two very similar cases.

If $\phi = \mathbf{X} \vdash x = y$, then we have $\hat{l}(x) = \hat{l}(y)$ because we know $f\hat{l}(x) = f\hat{l}(y)$ by (2.39) and f is injective.

If $\phi = \mathbf{X} \vdash x =_{\varepsilon} y$, then we have $d_{\mathbf{Y}}(\hat{l}(x), \hat{l}(y)) = d_{\mathbf{Z}}(f\hat{l}(x), f\hat{l}(y)) \leq \varepsilon$, where the equation holds because f is an isometry and the inequality holds by (2.39). \square

Corollary 2.46. *Let $f : \mathbf{Y} \rightarrow \mathbf{Z}$ be an isometric embedding between L-spaces. If \mathbf{Z} belongs to \mathbf{GMet} , then so does \mathbf{Y} . In particular, all the subspaces of a generalized metric space are also generalized metric spaces.³⁰⁵*

³⁰² This name is relatively rare because when dealing with metric spaces, the separation axiom implies that an isometry is automatically injective. This is also true for partial orders, where the name *order embedding* is common [DPo2, Definition 1.34.(ii)].

³⁰³ This is a general argument showing that any non-expansive function with a right inverse is an isometry, it is also an isometric embedding because a right inverse in \mathbf{Set} implies injectivity.

³⁰⁴ This is stronger because we have just shown the inverse of an isomorphism is an isometric embedding.

³⁰⁵ Both parts are immediate. The first follows from applying (2.38) to all ϕ in \hat{E} , the class of quantitative equations defining \mathbf{GMet} . The second follows from the inclusion of a subspace being an isometric embedding.

Example 2.47. Corollary 2.46 can be useful to identify some properties of L-spaces that cannot be modelled with quantitative equations. Here are a few of examples.

1. A binary relation $R \subseteq X \times X$ is called **total** if for every $x \in X$, there exists $y \in X$ such that $(x, y) \in R$. Let **TotRel** be the full subcategory of **BSpa** containing only total relations. Is **TotRel** equal to $\mathbf{GMet}(B, \hat{E})$ for some \hat{E} ? The existential quantification in the definition of total seems hard to simulate with a quantitative equation, but this is not a guarantee that maybe several equations cannot interact in such a counter-intuitive way.

In order to prove that no class \hat{E} defines total relations (i.e. $\mathbf{X} = \hat{E}$ if and only if the relation corresponding to $d_{\mathbf{X}}$ is total), we can exhibit an example of a B-space that is total with a subspace that is not total. It follows that **TotRel** is not closed under taking subspaces, so it is not a category of generalized metric spaces by Corollary 2.46.³⁰⁶

Let \mathbf{N} be the B-space with carrier \mathbb{N} and B-relation $d_{\mathbf{N}}(n, m) = \perp \Leftrightarrow m = n + 1$ (the corresponding relation is the graph of the successor function). This space satisfies totality, but the subspace obtained by removing 1 is not total because $d_{\mathbf{N}}(0, n) = \perp$ only when $n = 1$.

This same example works to show that surjectivity³⁰⁷ cannot be defined via quantitative equations.

2. A very famous condition to impose on metric spaces is **completeness** (we do not need to define it here). Just as famous is the fact that \mathbb{R} with the Euclidean metric from Example 2.14 is complete but the subspace \mathbb{Q} is not. Thus, completeness cannot be defined via quantitative equations.³⁰⁸

With this characterization of isomorphisms, we can also show the forgetful functor $U : \mathbf{GMet} \rightarrow \mathbf{Set}$ is an isofibration which concretely means that if you have a bijection $f : X \rightarrow Y$ and a generalized metric d_Y on Y , then you can construct a generalized metric d_X on X such that $f : \mathbf{X} \rightarrow \mathbf{Y}$ is an isomorphism. Indeed, if you let $d_X(x, x') = d_Y(f(x), f(x'))$, then f is automatically a bijective isometry.³⁰⁹

Definition 2.48 (Isofibration). A functor $P : \mathbf{C} \rightarrow \mathbf{D}$ is called an **isofibration**³¹⁰ if for any isomorphism $f : X \rightarrow PY$ in \mathbf{D} , there is an isomorphism $g : X' \rightarrow Y$ such that $Pg = f$, in particular $PX' = X$.

Proposition 2.49. *The forgetful functor $U : \mathbf{GMet} \rightarrow \mathbf{Set}$ is an isofibration.*

We wonder now how to complete the conceptual diagram below.

$$\begin{array}{ccc} \text{isomorphism in } \mathbf{GMet} & \longleftrightarrow & \text{bijective isometries} \\ \text{??? in } \mathbf{GMet} & \longleftrightarrow & \text{isometric embeddings} \end{array}$$

Since isometric embeddings correspond to subspaces, one might think that they are the monomorphisms in **GMet**. Unfortunately, they are way more restrained. Any nonexpansive map that is injective is already a monomorphism. To prove this, we rely on the existence of a space \mathbf{A} that informally *can pick elements*.

³⁰⁶ Actually, we have only proven that **TotRel** cannot be defined as a subcategory of **BSpa** with quantitative equations. There may still be some convoluted way that $\mathbf{TotRel} \cong \mathbf{GMet}(L, \hat{E})$.

³⁰⁷ This condition is symmetric to totality: $R \subseteq X \times X$ is **surjective** if for every $y \in X$, there exists $x \in X$ such that $(x, y) \in R$.

³⁰⁸ Still with the caveat that the full subcategory of complete metric spaces might still be isomorphic to some $\mathbf{GMet}(L, \hat{E})$.

³⁰⁹ Clearly, it is the unique distance on X that works, and we know that \mathbf{X} belongs to **GMet** thanks to Corollary 2.46.

³¹⁰ This term seems to have been coined by Lack and Paoli in [Lac07, §3.1] or [LPo8, §6].

Proposition 2.50. *There is a generalized metric space \mathbb{H} on the set $\{*\}$ such that for any other space \mathbf{X} , any function $f : \{*\} \rightarrow X$ is a nonexpansive map $\mathbb{H} \rightarrow \mathbf{X}$.³¹¹*

Proof. In \mathbf{LSpa} , \mathbb{H} is easy to find, its L-relation is defined by $d_{\mathbb{H}}(*, *) = \top$. Indeed, any function $f : \{*\} \rightarrow X$ is nonexpansive because \top is the maximum value $d_{\mathbf{X}}$ can assign, so

$$d_{\mathbf{X}}(f(*), f(*)) \leq \top = d_{\mathbb{H}}(*, *).$$

Unfortunately, this L-space does not satisfy some quantitative equations (e.g. reflexivity $x \vdash x = \perp x$), so we cannot guarantee it belongs to \mathbf{GMet} .

Recall that $\mathbf{1}$ is a generalized metric space on the same set $\{*\}$, but with $d_{\mathbf{1}}(*, *) = \perp$. However, in many cases, $\mathbf{1}$ is not the right candidate either because if every function $f : \{*\} \rightarrow X$ is nonexpansive from $\mathbf{1}$ to \mathbf{X} , it means $d_{\mathbf{X}}(x, x) = \perp$ for all $x \in X$, which is not always the case.³¹²

We have two L-spaces at the extremes of a range of L-spaces $\{(\{*\}, d_{\varepsilon})\}_{\varepsilon \in \mathbf{L}}$, where the L-relation d_{ε} sends $(*, *)$ to ε . At one extreme, we are guaranteed to be in \mathbf{GMet} , but we are too restricted, and at the other extreme we might not belong to \mathbf{GMet} . Getting inspiration from the [intermediate value theorem](#), we can attempt to find a middle ground, namely, a value $\varepsilon \in \mathbf{L}$ such that setting $d_{\mathbb{H}}(*, *) = \varepsilon$ yields a space that lives in \mathbf{GMet} but is not too restricted.

One natural thing to do is to take the biggest quantity ε such that d_{ε} is a generalized metric. In other words, we take the least restricted space that is in \mathbf{GMet} . Formally,

$$d_{\mathbb{H}}(*, *) = \sup \{ \varepsilon \in \mathbf{L} \mid (\{*\}, d_{\varepsilon}) \models \hat{E} \}.$$

It remains to check that any function $f : \{*\} \rightarrow X$ is nonexpansive from \mathbb{H} to $\mathbf{X} \in \mathbf{GMet}$. Consider the image of f seen as a subspace of \mathbf{X} . By Corollary 2.46, it belongs to \mathbf{GMet} and hence satisfies \hat{E} . Moreover, it is clearly isomorphic to the L-space $(\{*\}, d_{\varepsilon})$ with $\varepsilon = d_{\mathbf{X}}(f(*), f(*))$, which means that L-space satisfies \hat{E} as well (by Corollary 2.46 again). We conclude that $d_{\mathbf{X}}(f(*), f(*)) \leq d_{\mathbb{H}}(*, *)$.³¹³ \square

Proposition 2.51. *In \mathbf{GMet} , monomorphisms are precisely the injective nonexpansive maps.*

Proof. We show a morphism $f : \mathbf{X} \rightarrow \mathbf{Y}$ is monic if and only if it is injective.

(\Rightarrow) Let $x, x' \in X$ be such that $f(x) = f(x')$, and identify these elements with functions $x, x' : \{*\} \rightarrow X$ sending $*$ to x and x' respectively. By Proposition 2.50, we get two nonexpansive maps $x, x' : \mathbb{H} \rightarrow \mathbf{X}$. Post-composing by f , we find that $f \circ x = f \circ x'$ because they both send $*$ to $f(x) = f(x')$. By monicity of f , we find that $x = x'$ (as morphisms and hence as elements of X). We conclude f is injective.

(\Leftarrow) Suppose that $f \circ g = f \circ h$ for some nonexpansive maps $g, h : \mathbf{Z} \rightarrow \mathbf{X}$. Applying the forgetful functor $U : \mathbf{GMet} \rightarrow \mathbf{Set}$, we find that $f \circ g = f \circ h$ also as functions. Since Uf is monic (i.e. injective), Ug and Uh must be equal, and since U is faithful, we obtain $g = h$. \square

It remains to give a categorical characterization of isometric embeddings. This will rely on a well-known³¹⁴ abstract notion that we define here for completeness.

³¹¹ In category theory speak, \mathbb{H} is a representing object of the forgetful functor $U : \mathbf{GMet} \rightarrow \mathbf{Set}$.

³¹² It is equivalent to satisfying reflexivity.

³¹³ As a bonus, one could check that for any $\varepsilon \in \mathbf{L}$ that is smaller than $d_{\mathbb{H}}(*, *)$, $(\{*\}, d_{\varepsilon})$ also belongs to \mathbf{GMet} (using Lemma 2.37).

³¹⁴ While it is well-known, especially to those familiar with fibered category theory, it does not usually fit in a basic category theory course.

Definition 2.52 (Cartesian morphism). Let $F : \mathbf{C} \rightarrow \mathbf{D}$ be a functor, and $f : A \rightarrow B$ be a morphism in \mathbf{C} . We say f is a **cartesian morphism** (relative to F) if for every morphism $g : X \rightarrow B$ and factorization $Fg = Ff \circ u$, there exists a unique morphism $\hat{u} : X \rightarrow A$ with $F\hat{u} = u$ satisfying $x = f \circ \hat{u}$. This can be summarized (without the quantifiers) in the diagram below.

$$\begin{array}{ccc}
 X & & FX \\
 \hat{u} \downarrow & \searrow g & u \downarrow \searrow Fg \\
 A & \xrightarrow{f} & B & \xrightarrow{F} & FA & \xrightarrow{Ff} & FB
 \end{array}$$

Example 2.53 (in \mathbf{GMet}). Let us unroll this in the important case for us, when F is the forgetful functor $U : \mathbf{GMet} \rightarrow \mathbf{Set}$. A nonexpansive map $f : \mathbf{A} \rightarrow \mathbf{B}$ is a cartesian morphism if for any nonexpansive map $g : \mathbf{X} \rightarrow \mathbf{B}$, all functions $u : \mathbf{X} \rightarrow \mathbf{A}$ satisfying $g = f \circ u$ are nonexpansive maps $u : \mathbf{X} \rightarrow \mathbf{A}$.³¹⁵

We can turn this around into an equivalent definition. The morphism $f : \mathbf{A} \rightarrow \mathbf{B}$ is cartesian if for all functions $u : \mathbf{X} \rightarrow \mathbf{A}$, $f \circ u$ being nonexpansive from \mathbf{X} to \mathbf{B} implies u is nonexpansive from \mathbf{X} to \mathbf{A} .³¹⁶ In [AHS06, Definition 8.6], f is also called an *initial morphism*.

Proposition 2.54. *A morphism $f : \mathbf{A} \rightarrow \mathbf{B}$ in \mathbf{GMet} is an isometric embedding if and only if it is monic and cartesian.*

Proof. By Proposition 2.51, being an isometric embedding is equivalent to being a monomorphism (i.e. being injective) and being an isometry. Therefore, it is enough to show that when f is injective, isometry \iff cartesian.

(\implies) Suppose f is an isometry, and let $u : \mathbf{X} \rightarrow \mathbf{A}$ be a function such that $f \circ u$ is nonexpansive from $\mathbf{X} \rightarrow \mathbf{B}$, we need to show u is nonexpansive from $\mathbf{X} \rightarrow \mathbf{A}$.³¹⁷ This is true because for all $x, x' \in \mathbf{X}$,

$$d_{\mathbf{A}}(u(x), u(x')) = d_{\mathbf{B}}(fu(x), fu(x')) \leq d_{\mathbf{X}}(x, x'),$$

where the equation follows from f being an isometry, and the inequality from nonexpansiveness of $f \circ u$.

(\impliedby) Suppose f is cartesian. For any $a, a' \in \mathbf{A}$, we know that $d_{\mathbf{B}}(f(a), f(a')) \leq d_{\mathbf{A}}(a, a')$, but we still need to show the converse inequality. Let \mathbf{X} be the subspace of \mathbf{B} containing only the image of a and a' (its carrier is $\{f(a), f(a')\}$), and $u : \mathbf{X} \rightarrow \mathbf{A}$ be the function sending $f(a)$ to a and $f(a')$ to a' .³¹⁸ Notice that $f \circ u$ is the inclusion of \mathbf{X} in \mathbf{B} which is nonexpansive. Because f is cartesian, u must then be nonexpansive from \mathbf{X} to \mathbf{A} which implies

$$d_{\mathbf{A}}(a, a') = d_{\mathbf{A}}(u(f(a)), u(f(a'))) \leq d_{\mathbf{X}}(f(a), f(a')) = d_{\mathbf{B}}(f(a), f(a')).$$

We conclude that f is an isometry. □

Corollary 2.55. *If the composition $\mathbf{A} \xrightarrow{f} \mathbf{B} \xrightarrow{g} \mathbf{C}$ is an isometric embedding, then f is an isometric embedding.³¹⁹*

³¹⁵ We do not bother to write \hat{u} as it is automatically unique with underlying function u because U is faithful.

³¹⁶ If $f \circ u$ is nonexpansive from \mathbf{X} to \mathbf{B} , then it is equal to g for some $g : \mathbf{X} \rightarrow \mathbf{B}$ which yields $u : \mathbf{X} \rightarrow \mathbf{A}$ being nonexpansive.

³¹⁷ We use the second definition of cartesian from Example 2.53.

³¹⁸ We use the injectivity of f here.

³¹⁹ With the characterization of Proposition 2.54, this abstractly follows from [AHS06, Proposition 8.9]. We give the concrete proof anyways.

Proof. It is a standard result that if $g \circ f$ is monic then so is f . Even more standard for injectivity. Now, if $g \circ f$ is an isometry, we have for any $a, a' \in A$,³²⁰

$$d_{\mathbf{A}}(a, a') = d_{\mathbf{C}}(gf(a), gf(a')) \leq d_{\mathbf{B}}(f(a), f(a')) \leq d_{\mathbf{A}}(a, a'),$$

and we conclude that $d_{\mathbf{A}}(a, a') = d_{\mathbf{B}}(f(a), f(a'))$, hence f is an isometry. \square

The question of concretely characterizing epimorphisms is harder to settle. We can do it for **LSpa**, but not for an arbitrary **GMet**.

Proposition 2.56. *In **LSpa**, a morphism $f : \mathbf{X} \rightarrow \mathbf{A}$ is epic if and only if it is surjective.*

Proof. (\Rightarrow) Given any $a \in A$, we define the L-space \mathbf{A}_a to be \mathbf{A} with an additional copy of a with all the same distances. Namely, the carrier is $A + \{*_a\}$, for any $a' \in A$, $d_{\mathbf{A}_a}(*_a, a') = d_{\mathbf{A}}(a, a')$ and $d_{\mathbf{A}_a}(a', *_a) = d_{\mathbf{A}}(a', a)$, and all the other distances are as in \mathbf{A} .³²¹

If $f : \mathbf{X} \rightarrow \mathbf{A}$ is not surjective, then pick $a \in A$ that is not in the image of f , and define two functions $g_a, g_* : A \rightarrow A + \{*_a\}$ that act as identity on all A except a where $g_a(a) = a$ and $g_*(a) = *_a$. By construction, both g_a and g_* are nonexpansive from \mathbf{A} to \mathbf{A}_a and $g_a \circ f = g_* \circ f$. Since $g_a \neq g_*$, f cannot be epic, and we have proven the contrapositive of the forward implication.

(\Leftarrow) Suppose that $g, g' : \mathbf{A} \rightarrow \mathbf{B}$ are morphisms in **LSpa** such that $g \circ f = g' \circ f$. Apply the forgetful functor to get $Ug \circ Uf = Ug' \circ Uf$, and since U is epic in **Set**, we know $Ug = Ug'$. Since U is faithful, we conclude that $g = g'$.³²² \square

The standard example to show that Proposition 2.56 does not generalize to an arbitrary **GMet** is the inclusion of \mathbb{Q} into \mathbb{R} with the Euclidean metric inside **Met**. It is not surjective, but it is epic because any nonexpansive function from \mathbb{R} is determined by its image on the rationals.³²³

In Lemma 1.21, we saw that surjective homomorphisms preserve the satisfaction of classical equations. The “quantitative” version of this result is not that surjective nonexpansive maps preserve satisfaction of quantitative equations (see Remark 2.58). The result in the classical case was proven using the fact that for a homomorphism $h : \mathbb{A} \rightarrow \mathbb{B}$, $\mathbb{A} \models^t \phi \implies \mathbb{B} \models^{h \circ t} \phi$, and we proved a version of this for L-spaces in Lemma 2.37. However, the proof of Lemma 1.21 also used the fact that a surjective function always has a right inverse. This is not true in L-spaces, but we can still prove a weaker result (restricting to surjective nonexpansive maps with a right inverse).

Lemma 2.57. *Let $f : \mathbf{A} \rightarrow \mathbf{B}$ be a split epimorphism between L-spaces and ϕ a quantitative equation, then*

$$\mathbf{A} \models \phi \implies \mathbf{B} \models \phi. \quad (2.40)$$

Proof. Let $g : \mathbf{B} \rightarrow \mathbf{A}$ be the right inverse of f (i.e. $f \circ g = \text{id}_{\mathbf{B}}$) and \mathbf{X} be the context of ϕ .³²⁴ Any nonexpansive assignment $\hat{t} : \mathbf{X} \rightarrow \mathbf{B}$ yields an assignment $g \circ \hat{t} : \mathbf{X} \rightarrow \mathbf{A}$. By hypothesis, we know that \mathbf{A} satisfies ϕ for this particular assignment, namely,

$$\mathbf{A} \models^{g \circ \hat{t}} \phi. \quad (2.41)$$

Now, we can apply Lemma 2.37 with $f : \mathbf{A} \rightarrow \mathbf{B}$ to obtain $\mathbf{B} \models^{f \circ g \circ \hat{t}} \phi$, and since $f \circ g = \text{id}_{\mathbf{B}}$, we conclude $\mathbf{B} \models^{\hat{t}} \phi$. \square

³²⁰ The equation holds by hypothesis that $g \circ f$ is an isometry and the two inequalities hold by nonexpansiveness of g and f .

³²¹ This construction is already impossible to do in an arbitrary **GMet**. For instance, if \mathbf{A} satisfies $x =_0 y \vdash x = y$, then \mathbf{A}_a does not because $d_{\mathbf{A}_a}(a, *_a) = 0$.

³²² This direction works in an arbitrary **GMet**, that is, surjections are epic in any **GMet**.

³²³ For any $r \in \mathbb{R}$, you can always find $q_n \in \mathbb{Q}$ such that $d(q_n, r) \leq \frac{1}{n}$, hence $d_{\mathbf{A}}(f(q_n), f(r)) \leq \frac{1}{n}$ for any nonexpansive $f : (\mathbb{R}, d) \rightarrow \mathbf{A}$. We infer that $f(r)$ is determined by the values of all $f(q_n)$.

³²⁴ Note that we already argued in Footnote 303 that the right inverse implies g is an isometric embedding. Then we could conclude by Corollary 2.46. The proof given here is essentially the same.

Remark 2.58. It is not true in general that the image $f(A)$ of a nonexpansive function $f : \mathbf{A} \rightarrow \mathbf{B}$ (seen as a subspace of \mathbf{B}) satisfies the same equations as \mathbf{A} . For instance,³²⁵ let \mathbf{A} contain two points $\{a, b\}$ all at distance $1 \in [0, \infty]$ from each other (even from themselves). The $[0, \infty]$ -relation is symmetric so it satisfies for all $\varepsilon \in [0, 1]$. $y =_\varepsilon x \vdash x =_\varepsilon y$. If we define \mathbf{B} with the same points and distances except $d_{\mathbf{B}}(a, b) = 0.5$, then the identity function is nonexpansive from \mathbf{A} to \mathbf{B} , but its image is \mathbf{B} in which the distance is not symmetric.

Lemma 2.57 is basically a dual of Lemma 2.45 because isometric embeddings are split monomorphisms, so we do not get additional examples of properties that cannot be expressed with quantitative equations.³²⁶ Combined with Proposition 2.38, we showed that the categories \mathbf{GMet} are closed under subspaces and products. We will have a converse as in Birkhoff’s variety theorem (Theorem 1.29), but we postpone it to Theorem 3.65.

Theorem 2.59. *The subcategory $\mathbf{GMet} \subseteq \mathbf{LSpa}$ is closed under subspaces and products.*

Discrete Spaces

The forgetful functor $U : \mathbf{GMet} \rightarrow \mathbf{Set}$ has a left adjoint. Its concrete description is too involved, so we will prove this later in Corollary 3.58, but for the special case of \mathbf{LSpa} , we can prove it now.


Proposition 2.60. *The forgetful functor $U : \mathbf{LSpa} \rightarrow \mathbf{Set}$ has a left adjoint.*

Proof. For any set X , we define the **discrete space** \mathbf{X}_\top to be the set X equipped with the L-relation $d_\top : X \times X \rightarrow \mathbb{L}$ sending any pair to \top .³²⁷

For any L-space \mathbf{A} and function $f : X \rightarrow A$, the function f is nonexpansive from \mathbf{X}_\top to \mathbf{A} , thus \mathbf{X}_\top is the free object on X (relative to U).

We conclude there is a functor $F : \mathbf{Set} \rightarrow \mathbf{LSpa}$ sending X to \mathbf{X}_\top that is left adjoint to $U : \mathbf{LSpa} \rightarrow \mathbf{Set}$.³²⁸ □

³²⁵ Here is a graphical depiction:



³²⁶ In theory, duality may help in some settings, but I find isometric embeddings are easier to grasp.

³²⁷ When we talk about the discrete generalized metric space on X , we mean the space FX where $F : \mathbf{Set} \rightarrow \mathbf{GMet}$ is the left adjoint of $U : \mathbf{GMet} \rightarrow \mathbf{Set}$ we will describe in Corollary 3.58.

³²⁸ This follows from an abstract categorical argument, see e.g. [Awo10, Proposition 9.4].

3 Universal Quantitative Algebra

Saxophone concerto E minor OP 88

Deluxe

3.1 Quantitative Algebras	88
3.2 Free Quantitative Algebras	99
3.3 Quantitative Equational Logic	118
3.4 Quantitative Alg. Presentations	125
3.5 Lifting Presentations	129

For a comprehensive introduction to the concepts and themes explored in this chapter, please refer to §0.3. Here, we only give a brief overview.

It is time to combine what we learned about universal algebra in Chapter 1 and about generalized metric spaces in Chapter 2 to develop universal quantitative algebra. We follow an outline similar to that of Chapter 1 for definitions, results, and proofs, and we give some examples (reusing those of the previous chapters) throughout this chapter.

Outline: In §3.1 and §1.3, we define quantitative algebras and quantitative equations over a signature, and we explain how to construct the free quantitative algebras. In §3.3, we give the rules for quantitative equational logic to derive quantitative equations from other quantitative equations, and we show it is sound and complete. In §3.4, we define presentations for monads on generalized metric spaces, and we give some examples. In §3.5, we show that any monad lifting of a **Set** monad with an algebraic presentation to **GMet** can also be presented.

In the sequel and unless otherwise stated, Σ is an arbitrary signature, L is an arbitrary complete lattice, and **GMet** is an arbitrary category of generalized metric spaces. We will write $\hat{E}_{\mathbf{GMet}}$ for a class of quantitative equations over L (Definition 2.23) such that $\mathbf{GMet} = \mathbf{GMet}(L, \hat{E}_{\mathbf{GMet}})$.

3.1 Quantitative Algebras

Definition 3.1 (Quantitative algebra). A **quantitative Σ -algebra** (or just quantitative algebra)³²⁹ is a set A equipped with a Σ -algebra structure $(A, \llbracket - \rrbracket_A) \in \mathbf{Alg}(\Sigma)$ and a generalized metric space structure $(A, d_A) \in \mathbf{GMet}$. We will switch between using the single symbol \hat{A} or the triple $(A, \llbracket - \rrbracket_A, d_A)$ when referring to a quantitative algebra, we will also write \mathbb{A} for the **underlying Σ -algebra**, \mathbf{A} for the underlying space, and A for the underlying set.

A **homomorphism** from \hat{A} to \hat{B} is a function $h : A \rightarrow B$ between the underlying sets of \hat{A} and \hat{B} that is both a homomorphism $h : \mathbb{A} \rightarrow \mathbb{B}$ and a nonexpansive function $h : \mathbf{A} \rightarrow \mathbf{B}$. We sometimes emphasize and call h a nonexpansive homomorphism.³³⁰ The identity maps $\text{id}_A : A \rightarrow A$ and the composition of two homomorphisms are always homomorphisms, therefore we have a category whose

³²⁹ We may also simply write algebra.

³³⁰ We will not distinguish between a nonexpansive homomorphism $h : \hat{A} \rightarrow \hat{B}$ and its underlying homomorphism or nonexpansive function or function. We may write Uh with U being the appropriate forgetful functor when necessary.

objects are quantitative algebras and morphisms are nonexpansive homomorphisms. We denote it by $\mathbf{QAlg}(\Sigma)$.

This category is concrete over \mathbf{Set} , $\mathbf{Alg}(\Sigma)$, \mathbf{GMet} with forgetful functors:

- $U : \mathbf{QAlg}(\Sigma) \rightarrow \mathbf{Set}$ sends a quantitative algebra \hat{A} to its underlying set A and a nonexpansive homomorphism to the underlying function between carriers.
- $U : \mathbf{QAlg}(\Sigma) \rightarrow \mathbf{Alg}(\Sigma)$ sends \hat{A} to its underlying algebra A and a nonexpansive homomorphism to the underlying homomorphism.
- $U : \mathbf{QAlg}(\Sigma) \rightarrow \mathbf{GMet}$ sends \hat{A} to its underlying space A and a nonexpansive homomorphism to the underlying nonexpansive function.

One can quickly check that the following diagram commutes, and that it yields an alternative definition of $\mathbf{QAlg}(\Sigma)$ as a pullback of categories.³³¹ We can also mention there is another forgetful functor $U : \mathbf{QAlg}(\Sigma) \rightarrow \mathbf{LSpa}$ obtained by composing $U : \mathbf{QAlg}(\Sigma) \rightarrow \mathbf{GMet}$ with the inclusion $\mathbf{GMet} \rightarrow \mathbf{LSpa}$.

$$\begin{array}{ccc}
 \mathbf{QAlg}(\Sigma) & \xrightarrow{U} & \mathbf{GMet} \\
 \downarrow U & \lrcorner & \downarrow U \\
 & U & \\
 \mathbf{Alg}(\Sigma) & \xrightarrow{U} & \mathbf{Set}
 \end{array}$$

Example 3.2. Since a quantitative algebra is just an algebra and a generalized metric space on the same set, we can find simple examples by combining pieces we have already seen.

1. In Example 1.4, we saw that an algebra for the signature $\Sigma = \{p:0\}$ is just a pair (X, x) comprising a set X with a distinguished point $x \in X$. In Example 2.14, we discussed the \mathbb{N}_∞ -space (H, d) where H is the set of humans and d is the collaboration distance. We can therefore consider the quantitative Σ -algebras $(H, \text{Paul Erdős}, d)$, which is the set of all humans with Paulo Erdős as a distinguished point and the collaboration distance.³³²
2. In Example 1.4, we saw the $\{f:1\}$ -algebra \mathbb{Z} where f is interpreted as adding 1. On top of that, we consider the B-relation corresponding to the partial order \leq on \mathbb{Z} : $d_\leq : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{B}$ that sends (n, m) to \perp if and only if $n \leq m$. We get a quantitative algebra $(\mathbb{Z}, - + 1, d_\leq)$.³³³
3. In Example 2.14, we saw that \mathbb{R} equipped with the Euclidean distance d is a metric space, i.e. an object of $\mathbf{GMet} = \mathbf{Met}$. The addition of real numbers is the most natural interpretation of $\Sigma = \{+ : 2\}$, thus we get a quantitative algebra $(\mathbb{R}, +, d)$.

Remark 3.3. Already here, we covered three examples that are not possible with the original (and predominant in the literature) definition of quantitative algebras [MPP16, Definition 3.1]. The first two are not possible because the base category is not \mathbf{Met} . The third is not possible even if it deals with metric spaces.

³³¹ We do not spend time making this precise, but it [post-rigorously](#) makes the case for universal quantitative algebra as a straightforward combination of universal algebra and generalized metric spaces.

³³² Note that \mathbf{GMet} is instantiated as $\mathbb{N}_\infty \mathbf{Spa}$, i.e. $\mathbb{L} = \mathbb{N}_\infty$ and $\hat{E}_{\mathbf{GMet}} = \emptyset$.

³³³ This time, \mathbf{GMet} is instantiated as \mathbf{Poset} with $\mathbb{L} = \mathbb{B}$ and $\hat{E}_{\mathbf{GMet}} = \hat{E}_{\mathbf{Poset}}$ as defined after Definition 2.32.

Indeed, as already noted in [Adá22, Remark 3.1.(2)], the addition of real numbers is not a nonexpansive function $(\mathbb{R}, d) \times (\mathbb{R}, d) \rightarrow (\mathbb{R}, d)$, where \times denotes the categorical product because,³³⁴ recalling Corollary 2.39, we have

$$(d \times d)((1, 1), (2, 2)) = \sup\{d(1, 2), d(1, 2)\} = 1 < 2 = d(2, 4) = d(1 + 1, 2 + 2).$$

Here are two more compelling examples from the original paper [MPP16].

Example 3.4 (Hausdorff). In Example 2.17, we defined the Hausdorff distance d^\dagger on $\mathcal{P}_{\text{ne}}X$ that depends on an L-relation $d : X \times X \rightarrow L$. In Example 1.78, we described a $\Sigma_{\mathcal{S}}$ -algebra structure on $\mathcal{P}_{\text{ne}}X$ (interpreting \oplus as union). Combining these, we get a quantitative $\Sigma_{\mathcal{S}}$ -algebra $(\mathcal{P}_{\text{ne}}X, \cup, d^\dagger)$ for any L-space (X, d) .

If we know that (X, d) satisfies some quantitative equations in $\hat{E}_{\mathbf{GMet}}$, we can sometimes prove that $(\mathcal{P}_{\text{ne}}X, d^\dagger)$ does too. For instance, picking $L = [0, 1]$ or $L = [0, \infty]$, $\mathbf{GMet} = \mathbf{Met}$, and $\hat{E}_{\mathbf{GMet}} = \hat{E}_{\mathbf{Met}}$, one can show that if (X, d) belongs to \mathbf{Met} , then so does $(\mathcal{P}_{\text{ne}}X, d^\dagger)$, and we still get a quantitative $\Sigma_{\mathcal{S}}$ -algebra $(\mathcal{P}_{\text{ne}}X, \cup, d^\dagger)$, now over \mathbf{Met} .³³⁵

Example 3.5 (Kantorovich). Given a L-relation $d : X \times X \rightarrow [0, 1]$, we define the **Kantorovich distance** d_K on $\mathcal{D}X$ as follows:³³⁶ for all $\varphi, \psi \in \mathcal{D}X$,

$$d_K(\varphi, \psi) = \inf \left\{ \sum_{(x, x')} \tau(x, x')d(x, x') \mid \tau \in \mathcal{D}(X \times X), \mathcal{D}\pi_1(\tau) = \varphi, \mathcal{D}\pi_2(\tau) = \psi \right\}.$$

The distributions τ above range over **couplings** of φ and ψ , i.e. distributions over $X \times X$ whose marginals are φ and ψ . Thus, what d_K does, in words, is computing the average distance according to all couplings, and then taking the smallest one.

In Example 1.79, we gave a $\Sigma_{\mathbf{CA}}$ -algebra structure on $\mathcal{D}X$ (interpreting $+_p$ as convex combination). Combining the algebra and the $[0, 1]$ -space, we get a quantitative $\Sigma_{\mathbf{CA}}$ -algebra $(\mathcal{D}X, \llbracket - \rrbracket_{\mathcal{D}X}, d_K)$. Once again, we can prove that if (X, d) is a metric space, then so is $(\mathcal{D}X, d_K)$, and we obtain a quantitative algebra $(\mathcal{D}X, \llbracket - \rrbracket_{\mathcal{D}X}, d_K)$ over \mathbf{Met} .³³⁷

Unlike the first examples, the interpretations in $(\mathcal{P}_{\text{ne}}X, \cup, d^\dagger)$ and $(\mathcal{D}X, \llbracket - \rrbracket_{\mathcal{D}X}, d_K)$ are nonexpansive with respect to the product distance. Concretely,

$$\forall S, S', T, T' \in \mathcal{P}_{\text{ne}}X, \quad d^\dagger(S \cup S', T \cup T') \leq \max \{d^\dagger(S, T), d^\dagger(S', T')\} \quad (3.1)$$

$$\forall \varphi, \varphi', \psi, \psi' \in \mathcal{D}X, \quad d_K(p\varphi + \bar{p}\varphi', p\psi + \bar{p}\psi') \leq \max \{d_K(\varphi, \psi), d_K(\varphi', \psi')\}. \quad (3.2)$$

The initial motivation to remove this requirement and arrive at Definition 3.1³³⁸ came from a variant of the Kantorovich distance called the **Łukaszyk–Karmowski** (ŁK for short) distance [Łuko4, Eq. (21)] which sends $\varphi, \psi \in \mathcal{D}X$ to

$$d_{\text{ŁK}}(\varphi, \psi) = \sum_{(x, x')} \varphi(x)\psi(x')d(x, x'). \quad (3.3)$$

In words, instead of looking at all the couplings to find the best one, we only look at the independent coupling $\tau(x, x') = \varphi(x)\psi(x')$.³³⁹ In particular, it coincides with

³³⁴ In [MPP16], the interpretation of an n -ary operation symbol is required to be a nonexpansive map from the n -wise product of the carrier to the carrier.

³³⁵ This is the quantitative algebra denoted by $\Pi[M]$ in [MPP16, Theorem 9.2].

³³⁶ This lifting of a distance on X to a distance on $\mathcal{D}X$ is well-known in optimal transport theory [Vil09]. You can find a well-written concise description of d_K in [BBKK18, §2.1] in the case $L = [0, \infty]$ where it is denoted d^{LD} . They also give a dual description as we did for the Hausdorff distance in Example 2.17, but the strong duality result ($d^{\text{LD}} = d^{\text{LD}}$) does not hold in general.

³³⁷ This is the quantitative algebra denoted by $\Pi[M]$ in [MPP16, Theorem 10.4].

³³⁸ Which imposes no further relation between the Σ -algebra and the L-space other than being on the same set.

³³⁹ The ŁK distance is easier to compute than the Kantorovich distance since there is no optimization. It is the reason why it was considered in [CKPR21] for an application to reinforcement learning.

the Kantorovich distance on Dirac distributions since the independent coupling of δ_x and δ_y is the only coupling, we obtain

$$d_K(\delta_x, \delta_y) = d_{\mathbb{L}K}(\delta_x, \delta_y) = d(x, y).$$

We can show that convex combination is not nonexpansive with respect to the product of the $\mathbb{L}K$ distance, namely, there exists a $[0, 1]$ -space (X, d) , distributions $\varphi, \varphi', \psi, \psi' \in \mathcal{D}X$, and $p \in (0, 1)$ such that

$$d_{\mathbb{L}K}(p\varphi + \bar{p}\varphi', p\psi + \bar{p}\psi') > \sup \{d_{\mathbb{L}K}(\varphi, \psi), d_{\mathbb{L}K}(\varphi', \psi')\}.$$

Take $X = \{x, y\}$ with $d(x, y) = d(y, x) = 1$ and the self-distances being 0,³⁴⁰ then for any $p \in (0, 1)$,

$$\begin{aligned} d_{\mathbb{L}K}(p\delta_x + \bar{p}\delta_y, p\delta_x + \bar{p}\delta_y) &= p^2d(x, x) + p\bar{p}d(x, y) + \bar{p}pd(y, x) + \bar{p}^2d(y, y) \\ &= 2p\bar{p} \\ &> 0 \\ &= \sup \{0, 0\} \\ &= \sup \{d_{\mathbb{L}K}(\delta_x, \delta_x), d_{\mathbb{L}K}(\delta_y, \delta_y)\}. \end{aligned}$$

Therefore, $(\mathcal{D}X, \llbracket - \rrbracket_{\mathcal{D}X}, d_{\mathbb{L}K})$ is always a quantitative algebra in the sense of Definition 3.1, but not always in the sense of [MPP16, Definition 3.1].³⁴¹

We now turn to subalgebras and products.

Definition 3.6 (Subalgebra). Given $\hat{\mathbb{A}} \in \mathbf{QAlg}(\Sigma)$, a **subalgebra** of $\hat{\mathbb{A}}$ is a quantitative algebra $\hat{\mathbb{B}}$ such that \mathbb{B} is a subalgebra of \mathbb{A} , and \mathbf{B} is a subspace of \mathbf{A} . Explicitly, it is a subset $B \subseteq A$ that is closed under the operations in Σ , namely, for any $\text{op} : n \in \Sigma$ and $b_1, \dots, b_n \in B$, $\llbracket \text{op} \rrbracket_A(b_1, \dots, b_n) \in B$, and it is equipped with an L-relation $d_{\mathbf{B}}$ satisfying $d_{\mathbf{B}}(b, b') = d_{\mathbf{A}}(b, b')$ for all $b, b' \in B \subseteq A$. It quickly follows that $\llbracket - \rrbracket_B : \Sigma(B) \rightarrow B$ can be defined as a (co)restriction of $\llbracket - \rrbracket_A$, making $\hat{\mathbb{B}} = (B, \llbracket - \rrbracket_B, d_{\mathbf{B}})$ into a quantitative Σ -algebra and the inclusion $B \hookrightarrow A$ into a nonexpansive homomorphism.³⁴²

Since both forgetful functors from $\mathbf{Alg}(\Sigma)$ and \mathbf{GMet} to \mathbf{Set} preserve products, we can easily infer that the categorical product of quantitative algebras is the product of the underlying algebras and spaces.

Lemma 3.7. Let $\{\hat{\mathbb{A}}_i = (A_i, \llbracket - \rrbracket_i, d_i) \mid i \in I\}$ be a family of quantitative algebras indexed by I . We define the quantitative algebra $\hat{\mathbb{A}} = (A, \llbracket - \rrbracket_A, d)$ with

$$\mathbb{A} = (A, \llbracket - \rrbracket_A) = \prod_{i \in I} \mathbb{A}_i \quad \text{and} \quad \mathbf{A} = (A, d) = \prod_{i \in I} \mathbf{A}_i.$$

Then $\hat{\mathbb{A}}$ is the product $\prod_{i \in I} \hat{\mathbb{A}}_i$ with $\pi_i : \hat{\mathbb{A}} \rightarrow \hat{\mathbb{A}}_i$ being the projection of the cartesian product.³⁴³

Proof. First, we recall from (the proofs of) Lemma 1.8 and Corollary 2.39 that the underlying set of the products of algebras and of spaces is the cartesian product of the

³⁴⁰ We gave another example in [MSV22, Lemma 5.3].

³⁴¹ In fact, even if d is a metric, $d_{\mathbb{L}K}$ is not a metric (by the example above, self-distances are not always 0, so it does not satisfy $x \vdash x =_0 x$). That is another reason why [MPP16] does not apply.

³⁴² Combining our intuitions from Remark 1.6 and Definition 2.43, we find that inclusions of subalgebras are, up to isomorphisms, precisely the isometric injective homomorphisms.

³⁴³ This follows more abstractly from [For22, Proposition 3.2.1(2)] when we see $\mathbf{QAlg}(\Sigma)$ as the pullback of $\mathbf{Alg}(\Sigma)$ and \mathbf{GMet} (with two forgetful functors to \mathbf{Set} that preserve products), and we use that $U : \mathbf{GMet} \rightarrow \mathbf{Set}$ is an isofibration (Proposition 2.49). From [For22, Proposition 3.2.1(1)], it also follows that $U : \mathbf{QAlg}(\Sigma) \rightarrow \mathbf{Alg}(\Sigma)$ is an isofibration.

underlying sets, hence $A = \prod_{i \in I} A_i$ is the carrier of both \mathbb{A} and \mathbf{A} , so $\hat{\mathbb{A}}$ belongs to $\mathbf{QAlg}(\Sigma)$. Moreover, we also showed that each π_i is both a homomorphism $\mathbb{A} \rightarrow \mathbb{A}_i$ and a nonexpansive map $\mathbf{A} \rightarrow \mathbf{A}_i$, thus they are all morphisms in $\mathbf{QAlg}(\Sigma)$.

If $f_i : \hat{\mathbb{X}} \rightarrow \hat{\mathbb{A}}_i$ is a family of nonexpansive homomorphisms, we saw in the proofs of Lemma 1.8 and Proposition 2.36 that the pairing of functions $\langle f_i \rangle_{i \in I}$ is a homomorphism $\hat{\mathbb{X}} \rightarrow \hat{\mathbb{A}}$ in $\mathbf{Alg}(\Sigma)$ and a nonexpansive map $\mathbf{X} \rightarrow \mathbf{A}$ in \mathbf{GMet} , and it satisfies $\pi_i \circ \langle f_i \rangle_{i \in I} = f_i$. It is unique because the forgetful functors are faithful. We conclude that $\hat{\mathbb{A}}$ is the product of the $\hat{\mathbb{A}}_i$ s. \square

Quantitative Equations

Now, in order to get back the expressiveness of the original framework, we need a way to impose this property of nonexpansiveness with respect to the product distance, and we also need a way to impose other properties like the fact that \oplus should be interpreted as a commutative operation. We achieve both things at once with the following definition.

Definition 3.8 (Quantitative Equation). A **quantitative equation** (over Σ and \mathbf{L}) is a tuple comprising an \mathbf{L} -space \mathbf{X} called the **context**,³⁴⁴ two terms $s, t \in \mathcal{T}_\Sigma \mathbf{X}$, and optionally a quantity $\varepsilon \in \mathbf{L}$. We write these as $\mathbf{X} \vdash s = t$ when no ε is given or $\mathbf{X} \vdash s =_\varepsilon t$ when it is given.

A quantitative algebra $\hat{\mathbb{A}}$ **satisfies** a quantitative equation³⁴⁵

- $\mathbf{X} \vdash s = t$ if for any nonexpansive assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$, $\llbracket s \rrbracket_A^{\hat{\iota}} = \llbracket t \rrbracket_A^{\hat{\iota}}$.
- $\mathbf{X} \vdash s =_\varepsilon t$ if for any nonexpansive assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$, $d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{\iota}}, \llbracket t \rrbracket_A^{\hat{\iota}}) \leq \varepsilon$.

We use ϕ and ψ to refer to a quantitative equation, and we sometimes simply call them equations. We write $\hat{\mathbb{A}} \models \phi$ when $\hat{\mathbb{A}}$ satisfies ϕ ,³⁴⁶ and we also write $\hat{\mathbb{A}} \models^{\hat{\iota}} \phi$ when the equality $\llbracket s \rrbracket_A^{\hat{\iota}} = \llbracket t \rrbracket_A^{\hat{\iota}}$ or the bound $d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{\iota}}, \llbracket t \rrbracket_A^{\hat{\iota}}) \leq \varepsilon$ holds for a particular assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$ (and not necessarily for all assignments).

Our overloading of the terminology *quantitative equation* (recall Definition 2.23) is practically harmless because a quantitative equation from Chapter 2 $\mathbf{X} \vdash x = y$ (or $\mathbf{X} \vdash x =_\varepsilon y$) can be seen as the new kind of quantitative equation by viewing x and y as terms via the embedding η_X^Σ . Formally, since $\llbracket \eta_X^\Sigma(x) \rrbracket_A^{\hat{\iota}} = \hat{\iota}(x)$ for any $x \in X$ and $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$,³⁴⁷

$$\begin{aligned} \mathbf{A} \models \mathbf{X} \vdash x = y &\iff \hat{\mathbb{A}} \models \mathbf{X} \vdash \eta_X^\Sigma(x) = \eta_X^\Sigma(y) \\ \mathbf{A} \models \mathbf{X} \vdash x =_\varepsilon y &\iff \hat{\mathbb{A}} \models \mathbf{X} \vdash \eta_X^\Sigma(x) =_\varepsilon \eta_X^\Sigma(y). \end{aligned} \quad (3.4)$$

In particular, since we assumed the underlying space of any $\hat{\mathbb{A}} \in \mathbf{QAlg}(\Sigma)$ to be a generalized metric space, we can say that $\hat{\mathbb{A}} \models \phi$ for any $\phi \in \hat{E}_{\mathbf{GMet}}$.³⁴⁸ Another consequence is that over the empty signature $\Sigma = \emptyset$, the quantitative equations from Definition 2.23 and Definition 3.8 are the same.

Furthermore, the new quantitative equations also generalize the equations of universal algebra (Definition 1.16). Indeed, given an equation $X \vdash s = t$, we construct

³⁴⁴ Note that even with algebras in \mathbf{GMet} , the context is in \mathbf{LSpa} . This differs slightly from [AFMS21, FMS21].

³⁴⁵ Formally, we would need to write $\llbracket - \rrbracket_A^{U\hat{\iota}}$ instead of $\llbracket - \rrbracket_A^{\hat{\iota}}$ because $U\hat{\iota} : X \rightarrow A$ is the assignment we use to interpret the terms.

³⁴⁶ As usual, satisfaction generalizes to classes of quantitative equations, i.e. if \hat{E} is a classes of quantitative equations, $\hat{\mathbb{A}} \models \hat{E}$ means $\hat{\mathbb{A}} \models \phi$ for all $\phi \in \hat{E}$.

³⁴⁷ Later on, we will seldom distinguish between x and $\eta_X^\Sigma(x)$ and write the former for simplicity.

³⁴⁸ We implicitly see the equations in $\hat{E}_{\mathbf{GMet}}$ as the new kind of equations from Definition 3.8.

the quantitative equation $\mathbf{X}_\top \vdash s = t$ where the new context is the discrete space on the old context. We show that

$$\mathbb{A} \models X \vdash s = t \iff \hat{\mathbb{A}} \models \mathbf{X}_\top \vdash s = t. \quad (3.5)$$

By Proposition 2.60, any assignment $\iota : X \rightarrow A$ is nonexpansive from \mathbf{X}_\top to \mathbf{A} . Any nonexpansive assignment $\hat{\iota} : \mathbf{X}_\top \rightarrow \mathbf{A}$ also yields an assignment $X \rightarrow A$ by applying the forgetful functor U since the carrier of \mathbf{X}_\top is X . Therefore, the interpretations of s and t coincide under all assignments if and only if they coincide under all nonexpansive assignments.

Remark 3.9. The name quantitative equation is already used in, e.g. [MPP16, MPP17, Adá22, ADV23b] for a fairly restricted subsets of what we call quantitative equation. They use it to refer to our quantitative equations with a quantity and a discrete context, and they call our unrestricted quantitative equations *basic quantitative inferences*. We believe the judgments of Definition 3.8 are a better generalization of equations in equational logic to the quantitative setting, hence we propose to call those quantitative equations rather than following the custom in the quantitative algebra literature.

It is hard to argue objectively for this choice since equations are so prevalent in mathematics, and hence the question of what a quantitative equation should be has many good answers. We can mention three arguments in favor of our terminology:

- The quantitative equations of this chapter are a straightforward combination of the classical equations (Definition 1.16) and the quantitative equations for L-spaces (Definition 2.23) as witnessed by (3.4) and (3.5). This might seem circular because we already called the judgments for L-spaces quantitative equations, but if you restrict these quantitative equations to those with a discrete context as in [MPP16], you get essentially four possibilities:³⁴⁹

$$x \vdash x = x \quad x \vdash x =_\varepsilon x \quad x, y \vdash x = y \quad x, y \vdash x =_\varepsilon y.$$

With only these options, you can only define L-spaces with global upper bound on distances or on self-distances, or with at most one point.

- In §3.3, we will introduce quantitative equational logic which greatly resembles equational logic, I hope Examples 3.70 and 3.71 will convince you of that. Now, restricting to only discrete contexts does not break this connection,³⁵⁰ but it is compelling that we can have an extremely similar logic even with stronger judgments.
- There is a straightforward generalization of abstract equations (Definition 1.50) that corresponds to quantitative equations as we define them (see Propositions 3.62 and 3.63 that mirror Propositions 1.51 and 1.52).

Let us get to more interesting examples of quantitative equations now.³⁵¹

Example 3.10 (Quasi-commutativity). Let $+: 2 \in \Sigma$ be a binary operation symbol. As shown above, to ensure $+$ is interpreted as a commutative operation in a quantitative

³⁴⁹ Written in syntactic sugar.

³⁵⁰ For example, the authors of [BV05] develop a *fuzzy equational logic* that is extremely similar to equational logic, and allows reasoning about fuzzy relations (basically distances but with a different attitude). Their judgments (implicitly) have discrete contexts only.

³⁵¹ More examples are in the papers we cited in the introduction when we talked about universal algebra on partial orders and on metric spaces. In particular, there is a long list in [AFMS21, Example 3.19], where **GMet** is instantiated as **Poset**.

algebra, we can use the quantitative equation $\mathbf{X}_\top \vdash x + y = y + x$ where $X = \{x, y\}$. In fact, using the same syntactic sugar as we did in Chapter 2 to avoid explicitly describing all the context, we can write $x, y \vdash x + y = y + x$ ³⁵² which looks exactly like the classical equation for commutativity.

Since the context can be any L-space, we can now add some nuance to the commutativity property. For instance, we can guarantee that $+$ is commutative only between elements that are close to each other with $x =_\varepsilon y \vdash x + y = y + x$ where $\varepsilon \in \mathbf{L}$ is fixed.³⁵³ Unrolling the syntactic sugar, the context is the L-space \mathbf{X} containing two points x and y with $d_{\mathbf{X}}(x, y) = \varepsilon$ and all other distances being \top . Therefore, a nonexpansive assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$ is a choice of two elements $\hat{\iota}(x)$ and $\hat{\iota}(y)$ with $d_{\mathbf{A}}(\hat{\iota}(x), \hat{\iota}(y)) \leq \varepsilon$, and no other constraint. We conclude that $\hat{\mathbf{A}}$ satisfies $x =_\varepsilon y \vdash x + y = y + x$ if and only if $\llbracket + \rrbracket_{\mathbf{A}}(a, b) = \llbracket + \rrbracket_{\mathbf{A}}(b, a)$ whenever $d_{\mathbf{A}}(a, b) \leq \varepsilon$.

Another possible variant on commutativity is $x =_\perp y \vdash x + y = y + x$. This means $+$ is guaranteed to be commutative only on elements which have a self-distance of \perp . For instance, in distributions with the LK distance, $d_{\text{LK}}(\varphi, \varphi) = 0$ only when the elements in the support of φ are all at distance 0 from each other. In particular, when d is a metric, $d_{\text{LK}}(\varphi, \varphi) = 0$ if and only if φ is a Dirac distribution. So that quantitative equation would ensure commutativity only on Dirac distributions.

Remark 3.11. Note that our syntactic sugar now allows terms that are not variables in the conclusion, but it does not allow them in the premises. This is in contrast with the quantitative inferences of [MPP16] as they allow arbitrary terms in the premises. Thus, when the signature is not empty, our quantitative equations cannot correspond to their quantitative inferences. The authors had realized the restriction to variables was valuable, and sometimes necessary.³⁵⁴ They call the restricted judgments basic quantitative inferences (they also require a finite set of premises). Following [MSV23, Lemma 8.4 and §9.1], one could prove that our quantitative equations are equivalent to quantitative inferences whose premises only contain variables.³⁵⁵

Example 3.12 (Nonexpansiveness).³⁵⁶ We can translate (3.1) and (3.2) into the following (family of) quantitative equations.

$$\forall \varepsilon, \varepsilon' \in \mathbf{L}, \quad x =_\varepsilon y, x' =_{\varepsilon'} y' \vdash x \oplus x' =_{\max\{\varepsilon, \varepsilon'\}} y \oplus y' \quad (3.6)$$

$$\forall \varepsilon, \varepsilon' \in \mathbf{L}, \quad x =_\varepsilon y, x' =_{\varepsilon'} y' \vdash x +_p x' =_{\max\{\varepsilon, \varepsilon'\}} y +_p y' \quad (3.7)$$

The quantitative algebra from Example 3.4 satisfies (3.6), and the one from Example 3.5 satisfies (3.7), but its variant with the LK distance does not satisfy (3.7).

In general, if we want an n -ary operation symbol $\text{op} \in \Sigma$ to be interpreted as a nonexpansive map $\mathbf{A}^n \rightarrow \mathbf{A}$, we can impose the equations³⁵⁷

$$\forall \{\varepsilon_i\}_{i \in I} \subseteq \mathbf{L}, \quad \{x_i =_{\varepsilon_i} y_i \mid 1 \leq i \leq n\} \vdash \text{op}(x_1, \dots, x_n) =_{\max_i \varepsilon_i} \text{op}(y_1, \dots, y_n). \quad (3.8)$$

Example 3.13 (L -nonexpansiveness). In most papers on quantitative algebras this property is called “nonexpansiveness of the operations”. In [MSV22], we remarked this can be ambiguous because one could consider a different distance on n -tuples of inputs rather than the product distance. We then presented quantitative algebras for *lifted signature* which can deal with more general operations.

³⁵² Whenever we will write $x_1, \dots, x_n \vdash s = t$, we will mean $\mathbf{X}_\top \vdash s = t$ where $X = \{x_1, \dots, x_n\}$, and similarly for $=_\varepsilon$.

³⁵³ This example comes from [ADV23a, Example 8.3].

³⁵⁴ Barr made a similar observation in [Bar92].

³⁵⁵ See also [ADV23a, Remark 8.20 and Construction 8.21].

³⁵⁶ A similar example is detailed, in the context of ordered algebras, in [AFMS21, Examples 3.19.(2) and 3.19.(3)]. They call algebras **coherent** when all operations are nonexpansive in this sense.

³⁵⁷ This is an axiom in the logic of [MPP16]. It is not in our formulation of quantitative equational logic. Thus, the algebras of Mardare et al. are all coherent in the terminology of [AFMS21, Definition 3.2].

In a lifted signature, each operation symbol $\text{op} : n \in \Sigma$ comes with an assignment $(A, d) \mapsto (A^n, L_{\text{op}}(d))$ (on generalized metric spaces) which specifies the distance $L_{\text{op}}(d)$ on n -tuples that needs to be considered. We say that the interpretation $\llbracket \text{op} \rrbracket_A$ is L_{op} -nonexpansive when it is a nonexpansive map $\llbracket \text{op} \rrbracket_A : (A^n, L_{\text{op}}(d)) \rightarrow (A, d)$.³⁵⁸ We can also express L_{op} -nonexpansiveness with a family of quantitative equations like we did in Example 3.12:³⁵⁹

$$\forall \mathbf{X} \in \mathbf{GMet}, \forall x, y \in X^n, \quad \mathbf{X} \vdash \text{op}(x_1, \dots, x_n) =_{L_{\text{op}}(d_{\mathbf{X}})(x, y)} \text{op}(y_1, \dots, y_n). \quad (3.9)$$

If an algebra $\hat{\mathbf{A}}$ satisfies these equations, then in particular, for all $a, b \in A^n$, it satisfies $\mathbf{A} \vdash \text{op}(a_1, \dots, a_n) =_{L_{\text{op}}(d_{\mathbf{A}})(a, b)} \text{op}(b_1, \dots, b_n)$ under the assignment $\text{id}_A : \mathbf{A} \rightarrow \mathbf{A}$. This means

$$d_{\mathbf{A}}(\llbracket \text{op} \rrbracket_A(a_1, \dots, a_n), \llbracket \text{op} \rrbracket_A(b_1, \dots, b_n)) \leq L_{\text{op}}(d_{\mathbf{A}})(a, b),$$

so we conclude that $\llbracket \text{op} \rrbracket_A : (A^n, L_{\text{op}}(d_{\mathbf{A}})) \rightarrow \mathbf{A}$ is nonexpansive.

Now, we still have to show that L_{op} -nonexpansiveness is the only consequence of (3.9). This requires an assumption on L_{op} that morally says the distance between two tuples x and y in $(X^n, L_{\text{op}}(d_{\mathbf{X}}))$ depends only on the distances between the coordinates x_1, \dots, x_n and y_1, \dots, y_n in \mathbf{X} .³⁶⁰ We refer to [MSV22] for more details, in particular Definitions 3.1 and 3.2 give the condition on L_{op} . Briefly, we need L_{op} to be a functor that preserves isometric embeddings.

As a particular case, one can take $L_{\text{op}}(d)$ to be the product distance and recover the original nonexpansiveness of Example 3.12. Another interesting instance is taking $L_{\text{op}}(d)$ to be the discrete distance (in case $\mathbf{GMet} = \mathbf{LSpa}$, $\forall x, y \in X^n, L_{\text{op}}(d)(x, y) = \top$), then (3.9) becomes trivial as we will see in Lemma 3.34. Intuitively, it is because any function from the discrete space on A^n to \mathbf{A} is nonexpansive.

Example 3.14 (Convexity). The quantitative algebra $(\mathcal{D}X, \llbracket - \rrbracket_{\mathcal{D}X}, d_K)$ satisfies another family of quantitative equations that is stronger than (3.7):³⁶¹

$$\forall \varepsilon, \varepsilon' \in \mathbf{L}, \quad x =_{\varepsilon} y, x' =_{\varepsilon'} y' \vdash x +_p x' =_{p\varepsilon + p\varepsilon'} y +_p y'. \quad (3.10)$$

This property of $\llbracket +_p \rrbracket_{\mathcal{D}X}$ is called convexity in e.g. [MV20, Definition 30].

As a sanity check for our definitions, we can verify that homomorphisms preserve the satisfaction of quantitative equations.³⁶²

Lemma 3.15. *Let ϕ be an equation with context \mathbf{X} . If $h : \hat{\mathbf{A}} \rightarrow \hat{\mathbf{B}}$ is a homomorphism and $\hat{\mathbf{A}} \models^{\hat{\iota}} \phi$ for an assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$, then $\hat{\mathbf{B}} \models^{h \circ \hat{\iota}} \phi$.*

Proof. We have two very similar cases. Let ϕ be the equation $\mathbf{X} \vdash s = t$, we have

$$\begin{aligned} \hat{\mathbf{A}} \models^{\hat{\iota}} \phi &\iff \llbracket s \rrbracket_{\hat{\mathbf{A}}}^{\hat{\iota}} = \llbracket t \rrbracket_{\hat{\mathbf{A}}}^{\hat{\iota}} && \text{definition of } \models \\ &\implies h(\llbracket s \rrbracket_{\hat{\mathbf{A}}}^{\hat{\iota}}) = h(\llbracket t \rrbracket_{\hat{\mathbf{A}}}^{\hat{\iota}}) \\ &\implies \llbracket s \rrbracket_{\hat{\mathbf{B}}}^{h \circ \hat{\iota}} = \llbracket t \rrbracket_{\hat{\mathbf{B}}}^{h \circ \hat{\iota}} && \text{by (1.12)} \\ &\iff \hat{\mathbf{B}} \models^{h \circ \hat{\iota}} \phi. && \text{definition of } \models \end{aligned}$$

³⁵⁸ See [MSV22, Definitions 3.4 and 3.6].

³⁵⁹ This is the L -NE rule of [MSV22, Definition 3.11], but it has been written more cleanly with quantitative equations with contexts.

³⁶⁰ This is the case for nonexpansiveness with respect to the product distance. In fact, the only distances that matter there are the pairwise $d_{\mathbf{X}}(x_i, y_i)$ for all i . For L_{op} -nonexpansiveness, the other distances like $d_{\mathbf{X}}(x_1, x_1)$ or $d_{\mathbf{X}}(y_3, x_1)$ may be important, but never $d_{\mathbf{X}}(x, z)$ for some fresh z .

³⁶¹ Instead of taking the maximum between ε and ε' , we take their convex combination, and since the former is always larger than the latter, (3.10) is stronger than (3.7).

³⁶² Just like we did in Lemma 1.20 for \mathbf{Set} and Lemma 2.37 for \mathbf{LSpa} . In fact, the proofs are very similar.

Let ϕ be the equation $\mathbf{X} \vdash s =_\varepsilon t$, we have

$$\begin{aligned}
\hat{\mathbf{A}} \models^{\hat{t}} \phi &\iff d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{t}}, \llbracket t \rrbracket_A^{\hat{t}}) \leq \varepsilon && \text{definition of } \models \\
&\implies d_{\mathbf{A}}(h(\llbracket s \rrbracket_A^{\hat{t}}), h(\llbracket t \rrbracket_A^{\hat{t}})) \leq \varepsilon \\
&\implies d_{\mathbf{A}}(\llbracket s \rrbracket_B^{h\circ\hat{t}}, \llbracket t \rrbracket_B^{h\circ\hat{t}}) \leq \varepsilon && \text{by (1.12)} \\
&\iff \hat{\mathbf{B}} \models^{h\circ\hat{t}} \phi. && \text{definition of } \models \quad \square
\end{aligned}$$

Definition 3.16 (Quantitative variety). Given a class \hat{E} of quantitative equations, a (Σ, \hat{E}) -**algebra** is a quantitative Σ -algebra that satisfies \hat{E} . We define $\mathbf{QAlg}(\Sigma, \hat{E})$, the category of (Σ, \hat{E}) -algebras, to be the full subcategory of $\mathbf{QAlg}(\Sigma)$ containing only those algebras that satisfy \hat{E} . A **quantitative variety** is a category equal to $\mathbf{QAlg}(\Sigma, \hat{E})$ for some class of quantitative equations \hat{E} .

There are many forgetful functors obtained by composing the forgetful functors from $\mathbf{QAlg}(\Sigma)$ with the inclusion functor $\mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{QAlg}(\Sigma)$:

- $U : \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{Set} = \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{QAlg}(\Sigma) \xrightarrow{U} \mathbf{Set}$
- $U : \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{Alg}(\Sigma) = \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{QAlg}(\Sigma) \xrightarrow{U} \mathbf{Alg}(\Sigma)$
- $U : \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{GMet} = \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{QAlg}(\Sigma) \xrightarrow{U} \mathbf{GMet}$
- $U : \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{LSpa} = \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{QAlg}(\Sigma) \xrightarrow{U} \mathbf{LSpa}$

Remark 3.17. Compared to the usage of the term *variety* in the literature (e.g. [MPP17, Adá22, ADV23b]), our quantitative varieties are more general, even when $\mathbf{GMet} = \mathbf{Met}$. First, we do not constrain our operations to be interpreted as nonexpansive maps from the product as the other authors do. Second, we do not restrict the size of the context of the equations in \hat{E} as is done in loc. cit.³⁶³

Example 3.18. 1. With $\Sigma = \{p : 0\}$, we now have a lot more quantitative varieties than we had varieties in Example 1.27. Even restricting to a discrete context, we have the following quantitative equations where ε ranges over L :³⁶⁴

$$\begin{array}{cccccc}
\vdash p = p & x \vdash x = x & x \vdash p = x & & x, y \vdash x = y & \\
\vdash p =_\varepsilon p & x \vdash x =_\varepsilon x & x \vdash p =_\varepsilon x & x \vdash x =_\varepsilon p & x, y \vdash x =_\varepsilon y &
\end{array}$$

The meaning of the first row does not change from Example 1.27, and the meaning of the second row can be inferred by replacing equality between terms with distance between terms. For example, $\vdash p =_\varepsilon p$ says that the self-distance of the interpretation of the constant p is at most ε . Classifying the quantitative varieties for this signature would require a lot more work than for the classical varieties.³⁶⁵

³⁶³ Their restrictions are subtler than just putting an upper bound on the cardinality of the underlying set of the context.

³⁶⁴ The first row comes from the classical case, and the second row replaces equality with equality up to ε ($=_\varepsilon$). The only difference being that $p =_\varepsilon x$ and $x =_\varepsilon p$ are not equivalent, so we need two distinct equations.

³⁶⁵ Although I think it is feasible, tedious but feasible.

2. When $\Sigma = \emptyset$, we mentioned that the quantitative equations are those of Chapter 2, so $\mathbf{QAlg}(\emptyset, \hat{E})$ is the subcategory of L-spaces that satisfy \hat{E} . In particular, the category \mathbf{GMet} is a quantitative variety as it equals $\mathbf{QAlg}(\emptyset, \hat{E}_{\mathbf{GMet}})$.
3. If \hat{E} contains the equations in $E_{\mathbf{CA}}$ and the equations in (3.10), then $\mathbf{QAlg}(\Sigma_{\mathbf{CA}}, \hat{E})$ is the category of convex algebras equipped with a convex metric [MV20, Definition 30] and nonexpansive homomorphisms.

We will not prove a generalization of Birkhoff's variety theorem (Theorem 1.29) for quantitative varieties, but we can prove one direction of it.³⁶⁶ We first have a result combining Lemmas 1.21 and 2.57.

Lemma 3.19. *Let ϕ be a quantitative equation and $h : \hat{\mathbf{A}} \rightarrow \hat{\mathbf{B}}$ be a homomorphism such that $Uh : \mathbf{A} \rightarrow \mathbf{B}$ is a split epimorphism, then $\hat{\mathbf{A}} \models \phi$ implies $\hat{\mathbf{B}} \models \phi$.³⁶⁷*

Proof. Let $h^{-1} : \mathbf{B} \rightarrow \mathbf{A}$ be the right inverse of Uh . It is not necessarily a homomorphism,³⁶⁸ but it is a nonexpansive map. Therefore, with \mathbf{X} being the context of ϕ , any assignment $\hat{t} : \mathbf{X} \rightarrow \mathbf{B}$ can be composed with h^{-1} to get a nonexpansive assignment $h^{-1} \circ \hat{t} : \mathbf{X} \rightarrow \mathbf{A}$. By hypothesis, $\hat{\mathbf{A}} \models^{h^{-1} \circ \hat{t}} \phi$ which implies $\hat{\mathbf{B}} \models^{\hat{t}} \phi$ by Lemma 3.15. By construction, $h \circ h^{-1} = \text{id}_{\mathbf{B}}$, so we conclude that \mathbf{B} satisfies ϕ . \square

Next, we prove a result combining Lemmas 1.22 and 2.45.

Lemma 3.20. *Let ϕ be a quantitative equation and $h : \hat{\mathbf{A}} \rightarrow \hat{\mathbf{B}}$ be a homomorphism such that $Uh : \mathbf{A} \rightarrow \mathbf{B}$ is an isometric embedding, then $\hat{\mathbf{B}} \models \phi$ implies $\hat{\mathbf{A}} \models \phi$.³⁶⁹*

Proof. With \mathbf{X} being the context of ϕ , any assignment $\hat{t} : \mathbf{X} \rightarrow \mathbf{A}$ can be composed with h to get a nonexpansive assignment $h \circ \hat{t} : \mathbf{X} \rightarrow \mathbf{B}$. By hypothesis, $\hat{\mathbf{B}} \models^{h \circ \hat{t}} \phi$, and we resolve the two similar cases using the fact that $\llbracket - \rrbracket_{\mathbf{B}}^{h \circ \hat{t}} = h(\llbracket - \rrbracket_{\mathbf{A}}^{\hat{t}})$ (1.12).

If $\phi = \mathbf{X} \vdash s = t$, then we have $h(\llbracket s \rrbracket_{\mathbf{A}}^{\hat{t}}) = h(\llbracket t \rrbracket_{\mathbf{A}}^{\hat{t}})$ by satisfaction in $\hat{\mathbf{B}}$ and (1.12). Since h is injective, we conclude that $\llbracket s \rrbracket_{\mathbf{A}}^{\hat{t}} = \llbracket t \rrbracket_{\mathbf{A}}^{\hat{t}}$.

If $\phi = \mathbf{X} \vdash s =_{\varepsilon} t$, then we have $d_{\mathbf{B}}(h(\llbracket s \rrbracket_{\mathbf{A}}^{\hat{t}}), h(\llbracket t \rrbracket_{\mathbf{A}}^{\hat{t}})) \leq \varepsilon$ by satisfaction in $\hat{\mathbf{B}}$ and (1.12). Since h is an isometry, we conclude that $d_{\mathbf{A}}(\llbracket s \rrbracket_{\mathbf{A}}^{\hat{t}}, \llbracket t \rrbracket_{\mathbf{A}}^{\hat{t}}) \leq \varepsilon$.

This works for all assignments, so we get $\hat{\mathbf{A}} \models \phi$. \square

Finally, we prove that products of quantitative algebras inside a quantitative variety are also computed by taking the product of underlying algebras and spaces (combining Lemma 1.8 and Proposition 2.38).³⁷⁰

Lemma 3.21. *Let ϕ be a quantitative equation with context \mathbf{X} , $\{\hat{\mathbf{A}}_i = (A_i, \llbracket - \rrbracket_i, d_i) \mid i \in I\}$ be a family of quantitative algebras indexed by I , and $\hat{\mathbf{A}} = \prod_{i \in I} \hat{\mathbf{A}}_i$ be their product as described in Lemma 3.7. For any assignment $\hat{t} : \mathbf{X} \rightarrow \mathbf{A}$,*

$$\hat{\mathbf{A}} \models^{\hat{t}} \phi \Leftrightarrow \forall i \in I, \hat{\mathbf{A}}_i \models^{\pi_i \circ \hat{t}} \phi. \quad (3.11)$$

Consequently, if every $\hat{\mathbf{A}}_i$ satisfies ϕ , then so does $\hat{\mathbf{A}}$.³⁷¹

Proof. Because each π_i is a homomorphism, we can use Lemma 3.15 for the forward direction (\Rightarrow). For the converse (\Leftarrow), let us prove two similar cases.

If $\phi = \mathbf{X} \vdash s = t$, then we have $\pi_i(\llbracket s \rrbracket_{\hat{\mathbf{A}}}^{\hat{t}}) = \pi_i(\llbracket t \rrbracket_{\hat{\mathbf{A}}}^{\hat{t}})$ by satisfaction under each $\pi_i \circ \hat{t}$ and by (1.12). This means that the interpretations under \hat{t} of s and t agree on all coordinates, hence they must coincide, i.e. $\hat{\mathbf{A}} \models^{\hat{t}} \phi$.

If $\phi = \mathbf{X} \vdash s =_{\varepsilon} t$, then we have $d_i(\pi_i(\llbracket s \rrbracket_{\hat{\mathbf{A}}}^{\hat{t}}), \pi_i(\llbracket t \rrbracket_{\hat{\mathbf{A}}}^{\hat{t}})) \leq \varepsilon$ by satisfaction under each $\pi_i \circ \hat{t}$ and by (1.12). By definition of $d_{\hat{\mathbf{A}}}$ as a supremum of the d_i s, this means $d_{\hat{\mathbf{A}}}(\llbracket s \rrbracket_{\hat{\mathbf{A}}}^{\hat{t}}, \llbracket t \rrbracket_{\hat{\mathbf{A}}}^{\hat{t}}) \leq \varepsilon$, i.e. $\hat{\mathbf{A}} \models^{\hat{t}} \phi$. \square

Combining these last three results yields one direction for a possible characterization of quantitative varieties. We introduce some terminology first.

³⁶⁶ We give a proof sketch of the converse for an empty signature in Theorem 3.65.

³⁶⁷ c.f. [MSV23, Lemma 6.2].

³⁶⁸ c.f. Footnote 324 which applies only when $\Sigma = \emptyset$.

³⁶⁹ Up to isomorphism, $\hat{\mathbf{A}}$ is a subalgebra of $\hat{\mathbf{B}}$.

³⁷⁰ i.e. quantitative varieties are closed under products.

³⁷¹ This readily follows from (3.11) because for any nonexpansive assignment $\hat{t} : \mathbf{X} \rightarrow \mathbf{A}$, every $\pi_i \circ \hat{t}$ is a nonexpansive assignment $\mathbf{X} \rightarrow \mathbf{A}_i$. Then, by hypothesis $\hat{\mathbf{A}}_i \models^{\pi_i \circ \hat{t}} \phi$ holds for every $i \in I$, and we conclude that $\hat{\mathbf{A}}$ satisfies ϕ by (3.11).

Definition 3.22 (Reflexive homomorphism). A homomorphism $h : \hat{\mathbf{A}} \rightarrow \hat{\mathbf{B}}$ is called **reflexive** if its underlying nonexpansive map $h : \mathbf{A} \rightarrow \mathbf{B}$ is a split epimorphism. Equivalently, for any subspace $\mathbf{B}' \subseteq \mathbf{B}$, there is a subspace $\mathbf{A}' \subseteq \mathbf{A}$ such that $h(\mathbf{A}') = \mathbf{B}'$ and the (co)restriction $h : \mathbf{A}' \rightarrow \mathbf{B}'$ is an isomorphism.³⁷² In particular, h is surjective (take $\mathbf{B}' = \mathbf{B}$), and we call $\hat{\mathbf{B}}$ a **reflexive homomorphic image** of $\hat{\mathbf{A}}$.

When the signature is empty, reflexive homomorphisms are just split epimorphisms, so as we argued in Footnote 303, a reflexive homomorphic image is always a subalgebra. This is not true in general because while h has a right inverse in \mathbf{GMet} , nothing guarantees it belongs to $\mathbf{QAlg}(\Sigma)$, i.e. that it is a homomorphism. Still, the following theorem generalizing one direction of Birkhoff's variety theorem (Theorem 1.29) also generalizes Theorem 2.59.

Theorem 3.23. *For any class of quantitative equations \hat{E} , the category $\mathbf{QAlg}(\Sigma, \hat{E})$ is closed under reflexive homomorphic images, subalgebras, and products.³⁷³*

Remark 3.24. There are already variety theorems for quantitative varieties in the original setting of Mardare et al. [MPP17, Theorem 3.11], and with arbitrary (not nonexpansive) operations [JMU24, Theorem 4.16], but they both put some size conditions on the context of quantitative equations. With this limitation, they can characterize some collections of quantitative varieties as the subcategories of $\mathbf{QAlg}(\Sigma)$ that are closed under c -reflexive homomorphic images, subalgebras, and products. This suggests that the converse of Theorem 3.23 should hold, but some work is still required.

Definition 3.25 (Quantitative algebraic theory). Given a class \hat{E} of quantitative equations over Σ and L , the **quantitative algebraic theory** generated by \hat{E} , denoted by $\Omega\mathfrak{Th}(\hat{E})$, is the class of quantitative equations that are satisfied in all (Σ, \hat{E}) -algebras:³⁷⁴

$$\Omega\mathfrak{Th}(\hat{E}) = \{ \phi \mid \forall \hat{\mathbf{A}} \in \mathbf{QAlg}(\Sigma, \hat{E}), \hat{\mathbf{A}} \models \phi \}.$$

Equivalently, $\Omega\mathfrak{Th}(\hat{E})$ contains the equations that are semantically entailed by \hat{E} ,³⁷⁵ namely $\phi \in \Omega\mathfrak{Th}(\hat{E})$ if and only if

$$\forall \hat{\mathbf{A}} \in \mathbf{QAlg}(\Sigma), \hat{\mathbf{A}} \models \hat{E} \implies \hat{\mathbf{A}} \models \phi. \quad (3.12)$$

We will see in §3.3 how to find which quantitative equations are entailed by others.

We call a class of quantitative equations a quantitative algebraic theory if it is generated by some class \hat{E} .

We will see twice³⁷⁶ that the algebraic reasoning we are used to from Chapter 1 is embedded in quantitative algebraic reasoning. In particular, Example 1.31 which showed some equations which belong to the algebraic theory of commutative monoids can be read *unchanged* to find quantitative equations that belong to the quantitative algebraic theory of commutative monoids. Let us give another example that deals with quantities.

Example 3.26. We mentioned in Example 3.14 that the equations for convexity (3.10) are *stronger* than the equations for nonexpansiveness with respect to the

³⁷² In [MPP17], the authors introduced c -reflexive homomorphisms parametrized by a cardinal c . The definition changes by restricting the quantification of subspaces \mathbf{B}' to those with cardinality smaller than c . One easily checks that reflexive homomorphisms are precisely the homomorphisms that are c -reflexive for every c .

³⁷³ This quickly follows from Lemmas 3.19–3.21.

³⁷⁴ Again $\Omega\mathfrak{Th}(\hat{E})$ is never a set (recall Definition 1.30).

³⁷⁵ As in the classical case, $\Omega\mathfrak{Th}(\hat{E})$ contains all of \hat{E} but also many more equations like $x \vdash x = x$ or $x =_{\varepsilon} y \vdash x =_{\varepsilon} y$. Furthermore, $\Omega\mathfrak{Th}(\hat{E})$ contains all the quantitative equations in $\hat{E}_{\mathbf{GMet}}$ because the underlying spaces of algebras in $\mathbf{QAlg}(\Sigma, \hat{E})$ belong to \mathbf{GMet} .

³⁷⁶ In Examples 3.70 and 3.71.

product distance (3.7). Formally what this means is that if \hat{E} contains (3.10), then the interpretation of $+_p$ in a $(\Sigma_{\mathbf{CA}}, \hat{E})$ -algebra $\hat{\mathbf{A}}$ will be a nonexpansive map $\mathbf{A} \times \mathbf{A} \rightarrow \mathbf{A}$, hence $\hat{\mathbf{A}}$ will satisfy (3.7). Concisely, the equations of (3.7) belong to $\mathcal{Q}\mathfrak{Th}(\hat{E})$.

3.2 Free Quantitative Algebras

We turn to the construction of free algebras, and we start with a simple example.

Example 3.27 (Free metric). We already have some intuitions about terms and equations from Example 1.32, thus we consider an empty signature in order to focus on the new contexts and quantities. For \hat{E} , let us take the set of equations defining a metric space (with $\mathbf{L} = [0, 1]$),³⁷⁷ so that $\mathbf{QAlg}(\emptyset, \hat{E}) = \mathbf{Met}$.

Now we wonder, given an \mathbf{L} -space \mathbf{X} , what is the free metric space on it? Rehashing Definition 1.47, we want to find a metric space \mathbf{FX} and a nonexpansive map $\eta : \mathbf{X} \rightarrow \mathbf{FX}$ such that any nonexpansive map from \mathbf{X} to a metric space \mathbf{A} factors through η uniquely. Of course, if \mathbf{X} is already a metric space, then taking $\mathbf{FX} = \mathbf{X}$ and $\eta = \text{id}_{\mathbf{X}}$ works. Otherwise, we can look at what prevents $d_{\mathbf{X}}$ from being a metric.

For instance, if \mathbf{X} does not satisfy $\vdash x =_0 x$, it means there is some $x \in X$ such that $d_{\mathbf{X}}(x, x) > 0$. Inside \mathbf{FX} , we know that the distance between $\eta(x)$ and $\eta(x)$ must be 0. Note that if \mathbf{A} is a metric space and $f : \mathbf{X} \rightarrow \mathbf{A}$ is nonexpansive, we know that $d_{\mathbf{A}}(f(x), f(x)) = 0$ too, so sending $\eta(x)$ to $f(x)$ will not be a problem.

For a second example, suppose $d_{\mathbf{X}}$ is not symmetric, without loss of generality $d_{\mathbf{X}}(x, y) < d_{\mathbf{X}}(y, x)$ for some $x, y \in X$. We know that $d_{\mathbf{FX}}(\eta(x), \eta(y)) = d_{\mathbf{FX}}(\eta(y), \eta(x))$, but what value should it be? To ensure that η is nonexpansive, this value must be at most $d_{\mathbf{X}}(x, y)$, but why not smaller? If this lack of symmetry is the only thing preventing $d_{\mathbf{X}}$ from being a metric (i.e. defining d' everywhere like $d_{\mathbf{X}}$ except $d'(x, y) = d'(y, x)$ yields a metric), we cannot make $d_{\mathbf{FX}}(x, y)$ smaller, because the identity function $\text{id}_{\mathbf{X}}$ would be a nonexpansive map $\mathbf{X} \rightarrow (X, d')$ that does not factor through η (since $d'(x, y) > d_{\mathbf{FX}}(\eta(x), \eta(y))$). In fact, you can check that $\mathbf{FX} = (X, d')$ with $\eta = \text{id}_{\mathbf{X}}$ is the free metric space on \mathbf{X} because our definition of d' fixed the only problem with $d_{\mathbf{X}}$.

In general, for any $x, y \in X$, we want $d_{\mathbf{FX}}(\eta(x), \eta(y))$ to be as large as possible while guaranteeing that $d_{\mathbf{FX}}$ is a metric and η is nonexpansive, but it is not always that simple. The complexity comes from the possible interactions between different equations in \hat{E} . Say you have $d_{\mathbf{X}}(x, z) > d_{\mathbf{X}}(x, y) + d_{\mathbf{X}}(y, z)$ so the triangle inequality does not hold, hence you try to fix this by lowering $d_{\mathbf{FX}}(\eta x, \eta z)$ down exactly to $d_{\mathbf{FX}}(\eta x, \eta y) + d_{\mathbf{FX}}(\eta y, \eta z)$.³⁷⁸ Then, to ensure symmetry, you need to lower $d_{\mathbf{FX}}(z, x)$ down to that same value, but after that you may need to lower $d_{\mathbf{FX}}(x, y)$ so that it is not bigger than the new value of $d_{\mathbf{FX}}(y, z) + d_{\mathbf{FX}}(z, x)$. In the end, you can end up back with $d_{\mathbf{FX}}(x, z) > d_{\mathbf{FX}}(x, y) + d_{\mathbf{FX}}(y, z)$, so you have to do another round of fixes.

Intuitively, \mathbf{FX} is the space you obtain by iterating this process (possibly for infinitely many steps) and looking at the limit. We will give a rigorous description in the case of a more general signature,³⁷⁹ but we want to point out now that this

³⁷⁷ As a reminder, \hat{E} contains

$$\begin{aligned} \forall \varepsilon \in [0, 1], \quad & y =_{\varepsilon} x \vdash x =_{\varepsilon} y \\ & \vdash x =_0 x \\ & x =_0 y \vdash x = y \\ \forall \varepsilon, \delta \in [0, 1], \quad & x =_{\varepsilon} y, y =_{\delta} z \vdash x =_{\varepsilon+\delta} z. \end{aligned}$$

³⁷⁸ Let us not write η each time for better readability, this is a bit informal as we will see that η is not necessarily injective.

³⁷⁹ This is the construction of free quantitative algebras that starts in the next paragraph.

process does not deal only with distances, it can also force some equations. For example, if $d_X(x, y) = 0$ with $x \neq y$ at the start, you will end up with $\eta(x) = \eta(y)$ inside FX .

Fix a class \hat{E} of quantitative equations over Σ and L . For any generalized metric space \mathbf{X} , we can define a binary relation $\equiv_{\hat{E}}$ and an L-relation $d_{\hat{E}}$ on Σ -terms as follows:³⁸⁰ for any $s, t \in \mathcal{T}_{\Sigma}X$,

$$s \equiv_{\hat{E}} t \iff \mathbf{X} \vdash s = t \in \mathcal{Q}\mathfrak{Th}(\hat{E}) \text{ and } d_{\hat{E}}(s, t) = \inf\{\varepsilon \mid \mathbf{X} \vdash s =_{\varepsilon} t \in \mathcal{Q}\mathfrak{Th}(\hat{E})\}. \quad (3.13)$$

The definition of $\equiv_{\hat{E}}$ is completely analogous to what we did in the classical case (1.24). The definition of $d_{\hat{E}}$ is new but it also looks like how we defined an L-relation from an L-structure in Proposition 2.21. In fact, we can also prove a counterpart to (2.8), giving us an equivalent definition of $d_{\hat{E}}$: for any $s, t \in \mathcal{T}_{\Sigma}X$ and $\varepsilon \in L$,³⁸¹

$$d_{\hat{E}}(s, t) \leq \varepsilon \iff \mathbf{X} \vdash s =_{\varepsilon} t \in \mathcal{Q}\mathfrak{Th}(\hat{E}). \quad (3.14)$$

Proof of (3.14). (\Leftarrow) holds directly by definition of infimum. For (\Rightarrow), we need to show that any (Σ, \hat{E}) -algebra satisfies $\mathbf{X} \vdash s =_{\varepsilon} t$. Let $\hat{A} \in \mathbf{QAlg}(\Sigma, \hat{E})$ and $\hat{t}: \mathbf{X} \rightarrow \mathbf{A}$ be a nonexpansive assignment. We know that for every δ such that $\mathbf{X} \vdash s =_{\delta} t \in \mathcal{Q}\mathfrak{Th}(\hat{E})$, $d_{\mathbf{A}}(\llbracket s \rrbracket_{\hat{A}}, \llbracket t \rrbracket_{\hat{A}}) \leq \delta$, thus

$$d_{\mathbf{A}}(\llbracket s \rrbracket_{\hat{A}}, \llbracket t \rrbracket_{\hat{A}}) \leq \inf\{\delta \mid \mathbf{X} \vdash s =_{\delta} t \in \mathcal{Q}\mathfrak{Th}(\hat{E})\} = d_{\hat{E}}(s, t) \leq \varepsilon.$$

We conclude that $\hat{A} \models^{\hat{t}} \mathbf{X} \vdash s =_{\varepsilon} t$, and we are done since \hat{A} and \hat{t} were arbitrary. \square

When we were not dealing with distances, we only had to prove that the relation \equiv_E defined between terms was a congruence (Lemma 1.33), and then we were able to construct the term algebra by quotienting the set of terms and interpreting the operation symbols syntactically. Here we have to prove a bit more, namely that $d_{\hat{E}}$ is invariant under $\equiv_{\hat{E}}$ so the L-relation restricts to the quotient, and that the resulting L-space is a generalized metric space.

Let us decompose this in several small lemmas. We also collect here some more lemmas that look similar, many of which will be part of the proof of soundness when we introduce quantitative equational logic.³⁸² Let $\mathbf{X} \in \mathbf{LSpa}$ and $\hat{A} \in \mathbf{QAlg}(\Sigma)$ be universally quantified in all these lemmas.

First, Lemmas 3.28–3.31 say that $\equiv_{\hat{E}}$ is an equivalence relation and a congruence.³⁸³

Lemma 3.28. *For any $t \in \mathcal{T}_{\Sigma}X$, \hat{A} satisfies $\mathbf{X} \vdash t = t$.*

Proof. Obviously, $\llbracket t \rrbracket_{\hat{A}} = \llbracket t \rrbracket_{\hat{A}}$ holds for all $\hat{t}: \mathbf{X} \rightarrow \mathbf{A}$. \square

Lemma 3.29. *For any $s, t \in \mathcal{T}_{\Sigma}X$, if \hat{A} satisfies $\mathbf{X} \vdash s = t$, then \hat{A} satisfies $\mathbf{X} \vdash t = s$.*

Proof. If $\llbracket s \rrbracket_{\hat{A}} = \llbracket t \rrbracket_{\hat{A}}$ holds for all \hat{t} , then $\llbracket t \rrbracket_{\hat{A}} = \llbracket s \rrbracket_{\hat{A}}$ holds too. \square

Lemma 3.30. *For any $s, t, u \in \mathcal{T}_{\Sigma}X$, if \hat{A} satisfies $\mathbf{X} \vdash s = t$ and $\mathbf{X} \vdash t = u$, then \hat{A} satisfies $\mathbf{X} \vdash s = u$.*

³⁸⁰ The notation for $\equiv_{\hat{E}}$ and $d_{\hat{E}}$ should really depend on the space \mathbf{X} , but we prefer to omit this for better readability.

³⁸¹ In words, $d_{\hat{E}}$ assigns a distance below ε to s and t if and only if their interpretations in each (Σ, \hat{E}) -algebras are always at a distance below ε .

³⁸² We were less explicit back then, but that is what happened with Lemma 1.33 and soundness of equational logic.

³⁸³ The proofs are exactly the same as for Lemma 1.33 because $\equiv_{\hat{E}}$ does not involve distances.

Proof. If $\llbracket s \rrbracket_A^{\hat{l}} = \llbracket t \rrbracket_A^{\hat{l}}$ and $\llbracket t \rrbracket_A^{\hat{l}} = \llbracket u \rrbracket_A^{\hat{l}}$ holds for all \hat{l} , then $\llbracket s \rrbracket_A^{\hat{l}} = \llbracket u \rrbracket_A^{\hat{l}}$ holds too. \square

Lemma 3.31. *For any $\text{op} : n \in \Sigma$, $s_1, \dots, s_n, t_1, \dots, t_n \in \mathcal{T}_\Sigma X$, if $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash s_i = t_i$ for all $1 \leq i \leq n$, then $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash \text{op}(s_1, \dots, s_n) = \text{op}(t_1, \dots, t_n)$.*

Proof. For any assignment $\hat{l} : \mathbf{X} \rightarrow \mathbf{A}$, we have $\llbracket s_i \rrbracket_A^{\hat{l}} = \llbracket t_i \rrbracket_A^{\hat{l}}$ for all i . Hence,

$$\begin{aligned} \llbracket \text{op}(s_1, \dots, s_n) \rrbracket_A^{\hat{l}} &= \llbracket \text{op} \rrbracket_A(\llbracket s_1 \rrbracket_A^{\hat{l}}, \dots, \llbracket s_n \rrbracket_A^{\hat{l}}) && \text{by (1.9)} \\ &= \llbracket \text{op} \rrbracket_A(\llbracket t_1 \rrbracket_A^{\hat{l}}, \dots, \llbracket t_n \rrbracket_A^{\hat{l}}) && \forall i, \llbracket s_i \rrbracket_A^{\hat{l}} = \llbracket t_i \rrbracket_A^{\hat{l}} \\ &= \llbracket \text{op}(s_1, \dots, s_n) \rrbracket_A^{\hat{l}}. && \text{by (1.9)} \end{aligned} \quad \square$$

Lemmas 3.32 and 3.33 mean that $d_{\hat{\mathbf{E}}}$ is well-defined on equivalence classes of $\equiv_{\hat{\mathbf{E}}}$, namely, $d_{\hat{\mathbf{E}}}(s, t) = d_{\hat{\mathbf{E}}}(s', t')$ whenever $s \equiv_{\hat{\mathbf{E}}} s'$ and $t \equiv_{\hat{\mathbf{E}}} t'$.³⁸⁴

Lemma 3.32. *For any $s, t, t' \in \mathcal{T}_\Sigma X$ and $\varepsilon \in \mathbf{L}$, if $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash s =_\varepsilon t$ and $\mathbf{X} \vdash t = t'$, then $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash s =_\varepsilon t'$.*

Proof. For any $\hat{l} : \mathbf{X} \rightarrow \mathbf{A}$, we have $d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{l}}, \llbracket t \rrbracket_A^{\hat{l}}) \leq \varepsilon$ and $\llbracket t \rrbracket_A^{\hat{l}} = \llbracket t' \rrbracket_A^{\hat{l}}$, thus

$$d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{l}}, \llbracket t' \rrbracket_A^{\hat{l}}) = d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{l}}, \llbracket t \rrbracket_A^{\hat{l}}) \leq \varepsilon. \quad \square$$

Lemma 3.33. *For any $s, s', t \in \mathcal{T}_\Sigma X$ and $\varepsilon \in \mathbf{L}$, if $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash s =_\varepsilon t$ and $\mathbf{X} \vdash s = s'$, then $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash s' =_\varepsilon t$.*

Proof. Symmetric argument to the previous proof. \square

Lemmas 3.34–3.37 will correspond to other rules in quantitative equational logic, and they will be explained in more details in §3.3.

Lemma 3.34. *For any $s, t \in \mathcal{T}_\Sigma X$, $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash s =_\top t$.*

Proof. By definition of \top (the supremum of all \mathbf{L}), for any \hat{l} , $d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{l}}, \llbracket t \rrbracket_A^{\hat{l}}) \leq \top$. \square

Lemma 3.35. *For any $x, x' \in X$, if $d_{\mathbf{X}}(x, x') = \varepsilon$, then $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash x =_\varepsilon x'$.*

Proof. For any nonexpansive $\hat{l} : \mathbf{X} \rightarrow \mathbf{A}$, we have³⁸⁵

$$d_{\mathbf{A}}(\llbracket x \rrbracket_A^{\hat{l}}, \llbracket x' \rrbracket_A^{\hat{l}}) = d_{\mathbf{A}}(\hat{l}(x), \hat{l}(x')) \leq d_{\mathbf{X}}(x, x') = \varepsilon. \quad \square$$

Lemma 3.36. *For any $s, t \in \mathcal{T}_\Sigma X$ and $\varepsilon, \varepsilon' \in \mathbf{L}$, if $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash s =_\varepsilon t$ and $\varepsilon \leq \varepsilon'$, then $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash s =_{\varepsilon'} t$.³⁸⁶*

Proof. For any $\hat{l} : \mathbf{X} \rightarrow \mathbf{A}$, we have $d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{l}}, \llbracket t \rrbracket_A^{\hat{l}}) \leq \varepsilon \leq \varepsilon'$. \square

Lemma 3.37. *For any $s, t \in \mathcal{T}_\Sigma X$ and $\{\varepsilon_i\}_{i \in I} \subseteq \mathbf{L}$, if $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash s =_{\varepsilon_i} t$ for all $i \in I$, then $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash s =_\varepsilon t$ with $\varepsilon = \inf_{i \in I} \varepsilon_i$.*

Proof. For any \hat{l} and for all $i \in I$, we have $d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{l}}, \llbracket t \rrbracket_A^{\hat{l}}) \leq \varepsilon_i$ by hypothesis. By definition of infimum, this means $d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{l}}, \llbracket t \rrbracket_A^{\hat{l}}) \leq \inf_{i \in I} \varepsilon_i = \varepsilon$. \square

This shall take care of all except two rules in quantitative equational logic which we will get to in no time. The following result is a generalization of Lemma 2.30, and it morally says that $\mathcal{T}_\Sigma f$ is well-defined and nonexpansive when f is nonexpansive.

³⁸⁴ By Lemmas 3.29 and 3.32, if $t \equiv_{\hat{\mathbf{E}}} t'$, then

$$\mathbf{X} \vdash s =_\varepsilon t \iff \mathbf{X} \vdash s =_\varepsilon t'.$$

By Lemmas 3.29 and 3.33, if $s \equiv_{\hat{\mathbf{E}}} s'$, then

$$\mathbf{X} \vdash s =_\varepsilon t' \iff \mathbf{X} \vdash s' =_\varepsilon t'.$$

Combining these with (3.14), we get

$$d_{\hat{\mathbf{E}}}(s, t) \leq \varepsilon \iff d_{\hat{\mathbf{E}}}(s', t') \leq \varepsilon,$$

for all $\varepsilon \in \mathbf{L}$, and we conclude $d_{\hat{\mathbf{E}}}(s, t) = d_{\hat{\mathbf{E}}}(s', t')$.

³⁸⁵ The equation holds by definition of $\llbracket - \rrbracket_A^{\hat{l}}$ on variables, and the inequality holds by definition of non-expansiveness.

³⁸⁶ In words, if the interpretations of s and t are at distance at most ε , then they are also at distance at most ε' when $\varepsilon \leq \varepsilon'$.

Lemma 3.38. *Let $f : \mathbf{X} \rightarrow \mathbf{Y}$ be a nonexpansive map. If \mathbf{A} satisfies $\mathbf{X} \vdash s = t$ (resp. $\mathbf{X} \vdash s =_\varepsilon t$), then \mathbf{A} satisfies $\mathbf{Y} \vdash \mathcal{T}_\Sigma f(s) = \mathcal{T}_\Sigma f(t)$ (resp. $\mathbf{Y} \vdash \mathcal{T}_\Sigma f(s) =_\varepsilon \mathcal{T}_\Sigma f(t)$).³⁸⁷*

Proof. Any nonexpansive assignment $\hat{\iota} : \mathbf{Y} \rightarrow \mathbf{A}$, yields a nonexpansive assignment $\hat{\iota} \circ f : \mathbf{X} \rightarrow \mathbf{A}$. Moreover, by functoriality of \mathcal{T}_Σ , we have

$$\llbracket - \rrbracket_A^{\hat{\iota} \circ f} \stackrel{(1.10)}{=} \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma(\hat{\iota} \circ f) = \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma \hat{\iota} \circ \mathcal{T}_\Sigma f = \llbracket \mathcal{T}_\Sigma f(-) \rrbracket_A^{\hat{\iota}}.$$

By hypothesis, we have

$$\mathbf{A} \models^{\hat{\iota} \circ f} \mathbf{X} \vdash s = t \quad (\text{resp. } \mathbf{A} \models^{\hat{\iota} \circ f} \mathbf{X} \vdash s =_\varepsilon t),$$

which means

$$\begin{aligned} \llbracket \mathcal{T}_\Sigma f(s) \rrbracket_A^{\hat{\iota}} &= \llbracket s \rrbracket_A^{\hat{\iota} \circ f} = \llbracket t \rrbracket_A^{\hat{\iota} \circ f} = \llbracket \mathcal{T}_\Sigma f(t) \rrbracket_A^{\hat{\iota}} \\ \text{resp. } d_{\mathbf{A}}(\llbracket \mathcal{T}_\Sigma f(s) \rrbracket_A^{\hat{\iota}}, \llbracket \mathcal{T}_\Sigma f(t) \rrbracket_A^{\hat{\iota}}) &= d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{\iota} \circ f}, \llbracket t \rrbracket_A^{\hat{\iota} \circ f}) \leq \varepsilon. \end{aligned}$$

Thus, we conclude

$$\mathbf{A} \models^{\hat{\iota}} \mathbf{Y} \vdash \mathcal{T}_\Sigma f(s) = \mathcal{T}_\Sigma f(t) \quad (\text{resp. } \mathbf{A} \models^{\hat{\iota}} \mathbf{Y} \vdash \mathcal{T}_\Sigma f(s) =_\varepsilon \mathcal{T}_\Sigma f(t)). \quad \square$$

Let us end our list of small results with Lemmas 3.39–3.41 which are for later.

Lemma 3.39. *For any $s, t \in \mathcal{T}_\Sigma X$ if $\hat{\mathbf{A}}$ satisfies $\mathbf{X}_\top \vdash s = t$, then $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash s = t$, and for any $\varepsilon \in \mathbb{L}$, if $\hat{\mathbf{A}}$ satisfies $\mathbf{X}_\top \vdash s =_\varepsilon t$, then $\hat{\mathbf{A}}$ satisfies $\mathbf{X} \vdash s =_\varepsilon t$.³⁸⁸*

Proof. For any nonexpansive assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$, you can pre-compose it with $\text{id}_X : \mathbf{X}_\top \rightarrow \mathbf{X}$ (which is nonexpansive) without changing the interpretation of terms: $\llbracket s \rrbracket_A^{\hat{\iota}} = \llbracket s \rrbracket_A^{\hat{\iota} \circ \text{id}_X}$. By hypothesis, we know that $\hat{\mathbf{A}}$ satisfies $s = t$ (resp. $s =_\varepsilon t$) under the nonexpansive assignment $\hat{\iota} \circ \text{id}_X : \mathbf{X}_\top \rightarrow \mathbf{A}$, and we conclude $\hat{\mathbf{A}}$ also satisfies $s = t$ (resp. $s =_\varepsilon t$) under the assignment $\hat{\iota}$. \square

Lemma 3.40. *For any $s, t \in \mathcal{T}_\Sigma X$, if \mathbf{A} satisfies $\mathbf{X} \vdash s = t$, then $\hat{\mathbf{A}}$ satisfies $\mathbf{X}_\top \vdash s = t$.³⁸⁹*

Proof. Any nonexpansive assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$ is in particular an assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$, thus $\llbracket s \rrbracket_A^{\hat{\iota}} = \llbracket t \rrbracket_A^{\hat{\iota}}$ hold by hypothesis that \mathbf{A} satisfies $\mathbf{X} \vdash s = t$. \square

Lemma 3.41. *For any $s, t \in \mathcal{T}_\Sigma X$, if $\hat{\mathbf{A}}$ satisfies $\mathbf{X}_\top \vdash s = t$, then \mathbf{A} satisfies $\mathbf{X} \vdash s = t$.³⁹⁰*

Proof. This follows by definition of the discrete space \mathbf{X}_\top . Indeed, any assignment $\iota : \mathbf{X} \rightarrow \mathbf{A}$ is the underlying function of a nonexpansive assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$, and since $\hat{\mathbf{A}}$ satisfies $s = t$ under $\hat{\iota}$ by hypothesis, \mathbf{A} satisfies $s = t$ under ι . \square

We can now get back to the equality $\equiv_{\hat{E}}$ and distance $d_{\hat{E}}$ between terms, and define the underlying space of the quantitative term algebra.

Since $\equiv_{\hat{E}}$ is an equivalence relation for any \mathbf{X} , we can consider the set $\mathcal{T}_\Sigma X / \equiv_{\hat{E}}$ of **terms modulo \hat{E}** .³⁹¹ We denote with $[-]_{\hat{E}} : \mathcal{T}_\Sigma X \rightarrow \mathcal{T}_\Sigma X / \equiv_{\hat{E}}$ the canonical quotient map, and by Lemmas 3.32 and 3.33, we can define an L-relation on terms modulo \hat{E} by factoring $d_{\hat{E}}$ through $[-]_{\hat{E}}$. We obtain the L-relation $d_{\hat{E}}$ as the unique function making the triangle below commute.³⁹²

³⁸⁷ Note that when s and t are variables, we get back Lemma 2.30.

³⁸⁸ In words, if $\hat{\mathbf{A}}$ satisfies an equation where the context is the discrete space on X , then $\hat{\mathbf{A}}$ satisfies that same equation with the context replaced by any other L-space on X . This is also a special case of Lemma 3.38 where $f : \mathbf{X}_\top \rightarrow \mathbf{X}$ is the identity map.

³⁸⁹ In words, if the underlying classical algebra satisfies an equation, then so does the quantitative algebra where the context can be endowed with any L-relation.

³⁹⁰ Combining Lemmas 3.40 and 3.41, we find

$$\mathbf{A} \models \mathbf{X} \vdash s = t \iff \hat{\mathbf{A}} \models \mathbf{X}_\top \vdash s = t. \quad (3.15)$$

This will be useful when comparing equational logic and quantitative equational logic in Example 3.71.

³⁹¹ Keep in mind that for different L-relations on X , we may get different equivalence relations on $\mathcal{T}_\Sigma X$, but we do not make this explicit in the notation $\equiv_{\hat{E}}$.

³⁹² We used the same symbol, because the first $d_{\hat{E}}$ was only used to define this new $d_{\hat{E}}$.

$$\begin{array}{ccc}
 \mathcal{T}_\Sigma X \times \mathcal{T}_\Sigma X & \xrightarrow{d_{\hat{E}}} & \mathbf{L} \\
 \downarrow [-]_{\hat{E}} \times [-]_{\hat{E}} & \searrow^{d_{\hat{E}}} & \\
 \mathcal{T}_\Sigma X / \equiv_{\hat{E}} \times \mathcal{T}_\Sigma X / \equiv_{\hat{E}} & &
 \end{array} \quad (3.16)$$

We write $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$ for the resulting L-space $(\mathcal{T}_\Sigma X / \equiv_{\hat{E}}, d_{\hat{E}})$. We still have an alternative definition analog to (3.14) for the new L-relation $d_{\hat{E}}$.³⁹³

$$d_{\hat{E}}([s]_{\hat{E}}, [t]_{\hat{E}}) \leq \varepsilon \iff \mathbf{X} \vdash s =_\varepsilon t \in \Omega \mathfrak{Th}(\hat{E}). \quad (3.17)$$

This will be the carrier of the term algebra on \mathbf{X} , so we need to prove that $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$ belongs to \mathbf{GMet} . We rely on the following generalization of Lemma 1.45. It essentially states that satisfaction of quantitative equations is preserved by substitutions that are nonexpansive. This result will also take care of the last two rules of quantitative equational logic.

Lemma 3.42. *Let \mathbf{Y} be an L-space and $\sigma : Y \rightarrow \mathcal{T}_\Sigma X$ be an assignment such that³⁹⁴*

$$\forall y, y' \in Y, \quad \mathbf{X} \vdash \sigma(y) =_{d_{\mathbf{Y}}(y, y')} \sigma(y') \in \Omega \mathfrak{Th}(\hat{E}), \quad (3.18)$$

and $\hat{\mathbf{A}}$ a (Σ, \hat{E}) -algebra. If $\hat{\mathbf{A}}$ satisfies $\mathbf{Y} \vdash s = t$ (resp. $\mathbf{Y} \vdash s =_\varepsilon t$), then it also satisfies $\mathbf{X} \vdash \sigma^(s) = \sigma^*(t)$ (resp. $\mathbf{X} \vdash \sigma^*(s) =_\varepsilon \sigma^*(t)$).*

Proof. Let $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$ be a nonexpansive assignment, we need to show $\llbracket \sigma^*(s) \rrbracket_A^{\hat{\iota}} = \llbracket \sigma^*(t) \rrbracket_A^{\hat{\iota}}$ (resp. $d_{\mathbf{A}}(\llbracket \sigma^*(s) \rrbracket_A^{\hat{\iota}}, \llbracket \sigma^*(t) \rrbracket_A^{\hat{\iota}}) \leq \varepsilon$). Just like in Lemma 1.45, we define the assignment $\hat{\iota}_\sigma : Y \rightarrow \mathbf{A}$ that sends $y \in Y$ to $\llbracket \sigma(y) \rrbracket_A^{\hat{\iota}}$, and we had already proven $\llbracket - \rrbracket_A^{\hat{\iota}_\sigma} = \llbracket \sigma^*(-) \rrbracket_A^{\hat{\iota}}$. Now, it is enough to show $\hat{\iota}_\sigma$ is nonexpansive $\mathbf{Y} \rightarrow \mathbf{A}$ ³⁹⁵ and the lemma will follow because by hypothesis, $\llbracket s \rrbracket_A^{\hat{\iota}_\sigma} = \llbracket t \rrbracket_A^{\hat{\iota}_\sigma}$ (reps. $d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{\iota}_\sigma}, \llbracket t \rrbracket_A^{\hat{\iota}_\sigma}) \leq \varepsilon$).

For any $y, y' \in Y$, we have

$$d_{\mathbf{A}}(\hat{\iota}_\sigma(y), \hat{\iota}_\sigma(y')) = d_{\mathbf{A}}(\llbracket \sigma(y) \rrbracket_A^{\hat{\iota}}, \llbracket \sigma(y') \rrbracket_A^{\hat{\iota}}) \leq d_{\mathbf{Y}}(y, y'),$$

where the equation holds by definition of $\hat{\iota}_\sigma$, and the inequality holds because $\hat{\mathbf{A}}$ belongs to $\mathbf{QAlg}(\Sigma, \hat{E})$ and hence satisfies $\mathbf{X} \vdash \sigma(y) =_{d_{\mathbf{Y}}(y, y')} \sigma(y') \in \Omega \mathfrak{Th}(\hat{E})$ (in particular under the nonexpansive assignment $\hat{\iota}$). Hence $\hat{\iota}_\sigma$ is nonexpansive. \square

Lemma 3.43. *For any L-space \mathbf{X} and any quantitative equation $\phi \in \hat{E}_{\mathbf{GMet}}$, $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \models \phi$.*

Proof. We mentioned in Footnote 375 that $\phi \in \Omega \mathfrak{Th}(\hat{E})$ because the carriers of (Σ, \hat{E}) -algebras are generalized metric spaces, so any (Σ, \hat{E}) -algebra $\hat{\mathbf{A}}$ satisfies it.

Let $\hat{\iota} : \mathbf{Y} \rightarrow \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$ be a nonexpansive assignment. By the axiom of choice,³⁹⁶ there is a function $\sigma : Y \rightarrow \mathcal{T}_\Sigma X$ satisfying $[\sigma(y)]_{\hat{E}} = \hat{\iota}(y)$ for all $y \in Y$. This assignment satisfies (3.18) because for all $y, y' \in Y$, (3.17) yields

$$d_{\hat{E}}([\sigma(y)]_{\hat{E}}, [\sigma(y')]_{\hat{E}}) \leq d_{\mathbf{Y}}(y, y') \stackrel{(3.17)}{\iff} \mathbf{X} \vdash \sigma(y) =_{d_{\mathbf{Y}}(y, y')} \sigma(y') \in \Omega \mathfrak{Th}(\hat{E}),$$

and the L.H.S. holds because $\hat{\iota}$ is nonexpansive.

³⁹³ In particular, the quotient map is nonexpansive:

$$[-]_{\hat{E}} : (\mathcal{T}_\Sigma X, d_{\hat{E}}) \rightarrow \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}.$$

³⁹⁴ By combining (3.18) with (3.14) we find that σ is a nonexpansive map $\mathbf{Y} \rightarrow (\mathcal{T}_\Sigma X, d_{\hat{E}})$, and any such nonexpansive map satisfies (3.18). We explicitly write (3.18) to better emulate the corresponding rules in quantitative equational logic.

³⁹⁵ Something we did not have to do in the classical case.

³⁹⁶ Choice implies the quotient map $[-]_{\hat{E}}$ has a right inverse $r : \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \rightarrow \mathcal{T}_\Sigma X$, and we set $\sigma = r \circ \hat{\iota}$.

Therefore, if ϕ has the shape $\mathbf{Y} \vdash y = y'$ (resp. $\mathbf{Y} \vdash y =_\varepsilon y'$), by Lemma 3.42, all (Σ, \hat{E}) -algebras satisfy $\mathbf{X} \vdash \sigma(y) = \sigma(y')$ (resp. $\mathbf{X} \vdash \sigma(y) =_\varepsilon \sigma(y')$). By definition of $\equiv_{\hat{E}}$ (resp. by definition of $d_{\hat{E}}$ (3.17)), we have

$$\hat{i}(y) = [\sigma(y)]_{\hat{E}} = [\sigma(y')]_{\hat{E}} = \hat{i}(y') \quad (\text{resp. } d_{\hat{E}}(\hat{i}(y), \hat{i}(y')) = d_{\hat{E}}([\sigma(y)]_{\hat{E}}, [\sigma(y')]_{\hat{E}}) \leq \varepsilon),$$

which means $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$ satisfies ϕ under \hat{i} . Since \hat{i} and ϕ were arbitrary, we conclude $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$ satisfies all of $\hat{E}_{\mathbf{GMet}}$, i.e. it is a generalized metric space. \square

As for **Set**, we obtain a functor $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} : \mathbf{GMet} \rightarrow \mathbf{GMet}$ ³⁹⁷ by setting $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} f$ equal to the unique function making (3.19) commute. Concretely, we have $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} f([t]_{\hat{E}}) = [\mathcal{T}_{\Sigma} f(t)]_{\hat{E}}$ which is well-defined by one part of Lemma 3.38.

$$\begin{array}{ccc} \mathcal{T}_{\Sigma} X & \xrightarrow{[-]_{\hat{E}}} & \mathcal{T}_{\Sigma} X / \equiv_{\hat{E}} \\ \mathcal{T}_{\Sigma} f \downarrow & & \downarrow \widehat{\mathcal{T}}_{\Sigma, \hat{E}} f \\ \mathcal{T}_{\Sigma} Y & \xrightarrow{[-]_{\hat{E}}} & \mathcal{T}_{\Sigma} Y / \equiv_{\hat{E}} \end{array} \quad (3.19)$$

Although we do have to check that $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} f$ is nonexpansive whenever f is, and we use the other part of Lemma 3.38.

Lemma 3.44. *If $f : \mathbf{X} \rightarrow \mathbf{Y}$ is nonexpansive, then so is $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} f : \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \rightarrow \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{Y}$.*

Proof. For any $s, t \in \mathcal{T}_{\Sigma} X$, we have

$$\begin{aligned} d_{\hat{E}}([s]_{\hat{E}}, [t]_{\hat{E}}) \leq \varepsilon &\iff \mathbf{X} \vdash s =_\varepsilon t \in \Omega \mathfrak{Th}(\hat{E}) && \text{by (3.17)} \\ &\implies \mathbf{X} \vdash \mathcal{T}_{\Sigma} f(s) =_\varepsilon \mathcal{T}_{\Sigma} f(t) \in \Omega \mathfrak{Th}(\hat{E}) && \text{Lemma 3.38} \\ &\iff d_{\hat{E}}([\mathcal{T}_{\Sigma} f(s)]_{\hat{E}}, [\mathcal{T}_{\Sigma} f(t)]_{\hat{E}}) \leq \varepsilon && \text{by (3.17)} \\ &\iff d_{\hat{E}}(\widehat{\mathcal{T}}_{\Sigma, \hat{E}} f[s]_{\hat{E}}, \widehat{\mathcal{T}}_{\Sigma, \hat{E}} f[t]_{\hat{E}}) \leq \varepsilon. && \text{by (3.19)} \end{aligned}$$

Therefore, $d_{\hat{E}}(\widehat{\mathcal{T}}_{\Sigma, \hat{E}} f[s]_{\hat{E}}, \widehat{\mathcal{T}}_{\Sigma, \hat{E}} f[t]_{\hat{E}}) \leq d_{\hat{E}}([s]_{\hat{E}}, [t]_{\hat{E}})$. \square

We may now define the interpretation of operation symbols syntactically to obtain the quantitative term algebra.

Definition 3.45 (Quantitative term algebra, semantically). The **quantitative term algebra** for (Σ, \hat{E}) on \mathbf{X} is the quantitative Σ -algebra whose underlying space is $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$ and whose interpretation of $\text{op} : n \in \Sigma$ is defined by³⁹⁸

$$\llbracket \text{op} \rrbracket_{\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}}([t_1]_{\hat{E}}, \dots, [t_n]_{\hat{E}}) = [\text{op}(t_1, \dots, t_n)]_{\hat{E}}. \quad (3.20)$$

We denote this algebra by $\widehat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{X}$ or simply $\widehat{\mathbb{T}} \mathbf{X}$.

This should feel very familiar to what we did in Definition 1.34.³⁹⁹ In particular, we still have that $[-]_{\hat{E}}$ is a homomorphism from $\mathcal{T}_{\Sigma} X$ to the underlying algebra of $\widehat{\mathbb{T}} \mathbf{X}$. Indeed, we can put $h = [-]_{\hat{E}}$ in (1.2) to get (3.20), or show that (3.21) commutes (recall Footnote 75).

$$\begin{array}{ccc} \mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma} X & \xrightarrow{\mathcal{T}_{\Sigma} [-]_{\hat{E}}} & \mathcal{T}_{\Sigma} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \\ \mu_{\Sigma}^{\mathcal{T}} \downarrow & & \downarrow \llbracket [-] \rrbracket_{\widehat{\mathbb{T}} \mathbf{X}} \\ \mathcal{T}_{\Sigma} X & \xrightarrow{[-]_{\hat{E}}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \end{array} \quad (3.21)$$

³⁹⁷ In fact, we defined a functor $\mathbf{LSpa} \rightarrow \mathbf{GMet}$, but we are interested in its restriction to \mathbf{GMet} .

³⁹⁸ This is well-defined by Lemma 3.31.

³⁹⁹ In fact, we can make the connection more precise, $\mathbb{T} \mathbf{X}$ is constructed by quotienting $\mathcal{T}_{\Sigma} X$ by the congruence \equiv_E , and (the underlying algebra of) $\widehat{\mathbb{T}} \mathbf{X}$ by quotienting $\mathcal{T}_{\Sigma} X$ by the congruence $\equiv_{\hat{E}}$ (see Remark 1.35).

While (3.21) is a diagram in **Set**, we write $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$ instead of the underlying set $\mathcal{T}_{\Sigma} X / \equiv_{\hat{E}}$ for better readability. We will keep this habit.

Your intuition for $\llbracket - \rrbracket_{\widehat{\mathcal{T}} \mathbf{X}}$ (the interpretation of arbitrary terms) should be exactly the same as the one for $\llbracket - \rrbracket_{\mathcal{T} X}$ in classical universal algebra: it takes a term in $\mathcal{T}_{\Sigma} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$, replaces the leaves with a representative term, and gives back the equivalence class of the resulting term. We can also use it to define an analog to flattening.⁴⁰⁰ For any space \mathbf{X} , let $\widehat{\mu}_{\mathbf{X}}^{\Sigma, \hat{E}}$ be the unique function making (3.22) commute.

$$\begin{array}{ccc}
 \mathcal{T}_{\Sigma} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} & \xrightarrow{\llbracket - \rrbracket_{\widehat{\mathcal{T}} \mathbf{X}}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \\
 \searrow [-]_{\hat{E}} & & \nearrow \widehat{\mu}_{\mathbf{X}}^{\Sigma, \hat{E}} \\
 & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} &
 \end{array} \quad (3.22)$$

⁴⁰⁰ Just as we did in (1.30).

Let us show that $\widehat{\mu}_{\mathbf{X}}^{\Sigma, \hat{E}}$ is nonexpansive and natural.

Lemma 3.46. *For any space \mathbf{X} , $\widehat{\mu}_{\mathbf{X}}^{\Sigma, \hat{E}}$ is a nonexpansive map $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \rightarrow \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$.*

Proof. Let $[s]_{\hat{E}}, [t]_{\hat{E}} \in \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$ be such that $d_{\hat{E}}([s]_{\hat{E}}, [t]_{\hat{E}}) \leq \varepsilon$. By (3.17), this means

$$\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \vdash s =_{\varepsilon} t \in \Omega \mathfrak{Th}(\hat{E}), \quad (3.23)$$

namely, the distance between interpretations of s and t is bounded above by ε in all (Σ, \hat{E}) -algebras. We need to show $d_{\hat{E}}(\llbracket s \rrbracket_{\widehat{\mathcal{T}} \mathbf{X}}, \llbracket t \rrbracket_{\widehat{\mathcal{T}} \mathbf{X}}) \leq \varepsilon$, or using (3.22),

$$d_{\hat{E}}(\llbracket s \rrbracket_{\widehat{\mathcal{T}} \mathbf{X}}, \llbracket t \rrbracket_{\widehat{\mathcal{T}} \mathbf{X}}) \leq \varepsilon. \quad (3.24)$$

We want to use (3.17) again to reduce that inequality to a bound on distances between interpretations, but that requires choosing representatives for $\llbracket s \rrbracket_{\widehat{\mathcal{T}} \mathbf{X}}, \llbracket t \rrbracket_{\widehat{\mathcal{T}} \mathbf{X}} \in \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$.

Instead of choosing them naively, let $s', t' \in \mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma} X$ be such that $\mathcal{T}_{\Sigma} [-]_{\hat{E}}(s') = s$ and $\mathcal{T}_{\Sigma} [-]_{\hat{E}}(t') = t$. In words, s' and t' are the same as s and t where equivalence classes at the leaves are replaced representative terms.⁴⁰¹ Commutativity of (3.21) implies $[\mu_{\mathbf{X}}^{\Sigma}(s')]_{\hat{E}} = \llbracket s \rrbracket_{\widehat{\mathcal{T}} \mathbf{X}}$ and similarly for t . We can now use (3.17) to infer that proving (3.24) is equivalent to proving

$$\mathbf{X} \vdash \mu_{\mathbf{X}}^{\Sigma}(s') =_{\varepsilon} \mu_{\mathbf{X}}^{\Sigma}(t') \in \Omega \mathfrak{Th}(\hat{E}). \quad (3.25)$$

This means we need to show that, for all $\hat{A} \in \mathbf{QAlg}(\Sigma, \hat{E})$ and $\hat{t} : \mathbf{X} \rightarrow \hat{A}$, $d_{\hat{A}}(\llbracket \mu_{\mathbf{X}}^{\Sigma}(s') \rrbracket_{\hat{A}}, \llbracket \mu_{\mathbf{X}}^{\Sigma}(t') \rrbracket_{\hat{A}}) \leq \varepsilon$.

We already know by (3.23) that for all $\hat{\sigma} : \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \rightarrow \hat{A}$, $d_{\hat{A}}(\llbracket s \rrbracket_{\hat{A}}^{\hat{\sigma}}, \llbracket t \rrbracket_{\hat{A}}^{\hat{\sigma}}) \leq \varepsilon$, so it suffices to find, for each $\hat{t} : \mathbf{X} \rightarrow \hat{A}$, a nonexpansive assignment $\hat{\sigma}_{\hat{t}} : \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \rightarrow \hat{A}$ such that

$$\llbracket \mu_{\mathbf{X}}^{\Sigma}(s') \rrbracket_{\hat{A}}^{\hat{\sigma}_{\hat{t}}} = \llbracket s \rrbracket_{\hat{A}}^{\hat{\sigma}_{\hat{t}}} \text{ and } \llbracket \mu_{\mathbf{X}}^{\Sigma}(t') \rrbracket_{\hat{A}}^{\hat{\sigma}_{\hat{t}}} = \llbracket t \rrbracket_{\hat{A}}^{\hat{\sigma}_{\hat{t}}}. \quad (3.26)$$

We define $\hat{\sigma}_{\hat{t}} : \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \rightarrow \hat{A}$ to be the unique function making (3.27) commute.⁴⁰²

$$\begin{array}{ccc}
 \mathcal{T}_{\Sigma} X & \xrightarrow{\mathcal{T}_{\Sigma} \hat{t}} & \mathcal{T}_{\Sigma} A \\
 \downarrow [-]_{\hat{E}} & & \downarrow \llbracket - \rrbracket_A \\
 \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} & \xrightarrow{\hat{\sigma}_{\hat{t}}} & A
 \end{array} \quad (3.27)$$

⁴⁰¹ Since s and t have finitely many leaves, we are only doing finitely many choices of representatives.

⁴⁰² It exists because \hat{A} satisfies all the equations in $\Omega \mathfrak{Th}(\hat{E})$ so if $s \equiv_{\hat{E}} t$ then

$$\llbracket \mathcal{T}_{\Sigma} \hat{t}(s) \rrbracket_A \stackrel{(1.10)}{=} \llbracket s \rrbracket_A^{\hat{t}} = \llbracket t \rrbracket_A^{\hat{t}} \stackrel{(1.10)}{=} \llbracket \mathcal{T}_{\Sigma} \hat{t}(t) \rrbracket_A.$$

First, $\hat{\sigma}_i$ is a nonexpansive map $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \rightarrow \mathbf{A}$ because for any $[u]_{\hat{E}}, [v]_{\hat{E}} \in \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$,

$$d_{\mathbf{A}}(\hat{\sigma}_i[u]_{\hat{E}}, \hat{\sigma}_i[v]_{\hat{E}}) \stackrel{(3.27)}{=} d_{\mathbf{A}}([\mathcal{T}_{\Sigma} i(u)]_A, [\mathcal{T}_{\Sigma} i(v)]_A) \stackrel{(1.10)}{=} d_{\mathbf{A}}([\![u]\!]_A, [\![v]\!]_A) \leq d_{\hat{E}}([u]_{\hat{E}}, [v]_{\hat{E}}),$$

where the inequality holds by definition of $d_{\hat{E}}$ and because $\hat{\mathbf{A}}$ satisfies all the equations in $\Omega\mathfrak{Th}(\hat{E})$.

Second, we can prove that

$$[\![-]\!]_A \circ \mu_{\mathbf{X}}^{\Sigma} = [\![-]\!]_A^{\hat{\sigma}_i} \circ \mathcal{T}_{\Sigma}[-]_{\hat{E}}, \tag{3.28}$$

which implies (3.26) holds (by applying both sides of (3.28) to s' and t'). We pave the following diagram.

$$\begin{array}{ccc}
 \mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma} \mathbf{X} & \xrightarrow{\mathcal{T}_{\Sigma}[-]_{\hat{E}}} & \mathcal{T}_{\Sigma} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \\
 \downarrow \mu_{\mathbf{X}}^{\Sigma} & \searrow \mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma} i & \swarrow \mathcal{T}_{\Sigma} \hat{\sigma}_i \\
 & \mathcal{T}_{\Sigma} \mathbf{X} & \xrightarrow{\mathcal{T}_{\Sigma}[-]_A} \mathcal{T}_{\Sigma} \mathbf{A} \\
 & \swarrow & \searrow \\
 \mathcal{T}_{\Sigma} \mathbf{X} & \xrightarrow{[\![-]\!]_A} & \mathbf{A}
 \end{array}
 \tag{3.29}$$

□

Showing (3.29) commutes:

(a) Apply \mathcal{T}_{Σ} to (3.27).

(b) By (1.15).

(c) By (1.10).

Lemma 3.47. *The family of maps $\hat{\mu}_{\mathbf{X}}^{\Sigma, \hat{E}} : \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \rightarrow \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$ is natural in \mathbf{X} .*⁴⁰³

Proof. We need to prove that for any function $f : \mathbf{X} \rightarrow \mathbf{Y}$, the square below commutes.

$$\begin{array}{ccc}
 \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} & \xrightarrow{\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} f} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{Y} \\
 \hat{\mu}_{\mathbf{X}}^{\Sigma, \hat{E}} \downarrow & & \downarrow \hat{\mu}_{\mathbf{Y}}^{\Sigma, \hat{E}} \\
 \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} & \xrightarrow{\widehat{\mathcal{T}}_{\Sigma, \hat{E}} f} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{Y}
 \end{array}
 \tag{3.30}$$

⁴⁰³ We will (for posterity) reproduce the proof we did for Proposition 1.38, but it is important to note that nothing changes except the notation which now has lots of little hats.

We can pave the following diagram.

$$\begin{array}{ccccc}
 \mathcal{T}_{\Sigma} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} & \xrightarrow{[\![-]\!]_{\hat{E}}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} & \xrightarrow{\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} f} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{Y} \\
 \downarrow [\![-]\!]_{\hat{E}} & \searrow \mathcal{T}_{\Sigma} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} f & \swarrow [\![-]\!]_{\hat{E}} & \swarrow [\![-]\!]_{\hat{E}} & \downarrow \hat{\mu}_{\mathbf{Y}}^{\Sigma, \hat{E}} \\
 & \mathcal{T}_{\Sigma} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{Y} & & & \\
 & \swarrow & \searrow & \searrow & \\
 \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} & \xrightarrow{\hat{\mu}_{\mathbf{X}}^{\Sigma, \hat{E}}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} & \xrightarrow{\widehat{\mathcal{T}}_{\Sigma, \hat{E}} f} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{Y}
 \end{array}$$

All of (a), (b) and (d) commute by definition. In more details, (a) is an instance of (3.19) with \mathbf{X} replaced by $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X}$, \mathbf{Y} by $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{Y}$ and f by $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} f$, and both (b) and (d) are

instances of (3.22). To show (c) commutes, we draw another diagram that looks like a cube and where (c) is the front face. We can show all the other faces commute, and then use the fact that $\mathcal{T}_\Sigma[-]_{\hat{E}}$ is surjective (i.e. epic) to conclude that the front face must also commute.⁴⁰⁴

$$\begin{array}{ccccc}
 \mathcal{T}_\Sigma \mathcal{T}_\Sigma X & \xrightarrow{\mathcal{T}_\Sigma \mathcal{T}_\Sigma f} & \mathcal{T}_\Sigma \mathcal{T}_\Sigma Y & & \\
 \downarrow \mu_X^\Sigma & \searrow \mathcal{T}_\Sigma[-]_{\hat{E}} & \downarrow \mu_Y^\Sigma & \searrow \mathcal{T}_\Sigma[-]_{\hat{E}} & \\
 \mathcal{T}_\Sigma X & \xrightarrow{\mathcal{T}_\Sigma f} & \mathcal{T}_\Sigma Y & & \\
 \downarrow [-]_{\hat{E}} & \searrow \llbracket - \rrbracket_{\hat{\mathbb{T}}X} & \downarrow [-]_{\hat{E}} & \searrow \llbracket - \rrbracket_{\hat{\mathbb{T}}Y} & \\
 \hat{\mathcal{T}}_{\Sigma, \hat{E}} X & \xrightarrow{\hat{\mathcal{T}}_{\Sigma, \hat{E}} f} & \hat{\mathcal{T}}_{\Sigma, \hat{E}} Y & &
 \end{array}
 \quad (3.31)$$

The first diagram we paved implies (1.31) commutes because $[-]_{\hat{E}}$ is surjective. \square

From the front face of the cube above, we find that for any $f : X \rightarrow Y$, $\hat{\mathcal{T}}_{\Sigma, \hat{E}} f$ is a homomorphism between the underlying algebras of $\hat{\mathbb{T}}X$ and $\hat{\mathbb{T}}Y$. We already showed $\hat{\mathcal{T}}_{\Sigma, \hat{E}} f$ is nonexpansive in Lemma 3.44, thus it is a homomorphism between the quantitative algebras $\hat{\mathbb{T}}X$ and $\hat{\mathbb{T}}Y$.

Lemma 3.48. *For any nonexpansive map $f : X \rightarrow Y$, $\hat{\mathcal{T}}_{\Sigma, \hat{E}} f$ is a nonexpansive homomorphism $\hat{\mathbb{T}}X \rightarrow \hat{\mathbb{T}}Y$.*

We now prove generalizations of results from Chapter 1 in order to show that $\hat{\mathbb{T}}X$ is not just a quantitative Σ -algebra but a (Σ, \hat{E}) -algebra.

We can prove, analogously to Lemma 1.39, that for any $\hat{A} \in \mathbf{QAlg}(\Sigma, \hat{E})$, $\llbracket - \rrbracket_A$ is a homomorphism between $\hat{\mathbb{T}}A$ and \hat{A} .

Lemma 3.49. *For any (Σ, \hat{E}) -algebra \hat{A} , the square (3.32) commutes, and $\llbracket - \rrbracket_A$ is a nonexpansive map $\hat{\mathcal{T}}_{\Sigma, \hat{E}} A \rightarrow \hat{A}$.⁴⁰⁵*

$$\begin{array}{ccc}
 \mathcal{T}_\Sigma \hat{\mathcal{T}}_{\Sigma, \hat{E}} A & \xrightarrow{\mathcal{T}_\Sigma \llbracket - \rrbracket_A} & \mathcal{T}_\Sigma A \\
 \llbracket - \rrbracket_{\hat{\mathbb{T}}A} \downarrow & & \downarrow \llbracket - \rrbracket_A \\
 \hat{\mathcal{T}}_{\Sigma, \hat{E}} A & \xrightarrow{\llbracket - \rrbracket_A} & A
 \end{array}
 \quad (3.32)$$

Proof. For the commutative square, we can reuse the proof of Lemma 1.39.

Consider the following diagram that we can view as a triangular prism whose front face is (3.32). Both triangles commute by Footnote 405, the square face at the back and on the left commutes by (3.21), and the square face at the back and on the right commutes by (1.14). With the same trick as in the proof of Lemma 3.47 using the surjectivity of $\mathcal{T}_\Sigma[-]_{\hat{E}}$, we conclude that the front face commutes.⁴⁰⁶

⁴⁰⁴ In more details, the left and right faces commute by (3.21), the bottom and top faces commute by (3.19), and the back face commutes by (1.8).

The function $\mathcal{T}_\Sigma[-]_{\hat{E}}$ is surjective (i.e. epic) because $[-]_{\hat{E}}$ is (it is a canonical quotient map) and functors on \mathbf{Set} preserve epimorphisms (if we assume the axiom of choice). Thus, it suffices to show that $\mathcal{T}_\Sigma[-]_{\hat{E}}$ pre-composed with the bottom path or the top path of the front face gives the same result.

Now it is just a matter of going around the cube using the commutativity of the other faces. Here is the complete derivation (we write which face was used as justifications for each step).

$$\begin{aligned}
 & \hat{\mathcal{T}}_{\Sigma, \hat{E}} f \circ \llbracket - \rrbracket_{\hat{\mathbb{T}}X} \circ \mathcal{T}_\Sigma[-]_{\hat{E}} && \text{left} \\
 & = \hat{\mathcal{T}}_{\Sigma, \hat{E}} f \circ [-]_{\hat{E}} \circ \mu_X^\Sigma && \text{bottom} \\
 & = [-]_{\hat{E}} \circ \mathcal{T}_\Sigma f \circ \mu_X^\Sigma && \text{back} \\
 & = [-]_{\hat{E}} \circ \mu_Y^\Sigma \circ \mathcal{T}_\Sigma \mathcal{T}_\Sigma f && \text{right} \\
 & = \llbracket - \rrbracket_{\hat{\mathbb{T}}Y} \circ \mathcal{T}_\Sigma[-]_{\hat{E}} \circ \mathcal{T}_\Sigma \mathcal{T}_\Sigma f && \text{right} \\
 & = \llbracket - \rrbracket_{\hat{\mathbb{T}}Y} \circ \mathcal{T}_\Sigma \hat{\mathcal{T}}_{\Sigma, \hat{E}} f \circ \mathcal{T}_\Sigma[-]_{\hat{E}} && \text{top}
 \end{aligned}$$

⁴⁰⁵ We use the same convention as in (1.34) and write $\llbracket - \rrbracket_A$ for both maps $\mathcal{T}_\Sigma A \rightarrow A$ and $\hat{\mathcal{T}}_{\Sigma, \hat{E}} A \rightarrow A$. Recall the latter is well-defined because whenever $[s]_{\hat{E}} = [t]_{\hat{E}}$, \hat{A} must satisfy $A \vdash s = t$, and in particular under the assignment $\text{id}_A : A \rightarrow A$, this yields $\llbracket s \rrbracket_A = \llbracket t \rrbracket_A$.

⁴⁰⁶ Here is the complete derivation.

$$\begin{aligned}
 & \llbracket - \rrbracket_A \circ \llbracket - \rrbracket_{\hat{\mathbb{T}}A} \circ \mathcal{T}_\Sigma[-]_{\hat{E}} && \text{left} \\
 & = \llbracket - \rrbracket_A \circ [-]_{\hat{E}} \circ \mu_A^\Sigma && \text{bottom} \\
 & = \llbracket - \rrbracket_A \circ \mu_A^\Sigma && \text{right} \\
 & = \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma \llbracket - \rrbracket_A && \text{right} \\
 & = \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma[-]_{\hat{E}} && \text{top}
 \end{aligned}$$

Then, since $\mathcal{T}_\Sigma[-]_{\hat{E}}$ is epic, we conclude that $\llbracket - \rrbracket_A \circ \llbracket - \rrbracket_{\hat{\mathbb{T}}A} = \llbracket - \rrbracket_A \circ \mathcal{T}_\Sigma \llbracket - \rrbracket_A$.

$$\begin{array}{ccccc}
 & & \mathcal{T}_\Sigma \mathcal{T}_\Sigma A & & \\
 & \mathcal{T}_\Sigma [-]_{\hat{E}} \swarrow & \downarrow & \searrow \mathcal{T}_\Sigma [-]_A & \\
 \mathcal{T}_\Sigma \mathcal{T}_{\Sigma, \hat{E}} A & \xrightarrow{\quad} & \mathcal{T}_\Sigma A & \xrightarrow{\quad} & \mathcal{T}_\Sigma A \\
 \downarrow \llbracket - \rrbracket_{\hat{\mathbb{T}}A} & & \downarrow \mu_A^\Sigma & & \downarrow \llbracket - \rrbracket_A \\
 & \swarrow [-]_{\hat{E}} & \mathcal{T}_\Sigma A & \searrow \llbracket - \rrbracket_A & \\
 \mathcal{T}_{\Sigma, \hat{E}} A & \xrightarrow{\quad} & A & \xrightarrow{\quad} & A \\
 & & \llbracket - \rrbracket_A & &
 \end{array}$$

For nonexpansiveness, if $d_{\hat{E}}([s]_{\hat{E}}, [t]_{\hat{E}}) \leq \varepsilon$, then by (3.17), $\mathbf{A} \vdash s =_\varepsilon t$ belongs to $\Omega\mathfrak{T}\mathfrak{h}(\hat{E})$ which means $\hat{\mathbf{A}}$ must satisfy that equation, and in particular under the assignment $\text{id}_A : \mathbf{A} \rightarrow \mathbf{A}$, this yields $d_A(\llbracket s \rrbracket_A, \llbracket t \rrbracket_A) \leq \varepsilon$. \square

We can prove, analogously to Lemma 1.40, that for any \mathbf{X} , $\hat{\mu}_X^{\Sigma, \hat{E}}$ is a homomorphism from $\hat{\mathbb{T}}\mathbb{T}\mathbf{X}$ to $\hat{\mathbb{T}}\mathbf{X}$.

Lemma 3.50. *For any generalized metric space \mathbf{X} , the following square commutes, and $\hat{\mu}_X^{\Sigma, \hat{E}}$ is a nonexpansive map $\hat{\mathbb{T}}_{\Sigma, \hat{E}} \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{X} \rightarrow \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{X}$.*

$$\begin{array}{ccc}
 \mathcal{T}_\Sigma \hat{\mathbb{T}}_{\Sigma, \hat{E}} \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{X} & \xrightarrow{\mathcal{T}_\Sigma \hat{\mu}_X^{\Sigma, \hat{E}}} & \mathcal{T}_\Sigma \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{X} \\
 \llbracket - \rrbracket_{\hat{\mathbb{T}}\mathbb{T}\mathbf{X}} \downarrow & & \downarrow \llbracket - \rrbracket_{\hat{\mathbb{T}}\mathbf{X}} \\
 \hat{\mathbb{T}}_{\Sigma, \hat{E}} \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{X} & \xrightarrow{\hat{\mu}_X^{\Sigma, \hat{E}}} & \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{X}
 \end{array} \quad (3.33)$$

Proof. We already showed nonexpansiveness in Lemma 3.46. For the commutative square, we can reuse the argument of Lemma 1.40 and add the little hats.

We prove it exactly like Lemma 3.49 with the following diagram.⁴⁰⁷

$$\begin{array}{ccccc}
 & & \mathcal{T}_\Sigma \mathcal{T}_\Sigma \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{A} & & \\
 & \mathcal{T}_\Sigma [-]_{\hat{E}} \swarrow & \downarrow & \searrow \mathcal{T}_\Sigma \llbracket - \rrbracket_{\hat{\mathbb{T}}A} & \\
 \mathcal{T}_\Sigma \hat{\mathbb{T}}_{\Sigma, \hat{E}} \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{A} & \xrightarrow{\mathcal{T}_\Sigma \mu_A^{\Sigma, \hat{E}}} & \mathcal{T}_\Sigma \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{A} & \xrightarrow{\quad} & \mathcal{T}_\Sigma \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{A} \\
 \downarrow \llbracket - \rrbracket_{\hat{\mathbb{T}}\hat{\mathbf{A}}} & & \downarrow \mu_{\hat{\Sigma, \hat{E}} \mathbf{A}}^\Sigma & & \downarrow \llbracket - \rrbracket_{\hat{\mathbb{T}}A} \\
 & \swarrow [-]_{\hat{E}} & \mathcal{T}_\Sigma \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{A} & \searrow \llbracket - \rrbracket_{\hat{\mathbb{T}}A} & \\
 \hat{\mathbb{T}}_{\Sigma, \hat{E}} \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{A} & \xrightarrow{\quad} & \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{A} & \xrightarrow{\quad} & \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{A} \\
 & & \hat{\mu}_A^{\Sigma, \hat{E}} & &
 \end{array}$$

\square

Of course, paired with the flattening we also have a map $\hat{\eta}_A^{\Sigma, \hat{E}}$ which sends elements $a \in A$ to the equivalence class containing a seen as a trivial term, namely,

$$\hat{\eta}_A^{\Sigma, \hat{E}} = \mathbf{A} \xrightarrow{\eta_A^\Sigma} \mathcal{T}_\Sigma A \xrightarrow{[-]_{\hat{E}}} \hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{A}. \quad (3.34)$$

We need to show $\hat{\eta}_A^{\Sigma, \hat{E}}$ is nonexpansive and natural in \mathbf{A} .

⁴⁰⁷ The top and bottom faces commute by definition of $\hat{\mu}_A^{\Sigma, \hat{E}}$ (3.22), the back-left face by (3.21), and the back-right face by (1.14).

Then, $\mathcal{T}_\Sigma [-]_{\hat{E}}$ is epic, so the following derivation suffices.

$$\begin{aligned}
 & \hat{\mu}_A^{\Sigma, \hat{E}} \circ \llbracket - \rrbracket_{\hat{\mathbb{T}}\hat{\mathbf{A}}} \circ \mathcal{T}_\Sigma [-]_{\hat{E}} && \\
 & = \hat{\mu}_A^{\Sigma, \hat{E}} \circ [-]_{\hat{E}} \circ \mu_{\hat{\Sigma, \hat{E}} \mathbf{A}}^\Sigma && \text{left} \\
 & = \llbracket - \rrbracket_{\hat{\mathbb{T}}A} \circ \mu_{\hat{\Sigma, \hat{E}} \mathbf{A}}^\Sigma && \text{bottom} \\
 & = \llbracket - \rrbracket_{\hat{\mathbb{T}}A} \circ \mathcal{T}_\Sigma \llbracket - \rrbracket_{\hat{\mathbb{T}}A} && \text{right} \\
 & = \llbracket - \rrbracket_{\hat{\mathbb{T}}A} \circ \mathcal{T}_\Sigma \hat{\mu}_A^{\Sigma, \hat{E}} \circ \mathcal{T}_\Sigma [-]_{\hat{E}} && \text{top}
 \end{aligned}$$

Lemma 3.51. For any space \mathbf{A} , $\widehat{\eta}_{\mathbf{A}}^{\Sigma, \hat{E}}$ is a nonexpansive map $\mathbf{A} \rightarrow \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{A}$.

Proof. This is a direct consequence of Lemma 3.35. For any $a, a' \in X$ and $\varepsilon \in L$,

$$\begin{aligned} d_{\mathbf{A}}(a, a') \leq \varepsilon &\implies \mathbf{A} \vdash a =_{\varepsilon} a' \in \Omega \mathfrak{Th}(\hat{E}) && \text{by Lemma 3.35} \\ &\iff d_{\hat{E}}([a]_{\hat{E}}, [a']_{\hat{E}}) \leq \varepsilon. && \text{by (3.17)} \end{aligned}$$

Therefore, $d_{\hat{E}}([a]_{\hat{E}}, [a']_{\hat{E}}) \leq d_{\mathbf{A}}(a, a')$. \square

Lemma 3.52. For any nonexpansive map $f : \mathbf{A} \rightarrow \mathbf{B}$, the following square commutes.⁴⁰⁸

$$\begin{array}{ccc} \mathbf{A} & \xrightarrow{\widehat{\eta}_{\mathbf{A}}^{\Sigma, \hat{E}}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{A} \\ f \downarrow & & \downarrow \widehat{\mathcal{T}}_{\Sigma, \hat{E}} f \\ \mathbf{B} & \xrightarrow{\widehat{\eta}_{\mathbf{B}}^{\Sigma, \hat{E}}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{B} \end{array} \quad (3.35)$$

Proof. We pave the following diagram (in \mathbf{Set} , but that is enough since $U : \mathbf{GMet} \rightarrow \mathbf{Set}$ is faithful).

$$\begin{array}{ccccc} \mathbf{A} & \xrightarrow{\widehat{\eta}_{\mathbf{A}}^{\Sigma, \hat{E}}} & & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{A} & \\ \downarrow f & \searrow \eta_{\mathbf{A}}^{\Sigma} & \text{(a)} & \searrow [-]_{\hat{E}} & \downarrow \widehat{\mathcal{T}}_{\Sigma, \hat{E}} f \\ & & \mathcal{T}_{\Sigma} \mathbf{A} & & \\ & & \downarrow \mathcal{T}_{\Sigma} f & \text{(c)} & \\ & & \mathcal{T}_{\Sigma} \mathbf{B} & & \\ & \nearrow \eta_{\mathbf{B}}^{\Sigma} & \text{(d)} & \nearrow [-]_{\hat{E}} & \\ \mathbf{B} & \xrightarrow{\widehat{\eta}_{\mathbf{B}}^{\Sigma, \hat{E}}} & & \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{B} & \end{array} \quad (3.36)$$

Showing (3.36) commutes:
 (a) Definition of $\widehat{\eta}_{\mathbf{A}}^{\Sigma, \hat{E}}$ (3.34).
 (b) Naturality of η^{Σ} (1.6).
 (c) Definition of $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} f$ (3.19).
 (d) Definition of $\widehat{\eta}_{\mathbf{B}}^{\Sigma, \hat{E}}$ (3.34).

\square

We also have the following technical lemma and its corollary analogous to Lemma 1.41 and Lemma 1.42.

Lemma 3.53. For any generalized metric space \mathbf{X} , $\llbracket [-] \rrbracket_{\widehat{\mathbf{T}}\mathbf{X}}^{\Sigma, \hat{E}} = [-]_{\hat{E}}$.⁴⁰⁹

Proof. We proceed by induction. For the base case, we have

$$\begin{aligned} \llbracket \eta_{\mathbf{X}}^{\Sigma}(x) \rrbracket_{\widehat{\mathbf{T}}\mathbf{X}}^{\Sigma, \hat{E}} &= \llbracket \mathcal{T}_{\Sigma} \widehat{\eta}_{\mathbf{X}}^{\Sigma, \hat{E}}(\eta_{\mathbf{X}}^{\Sigma}(x)) \rrbracket_{\widehat{\mathbf{T}}\mathbf{X}} && \text{by (1.10)} \\ &= \llbracket \mathcal{T}_{\Sigma} [-]_{\hat{E}}(\mathcal{T}_{\Sigma} \eta_{\mathbf{X}}^{\Sigma}(\eta_{\mathbf{X}}^{\Sigma}(x))) \rrbracket_{\widehat{\mathbf{T}}\mathbf{X}} && \text{Lemma 1.11} \\ &= \llbracket \mathcal{T}_{\Sigma} [-]_{\hat{E}}(\eta_{\mathcal{T}_{\Sigma} \mathbf{X}}^{\Sigma}(\eta_{\mathbf{X}}^{\Sigma}(x))) \rrbracket_{\widehat{\mathbf{T}}\mathbf{X}} && \text{by (1.6)} \\ &= \llbracket \eta_{\mathcal{T}_{\Sigma, \hat{E}} \mathbf{X}}^{\Sigma}([\eta_{\mathbf{X}}^{\Sigma}(x)]_{\hat{E}}) \rrbracket_{\widehat{\mathbf{T}}\mathbf{X}} && \text{by (1.6)} \\ &= [\eta_{\mathbf{X}}^{\Sigma}(x)]_{\hat{E}} && \text{by (1.29)} \end{aligned}$$

For the inductive step, if $t = \text{op}(t_1, \dots, t_n)$, we have

$$\llbracket t \rrbracket_{\widehat{\mathbf{T}}\mathbf{X}}^{\Sigma, \hat{E}} = \llbracket \mathcal{T}_{\Sigma} \widehat{\eta}_{\mathbf{X}}^{\Sigma, \hat{E}}(t) \rrbracket_{\widehat{\mathbf{T}}\mathbf{X}} \quad \text{by (1.10)}$$

⁴⁰⁹ The proof is identical to that of Lemma 1.41.

⁴⁰⁸ Naturality of $\eta^{\Sigma, \hat{E}}$ was easier in \mathbf{Set} because it is the vertical composition of two natural transformations, η^{Σ} and $[-]_{\hat{E}}$, which do not have counterparts in \mathbf{GMet} .

$$\begin{aligned}
 &= \llbracket \mathcal{T}_\Sigma \widehat{\eta}_X^{\Sigma, \hat{E}}(\text{op}(t_1, \dots, t_n)) \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}} \\
 &= \llbracket \text{op}(\mathcal{T}_\Sigma \widehat{\eta}_X^{\Sigma, \hat{E}}(t_1), \dots, \mathcal{T}_\Sigma \widehat{\eta}_X^{\Sigma, \hat{E}}(t_n)) \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}} && \text{by (1.5)} \\
 &= \llbracket \text{op} \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}} \left(\llbracket \mathcal{T}_\Sigma \widehat{\eta}_X^{\Sigma, \hat{E}}(t_1) \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}}, \dots, \llbracket \mathcal{T}_\Sigma \widehat{\eta}_X^{\Sigma, \hat{E}}(t_n) \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}} \right) && \text{by (1.29)} \\
 &= \llbracket \text{op} \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}} ([t_1]_{\hat{E}}, \dots, [t_n]_{\hat{E}}) && \text{I.H.} \\
 &= [\text{op}(t_1, \dots, t_n)]_{\hat{E}}. && \text{by (3.20)} \quad \square
 \end{aligned}$$

We get that for any quantitative equation ϕ with context \mathbf{X} , ϕ belongs to $\Omega\mathfrak{Th}(\hat{E})$ if and only if the algebra $\widehat{\mathbb{T}}_{\Sigma, \hat{E}}\mathbf{X}$ satisfies it under the assignment $\widehat{\eta}_X^{\Sigma, \hat{E}}$.

Lemma 3.54. *Let ϕ be an equation with context \mathbf{X} , $\phi \in \Omega\mathfrak{Th}(\hat{E})$ if and only if $\widehat{\mathbb{T}}\mathbf{X} \models_{\widehat{\eta}_X^{\Sigma, \hat{E}}} \phi$.⁴¹⁰*

Proof. We have two cases to show.

- $\mathbf{X} \vdash s = t \in \Omega\mathfrak{Th}(\hat{E})$ if and only if $\widehat{\mathbb{T}}\mathbf{X} \models_{\widehat{\eta}_X^{\Sigma, \hat{E}}} \mathbf{X} \vdash s = t$, and
- $\mathbf{X} \vdash s =_\varepsilon t \in \Omega\mathfrak{Th}(\hat{E})$ if and only if $\widehat{\mathbb{T}}\mathbf{X} \models_{\widehat{\eta}_X^{\Sigma, \hat{E}}} \mathbf{X} \vdash s =_\varepsilon t$.

By Lemma 3.53,

$$\llbracket [s] \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}}^{\widehat{\eta}_X^{\Sigma, \hat{E}}} = [s]_{\hat{E}} \text{ and } \llbracket [t] \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}}^{\widehat{\eta}_X^{\Sigma, \hat{E}}} = [t]_{\hat{E}}, \quad (3.37)$$

then by using definitions, we have (as desired)

$$\begin{aligned}
 \mathbf{X} \vdash s = t \in \Omega\mathfrak{Th}(\hat{E}) &\stackrel{(3.13)}{\iff} [s]_{\hat{E}} = [t]_{\hat{E}} \stackrel{(3.37)}{\iff} \llbracket [s] \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}}^{\widehat{\eta}_X^{\Sigma, \hat{E}}} = \llbracket [t] \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}}^{\widehat{\eta}_X^{\Sigma, \hat{E}}} \\
 \mathbf{X} \vdash s =_\varepsilon t \in \Omega\mathfrak{Th}(\hat{E}) &\stackrel{(3.17)}{\iff} d_{\hat{E}}(\llbracket [s] \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}}^{\widehat{\eta}_X^{\Sigma, \hat{E}}}, \llbracket [t] \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}}^{\widehat{\eta}_X^{\Sigma, \hat{E}}}) \leq \varepsilon \stackrel{(3.37)}{\iff} d_{\hat{E}}(\llbracket [s] \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}}^{\widehat{\eta}_X^{\Sigma, \hat{E}}}, \llbracket [t] \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}}^{\widehat{\eta}_X^{\Sigma, \hat{E}}}) \leq \varepsilon. \quad \square
 \end{aligned}$$

The next result, analogous to Lemma 1.43, tells us that $\widehat{\eta}^{\Sigma, \hat{E}}$ and $\widehat{\mu}^{\Sigma, \hat{E}}$ interact together like the unit and multiplication of a monad.

Lemma 3.55. *The following diagram commutes.⁴¹¹*

$$\begin{array}{ccccc}
 \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} & \xrightarrow{\widehat{\eta}_{\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}}^{\Sigma, \hat{E}}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} & \xleftarrow{\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\widehat{\eta}_{\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}}^{\Sigma, \hat{E}}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} \\
 & \searrow \text{id}_{\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}} & \downarrow \widehat{\mu}_{\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}}^{\Sigma, \hat{E}} & \swarrow \text{id}_{\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}} & \\
 & & \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} & &
 \end{array}$$

Proof. For the triangle on the left, we pave the following diagram.

$$\begin{array}{ccc}
 & \widehat{\eta}_{\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}}^{\Sigma, \hat{E}} & \\
 & \curvearrowright \text{(a)} & \\
 \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} & \xrightarrow{\widehat{\eta}_{\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}}^{\Sigma, \hat{E}}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} \xrightarrow{[-]_{\hat{E}}} \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} \\
 & \searrow \text{(b)} \llbracket [-] \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}} & \downarrow \widehat{\mu}_{\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}}^{\Sigma, \hat{E}} \\
 & \searrow \text{id}_{\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} \\
 & & \text{(c)} \downarrow \widehat{\mu}_{\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}}^{\Sigma, \hat{E}}
 \end{array} \quad (3.38)$$

⁴¹⁰ Once again, we are only adapting the argument from the proof of Lemma 1.42.

⁴¹¹ We reuse the proof of Lemma 1.43, although when using naturality of $[-]_{\hat{E}}$ in **Set**, we replace it by (3.19) which is not formally a naturality property (because \mathcal{T}_Σ is not a functor on **GMet**).

Showing (3.38) commutes:

- (a) Definition of $\widehat{\eta}_X^{\Sigma, \hat{E}}$ (3.34).
- (b) Definition of $\llbracket [-] \rrbracket_{\widehat{\mathbb{T}}\mathbf{X}}$ (1.29).
- (c) Definition of $\widehat{\mu}_X^{\Sigma, \hat{E}}$ (3.22).

For the triangle on the right, we show that $[-]_{\hat{E}} = \hat{\mu}_X^{\Sigma, \hat{E}} \circ \hat{\tau}_{\Sigma, \hat{E}} \hat{\eta}_X^{\Sigma, \hat{E}} \circ [-]_{\hat{E}}$ by paving (3.39), and we can conclude since $[-]_{\hat{E}}$ is epic that $\text{id}_{\hat{\tau}_{\Sigma, \hat{E}} X} = \hat{\mu}_X^{\Sigma, \hat{E}} \circ \hat{\tau}_{\Sigma, \hat{E}} \hat{\eta}_X^{\Sigma, \hat{E}}$.

$$\begin{array}{ccccc}
 & & \hat{\tau}_{\Sigma, \hat{E}} \hat{\eta}_X^{\Sigma, \hat{E}} & & \\
 & & \curvearrowright & & \\
 \mathcal{T}_{\Sigma} X & \xrightarrow{[-]_{\hat{E}}} & \hat{\tau}_{\Sigma, \hat{E}} X & \xrightarrow{\hat{\tau}_{\Sigma, \hat{E}} \hat{\eta}_X^{\Sigma, \hat{E}}} & \hat{\tau}_{\Sigma, \hat{E}} \mathcal{T}_{\Sigma} X & \xrightarrow{\hat{\tau}_{\Sigma, \hat{E}} [-]_{\hat{E}}} & \hat{\tau}_{\Sigma, \hat{E}} \hat{\tau}_{\Sigma, \hat{E}} X \\
 & \searrow & \text{(b)} & \uparrow & \text{(c)} & \nearrow & \\
 & & \hat{\tau}_{\Sigma} \hat{\eta}_X^{\Sigma} & & \mathcal{T}_{\Sigma} \mathcal{T}_{\Sigma} X & \xrightarrow{\hat{\tau}_{\Sigma} [-]_{\hat{E}}} & \hat{\tau}_{\Sigma} \hat{\tau}_{\Sigma, \hat{E}} X \\
 & \searrow & \text{(d)} & \downarrow & \text{(e)} & \nearrow & \\
 & & \text{id}_{\mathcal{T}_{\Sigma} X} & & \mathcal{T}_{\Sigma} X & \xrightarrow{[-]_{\hat{E}}} & \hat{\tau}_{\Sigma, \hat{E}} X \\
 & & & & & & \downarrow \hat{\mu}_X^{\Sigma, \hat{E}} \\
 & & & & & & \hat{\tau}_{\Sigma, \hat{E}} X
 \end{array} \quad (3.39)$$

Showing (3.39) commutes:

- (a) Definition of $\hat{\eta}_X^{\Sigma, \hat{E}}$ and functoriality of $\hat{\tau}_{\Sigma, \hat{E}}$.
- (b) ‘‘Naturality’’ of $[-]_{\hat{E}}$ (3.19).
- (c) By (3.19) again.
- (d) Definition of $\hat{\mu}_X^{\Sigma}$ (1.7).
- (e) By (3.21).
- (f) By (3.22).

□

Finally, we can show that $\hat{\mathbb{T}}_{\Sigma, \hat{E}} X$ is (Σ, \hat{E}) -algebra (analogous to Proposition 1.46).

Proposition 3.56. *For any space \mathbf{A} , the term algebra $\hat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{A}$ satisfies all the equations in \hat{E} .*

Proof. Let $\phi \in \hat{E}$ be an equation with context \mathbf{X} and $\hat{\iota} : \mathbf{X} \rightarrow \hat{\tau}_{\Sigma, \hat{E}} \mathbf{A}$ be a nonexpansive assignment. We factor $\hat{\iota}$ into⁴¹²

$$\hat{\iota} = X \xrightarrow{\hat{\eta}_X^{\Sigma, \hat{E}}} \hat{\tau}_{\Sigma, \hat{E}} X \xrightarrow{\hat{\tau}_{\Sigma, \hat{E}} \hat{\iota}} \hat{\tau}_{\Sigma, \hat{E}} \hat{\tau}_{\Sigma, \hat{E}} \mathbf{A} \xrightarrow{\hat{\mu}_{\mathbf{A}}^{\Sigma, \hat{E}}} \hat{\tau}_{\Sigma, \hat{E}} \mathbf{A}.$$

Now, Lemma 3.54 says that ϕ is satisfied in $\hat{\mathbb{T}}X$ under the assignment $\hat{\eta}_X^{\Sigma, \hat{E}}$. We also know by Lemma 3.15 that homomorphisms preserve satisfaction, so we can apply it twice using the facts that $\hat{\tau}_{\Sigma, \hat{E}} \hat{\iota}$ and $\hat{\mu}_{\mathbf{A}}^{\Sigma, \hat{E}}$ are homomorphisms (the former was shown after Lemma 3.47 and the latter in Lemma 3.50) to conclude that $\hat{\mathbb{T}}\mathbf{A}$ satisfies ϕ under $\hat{\mu}_{\mathbf{A}}^{\Sigma, \hat{E}} \circ \hat{\tau}_{\Sigma, \hat{E}} \hat{\iota} \circ \hat{\eta}_X^{\Sigma, \hat{E}} = \hat{\iota}$. □

We end this section just like we ended §1.3 by showing that $\hat{\mathbb{T}}X$ is the free (Σ, \hat{E}) -algebra.⁴¹³

Theorem 3.57. *For any space \mathbf{X} , the term algebra $\hat{\mathbb{T}}X$ is the free (Σ, \hat{E}) -algebra on \mathbf{X} .*

Proof. Note that the morphism witnessing freeness of $\hat{\mathbb{T}}X$ is $\hat{\eta}_X^{\Sigma, \hat{E}} : \mathbf{X} \rightarrow \hat{\tau}_{\Sigma, \hat{E}} X$.⁴¹⁴

Let $\hat{\mathbf{A}}$ be another (Σ, \hat{E}) -algebra and $f : \mathbf{X} \rightarrow \hat{\mathbf{A}}$ a nonexpansive function. We claim that $f^* = \llbracket - \rrbracket_{\hat{\mathbf{A}}} \circ \hat{\tau}_{\Sigma, \hat{E}} f$ is the unique homomorphism making the following commute.

$$\begin{array}{ccc}
 \text{in } \mathbf{GMet} & & \text{in } \mathbf{QAlg}(\Sigma, \hat{E}) \\
 \mathbf{X} \xrightarrow{\hat{\eta}_X^{\Sigma, \hat{E}}} \hat{\tau}_{\Sigma, \hat{E}} X & & \hat{\mathbb{T}}X \\
 \searrow f & \swarrow U & \downarrow f^* \\
 & & \hat{\mathbf{A}}
 \end{array}$$

⁴¹² This factoring is correct because

$$\begin{aligned}
 \hat{\iota} &= \text{id}_{\hat{\tau}_{\Sigma, \hat{E}} \mathbf{A}} \circ \hat{\iota} \\
 &= \hat{\mu}_{\mathbf{A}}^{\Sigma, \hat{E}} \circ \hat{\eta}_{\hat{\tau}_{\Sigma, \hat{E}} \mathbf{A}}^{\Sigma, \hat{E}} \circ \hat{\iota} && \text{Lemma 3.55} \\
 &= \hat{\mu}_{\mathbf{A}}^{\Sigma, \hat{E}} \circ \hat{\tau}_{\Sigma, \hat{E}} \hat{\iota} \circ \hat{\eta}_X^{\Sigma, \hat{E}}. && \text{naturality of } \hat{\eta}_X^{\Sigma, \hat{E}}
 \end{aligned}$$

⁴¹³ In both [MSV22] and [MSV23], we constructed the free algebra using quantitative equational logic. This is an alternative proof that does not rely on the logic.

⁴¹⁴ As expected, the proof goes exactly like for Proposition 1.49 except for dealing with nonexpansiveness at the end.

First, f^* is a homomorphism because it is the composite of two homomorphisms $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} f$ (by Lemma 3.48) and $\llbracket - \rrbracket_A$ (by Lemma 3.49 since $\hat{\mathbb{A}}$ satisfies \hat{E}). Next, the triangle commutes by the following derivation.

$$\begin{aligned}
 \llbracket - \rrbracket_A \circ \widehat{\mathcal{T}}_{\Sigma, \hat{E}} f \circ \widehat{\eta}_{\mathbf{X}}^{\Sigma, \hat{E}} &= \llbracket - \rrbracket_A \circ \widehat{\eta}_{\hat{\mathbb{A}}}^{\Sigma, \hat{E}} \circ f && \text{by (3.35)} \\
 &= \llbracket - \rrbracket_A \circ [-]_{\hat{E}} \circ \eta_{\hat{\mathbb{A}}}^{\Sigma} \circ f && \text{definition of } \widehat{\eta}^{\Sigma, \hat{E}} \\
 &= \llbracket - \rrbracket_A \circ \eta_{\hat{\mathbb{A}}}^{\Sigma} \circ f && \text{Footnote 405} \\
 &= f && \text{definition of } \llbracket - \rrbracket_A \text{ (3.20)}
 \end{aligned}$$

Finally, uniqueness follows from the inductive definition of $\widehat{\mathbb{T}\mathbf{X}}$ and the homomorphism property. Briefly, if we know the action of a homomorphism on equivalence classes of terms of depth 0, we can infer all of its action because all other classes of terms can be obtained by applying operation symbols.⁴¹⁵

It remains to show that $f^* : \widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{X} \rightarrow \mathbf{A}$ is nonexpansive. This follows by the following derivation, where we implicitly use nonexpansiveness of f in the second step, where f is used as a nonexpansive assignment.

$$\begin{aligned}
 d_{\hat{E}}([s]_{\hat{E}}, [t]_{\hat{E}}) \leq \varepsilon &\iff \mathbf{X} \vdash s =_{\varepsilon} t \in \Omega \mathfrak{T} \mathfrak{h}(\hat{E}) && \text{by (3.17)} \\
 &\implies d_{\mathbf{A}}(\llbracket s \rrbracket_A^f, \llbracket t \rrbracket_A^f) \leq \varepsilon && \hat{\mathbb{A}} \in \mathbf{QAlg}(\Sigma, \hat{E}) \\
 &\iff d_{\mathbf{A}}(\llbracket \mathcal{T}_{\Sigma} f(s) \rrbracket_A, \llbracket \mathcal{T}_{\Sigma} f(t) \rrbracket_A) && \text{by (1.10)} \\
 &\iff d_{\mathbf{A}}(\llbracket [\mathcal{T}_{\Sigma} f(s)]_{\hat{E}} \rrbracket_A, \llbracket [\mathcal{T}_{\Sigma} f(t)]_{\hat{E}} \rrbracket_A) && \text{Footnote 405} \\
 &\iff d_{\mathbf{A}}(\llbracket \widehat{\mathcal{T}}_{\Sigma, \hat{E}} f[s]_{\hat{E}} \rrbracket_A, \llbracket \widehat{\mathcal{T}}_{\Sigma, \hat{E}} f[t]_{\hat{E}} \rrbracket_A) && \text{by (3.19)} \\
 &\iff d_{\mathbf{A}}(f^*[s]_{\hat{E}}, f^*[t]_{\hat{E}}) && \text{definition of } f^* \quad \square
 \end{aligned}$$

Since we have a free (Σ, \hat{E}) -algebra $\widehat{\mathbb{T}\mathbf{X}}$ for every generalized metric space \mathbf{X} , we get a left adjoint to $U : \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{GMet}$. This automatically yields a monad structure on $\widehat{\mathcal{T}}_{\Sigma, \hat{E}}$ that we will study after developing quantitative equational logic. Before that, we make use of a special case of the adjunction above.

Corollary 3.58. *The forgetful functor $U : \mathbf{GMet} \rightarrow \mathbf{Set}$ has a left adjoint.*

Proof. The following adjoints compose to yield a left adjoint to $U : \mathbf{GMet} \rightarrow \mathbf{Set}$.⁴¹⁶

$$\begin{array}{ccc}
 & & U \\
 & \curvearrowright & \\
 \mathbf{GMet} & \xrightarrow{\tau} & \mathbf{LSpa} \xrightarrow{\tau} \mathbf{Set} \\
 & \xleftarrow{\tau} & \xleftarrow{\tau}
 \end{array} \quad \square$$

Example 3.59 (Discrete metric). To make this more concrete, one can wonder what is the free metric space on a set X (with $\mathbf{L} = [0, 1]$). According to the diagram above, we first need to construct the discrete space \mathbf{X}_{\top} on X , then construct the free metric space on \mathbf{X}_{\top} . We know how to do the first step (Proposition 2.60), and the second step is also fairly easy to do.⁴¹⁷ The only thing that prevents \mathbf{X}_{\top} from being a metric is reflexivity, i.e. $d_{\top}(x, x) = 1 \neq 0$. If we define $d_{\mathbf{X}}$ just like d_{\top} except with $d_{\mathbf{X}}(x, x) = 0$, then it is a metric,⁴¹⁸ and $(X, d_{\mathbf{X}})$ is the free metric space over X .

Corollary 3.58 applies to any category \mathbf{GMet} , so we can always construct the discrete generalized metric on a set.

⁴¹⁵ Formally, let $f, g : \widehat{\mathbb{T}\mathbf{X}} \rightarrow \hat{\mathbb{A}}$ be two homomorphisms such that for any $x \in X$, $f[x]_{\hat{E}} = g[x]_{\hat{E}}$, then, we can show that $f = g$. For any $t \in \widehat{\mathcal{T}}_{\Sigma} X$, we showed in Lemma 3.53 that $[t]_{\hat{E}} = \llbracket t \rrbracket_{\widehat{\mathbb{T}\mathbf{X}}}^{\Sigma, \hat{E}}$. Then using (1.12), we have

$$f[t]_{\hat{E}} = \llbracket t \rrbracket_A^{f \circ \widehat{\eta}_{\mathbf{X}}^{\Sigma, \hat{E}}} = \llbracket t \rrbracket_A^{g \circ \widehat{\eta}_{\mathbf{X}}^{\Sigma, \hat{E}}} = g[t]_{\hat{E}},$$

where the second inequality follows by hypothesis that f and g agree on equivalence classes of terms of depth 0.

⁴¹⁶ The adjunction between \mathbf{LSpa} and \mathbf{Set} was described in Proposition 2.60. The adjunction between \mathbf{GMet} and \mathbf{LSpa} is the one we just obtained via Theorem 3.57 that we instantiate with $\mathbf{GMet} = \mathbf{QAlg}(\emptyset, \hat{E}_{\mathbf{GMet}})$ (recall Example 3.18).

⁴¹⁷ Even though we said in Example 3.27 that the free metric space on an arbitrary \mathbf{X} is harder to describe.

⁴¹⁸ Identity of indiscernibles and symmetry hold because $d_{\mathbf{X}}(x, y) = d_{\mathbf{X}}(y, x) = 1$ when $x \neq y$. The triangle inequality holds because

$$d_{\mathbf{X}}(x, z) = 1 \leq 1 + 1 = d_{\mathbf{X}}(x, y) + d_{\mathbf{X}}(y, z).$$

With the help of quantitative algebraic theories and free algebras, we can now define coproducts inside **GMet**.

Corollary 3.60. *The category **GMet** has coproducts.*

Proof. We will only do the case of binary coproducts for exposition's sake, but the proof can be adapted to arbitrary families. For any generalized metric space \mathbf{A} , the quantitative algebraic theory of \mathbf{A} is generated by the signature $\Sigma_{\mathbf{A}} = \{a : 0 \mid a \in A\}$ and the quantitative equations⁴¹⁹

$$\hat{E}_{\mathbf{A}} = \left\{ \vdash a =_{d_{\mathbf{A}}(a,a')} a' \mid a, a' \in A \right\}.$$

A $(\Sigma_{\mathbf{A}}, \hat{E}_{\mathbf{A}})$ -algebra $\hat{\mathbf{B}}$ is a generalized metric space \mathbf{B} equipped with an interpretation $\llbracket a \rrbracket_{\mathbf{B}}$ for every $a \in A$ such that $d_{\mathbf{B}}(\llbracket a \rrbracket_{\mathbf{B}}, \llbracket a' \rrbracket_{\mathbf{B}}) \leq d_{\mathbf{A}}(a, a')$ for every $a, a' \in A$. Equivalently, all the interpretations can be seen as a single nonexpansive map $\llbracket - \rrbracket_{\mathbf{B}} : \mathbf{A} \rightarrow \mathbf{B}$. Therefore, $\mathbf{QAlg}(\Sigma_{\mathbf{A}}, \hat{E}_{\mathbf{A}})$ is the coslice category \mathbf{A}/\mathbf{GMet} .

Given another space \mathbf{A}' , if we combine the theories of \mathbf{A} and \mathbf{A}' with no additional equations, we get the category $\mathbf{QAlg}(\Sigma_{\mathbf{A}} + \Sigma_{\mathbf{A}'}, \hat{E}_{\mathbf{A}} + \hat{E}_{\mathbf{A}'})$ of spaces \mathbf{B} equipped with two nonexpansive maps $\llbracket - \rrbracket_{\mathbf{B}} : \mathbf{A} \rightarrow \mathbf{B}$ and $\llbracket - \rrbracket'_{\mathbf{B}} : \mathbf{A}' \rightarrow \mathbf{B}$. This category has an initial object, the free algebra on the initial generalized metric space from Proposition 2.41. Moreover, this category can be equivalently described as the comma category $[\mathbf{A}, \mathbf{A}'] \downarrow \text{id}_{\mathbf{GMet}}$ where $[\mathbf{A}, \mathbf{A}'] : \mathbf{1} + \mathbf{1} \rightarrow \mathbf{GMet}$ is the constant functor sending the two objects in the domain to \mathbf{A} and \mathbf{A}' respectively.⁴²⁰ The initial object of this category (we just showed it exists) is the coproduct $\mathbf{A} + \mathbf{A}'$ (by definition of coproducts and comma categories). \square

⁴¹⁹ Note that a and a' are seen as constants, not variables, so the context of these equations is the empty L-space.

⁴²⁰ The category $\mathbf{1} + \mathbf{1}$ has two objects, their identity morphisms, and that is it.

Abstract Quantitative Equations

We finish this section like we finished §1.3: by giving an equivalent definition for quantitative equations. Recall that an abstract equation is a surjective homomorphism $e : \mathcal{T}_{\Sigma}X \twoheadrightarrow \mathbb{Y}$ in $\mathbf{Alg}(\Sigma)$. To generalize, we clearly want to replace $\mathbf{Alg}(\Sigma)$ with $\mathbf{QAlg}(\Sigma)$, but it is less clear how to replace $\mathcal{T}_{\Sigma}X$ and surjective homomorphisms.

Examining the proofs of Propositions 1.51 and 1.52, we can recognize two important properties of $\mathcal{T}_{\Sigma}X$:

- $\mathcal{T}_{\Sigma}X$ is the free algebra over X , and
- given an algebra $\mathbf{A} \in \mathbf{Alg}(\Sigma)$ and an assignment $\iota : X \rightarrow A$, $\llbracket - \rrbracket'_{\mathbf{A}} : \mathcal{T}_{\Sigma}X \rightarrow A$ is the unique homomorphism satisfying $\llbracket - \rrbracket'_{\mathbf{A}} \circ \eta_X^{\Sigma} = \iota$.

Let $\hat{\mathcal{T}}_{\Sigma}X = \hat{\mathcal{T}}_{\Sigma, \emptyset}X$ denote the free quantitative Σ -algebra and its underlying space, its elements are not necessarily terms over X because \equiv_{\emptyset} might be non-trivial even when generated by no quantitative equations.⁴²¹ To replace $\llbracket - \rrbracket'_{\mathbf{A}}$, we want, for every quantitative algebra $\hat{\mathbf{A}}$ and nonexpansive assignment $\hat{\iota} : X \rightarrow \mathbf{A}$, a homomorphism $\hat{\mathcal{T}}_{\Sigma}X \rightarrow \hat{\mathbf{A}}$ in $\mathbf{QAlg}(\Sigma)$ that acts like $\hat{\iota}$ on (equivalence classes) of variables. It exists

⁴²¹ For an extreme example, if $\hat{E}_{\mathbf{GMet}}$ contains $x, y \vdash x = y$, then any generalized metric space must be empty or a singleton, hence $\hat{\mathcal{T}}_{\Sigma}X$ contains one equivalence class of terms (unless X is empty and Σ has no constants, then $\hat{\mathcal{T}}_{\Sigma}X$ is empty).

and is unique by the universal property of $\widehat{\mathcal{T}}_\Sigma \mathbf{X}$, we denote it by \hat{i}^\sharp :

$$\begin{array}{ccc}
 \mathbf{X} & \xrightarrow{\hat{\eta}_\mathbf{X}^{\Sigma, \emptyset}} & \widehat{\mathcal{T}}_\Sigma \mathbf{X} \\
 \searrow \hat{i} & & \downarrow \hat{i}^\sharp \\
 & & \mathbf{A}
 \end{array}
 \quad \leftarrow u \quad
 \begin{array}{ccc}
 & & \widehat{\mathcal{T}}_\Sigma \mathbf{X} \\
 & & \downarrow \hat{i}^\sharp \\
 & & \hat{\mathbf{A}}
 \end{array}
 \quad (3.40)$$

Now, how should surjective homomorphisms in $\mathbf{Alg}(\Sigma)$ be generalized in $\mathbf{QAlg}(\Sigma)$? The instinctive answer for a category theorist would be homomorphisms in $\mathbf{QAlg}(\Sigma)$ such that the underlying morphism in \mathbf{GMet} is an epimorphism.⁴²² However, we will see that the right choice is actually still surjective homomorphisms.

Definition 3.61. An **abstract quantitative equation** is a surjective nonexpansive homomorphism $e : \widehat{\mathcal{T}}_\Sigma \mathbf{X} \rightarrow \hat{\mathbf{Y}}$ in $\mathbf{QAlg}(\Sigma)$.⁴²³ We say that a quantitative algebra $\hat{\mathbf{A}}$ **satisfies** e if for any nonexpansive assignment $\hat{i} : \mathbf{X} \rightarrow \mathbf{A}$, the homomorphism \hat{i}^\sharp factors through e in $\mathbf{QAlg}(\Sigma)$:

$$\hat{i}^\sharp = \widehat{\mathcal{T}}_\Sigma \mathbf{X} \xrightarrow{e} \hat{\mathbf{Y}} \xrightarrow{h} \hat{\mathbf{A}}.$$

We say that $\hat{\mathbf{A}}$ satisfies a class of abstract quantitative equations if it satisfies all of its elements.

We now show how abstract quantitative equations and quantitative equations have the same expressive power. The intuition and proofs are very similar to the classical case.

Proposition 3.62. *If \hat{E} is a class of abstract quantitative equations, then there is a class \hat{E}° of quantitative equations such that $\hat{\mathbf{A}}$ satisfies \hat{E} if and only if it satisfies \hat{E}° .*

Proof. We construct \hat{E}° similarly to the classical case, but we have to add the distance information too:⁴²⁴

$$\begin{aligned}
 \hat{E}^\circ = & \left\{ \mathbf{X} \vdash s = t \mid e : \widehat{\mathcal{T}}_\Sigma \mathbf{X} \rightarrow \hat{\mathbf{Y}} \in \hat{E}, s, t \in \mathcal{T}_\Sigma \mathbf{X}, e[s]_\emptyset = e[t]_\emptyset \right\} \\
 & \cup \left\{ \mathbf{X} \vdash s =_\varepsilon t \mid e : \widehat{\mathcal{T}}_\Sigma \mathbf{X} \rightarrow \hat{\mathbf{Y}} \in \hat{E}, s, t \in \mathcal{T}_\Sigma \mathbf{X}, d_{\mathbf{Y}}(e[s]_\emptyset, e[t]_\emptyset) \leq \varepsilon \right\}.
 \end{aligned}$$

We will show that $\hat{\mathbf{A}}$ satisfies \hat{E} if and only if it satisfies \hat{E}° .

(\Rightarrow) Suppose $e : \widehat{\mathcal{T}}_\Sigma \mathbf{X} \rightarrow \hat{\mathbf{Y}}$ belongs to \hat{E} and fix $s, t \in \mathcal{T}_\Sigma \mathbf{X}$. For any nonexpansive assignment $\hat{i} : \mathbf{X} \rightarrow \mathbf{A}$, the factorization $\hat{i}^\sharp = h \circ e$ implies that $\llbracket - \rrbracket_A^{\hat{i}}$ factors through $e[-]_\emptyset$. Indeed, let us look at (3.41).⁴²⁵

$$\begin{array}{ccccccc}
 & & \hat{\eta}_\mathbf{X}^{\Sigma, \emptyset} & & & & \\
 & & \text{(a)} & & & & \\
 \mathbf{X} & \xrightarrow{\hat{\eta}_\mathbf{X}^\Sigma} & \mathcal{T}_\Sigma \mathbf{X} & \xrightarrow{[-]_\emptyset} & \widehat{\mathcal{T}}_\Sigma \mathbf{X} & \xrightarrow{e} & \hat{\mathbf{Y}} \\
 & \searrow & \downarrow & \searrow & \downarrow & \downarrow & \downarrow h \\
 & & & \text{(c)} & & \text{(d)} & \\
 & & & \llbracket - \rrbracket_A^{\hat{i}} & & \hat{i}^\sharp & \\
 & & & & & & \hat{\mathbf{A}} \\
 & \searrow \hat{i} & & & & & \\
 & & & & & &
 \end{array}
 \quad (3.41)$$

⁴²² Because surjective functions in \mathbf{Set} are precisely the epimorphisms.

⁴²³ We reiterate that the terminology comes from [JMU24], the following arguments are inspired by their proof of [JMU24, Theorem 4.16].

⁴²⁴ One small change is that abstract quantitative equations are functions taking equivalence classes of terms as inputs rather than just terms. Recall that $[t]_\emptyset$ denotes the equivalence class of t in $\widehat{\mathcal{T}}_\Sigma \mathbf{X}$, and we write $e[t]_\emptyset$ instead of the more clunky $e(\{t\}_\emptyset)$.

⁴²⁵ We are implicitly going back and forth between the categories \mathbf{Set} , \mathbf{GMet} , $\mathbf{Alg}(\Sigma)$, and $\mathbf{QAlg}(\Sigma)$.

The triangles (a) and (b) commute by definition of $\widehat{\eta}_{\mathbf{X}}^{\Sigma, \emptyset}$ and $\llbracket - \rrbracket_A^{\hat{t}}$ respectively. The triangle combining (a), (b), and (c) commutes by definition of $\hat{t}^{\#}$, and since $\hat{t}^{\#} \circ [-]_{\emptyset}$ is a homomorphism in $\mathbf{Alg}(\Sigma)$,⁴²⁶ by the universal property of $\widehat{\mathcal{T}}_{\Sigma} X$ (uniqueness of $\llbracket - \rrbracket_A^{\hat{t}}$), the triangle (c) commutes as well. The hypothesis that (d) commutes means that $\llbracket - \rrbracket_A^{\hat{t}} = h \circ e[-]_{\emptyset}$.

Therefore, if $\mathbf{X} \vdash s = t$ belongs to \hat{E}° because $e[s]_{\emptyset} = e[t]_{\emptyset}$, then $\llbracket s \rrbracket_A^{\hat{t}} = \llbracket t \rrbracket_A^{\hat{t}}$ must hold for all \hat{t} ,⁴²⁷ so $\hat{\mathbf{A}} \models \mathbf{X} \vdash s = t$. Similarly, if $\mathbf{X} \vdash s =_{\varepsilon} t$ belongs to \hat{E}° because $d_Y(e[s]_{\emptyset}, e[t]_{\emptyset}) \leq \varepsilon$, then $d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{t}}, \llbracket t \rrbracket_A^{\hat{t}}) \leq \varepsilon$ must hold for all \hat{t} ,⁴²⁸ so $\hat{\mathbf{A}} \models \mathbf{X} \vdash s =_{\varepsilon} t$. This works for every $e \in \hat{E}$, so we conclude that $\hat{\mathbf{A}}$ satisfies all the quantitative equations in \hat{E}° .

(\Leftarrow) Let $e : \widehat{\mathcal{T}}_{\Sigma} \mathbf{X} \rightarrow \hat{Y}$ belong to \hat{E} . For any nonexpansive assignment $\hat{t} : \mathbf{X} \rightarrow \mathbf{A}$, since e is surjective, we can define a function $h : Y \rightarrow A$ by $h(y) = \llbracket t_y \rrbracket_A^{\hat{t}}$ with t_y a representative of an equivalence class in $e^{-1}(y)$. In other words, t_y is an element in the preimage of y under $e[-]_{\emptyset}$. Surjectivity of e means h is defined on all Y , and the choice of t_y does not matter because if t'_y belongs to an equivalence class in $e^{-1}(y)$, then $e[t_y]_{\emptyset} = y = e[t'_y]_{\emptyset}$ implies $\llbracket t_y \rrbracket_A^{\hat{t}} = \llbracket t'_y \rrbracket_A^{\hat{t}}$ because $\hat{\mathbf{A}} \models \hat{E}^{\circ}$.⁴²⁹

Our definition of h ensures $\hat{t}^{\#} = h \circ e$ since any element T in $\widehat{\mathcal{T}}_{\Sigma} \mathbf{X}$ is equal to $[t_y]_{\emptyset}$ for some term t_y in the equivalence class of $e^{-1}(y)$ for some $y \in Y$, which yields⁴³⁰

$$(h \circ e)(T) = (h \circ e)[t_y]_{\emptyset} = h(y) = \llbracket t_y \rrbracket_A^{\hat{t}} \stackrel{(*)}{=} \hat{t}^{\#}[t_y]_{\emptyset} = \hat{t}^{\#}(T).$$

It remains to show that h is a nonexpansive homomorphism.

For any $y, y' \in Y$, pick t_y and $t_{y'}$ such that $e[t_y]_{\emptyset} = y$ and $e[t_{y'}]_{\emptyset} = y'$. Since $d_Y(e[t_y]_{\emptyset}, e[t_{y'}]_{\emptyset}) \leq d_Y(y, y')$, the quantitative equation $\mathbf{X} \vdash t_y =_{d_Y(y, y')} t_{y'}$ belongs to \hat{E}° , so $\hat{\mathbf{A}}$ satisfies it by hypothesis. Therefore,

$$d_{\mathbf{A}}(h(y), h(y')) = d_{\mathbf{A}}(\llbracket t_y \rrbracket_A^{\hat{t}}, \llbracket t_{y'} \rrbracket_A^{\hat{t}}) \leq d_Y(y, y') \leq \varepsilon.$$

We conclude that h is nonexpansive. The argument for h preserving operations is copied from the proof of Proposition 1.51, replacing e with $e[-]_{\emptyset}$ which is a homomorphism in $\mathbf{Alg}(\Sigma)$ by (3.21). For any $\text{op} : n \in \Sigma$ and $y_1, \dots, y_n \in Y$, pick t_i in the preimage of y_i under $e[-]_{\emptyset}$, then we have

$$\begin{aligned} h(\llbracket \text{op} \rrbracket_Y(y_1, \dots, y_n)) &= h(\llbracket \text{op} \rrbracket_Y(e[t_1]_{\emptyset}, \dots, e[t_n]_{\emptyset})) && \text{definition of } t_i \\ &= h \circ e[\text{op}(t_1, \dots, t_n)]_{\emptyset} && e[-]_{\emptyset} \text{ is a homomorphism} \\ &= \llbracket \text{op}(t_1, \dots, t_n) \rrbracket_A^{\hat{t}} && \text{definition of } h \\ &= \llbracket \text{op} \rrbracket_A(\llbracket t_1 \rrbracket_A^{\hat{t}}, \dots, \llbracket t_n \rrbracket_A^{\hat{t}}) && \text{by (1.15)} \\ &= \llbracket \text{op} \rrbracket_A(h \circ e[t_1]_{\emptyset}, \dots, h \circ e[t_n]_{\emptyset}) && \text{definition of } h \\ &= \llbracket \text{op} \rrbracket_A(h(y_1), \dots, h(y_n)). && \text{definition of } t_i \quad \square \end{aligned}$$

Proposition 3.63. *If \hat{E} is a class of quantitative equations, then there is a class \hat{E}^{\bullet} of abstract quantitative equations such that $\hat{\mathbf{A}}$ satisfies \hat{E} if and only if it satisfies \hat{E}^{\bullet} .*

Proof. Given a quantitative equation ϕ with context \mathbf{X} , we let ϕ^{\bullet} be the homomorphism $\widehat{\mathcal{T}}_{\Sigma} \mathbf{X} \rightarrow \widehat{\mathcal{T}}_{\Sigma, \{\phi\}} \mathbf{X}$ that uniquely makes the following diagram commute:⁴³¹

⁴²⁶ Since $\hat{t}^{\#}$ is a homomorphism in $\mathbf{QAlg}(\Sigma)$, the underlying function is a homomorphism in $\mathbf{Alg}(\Sigma)$, and $[-]_{\emptyset}$ is a homomorphism by (3.21).

⁴²⁷ Because

$$\llbracket s \rrbracket_A^{\hat{t}} = h(e[s]_{\emptyset}) = h(e[t]_{\emptyset}) = \llbracket t \rrbracket_A^{\hat{t}}.$$

⁴²⁸ Because h is nonexpansive so

$$\begin{aligned} d_{\mathbf{A}}(\llbracket s \rrbracket_A^{\hat{t}}, \llbracket t \rrbracket_A^{\hat{t}}) &= d_{\mathbf{A}}(h(e[s]_{\emptyset}), h(e[t]_{\emptyset})) \\ &\leq d_Y(e[s]_{\emptyset}, e[t]_{\emptyset}) \\ &\leq \varepsilon. \end{aligned}$$

⁴²⁹ By definition, \hat{E}° contains $\mathbf{X} \vdash t_y = t'_y$ because $e[t_y]_{\emptyset} = e[t'_y]_{\emptyset}$.

⁴³⁰ The equation marked $(*)$ holds by (c) commuting in (3.41).

⁴³¹ The existence and uniqueness of ϕ^{\bullet} are consequences of $\widehat{\mathcal{T}}_{\Sigma} \mathbf{X}$ being the free object in $\mathbf{QAlg}(\Sigma)$ generated by \mathbf{X} , and $\widehat{\mathcal{T}}_{\Sigma, \{\phi\}}$ being the underlying space of $\widehat{\mathcal{T}}_{\Sigma, \{\phi\}} \mathbf{X} \in \mathbf{QAlg}(\Sigma)$.

$$\begin{array}{ccc}
 \mathbf{X} & \xrightarrow{\widehat{\eta}_{\mathbf{X}}^{\Sigma, \emptyset}} & \widehat{\mathcal{T}}_{\Sigma} \mathbf{X} \\
 & \searrow \widehat{\eta}_{\mathbf{X}}^{\Sigma, \{\phi\}} & \downarrow \phi^{\bullet} \\
 & & \widehat{\mathcal{T}}_{\Sigma, \{\phi\}} \mathbf{X}
 \end{array}$$

We first need to argue that ϕ^{\bullet} is surjective. We show something stronger:

$$\forall t \in \mathcal{T}_{\Sigma} \mathbf{X}, \quad \phi^{\bullet}[t]_{\emptyset} = [t]_{\{\phi\}}. \quad (3.42)$$

We proceed by induction. The base case follows from the triangle above. For the inductive step, suppose $t = \text{op}(t_1, \dots, t_n)$, using that ϕ^{\bullet} is a homomorphism and the definition of $[-]_{-}$, we have

$$\begin{aligned}
 \phi^{\bullet}[t]_{\emptyset} &= \phi^{\bullet}(\llbracket \text{op} \rrbracket_{\widehat{\mathcal{T}}_{\Sigma, \mathbf{X}}}([t_1]_{\emptyset}, \dots, [t_n]_{\emptyset})) && \text{by (3.20)} \\
 &= \llbracket \text{op} \rrbracket_{\widehat{\mathcal{T}}_{\Sigma, \{\phi\}} \mathbf{X}}(\phi^{\bullet}[t_1]_{\emptyset}, \dots, \phi^{\bullet}[t_n]_{\emptyset}) && \phi^{\bullet} \text{ is a homomorphism} \\
 &= \llbracket \text{op} \rrbracket_{\widehat{\mathcal{T}}_{\Sigma, \{\phi\}} \mathbf{X}}([t_1]_{\{\phi\}}, \dots, [t_n]_{\{\phi\}}) && \text{I.H.} \\
 &= [t]_{\{\phi\}}. && \text{by (3.20)}
 \end{aligned}$$

It follows that ϕ^{\bullet} is surjective.⁴³²

Now, we show that an algebra $\widehat{\mathbf{A}}$ satisfies ϕ if and only if it satisfies ϕ^{\bullet} .

(\Rightarrow) If $\widehat{\mathbf{A}} \models \phi$, then for any assignment $\hat{i} : \mathbf{X} \rightarrow \mathbf{A}$, we have the following unique factorization because $\widehat{\mathcal{T}}_{\Sigma, \{\phi\}} \mathbf{X}$ is the free $(\Sigma, \{\phi\})$ -algebra, and $\widehat{\mathbf{A}} \in \mathbf{QAlg}(\Sigma, \{\phi\})$:

$$\begin{array}{ccc}
 \mathbf{X} & \xrightarrow{\widehat{\eta}_{\mathbf{X}}^{\Sigma, \{\phi\}}} & \widehat{\mathcal{T}}_{\Sigma, \{\phi\}} \mathbf{X} \\
 & \searrow \hat{i} & \downarrow h \\
 & & \widehat{\mathbf{A}}
 \end{array} \quad (3.43)$$

We can further decompose (3.43) with another factorization.

$$\begin{array}{ccccc}
 & & \widehat{\eta}_{\mathbf{X}}^{\Sigma, \{\phi\}} & & \\
 & & \curvearrowright & & \\
 \mathbf{X} & \xrightarrow{\widehat{\eta}_{\mathbf{X}}^{\Sigma, \emptyset}} & \widehat{\mathcal{T}}_{\Sigma} \mathbf{X} & \xrightarrow{\phi^{\bullet}} & \widehat{\mathcal{T}}_{\Sigma, \{\phi\}} \mathbf{X} \\
 & & \searrow \hat{i}^{\sharp} & \searrow \hat{i}^{\sharp} & \downarrow h \\
 & & & & \widehat{\mathbf{A}} \\
 & \searrow \hat{i} & & & \\
 & & & &
 \end{array} \quad (3.44)$$

The triangles (a) and (b) commute by definition of ϕ^{\bullet} and \hat{i}^{\sharp} respectively. Also, \hat{i}^{\sharp} is the unique homomorphism making (b) commute. Since the triangle combining (a), (b), and (c) commutes by (3.43), the composite $h \circ \phi^{\bullet}$ also makes (b) commute, so it is equal to \hat{i}^{\sharp} , i.e. (c) commutes. This gives the desired factorization $\hat{i}^{\sharp} = h \circ \phi^{\bullet}$, thus $\widehat{\mathbf{A}}$ satisfies ϕ^{\bullet} .

(\Leftarrow) Suppose $\widehat{\mathbf{A}}$ satisfies ϕ^{\bullet} , we consider two cases separately. If $\phi = \mathbf{X} \vdash s = t$, then we have the following derivation for any $\hat{i} : \mathbf{X} \rightarrow \mathbf{A}$:

$$\mathbf{X} \vdash s = t \in \Omega \mathfrak{Th}(\{\phi\}) \implies [s]_{\{\phi\}} = [t]_{\{\phi\}} \quad \text{definition of } [-]_{\{\phi\}}$$

⁴³² Any element of $\widehat{\mathcal{T}}_{\Sigma, \{\phi\}}$ is an equivalence class $[t]_{\{\phi\}}$ for some $t \in \mathcal{T}_{\Sigma} \mathbf{X}$, hence it is the image of $[t]_{\emptyset}$ under ϕ^{\bullet} .

$$\begin{aligned}
 &\implies \phi^\bullet[s]_\emptyset = \phi^\bullet[t]_\emptyset && \text{by (3.42)} \\
 &\implies \hat{t}^\sharp[s]_\emptyset = \hat{t}^\sharp[t]_\emptyset && \text{by } \hat{t}^\sharp = h \circ \phi^\bullet \\
 &\implies \llbracket s \rrbracket_A^\hat{t} = \llbracket t \rrbracket_A^\hat{t}. && \text{by (c) in (3.41)}
 \end{aligned}$$

Thus, $\hat{\mathbb{A}} \models \phi$. If $\phi = \mathbf{X} \vdash s =_\varepsilon t$, we have almost the same derivation for all $\hat{t} : \mathbf{X} \rightarrow \mathbf{A}$.

$$\begin{aligned}
 \mathbf{X} \vdash s =_\varepsilon t \in \Omega\mathfrak{Ih}(\{\phi\}) &\implies d_{\{\phi\}}([s]_{\{\phi\}}, [t]_{\{\phi\}}) \leq \varepsilon && \text{by (3.14)} \\
 &\implies d_{\{\phi\}}(\phi^\bullet[s]_\emptyset, \phi^\bullet[t]_\emptyset) \leq \varepsilon && \text{by (3.42)} \\
 &\implies d_{\mathbf{A}\{\phi\}}(h(\phi^\bullet[s]_\emptyset), h(\phi^\bullet[t]_\emptyset)) \leq \varepsilon && h \text{ is nonexpansive} \\
 &\implies d_{\mathbf{A}}(\hat{t}^\sharp[s]_\emptyset, \hat{t}^\sharp[t]_\emptyset) \leq \varepsilon && \text{by } \hat{t}^\sharp = h \circ \phi^\bullet \\
 &\implies d_{\mathbf{A}}(\llbracket s \rrbracket_A^\hat{t}, \llbracket t \rrbracket_A^\hat{t}). && \text{by (c) in (3.41)}
 \end{aligned}$$

Thus, $\hat{\mathbb{A}} \models \phi$.

Now, given a class \hat{E} of quantitative equations, $\hat{\mathbb{A}}$ satisfies \hat{E} if and only if it satisfies $\hat{E}^\bullet = \{\phi^\bullet \mid \phi \in \hat{E}\}$. \square

Let us give an example showcasing why surjective homomorphisms were the right choice rather than homomorphisms with underlying epimorphism in \mathbf{GMet} .

Example 3.64. Let \mathbf{GMet} be the category \mathbf{Met} of metric spaces and Σ be empty, so $\mathbf{QAlg}(\Sigma) = \mathbf{Met}$. We consider the epimorphism $e : \widehat{\mathcal{T}}_\emptyset \mathbf{Q} = \mathbf{Q} \rightarrow \mathbb{R}$ that is the injection of the rationals into the reals (with the Euclidean metric).⁴³³ It is an epimorphism because nonexpansive maps out of \mathbb{R} are determined by their value on \mathbf{Q} (see Footnote 323). The subcategory of quantitative algebras that satisfy e is not closed under subalgebras (in this case, they are subspaces) because \mathbb{R} satisfies e but \mathbf{Q} does not. Hence, this subcategory is not a quantitative variety.

To show \mathbf{Q} does not satisfy e , take the identity assignment $\hat{t} : \mathbf{Q} \rightarrow \mathbf{Q}$, and note that $\hat{t}^\sharp = \hat{t}$ does not factor through e because any nonexpansive (hence continuous) function $\mathbb{R} \rightarrow \mathbf{Q}$ is constant.⁴³⁴

To show \mathbb{R} satisfies e , let $\hat{t} : \mathbf{Q} \rightarrow \mathbb{R}$ be a nonexpansive assignment. We extend \hat{t} to $h : \mathbb{R} \rightarrow \mathbb{R}$ in the canonical way setting $h(r) = \lim_n \hat{t}(r_n)$, where $\{r_n\}_{n \in \mathbb{N}}$ is a sequence of rationals converging to r . Note that h extends \hat{t} because if $r \in \mathbf{Q}$, $h(r) = \lim_n \hat{t}(r) = \hat{t}(r)$. The choice of sequence does not matter because \hat{t} is nonexpansive hence continuous. It also follows from nonexpansiveness of $\hat{t} : \mathbf{Q} \rightarrow \mathbb{R}$ and continuity of the Euclidean distance that $d(\lim_n \hat{t}(r_n), \lim_n \hat{t}(r'_n)) \leq d(\lim_n r_n, \lim_n r'_n)$, hence the extension $h : \mathbb{R} \rightarrow \mathbb{R}$ is also nonexpansive. We obtain the factorization $\hat{t} = \hat{t}^\sharp = h \circ e$.

With the equivalence between quantitative equations and abstract quantitative equations, we can use abstract results from the literature to prove the variety theorem for $\mathbf{GMet} = \mathbf{LSpa}$ and $\Sigma = \emptyset$.

Theorem 3.65. *A subcategory \mathbf{K} of \mathbf{LSpa} is closed under subspaces (up to isomorphisms) and products if and only if it is a quantitative variety $\mathbf{QAlg}(\emptyset, \hat{E})$.*

*Proof sketch.*⁴³⁵ We already showed the right-to-left direction in Theorem 2.59, noting that by definition, $\mathbf{QAlg}(\emptyset, \hat{E})$ is a category \mathbf{GMet} .

⁴³³ Note that $\widehat{\mathcal{T}}_\emptyset$ is the identity functor on \mathbf{Met} .

⁴³⁴ For an interval $[a, b] \subseteq \mathbb{R}$, the extreme and intermediate value theorems imply that the image of any continuous function $f : [a, b] \rightarrow \mathbb{R}$ is a closed interval. Hence, if f is valued in the rationals, the image of f can only be a single rational number, i.e. f is constant.

⁴³⁵ We give only a sketch because we use some results that would require more background to fully detail the proofs. Nevertheless, we provide all the necessary steps.

For the converse, we first need to know that \mathbf{LSpa} has a factorization system $(\mathcal{E}, \mathcal{M})$ [JMU24, §2] (also called factorization structure in [AHS06, Definition 14.1]). This is shown in [JMU24, Lemma 3.16], where

$$\begin{aligned}\mathcal{E} &= \{e : \mathbf{A} \rightarrow \mathbf{B} \in \mathbf{LSpa} \mid e \text{ is surjective}\} \text{ and} \\ \mathcal{M} &= \{i : \mathbf{A} \rightarrow \mathbf{B} \in \mathbf{LSpa} \mid i \text{ is an isometric embedding}\}.\end{aligned}$$

Then, we use [AHS06, Theorem 16.8] to show that, since \mathbf{K} is closed under subspaces (up to isomorphisms)⁴³⁶ and products, it must be \mathcal{E} -reflective [AHS06, Definition 16.1]. Next, by [AHS06, Theorem 16.14], \mathbf{K} is \mathcal{E} -implicational [AHS06, Definition 16.12].

Unrolling the definition of \mathcal{E} -implicational and translating to our terminology, it means there is a class of abstract quantitative equations \hat{E} such that $\hat{A} \in \mathbf{K}$ if and only if \hat{A} satisfies \hat{E} . In this step, we use the fact that $\widehat{\mathcal{T}}_{\emptyset}$ is the identity functor, it means that

1. any surjective nonexpansive map⁴³⁷ is an abstract quantitative equation, and
2. any nonexpansive assignment is also a homomorphism (for the empty signature), hence satisfaction for abstract quantitative equations coincides with satisfaction of implications as defined in [AHS06, Definition 16.12.(2)].

Finally, by Proposition 3.62, \mathbf{K} is a quantitative variety $\mathbf{QAlg}(\emptyset, \hat{E}^\circ)$. □

3.3 Quantitative Equational Logic

It is now time to introduce quantitative equational logic (QEL), which you can think of as both a generalization and an extension of equational logic. It is a generalization because it is parametrized by a complete lattice L , and when instantiating $L = 1$, we get back equational logic as we will explain in Example 3.70. It is an extension because all the rules of equational logic are valid in QEL when replacing the contexts with discrete spaces as we will explain in Example 3.71. Figure 3.1 displays the inference rules of **quantitative equational logic**. The notion of **derivation** is straightforwardly adapted from Definition 1.53, the crucial difference is that proof trees can now be infinitely branching.⁴³⁸

Given a class of quantitative equations \hat{E} , we denote by $\Omega\mathfrak{Th}'(\hat{E})$ the class of quantitative equations that can be proven from \hat{E} in quantitative equational logic, in other words, $\phi \in \Omega\mathfrak{Th}'(\hat{E})$ if and only if there is a derivation of ϕ in QEL with axioms \hat{E} .

Our goal now is to prove that $\Omega\mathfrak{Th}'(\hat{E}) = \Omega\mathfrak{Th}(\hat{E})$. We say that QEL is sound and complete for (Σ, \hat{E}) -algebras. Less concisely, soundness means that whenever QEL proves an equation ϕ with axioms \hat{E} , ϕ is satisfied by all (Σ, \hat{E}) -algebras, and completeness says that whenever an equation ϕ is satisfied by all (Σ, \hat{E}) -algebras, there is a derivation of ϕ in QEL with axioms \hat{E} .

Just like for equational logic, all the rules in Figure 3.1 are sound for any fixed quantitative algebra meaning that if \hat{A} satisfies the equations on top of a rule, it must satisfy the conclusion of that rule. Let us explain the rules as we prove soundness.

⁴³⁶This is the same as being closed under \mathcal{M} -subobjects.

⁴³⁷i.e. any implication in \mathcal{E} following [AHS06]

⁴³⁸This is necessary due to the rules SUB , SUBQ , and CONT , which can have infinitely many quantitative equations as hypotheses.

$$\begin{array}{c}
 \frac{}{\mathbf{X} \vdash t = t} \text{REFL} \quad \frac{\mathbf{X} \vdash s = t}{\mathbf{X} \vdash t = s} \text{SYMM} \quad \frac{\mathbf{X} \vdash s = t \quad \mathbf{X} \vdash t = u}{\mathbf{X} \vdash s = u} \text{TRANS} \\
 \\
 \frac{\text{op} : n \in \Sigma \quad \forall 1 \leq i \leq n, \mathbf{X} \vdash s_i = t_i}{\mathbf{X} \vdash \text{op}(s_1, \dots, s_n) = \text{op}(t_1, \dots, t_n)} \text{CONG} \\
 \\
 \frac{\sigma : Y \rightarrow \mathcal{T}_\Sigma X \quad \mathbf{Y} \vdash s = t \quad \forall y, y' \in Y, \mathbf{X} \vdash \sigma(y) =_{d_{\mathbf{Y}}(y, y')} \sigma(y')}{\mathbf{X} \vdash \sigma^*(s) = \sigma^*(t)} \text{SUB} \\
 \\
 \frac{}{\mathbf{X} \vdash s =_\top t} \text{TOP} \quad \frac{d_{\mathbf{X}}(x, x') = \varepsilon}{\mathbf{X} \vdash x =_\varepsilon x'} \text{VARS} \quad \frac{\mathbf{X} \vdash s =_\varepsilon t \quad \varepsilon \leq \varepsilon'}{\mathbf{X} \vdash s =_{\varepsilon'} t} \text{MON} \\
 \\
 \frac{\forall i, \mathbf{X} \vdash s =_{\varepsilon_i} t \quad \varepsilon = \inf_i \varepsilon_i}{\mathbf{X} \vdash s =_\varepsilon t} \text{CONT} \quad \frac{\phi \in \hat{E}_{\text{GMet}}}{\phi} \text{GMET} \\
 \\
 \frac{\mathbf{X} \vdash s = t \quad \mathbf{X} \vdash s =_\varepsilon u}{\mathbf{X} \vdash t =_\varepsilon u} \text{COMPL} \quad \frac{\mathbf{X} \vdash s = t \quad \mathbf{X} \vdash u =_\varepsilon s}{\mathbf{X} \vdash u =_\varepsilon t} \text{COMPR} \\
 \\
 \frac{\sigma : Y \rightarrow \mathcal{T}_\Sigma X \quad \mathbf{Y} \vdash s =_\varepsilon t \quad \forall y, y' \in Y, \mathbf{X} \vdash \sigma(y) =_{d_{\mathbf{Y}}(y, y')} \sigma(y')}{\mathbf{X} \vdash \sigma^*(s) =_\varepsilon \sigma^*(t)} \text{SUBQ}
 \end{array}$$

Figure 3.1: Rules of quantitative equational logic over the signature Σ and the complete lattice L , where \mathbf{X} and \mathbf{Y} can be any L -space, s, t, u, s_i and t_i can be any term in $\mathcal{T}_\Sigma X$, and $\varepsilon, \varepsilon'$ and ε_i range over L . As indicated in the premises of the rules CONG, SUB, and SUBQ, they can be instantiated for any n -ary operation symbol and for any function σ respectively.

The first four rules say that equality is an equivalence relation that is preserved by the operations, we showed they were sound in Lemmas 3.28–3.31. More formally, we can define (for any \mathbf{X}) a binary relation \equiv'_ε on Σ -terms⁴³⁹ that contains the pair (s, t) whenever $\mathbf{X} \vdash s = t$ can be proven in QEL (c.f. (3.13)): for any $s, t \in \mathcal{T}_\Sigma X$,

$$s \equiv'_\varepsilon t \iff \mathbf{X} \vdash s = t \in \Omega \mathfrak{I} h'(\hat{E}). \quad (3.45)$$

Then, REFL, SYMM, TRANS, and CONG make \equiv'_ε a congruence relation.

Lemma 3.66. *For any L -space \mathbf{X} , the relation \equiv'_ε is reflexive, symmetric, transitive, and for any $\text{op} : n \in \Sigma$ and $s_1, \dots, s_n, t_1, \dots, t_n \in \mathcal{T}_\Sigma X$,⁴⁴⁰*

$$\forall 1 \leq i \leq n, s_i \equiv'_\varepsilon t_i \implies \text{op}(s_1, \dots, s_n) \equiv'_\varepsilon \text{op}(t_1, \dots, t_n). \quad (3.46)$$

We denote with $\lambda - \int_{\hat{E}}$ the canonical quotient map $\mathcal{T}_\Sigma X \rightarrow \mathcal{T}_\Sigma X / \equiv'_\varepsilon$.

Skipping SUB for now, the TOP rule says that \top is an upper bound for all distances since it is the maximum element of L . We showed it is sound in Lemma 3.34.

The VARS rule is, in a sense, the quantitative version of REFL. It reflects the fact that assignments of variables are nonexpansive with respect to the distance in the context. Indeed, $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$ is nonexpansive precisely when, for all $x, x' \in X$,

$$d_{\mathbf{A}}(\hat{\iota}(x), \hat{\iota}(x')) = d_{\mathbf{A}}(\llbracket x \rrbracket_A^{\hat{\iota}}, \llbracket x' \rrbracket_A^{\hat{\iota}}) \leq d_{\mathbf{X}}(x, x').$$

How is this related to REFL? Letting $t = x \in X$, REFL says that for any assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$, $\hat{\iota}(x) = \hat{\iota}(x)$. This seems trivial, but it hides a deeper fact that the

⁴³⁹ Again, we omit the L -space \mathbf{X} from the notation.

⁴⁴⁰ i.e. \equiv'_ε is a congruence on the Σ -algebra $\mathcal{T}_\Sigma X$ defined in Remark 1.24.

assignment must be deterministic (a functional relation) as it cannot assign two different values to the same input.⁴⁴¹ So just like REFL imposes the constraint of determinism on assignments, VARS imposes nonexpansiveness. We showed VARS is sound in Lemma 3.35.

The rules MON and CONT should remind you of the definition of L-structures (Definition 2.19). Very briefly, they ensure that equipping the set of terms over X with the relations $R_\varepsilon^X \subseteq \mathcal{T}_\Sigma X \times \mathcal{T}_\Sigma X$ defined by

$$s R_\varepsilon^X t \iff \mathbf{X} \vdash s =_\varepsilon t \in \Omega\mathfrak{Th}'(\hat{E}), \quad (3.47)$$

yields an L-structure.⁴⁴² We showed they are sound in Lemmas 3.36 and 3.37. Note that TOP is an instance of CONT with the empty index set (recall that $\top = \inf \emptyset$).

The soundness of GMET is a consequence of (3.4) and the definition of quantitative algebra which requires the underlying space to satisfy all the equations in $\hat{E}_{\mathbf{GMet}}$.

COMPL and COMPR guarantee that the L-structure we just defined factors through the quotient $\mathcal{T}_\Sigma X / \equiv_{\hat{E}}$.⁴⁴³ We showed they are sound in Lemmas 3.32 and 3.33. In the presence of a symmetry axiom, only one of them would be sufficient.

Finally, we get to the substitutions SUB and SUBQ, they are the same except for replacing $=$ with $=_\varepsilon$. Recall that the substitution rule in equational logic is

$$\frac{\sigma : Y \rightarrow \mathcal{T}_\Sigma X \quad Y \vdash s = t}{\mathbf{X} \vdash \sigma^*(s) = \sigma^*(t)},$$

which morally means that variables in the context Y are universally quantified. In SUB and SUBQ, there is an additional condition on σ which arises because the variables in Y are *not* universally quantified: an assignment $Y \rightarrow A$ is considered in the definition of satisfaction only if it is nonexpansive from \mathbf{Y} to \mathbf{A} .⁴⁴⁴

We proved SUB and SUBQ are sound in Lemma 3.42, and we can compare with the proof of soundness of SUB in equational logic (Lemma 1.45) to find the same key argument: the interpretation of $\sigma^*(t)$ under some assignment \hat{t} is equal to the interpretation of t under the assignment \hat{t}_σ sending y to the interpretation of $\sigma(y)$ under \hat{t} . Since satisfaction for quantitative algebras only deals with nonexpansive assignments, we needed to check that \hat{t}_σ is nonexpansive whenever \hat{t} is, and this was true thanks to the conditions on σ . Let us give an illustrative example of why the extra conditions are necessary.

Example 3.67. We work over $L = [0, 1]$, $\mathbf{GMet} = \mathbf{Met}$, $\Sigma = \emptyset$, and $\hat{E} = \emptyset$. Let $\mathbf{Y} = \{y_0, y_1\}$ with $d_Y(y_0, y_1) = d_Y(y_1, y_0) = \frac{1}{2}$ and $\mathbf{X} = \{x_0, x_1\}$ with $d_X(x_0, x_1) = d_X(x_1, x_0) = 1$.⁴⁴⁵ We consider the algebra $\hat{\mathbf{A}}$ whose underlying space is $\mathbf{A} = \mathbf{X}$ (since Σ is empty that is the only data required to define an algebra). It satisfies the equation $\mathbf{Y} \vdash y_0 = y_1$ because any nonexpansive assignment of \mathbf{Y} into \mathbf{A} must identify y_0 and y_1 (there are no distinct points with distance less than $\frac{1}{2}$).

Take the substitution $\sigma : Y \rightarrow \mathcal{T}_\Sigma X$ defined by $y_0 \mapsto x_0$ and $y_1 \mapsto x_1$, we can check $\hat{\mathbf{A}}$ does not satisfy $\mathbf{X} \vdash \sigma^*(y_0) = \sigma^*(y_1)$.⁴⁴⁶ This means that σ cannot satisfy the extra conditions in SUB. Indeed, $\hat{\mathbf{A}}$ does not satisfy $\mathbf{X} \vdash \sigma(y_0) =_{\frac{1}{2}} \sigma(y_1)$ (take the assignment id_X again).

⁴⁴¹ A similar thing happens for CONG which says that the interpretations of operation are deterministic (both in equational logic and QEL). In [MPP16], the logic has a rule NEXP which morally says that the interpretations of operations are nonexpansive too, i.e. NEXP is to CONG what VARS is to REFL. We said more on our choice to omit NEXP in §0.3.

⁴⁴² Monotonicity and continuity hold by MON and CONT respectively. This is where the rules' names come from. These rules were given several names in the literature, like MAX instead of MON, and ARCH instead of CONT.

⁴⁴³ i.e. the following relation is well-defined:

$$\llbracket s \rrbracket_{\hat{E}} R_\varepsilon^X \llbracket t \rrbracket_{\hat{E}} \iff \mathbf{X} \vdash s =_\varepsilon t \in \Omega\mathfrak{Th}'(\hat{E}), \quad (3.48)$$

⁴⁴⁴ Put differently, the variables are universally quantified subject to certain constraints on their distances relative to the context \mathbf{Y} .

⁴⁴⁵ We can see both \mathbf{Y} and \mathbf{X} as subspace of $[0, 1]$ with the Euclidean metric, where e.g. y_0 is embedded as 0 and y_1 as $\frac{1}{2}$, and x_0 is embedded as 0 and x_1 as 1.

⁴⁴⁶ That equation is $\mathbf{X} \vdash x_0 = x_1$ and with the assignment $\text{id}_X : \mathbf{X} \rightarrow \mathbf{X} = \mathbf{A}$, we have

$$\llbracket x_0 \rrbracket_{\mathbf{A}}^{\text{id}_X} = x_0 \neq x_1 = \llbracket x_1 \rrbracket_{\mathbf{A}}^{\text{id}_X}.$$

Remark 3.68. The substitution rule in the original paper [MPP16, (Subst) in Definition 2.1] is

$$\frac{\{s_i =_{\varepsilon_i} t_i\} \vdash s =_{\varepsilon} t}{\{\sigma^*(s_i) =_{\varepsilon_i} \sigma^*(t_i)\} \vdash \sigma^*(s) =_{\varepsilon} \sigma^*(t)}.$$

This cannot easily be translated into our framework because it has to work with quantitative inferences that are not basic (Remark 3.11). Indeed, even if the top inference is basic (i.e. each s_i and t_i are variables), the bottom one will not be basic when σ sends these variables to complex terms. In this sense, we can say that our quantitative equational logic is closed under basic quantitative inferences,⁴⁴⁷ while theirs is not.

This is an advantage of our presentation with respect to its comparison with equational logic. Indeed, non-basic quantitative inferences are a better analog for implications in implicative logic [Wec92, §3.3, Definition 1]. For example, you can model cancellative monoids, with something like $x + y =_0 x + z \vdash y =_0 z$, and they are a canonical example of structures not captured by universal algebra.

By proving each rule is sound, we have shown that QEL is sound.

Theorem 3.69 (Soundness). *If $\phi \in \Omega\mathfrak{Th}'(\hat{E})$, then $\phi \in \Omega\mathfrak{Th}(\hat{E})$.*

Let us explain how to recover equational logic from quantitative equational logic in two different ways.

Example 3.70 (Recovering equational logic I). In Example 2.20, we saw that $\mathbf{1Spa}$ is the category **Set**. Here we show that QEL over the complete lattice $\mathbf{1}$ with $\hat{E}_{\mathbf{GMet}} = \emptyset$ is the same thing as equational logic. First, what is a quantitative equation ϕ over $\mathbf{1}$? Since the context is a $\mathbf{1}$ -space, it is just a set,⁴⁴⁸ and furthermore, since $\mathbf{1}$ contains a single element (which we call \top here, but it is equal to \perp) ϕ is either

$$X \vdash s = t \quad \text{or} \quad X \vdash s =_{\top} t.$$

Now, the second equation always belongs to $\Omega\mathfrak{Th}'(\hat{E})$ for any \hat{E} by **TOP**. Therefore, the rules whose conclusions have an equation with a quantity (all but the first five) can be replaced by **TOP**. The remaining rules are exactly those of equational logic except the substitution rule which has some additional constraints. The latter require proving only equations with quantities which we can always do with **TOP**.

Thus, we can infer that for any \hat{E} , the equations without quantities in $\Omega\mathfrak{Th}'(\hat{E})$ are exactly the equations in $\mathfrak{Th}'(E)$, where E contains the quantitative equations without quantities of \hat{E} seen as equations.⁴⁴⁹

If we had naively generalized the original logic of [MPP16] by replacing $[0, \infty]$ with an arbitrary complete lattice, this instantiation to $\mathbf{1}$ would not have been equivalent to equational logic. Indeed, as we explained in §0.3, the judgments of [MPP16], called quantitative inferences, are more general than quantitative equations, and they can express properties which cannot be expressed with equations.⁴⁵⁰

Example 3.71 (Recovering equational logic II). There is a less trivial way to see that equational reasoning faithfully embeds into quantitative equational reasoning.

⁴⁴⁷ Recall that basic quantitative inferences correspond to quantitative equations.

⁴⁴⁸ In other words, X and \mathbf{X} are the same thing.

⁴⁴⁹ i.e. $E = \{X \vdash s = t \mid \mathbf{X} \vdash s = t \in \hat{E}\}$

⁴⁵⁰ The standard example of left-cancellability of a binary operation would be expressed with the quantitative inference

$$x \cdot y = x \cdot z \vdash y = z,$$

but it cannot be expressed with equations. Quantitative inferences are better quantitative versions of the implications in [Wec92, §3.3, Definition 1].

We are back to the general case of L being an arbitrary complete lattice and \hat{E}_{GMet} being possibly non-empty. Let E be a class of classical equations, and let \hat{E} contain every equation in E seen as a quantitative equation with its context being the discrete space, i.e.

$$\hat{E} = \{X_{\top} \vdash s = t \mid X \vdash s = t \in E\}. \quad (3.49)$$

Claim. If $X \vdash s = t \in \mathfrak{Th}'(E)$, then $X_{\top} \vdash s = t \in \Omega\mathfrak{Th}'(\hat{E})$.⁴⁵¹

Proof 1. You can show by induction that a derivation of $X \vdash s = t$ in equational logic with axioms E can be transformed into a derivation of $X_{\top} \vdash s = t$ in QEL with axioms \hat{E} . The base cases are handled by the definition of \hat{E} and the rule **REFL** in QEL instantiated with the discrete spaces which perfectly emulates the rule **REFL** in equational logic.

For the inductive step, the rules **SYMM**, **TRANS**, and **CONG** in equational logic all have perfect counterparts in QEL. The substitution rule needs a bit more work. If the last rule in the derivation in equational logic is

$$\frac{\sigma : Y \rightarrow \mathcal{T}_{\Sigma} X \quad Y \vdash s = t}{X_{\top} \sigma^*(s) = \sigma^*(t)} \text{SUB},$$

then by induction hypothesis, there is a derivation of $Y_{\top} \vdash s = t$ in QEL. We obtain the following derivation noting that for all $y, y' \in Y$, $d_{\top}(y, y') = \top$.

$$\frac{\sigma : Y \rightarrow \mathcal{T}_{\Sigma} X \quad \frac{\text{I.H.}}{Y_{\top} \vdash s = t} \quad \frac{\forall y, y' \in Y, X_{\top} \vdash \sigma(y) =_{d_{\top}(y, y')} \sigma(y')}{\text{TOP}}}{X_{\top} \vdash \sigma^*(s) = \sigma^*(t)} \text{SUB}$$

□

Proof 2. The proof above reasoning on derivations is useful to get familiar with QEL, but there is a faster *semantic* proof that relies on completeness. By soundness and completeness,⁴⁵² it is enough to prove that if $X \vdash s = t \in \mathfrak{Th}(E)$, then $X_{\top} \vdash s = t \in \Omega\mathfrak{Th}(\hat{E})$. This follows from the equivalence (3.15) (which was easy to prove):

$$\hat{\mathbb{A}} \models \hat{E} \stackrel{(3.15)}{\iff} \mathbb{A} \models E \stackrel{(1.21)}{\iff} \mathbb{A} \models X \vdash s = t \stackrel{(3.15)}{\iff} \hat{\mathbb{A}} \models X_{\top} \vdash s = t. \quad \square$$

This second proof also points to a stronger version of the claim that we state as a lemma for future use.

Lemma 3.72. *Let E be a class of classical equations and \hat{E} be defined as in (3.49). If $X \vdash s = t \in \mathfrak{Th}'(E)$, then, for any L -space \mathbf{X} with carrier X , $X_{\top} \vdash s = t \in \Omega\mathfrak{Th}'(\hat{E})$.⁴⁵³*

Let us get back to our goal of showing QEL is complete. We follow the proof sketch of completeness for equational logic.⁴⁵⁴ We define a quantitative algebra exactly like $\hat{\mathbb{T}}\mathbf{X}$ but using the equality relation and L -relation induced by $\Omega\mathfrak{Th}'(\hat{E})$ instead of $\Omega\mathfrak{Th}(\hat{E})$, and then we show it satisfies \hat{E} which, by construction, will imply $\Omega\mathfrak{Th}(\hat{E}) \subseteq \Omega\mathfrak{Th}'(\hat{E})$.

⁴⁵¹ Depending on the equations inside \hat{E}_{GMet} , it is possible that $\Omega\mathfrak{Th}'(\hat{E})$ contains more equations without quantities than $\mathfrak{Th}'(E)$. Nevertheless, we show that everything you can prove in equational logic can also be proven in QEL.

⁴⁵² Of both equational logic (Theorems 1.55 and 1.60) and QEL (Theorems 3.69 and 3.76).

⁴⁵³ Follow the second proof above but instead of the second use of (3.15), use Lemma 3.40. (This requires assuming $\Omega\mathfrak{Th}(\hat{E}) = \Omega\mathfrak{Th}'(\hat{E})$ which we prove soon.)

⁴⁵⁴ Our proof of completeness for the logic in [MSV22] seems more complex (in my opinion), but it morally follows the same sketch. It is obfuscated however by the fact that [MSV22] did not deal with contexts, instead we were using what we now call syntactic sugar to describe quantitative equations.

Definition 3.73 (Quantitative term algebra, syntactically). The *new* quantitative term algebra for (Σ, \hat{E}) on \mathbf{X} is the quantitative Σ -algebra whose underlying space is $\mathcal{T}_\Sigma X / \equiv'_{\hat{E}}$ equipped with the L-relation corresponding to the L-structure defined in (3.48),⁴⁵⁵ and whose interpretation of $\text{op} : n \in \Sigma$ is defined by⁴⁵⁶

$$\llbracket \text{op} \rrbracket_{\hat{\mathbb{T}}' \mathbf{X}} (\lambda t_1 \int_{\hat{E}}, \dots, \lambda t_n \int_{\hat{E}}) = \lambda \text{op}(t_1, \dots, t_n) \int_{\hat{E}}. \quad (3.51)$$

We denote this algebra by $\hat{\mathbb{T}}'_{\Sigma, \hat{E}} \mathbf{X}$ or simply $\hat{\mathbb{T}}' \mathbf{X}$.

We will prove this alternative definition of the term algebra coincides with $\hat{\mathbb{T}} \mathbf{X}$. First, we have to show that $\hat{\mathbb{T}}' \mathbf{X}$ belongs to $\mathbf{QAlg}(\Sigma, \hat{E})$ like we did for $\hat{\mathbb{T}} \mathbf{X}$ in Proposition 3.56, and we state a technical lemma before that.

Lemma 3.74. *Let $\iota : Y \rightarrow \mathcal{T}_\Sigma X / \equiv'_{\hat{E}}$ be any assignment. For any function $\sigma : Y \rightarrow \mathcal{T}_\Sigma X$ satisfying $\lambda \sigma(y) \int_{\hat{E}} = \iota(y)$ for all $y \in Y$, we have $\llbracket - \rrbracket_{\hat{\mathbb{T}}' \mathbf{X}}^{\iota} = \lambda \sigma^*(-) \int_{\hat{E}}$.⁴⁵⁷*

Proposition 3.75. *For any space \mathbf{X} , $\hat{\mathbb{T}}' \mathbf{X}$ satisfies all the equations in \hat{E} .*

Proof. Let $\mathbf{Y} \vdash s = t$ (resp. $\mathbf{Y} \vdash s =_\varepsilon t$) belong to \hat{E} and $\hat{\iota} : \mathbf{Y} \rightarrow (\mathcal{T}_\Sigma X / \equiv'_{\hat{E}}, d'_{\hat{E}})$ be a nonexpansive assignment. By the axiom of choice,⁴⁵⁸ there is a function $\sigma : Y \rightarrow \mathcal{T}_\Sigma X$ satisfying $\lambda \sigma(y) \int_{\hat{E}} = \hat{\iota}(y)$ for all $y \in Y$. Thanks to Lemma 3.74, it is enough to show $\lambda \sigma^*(s) \int_{\hat{E}} = \lambda \sigma^*(t) \int_{\hat{E}}$ (resp. $d'_{\hat{E}}(\lambda \sigma^*(s) \int_{\hat{E}}, \lambda \sigma^*(t) \int_{\hat{E}}) \leq \varepsilon$).⁴⁵⁹

Equivalently, by definition of $\lambda - \int_{\hat{E}}$ and $\Omega \mathfrak{H}'(\hat{E})$, we can just exhibit a derivation of $\mathbf{X} \vdash \sigma^*(s) = \sigma^*(t)$ (resp. $\mathbf{X} \vdash \sigma^*(s) =_\varepsilon \sigma^*(t)$) in QEL with axioms \hat{E} . That equation can be proven with the SUB (resp. SUBQ) rule instantiated with $\sigma : Y \rightarrow \mathcal{T}_\Sigma X$ and the equation $\mathbf{Y} \vdash s = t$ (resp. $\mathbf{Y} \vdash s =_\varepsilon t$) which is an axiom, but we need derivations showing σ satisfies the side conditions of the substitution rules. This follows from nonexpansiveness of $\hat{\iota}$ because for any $y, y' \in Y$, we know that

$$d_{\hat{E}}(\lambda \sigma(y) \int_{\hat{E}}, \lambda \sigma(y') \int_{\hat{E}}) = d_{\hat{E}}(\hat{\iota}(y), \hat{\iota}(y')) \leq d_{\mathbf{Y}}(y, y'),$$

which means by (3.50) that $\mathbf{X} \vdash \sigma(y) =_{d_{\mathbf{Y}}(y, y')} \sigma(y')$ belongs to $\Omega \mathfrak{H}'(\hat{E})$. \square

Completeness of quantitative equational logic readily follows.

Theorem 3.76 (Completeness). *If $\phi \in \Omega \mathfrak{H}(\hat{E})$, then $\phi \in \Omega \mathfrak{H}'(\hat{E})$.*

Proof. Let $\phi \in \Omega \mathfrak{H}(\hat{E})$ and \mathbf{X} be its context. By Proposition 3.75 and definition of $\Omega \mathfrak{H}(\hat{E})$, we know that $\hat{\mathbb{T}}' \mathbf{X} \models \phi$. In particular, $\hat{\mathbb{T}}' \mathbf{X}$ satisfies ϕ under the assignment

$$\hat{\iota} = \mathbf{X} \xrightarrow{\eta_{\hat{\mathbf{X}}}^{\Sigma}} \mathcal{T}_\Sigma X \xrightarrow{\lambda - \int_{\hat{E}}} \mathcal{T}_\Sigma X / \equiv'_{\hat{E}},$$

which is nonexpansive by VARS.⁴⁶⁰

Moreover with $\sigma = \eta_{\hat{\mathbf{X}}}^{\Sigma}$, we can show σ satisfies the hypothesis of Lemma 3.74 and $\sigma^* = \text{id}_{\mathcal{T}_\Sigma X}$,⁴⁶¹ thus we conclude

- if $\phi = \mathbf{X} \vdash s = t : \lambda s \int_{\hat{E}} = \llbracket s \rrbracket_{\hat{\mathbb{T}}' \mathbf{X}}^{\hat{\iota}} = \llbracket t \rrbracket_{\hat{\mathbb{T}}' \mathbf{X}}^{\hat{\iota}} = \lambda t \int_{\hat{E}}$, and
- if $\phi = \mathbf{X} \vdash s =_\varepsilon t : d'_{\hat{E}}(\lambda s \int_{\hat{E}}, \lambda t \int_{\hat{E}}) = d'_{\hat{E}}(\llbracket s \rrbracket_{\hat{\mathbb{T}}' \mathbf{X}}^{\hat{\iota}}, \llbracket t \rrbracket_{\hat{\mathbb{T}}' \mathbf{X}}^{\hat{\iota}}) \leq \varepsilon$.

⁴⁵⁵ Explicitly, it is the L-relation $d'_{\hat{E}}$ that satisfies

$$d'_{\hat{E}}(\lambda s \int_{\hat{E}}, \lambda t \int_{\hat{E}}) \leq \varepsilon \iff \mathbf{X} \vdash s =_\varepsilon t \in \Omega \mathfrak{H}'(\hat{E}). \quad (3.50)$$

⁴⁵⁶ This is well-defined (i.e. invariant under change of representative) by (3.46).

⁴⁵⁷ The proof goes exactly as in the classical case (Lemma 1.58). We do not even need to ask ι to be nonexpansive, but we will use the result with a nonexpansive assignment.

⁴⁵⁸ Choice implies the quotient map $\lambda - \int_{\hat{E}}$ has a right inverse $r : \mathcal{T}_\Sigma X / \equiv'_{\hat{E}} \rightarrow \mathcal{T}_\Sigma X$, and we set $\sigma = r \circ \hat{\iota}$.

⁴⁵⁹ By Lemma 3.74, it implies

$$\llbracket s \rrbracket_{\hat{\mathbb{T}}' \mathbf{X}}^{\hat{\iota}} = \lambda \sigma^*(s) \int_{\hat{E}} = \lambda \sigma^*(t) \int_{\hat{E}} = \llbracket t \rrbracket_{\hat{\mathbb{T}}' \mathbf{X}}^{\hat{\iota}}$$

$$\text{resp. } d'_{\hat{E}}(\llbracket s \rrbracket_{\hat{\mathbb{T}}' \mathbf{X}}^{\hat{\iota}}, \llbracket t \rrbracket_{\hat{\mathbb{T}}' \mathbf{X}}^{\hat{\iota}}) = d'_{\hat{E}}(\lambda \sigma^*(s) \int_{\hat{E}}, \lambda \sigma^*(t) \int_{\hat{E}}) \leq \varepsilon$$

and since $\hat{\iota}$ was arbitrary, we conclude that $\hat{\mathbb{T}}' \mathbf{X}$ satisfies $\mathbf{Y} \vdash s = t$ (resp. $\mathbf{Y} \vdash s =_\varepsilon t$).

⁴⁶⁰ Explicitly, VARS means $\mathbf{X} \vdash x =_{d_{\mathbf{X}}(x, x')} x'$ belongs to $\Omega \mathfrak{H}'(\hat{E})$, hence, (3.50) implies

$$d'_{\hat{E}}(\lambda x \int_{\hat{E}}, \lambda x' \int_{\hat{E}}) \leq d_{\mathbf{X}}(x, x').$$

⁴⁶¹ We defined $\hat{\iota}$ precisely to have $\lambda \eta_{\hat{\mathbf{X}}}^{\Sigma}(x) \int_{\hat{E}} = \hat{\iota}(x)$. To show $\sigma^* = \eta_{\hat{\mathbf{X}}}^{\Sigma}$ is the identity, use (1.38) and the fact that $\mu^{\Sigma} \cdot \eta^{\Sigma} \mathcal{T}_\Sigma = \mathbb{1}_{\mathcal{T}_\Sigma}$ (it holds by definition (1.7)).

By definition of $\equiv_{\hat{E}}$ (3.45) and $d'_{\hat{E}}$ (3.50), this implies $\mathbf{X} \vdash s = t$ (resp. $\mathbf{X} \vdash s =_{\varepsilon} t$) belongs to $\Omega\mathfrak{Th}'(\hat{E})$. \square

Note that because $\widehat{\mathbb{T}}\mathbf{X}$ and $\widehat{\mathbb{T}}'\mathbf{X}$ were defined in the same way in terms of $\Omega\mathfrak{Th}(\hat{E})$ and $\Omega\mathfrak{Th}'(\hat{E})$ respectively, and since we have proven the latter to be equal, we obtain that $\widehat{\mathbb{T}}\mathbf{X}$ and $\widehat{\mathbb{T}}'\mathbf{X}$ are the same quantitative algebra. In the sequel, we will work with $\widehat{\mathbb{T}}\mathbf{X}$ mostly, but we may use the facts that $s \equiv_{\hat{E}} t$ (resp. $d'_{\hat{E}}(s, t) \leq \varepsilon$) if and only if there is a derivation of $\mathbf{X} \vdash s = t$ (resp. $\mathbf{X} \vdash s =_{\varepsilon} t$) in QEL.⁴⁶²

Remark 3.77. Mirroring Remark 1.61, we would like to say that the axiom of choice was not necessary in the proofs above. Unfortunately, this situation is more delicate, and I do not know for sure that we can avoid using choice (although I expect we can).

At first, you might think that since terms are still finite, we can still restrict the context to the free variables which is finite. Unfortunately, even if $x \in \text{FV}\{s, t\}$ and $y \notin \text{FV}\{s, t\}$, it is possible that the distance between x and y in the context is necessary to state the right property. Here is an example that we carry with $\mathbf{GMet} = [0, 1]\mathbf{Spa}$, $\Sigma = \emptyset$, and \hat{E} defining discrete metrics:⁴⁶³

$$\hat{E} = \{x =_{\varepsilon} y \vdash x = y \mid 1 \neq \varepsilon \in \mathbb{L}\} \cup \{x = y \vdash x =_0 y\}.$$

Let $\mathbf{X} = \{x, z\}$ and $\mathbf{Y} = \{x, y, z\}$ with the following distances (\mathbf{X} is a subspace of \mathbf{Y}):

$$\begin{array}{c} \overset{0}{\curvearrowright} \\ x \end{array} \xrightarrow{\frac{1}{2}} \begin{array}{c} \overset{0}{\curvearrowright} \\ y \end{array} \xrightarrow{\frac{1}{2}} \begin{array}{c} \overset{0}{\curvearrowright} \\ z \end{array}$$

The equation $\mathbf{Y} \vdash x = z$ belongs to $\Omega\mathfrak{Th}(\hat{E})$. Indeed, if $\mathbf{A} \models \hat{E}$, then $d_{\mathbf{A}}(a, b) \leq \frac{1}{2}$ implies $a = b$, so any nonexpansive assignment $\hat{\iota} : \mathbf{Y} \rightarrow \mathbf{A}$ must identify x and y , and y and z , hence $\hat{\iota}(x) = \hat{\iota}(z)$. However, the equation $\mathbf{X} \vdash x = z$ is not in $\Omega\mathfrak{Th}(\hat{E})$ because you can have $d_{\mathbf{A}}(\hat{\iota}(x), \hat{\iota}(z)) \leq 1$ without $\hat{\iota}(x) = \hat{\iota}(z)$.

This shows that some variables in the context which are not used in the terms of the equation (in this instance y) might still be important. One may still wonder whether it is possible to restrict the contexts to be finite or countable.⁴⁶⁴ I do not know if that is true, but I expect that countable contexts are enough and that finite contexts are not.

In summary, while there can be an analog to the derivable ADD rule in equational logic,⁴⁶⁵ the obvious counterpart to the DEL rule is not even sound. Recalling that ADD and DEL were both derivable by using SUB in equational logic, this also explains the need for additional premises in the SUB and SUBQ rules of quantitative equational logic (c.f. Example 3.67).

Let us highlight one last feature of quantitative equational logic: the rule GMET defining what kind of generalized metric spaces are considered is independent of all the other rules.⁴⁶⁶ As a consequence, and we give more details in [MSV23, §8], you can choose to work over \mathbf{LSpa} all the time and add the equations in $\hat{E}_{\mathbf{GMet}}$ as axioms in \hat{E} anytime you wish to restrict to algebras whose carriers are generalized metric spaces. Written a bit ambiguously,⁴⁶⁷

⁴⁶² i.e. when proving that an equation holds in some theory $\Omega\mathfrak{Th}(\hat{E})$, we can either use the rules of QEL or the several lemmas from §3.2 which are the semantic counterparts to the inference rules.

⁴⁶³ When $d_{\mathbf{A}}(a, b)$ is not 1, it must be that $a = b$ by the first set of equations, by the second set, it must be that $d_{\mathbf{A}}(a, b) = 0$. Under such constraints \mathbf{A} must be the discrete metric on A that we described in Example 3.59, so $\mathbf{QAlg}(\emptyset, \hat{E})$ is the category of discrete metrics.

⁴⁶⁴ i.e. for any equation ϕ , is there a set of equations \hat{E}_{ϕ} with finite (or countable) contexts such that

$$\hat{\mathbf{A}} \models \phi \iff \hat{\mathbf{A}} \models \hat{E}_{\phi}.$$

⁴⁶⁵ When adding a variable y to a context \mathbf{X} , you put y at distance \top from all other variables.

⁴⁶⁶ Although it was less explicit because only \mathbf{Met} was considered, this was already a feature of the logic in [MPP16].

⁴⁶⁷ What we really mean is that on the left, \mathbf{QAlg} and $\Omega\mathfrak{Th}$ are the operators we described with the parameter \mathbf{GMet} built in, and on the right, they are the same operators instantiated with \mathbf{LSpa} instead.

$$\mathbf{QAlg}(\Sigma, \hat{E}) = \mathbf{QAlg}(\Sigma, \hat{E} \cup \hat{E}_{\mathbf{GMet}}) \quad \text{and} \quad \Omega\mathfrak{Th}(\hat{E}) = \Omega\mathfrak{Th}(\hat{E} \cup \hat{E}_{\mathbf{GMet}}). \quad (3.52)$$

3.4 Quantitative Algebraic Presentations

In order to obtain a more categorical understanding of quantitative algebras, a first step is to show that the functor $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} : \mathbf{GMet} \rightarrow \mathbf{GMet}$ we constructed is a monad.

Proposition 3.78. *The functor $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} : \mathbf{GMet} \rightarrow \mathbf{GMet}$ defines a monad on \mathbf{GMet} with unit $\widehat{\eta}^{\Sigma, \hat{E}}$ and multiplication $\widehat{\mu}^{\Sigma, \hat{E}}$. We call it the **term monad** for (Σ, \hat{E}) .*

Proof. A first proof uses a standard result of category theory. Since we showed that $\widehat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{A}$ is the free (Σ, \hat{E}) -algebra on \mathbf{A} for every space \mathbf{A} (Theorem 3.57), we obtain a monad sending \mathbf{A} to the underlying space of $\widehat{\mathbb{T}}_{\Sigma, \hat{E}} \mathbf{A}$, i.e. $\widehat{\mathcal{T}}_{\Sigma, \hat{E}} \mathbf{A}$.⁴⁶⁸

One could also follow the proof we gave for \mathbf{Set} and explicitly show that $\widehat{\eta}^{\Sigma, \hat{E}}$ and $\widehat{\mu}^{\Sigma, \hat{E}}$ obey the laws for the unit and multiplication (most of the work having been done earlier in this chapter). \square

What is arguably more important is that quantitative (Σ, \hat{E}) -algebras on a space \mathbf{A} correspond to $\widehat{\mathcal{T}}_{\Sigma, \hat{E}}$ -algebras on \mathbf{A} .⁴⁶⁹ We construct an isomorphism between $\mathbf{QAlg}(\Sigma, \hat{E})$ and $\mathbf{EM}(\widehat{\mathcal{T}}_{\Sigma, \hat{E}})$ using the isomorphism $P : \mathbf{Alg}(\Sigma) \cong \mathbf{EM}(\mathcal{T}_{\Sigma}) : P^{-1}$ that we defined in Proposition 1.70,⁴⁷⁰ the forgetful functor $U : \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{Alg}(\Sigma)$ that sends \hat{A} to the underlying algebra A , and the functor $\mathbf{EM}(\widehat{\mathcal{T}}_{\Sigma, \hat{E}}) \rightarrow \mathbf{EM}(\mathcal{T}_{\Sigma})$ we define below.

Lemma 3.79. *For any $\widehat{\mathcal{T}}_{\Sigma, \hat{E}}$ -algebra (A, α) , the map $U\alpha \circ [-]_{\hat{E}} : \mathcal{T}_{\Sigma} A \rightarrow A$ is a \mathcal{T}_{Σ} -algebra. Furthermore, this defines a functor $U^{[-]_{\hat{E}}} : \mathbf{EM}(\widehat{\mathcal{T}}_{\Sigma, \hat{E}}) \rightarrow \mathbf{EM}(\mathcal{T}_{\Sigma})$.*

Proof. Apply Proposition 1.82 after checking that $(U, [-]_{\hat{E}})$ is a lax monad morphism from \mathcal{T}_{Σ} to $\widehat{\mathcal{T}}_{\Sigma, \hat{E}}$.⁴⁷¹ \square

Theorem 3.80. *There is an isomorphism $\mathbf{QAlg}(\Sigma, \hat{E}) \cong \mathbf{EM}(\widehat{\mathcal{T}}_{\Sigma, \hat{E}})$.*

Proof. In the diagram below, we already have the functors drawn with solid arrows, and we want to construct \widehat{P} and \widehat{P}^{-1} drawn with dashed arrows before proving they are inverses to each other.

$$\begin{array}{ccc} \mathbf{QAlg}(\Sigma, \hat{E}) & \xleftarrow[\widehat{P}^{-1}]{\widehat{P}} & \mathbf{EM}(\widehat{\mathcal{T}}_{\Sigma, \hat{E}}) \\ U \downarrow & & \downarrow U^{[-]_{\hat{E}}} \\ \mathbf{Alg}(\Sigma) & \xleftarrow[p^{-1}]{P} & \mathbf{EM}(\mathcal{T}_{\Sigma}) \end{array} \qquad \begin{array}{ccc} \mathbf{QAlg}(\Sigma, \hat{E}) & \xleftarrow[\widehat{P}^{-1}]{\widehat{P}} & \mathbf{EM}(\widehat{\mathcal{T}}_{\Sigma, \hat{E}}) \\ & \searrow U & \swarrow U^{\widehat{\mathcal{T}}_{\Sigma, \hat{E}}} \\ & \mathbf{GMet} & \end{array}$$

A (meaningful) sidequest for us is to make the diagrams above commute, namely, the underlying \mathcal{T}_{Σ} -algebra of $\widehat{P}\hat{A}$ should be PA and the underlying space of $\widehat{P}\hat{A}$ should be the underlying space of \hat{A} , and similarly for \widehat{P}^{-1} . It turns out this completely determines our functors, up to some quick checks. We will move between spaces and their underlying sets without indicating it by $U : \mathbf{GMet} \rightarrow \mathbf{Set}$.

⁴⁶⁸ The unit is automatically $\widehat{\eta}^{\Sigma, \hat{E}}$, but some computations are needed to show the multiplication is $\widehat{\mu}^{\Sigma, \hat{E}}$.

⁴⁶⁹ i.e. $U : \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{GMet}$ is monadic.

⁴⁷⁰ Take the statement of Proposition 1.70 with $E = \emptyset$.

⁴⁷¹ The appropriate diagrams (1.61) and (1.62) commute by (3.34) and a combination of (3.21) and (3.22).

Given $\hat{\mathbf{A}} \in \mathbf{QAlg}(\Sigma, \hat{E})$, we look at the underlying Σ -algebra \mathbf{A} , apply P to it to get $\alpha_{\mathbf{A}} : \mathcal{T}_{\Sigma}A \rightarrow \mathbf{A}$ which sends a term t to its interpretation $\llbracket t \rrbracket_{\mathbf{A}}$, and we need to check that it factors through $[-]_{\hat{E}}$ and a nonexpansive map $\hat{\alpha}_{\hat{\mathbf{A}}}$ as in (3.53).

First, $\alpha_{\mathbf{A}}$ is well-defined on terms modulo \hat{E} because if $s \equiv_{\hat{E}} t$, then $\hat{\mathbf{A}}$ satisfies $\mathbf{A} \vdash s = t \in \Omega\mathfrak{Th}(\hat{E})$, and this in turn means (taking the assignment $\text{id}_{\mathbf{A}} : \mathbf{A} \rightarrow \mathbf{A}$):

$$\alpha_{\mathbf{A}}(s) = \llbracket s \rrbracket_{\mathbf{A}} = \llbracket s \rrbracket_{\mathbf{A}}^{\text{id}_{\mathbf{A}}} = \llbracket t \rrbracket_{\mathbf{A}}^{\text{id}_{\mathbf{A}}} = \llbracket t \rrbracket_{\mathbf{A}} = \alpha_{\mathbf{A}}(t).$$

Next, the factor we obtain $\hat{\alpha}_{\hat{\mathbf{A}}} : \mathcal{T}_{\Sigma}A / \equiv_{\hat{E}} \rightarrow \mathbf{A}$ is nonexpansive from $\hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{A}$ to \mathbf{A} . Indeed, if $d_{\hat{E}}(\llbracket s \rrbracket_{\hat{E}}, \llbracket t \rrbracket_{\hat{E}}) \leq \varepsilon$, then $\hat{\mathbf{A}}$ satisfies $\mathbf{A} \vdash s =_{\varepsilon} t \in \Omega\mathfrak{Th}(\hat{E})$, and this means:

$$d_{\mathbf{A}}(\hat{\alpha}_{\hat{\mathbf{A}}}[\llbracket s \rrbracket_{\hat{E}}], \hat{\alpha}_{\hat{\mathbf{A}}}[\llbracket t \rrbracket_{\hat{E}}]) = d_{\mathbf{A}}(\alpha_{\mathbf{A}}(s), \alpha_{\mathbf{A}}(t)) = d_{\mathbf{A}}(\llbracket s \rrbracket_{\mathbf{A}}, \llbracket t \rrbracket_{\mathbf{A}}) = d_{\mathbf{A}}(\llbracket s \rrbracket_{\mathbf{A}}^{\text{id}_{\mathbf{A}}}, \llbracket t \rrbracket_{\mathbf{A}}^{\text{id}_{\mathbf{A}}}) \leq \varepsilon.$$

Finally, if $h : \hat{\mathbf{A}} \rightarrow \hat{\mathbf{B}}$ is a homomorphism, then by definition it is nonexpansive $\mathbf{A} \rightarrow \mathbf{B}$ and it commutes with $[-]_{\mathbf{A}}$ and $[-]_{\mathbf{B}}$. The latter means it commutes with $\alpha_{\mathbf{A}}$ and $\alpha_{\mathbf{B}}$, which in turn means it commutes with $\hat{\alpha}_{\hat{\mathbf{A}}}$ and $\hat{\alpha}_{\hat{\mathbf{B}}}$ because $[-]_{\hat{E}}$ is epic (see (3.54)). We obtain our functor $\hat{P} : \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{EM}(\hat{\mathcal{T}}_{\Sigma, \hat{E}})$.

Given a $\hat{\mathcal{T}}_{\Sigma, \hat{E}}$ -algebra $\hat{\alpha} : \hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{A} \rightarrow \mathbf{A}$, we look at the \mathcal{T}_{Σ} -algebra

$$U^{[-]_{\hat{E}}}\hat{\alpha} = U\hat{\alpha} \circ [-]_{\hat{E}} : \mathcal{T}_{\Sigma}A \rightarrow \mathbf{A}$$

obtained via Lemma 3.79, then we apply P^{-1} to get the Σ -algebra $(A, [-]_{U^{[-]_{\hat{E}}}\hat{\alpha}})$. Since $\mathbf{A} = (A, d_{\mathbf{A}})$ is a generalized metric space (because $\hat{\alpha}$ belongs to $\mathbf{EM}(\hat{\mathcal{T}}_{\Sigma, \hat{E}})$), we obtain a quantitative algebra $\hat{\mathbf{A}}_{\hat{\alpha}} = (A, [-]_{U^{[-]_{\hat{E}}}\hat{\alpha}}, d_{\mathbf{A}})$, and we need to check it satisfies the equations in \hat{E} .

Recall from the proof of Proposition 1.70 that interpreting terms in $\hat{\mathbf{A}}_{\hat{\alpha}}$ is the same thing as applying $U^{[-]_{\hat{E}}}\hat{\alpha} = U\hat{\alpha} \circ [-]_{\hat{E}}$. Therefore, given any L-space \mathbf{X} , nonexpansive assignment $\hat{\iota} : \mathbf{X} \rightarrow \mathbf{A}$, and $t \in \mathcal{T}_{\Sigma}X$, we have

$$\llbracket t \rrbracket_{U^{[-]_{\hat{E}}}\hat{\alpha}}^{\hat{\iota}} \stackrel{(1.10)}{=} \llbracket \mathcal{T}_{\Sigma}\hat{\iota}(t) \rrbracket_{U^{[-]_{\hat{E}}}\hat{\alpha}} = \hat{\alpha}[\mathcal{T}_{\Sigma}\hat{\iota}(t)]_{\hat{E}}.$$

Now, if $\mathbf{X} \vdash s = t \in \hat{E}$, we also have $\mathbf{A} \vdash \mathcal{T}_{\Sigma}\hat{\iota}(s) = \mathcal{T}_{\Sigma}\hat{\iota}(t) \in \Omega\mathfrak{Th}(\hat{E})$ by Lemma 3.38, which means

$$\llbracket s \rrbracket_{U^{[-]_{\hat{E}}}\hat{\alpha}}^{\hat{\iota}} = \hat{\alpha}[\mathcal{T}_{\Sigma}\hat{\iota}(s)]_{\hat{E}} = \hat{\alpha}[\mathcal{T}_{\Sigma}\hat{\iota}(t)]_{\hat{E}} = \llbracket t \rrbracket_{U^{[-]_{\hat{E}}}\hat{\alpha}}^{\hat{\iota}}.$$

Similarly for $\mathbf{X} \vdash s =_{\varepsilon} t \in \hat{E}$, Lemma 3.38 means $\mathbf{A} \vdash \mathcal{T}_{\Sigma}\hat{\iota}(s) =_{\varepsilon} \mathcal{T}_{\Sigma}\hat{\iota}(t) \in \Omega\mathfrak{Th}(\hat{E})$, so⁴⁷²

$$d_{\mathbf{A}}(\llbracket s \rrbracket_{U^{[-]_{\hat{E}}}\hat{\alpha}}^{\hat{\iota}}, \llbracket t \rrbracket_{U^{[-]_{\hat{E}}}\hat{\alpha}}^{\hat{\iota}}) = d_{\mathbf{A}}(\hat{\alpha}[\mathcal{T}_{\Sigma}\hat{\iota}(s)]_{\hat{E}}, \hat{\alpha}[\mathcal{T}_{\Sigma}\hat{\iota}(t)]_{\hat{E}}) \leq d_{\hat{E}}([\mathcal{T}_{\Sigma}\hat{\iota}(s)]_{\hat{E}}, [\mathcal{T}_{\Sigma}\hat{\iota}(t)]_{\hat{E}}) \leq \varepsilon.$$

Finally, if $h : (\mathbf{A}, \hat{\alpha}) \rightarrow (\mathbf{B}, \hat{\beta})$ is $\hat{\mathcal{T}}_{\Sigma, \hat{E}}$ -homomorphism, then by definition, it is nonexpansive $\mathbf{A} \rightarrow \mathbf{B}$, and by Lemma 3.79 it commutes with $U^{[-]_{\hat{E}}}\hat{\alpha}$ and $U^{[-]_{\hat{E}}}\hat{\beta}$ which means it is a homomorphism of the underlying algebras of $\hat{\mathbf{A}}_{\hat{\alpha}}$ and $\hat{\mathbf{B}}_{\hat{\beta}}$. We conclude it is also a homomorphism between the quantitative algebras $\hat{\mathbf{A}}_{\hat{\alpha}}$ and $\hat{\mathbf{B}}_{\hat{\beta}}$.⁴⁷³ We obtain our functor $\hat{P}^{-1} : \mathbf{EM}(\hat{\mathcal{T}}_{\Sigma, \hat{E}}) \rightarrow \mathbf{QAlg}(\Sigma, \hat{E})$.

The diagrams at the start of the proof commute by construction, and P and P^{-1} are inverses by Proposition 1.70. That is enough to conclude that \hat{P} and \hat{P}^{-1} are also inverses. Indeed, by commutativity of the triangle, \hat{P} and \hat{P}^{-1} preserve the

$$\begin{array}{ccc} \mathcal{T}_{\Sigma}A & \xrightarrow{\alpha_{\mathbf{A}}} & \mathbf{A} \\ & \searrow [-]_{\hat{E}} & \nearrow \hat{\alpha}_{\hat{\mathbf{A}}} \\ & \hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{A} & \end{array} \quad (3.53)$$

$$\begin{array}{ccccc} \mathcal{T}_{\Sigma}A & \xrightarrow{\mathcal{T}_{\Sigma}h} & \mathcal{T}_{\Sigma}B & & \\ & \searrow \alpha_{\mathbf{A}} & \searrow \alpha_{\mathbf{B}} & & \\ & & \mathbf{A} & \xrightarrow{h} & \mathbf{B} \\ & \searrow [-]_{\hat{E}} & \searrow [-]_{\hat{E}} & \nearrow \hat{\alpha}_{\hat{\mathbf{A}}} & \nearrow \hat{\alpha}_{\hat{\mathbf{B}}} \\ & & \hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{A} & \xrightarrow{\hat{\mathcal{T}}_{\Sigma, \hat{E}}h} & \hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{B} \end{array} \quad (3.54)$$

The top face of the prism in (3.54) commutes because h is a homomorphism, the back face commutes by (3.19), and the side faces commute by (3.53). Thus, the bottom face commutes because $[-]_{\hat{E}}$ is epic.

⁴⁷² The first inequality holds by nonexpansiveness of $\hat{\alpha}$ and the second by definition of $d_{\hat{E}}$ (3.17).

⁴⁷³ Recall that homomorphisms between quantitative algebras are just nonexpansive homomorphisms.

underlying spaces, and if we fix a space \mathbf{A} , the forgetful functors U and $U^{[-]_{\hat{E}}}$ are injective.⁴⁷⁴ Then, still with a fixed space \mathbf{A} , by commutativity of the square, we have

$$\begin{aligned} U\hat{P}^{-1}\hat{P}\hat{A} &= P^{-1}U^{[-]_{\hat{E}}}\hat{P}\hat{A} = P^{-1}PU\hat{A} = U\hat{A}, \text{ and} \\ U^{[-]_{\hat{E}}}\hat{P}\hat{P}^{-1}\hat{a} &= PU\hat{P}^{-1}\hat{a} = PP^{-1}U^{[-]_{\hat{E}}}\hat{a} = U^{[-]_{\hat{E}}}\hat{a}, \end{aligned}$$

with which we can conclude by injectivity of U and $U^{[-]_{\hat{E}}}$. \square

Remark 3.81. We followed the proof of [MSV22] which does not rely on monadicity theorems (c.f. Remark 1.71).⁴⁷⁵ To show that $U : \mathbf{QAlg}(\Sigma, \hat{E}) \rightarrow \mathbf{GMet}$ is (strictly) monadic, it would be enough to check that the isomorphism we constructed above is the comparison functor.

This motivates the following definition.

Definition 3.82 (GMet presentation). Let M be a monad on \mathbf{GMet} , a **quantitative algebraic presentation** of M is signature Σ and a class of quantitative equations \hat{E} along with a monad isomorphism $\rho : \hat{\mathcal{T}}_{\Sigma, \hat{E}} \cong M$. We also say M is presented by (Σ, \hat{E}) . By Proposition 1.76 and Theorem 3.80, this is equivalent to having an isomorphism $\mathbf{EM}(\hat{\mathcal{T}}_{\Sigma, \hat{E}}) \cong \mathbf{QAlg}(\Sigma, \hat{E})$ that commutes with the forgetful functors.

Example 3.83 (Hausdorff). We saw in Example 1.78 that the monad \mathcal{P}_{ne} on \mathbf{Set} is presented by the theory of semilattices. In this example,⁴⁷⁶ we define the theory of quantitative semilattices and show it presents a monad which sends (X, d) to $\mathcal{P}_{\text{ne}}X$ equipped with the Hausdorff distance d^\dagger .

A **quantitative semilattice** is a semilattice (i.e. a $(\Sigma_{\mathcal{S}}, E_{\mathcal{S}})$ -algebra) equipped with an L-relation such that the interpretation of the semilattice operation is nonexpansive with respect to the product distance. Equivalently, it is a quantitative $\Sigma_{\mathcal{S}}$ -algebra that satisfies $\hat{E}_{\mathcal{S}}$ which contains:⁴⁷⁷

$$\begin{aligned} x \vdash x &= x \oplus x \\ x, y \vdash x \oplus y &= y \oplus x \\ x, y, z \vdash x \oplus (y \oplus z) &= (x \oplus y) \oplus z \\ \forall \varepsilon, \varepsilon' \in \mathbf{L}, \quad x =_{\varepsilon} y, x' =_{\varepsilon'} y' \vdash x \oplus x' &=_{\max\{\varepsilon, \varepsilon'\}} y \oplus y' \end{aligned}$$

We can give an alternative description of the free quantitative semilattice.

Lemma 3.84. *The free quantitative semilattice on (X, d) is $\hat{\mathbb{P}}_{(X, d)} = (\mathcal{P}_{\text{ne}}X, \cup, d^\dagger)$.*⁴⁷⁸

Proof. We know from Example 1.78 that $(\mathcal{P}_{\text{ne}}X, \cup)$ is the free semilattice and hence satisfies $E_{\mathcal{S}}$, thus by Lemma 3.40, $\hat{\mathbb{P}}_{(X, d)}$ satisfies the first three equations above. We already mentioned that $\hat{\mathbb{P}}_{(X, d)}$ satisfies (3.6) because it satisfies (3.1).⁴⁷⁹ Thus, $\hat{\mathbb{P}}_{(X, d)}$ is a quantitative semilattice.

Let \hat{A} be a quantitative semilattice and $f : (X, d) \rightarrow \hat{A}$ be a nonexpansive map. By Lemma 3.41, \hat{A} is a semilattice, hence the universal property of the free semilattice gives a unique homomorphism of $(\Sigma_{\mathcal{S}}, E_{\mathcal{S}})$ -algebras $f^* : (\mathcal{P}_{\text{ne}}X, \cup) \rightarrow \hat{A}$ such that

⁴⁷⁴ For U , it is clear because it only forgets the L-relation. For $U^{[-]_{\hat{E}}}$, it is also not too hard to see, and it is because $U : \mathbf{GMet} \rightarrow \mathbf{Set}$ is faithful and $[-]_{\hat{E}}$ is epic.

⁴⁷⁵ For a proof that does, see [MSV23, Theorems 6.3 and 8.11] where we showed strict monadicity for $[0, 1]$ -spaces first, then for generalized metric spaces using (3.52), and the cancellability of monadicity [Bou92, Proposition 5].

⁴⁷⁶ We adapted it from [MPP16, §9.1].

⁴⁷⁷ The first three equations are those of $E_{\mathcal{S}}$ seen with the discrete context as in Example 3.71. The last row is (3.6) which enforces the nonexpansiveness property of $[\oplus]$.

⁴⁷⁸ This corresponds to [MPP16, Theorem 9.3].

⁴⁷⁹ We did not give a proof for (3.1).

$f^*({x}) = f(x)$ for all $x \in X$. It remains to show that f^* is a nonexpansive map $(\mathcal{P}_{\text{ne}}X, d^\uparrow) \rightarrow \mathbf{A}$.⁴⁸⁰

Let $S, T \in \mathcal{P}_{\text{ne}}X$, $C \in \mathcal{P}_{\text{ne}}(X \times X)$ be a coupling for S and T , and suppose C is ordered with $C = \{c_1, \dots, c_n\}$. In particular, we have $S = \pi_1(c_1) \cup \dots \cup \pi_1(c_n)$ and $T = \pi_2(c_1) \cup \dots \cup \pi_2(c_n)$. Since f^* is a homomorphism of semilattices, this implies

$$\begin{aligned} f^*(S) &= f(\pi_1(c_1)) \llbracket \oplus \rrbracket_A \cdots \llbracket \oplus \rrbracket_A f(\pi_1(c_n)), \text{ and} \\ f^*(T) &= f(\pi_2(c_1)) \llbracket \oplus \rrbracket_A \cdots \llbracket \oplus \rrbracket_A f(\pi_2(c_n)). \end{aligned}$$

Now, we can use the fact that $\hat{\mathbf{A}}$ satisfies the equations in (3.6) n times in the first step of the following derivation.

$$\begin{aligned} d_{\mathbf{A}}(f^*(S), f^*(T)) &\leq \max_{1 \leq i \leq n} d_{\mathbf{A}}(f(\pi_1(c_i)), f(\pi_2(c_i))) && \text{by (3.6)} \\ &\leq \max_{1 \leq i \leq n} d(\pi_1(c_i), \pi_2(c_i)) && f \text{ nonexpansive} \\ &\leq d^\downarrow(S, T) && \text{definition of } d^\downarrow \\ &= d^\uparrow(S, T) && \text{Lemma 2.18} \end{aligned}$$

We conclude that f^* is a homomorphism between the quantitative algebras $\hat{\mathbf{P}}_{(X,d)}$ and $\hat{\mathbf{A}}$. The uniqueness follows from it being unique as a homomorphism of semilattices and the faithfulness of $U : \mathbf{QAlg}(\Sigma_S, \hat{E}_S) \rightarrow \mathbf{Alg}(\Sigma_S)$. \square

Since $\hat{\mathbf{T}}(X, d)$ is also the free quantitative semilattice on (X, d) by Theorem 3.57 and free objects are unique by Proposition 1.48, there is an isomorphism of quantitative algebras $\rho_{(X,d)} : \hat{\mathbf{T}}(X, d) \cong \hat{\mathbf{P}}_{(X,d)}$. After some abstract categorical arguments we do not reproduce, one finds that ρ is a monad isomorphism $\hat{\mathbf{T}}_{\Sigma_S, \hat{E}_S} \cong \mathcal{P}_{\text{ne}}^\uparrow$, where $\mathcal{P}_{\text{ne}}^\uparrow : \mathbf{GMet} \rightarrow \mathbf{GMet}$ sends (X, d) to $(\mathcal{P}_{\text{ne}}X, d^\uparrow)$ and its unit and multiplication act just like those of \mathcal{P}_{ne} .⁴⁸¹

The second example of presentation is from [MPP16, §10.1].

Example 3.85 (Kantorovich). We saw in Example 1.79 that the monad \mathcal{D} on \mathbf{Set} is presented by the theory of convex algebras. Let $\mathbf{L} = [0, \infty]$ and $\mathbf{GMet} = \mathbf{Met}$. The theory of **quantitative convex algebras** is generated by $\hat{E}_{\mathbf{CA}}$ which contains the equations of $E_{\mathbf{CA}}$ seen as quantitative equations (as explained in Example 3.71) and the quantitative equations for convexity (3.10).⁴⁸²

Let $(\mathcal{D}X, \llbracket - \rrbracket_{\mathcal{D}X})$ be the free convex algebra, where $+_p$ is interpreted as convex combination of distributions (1.60). Thanks to Lemma 3.40, we know that for any metric d on X , we can equip $\mathcal{D}X$ with the Kantorovich distance d_K and obtain a quantitative algebra $(\mathcal{D}X, \llbracket - \rrbracket_{\mathcal{D}X}, d_K)$ that satisfies the equations of convex algebras (seen with a discrete context). Moreover, with Example 3.14 we can infer that $(\mathcal{D}X, \llbracket - \rrbracket_{\mathcal{D}X}, d_K)$ is a quantitative convex algebra (i.e. it also satisfies (3.10)). In [MPP16, Theorem 10.5], the authors show that, along with the map $\eta_X^{\mathcal{D}} : (X, d) \rightarrow (\mathcal{D}X, d_K)$ sending x to the Dirac distribution on x , it is the free quantitative convex algebra on (X, d) .

⁴⁸⁰ Actually, you also have to prove that $\eta : (X, d) \rightarrow (\mathcal{P}_{\text{ne}}X, d^\uparrow)$ sending x to $\{x\}$ is nonexpansive. This is easy to check.

⁴⁸¹ This monad is famous independently of quantitative algebras, variations of it were studied in, e.g. [ACT10, §4], [Tho12, §4], [BBKK18, Example 8.3], and [DFM23, §6].

⁴⁸² As a reminder, $\hat{E}_{\mathbf{CA}}$ contains

$$\begin{aligned} x \vdash x &= x +_p x \\ x, y \vdash x +_p y &= y +_{1-p} x \\ x, y, z \vdash (x +_p y) +_q z &= x +_{pq} + (y +_{\frac{p(1-q)}{1-pq}} z) \\ x =_\varepsilon y, x' =_{\varepsilon'} y' \vdash x +_p x' &=_{p\varepsilon + \bar{p}\varepsilon'} y +_p y' \end{aligned}$$

We can conclude that $(\Sigma_{\mathbf{CA}}, \hat{E}_{\mathbf{CA}})$ presents a monad $\mathcal{D}_K : \mathbf{Met} \rightarrow \mathbf{Met}$ which sends (X, d) to $(\mathcal{D}X, d_K)$ and whose unit and multiplication act just like those of the \mathbf{Set} monad \mathcal{D} .⁴⁸³

Here is one last example.

Example 3.86 (Maybe). We saw in Example 1.74 that the maybe monad on \mathbf{Set} is presented by the theory of $\Sigma = \{p : 0\}$ with no equations. Let us generalize this to the maybe monad on \mathbf{GMet} .⁴⁸⁴ We saw in Corollary 3.60 that $\mathbf{QAlg}(\Sigma, \hat{E}_1) \cong \mathbf{1}/\mathbf{GMet}$, where \hat{E}_1 contains the single equation $\vdash p =_\varepsilon p$ with ε being the self-distance of the unique element in $\mathbf{1}$, are the same thing as objects in the coslice. This isomorphism commutes with the forgetful functors to \mathbf{GMet} ,⁴⁸⁵ and we get that the monad $\widehat{\mathcal{T}}_{\Sigma, \hat{E}_1}$ obtained via the existence of free algebras is isomorphic to the monad $- + \mathbf{1}$ which is obtained via the existence of free objects in $\mathbf{1}/\mathbf{GMet}$.⁴⁸⁶

3.5 Lifting Presentations

Most examples of \mathbf{GMet} presentations in the literature, e.g. [MPP16, MV20, MSV21, MSV22], including Examples 3.83, 3.85, and 3.86, are built on top of a \mathbf{Set} presentation. In summary, there is a monad M on \mathbf{Set} with a known algebraic presentation (Σ, E) (e.g. \mathcal{P}_{ne} and semilattices or \mathcal{D} and convex algebras) and a lifting of every space (X, d) to a space (MX, \hat{d}) . Then, a quantitative algebraic theory (Σ, \hat{E}) over the same signature is generated by counterparts to the equations in E as well as new quantitative equations to model the liftings. Finally, it is shown how the theory axiomatizes the lifting, namely, the \mathbf{GMet} monad induced by the theory is isomorphic to a monad whose action on objects is the assignment $(X, d) \mapsto (MX, \hat{d})$.

In this section, we prove Theorem 3.98 which makes this process more automatic and gives necessary and sufficient conditions for when it can actually be done. Throughout, we fix a monad (M, η, μ) on \mathbf{Set} and an algebraic theory (Σ, E) presenting M via an isomorphism $\rho : \mathcal{T}_{\Sigma, E} \cong M$. We first give multiple definitions to make precise what we mean by *lifting*.

Definition 3.87 (Liftings). We have three different notions of lifting that we introduce from weakest to strongest. The last one coincides with the liftings defined in [Bec69].

- A **mere lifting** of M to \mathbf{GMet} is an assignment $(X, d_X) \mapsto (MX, \hat{d}_X)$ defining a generalized metric on MX for every generalized metric on X .⁴⁸⁷
- A **functor lifting** of M to \mathbf{GMet} is a functor $\widehat{M} : \mathbf{GMet} \rightarrow \mathbf{GMet}$ that makes the square below commute.

$$\begin{array}{ccc} \mathbf{GMet} & \xrightarrow{\widehat{M}} & \mathbf{GMet} \\ u \downarrow & & \downarrow u \\ \mathbf{Set} & \xrightarrow{M} & \mathbf{Set} \end{array} \quad (3.55)$$

Note in particular that for every space \mathbf{X} , the carrier of $\widehat{M}\mathbf{X}$ is $M\mathbf{X}$, so we obtain a mere lifting $\mathbf{X} \mapsto \widehat{M}\mathbf{X}$. Furthermore, given a nonexpansive map $f : \mathbf{X} \rightarrow \mathbf{Y}$, the underlying function of $\widehat{M}f$ is Mf , i.e. $Mf : \widehat{M}\mathbf{X} \rightarrow \widehat{M}\mathbf{Y}$ is nonexpansive.

⁴⁸³ This monad is famous independently of quantitative algebras, variations of it were studied in, e.g. [vBo5, §5], [MMM12], [BBKK18, Example 8.4], and [FP19].

⁴⁸⁴ It exists because \mathbf{GMet} has a terminal object (Proposition 2.35) and coproducts (Corollary 3.60).

⁴⁸⁵ The functor $U : \mathbf{1}/\mathbf{GMet} \rightarrow \mathbf{GMet}$ sends the pair $(\mathbf{X}, f : \mathbf{1} \rightarrow \mathbf{X})$ to \mathbf{X} .

⁴⁸⁶ You need to check that $\mathbf{X} + \mathbf{1}$ is indeed the free object on \mathbf{X} in this coslice.

⁴⁸⁷ The name *lifting* more commonly refers to what we call functor lifting or monad lifting which require more conditions than a mere lifting, hence the name *mere lifting*.

In fact, if we have a mere lifting $(X, d_X) \mapsto (MX, \widehat{d}_X)$ such that for every nonexpansive map $f : \mathbf{X} \rightarrow \mathbf{Y}$, $Mf : (MX, \widehat{d}_X) \rightarrow (MY, \widehat{d}_Y)$ is nonexpansive, we automatically get a functor lifting \widehat{M} whose action on objects is given by the mere lifting.⁴⁸⁸ We conclude that functor liftings are just mere liftings with that additional condition.

- A **monad lifting** of M to \mathbf{GMet} is a monad $(\widehat{M}, \widehat{\eta}, \widehat{\mu})$ on \mathbf{GMet} such that \widehat{M} is a functor lifting of M and furthermore $U\widehat{\eta} = \eta U$ and $U\widehat{\mu} = \mu U$. These two equations mean that the underlying functions of the unit and multiplication $\widehat{\eta}_X$ and $\widehat{\mu}_X$ are η_X and μ_X for any space \mathbf{X} .⁴⁸⁹ In particular, the maps

$$\eta_X : \mathbf{X} \rightarrow \widehat{M}\mathbf{X} \quad \text{and} \quad \mu_X : \widehat{M}\widehat{M}\mathbf{X} \rightarrow \widehat{M}\mathbf{X}$$

are nonexpansive for every \mathbf{X} . In fact, since U is faithful, that completely determines $\widehat{\eta}_X$ and $\widehat{\mu}_X$, and we conclude as before that a monad lifting is just a mere lifting with three additional conditions:

1. $Mf : (MX, \widehat{d}_X) \rightarrow (MY, \widehat{d}_Y)$ is nonexpansive if $f : \mathbf{X} \rightarrow \mathbf{Y}$ is nonexpansive,
2. $\eta_X : (X, d_X) \rightarrow (MX, \widehat{d}_X)$ is nonexpansive for every \mathbf{X} , and
3. $\mu_X : (MMX, \widehat{d}_X) \rightarrow (MX, \widehat{d}_X)$ is nonexpansive for every \mathbf{X} .

In practice, when defining a monad lifting, we will define a mere lifting and check Items 1–3. Let us give an example.

Example 3.88. Given an L-space (X, d) , we define an L-relation \widehat{d} on $\mathcal{P}_{\text{ne}}X$ as follows: for any non-empty finite $S, S' \subseteq X$,

$$\widehat{d}(S, S') = \begin{cases} \perp & S = S' \\ d(x, y) & S = \{x\} \text{ and } S' = \{y\} \\ \top & \text{otherwise} \end{cases} \quad (3.56)$$

Instantiating \mathbf{GMet} with the category of L-spaces that satisfy reflexivity ($x \vdash x = \perp$), (3.56) defines a mere lifting of \mathcal{P}_{ne} to \mathbf{GMet} given by $(X, d) \mapsto (\mathcal{P}_{\text{ne}}X, \widehat{d})$.⁴⁹⁰ Viewing \mathcal{P}_{ne} as modelling nondeterminism, this lifting could model a system where nondeterministic processes cannot be meaningfully compared (they are put at maximum distance) unless the sets of possible outcomes are the same (distance is minimal) or both processes are deterministic (distance is inherited from the distance between the only possible outcomes).

We show this is a monad lifting of $(\mathcal{P}_{\text{ne}}, \eta, \mu)$,⁴⁹¹ with Lemmas 3.89–3.91.

Lemma 3.89. *If $f : (X, d) \rightarrow (Y, \Delta)$ is nonexpansive, then so is the direct image function $\mathcal{P}_{\text{ne}}f : (\mathcal{P}_{\text{ne}}X, \widehat{d}) \rightarrow (\mathcal{P}_{\text{ne}}Y, \widehat{\Delta})$.*⁴⁹²

Proof. Let $S, S' \in \mathcal{P}_{\text{ne}}X$. If $S = S'$, then $f(S) = f(S')$, so

$$\widehat{\Delta}(f(S), f(S')) = \perp \leq \perp = \widehat{d}(S, S').$$

If $S = \{x\}$ and $S' = \{y\}$, then $f(S) = \{f(x)\}$ and $f(S') = \{f(y)\}$, so⁴⁹³

⁴⁸⁸ The action on morphisms is prescribed by (3.55), namely, the underlying function of $\widehat{M}f$ is Mf which is nonexpansive by hypothesis, and since U is faithful, that determines $\widehat{M}f$.

⁴⁸⁹ In summary, the description of a monad M and its monad lifting \widehat{M} are exactly the same after forgetting about distances. In particular, the action of \widehat{M} on morphisms does not depend on the distances at the source or the target, and similarly, the unit and multiplication maps do not depend on the distance of the space.

⁴⁹⁰ We need reflexivity to ensure the first and second cases do not clash. You can also check that whenever d is a metric space, \widehat{d} is as well, so we get a mere lifting of \mathcal{P}_{ne} to \mathbf{Met} as well.

⁴⁹¹ The unit and multiplication of \mathcal{P}_{ne} were defined in Example 1.64.

⁴⁹² We write $f(S)$ instead of $\mathcal{P}_{\text{ne}}f(S)$ for better readability.

⁴⁹³ The inequality holds because f is nonexpansive.

$$\widehat{\Delta}(f(S), f(S')) = \Delta(f(x), f(y)) \leq d(x, y) = \widehat{d}(S, S').$$

Otherwise, $\widehat{d}(S, S') = \top$ and $\widehat{\Delta}(f(S), f(S'))$ is always less or equal to \top . \square

Lemma 3.90. *For any (X, d) , the map $\eta_X : (X, d) \rightarrow (\mathcal{P}_{\text{ne}}X, \widehat{d})$ is nonexpansive.*

Proof. Recall that $\eta_X(x) = \{x\}$. For any $x, y \in X$, $\widehat{d}(\{x\}, \{y\}) = d(x, y)$, so η_X is even an isometry. \square

Lemma 3.91. *For any (X, d) , the map $\mu_X : (\mathcal{P}_{\text{ne}}\mathcal{P}_{\text{ne}}X, \widehat{\widehat{d}}) \rightarrow (\mathcal{P}_{\text{ne}}X, \widehat{d})$ is nonexpansive.*

Proof. Recall that $\mu_X(F) = \cup F$ and let $F, F' \in \mathcal{P}_{\text{ne}}\mathcal{P}_{\text{ne}}X$. The case $F = F'$ is dealt with like in Lemma 3.89, it implies $\cup F = \cup F'$, hence the distances on both sides are \perp . If $F = \{S\}$ and $F' = \{S'\}$, $\cup F = S$ and $\cup F' = S'$, then

$$\widehat{d}(\mu_X(F), \mu_X(F')) = \widehat{d}(S, S') = \widehat{\widehat{d}}(\{S\}, \{S'\}).$$

Otherwise, $\widehat{\widehat{d}}(\{S\}, \{S'\}) = \top$, so the inequality holds because $\widehat{d}(\mu_X(F), \mu_X(F'))$ is always less or equal to \top . \square

Here is an example of a functor lifting that is not a monad lifting.

Example 3.92. The **total variation** distance is a metric defined on probability distributions. For any X , we define $\text{tv} : \mathcal{D}X \times \mathcal{D}X \rightarrow [0, 1]$ by, for any $\varphi, \psi \in \mathcal{D}X$,⁴⁹⁴

$$\text{tv}(\varphi, \psi) = \sup_{S \subseteq X} |\varphi(S) - \psi(S)|.$$

Even though the assignment $(X, d) \mapsto (\mathcal{D}X, \text{tv})$ is a mere lifting of the monad \mathcal{D} to **Met**, namely, $(\mathcal{D}X, \text{tv})$ is a metric whenever (X, d) is, it is not a monad lifting. One can show that Mf is nonexpansive whenever f is, so it is a functor lifting, and even that the multiplication is always nonexpansive, but the unit is not because if $x \neq y \in X$ are points at distance $d(x, y) < 1$, then $\text{tv}(\delta_x, \delta_y) = 1 > d(x, y)$.

Many monads of interest on different **GMet** categories are monad liftings of **Set** monads which have an algebraic presentation. We already mentioned the Hausdorff and Kantorovich monad liftings in Examples 3.83 and 3.85, but there is also a combination of the two: the Hausdorff–Kantorovich monad lifting of the convex sets of distributions monad [MV20] to **Met**. In [MSV21], we further combined these with the maybe monad on **Met**. Another example is the formal ball monad on quasi-metric spaces [GL19] which is a monad lifting of a writer monad on **Set**. All of these happen to have a quantitative algebraic presentation,⁴⁹⁵ and we will show that this is not a coincidence.

Given a monad lifting \widehat{M} , we know that it acts on sets just like M does, and that can be described algebraically through the presentation $\rho : \mathcal{T}_{\Sigma, E} \cong M$. This can help to understand how \widehat{M} acts on distances. For any space \mathbf{X} , we see the distance $\widehat{d}_{\mathbf{X}}$ on $M\mathbf{X}$ as a distance \widehat{d} on terms modulo E via the bijection $\rho_{\mathbf{X}}$:⁴⁹⁶

$$\widehat{d}([s]_E, [t]_E) = \widehat{d}_{\mathbf{X}}(\rho_{\mathbf{X}}[s]_E, \rho_{\mathbf{X}}[t]_E).$$

⁴⁹⁴ Since φ and ψ have finite support, we can restrict the quantification of the supremum to finite subsets of X , or even to subsets of the union of the supports of φ and ψ . Also, both $\varphi(S)$ and $\psi(S)$ are at most in $[0, 1]$, so $\text{tv}(\varphi, \psi)$ also takes values in $[0, 1]$.

⁴⁹⁵ Goubault-Larrecq does not talk about quantitative algebras in [GL19], but the quantitative writer monad of [BMP21, §4.3.2] has a presentation which can easily be adapted to present the monad of [GL19].

⁴⁹⁶ Recall Proposition 2.49.

Can we find some quantitative equations \hat{E} that axiomatize \hat{d} , i.e. such that $d_{\hat{E}}$ and \hat{d} are isomorphic (uniformly for all \mathbf{X})?

First of all, for the distances to be isomorphic, they need to be on the same set, namely, we need to have $\mathcal{T}_{\Sigma}X/\equiv_E \cong \mathcal{T}_{\Sigma}X/\equiv_{\hat{E}}$, or equivalently, $s \equiv_E t \iff s \equiv_{\hat{E}} t$. At once, this removes some options for which equations to add in \hat{E} . For instance, we cannot add $\mathbf{X} \vdash s = t$ if $X \vdash s = t$ does not already belong to $\mathfrak{Th}(E)$. Conversely, if $X \vdash s = t \in \mathfrak{Th}(E)$, we need to ensure $\mathbf{X} \vdash s = t$ belongs to $\Omega\mathfrak{Th}(\hat{E})$. We can do this by adding $\mathbf{X} \vdash s = t$ to \hat{E} thanks to Example 3.71.

After that, we will have to add quantitative equations with quantities to axiomatize \hat{d} , but we have to be careful not to break the equivalence we just obtained between \equiv_E and $\equiv_{\hat{E}}$. For instance, if $\mathbf{GMet} = \mathbf{Met}$, $f : 1 \in \Sigma$ and $E = \emptyset$, then we cannot have $x =_{\frac{1}{2}} y \vdash fx =_0 fy \in \hat{E}$, because using the equation $x =_0 y \vdash x = y$ that defines \mathbf{Met} , we could conclude that $x =_{\frac{1}{2}} y \vdash fx = fy$ belongs to $\Omega\mathfrak{Th}(\hat{E})$, which means $fx \equiv_{\hat{E}} fy$ whenever $d_{\mathbf{X}}(x, y) \leq \frac{1}{2}$ while $fx \not\equiv_E fy$.

The relation between \hat{E} and E seems to mimic our intuition about mere liftings. We say that \hat{E} extends E .

Definition 3.93 (Extension). Given a class E of equations over Σ and a class \hat{E} of quantitative equations over Σ , we say that \hat{E} is an **extension** of E if for all $\mathbf{X} \in \mathbf{GMet}$ and $s, t \in \mathcal{T}_{\Sigma}X$,

$$X \vdash s = t \in \mathfrak{Th}(E) \iff \mathbf{X} \vdash s = t \in \Omega\mathfrak{Th}(\hat{E}). \quad (3.57)$$

Remark 3.94. Let us make two delicate points on the quantification of \mathbf{X} in (3.57).

First, it happens *before* the equivalence. This means that equalities⁴⁹⁷ that hold in $\mathcal{T}_{\Sigma, E}X$ coincide with the equalities that hold in $\hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}$ for each \mathbf{X} individually. In particular, if \mathbf{X} and \mathbf{X}' are spaces on the same set X , then the equalities that hold in $\hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}$ and $\hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}'$ coincide. This intuitively corresponds to the fact that the action of $\hat{\mathcal{T}}_{\Sigma, \hat{E}}$ does not depend on distances.

If instead of (3.57) we had the following equivalence with the quantification inside,

$$X \vdash s = t \in \mathfrak{Th}(E) \iff \forall \mathbf{X} \in \mathbf{GMet}, \mathbf{X} \vdash s = t \in \Omega\mathfrak{Th}(\hat{E}),$$

then the equalities in $\mathcal{T}_{\Sigma, E}X$ would be those that hold in all $\hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}$ (for all spaces \mathbf{X} with carrier X). In particular, $\hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}$ and $\hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}'$ could have different equivalence classes. That is not desirable when defining a mere lifting.

Second, even though the context of a quantitative equation can be any L-space, \mathbf{X} is only quantified over generalized metric spaces here. This implies that the equivalence classes of $\hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}$ and $\hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}'$ may be different if $d_{\mathbf{X}}$ and $d_{\mathbf{X}'}$ are two different L-relations on X . This does not contradict our intuition about liftings because we only care about the action of $\hat{\mathcal{T}}_{\Sigma, \hat{E}}$ on L-spaces that belong to \mathbf{GMet} .

For instance, let $\Sigma = \{f : 1\}$, $E = \emptyset$, $\hat{E} = \emptyset$, and \mathbf{GMet} be defined by the equation $x =_{\perp} y \vdash x = x$. If $X = \{x, y\}$ and $d_{\mathbf{X}}(x, y) = \perp$, then $\mathbf{X} \vdash fx = fy$ belongs to $\Omega\mathfrak{Th}(\hat{E})$ while $fx \not\equiv_E fy$.⁴⁹⁸ Still, it makes sense that \hat{E} extend E since both have no equations.

It turns out that extensions are stronger than mere liftings because we can show the monad we constructed via terms modulo \hat{E} is a monad lifting of $\mathcal{T}_{\Sigma, E}$.

⁴⁹⁷ This is not a formal term: by *equalities that hold*, we mean which Σ -terms are in the same equivalence class.

⁴⁹⁸ Here is the derivation (the application of \mathbf{GMet} implicitly uses the fact that $x =_{\perp} y \vdash x = x$ is syntactic sugar for $\mathbf{X} \vdash x =_{\perp} y$):

$$\frac{\frac{}{\mathbf{X} \vdash x = y} \text{GMET}}{\mathbf{X} \vdash fx = fy} \text{CONG}$$

Proposition 3.95. *If \hat{E} is an extension of E , then $\widehat{\mathcal{T}}_{\Sigma, \hat{E}}$ is a monad lifting of $\mathcal{T}_{\Sigma, E}$.*

Proof. We need to check the following three equations where $U : \mathbf{GMet} \rightarrow \mathbf{Set}$ is the forgetful functor:

$$U\widehat{\mathcal{T}}_{\Sigma, \hat{E}} = \mathcal{T}_{\Sigma, E}U \quad U\widehat{\eta}^{\Sigma, \hat{E}} = \eta^{\Sigma, E}U \quad U\widehat{\mu}^{\Sigma, \hat{E}} = \mu^{\Sigma, E}U.$$

First, we have to show that for any space \mathbf{X} , $U\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} = \mathcal{T}_{\Sigma, E}U\mathbf{X}$. By definitions, the L.H.S. is $\mathcal{T}_{\Sigma}X/\equiv_{\hat{E}}$ and the R.H.S. is $\mathcal{T}_{\Sigma}X/\equiv_E$, so it boils down to showing that for all $s, t \in \mathcal{T}_{\Sigma}X$, $s \equiv_{\hat{E}} t \iff s \equiv_E t$. This readily follows from the definitions of $\equiv_{\hat{E}}$ and \equiv_E , and from (3.57):⁴⁹⁹

$$s \equiv_{\hat{E}} t \xleftrightarrow{(3.13)} \mathbf{X} \vdash s = t \in \Omega\mathfrak{H}(\hat{E}) \xleftrightarrow{(3.57)} \mathbf{X} \vdash s = t \in \mathfrak{H}(E) \xleftrightarrow{(1.24)} s \equiv_E t.$$

Next, we have to show that $U\widehat{\mathcal{T}}_{\Sigma, \hat{E}}f = \mathcal{T}_{\Sigma, E}f$ for any $f : \mathbf{X} \rightarrow \mathbf{Y}$. This is done rather quickly by comparing their definitions, they make the same squares (1.26) and (3.19) commute now that we know $\equiv_{\hat{E}}$ and \equiv_E coincide.

This takes care of the first equation, and the other two are done very similarly, we compare the definitions of $\widehat{\eta}^{\Sigma, \hat{E}}$ and $\eta^{\Sigma, E}$ (resp. $\widehat{\mu}^{\Sigma, \hat{E}}$ and $\mu^{\Sigma, E}$) and conclude they are the same when $\equiv_{\hat{E}}$ and \equiv_E coincide.⁵⁰⁰ \square

So if we are able to construct an extension \hat{E} of E , we can obtain a monad lifting of M by passing through the isomorphism $\rho : \mathcal{T}_{\Sigma, E} \cong M$.

Corollary 3.96. *If M is presented by (Σ, E) , and \hat{E} is an extension of E , then \hat{E} presents a monad lifting of M .*

Proof. We first construct a monad lifting of (M, η, μ) . For any space \mathbf{X} , we have an isomorphism $\rho_X^{-1} : MX \rightarrow \mathcal{T}_{\Sigma, E}X$, and a generalized metric $d_{\hat{E}}$ on $\mathcal{T}_{\Sigma, E}$ (since the underlying set of $\widehat{\mathcal{T}}_{\Sigma, \hat{E}}$ is $\mathcal{T}_{\Sigma, E}$ by Proposition 3.95). We can define a generalized metric \widehat{d}_X on MX as we have done for Proposition 2.49 to guarantee that $\rho_X^{-1} : (MX, \widehat{d}_X) \rightarrow \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X}$ is an isomorphism:⁵⁰¹

$$\widehat{d}_X(m, m') = d_{\hat{E}}(\rho_X^{-1}(m), \rho_X^{-1}(m')). \quad (3.58)$$

This yields a mere lifting $(X, d_X) \mapsto (MX, \widehat{d}_X)$.

In order to show this is a monad lifting, we use the following diagrams (quantified for all $\mathbf{X} \in \mathbf{GMet}$ and nonexpansive $f : \mathbf{X} \rightarrow \mathbf{Y}$) which commute because ρ is a monad isomorphism with inverse ρ^{-1} .⁵⁰²

$$\begin{array}{ccc} (MX, \widehat{d}_X) & \xrightarrow{\rho_X^{-1}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} \\ Mf \downarrow & & \downarrow \mathcal{T}_{\Sigma, E}f \\ (MY, \widehat{d}_Y) & \xleftarrow{\rho_Y} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{Y} \end{array} \quad \begin{array}{ccc} \mathbf{X} & \xrightarrow{\eta_X^{\Sigma, E}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} \\ & \searrow \eta_X & \downarrow \rho_X \\ & & (MX, \widehat{d}_X) \end{array}$$

$$\begin{array}{ccc} (MMX, \widehat{d}_X) & \xrightarrow{\rho_{MX}^{-1}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}}(X, \widehat{d}_X) & \xrightarrow{\mathcal{T}_{\Sigma, E}\rho_X^{-1}} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} \\ \mu_X \downarrow & & & & \downarrow \mu_X^{\Sigma, E} \\ (MX, \widehat{d}_X) & \xleftarrow{\rho_X} & \widehat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} & & \end{array}$$

⁴⁹⁹ Note again the importance of being able to do this for each \mathbf{X} individually.

⁵⁰⁰ We defined $\widehat{\eta}^{\Sigma, \hat{E}}$ in (3.34), $\eta^{\Sigma, E}$ in Footnote 124, $\widehat{\mu}^{\Sigma, \hat{E}}$ in (3.22), and $\mu^{\Sigma, E}$ in (1.35).

⁵⁰¹ In words, the distance between m and m' in MX is computed by viewing them as (equivalence classes of) terms in $\mathcal{T}_{\Sigma}X$, then using the distance between them given by $d_{\hat{E}}$.

⁵⁰² The first holds by naturality, the second by (1.54), and the third by (1.55). Moreover, all the functions in these diagrams are nonexpansive (with the sources and targets as drawn) by previous results:

- We just showed the components of ρ are isometries.
- We showed $\mathcal{T}_{\Sigma, E}f$ is the underlying function of $\widehat{\mathcal{T}}_{\Sigma, \hat{E}}f$ because $\widehat{\mathcal{T}}_{\Sigma, \hat{E}}$ is a monad lifting of $\mathcal{T}_{\Sigma, E}$ (Proposition 3.95), so $\mathcal{T}_{\Sigma, E}f$ is nonexpansive when f is nonexpansive.
- By the previous two points, $\mathcal{T}_{\Sigma, E}\rho_X^{-1}$ is nonexpansive.
- Again since $\widehat{\mathcal{T}}_{\Sigma, \hat{E}}$ is a monad lifting of $\mathcal{T}_{\Sigma, E}$, $\eta_X^{\Sigma, E}$ and $\mu_X^{\Sigma, E}$ are nonexpansive.

These show (detailed in the footnote) that Mf , η_X and μ_X are compositions of nonexpansive maps, and hence are nonexpansive. We obtain a monad lifting \widehat{M} of M to \mathbf{GMet} which sends (X, d_X) to (MX, \widehat{d}_X) .

It remains to show that \widehat{M} is presented by (Σ, \widehat{E}) . By construction, we have the isomorphism $\widehat{\rho}_X : \widehat{\mathcal{T}}_{\Sigma, \widehat{E}} \mathbf{X} \rightarrow \widehat{M}\mathbf{X}$ whose underlying function is ρ_X for every \mathbf{X} . The fact that $\widehat{\rho}$ is a monad morphism follows from the facts that ρ is a monad morphism, and that $U : \mathbf{GMet} \rightarrow \mathbf{Set}$ is faithful so it reflects commutativity of diagrams.⁵⁰³ \square

Now, we would like to have a converse to Corollary 3.96. Namely, if $(X, d_X) \mapsto (MX, \widehat{d}_X)$ is given by a monad lifting \widehat{M} of M to \mathbf{GMet} , our goal is to construct an extension \widehat{E} of E such that the monad lifting corresponding to \widehat{E} (given in Corollary 3.96) is \widehat{M} . There is no obvious reason this is even possible, maybe \widehat{M} is a monad lifting that has no quantitative algebraic presentation.⁵⁰⁴ Our next theorem shows that such an \widehat{E} always exists. In fact, it is constructed very naively.

As discussed in Example 3.71, when \widehat{E} contains all the quantitative equations in

$$\widehat{E}_1 = \{ \mathbf{X} \vdash s = t \mid X \vdash s = t \in E \}, \quad (3.59)$$

then we have at least one direction of (3.57), namely, that $X \vdash s = t \in \mathfrak{Th}(E)$ implies $\mathbf{X} \vdash s = t \in \Omega\mathfrak{Th}(\widehat{E})$ for all \mathbf{X} and $s, t \in \mathcal{T}_X$.⁵⁰⁵ Next, we include in \widehat{E} all the possible equations $\mathbf{X} \vdash s =_\varepsilon t$ where ε is the distance between s and t when viewed inside $\widehat{M}\mathbf{X}$ (via ρ_X),⁵⁰⁶ namely, $\widehat{E}_2 \subseteq \widehat{E}$ where

$$\widehat{E}_2 = \{ \mathbf{X} \vdash s =_\varepsilon t \mid \mathbf{X} \in \mathbf{GMet}, s, t \in \mathcal{T}_X, \varepsilon = \widehat{d}_X(\rho_X[s]_E, \rho_X[t]_E) \}. \quad (3.60)$$

This is a very large bunch of equations (it is not even a set), but it leaves no stone unturned, meaning that the distance computed by \widehat{E} will always be smaller than the distance in $\widehat{M}\mathbf{X}$. Indeed, for any $m, m' \in MX$, letting $s, t \in \mathcal{T}_X$ be such that $\rho_X[s]_E = m$ and $\rho_X[t]_E = m'$ (by surjectivity of ρ_X), we have⁵⁰⁷

$$\begin{aligned} \widehat{d}_X(m, m') \leq \varepsilon &\implies \mathbf{X} \vdash s =_\varepsilon t \in \Omega\mathfrak{Th}(\widehat{E}) \\ &\iff d_{\widehat{E}}([s]_E, [t]_E) \leq \varepsilon \\ &\iff d_{\widehat{E}}(\rho_X^{-1}(m), \rho_X^{-1}(m')) \leq \varepsilon. \end{aligned}$$

In order to conclude that $\widehat{E} = \widehat{E}_1 \cup \widehat{E}_2$ presents \widehat{M} , we need to show that \widehat{E} is an extension of E , i.e. the other direction of (3.57), and that the monad lifting defined in Corollary 3.96 coincides with \widehat{M} , i.e. the converse implication of the previous derivation holds. We will prove these by constructing a (family of) special algebras in $\mathbf{QAlg}(\Sigma, \widehat{E})$.⁵⁰⁸

For any generalized metric space \mathbf{A} , we denote by \mathbf{MA} the quantitative Σ -algebra $(MA, \llbracket - \rrbracket_{\mu_A}, \widehat{d}_A)$, where

- (MA, \widehat{d}_A) is the space obtained by applying \widehat{M} to \mathbf{A} , and
- $(MA, \llbracket - \rrbracket_{\mu_A})$ is the Σ -algebra obtained by applying the isomorphism $\mathbf{Alg}(\Sigma, E) \cong \mathbf{EM}(M)$ (from the presentation) to the M -algebra (MA, μ_A) (from Example 1.69).

We can show that \mathbf{MA} belongs to $\mathbf{QAlg}(\Sigma, \widehat{E}_1 \cup \widehat{E}_2)$.

⁵⁰³ Let us detail the argument for naturality, the others would follow the same pattern. We need to show that $\widehat{\rho}_Y \circ \widehat{M}f = \widehat{M}f \circ \widehat{\rho}_X$. Applying U , we get $\rho_Y \circ Mf = Mf \circ \rho_X$ which is true because ρ is natural, hence $U(\widehat{\rho}_Y \circ \widehat{M}f) = U(\widehat{M}f \circ \widehat{\rho}_X)$. Since U is faithful, and the desired equation holds.

⁵⁰⁴ Or maybe \widehat{M} has a presentation that is not an extension of E , but our informal discussion leading to the definition of extensions indicates that is less probable.

⁵⁰⁵ We use Lemma 3.72.

⁵⁰⁶ We are essentially doing the opposite of (3.58).

⁵⁰⁷ The implication follows because by definition, \widehat{E} will contain $\mathbf{X} \vdash s =_{d_X(m, m')} t$, hence by the **MON** rule, we will have $\mathbf{X} \vdash s =_\varepsilon t \in \Omega\mathfrak{Th}(\widehat{E})$. The first equivalence is (3.17), and the second holds because ρ_X^{-1} is the inverse of ρ_X .

⁵⁰⁸ It turns out (after the rest of the proof) we are constructing the free algebra over \mathbf{A} , but we feel it is not necessary to make that explicit.

Lemma 3.97. For all $\phi \in \hat{E}_1 \cup \hat{E}_2$, $\mathbb{M}\mathbf{A} \models \phi$.

Proof. If $\phi = \mathbf{X} \vdash s = t \in \hat{E}_1$, then by construction $(MA, \llbracket - \rrbracket_{\mu_A})$ satisfies $X \vdash s = t \in E$. So $\mathbb{M}\mathbf{A}$ satisfies ϕ by Lemma 3.40.

Suppose now that $\phi = \mathbf{X} \vdash s =_\varepsilon t \in \hat{E}_2$ with $\varepsilon = \widehat{d}_{\mathbf{X}}(\rho_X[s]_E, \rho_X[t]_E)$. A bit of unrolling⁵⁰⁹ shows that for an assignment $\iota : X \rightarrow MA$, the interpretation $\llbracket - \rrbracket_{\mu_A}^\iota$ is the composite

$$\mathcal{T}_\Sigma X \xrightarrow{\overline{\mathcal{T}}_\Sigma \iota} \mathcal{T}_\Sigma MA \xrightarrow{[-]_E} \mathcal{T}_{\Sigma, E} MA \xrightarrow{\rho_{MA}} MMA \xrightarrow{\mu_A} MA.$$

For later use, we apply the naturality of $[-]_E$ (1.26) and ρ to rewrite the composite as

$$\llbracket - \rrbracket_{\mu_A}^\iota = \mathcal{T}_\Sigma X \xrightarrow{[-]_E} \mathcal{T}_{\Sigma, E} X \xrightarrow{\rho_X} MX \xrightarrow{M\iota} MMA \xrightarrow{\mu_A} MA. \quad (3.61)$$

We conclude that $\mathbb{M}\mathbf{A} \models \phi$ with the following derivation which holds for all nonexpansive $\hat{\iota} : \mathbf{X} \rightarrow \widehat{M}\mathbf{A}$.⁵¹⁰

$$\begin{aligned} \widehat{d}_{\mathbf{A}}(\llbracket [s] \rrbracket_{\mu_A}^\iota, \llbracket [t] \rrbracket_{\mu_A}^\iota) &= \widehat{d}_{\mathbf{A}}(\mu_A(M\hat{\iota}(\rho_X[s]_E)), \mu_A(M\hat{\iota}(\rho_X[t]_E))) \quad \text{by (3.61)} \\ &\leq \widehat{d}_{\mathbf{A}}(M\hat{\iota}(\rho_X[s]_E), M\hat{\iota}(\rho_X[t]_E)) \quad \mu_A \text{ is nonexpansive} \\ &\leq \widehat{d}_{\mathbf{X}}(\rho_X[s]_E, \rho_X[t]_E) \quad M\hat{\iota} \text{ is nonexpansive} \\ &= \varepsilon \quad \square \end{aligned}$$

Theorem 3.98. Let \widehat{M} be a monad lifting of M to \mathbf{GMet} , and $\hat{E} = \hat{E}_1 \cup \hat{E}_2$. Then, \hat{E} is an extension of E and it presents \widehat{M} .

Proof. We already showed the forward implication of (3.57) when we defined \hat{E}_1 (3.59). For the converse, suppose that $\mathbf{X} \vdash s = t \in \Omega\mathfrak{Th}(\hat{E})$, we saw in Lemma 3.97 that $\mathbb{M}\mathbf{X}$ satisfies $\mathbf{X} \vdash s = t$. Taking the assignment $\eta_X : \mathbf{X} \rightarrow \widehat{M}\mathbf{X}$ which is nonexpansive because \widehat{M} is a monad lifting, we have $\llbracket [s] \rrbracket_{\mu_X}^{\eta_X} = \llbracket [t] \rrbracket_{\mu_X}^{\eta_X}$. Using (3.61) and the monad law $\mu_X \circ M\eta_X = \text{id}_{MX}$ (left triangle in (1.43)), we find

$$\rho_X[s]_E = \mu_X(M\eta_X(\rho_X[s]_E)) = \llbracket [s] \rrbracket_{\mu_X}^{\eta_X} = \llbracket [t] \rrbracket_{\mu_X}^{\eta_X} = \mu_X(M\eta_X(\rho_X[t]_E)) = \rho_X[t]_E.$$

Finally, since ρ_X is a bijection, we have $[s]_E = [t]_E$, i.e. $\mathbf{X} \vdash s = t \in \mathfrak{Th}(E)$.

We already showed that $\widehat{d}_{\mathbf{X}}(m, m') \geq d_{\hat{E}}(\rho_X^{-1}(m), \rho_X^{-1}(m'))$ when defining \hat{E}_2 . For the converse, let $m = \rho_X[s]_E$ and $m' = \rho_X[t]_E$ for some $s, t \in \mathcal{T}_\Sigma X$ and suppose that $d_{\hat{E}}([s]_E, [t]_E) \leq \varepsilon$, or equivalently by (3.17), that $\mathbf{X} \vdash s =_\varepsilon t \in \Omega\mathfrak{Th}(\hat{E})$. As above, Lemma 3.97 says that $\mathbb{M}\mathbf{X}$ satisfies that equation. Taking the assignment $\eta_X : \mathbf{X} \rightarrow \widehat{M}\mathbf{X}$ which is nonexpansive because \widehat{M} is a monad lifting, we have⁵¹¹

$$\widehat{d}_{\mathbf{X}}(m, m') = \widehat{d}_{\mathbf{X}}(\rho_X[s]_E, \rho_X[t]_E) = \widehat{d}_{\mathbf{X}}(\llbracket [s] \rrbracket_{\mu_X}^{\eta_X}, \llbracket [t] \rrbracket_{\mu_X}^{\eta_X}) \leq \varepsilon.$$

Comparing with (3.58), we conclude that \widehat{M} is exactly the monad lifting from Corollary 3.96. In particular, \hat{E} presents \widehat{M} via $\hat{\rho}$ whose component at \mathbf{X} is ρ_X . \square

Remark 3.99. A deeper result hides behind the last line. It follows from our constructions that if you start from an extension \hat{E} , build a monad lifting \widehat{M} from \hat{E} with

⁵⁰⁹ Look at the definition of P^{-1} in Proposition 1.70, in particular what we proved in Footnote 179, and the definition of $-\rho$ in (1.59).

⁵¹⁰ Our hypothesis that \widehat{M} is a monad lifting yields nonexpansiveness of μ_A and $M\hat{\iota}$.

⁵¹¹ The second inequality holds again by (3.61) and (1.43).

Corollary 3.96, then build an extension \hat{E}' from \hat{M} with Theorem 3.98, you obtain two *equivalent* classes of equations, i.e. $\Omega\mathfrak{Th}(\hat{E}) = \Omega\mathfrak{Th}(\hat{E}')$. Similarly, if you start with a monad lifting \hat{M} , then build an extension \hat{E} , then build a monad lifting \hat{M}' , then $\hat{M} = \hat{M}'$.⁵¹²

This does not yield a bijection but almost. If you restrict extensions of E to those that are quantitative algebraic theories,⁵¹³ then you get a bijection with monad liftings of M .

I believe it is a simple exercise in categorical logic to transform this remark into an (dual) equivalence of categories.⁵¹⁴ A slightly more challenging task would be to allow M and E to vary to get a (fibered) equivalence.

When constructing the extension $\hat{E} = \hat{E}_1 \cup \hat{E}_2$, \hat{E}_1 can be fairly small since it has the size of E , but \hat{E}_2 as defined is always huge (not even a set). In theory, some results in the literature could allow us to restrict the size of contexts to be of a moderate size only with mild size conditions on L and $\hat{E}_{\mathbf{GMet}}$.⁵¹⁵ In practice, we can sometimes find some simple set of quantitative equations which will be equivalent to \hat{E}_2 (when \hat{E}_1 is present), and we give a couple of examples below. They require some *clever* arguments that depend on the application, but there may be room for optimization in the definition of \hat{E}_2 .

Example 3.100 (Trivial Lifting of \mathcal{P}_{ne}). Recall the monad lifting of \mathcal{P}_{ne} to $\mathbf{GMet} = \mathbf{QAlg}(\emptyset, \{x \vdash x =_{\perp} x\})$ from Example 3.88. Let us denote it by $\hat{\mathcal{P}}$, and its action on objects by $(X, d) \mapsto (\mathcal{P}_{\text{ne}}X, \widehat{d_X})$.⁵¹⁶ We also denote with ρ the monad isomorphism witnessing that \mathcal{P}_{ne} is presented by the theory of semilattices (Σ_S, E_S) (recall Example 1.78). By Theorem 3.98, there is a quantitative algebraic presentation for $\hat{\mathcal{P}}$ given by⁵¹⁷

$$\hat{E}_1 = \{\mathbf{X} \vdash s = t \mid X \vdash s = t \in E_S\} \text{ and } \hat{E}_2 = \left\{ \mathbf{X} \vdash s =_{\varepsilon} t \mid \varepsilon = \widehat{d_X} \left(\rho_X[s]_{E_S}, \rho_X[t]_{E_S} \right) \right\}.$$

We claim that the equations in \hat{E}_1 are enough, namely, $\Omega\mathfrak{Th}(\hat{E}_1 \cup \hat{E}_2) = \Omega\mathfrak{Th}(\hat{E}_1)$.

First, since $\hat{E}_1 \subseteq \hat{E}_1 \cup \hat{E}_2$, we infer that $\Omega\mathfrak{Th}(\hat{E}_1) \subseteq \Omega\mathfrak{Th}(\hat{E}_1 \cup \hat{E}_2)$.⁵¹⁸

Second, recall from Lemma 3.72 that with the equations in \hat{E}_1 , we can already prove all the equations in the theory of semilattices. This means that for any $\mathbf{X} \vdash s =_{\varepsilon} t \in \hat{E}_2$ with $\varepsilon = \widehat{d_X} \left(\rho_X[s]_{E_S}, \rho_X[t]_{E_S} \right)$, we have the three following cases.

- If $[s]_{E_S} = [t]_{E_S}$ and $\varepsilon = \perp$, i.e. s and t represent the same subset of X , then the equation $X \vdash s = t$ is in $\mathfrak{Th}(E_S)$ which implies $\mathbf{X} \vdash s = t$ is in $\Omega\mathfrak{Th}(\hat{E}_1)$. It follows by the following derivation that $\mathbf{X} \vdash s =_0 t \in \Omega\mathfrak{Th}(\hat{E}_1)$ as desired.⁵¹⁹

$$\frac{\mathbf{X} \vdash s = t \quad \frac{\sigma = x \mapsto s \quad \frac{x \vdash x =_{\perp} x \quad \mathbf{GMET}}{\mathbf{X} \vdash s =_{\top} s} \text{TOP}}{\mathbf{X} \vdash s =_{\perp} s} \text{SUBQ}}{\mathbf{X} \vdash s =_{\perp} t} \text{COMPR}$$

- If $[s]_{E_S} = [x]_{E_S}$ and $[t]_{E_S} = [y]_{E_S}$ for some $x, y \in X$ and $\varepsilon = d_X(x, y)$, then the equations $X \vdash s = x$ and $X \vdash y = t$ are in $\mathfrak{Th}(E_S)$ which implies $\mathbf{X} \vdash s = x$ and $\mathbf{X} \vdash y = t$ are in $\Omega\mathfrak{Th}(\hat{E}_1)$. Furthermore, Lemma 3.35 implies $\mathbf{X} \vdash x =_{\varepsilon} y \in \Omega\mathfrak{Th}(\hat{E}_1)$, and finally by Lemmas 3.32 and 3.33, $\mathbf{X} \vdash s =_{\varepsilon} t$ also belongs to $\Omega\mathfrak{Th}(\hat{E}_1)$ as desired.

⁵¹² We have equality on the nose because monad liftings are defined with equality on the nose. One can probably relax the definition of monad lifting to be up to isomorphisms without breaking this correspondence.

⁵¹³ i.e. they are *saturated*, you cannot add more quantitative equations without changing the algebras

⁵¹⁴ c.f. a similar result proven in [ADV22, Theorem 49] in the case of $\mathbf{GMet} = \mathbf{Poset}$.

⁵¹⁵ I will not write the proofs because I am not confident enough with the literature on accessible and presentable categories, but I believe [FMS21, Propositions 3.8 and 3.9] make it possible to adapt the arguments of Remark 1.61 replacing \aleph_0 with a different cardinal (we implicitly used \aleph_0 because $\lambda < \aleph_0 \Leftrightarrow \lambda$ finite).

⁵¹⁶ The distance $\widehat{d_X}$ was defined in (3.56).

⁵¹⁷ We are concise in the quantifications for \hat{E}_2 .

⁵¹⁸ There are two ways to understand this. Semantically, the equations that are satisfied by all algebras in $\mathbf{QAlg}(\Sigma, \hat{E}_1)$ are also satisfied by all algebras in $\mathbf{QAlg}(\Sigma, \hat{E}_1 \cup \hat{E}_2)$ because the second category is contained in the first. Syntactically, if you have more axioms, you can prove more things.

⁵¹⁹ Recall that the context of $x \vdash x =_{\perp} x$, after unrolling the syntactic sugar, is the L-space with x at distance \top from itself, so we only need to prove $\sigma(x)$ is also at distance \top from itself (we do it with Top).

- Otherwise, $\varepsilon = \top$, so $\mathbf{X} \vdash s =_\varepsilon t$ belongs to $\Omega\mathfrak{Th}(\hat{E}_1)$ by Lemma 3.34.

We have shown that $\hat{E}_2 \subseteq \Omega\mathfrak{Th}(\hat{E}_1)$, and it follows that $\Omega\mathfrak{Th}(\hat{E}_1 \cup \hat{E}_2) \subseteq \Omega\mathfrak{Th}(\hat{E}_1)$.⁵²⁰

In conclusion, we found that $\hat{\mathcal{P}}$ is presented by the equations in \hat{E}_1 which we rewrite below:

$$x \vdash x = x \oplus x \quad x, y \vdash x \oplus y = y \oplus x \quad x, y, z \vdash x \oplus (y \oplus z) = (x \oplus y) \oplus z.$$

Remark 3.101. Compared to the presentation of $\mathcal{P}_{\text{ne}}^\uparrow$, we simply removed (3.6). These quantitative equations are included in the theory by default in the framework of [MPP16] because they only consider quantitative algebras with interpretations of operations that are nonexpansive with respect to the product metric (see Example 3.12). It is then natural to ask whether the monad lifting $\hat{\mathcal{P}}$ we defined can be presented by a quantitative algebraic theory in the sense of [MPP16]. The answer is negative because of a property that all monads presented by such theories share: they are enriched over $(\mathbf{Met}, \otimes, \mathbf{1})$ ⁵²¹

The monad $\hat{\mathcal{P}}$ is not enriched because it does not satisfy (see [ADV23b, Example 7.(1)])

$$\forall f, g : (X, d) \rightarrow (Y, \Delta), \sup_{x \in X} \Delta(f(x), g(x)) \geq \sup_{S \in \mathcal{P}X} \hat{\Delta}(f(S), g(S)).$$

Let f be the identity function on $[0, \frac{1}{2}]$ and g be the squaring function, then the left hand side is at most $\frac{1}{2}$ (Δ is bounded by $\frac{1}{2}$), and the right hand side is 1 as witnessed by $S = \{0, \frac{1}{2}\}$: $f(S) = S$ and $g(S) = \{0, \frac{1}{4}\}$, so $\hat{\Delta}(f(S), g(S)) = 1$.

This enrichment property is also shared by the free algebra monads of [FMS21], as they prove in Corollary 4.14, so in this direction, our framework is more general than theirs.

In a sense, $\hat{\mathcal{P}}$ can be seen as a *trivial* monad lifting of \mathcal{P}_{ne} because we simply viewed the equations presenting \mathcal{P}_{ne} as quantitative equations as we did in (3.49), and we added nothing else. After this example, you may want to conjecture that whenever \hat{E} is constructed from E like that, then \hat{E} presents a monad lifting of the $\mathcal{T}_{\Sigma, E}$, or equivalently thanks to Corollary 3.96 and Theorem 3.98, \hat{E} is an extension of E . That is not true. We showed in [MSV21, Theorem 44] that \hat{E} can sometimes prove more equations than E . This implies $U\hat{\mathcal{T}}_{\Sigma, \hat{E}}\mathbf{X} \neq \mathcal{T}_{\Sigma, E}X$, so $\hat{\mathcal{T}}_{\Sigma, \hat{E}}$ is not a monad lifting of $\mathcal{T}_{\Sigma, E}$.

We end this chapter with a final example, the one that motivated a lot of ideas in this manuscript.

Example 3.102 (LK). The LK distance on probability distributions defined in (3.3) defines a mere lifting $(X, d) \mapsto (\mathcal{D}X, d_{\text{LK}})$ of \mathcal{D} to $\mathbf{GMet} = [0, 1]\mathbf{Spa}$.⁵²² We show this is a monad lifting of (\mathcal{D}, η, μ) (as defined in Example 1.65) with Lemmas 3.103–3.105.

Lemma 3.103. *If $f : (X, d) \rightarrow (Y, \Delta)$ is nonexpansive, then so is $\mathcal{D}f : (\mathcal{D}X, d_{\text{LK}}) \rightarrow (\mathcal{D}Y, \Delta_{\text{LK}})$.*

Proof. Let $\varphi, \psi \in \mathcal{D}X$, we have

$$d_{\text{LK}}(\mathcal{D}f(\varphi), \mathcal{D}f(\psi))$$

⁵²⁰ Again, there are two different ways to understand this. Semantically, if all algebras in $\mathbf{QAlg}(\Sigma, \hat{E}_1)$ satisfy \hat{E}_2 , then $\mathbf{QAlg}(\Sigma, \hat{E}_1)$ and $\mathbf{QAlg}(\Sigma, \hat{E}_1 \cup \hat{E}_2)$ are the same categories. Syntactically, in any derivation with axioms $\hat{E}_1 \cup \hat{E}_2$, you can replace each axiom in \hat{E}_2 by a derivation using only axioms in \hat{E}_1 .

⁵²¹ See [ADV23a, after Corollary 4.19].

⁵²² Of course, you can take $[0, \infty]\mathbf{Spa}$ as well. You can also show that this mere lifting preserves the satisfaction of all the equations defining metric spaces except reflexivity ($x \vdash x =_0 x$). Indeed, we have $d_{\text{LK}}(\varphi, \varphi) = 0$ if and only if $d(x, y) = 0$ for all $x, y \in \text{supp}(\varphi)$ (if d is reflexive, this forces $\varphi = \delta_x$). This means you can take \mathbf{GMet} to be the category of diffuse metric spaces as we did in [MSV22, §5.3].

$$\begin{aligned}
&= \sum_{(y,y')} \mathcal{D}f(\varphi)(y)\mathcal{D}f(\psi)(y')\Delta(y,y') \\
&= \sum_{(y,y')} \left(\sum_{x \in f^{-1}(y)} \varphi(x) \right) \left(\sum_{x' \in f^{-1}(y')} \psi(x') \right) \Delta(y,y') \quad \text{definition of } \mathcal{D}f \text{ (1.45)} \\
&= \sum_{(y,y')} \sum_{x \in f^{-1}(y)} \sum_{x' \in f^{-1}(y')} \varphi(x)\psi(x')\Delta(y,y') \\
&= \sum_{(x,x')} \varphi(x)\psi(x')\Delta(f(x),f(x')) \\
&\leq \sum_{(x,x')} \varphi(x)\psi(x')d(f(x),f(x')) \quad f \text{ is nonexpansive} \\
&= d_{\text{LK}}(\varphi,\psi). \quad \text{definition of } d_{\text{LK}} \quad \square
\end{aligned}$$

Lemma 3.104. For any (X, d) , the map $\eta_X : (X, d) \rightarrow (\mathcal{D}X, d_{\text{LK}})$ is nonexpansive.

Proof. For any $a, a' \in X$, we have⁵²³

⁵²³ Notice that η_X is even an isometric embedding.

$$d_{\text{LK}}(\delta_a, \delta_{a'}) \stackrel{(3.3)}{=} \sum_{(x,x')} \delta_a(x)\delta_{a'}(x')d(x,x') = \delta_a(a)\delta_{a'}(a')d(a,a') = d(a,a'). \quad \square$$

Lemma 3.105. For any (X, d) , the map $\mu_X : (\mathcal{D}\mathcal{D}X, d_{\text{LK}\text{LK}}) \rightarrow (\mathcal{D}X, d_{\text{LK}})$ is nonexpansive.

Proof. For any $\Phi, \Psi \in \mathcal{D}\mathcal{D}X$, we have

$$\begin{aligned}
d_{\text{LK}}(\mu_X\Phi, \mu_X\Psi) &\stackrel{(3.3)}{=} \sum_{(x,x')} \mu_X\Phi(x)\mu_X\Psi(x')d(x,x') \\
&\stackrel{(1.46)}{=} \sum_{(x,x')} \left(\sum_{\varphi \in \text{supp}(\Phi)} \Phi(\varphi)\varphi(x) \right) \left(\sum_{\psi \in \text{supp}(\Psi)} \Psi(\psi)\psi(x') \right) d(x,x') \\
&= \sum_{(x,x')} \sum_{(\varphi,\psi)} \Phi(\varphi)\varphi(x)\Psi(\psi)\psi(x')d(x,x') \\
&= \sum_{(\varphi,\psi)} \Phi(\varphi)\Psi(\psi) \left(\sum_{(x,x')} \varphi(x)\psi(x')d(x,x') \right) \\
&\stackrel{(3.3)}{=} \sum_{\varphi,\psi} \Phi(\varphi)\Psi(\psi)d_{\text{LK}}(\varphi,\psi) \\
&\stackrel{(3.3)}{=} d_{\text{LK}\text{LK}}(\Phi,\Psi) \quad \square
\end{aligned}$$

Let us denote this monad lifting by \mathcal{D}_{LK} . In [MSV22, §5.3], we gave a relatively simple quantitative algebraic presentation for \mathcal{D}_{LK} , but Theorem 3.98 will help us find a simpler one. Since, by Example 1.79, the theory of convex algebras generated by $(\Sigma_{\text{CA}}, E_{\text{CA}})$ presents \mathcal{D} (via a monad isomorphism that we write ρ), the theorem gives us a theory presenting \mathcal{D}_{LK} generated by $\hat{E}_1 \cup \hat{E}_2$ where

$$\hat{E}_1 = \{ \mathbf{X}_{\top} \vdash s = t \mid X \vdash s = t \in E_{\text{CA}} \} \text{ and}$$

$$\hat{E}_2 = \left\{ (X, d) \vdash s =_\varepsilon t \mid \varepsilon = d_{\text{LK}} \left(\rho_X[s]_{E_{\text{CA}}}, \rho_X[t]_{E_{\text{CA}}} \right) \right\}.$$

In order to simplify \hat{E}_2 , we rely on two property that d_{LK} has (one symmetric to the other) : for any $\varphi, \varphi', \psi \in \mathcal{D}X$ and $p \in [0, 1]$,

$$d_{\text{LK}}(p\varphi + \bar{p}\varphi', \psi) = pd_{\text{LK}}(\varphi, \psi) + \bar{p}d_{\text{LK}}(\varphi', \psi) \text{ and} \quad (3.62)$$

$$d_{\text{LK}}(\varphi, p\varphi + \bar{p}\varphi') = pd_{\text{LK}}(\varphi, \psi) + \bar{p}d_{\text{LK}}(\varphi, \psi'). \quad (3.63)$$

Intuitively, this means that we can compute the distance between s and t by decomposing the terms into their variables, computing simple distances, then combining them to get back to s and t .⁵²⁴ Formally, we only need to keep the quantitative equations in \hat{E}_2 that belong to⁵²⁵

$$\begin{aligned} \hat{E}'_2 = & \{ x =_{\varepsilon_1} y, x =_{\varepsilon_2} z \vdash x =_{p\varepsilon_1 + \bar{p}\varepsilon_2} y +_p z \mid \varepsilon_1, \varepsilon_2 \in [0, 1], p \in (0, 1) \} \\ & \cup \{ y =_{\varepsilon_1} x, z =_{\varepsilon_2} x \vdash y +_p z =_{p\varepsilon_1 + \bar{p}\varepsilon_2} x \mid \varepsilon_1, \varepsilon_2 \in [0, 1], p \in (0, 1) \}. \end{aligned}$$

We will prove that for any $\hat{A} \in \mathbf{QAlg}(\Sigma_{\text{CA}})$, $\hat{A} \models \hat{E}_1 \cup \hat{E}'_2$ implies $\hat{A} \models \hat{E}_1 \cup \hat{E}_2$.⁵²⁶ Suppose $\hat{A} \models \hat{E}_1 \cup \hat{E}'_2$, we proceed by induction on the structure of s and t to show that \hat{A} satisfies $(X, d) \vdash s =_\varepsilon t$, where $\varepsilon = d_{\text{LK}} \left(\rho_X[s]_{E_{\text{CA}}}, \rho_X[t]_{E_{\text{CA}}} \right)$.

If s and t are variables, then $\rho_X[s]_{E_{\text{CA}}} = \delta_x$ and $\rho_X[t]_{E_{\text{CA}}} = \delta_y$ for some $x, y \in X$, thus $\varepsilon = d(x, y)$ and $(X, d) \vdash x =_{d(x, y)} y$ is satisfied by \hat{A} (by 3.35).

Otherwise, without loss of generality,⁵²⁷ we write $t = t_1 +_p t_2$, and let $\varepsilon_i = d_{\text{LK}} \left(\rho_X[s]_{E_{\text{CA}}}, \rho_X[t_i] \right)$. By the induction hypothesis, $\hat{A} \models (X, d) \vdash s =_{\varepsilon_i} t_i$ for $i = 1, 2$. Then, we define a substitution map $\sigma : \{x, y, z\} \rightarrow \mathcal{T}_\Sigma X$ with $x \mapsto s$, $y \mapsto t_1$ and $z \mapsto t_2$, and since \hat{A} satisfies $x =_{\varepsilon_1} y, x =_{\varepsilon_2} z \vdash x =_{p\varepsilon_1 + \bar{p}\varepsilon_2} y +_p z \in \hat{E}'_2$, we can apply Lemma 3.42 to conclude \hat{A} satisfies $(X, d) \vdash s =_{\varepsilon'} t$ with

$$\begin{aligned} \varepsilon' &= pd_{\text{LK}} \left(\rho_X[s]_{E_{\text{CA}}}, \rho_X[t_1] \right) + \bar{p}d_{\text{LK}} \left(\rho_X[s]_{E_{\text{CA}}}, \rho_X[t_2] \right) \\ &= d_{\text{LK}} \left(\rho_X[s]_{E_{\text{CA}}}, p\rho_X[t_1] + \bar{p}\rho_X[t_2] \right) && \text{by (3.62)} \\ &= d_{\text{LK}} \left(\rho_X[s]_{E_{\text{CA}}}, \rho_X[t_1 +_p t_2] \right) \\ &= d_{\text{LK}} \left(\rho_X[s]_{E_{\text{CA}}}, \rho_X[t]_{E_{\text{CA}}} \right) = \varepsilon. \end{aligned}$$

We conclude that $\hat{E}_1 \cup \hat{E}'_2$ presents \mathcal{D}_{LK} .

⁵²⁴ This is very similar to what happens for the Kantorovich distance and (3.10).

⁵²⁵ If you have symmetry ($x =_\varepsilon y \vdash y =_\varepsilon x$) as an axiom in \mathbf{GMet} already, you can keep only one of these sets.

⁵²⁶ It follows that $\Omega\mathfrak{Th}(\hat{E}_1 \cup \hat{E}'_2) = \Omega\mathfrak{Th}(\hat{E}_1 \cup \hat{E}_2)$ because we already have the \supseteq inclusion as explained in Footnote 520.

⁵²⁷ If s is a term of depth > 0 but t is a variable, you decompose s instead, and then you have to use a symmetric argument.

4 Conclusion

In [MPP16], the authors introduced a theoretical framework to reason algebraically about distances inside a metric space. We have made adjustments to their proposal with two main goals in mind:

1. replace metrics with a more general notion of distance, and
2. tighten the relationship with classical universal algebra.

The result is a theory of quantitative algebras which are algebras $(A, \llbracket - \rrbracket_A)$ paired with a distance function $d : A \times A \rightarrow L$ valued in a complete lattice, and no hardcoded constraint on the interaction between $\llbracket - \rrbracket_A$ and d , in contrast with the nonexpansiveness requirement (0.1) of [MPP16].⁵²⁸

We introduced a sound and complete deduction system (Figure 3.1) generalizing Birkhoff's equational logic. The judgments are quantitative equations, a closer analog to classical equations than the judgments of [MPP16].

We gave a construction for free quantitative (Σ, \hat{E}) -algebras (Theorem 3.57) relative to any class \hat{E} of quantitative equations, following that of free classical algebras (Proposition 1.49) almost to the T. This yielded a monad $\widehat{\mathcal{T}}_{\Sigma, \hat{E}}$ on the category of generalized metric spaces **GMet**.

We showed that algebras for the monad $\widehat{\mathcal{T}}_{\Sigma, \hat{E}}$ coincide with the (Σ, \hat{E}) -algebras (Theorem 3.80), justifying a search for quantitative algebraic presentations for monads on **GMet**, of which we gave several examples (Examples 3.83, 3.85, 3.86, 3.100, and 3.102).

Finally, we gave a sufficient condition for a distance on Σ -terms to be axiomatized with a quantitative algebraic theory (Theorem 3.98). More precisely, if M is a monad on **Set** with an algebraic presentation (Σ, E) , and \widehat{M} is a monad lifting of M to **GMet**, then we constructed a quantitative algebraic theory \hat{E} that extends E and gives a presentation for \widehat{M} .

4.1 Future Work

We mention some lines of questioning that need further investigation.

⁵²⁸ It is still possible to enforce (0.1) and variants with supplementary axioms (see (3.8) and (3.9)).

Examples

In the original paper on quantitative algebras [MPP16], the authors gave theories axiomatizing the Hausdorff distance (Example 3.83) and the Kantorovich distance (Example 1.79). I think these are amazing examples to showcase the potential of quantitative algebraic reasoning, and I would like to find more. Several papers like [BMPP18, MV20, BMPP21, MSV21, MSV22, ?] contain additional examples, and most of them follow the leitmotif discussed in §3.5, namely, they are built on top of a classical algebraic theory. I believe that Theorem 3.98 will accelerate the process of developing similar examples, but some efforts are still needed.⁵²⁹

Examples in [MV20, MSV21] are of particular interest to me because they deal with combining quantitative algebraic theories and their corresponding monads. For instance, the main ingredients in [MV20] are

- The algebraic theories E_S and E_{CA} presenting the monads \mathcal{P}_{ne} and \mathcal{D} respectively.
- An equation ϕ such that $E_S \cup E_{CA} \cup \{\phi\}$ presents the monad \mathcal{C} of convex sets of distributions [MV20, Definition 5].⁵³⁰
- The extension of E_S presenting the Hausdorff monad lifting \mathcal{P}_{ne}^\dagger on **Met**.
- The extension of E_{CA} presenting the Kantorovich monad lifting \mathcal{D}_K on **Met**.

Then, they show that the union of these extensions with ϕ seen as a quantitative equation (recall Example 3.71) presents a monad lifting of \mathcal{C} on **Met**. It would be interesting to make this result more abstract and work with any theories, extensions, and monads, but we found a counterexample that breaks the general pattern in [MSV21, Theorem 44], so some work is required to identify abstractly why it breaks.

Quantitative Diagrammatic Reasoning

Diagrammatic reasoning is another generalization of algebraic reasoning that has been popular in recent years. Using string diagrams in particular, people have axiomatized languages for quantum processes [CK17], stochastic processes [Fri20], machine learning models [CGG⁺22], satisfaction of Boolean formulas [GPZ23], finite state automata [PZ23], and more. There is a gap in the literature on the combination of quantitative and diagrammatic reasoning. I am aware of only one paper [KTW17] going in this direction.

HSP Theorems

We mentioned in the introduction that Birkhoff's HSP theorem [Bir35] is a celebrated result in universal algebra. In [MPP17], the authors proved a variant of this theorem for the quantitative algebras in the original paper [MPP16]. The question of how to adapt their methods to our new framework is still open.⁵³¹ We can also mention other variants of the HSP theorem in similar settings that are proven (with concrete methods) in [Wea95, BV05, Hin16, Hin17, Ros24].

⁵²⁹ I planned to include a chapter in this thesis with detailed examples and non-examples to help others in this search, but I ran out of time.

⁵³⁰ The monad \mathcal{C} combines \mathcal{P}_{ne} and \mathcal{D} in a sense made precise in [GP20].

⁵³¹ After some unsuccessful attempts during my PhD.

In the process of abstracting universal algebra away from the category of sets, several abstract HSP theorems were proven (see, e.g. [BH76, Man76, Bar94, Bar02, ARV11, MU19]). In [MU19], Milius and Urbat prove one such result and apply it to the quantitative algebras of [MPP16]. They obtain a generalization of Mardare et al.'s result from [MPP17]. Rosický proves a similar result in [Ros24] using abstract results from [Man76]. In [JMU24], the authors apply Milius and Urbat's result to a new class of algebras that are a mix between [FMS21]'s and [MSV22]'s, and it should apply to the quantitative algebras presented in this thesis,⁵³² but careful checks are needed (see Footnote 372 and Remark 3.24).

There are other theoretical results that followed Mardare et al.'s introduction of quantitative algebras which could be generalized to the present work. I am most interested in their work on combining theories and monads [BMPP18, BMPP21], and in the characterization of monads which can be presented by a quantitative algebraic theory [AFMS21, FMS21, Adá22, ADV23b].

Partial Operations

In classical universal algebra, a signature Σ is a set of operation symbols each equipped with an arity in \mathbb{N} . Then, the interpretation of an n -ary operation is a function $\llbracket \text{op} \rrbracket_A : A^n \rightarrow A$, where A^n is the n -wise cartesian product. We can also see A^n as an exponential, namely, the set of functions from $\{1, \dots, n\}$ to A , and this point of view is often used when generalizing algebraic reasoning.

For instance, in [FMS21], the arity of an operation is allowed to be an arbitrary generalized metric space on $[n] = \{1, \dots, n\}$.⁵³³ Then, the interpretation of a $([n], d)$ -ary operation symbol is a nonexpansive map $\llbracket \text{op} \rrbracket_A : \mathbf{A}^{([n], d)} \rightarrow \mathbf{A}$. The definition of $\mathbf{A}^{([n], d)}$ is out of scope (it is not an exponential in the sense of cartesian closed categories), but it is a generalized metric on the set of nonexpansive maps $([n], d) \rightarrow \mathbf{A}$. This has two notable consequences.

1. The carrier of $\mathbf{A}^{([n], d)}$ does not necessarily contain all the functions from $[n]$ to A , so $\llbracket \text{op} \rrbracket_A$ may not be applicable to all n -tuples of elements in A . Hence, we can see it as a partial function $A^n \rightarrow A$. Not all partial functions of this type arise as nonexpansive maps from $\mathbf{A}^{([n], d)}$ even if we let d vary. In particular, the partiality of operations depends on the distance of the carrier \mathbf{A} .⁵³⁴
2. When d is the discrete generalized metric on $[n]$ (recall Example 3.59), the carrier of $\mathbf{A}^{([n], d)}$ is all of A^n , and the nonexpansiveness of $\llbracket \text{op} \rrbracket_A$ translates to the original requirement (0.1) of [MPP16].⁵³⁵

It is not known how to keep the flexibility of Item 1 to deal with *partial* operations without the constraint of Item 2. Namely, $\llbracket \text{op} \rrbracket_A$ should be a function from the carrier of $\mathbf{A}^{([n], d)}$ to the carrier of \mathbf{A} that is not necessarily nonexpansive. This would combine the generality of both [FMS21]'s and our algebras.

⁵³² They consider arbitrary relational structures like in [FMS21], but the arities are restricted to be natural numbers only, so operations are not partial. They do not require operations to be nonexpansive in the sense of (0.1), but they achieve this with lifted signatures like in [MSV22].

⁵³³ We are simplifying to keep things light and closer to our work. They actually allow infinite arities and arbitrary relational structures.

⁵³⁴ e.g. if $d_{\mathbf{A}}(a, a') = \perp$ for all $a, a' \in A$, then the carrier of $\mathbf{A}^{([n], d)}$ is always A^n .

⁵³⁵ Briefly, it is because the distance between two functions $f, g : ([n], d) \rightarrow \mathbf{A}$ is

$$d(f, g) = \sup_{i \in [n]} d_{\mathbf{A}}(f(i), g(i)),$$

which is the coordinatewise maximum distance when viewing f and g as tuples.

Functorial Semantics

Our restriction to discrete arities is also an obstacle in the search for a functorial semantics of quantitative algebras. In his thesis [Law63], Lawvere gave a novel account of universal algebra based on functors. Given a signature Σ and equations E , a syntactic category $\mathcal{S}_{\Sigma,E}$ is constructed such that functors $M : \mathcal{S}_{\Sigma,E} \rightarrow \mathbf{Set}$ that preserve products correspond to (Σ, E) -algebras, and natural transformations between such functors correspond to homomorphisms. This led to the first proof of the correspondence (equivalence of categories) between finitary monads and varieties in [Lin66].

People also investigated what happens when \mathbf{Set} is replaced with another category, and what happens when using enriched categories. Plenty of so-called *monad-theory correspondences* were worked out under this lens throughout many papers including [Dub70, Gra75, BD80, Bur81, KP93, GP98, Pow99, Rob02, HP06, NP09, LP09, LR11, LW16, GP18, BG19, LP23, Ros24].

Since \mathbf{GMet} has finite products, the first naive attempt to define algebras over generalized metric spaces could be to simply replace \mathbf{Set} with \mathbf{GMet} in Lawvere's account. The category of finite product-preserving functors $\mathcal{S}_{\Sigma,E} \rightarrow \mathbf{Met}$ coincides with the category of quantitative algebras defined in [MPP16]⁵³⁶ that satisfy the equations in E translated to quantitative equations with the discrete context (see Example 3.71). However, there is no obvious way to construct a syntactic category starting from quantitative equations.

Since \mathbf{Met} is symmetric monoidal closed [ADV23b, Example 3.(2)] and locally \aleph_1 -presentable [LR17, Example 4.5.(3)], the theory of enriched Lawvere theories in [Pow99] applies,⁵³⁷ and it allows handling quantitative equations. However, [Pow99] allows arities to be non-discrete spaces, so any quantitative algebraic theory (again in the sense of [MPP16]) yields an enriched Lawvere theory, but not vice-versa.

In [HP06], the authors study discrete Lawvere theories, which are essentially enriched Lawvere theories where arities are required to be discrete. Unfortunately, this is too restricted. Namely, a discrete Lawvere theory can be translated into a quantitative algebraic theory of [MPP16] (with nonexpansive operations), but not vice-versa. In a nutshell, the problem is that while terms in $\mathcal{T}_{\mathbb{E}}X$ are defined independently of the distance on X (i.e. arities are discrete), the distance and even the equality between terms ($d_{\mathbb{E}}$ and $\equiv_{\mathbb{E}}$) can vary when different distances on X are considered. Recently, Rosický proposed a solution for this in [Ros24], but there remains a small gap with Mardare et al.'s algebras because operations in [Ros24] can have countably infinite (discrete) arities.

There remains the problem of allowing operations that are not necessarily non-expansive as we do in Definition 3.1. In [AFMS21, §5], they say that models of Nishizawa and Power's \mathbf{Poset} -Lawvere theories for \mathbf{Set} [NP09] are algebras where operations can be interpreted as arbitrary functions. Unfortunately, their arities are not necessarily discrete so their monad-theory correspondence [AFMS21, Theorem 5.9] (proven concretely in [AFMS21, Corollary 4.5]) is not enough.⁵³⁸

⁵³⁶ i.e. operations are interpreted as nonexpansive maps from the product of spaces:

$$[\text{op}]_A : \mathbf{A}^n \rightarrow \mathbf{A}.$$

⁵³⁷ One also needs that \mathbf{Met} is locally \aleph_1 -presentable as a symmetric monoidal closed category. It would be interesting to determine when a category \mathbf{GMet} has these properties.

⁵³⁸ Even before considering generalizing from \mathbf{Poset} to \mathbf{GMet} which is difficult because of the question of local presentability.

Applications

Many ad-hoc methods for combining algebraic reasoning with various structures like metrics or orders to reason about program semantics already exist in the literature.⁵³⁹ Our abstract framework could allow viewing several of these examples under the same lens, and facilitate the discovery of new similar methods.

With applications in mind, we can mention term rewriting systems [BKdV03] which are a popular approach to compute *in an actual computer* with classical equations. Gavazzo and Di Florio gave a very elegant account of quantitative rewriting systems in [GD23]. It seems our approaches are complementary because they replaced $[0, \infty]$ with an arbitrary quantale (a kind of complete lattice), and they also rework the nonexpansiveness assumption (0.1) in [GD23, §6].

⁵³⁹ See e.g. [CPV16, BBKK18, BBLM18a, BBLM18b, DLHLP22, Sch22a, Sch22b].

Bibliography

- [ABH⁺12] Jiří Adámek, Filippo Bonchi, Mathias Hülsbusch, Barbara König, Stefan Milius, and Alexandra Silva. A coalgebraic perspective on minimization and determinization. In Lars Birkedal, editor, *Foundations of Software Science and Computational Structures*, page 58–73, Berlin, Heidelberg, 2012. Springer Berlin Heidelberg.
- [ACT10] Andrei Akhvediani, Maria Manuel Clementino, and Walter Tholen. On the categorical meaning of hausdorff and gromov distances, i. *Topology and its Applications*, 157(8):1275–1295, 2010. Advances in Set-Theoretic Topology: Proceedings of the Conference in Honour of Professor Tsugunori Nogura on his Sixtieth Birthday (9–19 June 2008, Erice, Sicily, Italy).
- [Adá22] Jiří Adámek. Varieties of quantitative algebras and their monads. In *Proceedings of the 37th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS '22*, New York, NY, USA, 2022. Association for Computing Machinery.
- [ADV22] Jiří Adámek, Matej Dostál, and Jiří Velebil. A categorical view of varieties of ordered algebras. *Math. Struct. Comput. Sci.*, 32(4):349–373, 2022.
- [ADV23a] Jiří Adámek, Matej Dostál, and Jiří Velebil. Quantitative algebras and a classification of metric monads, 2023. <https://arxiv.org/abs/2210.01565v4>.
- [ADV23b] Jiří Adámek, Matej Dostál, and Jiří Velebil. Strongly finitary monads for varieties of quantitative algebras. In Paolo Baldan and Valeria de Paiva, editors, *10th Conference on Algebra and Coalgebra in Computer Science, CALCO 2023, June 19–21, 2023, Indiana University Bloomington, IN, USA*, volume 270 of *LIPICs*, pages 10:1–10:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2023.
- [AFMS21] Jiří Adámek, Chase Ford, Stefan Milius, and Lutz Schröder. Finitary monads on the category of posets. *Math. Struct. Comput. Sci.*, 31(7):799–821, 2021.
- [AHS06] Jiří Adámek, Horst Herrlich, and George E Strecker. Abstract and concrete categories; the joy of cats 2004. *Reprints in Theory and Applications of Categories*, 17:1–507, 2006. Originally published by: John Wiley and Sons, New York, 1990.
- [AMMU15] Jiří Adámek, Stefan Milius, Lawrence S. Moss, and Henning Urbat. On finitary functors and their presentations. *J. Comput. Syst. Sci.*, 81(5):813–833, 2015.
- [ANR85] Jiří Adámek, Evelyn Nelson, and Jan Reiterman. The birkhoff variety theorem for continuous algebras. *Algebra Universalis*, 20(3):328–350, 10 1985.
- [ARV11] J. Adámek, J. Rosický, and E. M. Vitale. *Algebraic theories. A categorical introduction to general algebra. With a foreword by F. W. Lawvere*, volume 184 of *Camb. Tracts Math.* Cambridge: Cambridge University Press, 2011.
- [AW20] Alessandro Aldini and Herbert Wiklicky, editors. *Proceedings 16th Workshop on Quantitative Aspects of Programming Languages and Systems, QAPL@ETAPS 2019, Prague, Czech Republic, 7th April 2019*, volume 312 of *EPTCS*, 2020.
- [Awo10] Steve Awodey. *Category Theory*. Oxford University Press, 2010.
- [Bar92] Michael Barr. Hsp type theorems in the category of posets. In Stephen Brookes, Michael Main, Austin Melton, Michael Mislove, and David Schmidt, editors, *Mathematical Foundations of Programming Semantics*, pages 221–234, Berlin, Heidelberg, 1992. Springer Berlin Heidelberg.

- [Bar94] Michael Barr. Functorial semantics and HSP type theorems. *Algebra Univers.*, 31(2):223–251, 1994.
- [Bar02] Michael Barr. HSP subcategories of Eilenberg-Moore algebras. *Theory Appl. Categ.*, 10:461–468, 2002.
- [Bar06] Michael Barr. Relational algebras. In *Reports of the Midwest Category Seminar IV*, page 39–55. Springer, 2006.
- [Bau19] Andrej Bauer. What is algebraic about algebraic effects and handlers?, 2019.
- [BB92] E. Burroni and A. Burroni. Structures algébriques : thème et variations. *Cahiers de Topologie et Géométrie Différentielle Catégoriques*, 33(3):207–216, 1992.
- [BBKK18] Paolo Baldan, Filippo Bonchi, Henning Kerstan, and Barbara König. Coalgebraic behavioral metrics. *Log. Methods Comput. Sci.*, 14(3), 2018.
- [BBLM18a] Giorgio Bacci, Giovanni Bacci, Kim G. Larsen, and Radu Mardare. A Complete Quantitative Deduction System for the Bisimilarity Distance on Markov Chains. *Logical Methods in Computer Science*, Volume 14, Issue 4, November 2018.
- [BBLM18b] Giorgio Bacci, Giovanni Bacci, Kim G. Larsen, and Radu Mardare. Complete axiomatization for the total variation distance of markov chains. *Electronic Notes in Theoretical Computer Science*, 336:27–39, 2018. The Thirty-third Conference on the Mathematical Foundations of Programming Semantics (MFPS XXXIII).
- [BBvT12] Branka Budimirović, Vjekoslav Budimirović, Branimir Šešelja, and Andreja Tepavčević. Fuzzy equational classes. In *2012 IEEE International Conference on Fuzzy Systems*, pages 1–6, 2012.
- [BD80] Francis Borceux and Brian Day. Universal algebra in a closed category. *Journal of Pure and Applied Algebra*, 16(2):133–147, 1980.
- [BE99] Nuno Barreiro and Thomas Ehrhard. Quantitative semantics revisited. In Jean-Yves Girard, editor, *Typed Lambda Calculi and Applications*, pages 40–53, Berlin, Heidelberg, 1999. Springer Berlin Heidelberg.
- [Bec69] Jon Beck. Distributive laws. In B. Eckmann, editor, *Seminar on Triples and Categorical Homology Theory*, page 119–140, Berlin, Heidelberg, 1969. Springer Berlin Heidelberg.
- [Ben68] Jean Benabou. Structures algébriques dans les catégories. *Cahiers de topologie et géométrie différentielle*, 10(1):1–126, 1968.
- [BG19] John Bourke and Richard Garner. Monads and theories. *Advances in Mathematics*, 351:1024–1071, 2019.
- [BH76] B. Banaschewski and Horst Herrlich. Subcategories defined by implications. *Houston J. Math.*, 2:149–171, 1976.
- [BHKR15] Marcello M. Bonsangue, Helle Hvid Hansen, Alexander Kurz, and Jurriaan Rot. Presenting Distributive Laws. *Logical Methods in Computer Science*, Volume 11, Issue 3, August 2015.
- [Bir33] Garrett Birkhoff. On the combination of subalgebras. *Mathematical Proceedings of the Cambridge Philosophical Society*, 29(4):441–464, 1933.
- [Bir35] Garrett Birkhoff. On the structure of abstract algebras. *Mathematical Proceedings of the Cambridge Philosophical Society*, 31(4):433–454, 1935.
- [BKdV03] Marc Bezem, Jan Willem Klop, and Roel de Vrijer, editors. *Terese. Term rewriting systems*, volume 55 of *Camb. Tracts Theor. Comput. Sci.* Cambridge: Cambridge University Press, 2003.
- [Blo76] Stephen L. Bloom. Varieties of ordered algebras. *Journal of Computer and System Sciences*, 13(2):200–212, 1976.
- [BMPP18] Giorgio Bacci, Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. An algebraic theory of markov processes. In Anuj Dawar and Erich Grädel, editors, *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2018, Oxford, UK, July 09-12, 2018*, page 679–688. ACM, 2018.
- [BMPP21] Giorgio Bacci, Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. Tensor of quantitative equational theories. In Fabio Gadducci and Alexandra Silva, editors, *9th Conference on Algebra and Coalgebra in Computer Science, CALCO 2021, August 31 to September 3, 2021, Salzburg, Austria*, volume 211 of *LIPICs*, page 7:1–7:17. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.

- [Bor94] Francis Borceux. *Handbook of Categorical Algebra*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 1994.
- [Bou92] Dominique Bourn. Low dimensional geometry of the notion of choice. In *Category Theory 1991, CMS Conference Proceedings*, volume 13, page 55–73, 1992.
- [BP98] Sergey Brin and Lawrence Page. The anatomy of a large-scale hypertextual web search engine. *Computer Networks and ISDN Systems*, 30(1):107–117, 1998. Proceedings of the Seventh International World Wide Web Conference.
- [BP15] Filippo Bonchi and Damien Pous. Hacking nondeterminism with induction and coinduction. *Commun. ACM*, 58(2):87–95, jan 2015.
- [Bra00] A. Branciari. A fixed point theorem of Banach-Caccioppoli type on a class of generalized metric spaces. *Publ. Math. Debr.*, 57(1-2):31–37, 2000.
- [BSS21] Filippo Bonchi, Alexandra Silva, and Ana Sokolova. Distribution bisimilarity via the power of convex algebras. *Logical Methods in Computer Science*, Volume 17, Issue 3, July 2021.
- [BSV19] Filippo Bonchi, Ana Sokolova, and Valeria Vignudelli. The theory of traces for systems with nondeterminism and probability. In *34th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2019, Vancouver, BC, Canada, June 24-27, 2019*, page 1–14. IEEE, 2019.
- [BSV22] Filippo Bonchi, Ana Sokolova, and Valeria Vignudelli. The theory of traces for systems with nondeterminism, probability, and termination. *Logical Methods in Computer Science*, Volume 18, Issue 2, June 2022.
- [Bur81] Albert Burroni. Algèbres graphiques (sur un concept de dimension dans les langages formels). *Cahiers de topologie et géométrie différentielle*, 22(3):249–265, 1981.
- [Bur82] Albert Burroni. Algèbres graphiques (suites : Les bidules). *Diagrammes*, 7:B1–B8, 1982.
- [BV05] Radim Bělohlávek and Vilém Vychodil. *Fuzzy equational logic*, volume 186 of *Stud. Fuzziness Soft Comput.* Berlin: Springer, 2005.
- [BvBR98] Marcello M. Bonsangue, Franck van Breugel, and Jan J. M. M. Rutten. Generalized metric spaces: Completion, topology, and powerdomains via the yoneda embedding. *Theor. Comput. Sci.*, 193(1-2):1–51, 1998.
- [BW05] Michael Barr and Charles Wells. Toposes, triples and theories. *Reprints in Theory and Applications of Categories*, 12:1–287, 2005. Originally published by: John Wiley and Sons, New York, 1990.
- [CDL15] Raphaëlle Crubille and Ugo Dal Lago. Metric reasoning about λ -terms: The affine case. In *Proceedings of the 2015 30th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*, LICS '15, page 633–644, USA, 2015. IEEE Computer Society.
- [CDL17] Raphaëlle Crubillé and Ugo Dal Lago. Metric reasoning about λ -terms: The general case. In Hongseok Yang, editor, *Programming Languages and Systems*, pages 341–367, Berlin, Heidelberg, 2017. Springer Berlin Heidelberg.
- [CGG⁺22] Geoffrey S. H. Cruttwell, Bruno Gavranović, Neil Ghani, Paul Wilson, and Fabio Zanasi. Categorical foundations of gradient-based learning. In Ilya Sergey, editor, *Programming Languages and Systems*, pages 1–28, Cham, 2022. Springer International Publishing.
- [CH06] Maria Manuel Clementino and Dirk Hofmann. Exponentiation in v -categories. *Topology and its Applications*, 153(16):3113–3128, 2006. Special Issue: Aspects of Contemporary Topology.
- [Che16] Eugenia Cheng. *How to Bake PI*. Basic Books, 5 2016.
- [CHN11] Petr Cintula, Petr Hájek, and Carles Noguera, editors. *Handbook of mathematical fuzzy logic. Volume 1*, volume 37 of *Stud. Log. (Lond.)*. London: College Publications, 2011.

- [CK17] Bob Coecke and Aleks Kissinger. *Picturing quantum processes. A first course in quantum theory and diagrammatic reasoning*. Cambridge: Cambridge University Press, 2017.
- [CKPR21] Pablo Samuel Castro, Tyler Kastner, Prakash Panangaden, and Mark Rowland. Mico: Improved representations via sampling-based state similarity for markov decision processes. *Advances in Neural Information Processing Systems*, 34:30113–30126, 2021.
- [CM22a] Davide Castelnovo and Marino Miculan. Fuzzy Algebraic Theories. In Florin Manea and Alex Simpson, editors, *30th EACSL Annual Conference on Computer Science Logic (CSL 2022)*, volume 216 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 13:1–13:17, Dagstuhl, Germany, 2022. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [CM22b] Alex Citkin and Alexei Muravitsky. *Consequence Relations: An Introduction to the Lindenbaum-Tarski Method*. Oxford University Press, 07 2022. Available at <https://arxiv.org/abs/2106.10966>.
- [CMS23] Christopher Conlon, Julie Holland Mortimer, and Paul Sarkis. Estimating preferences and substitution patterns from second-choice data alone. Working Paper, Accessed on: 2023-12-08, 2023.
- [Col24] Thomas Colcombet. knowledge. \LaTeX package, 2024. <https://ctan.org/pkg/knowledge>.
- [Con17] Gabriel Conant. Distance structures for generalized metric spaces. *Annals of Pure and Applied Logic*, 168(3):622–650, 2017.
- [CPV16] Konstantinos Chatzikokolakis, Catuscia Palamidessi, and Valeria Vignudelli. Up-To Techniques for Generalized Bisimulation Metrics. In *27th International Conference on Concurrency Theory (CONCUR 2016)*, volume 59 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 35:1–35:14, Québec City, Canada, August 2016.
- [Cur29] Haskell B. Curry. An analysis of logical substitution. *American Journal of Mathematics*, 51(3):363–384, 1929.
- [DFM23] Francesco Dagnino, Amin Farjudian, and Eugenio Moggi. Robustness in metric spaces over continuous quantales and the hausdorff-smyth monad. In Erika Ábrahám, Clemens Dubslaff, and Silvia Lizeth Tapia Tarifa, editors, *Theoretical Aspects of Computing – ICTAC 2023*, page 313–331, Cham, 2023. Springer Nature Switzerland.
- [DGY19] Ugo Dal Lago, Francesco Gavazzo, and Akira Yoshimizu. Differential logical relations, part i: The simply-typed case. In Christel Baier, Ioannis Chatzigiannakis, Paola Flocchini, and Stefano Leonardi, editors, *46th International Colloquium on Automata, Languages, and Programming (ICALP 2019)*, volume 132 of *Leibniz International Proceedings in Informatics (LIPIcs)*, page 111:1–111:14, Dagstuhl, Germany, 2019. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [Die80] Yves Diers. Categories multialgebriques. *Arch. Math.*, 34:193–209, 1980.
- [DLHLP22] Ugo Dal Lago, Furio Honsell, Marina Lenisa, and Paolo Pistone. On Quantitative Algebraic Higher-Order Theories. In Amy P. Felty, editor, *7th International Conference on Formal Structures for Computation and Deduction (FSCD 2022)*, volume 228 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 4:1–4:18, Dagstuhl, Germany, 2022. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [DLPS07] Christian Delhommé, Claude Laflamme, Maurice Pouzet, and Norbert Sauer. Divisibility of countable metric spaces. *European Journal of Combinatorics*, 28(6):1746–1769, 2007.
- [DP02] B. A. Davey and H. A. Priestley. *Introduction to Lattices and Order*. Cambridge University Press, 2 edition, 2002.
- [DPS18] Fredrik Dahlqvist, Louis Parlant, and Alexandra Silva. Layer by layer – combining monads. In Bernd Fischer and Tarmo Uustalu, editors, *Theoretical Aspects of Computing – ICTAC 2018*, page 153–172, Cham, 2018. Springer International Publishing.
- [Dub70] E. J. Dubuc. Enriched semantics-structure (meta) adjointness. *Revista de la Unión Matemática Argentina*, 25:5–26, 1970.
- [Dwo06] Cynthia Dwork. Differential privacy. In Michele Bugliesi, Bart Preneel, Vladimiro Sassone, and Ingo Wegener, editors, *Automata, Languages and Programming*, pages 1–12, Berlin, Heidelberg, 2006. Springer Berlin Heidelberg.

- [EE86] Paul Erdős and Marcel Erné. Clique numbers of graphs. *Discrete Mathematics*, 59(3):235–242, 1986.
- [EGP07] Marcel Erné, Mai Gehrke, and Aleš Pultr. Complete congruences on topologies and down-set lattices. *Applied Categorical Structures*, 15(1):163–184, Apr 2007.
- [EM45] Samuel Eilenberg and Saunders MacLane. General theory of natural equivalences. *Transactions of the American Mathematical Society*, 58(2):231–294, 1945.
- [EM65] Samuel Eilenberg and John C Moore. Adjoint functors and triples. *Illinois Journal of Mathematics*, 9(3):381–398, 1965.
- [Esc99] Martin Escardo. A metric model of pcf. In *Workshop on Realizability Semantics and Applications*, volume 417, page 418. Citeseer, 1999.
- [FGSW21] Uli Fahrenberg, Mai Gehrke, Luigi Santocane, and Michael Winter, editors. *Relational and Algebraic Methods in Computer Science*, Lecture Notes in Computer Science. Springer Cham, 2021.
- [FH11] Marcelo Fiore and Chung-Kil Hur. On the mathematical synthesis of equational logics. *Logical Methods in Computer Science*, Volume 7, Issue 3, September 2011.
- [Fla97] R. C. Flagg. Quantales and continuity spaces. *Algebra Universalis*, 37(3):257–276, Jun 1997.
- [FMS21] Chase Ford, Stefan Milius, and Lutz Schröder. Behavioural preorders via graded monads. In *Proceedings of the 36th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS '21*, New York, NY, USA, 2021. Association for Computing Machinery.
- [FMS21] Chase Ford, Stefan Milius, and Lutz Schröder. Monads on Categories of Relational Structures. In Fabio Gadducci and Alexandra Silva, editors, *9th Conference on Algebra and Coalgebra in Computer Science (CALCO 2021)*, volume 211 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 14:1–14:17, Dagstuhl, Germany, 2021. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [For22] Simon Forest. An extension of batanin’s approach to globular algebras, 2022.
- [FP19] Tobias Fritz and Paolo Perrone. A probability monad as the colimit of spaces of finite samples. *Theory and Applications of Categories*, 34(7):170–220, 2019.
- [Fré06] Maurice Fréchet. Sur quelques points du calcul fonctionnel. *Rend. Circ. Mat. Palermo*, 22:1–74, 1906.
- [Fri20] Tobias Fritz. A synthetic approach to markov kernels, conditional independence and theorems on sufficient statistics. *Advances in Mathematics*, 370:107239, 2020.
- [FSW⁺23] Jonas Forster, Lutz Schröder, Paul Wild, Harsh Beohar, Sebastian Gurke, Barbara König, and Karla Messing. Graded semantics and graded logics for eilenberg-moore coalgebras. *arXiv preprint arXiv:2307.14826*, 2023.
- [GD23] Francesco Gavazzo and Cecilia Di Florio. Elements of quantitative rewriting. *Proc. ACM Program. Lang.*, 7(POPL), jan 2023.
- [GHK⁺03] Gerhard Gierz, Karl Hofmann, Klaus Keimel, Jimmie Lawson, Michael Mislove, and Dana S. Scott. *Continuous lattices and domains*, volume 93 of *Encycl. Math. Appl.* Cambridge: Cambridge University Press, 2003.
- [Gir82] Michèle Giry. A categorical approach to probability theory. In B. Banaschewski, editor, *Categorical Aspects of Topology and Analysis*, page 68–85, Berlin, Heidelberg, 1982. Springer Berlin Heidelberg.
- [Gir88] Jean-Yves Girard. Normal functors, power series and λ -calculus. *Annals of Pure and Applied Logic*, 37(2):129–177, 1988.
- [GL19] Jean Goubault-Larrecq. Formal ball monads. *Topology and its Applications*, 263:372–391, 2019.
- [GMS⁺23] Sergey Goncharov, Stefan Milius, Lutz Schröder, Stelios Tsampas, and Henning Urbat. Towards a higher-order mathematical operational semantics. *Proc. ACM Program. Lang.*, 7(POPL), jan 2023.
- [GP98] R. Gordon and A.J. Power. Algebraic structure for bicategory enriched categories. *Journal of Pure and Applied Algebra*, 130(2):119–132, 1998.

- [GP18] Richard Garner and John Power. An enriched view on the extended finitary monad–Lawvere theory correspondence. *Logical Methods in Computer Science*, Volume 14, Issue 1, February 2018.
- [GP20] Alexandre Goy and Daniela Petrişan. Combining probabilistic and non-deterministic choice via weak distributive laws. In *Proceedings of the 35th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS '20*, page 454–464, New York, NY, USA, 2020. Association for Computing Machinery.
- [GP21] Guillaume Geoffroy and Paolo Pistone. A partial metric semantics of higher-order types and approximate program transformations. In *CSL 2021 - Computer Science Logic*, Lubjana, Slovenia, January 2021.
- [GPA21] Alexandre Goy, Daniela Petrişan, and Marc Aiguier. Powerset-like monads weakly distribute over themselves in toposes and compact hausdorff spaces. In Nikhil Bansal, Emanuela Merelli, and James Worrell, editors, *48th International Colloquium on Automata, Languages, and Programming (ICALP 2021)*, volume 198 of *Leibniz International Proceedings in Informatics (LIPIcs)*, page 132:1–132:14, Dagstuhl, Germany, 2021. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [GPR16] Mai Gehrke, Daniela Petrişan, and Luca Reggio. The schützenberger product for syntactic spaces. In Ioannis Chatzigiannakis, Michael Mitzenmacher, Yuval Rabani, and Davide Sangiorgi, editors, *43rd International Colloquium on Automata, Languages, and Programming (ICALP 2016)*, volume 55 of *Leibniz International Proceedings in Informatics (LIPIcs)*, page 112:1–112:14, Dagstuhl, Germany, 2016. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [GPZ23] Tao Gu, Robin Piedeleu, and Fabio Zanasi. A Complete Diagrammatic Calculus for Boolean Satisfiability. *Electronic Notes in Theoretical Informatics and Computer Science*, Volume 1 - Proceedings of MFPS XXXVIII, February 2023.
- [Gra75] John W. Gray. Remarks on \mathcal{V} -algebraic theories. In *Deuxième colloque sur l'algèbre des catégories. Amiens-1975. Résumé des conférences*, volume 16, pages 240–243. Dunod éditeur, publié avec le concours du CNRS, 1975.
- [Gra88] Steven K. Graham. Closure properties of a probabilistic domain construction. In M. Main, A. Melton, M. Mislove, and D. Schmidt, editors, *Mathematical Foundations of Programming Language Semantics*, pages 213–233, Berlin, Heidelberg, 1988. Springer Berlin Heidelberg.
- [GS21] Fabio Gadducci and Alexandra Silva, editors. *LIPIcs, Volume 211, CALCO 2021, Complete Volume*, volume 211 of *Leibniz International Proceedings in Informatics (LIPIcs)*, Dagstuhl, Germany, 2021. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [GTWW77] J. A. Goguen, J. W. Thatcher, E. G. Wagner, and J. B. Wright. Initial algebra semantics and continuous algebras. *J. ACM*, 24(1):68–95, jan 1977.
- [Hau14] Felix Hausdorff. *Grundzüge der Mengenlehre*. 1914.
- [HHL22] André Hirschowitz, Tom Hirschowitz, and Ambroise Lafont. Modules over monads and operational semantics (expanded version). *Logical Methods in Computer Science*, Volume 18, Issue 3, August 2022.
- [Hin16] Wataru Hino. Varieties of metric and quantitative algebras, 2016.
- [Hin17] Wataru Hino. Continuous varieties of metric and quantitative algebras, 2017.
- [HP06] Martin Hyland and John Power. Discrete lawvere theories and computational effects. *Theoretical Computer Science*, 366(1):144–162, 2006. Algebra and Coalgebra in Computer Science.
- [HP07] Martin Hyland and John Power. The category theoretic understanding of universal algebra: Lawvere theories and monads. *Electronic Notes in Theoretical Computer Science*, 172:437–458, 2007. Computation, Meaning, and Logic: Articles dedicated to Gordon Plotkin.
- [HPP06] Martin Hyland, Gordon Plotkin, and John Power. Combining effects: Sum and tensor. *Theoretical Computer Science*, 357(1):70–99, 2006. Clifford Lectures and the Mathematical Foundations of Programming Semantics.
- [HR13] Dirk Hofmann and C.D. Reis. Probabilistic metric spaces as enriched categories. *Fuzzy Sets and Systems*, 210:1–21, 2013. Theme : Topology and Algebra.

- [HS00] Pascal Hitzler and Anthony Karel Seda. Dislocated topologies. *J. Electr. Eng*, 51(12):3–7, 2000.
- [HST14] Dirk Hofmann, Gavin J Seal, and Walter Tholen. *Monoidal Topology: A Categorical Approach to Order, Metric, and Topology*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2014.
- [Jac16] Bart Jacobs. *Introduction to Coalgebra: Towards Mathematics of States and Observation*. Cambridge Tracts in Theoretical Computer Science. Cambridge University Press, 2016.
- [JMU24] Jan Jurka, Stefan Milius, and Henning Urbat. Algebraic reasoning over relational structures. *CoRR*, abs/2401.08445, 2024. <https://arxiv.org/abs/2401.08445v2>.
- [JP89] C. Jones and G. Plotkin. A probabilistic powerdomain of evaluations. In *Proceedings of the Fourth Annual Symposium on Logic in Computer Science*, page 186–195. IEEE Press, 1989.
- [KP93] G.M. Kelly and A.J. Power. Adjunctions whose counits are coequalizers, and presentations of finitary enriched monads. *Journal of Pure and Applied Algebra*, 89(1):163–179, 1993.
- [KP22] H. Kunzi and H. Pajoohesh. b-metrics. *Topology and its Applications*, 309:107905, 2022. Remembering Hans-Peter Kunzi (1955-2020).
- [KS18] Bartek Klin and Julian Salamanca. Iterated covariant powerset is not a monad. *Electronic Notes in Theoretical Computer Science*, 341:261–276, 2018. Proceedings of the Thirty-Fourth Conference on the Mathematical Foundations of Programming Semantics (MFPS XXXIV).
- [KTW17] Aleks Kissinger, Sean Tull, and Bas Westerbaan. Picture-perfect quantum key distribution, 2017.
- [KV14] Delia Kesner and Daniel Ventura. Quantitative types for the linear substitution calculus. In Josep Diaz, Ivan Lanese, and Davide Sangiorgi, editors, *Theoretical Computer Science*, pages 296–310, Berlin, Heidelberg, 2014. Springer Berlin Heidelberg.
- [KV17] Alexander Kurz and Jiří Velebil. Quasivarieties and varieties of ordered algebras: regularity and exactness. *Mathematical Structures in Computer Science*, 27(7):1153–1194, 2017.
- [KWG15] Bil Kleb, Bill Wood, and Kevin Godby. `tufte-latex`. L^AT_EX package, 2015. <https://ctan.org/pkg/tufte-latex>.
- [Kwio7] Marta Kwiatkowska. Quantitative verification: models techniques and tools. In *Proceedings of the the 6th Joint Meeting of the European Software Engineering Conference and the ACM SIGSOFT Symposium on The Foundations of Software Engineering, ESEC-FSE '07*, page 449–458, New York, NY, USA, 2007. Association for Computing Machinery.
- [KyKK⁺21] Yuichi Komorida, Shin ya Katsumata, Clemens Kupke, Jurriaan Rot, and Ichiro Hasuo. Expressivity of quantitative modal logics : Categorical foundations via codensity and approximation. In *2021 36th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*, page 1–14, 2021.
- [Lac07] S. Lack. Homotopy-theoretic aspects of 2-monads. *Journal of Homotopy and Related Structures*, 2(2):229–260, 2007.
- [Law62] F. W. Lawvere. The category of probabilistic mappings. *preprint*, 1962.
- [Law63] F. W. Lawvere. Functorial semantics of algebraic theories. *Proc. Natl. Acad. Sci. USA*, 50:869–872, 1963.
- [Lawo2] F William Lawvere. Metric spaces, generalized logic, and closed categories. *Reprints in Theory and Applications of Categories*, 1:1–37, 2002. Originally published in *Rendiconti del seminario matematico e fisico di Milano*, XLIII (1973).
- [Lin66] F. E. J. Linton. Some aspects of equational categories. In S. Eilenberg, D. K. Harrison, S. MacLane, and H. Röhrh, editors, *Proceedings of the Conference on Categorical Algebra*, page 84–94, Berlin, Heidelberg, 1966. Springer Berlin Heidelberg.
- [Lin69] F. E. J. Linton. An outline of functorial semantics. In B. Eckmann, editor, *Seminar on Triples and Categorical Homology Theory*, page 7–52, Berlin, Heidelberg, 1969. Springer Berlin Heidelberg.
- [LJ11] Weisi Lin and C.-C. Jay Kuo. Perceptual visual quality metrics: A survey. *Journal of Visual Communication and Image Representation*, 22(4):297–312, 2011.

- [LMMP13] Jim Laird, Giulio Manzonetto, Guy McCusker, and Michele Pagani. Weighted relational models of typed lambda-calculi. In *Proceedings of the 2013 28th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS '13*, page 301–310, USA, 2013. IEEE Computer Society.
- [LPo8] Stephen Lack and Simona Paoli. 2-nerves for bicategories. *K-Theory: interdisciplinary journal for the development, application and influence of K-theory in the mathematical sciences*, 38(2):153–175, January 2008.
- [LPo9] Stephen Lack and John Power. Gabriel—ulmer duality and lawvere theories enriched over a general base. *Journal of Functional Programming*, 19(3–4):265–286, 2009.
- [LP23] Rory B. B. Lucyshyn-Wright and Jason Parker. Diagrammatic presentations of enriched monads and varieties for a subcategory of arities. *Appl. Categorical Struct.*, 31(5):40, 2023.
- [LR11] Stephen Lack and Jiří Rosický. Notions of Lawvere theory. *Appl. Categ. Structures*, 19(1):363–391, 2011.
- [LR17] Michael Lieberman and Jiří Rosický. Metric abstract elementary classes as accessible categories. *The Journal of Symbolic Logic*, 82(3):1022–1040, 2017.
- [LS12] Stephen Lack and Michael Shulman. Enhanced 2-categories and limits for lax morphisms. *Advances in Mathematics*, 229(1):294–356, 2012.
- [Łuk04] Szymon Łukaszyk. A new concept of probability metric and its applications in approximation of scattered data sets. *Computational Mechanics*, 33:299–304, 2004.
- [LW16] Rory B. B. Lucyshyn-Wright. Enriched algebraic theories and monads for a system of arities. *Theory and Applications of Categories*, 31(5):101–137, 2016. Published 2016-01-31.
- [LY16] Shou Lin and Ziqiu Yun. *Generalized metric spaces and mappings*, volume 6 of *Atlantis Studies in Mathematics*. Atlantis Press, [Paris], chinese edition, 2016. With a foreword by Alexander V. Arhangel'skiĭ, With a preface by Guoshi Gao.
- [Mac71] Saunders Mac Lane. *Categories for the Working Mathematician*. Springer-Verlag, 2nd edition, 1971.
- [Man76] Ernest G. Manes. *Algebraic theories*, volume 26 of *Grad. Texts Math*. Springer, Cham, 1976.
- [Mat94] S. G. Matthews. Partial metric topology. *Annals of the New York Academy of Sciences*, 728(1):183–197, 1994.
- [MB99] Saunders Mac Lane and Garrett Birkhoff. *Algebra*, volume 330. American Mathematical Soc., 1999.
- [Mim20] Samuel Mimram. *PROGRAM = PROOF*. 2020. Available at <https://www.lix.polytechnique.fr/Labo/Samuel.Mimram/teaching/INF551/course.pdf>.
- [MMM12] Annabelle McIver, Larissa Meinicke, and Carroll Morgan. A kantrovich-monadic powerdomain for information hiding, with probability and nondeterminism. In *2012 27th Annual IEEE Symposium on Logic in Computer Science*, pages 461–470, 2012.
- [Mog89] E. Moggi. Computational lambda-calculus and monads. In *Proceedings. Fourth Annual Symposium on Logic in Computer Science*, pages 14,15,16,17,18,19,20,21,22,23, Los Alamitos, CA, USA, jun 1989. IEEE Computer Society.
- [Mog91] Eugenio Moggi. Notions of computation and monads. *Information and Computation*, 93(1):55–92, 1991. Selections from 1989 IEEE Symposium on Logic in Computer Science.
- [MPP17] Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. On the axiomatizability of quantitative algebras. In *32nd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2017, Reykjavik, Iceland, June 20-23, 2017*, page 1–12. IEEE Computer Society, 2017.
- [MPP18] Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. Free complete wasserstein algebras. *Log. Methods Comput. Sci.*, 14(3), 2018.

- [MPP21] Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. Fixed-points for quantitative equational logics. In *36th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2021, Rome, Italy, June 29 - July 2, 2021*, page 1–13. IEEE, 2021.
- [MPP16] Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. Quantitative algebraic reasoning. In Martin Grohe, Eric Koskinen, and Natarajan Shankar, editors, *Proceedings of the 31st Annual ACM/IEEE Symposium on Logic in Computer Science, LICS '16, New York, NY, USA, July 5-8, 2016*, page 700–709. ACM, 2016.
- [MSV21] Matteo Mio, Ralph Sarkis, and Valeria Vignudelli. Combining nondeterminism, probability, and termination: Equational and metric reasoning. In *2021 36th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*, page 1–14, 2021.
- [MSV22] Matteo Mio, Ralph Sarkis, and Valeria Vignudelli. Beyond nonexpansive operations in quantitative algebraic reasoning. In *Proceedings of the 37th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS '22, New York, NY, USA, 2022*. Association for Computing Machinery.
- [MSV23] Matteo Mio, Ralph Sarkis, and Valeria Vignudelli. Universal quantitative algebra for fuzzy relations and generalised metric spaces, July 2023.
- [MU19] Stefan Milius and Henning Urbat. Equational axiomatization of algebras with structure. In Mikolaj Bojanczyk and Alex Simpson, editors, *Foundations of Software Science and Computation Structures - 22nd International Conference, FOSSACS 2019, Held as Part of the European Joint Conferences on Theory and Practice of Software, ETAPS 2019, Prague, Czech Republic, April 6-11, 2019, Proceedings*, volume 11425 of *Lecture Notes in Computer Science*, pages 400–417. Springer, 2019.
- [MV20] Matteo Mio and Valeria Vignudelli. Monads and quantitative equational theories for nondeterminism and probability. In Igor Konnov and Laura Kovács, editors, *31st International Conference on Concurrency Theory, CONCUR 2020, September 1-4, 2020, Vienna, Austria (Virtual Conference)*, volume 171 of *LIPICs*, page 28:1–28:18. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- [Mé11] Facundo Mémoli. Gromov–wasserstein distances and the metric approach to object matching. *Foundations of Computational Mathematics*, 11(4):417–487, Aug 2011.
- [Ngu10] L. Nguyen Van Thé. *Structural Ramsey theory of metric spaces and topological dynamics of isometry groups*. American Mathematical Soc., 2010.
- [NP09] Koki Nishizawa and John Power. Lawvere theories enriched over a general base. *Journal of Pure and Applied Algebra*, 213(3):377–386, 2009.
- [Par22] Jason Parker. Extensivity of categories of relational structures. *Theory and Applications of Categories*, 38(23):898–912, 2022.
- [Par23] Jason Parker. Exponentiability in categories of relational structures. *Theory and Applications of Categories*, 39(16):493–518, 2023.
- [Pis21] Paolo Pistone. On generalized metric spaces for the simply typed lambda-calculus. In *Proceedings of the 36th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS '21, New York, NY, USA, 2021*. Association for Computing Machinery.
- [Pow99] John Power. Enriched lawvere theories. *Theory and Applications of Categories*, 6(7):83–93, 1999.
- [PPo1a] Gordon Plotkin and John Power. Adequacy for algebraic effects. In Furio Honsell and Marino Miculan, editors, *Foundations of Software Science and Computation Structures*, page 1–24, Berlin, Heidelberg, 2001. Springer Berlin Heidelberg.
- [PPo1b] Gordon Plotkin and John Power. Semantics for algebraic operations. In *MFPS 2001. Papers from the 17th conference on the mathematical foundations of programming semantics, Aarhus University, Aarhus, Denmark, May 23–26, 2001.*, pages 332–345. Amsterdam: Elsevier, 2001.
- [PPo2] Gordon Plotkin and John Power. Notions of computation determine monads. In Mogens Nielsen and Uffe Engberg, editors, *Foundations of Software Science and Computation Structures*, pages 342–356, Berlin, Heidelberg, 2002. Springer Berlin Heidelberg.

- [PRSW20] Louis Parlant, Jurriaan Rot, Alexandra Silva, and Bas Westerbaan. Preservation of Equations by Monoidal Monads. In Javier Esparza and Daniel Král', editors, *45th International Symposium on Mathematical Foundations of Computer Science (MFCS 2020)*, volume 170 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 77:1–77:14, Dagstuhl, Germany, 2020. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [PS21] Daniela Petrişan and Ralph Sarkis. Semialgebras and weak distributive laws. In Ana Sokolova, editor, *Proceedings 37th Conference on Mathematical Foundations of Programming Semantics, MFPS 2021, Hybrid: Salzburg, Austria and Online, 30th August - 2nd September, 2021*, volume 351 of *EPTCS*, pages 218–241, 2021.
- [PZ23] Robin Piedeleu and Fabio Zanasi. A Finite Axiomatisation of Finite-State Automata Using String Diagrams. *Logical Methods in Computer Science*, Volume 19, Issue 1, February 2023.
- [QIF20] *The Science of Quantitative Information Flow*. Information Security and Cryptography. Springer Cham, 1 edition, 2020.
- [Rie13] Emily Riehl. Monoidal algebraic model structures. *Journal of Pure and Applied Algebra*, 217(6):1069–1104, 2013.
- [Rie17] Emily Riehl. *Category Theory in Context*. Dover Publications, 2017.
- [Robo2] Edmund Robinson. Variations on algebra: Monadicity and generalisations of equational theories. *Formal Aspects of Computing*, 13(3):308–326, 07 2002.
- [Ros21] Jiří Rosický. Metric monads. *Mathematical Structures in Computer Science*, 31(5):535–552, 2021.
- [Ros24] Jiří Rosický. Discrete equational theories. *Mathematical Structures in Computer Science*, 34(2):147–160, 2024.
- [RPo2] Norman Ramsey and Avi Pfeffer. Stochastic lambda calculus and monads of probability distributions. *SIGPLAN Not.*, 37(1):154–165, jan 2002.
- [RT23] Jiří Rosický and Giacomo Tendas. Enriched universal algebra, 2023.
- [Rut96] J.J.M.M. Rutten. Elements of generalized ultrametric domain theory. *Theoretical Computer Science*, 170(1):349–381, 1996.
- [RZHE24] Aloïs Rosset, Maaïke Zwart, Helle Hvid Hansen, and Jörg Endrullis. Correspondence between composite theories and distributive laws, 2024.
- [SB06] H.R. Sheikh and A.C. Bovik. Image information and visual quality. *IEEE Transactions on Image Processing*, 15(2):430–444, 2006.
- [Sch22a] Todd Schmid. A (co)algebraic framework for ordered processes, 2022.
- [Sch22b] Todd Schmid. Presenting with quantitative inequational theories, 2022.
- [Sco76] Dana Scott. Data types as lattices. *SIAM Journal on Computing*, 5(3):522–587, 1976.
- [SS71] Dana Scott and Christopher Strachey. Toward a mathematical semantics for computer languages. In *Proceedings of the Symposium on Computers and Automata*, volume 21, 1971.
- [Sta17] Sam Staton. Commutative semantics for probabilistic programming. In Hongseok Yang, editor, *Programming Languages and Systems*, pages 855–879, Berlin, Heidelberg, 2017. Springer Berlin Heidelberg.
- [Str72] Ross Street. The formal theory of monads. *Journal of Pure and Applied Algebra*, 2(2):149–168, 1972.
- [SW18] Ana Sokolova and Harald Woracek. Termination in convex sets of distributions. *Logical Methods in Computer Science*, Volume 14, Issue 4, November 2018.
- [Tho12] Walter Tholen. Kleisli enriched. *Journal of Pure and Applied Algebra*, 216(8):1920–1931, 2012. Special Issue devoted to the International Conference in Category Theory ‘CT2010’.
- [TP97] Daniele Turi and Gordon D. Plotkin. Towards a mathematical operational semantics. In *Proceedings of the Twelfth Annual IEEE Symposium on Logic in Computer Science (LICS 1997)*, pages 280–291. IEEE Computer Society Press, June 1997.

- [vB01] Franck van Breugel. An introduction to metric semantics: operational and denotational models for programming and specification languages. *Theoretical Computer Science*, 258(1):1–98, 2001.
- [vB05] Franck van Breugel. The metric monad for probabilistic nondeterminism. Available at <http://www.cse.yorku.ca/~franck/research/drafts/monad.pdf>, 2005.
- [vBW01] Franck van Breugel and James Worrell. Towards quantitative verification of probabilistic transition systems. In Fernando Orejas, Paul G. Spirakis, and Jan van Leeuwen, editors, *Automata, Languages and Programming*, page 421–432, Berlin, Heidelberg, 2001. Springer Berlin Heidelberg.
- [Vig23] Ignacio Viglizzo. Basic constructions in the categories of sets, sets with a binary relation on them, preorders, and posets, June 2023.
- [Vil09] Cédric Villani. *Optimal transport: old and new*, volume 338 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 2009. Old and new.
- [VK11] Jiří Velebil and Alexander Kurz. Equational presentations of functors and monads. *Mathematical Structures in Computer Science*, 21(2):363–381, 2011.
- [vN51] John von Neumann. Various techniques used in connection with random digits. In G. E. Forsythe, editor, *National Bureau of Standards Applied Math Series*, volume 12, pages 36–38. National Bureau of Standards, 1951. Reprinted in von Neumann’s *Collected Works*, Volume 5, Pergamon Press, 1963, pp. 768–770.
- [VW06] Daniele Varacca and Glynn Winskel. Distributing probability over non-determinism. *Mathematical Structures in Computer Science*, 16(1):87–113, 2006.
- [Wea93] Nikolai Weaver. Generalized varieties. *Algebra Universalis*, 30(1):27–52, 03 1993.
- [Wea95] Nikolai Weaver. Quasi-varieties of metric algebras. *Algebra Universalis*, 33(1):1–9, 03 1995.
- [Wec92] Wolfgang Wechler. *Universal algebra for computer scientists*, volume 25 of *EATCS Monogr. Theoret. Comput. Sci.* Berlin etc.: Springer-Verlag, 1992.
- [Wil31a] Wallace Alvin Wilson. On quasi-metric spaces. *American Journal of Mathematics*, 53(3):675–684, 1931.
- [Wil31b] Wallace Alvin Wilson. On semi-metric spaces. *American Journal of Mathematics*, 53(2):361–373, 1931.
- [WS20] Paul Wild and Lutz Schröder. Characteristic logics for behavioural metrics via fuzzy lax extensions. In Igor Konnov and Laura Kovács, editors, *31st International Conference on Concurrency Theory (CONCUR 2020)*, volume 171 of *Leibniz International Proceedings in Informatics (LIPIcs)*, page 27:1–27:23, Dagstuhl, Germany, 2020. Schloss Dagstuhl – Leibniz-Zentrum für Informatik.
- [ZK22] Linpeng Zhang and Benjamin Lucien Kaminski. Quantitative strongest post: a calculus for reasoning about the flow of quantitative information. *Proc. ACM Program. Lang.*, 6(OOPSLA1), apr 2022.
- [ZM22] Maaïke Zwart and Dan Marsden. No-Go Theorems for Distributive Laws. *Logical Methods in Computer Science*, Volume 18, Issue 1, January 2022.